

ASYMPTOTICS OF PARTIAL DENSITY FUNCTION VANISHING ALONG SMOOTH SUBVARIETY

JINGZHOU SUN

ABSTRACT. We study the asymptotic of the partial density function associated to holomorphic section of a positive line bundle vanishing to high orders along a fixed smooth subvariety. Assuming local torus-action-invariance, we describe the forbidden region, generalizing the result of Ross-Singer.

CONTENTS

1. Introduction	1
2. One dimensional model	3
3. proof of the main theorem	5
References	8

1. INTRODUCTION

Let (X, L) be a polarized compact Kähler manifold of dimension m . We endow L with a Hermitian metric h with positive curvature. And we use $\omega = \frac{i}{2}\Theta_h$ as the Kähler form. By abuse of notation, we still use h to denote the induced metric on the k -th power L^k . Then we have a Hermitian inner product on $H^0(X, L^k)$, defined by

$$\langle s_1, s_2 \rangle = \int_X h(s_1, s_2) \frac{\omega^m}{m!}$$

Let $\{s_i\}$ be an orthonormal basis of $\mathcal{H}_k = H^0(X, L^k)$. Then the on-diagonal Bergman kernel, also called density of states function, is defined as

$$\rho_k(z) = \sum |s_i(z)|_h^2.$$

The on-diagonal Bergman kernel ρ_k has very nice asymptotic expansion by the results of Tian, Zelditch, Lu, etc. [29, 30, 14, 6, 15]. The asymptotic expansion has been found very useful, making ρ_k an important bridge connecting Kähler geometry to algebraic geometry (see for example [8, 10]).

Let $V \subset X$ be a closed submanifold of dimension n . Let $\delta > 0$ be a small number. We denote by $\mathcal{H}_k^{\delta k} \subset \mathcal{H}_k$ the subspace consisting of holomorphic sections whose vanishing orders

Date: October 13, 2020.

The author is partially supported by NNSF of China no.11701353 and the STU Scientific Research Foundation for Talents no.130/760181.

along V are no less than δk . The partial density function ρ_k^δ is then defined as the density of states function for $\mathcal{H}_k^{\delta k}$:

$$\rho_k^\delta(z) = \sum |s_i(z)|_h^2,$$

where $\{s_i\}$ is an orthonormal basis of $\mathcal{H}_k^{\delta k}$. When V is a divisor, the partial density functions have been studied by Zelditch-Zhou [31, 32], Ross-Singer[16] and Coman-Marinescu [7] etc. It was first shown by Shiffman-Zelditch [22] in the toric case, that there is a ‘forbidden region’ in which the partial density function is exponentially small. Berman [1] showed that there is an open forbidden region R that asymptotically the partial density function is exponentially small on compact subsets and is equal to the usual density function on compact subsets of the complement of \bar{R} . In particular, Ross-Singer[16] studied the case when V is smooth divisor, plus the condition that the line bundle and the metric is S^1 -symmetric in a neighborhood U of V . Let $\mu : U \rightarrow \mathbb{R}$ be the Hamiltonian of the S^1 -action, normalized so that $\mu^{-1}(0) = V$. Then Ross-Singer proved a distributional asymptotic expansion for the partial density function and showed that the forbidden region is $\mu^{-1}[0, \delta]$:

Theorem 1.1 ([16]). *For sufficiently small δ , we have*

$$\rho_k^\delta \sim \begin{cases} O(k^{-\infty}) & \text{on } \mu^{-1}[0, \delta] \\ \rho_k + O(k^{-\infty}) & \text{on } X \setminus \mu^{-1}[0, \delta] \end{cases},$$

where \sim means the equality holds on any given compact subset of $\mu^{-1}[0, \delta]$ and $X \setminus \mu^{-1}[0, \delta]$ respectively.

The term $k^{-\infty}$ in the theorem means a quantity depending on k that is $O(k^{-m})$ as $k \rightarrow \infty$, for all $m \geq 0$. We will adopt the notation used by Donaldson and denote $k^{-\infty}$ by $\varepsilon(k)$. So, in order to avoid confusion, we use δ for the partial density function instead of ε used in [16].

In this article, we generalize Ross-Singer’s result to the case when V is a smooth subvariety. Let $T^{m-n} = (S^1)^{m-n}$ denote the $(m-n)$ -dimensional torus. We assume that there is a neighborhood U of V that admits a holomorphic T^{m-n} -action. We also assume the line bundle L and the metric h , hence ω , are invariant under the T^{m-n} -action.

Let $\mu : U \rightarrow \mathbb{R}^{m-n}$ be the moment map of the torus action, normalized so that $\mu(V) = 0$. We denote by μ_i the i -th component of μ . Then $\nu = \sum_{i=1}^{m-n} \mu_i : U \rightarrow \mathbb{R}$ is the moment map for the diagonal S^1 -action. Our main result is that the forbidden region is $\nu^{-1}([0, \delta])$:

Theorem 1.2. *For sufficiently small δ , we have*

$$\rho_k^\delta \sim \begin{cases} \varepsilon(k) & \text{on } \nu^{-1}[0, \delta] \\ \rho_k + \varepsilon(k) & \text{on } X \setminus \nu^{-1}[0, \delta] \end{cases}$$

The proof uses the techniques we developed in [25, 28, 26]. The idea of the proof is similar to that in [16]. Unfortunately, we have not been able to give an asymptotic expansion for the renormalized partial Bergman kernel as Ross-Singer did in [16]. It is also very interesting to know if there is still a forbidden region and what the boundary is, if the assumption of the torus-invariance is removed.

We would like to compare the partial density function to the logarithmic Bergman kernel studied by the author in [27], for which the vanishing order is ≥ 1 instead of $\geq \delta k$. In [27], we showed that the asymptotics near the subvariety depend on the distance function to the

subvariety. When δ is small enough, the moment function ν is comparable with the distance function squared. One can see the similarity between them.

The asymptotics of off-diagonal Bergman kernel have been extensively used in the value distribution theory of sections of line bundles by Shiffman-Zelditch and others[2, 3, 4, 5, 18, 19, 20, 21, 22, 23, 24, 11, 12, 13]. It would be very interesting and important to study the asymptotics of the partial off-diagonal Bergman kernel vanishing along a general subvariety.

The structure of this article is as follows. We will first prove the key estimates on the complex plane. Then we apply the key lemma to the neighborhood U . Then with the help of the Hörmander's technique, we show that what we get on U goes directly to X , hence proving the main theorem.

Acknowledgements. The author would also like to thank Professor Song Sun and Professor Chengjie Yu for many very helpful discussions.

2. ONE DIMENSIONAL MODEL

Let $\omega = f\sqrt{-1}dz \wedge d\bar{z}$ be a smooth Kähler form on \mathbb{C} invariant under the natural S^1 -action. Let $\varphi(|z|)$ be a smooth potential of ω . By a linear change of coordinates, we can assume that $\varphi = |z|^2 + O(|z|^4)$. Let $\mu : \mathbb{C} \rightarrow \mathbb{R}$ be the moment map of the S^1 -action normalized so that $\mu(0) = 0$. We will use the notations $r = |z|$, $x = |z|^2$ and $t = -\log x$. So $\varphi_{tt} = x\Delta\varphi = xf$. Then

$$\mu = \int_0^x f dx.$$

And since $\lim_{t \rightarrow \infty} \varphi_t = 0$, we have

$$\begin{aligned} \varphi_t &= - \int_t^\infty \varphi_{tt} dt \\ &= - \int_x^0 f dx \\ &= -\mu \end{aligned}$$

Let $a \geq 0$ be an integer, $f_1(|z|) > 0$ be a smooth function, we consider the k -th norm of the holomorphic function z^a :

$$I_a = \int_{\mathbb{C}} e^{-k\varphi} |z|^{2a} f_1 \sqrt{-1} dz \wedge d\bar{z} = 2\pi \int_{-\infty}^\infty e^{-k\varphi - (a+1)t + \log f_1} dt.$$

Denote by $g_a(t) = -k\varphi - (a+1)t + \log f_1$. Then $g'_a = -k\varphi_t - (a+1) + (\log f_1)_t$ and $g''_a = -k\varphi_{tt} + (\log f_1)_{tt}$. Since $f_1 = A + O(|z|^2)$, $A > 0$, we have $(\log f_1)_t = O(\varphi_t)$ and $(\log f_1)_{tt} = O(\varphi_{tt})$. So for k large enough, g_a is a concave function of t which attains its only maximum at t_a satisfying

$$\varphi_t(t_a) = -\frac{a+1}{k} + \frac{(\log f_1)_t(t_a)}{k}.$$

Namely,

$$\mu(t_a) = \frac{a+1}{k} + O\left(\frac{x_a}{k}\right),$$

when $x_a < 1$.

We recall the following basic lemma used in [25]:

Lemma 2.1. *Let $f(x)$ be a concave function. Suppose $f'(x_0) < 0$, then we have*

$$\int_{x_0}^{\infty} e^{f(x)} dx \leq \frac{e^{f(x_0)}}{-f'(x_0)}$$

Let $0 < R_1 < R < 1$ be two fixed numbers. When $\mu(t_a) > R$, we have $a > Rk - C$ for some C independent of k , then $g'_a < 0$ when $\mu \leq R$. Let $\mu(t_R) = R$ and $\mu(t_{R_1}) = R_1$, then we let $t_m = \frac{t_R + t_{R_1}}{2}$. Clearly, $|g'_a(t_m)| > C_1 k$ for some $C_1 > 0$ depending on R and R_1 but independent of k and a . By the concavity of g_a , we have

$$e^{g_a(t_m) - g_a(t_{R_1})} = \varepsilon(k)$$

Therefore, by lemma 2.1, we have

$$\int_{\mu < R_1} e^{g_a(t)} dt = \varepsilon(k) \int_{R_1 \leq \mu \leq R} e^{g_a(t)} dt.$$

When $\mu(t_a) \leq R$, depending on a , there are two cases .

When $a \geq \sqrt{k}$, $g''_a(t_a)$ is large, so we can use Laplace's method to estimate the integral, namely

$$I_a \approx \frac{(2\pi)^{3/2}}{\sqrt{g''_a(t_a)}}.$$

More importantly, the mass of the measure $e^{g_a(t)} dt$ is concentrated in a small neighborhood of t_a . More precisely, if $k\varphi_{tt}(t_a)\delta_a^2 > 2\log^2 k$, then by lemma 2.1, we have

$$I_a = (1 + \varepsilon(k)) \int_{t_a - \delta_a}^{t_a + \delta_a} e^{g_a(t)} dt.$$

It is easy to see that there is a constant $C_2 > 0$ independent of k such that if $\delta_a > C_2 \frac{\log k}{\sqrt{a}}$, then the condition $k\varphi_{tt}(t_a)\delta_a^2 > 2\log^2 k$ is satisfied.

When $a < \sqrt{k}$, the term $k\varphi_{tt}(t_a)$ may be too small to use Laplace's method. We have the following

$$\begin{aligned} \int_{x < k^{-1/4}} e^{g_a(t)} dt &\geq e^{-(a+1)} e^{-(a+1)e} e^{g_a(t_a)} \\ &\geq e^{-(a+1)} e^{-(a+1)e} e^{-(a+1) + a \log \frac{a+1}{k}} \\ &\geq e^{-5(a+1)} \left(\frac{a+1}{k}\right)^a \end{aligned}$$

When $x = k^{-1/4}$, $\frac{e^{g_a(\log k^{1/4})}}{e^{-5(a+1)} \left(\frac{a+1}{k}\right)^a} = \varepsilon(k)$. So by lemma 2.1, we have for $a < \sqrt{k}$,

$$\int_{\mu \geq 2k^{-1/4}} e^{g_a(t)} dt \leq \int_{x \geq k^{-1/4}} e^{g_a(t)} dt = \varepsilon(k) I_a$$

In summary, we have proved the following

Lemma 2.2. *For fixed $0 < R_1 < R < 1$, there exist constants $C, C_1 > 0$ independent of k , such that for k large enough we have the following estimations:*

1) If $a > Rk - C$,

$$\int_{\mu < R_1} e^{g_a(t)} dt = \varepsilon(k) \int_{R_1 \leq \mu \leq R} e^{g_a(t)} dt,$$

where the $\varepsilon(k)$ term is independent of a .

2) If $\sqrt{k} \leq a \leq Rk - C$,

$$\int_{|\mu - \frac{a}{k}| \geq C_1 \frac{\sqrt{a} \log k}{k}} e^{g_a(t)} dt = \varepsilon(k) \int_{|\mu - \frac{a}{k}| < C_1 \frac{\sqrt{a} \log k}{k}} e^{g_a(t)} dt,$$

where the $\varepsilon(k)$ term is independent of a .

3) If $a < \sqrt{k}$,

$$\int_{\mu \geq 2k^{-1/4}} e^{g_a(t)} dt = \varepsilon(k) \int_{\mu < 2k^{-1/4}} e^{g_a(t)} dt,$$

where the $\varepsilon(k)$ term is independent of a .

3. PROOF OF THE MAIN THEOREM

By assumption, there is a neighborhood U of V that is covered with charts that admit standard coordinates (z, w) so that $V = \{z = 0\}$, where $z = (z_1, \dots, z_{m-n}) \in \mathbb{C}^{m-n}$ and $w = (w_1, \dots, w_n) \in \mathbb{C}^n$, and

$$\lambda \cdot (z, w) = (\lambda z, w),$$

for $\lambda = (\lambda_1, \dots, \lambda_{m-n})$, where $\lambda z = (\lambda_1 z_1, \dots, \lambda_{m-n} z_{m-n})$. We have the following lemma,

Lemma 3.1. *There is a T^{m-n} -invariant biholomorphism Ψ from the neighborhood U to an open neighborhood of V in the total space of the vector bundle $L_1 \oplus L_2 \cdots \oplus L_{m-n}$, where each L_i is a holomorphic line bundle.*

The proof is very similar to that of lemma 5.3 in [17], and for the convenience of the reader, we include a proof here.

Proof. Let (z_α, w_α) and (z_β, w_β) be two sets of standard coordinates. Then the transition functions between them are necessarily of the form

$$z_\beta = \lambda_{\alpha\beta}(z_\alpha, w_\alpha) z_\alpha, \quad w_\beta = \tau_{\alpha\beta}(z_\alpha, w_\alpha)$$

where $\lambda_{\alpha\beta}$ and $\tau_{\alpha\beta}$ are holomorphic and take values in $\mathbb{C}^{*(m-n)}$ and $\mathbb{C}^{(m-n)}$ respectively. We understand the torus T^{m-n} -action on the w -part is trivial. Then both $\lambda_{\alpha\beta}$ and $\tau_{\alpha\beta}$ are T^{m-n} -invariant, so their dependence on z_α are trivial. So the transition functions

$$z_\beta = \lambda_{\alpha\beta}(w_\alpha) z_\alpha$$

defines a holomorphic vector bundle E of rank $m - n$ over V , with structure group T^{m-n} . Therefore, E decomposes as $E = L_1 \oplus L_2 \cdots \oplus L_{m-n}$, and U is biholomorphic to an open neighborhood of V in E . Clearly, the biholomorphism is T^{m-n} -invariant. \square

We consider U as an open set in $\oplus L_i$ and denote by $\pi : U \rightarrow V$ the projection. Then for A small enough, U contains the poly-disc bundle $U_A = \cap_{1 \leq i \leq m-n} \mu_i^{-1}(x < A)$. For each $p \in V$, the fiber of U_A is an open poly-disc D_A in \mathbb{C}^{m-n} . By the torus-invariance, the restriction φ_p of φ depends only on $|z_1|, \dots, |z_{m-n}|$, so does the restriction ω_p of ω . So the monomials $z^a = \prod z_i^{a_i}$ are all orthogonal to each other on D_p , where $a = (a_1, \dots, a_{m-n})$ is an $(m-n)$ -tuple of non-negative integers.

Lemma 3.2. *Let $W_\delta = \{q \in D_p | \nu(q) < \delta - k^{1/4}\}$. When δ is small enough, for each holomorphic function $f = \sum_{\sum a_i \geq \delta k} c_a z^a$, we have*

$$\int_{W_\delta} |f|^2 e^{-k\varphi_p} \omega_p^{m-n} = \varepsilon(k) \int_{D_p} |f|^2 e^{-k\varphi_p} \omega_p^{m-n}$$

Proof. By orthogonality, $\int |f|^2 e^{-k\varphi_p} \omega_p^{m-n} = \sum_{\sum a_i \geq \delta k} |c_a|^2 \int |z^a|^2 e^{-k\varphi_p} \omega_p^{m-n}$. So it suffices to prove the lemma for the monomials z^a satisfying $\sum a_i \geq \delta k$.

By making A a little smaller, a compactness argument shows that the constants C and C_1 in lemma 2.2 can be chosen to be the same for all z_i -discs. For each point p in the open set, at least one $1 \leq i \leq m-n$ satisfies the condition that $\mu_i < \frac{a_i}{k} - k^{-3/8}$. Therefore, lemma 2.2 shows that the mass of z^a in W is $\varepsilon(k)$ relative to the mass in $D_A \setminus W_\delta$. And the theorem is proved □

As a corollary, we can prove the part about the forbidden region of theorem 1.2.

Proof of the first half of theorem 1.2. We can take an arbitrary orthonormal basis $\{s_i\}$ for the space $\mathcal{H}_k^{\delta k}$. Then the lemma above shows that each s_i has L^2 -mass of the size $\varepsilon(k)$ in the region W_δ . We recall that for each point $p \in X$, there is a unit section s_p , called the peak section, satisfying $|s_p(p)|_h^2 = \rho_k(p)$ and $s_p \perp s$ as long as $s(p) = 0$.

We take the inner product of s_i with the peak sections s_p for each $p \in W$

$$|s_i(p)| = |\langle s_i, s_p \rangle| \leq \sqrt{\rho_k(p)} \leq \|s_i\| \sqrt{\rho_k(p)}$$

we see that the point-wise norm $|s_i(p)|$ is also $\varepsilon(k)$. Since the dimension of $\mathcal{H}_k^{\delta k}$ is $O(k^m)$. We get the conclusion. □

We denote by T_i the i -th component of T^{m-n} . For each L_i , we fix a T_i -invariant section s_i such that $\mu_i^{-1}(0) = \{s_i = 0\}$. So we can choose local frame e_i of L_i , so that $s_i = z_i e_i$ in the standard torus-invariance coordinates.

For each $s \in H^0(V, kL - \otimes L_i^{a_i})$, where $a_i \geq 0$, $\pi^* s \in H^0(U, kL - \otimes L_i^{a_i})$, so $\tilde{s} = \pi^* s \otimes \prod s_i^{a_i} \in H^0(U, L^k)$. We denote by G_δ the span of all such sections for $\sum_{i=1}^{m-n} a_i < \delta k$. Then clearly, for each section $\beta \in \mathcal{H}_k^{\delta k}$, we have $\beta|_U \perp G_\delta$. Moreover, since each \tilde{s} is represented by a holomorphic function of the form

$$\sum_{\sum_{i=1}^{m-n} a_i < \delta k} f_a(w) z^a,$$

by lemma 2.2, each section $\tilde{s} \in G_\delta$ has its mass concentrated within the set $Y_\delta = \{q | \nu(q) \leq \delta + 2(m-n)k^{-1/4}\}$ and decays fast as ν gets bigger. More precisely, the measure outside Y_δ is $\varepsilon(k)$ relative to that inside Y_δ . To connect G_δ to $(\mathcal{H}_k^{\delta k})^\perp$, we need to use Hörmander's L^2 technique. The following lemma is well-known, see for example [29].

Lemma 3.3. *Suppose (M, g) is a complete Kähler manifold of complex dimension n , \mathcal{L} is a line bundle on M with hermitian metric h . If*

$$\langle -2\pi i\Theta_h + Ric(g), v \wedge \bar{v} \rangle_g \geq C|v|_g^2$$

for any tangent vector v of type $(1, 0)$ at any point of M , where $C > 0$ is a constant and Θ_h is the curvature form of h . Then for any smooth \mathcal{L} -valued $(0, 1)$ -form α on M with $\bar{\partial}\alpha = 0$ and $\int_M |\alpha|^2 dV_g$ finite, there exists a smooth \mathcal{L} -valued function β on M such that $\bar{\partial}\beta = \alpha$ and

$$\int_M |\beta|^2 dV_g \leq \frac{1}{C} |\alpha|^2 dV_g$$

where dV_g is the volume form of g and the norms are induced by h and g .

Now let $\chi \in C_0^\infty(U)$ be a non-negative cut-off function, that equals 1 in a neighborhood of the subset $\{p \in U | \nu(p) \leq \delta + c\}$ for some small $c > 0$. Then for each $\tilde{s} \in G_\delta$ with $\|\tilde{s}\|^2 = 1$, we take the orthogonal projection of $\chi\tilde{s}$ onto \mathcal{H}_k . The way to do so is to solve the $\bar{\partial}$ -equation

$$\bar{\partial}v = \bar{\partial}\chi \otimes \tilde{s}.$$

And since $|\tilde{s}|$ is $\varepsilon(k)$ on the place where $\bar{\partial}\chi \neq 0$, we can apply the lemma above to see that the global holomorphic section $\chi\tilde{s} - v \in \mathcal{H}_k$ has $\varepsilon(k)$ L^2 -mass outside U and the L^2 -mass of $\chi\tilde{s} - v - \tilde{s}$ within U is also $\varepsilon(k)$. So for each $f \in \mathcal{H}_k^{\delta k}$ with $\|f\|^2 = 1$, we have

$$|\langle f, \chi\tilde{s} - v \rangle| = \varepsilon(k)$$

We then take the orthogonal projection of $\chi\tilde{s} - v$ onto $(\mathcal{H}_k^{\delta k})^\perp$ and denote by \bar{s} the section obtained. So we have

$$\int_U |\bar{s} - \tilde{s}|^2 e^{-k\varphi} \omega^m = \varepsilon(k).$$

We denote by J the map we just constructed from G_δ to $(\mathcal{H}_k^{\delta k})^\perp$. So J is almost an isometry, namely

$$\|J(\tilde{s})\|^2 = (1 + \varepsilon(k)) \|\tilde{s}\|^2.$$

Therefore, if we start with an orthonormal basis of G_δ , we obtain an almost orthonormal basis of $(\mathcal{H}_k^{\delta k})^\perp$. And then after a Gram-Schmidt process, we get an orthonormal basis. We denote by ρ_G the density function of G_δ , considered as a function on X by extension by zero. Since the density function of $(\mathcal{H}_k^{\delta k})^\perp$ is $\rho_k - \rho_k^\delta$, we have

$$\int_X (\rho_k - \rho_k^\delta - \rho_G) \omega^m = \varepsilon(k).$$

Since, ρ_G has $\varepsilon(k)$ mass outside Y_δ , we have

$$\int_{X \setminus Y_\delta} (\rho_k - \rho_k^\delta) \omega^m = \varepsilon(k).$$

Recall that by a local construction of the peak section s_p for any $p \in X$ (see for example [9]), $|s_p|$ decays exponentially away from p . More precisely, $|s_p(q)| = \varepsilon(k)$, when $d(p, q) > \frac{\log k}{\sqrt{k}}$. When δ is small enough, ν is comparable with the distance function $d^2(p, q)$, so if we denote by $Y'_\delta = \{q | \nu(q) \leq \delta + 3(m - n)k^{-1/4}\}$, then for any unit $s \in (\mathcal{H}_k^{\delta k})^\perp$, we have

$$|\langle s, s_p \rangle| = \varepsilon(k),$$

for $p \in X \setminus Y'_\delta$. Therefore $|s(p)| = \varepsilon(k)$, $p \in X \setminus Y'_\delta$. Since $\lim_{k \rightarrow \infty} Y'_\delta = \{q | \nu(q) \leq \delta\}$, this implies the second part of theorem 1.2.

REFERENCES

- [1] Robert Berman. Bergman kernels and equilibrium measures for ample line bundles. *arXiv: Complex Variables*, 2007.
- [2] Pavel Bleher, Bernard Shiffman, and Steve Zelditch. Poincaré-Lelong approach to universality and scaling of correlations between zeros. *Comm. Math. Phys.*, 208(3):771–785, 2000.
- [3] Pavel Bleher, Bernard Shiffman, and Steve Zelditch. Universality and scaling of correlations between zeros on complex manifolds. *Invent. Math.*, 142(2):351–395, 2000.
- [4] Pavel Bleher, Bernard Shiffman, and Steve Zelditch. Correlations between zeros and supersymmetry. *Comm. Math. Phys.*, 224(1):255–269, 2001.
- [5] Pavel Bleher, Bernard Shiffman, and Steve Zelditch. Universality and scaling of zeros on symplectic manifolds. In *Random matrix models and their applications*, volume 40 of *Math. Sci. Res. Inst. Publ.*, pages 31–69. Cambridge Univ. Press, Cambridge, 2001.
- [6] David Catlin. The Bergman kernel and a theorem of Tian. In *Analysis and geometry in several complex variables (Katata, 1997)*, Trends Math., pages 1–23. Birkhäuser Boston, Boston, MA, 1999.
- [7] Dan Coman and George Marinescu. On the first order asymptotics of partial bergman kernels. *Annales de la Faculté des sciences de Toulouse : Mathématiques*, Ser. 6, 26(5):1193–1210, 2017.
- [8] Simon Donaldson. Scalar curvature and projective embeddings, I. *Journal of Differential Geometry*, 59(3):479–522, 2001.
- [9] Simon Donaldson. Algebraic families of constant scalar curvature Kähler metrics. In *Surveys in differential geometry 2014. Regularity and evolution of nonlinear equations*, volume 19 of *Surv. Differ. Geom.*, pages 111–137. Int. Press, Somerville, MA, 2015.
- [10] Simon Donaldson and Song Sun. Gromov-Hausdorff limits of Kähler manifolds and algebraic geometry, II. *J. Differential Geom.*, 107(2):327–371, 2017.
- [11] Michael R. Douglas, Bernard Shiffman, and Steve Zelditch. Critical points and supersymmetric vacua. I. *Comm. Math. Phys.*, 252(1-3):325–358, 2004.
- [12] Michael R. Douglas, Bernard Shiffman, and Steve Zelditch. Critical points and supersymmetric vacua. II. Asymptotics and extremal metrics. *J. Differential Geom.*, 72(3):381–427, 2006.
- [13] Michael R. Douglas, Bernard Shiffman, and Steve Zelditch. Critical points and supersymmetric vacua. III. String/M models. *Comm. Math. Phys.*, 265(3):617–671, 2006.
- [14] Zhiqin Lu. On the lower order terms of the asymptotic expansion of Tian-Yau-Zelditch. *Amer.J.math*, 122(2):235–273, 2000.
- [15] Xiaonan Ma and George Marinescu. *Holomorphic Morse inequalities and Bergman kernels*, volume 254 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2007.
- [16] Julius Ross and Michael Singer. Asymptotics of partial density functions for divisors. *The Journal of Geometric Analysis*, 27(3):1803–1854, Jul 2017.
- [17] Julius Ross and Michael Singer. Asymptotics of partial density functions for divisors. *J. Geom. Anal.*, 27(3):1803–1854, 2017.
- [18] Bernard Shiffman. Convergence of random zeros on complex manifolds. *Sci. China Ser. A*, 51(4):707–720, 2008.
- [19] Bernard Shiffman and Steve Zelditch. Distribution of zeros of random and quantum chaotic sections of positive line bundles. *Comm. Math. Phys.*, 200(3):661–683, 1999.
- [20] Bernard Shiffman and Steve Zelditch. Asymptotics of almost holomorphic sections of ample line bundles on symplectic manifolds. *J. Reine Angew. Math.*, 544:181–222, 2002.
- [21] Bernard Shiffman and Steve Zelditch. Random polynomials of high degree and Levy concentration of measure. *Asian J. Math.*, 7(4):627–646, 2003.
- [22] Bernard Shiffman and Steve Zelditch. Random polynomials with prescribed Newton polytope. *J. Amer. Math. Soc.*, 17(1):49–108, 2004.
- [23] Bernard Shiffman, Steve Zelditch, and Qi Zhong. Random zeros of complex manifolds: conditional expectations. *J. Inst. Math. Jussieu*, 10(3):753–783, 2011.

- [24] Bernard Shiffman, Steve Zelditch, and Scott Zrebiec. Overcrowding and hole probabilities for random zeros on complex manifolds. *Indiana Univ. Math. J.*, 57(5):1977–1997, 2008.
- [25] J. Sun and S. Sun. Projective embedding of log Riemann surfaces and K-stability. *to appear in J Geom Anal*, May 2020.
- [26] Jingzhou Sun. Estimations of the Bergman kernel of the punctured disk. *ArXiv e-prints*, 2017.
- [27] Jingzhou Sun. Logarithmic bergman kernel and conditional expectation of gaussian holomorphic fields. *arXiv: Complex Variables*, 2019.
- [28] Jingzhou Sun. Projective embedding of pairs and logarithmic K-stability. *Math. Ann.*, 375(3-4):1307–1336, 2019.
- [29] Gang Tian. On a set of polarized Kähler metrics on algebraic manifolds. *Journal of Differential Geometry*, 32(1990):99–130, 1990.
- [30] Steve Zelditch. Szego kernels and a theorem of Tian. *International Mathematics Research Notices*, 6(6):317–331, 2000.
- [31] Steve Zelditch and Peng Zhou. Interface asymptotics of partial bergman kernels on s^1 -symmetric kaehler manifolds. *math arxiv*, arXiv:1604.06655.
- [32] Steve Zelditch and Peng Zhou. Central limit theorem for spectral partial Bergman kernels. *Geom. Topol.*, 23(4):1961–2004, 2019.

DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU CITY, GUANGDONG 515063, CHINA
Email address: jzsun@stu.edu.cn