

SIGN STABILITY OF MAPPING CLASSES ON MARKED SURFACES I: EMPTY BOUNDARY CASE

TSUKASA ISHIBASHI AND SHUNSUKE KANO

ABSTRACT. For a mapping class on a punctured surface, we prove that the pseudo-Anosov property is equivalent to the sign stability introduced in [IK21]. In particular we conclude that the algebraic entropies of the cluster \mathcal{A} - and \mathcal{X} -transformations induced by a mutation loop given by a pseudo-Anosov mapping class both coincide with its topological entropy.

CONTENTS

1. Introduction	2
1.1. Sign stability and the pseudo-Anosov property	4
1.2. Sign stability of Dehn twists	4
1.3. The topological and algebraic entropies	5
Organization of the paper	5
Acknowledgements	6
2. Laminations on a marked surface	6
2.1. The space of \mathcal{A} -laminations	7
2.2. The space of \mathcal{X} -laminations	11
2.3. Ensemble map and the space of \mathcal{U} -laminations	15
Exchange matrix	15
2.4. Measured foliations	17
2.5. Measured geodesic laminations	22
3. Mutation loops and their representation paths in the exchange graph	27
3.1. The labeled exchange graph	27
3.2. Mutation loops and the cluster modular group	31
4. Measured laminations from the cluster algebraic point of view	35
4.1. The cluster ensemble associated with a marked surface	35
4.2. The cluster modular group associated with a marked surface	39
5. Sign stability of mutation loops	42
5.1. PL maps and their presentation matrices	43
5.2. Sign stability	44
5.3. Uniform sign stability	49

5.4. Relation between the uniform sign stability and the cluster-pseudo-Anosov property	51
6. Sign stability of Dehn twists	51
7. Sign stability of pseudo-Anosov mapping classes: empty boundary case	54
7.1. Sign stability and the pA property	55
7.2. NS dynamics on the space of \mathcal{X} -laminations and the pA property	57
8. Topological and algebraic entropies	64
8.1. Topological entropy	64
8.2. Algebraic entropy	65
8.3. Comparison of topological and algebraic entropies	67
Appendix A. Review on the cluster ensembles	67
A.1. Seed patterns	68
A.2. Cluster varieties	69
A.3. Tropicalizations of the cluster varieties	73
References	74

1. INTRODUCTION

The *mapping class group* of an oriented closed surface Σ is the group of isotopy classes of orientation-preserving diffeomorphisms on Σ . It naturally acts on the Teichmüller space $T(\Sigma)$ of Σ properly discontinuously, and the quotient orbifold is the celebrated moduli space of Riemann surfaces. This action extends continuously to the Thurston compactification of the Teichmüller space, which is obtained by attaching the space of *projective measured geodesic laminations* at infinity. In terms of certain fixed point properties of this action, each mapping class is classified into three types: periodic, reducible, and pseudo-Anosov (often abbreviated as “pA”) [Th88]. This result is called the *Nielsen–Thurston classification*, which gives a beautiful generalization of the elliptic-parabolic-hyperbolic trichotomy for elements of the modular group $PSL_2(\mathbb{Z})$.

For a punctured surface Σ (*i.e.*, a closed surface equipped with a finite set of marked points), the *decorated Teichmüller theory* [Pen87] provides a combinatorial tool for study of the Teichmüller–Thurston theory mentioned above. More precisely, there are two variants $\tilde{T}(\Sigma)$ and $\hat{T}(\Sigma)$ of the Teichmüller space called the *decorated* and *enhanced Teichmüller spaces*, respectively. These spaces have distinguished charts associated with ideal triangulations, and the action of the mapping class group has a rational expression in terms of these coordinate systems. A similar construction applies for the spaces $\widetilde{\mathcal{ML}}(\Sigma)$ and $\widehat{\mathcal{ML}}(\Sigma)$ of *decorated* and *enhanced measured geodesic laminations* (or equivalently, *measured foliations*), which have piecewise-linear (PL for short) coordinate systems associated with ideal triangulations [PP93, BKS11]. The actions of the mapping class group on these spaces of laminations are expressed as PL maps in these coordinates, which are

the *tropical analogues* of the expressions of the actions on the corresponding Teichmüller spaces. The relations between the spaces mentioned above are summarized in the following diagram:

$$\begin{array}{ccc} \widetilde{T}(\Sigma) & & \widetilde{\mathcal{ML}}(\Sigma) \\ \downarrow & & \downarrow \\ T(\Sigma) & \hookrightarrow & \widehat{T}(\Sigma), \quad \mathcal{ML}_0(\Sigma) \hookrightarrow \widehat{\mathcal{ML}}(\Sigma). \end{array}$$

where $\mathcal{ML}_0(\Sigma)$ denotes the space of measured geodesic laminations with compact support. In the last section in their paper [PP93], Papadopoulos and Penner raised the following question:

Problem 1.1 (Papadopoulos–Penner [PP93]). Characterize the Nielsen–Thurston type of a mapping class in terms of its rational or PL coordinate expressions.

Nowadays the combinatorial structures of the Teichmüller and lamination spaces mentioned above are recognized in terms of the *cluster algebra*, which is introduced by Fomin–Zelevinsky [FZ02] and Fock–Goncharov [FG06a]. In the language of the cluster algebra, there exists a seed pattern \mathbf{s}_Σ associated with a punctured surface Σ formed by the ideal (or tagged) triangulations, and the Teichmüller and lamination spaces are uniformly reconstructed from the associated *cluster varieties* \mathcal{A}_Σ and \mathcal{X}_Σ , as follows [FG07, FST08]:

$$\begin{aligned} \mathcal{A}_\Sigma(\mathbb{R}_{>0}) &\cong \widetilde{T}(\Sigma), & \mathcal{X}_\Sigma(\mathbb{R}_{>0}) &\cong \widehat{T}(\Sigma), \\ \mathcal{A}_\Sigma(\mathbb{R}^{\text{trop}}) &\cong \widetilde{\mathcal{ML}}(\Sigma), & \mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}}) &\cong \widehat{\mathcal{ML}}(\Sigma). \end{aligned}$$

Here the *positive structures* of the cluster varieties allow us to consider their \mathbb{P} -valued points for any semifield $\mathbb{P} = (\mathbb{P}, \oplus, \cdot)$, where $\mathbb{R}_{>0} = (\mathbb{R}_{>0}, +, \cdot)$ and $\mathbb{R}^{\text{trop}} = (\mathbb{R}, \min, +)$ is the tropical semifield. The natural actions of the mapping class group $MC(\Sigma)$ on these spaces coincides with the one through an embedding $MC(\Sigma) \rightarrow \Gamma_\Sigma$ into the *cluster modular group*, which is a group consisting of *mutation loops* and acting on the cluster varieties. Based on this correspondence, the first author studied an analogue of the Nielsen–Thurston classification for cluster modular groups in [Ish19] by introducing three types for mutation loops: periodic, cluster-reducible, cluster-pA. We have a nice reduction construction called the *cluster reduction* which eventually produces a cluster-pA mutation loop, and moreover the three types can be characterized by certain fixed point properties of the action on the *tropical compactification* [FG16, Le16] of the spaces $\mathcal{A}_\mathbf{s}(\mathbb{R}_{>0})$ and $\mathcal{X}_\mathbf{s}(\mathbb{R}_{>0})$ similarly to the original Nielsen–Thurston’s spirit. However, one unsatisfactory point is that the cluster-pA property is weaker than the original pA property ([Ish19, Example 3.13]), and an analogue of the *stretch factor* of a pA mapping class is also missing. Our aim in this paper is to study a more suitable class of mutation loops which generalizes pA mapping classes.

1.1. Sign stability and the pseudo-Anosov property. A new property of mutation loops called the *sign stability* is introduced in [IK21]. It is defined as a certain stability property of the PL expression of the action of a mutation loop on the \mathbb{R}^{trop} -valued point $\mathcal{X}_s(\mathbb{R}^{\text{trop}})$ by mimicking the convergence of *RLS sequences* of train track splittings for pA mapping classes [PP87]. A sign-stable mutation loop comes with a numerical invariant called the *cluster stretch factor*, defined as the Perron–Frobenius eigenvalue of the stable presentation matrix (Definition 5.12). Our assertion is that the sign stability is a correct generalization of the pA property.

Precisely speaking, the sign stability does depend on the choices of a representation path γ and an $\mathbb{R}_{>0}$ -invariant subset $\Omega \subset \mathcal{X}_s(\mathbb{R}^{\text{trop}})$ on which the PL dynamical system stabilizes to a linear one. We can consider in a sense “strongest” version of sign stability called the *uniform sign stability* (Definition 5.17), which is genuinely a property of a mutation loop. In this paper, we verify that the uniform sign stability is indeed equivalent to the pA property for the mutation loops arising from mapping classes on a punctured surface.

Theorem 1.2 (Theorem 7.1). *Let Σ be a punctured surface. Then a mapping class $\phi \in MC(\Sigma) \subset \Gamma_\Sigma$ is pseudo-Anosov if and only if the corresponding mutation loop is uniformly sign-stable. Moreover, the associated cluster stretch factor λ_ϕ coincides with the stretch factor of the pseudo-Anosov mapping class ϕ .*

The second statement shows that our concept of the cluster stretch factor also correctly generalize the stretch factor of a pA mapping class. The definition of the former as the spectral radius of the stable presentation matrix also quite resembles the description of the latter as the spectral radius of the *transition matrix* for the train track splittings [PH, BH95]. We also explain certain relations between the sign stability and the Nielsen–Thurston classification of [Ish19] in Section 5.4.

Since the uniform sign stability is written in terms of the mutation loop (or equivalently, the collection of coordinate expressions) associated with a mapping class, this theorem can be served as an answer to Problem 1.1. We prove a slightly generalized result for mapping classes with reflections (Corollary 7.7).

1.2. Sign stability of Dehn twists. We also verify that a non-uniform version of sign stability holds for specific representation paths of Dehn twists.

The existence of such a sign-stable representation path is useful enough to compute the algebraic entropies of the induced cluster \mathcal{A} - and \mathcal{X} -transformations as the logarithm of the cluster stretch factor, as demonstrated in [IK21]. We conclude that a Dehn twist satisfying a mild condition admits a representation path which is sign-stable on a certain $\mathbb{R}_{>0}$ -invariant set and its cluster stretch factor is 1 (Proposition 6.1), and hence the entropies of the induced cluster transformations are 0 (Corollary 8.8). On the other hand, we observe that a Dehn twist also has a representation path which is not sign-stable on the same $\mathbb{R}_{>0}$ -invariant set (Example 6.3). This gives a counterexample to the

naive conjecture [IK21, Conjecture 1.3] which asserted the independence of sign stability on representation paths.

1.3. The topological and algebraic entropies. As an application of our study of sign stability, we compute the algebraic entropies of the cluster \mathcal{A} - and \mathcal{X} -transformations induced by mapping classes ϕ , based on the main result of [IK21]. Moreover we compare them with the *topological entropy* [AKM65] $\mathcal{E}_\phi^{\text{top}}$.

For a mutation loop $\phi \in \Gamma_s$, let \mathcal{E}_ϕ^a (resp. \mathcal{E}_ϕ^x) denote the algebraic entropy of the cluster \mathcal{A} - (resp. \mathcal{X} -)transformation induced by ϕ .

Theorem 1.3 (Corollaries 8.8 and 8.9). *We have the following comparison results:*

- (1) *For a simple closed curve C on a marked surface Σ satisfying the condition (6.1), we have*

$$\mathcal{E}_{t_C}^a = \mathcal{E}_{t_C}^x = \mathcal{E}_{t_C}^{\text{top}} = 0.$$

- (2) *For a punctured surface Σ and a pA mutation loop $\phi \in \Gamma_\Sigma$, we have*

$$\mathcal{E}_\phi^a = \mathcal{E}_\phi^x = \mathcal{E}_\phi^{\text{top}} = \log \lambda_\phi.$$

Here $\lambda_\phi > 1$ is the cluster stretch factor (=stretch factor) of ϕ .

Thus we get a complete agreement of the two kinds of entropies associated with these mapping classes. Note that the cluster variety \mathcal{A}_Σ (resp. \mathcal{X}_Σ) is birationally isomorphic to the moduli space of decorated twisted SL_2 - (resp. framed PGL_2 -)local systems on Σ [FG06a], and \mathcal{E}_ϕ^a (resp. \mathcal{E}_ϕ^x) is the algebraic entropy of the natural action of a mapping class on that moduli space. It should be compared with the work of Hadari [Had11], where the algebraic entropies of the actions of a mapping class on various character varieties are studied.

Organization of the paper. A reader familiar with the relation between the Teichmüller–Thurston theory and the cluster algebra may skip to the Section 5.

In Section 2, we recall the two spaces $\mathcal{A}_\Sigma(\mathbb{R}^{\text{trop}})$ and $\mathcal{X}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}})$ of measured laminations. In the first three subsections we give the combinatorial definition of these spaces as certain completions of the set of rational laminations, following Fock–Goncharov [FG07]. Then the geometric description of these spaces in terms of the *measured foliations* and *measured geodesic laminations* are given.

In Section 3, the definition of the labeled exchange graph, the cluster modular group and the sign stability are recalled. Since we need to carefully distinguish mutation loops and their representation paths, we give a detailed reformulation of these notions. A simplified definition given in [IK21] may help the reader to follow the discussion. Basic definitions on the cluster varieties are collected in Appendix A.

In Section 4, we define the seed patterns associated with a marked surface. A dictionary between the lamination theory and the cluster algebra is provided. Theorem 4.5 gives a slight refinement of known results on the cluster modular group.

In Section 5, we introduce basic notions related to the sign stability. The relation to the Nielsen–Thurston classification in [Ish19] is also explained in Section 5.4. In particular, we see that the sign stability is preserved under the cluster reduction and that the uniformly sign-stable mutation loop is cluster-pA.

In Section 6, we study the sign stability of cluster Dehn twists. The contents in this section may be regarded as a mild introduction to the sign stability.

In Section 7, we investigate the uniform sign stability of pA mutation loops on a punctured surface and prove Theorem 1.2.

Acknowledgements. The authors are grateful to Hidetoshi Masai for informing us of the modern results on the North-South dynamics on the lamination spaces. T. I. would like to express his gratitude to his former supervisor Nariya Kawazumi for his continuous guidance and encouragement in the earlier stage of this work. S. K. would like to thank Yoshihiko Mitsumatsu. The strategy of the proof of Lemma 7.6 is inspired by him. S. K. is also deeply grateful to his supervisor Yuji Terashima for his advice and giving him a lot of knowledge.

T. I. is partially supported by JSPS KAKENHI Grant Number 18J13304 and the Program for Leading Graduate Schools, MEXT, Japan.

2. LAMINATIONS ON A MARKED SURFACE

In this section we recall the concepts around the laminations following [FG07]. In the first two sections, two spaces of laminations are introduced, which will be identified with the tropical cluster varieties $\mathcal{A}_\Sigma(\mathbb{R}^{\text{trop}})$ and $\mathcal{X}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}})$ (see Section 4). An ensemble structure between these spaces are recalled in Section 2.3. In Sections 2.4 and 2.5, we review geometric models for the laminations called the *measured foliations* and the *measured geodesic laminations* respectively, which allow us to analyze the irrational points of the tropical cluster varieties.

A marked surface Σ is a compact oriented surface with a fixed non-empty finite set of *marked points* on it. A marked point is called a *puncture* if it lies in the interior of Σ , and a *special point* otherwise. Let $P = P(\Sigma)$ (resp. $M_\partial = M_\partial(\Sigma)$) denote the set of punctures (resp. special points). A marked surface is called a *punctured surface* if it has empty boundary (and hence $M_\partial = \emptyset$). We denote by g the genus of Σ , h the number of punctures, and b the number of boundary components in the sequel. We always assume the following conditions:

- (S1) Each boundary component (if exists) has at least one marked point.
- (S2) $3(2g - 2 + h + b) + 2|M_\partial| > 0$.
- (S3) If $g = 0$ and $b = 0$, then $h \geq 4$.

An *ideal arc* in Σ is the isotopy class of a non-contractible curve α such that $\partial\alpha \subset P \cup M_\partial$. An *ideal triangulation* Δ of Σ is a collection of ideal arcs such that

- each pair of ideal arcs in Δ can intersect only at their endpoints;

- each region complementary to Δ is a triangle whose edges (resp. vertices) are ideal arcs (resp. marked points).

The conditions (S1) and (S2) ensure that such an ideal triangulation Δ exists, and in particular the number $3(2g - 2 + h + b) + 2|M_\partial| > 0$ in (S2) gives the number of ideal arcs in Δ . For an ideal triangulation Δ , let $\Delta_f \subset \Delta$ denote the subset consisting of boundary arcs and hence $\Delta_{\text{uf}} := \Delta \setminus \Delta_f$ is the subset consisting of interior ideal arcs.

In the sequel, by a *curve* we mean a simple curve γ in Σ which satisfies $\partial\gamma = \emptyset$ (closed) or $\partial\gamma \subset P \cup (\partial\Sigma \setminus M_\partial)$. Isotopies of curves are considered within this class. Such a curve γ is said to be

- *peripheral*¹ if it is either isotopic to a puncture $m \in P$ or an interval in $\partial\Sigma$ which contains exactly one special point $m \in M_\partial$ (in each case, called a peripheral curve *around* m);
- *contractible* if it is isotopic to a point other than a puncture;
- *compact* if it is closed or $\partial\gamma \subset \partial\Sigma \setminus M_\partial$.

A *rational weighted curve* is a pair (γ, w) of a curve in the above sense and a rational number $w \in \mathbb{Q}$.

2.1. The space of \mathcal{A} -laminations.


Definition 2.1. A *rational \mathcal{A} -lamination* (or a *rational bounded lamination*) on Σ is a collection of mutually disjoint compact rational weighted curves in Σ such that the weight of a non-peripheral curve is positive, subject to the following equivalence relations (see Figure 1):

- (1) A collection containing a curve of weight zero is equivalent to the collection with this curve removed.
- (2) A collection containing a contractible curve is equivalent to the collection with this curve removed.
- (3) A collection containing two isotopic curves with weights u and v is equivalent to the collection with one of these curves removed and with the weight $u + v$ on the other.

We write a rational \mathcal{A} -lamination as $L = \bigsqcup_j (\gamma_j, w_j)$, where each (γ_j, w_j) is a rational weighted curve in that collection. We call each curve γ_j a *leaf* of L . Let $\mathcal{L}^a(\Sigma, \mathbb{Q})$ denote the set of rational \mathcal{A} -laminations on Σ . We are going to give a certain global coordinate on this space, associated with an ideal triangulation.

Following [FG07], we define a coordinate function \mathbf{a}_α on $\mathcal{L}^a(\Sigma, \mathbb{Q})$ for each ideal arc α . For $L = \bigsqcup_j (\gamma_j, w_j) \in \mathcal{L}^a(\Sigma, \mathbb{Q})$, represent the leaves γ_j and the ideal arc α so that the

¹It is called “special” in [FG07].

FIGURE 1. An example of a rational \mathcal{A} -lamination.

intersections of these curves are minimal. Then we define

$$\mathbf{a}_\alpha(L) := \frac{1}{2} \sum_j w_j \text{Int}(\alpha, \gamma_j),$$

where Int denote the geometric intersection number (*i.e.*, it just counts the number of intersections of two curves). Then we have the following:

Proposition 2.2 ([FG07, Section 3.2]). *For any ideal triangulation Δ of Σ , the map*

$$\mathbf{a}_\Delta : \mathcal{L}^a(\Sigma, \mathbb{Q}) \xrightarrow{\sim} \mathbb{Q}^\Delta, \quad L \mapsto \{\mathbf{a}_\alpha(L)\}_{\alpha \in \Delta}$$

gives a bijection. Moreover, if an ideal triangulation $\Delta' = f_\kappa(\Delta)$ is obtained by the flip along an arc $\kappa \in \Delta_{\text{uf}}$, then the coordinate transformation $\mathbf{a}_{\Delta'} \circ \mathbf{a}_\Delta^{-1}$ is given as in Figure 2.

In particular, the coordinate transformation $\mathbf{a}_{\Delta'} \circ \mathbf{a}_\Delta^{-1}$ is a Lipschitz map with respect to the Euclidean metric on $\mathbb{Q}^\Delta \cong \mathbb{Q}^{3(2g-2+h+b)+2|M_\partial|}$. Let $\mathcal{L}^a(\Sigma, \mathbb{R})$ be the corresponding metric completion of $\mathcal{L}^a(\Sigma, \mathbb{Q})$, which does not depend on a specific coordinate system. Each coordinate system \mathbf{a}_Δ is extended to a homeomorphism $\mathbf{a}_\Delta : \mathcal{L}^a(\Sigma, \mathbb{R}) \xrightarrow{\sim} \mathbb{R}^\Delta$ (still denoted by the same symbol). We call an element of $\mathcal{L}^a(\Sigma, \mathbb{R})$ a *real \mathcal{A} -lamination*.




FIGURE 2. The coordinate transformation for a flip. Here the transformation rule is still valid even when some of edges are identified.

Thus we get a PL manifold $\mathcal{L}^a(\Sigma, \mathbb{R})$ equipped with a distinguished collection $\{\mathbf{a}_\Delta\}_\Delta$ of global charts parametrized by ideal triangulations. The space $\mathcal{L}^a(\Sigma, \mathbb{R})$ has the following additional structures.

Action of the group $\mathbb{R}^{h+|M_\partial|}$. For each marked point $m \in P \cup M_\partial$, a rational number $u \in \mathbb{Q}$ and a rational \mathcal{A} -lamination $L \in \mathcal{L}^a(\Sigma, \mathbb{Q})$, let $r_m(u, L)$ be the lamination obtained by adding a peripheral curve with weight u around the marked point m to the lamination L . Then we get an action $r_m : \mathbb{Q} \times \mathcal{L}^a(\Sigma, \mathbb{Q}) \rightarrow \mathcal{L}^a(\Sigma, \mathbb{Q})$. For an ideal triangulation Δ , this action is expressed in the coordinate as

$$\mathbf{a}'_\alpha = \begin{cases} \mathbf{a}_\alpha + u & \text{if both ends of } \alpha \text{ are incident to } m, \\ \mathbf{a}_\alpha + \frac{1}{2}u & \text{if one of the ends of } \alpha \text{ is incident to } m, \\ \mathbf{a}_\alpha & \text{otherwise.} \end{cases}$$

Here $\mathbf{a}_\alpha := \mathbf{a}_\alpha(L)$ and $\mathbf{a}'_\alpha := \mathbf{a}_\alpha(r_m(u, L))$ for $\alpha \in \Delta$. In particular the action r_m is Lipschitz (indeed, \mathbb{Q} -linear) and thus extended to an action

$$r_m : \mathbb{R} \times \mathcal{L}^a(\Sigma, \mathbb{R}) \rightarrow \mathcal{L}^a(\Sigma, \mathbb{R}). \quad (2.1)$$

Since these actions for different marked points commute with each other, we get an action $r := (r_m)_{m \in P \cup M_\partial}$ of $\mathbb{R}^{h+|M_\partial|}$ on $\mathcal{L}^a(\Sigma, \mathbb{R})$.

Tropicalized Goncharov–Shen potentials. For each marked point $m \in P \cup M_\partial$ and a rational \mathcal{A} -lamination $L \in \mathcal{L}^a(\Sigma, \mathbb{Q})$, let $W_m(L)$ be the half of the total weight of the peripheral curves of L around m . For an ideal triangulation Δ , the function W_m is expressed as follows:

$$W_{\Delta, m} := W_m \circ \mathbf{a}_\Delta^{-1} = \min\{\mathbf{a}_{\alpha_{m,t}^1} + \mathbf{a}_{\alpha_{m,t}^2} - \mathbf{a}_{\alpha_{m,t}^3}\}.$$

Here the minimum is taken over all the triangles t of Δ which has the marked point m as one of its vertices, and the ideal arcs $\alpha_{m,t}^k$ for $k = 1, 2, 3$ is defined in Figure 3. In particular the function W_m is Lipschitz and thus extended to a PL function $W_m : \mathcal{L}^a(\Sigma, \mathbb{R}) \rightarrow \mathbb{R}$. We call these functions the *tropicalized Goncharov–Shen potentials*. The relation

$$W_m(r_{m'}(u, L)) = W_m(L) + \delta_{mm'}u$$

is clear from their geometric definitions.




FIGURE 3.

Mapping class group action. Let $MC(\Sigma)$ be the *mapping class group* of Σ , which consists of the isotopy classes of diffeomorphisms of Σ which preserve the set M of marked points. If Σ is a once-punctured torus, then we redefine $MC(\Sigma)$ as the quotient of that group by the hyperelliptic involution. An element of $MC(\Sigma)$ is called a *mapping class*. Note that a mapping class is allowed to permute boundary components with the same number of marked points and/or rotate them. A permutation of punctures is also allowed.

A mapping class naturally acts on the homotopy classes of curves on Σ , preserving the peripheral/contractible/compact properties. Thus the mapping class group naturally acts on the space $\mathcal{L}^a(\Sigma, \mathbb{Q})$. This action is expressed in terms of the charts \mathbf{a}_Δ as follows. For a mapping class $\phi \in MC(\Sigma)$, take a sequence f_ϕ of flips from Δ to $\phi^{-1}(\Delta)$. Such a sequence is known to always exist: see, for instance, [Pen]. Then we have the coordinate transformation $f_\phi^a := \mathbf{a}_{\phi^{-1}(\Delta)} \circ \mathbf{a}_\Delta^{-1} : \mathbb{Q}^\Delta \rightarrow \mathbb{Q}^{\phi^{-1}(\Delta)}$ and the commutative diagram

$$\begin{array}{ccccc} \mathbb{Q}^\Delta & \xrightarrow{f_\phi^a} & \mathbb{Q}^{\phi^{-1}(\Delta)} & \xrightarrow{\phi^*} & \mathbb{Q}^\Delta \\ \mathbf{a}_\Delta \uparrow & & \mathbf{a}_{\phi^{-1}(\Delta)} \uparrow & & \mathbf{a}_\Delta \uparrow \\ \mathcal{L}^a(\Sigma, \mathbb{Q}) & \xlongequal{\quad} & \mathcal{L}^a(\Sigma, \mathbb{Q}) & \xrightarrow{\phi} & \mathcal{L}^a(\Sigma, \mathbb{Q}). \end{array} \quad (2.2)$$

Here the map $\phi^* : \mathbb{Q}^{\phi^{-1}(\Delta)} \xrightarrow{\sim} \mathbb{Q}^\Delta$ is the pull-back induced by $\phi^{-1} : \Delta \rightarrow \phi^{-1}(\Delta)$. Hence the coordinate expression of the action of ϕ on $\mathcal{L}^a(\Sigma, \mathbb{Q})$ is given by the composition $\phi^a := \phi^* \circ f_\phi^a$, which is a PL map. In particular it is extended to a PL action on $\mathcal{L}^a(\Sigma, \mathbb{R})$.

Coordinates associated with tagged triangulations. In general, there exists a flip producing a *self-folded triangle* as shown in Figure 4. In this case the coordinate transformation does not agree with the tropical cluster \mathcal{A} -transformation, and what is worse, we cannot perform a flip anymore along the edge enclosed by the loop. These features are unsatisfactory in comparison with the cluster algebra (see Section 3), so we are lead to introduce *tagged triangulations* to fix them. We follow the description given by Bridgeland–Smith [BS15].




FIGURE 4. The coordinate transformation for a flip producing a self-folded triangle.

A *signed triangulation* is a pair (Δ, ς) of an ideal triangulation Δ and a tuple $\varsigma \in \{+, -\}^P$ of signs. Two signed triangulations are equivalent if the underlying ideal triangulations are the same and the signatures differ only on univalent punctures. The equivalence classes are called *tagged triangulations*. We regard an ideal triangulation as a signed triangulation with constant signature $\varsigma_p = +$ for all $p \in P$, and as the corresponding tagged triangulation. The notion of flips of ideal triangulations are obviously extended to those of signed triangulations, and hence of tagged triangulations.

A *tagged arc* is an ideal arc each of whose end is labeled as “plain” or “notched”. In the pictures, the plain tag is omitted and the notched tag is indicated by the symbol \bowtie . For a tagged triangulations (Δ, ς) and an arc $\alpha \in \Delta$, we associate a tagged arc $t_\varsigma(\alpha)$ by the following rules:

- (a) If α is not a loop enclosing a once-punctured monogon, then the underlying arc of $t_\varsigma(\alpha)$ is α itself and an end of e incident to a puncture p is tagged notched if and only if $\varsigma_p = -$.
- (b) If α is a loop based at a marked point m and enclosing a once-punctured monogon with the central puncture p , then the underlying arc of $t_\varsigma(\alpha)$ is the arc inside the monogon connecting m with p . The tag of the end incident to m is determined as before (notched when $\varsigma_m = -$), while the end incident to p is tagged notched if and only if $\varsigma_p = +$.

Then the family $\{t_\varsigma(\alpha)\}_{\alpha \in \Delta}$ gives a tagged triangulation in the sense of Fomin–Shapiro–Thurston [FST08], and indeed the two definitions are equivalent. See [BS15, Section 8.3].

Now let us define a coordinate function $\mathbf{a}_{t_\varsigma(\alpha)}$ on $\mathcal{L}^a(\Sigma, \mathbb{Q})$ for a tagged arc $t_\varsigma(\alpha)$ associated with an ideal arc α and a signature $\varsigma \in \{+, -\}^P$. For $L \in \mathcal{L}^a(\Sigma, \mathbb{Q})$, let $\tau_\varsigma(L)$ be the rational \mathcal{A} -lamination obtained from L by multiplying the sign ς_p to the weight assigned to the peripheral leaf around p (if exists) for each puncture $p \in P$. Then define

$$\mathbf{a}_{t_\varsigma(\alpha)}(L) := \mathbf{a}_\alpha(\tau_\varsigma(L)).$$

Then for a tagged triangulation (Δ, ς) , we get a coordinate system

$$\mathbf{a}_{(\Delta, \varsigma)} := \{\mathbf{a}_{t_\varsigma(\alpha)}\}_{\alpha \in \Delta} : \mathcal{L}^a(\Sigma, \mathbb{Q}) \rightarrow \mathbb{Q}^\Delta.$$

It is not hard to see that $\mathbf{a}_{\alpha'} = \mathbf{a}_\beta + \mathbf{a}_{t_\varsigma(\alpha)}$ in the situation of Figure 4, which leads to the correct cluster transformation (Proposition 2.7).

2.2. The space of \mathcal{X} -laminations.

Definition 2.3. A *rational \mathcal{X} -lamination* (or a *rational unbounded lamination*) on Σ consists of a collection of mutually disjoint rational weighted curves in Σ such that all weights are positive, and a tuple $\sigma_p \in \{+, 0, -\}$ of signs assigned to punctures $p \in P$ satisfying $\sigma_p \in \{+, -\}$ if some of the curves are incident to p , and otherwise $\sigma_p = 0$. Such data is considered modulo the following equivalence relations (see Figure 5):




FIGURE 5. An example of a rational \mathcal{X} -lamination. Here $\sigma_1, \sigma_2 \in \{+, -\}$.

- (1) A collection containing a peripheral or contractible curve is equivalent to the collection with this curve removed.
- (2) A collection containing two isotopic curves with weights u and v is equivalent to the collection with one of these curves removed and with the weight $u + v$ on the other.

We write a rational \mathcal{X} -lamination as $\widehat{L} = (L, \sigma_L)$, where $L = \bigsqcup (\gamma_j, w_j)$ is the underlying collection of rational weighted curves (γ_j, w_j) , and $\sigma_L = (\sigma_p)_{p \in P}$ is the tuple of signs which we call the *lamination signature*. We call each curve γ_j a *leaf* of \widehat{L} . Let $\mathcal{L}^x(\Sigma, \mathbb{Q})$ denote the set of rational \mathcal{X} -laminations.

Given an ideal triangulation Δ of Σ , we define a coordinate system on $\mathcal{L}^x(\Sigma, \mathbb{Q})$ following [FG07]. For $\widehat{L} = (L, \sigma_L) \in \mathcal{L}^x(\Sigma, \mathbb{Q})$, represent the leaves γ_j of L and the ideal arcs in Δ so that the intersections of these curves are minimal. After that we deform each interior ideal arc γ_j in L as follows: if the sign at an endpoint of γ_j is positive (resp. negative), then replace the corresponding end of γ_j by a curve which spirals around that endpoint in the clockwise (resp. counter-clockwise) direction (see Figure 6).




FIGURE 6. Sign and the spiralling direction.

For each interior arc $\alpha \in \Delta_{\text{uf}}$, let $\widehat{\text{Int}}_{\Delta}(\alpha, \gamma_j)$ be the intersection number given by the sum of the following contributions: an intersection as in the left (resp. right) of Figure 7 contributes as $+1$ (resp. -1), and the others 0. Note that $\widehat{\text{Int}}_{\Delta}(\alpha, \gamma_j)$ is a finite integer,

while the ordinary intersection number $\text{Int}(\alpha, \gamma_j)$ can be infinite due to spirals. When α is not the interior edge of a self-folded triangle, we define

$$\mathbf{x}_\alpha^\Delta(\widehat{L}) := \sum_j w_j \widehat{\text{Int}}_\Delta(\alpha, \gamma_j).$$

Note that peripheral or contractible curves have no contribution to these coordinates, which is the reason for the equivalence relation (1) in Definition 2.3. When α is the interior edge of a self-folded triangle, let β be the loop enclosing this self-folded triangle and define

$$\mathbf{x}_\alpha^\Delta(\widehat{L}) := \sum_j w_j (\widehat{\text{Int}}_\Delta(\alpha, \gamma_j) + \widehat{\text{Int}}_\Delta(\beta, \gamma_j)).$$

This modification is made so that the coordinate transformation agrees with the tropical cluster \mathcal{X} -transformation: see Figure 9 and Proposition 2.7.




FIGURE 7. Contribution of a intersection to $\widehat{\text{Int}}_\Delta(\alpha, \gamma_j)$.

Proposition 2.4 ([FG07, Section 3.1]). *For any ideal triangulation Δ of Σ , the map*

$$\mathbf{x}_\Delta : \mathcal{L}^x(\Sigma, \mathbb{Q}) \xrightarrow{\sim} \mathbb{Q}^{\Delta_{\text{uf}}}, \quad \widehat{L} \mapsto \{\mathbf{x}_\alpha^\Delta(\widehat{L})\}_{\alpha \in \Delta_{\text{uf}}}$$

gives a bijection. Moreover if an ideal triangulation $\Delta' = f_\kappa(\Delta)$ is obtained by the flip along an arc $\kappa \in \Delta_{\text{uf}}$, the coordinate transformation $\mathbf{x}_{\Delta'} \circ \mathbf{x}_\Delta^{-1}$ is given as in Figure 8 and Figure 9².

Since the coordinate transformation $\mathbf{x}_{\Delta'} \circ \mathbf{x}_\Delta^{-1}$ is a Lipschitz map with respect to the Euclidean metric on $\mathbb{Q}^{\Delta_{\text{uf}}} \cong \mathbb{Q}^{3(2g-2+h+b)+|M_\partial|}$, we can consider the corresponding metric completion $\mathcal{L}^x(\Sigma, \mathbb{R})$ as before. Each coordinate system \mathbf{x}_Δ is extended to a homeomorphism $\mathbf{x}_\Delta : \mathcal{L}^x(\Sigma, \mathbb{R}) \xrightarrow{\sim} \mathbb{R}^{\Delta_{\text{uf}}}$ (still denoted by the same symbol). We call an element of $\mathcal{L}^x(\Sigma, \mathbb{R})$ a *real \mathcal{X} -lamination*.

Thus we get a PL manifold $\mathcal{L}^x(\Sigma, \mathbb{R})$ equipped with a distinguished collection $\{\mathbf{x}_\Delta\}_\Delta$ of global charts parametrized by ideal triangulations.

²The transformation formula shown in Figure 8 holds in the universal cover. Applying this formula in the case shown in Figure 9, we have $\widehat{\text{Int}}_{\Delta'}(\beta, -) = \mathbf{x}_0 + \mathbf{x}_3$. Thus we get $\mathbf{x}_3^{\Delta'} = \widehat{\text{Int}}_{\Delta'}(\alpha, -) + \widehat{\text{Int}}_{\Delta'}(\beta, -) = -\mathbf{x}_0 + (\mathbf{x}_0 + \mathbf{x}_3) = \mathbf{x}_3$.




FIGURE 8. The coordinate transformation for a flip. The formula is still valid when some of the edges are identified as $x_1 = x_3$ and/or $x_2 = x_4$. For the case $x_1 = x_2$ or $x_3 = x_4$, see Figure 9.




FIGURE 9. The coordinate transformation for a flip producing a self-folded triangle.

Example 2.5 (A collection of real weighted curves). Let $(\gamma_i)_i$ be a collection of mutually disjoint curves, and $(w_i)_i$ a collection of non-negative weights. Further by specifying the sign $\sigma_p \in \{+, -\}$ for each puncture $p \in P$ to which some of the curves γ_i incident, we get a real \mathcal{X} -lamination. Indeed, it can be approximated by rational laminations by taking a sequence of rational numbers that approximates w_i . However, not all the real \mathcal{X} -laminations arise in this way. See Section 2.5 for a geometric model of general real \mathcal{X} -laminations.

The space $\mathcal{L}^x(\Sigma, \mathbb{R})$ has the following additional structures.

Tropicalized Casimir functions. For each puncture $p \in P$ and a rational \mathcal{X} -lamination $\widehat{L} = (L, \sigma_L) \in \mathcal{L}^x(\Sigma, \mathbb{Q})$, let $\theta_p(\widehat{L})$ be the total weight of the curves incident to the puncture p . Here the weight of a curve whose endpoints are both incident to p is counted twice. We call these functions the *tropicalized Casimir functions*. For an ideal triangulation Δ , the function θ_p is expressed as

$$\theta_{\Delta,p} := \theta_p \circ x_{\Delta}^{-1} = \sum_{\alpha} x_{\alpha}. \tag{2.3}$$

Here the sum is taken over all the ideal arcs α in Δ_{uf} which is incident to the puncture p , and counted twice if both of its endpoints are incident to p . The lamination signature σ_L coincides with the absolute value of $\theta_p(\widehat{L})$. The tropicalized Casimir functions are clearly extended to PL functions $\theta_p : \mathcal{L}^x(\Sigma, \mathbb{R}) \rightarrow \mathbb{R}$. Thus we get $\theta := (\theta_p)_{p \in P} : \mathcal{L}^x(\Sigma, \mathbb{R}) \rightarrow \mathbb{R}^h$.

Mapping class group action. The mapping class group $MC(\Sigma)$ naturally acts on the space $\mathcal{L}^x(\Sigma, \mathbb{Q})$. Here the lamination signatures are permuted according to the natural projection $MC(\Sigma) \rightarrow \text{Aut}(P) \cong \mathfrak{S}_h$. The description of this action in terms of the charts \mathbf{x}_Δ is similar to the \mathcal{A} -case. For a mapping class $\phi \in MC(\Sigma)$, let f_ϕ be a sequence of flips from Δ to $\phi^{-1}(\Delta)$ and consider the corresponding coordinate transformation $f_\phi^x := \mathbf{x}_{\phi^{-1}(\Delta)} \circ \mathbf{x}_\Delta^{-1}$. Then the action is given by the PL map $\phi^x := \phi^* \circ f_\phi^x$. It is extended to a PL action on $\mathcal{L}^x(\Sigma, \mathbb{R})$.

Coordinates associated with tagged triangulations. For a tagged triangulation (Δ, ς) , we associate a coordinate system $\mathbf{x}_{(\Delta, \varsigma)} = \{\mathbf{x}_\alpha^{(\Delta, \varsigma)}\}_{\alpha \in \Delta_{\text{uf}}}$ on $\mathcal{L}^x(\Sigma, \mathbb{Q})$ as follows. For $\widehat{L} = (L, \sigma_L) \in \mathcal{L}^x(\Sigma, \mathbb{Q})$, set $\tau_\varsigma(\widehat{L}) := (L, \varsigma \cdot \sigma_L)$. Here the dot means the component-wise multiplication. Then define

$$\mathbf{x}_\alpha^{(\Delta, \varsigma)}(\widehat{L}) := \mathbf{x}_\alpha^\Delta(\tau_\varsigma(\widehat{L})).$$

2.3. Ensemble map and the space of \mathcal{U} -laminations. The two spaces $\mathcal{L}^a(\Sigma, \mathbb{R})$ and $\mathcal{L}^x(\Sigma, \mathbb{R})$ discussed above are related as follows. Given a rational \mathcal{A} -lamination $L \in \mathcal{L}^a(\Sigma, \mathbb{Q})$, let $p_\Sigma(L)$ be the collection of curves obtained from L by forgetting all the peripheral curves in L . Then $p_\Sigma(L)$ can be naturally regarded as a rational \mathcal{X} -lamination, since it has no leaves incident to punctures and thus we do not need to specify the lamination signature. Thus we get a map $p_\Sigma : \mathcal{L}^a(\Sigma, \mathbb{Q}) \rightarrow \mathcal{L}^x(\Sigma, \mathbb{Q})$, which is called the *ensemble map*.

Let $\mathcal{L}^u(\Sigma, \mathbb{Q}) \subset \mathcal{L}^x(\Sigma, \mathbb{Q})$ be the image of the ensemble map, which consists of the rational \mathcal{X} -laminations only with compact curves. We call an element of $\mathcal{L}^u(\Sigma, \mathbb{Q})$ a *rational \mathcal{U} -lamination*. The \mathcal{U} -laminations are characterized as follows:

Lemma 2.6. *We have $\mathcal{L}^u(\Sigma, \mathbb{Q}) = \bigcup_{p \in P} \theta_p^{-1}(0)$.*

In particular, the corresponding metric completion $\mathcal{L}^u(\Sigma, \mathbb{R})$ is a closed subset of $\mathcal{L}^x(\Sigma, \mathbb{R})$.

Exchange matrix. In order to express the ensemble map in the coordinate systems associated with a tagged triangulation (Δ, ς) , let us recall the *exchange matrix* $B^\Delta = (b_{\alpha\beta}^\Delta)_{\alpha, \beta \in \Delta}$ defined as follows (independent of the signature ς). For an arc $\alpha \in \Delta$, let $\pi_\Delta(\alpha)$ be the arc defined as follows: if α is the interior edge of a self-folded triangle, then $\pi_\Delta(\alpha)$ is the encircling edge; otherwise $\pi_\Delta(\alpha) := \alpha$. For a non-self-folded triangle t of Δ ,

we define a matrix $B(t) = (b_{\alpha\beta}^\Delta(t))_{\alpha,\beta \in \Delta}$ by

$$b_{\alpha\beta}^\Delta(t) := \begin{cases} 1 & \text{if } t \text{ has } \pi_\Delta(\alpha) \text{ and } \pi_\Delta(\beta) \text{ as its consecutive edges in the clockwise order,} \\ -1 & \text{if the same holds with the counter-clockwise order,} \\ 0 & \text{otherwise.} \end{cases}$$

Then we set $B^\Delta := \sum_t B(t)$, where t runs over all non-self-folded triangles of Δ . Then the formulae given in Propositions 2.2 and 2.4 are summarized as follows:

Proposition 2.7. *For a flip $(\Delta, \varsigma) \xrightarrow{f_\alpha} (\Delta', \varsigma')$ of tagged triangulations, the followings hold:*

(1) *On the space $\mathcal{L}^a(\Sigma, \mathbb{R})$ of real \mathcal{A} -laminations,*

$$\mathbf{a}_{\kappa'} = -\mathbf{a}_\kappa + \max \left\{ \sum_{\beta \in \Delta} [b_{\kappa\beta}^\Delta]_+ \mathbf{a}_\beta, \sum_{\beta \in \Delta} [-b_{\kappa\beta}^\Delta]_+ \mathbf{a}_\beta \right\}$$

and $\mathbf{a}'_\alpha = \mathbf{a}_\alpha$ for $\alpha \neq \kappa, \kappa'$.

(2) *On the space $\mathcal{L}^x(\Sigma, \mathbb{R})$ of real \mathcal{X} -laminations, $\mathbf{x}_{\kappa'}^{(\Delta', \varsigma')} = -\mathbf{x}_\kappa^{(\Delta', \varsigma')}$ and*

$$\mathbf{x}_\alpha^{(\Delta', \varsigma')} = \mathbf{x}_\alpha^{(\Delta, \varsigma)} - b_{\alpha\kappa}^\Delta \max \left\{ 0, -\text{sgn}(b_{\alpha\kappa}^\Delta) \mathbf{x}_\kappa^{(\Delta', \varsigma')} \right\}$$

for $\alpha \neq \kappa, \kappa'$.

Moreover, the exchange matrices B^Δ and $B^{\Delta'}$ are related by the matrix mutation (A.2) when we regard the κ' -th column as the κ -th column.

Using the exchange matrix, the ensemble map is expressed in the coordinate systems associated with a tagged triangulation (Δ, ς) as

$$p_\Sigma^* \mathbf{x}_\alpha^{(\Delta, \varsigma)} = \sum_\beta b_{\alpha\beta}^\Delta \mathbf{a}_\beta^{(\Delta, \varsigma)}. \quad (2.4)$$

In particular we see that the ensemble map is extended to a map $p_\Sigma : \mathcal{L}^a(\Sigma, \mathbb{R}) \rightarrow \mathcal{L}^x(\Sigma, \mathbb{R})$, whose image coincides with $\mathcal{L}^u(\Sigma, \mathbb{R})$. By its definition, the ensemble map is a principal $\mathbb{R}^{h+|M_\partial|}$ -bundle with respect to the action of the latter introduced in Section 2.1. These relations are summarized in the following ‘‘exact sequence’’:

$$0 \rightarrow \mathbb{R}^{h+|M_\partial|} \rightarrow \mathcal{L}^a(\Sigma, \mathbb{R}) \xrightarrow{p_\Sigma} \mathcal{L}^x(\Sigma, \mathbb{R}) \xrightarrow{\theta} \mathbb{R}^h \rightarrow 0 \quad (2.5)$$

This sequence is indeed induced from an exact sequence in the category of finite rank lattices: see Section 3. All maps in (2.5) are $MC(\Sigma)$ -equivariant.

Notice that we can also regard $\mathcal{L}^u(\Sigma, \mathbb{Q})$ as a subspace of $\mathcal{L}^a(\Sigma, \mathbb{Q})$, which consists of the rational \mathcal{A} -laminations without peripheral leaves (and similarly for their completions). This is clear from their geometric definitions, while from an algebraic point of view, it amounts to consider a section of the ensemble map given by $\bigcap_{m \in P \cup M_\partial} W_m^{-1}(0)$. In other words, the tropicalized Goncharov–Shen potentials provide a splitting

$$\mathcal{L}^a(\Sigma, \mathbb{R}) \cong \mathcal{L}^u(\Sigma, \mathbb{R}) \times \mathbb{R}^{h+|M_\partial|}. \quad (2.6)$$

Correspondingly, the mapping class group action is decomposed as follows:

Proposition 2.8. *Let $\sigma_\Sigma : MC(\Sigma) \rightarrow \mathfrak{S}_{h+|M_\partial|}$ be the permutation action of the mapping class group on the marked points, and $\sigma_\Sigma^{\text{uf}} : MC(\Sigma) \rightarrow \mathfrak{S}_h$ its restriction to the set of punctures.*

(1) *Under the splitting (2.6), the action of $MC(\Sigma)$ on $\mathcal{L}^a(\Sigma, \mathbb{R})$ is splitted into the product of the following two actions:*

- *the induced action on $\mathcal{L}^u(\Sigma, \mathbb{R})$, and*
- *the action σ_Σ on $\mathbb{R}^{h+|M_\partial|}$.*

(2) *We have the following commutative diagram*

$$\begin{array}{ccc} \mathcal{L}^x(\Sigma, \mathbb{R}) & \xrightarrow{\phi^x} & \mathcal{L}^x(\Sigma, \mathbb{R}) \\ \theta \downarrow & & \downarrow \theta \\ \mathbb{R}^h & \xrightarrow{\sigma_\Sigma^{\text{uf}}(\phi)} & \mathbb{R}^h \end{array}$$

for all $\phi \in MC(\Sigma)$.

2.4. Measured foliations. Here we recall a geometric description of the real \mathcal{A} -laminations by *measured foliations* [FLP], which in particular provides us a concrete realization of an irrational point as a geometric object on Σ . A nice survey is found in [CP07]. The *Nielsen–Thurston classification* of mapping classes is originally stated in this setting. Moreover, Proposition 2.14 will be a key ingredient in the study of sign stability of pA mapping classes.

Let Σ be a punctured surface, *i.e.*, a marked surface with empty boundary. Let Σ° be the compact surface obtained from Σ by removing a small open disk D_p around each puncture $p \in P$. Let \mathcal{F} be a foliation on Σ° with isolated singularities, satisfying the following conditions ³:

- each boundary component is a leaf of \mathcal{F} ;
- a singularity of \mathcal{F} must be a k -pronged singularity with $k \geq 3$ in the interior, or $k \geq 2$ on the boundary.

Namely, each point $x \in \Sigma^\circ$ has a neighborhood on which \mathcal{F} restricts to either of the local models shown in Figure 10. A *transverse measure* of \mathcal{F} is an assignment μ of a positive number $\mu(\alpha) > 0$ to each arc $\alpha : [0, 1] \rightarrow \Sigma^\circ$ transverse to the leaves of \mathcal{F} , which satisfies the following conditions:

Horizontal invariance: If two transverse arcs α, β are isotopic through transverse arcs whose endpoints remain in the same leaves, then $\mu(\alpha) = \mu(\beta)$.

Sigma-additivity: If a transverse arc α is a concatenation of a countable family of transverse arcs $(\alpha_i)_{i=1}^\infty$, then $\mu(\alpha) = \sum_{i=1}^\infty \mu(\alpha_i)$.

³We are going to consider the measured foliations with compact support in the terminology of [CP07].




FIGURE 10. Local models of foliations. Top row: around an interior point x . Bottom row: around a boundary point x .

Such a pair (\mathcal{F}, μ) is called a *measured foliation* on Σ° . There is an equivalence relation on measured foliations generated by isotopy and *Whitehead collapses* as illustrated in Figure 11, and the set of the equivalence classes of measured foliations (including the empty foliation \emptyset) is denoted by $\mathcal{MF}(\Sigma)$.




FIGURE 11. Whitehead collapsing

Let us briefly mention to the natural topology on $\mathcal{MF}(\Sigma)$. Let $\mathcal{S}(\Sigma)$ denote the set of homotopy classes of non-peripheral simple closed curves on Σ° . Then it is known that each class $[c] \in \mathcal{S}(\Sigma)$ has a representative $c_{\mathcal{F}} \in [c]$ which attains the minimum measure for μ in that class for each $(\mathcal{F}, \mu) \in \mathcal{MF}(\Sigma)$. Moreover if two measured foliations (\mathcal{F}_1, μ_1) and (\mathcal{F}_2, μ_2) are equivalent, then we have $\mu_1(c_{\mathcal{F}_1}) = \mu_2(c_{\mathcal{F}_2})$. See [FLP, Section 5.3] for the proofs of these statements. Thus we get a well-defined map

$$I_*^{\text{MF}} : \mathcal{MF}(\Sigma) \rightarrow \mathbb{R}_{>0}^{\mathcal{S}(\Sigma)}, \quad [\mathcal{F}, \mu] \mapsto I_{(\mathcal{F}, \mu)}^{\text{MF}}$$

with $I_{(\mathcal{F}, \mu)}^{\text{MF}}([c]) := \mu(c_{\mathcal{F}})$ for $[c] \in \mathcal{S}(\Sigma)$. The empty foliation is identified with the 0-vector in the image. The map I_*^{MF} is known to be injective ([FLP, Theorem 6.13]), and hence it pulls-back the weak topology on $\mathbb{R}_{>0}^{\mathcal{S}(\Sigma)}$ to $\mathcal{MF}(\Sigma)$.

Enlarging procedure. The set $\mathcal{L}^u(\Sigma, \mathbb{Q})$ can be regarded as a subset of $\mathcal{MF}(\Sigma)$, as follows. The *enlarging procedure* explained in [FLP, Section 5.4] gives a way to enlarge a measured foliation (\mathcal{F}_0, μ_0) defined on a compact sub-surface $S_0 \subset \text{int } \Sigma^\circ$ to that on the entire surface Σ° . Roughly speaking, it is done by choosing a fatgraph spine of the complement of S_0 , and stretching (\mathcal{F}_0, μ_0) without changing the measure until its boundary leaves are “squeezed” into the fatgraph spine piecewise-smoothly. The equivalence class of the resulting measured foliation does not depend on the choice of a fatgraph spine. See [FLP, Section 5.4] for a detail.

Let $L = \bigcup_{j=1}^n (\gamma_j, w_j) \in \mathcal{L}^u(\Sigma, \mathbb{Q})$ be a rational \mathcal{A} -lamination without peripheral leaves. Define S_0 to be the union of mutually disjoint tubular neighborhoods $\mathcal{N}(\gamma_j)$ of the curves γ_j for $j = 1, \dots, n$. Let \mathcal{F}_0 be the non-singular foliation of S_0 by closed curves parallel to γ_j . Setting the measure of the transverse arc connecting the two boundary components of the annulus $\mathcal{N}(\gamma_j)$ to be the weight $w_j \in \mathbb{Q}_{>0}$, we get a measured foliation (\mathcal{F}_0, μ_0) on S_0 . Then using the enlarging procedure explained above, we get a measured foliation $(\mathcal{F}, \mu) = (\mathcal{F}, \mu)_L$ on Σ° . Thus we get a well-defined map

$$\mathcal{L}^u(\Sigma, \mathbb{Q}) \rightarrow \mathcal{MF}(\Sigma), \quad L \mapsto [(\mathcal{F}, \mu)_L]. \quad (2.7)$$

Decorated measured foliations. In order to include a general rational \mathcal{A} -lamination, we define the space of *decorated measured foliations* to be the trivial bundle $\widetilde{\mathcal{MF}}(\Sigma) := \mathcal{MF}(\Sigma) \times \mathbb{R}^P$ [PP93]. One can think of an element $([\mathcal{F}, \mu], (c_p)_{p \in P}) \in \widetilde{\mathcal{MF}}(\Sigma)$ as a measured foliation on the original punctured surface Σ which restricts to $[\mathcal{F}, \mu]$ on Σ° and has non-singular peripheral leaves foliating the punctured disk $D_p \setminus \{p\}$, and the “measure” of a transverse arc connecting the puncture p to the boundary of D_p is given by c_p . Note that c_p can take a negative value. Then we have a map

$$j_{\mathbb{Q}}^{\text{MF}} : \mathcal{L}^u(\Sigma, \mathbb{Q}) \rightarrow \widetilde{\mathcal{MF}}(\Sigma), \quad L \mapsto ([(\mathcal{F}, \mu)_L], (W_p(L)/2)_{p \in P}). \quad (2.8)$$

We have the following coordinate system on $\widetilde{\mathcal{MF}}(\Sigma)$ associated with an ideal triangulation Δ of Σ . Note that $\Delta_{\text{uf}} = \Delta$, since Σ is now a punctured surface. For an arc $\alpha \in \Delta$ and a decorated measured foliation $([\mathcal{F}, \mu], \mathbf{c} = (c_p)_p) \in \widetilde{\mathcal{MF}}(\Sigma)$, define

$$\tilde{\mathbf{a}}_\alpha([\mathcal{F}, \mu], \mathbf{c}) := \frac{1}{2}(I_{(\mathcal{F}, \mu)}^{\text{MF}}(\alpha) + c_{p_+(\alpha)} + c_{p_-(\alpha)}).$$

Here $p_\pm(\alpha) \in P$ denotes the punctures to which α incident (may not be distinct). Then we get a coordinate system $\tilde{\mathbf{a}}_\Delta := (\tilde{\mathbf{a}}_\alpha)_{\alpha \in \Delta} : \widetilde{\mathcal{MF}}(\Sigma) \rightarrow \mathbb{R}^\Delta$.

Theorem 2.9 (Papadopoulos–Penner [PP93]). *The map $\tilde{\mathbf{a}}_\Delta : \widetilde{\mathcal{MF}}(\Sigma) \xrightarrow{\sim} \mathbb{R}^\Delta$ gives a homeomorphism for any ideal triangulation Δ of Σ . Moreover the following diagram commutes:*

$$\begin{array}{ccc} \mathcal{L}^u(\Sigma, \mathbb{Q}) & \xrightarrow{j_{\mathbb{Q}}^{\text{MF}}} & \widetilde{\mathcal{MF}}(\Sigma) \\ & \searrow a_\Delta & \downarrow \tilde{\mathbf{a}}_\Delta \\ & & \mathbb{R}^\Delta. \end{array}$$

In particular, the map (2.8) extends to a homeomorphism $j^{\text{MF}} : \mathcal{L}^a(\Sigma, \mathbb{R}) \xrightarrow{\sim} \widetilde{\mathcal{MF}}(\Sigma)$, whose restriction gives a homeomorphism $\mathcal{L}^u(\Sigma, \mathbb{R}) \xrightarrow{\sim} \mathcal{MF}(\Sigma)$.

In particular, we can identify the two functions \mathbf{a}_α and $\tilde{\mathbf{a}}_\alpha$ via the homeomorphism $\mathcal{L}^a(\Sigma, \mathbb{R}) \cong \widetilde{\mathcal{MF}}(\Sigma)$. A crucial point in the proof of Theorem 2.9 is the existence of a representative $(\mathcal{F}_\Delta, \mu_\Delta)$ in each equivalence class $[\mathcal{F}, \mu] \in \mathcal{MF}(\Sigma)$ associated with an ideal triangulation Δ , as shown in Figure 12. It is a variant of the representative associated with a “train track” which is useful in the theory of measured foliations: see [PP93] and [PH] for details.




FIGURE 12. Local model of $(\mathcal{F}_\Delta, \mu_\Delta)$.

Shear coordinates of measured foliations. Using the representative associated with an ideal triangulation, we can also describe the coordinates \mathbf{x}_Δ on $\mathcal{L}^u(\Sigma, \mathbb{R})$ in terms of measured foliations. Let $(\mathcal{F}_\Delta, \mu_\Delta)$ be a representative associated with Δ of an equivalence




FIGURE 13. Foliation shear coordinate.

class $[\mathcal{F}, \mu] \in \mathcal{MF}(\Sigma)$. For an edge $\alpha \in \Delta_{\text{uf}}$, each of the two triangles sharing α contains a singularity near α . Define $\tilde{\mathbf{x}}_\alpha([\mathcal{F}, \mu])$ to be the signed μ_Δ -measure of a transverse arc connecting the two singular leaves that emanate from these singular points, where the sign is positive (resp. negative) if one of the singular leaves is found on the left (resp. right) side of the other one. The resulting function $\tilde{\mathbf{x}}_\alpha : \mathcal{MF}(\Sigma) \rightarrow \mathbb{R}$ is called the *foliation-shear coordinate* along α .

Note that any measured foliation in the class $[\mathcal{F}, \mu]$ only with trivalent (resp. univalent) singularities in the interior (resp. on the boundary) satisfying $\tilde{x}_\alpha([\mathcal{F}, \mu]) = 0$ for some $\alpha \in \Delta_{\text{uf}}$ necessarily have a segment joining two singular points, which is called a *saddle connection*. Indeed, the representative $(\mathcal{F}_\Delta, \mu_\Delta)$ has such a segment by the definition of the foliation-shear coordinates and the Whitehead move preserves this property. The following lemma is clear from the definitions:

Lemma 2.10. *The following diagram commutes:*

$$\begin{array}{ccc} \mathcal{L}^u(\Sigma, \mathbb{Q}) & \xrightarrow{j_{\mathbb{Q}}^{\text{MF}}} & \mathcal{MF}(\Sigma) \\ & \searrow \times_\Delta & \downarrow \tilde{x}_\Delta \\ & & \mathbb{R}^\Delta. \end{array}$$

Here the horizontal map is (2.7).

In particular, we can identify the two functions x_α and \tilde{x}_α via the homeomorphism $j^{\text{MF}} : \mathcal{L}^u(\Sigma, \mathbb{R}) \xrightarrow{\sim} \mathcal{MF}(\Sigma)$.

Nielsen–Thurston classification of mapping classes. Let Σ be a punctured surface. Note that the group $\mathbb{R}_{>0}$ acts on $\mathcal{MF}(\Sigma)$ by multiplying a common factor to the measures of transverse arcs. A mapping class $\phi \in MC(\Sigma)$ is said to be

- *periodic* if it is finite order;
- *reducible* if it fixes a *multicurve*, which is the isotopy class of a collection of mutually disjoint simple closed curves in Σ neither peripheral nor contractible;
- *pseudo-Anosov* (“*pA*” for short) if there exists a real number $\lambda > 1$ and two measured foliations $(\mathcal{F}_\pm, \mu_\pm)$ on Σ which shares the singular points and transverse to each other off the singularities, satisfying $\phi^{\pm 1}([\mathcal{F}_\pm, \mu_\pm]) = \lambda^{\pm 1}[\mathcal{F}_\pm, \mu_\pm]$.

Note that a multicurve can be regarded as a rational \mathcal{U} -lamination with weight 1. The numerical invariant $\lambda > 1$ associated with a pseudo-Anosov mapping class ϕ is called the *stretch factor*. We call the pair $([\mathcal{F}_+, \mu_+], [\mathcal{F}_-, \mu_-])$ the *pA pair* of ϕ .

Theorem 2.11 (Nielsen–Thurston classification, e.g. [FLP]). *Each mapping class on a punctured surface is either periodic, reducible or pseudo-Anosov. Moreover, a pseudo-Anosov mapping class is neither periodic nor reducible.*

Indeed, this classification theorem is based on Brouwer’s fixed point theorem, applied to the action of a mapping class on the Thurston compactification of the Teichmüller space of Σ . The boundary is given by the space $\mathcal{PMF}(\Sigma) := (\mathcal{MF}(\Sigma) \setminus \{0\})/\mathbb{R}_{>0}$ of projective measured foliations. A mapping class is pseudo-Anosov if and only if it fixes the projective class of an *arational* foliation *i.e.*, a measured foliation with no closed curves which is the union of segments joining singular points (such a segment is called a *saddle connection*). In this case, both of the pA pair are arational.

Example 2.12. When Σ is a once-punctured torus, the mapping class group is isomorphic to $PSL_2(\mathbb{Z}) = SL_2(\mathbb{Z})/\{\pm 1\}$, where the quotient by $\{\pm 1\}$ corresponds to that by the hyperelliptic involution. In this case, the Nielsen–Thurston classification boils down to the well-known trichotomy elliptic/parabolic/hyperbolic for Möbius transformations which act on the compactified upper half plane $\overline{\mathbb{H}^2} = \mathbb{H}^2 \cup S_\infty^1$. The space $\mathcal{PMF}(\Sigma)$ is identified with $S_\infty^1 = \mathbb{R} \cup \{\infty\}$, where arational foliations correspond to irrational numbers.

We are going to characterize the classes of measured foliations appearing in the Nielsen–Thurston classification in terms of the shear coordinates.

Lemma 2.13 (e.g. [FLP, Corollary 9.2]). *If (\mathcal{F}, μ) is an arational foliation on Σ , then there exists an equivalent measured foliation (\mathcal{F}', μ') only with trivalent (resp. univalent) singularities in the interior (resp. on the boundary), and with no saddle connections. Such a measured foliation is unique up to isotopy in the equivalence class.*

As a consequence, we get the following:

Proposition 2.14. *Let $[\mathcal{F}, \mu] \in \mathcal{MF}(\Sigma)$ be a class of a measured foliation on a punctured surface Σ .*


- (1) *If the class $[\mathcal{F}, \mu]$ admits an arational representative, then it satisfies $\tilde{\chi}_\alpha([\mathcal{F}, \mu]) \neq 0$ for any ideal triangulation Δ and any arc $\alpha \in \Delta$.*
- (2) *If the class $[\mathcal{F}, \mu]$ lies in the image of the embedding (2.7), then it satisfies $\tilde{\chi}_\alpha([\mathcal{F}, \mu]) = 0$ for some ideal triangulation Δ and an arc $\alpha \in \Delta$. In particular this holds for a multicurve.*

Proof. (1): As noted before, the condition $\tilde{\chi}_\alpha([\mathcal{F}, \mu]) = 0$ for some $\alpha \in \Delta$ implies the existence of a saddle connection. Then the above lemma implies that such a measured foliation is never arational. Thus the part (1) is proved.

(2): Let $L \in \mathcal{L}^u(\Sigma, \mathbb{Q})$ be a rational lamination on Σ , and take a pants decomposition \mathcal{R} of Σ which contains the support of L . Then there is a component P of $\Sigma \setminus \bigcup \mathcal{R}$ which contains some punctures. In particular, one can find an ideal arc $\alpha \subset P \subset \Sigma$ both of whose endpoints are the punctures in P . See Figure 14. Let Δ be an ideal triangulation containing α . Then we have $\tilde{\chi}_\alpha(j_{\mathbb{Q}}^{\text{MF}}(L)) = \chi_\alpha(L) = 0$ by Lemma 2.10, since the support of L and α are disjoint. \blacksquare

Definition 2.15. We say that a class $[\mathcal{F}, \mu] \in \mathcal{MF}(\Sigma)$ of a measured foliation on a punctured surface Σ is \mathcal{X} -filling if it satisfies $\tilde{\chi}_\alpha([\mathcal{F}, \mu]) \neq 0$ for any ideal triangulation Δ and any arc $\alpha \in \Delta$.

2.5. Measured geodesic laminations. As a geometric model of the real \mathcal{X} -laminations, we recall the *measured geodesic laminations*. See [PH] for details. We follow the description given by [BB09]. This model is suited for the cut-and-paste operations of marked surfaces, which are useful in Section 7 and the accompanying paper [IK20].

FIGURE 14. Examples of the ideal arc α .

Let Σ be a punctured surface. Fix a complete hyperbolic structure of finite area on the bordered surface $\Sigma^\circ = \Sigma \setminus \bigsqcup_{p \in P} D_p$ with geodesic boundary, and denote by F the resulting hyperbolic surface. More precisely, it is such a hyperbolic surface F together with a homeomorphism marking $\Sigma^\circ \rightarrow F$. Note that each puncture $p \in P$ in Σ corresponds to a geodesic boundary $\partial_p \subset \partial F$. A *geodesic lamination* on F is a closed subset $G \subset F$ equipped with a partition by mutually disjoint complete geodesics. A *transverse measure* μ on G is an assignment of a number $\mu(\alpha) \geq 0$ to each transverse arc $\alpha \subset F$ to G satisfying the following conditions:

- (1) $\mu(\alpha) > 0$ if and only if $\alpha \cap G \neq \emptyset$.
- (2) If α_1 and α_2 are isotopic through transverse arcs, then $\mu(\alpha_1) = \mu(\alpha_2)$.
- (3) $\mu(\alpha)$ is σ -additive in the sense that $\mu(\alpha) = \sum_{i=1}^{\infty} \mu(\alpha_i)$ for any countable partition $\alpha = \bigcup_{i=1}^{\infty} \alpha_i$.

Here a transverse arc to G is a simple arc α such that $\partial\alpha \subset \text{int } F \setminus G$ and transverse to each leaf of G . Such a pair (G, μ) is called a *measured geodesic lamination* on F . Let $\widehat{\mathcal{ML}}(F)$ denote the space of measured geodesic laminations on F . Here are some basic results on the structure of a measured lamination:

Lemma 2.16 ([BKS11, Lemma 2.1]). *Let G be a geodesic lamination on F . Then for each boundary component $\partial_p \subset \partial F$ for $p \in P$, there exists an ϵ -neighborhood U_p such that every leaf of G intersecting U_p must spiral around ∂_p (Figure 15). Moreover, the leaves in $U_p \cap G$ are locally isolated.*

We say that a geodesic lamination G is *compact* if it is disjoint from a neighborhood of the boundary of F .

Lemma 2.17 ([BKS11, Lemma 2.2], [PH, Corollary 1.7.3]). *Any measured lamination (G, μ) on F splits as the union of sub-laminations*

$$G = B \cup S \cup G_1 \cup \cdots \cup G_k,$$

where B is the union of leaves that enter any neighborhood of the boundary of F , S is a finite union of closed geodesics, and G_j is a compact lamination whose leaves are bi-infinite geodesics which are dense in G_j for $j = 1, \dots, k$.




FIGURE 15. Left: Two spiralling patterns around ∂_p . Right: Lifts in the universal cover $\tilde{F} = \mathbb{H}^2$. Each region separated by dashed lines is a fundamental domain.

The compact part $S \cup G_1 \cup \cdots \cup G_k$ is classically well-studied: for example, see [PH, CB].

Note that each leaf g in the non-compact part B determines a properly embedded arc in the surface $\overline{F \setminus \bigcup_{p \in P} U_p}$. We call its homotopy class the *homotopy class of g* , which is independent of the neighborhoods U_p in Lemma 2.16. We define the signature $\sigma_G : P \rightarrow \{+, 0, -\}$ of G by $\sigma_G(p) := +$ (resp. $\sigma_G(p) := -$) if some leaves of G spiral positively (resp. negatively) as in the top (resp. bottom) in Figure 15 around the boundary component ∂_p , and $\sigma_G(p) := 0$ otherwise.

Lemma 2.18 ([BKS11, Lemma 2.3], [BB09, Proposition 3.18]). *The support of the non-compact part B of a measured geodesic lamination (G, μ) on F is uniquely determined by its homotopy class and the signature σ_G .*

This lemma tells us that the non-compact part of any measured geodesic lamination is controlled by a combinatorial data similar to that we studied in Section 2.2. In particular, every rational \mathcal{X} -lamination gives rise to a measured geodesic lamination on F :

Corollary 2.19. *Let F be any hyperbolic structure on Σ° as in the beginning of the current subsection. Then we have a canonical inclusion*

$$j_{\mathbb{Q}}^{\text{ML}}(F) : \mathcal{L}^x(\Sigma, \mathbb{Q}) \hookrightarrow \widehat{\mathcal{ML}}(F). \quad (2.9)$$

Proof. Let $\widehat{L} = (L, \sigma_L)$ be a rational \mathcal{X} -lamination, where $L = \{(\gamma_i, w_i)\}_i$ is the underlying collection of rational weighted curves. For each closed leaf γ_i , take its unique geodesic representative g_i in F . They give a compact geodesic lamination. Each ideal leaf γ_i

gives a unique bi-infinite geodesic g_i in F , whose spiralling direction is determined by the signature σ_L . Let $j_{\mathbb{Q}}^{\text{ML}}(\widehat{L})$ be the measured geodesic lamination whose support is the union of the geodesics g_i , and the measure assigns the number w_i to an arc transversely intersect only with the leaf g_i . The map (2.9) is injective by Lemma 2.18. \blacksquare

We are going to show that the inclusion extends to a bijection $\mathcal{L}^x(\Sigma, \mathbb{R}) \cong \widehat{\mathcal{ML}}(F)$. For this, we need to discuss the topology on the latter space.

Recall the set $\mathcal{S}(\Sigma)$ of homotopy classes of non-peripheral simple closed curves on Σ° , and consider the (*enhanced*) *marked measure spectrum*

$$I_*^{\text{ML}} : \widehat{\mathcal{ML}}(F) \rightarrow \mathbb{R}^P \times \mathbb{R}_{\geq 0}^{\mathcal{S}(\Sigma)}, \quad (G, \mu) \mapsto I_{(G, \mu)}^{\text{ML}} \quad (2.10)$$

defined by

$$I_{(G, \mu)}^{\text{ML}}(\alpha) := \inf_{c \in \alpha} \mu(c)$$

for $\alpha \in \mathcal{S}(\Sigma)$ and

$$I_{(G, \mu)}^{\text{ML}}(p) := \sigma_G(p) \cdot \inf_{c \in [\partial_p]} \mu(c)$$

for $p \in P$. Here $\mu(c)$ means the measure of the curve in F obtained by applying the marking homeomorphism $\Sigma^\circ \rightarrow F$ to $c \subset \Sigma^\circ$. We consider the weak topology on the space $\mathbb{R}^P \times \mathbb{R}_{> 0}^{\mathcal{S}(\Sigma)}$. The following is a slight reformulation of [BB09, Proposition 3.22]:

Proposition 2.20. (1) *The map (2.10) is injective, and the image is homeomorphic to $\mathbb{R}^{6g-6+3h}$.*

(2) *For every pants decomposition \mathcal{R} of Σ° , associated is a finite subset $\mathcal{S}_{\mathcal{R}} \subset \mathcal{S}(\Sigma)$ such that the composition*

$$I_*^{\mathcal{R}} : \widehat{\mathcal{ML}}(F) \xrightarrow{I_*^{\text{ML}}} \mathbb{R}^P \times \mathbb{R}_{\geq 0}^{\mathcal{S}(\Sigma)} \rightarrow \mathbb{R}^P \times \mathbb{R}_{\geq 0}^{\mathcal{S}_{\mathcal{R}}}$$

with the coordinate projection is still injective. By varying the pants decomposition \mathcal{R} , we get a PL atlas on $\widehat{\mathcal{ML}}(F)$.

(3) *If we endow $\widehat{\mathcal{ML}}(F)$ with the subspace topology, then the image of the inclusion (2.9) is dense.*

Now we have:

Theorem 2.21. *The inclusion (2.9) extends to a PL isomorphism*

$$j^{\text{ML}}(F) : \mathcal{L}^x(\Sigma, \mathbb{R}) \xrightarrow{\sim} \widehat{\mathcal{ML}}(F). \quad (2.11)$$

Although this theorem seems to be well-known to specialists, we give a proof here for completeness.

Proof. We claim that the composition $I_*^{\mathcal{R}} \circ j_{\mathbb{Q}}^{\text{ML}}(F)$ is a PL map for any pants decomposition \mathcal{R} . For this, consider the intersection pairing [FG06a, Section 12]

$$\mathcal{I}_{\Sigma} : \mathcal{L}^a(\Sigma, \mathbb{Q}) \times \mathcal{L}^x(\Sigma, \mathbb{Q}) \rightarrow \mathbb{Q}$$

given by the weighted sum of the minimal intersection numbers between the leaves. Fixing an \mathcal{A} -lamination $L \in \mathcal{L}^a(\Sigma, \mathbb{Q})$, the function $\mathcal{I}_L := \mathcal{I}_\Sigma(L, -)$ is a PL function by [FG06a, Theorem 12.2 (2)], as the tropical limit of the trace function along L . Regarding $\alpha \in \mathcal{S}(\Sigma)$ as a rational \mathcal{A} -lamination, the following diagram commutes:

$$\begin{array}{ccc} \mathcal{L}^x(\Sigma, \mathbb{Q}) & & \\ j_{\mathbb{Q}}^{\text{ML}}(F) \downarrow & \searrow \mathcal{I}_\alpha & \\ \widehat{\mathcal{ML}}(F) & \xrightarrow{I_*^{\text{ML}}(\alpha)} & \mathbb{R}. \end{array}$$

We have a similar diagram for boundary curves, regarding them as rational \mathcal{A} -laminations with weight -1 . Hence the map $I_*^{\mathcal{R}} \circ j_{\mathbb{Q}}^{\text{ML}}(F)$ is PL for any \mathcal{R} , and in particular Lipschitz continuous. Thus it can be extended to a PL homeomorphism from the completion $\mathcal{L}^x(\Sigma, \mathbb{R})$, which is surjective onto the image $I_*^{\mathcal{R}}(\widehat{\mathcal{ML}}(F))$ by Brouwer's invariance of domain. Since the inverse of a PL map is also PL, the assertion follows. \blacksquare

Remark 2.22. The restriction of $j^{\text{ML}}(F)$ gives a PL homeomorphism from $\mathcal{L}^u(\Sigma, \mathbb{R}) \subset \mathcal{L}^x(\Sigma, \mathbb{R})$ onto the subspace $\mathcal{ML}_0(F) \subset \widehat{\mathcal{ML}}(F)$ of compact measured laminations. Comparing with the discussion in the previous section, we get the commutative diagram for any hyperbolic structure F :

$$\begin{array}{ccc} & \mathcal{MF}(\Sigma) & \\ j^{\text{MF}} \nearrow & & \searrow I_*^{\text{MF}} \\ \mathcal{L}^u(\Sigma, \mathbb{R}) & & \mathbb{R}_{\geq 0}^{\mathcal{S}(\Sigma)} \\ j^{\text{ML}}(F) \searrow & & \nearrow I_*^{\text{ML}} \\ & \mathcal{ML}_0(F) & \end{array}$$

The well-known correspondence $\mathcal{MF}(\Sigma) \cong \mathcal{ML}_0(F)$ (see, for instance, [PH, Epilogue]) fits into this diagram.

Remark 2.23. In view of Theorem 2.21 and the previous remark, Lemma 2.17 implies that any real \mathcal{X} -lamination $\widehat{L} \in \mathcal{L}^x(\Sigma, \mathbb{R})$ can be uniquely decomposed as $\widehat{L} = L_u \sqcup \widehat{L}_+$, where $L_u \in \mathcal{L}^u(\Sigma, \mathbb{R})$ and \widehat{L}_+ is a collection of real weighted ideal arcs. With respect to this decomposition, we have

$$\mathbf{x}_\alpha^{(\Delta, \varsigma)}(\widehat{L}) = \mathbf{x}_\alpha^{(\Delta, \varsigma)}(L_u) + \mathbf{x}_\alpha^{(\Delta, \varsigma)}(\widehat{L}_+)$$

for any tagged triangulation (Δ, ς) and $\alpha \in \Delta$. Indeed, the formula holds for all rational \mathcal{X} -laminations by the leaf-wise definition of the coordinates, and thus holds in general by continuity. Also note that \widehat{L}_+ belongs to the geometric realization of the Fock–Goncharov cluster complex [GHKK18]: see Section 5.3 below.

General marked surfaces. Let Σ be a marked surface possibly with non-empty boundary. Let $D^\times\Sigma$ denote the punctured surface obtained by gluing two copies of Σ along their corresponding boundary components, so that each pair of special points matches to produce a new puncture of $D^\times\Sigma$. The set of punctures of $D^\times\Sigma$ is given by $D^\times P := \{(p_+, p_-) \mid p \in P\} \sqcup \{p_m \mid m \in M_\partial\}$. The surface $D^\times\Sigma$ is naturally equipped with an orientation-reversing involution $\iota : D^\times\Sigma \rightarrow D^\times\Sigma$, which interchanges the two copies of Σ .

Let $D^\times F$ be a complete hyperbolic structure of finite area on the double $D^\times\Sigma$ with geodesic boundary, which is ι -invariant. Then the restriction F of $D^\times F$ to a copy of Σ is a hyperbolic structure where the punctured boundary $\partial\Sigma \setminus M_\partial$ is represented totally geodesically [Pen04]. Let $\widehat{\mathcal{ML}}(F) \subset \widehat{\mathcal{ML}}(D^\times F)$ be the subset of ι -invariant measured geodesic laminations. Notice that the ι -invariance prevents each leaf from spiralling around any boundary components arising from the punctures p_m , $m \in M_\partial$. Then one can follow the previous discussions with the ι -invariance in mind, and get the following:

Theorem 2.24. *For any marked surface Σ and an ι -invariant hyperbolic structure $D^\times F$ on $D^\times\Sigma$, we have a PL isomorphism*

$$j^{\text{ML}}(F) : \mathcal{L}^x(\Sigma, \mathbb{R}) \xrightarrow{\sim} \widehat{\mathcal{ML}}(F).$$

Remark 2.25. As in the punctured case, let $\mathcal{ML}_0(F) \subset \widehat{\mathcal{ML}}(F)$ denote the subspace consisting of ι -invariant compact measured laminations on $D^\times\Sigma$. Then similarly to Remark 2.22, $j^{\text{ML}}(F)$ restricts to a PL isomorphism $\mathcal{L}^u(\Sigma, \mathbb{R}) \rightarrow \mathcal{ML}_0(F)$.

We also have a decomposition $\widehat{L} = L_u \sqcup \widehat{L}^+$ of real \mathcal{X} -laminations similar to the one mentioned in Remark 2.23 for a general marked surfaces, where $L_u \in \mathcal{ML}_0(F)$ and the lamination \widehat{L}_+ belongs to the geometric realization of the Fock–Goncharov cluster complex, which is the restriction to Σ of a collection of real weighted ideal arcs in the double $D^\times\Sigma$.

3. MUTATION LOOPS AND THEIR REPRESENTATION PATHS IN THE EXCHANGE GRAPH

Here we introduce the notion of “mutation sequences” more carefully than the literature, in order to study their sign stability correctly. For the basic concepts related to the cluster ensembles, see Appendix A. In our formulation, they are understood as edge paths in a graph \mathbb{E}_I defined below. As a quotient graph of \mathbb{E}_I the *labeled exchange graph* $\mathbb{E}\text{ch}_s$ is defined, where the edge paths in the latter are intrinsic to the given seed pattern.

An element of the cluster modular group, called a *mutation loop*, is a certain equivalence class of an edge path in the labeled exchange graph. A quick look at the type A_2 example Example 3.11 is recommended.

3.1. The labeled exchange graph.

Labeled seed patterns. Fix a finite set $I = I_{\text{uf}} \sqcup I_{\text{f}}$ as in Appendix A. We are going to include an effect of permutations of indices in the seed pattern. Let \mathfrak{S}_I denote the group which consists of permutations which do not mixup I_{uf} and I_{f} :

$$\mathfrak{S}_I := \mathfrak{S}_{I_{\text{uf}}} \times \mathfrak{S}_{I_{\text{f}}}.$$

Let \mathbb{E}_I be the graph with

- vertices given by the pairs (t, σ) for $t \in \mathbb{T}_{I_{\text{uf}}}$ and $\sigma \in \mathfrak{S}_I$, and
- edges of the following two types:
 - (*horizontal edges*) For each $t \xrightarrow{k} t'$ in $\mathbb{T}_{I_{\text{uf}}}$ and $\sigma \in \mathfrak{S}_I$, there is an edge $(t, \sigma) \xrightarrow{\sigma(k)} (t', \sigma)$.
 - (*vertical edges*) For each vertex $t \in \mathbb{T}_{I_{\text{uf}}}$, $\sigma \in \mathfrak{S}_I$ and a transposition $(i j) \in \mathfrak{S}_I$, there is an edge $(t, \sigma) \xrightarrow{(i j)} (t, (i j)\sigma)$.

We have a natural projection $\pi_{\mathbb{T}} : \mathbb{E}_I \rightarrow \mathbb{T}_{I_{\text{uf}}}$ forgetting the vertical edges and mapping a horizontal edge $(t, \sigma) \xrightarrow{\sigma(k)} (t', \sigma)$ to an edge $t \xrightarrow{k} t'$, whose fiber $\pi^{-1}(t)$ is isomorphic to the Cayley graph of \mathfrak{S}_I with the generating set as the transpositions in \mathfrak{S}_I . A seed pattern $\mathbf{s} : \mathbb{T}_{I_{\text{uf}}} \ni t \mapsto (N^{(t)}, B^{(t)})$ lifts to a *labeled seed pattern* $\tilde{\mathbf{s}} : \mathbb{E}_I \ni (t, \sigma) \mapsto (N^{(t, \sigma)}, B^{(t, \sigma)})$ defined as

$$N^{(t, \sigma)} := \bigoplus_{i \in I} \mathbb{Z}e_i^{(t, \sigma)}, \quad B^{(t, \sigma)} := \sigma.B^{(t)}.$$

Here $\sigma.A := (a_{\sigma^{-1}(i), \sigma^{-1}(j)})_{i, j \in I}$ for an $I \times I$ -matrix $A = (a_{ij})_{i, j \in I}$. As an ordinary seed pattern, $N^{(t, \sigma)}$ has a skew-symmetric bilinear form $\{-, -\} : N^{(t, \sigma)} \times N^{(t, \sigma)} \rightarrow \mathbb{Z}$ defined by $\{e_{\sigma(i)}^{(t, \sigma)}, e_{\sigma(j)}^{(t, \sigma)}\} := b_{ij}^{(t)}$. By definition, we have a natural seed isomorphism

$$\psi_{\sigma}^* : (N^{(t)}, B^{(t)}) \longrightarrow (N^{(t, \sigma)}, B^{(t, \sigma)}); \quad e_i^{(t)} \mapsto e_{\sigma(i)}^{(t, \sigma)}.$$

The two lattices $N^{(t, \sigma)}$ and $N^{(t, \sigma')}$ with the common underlying vertex $t \in \mathbb{T}_{I_{\text{uf}}}$ are related by the seed isomorphism

$$j_{\sigma'\sigma^{-1}}^* : (N^{(t, \sigma')}, B^{(t, \sigma')}) \longrightarrow (N^{(t, \sigma)}, B^{(t, \sigma)}); \quad e_{\sigma'(i)}^{(t, \sigma')} \mapsto e_{\sigma(i)}^{(t, \sigma)}.$$

When $\sigma' = \rho\sigma$ for some $\rho \in \mathfrak{S}_I$, we simply write $\rho^* := j_{\sigma'\sigma^{-1}}^*$. In particular, we have $\rho^*(e_k^{(t, \sigma')}) = e_{\rho^{-1}(k)}^{(t, \sigma)}$.

In the same way as the ordinary seed pattern, we define $\mathcal{X}_{(t, \sigma)}^{\text{uf}} := T_{M_{\text{uf}}^{(t, \sigma)}}$ and $\mathcal{A}_{(t, \sigma)} := T_{N^{(t, \sigma)}}$. For each vertex $(t, \sigma) \in \mathbb{E}_I$ and $(z, \mathcal{Z}) = (a, \mathcal{A})$ or $(x, \mathcal{X}^{\text{uf}})$, these tori are related by the following (birational) morphisms:

- (1) for a vertical edge $(t, \sigma) \xrightarrow{(i j)} (t, (i j)\sigma)$, an isomorphism $(i j)^z : \mathcal{Z}_{(t, \sigma)} \xrightarrow{\sim} \mathcal{Z}_{(t, (i j)\sigma)}$ induced by the seed isomorphism $(i j)^*$.

- (2) for a horizontal edge $(t, \sigma) \xrightarrow{\sigma(k)} (t', \sigma)$, a cluster transformation $\mu_{\sigma(k)}^z : \mathcal{Z}_{(t, \sigma)} \rightarrow \mathcal{Z}_{(t', \sigma)}$ defined so that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{Z}_{(t, \sigma)} & \xrightarrow{\mu_{\sigma(k)}^z} & \mathcal{Z}_{(t', \sigma)} \\ \psi_{\sigma}^z \downarrow & & \downarrow \psi_{\sigma}^z \\ \mathcal{Z}_{(t)} & \xrightarrow{\mu_k^z} & \mathcal{Z}_{(t')}. \end{array}$$

Here ψ_{σ}^z is an isomorphism induced by the seed isomorphism ψ_{σ}^* .

Definition 3.1 (the cluster transformations associated with an edge path). For an edge path $\gamma : (t, \sigma) \rightarrow (t', \sigma')$ in \mathbb{E}_I , we define the *cluster transformation* $\mu_{\gamma}^z : \mathcal{Z}_{(t, \sigma)} \rightarrow \mathcal{Z}_{(t', \sigma')}$ associated with γ to be the composition of the birational maps associated to the edges it traverses for $(z, \mathcal{Z}) = (a, \mathcal{A}), (x, \mathcal{X}^{\text{uf}})$.

It only depends on the endpoints of γ , thanks to the following lemma:

Lemma 3.2 (Naturality relation). *For each $k \in I_{\text{uf}}$ and two permutations $\sigma, \sigma' \in \mathfrak{S}_I$ which are related by $\sigma' = \rho\sigma$ for some $\rho \in \mathfrak{S}_I$, we have the following commutative diagram:*

$$\begin{array}{ccc} (t, \sigma) & \xrightarrow{k} & (t', \sigma) \\ \rho \downarrow & & \downarrow \rho \\ (t, \sigma') & \xrightarrow{\rho(k)} & (t', \sigma') \end{array} \quad \xrightarrow{\mathcal{Z}_{\bullet}} \quad \begin{array}{ccc} \mathcal{Z}_{(t, \sigma)} & \xrightarrow{\mu_k^z} & \mathcal{Z}_{(t', \sigma)} \\ \rho^z \downarrow & \circlearrowleft & \downarrow \rho^z \\ \mathcal{Z}_{(t, \sigma')} & \xrightarrow{\mu_{\rho(k)}^z} & \mathcal{Z}_{(t', \sigma')} \end{array}$$

for $(z, \mathcal{Z}) = (a, \mathcal{A}), (x, \mathcal{X}^{\text{uf}})$, which we call the *naturality relation*. Here $(t, \sigma) \xrightarrow{\rho} (t, \sigma')$ is the concatenation of vertical edges $(t, \sigma) \xrightarrow{(i_1 \ j_1)} \dots \xrightarrow{(i_r \ j_r)} (t, \sigma')$ for an arbitrary expression $\rho = (i_r \ j_r) \cdots (i_1 \ j_1)$ in \mathfrak{S}_I , and ρ^z is the isomorphism induced by the seed isomorphism ρ^* .

We call the square in \mathbb{E}_I considered above the *naturality square*. For each $t \in \mathbb{T}_{I_{\text{uf}}}$, we can identify $\mathcal{Z}_{(t)}$ with $\bigcup_{\sigma \in \mathfrak{S}_I} \mathcal{Z}_{(t, \sigma)}$ via ψ_{σ} , where the latter is obtained by patching the tori $\mathcal{Z}_{(t, \sigma)}$ by isomorphisms σ^z . So we can identify $\mathcal{Z}_{\mathbf{s}}$ with $\bigcup_{(t, \sigma) \in \mathbb{E}_I} \mathcal{Z}_{(t, \sigma)}$.

The labeled exchange graph. For two vertices $(t, \sigma), (t', \sigma') \in \mathbb{E}_I$, we consider the following two equivalence relations:

- (1) We say that the vertices $(t, \sigma), (t', \sigma')$ are *\mathbf{s} -equivalent* and write $(t, \sigma) \sim_{\mathbf{s}} (t', \sigma')$ if $B^{(t, \sigma)} = B^{(t', \sigma')}$. This is equivalent to the condition that the following linear map is a seed isomorphism:

$$(N^{(t', \sigma')}, B^{(t', \sigma')}) \rightarrow (N^{(t, \sigma)}, B^{(t, \sigma)}); \quad e_i^{(t', \sigma')} \mapsto e_i^{(t, \sigma)}. \quad (3.1)$$

- (2) We write $(t, \sigma) \sim_{\text{triv}} (t', \sigma')$ if they are \mathbf{s} -equivalent and for some (and any) edge path $\gamma : (t, \sigma) \rightarrow (t', \sigma')$ in \mathbb{E}_I , the composition

$$\mathcal{Z}_{(t, \sigma)} \xrightarrow{\mu_{\gamma}^z} \mathcal{Z}_{(t', \sigma')} \xrightarrow{\sim} \mathcal{Z}_{(t, \sigma)} \quad (3.2)$$

is identity for $(z, \mathcal{Z}) = (a, \mathcal{A}), (x, \mathcal{X}^{\text{uf}})$. Here the second map is induced from the seed isomorphism (3.1). We call such a path γ *trivial*.

We denote the second equivalence class containing (t, σ) by $[t, \sigma]_{\text{triv}}$.

For two oriented edges $e_\nu = ((t_\nu, \sigma_\nu) \text{ --- } (t'_\nu, \sigma'_\nu))$ with $\nu = 1, 2$ in \mathbb{E}_I , we write $e_1 \sim_{\text{triv}} e_2$ if $(t_1, \sigma_1) \sim_{\text{triv}} (t_2, \sigma_2)$ and $(t'_1, \sigma'_1) \sim_{\text{triv}} (t'_2, \sigma'_2)$.

Definition 3.3 (The labeled exchange graph). The quotient graph

$$\mathbb{E}\text{xch}_{\mathbf{s}} := \mathbb{E}_I / \sim_{\text{triv}}$$

is called the *labeled exchange graph* of the seed pattern \mathbf{s} .

Note that each edge in \mathbb{E}_I is projected to an edge in $\mathbb{E}\text{xch}_{\mathbf{s}}$ with distinct endpoints, since two vertices connected by an edge in \mathbb{E}_I are not equivalent under \sim_{triv} . It is not hard to check that the relation \sim_{triv} respects the horizontal/vertical properties of an edge in \mathbb{E}_I , and preserves their labels:

$$\begin{aligned} (1) \text{ Horizontal edges: } & \begin{array}{ccc} (t_1, \sigma_1) & \xrightarrow{\sigma_1(k_1)} & (t'_1, \sigma_1) \\ \wr_{\text{triv}} & & \wr_{\text{triv}} \\ (t_2, \sigma_2) & \xrightarrow{\sigma_2(k_2)} & (t'_2, \sigma_2) \end{array} \implies \sigma_1(k_1) = \sigma_2(k_2) \in I_{\text{uf}}. \\ (2) \text{ Vertical edges: } & \begin{array}{ccc} (t_1, \sigma_1) & \xrightarrow{(i_1 \ j_1)} & (t_1, \sigma'_1) \\ \wr_{\text{triv}} & & \wr_{\text{triv}} \\ (t_2, \sigma_2) & \xrightarrow{(i_2 \ j_2)} & (t_2, \sigma'_2) \end{array} \implies (i_1 \ j_1) = (i_2 \ j_2) \in \mathfrak{S}_I. \end{aligned}$$

Therefore we can roughly say that $\mathbb{E}\text{xch}_{\mathbf{s}}$ is “locally” isomorphic to \mathbb{E}_I together with labeling of edges.

Remark 3.4. Since the relation \sim_{triv} respects the horizontal/vertical properties of an edge in \mathbb{E}_I , the vertical projection $\pi_{\mathbb{T}} : \mathbb{E}_I \rightarrow \mathbb{T}_{I_{\text{uf}}}$ induces a graph projection $\pi_{\mathbb{E}\text{x}} : \mathbb{E}\text{xch}_{\mathbf{s}} \rightarrow \mathbb{E}\text{xch}_{\mathbf{s}}$, and the latter graph $\mathbb{E}\text{xch}_{\mathbf{s}}$ is called the *exchange graph*:

$$\begin{array}{ccc} \mathbb{E}_I & \longrightarrow & \mathbb{E}\text{xch}_{\mathbf{s}} \\ \pi \downarrow & & \downarrow \pi_{\mathbb{E}\text{x}} \\ \mathbb{T}_{I_{\text{uf}}} & \longrightarrow & \mathbb{E}\text{xch}_{\mathbf{s}} \end{array}$$

Here note that the labels on the edges on $\mathbb{E}\text{xch}_{\mathbf{s}}$ do not descend to $\mathbb{E}\text{xch}_{\mathbf{s}}$: see Example 3.11 below.

Parametrization of cluster charts. Here is a way to parametrize the coordinate charts on the cluster varieties by the vertices of the labeled exchange graph $\mathbb{E}\text{xch}_{\mathbf{s}}$. For two vertices $(t, \sigma), (t', \sigma')$ such that $(t, \sigma) \sim_{\text{triv}} (t', \sigma')$, the two tori $\mathcal{Z}_{(t, \sigma)}$ and $\mathcal{Z}_{(t', \sigma')}$ are canonically identified in $\mathcal{Z}_{\mathbf{s}}$ by the cluster transformation μ_{γ}^z for a trivial path $\gamma : (t, \sigma) \rightarrow (t', \sigma')$. Define $\mathcal{Z}_{[t, \sigma]_{\text{triv}}} := \bigcup_{(t', \sigma') \in [t, \sigma]_{\text{triv}}} \mathcal{Z}_{(t', \sigma')}$, where the latter is obtained by patching the tori

by trivial cluster transformations, so that we have

$$\mathcal{Z}_{\mathfrak{s}} = \bigcup_{[t,\sigma]_{\text{triv}} \in \mathbb{E}\text{ch}_{\mathfrak{s}}} \mathcal{Z}_{[t,\sigma]_{\text{triv}}}.$$

Since it is more intrinsic and economical, we often use this parametrization of charts and the edge paths in $\mathbb{E}\text{ch}_{\mathfrak{s}}$ rather than those in \mathbb{E}_I . The graph $\mathbb{E}\text{ch}_{\mathfrak{s}}$ is a finite graph if and only if the seed pattern \mathfrak{s} is of finite type.

From the above discussions, the seed pattern $\tilde{\mathfrak{s}}$ on \mathbb{E}_I also descends to a “seed pattern” $\mathfrak{s}_{\text{Ex}} : \mathbb{E}\text{ch}_{\mathfrak{s}} \ni v \mapsto (N^{(v)}, B^{(v)})$ satisfying $B^{(v)} = \sigma.B^{(t)}$ for some $(t, \sigma) \in v$. We also write $e_i^{(v)} := e_i^{(t,\sigma)}$ and $f_i^{(v)} := f_i^{(t,\sigma)}$ for $i \in I$, and define the corresponding cluster coordinates by $X_i^{(v)} := \text{ch}_{e_i^{(v)}}$ and $A_i^{(v)} := \text{ch}_{f_i^{(v)}}$. For an edge path $\gamma : v_0 \rightarrow v$ in $\mathbb{E}\text{ch}_{\mathfrak{s}}$, the associated cluster transformation

$$\mu_{\gamma}^z : \mathcal{Z}_{(v_0)} \rightarrow \mathcal{Z}_{(v)} \quad (3.3)$$

is well-defined for $(z, \mathcal{Z}) = (a, \mathcal{A}), (x, \mathcal{X}^{\text{uf}})$.

Remark 3.5. Further we can consider a torus $\mathcal{Z}_v := \bigcup_{[t,\sigma]_{\text{triv}} \in \pi_{\text{Ex}}^{-1}(v)} \mathcal{Z}_{[t,\sigma]_{\text{triv}}}$ associated to a vertex $v \in \text{Exch}_{\mathfrak{s}}$, where the identifications are given by the isomorphisms σ^z . The cluster coordinates on the torus \mathcal{Z}_v is unordered. In the relation with the lamination theory, the tori \mathcal{Z}_v (resp. $\mathcal{Z}_{[t,\sigma]_{\text{triv}}}$) correspond to labeled (resp. unlabeled) tagged triangulations. See Section 4.1.

C- and G-matrices. Recall the *C*- and *G*-matrices from Appendix A.2. Fixing a vertex $(t_0, \sigma_0) \in \mathbb{E}_I$, the assignment of these matrices can be extended to \mathbb{E}_I by

$$C_{(t,\sigma)}^{\tilde{\mathfrak{s}};(t_0,\sigma_0)} := P_{\sigma} C_t^{\mathfrak{s};t_0} P_{\sigma_0}^{-1}, \quad G_{(t,\sigma)}^{\tilde{\mathfrak{s}};(t_0,\sigma_0)} := P_{\sigma} G_t^{\mathfrak{s};t_0} P_{\sigma_0}^{-1},$$

where P_{σ} denotes the presentation matrix of $\sigma \in \mathfrak{S}_N$, i.e., $P_{\sigma} = (\delta_{i,\sigma(j)})$. Then the separation formula [FZ07, Proposition 3.13, Corollary 6.4] tells us that the assignments

$$C_v^{v_0} := C_{(t,\sigma)}^{\tilde{\mathfrak{s}};(t_0,\sigma_0)}, \quad G_v^{v_0} := G_{(t,\sigma)}^{\tilde{\mathfrak{s}};(t_0,\sigma_0)}$$

for $v_0 := [t_0, \sigma_0]_{\text{triv}}$, $v = [t, \sigma]_{\text{triv}} \in \mathbb{E}\text{ch}_{\mathfrak{s}}$ are well-defined. This also follows from the relations (A.9), (A.10) and the fact that the tropical sign coincides with the sign at a point in the interior of the cone $\mathcal{C}_{(v_0)}^+$: see Section 5.2 below. From the tropical duality (A.11), we get

$$G_v^{v_0} = \check{C}_v^{v_0}. \quad (3.4)$$

3.2. Mutation loops and the cluster modular group. A mutation loop is defined to be an equivalence class of an edge path in $\mathbb{E}\text{ch}_{\mathfrak{s}}$. Note that the \mathfrak{s} -equivalence can be regarded as an equivalence relation among the vertices of the labeled exchange graph $\mathbb{E}\text{ch}_{\mathfrak{s}}$. Let $\gamma_{\nu} : v_{\nu} \rightarrow v'_{\nu}$ be an edge path in $\mathbb{E}\text{ch}_{\mathfrak{s}}$ such that $v_{\nu} \sim_{\mathfrak{s}} v'_{\nu}$ for $\nu = 1, 2$. We

say that γ_1 and γ_2 are \mathbf{s} -equivalent if there exists path $\delta : v_1 \rightarrow v_2$ such that the following diagram commutes:

$$\begin{array}{ccccc} \mathcal{Z}_{(v_1)} & \xrightarrow{\mu_{\gamma_1}^z} & \mathcal{Z}_{(v'_1)} & \xrightarrow{\sim} & \mathcal{Z}_{(v_1)} \\ \mu_{\delta}^z \downarrow & & & & \downarrow \mu_{\delta}^z \\ \mathcal{Z}_{(v_1)} & \xrightarrow{\mu_{\gamma_2}^z} & \mathcal{Z}_{(v'_2)} & \xrightarrow{\sim} & \mathcal{Z}_{(v_2)}. \end{array} \quad (3.5)$$

Here $(z, \mathcal{Z}) = (a, \mathcal{A}), (x, \mathcal{X}^{\text{uf}})$, $\mu_{\gamma_\nu}^z$ is the associated cluster transformation (3.3) and the isomorphism $\mathcal{Z}_{(v'_\nu)} \xrightarrow{\sim} \mathcal{Z}_{(v_\nu)}$ is induced by the seed isomorphism (3.1) for $\nu = 1, 2$. The \mathbf{s} -equivalence class containing an edge path γ is denoted by $[\gamma]_{\mathbf{s}}$. Note that the commutativity of the diagram (3.5) does not depend on the choice of a path δ by Lemma 3.2.

Definition 3.6 (mutation loops). A *mutation loop* is an \mathbf{s} -equivalence class of an edge path $\gamma : v \rightarrow v'$ in the labeled exchange graph $\text{Exch}_{\mathbf{s}}$ such that $v \sim_{\mathbf{s}} v'$. We call γ a *representation path* of the mutation loop $\phi = [\gamma]_{\mathbf{s}}$.

The concatenation of two paths $\gamma_1 : v_0 \rightarrow v_1$ and $\gamma_2 : v_1 \rightarrow v_2$ is denoted by $\gamma_1 * \gamma_2 : v_0 \rightarrow v_1 \rightarrow v_2$.

Definition 3.7 (cluster modular group). The group $\Gamma_{\mathbf{s}}$ consisting of all the mutation loops is called the *cluster modular group*. Here the multiplication of two mutation loops ϕ_1 and ϕ_2 is defined as $\phi_2 \cdot \phi_1 := [\gamma_1 * \gamma_2]_{\mathbf{s}}$, where γ_ν is a representation path of ϕ_ν for $\nu = 1, 2$ such that the terminal vertex of γ_1 coincides with the initial vertex of γ_2 .

Remark 3.8. Any two representation paths of a mutation loop are related to each other by a finite composition of the following two types of operations.

- (a) A replacement of a path γ_1 with another path γ_2 with the common initial vertex, the path δ in (3.5) being trivial. For example:
 - Deformation through a naturality square (Lemma 3.2). That is, a replacement of a subpath $v \xrightarrow{k} v' \xrightarrow{\rho} \rho.v'$ with the path $v \xrightarrow{\rho} \rho.v \xrightarrow{\rho(k)} \rho.v'$. Here we use the notation $\rho.v = [t, \rho\sigma]_{\text{triv}}$ for a vertex $v = [t, \sigma]_{\text{triv}}$ and a permutation $\rho \in \mathfrak{S}_I$.
 - An elimination or addition of a round trip $\delta' : v \xrightarrow{k} v' \xrightarrow{k} v$.
 - An elimination or addition of a path δ' corresponding to the standard $(h+2)$ -gon relation [FG09].
- (b) A replacement of a path $\gamma_1 : v_1 \rightarrow v'_1$ with another one $\gamma_2 : v_2 \rightarrow v'_2$ with a different initial vertex (i.e., $v_1 \neq v_2$), the paths $\delta : v_1 \xrightarrow{\mathbf{k}} v_2$ and $\delta' : v'_1 \xrightarrow{\mathbf{k}'} v'_2$ satisfying

$$\begin{array}{ccc} \mathcal{Z}_{(v_1)} & \xrightarrow{\mu_{\gamma_1}^z} & \mathcal{Z}_{(v'_1)} \\ \mu_{\delta}^z \downarrow & \circlearrowleft & \downarrow \mu_{\delta'}^z \\ \mathcal{Z}_{(v_2)} & \xrightarrow{\mu_{\gamma_2}^z} & \mathcal{Z}_{(v'_2)} \end{array}$$

being related as $\mathbf{k} = \mathbf{k}'$. In this case, the birational maps $\phi_{(v_1)}^z$ and $\phi_{(v_2)}^z$ are related by the conjugation of the map μ_δ^z .

Since a mutation loop $\phi \in \Gamma_{\mathbf{s}}$ admits a representation path in $\mathbb{E}\text{xch}_{\mathbf{s}}$ which starts from any vertex $v_0 \in \mathbb{E}\text{xch}_{\mathbf{s}}$ by Remark 3.8 (b), the definition of the multiplication of the cluster modular group is well-defined.

The cluster modular group acts on the cluster varieties $\mathcal{X}_{\mathbf{s}}^{\text{uf}}$ and $\mathcal{A}_{\mathbf{s}}$ as automorphisms, as follows. For $\phi \in \Gamma_{\mathbf{s}}$, take its representation path $\gamma : v_0 \rightarrow v$ in $\mathbb{E}\text{xch}_{\mathbf{s}}$. Then we have the following composite of birational isomorphisms:

$$\phi_{(v_0)}^z : \mathcal{Z}_{(v_0)} \xrightarrow{\mu_\gamma^z} \mathcal{Z}_{(v)} \xrightarrow{\sim} \mathcal{Z}_{(v_0)} \quad (3.6)$$

for $(z, \mathcal{Z}) = (a, \mathcal{A}), (x, \mathcal{X}^{\text{uf}})$. Here the second map is induced by the seed isomorphism (3.1). The mutation loop ϕ induces an automorphism ϕ^z on the cluster variety $\mathcal{Z}_{\mathbf{s}}$, which fits into the following commutative diagram:

$$\begin{array}{ccccc} \mathcal{Z}_{(v_0)} & \xrightarrow{\mu_\gamma^z} & \mathcal{Z}_{(v)} & \xrightarrow{\sim} & \mathcal{Z}_{(v_0)} \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{Z}_{\mathbf{s}} & \xlongequal{\quad} & \mathcal{Z}_{\mathbf{s}} & \xrightarrow{\phi^z} & \mathcal{Z}_{\mathbf{s}}. \end{array}$$

Here the vertical maps are coordinate embeddings given by definition of the cluster variety. Combining Lemma 3.2 and the commutative diagram (3.5), one can see that the induced automorphisms on the cluster variety $\mathcal{Z}_{\mathbf{s}}$ for different choices of representative paths coincide, and get a left action of $\Gamma_{\mathbf{s}}$ on $\mathcal{Z}_{\mathbf{s}}$. Since the birational map (3.6) only depends on the mutation loop ϕ and the choice of a vertex $v_0 \in \mathbb{E}\text{xch}_{\mathbf{s}}$ by Lemma 3.2, it is called the *coordinate expression* of ϕ with respect to the vertex v_0 .

Furthermore, we can identify $H_{\mathcal{Z}, \mathbf{s}}$ and $\bigcup_{v \in \mathbb{E}\text{xch}_{\mathbf{s}}} H_{\mathcal{Z}, (v)}$ by the same manner as cluster varieties $\mathcal{Z}_{\mathbf{s}}$, for $\mathcal{Z} = \mathcal{A}, \mathcal{X}$. In view of Proposition A.4, let a mutation loop $\phi = [\gamma]_{\mathbf{s}}$ act on the tori $H_{\mathcal{A}, \mathbf{s}}$ (and $H_{\mathcal{X}, \mathbf{s}}$) as follows:

$$\begin{array}{ccccc} H_{\mathcal{Z}, (v_0)} & \xrightarrow{\tilde{\mu}_\gamma^\epsilon} & H_{\mathcal{Z}, (v)} & \xrightarrow{\sim} & H_{\mathcal{Z}, (v_0)} \\ \downarrow & & \downarrow & & \downarrow \\ H_{\mathcal{Z}} & \xlongequal{\quad} & H_{\mathcal{Z}} & \xrightarrow{\phi^z} & H_{\mathcal{Z}}. \end{array}$$

Here $\tilde{\mu}_\gamma^\epsilon$ is the monomial isomorphism given by the composition of the monomial parts of labeled cluster transformations (with arbitrary choices of signs) associated to the edges traversed by γ , and the second map in the top row is induced by the seed isomorphism (3.1). As a direct consequence of Proposition A.4, we have:

Corollary 3.9. *Each map in the cluster exact sequence (A.7) is $\Gamma_{\mathbf{s}}$ -equivariant.*

The cluster modular group also acts on the labeled exchange graph as $\phi(v_0) := v$ for $v_0 \in \mathbb{E}\text{xch}_{\mathbf{s}}$ and $\phi \in \Gamma_{\mathbf{s}}$, where v is the terminal vertex of a representation path $\gamma : v_0 \rightarrow v$

of ϕ in $\mathbb{E}\text{ch}_{\mathbf{s}}$. One can check the following lemma by comparing the definitions of the \mathbf{s} -equivalence of paths in \mathbb{E}_I and the relation \sim_{triv} :

Lemma 3.10. *The action of $\Gamma_{\mathbf{s}}$ on $\mathbb{E}\text{ch}_{\mathbf{s}}$ is well-defined, simplicial and free.*

The cluster modular group action on $\mathbb{E}\text{ch}_{\mathbf{s}}$ descends to that on $\text{Exch}_{\mathbf{s}}$, where the latter action possibly has fixed points.

Example 3.11 (The seed pattern of type A_2). Here is a simple example. Let $I = I_{\text{uf}} := \{1, 2\}$. Since the tree $\mathbb{T}_{I_{\text{uf}}}$ is a bi-valent tree in this case, we identify its set of vertices with the set \mathbb{Z} of integers. A seed pattern $\mathbf{s} : \mathbb{T}_{I_{\text{uf}}} \ni t \mapsto (N^{(t)}, B^{(t)})$ satisfying

$$B^{(2m)} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad B^{(2m+1)} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

for all $m \in \mathbb{Z}$ is called a *seed pattern of type A_2* . The quiver representing the initial matrix $B^{(0)}$ is

$$\underset{\circ}{1} \longrightarrow \underset{\circ}{2}.$$

The corresponding graph \mathbb{E}_I is shown in Figure 16. We have two \mathbf{s} -equivalence classes $[t_0, e]_{\mathbf{s}}$ and $[t_0, (1\ 2)]_{\mathbf{s}}$, which are distinguished by two colors in the figure. For example, we have $[t_0, e]_{\mathbf{s}} = \{(t_{2m}, e), (t_{2m+1}, (1\ 2)) \mid m \in \mathbb{Z}\}$. One can verify that the path $\delta : (t_0, e) \xrightarrow{(1,2,1,2,1,(1\ 2))} (t_5, \sigma)$ is trivial (known as the *pentagon relation*), and we get $\mathbb{E}\text{ch}_{\mathbf{s}}$ as shown in Figure 17.

The image of the paths γ_1 , γ_2 , and γ'_2 shown in Figure 16 onto the labeled exchange graph give rise to mutation loops. For example, the cluster transformations associated with the paths γ_1 and γ_2 are given by

$$\begin{aligned} \mu_{\gamma_1}^z &:= (1\ 2)^z \mu_1^z : \mathcal{Z}_{(v_0)} \rightarrow \mathcal{Z}_{(\sigma.v_1)}, \\ \mu_{\gamma_2}^z &:= \mu_2^z \mu_1^z : \mathcal{Z}_{(v_0)} \rightarrow \mathcal{Z}_{(v_2)} \end{aligned}$$

for $(z, \mathcal{Z}) = (a, \mathcal{A}), (x, \mathcal{X})$. The coordinate expression of the mutation loop $\phi_1 := [\gamma_1]_{\mathbf{s}}$ with respect to the vertex v_0 is given by

$$\phi_1^z : \mathcal{Z}_{(v_0)} \xrightarrow{\mu_{\gamma_1}^z} \mathcal{Z}_{(\sigma.v_1)} \xrightarrow{\sim} \mathcal{Z}_{(v_0)},$$

which are explicitly expressed on the cluster coordinates as

$$(\phi_1^a)^*(A_1, A_2) = \left(A_2, \frac{1 + A_2}{A_1} \right), \quad (3.7)$$

$$(\phi_1^x)^*(X_1, X_2) = (X_2(1 + X_1), X_1^{-1}). \quad (3.8)$$

The composition of ϕ_1 and $\phi_2 := [\gamma_2]_{\mathbf{s}}$ is represented by the edge path $\gamma_1 * \gamma'_2$, where note that $[\gamma'_2]_{\mathbf{s}} = [\gamma_2]_{\mathbf{s}}$.

From the vertical projection we get a pentagon, which is the exchange graph Exch_{A_2} of type A_2 . The mutation loop ϕ_1 acts on this pentagon as a shift by one.




FIGURE 16. The graph \mathbb{E}_I of type A_2 .




FIGURE 17. The labeled and unlabeled exchange graphs of type A_2 . Here $v_0 := [t_0, e]_{\text{triv}} = [t_5, \sigma]_{\text{triv}}$ and $\sigma.v_0 = [t_0, \sigma]_{\text{triv}} = [t_5, e]_{\text{triv}}$.

4. MEASURED LAMINATIONS FROM THE CLUSTER ALGEBRAIC POINT OF VIEW

In this section, we review the connection between the theory of laminations and the cluster ensembles. These relations will enable us to bring the notion of *sign stability* [IK21] into the study of the mapping class group action.


4.1. The cluster ensemble associated with a marked surface. For a marked surface Σ , one can construct a seed pattern s_Σ canonically.

Let $I = I(\Sigma) := \{1, \dots, 3(2g - 2 + h + b) + 2|M_\partial|\}$, $I_{\text{uf}} = I_{\text{uf}}(\Sigma) := \{1, \dots, 3(2g - 2 + h + b) + |M_\partial|\}$ and $I_f = I_f(\Sigma) := I \setminus I_{\text{uf}}$ (hence $|I_f| = |M_\partial|$). A *labeled triangulation* consists of an ideal triangulation Δ and a bijection $\ell : I \rightarrow \Delta$ called a *labeling* such that $\ell(I_{\text{uf}}) = \Delta_{\text{uf}}$ (and hence $\ell(I_f) = \Delta_f$). A *labeled signed triangulation* is similarly defined. An equivalence relation among labeled signed triangulations is generated by the operation which alters the sign of a univalent puncture p and simultaneously transposes the labels of edges forming a self-folded triangle containing p . An equivalence class is called a *labeled tagged triangulation*.

Let us fix an “initial” labeled tagged triangulation $(\Delta_0, \varsigma_0, \ell_0)$. For a labeled tagged triangulation $(\Delta, \varsigma, \ell)$, we define a skew-symmetric matrix $B^{(\Delta, \varsigma, \ell)}$ by

$$B^{(\Delta, \varsigma, \ell)} := (b_{\ell(i), \ell(j)}^\Delta)_{i, j \in I}.$$

The quiver representing the matrix $B^{(\Delta, \varsigma, \ell)}$ locally looks like



In general, some of arrows may be merged to produce multiple arrows or canceled to each other, depending on the gluing pattern of triangles.

Let us fix a vertex t_0 of the regular tree $\mathbb{T}_{I_{\text{uf}}}$ and define the seed pattern $\mathbf{s}_\Sigma := \mathbf{s}_{t_0; B^{(\Delta_0, \varsigma_0, \ell_0)}}$ to be the seed pattern with the initial exchange matrix $B^{(\Delta_0, \varsigma_0, \ell_0)}$ (Lemma A.1). Then, we obtain cluster varieties and exchange graphs from the seed pattern \mathbf{s}_Σ . We abbreviate $Z_{\mathbf{s}_\Sigma}$ to Z_Σ for $Z = \mathcal{A}, \mathcal{X}^{\text{uf}}, H_{\mathcal{A}}, H_{\mathcal{X}}, \mathbb{E}\text{ch}$ and Ech . We reserve the notation Γ_Σ for a group slightly different from the cluster modular group $\Gamma_{\mathbf{s}_\Sigma}$, which will be defined in the next subsection.

Let $\text{Tri}^\bowtie(\Sigma)$ denote the graph whose vertices are tagged triangulations of Σ and adjacent tagged triangulations are related by a flip. This is the tagged version of the graph arising from the *Ptolemy groupoid* [Pen]. Flips of tagged triangulations can be upgraded to *labeled flips*

$$\mu_k : (\Delta, \varsigma, \ell) \rightarrow (\Delta', \varsigma, \ell') \quad (4.1)$$

for $k \in I_{\text{uf}}$, where Δ' is the tagged triangulation obtained by the flip along the arc $\ell(k)$, and ℓ' is defined by $\ell'(i) := \ell(i)$ for $i \neq k$ and by setting $\ell'(k)$ to be the new arc arising from the flip. Let $\text{Tri}^\bowtie(\Sigma)$ denotes the graph whose vertices are labeled tagged triangulation of Σ and adjacent labeled tagged triangulations are related by a labeled flip or the action of a transposition of labelings in $\mathfrak{S}_I = \mathfrak{S}_{|I_{\text{uf}}|} \times \mathfrak{S}_{|I_f|}$. Here the action of \mathfrak{S}_I on a labeling $\ell : I \rightarrow \Delta$ is defined as $\sigma.\ell(i) := \ell(\sigma^{-1}(i))$ for $\sigma \in \mathfrak{S}_I$ and $i \in I$. We will refer to an edge corresponding to a labeled flip (resp. the action of a transposition on a labeling)

as a *horizontal* (resp. *vertical*) edge. We denote by $\pi_{\text{Tri}} : \text{Tri}^{\boxtimes}(\Sigma) \rightarrow \text{Tri}^{\boxtimes}(\Sigma)$ the graph projection forgetting the labelings. By [FST08, Proposition 7.10 and Theorem 7.11] and [FT18], we obtain the following:

Theorem 4.1. *If Σ is not a once-punctured surface, there exists a unique graph isomorphism $\text{Tri}^{\boxtimes}(\Sigma) \xrightarrow{\sim} \text{Exch}_{\Sigma}$ such that*

- $(\Delta_0, \varsigma_0, \ell_0) \mapsto [t_0, e]_{\text{triv}}$, and
- if a vertex $(\Delta, \varsigma, \ell)$ is sent to $[t, \sigma]_{\text{triv}}$, then the horizontal edge given by a labeled flip $\mu_k : (\Delta, \varsigma, \ell) \rightarrow (\Delta', \varsigma, \ell')$ is sent to a horizontal edge $[t, \sigma]_{\text{triv}} \xrightarrow{k} [t', \sigma]_{\text{triv}}$, and the vertical edge given by the action of a transposition $(i\ j)$ on $(\Delta, \varsigma, \ell)$ is sent to the vertical edge $[t, \sigma]_{\text{triv}} \xrightarrow{(i\ j)} [t, (i\ j)\sigma]_{\text{triv}}$.

Moreover, it descends to a graph isomorphism $\text{Tri}^{\boxtimes}(\Sigma) \xrightarrow{\sim} \text{Exch}_{\Sigma}$:

$$\begin{array}{ccc} \text{Tri}^{\boxtimes}(\Sigma) & \xrightarrow{\sim} & \text{Exch}_{\Sigma} \\ \pi_{\text{Tri}} \downarrow & & \downarrow \pi_{\text{Ex}} \\ \text{Tri}^{\boxtimes}(\Sigma) & \xrightarrow{\sim} & \text{Exch}_{\Sigma}. \end{array}$$

If Σ is a once-punctured surface, then the graphs $\text{Tri}^{\boxtimes}(\Sigma)$ and $\text{Tri}^{\boxtimes}(\Sigma)$ respectively have two connected components, which are isomorphic to each other. The graph Exch_{Σ} (resp. Exch_{Σ}) is isomorphic to the connected component of $\text{Tri}^{\boxtimes}(\Sigma)$ containing $(\Delta_0, \varsigma_0, \ell_0)$ (resp. the component of $\text{Tri}^{\boxtimes}(\Sigma)$ containing (Δ_0, ς_0)) in the same way as above.


Example 4.2. Let Σ be a disk with 5 special points. When forgetting the frozen indices, it is well-known that the seed pattern \mathbf{s}_{Σ} coincides with the seed pattern of type A_2 . One can see the isomorphisms $\text{Tri}^{\boxtimes}(\Sigma) \cong \text{Exch}_{\Sigma}$ and $\text{Tri}^{\boxtimes}(\Sigma) \cong \text{Exch}_{\Sigma}$ by comparing Figure 18 with Figure 17.

Recall that the cluster varieties \mathcal{A}_{Σ} and $\mathcal{X}_{\Sigma}^{\text{uf}}$ are parametrized by Exch_{Σ} . The cluster interpretation of Proposition 2.7 is the following:

Theorem 4.3 (Fock–Goncharov, Fomin–Shapiro–Thurston). *We have canonical PL isomorphisms $\mathcal{L}^a(\Sigma, \mathbb{R}) \cong \mathcal{A}_{\Sigma}(\mathbb{R}^{\text{trop}})$ and $\mathcal{L}^x(\Sigma, \mathbb{R}) \cong \mathcal{X}_{\Sigma}^{\text{uf}}(\mathbb{R}^{\text{trop}})$. Moreover, the exact sequence (2.5) is isomorphic to the \mathbb{R}^{trop} -points of the cluster exact sequence (A.7):*

$$0 \rightarrow H_{\mathcal{A}, \Sigma}(\mathbb{R}^{\text{trop}}) \rightarrow \mathcal{A}_{\Sigma}(\mathbb{R}^{\text{trop}}) \xrightarrow{p^{\text{trop}}} \mathcal{X}_{\Sigma}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \xrightarrow{\theta^{\text{trop}}} H_{\mathcal{X}, \Sigma}(\mathbb{R}^{\text{trop}}) \rightarrow 0. \quad (4.2)$$

Proof. For each labeled tagged triangulation $(\Delta, \varsigma, \ell) \in \text{Tri}^{\boxtimes}(\Sigma)$, let $v = v[\Delta, \varsigma, \ell] \in \text{Exch}_{\Sigma}$ be the corresponding vertex under the isomorphism in Theorem 4.1. Define maps

FIGURE 18. The graphs $\text{Tri}^{\heartsuit}(\Sigma)$ and $\text{Tri}^{\heartsuit}(\Sigma)$ without frozen indices.

$\psi_{\Sigma}^a : \mathcal{L}^a(\Sigma, \mathbb{R}) \rightarrow \mathcal{A}_{\Sigma}(\mathbb{R}^{\text{trop}})$ and $\psi_{\Sigma}^x : \mathcal{L}^x(\Sigma, \mathbb{R}) \rightarrow \mathcal{X}_{\Sigma}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ so that the following diagrams commute:

$$\begin{array}{ccc}
 \mathcal{L}^a(\Sigma, \mathbb{R}) & \xrightarrow{\psi_{\Sigma}^a} & \mathcal{A}_{\Sigma}(\mathbb{R}^{\text{trop}}) & & \mathcal{L}^x(\Sigma, \mathbb{R}) & \xrightarrow{\psi_{\Sigma}^x} & \mathcal{X}_{\Sigma}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \\
 \mathfrak{a}(\Delta, \varsigma) \downarrow & & \uparrow \wr & & \mathfrak{x}(\Delta, \varsigma) \downarrow & & \uparrow \wr \\
 \mathbb{R}^{\Delta} & & \mathcal{A}_{(v)}(\mathbb{R}^{\text{trop}}) & & \mathbb{R}^{\Delta} & & \mathcal{X}_{(v)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \\
 \ell^* \downarrow & & \downarrow a_{(v)} & & \ell^* \downarrow & & \downarrow x_{(v)} \\
 \mathbb{R}^I & \xrightarrow{\iota} & \mathbb{R}^I & & \mathbb{R}^{I_{\text{uf}}} & \xrightarrow{\iota} & \mathbb{R}^{I_{\text{uf}}}
 \end{array}$$

Here $\mathcal{Z}_{(v)}(\mathbb{R}^{\text{trop}}) \xrightarrow{\sim} \mathcal{Z}_{\Sigma}(\mathbb{R}^{\text{trop}})$ is the tropicalization of the open embedding $\mathcal{Z}_{(v)} \subset \mathcal{Z}_{\Sigma}$ for $\mathcal{Z} = \mathcal{A}, \mathcal{X}^{\text{uf}}$, and $\iota((x_i)_{i \in I(\Sigma)}) := (-x_i)_{i \in I(\Sigma)}$. Then the maps ψ_{Σ}^a and ψ_{Σ}^x are independent of the choice of Δ in view of Proposition 2.7 and the fact that $\mathbb{R}^T = (\mathbb{R}, \max, +) \rightarrow \mathbb{R}^{\text{trop}}$, $x \mapsto -x$ gives a semifield isomorphism. See Remark A.7.

For the second statement, observe that the lattice $\ker p^* \subset N_{\text{uf}}^{(\Delta, \varsigma)}$ for a tagged triangulation (Δ, ς) is generated by the vectors

$$n(p) := \sum_{\substack{\alpha \in \Delta_{\text{uf}} \\ \text{incident to } p}} e_{\alpha}$$

associated with punctures $p \in P$, where α in Δ_{uf} is counted twice if both of its endpoints are incident to p . Comparing with (2.3), we get $(\theta^{\text{trop}})^* \text{ch}_{n(p)}^{\text{trop}} = \theta_{\Delta, p}$.

The lattice coker p^* is generated by the similarly-defined vectors

$$\ell(m) := \sum_{\substack{\alpha \in \Delta \\ \text{incident to } m}} e_\alpha$$

associated with marked points $m \in P \cup M_\partial$. It is not hard to see that the action $\mathbb{R} \times \mathcal{A}_\Sigma(\mathbb{R}^{\text{trop}}) \rightarrow \mathcal{A}_\Sigma(\mathbb{R}^{\text{trop}})$ induced by the vector $\ell(m)$ coincides with the action (2.1). ■

In particular, comparing Lemma 2.6 and Proposition A.4, we get a PL isomorphism $\mathcal{L}^u(\Sigma, \mathbb{R}) \cong \mathcal{U}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}})$. Henceforce we identify the connected component of $\text{Tri}^{\text{pl}}(\Sigma)$ (resp. $\text{Tri}^{\text{pl}}(\Sigma)$) described in Theorem 4.1 with the graph $\mathbb{E}xch_\Sigma$ (resp. Exch_Σ) without notice.

The following lemma is useful in the sequel. Recall Example 2.5. The following lemma can be directly confirmed from Figure 19.

Lemma 4.4. *For a tagged triangulation (Δ, ς) , the cone $\mathcal{C}_{(\Delta, \varsigma)}^+$ (resp. $\mathcal{C}_{(\Delta, \varsigma)}^-$) in $\mathcal{X}_{(\Delta, \varsigma)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ consists of the real \mathcal{X} -laminations $\widehat{L} = (L, \sigma_L = (\sigma_p))$ obtained from Δ by removing the boundary segments, assigning an arbitrary non-negative real weight to each interior arc, slightly moving each endpoint on a special point along the boundary against (resp. following) the orientation induced by Σ , and setting $\sigma_p := -\varsigma_p$ (resp. $\sigma_p := \varsigma_p$) for all $p \in P$.*




FIGURE 19. A lamination in $\mathcal{C}_{(\Delta, \varsigma)}^+$.

4.2. The cluster modular group associated with a marked surface. Let us describe the cluster modular group associated with the seed pattern \mathbf{s}_Σ . Hereafter we denote the seed pattern on $\mathbb{E}xch_\Sigma$ defined by

$$\mathbb{E}xch_\Sigma \ni (\Delta, \varsigma, \ell) \mapsto \left(N^{(\Delta, \varsigma, \ell)} := \bigoplus_{i \in I} \mathbb{Z}e_i^{(\Delta, \varsigma, \ell)}, B^{(\Delta, \varsigma, \ell)} \right)$$

by the same notation \mathbf{s}_Σ . For a mapping class $\phi \in MC(\Sigma)$ and a tagged triangulation $(\Delta, \varsigma, \ell) \in \mathbb{E}xch_\Sigma$, (the labeled version of) the sequence f_ϕ of flips determines a horizontal edge path from $(\Delta, \varsigma, \ell)$ to $(\phi^{-1}(\Delta, \varsigma), \ell')$. By concatenating a vertical edge path from

$(\phi^{-1}(\Delta, \varsigma), \ell')$ to $\phi^{-1}(\Delta, \varsigma, \ell) := (\phi^{-1}(\Delta, \varsigma), \phi^{-1} \circ \ell)$, we obtain an edge path in $\mathbb{E}xch_{\Sigma}$ which represents a mutation loop.

If Σ is not a once-punctured torus, then such a mutation loop conversely determines a mapping class, since a mapping class which fixes a labeled ideal triangulation is trivial. Thus we have an injective group homomorphism $MC(\Sigma) \rightarrow \Gamma_{s_{\Sigma}}$. When Σ is a once-punctured torus, the hyperelliptic involution fixes each labeled ideal triangulation and thus we also get an injective homomorphism $MC(\Sigma) \rightarrow \Gamma_{s_{\Sigma}}$. Here recall the modified definition of the mapping class group in this case as the quotient group by the hyperelliptic involution.

There are additional mutation loops arising from the reflections τ_{ς} for $\varsigma \in \{+, -\}^P$ when Σ is not a once-punctured surface. For any labeled tagged triangulation $(\Delta, \varsigma_{\Delta}, \ell)$ and $\varsigma \in \{+, -\}^P$ we can find an edge path in $\mathbb{E}xch_{\Sigma}$ from $(\Delta, \varsigma_{\Delta}, \ell)$ to $\tau_{\varsigma}^{-1}(\Delta, \varsigma_{\Delta}, \ell) = (\Delta, \varsigma \cdot \varsigma_{\Delta}, \ell)$. By the very definition of the exchange matrix, it represents a mutation loop.

In general, a pair (ϕ, ς) of a mapping class ϕ and $\varsigma \in \{+, -\}^P$ gives rise to a mutation loop, which is represented by a path from a labeled tagged triangulation $(\Delta, \varsigma_{\Delta}, \ell)$ to

$$(\phi, \tau_{\varsigma})^{-1}(\Delta, \varsigma, \ell) := (\phi^{-1}(\Delta), \varsigma \cdot \phi^* \varsigma_{\Delta}, \phi^{-1} \circ \ell), \quad (4.3)$$

where $(\phi^* \varsigma)_p := \varsigma_{\phi(p)}$ for $p \in P$. Define

$$\Gamma_{\Sigma} := \begin{cases} MC(\Sigma) & \text{if } \Sigma \text{ is a punctured surface with exactly one puncture,} \\ MC(\Sigma) \times (\mathbb{Z}/2)^P & \text{otherwise.} \end{cases}$$

Here in the second case, the group structure is given by

$$(\phi_1, \varsigma_1) \cdot (\phi_2, \varsigma_2) := (\phi_1 \phi_2, (\phi_2^* \varsigma_1) \varsigma_2)$$

for $(\phi_i, \varsigma_i) \in MC(\Sigma) \times (\mathbb{Z}/2)^P$. Now we have the following:

Theorem 4.5. *Let Σ be a marked surface. If Σ is not a sphere with 4 punctures, then the group Γ_{Σ} is isomorphic to the cluster modular group $\Gamma_{s_{\Sigma}}$. Otherwise, Γ_{Σ} is a subgroup of $\Gamma_{s_{\Sigma}}$ of index ≥ 2 .*

Proof. Except for the following cases, the statement is proved in [BS15]:

- (a) a sphere with 4 punctures;
- (b) a sphere with 5 punctures;
- (c) a once-punctured disk with 2 or 4 marked points on its boundary;
- (d) a twice-punctured disk with 2 marked points on its boundary;
- (e) an unpunctured disk with 4 marked points on its boundary;
- (f) an annulus with once marked point on each boundary component;
- (g) a once-punctured torus.

In the case (a), there is a mutation loop which is not contained in $MC(\Sigma) \times (\mathbb{Z}/2)^h$ (see [Gu11, Graph 2]). Nevertheless, the injectivity of $\Gamma_{\Sigma} \rightarrow \Gamma_{s_{\Sigma}}$ still holds (e.g., [ASS12]). The other cases are remedied as follows.

The case (b) was excluded only for the issue of existence of a non-degenerate potential of a quiver corresponding to an ideal triangulation, so it does not matter for our setting. In the cases (c) and (d), we have a mutation loop which is not contained in Γ_Σ when forgetting frozen indices. However, in our setting, those are not mutation loops since they do not preserve the frozen part of the exchange matrix. In the cases (e) and (f), the injectivity of $\Gamma_\Sigma \rightarrow \Gamma_{s_\Sigma}$ does not hold when forgetting frozen indices. It is also not an issue when considering frozen indices, since the relevant mutation loops are distinguished by their action on frozen coordinates.

In the case (g), the quotient of the mapping class group by the *hyperelliptic involution* is isomorphic to the automorphism group of the mapping class groupoid, which is nothing but the cluster modular group. See [Pen, Chapter 4, Section 4 and Chapter 5, Section 1]. ■

In view of this theorem, we regard an element of Γ_Σ as a mutation loop. A mutation loop $(\phi, \varsigma) \in \Gamma_\Sigma$ acts on the \mathcal{A} - and \mathcal{X} -laminations as $\phi \circ \tau_\varsigma$. Note that we have

$$(\phi, \varsigma)^r = (\phi^r, (\phi^{r-1})^* \varsigma \cdots \phi^* \varsigma \cdot \varsigma) \quad (4.4)$$

for all $r \in \mathbb{Z}$, and $(\phi, \varsigma)^{r_0} = (\phi^{r_0}, 1)$ for some $r_0 \in \mathbb{Z}_{>0}$. Hereafter we regard the mapping class group as a subgroup of the cluster modular group via the embedding i_Σ , and identify a mapping class ϕ with the associated mutation loop $i_\Sigma(\phi) = (\phi, 1)$. The image of a mutation loop under the projection $\Gamma_\Sigma \rightarrow MC(\Sigma)$ is called its *underlying mapping class*.

Finally, we have the following extension of Proposition 2.8 to the cluster modular group:

Proposition 4.6. *Let $\sigma_\Sigma : \Gamma_\Sigma \rightarrow \mathfrak{S}_{h+|M_\partial|} \times (\mathbb{Z}/2)^h$ be the action on $H_{\mathcal{A}}$ given by the permutation of marked points and the reflection of the signs of punctures. Let $\sigma_\Sigma^{\text{uf}} : \Gamma_\Sigma \rightarrow \mathfrak{S}_h \times (\mathbb{Z}/2)^h$ be the action on $H_{\mathcal{X}, \Sigma}$ given similarly by restricting to the action on the punctures and their signs.*

(1) *Then under the splitting $\mathcal{A}_\Sigma(\mathbb{R}^{\text{trop}}) \cong \mathcal{U}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}}) \times H_{\mathcal{A}, \Sigma}(\mathbb{R}^{\text{trop}})$ induced from (2.6), the action of Γ_Σ on $\mathcal{A}_\Sigma(\mathbb{R}^{\text{trop}})$ is decomposed into the direct product of the following two actions:*

- *the induced action on $\mathcal{U}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}})$, and*
- *the action σ_Σ on $H_{\mathcal{A}, \Sigma}(\mathbb{R}^{\text{trop}})$.*

(2) *We have the following commutative diagram:*

$$\begin{array}{ccc} \mathcal{X}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}}) & \xrightarrow{\phi^x} & \mathcal{X}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}}) \\ \theta \downarrow & & \downarrow \theta \\ H_{\mathcal{X}, \Sigma}(\mathbb{R}^{\text{trop}}) & \xrightarrow{\sigma_\Sigma^{\text{uf}}(\phi)} & H_{\mathcal{X}, \Sigma}(\mathbb{R}^{\text{trop}}). \end{array} \quad (4.5)$$

Example 4.7. Here are some simple examples of mutation loops obtained from mapping classes with reflections.

- (1) Let Σ be an annulus with one special point on each boundary component. Let C be a simple closed curve whose homotopy class generates $\pi_1(\Sigma) \cong \mathbb{Z}$, and $T_C \in MC(\Sigma)$ the (right-hand) Dehn twist along C . Choose a labeled triangulation (Δ, ℓ) as shown in the top left of Figure 20. Here $I(\Sigma) = \{1, 2, 3, 4\}$ and $I_{\text{uf}}(\Sigma) = \{1, 2\}$. Then one can see from the figure that the mutation loop $T_C = i_\Sigma(T_C) \in \Gamma_\Sigma$ is represented by the path

$$\gamma_C : (\Delta, \ell) \xrightarrow{1} (\Delta', \ell') \xrightarrow{(1\ 2)} T_C^{-1}(\Delta, \ell).$$

- (2) Here is an example of a reflection. Let Σ be a once-punctured disk with two special points on the boundary. Let p denote the central puncture, and $\tau_p \in \Gamma_\Sigma$ be the reflection corresponding to the sign $\varsigma_p = -$. Choose a labeled tagged triangulation $(\Delta, \varsigma_\Delta, \ell)$ (which comes from a labeled ideal triangulation) as shown in the top-left of Figure 21. Then one can see from the figure that the mutation loop τ_p is represented by the path

$$\gamma_p : (\Delta, \varsigma_\Delta, \ell) \xrightarrow{1} (\Delta', \varsigma_{\Delta'}, \ell') \xrightarrow{2} (\Delta'', \varsigma_{\Delta''}, \ell'') \xrightarrow{(1\ 2)} \tau_p^{-1}(\Delta, \varsigma_\Delta, \ell) = (\Delta, -\varsigma_\Delta, \ell).$$

Here recall the equivalence relation on the labeled signed triangulations. We could also adopt $(\Delta', \varsigma_{\Delta'}, \ell')$ as the initial triangulation: in that case τ_p is represented by the single transposition $(1\ 2)$.





FIGURE 20. The mutation loop given by the Dehn twist T_C .

5. SIGN STABILITY OF MUTATION LOOPS

In this section, we recall the notion of sign-stability of a representation path of a mutation loop from [IK21]. First we prepare some terminology for piecewise-linear maps (PL maps for short).

FIGURE 21. The mutation loop given by the reflection τ_p .

5.1. PL maps and their presentation matrices. For any vector space V , we have the translation isomorphism $t_x : V \xrightarrow{\sim} T_x V$, $v \mapsto \frac{d}{ds}|_{s=0} (x + sv)$ for any point $x \in V$. Let $\psi : V \rightarrow W$ be a PL map between two vector spaces. Here is a simple but useful observation:

Lemma 5.1. *Let $x \in V \setminus \{0\}$ be a point at which ψ is differentiable. Let $\mathcal{C} \subset V$ be a maximal dimensional cone containing x such that the restriction $\psi|_{\mathcal{C}}$ is linear. Then its linear extension $\psi \uparrow_{\mathcal{C}} : V \rightarrow W$ fits into the following commutative diagram:*

$$\begin{array}{ccc} V & \xrightarrow{\psi \uparrow_{\mathcal{C}}} & W \\ t_x \downarrow & & \downarrow t_{\psi(x)} \\ T_x V & \xrightarrow{(d\psi)_x} & T_{\psi(x)} W. \end{array}$$

In particular, the restriction $(d\psi)_x|_{t_x(\mathcal{C})}$ coincides with the restriction $\psi|_{\mathcal{C}}$ via the translations. See Figure 22.

Suppose further that V and W have fixed bases $(e_i)_{i=1}^n$ and $(e'_j)_{j=1}^m$, respectively. Based on the observation above, we use the following terminologies:

- The *presentation matrix of ψ at a differentiable point x* is the presentation matrix of the tangent map $(d\psi)_x : T_x V \rightarrow T_{\psi(x)} W$ with respect to the given bases.
- The *presentation matrix of ψ on the cone \mathcal{C}* is the presentation matrix of ψ at any interior point $x \in \mathcal{C}$. Equivalently, it is the presentation matrix of the linear extension $\psi \uparrow_{\mathcal{C}} : V \rightarrow W$.

The following lemma characterizes the eigenvectors of the presentation matrix in terms of tangent vectors, which will be useful for the study of spectral properties of a PL dynamical system:

Lemma 5.2. *Suppose that $n = \dim V = \dim W$, and consider the presentation matrix $E_x \in \text{Mat}_{n \times n}(\mathbb{R})$ of a PL map $\psi : V \rightarrow W$ at a differentiable point $x \in V$. Then for $\lambda \in \mathbb{R}$, the coordinate vector $\mathbf{v} = (v_i)_{i=1}^n \in \mathbb{R}^n$ of $v = \sum_i v_i e_i \in V$ satisfies $E_x \mathbf{v} = \lambda \mathbf{v}$ if and only if $(d\psi)_x(t_x(v)) = \lambda t_{\psi(x)}(v)$.*




FIGURE 22. Translations of a linear subspace $V_0 \subset V$ and a cone $\mathcal{C} \subset V$.

When $V = \mathcal{X}_{(v)}(\mathbb{R}^{\text{trop}})$ and $W = \mathcal{X}_{(v')}(\mathbb{R}^{\text{trop}})$ for some $v, v' \in \mathbb{E}\text{ch}_{\mathbf{s}}$, we always consider the bases $(f_i^{(v)})_{i \in I}$ and $(f_i^{(v')})_{i \in I}$ respectively, unless otherwise specified.

For a PL map $f : M \rightarrow N$ between two PL manifolds, note that the differentiability of f does depend on the choice of a chart.⁴ Under the situation that the PL atlases of M and N are specified, we say that f is *intrinsically differentiable* at $x \in M$ if its coordinate expression is differentiable at the corresponding point for any charts in the given atlases. When $M = \mathcal{X}_{\mathbf{s}}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ and $N = \mathcal{X}_{\mathbf{s}' }^{\text{uf}}(\mathbb{R}^{\text{trop}})$ for some seed patterns \mathbf{s} and \mathbf{s}' , we discuss the intrinsic differentiability with respect to the cluster atlases unless otherwise specified.

5.2. Sign stability. Now recall the tropical cluster \mathcal{X} -transformation $\mu_k : \mathcal{X}_{(v)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \rightarrow \mathcal{X}_{(v')}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ associated with an edge $v \xrightarrow{k} v'$ in $\mathbb{E}\text{ch}_{\mathbf{s}}$. See Appendix A.3 for our notational convention. For a real number $a \in \mathbb{R}$, let $\text{sgn}(a)$ denote its sign:

$$\text{sgn}(a) := \begin{cases} + & \text{if } a > 0, \\ 0 & \text{if } a = 0, \\ - & \text{if } a < 0. \end{cases}$$

⁴On the other hand, the notion of *tangentiability* is intrinsically defined: see, for instance, [Bon98]. We do not pursue this direction in this paper.

The following expression is useful in the sequel:

Lemma 5.3 ([IK21, Lemma 3.1]). *Fix a point $w \in \mathcal{X}_{(v)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$. Then the tropical cluster \mathcal{X} -transformation $\mu_k : \mathcal{X}_{(v)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \rightarrow \mathcal{X}_{(v')}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ is given by*

$$x_i^{(v')}(\mu_k(w)) = \begin{cases} -x_k^{(v)}(w) & \text{if } i = k, \\ x_i^{(v)}(w) + [\text{sgn}(x_k^{(v)}(w))b_{ik}^{(v)}]_+ x_k^{(v)}(w) & \text{if } i \neq k. \end{cases} \quad (5.1)$$

With this lemma in mind, we consider the half-spaces

$$\mathcal{H}_{k,\epsilon}^{x,(v)} := \{w \in \mathcal{X}_{(v)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \mid \epsilon x_k^{(v)}(w) \geq 0\}$$

for $k \in I_{\text{uf}}$, $\epsilon \in \{+, -\}$ and $v \in \mathbb{E}\text{Xch}_{\mathbf{s}}$. Then we have the following immediate corollary:

Corollary 5.4. *For $\epsilon \in \{+, -\}$, the tropical cluster \mathcal{X} -transformation $\mu_k : \mathcal{X}_{(v)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \rightarrow \mathcal{X}_{(v')}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ is differentiable at any point in $\text{int } \mathcal{H}_{k,\epsilon}^{x,(v)}$, and its presentation matrix is the unfrozen part of the matrix $E_{k,\epsilon}^{(v)}$ defined in (A.4).*

The unfrozen part of the matrix $E_{k,\epsilon}^{(v)}$ is denoted by the same symbol when no confusion can occur.

We are going to define the *sign* of a path in $\mathbb{E}\text{Xch}_{\mathbf{s}}$. In the sequel, we use the following notation.

Notation 5.5. (1) For an edge path $\gamma : v_0 \xrightarrow{k_0} v_1 \xrightarrow{k_1} \dots \xrightarrow{k_{m-1}} v_m$ in $\mathbb{E}\text{Xch}_{\mathbf{s}}$ and $i = 1, \dots, m$, let $\gamma_{\leq i} : v_0 \xrightarrow{(k_0, \dots, k_{i-1})} v_i$ be the sub-path of γ from v_0 to v_i . Let $\gamma_{\leq 0}$ be the constant path at v_0 . Let $(k_{i(0)}, \dots, k_{i(h-1)})$ be the subsequence of $\mathbf{k} = (k_0, \dots, k_{m-1})$ which corresponds to horizontal edges.

(2) Fixing the initial vertex v_0 , we simply denote the tropicalization of the coordinate expression (3.6) of ϕ with respect to v_0 by $\phi := \phi_{(v_0)} : \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \rightarrow \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$, when no confusion can occur.

Definition 5.6 (sign of a path). Let the notation as above, and fix a point $w \in \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$. The *sign* of γ at w is the sequence $\epsilon_\gamma(w) = (\epsilon_0, \dots, \epsilon_{h-1}) \in \{+, 0, -\}^h$ of signs defined by

$$\epsilon_\nu := \text{sgn}(x_{k_{i(\nu)}}^{(v_i(\nu))}(\mu_{\gamma_{\leq i(\nu)}}(w)))$$

for $\nu = 0, \dots, h-1$.

The next lemma expresses the heart of Definition 5.6.

Lemma 5.7. *Let $\epsilon = (\epsilon_0, \dots, \epsilon_{h-1})$ be the sign of a path γ at $w \in \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$. Then μ_γ is differentiable at w if and only if ϵ is strict i.e., $\epsilon \in \{+, -\}^h$. In this case, the presentation matrix of μ_γ at w is given by $E_\gamma^{\epsilon_\gamma(w)} := J_{k_m} \cdots J_{k_1}$, where*

$$J_{k_i} := \begin{cases} E_{k_{i(\nu)}, \epsilon_\nu}^{(v_i(\nu))} & \text{if } k_i = k_{i(\nu)} \text{ is a horizontal edge,} \\ P_{k_i} & \text{if } k_i \text{ is a vertical edge.} \end{cases}$$

Here P_σ denotes the presentation matrix for $\sigma \in \mathfrak{S}_I$.

The sign of a path is not invariant under the \mathbf{s} -equivalence. For example, its horizontal length is not preserved under some of the elementary operation (a) in Remark 3.8. Some of the operation (b) also preserves the sign in the following sense.

Lemma 5.8. (1) Let $\gamma : v_0 \xrightarrow{\mathbf{k}} v$ and $\gamma' : v'_0 \xrightarrow{\mathbf{k}} v'$ be two paths with the same labels in $\mathbb{E}\text{Xch}_{\mathbf{s}}$ such that the initial vertices v_0 and v'_0 are \mathbf{s} -equivalent. Then for any $w \in \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \cong \mathcal{X}_{(v'_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ we have $\epsilon_{\gamma}(w) = \epsilon_{\gamma'}(w)$, where the identification is via the seed isomorphism (3.1).

(2) Let $\gamma : v_0 \xrightarrow{\mathbf{k}} v$ and $\gamma' : \rho(v'_0) \xrightarrow{\rho(\mathbf{k})} \rho(v')$ for some $\rho \in \mathfrak{S}_I$ with $\rho(\mathbf{k}) := (\rho(k_0), \dots, \rho(k_{m-1}))$. Then for any $w \in \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ we have $\epsilon_{\gamma}(w) = \epsilon_{\gamma'}(\rho(w))$.

This lemma reduces some of redundancy for defining the signs.

Definition 5.9 (sign stability). Let γ be a path as above and suppose that $\phi := [\gamma]_{\mathbf{s}}$ is a mutation loop. Let $\Omega \subset \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ be a subset which is invariant under the rescaling action of $\mathbb{R}_{>0}$. Then we say that γ is *sign-stable* on Ω if there exists a sequence $\epsilon_{\gamma, \Omega}^{\text{stab}} \in \{+, -\}^h$ of strict signs such that for each $w \in \Omega \setminus \{0\}$, there exists an integer $n_0 \in \mathbb{N}$ such that

$$\epsilon_{\gamma}(\phi^n(w)) = \epsilon_{\gamma, \Omega}^{\text{stab}}$$

for all $n \geq n_0$. We call $\epsilon_{\gamma, \Omega}^{\text{stab}}$ the *stable sign* of γ on Ω .

For a mutation loop $\phi \in \Gamma_{\mathbf{s}}$ and a vertex $v_0 \in \mathbb{E}_I$, its presentation matrix at a differentiable point $w \in \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ is the presentation matrix $E_{\phi}^{(v_0)}(w) \in GL_N(\mathbb{Z})$ of the tangent map

$$(d\phi)_w : T_w \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \rightarrow T_{\phi(w)} \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$$

with respect to the basis $(f_i^{(v_0)})_{i \in I_{\text{uf}}}$ (recall the terminologies from Section 5.1). Note that a choice of a representation path $\gamma : v_0 \rightarrow v$ of ϕ gives a factorization $E_{\phi}^{(v_0)}(w) = E_{\gamma}^{\epsilon_{\gamma}(w)}$ into elementary matrices $E_{k_{i(\nu)}, \epsilon_{\nu}}^{(v_i(\nu))}$ and P_{k_i} as in Lemma 5.7. Thus the sign stability in particular implies that the presentation matrix of ϕ at each point $w \in \Omega$ stabilizes:

Corollary 5.10. Suppose γ is a path which defines a mutation loop $\phi = [\gamma]_{\mathbf{s}}$, and sign-stable on Ω . Then there exists an integral $N \times N$ -matrix $E_{\phi, \Omega}^{(v_0)} \in GL_N(\mathbb{Z})$ such that for each $w \in \Omega$, there exists an integer $n_0 \geq 0$ such that $E_{\phi}^{(v_0)}(\phi^n(w)) = E_{\phi, \Omega}^{(v_0)}$ for all $n \geq n_0$.

We call $E_{\phi, \Omega}^{(v_0)}$ the *stable presentation matrix* of ϕ with respect to the vertex v_0 . Notice that this matrix only depends on the mutation loop ϕ and the vertex v_0 , and defined when ϕ admits a sign-stable representation path starting from v_0 .

We are going to mention the Perron–Frobenius property of a sign-stable mutation loop. We say that a path $\gamma : (t_0, \sigma_0) \xrightarrow{\mathbf{k}=(k_0, \dots, k_{m-1})} (t_{m-1}, \sigma_{m-1})$ in \mathbb{E}_I with the horizontal subpath $(k_{i(0)}, \dots, k_{i(r-1)})$ is *fully-mutating* if

$$\langle \sigma_{m-1} \sigma_0^{-1} \rangle \cdot \{ \sigma_{i(0)}(k_{i(0)}), \dots, \sigma_{i(r-1)}(k_{i(r-1)}) \} = I_{\text{uf}}.$$

For example, the path $\gamma : (t_0, \sigma_0) \xrightarrow{1} (t_1, \sigma_0) \xrightarrow{\rho} (t_1, \rho\sigma_0)$ with the cyclic permutation $\rho := (1\ 2\ \cdots\ |I_{\text{uf}}|)$ is fully-mutating. A path in $\text{Exch}_{\mathfrak{s}}$ is said to be fully-mutating if it has a fully-mutating lift in \mathbb{E}_I .

An $\mathbb{R}_{\geq 0}$ -invariant set $\Omega \subset \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \cong \mathcal{X}_{\mathfrak{s}}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ is said to be *tame* if there exists $v \in \text{Exch}_{\mathfrak{s}}$ such that $\Omega \cap \text{int } \mathcal{C}_{(v)}^+ \neq \emptyset$. Here, $\mathcal{C}_{(v)}^{\pm} := \{w \in \mathcal{X}_{(v)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \mid \pm x_i^{(v)}(w) \geq 0, i \in I_{\text{uf}}\}$ for $v \in \text{Exch}_{\mathfrak{s}}$. In other words, $\Omega \subset \mathcal{X}_{\mathfrak{s}}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ intersects with the set $\mathfrak{U}_{\mathfrak{s}}$ defined in Lemma 5.19 below.

Theorem 5.11 (Perron–Frobenius property, [IK21, Theorem 3.12]). *Suppose γ is a path as above which represents a mutation loop $\phi = [\gamma]_{\mathfrak{s}}$, and sign-stable on a tame subset Ω . Then the spectral radius of the stable presentation matrix $E_{\phi, \Omega}^{(v_0)}$ is attained by a positive eigenvalue $\lambda_{\phi, \Omega} \geq 1$. Moreover if either γ is fully-mutating or $\lambda_{\phi, \mathcal{L}}^{(t_0)} > 1$, then one of the corresponding eigenvectors is given by the coordinate vector $\mathbf{x}(w_{\phi, \mathcal{L}}) \in \mathbb{R}^{I_{\text{uf}}}$ for some $w_{\phi, \mathcal{L}} \in \mathcal{C}_{\gamma}^{\text{stab}} \setminus \{0\}$.*

See [IK21, Remark 3.21] for a way to modify the proof of [IK21, Theorem 3.12] for non-horizontal paths.

Any path $\gamma : v_0 \rightarrow v$ has a constant sign in the interior of the cone $\mathcal{C}_{(v_0)}^+$ (resp. $\mathcal{C}_{(v_0)}^-$), which is given by the tropical sign (resp. that for the opposite seed pattern) [IK21, Lemma 3.12, Corollary 3.13]. In this sense the sign stability on the set

$$\Omega_{(v_0)}^{\text{can}} := \text{int } \mathcal{C}_{(v_0)}^+ \cup \text{int } \mathcal{C}_{(v_0)}^-$$

is most fundamental, and it turns out that it is sufficient for the computation of the algebraic entropy of cluster transformations: see Section 8.

Definition 5.12. If a mutation loop ϕ admits a representation path $\gamma : v_0 \rightarrow v$ which is sign-stable on the set $\Omega_{(v_0)}^{\text{can}}$, then we call $\lambda_{\phi} := \lambda_{\phi, \Omega_{(v_0)}^{\text{can}}}$ the *cluster stretch factor* of ϕ .

Remark 5.13 (Intrinsicity of the cluster stretch factor to the mutation loop). The quantity $\lambda_{\phi, \Omega_{(v_0)}^{\text{can}}}$ at once depends on the mutation loop ϕ and the vertex v_0 . If ϕ also admits a representation path $\gamma' : v'_0 \rightarrow v'$ which is sign-stable on $\Omega_{(v'_0)}^{\text{can}}$, then choosing a path $\delta : v_0 \rightarrow v'_0$, we get the relation

$$(d\phi_{(v'_0)})_{\mu_{\delta}(w)} \circ (d\mu_{\delta})_w = (d\mu_{\delta})_{\phi_{(v_0)}(w)} \circ (d\phi_{(v_0)})_w \quad (5.2)$$

for any point $w \in \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$. When $w = w_{\phi, \Omega}$, from Theorem 5.11 and Lemma 5.1 we get $\phi_{(v_0)}(w_{\phi, \Omega}) = \lambda_{\phi, \Omega_{(v_0)}^{\text{can}}} \cdot w_{\phi, \Omega}$ and in particular the points $w_{\phi, \Omega}$ and $\phi_{(v_0)}(w_{\phi, \Omega})$ have the same sign for any path δ . Denote the common presentation matrix of the PL map μ_{δ} at these points by M . Then (5.2) implies that

$$E_{\phi, \Omega_{(v'_0)}^{\text{can}}}^{(v'_0)} = M E_{\phi, \Omega_{(v_0)}^{\text{can}}}^{(v_0)} M^{-1},$$

hence the spectral radii $\lambda_{\phi, \Omega_{(v_0)}^{\text{can}}}$ and $\lambda_{\phi, \Omega_{(v'_0)}^{\text{can}}}$ are the same.

In Section 8, we will confirm that the following fundamental conjecture regarding the sign stability holds true for the mutation loops induced by mapping classes with reflections:

Conjecture 5.14 (Spectral duality conjecture). For any path γ which represents a mutation loop and a point $w \in \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ with strict sign $\epsilon_\gamma(w)$, the characteristic polynomials of the $I_{\text{uf}} \times I_{\text{uf}}$ matrices $E_\gamma^{\epsilon_\gamma(w)}$ and $\check{E}_\gamma^{\epsilon_\gamma(w)}$ are the same up to overall sign. In particular, the spectral radii of these matrices are the same.

Note that for an $N \times N$ -matrix E with $\det E = \pm 1$, the characteristic polynomials of the matrices E and \check{E} are the same up to overall sign if and only if the characteristic polynomial $P_E(\nu)$ of E is (anti-)palindromic: $P_E(\nu^{-1}) = \pm \nu^{-N} P_E(\nu)$. The conjecture has been proved for invertible exchange matrices [IK21, Proposition 3.11].

Example 5.15 (Kronecker quiver). Here is a simple but important family of examples of sign-stable mutation loops. Let $\ell \geq 2$ be an integer. Let $I := \{1, 2\}$ and $\mathbf{s} : t \mapsto (N^{(t)}, B^{(t)})$ a seed pattern such that

$$B^{(t_0)} = \begin{pmatrix} 0 & -\ell \\ \ell & 0 \end{pmatrix}$$

for a vertex $t_0 \in \mathbb{T}_I$. The quiver representing the initial matrix $B^{(t_0)}$ is

$$\begin{array}{ccc} \circ & \xleftarrow{\ell} & \circ \\ \text{1} & & \text{2} \end{array}$$

It is known that the exchange graph $\text{Exch}_{\mathbf{s}}$ is an infinite bivalent tree. Let us consider the path

$$\gamma : v_0 := [t_0, e]_{\text{triv}} \xrightarrow{1} v_1 \xrightarrow{(1\ 2)} (1\ 2).v_1,$$

in $\mathbb{E}\text{xch}_{\mathbf{s}}$ which represents a mutation loop $\phi := [\gamma]_{\mathbf{s}} \in \Gamma_{\mathbf{s}}$. On the interiors of the half-spaces $\mathcal{H}_{1,+}^{x,(v_0)}$ and $\mathcal{H}_{1,-}^{x,(v_0)}$, its action is represented respectively by

$$E_\gamma^{(+)} = \begin{pmatrix} \ell & 1 \\ -1 & 0 \end{pmatrix} \quad \text{and} \quad E_\gamma^{(-)} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

Then it turns out that the path γ is sign-stable on the entire space $\Omega := \mathcal{X}_{(v_0)}(\mathbb{R}^{\text{trop}})$ with the stable sign $\epsilon_\gamma^{\text{stab}} = (+)$. Indeed, one can verify that the cone

$$\mathcal{C} = \{x_1 \geq 0, x_1 + x_2 \geq 0\}$$

is ϕ -stable, and the orbit $(\phi^n(w))_{n \geq 0}$ of any point $w \in \mathcal{X}_{(v_0)}(\mathbb{R}^{\text{trop}}) \setminus \mathbb{R}_{\geq 0} L_\phi^-$ eventually enters this cone. Here $L_\phi^\pm := (1, (-\ell \pm \sqrt{\ell^2 - 4})/2)$ are the only eigenvectors of the PL action of ϕ with the eigenvalues $\lambda_\pm := (\ell \pm \sqrt{\ell^2 - 4})/2$, respectively. See Figure 23. The larger eigenvalue $\lambda_\phi := \lambda_+$ is the cluster stretch factor. Actually, the dynamical system determined by ϕ on the projectivized space $\mathbb{S}\mathcal{X}_{(v_0)}(\mathbb{R}^{\text{trop}}) := (\mathcal{X}_{(v_0)}(\mathbb{R}^{\text{trop}}) \setminus \{0\})/\mathbb{R}_{>0}$ has the following property: every projective orbit $([\phi^n(w)])_{n \geq 0}$ converges to the point $[L_\phi^+]$ and the orbit $([\phi^{-n}(w)])_{n \geq 0}$ does to $[L_\phi^-]$. When $\ell > 2$, that is, $[L_\phi^+] \neq [L_\phi^-]$, it is called the *North-South dynamics* (see Section 7 for a general definition).




FIGURE 23. An invariant cone (yellow region) and the eigenrays (red) of the mutation loop ϕ .

Remark 5.16 (Presentation matrices of tropical \mathcal{A} -transformations). The tropical cluster \mathcal{A} -transformation at a point $\tilde{w} \in \mathcal{A}_{(v)}(\mathbb{R}^{\text{trop}})$ can be written as follows:

$$a_i^{(v')}(\mu_k(\tilde{w})) = \begin{cases} -a_k^{(v)}(\tilde{w}) + \sum_{j \in I} [-\text{sgn}(x_k^{(v)}(p(\tilde{w})))b_{kj}^{(v)}]_+ a_j^{(v)}(\tilde{w}) & \text{if } i = k, \\ a_i^{(v)}(\tilde{w}) & \text{if } i \neq k. \end{cases} \quad (5.3)$$

It is presented by the matrix $\check{E}_{k,\epsilon}^{(v)}$ defined in (A.3), depending on the sign $\epsilon = \text{sgn}(x_k^{(v)}(p(\tilde{w})))$.

Hence, given a path $\gamma : v_0 \rightarrow v$, the presentation matrix of the tropical \mathcal{A} -transformation μ_γ^a at a point $\tilde{w} \in \mathcal{A}_{(v_0)}(\mathbb{R}^{\text{trop}})$ is determined by the sign of γ at the point $p(\tilde{w}) \in \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$. In particular, from the sign-stability on a set $\Omega \subset \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ with $\Omega \cap \mathcal{U}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \neq \emptyset$ we can know some dynamical behavior on the preimage $p^{-1}(\Omega)$ modulo the action on the $H_{\mathcal{A}}(\mathbb{R}^{\text{trop}})$ -direction. This will be shortly discussed in the forthcoming paper [IK20].

In fact, it turns out that the sign stability on the set $\Omega_{(v_0)}^{\text{can}}$ tells us more on the \mathcal{A} -transformations than one might expect. Although in the case that $\Omega_{(v_0)}^{\text{can}} \cap \mathcal{U}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) = \emptyset$ (e.g. for the seed pattern \mathbf{s}_Σ associated with a punctured surface Σ) we get no information in the sense mentioned above, the tropical duality (3.4) implies the stability of the recurrence relation for the G -matrices. This is one of the key observations for the computation of the algebraic entropy of cluster \mathcal{A} -transformations in [IK21].

5.3. Uniform sign stability. Let us consider the $\mathbb{R}_{>0}$ -invariant subset $\mathbb{R}_{>0} \cdot \mathcal{X}_s^{\text{uf}}(\mathbb{Q}^{\text{trop}}) \subset \mathcal{X}_s^{\text{uf}}(\mathbb{R}^{\text{trop}})$ and the corresponding sets $\mathbb{R}_{>0} \cdot \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{Q}^{\text{trop}}) \subset \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ in the coordinate charts.

Definition 5.17 (Uniform sign stability). A mutation loop $\phi \in \Gamma_{\mathfrak{s}}$ is said to be *uniformly sign-stable* if any representation path $\gamma : v_0 \rightarrow v$ of ϕ is sign-stable on $\mathbb{R}_{>0} \cdot \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{Q}^{\text{trop}})$.

Note that the uniform sign stability is genuinely a property of a mutation loop, rather than its representation paths. The uniform sign stability automatically implies the sign stability on a larger subset. For $v_0 \in \mathbb{E}\text{ch}_{\mathfrak{s}}$, let us consider the collection

$$\mathfrak{F}_{(v_0)}^{\pm} := \left\{ \mu_{\gamma(v)}^{-1}(\mathcal{C}_{(v)}^{\pm}) \mid v \in \mathbb{E}\text{ch}_{\mathfrak{s}} \right\}$$

of cones in $\mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$. Here $\gamma(v) : v_0 \rightarrow v$ is any edge path in $\mathbb{E}\text{ch}_{\mathfrak{s}}$. It forms a simplicial fan by [GHKK18, Theorem 2.13]. We call the support

$$|\mathfrak{F}_{(v_0)}^{\pm}| := \bigcup_{v \in \mathbb{E}\text{ch}_{\mathfrak{s}}} \mu_{\gamma(v)}^{-1}(\mathcal{C}_{(v)}^{\pm}) \subset \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$$

the (geometric realization of) the *Fock–Goncharov cluster complex*, according to [FG09, GHKK18]. It is known that the corresponding subsets $|\mathfrak{F}_{\mathfrak{s}}^{\pm}| \subset \mathcal{X}_{\mathfrak{s}}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ are independent of the choice of $v_0 \in \mathbb{E}\text{ch}_{\mathfrak{s}}$ [GHKK18, Theorem 2.13]. Therefore we can consider the canonically defined $\mathbb{R}_{>0}$ -invariant subset

$$\Omega_{\mathfrak{s}}^{\mathbb{Q}} := |\mathfrak{F}_{\mathfrak{s}}^{+}| \cup |\mathfrak{F}_{\mathfrak{s}}^{-}| \cup \mathbb{R}_{>0} \cdot \mathcal{X}_{\mathfrak{s}}^{\text{uf}}(\mathbb{Q}^{\text{trop}}) \subset \mathcal{X}_{\mathfrak{s}}^{\text{uf}}(\mathbb{R}^{\text{trop}})$$

and the corresponding sets $\Omega_{(v_0)}^{\mathbb{Q}} \subset \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ in the coordinate charts. Then [IK21, Proposition 3.14] tells us that the sign stability on $\mathbb{R}_{>0} \cdot \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{Q}^{\text{trop}})$ is equivalent to that on the set $\Omega_{(v_0)}^{\mathbb{Q}}$. Since $\Omega_{(v_0)}^{\text{can}} \subset \Omega_{(v_0)}^{\mathbb{Q}}$, a uniformly sign-stable mutation loop comes with its cluster stretch factor λ_{ϕ} .

Example 5.18. The mutation loop ϕ studied in Example 5.15 is uniformly sign-stable. Indeed, it suffices to show that the limit points L_{ϕ}^{\pm} have strict signs for every representation paths. In fact, their coordinate vectors in any cluster chart have no zero entries. To see this in the case $\ell > 2$, observe that there is no facets of the cluster complex in the second quadrant, and the rays $\mathbb{R}_{\geq 0} \cdot L_{\phi}^{\pm}$ themselves cannot be facets since they are irrational. If $\ell = 2$, then the ray $\mathbb{R}_{\geq 0} \cdot L_{\phi}^{+} = \mathbb{R}_{\geq 0} \cdot L_{\phi}^{-}$ is rational but do not coincides with any facets of the cluster complex. See Section 6 for a geometric description of this ray.

Common differentiable points. Recall that the PL automorphism on $\mathcal{X}_{\mathfrak{s}}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \rightarrow \mathcal{X}_{\mathfrak{s}}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ induced by a mutation loop $\phi \in \Gamma_{\mathfrak{s}}$ is intrinsically differentiable at a point $w \in \mathcal{X}_{\mathfrak{s}}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ if for any vertex $v_0 \in \mathbb{E}\text{ch}_{\mathfrak{s}}$, the PL automorphism $\phi_{(v_0)}$ on $\mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}})$ is differentiable at the point corresponding to w .

The following lemma shows that we have enough differentiable points:

Lemma 5.19. *Let $\phi \in \Gamma_{\mathfrak{s}}$ be a mutation loop. Then for any vertex $v_0 \in \mathbb{E}\text{ch}_{\mathfrak{s}}$, the PL automorphism $\phi_{(v_0)}$ is differentiable at any point in the open subset*

$$\mathfrak{U}_{(v_0)} := \bigcup_{v \in \mathbb{E}\text{ch}_{\mathfrak{s}}} \mu_{\gamma(v)}^{-1}(\text{int } \mathcal{C}_{(v)}^{+} \cup \text{int } \mathcal{C}_{(v)}^{-}) \subset \mathcal{X}_{(v_0)}^{\text{uf}}(\mathbb{R}^{\text{trop}}).$$

In particular, the PL action of ϕ on the PL manifold $\mathcal{X}_s^{\text{uf}}(\mathbb{R}^{\text{trop}})$ is intrinsically differentiable at any point in the corresponding Γ_s -invariant subset $\mathfrak{U}_s \subset \mathcal{X}_s^{\text{uf}}(\mathbb{R}^{\text{trop}})$.

5.4. Relation between the uniform sign stability and the cluster-pseudo-Anosov property. Recall that a mutation loop $\phi \in \Gamma_s$ is said to be *cluster-pseudo-Anosov* if no non-trivial power of ϕ fixes any point in $|\mathfrak{F}_s^+|$. This definition is equivalent to [Ish19, Definition 2.1], thanks to [GHKK18, Theorem 2.13].

Proposition 5.20. *A uniformly sign-stable mutation loop is cluster-pseudo-Anosov.*

Proof. Suppose that for a mutation loop ϕ , there exists an integer $m > 0$ and a point $w \in |\mathfrak{F}_s^+|$ such that $\phi^m(w) = w$. Take a vertex $v_0 \in \mathbb{E}\text{ch}_s$ such that $w \in \mathcal{C}_{(v_0)}^+$. Since the action of a mutation loop on the Fock–Goncharov cluster complex is simplicial, ϕ^m permutes the faces of the simplex $\mathbb{S}\mathcal{C}_{(v_0)}^+$ which contain w . Then there exists an integer $r > 0$ such that the power ϕ^{mr} fixes a face F_k of $\mathbb{S}\mathcal{C}_{(v_0)}^+$ for some $k \in I$.

Now we can choose a representation path γ of ϕ starting v_0 so that the first two edges are of the form $v_0 \xrightarrow{k} v_1 \xrightarrow{k} v_0$ by Remark 3.8 (a). Then the sign $\epsilon_\gamma(\phi^n(w))$ must contain a zero entry for all $n \in mr\mathbb{Z}_{>0}$, since the face F_k is contained in the hyperplane defined by $x_k^{(v_0)} = 0$. In particular the sign never stabilizes to a strict one, and hence γ is not sign-stable on $|\mathfrak{F}_s^+|$. Thus ϕ is not uniformly sign-stable. ■

6. SIGN STABILITY OF DEHN TWISTS

Here we discuss the sign stability of Dehn twists. Let C be a simple closed curve on a marked surface Σ satisfying the following condition:

There exists at least one marked point on each connected component of $\Sigma \setminus C$. (6.1)

On each side of C , choose a loop based at a marked point which is freely homotopic (forgetting the base-point) to the curve C so that these two loops form a tubular neighborhood $\mathcal{N}(C)$ of C . See Figure 24. Then $\mathcal{N}(C)$ can be regarded as a marked surface:




FIGURE 24. A tubular neighborhood $\mathcal{N}(C)$ with marked points.

an annulus with one marked point on each of its boundary component. Take a labeled triangulation of $\mathcal{N}(C)$ as shown in the top-left of Figure 20, and extend it arbitrarily to

a labeled triangulation (Δ_C, ℓ_C) of Σ . Then as we have seen in Example 4.7 (1), the path $\gamma_C : (\Delta_C, \ell_C) \xrightarrow{1} (\Delta', \ell') \xrightarrow{(1\ 2)} T_C^{-1}(\Delta_C, \ell_C)$ represents the mutation loop $T_C \in \Gamma_\Sigma$. Let us consider the $\mathbb{R}_{>0}$ -invariant set

$$\Omega_{(\Delta_C, \ell_C)}^{(1,2)} := \{\widehat{L} \in \mathcal{X}_{(\Delta_C, \ell_C)}^{\text{uf}}(\mathbb{R}^{\text{trop}}) \mid (x_1^{(\Delta_C, \ell_C)}(\widehat{L}), x_2^{(\Delta_C, \ell_C)}(\widehat{L})) \neq (0, 0)\},$$

which contains the subset $\Omega_{(\Delta_C, \ell_C)}^{\text{can}}$.

Proposition 6.1 (Sign stability of Dehn twists). *Let T_C be the Dehn twist along a simple closed curve C satisfying (6.1). Then the representation path $\gamma_C : (\Delta_C, \ell_C) \xrightarrow{1} (\Delta', \ell') \xrightarrow{(1\ 2)} T_C^{-1}(\Delta_C, \ell_C)$ is sign-stable on the $\mathbb{R}_{>0}$ -invariant set $\Omega_{(\Delta_C, \ell_C)}^{(1,2)}$. Its cluster stretch factor is 1.*

Proof. Since $b_{12} = -2$, the action of T_C on the coordinates $(x_1^{(\Delta_C, \ell_C)}(\widehat{L}), x_2^{(\Delta_C, \ell_C)}(\widehat{L}))$ is given by the example studied in Example 5.15 with $\ell = 2$. Since the sign of the path γ only concerns with the signs of these two coordinates, the first assertion follows. For the second statement, notice that the stable presentation matrix $E_{T_C}^{(\Delta_C, \ell_C)}$ is block-decomposed as follows:

$$E_{T_C}^{(\Delta_C, \ell_C)} = E_{\gamma_C}^{(+)} = \left(\begin{array}{cc|ccc} 2 & 1 & & & \\ -1 & 0 & & & \\ \hline & & * & & \\ & & & \ddots & \\ & & & & 1 \end{array} \right) \quad (6.2)$$

Here the first two directions correspond to $\{1, 2\}$. Then we get $\lambda_\phi = \rho(E_{T_C}^{(\Delta_C, \ell_C)}) = 1$ as desired. \blacksquare

Note that the limit point $L_{T_C} \in \mathcal{U}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}})$ given by $x_1^{(\Delta_C, \ell_C)}(L_{T_C}) = -x_2^{(\Delta_C, \ell_C)}(L_{T_C}) = 1$ and $x_j^{(\Delta_C, \ell_C)}(L_{T_C}) = 0$ for $j \notin \{1, 2\}$ is identified with the curve C (naturally regarded as a rational lamination) via the isomorphism given in Theorem 4.3.⁵

Remark 6.2. In fact, the $\mathbb{R}_{>0}$ -invariant set above is geometrically described as follows:

$$\Omega_{(\Delta_C, \ell_C)}^{(1,2)} = \{\widehat{L} \in \mathcal{X}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}}) \mid \mathcal{I}_C(\widehat{L}) \neq 0\} \sqcup \mathbb{R}_{>0}\{C\}.$$

Here $\mathcal{I}_C : \mathcal{X}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}}) \rightarrow \mathbb{R}$ is the PL function given by the weighted sum of the usual intersection number for a rational lamination, which can be continuously extended [FG06a, Corollary 12.2]. In the coordinates, it is given by

$$\mathcal{I}_C \circ x_{(\Delta_C, \ell_C)}^{-1} = \max \left\{ -\frac{x_1 + x_2}{2}, \frac{x_1 - x_2}{2}, \frac{x_1 + x_2}{2} \right\}.$$

Here $x_i := x_i^{(\Delta_C, \ell_C)}$ for $i = 1, 2$. Indeed, it can be obtained as the tropical limit of the trace function along C on the enhanced Teichmüller space [FG06a, Theorem 12.2].

⁵The lamination-shear coordinates of C are given by $x_{\ell_C(1)}^{\Delta_C}(C) = -x_{\ell_C(2)}^{\Delta_C}(C) = -1$ and $x_{\ell_C(j)}^{\Delta_C}(C) = 0$ for $j \notin \{1, 2\}$. See Figure 20.

Example 6.3. Let Σ be a once-punctured surface of genus 2. Let $C \subset \Sigma$ be the closed curve shown in Figure 25, and consider the Dehn twist T_C along C . A labeled triangulation (Δ, ℓ) of Σ is shown in the left of Figure 26. Note that the arcs 3 and 7 bounds a tubular neighborhood $\mathcal{N}(C)$ of C .




FIGURE 25. Once-punctured surface of genus 2. Here only some of the arcs in the labeled triangulation (Δ, ℓ) are shown.




FIGURE 26. Dehn twist along C .

Then the following path γ is a representation path for T_C :

$$\gamma : (\Delta, \ell) \xrightarrow{6} (\Delta', \ell') \xrightarrow{(2 \ 6)} T_C^{-1}(\Delta, \ell).$$

Proposition 6.1 says that this path is sign-stable on $\Omega_{(\Delta, \ell)}^{(6, 2)}$ with the stretch factor 1. One can verify that its stable sign is $(+)$.

Dependence of the sign stability on representation paths. As an aside, we give a counterexample to the following conjecture in [IK21] which asserts that the sign stability is an intrinsic property of a mutation loop:

Conjecture 6.4 ([IK21, Conjecture 1.3]). Let $\gamma_i : v_i \rightarrow v'_i$ ($i = 1, 2$) be two paths which represent the same mutation loop $\phi := [\gamma_1]_{\mathbf{s}} = [\gamma_2]_{\mathbf{s}}$. Then γ_1 is sign-stable if and only if γ_2 is.

Here is a counterexample. Let us consider the Dehn twist T_C as in Example 6.3. It has the following representation path:

$$\gamma' : (\Delta, \ell) \xrightarrow{(6,9,9)} (\Delta', \ell') \xrightarrow{(2,6)} T_C^{-1}(\Delta, \ell).$$

In other words, $T_C = [\gamma]_{\mathbf{s}_\Sigma} = [\gamma']_{\mathbf{s}_\Sigma}$. While the path γ is sign-stable on $\Omega_{(\Delta, \ell)}^{(6,2)}$, the path γ' is not. Indeed, its sign at $C \in \Omega_{(\Delta, \ell)}^{(6,2)}$ is $(-, 0, 0)$ and thus not strict. This is a quite general phenomenon: it can occur when the domain $\Omega = \Omega_{(\Delta, \ell)}^{(6,2)}$ of stability contains a non- \mathcal{X} -filling fixed point (recall Definition 2.15). What is worse, it has different strict signs at $L_1 := C \sqcup \delta_1$ and $L_2 := C \sqcup \delta_2$, where δ_i are closed curves shown in Figure 26. Since $C \in \Omega_{(\Delta, \ell)}^{(6,2)}$, $L_1, L_2 \in \Omega_{(\Delta, \ell)}^{(6,2)}$ and the signs of γ' at these points are $(+, +, +)$ and $(+, -, -)$, respectively.

7. SIGN STABILITY OF PSEUDO-ANOSOV MAPPING CLASSES: EMPTY BOUNDARY CASE

Let X be a compact Hausdorff space and $f : X \rightarrow X$ be a homeomorphism. We say that the discrete dynamical system (X, f) has *North-South dynamics* (*NS dynamics* for short) if there exist two distinct fixed points $x_f^+, x_f^- \in X$ of f such that for any point $x \in X$, the followings hold:

- if $x \neq x_f^-$, then $\lim_{n \rightarrow \infty} f^n(x) = x_f^+$;
- if $x \neq x_f^+$, then $\lim_{n \rightarrow \infty} f^{-n}(x) = x_f^-$.

The point x_f^+ (resp. x_f^-) is called the attracting (resp. repelling) point of f .

From this section onwards, we often abbreviate the labeled triangulation $(\Delta, \varsigma_\Delta)$ to Δ for simplicity. Note that in the case Σ has no boundary components, the seed pattern \mathbf{s}_Σ has no frozen indices (*i.e.*, those corresponding to boundary arcs), and hence $\mathcal{X}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}}) = \mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}})$ and $\mathcal{U}_\Sigma^{\text{uf}}(\mathbb{R}^{\text{trop}}) = \mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$. Therefore we will drop the symbol ‘uf’ in this section.

For $\mathcal{Z} \in \{\mathcal{A}, \mathcal{X}, \mathcal{U}\}$, let $\mathbb{S}\mathcal{Z}_\Sigma(\mathbb{R}^{\text{trop}}) := (\mathcal{Z}_\Sigma(\mathbb{R}^{\text{trop}}) \setminus \{0\})/\mathbb{R}_{>0}$. A subset of $\mathbb{S}\mathcal{Z}_\Sigma(\mathbb{R}^{\text{trop}})$ is identified with a scaling-invariant subset of $\mathcal{Z}_\Sigma(\mathbb{R}^{\text{trop}})$. Our aim in this section is to prove the following:

Theorem 7.1. *Let Σ be a punctured surface. For a mapping class $\phi \in MC(\Sigma) \subset \Gamma_\Sigma$, the following conditions are equivalent:*

- (1) *The mapping class ϕ is pseudo-Anosov.*
- (2) *The mutation loop ϕ is uniformly sign-stable.*
- (3) *The mapping class ϕ has NS dynamics on $\mathbb{S}\mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}})$ and the attracting and repelling points are \mathcal{X} -filling.*

In this case, the cluster stretch factor of the mutation loop ϕ coincides with the stretch factor λ_ϕ of the pA mapping class ϕ .

The similar statement for $\mathcal{A}_\Sigma(\mathbb{R}^{\text{trop}})$ as (3) is not equivalent to the others [IK20]. It is one of the reasons why we use the tropical \mathcal{X} -variety rather than the \mathcal{A} -variety in the definition of sign stability.

7.1. Sign stability and the pA property. We prove the equivalence of (1) and (2) in Theorem 7.1. First we recall the following classical theorem due to W.Thurston:

Theorem 7.2 ([Th88]). *Let Σ be a punctured surface and $\phi \in MC(\Sigma)$ a mapping class. Then the mapping class ϕ is pA if and only if the action of ϕ on $\mathbb{S}\mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$ has NS dynamics. Moreover, the attracting and repelling fixed points are given by the pA pair of ϕ .*

We call a mapping class *pure* if it fixes each puncture. Let $PMC(\Sigma)$ denote the normal subgroup consisting of the pure mapping classes, which fits into the following exact sequence:

$$1 \rightarrow PMC(\Sigma) \rightarrow MC(\Sigma) \xrightarrow{\sigma_\Sigma} \mathfrak{S}_h \rightarrow 1.$$

The following lemma is a direct consequence of the Ivanov's work [Iva].

Lemma 7.3. *If a pA mapping class ϕ is pure, then the action of ϕ on $\mathbb{S}\mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}})$ has NS dynamics, which is an extension of that on $\mathbb{S}\mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$.*

Proof. Let $D\Sigma$ be the closed surface obtained as the double of Σ . It is obtained by gluing two copies of the bordered surface $\Sigma^\circ = \Sigma \setminus \bigsqcup_{p \in P} D_p$ (see Section 2.4) with opposite orientations along the corresponding boundary components. The surface $D\Sigma$ is equipped with an orientation-reversing involution ι which interchanges the two copies of Σ° . A hyperbolic structure F as in Section 2.5 is canonically extended to an ι -invariant hyperbolic structure DF on $D\Sigma$. Let $\mathcal{ML}(DF)$ denote the space of measured geodesic laminations on DF . We can identify $\mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}}) \cong \widehat{\mathcal{ML}}(F)$ by Theorems 2.21 and 4.3.

For each sign $\sigma = (\sigma_p)_{p \in P} \in \{+, -\}^P$, let us consider the subset

$$\widehat{\mathcal{ML}}^{(\sigma)}(F) := \{(G, \mu) \in \widehat{\mathcal{ML}}(F) \mid \sigma_p \sigma_G(p) \geq 0 \text{ for } p \in P\},$$

which is preserved by the action of $PMC(\Sigma)$.

Fix $\sigma \in \{+, -\}^P$. Let us consider the embedding

$$D : \widehat{\mathcal{ML}}^{(\sigma)}(F) \hookrightarrow \mathcal{ML}(DF)$$

defined as follows. For each measured geodesic lamination in $\widehat{\mathcal{ML}}^{(\sigma)}(F)$, take its ι -invariant extension to the double DF . Then on the both sides of the boundary component ∂_p for $p \in P$ in DF , spiralling leaves come in pair related by the involution ι . We first straighten these leaves so that they hit the boundary component ∂_p transversely, and

then glue each ι -related pair of leaves together to get a measured geodesic lamination on the double DF . By Lemma 2.18, the resulting map D is injective. This construction is $PMC(\Sigma)$ -equivariant, where the action on $\mathcal{ML}(\Sigma)$ is through the group embedding

$$D : MC(\Sigma) \xrightarrow{\sim} MC(D\Sigma)^\iota \subset MC(D\Sigma).$$

Note that for each pA mapping class $\phi \in PMC(\Sigma)$, the mapping class $D\phi$ is “pure” in the sense of [Iva]. Let $L_\phi^+, L_\phi^- \in \mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$ be some representatives of the attracting and repelling points of the NS dynamics of $\phi \in PMC(\Sigma)$ on $\mathbb{S}\mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}}) = \mathcal{PML}(F)$, respectively. Then [Iva, Theorem A.1] (and a more detailed description given in [Iva, Lemma A.4]) tells us that

$$[D(\phi^n(\widehat{L}))] = [(D\phi)^n(D\widehat{L})] \xrightarrow{n \rightarrow \infty} [D(L_\phi^+)]$$

for any $\widehat{L} \in \mathcal{X}_\Sigma^{(\sigma)}(\mathbb{R}^{\text{trop}}) \setminus \mathbb{R}_{\geq 0}L_\phi^-$, where

$$\mathcal{X}_\Sigma^{(\sigma)}(\mathbb{R}^{\text{trop}}) := \{\widehat{L} \in \mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}}) \mid \sigma_p \theta_p(\widehat{L}) \geq 0, \text{ for } p \in P\} \cong \widehat{\mathcal{ML}}^{(\sigma)}(\Sigma).$$

Since D is an embedding and σ was arbitrary, the proof is completed. \blacksquare

Proof of (1) \iff (2) in Theorem 7.1. Let $\phi \in MC(\Sigma)$ be a pA mapping class and take a representation path $\gamma : (\Delta, \ell) \rightarrow \phi^{-1}(\Delta, \ell)$ in $\mathbb{E}\text{ch}_\Sigma$. By Theorem 7.2 the mapping class ϕ has NS dynamics on $\mathbb{S}\mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$, whose attracting and repelling points are written as $[L_\phi^+]$ and $[L_\phi^-]$, respectively. By Proposition 2.14(1), the sign $\epsilon_\gamma(L_\phi^+) =: \epsilon_\gamma^+$ of the point L_ϕ^+ is strict. Therefore, the cone

$$\mathcal{C}_{(\Delta, \ell)}^{\epsilon_\gamma^+} := \overline{\{\widehat{L} \in \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) \mid \epsilon_\gamma(\widehat{L}) = \epsilon_\gamma^+\}}$$

has the maximal dimension and $L_\phi^+ \in \text{int } \mathcal{C}_{(\Delta, \ell)}^{\epsilon_\gamma^+}$. Moreover, since $[L_\phi^+]$ is a fixed point of ϕ ,

$$\epsilon_{\gamma^r}(L_\phi^+) = \overbrace{(\epsilon_\gamma^+, \dots, \epsilon_\gamma^+)}^r =: \epsilon_{\gamma^r}^{+,r} \quad (7.1)$$

for any $r \in \mathbb{Z}_{>0}$. Hence $L_\phi^+ \in \text{int } \mathcal{C}_{(\Delta, \ell)}^{\epsilon_{\gamma^r}^{+,r}} \subset \text{int } \mathcal{C}_{(\Delta, \ell)}^{\epsilon_\gamma^+}$.

Take $r \in \mathbb{Z}_{>0}$ so that the mapping class ϕ^r is pure. It has NS dynamics on $\mathbb{S}\mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}})$ and its attracting and repelling points are $[L_\phi^+]$ and $[L_\phi^-]$ respectively by Lemma 7.3. Hence for any $\widehat{L} \in \mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}}) \setminus [L_\phi^-]$, there exists a positive integer $m_0 \in \mathbb{Z}$ such that $\phi^{mr}(\widehat{L}) \in \text{int } \mathcal{C}_{(\Delta, \ell)}^{\epsilon_{\gamma^r}^{+,r}}$ for all integer $m \geq m_0$. By (7.1), it implies that $\phi^n(\widehat{L}) \in \text{int } \mathcal{C}_{(\Delta, \ell)}^{\epsilon_\gamma^+}$ for all $n \geq m_0 r$. It is nothing but the sign stability of ϕ with the stable sign ϵ_γ^+ .

Conversely, assume that ϕ is not pA. Then by Nielsen–Thurston classification (Theorem 2.11), it is either periodic or reducible. In the periodic case, ϕ^r is the identity for some integer $r > 0$ so cannot be uniformly sign-stable.

In the reducible case, ϕ fixes a multicurve $L \in \mathcal{U}_\Sigma(\mathbb{Z}^{\text{trop}}) \subset \mathcal{X}_\Sigma(\mathbb{Z}^{\text{trop}})$. Then Proposition 2.14(2) tells us that there exists an ideal triangulation Δ and an interior arc $\alpha \in \Delta_{\text{uf}}$

such that $x_\alpha^\Delta(L) = 0$. By Remark 3.8, we can choose a representation path of the form

$$\gamma : (\Delta, \ell) \xrightarrow{\ell^{-1}(\alpha)} (\Delta', \ell) \xrightarrow{\ell^{-1}(\alpha)} (\Delta, \ell) - \dots$$

for some labeling ℓ (*i.e.*, change the basepoint to (Δ, ℓ) and insert a repeated flip along the arc α). Then $\epsilon_\gamma(L)$ is not a strict sign and hence ϕ cannot be uniformly sign-stable. \blacksquare

Remark 7.4. From the proof, the following weaker version of (2) is also equivalent to the conditions (1) and (2) in Theorem 7.1:

- (2') Any representation path $\gamma : (\Delta, \ell) \rightarrow (\Delta', \ell')$ of ϕ contained in the subgraph $\text{Tri}(\Sigma) \subset \text{Tri}^\square(\Sigma) \cong \mathbb{E}\text{ch}_\Sigma$ consisting of labeled ideal triangulations is sign-stable on $\Omega_{(\Delta, \ell)}^\mathbb{Q}$.

Let us record the argument used in the proof as follows:

Proposition 7.5. *Let \mathbf{s} be any seed pattern, and $\phi \in \Gamma_{\mathbf{s}}$ a mutation loop.*

- (i) *Assume that there exists an integer $r > 0$ such that the mutation loop ϕ^r has NS dynamics on $\mathbb{S}\mathcal{X}_{\mathbf{s}}(\mathbb{R}^{\text{trop}})$, and the attracting/repelling points $[L_\phi^\pm]$ are \mathcal{X} -filling and fixed by ϕ . Then ϕ is uniformly sign-stable.*
- (ii) *Conversely, if ϕ is uniformly sign-stable, then any positive power ϕ^r never have a fixed point $[L] \in \mathbb{S}\mathcal{X}_{\mathbf{s}}(\mathbb{Q}^{\text{trop}})$ which is not \mathcal{X} -filling. In particular, the mutation loop ϕ is cluster-pA.*

The statement (ii) is slightly stronger than Proposition 5.20, since a point contained in a face of $|\mathfrak{F}_{\mathbf{s}}^+|$ is not \mathcal{X} -filling but the converse is not true. For instance, the simple closed curve C in Figure 26 is not \mathcal{X} -filling but does not belong to any face of the Fock–Goncharov cluster complex.

7.2. NS dynamics on the space of \mathcal{X} -laminations and the pA property. Our aim here is to prove the equivalence of (1) and (3) in Theorem 7.1. From Theorem 7.2, the implication (3) \implies (1) is clear since the action of a mapping class on $\mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}})$ restricts to that on $\mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$. It remains to prove the implication (1) \implies (3). Although this statement seems to be well-known to specialists, we give a proof here based on the sign stability of a pA mapping class (*i.e.*, we are actually going to show (1) & (2) \implies (3)).

Let $\phi \in MC(\Sigma)$ be a pA mapping class. From the implication (1) \implies (2) in Theorem 7.1, any representation path $\gamma : (\Delta, \ell) \rightarrow \phi^{-1}(\Delta, \ell)$ of ϕ is sign-stable on $\Omega_{(\Delta, \ell)}^\mathbb{Q}$. Then for each point $\widehat{L} \in \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}})$, there exists a positive integer n_0 such that $\phi^n(\widehat{L}) \in \text{int } \mathcal{C}_\gamma^{\text{stab}}$ for all $n \geq n_0$, where $\mathcal{C}_\gamma^{\text{stab}} := \mathcal{C}_{(\Delta, \ell)}^{\epsilon_\gamma^+}$. Thus we have

$$\mathbf{x}_{(\Delta, \ell)}(\phi^n(\phi^{n_0}(\widehat{L}))) = (E_\gamma^{\epsilon_\gamma^+})^n \mathbf{x}_{(\Delta, \ell)}(\phi^{n_0}(\widehat{L}))$$

for all $n \geq 0$, where $\mathbf{x}_{(\Delta, \ell)}(w) := (x_\alpha^{(\Delta, \ell)}(w))_\alpha$ denotes the coordinate vector of a point $w \in \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}})$. In other words, the action of ϕ is a linear dynamical system in the stable range $n \geq n_0$.

Recall from Lemma 5.1 that the presentation matrix $E_\gamma^{e_\gamma^\dagger}$ also presents the tangent map $(d\phi)_L : T_L \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) \rightarrow T_{\phi(L)} \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}})$ for any $L \in \text{int } \mathcal{C}_\gamma^{\text{stab}}$, and the linear action of ϕ on the cone $\mathcal{C}_\gamma^{\text{stab}}$ is the same as $(d\phi)_L|_{t_L(\mathcal{C}_\gamma^{\text{stab}})}$. We are going to give a certain block-decomposition of this tangent map. Since the tropicalized Casimir map $\theta^{\text{trop}} : \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) \rightarrow \mathbb{R}^h$ is a linear map, for any differentiable point $L \in \mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$ we have an exact sequence

$$0 \rightarrow T_L \mathcal{U}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) \rightarrow T_L \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) \xrightarrow{(d\theta^{\text{trop}})_L} \mathbb{R}^h \rightarrow 0$$

and a similar sequence for tangent spaces at $\phi(L)$, by differentiating the cluster exact sequence (4.2) (or eq. (2.5)). From Proposition 4.6 we see that these sequences are $(d\phi)_L$ -equivariant, where the action on \mathbb{R}^h is given by $\sigma_\Sigma(\phi)$. Therefore by choosing sections $s_L : \mathbb{R}^h \rightarrow T_L \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}})$ and $s_{\phi(L)} : \mathbb{R}^h \rightarrow T_{\phi(L)} \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}})$, we get a block-decomposition of the form

$$(d\phi)_L = \begin{pmatrix} (d\phi^u)_L & * \\ 0 & \sigma_\Sigma(\phi) \end{pmatrix} \quad (7.2)$$

with respect to the direct sum decompositions $T_L \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) = T_L \mathcal{U}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) \oplus s_L(\mathbb{R}^h)$ and $T_{\phi(L)} \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) = T_{\phi(L)} \mathcal{U}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) \oplus s_{\phi(L)}(\mathbb{R}^h)$. Here ϕ^u is the restriction of the action of ϕ to $\mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$. By Theorem 7.1 we know that the linear action of ϕ^u on the cone $\mathcal{C}_\gamma^{\text{stab}} \cap \mathcal{U}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}})$ has NS dynamics. Interpreting this in terms of the corresponding presentation matrix, our assertion is reduced to the following lemma, which is just a matter of linear algebra:

Lemma 7.6. *Let N, N', h be positive integers satisfying $N = N' + h$ and let $E \in \text{GL}(N; \mathbb{R})$ be a matrix of the form*

$$E = \begin{pmatrix} A & B \\ 0 & D \end{pmatrix}, \quad A \in \text{GL}(N'; \mathbb{R}), \quad B \in \text{Mat}(h \times N'; \mathbb{R}), \quad D \in \text{GL}(h; \mathbb{R}).$$

Assume that

- (a) the matrix D has finite order s ;
- (b) there exists a cone $\mathcal{C} \subset \mathbb{R}^N$ which is invariant under the action of E ;
- (c) there exists a unique attracting point $[u_+] \in \mathbb{S}(\mathcal{C} \cap \mathbb{R}^{N'})$ for the action of A on $\mathbb{S}(\mathcal{C} \cap \mathbb{R}^{N'})$, where $u_+ \in \mathbb{R}^{N'}$ is a unit eigenvector of A with eigenvalue $\lambda > 1$.

Then for any point $[u] \in \mathbb{S}\mathcal{C}$, we have $E^n([u]) \rightarrow [u_+]$ as $n \rightarrow \infty$.

Proof. We have

$$E^n(w) = \left(A^n(u) + \sum_{k=0}^{n-1} A^{n-k-1} B D^k(v), D^n(v) \right)$$

for $w = (u, v) \in \mathbb{R}^N = \mathbb{R}^{N'} \times \mathbb{R}^h$ and $n \geq 0$. Fix any $\varepsilon > 0$.

For $u \in \mathcal{C} \cap \mathbb{R}^{N'}$, let $\mathbf{a}(u) := \lim_{n \rightarrow \infty} \|A^n(u)/\lambda^n\|$. From the assumption (c) it converges, and there exists an integer $n(u)$ such that

$$\left\| \frac{A^n(u)}{\lambda^n} - \mathbf{a}(u)u_+ \right\| < \varepsilon$$

for all $n \geq n(u)$. For any $w = (u, v) \in \mathcal{C} \subset \mathbb{R}^N = \mathbb{R}^{N'} \times \mathbb{R}^h$, we have

$$\begin{aligned} & \left\| \frac{E^n(w)}{\lambda^n} - \left(\mathbf{a}(u) + \sum_{k=0}^{n-1} \frac{\mathbf{a}(BD^k(v))}{\lambda^{k+1}} \right) u_+ \right\| \\ & \leq \left\| \frac{A^n(u)}{\lambda^n} - \mathbf{a}(u)u_+ \right\| + \sum_{k=0}^{n-1} \frac{1}{\lambda^{k+1}} \left\| \frac{A^{n-k-1}(BD^k(v))}{\lambda^{n-k-1}} - \mathbf{a}(BD^k(v))u_+ \right\| + \left\| \frac{D^n(v)}{\lambda^n} \right\|. \end{aligned}$$

The first term is ε -small for all $n \geq n(u)$. The third term is ε -small for all $n \geq n_1(v)$, where $n_1(v)$ is chosen so that $\lambda^{-n_1(v)} \max_{k=0, \dots, s-1} \|D^k(v)\| < \varepsilon$. In order to estimate the second term, let $n_2(v) := \max_{k=0, \dots, s-1} n(BD^k(v))$ and

$$M := \max_{m=0, \dots, n_2(v)-1} \max_{k=0, \dots, s-1} \left\| \frac{A^m(BD^k(v))}{\lambda^m} - \mathbf{a}(BD^k(v))u_+ \right\|.$$

For any $n > n_2(v)$, we have

$$\begin{aligned} & \sum_{k=0}^{n-1} \frac{1}{\lambda^{k+1}} \left\| \frac{A^{n-k-1}(BD^k(v))}{\lambda^{n-k-1}} - \mathbf{a}(BD^k(v))u_+ \right\| \\ & \leq \sum_{k=0}^{n-n_2(v)-1} \frac{1}{\lambda^{k+1}} \varepsilon + \sum_{k=n-n_2(v)}^{n-1} \frac{1}{\lambda^{k+1}} M \leq \frac{1}{\lambda-1} \varepsilon + \frac{M}{\lambda^n} \sum_{k=1}^{n_2(v)} \lambda^{k-1}. \end{aligned}$$

It is bounded from above by $(1/(\lambda-1) + 1)\varepsilon$ if moreover $n \geq n_3(v)$, where $n_3(v)$ is chosen so that $\lambda^{-n_3(v)} M \sum_{k=1}^{n_2(v)} \lambda^{k-1} < \varepsilon$. Summarizing, we get

$$\left\| \frac{E^n(w)}{\lambda^n} - \left(\mathbf{a}(u) + \sum_{k=0}^{n-1} \frac{\mathbf{a}(BD^k(v))}{\lambda^{k+1}} \right) u_+ \right\| < \left(3 + \frac{1}{\lambda-1} \right) \varepsilon$$

for all $n > \max\{n(u), n_1(v), n_2(v), n_3(v)\}$.

Observe that the infinite sum $\sum_{k=0}^{\infty} \mathbf{a}(BD^k(v))/\lambda^{k+1}$ converges since D has finite order, and thus we get

$$\left\| \frac{E^n(w)}{\lambda^n} - \left(\mathbf{a}(u) + \sum_{k=0}^{\infty} \frac{\mathbf{a}(BD^k(v))}{\lambda^{k+1}} \right) u_+ \right\| < \left(4 + \frac{1}{\lambda-1} \right) \varepsilon$$

for sufficiently large n . Hence $[E^n(w)] \rightarrow [u_+]$ as desired. \blacksquare

Proof of (1) \implies (3) in Theorem 7.1. Let $N := N(\Sigma)$, $N' := \dim \mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$, $\mathcal{C} := \mathcal{C}_\gamma^{\text{stab}}$, and E the block-decomposed matrix (7.2) (with a basis compatible with the decomposition). The assumptions are satisfied by Theorem 7.2, where $[u_+]$ is the coordinate vector of $[\mathcal{F}_+, \mu_+]$ and the corresponding eigenvalue is given by the stretch factor $\lambda_\phi > 1$. Hence ϕ has a unique attracting point in $\mathbb{S}\mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}})$.

By considering the action of ϕ^{-1} near the class $[\mathcal{F}_-, \mu_-]$, the same argument can be applied and we see that ϕ has a unique repelling point. Since the pA pair consists of arational foliations, the corresponding points in $\mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}})$ are \mathcal{X} -filling Proposition 2.14. Thus the assertion is proved. \blacksquare

Finally, we verify the coincidence of the stretch factor of the pA mapping class ϕ and the cluster stretch factor of the corresponding mutation loop ϕ . It is known that the former is equal to the spectral radius of the *transition matrix* of the *invariant train track* τ_ϕ of ϕ (see [BH95]). A train track τ is a certain trivalent graph on Σ equipped with a smoothing structure on each vertex, see Figure 27. For a detail, we refer the reader to [PH], or the appendix of [IK20]. Let $V(\tau) \subset \mathbb{R}^{\{\text{edges of } \tau\}}$ denote the cone of measures on τ (a linear subspace defined by the *switch conditions*), which is identified with a subcone of $\mathcal{MF}(\Sigma) \cong \mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$. It is said to be ϕ -invariant if $\phi(\tau)$ can be collapsed to τ preserving the smoothing structure at vertices. In particular, we have $L_\phi^+ \in V(\tau_\phi)$ and $L_\phi^+ \in V(\phi(\tau_\phi))$. The transition matrix is a certain linear isomorphism $T_\phi : \mathbb{R}^{\{\text{edges of } \phi(\tau_\phi)\}} \rightarrow \mathbb{R}^{\{\text{edges of } \tau_\phi\}}$ describing this collapsing such that $T_\phi(V(\phi(\tau_\phi))) \subset V(\tau_\phi)$ and $T_\phi|_{V(\phi(\tau_\phi))} = \phi^u|_{V(\phi(\tau_\phi))}$. Considering the exact sequence

$$0 \rightarrow T_{L_\phi^+} V(\phi(\tau_\phi)) \rightarrow T_{L_\phi^+} \mathbb{R}^{\{\text{edges of } \tau_\phi\}} \rightarrow T_{L_\phi^+} \mathbb{R}^{\{\text{edges of } \tau_\phi\}} / T_{L_\phi^+} V(\tau_\phi) \rightarrow 0$$

and taking a section of $T_{L_\phi^+} \mathbb{R}^{\{\text{edges of } \tau_\phi\}} / T_{L_\phi^+} V(\tau_\phi)$, we obtain a block decomposition of T_ϕ . The submatrix of T_ϕ corresponding to $T_{L_\phi^+} V(\phi(\tau_\phi)) \cong T_{L_\phi^+} \mathcal{U}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}})$ is nothing but $(d\phi^u)_{L_\phi^+}$ by taking a suitable basis [BBK12]. Thus, the spectral radius of $(d\phi^u)_{L_\phi^+}$, which is the cluster stretch factor by (7.2), coincides with the spectral radius of the transition matrix T_ϕ .




FIGURE 27. Switch condition.

The proof of Theorem 7.1 is completed.

Mapping classes with reflections. We say that a mutation loop $\phi \in \Gamma_\Sigma$ is *pseudo-Anosov* if its underlying mapping class is pseudo-Anosov. Then the following is a direct consequence of Proposition 7.5, Lemma 7.6 and (4.4):

Corollary 7.7. *Let Σ be a punctured surface. For a mutation loop $\phi \in \Gamma_\Sigma$, the following conditions are equivalent:*

- (1) *The mutation loop ϕ is pseudo-Anosov.*
- (2) *The mutation loop ϕ is uniformly sign-stable.*

(3) The mutation loop ϕ has NS dynamics on $\mathbb{S}\mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}})$, and its attracting and repelling points are \mathcal{X} -filling.

In this case, the cluster stretch factor of the mutation loop ϕ coincides with the stretch factor of the underlying pA mapping class of ϕ .

Proof. (1) \iff (2): This follows from Proposition 7.5 with a notice that the action of a mutation loop on $\mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$ coincides with that of its underlying mapping class.

(1) \iff (3): With the same notice, this follows from (4.4) and Lemma 7.6. \blacksquare

Remark 7.8. For a subset $P' \subset P$, let $\mathbb{E}\text{ch}_\Sigma^{P'} \subset \mathbb{E}\text{ch}_\Sigma$ be the subgraph consisting of labeled tagged triangulations $(\Delta, \varsigma_\Delta, \ell)$ such that $(\varsigma_\Delta)_p = +$ for all $p \in P'$. For any mutation loop $(\phi, \varsigma) \in \Gamma_\Sigma$ with $\varsigma_p = +$ for all $p \in P'$, the following condition is also equivalent to the conditions (1)–(3) in Corollary 7.7:

(2') Any representation path $\gamma : (\Delta, \ell) \rightarrow (\Delta', \ell')$ of ϕ contained in the subgraph $\mathbb{E}\text{ch}_\Sigma^{P'}$ is sign-stable on $\Omega_{(\Delta, \ell)}^\mathbb{Q}$.

Our results Theorem 7.1 and Corollary 7.7 is proved by taking two steps (1) \iff (2) and (1) \iff (3). Also note that under the condition (1), the condition (2) is equivalent to the following stronger condition:

(2[#]) The mutation loop ϕ is uniformly sign-stable and cluster stretch factor is greater than 1.

We expect that there is a direct proof of (2[#]) \iff (3). As a generalization, we conjecture the following:

Conjecture 7.9. Let \mathbf{s} be any seed pattern and $\phi \in \Gamma_{\mathbf{s}}$. Then ϕ is uniformly sign-stable with cluster stretch factor greater than 1 if and only if ϕ has NS dynamics on $\mathbb{S}\mathcal{X}_{\mathbf{s}}(\mathbb{R}^{\text{trop}})$ with \mathcal{X} -filling attracting/repelling points.

The stronger condition (2[#]) excludes the mutation loop studied in Example 5.15 for $\ell = 2$, which is uniformly sign-stable by Example 5.18 but it does not have NS dynamics.

Example 7.10. Here is an example of a pA mapping class. Let us observe the convergence of signs. A fourth-punctured sphere ($g = 0, h = 4$) with a labeled ideal triangulation (Δ, ℓ) is shown in Figure 28. Consider the mapping class $\phi := h_2 \circ h_5^{-1}$, where h_5^{-1} is the left half-twist along the arc $\ell(5)$ and h_2 is the right half-twist along $\ell(2)$. It is an example of a pA mapping class, which realizes the smallest stretch factor among those in $MC(\Sigma)$. See [FM, Section 15.1]. We show representation paths of h_2 and h_5^{-1} in Figure 29. One can verify that the concatenation of these paths is \mathbf{s}_Σ -equivalent to the path

$$\gamma : (\Delta, \ell) \xrightarrow{(1,5,3,2)} (\Delta', \ell') \xrightarrow{\sigma} \phi^{-1}(\Delta, \ell),$$

where $\sigma = (1\ 5\ 6) \circ (2\ 4\ 3) \in \mathfrak{S}_6$ and $(\Delta', \ell') \xrightarrow{\sigma} \phi^{-1}(\Delta, \ell)$ denotes the sequence of vertical edges corresponding to any reduced expression of σ . Thus, γ is a representation path of ϕ .




FIGURE 28. A fourth-punctured sphere with a labeled triangulation. A planar picture is shown in the right, regarding $\mathbb{S}^2 = \mathbb{R}^2 \cup \{\infty\}$.




FIGURE 29. The representation paths of h_2 and h_5^{-1} .

By taking the following steps, we can compute the coordinates of the representatives L_ϕ^\pm of the attracting/repelling points:

- (1) Compute the domains of linearity of the PL map $\phi^x : \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) \rightarrow \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}})$ and the corresponding presentation matrices.
- (2) Compute the eigenvectors of these matrices and find the ones which are contained in the corresponding domains of linearity. (Indeed, Theorem 7.1 (3) guarantees that there exist precisely two such eigenvectors.)

The point L_ϕ^+ (resp. L_ϕ^-) corresponds to an eigenvector whose eigenvalue is larger (resp. smaller) than 1. In this case, there are 16 domains of linearity of ϕ^x . By computing the eigenvectors of the corresponding presentation matrices, one obtains the coordinates of

L_ϕ^\pm as ⁶

$$\mathbf{x}_{(\Delta, \ell)}(L_\phi^\pm) = \pm \left(\frac{-1 + \sqrt{5}}{2}, -1, 1, \frac{1 - \sqrt{5}}{2}, 1, -1 \right).$$

Theorem 7.1 tells us that the signs at every orbit in $\mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}}) \setminus \mathbb{R}_{\geq 0} L_\phi^-$ converge to the stable sign $\epsilon_\gamma^{\text{stab}} := \epsilon_\gamma(L_\phi^+) = (+, -, -, +)$. For example, the convergent behavior of signs at the orbit of the point $l_\Delta^\pm \in \mathcal{C}_\Delta^\pm \subset \mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}})$ defined by $\mathbf{x}_{(\Delta, \ell)}(l_\Delta^\pm) = (\pm 1, \dots, \pm 1)$, see Lemma 4.4, is shown in Table 1.

TABLE 1. Orbit of signs for γ .

n	$\epsilon_\gamma(\phi^n(l_\Delta^+))$	$\epsilon_\gamma(\phi^n(l_\Delta^-))$
1	(+, +, +, +)	(-, -, -, -)
2	(+, -, +, +)	(-, -, -, +)
3	(+, -, -, +)	(+, -, -, +)
4	(+, -, -, +)	(+, -, -, +)
5	(+, -, -, +)	(+, -, -, +)
\vdots	\vdots	\vdots

For any reflection $\tau \in (\mathbb{Z}/2)^4$, the pA mutation loop (ϕ, τ) should indicate the same convergence behavior of signs at every orbit in $\mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}}) \setminus \mathbb{R}_{\geq 0} L_\phi^-$ by Corollary 7.7. For instance, consider the reflection $\tau_p \in \Gamma_\Sigma$ at the puncture $p \in P$ shown in Figure 28, which is represented by the path

$$\gamma_p : (\Delta, \ell) \xrightarrow{(4,2,1,5,2,4)} (\Delta'', \ell'') \xrightarrow{(1\ 5)} \tau_p^{-1}(\Delta, \ell).$$

One can verify that the path

$$\gamma' : (\Delta, \ell) \xrightarrow{(1,5,6,1,3,2)} (\Delta''', \ell''') \xrightarrow{\sigma'} (\tau_p \phi)^{-1}(\Delta, \ell)$$

is \mathbf{s}_Σ equivalent to $\gamma * \gamma_p$ with $\sigma' := (1\ 5) \circ \sigma$. The convergent behavior of signs at the orbit of $l_\Delta^\pm \in \mathcal{X}_\Sigma(\mathbb{R}^{\text{trop}})$ is shown in Table 2. In fact, the stable sign is given by $\epsilon_{\gamma'}^{\text{stab}} := \epsilon_{\gamma'}(L_\phi^+) = (+, -, +, -, -, +)$.

TABLE 2. Orbit of signs for γ' .

n	$\epsilon_\gamma(\phi^n(l_\Delta^+))$	$\epsilon_\gamma(\phi^n(l_\Delta^-))$
1	(+, +, +, -, +, +)	(-, -, -, -, -, -)
2	(+, -, +, -, +, +)	(+, -, +, -, -, +)
3	(+, -, +, -, -, +)	(+, -, +, -, -, +)
4	(+, -, +, -, -, +)	(+, -, +, -, -, +)
5	(+, -, +, -, -, +)	(+, -, +, -, -, +)
\vdots	\vdots	\vdots

⁶In this case, the attracting/repelling points L_ϕ^\pm are antipodal to each other although it is a very special situation.

8. TOPOLOGICAL AND ALGEBRAIC ENTROPIES

In this section, we compare the topological entropy of a pA mapping class and the algebraic entropies of the cluster transformations it induces. First of all, let us recall the definitions and fundamental results regarding these notions of entropies.

8.1. Topological entropy. Given an open cover \mathcal{U} of a compact topological space X , let $N(\mathcal{U})$ denote the minimal cardinality of its finite subcover. For open covers $\mathcal{U}_1, \dots, \mathcal{U}_n$ of X , we define their common refinement by

$$\bigvee_{i=1}^n \mathcal{U}_i := \{U_1 \cap \dots \cap U_n \mid U_i \in \mathcal{U}_i, i = 1, \dots, n\}.$$

For a continuous map $f : X \rightarrow X$ and an open cover \mathcal{U} of X , we define another open cover by

$$f^{-1}(\mathcal{U}) := \{f^{-1}(U) \mid U \in \mathcal{U}\}.$$

Definition 8.1 ([AKM65]). The topological entropy of a continuous map $f : X \rightarrow X$ with respect to an open cover \mathcal{U} of X is defined to be

$$\mathcal{E}^{\text{top}}(f, \mathcal{U}) := \lim_{n \rightarrow \infty} \log N\left(\bigvee_{i=0}^{n-1} f^{-i}(\mathcal{U})\right).$$

Then the topological entropy of f is defined to be

$$\mathcal{E}_f^{\text{top}} := \sup_{\mathcal{U}} \mathcal{E}^{\text{top}}(f, \mathcal{U}),$$

where \mathcal{U} runs over the all open covers of X .

For a mapping class $\phi \in MC(\Sigma)$, its topological entropy is the infimum of the topological entropies of its representatives:

$$\mathcal{E}_\phi^{\text{top}} := \inf_{f \in \phi} \mathcal{E}_f^{\text{top}}.$$

Theorem 8.2. *Let Σ be a marked surface and let $C \subset \Sigma^\circ$ be an essential simple closed curve. Then we have*

$$\mathcal{E}_{T_C}^{\text{top}} = 0.$$

Proof. Although this theorem seems to be well-known, we give an easy proof here since we could not find a reference. First, recall the following theorem giving the estimate of the topological entropy from above:

Theorem 8.3 ([Ito70]). *Let M be a compact n -dimensional Riemannian manifold and $f : M \rightarrow M$ be a C^1 -diffeomorphism. Then,*

$$\mathcal{E}_f^{\text{top}} \leq n \log \sup_{x \in M} \|df_x\|.$$

Here $\|\cdot\|$ denotes the operator norm with respect to the norms on the tangent spaces $T_x M$ and $T_{f(x)} M$ given by the Riemannian metric.

In order to use this, we fix an auxiliary Riemannian metric on Σ . Take a tubular neighborhood $\mathcal{N}(C)$ of C and an isometry

$$\alpha : A \rightarrow \mathcal{N}(C) \subset \Sigma$$

with an annulus $A := \mathbb{R}/2\pi\mathbb{Z} \times [0, 1]$ equipped with the standard Euclidean metric. Let $T : A \rightarrow A$ be an orientation-preserving diffeomorphism defined by $(\theta, t) \mapsto (\theta + b(t), t)$, where $b : [0, 1] \rightarrow [0, 2\pi]$ is a smooth monotonically increasing function such that $b'(0) = b'(1) = 0$. Then its tangent map is of the form

$$dT_{(\theta, t)} = \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}$$

for any $(\theta, t) \in A$. Then the identity-extension of the diffeomorphism $\alpha \circ T \circ \alpha^{-1}$ on $\mathcal{N}(C)$ represents the Dehn twist T_C along C . Then from Theorem 8.3 we get

$$\begin{aligned} \mathcal{E}_{T_C}^{\text{top}} &\leq \mathcal{E}_{\alpha \circ T \circ \alpha^{-1}}^{\text{top}} \\ &\leq 2 \log \sup_{x \in \Sigma} \|d(\alpha \circ T \circ \alpha^{-1})_x\| \\ &= 2 \log \sup_{x \in \mathcal{N}(C)} \|d(\alpha \circ T \circ \alpha^{-1})_x\| \\ &= 2 \log \sup_{y \in A} \|dT_y\| = 0 \end{aligned}$$

as desired. ■

For the topological entropy of a pA mapping class, we have the following classical result:

Theorem 8.4 (Thurston, [FLP]). *Let Σ be a punctured surface and $\phi \in MC(\Sigma)$ a pA mapping class with stretch factor $\lambda_\phi > 1$. Then its topological entropy is given by*

$$\mathcal{E}_\phi^{\text{top}} = \log \lambda_\phi.$$

8.2. Algebraic entropy. For a rational function $f(u_1, \dots, u_N)$ over \mathbb{Q} on N variables, write it as $f(u) = g(u)/h(u)$ for two polynomials g and h without common factors. Then the *degree* of f , denoted by $\deg f$, is defined to be the maximum of the degrees of the constituent polynomials g and h . For a homomorphism $\varphi^* : \mathbb{Q}(u_1, \dots, u_N) \rightarrow \mathbb{Q}(u_1, \dots, u_N)$ between the field of rational functions on N variables, let $\varphi_i := \varphi^*(u_i)$ for $i = 1, \dots, N$. Since $\mathbb{Q}(u_1, \dots, u_N)$ is the field of rational functions on the algebraic torus \mathbb{G}_m^N equipped with coordinate functions u_1, \dots, u_N , the homomorphism φ^* can be regarded as the pull-back action induced by a rational map $\varphi : \mathbb{G}_m^N \rightarrow \mathbb{G}_m^N$ between algebraic tori. We define the degree of φ to be the maximum of the degrees $\deg \varphi_1, \dots, \deg \varphi_N$ of components and denote it by $\deg(\varphi)$.

Definition 8.5 ([BV99]). The *algebraic entropy* $\mathcal{E}_\varphi^{\text{alg}}$ of a rational map $\varphi : \mathbb{G}_m^N \rightarrow \mathbb{G}_m^N$ is defined to be

$$\mathcal{E}_\varphi^{\text{alg}} := \limsup_{n \rightarrow \infty} \frac{1}{n} \log(\deg(\varphi^n)).$$

Note that the algebraic entropy is invariant under a conjugation by a birational map. Hence for a mutation loop ϕ and $(z, \mathcal{Z}) = (a, \mathcal{A})$ or (x, \mathcal{X}) , the algebraic entropy of the cluster transformation $\phi_{(v_0)}^z$ on the torus $\mathcal{Z}_{(v_0)}$ is independent of the choice of a vertex $v_0 \in \mathbb{E}\text{ch}_{\mathbf{s}}$. Therefore we simply write $\mathcal{E}_{\phi}^z := \mathcal{E}_{\phi_{(v_0)}^z}^{\text{alg}}$ for $z = a, x$.

Theorem 8.6 ([IK21, Corollary 1.2]). *Let \mathbf{s} be a seed pattern without frozen indices. Let $\phi = [\gamma]_{\mathbf{s}}$ be a mutation loop represented by a path $\gamma : v_0 \rightarrow v$ which is sign-stable on the set $\Omega_{(v_0)}^{\text{can}}$. Assuming that Conjecture 5.14 holds true for the stable sign of γ on $\Omega_{(v_0)}^{\text{can}}$, we have*

$$\mathcal{E}_{\phi}^a = \mathcal{E}_{\phi}^x = \log \lambda_{\phi},$$

where $\lambda_{\phi} \geq 1$ is the cluster stretch factor.

Actually, it is not hard to drop the assumption on the frozen indices, which will be discussed in [IK20]. For an estimate without the dependence on Conjecture 5.14, see [IK21, Theorem 1.1]. We give a partial confirmation of Conjecture 5.14 in the punctured surface case.

Proposition 8.7. *Let \mathbf{s}_{Σ} be the seed pattern associated with a punctured surface Σ . Then Conjecture 5.14 holds true for any path γ representing a mutation loop and any differentiable point in $\mathcal{U}_{\Sigma}(\mathbb{R}^{\text{trop}})$ of μ_{γ} .*

Proof. Let $\gamma : (\Delta, \ell) \rightarrow (\Delta', \ell')$ be a path in $\mathbb{E}\text{ch}_{\Sigma}$ which represents a mutation loop ϕ . During the proof, we restore the notations ϕ^a and ϕ^x to distinguish the actions on $\mathcal{A}_{\Sigma}(\mathbb{R}^{\text{trop}})$ and $\mathcal{X}_{\Sigma}(\mathbb{R}^{\text{trop}})$, respectively.

Let $L \in \mathcal{U}_{\Sigma}(\mathbb{R}^{\text{trop}})$ be a differentiable point with a strict sign $\epsilon := \epsilon_{\gamma}(L)$. Then E_{γ}^{ϵ} is the presentation matrix of the tangent map $(d\phi^x)_L$, which is block-decomposed as

$$(d\phi^x)_L = \begin{pmatrix} (d\phi^u)_L & * \\ 0 & \sigma_{\Sigma}(\phi) \end{pmatrix}$$

similarly to (7.2) with respect to the direct sum decomposition

$$T_L \mathcal{X}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) = T_L \mathcal{U}_{(\Delta, \ell)}(\mathbb{R}^{\text{trop}}) \oplus s_L(\mathbb{R}^h)$$

with a choice of a section s . Here ϕ^u is the restriction of the action ϕ^x to $\mathcal{U}_{\Sigma}(\mathbb{R}^{\text{trop}})$.

On the other hand, recall from Proposition 4.6 the Γ_{Σ} -equivariant PL splitting

$$\mathcal{A}_{\Sigma}(\mathbb{R}^{\text{trop}}) \cong \mathcal{U}_{\Sigma}(\mathbb{R}^{\text{trop}}) \times H_{\mathcal{A}, \Sigma}(\mathbb{R}^{\text{trop}})$$

given by the Goncharov–Shen potentials. We can replace L without changing its sign with a point where the Goncharov–Shen potentials are differentiable. Since the sign of the cluster \mathcal{A} -transformation is constant along the fiber of the ensemble map, $\tilde{E}_{\gamma}^{\epsilon}$ is the

presentation matrix of the tangent map $(d\phi^a)_{\tilde{L}}$ for any lift $\tilde{L} \in \mathcal{A}_\Sigma(\mathbb{R}^{\text{trop}})$ with $p(\tilde{L}) = L$. The latter is decomposed as

$$(d\phi^a)_{\tilde{L}} = \begin{pmatrix} (d\phi^a)_L & 0 \\ 0 & \sigma_\Sigma(\phi) \end{pmatrix}$$

with respect to the splitting. Thus the characteristic polynomials of the tangent maps $(d\phi^x)_L$ and $(d\phi^a)_{\tilde{L}}$ both coincide with the product of those of the tangent map $(d\phi^u)_L$ and the linear map $\sigma_\Sigma(\phi)$, as desired. \blacksquare

8.3. Comparison of topological and algebraic entropies. We obtain the following comparisons:

Corollary 8.8. *For a simple closed curve C on a marked surface Σ satisfying the condition (6.1), we have*

$$\mathcal{E}_{T_C}^a = \mathcal{E}_{T_C}^x = \mathcal{E}_{T_C}^{\text{top}} = 0.$$

Proof. For simplicity, first consider the case of a punctured surface. Consider the representation path γ_C studied in Proposition 6.1. Since its stable sign $\epsilon_\gamma^{\text{stab}} = (+)$ is the sign at the point $L_{T_C} \in \mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$, by Theorem 8.6 and Proposition 8.7 we get $\mathcal{E}_{T_C}^a = \mathcal{E}_{T_C}^x = 0$. Combining with Theorem 8.2, we get the desired assertion. As another proof applicable to the general case, we can verify the vanishing of the upper bound $\log R_\phi$ appeared in [IK21, Theorem 1.1] by a direct computation using (6.2). \blacksquare

Corollary 8.9. *For a punctured surface Σ and a pA mutation loop $\phi \in \Gamma_\Sigma$, we have*

$$\mathcal{E}_\phi^a = \mathcal{E}_\phi^x = \mathcal{E}_\phi^{\text{top}} = \log \lambda_\phi.$$

Here $\lambda_\phi > 1$ is the stretch factor (=cluster stretch factor) of the underlying mapping class of ϕ .

Proof. This is a direct consequence of Theorem 7.1, Theorem 8.6 and Proposition 8.7. Indeed, just note that the attracting point L_ϕ^+ giving the stable sign for any representation path is contained in $\mathcal{U}_\Sigma(\mathbb{R}^{\text{trop}})$. \blacksquare

APPENDIX A. REVIEW ON THE CLUSTER ENSEMBLES

In this section, we recall the theory of cluster ensembles following [FG09]. In this paper, we consider the \mathcal{X} -variety $\mathcal{X}_s^{\text{uf}}$ which has only the unfrozen coordinates.

We use the conventions in [IK21]. Fix a finite index set I . We also choose a (possibly empty) subset $I_f \subset I$, which we call the subset of *frozen indices*. The subset $I_{\text{uf}} := I \setminus I_f$ is called the subset of *unfrozen indices*. We always regard the index sets I , I_{uf} and I_f as finite ordered sets:

$$I = \underbrace{\{1, \dots, |I_{\text{uf}}|\}}_{I_{\text{uf}}}, \underbrace{\{|I_{\text{uf}}| + 1, \dots, |I_{\text{uf}}| + |I_f| = |I|\}}_{I_f}.$$

Then “mutations” will be allowed only for indices in I_{uf} .

A.1. Seed patterns. A *(Fock–Goncharov) seed* is a data (N, B) , where

- $N = \bigoplus_{i \in I} \mathbb{Z}e_i$ is a lattice with a basis $(e_i)_{i \in I}$,
- $B = (b_{ij})_{i, j \in I}$ is a skew-symmetric matrix with entries in \mathbb{Q} . We assume that the entry b_{ij} is integral unless $(i, j) \in I_f \times I_f$.

We call the lattice N the *seed lattice*, the matrix B the *exchange matrix*. We define a skew-symmetric bilinear form $\{-, -\} : N \times N \rightarrow \mathbb{Q}$ by $\{e_i, e_j\} := b_{ij}$.

Let $M = \text{Hom}(N, \mathbb{Z})$ be the dual lattice, and consider the sub-lattice $N_{\text{uf}} \subset N$ spanned by the basis vectors $(e_i)_{i \in I_{\text{uf}}}$. Then the bilinear form induces a linear map $p^* : N_{\text{uf}} \rightarrow M$, $n \mapsto \{n, -\}$ called the *ensemble map*⁷. The tuple $(N, \{-, -\}, (e_i)_i)$ is called a seed in [FG09]. We have the exact sequence

$$0 \rightarrow \ker p^* \rightarrow N_{\text{uf}} \xrightarrow{p^*} M \rightarrow \text{coker } p^* \rightarrow 0, \quad (\text{A.1})$$

which is called the *cluster exact sequence* [FG08].

Let $\mathbb{T}_{I_{\text{uf}}}$ be a regular $|I_{\text{uf}}|$ -valent tree whose edges are labeled by elements of I_{uf} , such that the edges incident to a common vertex have distinct labels. An assignment

$$\mathbf{s} : \mathbb{T}_{I_{\text{uf}}} \ni t \mapsto (N^{(t)} = \bigoplus_{i \in I} \mathbb{Z}e_i^{(t)}, B^{(t)})$$

of a seed to each vertex $t \in \mathbb{T}_{I_{\text{uf}}}$ is called a *(Fock–Goncharov) seed pattern* if for each edge $t \xrightarrow{k} t'$ of $\mathbb{T}_{I_{\text{uf}}}$ labeled by $k \in I_{\text{uf}}$, the exchange matrices $B^{(t)} = (b_{ij})$ and $B^{(t')} = (b'_{ij})$ are related by the *matrix mutation* at $k \in I_{\text{uf}}$ (write $B^{(t')} = \mu_k B^{(t)}$):

$$b'_{ij} = \begin{cases} -b_{ij} & \text{if } i = k \text{ or } j = k, \\ b_{ij} + [b_{ik}]_+ + [b_{kj}]_+ - [-b_{ik}]_+ - [-b_{kj}]_+ & \text{otherwise.} \end{cases} \quad (\text{A.2})$$

Here $[a]_+ := \max\{a, 0\}$ for $a \in \mathbb{R}$.

Two seed patterns \mathbf{s}, \mathbf{s}' are *isomorphic* if there exists an automorphism ϕ of the graph $\mathbb{T}_{I_{\text{uf}}}$ such that $\mathbf{s}(\phi(t)) = \mathbf{s}'(t)$ for all $t \in \mathbb{T}_{I_{\text{uf}}}$. We are going to study objects associated with an isomorphism class of seed patterns. A seed pattern is specified by choosing an “initial” exchange matrix:

Lemma A.1. *Let us fix a vertex $t_0 \in \mathbb{T}_{I_{\text{uf}}}$, and consider a skew-symmetric matrix B . Then there exists a seed pattern $\mathbf{s} = \mathbf{s}_{t_0; B} : \mathbb{T}_{I_{\text{uf}}} \ni t \mapsto (N^{(t)}, B^{(t)})$ such that $B^{(t_0)} = B$.*

We call $\mathbf{s}_{t_0; B}$ the *seed pattern with the initial exchange matrix B at the vertex t_0* . Note that its isomorphism class only depends on the mutation class (=the class for the equivalence relation generated by matrix mutations) of B .

It is useful to represent a skew-symmetric matrix $B = (b_{ij})_{i, j \in I}$ by a quiver Q . It has vertices parametrized by the set I and $|b_{ij}|$ arrows from i to j (resp. j to i) if $b_{ij} > 0$ (resp. $b_{ji} > 0$). Note that that the quiver Q has no loops and 2-cycles, and the matrix B can be reconstructed from such a quiver.

⁷Note that we do not have a canonical map $N \rightarrow M$: this point is carefully treated in [GHKK18, Appendix A].

Signed seed mutations. As a relation between the lattices assigned to t and t' , we consider two linear isomorphisms $(\tilde{\mu}_k^\epsilon)^* : N^{(t')} \xrightarrow{\sim} N^{(t)}$ which depend on a sign $\epsilon \in \{+, -\}$ and are given by

$$e_i^{(t')} \mapsto \begin{cases} -e_k^{(t)} & \text{if } i = k, \\ e_i^{(t)} + [\epsilon b_{ik}^{(t)}]_+ e_k^{(t)} & \text{if } i \neq k. \end{cases}$$

It induces a linear isomorphism $(\tilde{\mu}_k^\epsilon)^* : M^{(t')} \xrightarrow{\sim} M^{(t)}$ (denoted by the same symbol) which sends $f_i^{(t')}$ to the dual basis of $(\tilde{\mu}_k^\epsilon)^*(e_i^{(t')})$. Explicitly, it is given by

$$f_i^{(t')} \mapsto \begin{cases} -f_k^{(t)} + \sum_{j \in I} [-\epsilon b_{kj}^{(t)}]_+ f_j^{(t)} & \text{if } i = k, \\ f_i^{(t)} & \text{if } i \neq k. \end{cases}$$

We call each map $(\tilde{\mu}_k^\epsilon)^*$ the *signed seed mutation* at $k \in I_{\text{uf}}$.

The seed mutations are compatible with matrix mutations. Namely, for any edge $t \xrightarrow{k} t'$ in $\mathbb{T}_{I_{\text{uf}}}$, we have

$$\{\tilde{\mu}_k^*(e_i^{(t')}), \tilde{\mu}_k^*(e_j^{(t')})\} = b_{ij}^{(t')}.$$

For later discussions, we collect here some properties of the presentation matrices of the signed seed mutation and its dual, with respect to the seed bases. For an edge $t \xrightarrow{k} t'$ of $\mathbb{T}_{I_{\text{uf}}}$ and a sign $\epsilon \in \{+, -\}$, let us consider the matrices $\check{E}_{k,\epsilon}^{(t)} = (\check{E}_{ij})_{i,j \in I}$ and $E_{k,\epsilon}^{(t)} = (E_{ij})_{i,j \in I}$, given as follows:

$$\check{E}_{ij} := \begin{cases} 1 & \text{if } i = j \neq k, \\ -1 & \text{if } i = j = k, \\ [\epsilon b_{jk}^{(t)}]_+ & \text{if } i = k \text{ and } j \neq k, \\ 0 & \text{otherwise,} \end{cases} \quad (\text{A.3})$$

$$E_{ij} := \begin{cases} 1 & \text{if } i = j \neq k, \\ -1 & \text{if } i = j = k, \\ [-\epsilon b_{ki}^{(t)}]_+ & \text{if } j = k \text{ and } i \neq k, \\ 0 & \text{otherwise.} \end{cases} \quad (\text{A.4})$$

Then the matrix $\check{E}_{k,\epsilon}^{(t)}$ gives the presentation matrix of $(\tilde{\mu}_k^\epsilon)^* : N^{(t')} \xrightarrow{\sim} N^{(t)}$ with respect to the seed bases $(e_i^{(t')})$ and $(e_i^{(t)})$: $(\tilde{\mu}_k^\epsilon)^* e_i^{(t')} = \sum_{j \in I} e_j^{(t)} (\check{E}_{k,\epsilon}^{(t)})_{ji}$. Similarly the matrix $E_{k,\epsilon}^{(t)}$ gives the presentation matrix of $(\tilde{\mu}_k^\epsilon)^* : M^{(t')} \xrightarrow{\sim} M^{(t)}$ with respect to the bases $(f_i^{(t')})$ and $(f_i^{(t)})$. These matrices satisfy the relation $(E_{k,\epsilon}^{(t)})^\top = (\check{E}_{k,\epsilon}^{(t)})^{-1}$. Having this in mind, we use the notation $\check{M} := (M^\top)^{-1}$ for any invertible matrix M .

A.2. Cluster varieties.

Seed tori. Let $\mathbb{G}_m := \text{Spec } \mathbb{Z}[z, z^{-1}]$ be the multiplicative group. A reader unfamiliar with this notation can recognize it as $\mathbb{G}_m(\mathbb{k}) = \mathbb{k}^*$ by substituting a field \mathbb{k} . We have an equivalence

$$\text{Lattices} \xrightarrow{\sim} \text{Tori}, \quad L \mapsto T_L := \text{Hom}(L^*, \mathbb{G}_m)$$

of the category of finite rank lattices and that of split algebraic tori. Here $L^* := \text{Hom}(L, \mathbb{Z})$ denotes the dual lattice of L . Note that an element $\ell^* \in L^*$ defines a character (*i.e.*, a morphism of algebraic tori) $\text{ch}_{\ell^*} : T_L \rightarrow \mathbb{G}_m$ by $\text{ch}_{\ell^*}(\phi) := \phi(\ell^*)$. Therefore the lattice L^* can be regarded as the lattice of characters on the torus T_L . Dually, the lattice L is identified with the lattice of cocharacters $\mathbb{G}_m \rightarrow T_L$.

We are going to see that a seed pattern $\mathbf{s} : \mathbb{T}_{I_{\text{uf}}} \ni t \mapsto (N^{(t)}, B^{(t)})$ gives rise to two families of algebraic tori related by birational maps. For each $t \in \mathbb{T}_{I_{\text{uf}}}$, we define a pair of tori $\mathcal{X}_{(t)}^{\text{uf}} := T_{M_{\text{uf}}^{(t)}}$ and $\mathcal{A}_{(t)} := T_{N^{(t)}}$ called the *seed \mathcal{X} - and \mathcal{A} -torus*, respectively. Here M_{uf} is the dual lattice of N_{uf} . The characters $X_i^{(t)} := \text{ch}_{e_i^{(t)}}$ ($i \in I_{\text{uf}}$) and $A_i^{(t)} := \text{ch}_{f_i^{(t)}}$ ($i \in I$) are called the *cluster \mathcal{X} - and \mathcal{A} -coordinates*, respectively. The ensemble map induces a monomial map $p_{(t)} : \mathcal{A}_{(t)} \rightarrow \mathcal{X}_{(t)}^{\text{uf}}$ given by

$$p_{(t)}^* X_i^{(t)} = \prod_{j \in I} (A_j^{(t)})^{b_{ij}^{(t)}}$$

for $i \in I_{\text{uf}}$. Setting $H_{\mathcal{A},(t)} := T_{(\ker p^*)^*}$ and $H_{\mathcal{X},(t)} := T_{(\text{coker } p^*)^*}$, we get the torus version of the cluster exact sequence (A.1)

$$1 \rightarrow H_{\mathcal{A},(t)} \rightarrow \mathcal{A}_{(t)} \xrightarrow{p_{(t)}} \mathcal{X}_{(t)}^{\text{uf}} \xrightarrow{\theta_{(t)}} H_{\mathcal{X},(t)} \rightarrow 1, \quad (\text{A.5})$$

which is obtained by applying the functor $L \mapsto T_L$. Note that the inclusion $H_{\mathcal{A},(t)} \rightarrow \mathcal{A}_{(t)}$ induces an action

$$\alpha_{(t)} : H_{\mathcal{A},(t)} \times \mathcal{A}_{(t)} \rightarrow \mathcal{A}_{(t)} \quad (\text{A.6})$$

using the multiplicative structure of $\mathcal{A}_{(t)}$, which makes the ensemble map $p_{(t)}$ a principal $H_{\mathcal{A},(t)}$ -bundle over its image $\mathcal{U}_{(t)}^{\text{uf}} := p_{(t)}(\mathcal{A}_{(t)})$.

Cluster transformations. For an edge $t \xrightarrow{k} t'$ of $\mathbb{T}_{I_{\text{uf}}}$, the signed mutation induces monomial isomorphisms $\tilde{\mu}_k^\epsilon : \mathcal{X}_{(t)}^{\text{uf}} \rightarrow \mathcal{X}_{(t')}^{\text{uf}}$ and $\tilde{\mu}_k^\epsilon : \mathcal{A}_{(t)} \rightarrow \mathcal{A}_{(t')}$, which are given by

$$(\tilde{\mu}_k^\epsilon)^* X_i' := \begin{cases} X_k^{-1} & \text{if } i = k, \\ X_i X_k^{[eb_{ik}^{(t)}]_+} & \text{if } i \neq k, \end{cases} \quad (\tilde{\mu}_k^\epsilon)^* A_i' := \begin{cases} A_k^{-1} \prod_{j \in I} A_j^{[-eb_{kj}^{(t)}]_+} & \text{if } i = k, \\ A_i & \text{if } i \neq k. \end{cases}$$

Here $X_i := X_i^{(t)}$, $A_i := A_i^{(t)}$, $X_i' := X_i^{(t')}$ and $A_i' := A_i^{(t')}$. Let $\mu_k^{\#, \epsilon}$ be the birational isomorphism on $\mathcal{X}_{(t)}^{\text{uf}}$ and $\mathcal{A}_{(t)}$ (denoted by a common symbol) given by

$$(\mu_k^{\#, \epsilon})^* X_i := X_i (1 + X_k^\epsilon)^{-b_{ik}^{(t)}} \quad \text{and} \quad (\mu_k^{\#, \epsilon})^* A_i := A_i (1 + (p_{(t)}^* X_k)^\epsilon)^{-\delta_{ik}}.$$

Composing these maps, we get the *cluster transformation* $\mu_k := \tilde{\mu}_k^\epsilon \circ \mu_k^{\#, \epsilon}$, which are explicitly given by

$$\mu_k^* X'_i = \begin{cases} X_k^{-1} & \text{if } i = k, \\ X_i (1 + X_k^{-\text{sgn}(b_{ik}^{(t)})})^{-b_{ik}^{(t)}} & \text{if } i \neq k \end{cases}$$

and

$$\mu_k^* A'_i = \begin{cases} A_k^{-1} (\prod_{j \in I} A_j^{[b_{kj}^{(t)}]_+} + \prod_{j \in I} A_j^{[-b_{kj}^{(t)}]_+}) & \text{if } i = k, \\ A_i & \text{if } i \neq k, \end{cases}$$

Here note that they do not depend on the sign ϵ , and it is easy to verify that they are involutive: $\mu_k \mu_k = \text{id}$. When we stress the distinction between the \mathcal{X} - and \mathcal{A} -transformations, we write μ_k^x and μ_k^a instead of μ_k . What we obtain are two families of tori related by the cluster transformations, see Figure 30.




FIGURE 30. The seed tori related by cluster transformations, where $(z, \mathcal{Z}) = (a, \mathcal{A}), (x, \mathcal{X}^{\text{uf}})$.

Definition A.2. The *cluster varieties* $\mathcal{A}_{\mathbf{s}}$ and $\mathcal{X}_{\mathbf{s}}^{\text{uf}}$ associated with a seed pattern $\mathbf{s} : t \mapsto (N^{(t)}, B^{(t)})$ are defined by patching the corresponding tori by cluster transformations:

$$\mathcal{A}_{\mathbf{s}} := \bigcup_{t \in \mathbb{T}_{I_{\text{uf}}}} \mathcal{A}_{(t)}, \quad \mathcal{X}_{\mathbf{s}}^{\text{uf}} := \bigcup_{t \in \mathbb{T}_{I_{\text{uf}}}} \mathcal{X}_{(t)}^{\text{uf}}.$$

From the definition, each $\mathcal{X}_{(t)}^{\text{uf}}$ is an open subvariety of $\mathcal{X}_{\mathbf{s}}^{\text{uf}}$. The pair $(\mathcal{X}_{(t)}^{\text{uf}}, (X_i^{(t)})_{i \in I_{\text{uf}}})$ is called the *cluster \mathcal{X} -chart* associated with $t \in \mathbb{T}_{I_{\text{uf}}}$. Similarly we call the pair $(\mathcal{A}_{(t)}, (A_i^{(t)})_{i \in I})$ *cluster \mathcal{A} -chart*.

We call the triple $(\mathcal{A}_{\mathbf{s}}, \mathcal{X}_{\mathbf{s}}^{\text{uf}}, p)$ the *cluster ensemble* associated to the seed pattern \mathbf{s} . For a cluster ensemble including frozen \mathcal{X} -coordinates, see [GHKK18, Appendix A]. We define the unfrozen part of the cluster \mathcal{A} -variety to be $\mathcal{A}_{\mathbf{s}}^{\text{uf}} := \bigcup_{t \in \mathbb{T}_{I_{\text{uf}}}} \mathcal{A}_{(t)}^{\text{uf}}$, where $\mathcal{A}_{(t)}^{\text{uf}} := T_{N_{\text{uf}}^{(t)}} \subset \mathcal{A}_{(t)}$ is a subtorus given by $A_i^{(t)} = 1$ for all $i \in I_{\text{f}}$.

Definition A.3. The tori $H_{\mathcal{A}, \mathbf{s}}$ and $H_{\mathcal{X}, \mathbf{s}}$ associated with a seed pattern $\mathbf{s} : t \mapsto (N^{(t)}, B^{(t)})$ are defined by patching the corresponding tori by the monomial parts of the cluster transformations:

$$H_{\mathcal{A}, \mathbf{s}} := \bigcup_{t \in \mathbb{T}_{I_{\text{uf}}}} H_{\mathcal{A}, (t)}, \quad H_{\mathcal{X}, \mathbf{s}} := \bigcup_{t \in \mathbb{T}_{I_{\text{uf}}}} H_{\mathcal{X}, (t)}.$$

Here notice that the identification is independent of the choice of a sign $\epsilon \in \{+, -\}$ for the monomial part $\tilde{\mu}_k^\epsilon$, since the difference between two signed mutations applied on $n = \sum_i c_i e_i^{(t)} \in N_{\text{uf}}^{(t)}$ is given by

$$\sum_i c_i (e_i^{(t)} + [b_{ik}^{(t)}]_+ e_k^{(t)}) - \sum_i c_i (e_i^{(t)} + [-b_{ik}^{(t)}]_+ e_k^{(t)}) = \sum_i c_i b_{ik}^{(t)} e_k^{(t)},$$

which vanishes when $n \in \ker p^*$. The cluster exact sequence (A.5) is preserved under cluster transformations:

Proposition A.4 ([FG09, Proposition 2.2, Lemma 2.7 and Lemma 2.10(a)]). *For each $t \in \mathbb{T}_{I_{\text{uf}}}$, the following diagram commutes:*

$$\begin{array}{ccccccc} H_{\mathcal{A},(t)} & \longrightarrow & \mathcal{A}_{(t)} & \xrightarrow{p(t)} & \mathcal{X}_{(t)}^{\text{uf}} & \xrightarrow{\theta(t)} & H_{\mathcal{X},(t)} \\ & & \parallel & & \downarrow \mu_k^x & & \parallel \\ & & \mu_k^a \downarrow & & \downarrow \mu_k^x & & \parallel \\ H_{\mathcal{A},(t)} & \longrightarrow & \mathcal{A}_{(t)} & \xrightarrow{p(t)} & \mathcal{X}_{(t)}^{\text{uf}} & \xrightarrow{\theta(t)} & H_{\mathcal{X},(t)} \end{array}$$

Thus we get a well-defined action $\alpha : H_{\mathcal{A},s} \times \mathcal{A}_s \rightarrow \mathcal{A}_s$ and a morphism $\theta : \mathcal{X}_s^{\text{uf}} \rightarrow H_{\mathcal{X},s}$. The exact sequence (A.5) implies that the ensemble map $p : \mathcal{A}_s \rightarrow \mathcal{X}_s^{\text{uf}}$ is a principal $H_{\mathcal{A},s}$ -bundle over its image $\mathcal{U}_s^{\text{uf}} := p(\mathcal{A}_s)$, and the variety $\mathcal{U}_s^{\text{uf}}$ coincides with $\theta^{-1}(e)$. The situation is summarized in the ‘‘exact sequence’’

$$1 \rightarrow H_{\mathcal{A},s} \rightarrow \mathcal{A}_s \xrightarrow{p} \mathcal{X}_s^{\text{uf}} \xrightarrow{\theta} H_{\mathcal{X},s} \rightarrow 1, \quad (\text{A.7})$$

which is also called the *cluster exact sequence*. When $I_{\text{uf}} = I$, it is known that the variety $\mathcal{U}_s = \mathcal{U}_s^{\text{uf}}$ is a symplectic leaf of the cluster Poisson structure on $\mathcal{X}_s = \mathcal{X}_s^{\text{uf}}$ [FG09, Proposition 2.9].

C- and G-matrices. Fixing a vertex $t_0 \in \mathbb{T}_{I_{\text{uf}}}$, we assign the *C-matrix* $C_t^{\mathbf{s};t_0} = (c_{ij}^{(t)})_{i,j \in I}$ to each vertex $t \in \mathbb{T}_{I_{\text{uf}}}$ by the following rule:

- (1) $C_{t_0}^{\mathbf{s};t_0} = \text{Id}$,
- (2) For each $t \xrightarrow{k} t'$ of $\mathbb{T}_{I_{\text{uf}}}$, the matrices $C_t^{\mathbf{s};t_0}$ and $C_{t'}^{\mathbf{s};t_0}$ are related by

$$c'_{ij} = \begin{cases} -c_{ij} & i = k, \\ c_{ij} + [c_{kj}]_+ [b_{ik}^{(t)}]_+ - [-c_{kj}]_+ [-b_{ik}^{(t)}]_+ & i \neq k. \end{cases} \quad (\text{A.8})$$

Here we write $c_{ij} := c_{ij}^{(t)}$ and $c'_{ij} := c_{ij}^{(t')}$. Its row vectors $\mathbf{c}_i^{(t)} = (c_{ij}^{(t)})_{j \in I}$ are called *c-vectors*. The following theorem was firstly conjectured in [FZ07], and proved in [DWZ10] for skew-symmetric case, and in [GHKK18] for skew-symmetrizable case:

Theorem A.5 (Sign-coherence theorem for *c*-vectors). *For any $t \in \mathbb{T}_{I_{\text{uf}}}$ and $i \in I$, $\mathbf{c}_i^{(t)} \in \mathbb{Z}_{\geq 0}^I$ or $\mathbf{c}_i^{(t)} \in \mathbb{Z}_{\leq 0}^I$.*

Following [NZ12], we define the *tropical sign* $\epsilon_i^{(t)}$ to be $+$ in the former case, and $-$ in the latter case. Using the tropical sign, the relation (A.8) is equivalent to

$$C_{t'}^{s;t_0} = E_{k, \epsilon_k^{(t)}}^{(t)} C_t^{s;t_0}, \quad (\text{A.9})$$

where $E_{k, \epsilon_k^{(t)}}^{(t)}$ is the matrix (A.4). Similarly, we assign the *G-matrix* $G_t^{s;t_0} = (g_{ij}^{(t)})_{i,j \in I}$ to each vertex $t \in \mathbb{T}_{I_{\text{uf}}}$ by the following rule:

- (1) $G_{t_0}^{s;t_0} = \text{Id}$,
- (2) For each $t \xrightarrow{k} t'$ of $\mathbb{T}_{I_{\text{uf}}}$, the matrices $G_t^{s;t_0}$ and $G_{t'}^{s;t_0}$ are related by

$$G_{t'}^{s;t_0} = \check{E}_{k, \epsilon_k^{(t)}}^{(t)} G_t^{s;t_0}. \quad (\text{A.10})$$

We refer to the row vectors $\mathbf{g}_i^{(t)} = (g_{ij}^{(t)})_{j \in I}$ of $G_t^{s;t_0}$ as *g-vectors*. The *C*- and *G*-matrices are related by the following one of the *tropical dualities* [NZ12]

$$G_t^{s;t_0} = \check{C}_t^{s;t_0}. \quad (\text{A.11})$$

A.3. Tropicalizations of the cluster varieties. Let $\mathbb{P} = (\mathbb{P}, \oplus, \cdot)$ be a semifield. For a torus T_L with a lattice L of finite rank, we define the *set of \mathbb{P} -points* to be $T_L(\mathbb{P}) := L \otimes_{\mathbb{Z}} \mathbb{P}$. A positive rational map $f : T_L \rightarrow T_{L'}$ naturally induces a map $f(\mathbb{P}) : T_L(\mathbb{P}) \rightarrow T_{L'}(\mathbb{P})$. In particular, the character $\text{ch}_{\ell^*} : T_L \rightarrow \mathbb{G}_m$ associated with a point $\ell^* \in L^*$ induces a group homomorphism $\text{ch}_{\ell^*}(\mathbb{P}) : T_L(\mathbb{P}) \rightarrow \mathbb{G}_m(\mathbb{P}) = \mathbb{P}$, which coincides with the evaluation map $L \otimes_{\mathbb{Z}} \mathbb{P} \rightarrow \mathbb{P}$, $\lambda \otimes p \mapsto \langle \ell^*, \lambda \rangle p$.

Applying this construction to seed tori $\mathcal{X}_{(t)}^{\text{uf}}$ and $\mathcal{A}_{(t)}$, we get $\mathcal{X}_{(t)}^{\text{uf}}(\mathbb{P}) = M_{\text{uf}}^{(t)} \otimes \mathbb{P}$ and $\mathcal{A}_{(t)} = N^{(t)} \otimes \mathbb{P}$ equipped with functions

$$x_i^{(t)} := \text{ch}_{e_i^{(t)}}(\mathbb{P}) : \mathcal{X}_{(t)}^{\text{uf}}(\mathbb{P}) \rightarrow \mathbb{P}, \quad a_i^{(t)} := \text{ch}_{f_i^{(t)}}(\mathbb{P}) : \mathcal{A}_{(t)}(\mathbb{P}) \rightarrow \mathbb{P}$$

which we call the *tropical cluster \mathcal{X} - and \mathcal{A} -coordinates*. Since cluster transformations are positive rational maps, they induce maps between these sets.

Definition A.6. We define the set of \mathbb{P} -valued points of the cluster \mathcal{X} -variety as $\mathcal{X}_s^{\text{uf}}(\mathbb{P}) := \bigsqcup_{t \in \mathbb{T}_{I_{\text{uf}}}} \mathcal{X}_{(t)}^{\text{uf}}(\mathbb{P}) / \sim$, where for each edge $t \xrightarrow{k} t'$, two points $x \in \mathcal{X}_{(t)}^{\text{uf}}(\mathbb{P})$ and $x' \in \mathcal{X}_{(t')}^{\text{uf}}(\mathbb{P})$ are identified if $x' = \mu_k(\mathbb{P})(x)$. Similarly we define $\mathcal{A}_s(\mathbb{P})$.

For $\mathbb{A} = \mathbb{Z}$ or \mathbb{R} , let $\mathbb{A}^{\text{trop}} := (\mathbb{A}, \min, +)$ be the *tropical semifield* over \mathbb{A} . When dealing with the \mathbb{A}^{trop} -valued points of cluster varieties, we simply denote the map $f(\mathbb{A}^{\text{trop}})$ induced by a positive map f by f^{trop} . The cluster transformations $\mu_k^z(\mathbb{A}^{\text{trop}})$ for $z = a, x$ are piecewise-linear (PL for short) maps, and the sets $\mathcal{A}_s(\mathbb{R}^{\text{trop}})$ and $\mathcal{X}_s^{\text{uf}}(\mathbb{R}^{\text{trop}})$ have canonical PL structures. We call these PL manifolds the *tropical cluster varieties*. Moreover we omit the symbol “ a ” and “ x ” from the superscript of cluster transformations when no confusion can occur.

Remark A.7. Similarly, we have the PL manifolds $\mathcal{A}_s(\mathbb{R}^T)$ and $\mathcal{X}_s^{\text{uf}}(\mathbb{R}^T)$ corresponding to the semifield $\mathbb{R}^T := (\mathbb{R}, \max, +)$. The semifield isomorphism $\mathbb{R}^T \rightarrow \mathbb{R}^{\text{trop}}$, $x \mapsto -x$ induces an isomorphism $\iota : \mathcal{Z}_s(\mathbb{R}^T) \rightarrow \mathcal{Z}_s(\mathbb{R}^{\text{trop}})$ of PL manifolds for $\mathcal{Z} = \mathcal{A}, \mathcal{X}^{\text{uf}}$.

REFERENCES

- [AKM65] R. L. Adler, A. G. Konheim, M. H. McAndrew, *Topological entropy*, Trans. Amer. Math. Soc. **114** (1965), 309–319.
- [ASS12] I. Assem, R. Schiffler and V. Shramchenko, *Cluster automorphisms*, Proc. London Math. Soc. **104** (2012) 1271–1302.
- [BB09] R. Benedetti, F. Bonsante, *(2+1) Einstein spacetimes of finite type*, Handbook of Teichmüller theory. Vol. II, 533–609, IRMA Lect. Math. Theor. Phys., 13, Eur. Math. Soc., Zürich, 2009.
- [BH95] M. Bestvina and M. Handel, *Train-tracks for surface homeomorphisms*, Topology **34** (1995), 109–140.
- [BBK12] J. Birman, P. Brinkmann and K. Kawamuro, *A polynomial invariant of pseudo-Anosov maps*, Journal of Topology and Analysis, **4** (2012), 13–47.
- [BKS11] F. Bonsante, K. Krasnov and J.-M. Schlenker, *Multi-black holes and earthquakes on Riemann surfaces with boundaries*, Int. Math. Res. Not. (2011), 487–552.
- [Bon98] F. Bonahon, *Variations of the boundary geometry of 3-dimensional hyperbolic convex cores*. J. Differential Geom. **50** (1998), 1–24.
- [BS15] T. Bridgeland and I. Smith, *Quadratic differentials as stability conditions*, Publ. Math. Inst. HautesEtudes Sci. **121** (2015), 155–278.
- [BV99] M. P. Bellon and C.-M. Viallet, *Algebraic entropy*, Comm. Math. Phys. **204** (1999), 425–437.
- [CB] A. J. Casson and S. A. Bleiler, *Automorphisms of surfaces after Nielsen and Thurston*, London Mathematical Society Student Texts, **9**. Cambridge University Press, Cambridge (1988), iv+105 pp.
- [CP07] L. Chekhov and R. C. Penner, *On quantizing Teichmüller and Thurston theories*, Handbook of Teichmüller theory. Vol. I, 579–645, IRMA Lect. Math. Theor. Phys., 11, Eur. Math. Soc., Zürich, 2007.
- [DWZ10] H. Derksen, J. Weyman and A. Zelevinsky, *Quivers with potentials and their representations II: applications to cluster algebras*, J. Amer. Math. Soc. **23** (2010), 749–790.
- [FG06a] V. V. Fock and A. B. Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, Publ. Math. Inst. Hautes Études Sci., **103** (2006), 1–211.
- [FG07] V. V. Fock and A. B. Goncharov, *Dual Teichmüller and lamination spaces*, Handbook of Teichmüller theory, Vol. I, 647–684; IRMA Lect. Math. Theor. Phys., 11, Eur. Math. Soc., Zürich, 2007.
- [FG08] V. V. Fock and A. B. Goncharov, *The quantum dilogarithm and representations of quantum cluster varieties*, Invent. Math. **175** (2009), 223–286.
- [FG09] V. V. Fock and A. B. Goncharov, *Cluster ensembles, quantization and the dilogarithm*, Ann. Sci. Éc. Norm. Supér., **42** (2009), 865–930.
- [FG16] V. V. Fock and A. B. Goncharov, *Cluster Poisson varieties at infinity*, Selecta Math. (N.S.) **22** (2016), 2569–2589.
- [FLP] A. Fathi, F. Laudenbach and V. Poénaru, *Thurston’s work on surfaces*, Mathematical Notes, **48**, Princeton University Press, Princeton, NJ, 2012, Translated from the 1979 French original by D. M. Kim and D. Margalit.
- [FM] B. Farb and D. Margalit, *A primer on mapping class groups*, Princeton Mathematical Series, **49**, Princeton University Press, Princeton, NJ, 2012. xiv+472 pp.
- [FST08] S. Fomin, M. Shapiro and D. Thurston, *Cluster algebras and triangulated surfaces. I. Cluster complexes*, Acta Math. **201** (2008), 83–146.
- [FT18] S. Fomin and D. Thurston *Cluster algebras and triangulated surfaces Part II: Lambda lengths*, Mem. Amer. Math. Soc. **255** (2018).

- [FZ02] S. Fomin and A. Zelevinsky, *Cluster algebras. I. Foundations*, J. Amer. Math. Soc. **15** (2002), 497–529.
- [FZ07] S. Fomin and A. Zelevinsky, *Cluster algebras. IV. Coefficients*, Compos. Math. **143** (2007), 112–164.
- [GHKK18] M. Gross, P. Hacking and S. Keel and M. Kontsevich, *Canonical bases for cluster algebras*, J. Amer. Math. Soc. **31** (2018), 497–608.
- [Gu11] W. Gu, *Graphs with non-unique decomposition and their associated surfaces*, arXiv:1112.1008.
- [Had11] A. Hadari, *Algebraic entropy and the action of mapping class groups on character varieties*, Adv. Math. **226** (2011), 3282–3296.
- [IK21] T. Ishibashi and S. Kano, *Algebraic entropy of sign-stable mutation loops*, to appear in Geometriae Dedicata, arXiv:1911.07587.
- [IK20] T. Ishibashi and S. Kano, *Sign stability of mapping classes on marked surfaces II: general case*, arXiv:2011.14320.
- [Ish19] T. Ishibashi, *On a Nielsen–Thurston classification theory for cluster modular groups*, Annales de l’Institut Fourier, **69** (2019), 515–560.
- [Ito70] S. Ito, *An estimate from above for the entropy and the topological entropy of a C^1 -diffeomorphism*, Proc. Japan Acad. **46** (1970), 226–230.
- [Iva] N. V. Ivanov, *Subgroups of Teichmüller modular groups*, Translations of Mathematical Monographs, **115**, Amer. Math. Soc., Providence, RI, 1992.
- [Le16] I. Le, *Higher laminations and affine buildings*, Geom. Topol. **20** (2016), 1673–1735.
- [NZ12] T. Nakanishi and A. Zelevinsky, *On tropical dualities in cluster algebras*, Algebraic groups and quantum groups, Contemp. Math., Amer. Math. Soc., Providence, RI, **565** (2012), 217–226.
- [Pen] R. C. Penner, *Decorated Teichmüller theory*, QGM Master Class Series, European Mathematical Society (EMS), Zürich, 2012.
- [Pen87] R. C. Penner, *The decorated Teichmüller space of punctured surfaces*, Comm. Math. Phys. **113** (1987), 299–339.
- [Pen04] R. C. Penner, *Decorated Teichmüller theory of bordered surfaces*, Comm. Anal. Geom. **12** (2004), 793–820.
- [PH] R. C. Penner and J. L. Harer, *Combinatorics of train tracks*, *Annals of Mathematics Studies*, **125**, Princeton University Press, Princeton, NJ, 1992.
- [PP87] A. Papadopoulos and R. C. Penner, *A characterization of pseudo-Anosov foliations*, Pacific J. Math., **130** (1987), 359–377.
- [PP93] A. Papadopoulos and R. C. Penner, *The Weil–Petersson symplectic structure at Thurston’s boundary*, Trans. Amer. Math. Soc. **335** (1993), 891–904.
- [Th88] W. Thurston, *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Amer. Math. Soc. (N.S.) **19** (1988), 417–431.

TSUKASA ISHIBASHI, RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KITASHIRAKAWA OIWAKE-CHO, SAKYO-KU, KYOTO 606-8502, JAPAN.

Email address: ishiba@kurims.kyoto-u.ac.jp

SHUNSUKE KANO, RESEARCH ALLIANCE CENTER FOR MATHEMATICAL SCIENCES, TOHOKU UNIVERSITY 6-3 AOBA, ARAMAKI, AOBA-KU, SENDAI, MIYAGI 980-8578, JAPAN.

Email address: s.kano@tohoku.ac.jp