

IDENTITIES AND BASES IN THE HYPOPLACTIC MONOID

ALAN J. CAIN, ANTÓNIO MALHEIRO, AND DUARTE RIBEIRO

ABSTRACT. This paper presents new results on the identities satisfied by the hypoplactic monoid. We show how to embed the hypoplactic monoid of any rank strictly greater than 2 (including infinite rank) into a direct product of copies of the hypoplactic monoid of rank 2. This confirms that all hypoplactic monoids of rank greater than or equal to 2 satisfy exactly the same identities. We then give a complete characterization of those identities, and prove that the variety generated by the hypoplactic monoid has finite axiomatic rank, by giving a finite basis for it.

1. INTRODUCTION

A (non-trivial) identity is a formal equality $u \approx v$, where u and v are words over some alphabet of variables, which is not of the form $u \approx u$. If a monoid is known to satisfy an identity, an important question is whether the set of identities it satisfies is finitely based, that is, if all these identities are consequences of those in some finite subset (see [37, 42]).

The plactic monoid plac , also known as the monoid of Young tableaux, is an important algebraic structure, first studied by Schensted [38] and Knuth [23], and later studied in depth by Lascoux and Schützenberger [27]. It is connected to many different areas of Mathematics, such as algebraic combinatorics, symmetric functions [29], crystal bases [4] and representation theory [14]. In particular, the question of identities in the plactic monoid has received a lot of attention recently [26, 20].

Finitely-generated polynomial-growth groups are virtually nilpotent and so satisfy identities [16]. In [40], it was possible to construct examples of finitely-generated polynomial-growth semigroups that do not satisfy identities, but it would be interesting to have a natural class of monoids with this property. One of the initial motivations for the study of the plactic monoids of finite rank, in this context, was to obtain a more natural example, although it is now known that they all satisfy identities. For example, the plactic monoid of rank 2 satisfies Adjan's identity $xyyxyxyyyx \approx xyxyxyyyx$, the shortest non-trivial identity satisfied by the bicyclic monoid [1]. The plactic monoid of rank 3 satisfies the identity

2020 *Mathematics Subject Classification.* Primary 08B05; Secondary 05E99, 20M05, 20M07, 20M32.

Key words and phrases. Hypoplactic monoid, variety, identities, equational basis, axiomatic rank.

This work is funded by National Funds through the FCT – Fundação para a Ciência e a Tecnologia, I.P., under the scope of the project UIDB/00297/2020 (Center for Mathematics and Applications) and the project PTDC/MAT-PUR/31174/2017.

The third author is funded by National Funds through the FCT – Fundação para a Ciência e a Tecnologia, I.P., under the scope of the studentship SFRH/BD/138949/2018.

$uvvuvu = uvvvuv$, where $u(x, y)$ and $v(x, y)$ are respectively the left and right side of Adjan's identity [26]. However, it does not satisfy Adjan's identity itself.

On the other hand, the plactic monoid of rank n does not satisfy any non-trivial identity of length less than or equal to n [6]. Thus, there is no "global" identity satisfied by the plactic monoid of every rank and, consequently, the infinite-rank plactic monoid. Recently, a tropical representation for the plactic monoid of rank n has been constructed [22], which implies that every plactic monoid of finite rank satisfies a non-trivial identity.

In terms of identities, there is some relationship between plactic monoids, tropical algebra, and the bicyclic monoid: by [10], the monoid of 2×2 upper-triangular tropical matrices (see, for example, [30]), the bicyclic monoid, and the plactic monoid of rank 2 satisfy precisely the same identities. The bicyclic monoid is not finitely based [39], so none of these monoids are. This can be shown by using polytopes to analyze identities holding in the bicyclic monoid [35].

Monoids related to the plactic monoid, such as the hypoplactic monoid **hypo** (the monoid of quasi-ribbon tableaux), sylvester monoid **sylv** (the monoid of binary search trees [18]), Baxter monoid (pairs of twin binary search trees, connected with Baxter permutations [15]) satisfy identities, and the shortest identities have been characterized [8]. These identities are independent of rank (except for rank 1, since the monoids of rank 1 are all isomorphic to the free monogenic monoid and thus commutative). Also related to the plactic monoid, by its growth type [11], the Chinese monoid [9] embeds into a direct product of copies of the bicyclic monoid [21]. Furthermore, the Chinese and plactic monoids of rank 2 coincide, hence, they satisfy the same identities.

This paper focuses on the hypoplactic monoid **hypo**, first studied in depth by Novelli [34], which stands as the analogue of the plactic monoid in the theory of non-commutative symmetric functions and quasi-symmetric functions (see [24, 25]). It is also known as the monoid of quasi-ribbon tableaux, combinatorial objects which index quasi-ribbon functions and that are computed from words over the free monoid, using the Krob-Thibon insertion algorithm. The hypoplactic monoid arises by factoring the free monoid by the hypoplactic congruence \equiv_{hypo} , which can be defined in several ways, one of which is $u \equiv_{\text{hypo}} v$ if and only if u and v yield the same quasi-ribbon tableaux. Similarly to the plactic monoid and the crystal graph, in the sense of Kashiwara, the hypoplactic monoid has been shown to accept a quasi-crystal structure, which interacts neatly with the combinatorics of the quasi-ribbon tableaux and the hypoplactic version of the Robinson-Schensted-Knuth correspondence [7].

The main goal of this paper is to present a systematic study of the identities satisfied by the hypoplactic monoid. The paper is structured as follows: Section 2 provides the necessary background on universal algebra and the hypoplactic monoid. In Section 3, we first show that the hypoplactic monoid of rank n , denoted by hypo_n , with $n \geq 2$, does not embed into the hypoplactic monoid of rank 2. However, we show how to embed it into a direct product of copies of hypo_2 . We also show how to embed the infinite rank hypoplactic monoid **hypo** into a direct product of infinite copies of hypo_2 . Thus, we prove that all these monoids generate the same variety, denoted by \mathbf{V}_{hypo} , and satisfy the same identities. We deduce the basis rank of \mathbf{V}_{hypo} . Section 4 gives a complete characterization of the identities satisfied by **hypo**, along with some immediate consequences. Furthermore, it also

presents a finite basis for \mathbf{V}_{hypo} , the variety generated by **hypo**, thus showing it has finite axiomatic rank.

2. PRELIMINARIES AND NOTATION

This section gives the necessary background on universal algebra (see [5, 31, 33, 2]), in the context of monoids, followed by the definition and essential facts about the hypoplactic monoid.

For the necessary background on semigroups and monoids, see [19]; for presentations, see [17]; for a general background on the plactic monoid, see [28, Chapter 5].

2.1. Varieties, identities and bases. Let X be a non-empty set, referred to as an *alphabet*. The free semigroup over the alphabet X , denoted by X^+ , is the set of all words over X , under the operation of word concatenation. If we include the empty word, denoted by ε , we obtain the free monoid over the alphabet X , denoted by X^* .

For any word $u \in X^*$, the *length* of u is denoted by $|u|$, and for any $x \in X$, the number of occurrences of x in u is denoted by $|u|_x$. Suppose $u = u_1 \cdots u_k$, where $u_i \in X$. For any $1 \leq i \leq j \leq k$, the word $u_i \cdots u_j$ is a *factor* of u . For any $i_1, \dots, i_m \in \{1, \dots, k\}$ such that $i_1 < \cdots < i_m$, the word $u_{i_1} \cdots u_{i_m}$ is a *subsequence* of u .

The *content* of u , denoted by $c(u)$, is the infinite tuple of non-negative integers, indexed by X , whose x -th element is $|u|_x$. The *support* of u , denoted by $\text{supp}(u)$, is the subset of X such that $x \in \text{supp}(u) \iff |u|_x \geq 1$. Notice that if two words share the same content, then they also share the same support.

A (*monoid*) *identity*, over an alphabet of variables X , is a formal equality $u \approx v$, where u and v are words in the free monoid X^* , and is non-trivial if $u \neq v$. We say a variable x *occurs* in $u \approx v$ if $x \in \text{supp}(u)$ or $x \in \text{supp}(v)$. The *rank* of an identity is the number of distinct variables which occur in it. We say that an identity *holds* in a monoid M (or that M satisfies the identity) if for every morphism $\psi : X^* \rightarrow M$ (also referred to as an *evaluation*), the equality $\psi(u) = \psi(v)$ holds in M . In other words, M satisfies the identity $u \approx v$ if equality in M holds under every substitution of the variables of u and v by elements of M .

An identity $u \approx v$ is *balanced* if $c(u) = c(v)$. Any identity satisfied by a monoid that contains a free monogenic submonoid, such as the aforementioned plactic and related monoids, must be balanced.

Let Σ be a nonempty set of identities, over an alphabet X . An identity $u \approx v$ is said to be a *consequence* of Σ if there exist $k \in \mathbb{N}$, words $w_1, \dots, w_k \in X^*$, and substitutions $\sigma_1, \dots, \sigma_{k-1}$ of variables by elements of X^* (that is, endomorphisms of X^*), such that $u = w_1$, $v = w_k$ and, for $1 \leq i < k$,

$$\begin{aligned} w_i &= r_i \sigma_i(p_i) s_i, \\ w_{i+1} &= r_i \sigma_i(q_i) s_i, \end{aligned}$$

for some $p_i, q_i, r_i, s_i \in X^*$ such that $p_i \approx q_i$ or $q_i \approx p_i$ is in Σ .

A class of monoids \mathbf{K} is a *monoid equational class* (or simply an *equational class*) if there exists a set of identities Σ such that \mathbf{K} is the class of all monoids that satisfy all identities in Σ . Dually, a set of identities Σ is an *equational theory* if there exists a class of monoids \mathbf{K} such that Σ is the set of identities satisfied by all monoids in \mathbf{K} . When this holds, Σ is called the *equational theory of \mathbf{K}* . Equational theories are closed under taking consequences.

On the other hand, a class of monoids \mathbf{K} is a *monoid variety* (or simply a *variety*) if it is closed under the taking of homomorphic images, submonoids and direct products. For any class of monoids \mathbf{K} , we say that a variety is generated by \mathbf{K} if it is the smallest variety containing \mathbf{K} , and denote it by $\mathbf{V}(\mathbf{K})$. Similarly, we say a variety \mathbf{K} is generated by a monoid M if it is the smallest variety containing M , and denote it by $\mathbf{V}(M)$.

The notions of equational class and variety coincide, as seen in the following theorem, known as Birkhoff's *HSP* Theorem:

Theorem 2.1 ([3]). *\mathbf{K} is an equational class if and only if \mathbf{K} is a variety.*

An important corollary of this theorem is that, for any monoid M , the identities satisfied by M must also be satisfied by all other monoids in $\mathbf{V}(M)$.

A set of identities \mathcal{B} is an *equational basis* (or simply *basis*) of a variety \mathbf{V} if the equational theory of \mathbf{V} consists of all consequences of \mathcal{B} . A variety \mathbf{V} is *finitely based* if it admits a finite basis. The *axiomatic rank* of \mathbf{V} is the least natural number $r_a(\mathbf{V})$ such that \mathbf{V} admits a basis \mathcal{B} , where the rank of each identity in \mathcal{B} does not exceed $r_a(\mathbf{V})$. If no such natural number exists, we say that \mathbf{V} has *infinite axiomatic rank*. Notice that if \mathbf{V} is finitely based, then it has finite axiomatic rank.

A variety \mathbf{V} is always generated by its \mathbf{V} -free monoid $\mathcal{F}_\omega(\mathbf{V})$ over an infinite alphabet. However, \mathbf{V} may also be generated by a \mathbf{V} -free monoid $\mathcal{F}_n(\mathbf{V})$ over a finite alphabet with n letters, for some $n \in \mathbb{N}$. In such a case, the least natural number $r_b(\mathbf{V})$ such that \mathbf{V} is generated by $\mathcal{F}_{r_b(\mathbf{V})}(\mathbf{V})$ is called the *basis rank* of \mathbf{V} . If \mathbf{V} is generated by a monoid with a finite number of generators, the minimal such number coincides with the basis rank of \mathbf{V} .

A monoid M is *residually finite* (or *finitely approximable*) if it is embeddable in a direct product of a family of finite monoids. If a monoid is both residually finite and finitely presented, then the word problem is solvable for such a monoid (see, for example, [13]).

2.2. The hypoplactic monoid. This subsection gives a brief overview of the hypoplactic monoid and its related combinatorial object and insertion algorithm, as well as results from [8]. For more information, see [34] and [7].

Let $\mathcal{A} = \{1 < 2 < 3 < \dots\}$ denote the set of positive integers, viewed as an infinite ordered alphabet, and let $\mathcal{A}_n = \{1 < \dots < n\}$ denote the set of the first n positive integers, viewed as a finite ordered alphabet.

A *quasi-ribbon tableau* is a (finite) grid of cells, aligned so that the leftmost cell in each row is below the rightmost cell of the previous row, filled with letters from \mathcal{A} , such that the entries in each row are weakly increasing from left to right, and the entries in each column are strictly increasing from top to bottom. An example of a quasi-ribbon tableau is

1	2	4	
		5	
	6	6	
		7	8

Observe that the same letter cannot appear in two different rows of a quasi-ribbon tableau.

The following algorithm allows us to insert a letter from \mathcal{A} into an existing quasi-ribbon tableau, in order to obtain a new quasi-ribbon tableau:

Algorithm 1: *Krob–Thibon algorithm.*

Input: A quasi-ribbon tableau T and a letter $a \in \mathcal{A}$.

Output: A quasi-ribbon tableau $T \leftarrow a$.

Method:


if there is no entry in T that is less than or equal to a , **then** output the tableau obtained by creating a new cell, labelled with a , and gluing T by its top-leftmost entry to the bottom of this new cell;

else let x be the right-most and bottom-most entry of T that is less than or equal to a . Separate T in two parts, such that one part is from the top left down to and including x . Put a new entry a to the right of x and glue the remaining part of T (below and to the right of x) onto the bottom of the new entry a .

Output the resulting tableau.

Let $u \in \mathcal{A}^*$. Using the insertion algorithm above, we can compute a unique quasi-ribbon tableau $P_{\text{hypo}}(u)$ from u : we start with the empty tableau and insert the letters of u , one-by-one from left-to-right – see Example 2.2.

Example 2.2. Computing $P_{\text{hypo}}(12654768)$:



We define the relation \equiv_{hypo} on \mathcal{A}^* as follows: For $u, v \in \mathcal{A}^*$,

$$u \equiv_{\text{hypo}} v \iff P_{\text{hypo}}(u) = P_{\text{hypo}}(v).$$

This relation is a congruence on \mathcal{A}^* , called the *hypoplactic congruence*. The factor monoid $\mathcal{A}^*/\equiv_{\text{hypo}}$ is the infinite-rank *hypoplactic monoid*, denoted by hypo . The congruence \equiv_{hypo} naturally restricts to a congruence on \mathcal{A}_n^* , and the factor monoid $\mathcal{A}_n^*/\equiv_{\text{hypo}}$ is the *hypoplactic monoid of rank n* , denoted by hypo_n .

It follows from the definition of \equiv_{hypo} that each element $[u]_{\text{hypo}}$ of hypo can be identified with the combinatorial object $P_{\text{hypo}}(u)$.

Recall that the content of u describes the number of occurrences of each letter of \mathcal{A} in u . It is immediate from the definition of the hypoplactic monoid that if $u \equiv_{\text{hypo}} v$, then $c(u) = c(v)$. Thus, we can define the content of an element of hypo as the content of any word which represents it. Furthermore, since $c(u) = c(v)$ implies that $\text{supp}(u) = \text{supp}(v)$, we can also define the support of an element of hypo as the support of any word which represents it.

Notice that hypo_n is a submonoid of hypo , for each $n \in \mathbb{N}$, and, for $n, m \in \mathbb{N}$, if $n \leq m$, then hypo_n is a submonoid of hypo_m .

Let $u \in \mathcal{A}_n^*$. Suppose $\text{supp}(u) = \{a_1 < \cdots < a_k\}$, for some $k \in \mathbb{N}$. We say u has an $a_{i+1}-a_i$ *inversion*, for $1 \leq i \leq k-1$, if it admits $a_{i+1}a_i$ as a subsequence. In other words, when reading u from left-to-right, there is at least an occurrence of a_{i+1} before the last occurrence of a_i . Notice that we only consider inversions of consecutive elements of the support of w .

Example 2.3. The word 31214 has 3-2 and 2-1 inversions, but no 4-3 inversion. On the other hand, 21341 has a 2-1 inversion, but no 4-3 nor 3-2 inversions.

The following characterization of the hypoplactic monoid is a consequence of [34, Subsection 4.2]. It arises as a corollary of Theorem 4.18 and Note 4.10 of [34]:

Proposition 2.4. *For $u, v \in \mathcal{A}_n^*$, we have that $u \equiv_{\text{hypo}} v$ if and only if u and v share exactly the same content and inversions.*

Attending to the previous result, we say that an element $[u]_{\text{hypo}}$ of hypo has an $a_{i+1}-a_i$ inversion if the word u itself, and hence any other word in $[u]_{\text{hypo}}$, has an $a_{i+1}-a_i$ inversion. This characterization will be extensively used throughout the rest of this paper.

The hypoplactic monoid can also be defined by the presentation $\langle \mathcal{A} \mid \mathcal{R}_{\text{hypo}} \rangle$, where

$$\begin{aligned} \mathcal{R}_{\text{hypo}} = & \{(acb, cab) : a \leq b < c\} \\ & \cup \{(bac, bca) : a < b \leq c\} \\ & \cup \{(cadb, acbd) : a \leq b < c \leq d\} \\ & \cup \{(bdac, dbca) : a < b \leq c < d\}. \end{aligned}$$

The first two defining relations are known as the *plactic relations*, while the remaining two are known as the *hypoplactic relations*. A presentation for the hypoplactic monoid of rank n , for some $n \in \mathbb{N}$, can be obtained by restricting generators and relations of the above presentation to generators in \mathcal{A}_n . Notice that these relations are length-preserving. As such, it is easy to see that the hypoplactic monoids are residually finite.

3. EMBEDDINGS

In this section, we prove that the hypoplactic monoids of rank greater than or equal to 2 satisfy the same identities. We do this by constructing embeddings of hypoplactic monoids of any rank greater than 2 into direct products of copies of the hypoplactic monoid of rank 2, as it is not possible to directly embed one into another. Thus, they generate the same variety and, by Birkhoff's Theorem, satisfy exactly the same identities. We also show that the basis rank of the variety generated by hypo is 2.

3.1. Non-existence of embedding into a hypoplactic monoid of lesser rank. It is not possible to directly embed a hypoplactic monoid of finite rank into a hypoplactic monoid of lesser rank:

Proposition 3.1. *For all $n > m \geq 1$, there is no embedding of hypo_n into hypo_m .*

Proof. First of all, notice that hypo_1 is isomorphic to the free monogenic monoid and hypo_n is noncommutative, for any $n \geq 2$. Thus, there is no embedding of hypo_n into hypo_1 .

On the other hand, if there exists an embedding of \mathbf{hypo}_n into \mathbf{hypo}_m , for some $n > m \geq 2$, then, since \mathbf{hypo}_m is a submonoid of \mathbf{hypo}_{n-1} , there must also exist an embedding of \mathbf{hypo}_n into \mathbf{hypo}_{n-1} . As such, we just need to prove that this second embedding cannot exist.

Suppose, in order to obtain a contradiction, that there exists $n \geq 3$ such that we have an embedding $\phi : \mathbf{hypo}_n \hookrightarrow \mathbf{hypo}_{n-1}$. Without loss of generality, suppose n is the smallest positive integer in such conditions.

Observe that $\text{supp}\left(\phi\left([1 \cdots (n-1)]_{\mathbf{hypo}_n}\right)\right) = \mathcal{A}_{n-1}$, that is, the image of the product of all generators of \mathbf{hypo}_n , except for n , has all the possible letters of \mathcal{A}_{n-1} . Indeed, if $\text{supp}\left(\phi\left([1 \cdots (n-1)]_{\mathbf{hypo}_n}\right)\right) \subsetneq \mathcal{A}_{n-1}$, we would be able to construct an embedding from the submonoid \mathbf{hypo}_{n-1} of \mathbf{hypo}_n , generated by all generators of \mathbf{hypo}_n except for n , into a submonoid of \mathbf{hypo}_{n-1} isomorphic to \mathbf{hypo}_{n-2} . This contradicts the minimality of n .

Hence, $\phi\left([1 \cdots (n-1)]_{\mathbf{hypo}_n}^2\right)$ has all the possible inversions of letters of \mathcal{A}_{n-1} . If we multiply this element by any other element of \mathbf{hypo}_{n-1} , either on the left or the right, we obtain the same result, by Proposition 2.4. Thus, we have that

$$\phi\left([n]_{\mathbf{hypo}_n} \cdot [1 \cdots (n-1)]_{\mathbf{hypo}_n}^2\right) = \phi\left([1 \cdots (n-1)]_{\mathbf{hypo}_n}^2 \cdot [n]_{\mathbf{hypo}_n}\right).$$

On the other hand, we have that

$$[n]_{\mathbf{hypo}_n} \cdot [1 \cdots (n-1)]_{\mathbf{hypo}_n}^2 \neq [1 \cdots (n-1)]_{\mathbf{hypo}_n}^2 \cdot [n]_{\mathbf{hypo}_n}.$$

This contradicts our hypothesis that ϕ is injective. Hence, for all $n \geq 2$, there is no embedding of \mathbf{hypo}_n into \mathbf{hypo}_{n-1} . As such, there is no embedding of \mathbf{hypo}_n into \mathbf{hypo}_m , for $n > m \geq 2$. \square

Corollary 3.2. *There is no embedding of \mathbf{hypo} into \mathbf{hypo}_n , for any $n \in \mathbb{N}$.*

Proof. If such an embedding existed, for some $n \in \mathbb{N}$, then, by restricting the embedding to the first $n+1$ generators of \mathbf{hypo} , we would obtain an embedding of \mathbf{hypo}_{n+1} into \mathbf{hypo}_n , which contradicts the previous proposition. \square

3.2. Embedding into a direct product of copies of the hypoplactic monoid of rank 2. Although it is not possible to directly embed a hypoplactic monoid of finite rank n into the hypoplactic monoid of rank 2, for $n > 2$, we now show how to embed \mathbf{hypo}_n into a direct product of copies of \mathbf{hypo}_2 . We also show how to embed \mathbf{hypo} into a direct product of infinitely many copies of \mathbf{hypo}_2 . For simplicity, we shall denote by M^k the direct product of k copies of a monoid M .

For any $i, j \in \mathcal{A}$, with $i < j$, define a map from \mathcal{A} to \mathbf{hypo}_2 in the following way: For any $a \in \mathcal{A}$,

$$a \mapsto \begin{cases} [1]_{\mathbf{hypo}_2} & \text{if } a = i; \\ [2]_{\mathbf{hypo}_2} & \text{if } a = j; \\ [21]_{\mathbf{hypo}_2} & \text{if } i < a < j; \\ [\varepsilon]_{\mathbf{hypo}_2} & \text{otherwise;} \end{cases}$$

and extend it to a homomorphism $\varphi_{ij} : \mathcal{A}^* \rightarrow \mathbf{hypo}_2$, in the usual way.

Note that $\varphi_{ij}(w)$ is the hypoplactic class of the word obtained from w by replacing any occurrence of i by 1; any occurrence of j by 2; any occurrence of an a , with $i < a < j$, by 21; and erasing any occurrence of any other element.

Lemma 3.3. φ_{ij} factors to give a homomorphism $\varphi_{ij} : \text{hypo} \longrightarrow \text{hypo}_2$.

Proof. Since hypo is given by the presentation $\langle \mathcal{A} \mid \mathcal{R}_{\text{hypo}} \rangle$, we just need to verify that both sides of the plactic and hypoplactic relations have the same image under φ_{ij} . Let $a, b, c, d \in \mathcal{A}$. Assume, without loss of generality, that φ_{ij} does not map any letter to $[\varepsilon]_{\text{hypo}_2}$.

If $a \leq b < c \leq d$, then either φ_{ij} maps at least one letter to $[21]_{\text{hypo}_2}$, or φ_{ij} maps a and b to $[1]_{\text{hypo}_2}$, and c and d to $[2]_{\text{hypo}_2}$. Thus,

$$\varphi_{ij}(acb) = \varphi_{ij}(cab) \quad \text{and} \quad \varphi_{ij}(acbd) = \varphi_{ij}(cadb),$$

since both classes in each side of the equalities have 2-1 inversions.

If $a < b \leq c$, then either φ_{ij} maps at least one letter to $[21]_{\text{hypo}_2}$, or φ_{ij} maps a to $[1]_{\text{hypo}_2}$, and b and c to $[2]_{\text{hypo}_2}$. Thus,

$$\varphi_{ij}(bac) = \varphi_{ij}(bca).$$

since both classes in each side of the equality have 2-1 inversions.

If $a < b \leq c < d$, then φ_{ij} maps b and c to $[21]_{\text{hypo}_2}$. Thus, by the same reasoning,

$$\varphi_{ij}(bdac) = \varphi_{ij}(dbca).$$

Hence, $\mathcal{R}_{\text{hypo}} \subseteq \ker \varphi_{ij}$. □

Let $w \in \mathcal{A}_n^*$, for some $n \geq 3$. Suppose $\text{supp}(w) = \{a_1 < \dots < a_k\}$, for some $k \in \mathbb{N}$. Observe that, ranging $1 \leq i < k$, we can get from the maps $\varphi_{a_i a_{i+1}}$ the number of occurrences of a_i and a_{i+1} in w . Furthermore, we can also check if w has an a_{i+1} - a_i inversion: Since no element $a \in \mathcal{A}$ such that $a_i < a < a_{i+1}$ occurs in w , each occurrence of 1 in (a word in) $\varphi_{a_i a_{i+1}}([w]_{\text{hypo}})$ corresponds to an occurrence of a_i in w and, similarly, each occurrence of 2 in $\varphi_{a_i a_{i+1}}([w]_{\text{hypo}})$ corresponds to an occurrence of a_{i+1} in w . Thus, $\varphi_{a_i a_{i+1}}([w]_{\text{hypo}})$ has a 2-1 inversion if and only if w has an a_{i+1} - a_i inversion. Hence, we get the following lemma:

Lemma 3.4. Let $u, v \in \mathcal{A}_n^*$. Then, $u \equiv_{\text{hypo}} v$ if and only if $\varphi_{ij}([u]_{\text{hypo}}) = \varphi_{ij}([v]_{\text{hypo}})$, for all $1 \leq i < j \leq n$.

Proof. The direct implication is trivial, since φ_{ij} is well-defined as a map, for all $1 \leq i < j \leq n$. The proof of the converse follows from the previous observations, as well as Proposition 2.4. □

For each $n \in \mathbb{N}$, with $n \geq 3$, let I_n be the index set

$$\{(i, j) : 1 \leq i < j \leq n\},$$

and let $I := \bigcup_{n \in \mathbb{N}} I_n$. Now, consider the map

$$\phi_n : \text{hypo}_n \longrightarrow \prod_{I_n} \text{hypo}_2,$$

whose (i, j) -th component is given by $\varphi_{ij}([w]_{\text{hypo}})$, for $w \in \mathcal{A}_n^*$ and $(i, j) \in I_n$.

Proposition 3.5. The map ϕ_n is an embedding.

Proof. It is clear that ϕ_n is a homomorphism. It follows from the definition of ϕ_n and Lemma 3.4 that, for any $u, v \in \mathcal{A}_n^*$, we have $u \equiv_{\text{hypo}_n} v$ if and only if $\phi_n([u]_{\text{hypo}}) = \phi_n([v]_{\text{hypo}})$, hence ϕ_n is an embedding. □

Thus, for each $n \in \mathbb{N}$, we can embed hypo_n into a direct product of $\binom{n}{2}$ copies of hypo_2 .

Similarly, we can embed hypo into a direct product of infinitely many copies of hypo_2 . Consider the map

$$\phi : \text{hypo} \longrightarrow \prod_I \text{hypo}_2,$$

whose (i, j) -th component is given by $\varphi_{ij}([w]_{\text{hypo}})$, for $w \in \mathcal{A}^*$ and $(i, j) \in I$.

Proposition 3.6. *The map ϕ is an embedding.*

Proof. It is clear that ϕ is a homomorphism. Notice that, for any word $w \in \mathcal{A}^*$, there must exist $n \in \mathbb{N}$ such that $w \in \mathcal{A}_n^*$. Furthermore, for $(i, j) \in I_n$, we have that the (i, j) -th component of $\phi([w]_{\text{hypo}})$ is equal to the (i, j) -th component of $\phi_n([w]_{\text{hypo}})$. Thus, for $u, v \in \mathcal{A}^*$, if $\phi([u]_{\text{hypo}}) = \phi([v]_{\text{hypo}})$, then $\phi_n([u]_{\text{hypo}}) = \phi_n([v]_{\text{hypo}})$, for some $n \in \mathbb{N}$ such that $u, v \in \mathcal{A}_n^*$.

It follows from Lemma 3.4 that, for any $u, v \in \mathcal{A}^*$, we have $u \equiv_{\text{hypo}} v$ if and only if $\phi([u]_{\text{hypo}}) = \phi([v]_{\text{hypo}})$, hence ϕ is an embedding. \square

As such, all hypoplactic monoids of rank strictly greater than 2 are in the variety generated by hypo_2 . Since hypo_2 is a submonoid of hypo and hypo_n , for any $n \geq 3$, they all generate the same variety, which we will denote by \mathbf{V}_{hypo} . Thus, by Birkhoff's Theorem, we have the following result:

Theorem 3.7. *For any $n \geq 2$, hypo and hypo_n satisfy exactly the same identities.*

Another consequence of \mathbf{V}_{hypo} being generated by hypo_2 is the following:

Proposition 3.8. *The basis rank of \mathbf{V}_{hypo} is 2.*

Proof. Since \mathbf{V}_{hypo} is generated by hypo_2 , and hypo_2 is defined by a presentation where the alphabet has two generators, then $r_b(\mathbf{V}_{\text{hypo}})$ is less than or equal to 2.

On the other hand, notice that any monoid generated by a single element is commutative. Since hypo is not commutative, \mathbf{V}_{hypo} cannot be generated by any monoid which is itself generated by a single element. As such, $r_b(\mathbf{V}_{\text{hypo}})$ is strictly greater than 1.

Hence, the basis rank of \mathbf{V}_{hypo} is 2. \square

4. IDENTITIES AND BASES

In this section, we obtain a complete characterization of the identities satisfied by hypo , a finite basis for \mathbf{V}_{hypo} and also its axiomatic rank.

4.1. Characterization of the identities satisfied by the hypoplactic monoid.

Due to Theorem 3.7, the identities satisfied by hypo and those satisfied by hypo_2 are exactly the same. We shall use hypo_2 to obtain a characterization of the identities satisfied by hypo .

Observe that, for each element of hypo_2 , there is at most one other distinct element of hypo_2 which has the same content as it. Indeed, for an element of hypo_2 , with fixed content and support $\{1, 2\}$, either it has a 2-1 inversion or not.

Theorem 4.1. *The identities $u \approx v$ satisfied by hypo are exactly the balanced identities such that, for any variables x, y that occur in u and v , u admits xy as a subsequence if and only if v does too.*

Proof. We first prove by contradiction that an identity satisfied by hypo_2 must satisfy the previously mentioned conditions. Suppose $u \approx v$ is an identity satisfied by hypo_2 . Since hypo contains the free monogenic submonoid, we know that any identity satisfied by hypo must be a balanced identity. Thus, we assume $u \approx v$ is a balanced identity.

Suppose, in order to obtain a contradiction, that there exist variables x, y occurring in u and v , such that u admits xy as a subsequence, but v does not. Observe that, since both x and y occur in v , then v must admit yx as a subsequence.

Then, taking the evaluation ψ of X in hypo_2 such that $\psi(x) = [2]_{\text{hypo}_2}$, $\psi(y) = [1]_{\text{hypo}_2}$ and $\psi(z) = [\varepsilon]_{\text{hypo}_2}$, for all other variables z , we have

$$\psi(u) = [2^\alpha 1^\beta]_{\text{hypo}_2} \text{ and } \psi(v) = [1^\beta 2^\alpha]_{\text{hypo}_2}.$$

Thus, by Proposition 2.4, we have that $\psi(u) \neq \psi(v)$, which contradicts our hypothesis that $u \approx v$ is an identity.

We now prove by contradiction that an identity which satisfies the previously mentioned conditions must also be satisfied by hypo_2 . Suppose that $u \approx v$ is a balanced identity, such that, for all variables x, y which occur in u and v , u admits xy as a subsequence if and only if v does too. Suppose, in order to obtain a contradiction, that there is some evaluation ψ of X in hypo_2 such that $\psi(u) \neq \psi(v)$.

Notice that, since $u \approx v$ is a balanced identity, then $\psi(u)$ and $\psi(v)$ have the same content. Since we assumed that $\psi(u) \neq \psi(v)$, then, by Proposition 2.4, we have that $\text{supp}(\psi(u)) = \text{supp}(\psi(v)) = \{1, 2\}$, and $\psi(u)$ has a 2-1 inversion and $\psi(v)$ does not, or vice-versa.

We assume, without loss of generality, that $\psi(u)$ has a 2-1 inversion and $\psi(v)$ does not. Note that $\psi(v)$, as a word class, has only one word, of the form $1^\alpha 2^\beta$, for some $\alpha, \beta \geq 1$. Then, v must be of the form $v = v_1 v_2$ or $v = v_1 z v_2$, with $z \in X$, where: for each variable x which occurs in v_1 , $\psi(x)$ has support $\{1\}$; for each variable y which occurs in v_2 , $\psi(y)$ has support $\{2\}$; and $\psi(z)$ has support $\{1, 2\}$.

Notice that z is a variable that occurs in neither v_1 nor v_2 , and that no variable occurs simultaneously in v_1 and v_2 . Also notice that, for any variables x , which occurs in v_1 , and y , which occurs in v_2 , v admits xz, xy and zy as subsequences (if there is a variable z in the previously mentioned conditions), but not zx, yx nor yz . Thus, by the conditions imposed upon $u \approx v$, u must be of the form $u = u_1 u_2$ or $u = u_1 z u_2$, where $c(u_1) = c(v_1)$ and $c(u_2) = c(v_2)$. Hence, $\psi(u_1) = \psi(v_1)$ and $\psi(u_2) = \psi(v_2)$, by the observations in the previous paragraph.

Thus, we either have

$$\psi(u) = \psi(u_1)\psi(u_2) = \psi(v_1)\psi(v_2) = \psi(v)$$

or

$$\psi(u) = \psi(u_1)\psi(z)\psi(u_2) = \psi(v_1)\psi(z)\psi(v_2) = \psi(v).$$

Thus, we have reached a contradiction, hence, there is no evaluation ψ of X in hypo_2 such that $\psi(u) \neq \psi(v)$. Thus, $u \approx v$ is an identity satisfied by hypo_2 . \square

With this characterization, we recover as a corollary the following result:

Corollary 4.2 ([8, Proposition 12]). *The following non-trivial identities are satisfied by hypo :*

$$\begin{aligned} xyxy &\approx xyxx \approx yxyx \approx yxyx; \\ xxyx &\approx xyxx. \end{aligned}$$

Furthermore, up to equivalence (that is, up to renaming variables or swapping both sides of the identities), these are the shortest non-trivial identities satisfied by **hypo**.

We also easily obtain some examples of important non-trivial identities satisfied by **hypo**:

Example 4.3. The following non-trivial identities are satisfied by **hypo**:

$$\begin{aligned} \text{(L)} \quad & xyzxty \approx yxzxtty; \\ \text{(M)} \quad & xzxytx \approx xzyxttx; \\ \text{(R)} \quad & xzytxy \approx xzytyx. \end{aligned}$$

It is well known that the set of all balanced identities is the equational theory of the variety **Com** of all commutative monoids, which is generated by the free monogenic monoid. On the other hand, the set J_2 of all identities $u \approx v$ where u and v share exactly the same subsequences of length at most 2 is the equational theory of the pseudovariety \mathbf{J}_2 , which, due to Eilenberg's correspondence (see [12, 36]), corresponds to the class of all piecewise testable languages of height 2 (see [41]). This pseudovariety is generated by \mathcal{C}_3 , the 5-element monoid of all order preserving and extensive transformations of the chain $1 < 2 < 3$ (see [43]). Thus, the equational theory of $\mathbf{V}(\mathcal{C}_3)$, the variety generated by \mathcal{C}_3 , is J_2 . It is easy to see that the equational theory of \mathbf{V}_{hypo} is the intersection of the set of all balanced identities and the set J_2 . As such, we have the following corollary of Theorem 4.1, suggested by the anonymous referee:

Corollary 4.4. \mathbf{V}_{hypo} is the varietal join $\mathbf{Com} \vee \mathbf{V}(\mathcal{C}_3)$, and is generated by the free monogenic monoid and the monoid \mathcal{C}_3 .

An alternative characterization of the identities satisfied by **hypo** is the following:

Corollary 4.5. The identities $u \approx v$ satisfied by **hypo** are balanced identities such that, for any variables x, y that occur in u and v , $u|_{x,y} \approx v|_{x,y}$ is an identity satisfied by **hypo**, where $u|_{x,y}$ and $v|_{x,y}$ are obtained from u and v , respectively, by eliminating every occurrence of a variable other than x or y .

Proof. The proof of the implication is trivial, as it follows from the definition of an identity satisfied by a monoid.

Suppose then that the identity $u \approx v$ is balanced and satisfies the following property: For any variables x, y that occur in u and v , $u|_{x,y} \approx v|_{x,y}$ is an identity satisfied by **hypo**.

By Theorem 4.1, to prove that $u \approx v$ is satisfied by **hypo**, we only need to show that u admits xy (or yx) as a subsequence if and only if v does too, for any variables x, y that occur in u and v .

Let x, y be variables that occur in u and v . Hence, by the hypothesis, we have that $u|_{x,y} \approx v|_{x,y}$ is an identity satisfied by **hypo**. By Theorem 4.1, we know that $u|_{x,y}$ admits xy (or yx) as a subsequence if and only if $v|_{x,y}$ does too.

Notice that $u|_{x,y}$ is the unique subsequence of u with the same number of occurrences of x and y . Therefore, we can conclude that $u|_{x,y}$ admits xy (or yx) as a subsequence if and only if u does too. The same can be stated about $v|_{x,y}$ and v .

Therefore, u admits xy (or yx) as a subsequence if and only if v does too. Hence, we conclude that $u \approx v$ is an identity satisfied by **hypo**. \square

It is very easy to verify if a balanced identity, over a two-letter alphabet, is satisfied by **hypo**, by the following complete characterization:

Corollary 4.6. *The non-trivial identities, over the two-letter alphabet $\{x, y\}$, satisfied by **hypo** are balanced identities such that neither side of the identity is of the form $x^a y^b$ or $y^b x^a$, for some $a, b \in \mathbb{N}$.*

Proof. Let us first suppose that $u \approx v$ be a non-trivial identity, over the two-letter alphabet $\{x, y\}$, satisfied by **hypo**. By the previous theorem, we know that this identity must be balanced. Suppose, without loss of generality, that $u = x^a y^b$, for some $a, b \in \mathbb{N}$. Then, since u does not admit yx as a subsequence, v cannot as well. Since $u \approx v$ is a balanced identity, we conclude that $v = x^a y^b = u$, which contradicts our hypothesis that $u \approx v$ is non-trivial.

Since this argument can be applied to all other possible cases, we conclude that neither side of the identity is of the form $x^a y^b$ or $y^b x^a$.

Conversely, let $u \approx v$ be a non-trivial, balanced identity, over the two-letter alphabet $\{x, y\}$, such that neither side of the identity is of the form $x^a y^b$ or $y^b x^a$, for some $a, b \in \mathbb{N}$. The, since $u \approx v$ is a non-trivial identity, both x and y must occur at least once in both u and v .

Observe that the only words over $\{x, y\}$, where both x and y occur, that do not admit yx as a subsequence, are words of the form $x^a y^b$. Similarly, the only words over $\{x, y\}$, where both x and y occur, that do not admit xy as a subsequence, are words of the form $y^b x^a$. Thus, both u and v admit xy and yx as subsequences. Hence, by the previous Theorem, $u \approx v$ is satisfied by **hypo**. \square

The following corollary will be important in the next subsection:

Corollary 4.7. *The shortest non-trivial identity, with n variables, satisfied by **hypo**, is of length $n + 2$.*

Proof. It is immediate, by the previous theorem, that for variables $x, a_1 \dots a_{n-1}$,

$$xa_1 \dots a_{n-1}xx \approx xx a_1 \dots a_{n-1}x$$

is an identity satisfied by **hypo**.

On the other hand, suppose that there exists a non-trivial identity $u \approx v$, with n variables, satisfied by **hypo**. Let \mathcal{X} be the set of variables which occur in u and v . By Theorem 4.1, if u admits a subsequence xy , for variables $x, y \in \mathcal{X}$, then v must also admit such a subsequence. Thus, we easily conclude that the identity cannot be of length n , otherwise it would be trivial. As such, there must be some variable z which occurs twice in u and v .

If $u \approx v$ is of length $n + 1$, then z occurs twice and all other variables a_1, \dots, a_{n-1} occur only once, in u and v . Assume, without loss of generality, that u admits the subsequence $a_1 \dots a_{n-1}$. Again, by Theorem 4.1, v must also admit this subsequence.

Let i be the greatest index such that u admits $a_i z$ as a subsequence, but not $z a_i$, and let j be the smallest index such that u admits $z a_j$ as a subsequence, but not $a_j z$. Since z occurs exactly twice in u , we can conclude that

$$u = a_1 \dots a_i z a_{i+1} \dots a_{j-1} z a_j \dots a_{n-1}.$$

Once again, by Theorem 4.1, v must also satisfy these conditions. Hence, we can conclude that $u = v$.

This contradicts the hypothesis that $u \approx v$ is a non-trivial identity satisfied by *hypo*. Hence, there is no non-trivial identity, with n variables, satisfied by *hypo*, of length $n + 1$. \square

4.2. The axiomatic rank of the variety generated by the hypoplactic monoid. Now, we prove that not only \mathbf{V}_{hypo} has finite axiomatic rank, it is also finitely based. We give a basis for \mathbf{V}_{hypo} with three identities, all of them over a four-letter alphabet, each of length 6. This basis is minimal, in the sense that no identity in this basis is a consequence of the others, and also that each identity is of minimal length, for identities satisfied by *hypo* over a four-letter alphabet. Furthermore, we also prove that there exists no basis for \mathbf{V}_{hypo} with only identities over an alphabet with at most three variables, thus showing that the axiomatic rank of \mathbf{V}_{hypo} is 4.

Theorem 4.8. \mathbf{V}_{hypo} admits a finite basis $\mathcal{B}_{\text{hypo}}$, with the following identities:

- (L) $xyzxty \approx yxzxty;$
(M) $xzxytx \approx xzytx;$
(R) $xzytxy \approx xzytx.$

Proof. Let $\mathcal{B}_{\text{hypo}}$ be the set of identities which contains the previously mentioned identities. Notice that these identities are the ones given in Example 4.3.

The following proof will be done by induction, in the following sense: First, we order identities by their length. Then, within a set of identities of the same length, we order them by the length of the common prefix of both sides of the identity. The induction will be first on the length of the identities, and then on the length of the suffix, that is, the length of the identity minus the length of the common prefix.

The base cases for the induction on the length of the identities are those given in Corollary 4.2, as they are, up to equivalence, the shortest non-trivial identities satisfied by *hypo*. Notice that these identities are consequences of $\mathcal{B}_{\text{hypo}}$: to show this, we just need to replace the variables z and t with the empty word, and, if necessary, rename the remaining variables.

The base cases for the induction on the length of the suffix, for identities of length n (with $n \geq 4$), are those identities of the form

$$wxy \approx wyx,$$

where w is a word of length $n - 2$ and x, y are variables. Observe that, since any identity $u \approx v$ satisfied by *hypo* is a balanced identity, there are no non-trivial identities, of length n , with a common prefix of length greater than $n - 2$, satisfied by *hypo*. Furthermore, since wxy admits a subsequence xy and wyx admits a subsequence yx , then x and y must both occur in w . Thus, w is of the form

$$w_1xw_2yw_3 \quad \text{or} \quad w_1yw_2xw_3,$$

for some words w_1, w_2, w_3 . Therefore, by replacing z with w_2 , and t by w_3 , and, if necessary, renaming x and y , we can immediately deduce this identity from the identity (R) of $\mathcal{B}_{\text{hypo}}$.

The idea of the proof of the induction step is that, for any identity $u \approx v$, we can apply identities of $\mathcal{B}_{\text{hypo}}$, finitely many times, to deduce a new identity $u^* \approx v$ from $u \approx v$, such that u^* is ‘‘closer’’ to v than u , in the sense that u^* and v have a common prefix which is strictly longer than the common prefix of u and v . More

formally, by “deducing” $u^* \approx v$ from $u \approx v$, using identities of $\mathcal{B}_{\text{hypo}}$, we mean showing that $u \approx u^*$ is a consequence of $\mathcal{B}_{\text{hypo}}$.

The technical part of the proof allows us to show that there is always a way to shuffle some variables of u in such a way that we obtain u^* . We show that these variables must occur several times in u , thus allowing us to apply the identities of $\mathcal{B}_{\text{hypo}}$ to shuffle u and obtain u^* .

By the induction hypothesis, we know that $u^* \approx v$ is a consequence of $\mathcal{B}_{\text{hypo}}$. Thus, we conclude that $u \approx v$ is also a consequence of $\mathcal{B}_{\text{hypo}}$.

Let $u \approx v$ be a non-trivial identity satisfied by *hypo*. Let \mathcal{X} be the set of variables which occur in u and v . Since $u \approx v$ is a non-trivial identity, we must have $u = wxu'$ and $v = yv'$, for some words $w, u', v' \in \mathcal{X}^*$ and variables x, y such that $x \neq y$. Notice that u' and v' cannot be the empty word, otherwise, we would have $u = wx$ and $v = wy$, which contradicts the fact that $c(u) = c(v)$.

On the other hand, since $c(xu') = c(yv')$, we have that y occurs in u' . Thus, to distinguish the leftmost y in u' , we have that

$$xu' = u_1ayu_2,$$

for some variable a and words u_1 and u_2 , such that y does not occur in u_1 and $a \neq y$. Once again, since $c(xu') = c(yv')$, we have that a occurs in v' . Thus, to distinguish the leftmost a in v' , we have that

$$v' = v_1av_2,$$

for some words v_1 and v_2 , such that a does not occur in v_1 . To sum up, we have that

$$u = wu_1ayu_2 \quad \text{and} \quad v = wyv_1av_2,$$

where y does not occur in u_1 and a does not occur in v_1 .

Notice that u admits ay as a subsequence, hence, v must also do so. Thus, either a occurs in w , or y occurs in v_2 , since a does not occur in v_1 . But if y occurs in v_2 , then it must also occur in u_2 , since $c(xu') = c(yv')$ and y does not occur in u_1 .

On the other hand, notice that v admits ya as a subsequence, hence, u must also do so. Thus, either y occurs in w , or a occurs in u_2 , since y does not occur in u_1 .

As such, we have four possible cases to look at:

Case 1. Both variables y and a occur in w . Then, we can deduce the word wu_1yau_2 from u , by applying the identity (R), renaming x to a .

Case 2. Both variables y and a occur in u_2 . Then, we can deduce the word wu_1yau_2 from u , by applying the identity (L), renaming x to a .

Case 3. Variable y occurs in both w and u_2 . Then, we can deduce the word wu_1yau_2 from u , by applying the identity (M), renaming x to y and y to a .

Case 4. Variable a occurs in both w and u_2 . Then, we can deduce the word wu_1yau_2 from u , by applying the identity (M), renaming x to a .

Observe that we can repeatedly apply this reasoning until we obtain a word of the form

$$u^* = wyu'',$$

for some word u'' , since the only restriction imposed on the variable a was that $a \neq y$. Thus, we have proven that, for any non-trivial identity $u \approx v$ satisfied by *hypo*, we can obtain a new word u^* from u such that the common prefix of u^* and v is strictly longer than the common prefix of u and v , by applying identities of $\mathcal{B}_{\text{hypo}}$ finitely many times.

By the induction method, we conclude that $u \approx v$ is a consequence of $\mathcal{B}_{\text{hypo}}$, thus proving that $\mathcal{B}_{\text{hypo}}$ is a basis for \mathbf{V}_{hypo} . \square

An immediate consequence of having a finite basis is the following:

Corollary 4.9. \mathbf{V}_{hypo} has finite axiomatic rank.

Now, in order to determine the axiomatic rank of \mathbf{V}_{hypo} , we first check if all identities in $\mathcal{B}_{\text{hypo}}$ are necessary in order to obtain a basis for \mathbf{V}_{hypo} . It is easy to see that right-zero semigroups satisfy identities (L) and (M), but not the identity (R), and left-zero semigroups satisfy (R) and (M), but not (L). Thus, (L) and (R) are not consequences of the other identities in $\mathcal{B}_{\text{hypo}}$.

On the other hand, let $S = \{1, a, b, c, 0\}$ be a semigroup with multiplication table:

	1	a	b	c	0
1	1	a	b	c	0
a	a	1	b	c	0
b	b	c	0	0	0
c	c	b	0	0	0
0	0	0	0	0	0

This example was produced using Mace4 [32]. It consists of a null semigroup $K = \{b, c, 0\}$ with a cyclic group $C_2 = \{1, a\}$ acting on it identically from the left and by permuting b and c on the right. Notice that $a^2 = 1$, $ba = c = ac$ and $ca = b = ab$.

S satisfies (L) and (R): If we substitute all variables with elements of C_2 , we get equality by commutativity. If, after substituting, we have more than one element of K on each side, both sides equal 0. So we are left with the case of substituting a variable that appears once (either z or t) with an element of K and everything else with elements of C_2 . But since z and t are in the same position on both sides of the identities, and whichever elements of C_2 are substituted for x and y commute, the action of the elements of C_2 on the element of K gives the same result. However, S does not satisfy (M): Taking the evaluation ϕ such that $\phi(y) = c$ and $\phi(x) = \phi(z) = \phi(t) = a$, we get

$$\phi(xzxytx) = aaacaa = c \neq b = aacaaa = \phi(xzyxtx).$$

Since no identity of $\mathcal{B}_{\text{hypo}}$ is a consequence of the other identities also in $\mathcal{B}_{\text{hypo}}$, we conclude that $\mathcal{B}_{\text{hypo}}$ is minimal, in the sense that it does not contain any proper subset which is also a basis for \mathbf{V}_{hypo} .

We now show that the identities (L) and (R) are required to be in any basis for \mathbf{V}_{hypo} which contains only identities over an alphabet with four variables:

Proposition 4.10. *The identity (L) is not a consequence of the set of non-trivial identities, satisfied by hypo, over an alphabet with four variables, excluding (L) itself and equivalent identities.*

Proof. Let $X := \{x, y, z, t\}$ and let \mathcal{S} be the set of all non-trivial identities, satisfied by hypo, over an alphabet with four variables, excluding (L) and equivalent identities. Suppose, in order to obtain a contradiction, that (L) is a consequence of \mathcal{S} . As such, there must exist a non-trivial identity $u \approx v$ in \mathcal{S} , and a substitution σ , such that

$$xyzxty = w_1\sigma(u)w_2,$$

where w_1, w_2 are words over X , and $\sigma(u) \neq \sigma(v)$. Notice that $u \approx v$ must be balanced, and that there must be at least two variables occurring in u and v , otherwise, $u \approx v$ would be a trivial identity.

Observe that if the substitution σ maps some variables occurring in u and v to the empty word, then $\sigma(u) = \sigma(u')$ and $\sigma(v) = \sigma(v')$, where u' and v' are words obtained by eliminating every occurrence of such variables in u and v , respectively. Hence, we have that

$$xyzxty = w_1\sigma(u')w_2.$$

Notice that $u' \approx v'$ is an identity satisfied by *hypo*, which cannot be trivial, otherwise we would have $\sigma(u) = \sigma(u') = \sigma(v') = \sigma(v)$. Thus, $u' \approx v'$ is also in \mathcal{S} . On the other hand, notice that σ does not map any variable occurring in u' and v' to the empty word. As such, any case where σ maps any variable to the empty word guarantees the existence of another case where it does not map any variable to the empty word.

Therefore, we can assume, without loss of generality, that σ does not map any variable to the empty word. Due to this, and since (L) is an identity where x and y occur two times, and t and z each occur one time, we have that each variable occurring in $u \approx v$ can occur at most two times, and only two variables can occur more than one time. Furthermore, by Corollary 4.7, which gives us a lower bound for the length of the identities, we have that $u \approx v$ is of length at least 4. Thus, up to renaming of variables, x and y occur exactly twice in $u \approx v$, and t and z can occur at most one time.

Suppose now, in order to obtain a contradiction, that $w_1 \neq \varepsilon$. Then, since $u \approx v$ is of length at least 4, we must have w_1 of length at most 2, that is, w_1 is either x or xy . Therefore, x can occur only once in $\sigma(u)$. But x and y occur twice in u , and σ does not map any variable to the empty word, hence, there must be at least two variables which occur twice in $\sigma(u)$. However, only x and y occur twice in $xyzxty$. We have reached a contradiction, hence, $w_1 = \varepsilon$. Using a similar argument, we can also conclude that $w_2 = \varepsilon$. Therefore, we have that

$$xyzxty = \sigma(u).$$

As such, we can immediately conclude that only up to three variables occur in $u \approx v$: If $u \approx v$ were to be a four-variable identity, then it would be of length 6, and σ would be simply renaming the variables, thus implying that $u \approx v$ was equivalent to (L), which contradicts our hypothesis.

Suppose that $u \approx v$ is a two-variable identity. Hence, it is of length 4 and x and y both occur twice in it. Since $xyzxty = \sigma(u)$, then either $\sigma(x)$ or $\sigma(y)$ must be a single variable, and the other must be of length 2. Since no variable occurs more than twice in $\sigma(u)$, this implies that three variables occur twice in $\sigma(u)$. But x and y are the only variables which occur twice in $xyzxty$. We have reached a contradiction, hence, $u \approx v$ is not a two-variable identity.

Then, $u \approx v$ must be a three-variable identity. Hence, it is of length 5, with x and y occurring twice and z occurring once in it. Notice that $\sigma(x)$ and $\sigma(y)$ must be single variables, otherwise, the length of $\sigma(u)$ would be greater than 6. These variables cannot be z or t , since they occur only once in $xyzxty$. Therefore, $\sigma(z)$ must be a factor of $xyzxty$ of length 2. But neither x nor y can occur in $\sigma(z)$, hence, this factor cannot exist, and subsequently, σ cannot exist.

As such, we can conclude that (L) is not a consequence of the set of non-trivial identities, satisfied by *hypo*, over an alphabet with four variables, excluding (L) itself and equivalent identities. \square

Parallel reasoning shows the analogous result for (R):

Proposition 4.11. *The identity (R) is not a consequence of the set of non-trivial identities, satisfied by *hypo*, over an alphabet with four variables, excluding (R) itself and equivalent identities.*

Therefore, we can conclude that \mathbf{V}_{hypo} does not admit any basis with only identities over an alphabet with two or three variables. In other words, we have that:

Corollary 4.12. *The axiomatic rank of \mathbf{V}_{hypo} is 4.*

Another consequence of the previous proposition is the following:

Corollary 4.13. *Any basis for \mathbf{V}_{hypo} with only identities over an alphabet with four variables must contain the identities (L) and (R), or equivalent identities.*

Furthermore, since (M) is not a consequence of (L) and (R), any basis for \mathbf{V}_{hypo} with only identities over an alphabet with four variables must contain at least three identities, one of which must be either (M), an equivalent identity, or an identity of which (M) is a consequence of.

5. ACKNOWLEDGMENT

The authors thank the anonymous referee for their careful reading of the paper and many helpful comments, in particular, for the suggestion of Corollary 4.4.

REFERENCES

- [1] Sergei I. Adjan, 'Defining relations and algorithmic problems for groups and semigroups', *Tr. Mat. Inst. Steklov* **85**, Am. Math. Soc.(translation) **152** (1967).
- [2] Clifford Bergman, *Universal algebra: Fundamentals and selected topics*, Chapman & Hall Pure and Applied Mathematics, Taylor & Francis, 2011.
- [3] Garrett Birkhoff, *On the structure of abstract algebras*, *Mathematical Proceedings of the Cambridge Philosophical Society* **31** (1935), no. 4, 433—454.
- [4] Daniel Bump and Anne Schilling, *Crystal bases: Representations and combinatorics*, World Scientific Publishing Company Pte. Limited, 2017.
- [5] S. Burris and H.P. Sankappanavar, *A course in universal algebra*, Graduate Texts in Mathematics, Springer New York, 1981.
- [6] Alan J. Cain, Georg Klein, Łukasz Kubat, António Malheiro, and Jan Okniński, *A note on identities in plactic monoids and monoids of upper-triangular tropical matrices*, ArXiv e-prints (2017).
- [7] Alan J. Cain and António Malheiro, *Crystallizing the hypoplactic monoid: from quasi-Kashiwara operators to the Robinson–Schensted-type correspondence for quasi-ribbon tableaux*, *Journal of Algebraic Combinatorics* **45** (2017), 475–524.
- [8] Alan J. Cain and António Malheiro, *Identities in plactic, hypoplactic, sylvester, Baxter, and related monoids*, *Electronic Journal of Combinatorics* **25** (2018), no. 3.
- [9] Julien Cassaigne, Marc Espie, Daniel Krob, Jean-Christophe Novelli, and Florent Hivert, *The Chinese monoid*, *International Journal of Algebra and Computation* **11** (2001), no. 03, 301–334.
- [10] Laure Daviaud, Marianne Johnson, and Mark Kambites, *Identities in upper triangular tropical matrix semigroups and the bicyclic monoid*, *Journal of Algebra* **501** (2018), 503–525.
- [11] Gérard Duchamp and Daniel Krob, *Plactic-growth-like monoids*, *Words, languages and combinatorics II. Proceedings of the 2nd international conference, Kyoto, Japan, August 25-28, 1992*, Singapore: World Scientific, 1994, pp. 124–142.

- [12] Samuel Eilenberg, *Automata, languages, and machines, volume B*, Pure and applied mathematics, Academic Press, 1976.
- [13] Trevor Evans, *Word problems*, Bull. Amer. Math. Soc. **84** (1978), no. 5, 789–802.
- [14] William Fulton, *Young tableaux: With applications to representation theory and geometry*, London Mathematical Society Student Texts, Cambridge University Press, 1996.
- [15] Samuele Giraudo, *Algebraic and combinatorial structures on pairs of twin binary trees*, Journal of Algebra **360** (2012), 115–157.
- [16] Mikhael Gromov, *Groups of polynomial growth and expanding maps (with an appendix by Jacques Tits)*, Publications Mathématiques de l’IHÉS **53** (1981), 53–78.
- [17] Peter M. Higgins, *Techniques of semigroup theory*, Oxford University Press, 1992.
- [18] Florent Hivert, Jean-Christophe Novelli, and Jean-Yves Thibon, *The algebra of binary search trees*, Theoretical Computer Science **339** (2005), no. 1, 129–165.
- [19] John M. Howie, *Fundamentals of semigroup theory*, LMS monographs, Clarendon Press, 1995.
- [20] Zur Izhakian, *Tropical plactic algebra, the cloaktic monoid, and semigroup representations*, Journal of Algebra **524** (2019), 290–366.
- [21] Joanna Jaszusńska and Jan Okniński, *Structure of Chinese algebras*, Journal of Algebra **346** (2011), no. 1, 31–81.
- [22] Marianne Johnson and Mark Kambites, *Tropical representations and identities of plactic monoids*, arXiv preprint arXiv:1906.03991 (2019).
- [23] Donald E. Knuth, *Permutations, matrices, and generalized Young tableaux.*, Pacific J. Math. **34** (1970), no. 3, 709–727.
- [24] Daniel Krob and Jean-Yves Thibon, *Noncommutative symmetric functions IV: Quantum linear groups and Hecke algebras at $q = 0$* , Journal of Algebraic Combinatorics **6** (1997), no. 4, 339–376.
- [25] ———, *Noncommutative symmetric functions V: A degenerate version of $u_q(gl_N)$* , International Journal of Algebra and Computation **09** (1999), no. 03–04, 405–430.
- [26] Łukasz Kubat and Jan Okniński, *Identities of the plactic monoid*, Semigroup Forum **90** (2015), no. 1, 100–112.
- [27] Alain Lascoux and Marcel P. Schützenberger, *Le monoïde plaxique*, Noncommutative structures in algebra and geometric combinatorics (Rome), Quaderni de "La Ricerca Scientifica", no. 109, CNR, 1981, pp. 129–156.
- [28] M. Lothaire, *Algebraic combinatorics on words*, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2002.
- [29] Ian G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford Classic Texts in the Physical Sciences, Clarendon Press, 1998.
- [30] Diane Maclagan and Bernd Sturmfels, *Introduction to Tropical Geometry*, Graduate Studies in Mathematics, no. 161, American Mathematical Society, 2015.
- [31] Anatoly I. Mal’cev, *Algebraic systems*, Grundlehren der mathematischen Wissenschaften, Springer Berlin Heidelberg, 2012.
- [32] William McCune, *Prover9 and Mace4*, <http://www.cs.unm.edu/~mccune/prover9/>, 2005–2010.
- [33] Ralph N. McKenzie, George F. McNulty, and Walter F. Taylor, *Algebras, lattices, varieties*, AMS Chelsea Publishing, American Mathematical Society, 2018.
- [34] Jean-Christophe Novelli, *On the hypoplactic monoid*, Discrete Mathematics **217** (2000), no. 1, 315–336.
- [35] Francis Pastijn, *Polyhedral convex cones and the equational theory of the bicyclic semigroup*, Journal of the Australian Mathematical Society **81** (2006), no. 1, 63–96.
- [36] Jean-Éric Pin, *Varieties of formal languages*, Foundations of Computer Science, North Oxford Academic, 1986.
- [37] Mark V. Sapir, *Combinatorial algebra: Syntax and semantics*, Springer Monographs in Mathematics, Springer International Publishing, 2014.
- [38] Craige Schensted, *Longest increasing and decreasing subsequences*, Canad. J. Math. **13** (1961), 179–191.
- [39] Lev M. Shneerson, *On the axiomatic rank of varieties generated by a semigroup or monoid with one defining relation*, Semigroup Forum **39** (1989), no. 1, 17–38.
- [40] Lev M. Shneerson, *Identities in finitely generated semigroups of polynomial growth*, Journal of Algebra **154** (1993), no. 1, 67–85.

- [41] Imre Simon, *Hierarchies of events with dot-depth one*, Ph.D. thesis, University of Waterloo, Waterloo, Ont., Canada, 1972.
- [42] Mikhail V. Volkov, *The finite basis problem for finite semigroups*, Sci. Math. Jpn **53** (2001), no. 1, 171–199.
- [43] ———, *Reflexive relations, extensive transformations and piecewise testable languages of a given height*, International Journal of Algebra and Computation **14** (2004), no. 05–06, 817–827.

CENTRO DE MATEMÁTICA E APLICAÇÕES, FACULDADE DE CIÊNCIAS E TECNOLOGIA, UNIVERSIDADE NOVA DE LISBOA, 2829–516 CAPARICA, PORTUGAL
Email address: `a.cain@fct.unl.pt`

DEPARTAMENTO DE MATEMÁTICA & CENTRO DE MATEMÁTICA E APLICAÇÕES, FACULDADE DE CIÊNCIAS E TECNOLOGIA, UNIVERSIDADE NOVA DE LISBOA, 2829–516 CAPARICA, PORTUGAL
Email address: `ajm@fct.unl.pt`

DEPARTAMENTO DE MATEMÁTICA & CENTRO DE MATEMÁTICA E APLICAÇÕES, FACULDADE DE CIÊNCIAS E TECNOLOGIA, UNIVERSIDADE NOVA DE LISBOA, 2829–516 CAPARICA, PORTUGAL
Email address: `dc.ribeiro@campus.fct.unl.pt`