

**ON THE MAXIMUM PRINCIPLES AND THE QUANTITATIVE VERSION
OF THE HOPF LEMMA FOR UNIFORMLY ELLIPTIC
INTEGRO-DIFFERENTIAL OPERATORS**

TOMASZ KLIMSIK AND TOMASZ KOMOROWSKI

ABSTRACT. In the present paper we prove estimates on subsolutions of the equation $-Av + c(x)v = 0$, $x \in D$, where $D \subset \mathbb{R}^d$ is a domain (i.e. an open and connected set) and A is an integro-differential operator of the Waldenfel's type, whose differential part satisfies the uniform ellipticity condition on compact sets. In general, the coefficients of the operator need not be continuous but only bounded and Borel measurable. Some of our results may be considered "quantitative" versions of the Hopf lemma, as they provide the lower bound on the outward normal directional derivative at the maximum point of a subsolution in terms of its value at the point. We shall also show lower bounds on the subsolution around its maximum point by the principal eigenfunction associated with A and the domain. Additional results, among them the weak and strong maximum principles, the weak Harnack inequality are also proven.

1. INTRODUCTION

The maximum principle and the Hopf lemma express some of the most fundamental properties of solutions of elliptic partial differential equations. Consider a differential operator

$$Lu(x) := \frac{1}{2} \sum_{i,j=1}^d q_{i,j}(x) \partial_{x_i x_j}^2 u(x) + \sum_{i=1}^d b_i(x) \partial_{x_i} u(x), \quad u \in C^2(\mathbb{R}^d) \quad (1.1)$$

that is *uniformly elliptic* on compact sets, i.e. the coefficients $q_{i,j}$, b_j , $i, j = 1, \dots, d$ are bounded, Borel measurable and the matrix $\mathbf{Q}(x) = [q_{i,j}(x)]_{i,j=1}^d$ is *uniformly positive definite* on compact sets, see (2.3) below.

The strong maximum principle, see e.g. [19, Theorem 9.6, p. 225], can be formulated as follows. Suppose that D is a domain (an open and connected set) and c is a non-negative, bounded and measurable function on D . Assume furthermore that a continuous function u , belonging to $W_{\text{loc}}^{2,d}(D)$ (i.e. its two generalized derivatives are L^d -integrable on compacts) and satisfying

$$-Lu(x) + c(x)u(x) \leq 0, \quad x \in D \quad (1.2)$$

admits a maximum $M \geq 0$ in D . Then, u has to be constant, i.e. $u(x) = M$, $x \in D$. A function satisfying (1.2) is called a *subsolution* of the equation $(-L + c)f = 0$ in D .

If ∂D - the boundary of the domain - is sufficiently regular, say C^2 smooth and u is non-constant, then the *Hopf lemma*, see e.g. [27, Theorem 7, p. 65], asserts that $\partial_{\mathbf{n}}u(\hat{x})$ - the directional derivative of u in the outward normal to ∂D - at the maximal point \hat{x} has to be strictly positive.

It has been shown in [7] that if ∂D is C^2 smooth and $\underline{c} := \inf_{x \in \bar{D}} c(x) > 0$, then there exists a constant $a > 0$, depending only on the ellipticity constant on \bar{D} , \underline{c} , the L^∞ bound on the coefficients of L and c , and D such that

$$\partial_{\mathbf{n}}u(\hat{x}) > au(\hat{x}) \quad (1.3)$$

Mathematics Subject Classification: Primary 35B50; Secondary 35J15, 35J08

Keywords: Integro-differential elliptic operator, maximum principle, the Hopf lemma, principal eigenvalue and eigenvector

for any u satisfying (1.2) that is not identically equal to 0. The above estimate could be viewed as a form of a *quantitative Hopf lemma*. An elementary example, using harmonic functions, shows that (1.3) is false if $c \equiv 0$.

One of the objectives of the present paper is to investigate an analogue of the estimate (1.3) and its generalizations for non-local operators. More precisely, we consider non-divergence form integro-differential operators

$$Au(x) = Lu(x) + Su(x), \quad u \in C^2(\mathbb{R}^d) \cap C_b(\mathbb{R}^d), \quad x \in \mathbb{R}^d, \quad (1.4)$$

where L is given by (1.1) and

$$Su(x) := \int_{\mathbb{R}^d} \left(u(x+y) - u(x) - \sum_{i=1}^d \frac{y_i \partial_{x_i} u(x)}{1+|y|^2} \right) N(x, dy), \quad x \in \mathbb{R}^d. \quad (1.5)$$

The coefficients of the operator L are as in the foregoing and the kernel N (sometimes referred to as the Lévy kernel) assigns to each $x \in \mathbb{R}^d$ a σ -finite Borel measure $N(x, \cdot)$, which satisfies

$$N_* := \sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d} \min\{1, |y|^2\} N(x, dy) < +\infty. \quad (1.6)$$

Such operators are sometimes called of the *Waldenfels type*, see e.g. [30, Chapter 10]. A sufficiently regular function $u : D \rightarrow \mathbb{R}$, see Section 2.3 for a rigorous definition, is called a *subsolution* (resp. *supersolution*) of equation

$$(-A + c)f = 0. \quad (1.7)$$

if $-Au(x) + c(x)u(x) \leq 0$ (resp. $-Au(x) + c(x)u(x) \geq 0$) in D . Throughout the paper we shall always assume that $c : D \rightarrow \mathbb{R}$ is nonnegative, bounded and Borel measurable.

Let

$$\mathcal{S}(D) := \bigcup_{x \in D} \mathcal{S}_x, \quad \text{where } \mathcal{S}_x := [x + y : y \in \text{supp } N(x, \cdot)]. \quad (1.8)$$

The set \mathcal{S}_x describes the "range of non-locality" of A at x , i.e. the evaluation of $Au(x)$ depends on the values of u in an arbitrarily small neighborhood of x , and in \mathcal{S}_x .

Then, the following version of the "quantitative Hopf lemma" holds, see Theorem 3.5 below for a precise statement.

Theorem 1.1. *Suppose that $\underline{c} > 0$, the domain D is bounded and satisfies the uniform interior ball condition (see Definition 2.2), u - a subsolution of (1.7) - attains its maximum over $\mathcal{S}(D) \cup \bar{D}$ at some $\hat{x} \in \partial D$ and is not identically equal to 0. Then, there exists a $a > 0$ that depends only on D , ellipticity constant, \underline{c} , the L^∞ norm of the coefficients and N_* such that estimate (1.3) holds*

We can relax the assumption that $\underline{c} > 0$. Namely, see Theorem 3.6 below, we admit that $\underline{c} = 0$ but $c(x)$ does not vanish a.e. in D . However, in that case we have to allow $a > 0$ in (1.3) to depend also on the operator A (in particular on the lower bound of its Green function), not just on some characteristics of its coefficients as in Theorem 1.1.

In our further generalizations of estimate (1.3), see Theorems 5.1 and 5.2 below, we compare the increment $u(\hat{x}) - u(x)$ with the value of the principal eigenfunction $\varphi_D(x)$ associated with the operator A on the domain D , i.e. the unique normalized eigenfunction that is strictly positive in D , see Theorem 4.7. It turns out (see Theorem 5.2) that, with no assumption on the regularity of D , the following bound holds

$$u(\hat{x}) - u(x) \geq \frac{\varphi_D(x)}{2e\|\varphi_D\|_\infty} \left[\frac{\underline{c}u(\hat{x})}{\lambda_D + \underline{c}} + \frac{\text{essinf}_D(Au - cu)}{\lambda_D + \|c\|_\infty} \right], \quad x \in D. \quad (1.9)$$

Here λ_D is the principal eigenvalue corresponding to φ_D and $\|\cdot\|_\infty$ is the usual supremum norm.

In fact, instead of the essential infimum of $Au - cu$ in (1.9), we can use its average over D with respect to some measure equivalent with the Lebesgue measure. Namely, see Theorem 5.1,

there exist $\psi(x)$ continuous, strictly positive in D and $\chi(x)$ strictly positive, Borel measurable such that any subsolution u , described in the foregoing, satisfies

$$u(\hat{x}) - u(x) \geq \frac{c\varphi_D(x)u(\hat{x})}{2e\|\varphi_D\|_\infty(\lambda_D + c)} + \psi(x) \int_D (Au - cu)\chi dx, \quad x \in D. \quad (1.10)$$

We have already mentioned that estimates (1.9) and (1.10) are obtained without making assumptions on the regularity of the domain D . However, if we suppose that both the uniform interior and exterior ball conditions hold, see Definitions 2.2 and 2.5, then we can choose $\psi(x) = \delta_D(x)$, where $\delta_D(x) := \text{dist}(x, D^c)$, see Theorem 5.4. Furthermore, if some regularity of the coefficients of A is allowed, then, in addition to the above choice of $\psi(x)$, we can admit also $\chi(x) = a\delta_D(x)$, with some $a > 0$, see Theorems 5.6 and 5.8.

It is also worthwhile to mention that the estimates (1.9) and (1.10) hold even without the uniform ellipticity hypothesis. They are related to ergodic properties of the Markov process, associated with the operator A via the respective martingale problem, see Section 4 below, e.g. its uniform conditional ergodicity, or intrinsic ultracontractivity of its transition semigroup. This shall be the topic of our forthcoming paper [21].

In addition to the results mentioned above, in the present paper we also obtain a weak maximum principle for operators of the form (1.4), see Proposition 3.1. Namely, if u is a subsolution of (1.7) in D , then

$$\sup_{x \in D} u(x) \leq \sup_{y \in (S(D) \setminus D) \cup \partial D} u^+(y). \quad (1.11)$$

This result holds without assuming that D is connected. Using the weak form of the maximum principle we establish its strong version and the Hopf lemma, see Theorem 3.4. In addition we prove also the Bony type maximum principle and a form of the weak Harnack inequality, see Theorems 3.3 and 5.9, respectively. The latter asserts the existence of a strictly positive, bounded, Borel measurable function χ on D such that for any open V , with \bar{V} compact and $\bar{V} \subset D$, there exists a constant $C > 0$ such that for any u - a non-negative supersolution to (1.7) - we have

$$\inf_{x \in V} u(x) \geq C \int_V \chi u dx. \quad (1.12)$$

In the case when D is regular, e.g. satisfies both the interior and exterior ball conditions (see Definitions 2.2 and 2.5) and the formal adjoint to A satisfies appropriate hypotheses, see Section 5.2, one can take $\chi(x) = \delta_D(x)$, $x \in D$, see Remarks 9.3 and 9.4 below, and conclude the "usual" weak Harnack inequality, see e.g. [19, Theorem 8.18, p. 194, or Theorem 9.22, p.246],

$$\inf_{x \in V} u(x) \geq C \int_V u dx. \quad (1.13)$$

Recall that the (strong) Harnack inequality, asserts the existence of $C > 0$, for which $\inf_{x \in V} u(x) \geq C \sup_{x \in V} u(x)$ for all non-negative supersolutions. This type of inequality need not hold in general for operators of the type (1.4) and for its validity some additional restrictions on the kernel N are needed, see [18, Section 7] for more details.

Let us briefly discuss the relation between our work and some results existing in the literature. The maximum principles both weak and strong, together with the Hopf lemma can be found e.g. in [9, Section I.4], [30, Section 10.2] and [33, Appendix C]. However, the versions of the maximum principles we are aware of do not cover the case considered in our paper. For example, the formulations of [30, 33] concern only the case when coefficients are C^∞ smooth, while [9] requires some regularity assumptions on the Lévy kernel, see Section I.2.3 and condition (I.4.1) *ibid.* The closest formulation to part 1) of our Theorem 3.4, we have encountered in the recent paper of Taira [32]. In *ibid.* the author has additionally assumed that $\mathcal{S}_x(D) = \bar{D}$ for $x \in D$ (no jumps outside \bar{D}) and that there exists a reference measure, with respect to which, all the Lévy kernels are absolutely continuous. The regularity of the resulting densities is also assumed. The domain D is to be of class $C^{1,1}$ and $u \in W^{2,p}(D)$ for some $p > d$.

Our formulation of the Hopf lemma, see part 2) of Theorem 3.4 and in particular Corollary 3.11, resembles somewhat Proposition 1.34 of [20], which actually uses the phrase "quantitative

Hopf lemma". The result of *ibid.* holds for harmonic functions on a ball B and asserts that there exists $a > 0$ such that for any bounded harmonic function u in B , we have

$$\partial_{\mathbf{n}}u(\hat{x}) \geq a(u(\hat{x}) - u(x_0)) \quad (1.14)$$

with $\hat{x} \in \partial B$ a maximum point of u in \bar{B} and x_0 - the center of the ball. The counterpart of this estimate for A -harmonic functions, provided that the operator has the Harnack property (see Definition 3.9), is stated in (3.8) below.

Estimate (1.10) of Theorem 5.1 can be compared with the Morel and Oswald version of the Hopf lemma proved by Brezis and Cabré in [11], see Lemma 3.2. It asserts that for a smooth and bounded domain D there exists $a > 0$ such that for any superharmonic function u , vanishing on the boundary, we have

$$u(x) \geq a\delta_D(x) \int_D (-\Delta u)(y)\delta_D(y) dy, \quad x \in D. \quad (1.15)$$

The estimate is sometimes referred to as the *uniform Hopf lemma* (see e.g. [16, 14, 26]).

We prove an analogue of (1.15), see Theorem 5.8, for C^2 -regular domains, $u(x)$ - a subsolution of (1.7) - and operator A of the form (1.4), whose coefficients of the second derivative, as well as the respective coefficients of its formal adjoint belong to the VMO class, see Definition 5.7. From (1.15) one can easily conclude the *weak Harnack inequality* (see e.g. [19, Theorem 9.22]) that is shown in the present paper in Theorem 5.9.

The principal tools used in our analysis are: the martingale problem associated with operator A , see Section 4.1, and the Feynman-Kac formula for the respective canonical process $(X_t)_{t \geq 0}$, see Proposition 4.6. They allow us to obtain a probabilistic representation for a subsolution of the equation (1.7), see Definition 2.1, in terms of the expectation of an appropriate functional of the process, see (4.10). We exploit various properties the process $(X_t)_{t \geq 0}$, such as its irreducibility and the strong Feller property, to derive our results.

The body of the paper is essentially divided into two parts. The first one, consisting of Sections 2 to 5, besides presenting some introductory material, is primarily devoted to the exposition of the results contained in the article. Their proofs have been placed in the second part, containing Sections 6 through 9. In more detail, in Section 2 we introduce the basic notation and definitions. The formulations of the main results are contained in Sections 3 and 5. In Section 4 we introduce some basic probabilistic tools used throughout the paper, such as: the already mentioned martingale problem associated with A , the (probabilistic) semigroup and resolvent corresponding to the canonical process, the gauge functional and the principal eigenvalue and eigenfunction associated with A and domain D . They are used in formulations of the results of Section 5. In Section 6 we present the proof of the Feynman-Kac formula, which is our basic tool in showing of virtually all the results of the paper. We have also placed in that section the proof of the weak maximum principle (Proposition 3.1) as it almost directly follows from the aforementioned formula. Section 7 discusses various properties of the resolvent that will be useful in our further arguments. Section 8 contains auxiliary facts and notions used in the proofs of our main results. Finally, the main results are proved in Section 9.

2. PRELIMINARIES

2.1. Basic notation. Given a metric space E we denote by $\mathcal{B}(E)$ its Borel σ -algebra. Let $B_b(E)$ ($B_b^+(E)$) be the space of all (non-negative) bounded Borel measurable functions and let $C_b(E)$ ($C_c(E)$) be the space of all bounded continuous (compactly supported) functions on E . Furthermore by $\mathcal{M}(E)$ we denote the set of all Borel measures on E . Suppose that $\mu, \nu \in \mathcal{M}(E)$. We say that μ dominates ν and write $\nu \ll \mu$ if all null sets for μ are also null for ν . We say that measures are equivalent and write $\mu \sim \nu$ iff $\mu \ll \nu$ and $\nu \ll \mu$.

Given a point $x \in E$ and $r > 0$ we let $B(x, r)$ be the open ball of radius r centered at x and $\bar{B}(x, r)$ be its closure. As it is customary for a given function $f : E \rightarrow \mathbb{R}$ we denote $\|f\|_\infty = \sup_{x \in E} |f(x)|$. For a subset $B \subset E$ we let $B^c := E \setminus B$ and \bar{B} be its closure.

Suppose that B is an arbitrary set. For functions $f, g : B \rightarrow [0, +\infty)$ we write $f \preceq g$ on B if there exists number $C > 0$, i.e., *constant*, such that

$$f(x) \leq Cg(x), \quad x \in B.$$

Furthermore, we write $f \approx g$ if both $f \preceq g$ and $g \preceq f$.

Assume now that $D \subset \mathbb{R}^d$ and $V \subset D$ are open. We shall write $V \Subset D$ if \bar{V} is compact and $\bar{V} \subset D$. Furthermore we let $C^m(D)$, $m \geq 1$ be the class of m -times continuously differentiable functions in D . By $C_0(D)$ we denote the subset of $C(D)$ that consists of functions extending continuously to \bar{D} by letting $f(x) \equiv 0$, $x \in \partial D$ - the boundary of D .

Denote by m_d the d -dimensional Lebesgue measure on \mathbb{R}^d . Sometimes we omit writing the subscript, when the dimension is obvious from the context. By dx we denote the volume element corresponding to the Lebesgue measure. A function $f \in B^+(D)$ is called non-trivial if $\int_D f dx > 0$.

For $p \in [1, +\infty)$ we denote by $L^p(D)$ ($L^p_{\text{loc}}(D)$) the space of functions that are integrable with their p -th power on D (any compact subset of D). By $W^{k,p}(D)$ ($W^{k,p}_{\text{loc}}(D)$) we denote the Sobolev space of functions whose k generalized derivatives belong to $L^p(D)$ ($L^p_{\text{loc}}(D)$). Given $f \in L^p(D)$ and $B \in \mathcal{B}(D)$ we denote by $\text{ess inf}_B f$ the usual essential infimum of f in B . We define the *essential limit inferior* of f at $x_0 \in D$ as

$$\liminf_{x \rightarrow x_0} \text{ess } f(x) := \lim_{r \rightarrow 0^+} \text{ess inf}_{B(x_0, r) \subset D} f. \quad (2.1)$$

2.2. Second-order, elliptic integro-differential operators. Consider an elliptic integro-differential operator of the form

$$Au(x) = Lu(x) + Su(x), \quad u \in C^2(\mathbb{R}^d) \cap C_b(\mathbb{R}^d), \quad x \in \mathbb{R}^d, \quad (2.2)$$

where L , S are a differential and integro-differential operators, given by (1.1) and (1.5) respectively. Throughout the paper we shall always assume that the coefficients satisfy the following hypotheses:

- A1) $q_{i,j} \in B_b(\mathbb{R}^d)$, $b_j \in B_b(\mathbb{R}^d)$, $i, j = 1, \dots, d$ and we suppose that the differential operator L is *uniformly elliptic* on compact sets, i.e. the matrix $\mathbf{Q}(x) = [q_{i,j}(x)]_{i,j=1}^d$ is *uniformly positive definite* on compact sets. The latter means that for any compact set $K \subset \mathbb{R}^d$ there exists $\gamma_K > 0$ such that

$$\gamma_K |\xi|^2 \leq \sum_{i,j=1}^d q_{i,j}(x) \xi_i \xi_j, \quad x \in K, \quad \xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d. \quad (2.3)$$

- A2) $N : \mathbb{R}^d \rightarrow \mathcal{M}(\mathbb{R}^d)$, called the *Lévy kernel*, is a σ -finite Borel measure valued function such that, the mapping $x \mapsto N(x, A)$ is Borel measurable for each $A \in \mathcal{B}(\mathbb{R}^d)$ and (1.6) holds.

We let

$$M_A := \sum_{i,j=1}^d \|q_{i,j}\|_{\infty} + \sum_{i=1}^d \|b_i\|_{\infty} + N_* \quad (2.4)$$

Suppose that D is an open set. Obviously in order to define $Au(x)$ for $x \in D$ it suffices to assume that $u \in C^2(D) \cap C_b(\mathbb{R}^d)$. In fact, Au can be defined as an element of $L^p(D)$, using (1.4)–(1.5), even if $u \in W^{2,p}_{\text{loc}}(D) \cap C_b(\mathbb{R}^d)$ when $p > d$. This is possible due to a well known estimate, see [8, Lemme 1, p. 361]: for any $p > d$ there exists $C > 0$ such that

$$\|U[v]\|_{L^p(\mathbb{R}^d)} \leq C \|\nabla^2 v\|_{L^p(\mathbb{R}^d)}, \quad v \in W^{2,p}(\mathbb{R}^d).$$

Here

$$U[v](x) := \sup_{|y|>0} |y|^{-2} \left| v(x+y) - v(x) - \sum_{i=1}^d y_i \partial_{x_i} v(x) \right|.$$

2.3. A subsolution, supersolution and solution of an elliptic type integro-differential equation.

Definition 2.1. Suppose that $c, g : \mathbb{R}^d \rightarrow \mathbb{R}$. A function $u : \mathbb{R}^d \rightarrow \mathbb{R}$ is called a *subsolution* of the equation

$$(-A + c)f = g \quad (2.5)$$

on an open set $D \subset \mathbb{R}^d$ if $u \in W_{\text{loc}}^{2,p}(D) \cap C_b(\mathbb{R}^d)$ for some $p > d$ and

$$-Au(x) + c(x)u(x) \leq g(x), \quad \text{for a.e. } x \in D. \quad (2.6)$$

We say that u is a *supersolution* of equation (1.7) if $-u$ is a subsolution. Furthermore, u is a *solution* if it is both sub- and supersolution.

Throughout the paper, we shall mostly deal with homogeneous equations, i.e. the case when $g = 0$.

Given a bounded and measurable function $c : D \rightarrow \mathbb{R}$ we let

$$\bar{c} := \sup_{x \in D} c(x), \quad \underline{c} := \inf_{x \in D} c(x) \quad \text{and} \quad \langle c \rangle_D := \int_D c \, dx. \quad (2.7)$$

Concerning the coefficient $c(\cdot)$ it is assumed to satisfy $c \in B_b(D)$ and either one of the following hypotheses:

A3) it is non-negative on D , or

A3') it is both non-negative on D and $\langle c \rangle_D > 0$, (i.e. $c \not\equiv 0$ a.e.), or

A3'') $\underline{c} > 0$.

Below we formulate some hypotheses concerning an open set D that shall be used in various results presented in the paper. Throughout the paper, unless stated otherwise (notably in Proposition 3.1) it is supposed to be a *domain*, i.e. a bounded, open and connected subset of \mathbb{R}^d .

Definition 2.2 (Interior ball condition). For a given domain D and $\hat{x} \in \partial D$, we say that the *interior ball condition* is satisfied at \hat{x} , if there exists B - an open ball of a positive radius - contained in D and such that $\hat{x} \in \partial D \cap \bar{B}$. Any ball B as above shall be called *an interior ball* for D at \hat{x} . We furthermore say that D satisfies the *interior ball condition* if the condition is satisfied at any point of its boundary. We let $\mathfrak{r}_D(\hat{x})$ denote the maximal radius, that is less than, or equal to 1, of an interior ball at the given $\hat{x} \in \partial D$. We say that D satisfies the *uniform interior ball condition* if it satisfies the interior ball condition and

$$\text{there exists } r_D \in (0, 1) \text{ such that } \mathfrak{r}_D(\hat{x}) \geq r_D, \text{ for all } \hat{x} \in \partial D. \quad (2.8)$$

Suppose D satisfies the interior ball condition at $\hat{x} \in \partial D$ and $B(y, r) \subset D$, with $r > 0$, is an interior ball at \hat{x} . Any vector of the form

$$\mathbf{n} := \frac{1}{r}(\hat{x} - y) \quad (2.9)$$

shall be called a *generalized unit outward normal vector* to ∂D at \hat{x} . Denote by $\mathfrak{n}(\hat{x})$ the set of all such vectors at $\hat{x} \in \partial D$.

Definition 2.3 (Lower outward normal derivative at a boundary point). Suppose that D satisfies the interior ball condition at $\hat{x} \in D$. Assume furthermore that $f : \mathbb{R}^d \rightarrow \mathbb{R}$. For any $\mathbf{n} \in \mathfrak{n}(\hat{x})$ define the *lower outward normal derivative* at \hat{x} as

$$\underline{\partial}_{\mathbf{n}} f(\hat{x}) := \liminf_{\substack{h \rightarrow 0+ \\ \hat{x} - h\mathbf{n} \in D}} \frac{f(\hat{x}) - f(\hat{x} - h\mathbf{n})}{h}. \quad (2.10)$$

Remark 2.4. If ∂D is C^2 smooth, then for any $\hat{x} \in \partial D$ the set $\mathfrak{n}(\hat{x})$ is a singleton and consists only of the unit outward normal vector to ∂D at \hat{x} . In addition, if $f : \mathbb{R}^d \rightarrow \mathbb{R}$ possesses a derivative at $\hat{x} \in \partial D$, then $\underline{\partial}_{\mathbf{n}} f = \partial_{\mathbf{n}} f$ - the outward normal derivative.

In some cases, see e.g. Theorem 5.4 below, we shall assume that the domain satisfies the exterior ball condition in the following sense.

Definition 2.5. For a given domain D and $\hat{x} \in \partial D$, we say that the *exterior ball condition* is satisfied at \hat{x} , if there exists B - an open ball of a positive radius - contained in D^c and such that $\hat{x} \in \partial D \cap \bar{B}$. Any ball B as above shall be called *an exterior ball* for D at \hat{x} . We furthermore say that D satisfies the *exterior ball condition* if the condition is satisfied at any point of its boundary.

3. MAXIMUM PRINCIPLES AND QUANTITATIVE HOPF LEMMAS

The present section is devoted to the formulation of the first part of the set of our main results. Their proofs are presented throughout Section 9.

3.1. Maximum principles and the Hopf lemma. Our first result is a form of a weak maximum principle.

Proposition 3.1. *[Weak maximum principle] Suppose that the function c satisfies A3). If u is a subsolution to (1.7) on a bounded, open set $D \subset \mathbb{R}^d$ then (cf (1.8))*

$$\sup_{x \in D} u(x) \leq \sup_{y \in (\mathcal{S}(D) \setminus D) \cup \partial D} u^+(y). \quad (3.1)$$

Moreover, if $c \equiv 0$, then u^+ in the right-hand side can be replaced by u .

The proof of the proposition is shown in Section 6.2.

Remark 3.2. Observe that the right-hand side of (3.1) reduces to $\sup_{y \in D^c} u^+(y)$ in the case $\mathcal{S}(D) = \mathbb{R}^d$ and to $\sup_{y \in \partial D} u^+(y)$ when $\mathcal{S}(D) = \overline{D}$.

Theorem 3.3 (Bony maximum principle). *Let D be an open subset of \mathbb{R}^d and $\mathcal{S}(D) \subset \overline{D}$. Suppose furthermore that $u \in W_{loc}^{2,p}(D) \cap C_b(\mathbb{R}^d)$ for some $p > d$ and $\hat{x} \in D$ is its maximum point in D . Then, cf (2.1),*

$$\liminf_{x \rightarrow \hat{x}} \text{ess } Au(x) \leq 0. \quad (3.2)$$

The proof of the theorem is presented in Section 9.6.

In an analogy to the case of elliptic differential operators, we have the strong maximum principle and Hopf lemma for a subsolution of (1.7). The formulation of the latter, given below, is of a bit more "quantitative" nature than usually encountered in the literature. To state it we introduce some notation. For $r > 0$ we let

$$D_r := [x \in D : \text{dist}(x, \partial D) > r]. \quad (3.3)$$

We also let $\mathcal{O}_D := [x \in \mathbb{R}^d : \text{dist}(x, D) < 1]$.

Theorem 3.4. *Suppose that the function c satisfies A3).*

- 1) *(The strong maximum principle) If u is a subsolution to (1.7) on a domain D that attains its maximum $M \geq 0$ over $\mathcal{S}(D) \cup \overline{D}$ at $\hat{x} \in D$, then $u \equiv M$ in D .*
- 2) *(The Hopf lemma) Assume that a domain D is bounded and satisfies the uniform interior ball condition, with constant $r_D \in (0, 1]$. Then, there exist $r \in (0, r_D/2]$, $a > 0$ depending only on M_A and $\gamma_{\overline{\mathcal{O}}_D}$ such that for any non-constant subsolution u to (1.7) that attains its non-negative maximum over $\mathcal{S}(D) \cup \overline{D}$ at $\hat{x} \in \partial D$, and any $\mathbf{n} \in \mathbf{n}(\hat{x})$ at $\hat{x} \in \partial D$ we have*

$$\partial_{\mathbf{n}} u(\hat{x}) \geq a \inf_{x \in D_r} (u(\hat{x}) - u(x)) > 0. \quad (3.4)$$

The proof of the theorem is given in Section 9.1.

3.2. Quantitative Hopf lemmas. Suppose that u is a subsolution of (1.7). If c satisfies either the hypothesis A3'), or A3''), then we can formulate a lower bound on the outward normal partial derivative of u at the maximum point in terms of its maximal value.

Theorem 3.5 (Quantitative Hopf Lemma I.A). *Suppose that function c satisfies hypothesis A3''). In addition, assume that a domain D is bounded and satisfies the uniform interior ball condition. Then, there exists $a > 0$ that depends only on r_D of (2.8), ellipticity constant $\gamma_{\overline{\mathcal{O}}_D}$, the lower bound \underline{c} and $M_A + \|c\|_{\infty}$ (cf (2.4)) such that for any u - a subsolution to (1.7) - that attains its maximum over $\mathcal{S}(D) \cup \overline{D}$ at $\hat{x} \in \partial D$ and is not identically equal to 0 we have*

$$\partial_{\mathbf{n}} u(\hat{x}) > au(\hat{x}). \quad (3.5)$$

The proof of the theorem is presented in Section 9.3.

A version of the result in the case when c is only assumed to be non-negative and not equal to 0 a.e. can be stated as follows.

Theorem 3.6 (Quantitative Hopf lemma I.B). *Suppose that function c satisfies A3'). In addition, assume that a bounded domain D satisfies the interior ball condition. Then, there exist a constant $r_0 > 0$, depending only on D , $\gamma_{\partial D}$ and $M_A + \|c\|_\infty$ (cf (2.4)), and a nondecreasing function $\rho_{c,A,D} : (0, +\infty) \rightarrow (0, +\infty)$ given by formula (8.26) below and depending only on A , c and D such that for u and $\hat{x} \in \partial D$ as in Theorem 3.5 we have*

$$\partial_{\mathbf{n}} u(\hat{x}) > a\rho_{c,A,D}(r_0 \wedge \mathfrak{r}(\hat{x}))u(\hat{x}). \quad (3.6)$$

Here $\mathfrak{r}(\cdot)$ is as in Definition 2.2.

The proof of the theorem is presented in Section 9.2.

Remark 3.7. At this point we wish to emphasize an important distinction between the conclusions of Theorems 3.5 and 3.6, see also Lemmas 8.7 and 8.8 below. Given a fixed region D , the constant a , appearing in the statements of both results, can be chosen uniformly for the class of operators of the form $A - c$, so long as their appropriate coefficients satisfy the uniform ellipticity condition and respective lower and upper bounds with the same constants. On the other hand, the function $\rho_{c,A,D}(\cdot)$, appearing in Theorem 3.6 and defined in (8.26), depends on the particular operator $-A + c$ through its Green function (see formula (8.23)) and we do not claim that it can be chosen uniformly with respect to the numerical bounds on the coefficients. It is the price we pay for weakening of the assumption on c . In Theorem 3.5 we have assumed that c is bounded away from zero (hypothesis A3''), while in Theorem 3.6 it sufficed only to suppose that c is merely non-trivial (hypothesis A3').

3.3. Harnack property and the Hopf lemma. For operators A having the *Harnack property* and A -harmonic functions we can replace the right hand of (3.4) by a constant multiplied by $u(\hat{x}) - u(x_0)$ for a fixed $x_0 \in D$. To rigorously formulate the result we start with the notion of an A -harmonic function.

Definition 3.8. Suppose that $p > d$ and $D \subset \mathbb{R}^d$ is open. We say that a function $u \in C_b(\mathbb{R}^d) \cap W_{loc}^{2,p}(D)$ is A -harmonic in D iff $Au = 0$ a.e. in D .

Definition 3.9. We say that A has the *Harnack property* in D iff for any $r \in (0, 1]$ there exists $C > 0$ such that for any $y_0 \in D$ satisfying $B(y_0, r) \subset D$ and a non-negative $u \in C_b(\mathbb{R}^d)$, which is A -harmonic in $B(y_0, r)$, we have

$$u(x) \leq Cu(y), \quad x, y \in B(y_0, r/2). \quad (3.7)$$

Remark 3.10. In general, the Harnack property need not hold for A . In fact, some additional hypotheses on the Lévy kernel N are needed. Some sufficient conditions on the kernel guaranteeing the Harnack property of A can be found in [18, Theorem 2.4, p. 25].

The following result is a straightforward conclusion from Theorem 3.4 and the Harnack property of A .

Corollary 3.11. *In addition to the assumptions made in Theorem 3.4 suppose that A has the Harnack property in a bounded domain D . Then, for any $x_0 \in D$ there exists $a > 0$ depending only on M_A , $\gamma_{\partial D}$, $\text{diam } D$ and $\delta_D(x_0) (= \text{dist}(x_0, \partial D))$ such that for any non-constant u , which is A -harmonic in D and attains its maximum in $\mathcal{S}(D) \cup \overline{D}$ at $\hat{x} \in \partial D$, we have*

$$\partial_{\mathbf{n}} u(\hat{x}) \geq a(u(\hat{x}) - u(x_0)) > 0. \quad (3.8)$$

Proof. By Theorem 3.4 we have estimate (3.4) that holds for some positive constants a and r depending only on M_A and $\gamma_{\partial D}$. Let

$$r' := \min \left\{ \frac{1}{2} \delta_D(x_0), r, 1 \right\}. \quad (3.9)$$

Then (3.4) holds also with D_r replaced by $D_{r'}$. Clearly D can be contained in a d -dimensional box of sidelength $\text{diam } D$. We can cover the box with

$$N = \left\{ \left[4 \text{diam } D \sqrt{d}/r' \right] + 1 \right\}^d$$

balls of radius $r'/4$, where $[\cdot]$ is the integer part of a number (the floor function). Let \mathcal{G} be the family of these balls which have a non-empty intersection with $\bar{D}_{r'}$. Obviously $2B \subset D$ for each $B \in \mathcal{G}$. Let $x \in D_{r'}$. Since D is connected there exist $n \geq 1$ and $B_1, \dots, B_n \in \mathcal{G}$ such that $x_0 \in B_1, x \in B_n$ and $B_i \cap B_{i+1} \neq \emptyset$. Let $y_i \in B_i \cap B_{i+1}, i = 1, \dots, n-1$. After an application of the Harnack inequality (3.7) to the non-negative harmonic function $u(\hat{x}) - u(x)$ on each $B_i, i = 1, \dots, n$, we conclude that

$$\inf_{y \in B_{i+1}} (u(\hat{x}) - u(y)) \geq C(r') \inf_{y \in B_i} (u(\hat{x}) - u(y)), \quad i = 1, \dots, n-1,$$

and

$$u(\hat{x}) - u(x) \geq C(r') \inf_{y \in B_n} (u(\hat{x}) - u(y)), \quad \inf_{y \in B_1} (u(\hat{x}) - u(y)) \geq C(r')(u(\hat{x}) - u(x_0)).$$

Here $1/C(r')$ is the constant appearing in (3.7) that corresponds to r' given in (3.9). Thus

$$u(\hat{x}) - u(x) \geq \min_{i=1, \dots, n} \inf_{y \in B_i} (u(\hat{x}) - u(y)) \geq [C(r') \wedge 1]^N (u(\hat{x}) - u(x_0)).$$

Taking the infimum over $x \in D_{r'}$ in the above inequality and using (3.4) we conclude (3.8). \square

4. PROBABILISTIC PRELIMINARIES

Throughout this section we shall assume that the hypotheses A1) – A3) hold.

4.1. On the martingale problem associated with operator A . Let \mathcal{D} be the space consisting of all functions $\omega : [0, +\infty) \rightarrow \mathbb{R}^d$, that are right continuous and possess the left limits for all $t > 0$ (cádlags), equipped with the Skorochod topology, see e.g. Section 12 of [3]. Define the canonical process $X_t(\omega) := \omega(t), \omega \in \mathcal{D}$ and its natural filtration (\mathcal{F}_t) , with $\mathcal{F}_t := \sigma(X_s, 0 \leq s \leq t)$.

Definition 4.1 (A solution of the martingale problem associated with the operator A). Suppose that μ is a Borel probability measure on \mathbb{R}^d . A Borel probability measure P_μ on \mathcal{D} is called a *solution of the martingale problem* associated with A with the initial distribution μ , cf [29], if

- i) $P_\mu[X_0 \in Z] = \mu[Z]$ for any Borel measurable $Z \subset \mathbb{R}^d$.
- ii) For every $f \in C_b^2(\mathbb{R}^d)$ - a C^2 -smooth function bounded with its two derivatives on \mathbb{R}^d - the process

$$M_t[f] := f(X_t) - f(X_0) - \int_0^t Af(X_r) dr, \quad t \geq 0 \tag{4.1}$$

is a (cádlag) martingale under measure P_μ with respect to natural filtration $(\mathcal{F}_t)_{t \geq 0}$. As usual we write $P_x := P_{\delta_x}, x \in \mathbb{R}^d$ and say that x is the initial condition. The expectations with respect to P_μ and P_x shall be denoted by \mathbb{E}_μ and \mathbb{E}_x , respectively.

Definition 4.2 (A strong Markovian solution of the martingale problem). We say that a family of Borel probability measures $(P_x)_{x \in \mathbb{R}^d}$ on \mathcal{D} is a strong Markovian solution to the martingale problem associated with A if:

- i) each P_x is a solution of the martingale problem associated with A , corresponding to the initial condition at x ,
- ii) the canonical process (X_t) is strongly Markovian with respect to the natural filtration (\mathcal{F}_t) and the family $(P_x)_{x \in \mathbb{R}^d}$,
- iii) the mapping $x \rightarrow P_x[C]$ is measurable for any Borel $C \subset \mathcal{D}$,
- iv) for any Borel probability measure μ on \mathbb{R}^d the probability measure

$$P_\mu(\cdot) := \int_{\mathbb{R}^d} P_x(\cdot) \mu(dx)$$

is a solution to the martingale problem associated with A with the initial distribution μ .

The following result concerning the existence of a strong Markov solution for a martingale problem has been proven in [1], see also [22].

Theorem 4.3. *Suppose that conditions (2.3) and (2.4) are satisfied. Then, the martingale problem associated with the operator A admits a strong Markovian solution.*

4.2. Analytic description of the canonical process and weak subsolution to (1.7).

4.2.1. *Exit time, transition probability semigroup, resolvent operator.* For a given domain D define the exit time $\tau_D : \mathcal{D} \rightarrow [0, +\infty]$ of the canonical process $(X_t)_{t \geq 0}$ from D as

$$\tau_D := \inf\{t > 0 : X_t \notin D\}. \quad (4.2)$$

It is a stopping time, i.e. for any $t \geq 0$ we have $[\tau_D \leq t] \in \mathcal{F}_t$, see Theorem I.10.7, p. 54 of [5] and Theorem IV.3.12, p. 181 of [17].

Proposition 4.4. *Suppose that D is a bounded domain. Then,*

$$\sup_{x \in D} \mathbb{E}_x \tau_D < +\infty. \quad (4.3)$$

Proof. See e.g. [18, Lemma 4]. □

We let

$$P_t^D f(x) := \mathbb{E}_x [f(X_t) 1_{[t < \tau_D]}] \quad t \geq 0, \quad f \in B_b(D). \quad (4.4)$$

This is a probabilistic formula for the semigroup generated by operator (1.4) on D with the null exterior condition.

For a positive $f \in B(D)$ and $\alpha \geq 0$ we let

$$R_\alpha^D f(x) := \int_0^\infty e^{-\alpha t} P_t^D f(x) dt = \mathbb{E}_x \left[\int_0^{\tau_D} e^{-\alpha r} f(X_r) dr \right], \quad x \in D. \quad (4.5)$$

Set $R^D := R_0^D$. We have $R_\alpha^D f(x) < +\infty$, $x \in D$ for $f \in B_b(D)$ and $\alpha > 0$.

4.2.2. The gauge function for $c(\cdot)$ and domain D .

Definition 4.5 (The gauge function for $c(\cdot)$ and domain D). The function

$$v_{c,D}(x) = \mathbb{E}_x e_c(\tau_D), \quad x \in \mathbb{R}^d, \quad (4.6)$$

where, the random functional,

$$e_c(t) := e^{-\int_0^t c(X_r) dr}, \quad t \geq 0, \quad (4.7)$$

is called the *gauge function* corresponding to $c(\cdot)$ and domain D , cf Section 4.3 of [12].

Let us denote

$$w_{c,D}(x) := 1 - v_{c,D}(x), \quad x \in \mathbb{R}^d. \quad (4.8)$$

Obviously $w_{c,D}(x) \geq 0$, $x \in \mathbb{R}^d$. In fact, as it turns out, under the assumptions made in the present paper $w_{c,D}(x)$ is strictly positive and continuous in D , see Lemma 8.5 below.

4.3. Feynman-Kac formula for (X_t) .

Proposition 4.6. *1) If $u \in W_{loc}^{2,p}(D)$ and $p > d$, then for any $V \Subset D$ and $t \geq 0$*

$$u(x) = \mathbb{E}_x [e_c(\tau_V \wedge t) u(X_{\tau_V \wedge t})] + \mathbb{E}_x \left[\int_0^{\tau_V \wedge t} e_c(s) (-Au + cu)(X_s) ds \right], \quad x \in V. \quad (4.9)$$

Moreover, (4.9) holds also with τ_V in place of $\tau_V \wedge t$.

2) If u is a subsolution of (1.7), then

$$u(x) = \mathbb{E}_x [e_c(\tau_D \wedge t)u(X_{\tau_D \wedge t})] + \mathbb{E}_x \left[\int_0^{\tau_D \wedge t} e_c(s)(-Au + cu)(X_s) ds \right], \quad x \in D. \quad (4.10)$$

for any $t \geq 0$. Moreover, if D is bounded, then (4.10) holds also with τ_D in place of $\tau_D \wedge t$.

The proof of this result is given in Section 6.1 below.

4.4. Principal eigenvalue and eigenfunction associated with A . The following result is the version of the Krein-Rutman theorem for the operators considered in Section 2.2.

Theorem 4.7. *Suppose that the domain D is bounded. Then, there exists a unique pair (φ_D, λ_D) , where $\varphi_D : D \rightarrow (0, +\infty)$ - a strictly positive continuous function - and $\lambda_D > 0$ are such that $\int_D \varphi_D dx = 1$ and*

$$e^{-\lambda_D t} \varphi_D(x) = P_t^D \varphi_D(x), \quad t \geq 0, x \in D. \quad (4.11)$$

The proof of the theorem is presented in Section 7

Definition 4.8. The function φ_D and λ_D are called respectively: the *principal eigenfunction* and its associate *principal eigenvalue* for the operator A , with the zero Dirichlet exterior condition on D^c , see e.g. [6].

5. GENERALIZATIONS OF THE QUANTITATIVE HOPF LEMMAS AND WEAK HARNACK ESTIMATES

Throughout this section we shall always assume that domain D is bounded.

5.1. Some generalizations of Theorems 3.5 and 3.6. Below, we formulate some generalizations of estimates (3.5) and (3.6). In general, we shall no longer assume that the boundary ∂D is regular enough so that the outward normal, appearing in the left hand sides of the aforementioned bounds, can be defined. It shall be replaced by the ratio $[u(\hat{x}) - u(x)]/\psi(x)$, $x \in D$. Here $\hat{x} \in \partial D$ and $\psi : D \rightarrow (0, +\infty)$ are the maximum point of the subsolution u and some suitably defined function (e.g. the principal eigenfunction associated with the operator), respectively. In case $\psi(x) \approx \delta_D(x)$ ($= \text{dist}(x, D^c)$), when $x \rightarrow \hat{x}$ (which may happen for sufficiently regular domains) these type of estimates lead to (3.5) and (3.6).

Theorem 5.1 (Quantitative Hopf lemma II.A). *Suppose that the assumption A3) holds. Then, there exist a continuous function $\psi : D \rightarrow (0, +\infty)$ and a Borel function $\chi : D \rightarrow (0, +\infty)$, such that for any subsolution $u(\cdot)$ of (1.7) satisfying*

$$u(\hat{x}) = \max_{x \in \mathcal{S}(D) \cup \bar{D}} u(x) \geq 0 \quad \text{for some } \hat{x} \in \partial D$$

we have

$$u(\hat{x}) - u(x) \geq \frac{c\varphi_D(x)u(\hat{x})}{2e\|\varphi_D\|_\infty(\lambda_D + c)} + \psi(x) \int_D (Au - cu)\chi dx, \quad x \in D. \quad (5.1)$$

The result is proved in Section 9.5.

Instead of the second term in the right hand side of (5.1) we can use a term depending on the principal eigenfunction, but then we have to replace the mean of $Au - cu$, with respect to some measure, by its essential infimum.

Theorem 5.2 (Quantitative Hopf lemma II.B). *Suppose that $u(\cdot)$ and \hat{x} are as in Theorem 5.1. Then, under the assumptions of the above theorem we have (cf (2.7))*

$$u(\hat{x}) - u(x) \geq \frac{\varphi_D(x)}{2e\|\varphi_D\|_\infty} \left[\frac{c u(\hat{x})}{\lambda_D + c} + \frac{\text{essinf}_D (Au - cu)}{\lambda_D + \|c\|_\infty} \right], \quad x \in D. \quad (5.2)$$

The result is proved in Section 9.4.

Remark 5.3. We stress the fact that in both Theorems 5.1 and 5.2 no assumption on ∂D is made, except that it is a boundary of a bounded domain.

5.2. Remarks on Theorem 5.1. It is interesting to compare Theorem 5.1 with Lemma 3.2 of [11], where a similar estimate is obtained for solutions of the homogeneous Dirichlet boundary problem for the Poisson equation, i.e.

$$-\Delta u = f \geq 0 \quad \text{and} \quad u(x) = 0, \quad x \in \partial D. \quad (5.3)$$

According to [11] if $f \in L^\infty$, ∂D is smooth, then there exists $a > 0$, depending only on D , such that for any u satisfying (5.3) we have

$$u(x) \geq \psi(x) \int_D f \chi \, dx, \quad x \in D, \quad (5.4)$$

where

$$\chi(x) = \delta_D(x) \quad \text{and} \quad \psi(x) = a\delta_D(x). \quad (5.5)$$

Note that in Theorem 5.1 no assumptions on smoothness of ∂D is made. In addition, the operator A is no longer just the laplacian. Instead, it suffices that hypotheses A1) and A2) hold. Obviously, in such a generality one cannot hope to show an analogue of (5.1), with the functions ψ , χ given by (5.5), see Remark 5.5 below. The estimate however holds, provided both the domain and the coefficients of A are sufficiently regular. We have the following.

Theorem 5.4. *Assume that D satisfies both the uniform interior ball and exterior ball conditions. Then, there exists $a > 0$ such that (5.1) holds with ψ replaced by $a\delta_D$ and χ unchanged.*

The proof of the theorem can be found in Section 9.8.

Remark 5.5. An example of a domain that satisfies both the interior and exterior ball conditions, but not in the uniform sense, see [2, Theorem 4], shows that even in the case when A is the laplacian one cannot use $\psi = a\delta_D$ in (5.1) (thus also in (5.4), with ψ as above) for some $a > 0$. On the other hand, the classical Hopf lemma, as formulated in [2, p. 82], is then satisfied.

We can further strengthen Theorem 5.4, allowing ψ and χ as in (5.5), if some additional assumptions on A are made. Consider the operator \hat{A} that is the formal adjoint (w.r.t. the Lebesgue measure) to A , i.e.

$$\int_{\mathbb{R}^d} v Au \, dx = \int_{\mathbb{R}^d} u \hat{A}v \, dx, \quad u, v \in C_c^2(\mathbb{R}^d). \quad (5.6)$$

Assume that it is of the form $\hat{A}_0 - \hat{c}$, where

$$\begin{aligned} \hat{A}_0 u(x) &= \frac{1}{2} \sum_{i,j=1}^d \hat{q}_{i,j}(x) \partial_{x_i, x_j}^2 u(x) + \sum_{i=1}^d \hat{b}_i(x) \partial_{x_i} u(x) \\ &+ \int_{\mathbb{R}^d} \left(u(x+y) - u(x) - \sum_{i=1}^d \frac{y_i \partial_{x_i} u(x)}{1+|y|^2} \right) \hat{N}(x, dy), \quad u \in C^2(D) \cap C_b(\mathbb{R}^d), \end{aligned} \quad (5.7)$$

with the coefficients $\hat{q}_{i,j}$, \hat{b}_i , \hat{N} satisfying A1), A2) and $\hat{c} \in B_b(\mathbb{R}^d)$. Define $\hat{\mathcal{S}}(D)$ using (1.8) for \hat{N} .

Let (\hat{P}_x) be a strong Markovian solution to the martingale problem corresponding to \hat{A}_0 . Then, by the strong Markov property

$$\hat{P}_t^{\hat{c}, D} f(x) := \hat{\mathbb{E}}_x [f(X_t) e_{\hat{c}}(t) \mathbf{1}_{\{t < \tau_D\}}], \quad x \in D, t \geq 0, f \in B_b(D) \quad (5.8)$$

defines a semigroup of operators $(\hat{P}_t^{\hat{c}, D})_{t \geq 0}$ on $B_b(D)$. Let $\hat{R}_\alpha^{\hat{c}, D}$ be the respective resolvent operator, defined by (4.5), where P_t^D is replaced by $\hat{P}_t^{\hat{c}, D}$. It can be defined for all $\alpha \geq \|\hat{c}^+\|_\infty$. Assume that

$$\int_D \hat{R}_\alpha^{\hat{c}, D} f g \, dx = \int_D f R_\alpha^D g \, dx, \quad f, g \in B^+(D), \alpha \geq \|\hat{c}^+\|_\infty. \quad (5.9)$$

The following result can be inferred from the quantitative Hopf lemma expressed in Theorem 5.1. Its proof is given in Section 9.8.

Theorem 5.6. *Assume that D satisfies both the uniform interior and exterior ball conditions and \hat{A} is as described above. Then, the conclusion of Theorem 5.1 holds with ψ and χ as in (5.5).*

In order to use Theorem 5.6 we need to verify that the resolvent $\hat{R}_\alpha^{\hat{c},D}$ satisfies (5.9). The equality holds, provided that both ∂D and coefficients of \hat{A} are sufficiently regular. Before we formulate some sufficient condition let us first recall the notion of a function of vanishing mean oscillation.

Definition 5.7 (Functions of vanishing mean oscillation). Given a function $f \in L^1_{\text{loc}}(\mathbb{R}^d)$ and $r > 0$ let us define

$$\eta(r) := \sup_{B \in \mathfrak{B}_r} \frac{1}{m_d(B)} \int_B |f(x) - f_B| dx.$$

Here \mathfrak{B}_r denotes the family of all balls of radius less than, or equal to r and $f_B := (1/m_d(B)) \int_B f(x) dx$. We say that a function is of vanishing mean oscillation (VMO) if $\sup_{r>0} \eta(r) < +\infty$, i.e. it is of bounded mean oscillation (BMO), and $\lim_{r \rightarrow 0} \eta(r) = 0$. Denote by *VMO* the class of all such functions.

The following result is shown in Section 9.9.

Theorem 5.8. *Suppose that ∂D is of C^2 class, the coefficients of both A and \hat{A}_0 satisfy hypotheses A1) and A2) and $\hat{c} \in B_b(\mathbb{R}^d)$. Furthermore, assume that $q_{i,j}, \hat{q}_{i,j}$ belong to *VMO* for all $i, j = 1, \dots, d$ and $\mathcal{S}(D) \cup \hat{\mathcal{S}}(D) \subset \bar{D}$. Then (5.9) holds and, as a result, the conclusion of Theorem 5.1 remain valid with ψ and χ as in (5.5).*

5.3. Weak Harnack inequality. We end this section with the result that can be interpreted as a weak form of the Harnack inequality.

Theorem 5.9. *Let D be a bounded domain in \mathbb{R}^d and c satisfies hypothesis A3). Then, there exists a bounded Borel measurable function $\chi : D \rightarrow (0, +\infty)$, such that for any $V \Subset D$ we can find a constant $C > 0$ such that for any u - a non-negative supersolution to (1.7) - we have*

$$\inf_{x \in V} u(x) \geq C \int_V u \chi dx. \quad (5.10)$$

The proof of the result is shown in Section 9.7.

A direct consequence of part 2) of Theorem 3.4 and Theorem 5.9 is the following.

Corollary 5.10. *Under the assumptions of part 2) of Theorem 3.4 and Theorem 5.9 for any $r > 0$ there exist $C > 0$ such that any non-constant subsolution u to (1.7), that attains its non-negative maximum over $\mathcal{S}(D) \cup \bar{D}$ at $\hat{x} \in \partial D$, and any $\mathbf{n} \in \mathbf{n}(\hat{x})$ at $\hat{x} \in \partial D$ satisfies (cf (3.3))*

$$\frac{\partial_{\mathbf{n}} u(\hat{x})}{r} \geq C \int_{D_r} (u(\hat{x}) - u(x)) \chi(x) dx > 0. \quad (5.11)$$

Here χ is as in the statement of Theorem 5.9.

6. PROOFS OF THE FEYNMAN-KAC FORMULA AND WEAK MAXIMUM PRINCIPLE

6.1. Proof of Proposition 4.6 - the Feynman-Kac formula. Suppose that $u \in W_{\text{loc}}^{2,p}(D) \cap C_b(\mathbb{R}^d)$ for some $p > d$ is a subsolution of (1.7). We can find $(u_n) \subset C_c^2(\mathbb{R}^d)$ such that $\|u - u_n\|_{W^{2,p}(V)} \rightarrow 0$ for any $V \Subset D$. This in particular implies that $u_n \rightarrow u$ and $\nabla u_n \rightarrow \nabla u$ uniformly on V , and $\|D^2 u_n - D^2 u\|_{L^p(V)} \rightarrow 0$ for any $V \Subset D$, as $n \rightarrow +\infty$. As a result for any such V , we have

$$\lim_{n \rightarrow +\infty} \|A u_n - A u\|_{L^p(V)} = 0. \quad (6.1)$$

By [29, Theorem 4.3],

$$M_t^n := u_n(X_t) - u_n(x) - \int_0^t A u_n(X_r) dr, \quad t \geq 0 \quad (6.2)$$

is a martingale under the measure P_x for every $x \in \mathbb{R}^d$. By the Itô formula, applied to $e_c(t)u_n(X_t)$, for any $V \in \mathcal{D}$ we get

$$\begin{aligned} u_n(x) &= e_c(\tau_V \wedge t)u_n(X_{\tau_V \wedge t}) + \int_0^{\tau_V \wedge t} e_c(r)(-Au_n + cu_n)(X_r) dr \\ &\quad - \int_0^{\tau_V \wedge t} e_c(r) dM_r^n, \quad t \geq 0, \quad P_x\text{-a.s.} \end{aligned}$$

Hence, we conclude that

$$u_n(x) = \mathbb{E}_x \left[e_c(\tau_V \wedge t)u_n(X_{\tau_V \wedge t}) \right] + \mathbb{E}_x \left[\int_0^{\tau_V \wedge t} e_c(s)(-Au_n + cu_n)(X_s) ds \right], \quad x \in V. \quad (6.3)$$

Recall the Krylov estimate

$$\sup_{x \in V} \mathbb{E}_x \left[\int_0^{\tau_V} |f(X_t)| dt \right] \leq C \|f\|_{L^d(V)}, \quad f \in L^d(V) \quad (6.4)$$

for some $C > 0$, see [1, Corollary p. 143], or [22, Thm III₁₄, (53)]. Letting $n \rightarrow \infty$ in (6.3) and using (6.4) we get

$$u(x) = \mathbb{E}_x \left[e_c(\tau_V \wedge t)u(X_{\tau_V \wedge t}) \right] + \mathbb{E}_x \left[\int_0^{\tau_V \wedge t} e_c(s)(-Au + cu)(X_s) ds \right], \quad x \in V. \quad (6.5)$$

Allowing then $t \rightarrow \infty$, using (6.4) and the fact that $P_x(\tau_V < \infty) = 1$, we conclude part 1) of the proposition.

Concerning the proof of part 2). Let $\{D_n\}$ be an increasing sequence of relatively compact open subsets of D such that $D_n \subset D_{n+1} \Subset D$ and $\bigcup_{n \geq 1} D_n = D$. Then the exit time τ_{D_n} increases to τ_D , as $n \rightarrow +\infty$. Taking D_n in place of V in (6.5) and letting $n \rightarrow \infty$, we get (using quasi-left continuity of the process, see Theorem IV.3.12, p. 181 of [17]) formula (4.10) for $x \in D$. Next, using the fact that $Au - cu \geq 0$ on D (since u is a subsolution of (1.7)), the fact that $P_x(\tau_D < \infty) = 1$, and the monotone convergence theorem, we may let $t \rightarrow \infty$ in (4.10). \square

6.2. Proof of Proposition 3.1 - the weak maximum principle. As an immediate application of the Feynman-Kac formula (Proposition 4.6) we show the weak maximum principle formulated in Proposition 3.1. Using formula (4.10), remembering that $-Au + cu \leq 0$, and letting $t \rightarrow +\infty$ we conclude that

$$u(x) \leq \mathbb{E}_x [e_c(\tau_D)u(X_{\tau_D})] \leq \mathbb{E}_x u^+(X_{\tau_D}), \quad x \in D. \quad (6.6)$$

Since D is bounded, cf (4.3), we have $\tau_D < +\infty$, P_x a.s. This in particular implies that X_{τ_D} is a.s. well defined.

Using the Ikeda-Watanabe formula, see [10, Remark 2.46, page 65], we obtain

$$\begin{aligned} &\mathbb{E}_{x_0} \left[\sum_{0 < s \leq \tau_D} \mathbf{1}_D(X_{s-}) \mathbf{1}_{\mathcal{S}(D)^c \setminus D}(X_s) \right] \\ &= \mathbb{E}_{x_0} \left[\int_{\mathbb{R}^d} \int_0^{\tau_D} \mathbf{1}_D(X_{s-}) \mathbf{1}_{\mathcal{S}(D)^c \setminus D}(y) N(X_{s-}, dy - X_{s-}) ds \right]. \end{aligned} \quad (6.7)$$

From the definition of $\mathcal{S}(D)$, cf (1.8), we conclude that the right-hand side of the above equation vanishes. Thus, $X_{\tau_D} \in \mathcal{S}(D) \setminus D$, if $X_{\tau_D} \neq X_{\tau_D-}$, or $X_{\tau_D} \in \partial D$, if otherwise. Summarizing, we have shown that

$$P_x(X_{\tau_D} \in (\mathcal{S}(D) \setminus D) \cup \partial D) = 1, \quad x \in D$$

and estimate (3.1) follows. When $c \equiv 0$ we get (6.6), with u^+ replaced by u . \square

7. PROPERTIES OF THE RESOLVENT - PROOF OF THEOREM 4.7

Suppose that D is a bounded domain. Recall that $R_\alpha^D f$ is defined in (4.5) for all $\alpha \geq 0$ and $f \in B^+(D)$. Using condition (4.3) we infer that the operator $R_\alpha^D : L^\infty(D) \rightarrow L^\infty(D)$ is bounded for any $\alpha \geq 0$. The operators $(R_\alpha^D)_{\alpha \geq 0}$ satisfy the resolvent identity

$$R_\alpha^D - R_\beta^D = (\beta - \alpha)R_\alpha^D R_\beta^D, \quad \alpha, \beta \geq 0. \quad (7.1)$$

Thanks to assumptions A1)-A2) there exists $C > 0$ such that

$$\sup_{x \in D} \mathbb{E}_x \left[\int_0^{\tau_D} |f(X_t)| dt \right] \leq C \|f\|_{L^d(D)}, \quad f \in L^d(D). \quad (7.2)$$

see (53) of [22], or Corollary 2 of [24]. The estimate implies that in fact R_α^D extends to the operator $R_\alpha^D : L^d(D) \rightarrow L^\infty(D)$ for each $\alpha \geq 0$. This in turn allows us to conclude that for any $\alpha \geq 0$ there exists $r_\alpha^D : D \times D \rightarrow [0, +\infty)$ such that

$$R_\alpha^D f(x) = \int_D r_\alpha^D(x, y) f(y) dy, \quad x \in D \quad (7.3)$$

and

$$\operatorname{esssup}_{x \in D} \|r_\alpha^D(x, \cdot)\|_{L^{d/(d-1)}(D)} \leq \|R_\alpha\|_{L^d \rightarrow L^\infty}. \quad (7.4)$$

The above implies that $R_\alpha^D : L^d(D) \rightarrow L^d(D)$ is compact for any $\alpha \geq 0$. This can be easily seen, as from any bounded sequence $(f_n) \subset L^d(D)$ we can select a subsequence (f_{n_k}) weakly converging to some f . Using (7.3) we conclude that $\lim_{k \rightarrow +\infty} R_\alpha^D f_{n_k}(x) = R_\alpha^D f(x)$ for a.e. $x \in D$. Since $(\|R_\alpha^D f_{n_k}\|_{L^\infty})$ is bounded, this implies the strong convergence of $(R_\alpha^D f_{n_k})$ to $R_\alpha^D f$ in $L^d(D)$.

By the support theorem of [18, Section 4], or by [23, Corollary 2] for any $x \in D$,

$$r_\alpha^D(x, \cdot) > 0, \quad m\text{-a.e. on } D, \alpha \geq 0. \quad (7.5)$$

By [24, Theorem 1] we infer that the resolvent R_α^D is strongly Feller for each $\alpha \geq 0$, i.e.

$$R_\alpha^D(B_b(D)) \subset C_b(D), \quad \alpha \geq 0. \quad (7.6)$$

Using the kernel $r_\alpha^D(\cdot, \cdot)$ we can define the adjoint (with respect to the Lebesgue measure) resolvent operator $\hat{R}_\alpha^D : L^{d/(d-1)}(D) \rightarrow L^{d/(d-1)}(D)$ given by

$$\hat{R}_\alpha^D \psi(y) = \int_D r_\alpha^D(x, y) \psi(x) dx, \quad y \in D, \psi \in L^{d/(d-1)}(D). \quad (7.7)$$

We say that it is strongly Feller, if the analogue of (7.6) holds for \hat{R}_α^D .

7.1. The proof of Theorem 4.7. We have already concluded that $R_\alpha^D : L^d(D) \rightarrow L^d(D)$ is compact for any $\alpha \geq 0$. In addition $\rho(R_1)$ - the spectral radius of R_1^D - belongs to $[0, 1]$. Moreover, by (7.5), operator R_1^D is *irreducible*, i.e.

$$\text{for any } f \in B^+(D) \text{ such that } \int_D f dx > 0 \text{ we have } R_1^D f(x) > 0, x \in D. \quad (7.8)$$

By virtue of the Jentzsch Theorem, see Theorem 6.6. of [28], the spectral radius $\rho(R_1^D)$ is positive and it equals to the (unique) simple eigenvalue of R_1^D . The corresponding eigenvector φ_D is strictly positive on D . With the help of the resolvent identity (7.1), applied to φ_D , with $\alpha + 1$ in place of α and $\beta = 1$, one can easily check that

$$R_{\alpha+1}^D \varphi_D(x) = \frac{\varphi_D(x)}{\rho^{-1}(R_1^D) + \alpha}, \quad \alpha > 0, x \in D.$$

Thus, inverting the Laplace transform, we obtain

$$e^{-t} P_t^D \varphi_D(x) = \exp\{-t\rho^{-1}(R_1^D)\} \varphi_D(x), \quad t \geq 0,$$

for each $x \in D$. This in turn leads to (4.11), with $\lambda_D := \rho^{-1}(R_1^D) - 1$. Clearly, $\lambda_D > 0$ as otherwise, this would contradict (4.3). \square

7.2. Irreducibility measure and its application. We recall the notion of an irreducibility measure associated with a non-negative kernel. Given $\alpha > 0$ we let $K_\alpha : D \times \mathcal{B}(D) \rightarrow [0, +\infty)$ be a substochastic kernel defined by

$$K_\alpha(x, B) := R_\alpha^D 1_B(x) \quad \text{for } (x, B) \in D \times \mathcal{B}(D).$$

Definition 7.1. (cf [25, Section 2.2]) Suppose that ν is a non-trivial Borel measure on D . We say that the kernel K_α is ν -irreducible, if for any $B \in \mathcal{B}(D)$ such that $\nu(B) > 0$ and $x \in D$ we have

$$K_\alpha^n(x, B) := \int_D K_\alpha(x, dy_1) \left(\int_D K_\alpha(y_1, dy_2) \dots \int_D K_\alpha(y_{n-1}, B) \right) > 0$$

for some $n \geq 1$. By convention $y_0 := x$. Measure ν is called then an *irreducibility measure* for K_α . The kernel is said to be irreducible if it is irreducible for some measure. A measure μ is called a *maximal irreducibility measure* for K_α if it is an irreducibility measure and any irreducibility measure ν for the kernel satisfies $\nu \ll \mu$.

Lemma 7.2. For each $\alpha \geq 0$ there exists a pair of functions $\bar{\psi}_\alpha, \bar{\phi}_\alpha \in B_b^+(D)$ such that both $\bar{\psi}_\alpha > 0$ and $\bar{\phi}_\alpha > 0$, m -a.e., and

$$R_\alpha^D f(x) \geq R_{\alpha+1}^D \bar{\psi}_\alpha(x) \int_D R_{\alpha+1}^D f \bar{\phi}_\alpha dx, \quad x \in D, \quad f \in B_b^+(D). \quad (7.9)$$

Proof. Estimate (7.8) implies that the kernel $K_\alpha(\cdot, \cdot)$ is m -irreducible. Here, as we recall m is the d -dimensional Lebesgue measure. By (7.3) we conclude also that $\int_D K_\alpha(x, \cdot) dx \ll m$. Using [25, part ii) of Proposition 2.4] we infer that the Lebesgue measure is in fact a maximal irreducible measure for the kernel. By [25, Theorem 2.1] there exist an integer $n \geq 1$, a function $\psi_\alpha : D \rightarrow [0, \infty)$ and a non-trivial Borel measure ν_α on D , the so called *small function* and *small measure*, such that $\int_D \psi_\alpha dx > 0$ and

$$\left(R_{\alpha+1}^D \right)^n f(x) \geq \psi_\alpha(x) \int_D f d\nu_\alpha, \quad f \in B^+(D), \quad x \in D. \quad (7.10)$$

We claim that in fact from (7.10) it follows that

$$R_\alpha^D f(x) \geq R_{\alpha+1}^D f(x) \geq \psi_\alpha(x) \int_D f d\nu_\alpha, \quad f \in B^+(D), \quad x \in D. \quad (7.11)$$

Indeed, we obviously have $R_{\alpha+1}^D f \leq R_\alpha^D f$ for any $f \in B^+(D)$. By the resolvent identity, see (7.1),

$$R_\alpha^D R_{\alpha+1}^D f = R_{\alpha+1}^D R_\alpha^D f = R_\alpha^D f - R_{\alpha+1}^D f \leq R_\alpha^D f \quad \text{for any } f \in B^+(D). \quad (7.12)$$

If $n \geq 2$ in (7.10), then we can write

$$\begin{aligned} R_\alpha^D \left(R_{\alpha+1}^D \right)^{n-2} f(x) &\geq \left(R_\alpha^D R_{\alpha+1}^D \right) \left(R_{\alpha+1}^D \right)^{n-2} f(x) \\ &\geq \left(R_{\alpha+1}^D \right)^n f(x) \geq \psi_\alpha(x) \int_D f d\nu_\alpha, \quad f \in B^+(D), \quad x \in D. \end{aligned} \quad (7.13)$$

Iterating this procedure, we conclude (7.11).

Applying (7.11) for $R_{\alpha+1}^D f$ in place of f , we get

$$R_\alpha^D f(x) \geq \psi_\alpha(x) \int_D f d\bar{\nu}_\alpha, \quad f \in B^+(D), \quad x \in D, \quad (7.14)$$

where

$$\bar{\nu}_\alpha(B) := \int_B R_{\alpha+1}^D 1_B d\nu_\alpha, \quad B \in \mathcal{B}(D).$$

Applying $R_{\alpha+1}^D$ to both sides of (7.14) and using again (7.12), we get

$$R_\alpha^D f(x) \geq \bar{\psi}_\alpha(x) \int_D f d\bar{\nu}_\alpha, \quad f \in B^+(D), \quad x \in D, \quad (7.15)$$

with $\bar{\psi}_\alpha := R_{\alpha+1}^D \psi_\alpha > 0$, m a.e. Applying $R_{\alpha+1}^D$ to both sides of (7.15) and using (7.12) we get

$$R_\alpha^D f(x) \geq R_{\alpha+1}^D \bar{\psi}_\alpha(x) \int_D f d\bar{\nu}_\alpha, \quad f \in B^+(D), \quad x \in D. \quad (7.16)$$

Invoking (7.8) we infer that $\bar{\nu}_\alpha \ll m$. Thus, there exists a non-trivial $\bar{\phi}_\alpha \in B_b^+(D)$ such that $d\bar{\nu}_\alpha = \bar{\phi}_\alpha dx$. By (7.5), $\bar{\phi}_\alpha$ is strictly positive, m -a.e.. Therefore, applying (7.16) with f replaced by $R_{\alpha+1}^D f$ we conclude (7.9). \square

7.3. Continuity of the resolvent to the boundary. Recall that a point $\hat{x} \in \partial D$ is called *Dirichlet regular* if $P_{\hat{x}}(\tau_D > 0) = 0$. If all the points of the boundary of D are Dirichlet regular, we say that D is Dirichlet regular. It is well known (see [13, Theorem 1.12]) that if $(R_\alpha^D)_{\alpha>0}$ is strongly Feller, i.e. (7.6) holds, and $\hat{x} \in \partial D$ is Dirichlet regular, then

$$\lim_{D \ni x \rightarrow \hat{x}} \mathbb{E}_x \tau_D = 0. \quad (7.17)$$

Therefore, for a Dirichlet regular D and $g \in B_b^+(D)$ function $R_\alpha^D g$ is continuous in \bar{D} , and

$$\lim_{D \ni x \rightarrow \hat{x}} R_\alpha^D g(x) = 0, \quad \text{for any } \hat{x} \in \partial D.$$

Proposition 7.3. *Suppose that D is bounded and conditions A1), A2) are in force. If D satisfies the exterior ball condition (see Definition 2.5), then D is Dirichlet regular.*

Proof. By [4, Proposition VII.3.3], to prove that $\hat{x} \in \partial D$ is Dirichlet regular it is enough to find a function ψ - the so called *barrier* - that is strictly positive, and superharmonic on D , i.e. $-A\psi \geq 0$ on D , such that

$$\lim_{\substack{x \rightarrow \hat{x} \\ x \in D}} \psi(x) = 0.$$

For $\hat{x} \in \partial D$ let $B(y_0, r)$ be the exterior ball to D at this point. Suppose that M is so large that $D \subset B(y_0, M)$. Define a compactly supported, smooth function $\psi : \mathbb{R}^d \rightarrow \mathbb{R}$, such that

$$\psi(x; \hat{x}) = c \left(\frac{1}{r^\sigma} - \frac{1}{|x - y_0|^\sigma} \right) \quad \text{for } \frac{r}{10} < |x - y_0| < M$$

for some $c, \sigma > 0$. It is elementary to check that ψ fulfills the requirements for a barrier, provided $c, \sigma > 0$ are sufficiently large. \square

8. SOME AUXILIARY RESULTS

8.1. An auxiliary function and its lower bound. Given $\bar{y} \in \mathbb{R}^d$ and $\beta > 0$ define

$$\eta(x; \beta, \bar{y}, r) = e^{-\beta|x-\bar{y}|^2} - e^{-\beta r^2}, \quad x \in \mathbb{R}^d. \quad (8.1)$$

We shall use this function in the proofs of the strong maximum principle and both the Hopf, and quantitative Hopf lemmas expressed in Theorems 3.4 – 3.6. For this purpose we shall prove various estimates of $\eta(x; \beta, \bar{y}, r)$ that are formulated in the present section as well as in Section 8.4.

Define open annuli

$$V_*(\bar{y}; r) = B(\bar{y}, r) \setminus \bar{B}(\bar{y}, r/2), \quad V^*(\bar{y}; r) = B(\bar{y}, 3r/2) \setminus \bar{B}(\bar{y}, r/2). \quad (8.2)$$

For given $y \in \mathbb{R}^d$ we let $y^2 := y \otimes y = [y_i y_j]_{i,j=1}^d$. We start with the following result.

Lemma 8.1. *Suppose that D is bounded satisfies the interior ball condition with $\mathfrak{r}(\cdot)$ as in Definition 2.2. Then, for any $K > 0$ there exist constants $r_0, C > 0$ depending only on $\gamma_{\partial D}, \|N\|_\infty, \|Q\|_\infty, \|b\|_\infty, \|c\|_\infty$ such that for every $\hat{x} \in \partial D$, any interior ball $B(\bar{y}, r)$ at \hat{x} (cf Definition 2.2) with $r \in (0, r_0 \wedge \mathfrak{r}(\hat{x}))$, we have*

$$(A - c)\eta(x; Cr^{-2}, \bar{y}, r) \geq K, \quad x \in V^*(\bar{y}; r). \quad (8.3)$$

Proof. Let $0 < r \leq 1$, $B(\bar{y}, r)$ be an interior ball in D at \hat{x} , and $V^* := V^*(\bar{y}; r)$ (cf (8.2)), see Fig. 1. Let $\eta(x) := \eta(x; \beta, \bar{y}, r)$, $x \in \mathbb{R}^d$ be given by (8.1). Then,

$$\begin{aligned}\nabla\eta(x) &= -2\beta(x - \bar{y})e^{-\beta|x - \bar{y}|^2}, \\ \nabla^2\eta(x) &= e^{-\beta|x - \bar{y}|^2} (4\beta^2(x - \bar{y})^2 - 2\beta I_d).\end{aligned}\tag{8.4}$$

With the notation $\text{Tr}(\mathbf{Q}(x))$ for the trace of $\mathbf{Q}(x)$, we can write

$$\sum_{i,j=1}^d \frac{1}{2} q_{i,j}(x) \eta_{x_i x_j}(x) = \beta e^{-\beta|x - \bar{y}|^2} \left[2\beta(x - \bar{y})^T \cdot \mathbf{Q}(x)(x - \bar{y}) - \text{Tr}(\mathbf{Q}(x)) \right]$$

and

$$\sum_{i=1}^d b_i(x) \eta_{x_i} = -2\beta e^{-\beta|x - \bar{y}|^2} b^T(x) \cdot (x - \bar{y}).$$




FIGURE 1

Using the uniform ellipticity assumption (2.3), we get

$$\begin{aligned}L\eta(x) &\geq \beta e^{-\beta|x - \bar{y}|^2} \{ 2\gamma_{\partial D} \beta |x - \bar{y}|^2 - (\text{Tr}(\mathbf{Q}(x)) - 2b^T(x) \cdot (x - \bar{y})) \} \\ &\geq \beta e^{-9\beta r^2/4} \left\{ \frac{\gamma_{\partial D} \beta r^2}{2} - \left(\|\text{Tr} \mathbf{Q}\|_\infty + \frac{3r}{2} \|b\|_\infty \right) \right\}, \quad x \in V^*.\end{aligned}\tag{8.5}$$

We shall choose $\beta > 0$, $r \in (0, 1]$ in such a way that

$$\beta r^2 = \gamma_* := \frac{4}{\gamma_{\partial D}} \left(\|\text{Tr} \mathbf{Q}\|_\infty + \frac{3}{2} \|b\|_\infty \right).\tag{8.6}$$

Then, remembering that $r \in (0, 1]$, we get

$$L\eta(x) \geq \frac{\beta \gamma_* \gamma_{\partial D}}{4} e^{-9\gamma_*/4}, \quad x \in V^*.\tag{8.7}$$

Using (1.5), we can write

$$S\eta = S_1\eta(x) + \int_{\mathbb{R}^d} \frac{\delta\eta(x; y)}{1 + |y|^2} N(x, dy),\tag{8.8}$$

where

$$\begin{aligned}S_1\eta(x) &:= \int_{\mathbb{R}^d} \frac{|y|^2 (\eta(x + y) - \eta(x))}{1 + |y|^2} N(x, dy), \\ \delta\eta(x; y) &:= \eta(x + y) - \eta(x) - \sum_{i=1}^d y_i \partial_{x_i} \eta(x).\end{aligned}\tag{8.9}$$

Using the fact that

$$\delta\eta(x; y) = \int_0^1 d\theta_1 \int_0^{\theta_1} \frac{d^2}{d\theta_2^2} \delta\eta(x; \theta_2 y) d\theta_2 = \int_0^1 d\theta_1 \int_0^{\theta_1} y \cdot (\nabla^2 \eta)(x; \theta_2 y) y d\theta_2$$

and subsequently formula (8.4) for $\nabla^2 \eta$ we conclude that for any $M > 10$

$$S\eta = \sum_{j=1}^4 S_j \eta, \quad (8.10)$$

where, cf (8.4),

$$\begin{aligned} S_2\eta(x) &:= -2\beta \int_0^1 d\theta_1 \int_0^{\theta_1} d\theta_2 \int_{|y| \leq Mr} e^{-\beta|x+\theta_2 y - \bar{y}|^2} \frac{|y|^2}{1+|y|^2} N(x, dy), \\ S_3\eta(x) &:= -2\beta \int_0^1 d\theta_1 \int_0^{\theta_1} d\theta_2 \int_{|y| > Mr} e^{-\beta|x+\theta_2 y - \bar{y}|^2} \frac{|y|^2}{1+|y|^2} N(x, dy), \\ S_4\eta(x) &:= 4\beta^2 \int_0^1 d\theta_1 \int_0^{\theta_1} d\theta_2 \int_{\mathbb{R}^d} e^{-\beta|x+\theta_2 y - \bar{y}|^2} \frac{[(x + \theta_2 y - \bar{y}) \cdot y]^2}{1+|y|^2} N(x, dy), \quad x \in V^*. \end{aligned} \quad (8.11)$$

We have $S_4\eta(x) \geq 0$, therefore

$$S\eta(x) \geq \sum_{j=1}^3 S_j \eta(x), \quad x \in V^*. \quad (8.12)$$

Furthermore, since $\|\eta\|_\infty = 1$ we get, see (1.6),

$$|S_1\eta(x)| \leq 2N_*, \quad x \in V^*. \quad (8.13)$$

In addition,

$$|S_2\eta(x)| \leq \beta n(Mr), \quad x \in V^*, \quad (8.14)$$

where

$$n(r) := \sup_{x \in V^*} N_{B(0,r)}(x).$$

Using assumption A2) and Dini's uniform convergence theorem we conclude that

$$\lim_{r \rightarrow 0^+} n(r) = 0. \quad (8.15)$$

Finally, we can write

$$|S_3\eta(x)| \leq 2\beta\Theta(x, Mr), \quad (8.16)$$

where

$$\Theta(x, r) := \int_0^1 d\theta_2 \int_{\theta_2}^1 d\theta_1 \int_{|y| > r} e^{-\beta|x+\theta_2 y - \bar{y}|^2} \frac{|y|^2}{1+|y|^2} N(x, dy).$$

We have

$$|x + \theta_2 y - \bar{y}| \geq \left(\theta_2 M - \frac{3}{2} \right) r \geq \frac{Mr}{2},$$

provided that

$$\frac{r}{2} + \frac{3}{2M} =: \theta_* \leq \theta_2 \leq 1.$$

We can estimate

$$\begin{aligned} \Theta(x, Mr) &\leq \int_0^{\theta_*} d\theta_2 \int_{\theta_2}^1 d\theta_1 \int_{|y| > Mr} \frac{|y|^2}{1+|y|^2} N(x, dy) \\ &\quad + \int_{\theta_*}^1 d\theta_2 \int_{\theta_2}^1 d\theta_1 e^{-\beta(Mr)^2/4} \int_{|y| > Mr} \frac{|y|^2}{1+|y|^2} N(x, dy) \\ &\leq \left(\theta_* + e^{-\beta(Mr)^2/4} \right) N_* \leq \left(\frac{r}{2} + \frac{3}{2M} + e^{-M^2\gamma_*/4} \right) N_* \\ &= \left(\frac{\sqrt{\gamma_*}}{2\sqrt{\beta}} + \frac{3}{2M} + e^{-M^2\gamma_*/4} \right) N_*. \end{aligned}$$

Using estimates (8.7) and (8.13) – (8.16) we get

$$\begin{aligned}
(A - c)\eta(x) &\geq \frac{\beta\gamma_*\gamma\bar{\sigma}_D}{4}e^{-9\gamma_*/4} - 2N_* - \beta n \left(\frac{M\sqrt{\gamma_*}}{\sqrt{\beta}} \right) \\
&\quad - 2 \left(\frac{\sqrt{\beta}\gamma_*}{2} + \frac{3\beta}{2M} + \beta e^{-M^2\gamma_*/4} \right) N_* - \|c\|_\infty \\
&= \beta \left\{ \frac{\gamma_*\gamma\bar{\sigma}_D}{4}e^{-9\gamma_*/4} - n \left(\frac{M\sqrt{\gamma_*}}{\sqrt{\beta}} \right) - 2 \left(\frac{\sqrt{\gamma_*}}{2\sqrt{\beta}} + \frac{3}{2M} + e^{-M^2\gamma_*/4} \right) N_* \right\} \\
&\quad - 2N_* - \|c\|_\infty.
\end{aligned} \tag{8.17}$$

Therefore, for a $\beta_0 \gg M^2 \gg 1$, we get that the right-hand side of the above inequality is greater than K on V^* . By (8.6) we conclude (8.3) with $r_0 = \sqrt{\gamma_*/\beta_0}$. \square

8.2. A weak subsolution of (2.5).

Definition 8.2 (Weak subsolution, supersolution and solution of (2.5)). Let $c, g \in B_b(\mathbb{R}^d)$. A function $u \in B_b(\mathbb{R}^d)$ is called a *weak subsolution* to (2.5) in D if for any $x \in D$,

$$u(x) \leq \mathbb{E}_x[e_c(t \wedge \tau_D)u(X_{t \wedge \tau_D})] + \mathbb{E}_x \left[\int_0^{t \wedge \tau_D} e_c(s)g(X_s) ds \right], \quad t \geq 0. \tag{8.18}$$

The notions of a supersolution and solution can be then introduced analogously as in Definition 2.1.

Remark 8.3. By Proposition 4.6 if $u \in C_b(\mathbb{R}^d) \cap W_{\text{loc}}^{2,d}(D)$ is a subsolution to (2.5) on D , then u is a weak subsolution to (2.5) on D .

The Feynman-Kac representation of solutions, see Proposition 4.6. holds also for weak subsolutions by their very definition. In particular, we have the following form of a weak maximum principle, that follows by the same argument as Proposition 3.1, see Section 6.2.

Proposition 8.4. *If u is a weak subsolution of (1.7) on a bounded and open set D , then (3.1) is in force.*

8.3. Properties of $w_{c,D}$. Recall that $w_{c,D}(\cdot)$ is given by (4.8). Obviously $0 \leq w_{c,D}(x) \leq 1$, $x \in D$. Define $\hat{w}_{c,D} := \mathbf{1}_D w_{c,D}$, and a martingale

$$M_t := \mathbb{E}_x[e_c(\tau_D)|\mathcal{F}_t], \quad t \geq 0.$$

where $e_c(t)$ is defined in (4.7). Suppose that $x \in D$. By the Markov property we have

$$M_t = e_c(\tau_D)\mathbf{1}_{[\tau_D \leq t]} + v_{c,D}(X_t)e_c(t)\mathbf{1}_{[\tau_D > t]},$$

cf (4.6). Hence,

$$\mathbf{1}_{[\tau_D > t \wedge \tau_D]} w_{c,D}(X_{t \wedge \tau_D}) = 1 - e_c^{-1}(t \wedge \tau_D)M_{t \wedge \tau_D}, \quad t \geq 0.$$

Since $\hat{w}_{c,D}(X_{t \wedge \tau_D}) = \mathbf{1}_{[\tau_D > t \wedge \tau_D]} w_{c,D}(X_{t \wedge \tau_D})$, we have

$$\hat{w}_{c,D}(X_{t \wedge \tau_D}) = 1 - e_c^{-1}(t \wedge \tau_D)M_{t \wedge \tau_D}, \quad t \geq 0. \tag{8.19}$$

Applying Itô's formula to $e_c^{-1}(t)M_t$ yields

$$e_c^{-1}(t)M_t = M_0 + \int_0^t e_c^{-1}(s) dM_s + \int_0^t e_c^{-1}(s)M_s c(X_s) ds. \tag{8.20}$$

Let N_t be the second term in the right-hand side of the above equality. It is a local martingale. Thus, substituting from (8.20) into (8.19) and then replacing t by $t \wedge \tau_V$, where $V \subset D$ is open, we get

$$\begin{aligned}
\hat{w}_{c,D}(X_{t \wedge \tau_V}) &= 1 - M_0 - N_{t \wedge \tau_V} - \int_0^{t \wedge \tau_V} M_s e_c^{-1}(s) c(X_s) ds \\
&= w_{c,D}(X_0) - N_{t \wedge \tau_V} - \int_0^{t \wedge \tau_V} (c v_{c,D})(X_s) ds, \quad t \geq 0.
\end{aligned} \tag{8.21}$$

Therefore, $\hat{w}_{c,D}$ is a weak solution to $-Af = cv_{c,D}$ in V . Applying the Itô formula to $\hat{w}_{c,D}(X_{t \wedge \tau_V})e_c(t \wedge \tau_V)$ we conclude also that $\hat{w}_{c,D}$ is also a weak solution to

$$-Af + cf = c, \quad \text{in any open } V \subset D. \quad (8.22)$$

From (8.21), we also get that

$$w_{c,D}(x) = R^D(cv_{c,D})(x) = R^D(c - cw_{c,D})(x), \quad x \in D. \quad (8.23)$$

By the very definition of $w_{c,D}$ we have

$$w_{c,D}(x) > 0 \quad \text{iff} \quad P_x \left(\int_0^{\tau_D} c(X_r) dr > 0 \right) > 0,$$

which in turn is equivalent to $R^D c(x) > 0$. The latter however is a direct consequence of (7.5). Thanks to (7.6) and (8.23) we conclude also that $w_{c,D}$ is continuous in D . Summarizing we have shown the following.

Lemma 8.5. *The function $w_{c,D}$ is strictly positive and continuous on D .*

The following result shall be used in the proof of estimate (8.31) formulated below.

Lemma 8.6. *Suppose that $A3''$) holds. Then for any open set V , compactly embedded in D , there exists $a_* > 0$ depending only on $\|N\|_\infty, \|Q\|_\infty, \|b\|_\infty, \underline{c} > 0$ and $r_* := \text{dist}(V, D^c) > 0$ such that*

$$\inf_{x \in V} w_{c,D}(x) \geq a_*. \quad (8.24)$$

Proof. Let $\bar{x} \in D$ and $r > 0$ be such that $B(\bar{x}, r) \subset D$. Observe that for any $t > 0$,

$$w_{c,D}(\bar{x}) \geq 1 - \mathbb{E}_{\bar{x}} e^{-\underline{c}\tau_D} \geq 1 - \mathbb{E}_{\bar{x}} e^{-\underline{c}\tau_{B(\bar{x}, r)}} \geq (1 - e^{-\underline{c}t}) P_{\bar{x}}(\tau_{B(\bar{x}, r)} > t).$$

Let $\rho_{\bar{x}, r} \in C_b^2(\mathbb{R}^d)$ be an arbitrary function taking values in $[0, 1]$ and satisfying

$$\rho_{\bar{x}, r}(\bar{x}) = 1, \quad 1 - \rho_{\bar{x}, r}(x) \geq a > 0, \quad x \in B^c(\bar{x}, r).$$

Since the process

$$M_t := 1 - \rho_{\bar{x}, r}(X_{t \wedge \tau_{B(\bar{x}, r)}}) + \int_0^{t \wedge \tau_{B(\bar{x}, r)}} A\rho_{\bar{x}, r}(X_s) ds, \quad t \geq 0$$

is a zero mean martingale we can write

$$\begin{aligned} P_{\bar{x}}(\tau_{B(\bar{x}, r)} \leq t) &\leq \frac{1}{a} \mathbb{E}_{\bar{x}} [(1 - \rho_{\bar{x}, r})(X_{t \wedge \tau_{B(\bar{x}, r)}})] = -\frac{1}{a} \mathbb{E}_{\bar{x}} \left[\int_0^{t \wedge \tau_{B(\bar{x}, r)}} A\rho_{\bar{x}, r}(X_s) ds \right] \\ &\leq \frac{1}{a} \mathbb{E}_{\bar{x}}(t \wedge \tau_{B(\bar{x}, r)}) \sup_{B(\bar{x}, r)} (A\rho_{\bar{x}, r})^- \leq \frac{t}{a} \sup_{B(\bar{x}, r)} (A\rho_{\bar{x}, r})^- \\ &\leq \frac{t}{a} C_A (\|\rho_{\bar{x}, r}\|_\infty + \|D^2 \rho_{\bar{x}, r}\|_\infty) \end{aligned} \quad (8.25)$$

for some constant C_A depending on the coefficients of the operator A .

Taking $r > 0$ sufficiently small, so that $V \Subset D_r$ and letting $\rho_{\bar{x}, r}(x) = e^{-|x - \bar{x}|^2 / r^2}$ for any $\bar{x} \in V$, we can choose t sufficiently small so that (8.24) is in force. \square

8.4. Some upper bounds on $\eta(x; \beta, \bar{y}, r)$. In the present section we prove some upper bounds on $\eta(x; \beta, \bar{y}, r)$ formulated with the help of function $w_{c,D}(x)$. They are instrumental in proving the quantitative Hopf lemmas expressed in Theorems 3.5 and 3.6.

Recall that $D_r = [x \in D : \text{dist}(x, \partial D) > r]$ for $r > 0$, cf (3.3). Let $r_* := \inf[r > 0 : D_r = \emptyset]$. Define

$$\rho_{c,A,D}(r) := \inf_{x \in D_r} w_{c,D}(x), \quad r \in (0, r_*). \quad (8.26)$$

We adopt the convention that $\rho_{c,A,D}(r) := \lim_{\xi \rightarrow r_*^-} \rho_{c,A,D}(\xi)$, if $r \geq r_*$. Clearly, $\rho_{c,A,D}$ is nondecreasing and takes values in $(0, 1]$. Since $w_{c,D}$ is a strictly positive and continuous on D , see Lemma 8.5, we conclude that $\rho_{c,A,D}(\cdot)$ is also strictly positive and continuous on $(0, +\infty)$. We start with the following.

Lemma 8.7. *Suppose that assumption $A3'$ is fulfilled and D is bounded and satisfies the interior ball condition. Let r_0, C be constants as in Lemma 8.1 stated for $K = 1$. Then, there exists a nondecreasing, strictly positive continuous function $\rho_{c,A,D} : (0, +\infty) \rightarrow (0, 1]$ depending only on c , of operator A and the domain D such that for any $\hat{x} \in \partial D$ and interior ball $B(\bar{y}, r)$ at \hat{x} (cf Definition 2.2) with $r \in (0, r_0 \wedge \tau(\hat{x})]$, we have*

$$\eta(x; Cr^{-2}, \bar{y}, r) \rho_{c,A,D}(r/2) \leq w_{c,D}(x), \quad x \in \bar{V}_*(\bar{y}; r). \quad (8.27)$$

Proof. By Lemma 8.1, for given $\hat{x} \in \partial D$, and any interior ball $B(\bar{y}, r)$ at \hat{x} (cf Definition 2.2) with $r \in (0, \tau(\hat{x}) \wedge r_0]$, function $\eta(\cdot) := \eta(\cdot; Cr^{-2}, \bar{y}, r)$ satisfies (8.3). This in particular means that $g := -A\eta < 0$ in $\bar{V}_*(\bar{y}; r)$. For any $\varepsilon \in (0, 1]$ we let $\eta_\varepsilon := \varepsilon\eta$. Clearly, $\eta_\varepsilon \leq 1$ (as $\eta \leq 1$) and

$$-A\eta_\varepsilon + c\eta_\varepsilon = c\eta_\varepsilon + \varepsilon g, \quad \text{in } V_*(\bar{y}; r).$$

By (8.22), $\hat{w}_{c,D}$ is a weak solution to

$$-Au + cu = c, \quad \text{in } V_*(\bar{y}; r).$$

Therefore, $\eta_\varepsilon - \hat{w}_{c,D}$ is a weak subsolution of (1.7) on $V_*(\bar{y}; r)$. Since $\eta \leq 1$ we have

$$\eta_\varepsilon(x) \leq \hat{w}_{c,D}(x), \quad x \in \bar{B}(\bar{y}, r/2), \quad (8.28)$$

with (cf (8.26))

$$\varepsilon := \inf_{D_{r/2}} w_{c,D} = \rho_{c,A,D}(r/2). \quad (8.29)$$

We also have $\eta_\varepsilon \leq 0 \leq \hat{w}_{c,D}$ on $\bar{B}^c(\bar{y}, r)$, therefore by Proposition 8.4 we have

$$\eta_\varepsilon(x) - \hat{w}_{c,D}(x) \leq \sup_{y \in V_*^c(\bar{y}; r)} [\eta_\varepsilon(y) - \hat{w}_{c,D}(y)] \leq 0, \quad x \in V_*(\bar{y}; r) \quad (8.30)$$

and the conclusion of the lemma follows. \square

Lemma 8.8. *Suppose that assumption $A3''$ is fulfilled and D is bounded and satisfies the interior ball condition. Then, for any $\hat{x} \in \partial D$ and any interior ball $B(\bar{y}, r)$ at \hat{x} there exist $C > 0$, depending only on $\gamma_{\partial D}, \|N\|_\infty, \|Q\|_\infty, \|b\|_\infty$, and $a > 0$ depending on the above parameters, radius r and $\underline{c} > 0$ such that*

$$a\eta(x; Cr^{-2}, \bar{y}, r) \leq w_{c,D}(x), \quad x \in \bar{V}_*(\bar{y}; r). \quad (8.31)$$

Proof. Suppose that $r > 0$ is as in the proof of Lemma 8.7. Thanks to Lemma 8.6 we can choose $a_* > 0$, depending on the same parameters as in the statement of the lemma, such that $\varepsilon \geq a_*$, where ε is defined in (8.29). This allows us to conclude (8.31) from (8.30). Recall that $\eta_\varepsilon = \varepsilon\eta$ in (8.28). \square

9. PROOFS OF THE MAIN RESULTS

9.1. Proof of Theorem 3.4.

Proof of part 1): the strong maximum principle. We start with the argument for the strong maximum principle. Recall that $u \in W_{\text{loc}}^{2,p}(D) \cap C_b(\mathbb{R}^d)$. The proof is carried out by a contradiction. We may assume that D is bounded, since otherwise we would consider an intersection of D with an open ball of a sufficiently large radius, that includes a maximal point. Suppose that there exist $P, Q \in D$ such that $u(P) = M \geq 0$ and $u(Q) < M$. In fact, see the proof of Theorem 3.5, p. 61 of [27], we can assume that there exists a ball $B(\bar{y}, 3r/2) \subset D$ such that $u(\hat{x}) = M$ for some $\hat{x} \in \partial B(\bar{y}, r)$ and $u(x) < M$ for all $x \in B(\bar{y}, r)$. By virtue of Lemma 8.1 we can assume that there exists also a function $\eta(\cdot; \beta, \bar{y}, r)$ such that (8.3) holds on $V^* := V^*(\bar{y}; r)$, cf (8.2). We can choose $\epsilon > 0$ sufficiently small so that

$$v(x) := u(x) + \epsilon\eta(x; \beta, \bar{y}, r) < M \quad \text{for } x \in \bar{B}(\bar{y}, r/2). \quad (9.1)$$

We also have $v(x) \leq M$ on $\bar{B}^c(\bar{y}, r)$ (since then $\eta(x; \beta, \bar{y}, r) < 0$). Since both $u(\cdot)$ and $\eta(\cdot)$ are subsolutions to (1.7) on V^* so is $v(\cdot)$. Let $\tilde{v}(x) := v(x) - M$. It is a subsolution to (1.7) on V^* . Therefore by Proposition 3.1 we have

$$\sup_{x \in V^*} \tilde{v}(x) \leq \sup_{x \in (V^*)^c} \tilde{v}(x) < 0,$$

which leads to a contradiction, as $0 = \tilde{v}(\hat{x})$ at $\hat{x} \in V^*$.

Proof of part 2): the Hopf lemma. Concerning the proof of (3.4) we essentially follow the classical proof of the Hopf lemma for a subsolution of a uniformly elliptic equation, as presented in e.g. [27], see Theorem 3.7, p. 65. Since D satisfies the interior ball condition, thanks to Lemma 8.1, we can find an interior ball $B(\bar{y}; r)$ at \hat{x} and $\beta > 0$ such that

$$(A - c)\eta(x; \beta, \bar{y}, r) > 0, \quad x \in \bar{V}_*,$$

where $V_* := V_*(\bar{y}; r)$. Thanks to the already proved strong maximum principle for u and our assumption that $u \not\equiv M$ on D , we know that $u(x) < M$ for $x \in B(\bar{y}; r/2)$. Since $\eta \leq 1$ on $B(\bar{y}; r/2) \subset D_{r/2}$, see (3.3), we have

$$v(x) := u(x) + \epsilon\eta(x; \beta, \bar{y}, r) < M \quad \text{for } x \in B(\bar{y}; r/2). \quad (9.2)$$

with

$$\epsilon := \inf_{y \in D_{r/2}} (M - u(y)). \quad (9.3)$$

We also have $v(x) < M$ for $x \in \bar{B}^c(\bar{y}, r)$ (as then $\eta \leq 0$). Therefore, by Proposition 3.1 applied to v on \bar{V}_* , we conclude that

$$v(x) \leq M = u(\hat{x}) = v(\hat{x}), \quad x \in \bar{V}_*,$$

which yields $\partial_{\mathbf{n}}v(\hat{x}) \geq 0$. Using (9.2) we get

$$\partial_{\mathbf{n}}u(\hat{x}) \geq 2\beta\epsilon r e^{-\beta r^2} > 0.$$

By the construction of $\eta(\cdot; \beta, \bar{y}, r)$, the parameters β, r depend only on $\gamma_{\bar{O}_D}, M_A$. This ends the proof of Theorem 3.4. \square

Remark 9.1. We follow the notation of the proof of Theorem 3.4. From the proof of part 2) of the theorem we conclude that

$$u(\hat{x}) - u(x) \geq \eta(x) \inf_{y \in D_{r/2}} (M - u(y)), \quad x \in \bar{V}_*, \quad (9.4)$$

where, for simplicity, we let $\eta(x) := \eta(x; \beta, \bar{y}, r)$.

Suppose that $r_0 > 0$ satisfies

$$2\beta r_0^2 < 1. \quad (9.5)$$

Recalling the definition of $\eta(\cdot)$ (see (8.1)) we infer from (9.4) that for any $r \in (0, r_0]$:

$$\begin{aligned} \frac{u(\hat{x}) - u(\hat{x} - tn(\hat{x}, \bar{y}))}{t} &\geq \frac{1}{t} \left[\eta(\hat{x} - tn(\hat{x}, \bar{y})) - \eta(\hat{x}) \right] \inf_{y \in D_{r/2}} (M - u(y)) \\ &\geq \partial_{\mathbf{n}}\eta(x)|_{x=\hat{x}-tn(\hat{x}, \bar{y})} \inf_{y \in D_{r/2}} (M - u(y)) \geq \beta r e^{-\beta r^2/4} \inf_{y \in D_{r/2}} (M - u(y)), \quad t \in (0, r/2], \end{aligned}$$

The last inequality follows from the concavity of $t \mapsto \eta(\hat{x} - tn(\hat{x}, \bar{y}))$ for $t \in [0, r/2]$, due to (9.5) (see (8.4)). Summarizing, we have shown that

$$\frac{u(\hat{x}) - u(\hat{x} - tn(\hat{x}, \bar{y}))}{t} \geq \beta r e^{-\beta r^2/4} \inf_{y \in D_{r/2}} (M - u(y)), \quad t \in (0, r/2], \quad (9.6)$$

where $n(\hat{x}, \bar{y}) := (\hat{x} - \bar{y})/|\hat{x} - \bar{y}|$.

The argument leading to (9.4) can be also used to obtain a lower bound on the resolvent operator applied to a non-negative function and we can formulate the following result.

Proposition 9.2. *Assume that a bounded domain D satisfies the uniform interior ball condition. Suppose that non-trivial $f \in B^+(D)$ is such that $R_\alpha^D f \in C_0(\bar{D})$ for some $\alpha \geq 0$. Then, there exists $a > 0$ such that*

$$R_\alpha^D f(x) \geq a \delta_D(x), \quad x \in D. \quad (9.7)$$

Proof. We use the notation from the proof of Theorem 3.4. It can be easily seen, by an application of the Itô formula, that $u = -R_\alpha^D f(x)$ is a weak subsolution of $(A - \alpha)v = 0$ on D . By virtue of (7.5) we have $u(x) < 0$, $x \in D$. We can invoke an analogue of (9.4) that also holds for weak subsolutions and conclude that there exists $a' > 0$ such that for any $\hat{x} \in \partial D$

$$\begin{aligned} R_\alpha^D f(\hat{x} - tn(\hat{x}, \bar{y})) &= -u(\hat{x} - tn(\hat{x}, \bar{y})) \geq a't \inf_{y \in D_{r/2}} (-u(y)) \\ &= a't \inf_{y \in D_{r/2}} R_\alpha^D f(y), \quad t \in (0, 1], \end{aligned} \quad (9.8)$$

where $n(\hat{x}, \bar{y}) := (\hat{x} - \bar{y})/|\hat{x} - \bar{y}|$. This, in turn, implies (9.7). \square

9.2. Proof of Theorem 3.6. Suppose that \hat{x} is as in the statement of Theorem 3.6 and u is a subsolution of (1.7). The conclusion of the theorem is obvious if $u(\hat{x}) < 0$ and follows directly from (3.4), if $u(\hat{x}) = 0$. Assume therefore that $u(\hat{x}) > 0$. Using the Feynman-Kac formula, see Proposition 4.6, we infer that for any $x \in D$

$$u(\hat{x}) - u(x) \geq u(\hat{x}) - \mathbb{E}_x [e_c(\tau_D) u(X_{\tau_D})] \geq u(\hat{x}) (1 - v_{c,D}(x)) = u(\hat{x}) w_{c,D}(x). \quad (9.9)$$

Invoking the lower bound on $w_{c,D}(x)$, contained in Lemma 8.7, we conclude (8.27) for any interior ball $B(\bar{y}, r)$ for D at \hat{x} , with $r \in (0, r_0 \wedge \mathfrak{r}(\hat{x}))$. Therefore, for \mathbf{n} that is the outward unit normal to $\partial B(\bar{y}, r)$ at \hat{x} , we have

$$\underline{\partial}_{\mathbf{n}} u(\hat{x}) \geq -\rho_{c,A,D}(r/2) u(\hat{x}) \partial_{\mathbf{n}} \exp \left\{ -\frac{C}{r^2} |x - \bar{y}|^2 \right\} \Big|_{x=\hat{x}} = \frac{2C}{r} e^{-C} \rho_{c,A,D}(r/2) u(\hat{x}). \quad (9.10)$$

The conclusion of Theorem 3.6 then follows. \square

9.3. Proof of Theorem 3.5. We start with (9.9). Invoking the lower bound on $w_{c,D}(x)$, provided in Lemma 8.8, we can find $B(\bar{y}, r)$ - an interior ball for D at \hat{x} - and $C, a > 0$ as in the statement of the lemma such that (8.31) holds. Therefore, for \mathbf{n} that is the outward unit normal to $\partial B(\bar{y}, r)$ at \hat{x} , we have

$$\underline{\partial}_{\mathbf{n}} u(\hat{x}) \geq -a u(\hat{x}) \partial_{\mathbf{n}} \exp \left\{ -\frac{C}{r^2} |x - \bar{y}|^2 \right\} \Big|_{x=\hat{x}} = 2 \frac{Ca}{r^2} r e^{-C} u(\hat{x}). \quad (9.11)$$

The conclusion of Theorem 3.5 then follows.

9.4. Proof of Theorem 5.2. Using the Feynman-Kac formula (Proposition 4.6) we can write

$$u(\hat{x}) - u(x) \geq (1 - \mathbb{E}_x e_c(\tau_D \wedge t)) u(\hat{x}) + \mathbb{E}_x \left[\int_0^{\tau_D \wedge t} e_c(s) (Au - cu)(X_s) ds \right], \quad t > 0. \quad (9.12)$$

Observe that

$$\left(1 - \mathbb{E}_x e_c(\tau_D \wedge t) \right) u(\hat{x}) \geq (1 - e^{-ct}) P_x(\tau_D > t) u(\hat{x}). \quad (9.13)$$

From (4.11) we have

$$e^{-\lambda_D t} \varphi_D(x) = \mathbb{E}_x [\varphi_D(X_t) \mathbf{1}_{\{t < \tau_D\}}] \leq \|\varphi_D\|_\infty P_x(\tau_D > t). \quad (9.14)$$

Substituting into (9.13) the lower bound on $P_x(\tau_D > t)$ obtained from (9.14) and maximizing over $t > 0$ we conclude that

$$u(\hat{x}) - u(x) \geq \frac{u(\hat{x})\varphi_D(x)}{\|\varphi_D\|_\infty} \cdot \frac{\underline{c}/\lambda_D}{(1 + \underline{c}/\lambda_D)^{1+\lambda_D/\underline{c}}} \geq \frac{u(\hat{x})\varphi_D(x)}{e\|\varphi_D\|_\infty} \cdot \frac{\underline{c}}{\lambda_D + \underline{c}}. \quad (9.15)$$

In the last inequality we have used an elementary bound $(1 + s)^{1/s} \leq e$ valid for all $s > 0$.

Furthermore, since $c(x) \geq 0$, we have $\bar{c} := \|c\|_\infty$ (cf (2.7)) and

$$\begin{aligned} \mathbb{E}_x \left[\int_0^{\tau_D \wedge t} e_c(r)(Au - cu)(X_r) dr \right] &\geq \text{essinf}_D(Au - cu) \mathbb{E}_x \left[\int_0^{\tau_D \wedge t} e^{-\bar{c}r} dr \right] \\ &\geq \frac{1}{\bar{c}}(1 - e^{-\bar{c}t}) \text{essinf}_D(Au - cu) P_x(\tau_D > t). \end{aligned} \quad (9.16)$$

Using again (9.14) to estimate $P_x(\tau_D > t)$ from below and maximizing over $t > 0$ we conclude that

$$\begin{aligned} u(\hat{x}) - u(x) &\geq \frac{\varphi_D(x)}{\|\varphi_D\|_\infty \bar{c}} \cdot \frac{\bar{c}/\lambda_D}{(1 + \bar{c}/\lambda_D)^{1+\lambda_D/\bar{c}}} \text{essinf}_D(Au - cu) \\ &\geq \frac{\varphi_D(x)}{e\|\varphi_D\|_\infty} \cdot \frac{1}{\lambda_D + \bar{c}} \text{essinf}_D(Au - cu). \end{aligned} \quad (9.17)$$

Estimate (5.2) follows easily from (9.15) and (9.17). \square

9.5. Proof of Theorem 5.1. We start with the estimate (9.12). The first term in the right hand side is bounded in the same way as in Section 9.4. To estimate the second term we note that, thanks to finiteness of the moment of the exit time (Proposition 4.4),

$$\lim_{t \rightarrow +\infty} \mathbb{E}_x \left[\int_0^{\tau_D \wedge t} e_c(s)(Au - cu)(X_s) ds \right] \geq R_{\bar{c}}^D((A - c)u). \quad (9.18)$$

By Lemma 7.2 there exist strictly positive, m -a.e. $\bar{\psi}, \bar{\phi} \in B_b^+(D)$ such that

$$R_{\bar{c}}^D f(x) \geq R_{\bar{c}+1}^D \bar{\psi}(x) \int_D R_{\bar{c}+1}^D f \bar{\phi} dx, \quad x \in D, \quad f \in B_b^+(D). \quad (9.19)$$

Thanks to the irreducibility and strong Feller properties of the resolvent, see (7.5) and (7.6), we conclude that $\psi := R_{\bar{c}+1}^D \bar{\psi}$ is strictly positive and continuous in D . By (7.5), the measure

$$\nu(A) = \int_D \bar{\phi} R_{\bar{c}+1}^D \mathbf{1}_A dx, \quad A \in \mathcal{B}(D)$$

is equivalent to m . Thus, $\chi := d\nu/dx$ is strictly positive in D (m -a.e.). Applying (9.19) we obtain

$$R_{\bar{c}}^D((A - c)u)(x) \geq \psi(x) \int_D (Au - cu)\chi dx, \quad x \in D \quad (9.20)$$

and (5.1) follows. \square

9.6. Proof of Theorem 3.3. Let $\varepsilon_0 > 0$ be such that $B(\hat{x}, 2\varepsilon_0) \subset D$. Set $V_\varepsilon = B(\hat{x}, \varepsilon)$. By the Feynman-Kac formula, see Proposition 4.6, for any $\varepsilon \in (0, \varepsilon_0]$,

$$u(\hat{x}) - \mathbb{E}_{\hat{x}}[u(X_{\tau_{V_\varepsilon}})] = -\mathbb{E}_{\hat{x}} \left[\int_0^{\tau_{V_\varepsilon}} (Au)(X_s) ds \right]. \quad (9.21)$$

Using the Ikeda Watanabe formula, see [10, Remark 2.46, page 65], we get

$$\begin{aligned} &\mathbb{E}_{\hat{x}} \left[\sum_{0 < s \leq \tau_D} \mathbf{1}_D(X_{s-}) \mathbf{1}_{\bar{D}^c}(X_s) \right] \\ &= \mathbb{E}_{\hat{x}} \left[\int_{\mathbb{R}^d} \int_0^{\tau_D} \mathbf{1}_D(X_{s-}) \mathbf{1}_{\bar{D}^c}(y) N(X_{s-}, dy - X_{s-}) ds \right] = 0. \end{aligned} \quad (9.22)$$

The last equality follows from the fact that $\mathcal{S}(D) \subset \bar{D}$. This implies that $\mathbb{P}_{\hat{x}}(X_{\tau_{V_\varepsilon}} \in \bar{D}) = 1$.

Since $\hat{x} \in D$ is a global maximal point of u in \bar{D} (by continuity of u), from (9.21), we obtain

$$\mathbb{E}_{\hat{x}} \left[\int_0^{\tau_{V_\varepsilon}} (Au)(X_s) ds \right] \leq 0, \quad \varepsilon \in (0, \varepsilon_0].$$

Combining the above with (7.3) (here $D := V_\varepsilon$ and $\alpha = 0$),

$$\left(\operatorname{ess\,inf}_{V_\varepsilon} Au \right) \int_{V_\varepsilon} r_0^{V_\varepsilon}(\hat{x}, y) dy \leq \mathbb{E}_{\hat{x}} \left[\int_0^{\tau_{V_\varepsilon}} (Au)(X_s) ds \right] \leq 0, \quad \varepsilon \in (0, \varepsilon_0].$$

By (7.5) we have $\int_{V_\varepsilon} r_0^{V_\varepsilon}(\hat{x}, y) dy > 0$. Therefore, $\operatorname{ess\,inf}_{V_\varepsilon} Au \leq 0$, $\varepsilon \in (0, \varepsilon_0]$, which proves the assertion of the theorem, cf (2.1). \square

9.7. Proof of Theorem 5.9. Using Lemma 7.2 we conclude that there exists a pair of functions $\bar{\psi}, \bar{\phi} \in B_b^+(D)$ such that $\bar{\psi} > 0$, $\bar{\phi} > 0$, m a.e. and

$$R_{\bar{c}+1}^D f(x) \geq R_{\bar{c}+2}^D \bar{\psi}(x) \int_D R_{\bar{c}+2}^D f \bar{\phi} dx, \quad x \in D, \quad f \in B_b^+(D). \quad (9.23)$$

Applying this inequality, with f replaced by $R_{\bar{c}}^D f$, and using (7.12) we get

$$R_{\bar{c}}^D f(x) \geq R_{\bar{c}+2}^D \bar{\psi}(x) \int_D R_{\bar{c}}^D f d\nu, \quad x \in D, \quad f \in B_b^+(D), \quad (9.24)$$

where

$$\nu(B) = \int_D \bar{\phi} R_{\bar{c}+2}^D \mathbf{1}_B dx, \quad B \in \mathcal{B}(D). \quad (9.25)$$

By (7.5), $\nu \sim m$. Therefore, we can write $\chi : D \rightarrow (0, +\infty)$ such that

$$\nu(B) = \int_B \chi(x) dx, \quad B \in \mathcal{B}(D), \quad \text{where } \chi(x) = \hat{R}_{\bar{c}+2}^D \bar{\phi} \quad (9.26)$$

and $\hat{R}_{\bar{c}+2}^D$ is defined by (7.7). By (7.5) and (7.6) we infer that $R_{\bar{c}+2}^D \bar{\psi}$ is continuous and strictly positive on D .

In the next step, we show that $R_{\bar{c}}^D f(x)$ in (9.24) can be replaced by w - any \bar{c} -excessive function with respect to $(P_t^D)_{t \geq 0}$, i.e. a non-negative function satisfying

$$P_t^D w(x) \leq e^{ct} w(x), \quad t \geq 0, x \in D \quad (9.27)$$

and

$$\lim_{t \rightarrow 0^+} e^{-ct} P_t^D w(x) = w(x), \quad x \in D. \quad (9.28)$$

More precisely, for any w , that is \bar{c} -excessive, we have

$$w(x) \geq R_{\bar{c}+2}^D \bar{\psi}(x) \int_D w d\nu, \quad x \in D. \quad (9.29)$$

Indeed, from [5, Proposition II.2.6] we conclude that, there exists a sequence of non-negative Borel functions $(f_n)_{n \geq 1}$ such that $(R_{\bar{c}}^D f_n(x))_{n \geq 1}$ is monotone increasing and

$$\lim_{n \rightarrow +\infty} R_{\bar{c}}^D f_n(x) = w(x), \quad x \in D. \quad (9.30)$$

We can write inequality (9.24) for each $R_{\bar{c}}^D f_n(x)$. Estimate (9.29) then follows by an application of the monotone convergence theorem and (9.30).

Coming back to the proof of (5.10), it follows from (9.29), provided we prove that u is \bar{c} -excessive. Indeed for any $V \Subset D$, by (9.24), we can write

$$u(x) \geq R_{\bar{c}}^D u(x) \geq C \int_V \chi u dx \quad (9.31)$$

with $C := \inf_{x \in V} R_{\bar{c}+2}^D \bar{\psi}(x) > 0$ and χ is defined in (9.26).

Since u is a supersolution to (1.7), by Proposition 4.6, we have

$$u(x) \geq \mathbb{E}_x \left[e_c(\tau_D \wedge t) u(X_{\tau_D \wedge t}) \right], \quad t \geq 0, x \in D.$$

Thus, by non-negativity of u ,

$$u(x) \geq e^{-ct} \mathbb{E}_x [u(X_t) \mathbf{1}_{[t < \tau_D]}] = e^{-ct} P_t^D u(x), \quad t \geq 0$$

and (9.27) holds. Equality (9.28) is a consequence of continuity of u , in \bar{D} . This ends the proof of Theorem 5.9. \square

Remark 9.3. Under the assumptions of Theorem 5.4 (resp. Theorem 5.6), using estimate (9.31) and the conclusion of Proposition 9.2, we can prove that in (5.10), $C := a \operatorname{dist}(V, D^c)$ (resp. $\nu := a \delta_D dx$, and $C := \operatorname{dist}(V, D^c)$) for some $a > 0$, depending only on D .

Remark 9.4. It is clear that if \hat{R}_α satisfy the strong Feller condition for some $\alpha > 0$, then $\chi \in C(D)$ and is strictly positive. As a result we conclude then the usual form of the weak Harnack inequality, i.e. that for any $V \Subset D$ we can find a constant $C > 0$ such that for any non-negative supersolution u we have

$$\inf_{x \in V} u(x) \geq C \int_V u dx. \quad (9.32)$$

According to [24, Theorem 1], a sufficient condition for the aforementioned strong Feller property of the adjoint resolvent is that the coefficients of the operator \hat{A}_0 , given by (5.7), satisfy the assumptions A1), A2) and A4).

9.8. Proofs of Theorems 5.4 and 5.6. Invoking the argument from the proofs of Theorem 5.1 and Lemma 7.2 we conclude that there exist strictly positive m -a.e. $\bar{\psi}_0, \bar{\phi}_0 \in B_b^+(D)$ such that

$$u(\hat{x}) - u(x) \geq \frac{\underline{c} \varphi_D(x) u(\hat{x})}{2e \|\varphi_D\|_\infty (\lambda_D + \underline{c})} + R_{\bar{c}+1}^D \bar{\psi}_0(x) \int_D R_{\bar{c}+1}^D (Au - cu) \bar{\phi}_0 dx, \quad x \in D. \quad (9.33)$$

By (7.5), measure

$$\nu(B) = \int_D \bar{\phi}_0 R_{\bar{c}+1}^D \mathbf{1}_B dx, \quad B \in \mathcal{B}(D)$$

is equivalent to m . Thus, $\phi_0 := d\nu/dx$ is strictly positive and

$$\int_D R_{\bar{c}+1}^D (Au - cu) \bar{\phi}_0 dx = \int_D (Au - cu) \phi_0 dx. \quad (9.34)$$

Since D satisfies the exterior ball condition and $R_{\bar{c}+1}^D$ is strongly Feller (see (7.6)), we have $R_{\bar{c}+1}^D \bar{\psi}_0 \in C_0(\bar{D})$, see Section 7.3. Using Proposition 9.2 we conclude that there exists $a_1 > 0$ such that $R_{\bar{c}+1}^D \bar{\psi}_0 \geq a_1 \delta_D$. This finishes the proof of Theorem 5.4.

Under assumptions of Theorem 5.6, we may write

$$\int_D R_{\alpha_*}^D (Au - cu) \bar{\phi}_0 dx = \int_D (Au - cu) \hat{R}_{\alpha_*}^{\hat{c}, D} \bar{\phi}_0 dx, \quad (9.35)$$

where $\alpha_* := (\bar{c} + 1) \vee (\|\hat{c}\|_\infty + 1)$. Invoking again Proposition 9.2 we conclude that there exists $a_2 > 0$ such that $\hat{R}_{\alpha_*}^{\hat{c}, D} \bar{\phi}_0 \geq a_2 \delta_D$ and the conclusion of the theorem follows. \square

9.9. Proof of Theorem 5.8. With no loss of generality we may and shall assume that $\hat{c} \leq 0$, as otherwise we could have considered $A - \alpha$ for a sufficiently large $\alpha > 0$.

Suppose that $p > d$. By [31, Theorem 1.2], the operators $(A, D(A))$ and $(\hat{A}, D(\hat{A}))$ generate C_0 -semigroups of contractions on $C_0(\bar{D})$, where

$$\begin{aligned} D(A) &= [u \in W^{2,p}(D) \cap C_0(\bar{D}) : Au \in C_0(\bar{D})], \\ D(\hat{A}) &= [u \in W^{2,p}(D) \cap C_0(\bar{D}) : \hat{A}u \in C_0(\bar{D})]. \end{aligned}$$

By the standard approximation we conclude that (5.6) holds for $u \in D(A), v \in D(\hat{A})$. Let $(G_\alpha)_{\alpha>0}, (\hat{G}_\alpha)_{\alpha>0}$ denote the semigroup resolvents generated by A, \hat{A} , respectively. Using the martingale problem we conclude that $G_\alpha = R_\alpha^D$ and $\hat{G}_\alpha = \hat{R}_\alpha^{\hat{c}, D}$, where as we recall $R_\alpha^D, \hat{R}_\alpha^{\hat{c}, D}$ are the resolvent corresponding to the probability transition semigroups (4.4) and (5.8). Thus,

$$R_\alpha^D(C_0(\bar{D})) \subset C_0(\bar{D}) \cap W^{2,p}(D) \quad \text{and} \quad \hat{R}_\alpha^{\hat{c}, D}(C_0(\bar{D})) \subset C_0(\bar{D}) \cap W^{2,p}(D)$$

for any $p > d$ and $\alpha \geq 0$. Let $(R_\alpha^D)^*$ denote the adjoint operator to R_α^D in $L^2(D; m)$. Then for any $w \in C_0(\overline{D})$, there exists $\eta \in D(A)$ such that $(-A + \alpha)\eta = w$. By using (5.6), for any $u \in C_0(\overline{D})$

$$\begin{aligned} \int_D w \hat{R}_\alpha^{\hat{c}, D} u \, dx &= \int_D (-A + \alpha)\eta \hat{R}_\alpha^{\hat{c}, D} u \, dx \\ &= \int_D \eta(-\hat{A} + \alpha)\hat{R}_\alpha^{\hat{c}, D} u \, dx = \int_D u\eta \, dx. \end{aligned}$$

On the other hand

$$\begin{aligned} \int_D u\eta \, dx &= \int_D u R_\alpha^D (-A + \alpha)\eta \, dx = \int_D (R_\alpha^D)^* u (-A + \alpha)\eta \, dx \\ &= \int_D (R_\alpha^D)^* u w \, dx = \int_D u R_\alpha^D w \, dx. \end{aligned}$$

Since $u, w \in C_0(\overline{D})$ were arbitrary, we get (5.9). \square

Acknowledgements. T. Klimsiak is supported by Polish National Science Centre: Grant No. 2017/25/B/ST1/00878. T. Komorowski acknowledges the support of the Polish National Science Centre: Grant No. 2016/23/B/ST1/00492. Both authors wish to express their gratitude to the referee for very careful reading of the manuscript and valuable remarks that lead to significant improvement of the presentation.

REFERENCES

- [1] Anulova, S.V., Pragarauskas, H.: *Weak Markov solutions of stochastic equations*. *Lithuanian Math. J.* **17**, (1977), 141–155
- [2] Beygmohammadi, M.; Sweers, G. *Hopf's boundary point lemma, the Krein-Rutman theorem and a special domain*. *Positivity* 18 (2014), no. 1, 81-94.
- [3] Billingsley, P., *Convergence of probability measures*. Second edition. Wiley Series in Probability and Statistics: Probability and Statistics. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1999.
- [4] J. Bliedner, W. Hansen, *Potential Theory*, Springer, Berlin, 1986.
- [5] Blumenthal, R. M.; Gettoor, R. K. *Markov processes and potential theory*. Pure and Applied Mathematics, Vol. 29 ,Academic Press, New York-London 1968.
- [6] Berestycki, H.; Nirenberg, L.; Varadhan, S. R. S. *The principal eigenvalue and maximum principle for second-order elliptic operators in general domains*. *Comm. Pure Appl. Math.* 47 (1994), no. 1, 47–92.
- [7] A. Bobrowski, T. Komorowski, *A quantitative Hopf-type maximum principle for subsolutions of elliptic PDEs*. *Discrete Contin. Dyn. Syst. Ser. S*. Volume **13**, Number12, (2020), 3495-3502 doi:10.3934/dcdss.2020248
- [8] Bony, J.-M. Problème de Dirichlet et semi-groupe fortement fellerien associés a un opérateur integro-differentiel. *C. R. Acad. Sci. Paris Ser. A-B* 265 (1967), A361-A364.
- [9] Bony, J.-M.; Courrége, P.; Priouret, P.: *Semi-groupes de Feller sur une variété à bord compacte et problèmes aux limites intégro-différentiels du second ordre donnant lieu au principe du maximum*. (French) *Ann. Inst. Fourier (Grenoble)* 18 (1968), no. fasc., fasc. 2, 369-521 (1969).
- [10] Böttcher, B., Schilling, R., Wang, J.: *Lévy Matters III. Lévy-Type Processes: Construction, Approximation and Sample Path Properties*. Lecture Notes in Mathematics, vol. 2099. Springer, Cham (2013)
- [11] Brezis, H; Cabré, X. *Some simple nonlinear PDE's without solutions*. *Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat.* (8) 1 (1998), no. 2, 223-262.
- [12] Chung, K. L.; Zhao, Z. X. *From Brownian motion to Schrödinger's equation*. Grundlehren der Mathematischen Wissenschaften, 312. Springer-Verlag, Berlin, 1995.
- [13] Dembinski, V.: *Regular points, transversal sets, and Dirichlet problem for standard processes*. *Z. Wahrsch. Verw. Gebiete* **66** (1984), 507–527.
- [14] Diaz, J. I.; Morel, J.-M.; Oswald, L.: *An elliptic equation with singular nonlinearity*. *Comm. Partial Differential Equations* 12 (1987), no. 12, 1333-1344.
- [15] Down, D.; Meyn, S. P.; Tweedie, R. L. *Exponential and uniform ergodicity of Markov processes*. *Ann. Probab.* 23 (1995), no. 4, 1671-1691.
- [16] El Berdan, N., Diaz, J.I., Rakotoson, J. M.: *The uniform Hopf inequality for discontinuous coefficients and optimal regularity in BMO for singular problems*. *J. Math. Anal. Appl.* 437 (2016), no. 1, 350-379.
- [17] Ethier, S. N.; Kurtz, T. G., *Markov processes. Characterization and convergence*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, 1986.

- [18] Foondun, M.: *Harmonic Functions for a Class of Integro-differential operators*. Potential Analysis, **31**, 21–44 (2009)
- [19] Gilbarg, D.; Trudinger, N. S. *Elliptic partial differential equations of second order*. Reprint of the 1998 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2001.
- [20] Han, Qing; Lin, Fanghua: *Elliptic partial differential equations*. Second edition. Courant Lecture Notes in Mathematics, 1. CIMS, New York; American Mathematical Society, Providence, RI, 2011.
- [21] T. Klimsiak, T. Komorowski *On the maximum principle for integro-differential operators of elliptic type - probabilistic approach*, arXiv:2102.08890 (2021).
- [22] Lepeltier, J.-P.; Marchal, B. *Problème des martingales et équations différentielles stochastiques associées à un opérateur intégral-différentiel*. Ann. Inst. H. Poincaré Sect. B (N.S.) 12 (1976), no. 1, 43–103.
- [23] Mikulevičius, R.; Pragarauskas, H. *Green measures of Itô processes*. (Russian) Litovsk. Mat. Sb. 27 (1987), no. 4, 711-723. English translation: Lithuanian Math. J. 27 (1987), no. 4, 325-334.
- [24] Mikulevičius, R.; Pragarauskas, H. *On Hölder continuity of solutions of certain integro-differential equations*. Ann. Acad. Sci. Fenn. Ser. A I Math. 13 (1988), no. 2, 231–238.
- [25] Nummelin, E.: *General Irreducible Markov Chains and Non-Negative Operator*. Cambridge Univ. Press (1984).
- [26] Orsina, L.; Ponce, A. C. *Hopf potentials for the Schrödinger operator*. Anal. PDE 11 (2018), no. 8, 2015-2047
- [27] Protter, M. H.; Weinberger, H. F. *Maximum principles in differential equations*. Corrected reprint of the 1967 original. Springer-Verlag, New York, 1984.
- [28] H. H. Schaefer, *Banach lattices and positive operators*, Springer-Verlag, New York, 1974.
- [29] Stroock, D.W.: *Diffusion processes associated with Lévy generators*. Probab. Theory Relat. Fields **32**, 209–244 (1975)
- [30] Taira, K.: *Boundary Value Problems and Markov Processes*. Lecture Notes in Math. 1499, 3-d edition, Springer Verlag (2020).
- [31] Taira, K. *Dirichlet problems with discontinuous coefficients and Feller semigroups*. Rendiconti del Circolo Matematico di Palermo Series 2 (2020) 69:287-323
- [32] Taira, K. *Ventcel boundary value problems for elliptic Waldenfels operators*. Boll. Unione Mat. Ital. 13 (2020), no. 2, 213-256
- [33] Taira, K. *Semigroups, boundary value problems and Markov processes*. Second edition. Springer Monographs in Mathematics. Springer, Heidelberg, 2014.

(Tomasz Klimsiak) INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, UL. ŚNIADECKICH 8, 00-656 WARSAW, POLAND, AND FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, NICOLAUS COPERNICUS UNIVERSITY, CHOPINA 12/18, 87-100 TORUN, POLAND, E-MAIL :tomas@mat.umk.pl

(Tomasz Komorowski) INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, UL. ŚNIADECKICH 8, 00-636 WARSAW, POLAND E-MAIL: tkomorowski@impan.pl