

Twisted boundary condition and Lieb-Schultz-Mattis ingappability for discrete symmetries

Yuan Yao¹ and Masaki Oshikawa^{1,2,3}

¹*Institute for Solid State Physics, The University of Tokyo, Kashiwa, Chiba 277-8581, Japan*

²*Kavli Institute for the Physics and Mathematics of the Universe (WPI),*

The University of Tokyo, Kashiwa, Chiba 277-8583, Japan

³*Trans-scale Quantum Science Institute, University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan*

(Dated: June 29, 2022)

We discuss quantum many-body systems with lattice translation and discrete on-site symmetries. We point out that, under a boundary condition twisted by a symmetry operation, there is an exact degeneracy of ground states if the unit cell forms a projective representation of the on-site discrete symmetry. Based on the quantum transfer matrix formalism, we show that, if the system is gapped, the ground-state degeneracy under the twisted boundary condition also implies a ground-state (quasi-)degeneracy under the periodic boundary conditions. This amounts to the Lieb-Schultz-Mattis type ingappability due to the on-site discrete symmetry, which was proposed and then proved in one dimension recently, in arbitrary dimensions.

Introduction.— In condensed matter and statistical physics, identifying various quantum phases is a central, but in general difficult task due to complicated interactions and strong correlations. Symmetries often play an important role in quantifying the phase. One of the most notable principles in quantum many-body systems is Lieb-Schultz-Mattis (LSM) theorem [1] and its generalizations [2–5], which state an “ingappability” — either the presence of gapless excitations above ground state(s) or a ground-state degeneracy in the large system size limit — of the systems possessing U(1) and translation symmetries with a fractional filling. Recently, stronger constraints due to higher symmetries such as the SU(2) [6, 7] and the SU(N) [8] symmetries are also investigated.

In these examples of the LSM-type theorems, a continuous symmetry of the Hamiltonian appears essential. However, recently, generalizations of the LSM theorem have been made for systems with only discrete symmetries [9–12]. This is so far limited to one dimension where the ground state of a generic quantum many-body Hamiltonian can be described by Matrix Product State (MPS) formulation [10] or satisfy a separability condition [11, 12].

On the other hand, from the field-theory perspective, in terms of quantum anomaly, ingappabilities related to discrete symmetries in two-dimensional systems are proposed in [13]. However, an explicit lattice approach is still lacking. Thus, it is of both practical and conceptual importances to study universal LSM-type ingappability constraints imposed by discrete symmetries in arbitrary dimensions

In this Letter, we discuss a generalization of the LSM ingappability for quantum many-body systems with discrete symmetries, in arbitrary dimension. The key notion is the robustness of the system to a twisting of the boundary condition by a (discrete) symmetry operation.

Translation invariant SU(2) spin chains with $\mathbb{Z}_2 \times$

\mathbb{Z}_2 symmetry under a symmetry-twisted boundary condition.—

To illustrate our point, let us first consider (1+1)d SU(2) spin chains as an example. The Hamiltonian is assumed to have the global (on-site) $\mathbb{Z}_2 \times \mathbb{Z}_2$ and the lattice translation symmetries. Considering a finite chain of length L with the periodic boundary condition (PBC), the symmetry generators are given by $R_{x,z}^\pi$ and T , where

$$R_x^\pi = \exp\left(i\pi \sum_{k=1}^L S_k^x\right); \quad R_z^\pi = \exp\left(i\pi \sum_{k=1}^L S_k^z\right), \quad (1)$$

$$T^{-1}\vec{S}_i T = \vec{S}_{i+1}, \quad \text{if } 1 \leq i < L; \quad T^{-1}\vec{S}_L T = \vec{S}_1, \quad (2)$$

where $\vec{S}_i \equiv [S_i^x, S_i^y, S_i^z]$ is the spin- s operator at the i -th site. A typical example of such a system is the XYZ chain:

$$\mathcal{H}_{XYZ} = \sum_i J_X S_i^x S_{i+1}^x + J_Y S_i^y S_{i+1}^y + J_Z S_i^z S_{i+1}^z. \quad (3)$$

Instead of PBC, now we introduce the following boundary condition, twisted by the symmetry generator R_z^π :

$$\vec{S}_{L+k} \equiv (R_z^\pi)^{-1} \vec{S}_k R_z^\pi, \quad k = 1, 2, \dots, L, \quad (4)$$

which is periodic up to a π -rotation around the z -axis. This symmetry-twisted boundary condition (STBC) can be implemented by introducing a “twisted” bond in any location, say between sites L and 1:

$$\begin{aligned} \mathcal{H}_{XYZ}^{\text{STBC}} = & \sum_{i=1}^{L-1} (J_X S_i^x S_{i+1}^x + J_Y S_i^y S_{i+1}^y + J_Z S_i^z S_{i+1}^z) \\ & - J_X S_L^x S_1^x - J_Y S_L^y S_1^y + J_Z S_L^z S_1^z. \end{aligned} \quad (5)$$

This Hamiltonian still possesses the global $\mathbb{Z}_2 \times \mathbb{Z}_2$ with respect to the same symmetry generators $R_{x,z}^\pi$. On the other hand, the lattice translation must be followed by

a “gauge transformation” to leave the Hamiltonian invariant. Namely, the symmetry generator of the lattice translation should be modified to

$$\tilde{T} = T e^{i\pi S_1^z}. \quad (6)$$

Now, among the three symmetry generators of the STBC Hamiltonian, the two satisfy the nontrivial commutation relation

$$R_x^\pi \tilde{T} = (-1)^{2s} \tilde{T} R_x^\pi. \quad (7)$$

The phase factor $(-1)^{2s}$, which essentially comes from

$$e^{\pi S_1^x} e^{\pi S_1^z} = (-1)^{2s} e^{\pi S_1^z} e^{\pi S_1^x}, \quad (8)$$

is nontrivial (-1) for an half-odd-integer spin s , while it is trivial (1) for an integer spin s . As a direct consequence of the nontrivial commutation relation, for a half-odd-integer spin s , all the eigenstates of the STBC Hamiltonian are doubly degenerate. In particular, the ground states are doubly degenerate. This is an exact and rigorous statement.

Generalization to higher dimensions and physical consequences.— This analysis can be extended to translation-invariant and $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetric quantum spin systems in higher dimensions. If we consider the system on a hypercubic lattice of the size $L_1 \times L_2 \times \dots \times L_d$, we can twist the boundary condition along 1 direction by the symmetry generator R_z^π . Such a system is invariant under the modified lattice translation operator

$$\tilde{T}_1 = T_1 \exp \left(i\pi \sum_{\vec{r}|r_1=1} S_{\vec{r}}^z \right), \quad (9)$$

where the summation is taken over the sites \vec{r} which has the first component $r_1 = 1$. Naively imposing periodic boundary conditions along the remaining 2 to d directions, we then find the commutation relation $R_x^\pi \tilde{T}_1 = (-1)^{2sA} \tilde{T}_1 R_x^\pi$, where $A = L_2 \times L_3 \times \dots \times L_d$ is the number of sites per the “cross section” perpendicular to the twist. All the eigenstates, including the ground states, must be exactly doubly degenerate under the STBC, if the spin s is half-odd-integer and A is odd (namely, all L_2, L_3, \dots, L_d are odd). This restriction on the system size can be removed by adopting the “tilted boundary condition” where the lattice translations are also included into STBC as follows [14]. Before twisting the boundary condition by R_z^π along the 1 direction, we tilt the lattice geometry such that completion of the $(i > 1)$ -th cycle of the torus effectively increases by \hat{x}_{i-1} , i.e.

$$(T_{i>1})^{-L_i} \vec{S}_{\vec{r}}(T_i)^{L_i} = \vec{S}_{\vec{r}+\hat{x}_{i-1}} \quad (10)$$

(\hat{x}_i the unit vector along the i -th direction) as sketched in FIG. (1). Then we obtain a nontrivial commutation relation

$$R_x^\pi \tilde{T}_1 = (-1)^{2s} \tilde{T}_1 R_x^\pi, \quad (11)$$




FIG. 1. W -twisted tilted boundary condition sketched for $d = 2$ with $W = R_z^\pi$ for $SU(2)$ spins and $W = W_N$ for $SU(N)$.

thereby the exact double degeneracy when s is half-odd-integer. It is also straightforward to generalize this to a more complicated systems with multiple spins within the unit cell. In this case, the exact double degeneracy holds if the total spin per unit cell is half-odd-integer, under the appropriate STBC.

Obviously, however, the above argument only applies to the system under the STBC. The question is then: is this degeneracy an unphysical artifact specific to the STBC, or is it of a physical significance? We argue that the latter must be the case. In fact, if the system is completely disordered without any order (conventional or topological), the system must be insensitive to the symmetry-twist of the boundary condition, and thus the ground-state degeneracy should not appear under STBC. The ground-state degeneracy under the STBC then implies some kind of ingappability (gapless excitations or degenerate ground states below a non-vanishing gap). Namely, we assert the following proposition.

Proposition: *When a quantum many-body system with the lattice translation invariance and a global (continuous or discrete) symmetry has a gapped spectrum with a unique ground state under the PBC, the system should also have a unique gapped ground state under the STBC.*

This proposition, together with the exact double degeneracy under the STBC implies that, a translation-invariant and $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetric quantum spin system with a half-odd-integer spin per unit cell cannot have a unique gapped ground state under the PBC as well. This amounts to be the generalization of the LSM theorem to the discrete $\mathbb{Z}_2 \times \mathbb{Z}_2$. For one dimension, this was already proved within the MPS framework in Ref. [10] and rigorously in Ref. [12], so the present argument gives an alternative physical interpretation. On the other hand, the present result naturally leads to the same statement in arbitrary dimensions, which has not been shown before.

When applied to a continuous $U(1)$ (on-site) symmetry, the STBC corresponds to insertion of an Aharonov-Bohm flux. This also shows that the present approach is a natural generalization of the LSM theorem from the $U(1)$ to discrete symmetries. More precisely, the above proposition applied to the case of $U(1)$ symmetry implies

that, if the gapped unique ground state is unique under the PBC, the ground state is gapped and unique for any value of Aharonov-Bohm flux inserted. This corresponds to the conjectured insensitivity of the excitation gap to the Aharonov-Bohm flux [4, 15] which leads to the LSM ingappability for systems with the U(1) symmetry. However, in this approach, we cannot constrain the number of the degenerate ground states (other than that it must be greater than 1).

Robustness of the spectrum to the symmetry-twist.— Now let us show that the proposition naturally follows from mild assumptions. We consider the partition function of the finite-size system of the size $L_1 \times L_2 \times \dots \times L_d$ with the PBC, at the inverse temperature β . Let us assume that it has a unique ground state below the gap Δ .

When the spectrum is gapped, Hastings and Koma proved that the correlation function of any local operator decays exponentially [16]. Since the system has a finite length scale (correlation length) ξ , we can expect the ground-state energy of the finite-size system is given in terms of the ground-state energy density ϵ_0 in the thermodynamic limit, with exponentially small corrections:

$$E_0 = \epsilon_0 V + O(e^{-L/\xi}), \quad (12)$$

where $L = \min(L_1, L_2, \dots, L_d)$, and $V = L_1 L_2 \dots L_d$ is the volume of the system. The correction might actually be of the form $O(L^\chi e^{-L/\xi})$ for some constant exponent χ . However, we can cover such a case with $O(e^{-L/\xi})$ by taking a larger value of ξ . Furthermore, the partition function $Z = \text{Tr} e^{-\beta \mathcal{H}}$ is expected to obey

$$\log Z(\beta, \{L_\alpha\}) \sim -\beta V \epsilon_0 + O(\beta e^{-L/\xi}) + O(V e^{-\beta \Delta}). \quad (13)$$

The first term $O(\beta e^{-L/\xi})$ simply comes from the exponentially small correction (12) to the ground-state energy. The second term $O(V e^{-\beta \Delta})$ represents the contribution from excited states. Here we assume that such a contribution is exponentially small per volume. Equation (13) implies that the entropy in the thermodynamic limit and zero-temperature limit taken simultaneously with $\beta \sim (L)^z$ for any “dynamical exponent” $z > 0$ is

$$S_0 = \lim_{\beta=(L)^z \rightarrow \infty} \left[\log Z(\beta, \{L_\alpha\}) - \beta \frac{\partial}{\partial \beta} \log Z(\beta, \{L_\alpha\}) \right] = 0. \quad (14)$$

Now let us consider the same system in terms of the quantum transfer matrix (QTM) [17] (For a recent application of QTM, see for example Ref. [18]). The partition function of the quantum system can be identified with the partition function of a classical statistical system in $d + 1$ dimensions by the Lie-Trotter-Suzuki decomposition [19]. QTM is the transfer matrix of this classical statistical system along a spatial direction [20]. Let $e^{-\tilde{H}}$

be the QTM along 1 direction. Using this, the partition function can be written as $Z = \text{Tr} e^{-L_1 \tilde{H}}$.

The asymptotic expression (13) implies that, the spectrum of the “QTM Hamiltonian” $\tilde{\mathcal{H}}$ must be also gapped, with $1/\xi$ being the “excitation gap”. The excitation gap Δ of the original Hamiltonian \mathcal{H} now corresponds to the correlation length in the imaginary time direction, which plays the role of one of the spatial coordinates. Eq. (13) and its consequence (14) implies that the “ground state” $|\tilde{0}\rangle$ of the QTM Hamiltonian $\tilde{\mathcal{H}}$ is also unique.

In the QTM formalism, the STBC is represented by an application of the symmetry operation \tilde{U}_s on the “state” at a fixed “time” (the coordinate r_1). The STBC corresponding to the inverse symmetry operation is represented by \tilde{U}_s^\dagger . Twisting the boundary condition by a symmetry operation in one place and its inverse elsewhere can be eliminated by a gauge transformation, leaving the partition function identical to the PBC. This implies the symmetry of the QTM

$$[\tilde{U}_s, \tilde{H}] = 0, \quad (15)$$

and the unitarity of the operator

$$\tilde{U}_s^\dagger \tilde{U}_s = \mathbb{1}. \quad (16)$$

Now let us consider the partition function of the system under the STBC. In the QTM formalism, it is given as

$$Z_s = \text{Tr} \left(\tilde{U}_s e^{-L_1 \tilde{H}} \right). \quad (17)$$

As in the case of the PBC, the partition function in the thermodynamic and zero-temperature limit is dominated by the “ground state” of the QTM. The symmetry of the QTM implies that the unique “ground state” of the QTM is an eigenstate of the operator \tilde{U}_s . Let $e^{i\theta_0}$ be the eigenvalue of \tilde{U}_s of the QTM “ground state” $|\tilde{0}\rangle$. Then

$$Z_s \sim e^{i\theta_0} e^{-\beta V \epsilon_0}, \quad (18)$$

in the thermodynamic/zero-temperature limit, up to exponentially small corrections. Since the Hamiltonian is Hermitian even under the STBC, the partition function must be real and positive, implying $\theta_0 = 0$. Thus STBC partition function is essentially identical to the PBC partition function up to exponentially small corrections, implying that the system has a unique gapped ground state also for STBC (since the entropy in the limit $\beta = (L)^z \rightarrow \infty$ is zero as in the case of PBC (14)). This is nothing but our proposition, and the ingappability of the translation-invariant and $\mathbb{Z}_2 \times \mathbb{Z}_2$ symmetric spin system with a half-odd-integer spin per unit cell follows.

General LSM ingappabilities and applications.— We can generalize the ingappabilities above to a more general class of unitary onsite symmetry groups. The projective nature of the onsite symmetry characterized by the commutator (8) directly results in the ingappabilities. $\mathbb{Z}_2 \times \mathbb{Z}_2$ can be generalized to an arbitrary unitary

symmetry G that can be centrally extended to G' , e.g., $G = G'/H$ with H a center subgroup of G' . Then each unit cell consists of tensor products of G' -“spins”. In the $SU(2)$ case, G' is arbitrary finite products of R_x^π and R_z^π treated as a subgroup of $SU(2)$ (rather than $PSU(2)$) and $H = \{1, (R_z^\pi)^2\}$.

Generalizing (R_x^π, R_z^π) and their relation (8) per unit cell, G' can possess an almost commuting pair [21–23] $v_{\vec{r}}$ and $w_{\vec{r}}$ satisfying:

$$v_{\vec{r}}w_{\vec{r}} = h_{V,W}w_{\vec{r}}v_{\vec{r}}, \quad (19)$$

where $h_{V,W} \in H$ is an \vec{r} -independent $U(1)$ phase due to the translation symmetry and Schur’s Lemma [24]. Parallel to the former discussions, we can twist and tilt the boundary condition by $W = \prod_{\vec{r}} w_{\vec{r}}$ and translations as in FIG. (1) and calculate the commutator between the modified translation

$$\mathcal{T}_1 = T_1 \prod_{\vec{r}|r_1=1} w_{\vec{r}}, \quad (20)$$

and $V = \prod_{\vec{r}} v_{\vec{r}}$ as

$$V\mathcal{T}_1 = h_{V,W}\mathcal{T}_1V. \quad (21)$$

Therefore, a nontrivial $h_{V,W} = \exp(i2\pi p/q) \neq 1$ (with p and q coprime) results in an exact q -fold degeneracy under STBC, which further implies an ingappability of the system under PBC, i.e. a unique gapped ground state is forbidden, by the Proposition of the spectrum insensitivity to the symmetry-twists argued by the QTM formalism. It can be summarized as

If a Hamiltonian on a d -dimensional periodic lattice possesses $G = G'/H$ symmetry and translational symmetries, a unique gapped ground state is strictly forbidden if there is a nontrivial almost commuting pair per unit cell, i.e. $h_{V,W} \neq 1$ in Eq. (19).

Since such a general result and Eq. (21) are abstract, a further example beyond the simplest $SU(2)$ case might be helpful. As a natural generalization of $SU(2)$ spins, the systems with the following $SU(N)$ degrees of freedom satisfy the $\mathfrak{su}(N)$ algebra:

$$[\mathcal{S}_{\alpha,\vec{r}}^\beta, \mathcal{S}_{\sigma,\vec{r}}^\rho] = \delta_{\vec{r},\vec{r}'} [\delta_\sigma^\beta \mathcal{S}_{\alpha,\vec{r}}^\rho - \delta_\alpha^\rho \mathcal{S}_{\sigma,\vec{r}}^\beta]. \quad (22)$$

Here $\{\mathcal{S}_{\beta,\vec{r}}^\alpha\}$ defined on the unit cell at \vec{r} are labelled by $1 \leq \alpha \leq N$ and $1 \leq \beta \leq N$, with a constraint $\sum_{\alpha=1}^N \mathcal{S}_\alpha^\alpha = 0$ [25, 26]. Additionally, the full global $SU(N)$ symmetry acts on the generators by $u \in SU(N)$:

$$\mathcal{S}_\beta^\alpha \rightarrow (u)_{\beta,\gamma} \mathcal{S}_\gamma^\rho (u^\dagger)_{\rho,\alpha} \quad (23)$$

which does nothing if the $SU(N)$ matrix u is in the \mathbb{Z}_N center of $SU(N)$: $u \propto 1$. Thus, the physical symmetry is only $PSU(N) \cong SU(N)/\mathbb{Z}_N$ if the fundamental degrees of freedom consist of $\{\mathcal{S}_\beta^\alpha\}$ only.

Experimental realizations of the $SU(N)$ symmetry have been proposed for ultracold atoms on optical lattices [27–34]. Let us consider a translation-invariant $SU(N)$ -spin Hamiltonian \mathcal{H} possessing $\mathbb{Z}_N \times \mathbb{Z}_N \subset PSU(N)$, which is generated by:

$$V_N = \prod_{\vec{r}} v_{N,\vec{r}}; \quad W_N = \prod_{\vec{r}} w_{N,\vec{r}}, \quad (24)$$

of which an explicit form together with a $SU(N)$ -analog of the XYZ model can be found in [35]. V_N (also known as the shift symmetry [36, 37]) and W_N reduce to R_x^π and R_z^π , respectively, when $N = 2$, but we only focus on the following essential relation

$$v_{N,\vec{r}}w_{N,\vec{r}} = \exp\left(i2\pi \frac{b}{N}\right) w_{N,\vec{r}}v_{N,\vec{r}}, \quad (25)$$

where b is the total number of Young-tableau (YT) boxes [24, 35] per unit cell as a generalization of $2s$ of a $SU(2)$ spin- s chain. Then, taking $v_{\vec{r}} = v_{N,\vec{r}}$ and $w_{\vec{r}} = w_{N,\vec{r}}$ in the general setting (20,21), we obtain

$$V\mathcal{T}_1 = \exp(i2\pi b/N)\mathcal{T}_1V, \quad (26)$$

which implies that the spin system has an exact $N/\text{gcd}(b, N)$ -fold degeneracy under STBC (“gcd”: the positive greatest common divisor) and, by the insensitivity of ingappability to boundary conditions, it cannot possess a unique gapped ground state under PBC if the total YT boxes per unit is indivisible by N .

Conclusions and conjectures.— We proposed a symmetry-twisted boundary condition in arbitrary dimensions, under which the energy spectrum of translation-symmetric lattice system is exactly degenerate due to a projective representation per unit cell characterized by almost commuting pairs. We further argue that such an exact degeneracy implies the ingappability of ground states under PBC by a proposition based on QTM formalism of partition functions. Our argument relies on hypotheses (12) and (13) on the scaling of the ground-state energy and free energy density of a gapped system in the low-temperature and thermodynamic limit. While we believe them to be reasonable for systems with short-range interactions only, they are not rigorously proved. We hope that the present work will stimulate further investigations in rigorous mathematics.

On the other hand, this QTM-type argument does not make a full use of the the number of degenerate ground states under STBC because only the case of a unique gapped ground state is excluded based on Eq. (18). Nevertheless, it is still reasonable to conjecture that the same number of the degenerate ground states is also held by the gapped ground states under PBC.

Furthermore, since our treatment of symmetry twisting in addition to the QTM argument (18) strongly depends on the unitarity of the symmetry, it cannot be directly applied to the anti-unitary symmetries, e.g. time reversal.

In contrast to Eq. (24), there is no general way to factorize an anti-unitary transformation in a local way due to an additional complex conjugation. Nevertheless, in one dimension, the description of matrix product states allow the complex conjugation to be assigned locally [10, 38], where we expect our method is still applicable. Higher-dimensional cases with anti-unitary symmetries are of future interest.

Both authors thank Hosho Katsura for useful discussions. Y. Y. was supported by JSPS fellowship. This work was supported in part by MEXT/JSPS KAKENHI Grant Nos. JP19J13783 (Y. Y.), JP17H06462 (M. O.), JP19H01808 (M. O.), and JST CREST Grant No. JP-MJCR19T2 (M. O.).

-
- [1] E. Lieb, T. Schultz, and D. Mattis, *Ann. Phys.* **16**, 407 (1961).
- [2] I. Affleck and E. H. Lieb, *Lett. Math. Phys.* **12**, 57 (1986).
- [3] M. Oshikawa, M. Yamanaka, and I. Affleck, *Phys. Rev. Lett.* **78**, 1984 (1997).
- [4] M. Oshikawa, *Phys. Rev. Lett.* **84**, 1535 (2000).
- [5] M. B. Hastings, *Phys. Rev. B* **69**, 104431 (2004).
- [6] S. C. Furuya and M. Oshikawa, *Phys. Rev. Lett.* **118**, 021601 (2017).
- [7] M. A. Metlitski and R. Thorngren, *Phys. Rev. B* **98**, 085140 (2018).
- [8] Y. Yao, C.-T. Hsieh, and M. Oshikawa, *Phys. Rev. Lett.* **123**, 180201 (2019).
- [9] X. Chen, Z.-C. Gu, and X.-G. Wen, *Phys. Rev. B* **83**, 035107 (2011).
- [10] H. Watanabe, H. C. Po, A. Vishwanath, and M. Zaletel, *Proc. Natl. Acad. Sci. USA* **112**, 14551 (2015).
- [11] Y. Ogata and H. Tasaki, arXiv preprint arXiv:1808.08740 (2018).
- [12] Y. Ogata, Y. Tachikawa, and H. Tasaki, arXiv preprint arXiv:2004.06458 (2020).
- [13] M. Cheng, M. Zaletel, M. Barkeshli, A. Vishwanath, and P. Bonderson, *Phys. Rev. X* **6**, 041068 (2016).
- [14] Y. Yao and M. Oshikawa, *Phys. Rev. X* **10**, 031008 (2020).
- [15] H. Watanabe, *Phys. Rev. B* **98**, 155137 (2018).
- [16] M. B. Hastings and T. Koma, *Commun. Math. Phys.* **265**, 781 (2006).
- [17] H. Betsuyaku, *Phys. Rev. Lett.* **53**, 629 (1984).
- [18] L. Piroli, B. Pozsgay, and E. Vernier, *Journal of Statistical Mechanics: Theory and Experiment* **2017**, 023106 (2017).
- [19] M. Suzuki, *Progr. Theor. Phys.* **56**, 1454 (1976).
- [20] The mapping to the classical statistical system is exact only in the limit of an infinite Trotter number. Here we assume that the QTM is well-defined in this limit.
- [21] C. Schweigert, *Nucl. Phys. B* **492**, 743 (1997).
- [22] A. Borel, R. Friedman, J. W. Morgan, and J. W. Morgan, *Almost commuting elements in compact Lie groups* (American Mathematical Soc., 2002).
- [23] V. G. Kac and A. V. Smilga, “Vacuum structure in supersymmetric yang–mills theories with any gauge group,” in *The Many Faces of the Superworld: Yuri Golfand Memorial Volume* (World Scientific, 2000) pp. 185–234.
- [24] H. Georgi, *Lie Algebras In Particle Physics* (Boca Raton: CRC Press, 2000).
- [25] I. Affleck and F. D. M. Haldane, *Phys. Rev. B* **36**, 5291 (1987).
- [26] I. Affleck, *Nucl. Phys. B* **305**, 582 (1988).
- [27] C. Wu, J.-P. Hu, and S.-C. Zhang, *Phys. Rev. Lett.* **91**, 186402 (2003).
- [28] C. Honerkamp and W. Hofstetter, *Phys. Rev. Lett.* **92**, 170403 (2004).
- [29] M. A. Cazalilla, A. Ho, and M. Ueda, *New J. Phys.* **11** (2009).
- [30] A. V. Gorshkov, M. Hermele, V. Gurarie, C. Xu, P. S. Julienne, J. Ye, P. Zoller, E. Demler, M. D. Lukin, and A. Rey, *Nature physics* **6**, 289 (2010).
- [31] S. Taie, R. Yamazaki, S. Sugawa, and Y. Takahashi, *Nature Physics* **8**, 825 (2012).
- [32] G. Pagano, M. Mancini, G. Cappellini, P. Lombardi, F. Schäfer, H. Hu, X.-J. Liu, J. Catani, C. Sias, and M. Inguscio, *Nature Physics* **10**, 198 (2014).
- [33] F. Scazza, C. Hofrichter, M. Höfer, P. De Groot, I. Bloch, and S. Fölling, *Nature Physics* **10**, 779 (2014).
- [34] X. Zhang, M. Bishof, S. Bromley, C. Kraus, M. Safronova, P. Zoller, A. M. Rey, and J. Ye, *science* **345**, 1467 (2014).
- [35] See *Supplemental Materials for an explicit form of V_N and W_N and a generalized XYZ model.*
- [36] H. Shimizu and K. Yonekura, *Phys. Rev. D* **97**, 105011 (2018).
- [37] Y. Tanizaki and Y. Kikuchi, *JHEP* **2017**, 102 (2017).
- [38] X. Chen and A. Vishwanath, *Phys. Rev. X* **5**, 041034 (2015).

SUPPLEMENTAL MATERIALS

An explicit form of $\mathbb{Z}_N \times \mathbb{Z}_N \subset \text{PSU}(N)$ and a generalized XYZ model

In this part, we explicitly give the operator form of $v_{N,\vec{r}}$ and $w_{N,\vec{r}}$ by $\text{SU}(N)$ -spin degrees of freedom as

$$\begin{aligned} v_{N,\vec{r}} &= \left[\exp\left(i\pi \frac{\mathcal{S}_{2,\vec{r}}^1 + \mathcal{S}_{1,\vec{r}}^2}{2}\right) \exp\left(-i\pi \frac{\mathcal{S}_{1,\vec{r}}^1 + \mathcal{S}_{2,\vec{r}}^2}{2}\right) \right] \dots \left[\exp\left(i\pi \frac{\mathcal{S}_{N,\vec{r}}^1 + \mathcal{S}_{1,\vec{r}}^N}{2}\right) \exp\left(-i\pi \frac{\mathcal{S}_{1,\vec{r}}^1 + \mathcal{S}_{N,\vec{r}}^N}{2}\right) \right]; \\ w_{N,\vec{r}} &= \exp\left(-\sum_{\alpha} i \frac{2\pi}{N} \alpha \mathcal{S}_{\alpha,\vec{r}}^{\alpha}\right). \end{aligned} \quad (27)$$

$V_N = \prod_{\vec{r}} v_{N,\vec{r}}$ and $W_N = \prod_{\vec{r}} w_{N,\vec{r}}$ reduce to R_x^{π} and R_z^{π} , respectively, when $N = 2$. They generate the $\mathbb{Z}_N \times \mathbb{Z}_N \subset \text{PSU}(N)$ in the (adjoint) representation:

$$\mathcal{S}_{\beta}^{\alpha} \rightarrow U^{\dagger} \mathcal{S}_{\beta}^{\alpha} U, \quad (28)$$

where U is any finite product of V_N and W_N . If the local Hilbert space per unit cell is a tensor product of these Yang-tableaux and we denote the total number of the boxes in this general tensor product by b , such as

$$\underbrace{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} \otimes \square \otimes \square \otimes \dots}_{b \text{ YT boxes}}, \quad (29)$$

we obtain

$$v_{N,\vec{r}} w_{N,\vec{r}} = \exp\left(i2\pi \frac{b}{N}\right) w_{N,\vec{r}} v_{N,\vec{r}}, \quad (30)$$

A general $\text{SU}(N)$ analog of $\text{SU}(2)$ XYZ chain (3) possessing $\mathbb{Z}_N \times \mathbb{Z}_N$ above takes the form as

$$\mathcal{H}_{N;XYZ} = \sum_{g \in \mathbb{Z}_N \times \mathbb{Z}_N; i} J_g \mathcal{S}_{\beta,i}^{\alpha} (g)_{\alpha\beta} (g^{\dagger})_{\lambda\varphi} S_{\varphi,i+1}^{\lambda} + \text{h.c.},$$

with all the dumb indices summed up.
