

An integral-free representation of the Dyson series using divided differences

Amir Kalev^{1,*} and Itay Hen^{2,3,†}

¹*Information Sciences Institute, University of Southern California, Arlington, VA 22203, USA*

²*Information Sciences Institute, University of Southern California, Marina del Rey, CA 90292, USA*

³*Department of Physics and Astronomy, and Center for Quantum Information Science & Technology, University of Southern California, Los Angeles, California 90089, USA*

The Dyson series is an infinite sum of multi-dimensional time-ordered integrals, which serves as a formal representation of the quantum time evolution operator in the interaction picture. Using the mathematical tool of divided differences, we introduce an alternative representation for the series that is entirely free from both time ordering and integrals. In this new formalism, the Dyson expansion is given as a sum of efficiently-computable divided differences of the exponential function, considerably simplifying the calculation of the Dyson expansion terms, while also allowing for time-dependent perturbation calculations to be performed directly in the Schrödinger picture. We showcase the utility of this novel representation by studying a number of use cases. We also discuss several immediate applications.

I. INTRODUCTION

The Dyson series [1] is one of the fundamental results of quantum scattering theory. It is a perturbative expansion of the quantum time evolution operator in the interaction picture, wherein each summand is formally represented as a multi-dimensional integral over a time-ordered product of the interaction-picture Hamiltonian at different points in time. As such, the series serves as a pivotal tool for studying transition properties of time-dependent quantum many-body systems [2]. The Dyson series also has a close relation to Feynman diagrams in quantum field theory [3] and to the Magnus expansion in the theory of first-order homogeneous linear differential equation for linear operators [4].

Despite its fundamental role in quantum theory, one of the key challenges in using the Dyson series in practical applications remains the evaluation of the multi-dimensional integrals over products of time-ordered operators, making the calculation of the terms in the series an exceedingly complicated task [5]. In this study, we derive an analytical closed-form expression for the summands of the Dyson series, by explicitly evaluating the Dyson integrals. We accomplish this by utilizing the machinery of ‘divided differences’ – a mathematical tool normally used for computing tables of logarithms and trigonometric functions and for calculating the coefficients in the interpolation polynomial in the Newton form [6–12].

The main technical contribution of this work is an alternative, yet equivalent, formulation of the Dyson series wherein the summands are given in terms of efficiently computable divided differences of the exponential function. We argue that our novel representation, which is devoid of integrals and time ordering, makes it a very useful tool in the study of perturbation effects in many-body quantum systems. In particular, this representation allows us to write an explicit, integral-free time-dependent perturbation expansion for the time evolution operator in the Schrödinger picture enabling us to carry perturbation calculations without the need to switch to the interaction picture. To showcase the utility of our formulation, we work out a number of examples, for which we explicitly calculate the first few terms of the series. We begin with a brief introduction of divided differences, followed by the derivation of the integral-free representation of the Dyson series.

* amirk@isi.edu

† itayhen@isi.edu

II. DIVIDED DIFFERENCES – A BRIEF OVERVIEW

The divided differences [7, 8], which will be a major ingredient in the derivation detailed in the next section, is a recursive division process. The divided differences of a function $f(\cdot)$ is defined as

$$f[x_0, \dots, x_q] \equiv \sum_{j=0}^q \frac{f(x_j)}{\prod_{k \neq j} (x_j - x_k)} \quad (1)$$

with respect to the multi-set¹ of input variables $\{x_0, \dots, x_q\}$. By definition the divided differences is invariant under permutation of the input variables. The above expression is ill-defined if some of the inputs have repeated values, in which case one must resort to the use of limits. For instance, in the case where $x_0 = x_1 = \dots = x_q = x$, the definition of divided differences reduces to:

$$f[x_0, \dots, x_q] = \frac{f^{(q)}(x)}{q!}, \quad (2)$$

where $f^{(n)}(\cdot)$ stands for the n -th derivative of $f(\cdot)$. Divided differences can alternatively be defined, and are often calculated, via the recursion relations

$$f[x_0, \dots, x_q] = \frac{f[x_1, \dots, x_q] - f[x_0, \dots, x_{q-1}]}{x_q - x_0}, \quad (3)$$

with the initial conditions $f[x_i] = f(x_i)$, for $i = 0, \dots, q$. As is evident from the above, the divided differences of any given function $f(\cdot)$ with q inputs can be calculated with $q(q-1)/2$ basic operations.

The divided differences of the exponential function $f(x) = e^{-itx}$, which will be the main focus of the present study, satisfy the following two identities which we will utilize in the next section (proofs are given in the Appendix):

Identity 1:

$$(-i)^q \int_0^t dt_q \cdots \int_0^{t_2} dt_1 e^{-i(\gamma_q t_q + \cdots + \gamma_1 t_1)} = e^{-it[x_0, x_1, \dots, x_{q-1}, 0]} \quad (4)$$

where $x_j = \sum_{k=j+1}^q \gamma_k$. This identity is a variant of what is often known as the Hermite-Genocchi formula [8].

Identity 2: Given an arbitrary multi-set of inputs $\{x_0, \dots, x_q\}$,

$$e^{-it[x_0, \dots, x_q]} = e^{-itx} e^{-it[\Delta_0, \dots, \Delta_q]}, \quad (5)$$

where x is an arbitrary constant and $\Delta_j = x_j - x$.

III. DIVIDED DIFFERENCES FORMULATION OF THE TIME-EVOLUTION OPERATOR

Often times, the most convenient approach to tackling the dynamics of time-dependent systems is through the use of (time-dependent) perturbation theory in the interaction-picture [13, 14] where the Hamiltonian is written as

$$H(t) = H_0 + V(t), \quad (6)$$

namely as a sum of a static (or ‘free’) Hamiltonian H_0 (that is assumed to have a known spectrum) and an additional time-dependent Hamiltonian, $V(t)$, that is treated as a perturbation to H_0 . The time-evolution

¹ A multi-set (also known as bag, mset) is a generalization of a set where repetition of elements matters.

operator in the interaction picture is given by $U_I(t) = \mathcal{T} \exp[-i \int_0^t H_I(t) dt]$ — a shorthand for the Dyson series

$$U_I(t) = \mathbb{1} + \sum_{q=1}^{\infty} \frac{(-i)^q}{q!} \int_0^t dt_1 \cdots \int_0^t dt_q \mathcal{T} [H_I(t_q) \cdots H_I(t_1)], \quad (7)$$

where \mathcal{T} denotes time-ordering [15] and $H_I(t) = e^{iH_0 t} V(t) e^{-iH_0 t}$. The Dyson series can also be re-written as

$$U_I(t) = \mathbb{1} + \sum_{q=1}^{\infty} (-i)^q \int_0^t dt_1 H_I(t_1) \cdots \int_0^{t_{q-1}} dt_q H_I(t_q) \quad (8)$$

and by relabeling the integration variables, the above equation takes the form:

$$U_I(t) = \sum_{q=0}^{\infty} (-i)^q \int_0^t dt_q \cdots \int_0^{t_2} dt_1 H_I(t_q) \cdots H_I(t_1), \quad (9)$$

where hereafter we will use the $q = 0$ term to symbolize the identity operator.

The operator $U_I(t)$ evolves the interaction-picture wave-function $|\psi_I(t)\rangle$ which is related to the Schrödinger-picture wave-function via $|\psi_I(t)\rangle = e^{iH_0 t} |\psi(t)\rangle$ (in our units, $\hbar = 1$). Similarly, the Schrödinger-picture time-evolution operator $U(t)$ is related to the interaction-picture operator via $U(t) = e^{-iH_0 t} U_I(t)$. In the next section we present an equivalent form for the Dyson series, Eq. (9), by systematically evaluating the integrals in the sum, writing $V(t)$ as a sum of exponentials in t .

A. Generalized permutation operator representation of the perturbation Hamiltonian

We begin by denoting the eigenstates and eigenenergies of the free Hamiltonian H_0 by $\mathcal{B} = \{|z\rangle\}$ and $\mathcal{E} = \{E_z\}$, respectively, such that $H_0|z\rangle = E_z|z\rangle$. (For simplicity we assume a discrete countable set of eigenstates and eigenenergies). We will refer to \mathcal{B} as the ‘computational basis’. Next, we write the perturbation Hamiltonian $V(t)$ as a sum of *generalized permutation operators* Π_i [16]:

$$V(t) = \sum_{i=0}^M \Pi_i(t) = \sum_{i=0}^M D_i(t) P_i, \quad (10)$$

where every generalized permutation operator is further expressed as a product of a (time dependent) diagonal (in the computational basis) operator D_i and a bona-fide permutation operator P_i . Specifically, the action of D_i and P_i on a computational basis states is given by $D_i|z\rangle = d_i(z)|z\rangle$, where $d_i(z)$ is in general a complex number, and $P_i|z\rangle = |z'\rangle$ for some $|z'\rangle \in \mathcal{B}$ depending on i and z . The $i = 0$ permutation operator will be reserved to the identity operator, that is, $P_0 = \mathbb{1}$. Armed with these notations, the action of a generalized permutation operator Π_i on a basis state $|z\rangle$ is given by $D_i P_i |z\rangle = d_i(z') |z'\rangle$, where z' depends on both the state z and the operator index i . We note that any Hamiltonian can be readily cast in the above form [11]. This representation, in terms of generalized permutation operators, was recently introduced in the context of quantum Monte Carlo simulations [9–11].

At this point, we write each diagonal operator, $D_i(t)$, in Eq. (10) as an exponential sum in t , that is,

$$D_i(t) = \sum_{k=1}^{K_i} e^{i\Lambda_i^{(k)} t} D_i^{(k)}, \quad (11)$$

where both $\Lambda_i^{(k)}$ and $D_i^{(k)}$ are (generally complex-valued) diagonal matrices and K_i denotes the number of terms in the decomposition of D_i . (For more details as to how to carry out this decomposition efficiently, see Refs. [17–19].) Thus, $V(t)$ can be written as

$$V(t) = \sum_{i=1}^M \sum_{k=1}^{K_i} e^{i\Lambda_i^{(k)} t} D_i^{(k)} P_i \quad (12)$$

(for simplicity, hereafter we fix $K_i = K \forall i$, though this assumption can be easily removed).

B. An integral-free form of the Dyson series

Using the expansion of $V(t)$, Eq. (12), the interaction-picture Hamiltonian $H_I(t)$ is written as

$$H_I(t) = e^{iH_0 t} V(t) e^{-iH_0 t} = \sum_{i=1}^M \sum_{k=1}^K e^{i(H_0 + \Lambda_i^{(k)})t} D_i^{(k)} P_i e^{-iH_0 t} \equiv \sum_{i=1}^M \sum_{k=1}^K e^{iH_0^{(i,k)} t} D_i^{(k)} P_i e^{-iH_0 t} \quad (13)$$

where we have defined $H_0^{(i,k)} \equiv H_0 + \Lambda_i^{(k)}$. With this form of $H_I(t)$ we can write the Dyson series for $U_I(t)$, Eq. (9), as

$$\begin{aligned} U_I(t) &= \sum_{q=0}^{\infty} (-i)^q \int_0^t dt_q \cdots \int_0^{t_2} dt_1 \left(\sum_{i_q=1}^M \sum_{k_q=1}^K e^{iH_0^{(i_q, k_q)} t_q} D_{i_q}^{(k_q)} P_{i_q} e^{-iH_0 t_q} \right) \\ &\cdots \left(\sum_{i_1=1}^M \sum_{k_1=1}^K e^{iH_0^{(i_1, k_1)} t_1} D_{i_1}^{(k_1)} P_{i_1} e^{-iH_0 t_1} \right) \\ &= \sum_{q=0}^{\infty} \sum_{\mathbf{i}_q} \sum_{\mathbf{k}_q} (-i)^q \int_0^t dt_q \cdots \int_0^{t_2} dt_1 \left(e^{iH_0^{(i_q, k_q)} t_q} D_{i_q}^{(k_q)} P_{i_q} e^{-iH_0 t_q} \cdots e^{iH_0^{(i_1, k_1)} t_1} D_{i_1}^{(k_1)} P_{i_1} e^{-iH_0 t_1} \right). \end{aligned} \quad (14)$$

$$(15)$$

where $\mathbf{i}_q = (i_1, i_2, \dots, i_q)$ and $\mathbf{k}_q = (k_1, k_2, \dots, k_q)$ are multi-indices. Acting with $U_I(t)$ on an arbitrary computational basis state $|z\rangle$, we get

$$\begin{aligned} U_I(t)|z\rangle &= \sum_{q=0}^{\infty} \sum_{\mathbf{i}_q} \sum_{\mathbf{k}_q} (-i)^q \int_0^t dt_q \cdots \int_0^{t_2} dt_1 e^{iE_z^{(i_q, k_q)} t_q} d_z^{(i_q, k_q)} e^{-iE_z^{(i_{q-1})} t_q} \cdots e^{iE_z^{(i_1, k_1)} t_1} d_z^{(i_1, k_1)} e^{-iE_z t_1} P_{\mathbf{i}_q} |z\rangle \\ &= \sum_{q=0}^{\infty} \sum_{\mathbf{i}_q} \sum_{\mathbf{k}_q} d_z^{(\mathbf{i}_q, \mathbf{k}_q)} \left((-i)^q \int_0^t dt_q \cdots \int_0^{t_2} dt_1 e^{-i(E_z^{(i_{q-1})} - E_z^{(i_q, k_q)}) t_q} \cdots e^{-i(E_z - E_z^{(i_1, k_1)}) t_1} \right) P_{\mathbf{i}_q} |z\rangle, \end{aligned} \quad (16)$$

where we have defined $P_{\mathbf{i}_q} = P_{i_q} \cdots P_{i_1}$ and $|z_{i_j}\rangle = P_{i_j} |z\rangle$ for every $j = 0, \dots, q$ (remembering that the value of z_{i_j} depends on that of z). Moreover, we define $E_z^{(i_j)} = \langle z_{i_j} | H_0 | z_{i_j} \rangle$ and similarly

$$E_z^{(i_j, k_j)} = \langle z_{i_j} | H_0^{(i_j, k_j)} | z_{i_j} \rangle = E_z^{(i_j)} + \lambda_z^{(i_j, k_j)}, \quad (17)$$

with $\lambda_z^{(i_j, k_j)} = \langle z_{i_j} | \Lambda_{i_j}^{(k_j)} | z_{i_j} \rangle$. In addition, we have denoted $d_z^{(i_j, k_j)} = \langle z_{i_j} | D_{i_j}^{(k_j)} | z_{i_j} \rangle$ and $d_z^{(\mathbf{i}_q, \mathbf{k}_q)} = \prod_{j=1}^q d_z^{(i_j, k_j)}$. Following Identity 1 [Eq. (4) of the previous section], the integrals appearing inside the parenthesis in Eq. (16) evaluate explicitly to the divided-differences exponential

$$(-i)^q \int_0^t dt_q \cdots \int_0^{t_2} dt_1 e^{-i(E_z^{(i_{q-1})} - E_z^{(i_q, k_q)}) t_q} \cdots e^{-i(E_z - E_z^{(i_1, k_1)}) t_1} = e^{-it[x_0, x_1, \dots, x_{q-1}, 0]}, \quad (18)$$

with

$$x_j = \sum_{\ell=j+1}^q E_z^{(i_{\ell-1})} - E_z^{(i_{\ell}, k_{\ell})} = E_z^{(i_j)} - E_z^{(i_q)} - \sum_{\ell=j+1}^q \lambda_z^{(i_{\ell}, k_{\ell})} \text{ for } j = 0, \dots, q-1 \quad (19)$$

(we have omitted the dependence of x_0, \dots, x_{q-1} on z , \mathbf{i}_q and on \mathbf{k}_q for a lighter notation). Importantly, the number $e^{-it[x_0, x_1, \dots, x_{q-1}, 0]}$ can be calculated efficiently, with computational complexity proportional to q^2 (see Ref. [20] for additional details). Using Eq. (18) we can write the action of $U_I(t)$ on an arbitrary computational basis state in an integral-free form as

$$U_I(t)|z\rangle = \sum_{q=0}^{\infty} \sum_{\mathbf{i}_q} \sum_{\mathbf{k}_q} d_z^{(\mathbf{i}_q, \mathbf{k}_q)} e^{-it[x_0, x_1, \dots, x_{q-1}, 0]} P_{\mathbf{i}_q} |z\rangle, \quad (20)$$

with the inputs to the divided-differences exponential given in Eq. (19). We thus arrive at:

$$U_I(t) = \sum_{q=0}^{\infty} \sum_z \sum_{\mathbf{i}_q} \sum_{\mathbf{k}_q} d_z^{(\mathbf{i}_q, \mathbf{k}_q)} e^{-it[x_0, x_1, \dots, x_{q-1}, 0]} P_{\mathbf{i}_q} |z\rangle \langle z| = \sum_{q=0}^{\infty} \sum_z \sum_{\mathbf{i}_q} \alpha_z^{(\mathbf{i}_q)} P_{\mathbf{i}_q} |z\rangle \langle z|, \quad (21)$$

where we have defined the (time-dependent) c-numbers

$$\alpha_z^{(\mathbf{i}_q)} = \sum_{\mathbf{k}_q} d_z^{(\mathbf{i}_q, \mathbf{k}_q)} e^{-it[x_0, x_1, \dots, x_{q-1}, 0]}. \quad (22)$$

Equation (21) is the main technical result of our paper. The equation is a reformulation of the Dyson series, Eq. (9), yet it includes only sums of efficiently-computable terms and is devoid of both integrals and time-ordering operators.

The current representation of $U_I(t)$ also allows us to obtain an integral-free time-dependent perturbation expansion of the the Schrödinger-picture time-evolution operator $U(t)$:

$$\begin{aligned} U(t) &= e^{-iH_0 t} U_I(t) = \sum_{q=0}^{\infty} \sum_z \sum_{\mathbf{i}_q} \alpha_z^{(\mathbf{i}_q)} e^{-iH_0 t} P_{\mathbf{i}_q} |z\rangle \langle z| \\ &= \sum_{q=0}^{\infty} \sum_z \sum_{\mathbf{i}_q} \alpha_z^{(\mathbf{i}_q)} e^{-itE_z^{(\mathbf{i}_q)}} P_{\mathbf{i}_q} |z\rangle \langle z| = \sum_{q=0}^{\infty} \sum_z \sum_{\mathbf{i}_q} \beta_z^{(\mathbf{i}_q)} P_{\mathbf{i}_q} |z\rangle \langle z|. \end{aligned} \quad (23)$$

where

$$\beta_z^{(\mathbf{i}_q)} = \alpha_z^{(\mathbf{i}_q)} e^{-itE_z^{(\mathbf{i}_q)}} = \sum_{\mathbf{k}_q} d_z^{(\mathbf{i}_q, \mathbf{k}_q)} e^{-it[y_0, y_1, \dots, y_q]}. \quad (24)$$

with

$$y_j = E_z^{(\mathbf{i}_j)} - \sum_{\ell=j+1}^q \lambda_z^{(\mathbf{i}_\ell, k_\ell)} \quad \text{for } j = 0, \dots, q. \quad (25)$$

We arrive at the last two equations by using Identity 2 [Eq. (5) given in the previous section].

Equation (23) allows us to carry out directly and explicitly, time-dependent perturbation calculations directly in the Schrödinger picture without invoking the interaction picture.

IV. USE CASES

In this section we illustrate, by examining few basic examples, the utility of Eq. (23) in a variety of settings, specifically in scenarios where the time-ordered integrals of the Dyson series are cumbersome to calculate or lead to unnecessary complicated expressions.

A. Transition amplitudes and Fermi's golden rule

Using the divided-differences series formulation of the time-evolution operator one may arrive at a rather simple expression for transition amplitudes, $A(z_{\text{in}} \rightarrow z_{\text{fin}}, t) = \langle z_{\text{fin}} | U(t) | z_{\text{in}} \rangle$, between an initial eigenstate $|z_{\text{in}}\rangle$ and a final eigenstate $|z_{\text{fin}}\rangle$ of the free Hamiltonian H_0 due to the perturbation effect of $V(t)$, cf. Eq. (6). Using Eq. (23) we find

$$A(z_{\text{in}} \rightarrow z_{\text{fin}}, t) = \sum_{q=0}^{\infty} \sum_{\mathbf{i}_q: \langle z_{\text{fin}} | P_{\mathbf{i}_q} | z_{\text{in}} \rangle = 1} \beta_{z_{\text{in}}}^{(\mathbf{i}_q)}, \quad (26)$$

where the second sum is over all multi-indices \mathbf{i}_q that obey $P_{\mathbf{i}_q} |z_{\text{in}}\rangle = |z_{\text{fin}}\rangle$.

For example, for the canonical (non-hermitian) case $H = H_0 + \gamma F e^{iEt}$ where γ is a perturbation parameter and F a permutation matrix such that $F|z_{\text{in}}\rangle = |z_{\text{fin}}\rangle$ and $F^2 = \mathbb{1}$, the zeroth-order term of the divided-differences series expansion corresponds to $P_{\mathbf{i}_0} = \mathbb{1}$, and thus vanishes (assuming $|z_{\text{in}}\rangle \neq |z_{\text{fin}}\rangle$). The first-order term of the series expansion ($q = 1$) is given by

$$A^{(1)}(z_{\text{in}} \rightarrow z_{\text{fin}}, t) = \gamma e^{-it[E_{z_{\text{in}}} + E, E_{z_{\text{fin}}}] = \gamma \frac{e^{-it(E_{z_{\text{in}}} + E)} - e^{-itE_{z_{\text{fin}}}}}{E_{z_{\text{in}}} + E - E_{z_{\text{fin}}}}, \quad (27)$$

in accordance with Fermi's golden rule [21]. An interesting observation is that while the divided-differences formulation is well-defined (by definition) even in the limit where $E_{z_{\text{in}}} + E - E_{z_{\text{fin}}} \rightarrow 0$, the explicit expression on right-hand-side of the equation above must be amended to account for the removable singularity of the expression.

Moreover, the present formulation allows us to immediately write an expression for transition amplitudes that are accurate to yet higher orders, that is,

$$A(z_{\text{in}} \rightarrow z_{\text{fin}}, t) = \gamma e^{-it[E_{z_{\text{in}}} + E, E_{z_{\text{fin}}}] + \gamma^3 e^{-it[E_{z_{\text{in}}} + 3E, E_{z_{\text{fin}}} + 2E, E_{z_{\text{in}}} + E, E_{z_{\text{fin}}}] + \gamma^5 e^{-it[E_{z_{\text{in}}} + 5E, E_{z_{\text{fin}}} + 4E, E_{z_{\text{in}}} + 3E, E_{z_{\text{fin}}} + 2E, E_{z_{\text{in}}} + E, E_{z_{\text{fin}}}] + \dots \quad (28)$$

This simple expression can be contrasted with the corresponding Dyson-series-based derivation and final expression which are considerably more cumbersome to derive and to calculate (see, e.g., Refs. [22, 23]).

B. Reduction to the time-independent case

In the special case where the Hamiltonian is time independent (TI), the formulation we derived above reduces to a more compact form. If time does not appear in any of the D_i operators, we can set $K = 1$ and $\Lambda_i^{(1)} = 0$ for all D_i . As a consequence, the y_j divided-differences inputs reduce to $y_j = E_z^{(i_j)}$ and so the time-evolution operator takes the simpler form

$$U_{\text{TI}}(t) = \sum_{q=0}^{\infty} \sum_{\mathbf{i}_q} \left(\sum_z D_z^{(\mathbf{i}_q)} e^{-it[E_z^{(i_0)}, E_z^{(i_1)}, \dots, E_z^{(i_q)}]} P_{\mathbf{i}_q} |z\rangle \langle z| \right), \quad (29)$$

where $D_z^{(\mathbf{i}_q)} = \prod_j d_z^{(i_j)}$. The above expression coincides with the expression worked out in Ref. [12], where time-independent Hamiltonians were considered (in a somewhat different context).

C. Analytical calculations of time-evolved states

The divided-differences formulation above allows us to readily compute analytical closed-form expressions in the Schrödinger picture for states evolved under a time-dependent Hamiltonian. We next work out a couple of examples where we demonstrate the ease with which the various terms in the expansion of the time evolution operator can be calculated analytically. Obtaining such expressions can be also useful, for example, in numerical simulation of perturbation effects.

1. A single-spin Hamiltonian

We first consider a simple example of a time-dependent problem, explicitly, the single-spin Hamiltonian $H(t) = aZ + be^{-\gamma t}X$ where Z and X are the Pauli- z and Pauli- x operators, respectively and a, b and γ are constants. We identify $H_0 = aZ$ and $V(t) = be^{-\gamma t}X$. The time-dependent component $V(t)$ contains a single permutation matrix, namely, $P_1 = X$ ($M = 1$) and the associated diagonal operator is $D_1 = be^{-\gamma t}\mathbb{1}$ has only one exponential term ($K = 1$). This further implies a single term at every expansion order q . In this case, the coefficients $\beta_z^{(q)}$, defined in Eq. (24), are given by

$$\beta_z^{(q)} = b^q e^{-it[y_0, \dots, y_q]} \quad \text{with} \quad y_j = (-1)^z (-1)^j a - i(q-j)\gamma, \quad \text{for } j = 0, \dots, q \quad (30)$$

[here $z = 0(1)$ corresponds to the $|0\rangle(|1\rangle)$ eigenstate of Z]. The q -th order permutation operator is $P_q = X^q$, i.e., $P_{2q} = \mathbb{1}$ and $P_{2q+1} = X$. The state of the system at time t given an initial state $|z\rangle$ is simply $|\psi(t)\rangle = U(t)|z\rangle = \sum_q \beta_z^{(q)} P_q |z\rangle = \left(\sum_q \beta_z^{(2q)}\right) |z\rangle + \left(\sum_q \beta_z^{(2q+1)}\right) |z\rangle$.

2. The time-dependent anharmonic oscillator

Next, we work out the expansion of the time-evolution operator for a particle moving under the influence of a harmonic potential perturbed by a periodic time-dependent anharmonic term, namely,

$$H(t) = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega^2 \hat{x}^2 + \frac{1}{4}\Gamma \cos(\Omega t) \hat{x}^4 \quad (31)$$

Identifying the static component as $H_0 = \hat{p}^2/2m + m\omega^2 \hat{x}^2/2$, we rewrite the Hamiltonian in terms of the harmonic annihilation-creation operators (recall that $\hbar = 1$)

$$\hat{a} = \sqrt{\frac{m\omega}{2}} \left(\hat{x} + \frac{i}{m\omega} \hat{p} \right) = \sum_n \sqrt{n} |n-1\rangle \langle n| \quad \text{and} \quad (32)$$

$$\hat{a}^\dagger = \sqrt{\frac{m\omega}{2}} \left(\hat{x} - \frac{i}{m\omega} \hat{p} \right) = \sum_n \sqrt{n+1} |n+1\rangle \langle n|, \quad (33)$$

where $|n\rangle$ with $n = 0, 1, \dots$ are the harmonic oscillator eigenstates whose energies are $E_n = \omega(n + \frac{1}{2})$. In terms of these, the Hamiltonian becomes

$$H(t) = \omega \left(\hat{a}^\dagger \hat{a} + \frac{1}{2} \right) + \frac{1}{4}\Gamma \cos(\Omega t) \left(\frac{1}{2m\omega} \right)^2 (\hat{a}^\dagger + \hat{a})^4. \quad (34)$$

To express $V(t) = \frac{1}{4}\Gamma \cos(\Omega t) \left(\frac{1}{2m\omega} \right)^2 (\hat{a}^\dagger + \hat{a})^4$ in terms of generalized permutation operators, using the harmonic oscillator eigenstates as basis states, we rewrite the anharmonic operator $(\hat{a}^\dagger + \hat{a})^4$ as $(\hat{a}^\dagger + \hat{a})^4 = \sum_{i \in \{0, \pm 2, \pm 4\}} D_i P_i$ where $P_i = \sum_n |n+i\rangle \langle n|$ and the matrix elements of the diagonal operators D_i are given by

$$\begin{aligned} (D_{-4})_{nn} &= \sqrt{n(n-1)(n-2)(n-3)}, \\ (D_{-2})_{nn} &= \sqrt{n(n-1)(4n-2)}, \\ (D_0)_{nn} &= 3(2n^2 + 2n + 1), \\ (D_2)_{nn} &= \sqrt{(n+1)(n+2)(4n+6)}, \\ (D_4)_{nn} &= \sqrt{(n+1)(n+2)(n+3)(n+4)}. \end{aligned} \quad (35)$$

In terms of the above operators, the Hamiltonian can be written as

$$H(t) = H_0 + \gamma \sum_i \sum_{k=\pm 1} e^{-ik\Omega t} D_i P_i \quad (36)$$

where $H_0 = \omega \sum_n (n + \frac{1}{2}) |n\rangle \langle n|$ and $\gamma = \frac{1}{8}\Gamma \left(\frac{1}{2m\omega} \right)^2$. Having cast the Hamiltonian in the appropriate form, it is straightforward to calculate the $\beta_n^{(i_q)}$ terms in the divided-differences expansion of $U(t)$, as per Eq. (24), as well as the corresponding permutation operators P_{i_q} . These are summarized in Table I. Together, these provide us with an easy way to compute the state of the system after time t . For an initial state $|n\rangle$, we get the analytic closed-form expression $|\psi(t)\rangle = U(t)|n\rangle = \sum_{i_q} \beta_n^{(i_q)} P_{i_q} |n\rangle$.

Having computed the coefficients, we may plot, as we do in Fig. 1(a), the population of every mode at various times. In the figure, the populations are given at $t = 0.04$ for $\omega = 1, \Omega = 2, \gamma = 0.02$ with the initial state $|n = 4\rangle$ and an expansion cutoff of $Q = 5$. To ascertain the accuracy of the Dyson series truncated

Order	\mathbf{i}_q	Coefficient $\beta_n^{(i_q)}$	$P_{\mathbf{i}_q}$	Comments
$q = 0$	$()$	e^{-itE_n}	$\mathbb{1}$	0-th order perturbation
$q = 1$	(-4)	$\gamma\sqrt{n(n-1)(n-2)(n-3)}(e^{-it[E_n+\Omega, E_{n-4}]} + e^{-it[E_n-\Omega, E_{n-4}]})$	P_{-4}	vanishes unless $n \geq 4$
	(-2)	$\gamma\sqrt{n(n-1)(4n-2)}(e^{-it[E_n+\Omega, E_{n-2}]} + e^{-it[E_n-\Omega, E_{n-2}]})$	P_{-2}	
	(0)	$3\gamma(2n^2 + 2n + 1)(e^{-it[E_n+\Omega, E_n]} + e^{-it[E_n-\Omega, E_n]})$	$P_0 = \mathbb{1}$	vanishes unless $n \geq 2$
	(2)	$\gamma\sqrt{(n+2)(n+1)(4n+6)}(e^{-it[E_n+\Omega, E_{n+2}]} + e^{-it[E_n-\Omega, E_{n+2}]})$	P_2	
	(4)	$\gamma\sqrt{(n+1)(n+2)(n+3)(n+4)}(e^{-it[E_n+\Omega, E_{n+4}]} + e^{-it[E_n-\Omega, E_{n+4}]})$	P_4	
$q = 2$	$(-4, -4)$	$\gamma^2\sqrt{n(n-1)\cdots(n-8)}$ $\times \sum_{k_1, k_2=\pm 1} e^{-it[E_n+(-1)^{k_1+k_2}\Omega, E_{n-4}+(-1)^{k_2}\Omega, E_{n-8}]}$	$P_{-8} = P_{-4}P_{-4}$	vanishes unless $n \geq 8$
	$(-4, -2)$	$\gamma^2\sqrt{n(n-1)\cdots(n-6)}$ $\times \sum_{k_1, k_2=\pm 1} e^{-it[E_n+(-1)^{k_1+k_2}\Omega, E_{n-4}+(-1)^{k_2}\Omega, E_{n-6}]}$	$P_{-6} = P_{-2}P_{-4}$	vanishes unless $n \geq 6$
	$(-4, 0)$	$3\gamma^2\sqrt{n(n-1)(n-2)(n-3)}(2n^2 - 14n + 25)$ $\times \sum_{k_1, k_2=\pm 1} e^{-it[E_n+(-1)^{k_1+k_2}\Omega, E_{n-4}+(-1)^{k_2}\Omega, E_{n-4}]}$	$P_{-4} = P_0P_{-4}$	vanishes unless $n \geq 4$
	...			

Table I. First $\beta_n^{(i_q)}$ terms in the expansion of the time-dependent anharmonic oscillator, Eq. (34). The energies E_n are the harmonic oscillator eigenenergies. Denoting the eigenenergies by $|n\rangle$, the permutation operators are given by $P_i = \sum_n |n+i\rangle\langle n|$. For an initial state $|n\rangle$, the state at time t is given by $|\psi(t)\rangle = U(t)|n\rangle = \sum_{\mathbf{i}_q} \beta_n^{(i_q)} P_{\mathbf{i}_q}|n\rangle$.

at different cutoff orders Q , we may also contrast the divided-differences expansion with exact-numerical results obtained via direct integration of the Schrödinger equation. In Fig. 1(b) we plot the infidelity $1 - |\langle\psi(t)|\psi_Q(t)\rangle|^2$ between the state $|\psi(t)\rangle$, the solution of the time-dependent Schrödinger equation of this system, and the state as obtained by using divided-differences expansion for $U(t)$, $|\psi_Q(t)\rangle = U_Q(t)|4\rangle$, that is, the state obtained by evolving $|4\rangle$ under $U(t)$ given by Eq. (23) as a function of evolution time t .




Figure 1. (a) Probabilities of the various modes at $t = 0.04$ for the anharmonic Hamiltonian, Eq. (34). Here, $\omega = 1, \Omega = 2, \gamma = 0.02$ and the initial state is $|n = 4\rangle$. The blue circles are obtained from numerical integration of the Schrödinger equation while the orange triangles correspond to the divided-differences expansion up to order $Q = 5$. (b) Infidelity $1 - |\langle\psi(t)|\psi_Q(t)\rangle|^2$ as a function of time. The various curves correspond to different truncation orders $Q = 0, \dots, 3$. As we expect, the higher the expansion order is, the better the approximation becomes.

V. SUMMARY AND DISCUSSION

In this work, we devised an alternative approach to time-dependent perturbation theory in quantum mechanics. We derived an expansion that is equivalent to the usual Dyson series but which includes only sums of closed-form analytical simple-to-calculate expressions rather than the usual Dyson multi-dimensional time-ordered integrals. This new formalism allowed us to write an integral-free perturbation expansion to

the time evolution operator. We illustrated the utility of our approach, we examined a number of use cases and calculated the series coefficients in a number of examples for which the usual Dyson series calculation is cumbersome, demonstrating the functionality and practicality of our approach.

Another area in which the divided-differences expansion can be applied, which we have not explored here, is quantum algorithms – algorithms designed to be executed on quantum computers. Specifically, the divided-differences expansion can be used in simulations of the time-evolution of quantum states evolving under time-dependent Hamiltonians (the time-independent variant of a Hamiltonian simulation algorithm which utilizes the off-diagonal series expansion was discussed in Ref. [12]). We have shown that within the divided-differences expansion, the time evolution operator can be written as a sum of generalized permutation matrices, equivalently a linear combination of unitary (LCU) operators. As such, this representation of the time-evolution operator lends itself naturally to the quantum LCU lemma [24] which provides a prescription for efficiently simulating such operators on quantum circuits. We leave that for future work.

We believe that the formulation introduced here will prove itself to be a powerful tool in the study of time-dependent quantum many-body systems.

ACKNOWLEDGMENTS

This work is supported by the U.S. Department of Energy (DOE), Office of Science, Basic Energy Sciences (BES) under Award No. DE-SC0020280.

-
- [1] F. J. Dyson, “The radiation theories of tomonaga, schwinger, and feynman,” *Phys. Rev.* **75**, 486–502 (1949).
 - [2] A. L. Fetter and J. D. Walecka, *Quantum Theory of Many Particle Systems* (Dover, New York, 2003).
 - [3] S. Weinberg, *The Quantum Theory of Fields* (Cambridge University Press, Cambridge York, 2005).
 - [4] S. Blanes, F. Casas, J.A. Oteo, and J. Ros, “The magnus expansion and some of its applications,” *Physics Reports* **470**, 151 – 238 (2009).
 - [5] C. Brouder, Å Mestre, and Patras F., “Tree expansion in time-dependent perturbation theory,” *Journal of Mathematical Physics* **51**, 072104 (2010).
 - [6] L.M. Milne-Thomson, *The Calculus of Finite Differences* (Macmillan and Company, limited, 1933).
 - [7] E. T. Whittaker and G. Robinson, “Divided differences,” in *The Calculus of Observations: A Treatise on Numerical Mathematics* (New York: Dover, New York, 1967).
 - [8] Carl de Boor, “Divided differences,” *Surveys in Approximation Theory* **1**, 46–69 (2005).
 - [9] Tameem Albash, Gene Wagenbreth, and Itay Hen, “Off-diagonal expansion quantum monte carlo,” *Phys. Rev. E* **96**, 063309 (2017).
 - [10] Itay Hen, “Off-diagonal series expansion for quantum partition functions,” *Journal of Statistical Mechanics: Theory and Experiment* **2018**, 053102 (2018).
 - [11] Lalit Gupta, Tameem Albash, and Itay Hen, “Permutation Matrix Representation Quantum Monte Carlo,” accepted for publication in *Journal of Statistical Mechanics: Theory and Experiment*, arXiv:1908.03740 (2019), arXiv:1908.03740 [cond-mat.stat-mech].
 - [12] Amir Kalev and Itay Hen, “Simulating Hamiltonian Dynamics with an Off-diagonal Series Expansion,” arXiv e-prints, arXiv:2006.02539 (2020), arXiv:2006.02539 [quant-ph].
 - [13] A. Messiah, *Quantum Mechanics*, Vol. 2 (North-Holland, Amsterdam, 1962).
 - [14] J. J. Sakurai, *Modern Quantum Mechanics* (Addison-Wesley, New York, 1994).
 - [15] S. Weinberg, *The quantum theory of fields*, Vol. 3 (Cambridge University Press, Cambridge, UK).
 - [16] David Joyner, *Adventures in group theory. Rubik’s cube, Merlin’s machine, and other mathematical toys* (Baltimore, MD: Johns Hopkins University Press, 2008).
 - [17] Gregory Beylkin and Lucas Monzón, “On approximation of functions by exponential sums,” *Applied and Computational Harmonic Analysis* **19**, 17 – 48 (2005).
 - [18] Gregory Beylkin and Lucas Monzón, “Approximation by exponential sums revisited,” *Applied and Computational Harmonic Analysis* **28**, 131 – 149 (2010), special Issue on Continuous Wavelet Transform in Memory of Jean Morlet, Part I.
 - [19] Gregory Beylkin, Martin J. Mohlenkamp, and Fernando Pérez, “Approximating a wavefunction as an unconstrained sum of slater determinants,” *Journal of Mathematical Physics* **49**, 032107 (2008), <https://doi.org/10.1063/1.2873123>.
 - [20] Lalit Gupta, Lev Barash, and Itay Hen, “Calculating the divided differences of the exponential function by addition and removal of inputs,” *Computer Physics Communications* **254**, 107385 (2020).

- [21] Merzbacher, Eugen, *Quantum Mechanics* (Wiley and Sons, 1998).
 [22] L.D. Landau and E.M. Lifshitz, *Quantum Mechanics. Non-Relativistic theory, Vol. III* (Pergamon Press Ltd., London, 1958).
 [23] L. I. Schiff, *Quantum Mechanics* (McGraw-Hill, 1955).
 [24] Dominic W. Berry, Andrew M. Childs, Richard Cleve, Robin Kothari, and Rolando D. Somma, “Simulating hamiltonian dynamics with a truncated taylor series,” *Phys. Rev. Lett.* **114**, 090502 (2015).

APPENDIX: PROOFS OF IDENTITY 1 AND IDENTITY 2

Proof of Identity 1: We start with left-hand-side of Eq. (4). Making a change of variables $t_j = ts_j$ we get

$$(-i)^q \int_0^t dt_q \cdots \int_0^{t_2} dt_1 e^{-i(\gamma_q t_q + \cdots + \gamma_1 t_1)} = (-it)^q \int_0^1 ds_q \cdots \int_0^{s_2} ds_1 e^{-it(\gamma_q s_q + \cdots + \gamma_1 s_1)}. \quad (37)$$

Next we prove by induction that

$$e^{-it[x_0, x_1, \dots, x_{q-1}, 0]} = (-it)^q \int_0^1 ds_q \cdots \int_0^{s_2} ds_1 e^{-it(\gamma_q s_q + \cdots + \gamma_1 s_1)} \quad (38)$$

where $x_j = \sum_{k=j+1}^q \gamma_k$.

Proving for $q = 1$: Starting from the left-hand-side of Eq. (38) we have

$$(-it) \int_0^1 ds_1 e^{-it\gamma_1 s_1} = \frac{(-it)}{\gamma_1} \int_0^{\gamma_1} d\xi e^{-it\xi} = \frac{1}{\gamma_1} \int_0^{\gamma_1} d\xi \frac{d}{d\xi} e^{-it\xi} = \frac{e^{-it\gamma_1} - 1}{\gamma_1} = e^{-it[0, \gamma_1]} = e^{-it[\gamma_1, 0]} \quad (39)$$

as required. In the first equality of (39) we changed the integration variable $\xi = s_1 \gamma_1$, so that ξ ranges from 0 to γ_1 , and $\gamma_1 ds_1 = d\xi$. Next, assuming the validity of Eq. (38), we prove that

$$(-it)^{q+1} \int_0^1 ds_{q+1} \cdots \int_0^{s_2} ds_1 e^{-it(\gamma_{q+1} s_{q+1} + \cdots + \gamma_1 s_1)} = e^{-it[x_0, x_1, \dots, x_q, 0]} \quad (40)$$

with $x_j = \sum_{k=j+1}^{q+1} \gamma_k$. Changing the integration variable s_1 to $\xi = \gamma_{q+1} s_{q+1} + \cdots + \gamma_1 s_1$, so that ξ ranges from $\gamma_{q+1} s_{q+1} + \cdots + \gamma_2 s_2$ to $\gamma_{q+1} s_{q+1} + \cdots + (\gamma_2 + \gamma_1) s_2$, and $\gamma_1 ds_1 = d\xi$ we get

$$\begin{aligned} & (-it)^{q+1} \int_0^1 ds_{q+1} \cdots \int_0^{s_2} ds_1 e^{-it(\gamma_{q+1} s_{q+1} + \cdots + \gamma_1 s_1)} \\ &= (-it)^{q+1} \frac{1}{\gamma_1} \int_0^1 ds_{q+1} \cdots \int_{\gamma_{q+1} s_{q+1} + \cdots + \gamma_2 s_2}^{\gamma_{q+1} s_{q+1} + \cdots + (\gamma_2 + \gamma_1) s_2} d\xi e^{-it\xi} \\ &= (-it)^q \frac{1}{\gamma_1} \int_0^1 ds_{q+1} \cdots \int_{\gamma_{q+1} s_{q+1} + \cdots + \gamma_2 s_2}^{\gamma_{q+1} s_{q+1} + \cdots + (\gamma_2 + \gamma_1) s_2} d\xi \frac{d}{d\xi} e^{-it\xi} \\ &= (-it)^q \frac{1}{\gamma_1} \int_0^1 ds_{q+1} \cdots \int_0^{s_3} ds_2 \left(e^{-it(\gamma_{q+1} s_{q+1} + \cdots + (\gamma_2 + \gamma_1) s_2)} - e^{-it(\gamma_{q+1} s_{q+1} + \cdots + \gamma_2 s_2)} \right) \\ &= \frac{1}{\gamma_1} \left(\underbrace{(-it)^q \int_0^1 ds_{q+1} \cdots \int_0^{s_3} ds_2 e^{-it(\gamma_{q+1} s_{q+1} + \cdots + (\gamma_2 + \gamma_1) s_2)}}_{(*)} \right. \\ & \quad \left. - \underbrace{(-it)^q \int_0^1 ds_{q+1} \cdots \int_0^{s_3} ds_2 e^{-it(\gamma_{q+1} s_{q+1} + \cdots + \gamma_2 s_2)}}_{(**)} \right), \end{aligned} \quad (41)$$

where from the induction assumption,

$$(*) = e^{-it[x_0, x_2, \dots, x_q, 0]}, \quad \text{and } (**) = e^{-it[x_1, x_2, \dots, x_q, 0]}, \quad (42)$$

with $x_j = \sum_{k=j+1}^{q+1} \gamma_k$. Therefore, we obtain,

$$\begin{aligned}
& (-it)^{q+1} \int_0^1 ds_{q+1} \cdots \int_0^{s_2} ds_1 e^{-it(\gamma_{q+1}s_{q+1} + \cdots + \gamma_1 s_1)} \\
&= \frac{e^{-it[x_0, x_2, \dots, x_q, 0]} - e^{-it[x_1, x_2, \dots, x_q, 0]}}{\gamma_1} = \frac{e^{-it[x_2, \dots, x_q, 0, x_0]} - e^{-it[x_2, \dots, x_q, 0, x_1]}}{x_0 - x_1} \\
&= e^{-it[x_2, \dots, x_q, 0, x_0, x_1]} = e^{-it[x_0, x_1, x_2, \dots, x_q, 0]}.
\end{aligned} \tag{43}$$

This concludes the proof.

Proof of Identity 2: We prove Eq. (5). By definition [8],

$$e^{-it[x_0, \dots, x_q]} = \sum_j \frac{e^{-itx_j}}{\prod_{k \neq j} (x_j - x_k)} \tag{44}$$

(assuming for now that all inputs are distinct). It follows then that

$$e^{-it[x_0, \dots, x_q]} = \sum_j \frac{e^{-it(\Delta_j + x)}}{\prod_{k \neq j} (\Delta_j - \Delta_k)} = e^{-itx} \sum_j \frac{e^{-it\Delta_j}}{\prod_{k \neq j} (\Delta_j - \Delta_k)} = e^{-itx} e^{-it[\Delta_0, \dots, \Delta_q]}. \tag{45}$$

This result holds for arbitrarily close inputs and can be easily generalized to the case where inputs have repeated values, as claimed.