

Hom-Jordan-Malcev-Poisson algebras

Taoufik Chtioui^{1 *}, Sami Mabrouk^{2 †}, Abdenacer Makhlouf^{3 ‡}

1. University of Sfax, Faculty of Sciences Sfax, BP 1171, 3038 Sfax, Tunisia

2. University of Gafsa, Faculty of Sciences Gafsa, 2112 Gafsa, Tunisia

3. Université de Haute Alsace, IRIMAS - Département de Mathématiques, F-68093 Mulhouse, France

Abstract

The purpose of this paper is to provide and study a Hom-type generalization of Jordan-Malcev-Poisson algebras, called Hom-Jordan-Malcev-Poisson algebras. We show that they are closed under twisting by suitable self-maps and give a characterization of admissible Hom-Jordan-Malcev-Poisson algebras. In addition, we introduce the notion of pseudo-Euclidian Hom-Jordan-Malcev-Poisson algebras and describe its T^* -extension. Finally, we generalize the notion of Lie-Jordan-Poisson triple system to the Hom setting and establish its relationships with Hom-Jordan-Malcev-Poisson algebras.

Key words: Hom-JMP-algebra, Hom-Malcev algebra, Hom-Jordan algebra, Hom-flexible algebras, admissible Hom-JMP algebra, Pseudo-Euclidian Hom-JMP algebra, Hom-Lie-Jordan-Poisson triple system.

Introduction

Non-associative algebras become important research field due to their importance in various problems related to physics and other branches of mathematics. The first instances of non-associative Hom-algebras appeared in the study of quasi-deformations of Lie algebras of vector fields. Hom-Lie algebras were first introduced by Hartwig, Larsson and Silvestrov in order to describe q -deformations of Witt and Virasoro algebras using σ -derivations (see [7]). The corresponding associative type objects, called Hom-associative algebras were introduced by Makhlouf and Silvestrov in [11]. Hom-alternative, Hom-Jordan and Hom-flexible algebras were introduced first in [10] and then considered as well as Hom-Malcev algebras in [18].

*E-mail: chtioui.taoufik@yahoo.fr

†E-mail: mabrouksami00@yahoo.fr

‡E-mail: Abdenacer.Makhlouf@uha.fr

Poisson algebras form an important class of non-associative algebras. They are used in many fields of mathematics and physics. They play a fundamental role in Poisson geometry, quantum groups, deformation theory, Hamiltonian mechanics and topological field theories. Poisson algebras are generalized in many ways. If we omit the commutativity of the associative structure, we get the class of non-commutative Poisson algebras. Another way to generalize this class is to replace the associative structure by a Jordan product and the Lie-bracket by a Malcev one. Hence, we obtain a new class of algebras called Jordan-Malcev-Poisson algebras (JMP-algebras) which are defined by a triple $A (A, [,], \circ)$ consisting of a linear equipped with a Malcev bracket and a Jordan structure satisfying the Leibniz rule:

$$[x, y \circ z] = [x, y] \circ z + y \circ [x, z].$$

They were introduced by Ait BenHaddou, Benayadi and Boulmane in [4]. Such algebras can be described in terms of a single bilinear operation, called admissible JMP algebras. This class contains alternative algebras. In the particular case where (A, \circ) is associative commutative, $(A, [,], \circ)$ becomes a Malcev Poisson algebra, a concept introduced first by Shestakov in [15].

The purpose of this paper is to study a twisted generalization of JMP algebras, called Hom-JMP algebras and some other algebraic structures (admissible Hom-JMP algebras). Next, we introduce and study pseudo-Euclidian Hom-JMP algebras, which are Hom-JMP algebras endowed with symmetric invariant non-degenerate bilinear forms. We provide a twist construction and extend the T^* -extension theory to this class of non-associative Hom-algebras. We also construct a generalized Hom-triple systems called Hom-Lie-Jordan-Poisson triple systems from admissible Hom-JMP algebras.

This paper is organized as follows: In section 1, we summarize the definitions and some key constructions of Hom-JMP algebras. In section 2, we study and highlight relationships between Hom-JMP algebras and admissible Hom-JMP algebras. In addition, it is shown that admissible Hom-JMP algebras are power Hom-associative. In Section 3, we introduce the notion of pseudo-Euclidian Hom-JMP algebras and describe its T^* -extension. Section 4 is devoted to the study of a Hom version of Lie-Jordan-Poisson triple system algebras and provide its connection with admissible Hom-JMP algebras.

1 Definitions and Preliminary Results

1.1 Basic definitions

In this section, we introduce Hom-Jordan-Malcev-Poisson algebras (Hom-JMP algebras) as a generalization of both Hom-Poisson algebras, Malcev-Poisson algebras and Jordan-Malcev-Poisson algebras (JMP algebras). We show that Hom-JMP algebras are closed under suitable twisting by weak morphisms.

Let us begin with the basic definitions regarding Hom-algebras. We work over a fixed commutative field \mathbb{K} of characteristic 0.

Definition 1.1. *Let (A, μ, α) be a Hom-algebra.*

1. The Hom-associator $as_A: A^{\otimes 3} \rightarrow A$ is defined as

$$as_A(x, y, z) = \mu(\mu(x, y), \alpha(z)) - \mu(\alpha(x), \mu(y, z)). \quad (1.1)$$

2. The Hom-algebra A is called a Hom-Jordan algebra if it is commutative and satisfies the Hom-Jordan identity

$$as_A(x^2, \alpha(y), \alpha(x)) = 0. \quad (1.2)$$

3. The Hom-Jacobiator $J_A: A^{\otimes 3} \rightarrow A$ is defined as

$$J_A(x, y, z) = \cup_{x,y,z} \mu(\mu(x, y), \alpha(z)), \quad (1.3)$$

where $\cup_{x,y,z}$ denotes the cyclic summation over x, y, z .

4. A Hom-Malcev algebra is a Hom-algebra $(A, [,], \alpha)$ such that $[,]$ is skewsymmetric and that the Hom-Malcev identity

$$J_A(\alpha(x), \alpha(y), [x, z]) = [J_A(x, y, z), \alpha^2(x)] \quad (1.4)$$

is satisfied for all $x, y, z \in A$.

5. A Hom-flexible algebra is a Hom-algebra (A, μ, α) satisfying:

$$as_A(x, y, z) + as_A(z, y, x) = 0,$$

or equivalently, $as_A(x, y, x) = 0$, for all $x, y, z \in A$.

6. A Hom-alternative algebra is a Hom-algebra (A, μ, α) satisfying

$$as_A(x, y, z) + as_A(y, x, z) = as_A(x, y, z) + as_A(x, z, y) = 0,$$

or equivalently, $as_A(x, x, y) = as_A(x, y, y) = 0$, for all $x, y, z \in A$.

Note that any Hom-alternative algebra is Hom-flexible.

Let (A, μ, α) be a Hom-algebra. Define the cyclic Hom-associator S_A by:

$$S_A(x, y, z) = \cup_{x,y,z} as_A(x, y, z).$$

Let us recall the definition of a JMP algebra [9].

Definition 1.2. A JMP algebra $(A, \{, \}, \circ)$ consists of a Malcev algebra $(A, \{, \})$ and a Jordan algebra (A, \circ) such that the Leibniz identity

$$\{x, y \circ z\} = \{x, y\} \circ z + y \circ \{x, z\}$$

is satisfied for all $x, y, z \in A$.

In a JMP algebra $(A, \{, \}, \circ)$, the bracket $\{, \}$ is called the Poisson bracket, and \circ is called the Jordan product. The Leibniz identity says that $\{x, -\}$ is a derivation with respect to the Jordan product.

Hom-Poisson algebras were first introduced in [12] by Makhlouf and Silvestrov. We now define the Hom-type generalization of a JMP algebra.

Definition 1.3. A Hom-JMP algebra $(A, \{, \}, \circ, \alpha)$ consists of a Hom-Malcev algebra $(A, \{, \}, \alpha)$ and a Hom-Jordan algebra (A, \circ, α) such that the Hom-Leibniz identity

$$\{\alpha(x), y \circ z\} = \{x, y\} \circ \alpha(z) + \alpha(y) \circ \{x, z\} \quad (1.5)$$

is satisfied for any $x, y, z \in A$.

In a Hom-JMP algebra $(A, \{, \}, \circ, \alpha)$, the operations $\{, \}$ and \circ are called the Hom-Poisson bracket and the Hom-Jordan product, respectively. By the skewsymmetry of the Hom-Poisson bracket $\{, \}$, the Hom-Leibniz identity is equivalent to

$$\{x \circ y, \alpha(z)\} = \{x, z\} \circ \alpha(y) + \alpha(x) \circ \{y, z\}. \quad (1.6)$$

A JMP algebra is exactly a Hom-JMP algebra with identity twisting map.

Let $(A, \{, \}_A, \circ_A, \alpha_A)$ and $(B, \{, \}_B, \circ_B, \alpha_B)$ be two Hom-JMP algebras. A weak morphism $f : A \rightarrow B$ is a linear map such that:

$$f\{, \}_A = \{, \}_B f^{\otimes 2} \quad \text{and} \quad f \circ_A = \circ_B f^{\otimes 2}.$$

A morphism $f : A \rightarrow B$ is a weak morphism such that $f \alpha_A = \alpha_B f$.

Note that a quadruple $(A, \{, \}, \circ, \alpha)$ is said multiplicative if and only if the twisting map $\alpha : A \rightarrow A$ is a morphism.

The following result says that Hom-JMP algebras are closed under twisting by weak self-morphisms.

Theorem 1.1. Let $(A, \{, \}, \circ, \alpha)$ be a Hom-JMP algebra and $\beta : A \rightarrow A$ be a weak morphism. Then

$$A_\beta = (A, \{, \}_\beta = \beta\{, \}, \circ_\beta = \beta \circ, \beta \alpha)$$

is also a Hom-JMP algebra. Moreover, if A is multiplicative and β is a morphism, then A_β is a multiplicative Hom-JMP algebra.

Proof. In [18] the author proved that $(A, \{, \}_\beta, \beta \alpha)$ is a Hom-Malcev algebra and $(A, \circ_\beta, \beta \alpha)$ is Hom-Jordan algebra.

It remains to show the Hom-Leibniz identity. Let $x, y, z \in A$, we know that

$$\{\alpha(x), y \circ z\} = \{x, y\} \circ \alpha(z) + \alpha(y) \circ \{x, z\}.$$

Now applying β^2 to the previous identity, we obtain

$$\{\beta^2 \alpha(x), \beta^2(y) \circ \beta^2(z)\} = \{\beta^2(x), \beta^2(y)\} \circ \beta^2 \alpha(z) + \beta^2 \alpha(y) \circ \{\beta^2(x), \beta^2(z)\}.$$

That is

$$\{\beta \alpha(x), y \circ_\beta z\}_\beta = \{x, y\}_\beta \circ_\beta \beta \alpha(z) + \beta \alpha(y) \circ_\beta \{x, z\}_\beta.$$

Therefore, A_β is a Hom-JMP algebra. □

Corollary 1.1. Let $(A, \{, \}, \circ)$ be a JMP algebra and $\alpha : A \rightarrow A$ be a JMP morphism. Then $(A, \{, \}_\alpha = \alpha\{, \}, \circ_\alpha = \alpha \circ, \alpha)$ is a multiplicative Hom-JMP algebra.

1.2 Admissible Hom-JMP algebras

Let (A, \cdot, α) be a Hom-algebra. One can define the two following new products:

$$[x, y] = x \cdot y - y \cdot x \text{ and } x \circ y = \frac{1}{2}(x \cdot y + y \cdot x), \text{ for all, } x, y, z \in A.$$

We will denote A^- , (respectively A^+) the algebra A with multiplication $[-, -]$, (respectively \circ).

Lemma 1.1. [18] *Let (A, \cdot, α) be Hom-flexible algebra. Then we have*

$$2S_A = J_{A^-}. \quad (1.7)$$

Lemma 1.2. [11] *A Hom-algebra (A, \cdot, α) is flexible if and only if*

$$[\alpha(x), y \circ z] = [x, y] \circ \alpha(z) + \alpha(y) \circ [x, z]. \quad (1.8)$$

Lemma 1.3. *Let (A, \cdot, α) be a Hom-flexible algebra. Then*

$$J_{A^-}(x^2, \alpha(y), \alpha(x)) = 0, \quad \forall x, y \in A, \text{ where } x^2 = x \cdot x = x \circ x. \quad (1.9)$$

Proof. Let $x, y \in A$. Then we have

$$\begin{aligned} & [[x^2, \alpha(y)], \alpha^2(x)] + [[\alpha(y), \alpha(x)], \alpha(x^2)] + \underbrace{[[\alpha(x), x^2], \alpha^2(y)]}_{=0} \\ &= 2[\alpha^2(x), [y, x] \circ \alpha(x)] + 2[[y, x], \alpha(x)] \circ \alpha^2(x) \\ &= 2\left\{ \alpha^2(x) \cdot ([y, x] \cdot \alpha(x)) + \alpha^2(x) \cdot (\alpha(x) \cdot [y, x]) - ([y, x] \cdot \alpha(x)) \cdot \alpha^2(x) - (\alpha(x) \cdot [y, x]) \cdot \alpha^2(x) \right. \\ & \quad \left. + ([y, x] \cdot \alpha(x)) \cdot \alpha^2(x) - (\alpha(x) \cdot [y, x]) \cdot \alpha^2(x) + \alpha^2(x) \cdot ([y, x] \cdot \alpha(x)) - \alpha^2(x) \cdot (\alpha(x) \cdot [y, x]) \right\} \\ &= 0. \end{aligned}$$

□

The following result gives a characterization of Hom-flexible algebras.

Proposition 1.1. *A Hom-algebra (A, \cdot, α) is flexible if and only if*

$$as_A(x, y, z) = \frac{1}{4}J_{A^-}(x, y, z) + \frac{1}{4}[\alpha(y), [z, x]] + as_{A^+}(x, y, z), \text{ for all, } x, y, z \in A. \quad (1.10)$$

Proof. If (A, \cdot, α) is Hom-flexible, then by Lemma 1.1, we have

$$\begin{aligned} & J_{A^-}(x, y, z) + [\alpha(y), [z, x]] + 4as_{A^+}(x, y, z) \\ &= 2S_A(x, y, z) + \alpha(y)(zx) - \alpha(y)(xz) - (zx)\alpha(y) + (xz)\alpha(y) + (xy)\alpha(z) \\ & \quad + (yx)\alpha(z) + \alpha(z)(xy) + \alpha(z)(yx) - \alpha(x)(yz) - \alpha(x)(zy) - (yz)\alpha(x) - (zy)\alpha(x) \end{aligned}$$

$$\begin{aligned}
&= 2S_A(x, y, z) - as_A(y, z, x) + as_A(y, x, z) - as_A(z, x, y) + as_A(x, z, y) + as_A(x, y, z) - as_A(z, y, x) \\
&= 4as_A(x, y, z).
\end{aligned}$$

Conversely, suppose that Eq. (1.10) holds, then

$$\begin{aligned}
as_A(x, y, x) &= \frac{1}{4}J_{A^-}(x, y, x) + \frac{1}{4}[\alpha(y), [x, x]] + as_{A^+}(x, y, x) \\
&= \frac{1}{4}([\![x, y]\!] + [\![y, x]\!] + [\![x, x]\!] + \alpha(y)) + (x \circ y) \circ \alpha(x) - \alpha(x) \circ (y \circ x) \\
&= 0. \text{ (Since } \circ \text{ is commutative).}
\end{aligned}$$

Hence A is Hom-flexible. □

Corollary 1.2. *Let (A, \cdot, α) be a Hom-flexible algebra. Then*

$$as_A(x^2, \alpha(y), \alpha(x)) = as_{A^+}(x^2, \alpha(y), \alpha(x)).$$

Proof. Straightforward. □

Definition 1.4. *A Hom-algebra (A, \cdot, α) is said to be an admissible Hom-JMP algebra if $(A, [\cdot], \circ, \alpha)$ is a Hom-JMP algebra.*

Remark 1.1. *Given a Hom-JMP algebra $(A, \{ \cdot \}, \circ, \alpha)$, then the vector space A endowed with the morphism α and the product defined by*

$$x \cdot y := \frac{1}{2}\{x, y\} + x \circ y,$$

is an admissible Hom-JMP algebra.

Proposition 1.2. *Let (A, \cdot, α) be an admissible Hom-JMP algebra and $\beta: A \rightarrow A$ be a weak morphism. Then*

$$A_\beta = (A, \cdot_\beta, \beta\alpha)$$

is also an admissible Hom-JMP algebra, where $x \cdot_\beta y = \beta(x) \cdot \beta(y)$. Moreover, if A is multiplicative and β is a morphism, then A_β is a multiplicative admissible Hom-JMP algebra.

Proof. straightforward. □

Example 1.1. *Every Hom-alternative algebra is an admissible Hom-JMP algebra.*

Remark 1.2. *Note that not all admissible Hom-JMP algebras are hom-alternative algebras. Indeed, let A be the three-dimensional algebra of basis $\{e_1, e_2, e_3\}$ defined by:*

★	e_1	e_2	e_3
e_1	0	e_2	$-e_3$
e_2	$-e_2$	0	e_1
e_3	e_3	$-e_1$	0

According to [4], A is an admissible JMP algebra. Let the morphism $\alpha : A \rightarrow A$ defined by:

$$\alpha(e_1) = e_1, \alpha(e_2) = \lambda e_2, \alpha(e_3) = \frac{1}{\lambda} e_3, \lambda \in \mathbb{K} \setminus \{0\}.$$

Then, in view of Proposition 1.2, $(A, \star_\alpha, \alpha)$ is an admissible Hom-JMP algebra. On the other hand,

$$as_A(e_2, e_3, e_3) = (e_2 \star_\alpha \star_\alpha e_3) \star_\alpha \alpha(e_3) - \alpha(e_2) \star_\alpha (e_3 \star_\alpha e_3) = -\frac{1}{\lambda^2} e_3 \neq 0.$$

Then A is not a Hom-alternative algebra.

Remark 1.3. Every admissible Hom-JMP algebra is Hom-flexible.

Theorem 1.2. Let (A, \cdot, α) be a Hom-flexible and Hom-Malcev admissible algebra. Then A is an admissible Hom-JMP algebra if and only if (A, \cdot, α) satisfies the identity

$$R_{\alpha^2(x)} L_{x^2} \alpha = L_{\alpha(x^2)} R_{\alpha(x)} \alpha, \quad \forall x \in A, \quad (1.11)$$

where L_x (respectively, R_x) is the left multiplication (respectively, the right multiplication) by x in the algebra (A, \cdot, α) .

Proof. Since (A, \cdot, α) is Hom-flexible and due to Corollary 1.2, we have

$$as_A(x^2, \alpha(y), \alpha(x)) = as_{A^+}(x^2, \alpha(y), \alpha(x)).$$

The rest of the proof follows immediately. \square

Example 1.2. Consider the five-dimensional Hom-algebra (A, \cdot, α) with basis $\{e_1, \dots, e_5\}$ and multiplication table:

\cdot	e_1	e_2	e_3	e_4	e_5
e_1	0	$e_5 + \frac{1}{2}e_4$	0	$\frac{\nu}{2}e_1$	0
e_2	$e_5 - \frac{1}{2}e_4$	0	0	$-\frac{\nu^{-1}}{2}e_2$	0
e_3	0	0	0	$\frac{\lambda}{2}e_3$	0
e_4	$-\frac{\nu}{2}e_1$	$\frac{\nu^{-1}}{2}e_2$	$-\frac{\lambda}{2}e_3$	$-e_5$	0
e_5	0	0	0	0	0

where $\alpha : A \rightarrow A$ is given by

$$\begin{aligned} \alpha(e_1) &= \nu e_1, & \alpha(e_2) &= \nu^{-1} e_2, & \alpha(e_3) &= \lambda e_3, \\ \alpha(e_4) &= e_4, & \alpha(e_5) &= e_5. \end{aligned}$$

D. Yau, in [18], proved that (A, \cdot, α) is Hom-flexible and Hom-Malcev admissible algebra. In addition, by a direct calculation, we can easily verify that it satisfies condition (1.11). Hence, using Theorem 1.2, we conclude that (A, \cdot, α) is an admissible Hom-JMP algebra.

Now we prove that multiplicative admissible Hom-JMP algebras are power Hom-associative. The power associativity of admissible JMP algebras is shown in [4] (Proposition 2.2). Let us begin by recalling the definition of a power Hom-associative algebra.

Definition 1.5. [17] Let (A, \cdot, α) be a Hom-algebra, $x \in A$, and n be a positive integer.

1. The n th Hom-power $x^n \in A$ is defined by

$$x^1 = x, \quad x^n = x^{n-1} \cdot \alpha^{n-2}(x), \quad n \geq 2. \quad (1.12)$$

2. A is called n th power Hom-associative if

$$x^n = \alpha^{n-i-1}(x^i) \cdot \alpha^{i-1}(x^{n-i}), \quad (1.13)$$

for all $x \in A$ and $i \in \{1, \dots, n-1\}$.

3. A is called power Hom-associative if A is n th power Hom-associative for all $n \geq 2$.

If the twisting map α is the identity map, then the n th power Hom-associativity becomes

$$x^n = x^i \cdot x^{n-i}. \quad (1.14)$$

The class of power Hom-associative algebras contains multiplicative right Hom-alternative algebras and non-commutative Hom-Jordan algebras. Other results for power Hom-associative algebras can be found in [17].

A well-known result of Albert [1] says that an algebra (A, \cdot) is power associative if and only if it is third and fourth power associative, i.e., the condition (1.14) holds for $n = 3, 4$. Moreover, for (1.14) to hold for $n = 3, 4$, it is necessary and sufficient that

$$(x \cdot x) \cdot x = x \cdot (x \cdot x) \quad \text{and} \quad ((x \cdot x) \cdot x) \cdot x = (x \cdot x) \cdot (x \cdot x), \quad \text{for all } x \in A.$$

The Hom-versions of these statements are also proved in [17]. More precisely, a multiplicative Hom-algebra (A, \cdot, α) is power Hom-associative if and only if it is third and fourth power Hom-associative, which in turn is equivalent to

$$x^3 = x^2 \cdot \alpha(x) = \alpha(x) \cdot x^2 \quad \text{and} \quad x^4 = x^3 \cdot \alpha^2(x) = \alpha(x^2) \cdot \alpha(x^2) \quad (1.15)$$

for all $x \in A$.

The following result is the Hom-version of Proposition 2.2 in [4].

Theorem 1.3. Every multiplicative admissible Hom-MJP algebra is power Hom-associative.

Proof. As discussed above, by a result in [17], it is sufficient to prove the two identities in (1.15). The Hom-flexibility implies that

$$0 = as_A(x, x, x) = x^2 \cdot \alpha(x) - \alpha(x) \cdot x^2,$$

which proves the first identity in (1.15). To prove the second equality in (1.15), since (A, \circ, α) is Hom-Jordan and due to Corollary 1.2, we get

$$\begin{aligned} 0 &= as_A(x^2, \alpha(x), \alpha(x)) \\ &= (x^2 \cdot \alpha(x)) \cdot \alpha^2(x) - \alpha(x^2) \cdot (\alpha(x) \cdot \alpha(x)) \\ &= x^3 \cdot \alpha^2(x) - \alpha(x^2) \cdot \alpha(x^2) \end{aligned}$$

We have proved the second identity in (1.15). □

2 Pseudo-Euclidian Hom-JMP algebras

In this section, we extend the notion of pseudo-Euclidian JMP algebra to Hom-JMP algebras and provide some properties. Let (A, \cdot, α) be a Hom-algebra and $B : A \times A \rightarrow \mathbb{K}$ be a bilinear form. B is called:

- (i) symmetric if $B(x, y) = B(y, x), \forall x, y \in A$.
- (ii) non-degenerate if $B(x, y) = 0, \forall y \in A \Rightarrow x = 0$ and if $B(x, y) = 0, \forall x \in A \Rightarrow y = 0$.
- (iii) invariant if $B(x \cdot y, z) = B(x, y \cdot z), \forall x, y, z \in A$.

In this case, (A, \cdot, α, B) will be called a pseudo-Euclidian Hom-algebra if, in addition:

$$B(\alpha(x), y) = B(x, \alpha(y)), \quad \forall x, y \in A.$$

Definition 2.1. Let $(A, \{, \}, \circ, \alpha)$ be a Hom-JMP algebra and $B : A \times A \rightarrow \mathbb{K}$ be a symmetric, non-degenerate and invariant bilinear form on A . We say that $(A, \{, \}, \circ, \alpha, B)$ is a pseudo-Euclidian Hom-JMP algebra if $(A, \{, \}, \alpha, B)$ and (A, \circ, α, B) are pseudo-Euclidian Hom-algebras.

Definition 2.2. A Hom-JMP algebra $(A, \{, \}, \circ, \alpha)$ is called Hom-pseudo-Euclidian if there exists (B, γ) , where B is a symmetric and non-degenerate bilinear form on A and $\gamma : A \rightarrow A$ is an homomorphism such that:

$$B(\alpha(x), y) = B(x, \alpha(y)); B(\{x, y\}, \gamma(z)) = B(\gamma(x), \{y, z\}); B(x \circ y, \gamma(z)) = B(\gamma(x), y \circ z), \quad (2.16)$$

for all $x, y, z \in A$.

Remark 2.1. Note that we recover pseudo-Euclidian Hom-JMP algebras when $\gamma = id$.

Let $(A, \{, \}, \circ, B)$ be a pseudo-Euclidian JMP algebra. We denote by $Aut_S(A, B)$ the set of symmetric automorphisms of A with respect to B , that is automorphisms $\beta : A \rightarrow A$ such that $B(\beta(x), y) = B(x, \beta(y)), \forall x, y \in A$.

Proposition 2.1. Let $(A, \{, \}, \circ, B)$ be a pseudo-Euclidian JMP algebra and $\alpha \in Aut_S(A, B)$. Then $A_\alpha = (A, \{, \}_\alpha, \circ_\alpha, \alpha, B_\alpha)$ is a pseudo-Euclidian Hom-JMP algebra, where for any $x, y \in A$

$$\{x, y\}_\alpha = \{\alpha(x), \alpha(y)\}; x \circ_\alpha y = \alpha(x) \circ \alpha(y); \text{ and } B_\alpha(x, y) = B(\alpha(x), y). \quad (2.17)$$

Proof. By Corollary 1.1, $(A, \{, \}_\alpha, \circ_\alpha, \alpha)$ is a Hom-JMP algebra.

The linear form B_α is nondegenerate since B is nondegenerate and α bijective. Now, let $x, y, z \in A$, then

$$\begin{aligned} B_\alpha(\{x, y\}_\alpha, z) &= B(\alpha(\{\alpha(x), \alpha(y)\}), z) \\ &= B(\{\alpha(x), \alpha(y)\}, \alpha(z)) \\ &= B(\alpha(x), \{\alpha(y), \alpha(z)\}) \\ &= B_\alpha(x, \{y, z\}_\alpha), \end{aligned}$$

and

$$\begin{aligned} B_\alpha(x \circ_\alpha y, z) &= B(\alpha(\alpha(x) \circ \alpha(y)), z) \\ &= B(\alpha(x) \circ \alpha(y), \alpha(z)) = B(\alpha(x), \alpha(y) \circ \alpha(z)) \\ &= B_\alpha(x, y \circ_\alpha z). \end{aligned}$$

On the other hand,

$$B_\alpha(x, y) = B(\alpha(x), y) = B(x, \alpha(y)) = B(\alpha(y), x) = B_\alpha(y, x),$$

and

$$B_\alpha(\alpha(x), y) = B(\alpha(\alpha(x)), y) = B(\alpha(x), \alpha(y)) = B_\alpha(x, \alpha(y)).$$

□

The following result allows to obtain new pseudo-Euclidian Hom-JMP algebras starting from multiplicative pseudo-Euclidian Hom-JMP algebras.

Proposition 2.2. *Let $(A, \{, \}, \circ, \alpha, B)$ be a pseudo-Euclidian Hom-JMP algebra. For any $n \geq 0$, the quadruple*

$$A_n = \left(A, \{, \}_n = \alpha^n \{, \}, \circ_n = \alpha^n \circ, \alpha^{n+1}, B_n \right), \quad (2.18)$$

where B_n is defined for $x, y \in A$ by $B_n(x, y) = B(\alpha^n(x), y)$, determines a pseudo-Euclidian Hom-JMP algebra.

Proof. straightforward. □

We provide here a construction of pseudo-Euclidian Hom-JMP algebra from an arbitrary Hom-JMP algebra (not necessarily pseudo-Euclidian).

Let $(A, \{, \}, \circ)$ be a JMP-algebra and A^* be the dual vector space of the underlying vector space of A . On the vector space $P = A \oplus A^*$, we define the following bracket $\{, \}_P$ and multiplication \circ_P by:

$$\{x + f, y + g\}_P := \{x, y\} + fad_y - gad_x, \quad (2.19)$$

$$(x + f) \circ_P (y + g) := x \circ y + fL_y + gL_x, \quad \forall (x, f), (y, g) \in P, \quad (2.20)$$

where $ad_x(y) = \{x, y\}$ and $L_x(y) = x \circ y$. Moreover, we consider the bilinear form B defined on P by:

$$B(x + f, y + g) = f(y) + g(x), \quad \forall (x, f), (y, g) \in P. \quad (2.21)$$

$(P, \{, \}_P, \circ_P, B)$ is pseudo-Euclidian JMP algebra algebras called the T^* -extension of A by means of A^* .

Proposition 2.3. *Let $(A, \{, \}, \circ)$ be a JMP algebra and $\alpha \in \text{Aut}(A)$. Then the endomorphism $\beta = \alpha + {}^t\alpha$ of P is an automorphism of P if and only if $\text{Im}(\alpha^2 - \text{Id}) \subseteq Z_J(A) \cap Z_M(A)$, where $Z_J(A)$ is the center of (A, \circ) and $Z_M(A)$ is the center of $(A, \{, \})$. Hence, if $\text{Im}(\alpha^2 - \text{Id}) \subseteq Z_J(A) \cap Z_M(A)$ then $(P, \{, \}_{P, \beta}, \circ_{P, \beta}, B_\beta)$ is a regular pseudo-Euclidian Hom-JMP algebra.*

Proof. Let $x, y \in A$ and $f, g \in A^*$.

$$\begin{aligned}\beta(\{x + f, y + g\}_P) &= \beta(\{x, y\} + fad_y - gad_x) \\ &= \alpha(\{x, y\}) + fad_y\alpha - gad_x\alpha,\end{aligned}$$

and

$$\begin{aligned}\{\beta(x + f), \beta(y + g)\}_P &= \{\alpha(x) + f\alpha, \alpha(y) + g\alpha\}_P \\ &= \{\alpha(x), \alpha(y)\} + f\alpha ad_{\alpha(y)} - g\alpha ad_{\alpha(x)},\end{aligned}$$

Then $\beta(\{x + f, y + g\}_P) = \{\beta(x + f), \beta(y + g)\}_P$ if and only if

$$\forall x, y \in A, \quad fad_y\alpha - gad_x\alpha = f\alpha ad_{\alpha(y)} - g\alpha ad_{\alpha(x)}.$$

That is for all $z \in A$

$$f(\{y, \alpha(z)\}) - g(\{x, \alpha(z)\}) = f(\alpha\{\alpha(y), z\}) - g(\alpha\{\alpha(x), z\}).$$

Hence, β is an automorphism of $(P, \{, \}_P)$ if and only if $f(\{x, \alpha(y)\}) = f(\alpha\{\alpha(x), y\})$, $\forall f \in A^*$ $\forall x, y \in A$, which is equivalent to $\{x, \alpha(y)\} = \alpha(\{\alpha(x), y\})$ $\forall x, y \in A$.

As a consequence, β is an automorphism of $(P, \{, \}_P)$ if and only if $\{\alpha^2(x) - x, \alpha(y)\} = 0$ $\forall x, y \in A$, ie. $Im(\alpha^2 - id) \subseteq Z_M(A)$, since $\alpha \in Aut(A)$.

Similarly, β is an automorphism of (P, \circ_P) if and only if $Im(\alpha^2 - id) \subseteq Z_J(A)$. Then $\beta \in Aut(P)$ if and only if $Im(\alpha^2 - id) \subseteq Z_M(A) \cap Z_J(A)$.

In the following we show that β is symmetric with respect to B . Indeed, let $x, y \in A$ and $f, g \in A^*$

$$\begin{aligned}B(\beta(x + f), y + g) &= B(\alpha(x) + f\alpha, y + g) \\ &= f(\alpha(y)) + g(\alpha(x)) \\ &= B(x + f, \alpha(y) + g\alpha) \\ &= B(x + f, \beta(y + g)).\end{aligned}$$

The last assertion is a consequence of the previous calculations and Proposition 2.1. \square

Corollary 2.1. *Let $(A, \{, \}, \circ)$ be a JMP algebra and $\theta \in Aut(A)$ such that $\theta^2 = id$ (θ is an involution). Then $(P, \{, \}_P, \circ_P, B_\beta)$ is a regular pseudo-Euclidian Hom-JMP algebra, where $\beta = \theta +^t \theta$.*

3 Hom-Lie-Jordan-Poisson Triple System

In this section, we generalize the notion of Lie-Jordan-Poisson triple system introduced in [4] to the Hom setting. We provide the relationships of this class of algebras with admissible Hom-JMP algebras. Finally, we endow it with a symmetric non-degenerate invariant bilinear form and give some key constructions.

Definition 3.1. A Hom-Lie triple system is a Hom-triple system $(L, [,], \alpha = (\alpha_1, \alpha_2))$ that satisfies the following conditions

(i) $[x, y, z] = -[y, x, z]$ (left skewsymmetry),

(ii) $[x, y, z] + [y, z, x] + [z, x, y] = 0$ (ternary Jacobi identity),

(iii) $[\alpha_1(x), \alpha_2(y), [u, v, w]] = [[x, y, u], \alpha_1(v), \alpha_2(w)] + [\alpha_1(u), [x, y, v], \alpha_2(w)] + [\alpha_1(u), \alpha_2(v), [x, y, w]],$

for all $u, v, w, x, y, z \in L$.

A particular situation, interesting for our setting, occurs when twisting maps α_i are all equal, that is $\alpha_1 = \alpha_2 = \alpha$, and $\alpha([x, y, z]) = [\alpha(x), \alpha(y), \alpha(z)]$, for all $x, y, z \in L$. The Hom-Lie triple system $(L, [,], \alpha)$ is said to be multiplicative.

Definition 3.2. A Hom-Lie-Jordan-Poisson triple system is a quadruple $(A, \{, \}, \circ, \alpha)$ such that

(i) (A, \circ, α) is a Hom Jordan algebra.

(ii) $(A, \{, \}, \alpha)$ is a multiplicative Hom-Lie triple system.

(iii) $\{\alpha(x), \alpha(y), z \circ t\} = \{x, y, z\} \circ \alpha(t) + \alpha(z) \circ \{x, y, t\}, \forall x, y, z, t \in A$.

In the case where (A, \circ, α) is a commutative Hom-associative algebra, the quadruple $(A, \{, \}, \circ, \alpha)$ is called a Hom-Lie-Poisson triple system.

Lemma 3.1. [3] Let $(A, \{, \}, \alpha)$ be a Hom-Malcev algebra. Then $(A, \{, \}, \alpha^2)$ is a multiplicative Hom-Lie triple system, where

$$\{x, y, z\} = 2\{\{x, y\}, \alpha(z)\} - \{\{y, z\}, \alpha(x)\} - \{\{z, x\}, \alpha(y)\}, \forall x, y, z \in A. \quad (3.22)$$

Proposition 3.1. Let $(A, \{, \}, \circ, \alpha)$ be a Hom-JMP algebra. Then the quadruple $(A, \{, \}, \circ_\alpha, \alpha^2)$ is a Hom-Lie-Jordan-Poisson triple system, where

$$\{x, y, z\} = 2\{\{x, y\}, \alpha(z)\} - \{\{y, z\}, \alpha(x)\} - \{\{z, x\}, \alpha(y)\}, \forall x, y, z \in A, \quad (3.23)$$

$$x \circ_\alpha y = \alpha(x) \circ \alpha(y), \forall x, y \in A. \quad (3.24)$$

Proof. Let $x, y, z, t \in A$, we have

$$\begin{aligned} \{\alpha^2(x), \alpha^2(y), z \circ_\alpha t\} &= 2\{\{\alpha^2(x), \alpha^2(y)\}, \alpha(z \circ_\alpha t)\} - \{\{\alpha^2(y), z \circ_\alpha t\}, \alpha^3(x)\} - \{\{z \circ_\alpha t, \alpha^2(x)\}, \alpha^3(y)\} \\ &= \alpha(2\{\{\alpha(x), \alpha(y)\}, \alpha(z) \circ \alpha(t)\} - \{\{\alpha(y), z \circ t\}, \alpha^2(x)\} - \{\{z \circ t, \alpha(x)\}, \alpha^2(y)\}) \\ &= \alpha(2\{\{x, y\}, \alpha(z)\} \circ \alpha^2(t) + 2\alpha^2(z) \circ \{\{x, y\}, \alpha(t)\} - \{\{y, z\} \circ \alpha(t), \alpha^2(x)\} \\ &\quad - \{\alpha(z) \circ \{y, t\}, \alpha^2(x)\} - \{\alpha(z) \circ \{t, x\}, \alpha^2(y)\} - \{\{z, x\} \circ \alpha(t), \alpha^2(y)\}) \\ &= \alpha(2\{\{x, y\}, \alpha(z)\} \circ \alpha^2(t) + 2\alpha^2(z) \circ \{\{x, y\}, \alpha(t)\} - \{\alpha(y), \alpha(z)\} \circ \{\alpha(t), \alpha(x)\} \\ &\quad - \{\{y, z\}, \alpha(x)\} \circ \alpha^2(t) - \{\alpha(z), \alpha(x)\} \circ \{\alpha(y), \alpha(t)\} - \alpha^2(z) \circ \{\{y, t\}, \alpha(x)\} \\ &\quad - \{\alpha(z), \alpha(y)\} \circ \{\alpha(t), \alpha(x)\} - \alpha^2(z) \circ \{\{t, x\}, \alpha(y)\}) \end{aligned}$$

$$\begin{aligned}
& - \{\alpha(z), \alpha(x)\} \circ \{\alpha(t), \alpha(y)\} - \{\{z, x\}, \alpha(y)\} \circ \alpha^2(t) \\
& = \alpha(\alpha^2(z) \circ (\{\{x, y\}, \alpha(t)\} - \{\{y, t\}, \alpha(x)\} - \{\{t, x\}, \alpha(y)\})) \\
& \quad + (2\{\{x, y\}, \alpha(z)\} - \{\{y, z\}, \alpha(x)\} - \{\{z, x\}, \alpha(y)\}) \circ \alpha^2(t) \\
& = \alpha^2(z) \circ_{\alpha} \{x, y, t\} + \{x, y, z\} \circ_{\alpha} \alpha^2(t).
\end{aligned}$$

□

Let $(A, \{, , \}, \alpha)$ be a multiplicative Hom-Lie triple system and $B : A \times A \rightarrow \mathbb{K}$ be a symmetric non-degenerate bilinear form. We say that B is invariant if

$$B(L(x, y)(z), t) = -B(z, L(x, y)(t)), \quad \forall x, y, z, t \in A,$$

where $L(x, y)(z) = \{x, y, z\}$. In this case $(A, \{, , \}, \alpha, B)$ will be called a pseudo-Euclidian Hom-Lie triple system if, in addition:

$$B(\alpha(x), y) = B(x, \alpha(y)), \quad \forall x, y \in A.$$

Definition 3.3. Let $(A, \{, , \}, \circ, \alpha)$ be a Hom-Lie-Jordan-Poisson triple system and $B : A \times A \rightarrow \mathbb{K}$ be a bilinear form on A . We say that $(A, \{, , \}, \circ, \alpha, B)$ is a pseudo-Euclidian Hom-Lie-Jordan-Poisson triple system if $(A, \{, , \}, \alpha, B)$ is a pseudo-Euclidian Hom-Lie triple system and (A, \circ, α, B) is a pseudo-Euclidian Hom-Jordan algebra.

Definition 3.4. A Hom-Lie-Jordan-Poisson triple system $(A, \{, , \}, \circ, \alpha)$ is called Hom-pseudo-Euclidian if there exists (B, γ) , where B is a symmetric and non-degenerate bilinear form on A and $\gamma : A \rightarrow A$ is a homomorphism such that:

$$\begin{aligned}
& B(\alpha(x), y) = B(x, \alpha(y)), \\
& B(L(x, y)(z), \gamma(t)) = -B(\gamma(z), L(x, y)(t)), \\
& B(x \circ y, \gamma(z)) = B(\gamma(x), y \circ z),
\end{aligned}$$

for all $x, y, z, t \in A$.

Corollary 3.1. Let $(A, \{, , \}, \circ, \alpha, B)$ be a pseudo-Euclidian Hom-JMP algebra. Then the 6-uplet $(A, \{, , \}, \circ_{\alpha}, \alpha^2, B, \alpha)$ is a Hom-pseudo-Euclidian Hom-Lie-Jordan-Poisson triple system, where $\{, , \}$ and \circ_{α} are defined in (3.23) and (3.24).

Proof. Let $x, y, z, t \in A$, we have

$$\begin{aligned}
B(\{x, y, z\}, \alpha(t)) & = B(2\{\{x, y\}, \alpha(z)\} - \{\{y, z\}, \alpha(x)\} - \{\{z, x\}, \alpha(y)\}, \alpha(t)) \\
& = B(2\{\{x, y\}, \alpha(z)\}, \alpha(t)) - B(\{\{y, z\}, \alpha(x)\}, \alpha(t)) - B(\{\{z, x\}, \alpha(y)\}, \alpha(t)) \\
& = -B(\alpha(z), 2\{\{x, y\}, \alpha(t)\}) - B(\{y, z\}, \{\alpha(x), \alpha(t)\}) - B(\{z, x\}, \{\alpha(y), \alpha(t)\}) \\
& = -B(\alpha(z), 2\{\{x, y\}, \alpha(t)\}) - B(\{\alpha(y), \alpha(z)\}, \{x, t\}) - B(\{\alpha(z), \alpha(x)\}, \{y, t\}) \\
& = -B(\alpha(z), 2\{\{x, y\}, \alpha(t)\}) - B(\alpha(z), \{\{x, t\}, \alpha(y)\}) - B(\alpha(z), \{\{t, y\}, \alpha(x)\}) \\
& = -B(\alpha(z), 2\{\{x, y\}, \alpha(t)\} - \{\{x, t\}, \alpha(y)\} - \{\{t, y\}, \alpha(x)\}) \\
& = -B(\alpha(z), \{x, y, t\}),
\end{aligned}$$

which end proof. □

References

- [1] A.A. Albert, On the power-associativity of rings, *Summa Brazil. Math.* 2 (1948) 21–32.
- [2] A.A. Albert, Power-associative rings, *Trans. Amer. Math. Soc.* 64 (1948) 552–593.
- [3] S. Attan, A. N. Issa, Hom-Lie Triple System and Hom-Bol Algebra Structures on Hom-Maltsev and Right Hom-Alternative Algebras. *International Journal of Mathematics and Mathematical Sciences* 2018 (2018).
- [4] M. Ait Ben Haddou, S. Benayadi and S. Boulmane, Malcev-Poisson-Jordan algebras. *Journal of Algebra and Its Applications* Vol. 15, No. 9 (2016) 1650159
- [5] S. Benayadi and A. Makhlouf, Hom-Lie algebras with symmetric invariant nondegenerate bilinear forms, *Journal of Geometry and Physics*, 76, pp 38–60 (2014)
- [6] M. Goze and E. Remm, Poisson algebras in terms of non-associative algebras, *J. Alg.* 320 (2008) 294–317.
- [7] J.T. Hartwig, D. Larsson, and S.D. Silvestrov, Deformations of Lie algebras using σ -derivations, *J. Alg.* 295 (2006) 314–361.
- [8] M. Kontsevich, Deformation quantization of Poisson manifolds, *Lett. Math. Phys.* 66 (2003) 157–216.
- [9] F. Kubo, Finite-dimensional non-commutative Poisson algebras, *J. Pure Appl. Alg.* 113 (1996) 307–314.
- [10] A. Makhlouf, Hom-alternative algebras and Hom-Jordan algebras, *International Elec. J of Alg.* Volume 8 (2010) 177–190
- [11] A. Makhlouf and S. Silvestrov, Hom-algebra structures, *J. Gen. Lie Theory Appl.* 2 (2008) 51–64.
- [12] A. Makhlouf and S. Silvestrov, Hom-algebras and Hom-coalgebras, *J. Alg. Appl.* 9 (2010) 1–37.
- [13] M. Markl and E. Remm, Algebras with one operation including Poisson and other Lie-admissible algebras, *J. Alg.* 299 (2006) 171–189.
- [14] P. Schaller and T. Strobl, Poisson structure induced (topological) field theories, *Mod. Phys. Lett. A* 9 (1994) 3129–3136.
- [15] I. P. Shestakov, Speciality problem for Malcev algebras and Poisson Malcev algebras, in *Nonassociative Algebra and its Applications, Lecture Notes in Pure and Applied Mathematics*, Vol. 211 (Dekker, New York, 2000), pp. 365–371.
- [16] I. Vaisman, *Lectures on the geometry of Poisson manifolds*, Birkhäuser, Basel, 1994.

- [17] D. Yau, power Hom-associative algebras, arXiv:1007.4118.
- [18] D. Yau, Hom-Malcev, Hom-alternative and Hom-Jordan algebras. International J. of Alg. Volume 11 (2012) 177–217.