

PROOF OF SOME SUPERCONGRUENCES CONCERNING TRUNCATED HYPERGEOMETRIC SERIES

CHEN WANG AND DIAN-WANG HU*

ABSTRACT. In this paper, we prove some supercongruences concerning truncated hypergeometric series. For example, we show that for any prime $p > 3$ and positive integer r ,

$$\sum_{k=0}^{p^r-1} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k \equiv p^r + \frac{7}{6} p^{r+3} B_{p-3} \pmod{p^{r+4}}$$

and

$$\sum_{k=0}^{(p^r-1)/2} (4k+1) \frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4} \equiv p^r + \frac{7}{6} p^{r+3} B_{p-3} \pmod{p^{r+4}},$$

where $(x)_k = x(x+1)\cdots(x+k-1)$ is the Pochhammer symbol and B_0, B_1, B_2, \dots are Bernoulli numbers. These two congruences confirm conjectures of Sun [Sci. China Math. 54 (2011), 2509–2535] and Guo [Adv. Appl. Math. 120 (2020), Art. 102078], respectively.

1. INTRODUCTION

For $m, n \in \mathbb{N} = \{0, 1, 2, \dots\}$, the truncated hypergeometric series ${}_{m+1}F_m$ is defined by

$${}_{m+1}F_m \left[\begin{matrix} x_0 & x_1 & \cdots & x_m \\ & y_1 & \cdots & y_m \end{matrix} \middle| z \right]_n = \sum_{k=0}^n \frac{(x_0)_k (x_1)_k \cdots (x_m)_k}{(y_1)_k \cdots (y_m)_k} \cdot \frac{z^k}{k!},$$

where $(x)_k = x(x+1)\cdots(x+k-1)$ is the Pochhammer symbol. During the past few decades, supercongruences concerning truncated hypergeometric series have been widely studied (cf. [2, 8, 10, 15, 16, 20, 21, 22, 23]).

In 2011, Sun [15] proposed some conjectural supercongruences which relate truncated hypergeometric series to Euler numbers and Bernoulli numbers (see [15] for the definitions of Euler numbers and Bernoulli numbers). For example, he conjectured that for any prime $p > 3$ we have

$$\sum_{k=0}^{(p-1)/2} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k \equiv p + 2(-1)^{(p-1)/2} p^3 E_{p-3} \pmod{p^4}, \quad (1.1)$$

2020 *Mathematics Subject Classification*. Primary 33C20, 11A07; Secondary 11B65, 05A10.

Key words and phrases. Truncated hypergeometric series, binomial coefficients, supercongruences, WZ pairs.

*Corresponding author.

and for any $r \in \mathbb{Z}^+$ we have

$$\sum_{k=0}^{p^r-1} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k \equiv p^r + \frac{7}{6} p^{r+3} B_{p-3} \pmod{p^{r+4}}, \quad (1.2)$$

where E_0, E_1, E_2, \dots are Euler numbers and B_0, B_1, B_2, \dots are Bernoulli numbers. Note that $ak+1 = (1+1/a)_k / (1/a)_k$. Thus the sums in (1.1) and (1.2) are actually the truncated hypergeometric series. In 2012, using the WZ method (cf. [12]), Guillera and Zudilin [2] proved that

$$\sum_{k=0}^{p-1} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k \equiv \sum_{k=0}^{(p-1)/2} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k \equiv p \pmod{p^3}, \quad (1.3)$$

which is (1.2) modulo p^3 with $r = 1$. In 2019, Mao and Zhang [10] confirmed (1.1) via a WZ pair found by Guillera and Zudilin [2]. The reader is referred to [17] for further conjectures involving the sums in (1.1) and (1.2).

Our first theorem confirms (1.2).

Theorem 1.1. *For any prime $p > 3$ and integer $r \geq 1$, we have*

$$\sum_{k=0}^{p^r-1} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k \equiv p^r + \frac{7}{6} p^{r+3} B_{p-3} \pmod{p^{r+4}}. \quad (1.4)$$

Using the same technique as the one used in the proof of Theorem 1.1 and using (1.1), we can also prove that for any prime $p > 3$ and positive integer r

$$\sum_{k=0}^{(p^r-1)/2} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k \equiv p^r + 2(-1)^{(p-1)/2} p^{r+2} E_{p-3} \pmod{p^{r+3}}.$$

It is worth mentioning that Guo and Schlosser [4] obtained two different q -analogues of (1.3), and similarly to (1.3), they conjectured that for any odd prime p

$$\sum_{k=0}^{(p+1)/2} (3k-1) \frac{\left(-\frac{1}{2}\right)_k^2 \left(\frac{1}{2}\right)_k}{(1)_k^3} 4^k \equiv p \pmod{p^3}$$

which has been confirmed by the first author [21] by extending it to the modulus p^4 case.

In 2011, as a refinement of the (C.2) supercongruence of Van Hamme [20], Long [8] proved that

$$\sum_{k=0}^{(p-1)/2} (4k+1) \frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4} \equiv p \pmod{p^4}. \quad (1.5)$$

Guo and Wang [5] obtained a generalization of (1.5). For any prime $p > 3$ and positive integer r , they proved that

$$\sum_{k=0}^{(p^r-1)/2} (4k+1) \frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4} \equiv p^r \pmod{p^{r+3}}. \quad (1.6)$$

Our next theorem confirms a conjecture of Guo [3, Conjecture 6.2] which extends (1.6) to the modulus p^{r+4} case.

Theorem 1.2. *Let $p > 3$ be a prime and r a positive integer. Then*

$$\sum_{k=0}^{(p^r-1)/2} (4k+1) \frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4} \equiv p^r + \frac{7}{6} p^{r+3} B_{p-3} \pmod{p^{r+4}}. \quad (1.7)$$

Note that Guo [3] proved that for any odd prime p and positive integer r

$$\sum_{k=0}^{(p^r-1)/2} (4k+1) \frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4} \equiv \sum_{k=0}^{p^r-1} (4k+1) \frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4} \pmod{p^{r+4}}.$$

Clearly, the two sums in (1.4) and (1.7) are the same modulo p^{r+4} . Guo [3, Conjecture 6.3] conjectured that it is also true for $p = 3$.

Theorem 1.3. *Let p be an odd prime and r a positive integer. Then*

$$\sum_{k=0}^{p^r-1} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k \equiv \sum_{k=0}^{(p^r-1)/2} (4k+1) \frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4} \pmod{p^{r+4}}. \quad (1.8)$$

Note that Guo [3, Conjecture 6.4] also conjectured a q -analogue of (1.8).

Our main strategy to prove Theorems 1.1–1.3 is using the WZ method (the reader is referred to [2, 12, 23] for further details and some well known WZ pairs). In fact, the case $r = 1$ is easy to deal with since the denominators appearing in the WZ pairs are not divisible by p . However, the case $r \geq 2$ is very sophisticated. In this case, we need to reduce the sums in (1.4) and (1.7) to the case $r = 1$ via some complicated calculation.

The paper is organized as follows. In both Sections 2 and 3, we shall first establish preliminary results which connect the case $r \geq 2$ with the case $r = 1$ and play important role in the proof of Theorem 1.3. Then we will use the preliminary results to prove Theorems 1.1 and 1.2. In the end of Section 3, we shall give the proof of Theorem 1.3.

2. PROOF OF THEOREM 1.1

We first establish the following result.

Theorem 2.1. *For any odd prime p and positive integer r we have*

$$\frac{1}{p^r} \sum_{k=0}^{p^r-1} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k \equiv \frac{1}{p} \sum_{k=0}^{p-1} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k \pmod{p^4}.$$

Define the multiple harmonic sum (cf. [19]) as follows:

$$H_n(s_1, s_2, \dots, s_r) = \sum_{1 \leq k_1 < k_2 < \dots < k_r \leq n} \frac{1}{k_1^{s_1} k_2^{s_2} \dots k_r^{s_r}},$$

where $n \geq r > 0$ and each s_i is a positive integer. Multiple harmonic sums have many congruence properties. For example, for any prime $p > s + 2$, Sun [14] proved that

$$H_{p-1}(s) \equiv \begin{cases} -\frac{s(s+1)}{2s+4} p^2 B_{p-s-2} \pmod{p^3} & \text{if } 2 \nmid s, \\ \frac{s}{s+1} p B_{p-s-1} \pmod{p^2} & \text{if } 2 \mid s; \end{cases} \quad (2.1)$$

for any $p > 5$, Kh. Hessami Pilehrood and T. Hessami Pilehrood [6, Lemma 3] proved that

$$H_{p-1}(1, 2) \equiv -\frac{3H_{p-1}(1)}{p^2} - \frac{5H_{p-1}(3)}{12} \pmod{p^3}. \quad (2.2)$$

Lemma 2.1. *For any odd prime p and positive integer r we have*

$$\begin{aligned} p^{2r} \sum_{n=1}^{p^r-1} \frac{1}{n^2} \binom{2n}{n} &\equiv p^2 \sum_{n=1}^{p-1} \frac{1}{n^2} \binom{2n}{n} \pmod{p^4}, \\ p^{2r} \sum_{n=1}^{p^r-1} \frac{H_{n-1}(1)}{n} \binom{2n}{n} &\equiv p^2 \sum_{n=1}^{p-1} \frac{H_{n-1}(1)}{n} \binom{2n}{n} \pmod{p^4}, \\ p^{3r} \sum_{n=1}^{p^r-1} \frac{H_{n-1}(1)^2}{n} \binom{2n}{n} &\equiv p^3 \sum_{n=1}^{p-1} \frac{H_{n-1}(1)^2}{n} \binom{2n}{n} \pmod{p^4}, \\ p^{3r} \sum_{n=1}^{p^r-1} \frac{H_{n-1}(2)}{n} \binom{2n}{n} &\equiv p^2 \sum_{n=1}^{p-1} \frac{H_{n-1}(2)}{n} \binom{2n}{n} \pmod{p^4}, \\ p^{3r} \sum_{n=1}^{p^r-1} \frac{H_{n-1}(1)}{n^2} \binom{2n}{n} &\equiv p^2 \sum_{n=1}^{p-1} \frac{H_{n-1}(1)}{n^2} \binom{2n}{n} \pmod{p^4}, \\ p^{2r} \sum_{k=1}^{(p^r-1)/2} \frac{1}{(2k-1)^2} &\equiv p^2 \sum_{k=1}^{(p-1)/2} \frac{1}{(2k-1)^2} \pmod{p^4}, \\ p^{3r} \sum_{k=1}^{(p^r-1)/2} \frac{H_{2k-2}(1)}{(2k-1)^2} &\equiv p^3 \sum_{k=1}^{(p-1)/2} \frac{H_{2k-2}(1)}{(2k-1)^2} \pmod{p^4}, \\ p^{3r} \sum_{k=1}^{(p^r-1)/2} \frac{H_{k-1}(1)}{(2k-1)^2} &\equiv p^3 \sum_{k=1}^{(p-1)/2} \frac{H_{k-1}(1)}{(2k-1)^2} \pmod{p^4}. \end{aligned}$$

Proof. We only prove the second congruence, since the other ones can be showed in a similar way. We shall finish the proof by induction on r .

Clearly, the second congruence holds for $r = 1$. Assume that it holds for $r = k > 1$. Now

$$\begin{aligned} p^{2k+2} \sum_{n=1}^{p^{k+1}-1} \frac{H_{n-1}(1)}{n} \binom{2n}{n} &= p^{2k+2} \sum_{\substack{n=1 \\ p \nmid n}}^{p^{k+1}-1} \frac{H_{n-1}(1)}{n} \binom{2n}{n} + p^{2k+2} \sum_{\substack{n=1 \\ p|n}}^{p^{k+1}-1} \frac{H_{n-1}(1)}{n} \binom{2n}{n} \\ &\equiv p^{2k+2} \sum_{\substack{n=1 \\ p|n}}^{p^{k+1}-1} \frac{H_{n-1}(1)}{n} \binom{2n}{n} \\ &= p^{2k+1} \sum_{n=1}^{p^k-1} \frac{H_{pn-1}(1)}{n} \binom{2pn}{pn} \pmod{p^4}, \end{aligned}$$

where we used the fact that $\text{ord}_p(H_{n-1}(1)) \geq -k$ for $1 \leq n \leq p^{k+1} - 1$. Note that

$$\text{ord}_p(p^{2k+1}/n) \geq k + 2 \geq 4 \text{ for } 1 \leq n \leq p^k.$$

Hence for $1 \leq n \leq p^k$ we have

$$\frac{p^{2k+1} H_{pn-1}(1)}{n} \equiv \frac{p^{2k+1}}{n} \sum_{\substack{j=1 \\ p|j}}^{pn-1} \frac{1}{j} = \frac{p^{2k} H_{n-1}(1)}{n} \pmod{p^4}$$

and

$$\frac{p^{2k} H_{n-1}(1)}{n} \equiv 0 \pmod{p^2}.$$

By the well-known Kazandzidis congruence (cf. [13, p. 380]) we have for any odd prime p

$$\binom{2pn}{pn} \equiv \binom{2n}{n} \pmod{p^2}.$$

Combining the above and by the induction hypothesis we arrive at

$$p^{2k+2} \sum_{n=1}^{p^{k+1}-1} \frac{H_{n-1}(1)}{n} \binom{2n}{n} \equiv p^{2k} \sum_{n=1}^{p^k-1} \frac{H_{n-1}(1)}{n} \binom{2n}{n} \equiv p^2 \sum_{n=1}^{p-1} \frac{H_{n-1}(1)}{n} \binom{2n}{n} \pmod{p^4}.$$

We are done. □

Proof of Theorem 2.1. As in [2], we shall use the following WZ pair

$$F(n, k) = (3n + 2k + 1) \frac{(\frac{1}{2})_n (\frac{1}{2} + k)_n^2}{(1)_n^3} 4^n$$

and

$$G(n, k) = -\frac{(\frac{1}{2})_n (\frac{1}{2} + k)_n^2}{(1)_{n-1}^3} 4^n.$$

Then we have

$$F(n, k - 1) - F(n, k) = G(n + 1, k) - G(n, k)$$

and

$$\sum_{k=0}^{p^r-1} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k = \sum_{n=0}^{p^r-1} F(n, 0).$$

Clearly,

$$\begin{aligned} \sum_{n=0}^{p^r-1} F(n, 0) &= \sum_{n=0}^{p^r-1} \sum_{k=1}^{(p^r-1)/2} (F(n, k-1) - F(n, k)) + \sum_{n=0}^{p^r-1} F\left(n, \frac{p^r-1}{2}\right) \\ &= \sum_{k=1}^{(p^r-1)/2} \sum_{n=0}^{p^r-1} (G(n+1, k) - G(n, k)) + \sum_{n=0}^{p^r-1} F\left(n, \frac{p^r-1}{2}\right) \\ &= \sum_{k=1}^{(p^r-1)/2} G(p^r, k) + \sum_{n=0}^{p^r-1} F\left(n, \frac{p^r-1}{2}\right), \end{aligned}$$

where the last follows from the fact that $G(0, k) = 0$. It suffices to show

$$\frac{1}{p^r} \sum_{n=0}^{p^r-1} F\left(n, \frac{p^r-1}{2}\right) \equiv \frac{1}{p} \sum_{n=0}^{p-1} F\left(n, \frac{p-1}{2}\right) \pmod{p^4} \quad (2.3)$$

and

$$\frac{1}{p^r} \sum_{k=1}^{(p^r-1)/2} G(p^r, k) \equiv \frac{1}{p} \sum_{k=1}^{(p-1)/2} G(p, k) \pmod{p^4}. \quad (2.4)$$

We first consider (2.3). It is easy to see that

$$F\left(n, \frac{p^r-1}{2}\right) = \begin{cases} \frac{p^{2r}(3n+p^r)}{4n^2} \binom{2n}{n} \frac{(1+\frac{p^r}{2})_{n-1}^2}{(1)_{n-1}^2} & \text{if } n \geq 1, \\ p^r & \text{if } n = 0. \end{cases}$$

Therefore,

$$\frac{1}{p^r} \sum_{n=1}^{p^r-1} F\left(n, \frac{p^r-1}{2}\right) = \frac{3p^r}{4} \sum_{n=1}^{p^r-1} \frac{(1+\frac{p^r}{2})_{n-1}^2}{(1)_{n-1}^2} \cdot \frac{\binom{2n}{n}}{n} + \frac{p^{2r}}{4} \sum_{n=1}^{p^r-1} \frac{(1+\frac{p^r}{2})_{n-1}^2}{(1)_{n-1}^2} \cdot \frac{\binom{2n}{n}}{n^2}.$$

For $1 \leq n \leq p^r-1$, it is clear that $\text{ord}_p(n) \leq r-1$. Note that

$$\frac{(1+\frac{p^r}{2})_{n-1}}{(1)_{n-1}} = 1 + \frac{p^r}{2} H_{n-1}(1) + \frac{p^{2r}}{4} H_{n-1}(1, 1) + \dots$$

and

$$\text{ord}_p\left(H_{n-1}\left(\overbrace{1, 1, \dots, 1}^{d \text{ s } 1}\right)\right) \geq -d(r-1).$$

Thus we have

$$\frac{(1 + \frac{p^r}{2})_{n-1}}{(1)_{n-1}} \equiv 1 + \frac{p^r}{2} H_{n-1}(1) + \frac{p^{2r}}{8} H_{n-1}(1)^2 - \frac{p^{2r}}{8} H_{n-1}(2) \pmod{p^3},$$

where we have used $H_{n-1}(1, 1) = (H_{n-1}(1)^2 - H_{n-1}(2))/2$. Now by Lemma 2.1,

$$\begin{aligned} \frac{1}{p^r} \sum_{n=1}^{p^r-1} F\left(n, \frac{p^r-1}{2}\right) &\equiv \frac{3p^r}{4} \sum_{n=1}^{p^r-1} \frac{\binom{2n}{n}}{n} \left(1 + \frac{p^r}{2} H_{n-1}(1) + \frac{p^{2r}}{8} H_{n-1}(1)^2 - \frac{p^{2r}}{8} H_{n-1}(2)\right)^2 \\ &\quad + \frac{p^{2r}}{4} \sum_{n=1}^{p^r-1} \frac{\binom{2n}{n}}{n^2} \left(1 + \frac{p^r}{2} H_{n-1}(1)\right)^2 \\ &\equiv \frac{3p^r}{4} \sum_{n=1}^{p^r-1} \frac{\binom{2n}{n}}{n} \left(1 + p^r H_{n-1}(1) + \frac{p^{2r}}{2} H_{n-1}(1)^2 - \frac{p^{2r}}{4} H_{n-1}(2)\right) \\ &\quad + \frac{p^{2r}}{4} \sum_{n=1}^{p^r-1} \frac{\binom{2n}{n}}{n^2} (1 + p^r H_{n-1}(1)) \\ &\equiv \frac{3p}{4} \sum_{n=1}^{p-1} \frac{\binom{2n}{n}}{n} \left(1 + p H_{n-1}(1) + \frac{p^2}{2} H_{n-1}(1)^2 - \frac{p^2}{4} H_{n-1}(2)\right) \\ &\quad + \frac{p^2}{4} \sum_{n=1}^{p-1} \frac{\binom{2n}{n}}{n^2} (1 + p H_{n-1}(1)) \pmod{p^4}, \end{aligned} \tag{2.5}$$

where in the last step we have used the fact (cf. [18, Theorem 1.3]) that

$$p^{r-1} \sum_{k=1}^{p^r-1} \frac{\binom{2k}{k}}{k} \equiv \begin{cases} 2 \pmod{p^3} & \text{if } p = 2, \\ 5 \pmod{p^3} & \text{if } p = 3, \\ \frac{8}{9} p^2 B_{p-3} \pmod{p^3} & \text{otherwise.} \end{cases} \tag{2.6}$$

Note that in (2.5) we actually obtain a result which is independent of r . Thus we have proved (2.3).

Now we consider (2.4). Clearly,

$$\frac{1}{p^r} \sum_{k=1}^{(p^r-1)/2} G(p^r, k) = -\frac{4p^{2r}}{16p^r} \binom{2p^r}{p^r}^3 \sum_{k=1}^{(p^r-1)/2} \frac{(\frac{1}{2} + p^r)_{k-1}^2}{(2k-1)^2 (\frac{1}{2})_{k-1}^2}.$$

By a similar argument as above, $\text{ord}_p(2k-1) \leq r-1$ and

$$\frac{(\frac{1}{2} + p^r)_{k-1}}{(\frac{1}{2})_{k-1}} \equiv 1 + 2p^r H_{2k-2}(1) - p^r H_{k-1}(1) \pmod{p^2}.$$

Therefore, by Lemma 2.1 we have

$$\begin{aligned} \frac{1}{p^r} \sum_{k=1}^{(p^r-1)/2} G(p^r, k) &\equiv -\frac{4p^{2r}}{16p^r} \binom{2p^r}{p^r}^3 \sum_{k=1}^{(p^r-1)/2} \frac{1 + 4p^r H_{2k-2}(1) - 2p^r H_{k-1}(1)}{(2k-1)^2} \\ &\equiv -\frac{4p^2}{16p^r} \binom{2p^r}{p^r}^3 \sum_{k=1}^{(p-1)/2} \frac{1 + 4p H_{2k-2}(1) - 2p H_{k-1}(1)}{(2k-1)^2} \pmod{p^4}. \end{aligned}$$

Note that

$$\sum_{k=1}^{(p-1)/2} \frac{1}{(2k-1)^2} = H_{p-1}(2) - \frac{1}{4} H_{(p-1)/2}(2) \equiv 0 \pmod{p}.$$

Hence we have

$$\sum_{k=1}^{(p-1)/2} \frac{1 + 4p H_{2k-2}(1) - 2p H_{k-1}(1)}{(2k-1)^2} \equiv 0 \pmod{p}.$$

Then from the Kazandzidis congruence and Fermat's little theorem, we immediately obtain

$$\frac{1}{p^r} \sum_{k=1}^{(p^r-1)/2} G(p^r, k) \equiv -2p^2 \binom{2p}{p}^3 \sum_{k=1}^{(p-1)/2} \frac{1 + 4p H_{2k-2}(1) - 2p H_{k-1}(1)}{(2k-1)^2} \pmod{p^4}. \quad (2.7)$$

This proves (2.4) since the right-hand side of the above congruence is independent of r .

The proof of Theorem 2.1 is now complete. \square

We are now in a position to prove Theorem 1.1. We need the following lemmas.

Lemma 2.2. *For any prime $p > 3$ we have*

$$\sum_{k=1}^{p-1} \frac{1}{k^3} \binom{2k}{k} \equiv -\frac{2H_{p-1}(1)}{p^2} \pmod{p}, \quad (2.8)$$

$$\sum_{k=1}^{p-1} \frac{H_k(2)}{k} \binom{2k}{k} \equiv \frac{2H_{p-1}(1)}{3p^2} \pmod{p}, \quad (2.9)$$

$$\sum_{k=1}^{p-1} \left(\frac{2}{k^2} - \frac{3H_k(1)}{k} \right) \binom{2k}{k} \equiv \frac{2H_{p-1}(1)}{p} \pmod{p^2}. \quad (2.10)$$

Proof. (2.8) was originally conjectured by Sun [15, Conjecture 1.1] and confirmed by Kh. Hessami Pilehrood and T. Hessami Pilehrood [6]. One may consult [15, Conjecture 1.1] for the modulus p^4 case of (2.8). We can directly verify (2.9) and (2.10) for $p = 5$. By [19] we know these two congruences hold for $p > 5$. \square

Lemma 2.3. *For any prime $p > 3$ we have*

$$\sum_{k=1}^{p-1} \left(\frac{3H_k(1)^2}{k} - \frac{4H_k(1)}{k^2} \right) \binom{2k}{k} \equiv \frac{6H_{p-1}(1)}{p^2} \pmod{p}.$$

Proof. As in [19], consider the WZ pair

$$F(n, k) = \frac{1}{k} \binom{n+k}{k} \quad \text{and} \quad G(n, k) = \frac{k}{(n+1)^2} \binom{n+k}{k}.$$

Then for any $n, k \in \mathbb{N}$ we have

$$F(n+1, k) - F(n, k) = G(n, k+1) - G(n, k).$$

Let $S_n = \sum_{k=1}^n F(n, k) H_k(1)^2$. Then

$$\begin{aligned} S_{n+1} - S_n &= \sum_{k=1}^{n+1} F(n+1, k) H_k(1)^2 - \sum_{k=1}^n F(n, k) H_k(1)^2 \\ &= F(n+1, n+1) H_{n+1}(1)^2 + \sum_{k=1}^n (F(n+1, k) - F(n, k)) H_k(1)^2 \\ &= F(n+1, n+1) H_{n+1}(1)^2 + \sum_{k=1}^n (G(n, k+1) - G(n, k)) H_k(1)^2 \\ &= F(n+1, n+1) H_{n+1}(1)^2 + \sum_{k=1}^n \left(G(n, k+1) H_k(1)^2 - G(n, k) H_{k-1}^2 \right. \\ &\quad \left. - \frac{2G(n, k) H_{k-1}}{k} - \frac{G(n, k)}{k^2} \right) \\ &= \frac{\binom{2n+2}{n+1} H_{n+1}(1)^2}{n+1} + \frac{\binom{2n+2}{n+1} H_n(1)^2}{2n+2} - 2 \sum_{k=1}^n \frac{G(n, k) H_k(1)}{k} + \sum_{k=1}^n \frac{G(n, k)}{k^2}. \end{aligned} \quad (2.11)$$

By [19, (16)] we have

$$\sum_{k=1}^n \frac{G(n, k)}{k^2} = \frac{1}{(n+1)^2} \sum_{k=1}^n \frac{1}{k} \binom{n+k}{k} = \frac{1}{(n+1)^2} \left(\frac{3}{2} \sum_{k=1}^n \frac{\binom{2k}{k}}{k} - H_n(1) \right). \quad (2.12)$$

From [1, (1.49)] we know that

$$\sum_{k=0}^n \binom{x+k}{k} = \binom{x+n+1}{n}.$$

Therefore,

$$\begin{aligned} \sum_{k=1}^n \frac{G(n, k) H_k(1)}{k} &= \frac{1}{(n+1)^2} \sum_{k=1}^n \binom{n+k}{k} \sum_{j=1}^k \frac{1}{j} = \frac{1}{(n+1)^2} \sum_{j=1}^n \frac{1}{j} \sum_{k=j}^n \binom{n+k}{k} \\ &= \frac{1}{(n+1)^2} \sum_{j=1}^n \frac{1}{j} \left(\binom{2n+1}{n} - \binom{n+j}{j-1} \right) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{(n+1)^2} \left(\frac{1}{2} \binom{2n+2}{n+1} H_n(1) - \frac{1}{n+1} \sum_{j=1}^n \binom{n+j}{j} \right) \\
&= \frac{1}{2(n+1)^2} \binom{2n+2}{n+1} H_n(1) - \frac{1}{2(n+1)^3} \binom{2n+2}{n+1} + \frac{1}{(n+1)^3}. \tag{2.13}
\end{aligned}$$

Substituting (2.12) and (2.13) into (2.11) we have

$$\begin{aligned}
S_{n+1} - S_n &= \frac{3 \binom{2n+2}{n+1} H_{n+1}(1)^2}{2n+2} - \frac{2 \binom{2n+2}{n+1} H_{n+1}(1)}{(n+1)^2} + \frac{5 \binom{2n+2}{n+1}}{2(n+1)^3} - \frac{H_{n+1}(1)}{(n+1)^2} \\
&\quad - \frac{1}{(n+1)^3} + \frac{3}{2(n+1)^2} \sum_{k=1}^n \frac{\binom{2k}{k}}{k}.
\end{aligned}$$

Now summing both sides over n from 0 to $p-2$ and noting that $S_0 = 0$ we have

$$\begin{aligned}
S_{p-1} &= \frac{3}{2} \sum_{n=1}^{p-1} \frac{\binom{2n}{n} H_n(1)^2}{n} - 2 \sum_{n=1}^{p-1} \frac{\binom{2n}{n} H_n(1)}{n^2} + \frac{5}{2} \sum_{n=1}^{p-1} \frac{\binom{2n}{n}}{n^3} - \sum_{k=1}^{p-1} \frac{H_n(1)}{n^2} - H_{p-1}(3) \\
&\quad + \frac{3}{2} \sum_{n=1}^{p-1} \frac{1}{n^2} \sum_{k=1}^{n-1} \frac{\binom{2k}{k}}{k}. \tag{2.14}
\end{aligned}$$

Clearly, for $1 \leq k \leq p-1$,

$$F(p-1, k) = \frac{1}{k} \binom{p-1+k}{k} \equiv \frac{1}{k} \binom{k-1}{k} = 0 \pmod{p}.$$

Thus we have

$$S_{p-1} = \sum_{k=1}^{p-1} F(p-1, k) H_k(1)^2 \equiv 0 \pmod{p}. \tag{2.15}$$

In view of (2.1) and (2.2) we have

$$\sum_{k=1}^{p-1} \frac{H_n(1)}{n^2} \equiv H_{p-1}(1, 2) \equiv -\frac{3H_{p-1}}{p^2} \pmod{p}. \tag{2.16}$$

Note that

$$\sum_{n=1}^{p-1} \frac{1}{n^2} \sum_{k=1}^{n-1} \frac{\binom{2k}{k}}{k} = \sum_{k=1}^{p-1} \frac{\binom{2k}{k}}{k} \sum_{n=k}^{p-1} \frac{1}{n^2} - \sum_{n=1}^{p-1} \frac{\binom{2n}{n}}{n^3} \equiv -\sum_{k=1}^{p-1} \frac{\binom{2k}{k} H_k(2)}{k} \pmod{p}. \tag{2.17}$$

Substituting (2.15)–(2.17) into (2.14) and using (2.8) and (2.9) we immediately obtain the desired result. \square

Lemma 2.4. *Let $p > 3$ be a prime. Then*

$$\sum_{k=1}^{(p-1)/2} \frac{1}{(2k-1)^2} \equiv \frac{1}{12} p B_{p-3} \pmod{p^2}, \tag{2.18}$$

$$\sum_{k=1}^{(p-1)/2} \frac{H_{2k-2}(1)}{(2k-1)^2} \equiv \frac{3}{8} B_{p-3} \pmod{p}, \quad (2.19)$$

$$\sum_{k=1}^{(p-1)/2} \frac{H_{k-1}}{(2k-1)^2} \equiv \frac{7}{8} B_{p-3} \pmod{p}. \quad (2.20)$$

Proof. By [14, Corollaries 5.1 and 5.2] we have

$$H_{p-1}(2) \equiv \frac{2}{3} p B_{p-3} \pmod{p^2} \quad \text{and} \quad H_{(p-1)/2}(2) \equiv \frac{7}{3} p B_{p-3} \pmod{p^2}.$$

It follows that

$$\sum_{k=1}^{(p-1)/2} \frac{1}{(2k-1)^2} = H_{p-1}(2) - \frac{1}{4} H_{(p-1)/2}(2) \equiv \frac{1}{12} p B_{p-3} \pmod{p^2}.$$

This proves (2.18).

Clearly,

$$\sum_{k=1}^{(p-1)/2} \frac{H_{2k-2}(1)}{(2k-1)^2} = \sum_{k=1}^{(p-1)/2} \frac{H_{2((p+1)/2-k)-2}(1)}{(2((p+1)/2-k)-1)^2} = \frac{1}{4} \sum_{k=1}^{(p-1)/2} \frac{H_{2k}}{k^2} \pmod{p},$$

where we used the fact $H_{p-1-k}(1) \equiv H_k(1) \pmod{p}$ for any $0 \leq k \leq p-1$. In view of [9, Lemma 2.4], we have

$$\sum_{k=1}^{(p-1)/2} \frac{H_{2k}(1)}{k^2} \equiv \frac{3}{2} B_{p-3} \pmod{p}. \quad (2.21)$$

This proves (2.19).

Note that

$$H_{(p-1)/2-k}(1) = H_{(p-1)/2}(1) - \sum_{j=1}^k \frac{1}{(p+1)/2-j} \equiv H_{(p-1)/2}(1) + 2H_{2k}(1) - H_k(1) \pmod{p}.$$

Therefore,

$$\begin{aligned} & \sum_{k=1}^{(p-1)/2} \frac{H_{k-1}(1)}{(2k-1)^2} = \sum_{k=1}^{(p-1)/2} \frac{H_{(p-1)/2-k}(1)}{(p-2k)^2} \\ & \equiv \frac{1}{4} \sum_{k=1}^{(p-1)/2} \frac{H_{(p-1)/2}(1) + 2H_{2k}(1) - H_k(1)}{k^2} \\ & \equiv \frac{1}{2} \sum_{k=1}^{(p-1)/2} \frac{H_{2k}(1)}{k^2} - \frac{1}{4} H_{(p-1)/2}(3) - \frac{1}{4} H_{(p-1)/2}(1, 2) \pmod{p}. \end{aligned}$$

It is known (cf. [7]) that

$$H_{(p-1)/2}(3) \equiv -2B_{p-3} \pmod{p} \quad \text{and} \quad H_{(p-1)/2}(1, 2) \equiv \frac{3}{2}B_{p-3} \pmod{p}.$$

Then (2.20) follows from the above two congruences and (2.21) immediately. \square

Proof of Theorem 1.1. By (2.5) and (2.7) we have

$$\begin{aligned} & \frac{1}{p^r} \sum_{k=0}^{p^r-1} (3k+1) \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} 4^k \\ \equiv & 1 + \frac{1}{4}p^2 \left(3 \sum_{n=1}^{p-1} \frac{\binom{2n}{n} H_n(1)}{n} - 2 \sum_{n=1}^{p-1} \frac{\binom{2n}{n}}{n^2} \right) + \frac{1}{8}p^3 \left(3 \sum_{n=1}^{p-1} \frac{\binom{2n}{n} H_n(1)^2}{n} - 4 \sum_{n=1}^{p-1} \frac{\binom{2n}{n} H_n(1)}{n^2} \right) \\ & + \frac{3}{4}p \sum_{n=1}^{p-1} \frac{\binom{2n}{n}}{n} + \frac{5}{16}p^3 \sum_{n=1}^{p-1} \frac{\binom{2n}{n}}{n^3} - \frac{3}{16}p^3 \sum_{n=1}^{p-1} \frac{\binom{2n}{n} H_n(2)}{n} - 2p^2 \sum_{k=1}^{(p-1)/2} \frac{1}{(2k-1)^2} \\ & - 8p^3 \sum_{k=1}^{(p-1)/2} \frac{H_{2k-2}(1)}{(2k-1)^2} + 4p^3 \sum_{k=1}^{(p-1)/2} \frac{H_{k-1}(1)}{(2k-1)^2} \pmod{p^4}. \end{aligned}$$

Then by (2.1), (2.6) and Lemmas 2.2–2.4 we have

$$\frac{1}{p^r} \sum_{n=0}^{p^r-1} (3n+1) \frac{\left(\frac{1}{2}\right)_n^3}{(1)_n^3} 4^n \equiv 1 + \frac{7}{6}p^3 B_{p-3} \pmod{p^4}.$$

The proof of Theorem 1.1 is now complete. \square

3. PROOF OF THEOREM 1.2

Similarly as in Section 2, we first establish the following result.

Theorem 3.1. *For any odd prime p and positive integer r we have*

$$\frac{1}{p^r} \sum_{k=0}^{(p^r-1)/2} (4k+1) \frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4} \equiv \frac{1}{p} \sum_{k=0}^{(p-1)/2} (4k+1) \frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4} \pmod{p^4}.$$

Lemma 3.1. *For any odd prime p and positive integer r we have*

$$p^r H_{p^r-1}(1) \equiv p H_{p-1}(1) \pmod{p^4}, \tag{3.1}$$

$$p^{2r} H_{p^r-1}(1, 1) \equiv p^2 H_{p-1}(1, 1) \pmod{p^4}, \tag{3.2}$$

$$p^{3r} H_{p^r-1}(1, 1, 1) \equiv p^3 H_{p-1}(1, 1, 1) \pmod{p^4}. \tag{3.3}$$

Proof. We first prove (3.1). Clearly, it holds for $r = 1$. Assume that it holds for $r = k > 1$. Then for $r = k + 1$ we have

$$\begin{aligned}
p^{k+1}H_{p^{k+1}-1}(1) &= p^{k+1} \sum_{\substack{j=1 \\ p \nmid j}}^{p^{k+1}-1} \frac{1}{j} + p^{k+1} \sum_{\substack{j=1 \\ p \nmid j}}^{p^{k+1}-1} \frac{1}{j} = p^k \sum_{j=1}^{p^k-1} \frac{1}{j} + p^{k+1} \sum_{l=0}^{p^k-1} \sum_{j=1}^{p-1} \frac{1}{lp+j} \\
&\equiv p^k \sum_{j=1}^{p^k-1} \frac{1}{j} + p^{k+1} \sum_{l=0}^{p^k-1} \sum_{j=1}^{p-1} \left(\frac{lp}{j^2} - \frac{1}{j} \right) \\
&= p^k H_{p^k-1}(1) + \frac{p^{2k+2}(p^k-1)}{2} H_{p-1}(2) - p^{2k+1} H_{p-1}(1) \\
&\equiv p^k H_{p^k-1}(1) \pmod{p^4}.
\end{aligned}$$

By the induction hypothesis we obtain (3.1).

(3.2) and (3.3) can be proved similarly by noting that

$$H_{p^r-1}(1, 1) = \frac{H_{p^r-1}(1)^2 - H_{p^r-1}(2)}{2}$$

and

$$H_{p^r-1}(1, 1, 1) = \frac{H_{p^r-1}(1)^3 - 3H_{p^r-1}(1)H_{p^r-1}(2) + 2H_{p^r-1}(3)}{6}.$$

□

Lemma 3.2. *For any odd prime p and positive integer r we have*

$$p^{3r} \sum_{k=0}^{(p^r-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} \equiv p^3 \sum_{k=0}^{(p-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} \pmod{p^4}.$$

Proof. Clearly, the congruence holds for $r = 1$.

Now suppose that $r \geq 2$. In view of the proof of [11, Theorem 1.1], for $0 < k < p^r/2$ we have

$$\frac{p^r}{k \binom{2k}{k}} \equiv 0 \pmod{p}. \quad (3.4)$$

Thus we obtain

$$\begin{aligned}
p^{3r} \sum_{k=0}^{(p^r-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} &= p^{3r} \sum_{\substack{k=0 \\ p \nmid 2k+1}}^{(p^r-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} + p^{3r} \sum_{\substack{k=0 \\ p \nmid 2k+1}}^{(p^r-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} \\
&\equiv p^{3r} \sum_{\substack{k=0 \\ p \nmid 2k+1}}^{(p^r-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} \pmod{p^4},
\end{aligned}$$

since by (3.4), $\text{ord}_p(p^{3r}/\binom{2k}{k}^2) \geq r+2 \geq 4$. It is routine to check that

$$p^{3r} \sum_{\substack{k=0 \\ p|2k+1}}^{(p^r-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} = p^{3r} \sum_{k=0}^{(p^{r-1}-3)/2} \frac{4^{(2k+1)p-1}}{(2k+1)^3 p^3 \binom{(2k+1)p-1}{((2k+1)p-1)/2}^2}.$$

Now

$$\binom{(2k+1)p-1}{((2k+1)p-1)/2} = \frac{\Gamma_p((2k+1)p)}{\Gamma_p(\frac{(2k+1)p+1}{2})^2} = -\frac{\Gamma_p((2k+1)p)}{\Gamma_p(\frac{(2k+1)p+1}{2})^2} \binom{2k}{k},$$

where $\Gamma_p(x)$ is the p -adic Gamma function (see [13, Chapter 7] for properties of this function). By (3.4), for $0 \leq k \leq (p^{r-1}-3)/2$ we have

$$\frac{p^{3r-3}}{(2k+1)^3 \binom{2k}{k}^2} = \frac{4p^{3r-3}}{(2k+1)(k+1)^2 \binom{2k+2}{k+1}^2} \equiv 0 \pmod{p^3}.$$

By Fermat's little theorem we have

$$4^{(2k+1)p-1} = 16^{kp} \cdot 4^{p-1} \equiv 16^k \pmod{p}.$$

Furthermore,

$$\frac{\Gamma_p((2k+1)p)^2}{\Gamma_p(\frac{(2k+1)p+1}{2})^4} \equiv \frac{\Gamma_p(0)^2}{\Gamma_p(\frac{1}{2})^4} = 1 \pmod{p}.$$

Combining the above we arrive at

$$p^{3r} \sum_{k=0}^{(p^r-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} \equiv p^{3r-3} \sum_{k=0}^{(p^{r-1}-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} \pmod{p^4}.$$

Then the desired result follows from induction on r . □

Proof of Theorem 3.1. We need the following pair which appeared in [22]

$$F(n, k) = (-1)^k (4n+1) \frac{\left(\frac{1}{2}\right)_n^3 \left(\frac{1}{2}\right)_{n+k}}{\left(1\right)_n^3 \left(1\right)_{n-k} \left(\frac{1}{2}\right)_k^2},$$

$$G(n, k) = (-1)^{k-1} \frac{4\left(\frac{1}{2}\right)_n^3 \left(\frac{1}{2}\right)_{n+k-1}}{\left(1\right)_{n-1}^3 \left(1\right)_{n-k} \left(\frac{1}{2}\right)_k^2}.$$

One may easily check that for any $n \in \mathbb{N}$ and $k \in \mathbb{Z}^+$

$$(2k-1)F(n, k-1) - 2kF(n, k) = G(n+1, k) - G(n, k).$$

Note that such pair is not a WZ pair. Nevertheless, it is also very useful as classical WZ pairs. For $m \in \mathbb{N}$, by induction on m and noting that

$$F(n, k) = 0 \quad \text{for } k > n$$

and

$$G(0, k) = 0 \quad \text{for } k > 0,$$

we have

$$\sum_{n=0}^m F(n, 0) = \frac{4^m}{\binom{2m}{m}} F(m, m) + \sum_{k=1}^m \frac{4^{k-1} G(m+1, k)}{(2k-1) \binom{2k-2}{k-1}}. \quad (3.5)$$

Taking $m = (p^r - 1)/2$ in (3.5) we have

$$\begin{aligned} \sum_{k=0}^{(p^r-1)/2} (2k+1) \frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4} &= \sum_{k=0}^{(p^r-1)/2} F(n, 0) \\ &= \frac{2^{p^r-1}}{\binom{p^r-1}{(p^r-1)/2}} F\left(\frac{p^r-1}{2}\right) + \sum_{k=1}^{(p^r-1)/2} \frac{4^{k-1}}{(2k-1) \binom{2k-2}{k-1}} G\left(\frac{p^r+1}{2}, k\right). \end{aligned}$$

It suffices to show

$$\frac{1}{p^r} \cdot \frac{2^{p^r-1}}{\binom{p^r-1}{(p^r-1)/2}} F\left(\frac{p^r-1}{2}\right) \equiv \frac{1}{p} \cdot \frac{2^{p-1}}{\binom{p-1}{(p-1)/2}} F\left(\frac{p-1}{2}\right) \pmod{p^4} \quad (3.6)$$

and

$$\frac{1}{p^r} \cdot \sum_{k=1}^{(p^r-1)/2} \frac{4^{k-1}}{(2k-1) \binom{2k-2}{k-1}} G\left(\frac{p^r+1}{2}, k\right) \equiv \frac{1}{p} \cdot \sum_{k=1}^{(p-1)/2} \frac{4^{k-1}}{(2k-1) \binom{2k-2}{k-1}} G\left(\frac{p+1}{2}, k\right) \pmod{p^4}. \quad (3.7)$$

We first consider (3.6). Note that

$$\frac{(1/2)_k}{(1)_k} = \frac{\binom{2k}{k}}{4^k}.$$

Thus we have

$$\begin{aligned} \frac{1}{p^r} \cdot \frac{2^{p^r-1}}{\binom{p^r-1}{(p^r-1)/2}} F\left(\frac{p^r-1}{2}\right) &= \frac{1}{p^r} \cdot \frac{(-1)^{(p^r-1)/2} 2^{p^r-1} (2p^r-1)}{\binom{p^r-1}{(p^r-1)/2}} \cdot \frac{\left(\frac{1}{2}\right)_{(p^r-1)/2}^3 \left(\frac{1}{2}\right)_{p^r-1}}{(1)_{(p^r-1)/2}^3 \left(\frac{1}{2}\right)_{(p^r-1)/2}^2} \\ &= \binom{\frac{1}{2}p^r-1}{p^r-1} \binom{2p^r-1}{p^r-1}. \end{aligned}$$

For any p -adic integer a , it is easy to see that

$$\begin{aligned} \binom{ap^r-1}{p^r-1} &= \frac{(1+(a-1)p^r)_{p^r-1}}{(1)_{p^r-1}} \\ &\equiv 1 + (a-1)p^r H_{p^r-1}(1) + (a-1)^2 p^{2r} H_{p^r-1}(1, 1) + (a-1)^3 p^{3r} H_{p^r-1}(1, 1, 1) \\ &\equiv 1 + (a-1)p H_{p-1}(1) + (a-1)^2 p^2 H_{p-1}(1, 1) + (a-1)^3 p^3 H_{p-1}(1, 1, 1) \\ &\equiv \binom{ap-1}{p-1} \pmod{p^4}, \end{aligned} \quad (3.8)$$

where we have used Lemma 3.1. Therefore,

$$\frac{1}{p^r} \cdot \frac{2^{p^r-1}}{\binom{p^r-1}{(p^r-1)/2}} F\left(\frac{p^r-1}{2}\right) \equiv \binom{\frac{1}{2}p-1}{p-1} \binom{2p-1}{p-1} \pmod{p^4}. \quad (3.9)$$

Then (3.6) follows by noting that the right-hand side of the above congruence is independent of r .

Below we consider (3.7). It is easy to see that

$$\begin{aligned} & \frac{1}{p^r} \cdot \sum_{k=1}^{(p^r-1)/2} \frac{4^{k-1}}{(2k-1)\binom{2k-2}{k-1}} G\left(\frac{p^r+1}{2}, k\right) \\ &= \frac{1}{p^r} \sum_{k=1}^{(p^r-1)/2} \frac{4^{k-1}}{(2k-1)\binom{2k-2}{k-1}} \cdot \frac{4(-1)^{k-1} \left(\frac{1}{2}\right)_{(p^r+1)/2}^3 \left(\frac{1}{2}\right)_{(p^r-1)/2+k}}{(1)_{(p^r-1)/2}^3 (1)_{(p^r+1)/2-k} \left(\frac{1}{2}\right)_k^2}. \end{aligned}$$

Note that

$$\left(\frac{1}{2}\right)_{(p^r-1)/2+k} = \left(\frac{1}{2}\right)_{(p^r-1)/2} \left(\frac{p^r}{2}\right)_k = \frac{p^r}{2} \left(\frac{1}{2}\right)_{(p^r-1)/2} \left(1 + \frac{p^r}{2}\right)_{k-1}$$

and

$$(1)_{(p^r+1)/2-k} = (-1)^{k-1} \frac{(1)_{(p^r-1)/2}}{\left(\frac{1}{2} - \frac{p^r}{2}\right)_{k-1}}.$$

Therefore,

$$\begin{aligned} & \frac{1}{p^r} \cdot \sum_{k=1}^{(p^r-1)/2} \frac{4^{k-1}}{(2k-1)\binom{2k-2}{k-1}} G\left(\frac{p^r+1}{2}, k\right) \\ &= \frac{p^{3r}}{16^{p^r-1}} \binom{p^r-1}{(p^r-1)/2}^4 \sum_{k=0}^{(p^r-3)/2} \frac{64^k \left(1 + \frac{p^r}{2}\right)_k \left(\frac{1}{2} - \frac{p^r}{2}\right)_k}{(2k+1)^3 \binom{2k}{k}^3 (1)_k^2}. \end{aligned}$$

By (3.4) we have

$$\frac{p^{3r}}{(2k+1)^3 \binom{2k}{k}^3} = \frac{8p^{3r}}{(k+1)^3 \binom{2k+2}{k+1}^3} \equiv 0 \pmod{p^3}.$$

Thus by Fermat's little theorem, (3.8) and Lemma 3.2 we further obtain

$$\begin{aligned} \frac{1}{p^r} \cdot \sum_{k=1}^{(p^r-1)/2} \frac{4^{k-1}}{(2k-1)\binom{2k-2}{k-1}} G\left(\frac{p^r+1}{2}, k\right) &\equiv p^{3r} \sum_{k=0}^{(p^r-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} \\ &\equiv p^3 \sum_{k=0}^{(p-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} \pmod{p^4}. \end{aligned} \quad (3.10)$$

This proves (3.7).

The proof of Theorem 3.1 is now complete. \square

Proof of Theorem 1.2. By (3.9) and (3.10) we arrive at

$$\frac{1}{p^r} \sum_{k=0}^{(p^r-1)/2} (4k+1) \frac{\left(\frac{1}{2}\right)_k^4}{(1)_k^4} \equiv \binom{\frac{1}{2}p-1}{p-1} \binom{2p-1}{p-1} + p^3 \sum_{k=0}^{(p-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} \pmod{p^4}. \quad (3.11)$$

By (3.8),

$$\binom{\frac{1}{2}p-1}{p-1} \equiv 1 - \frac{1}{2}pH_{p-1}(1) + \frac{1}{4}p^2H_{p-1}(1,1) - \frac{1}{8}p^3H_{p-1}(1,1,1) \pmod{p^4}$$

and

$$\binom{\frac{2}{p}-1}{p-1} \equiv 1 + pH_{p-1}(1) + p^2H_{p-1}(1,1) + p^3H_{p-1}(1,1,1) \pmod{p^4}.$$

It is known (cf. [7]) that

$$H_{p-1}(1,1) \equiv -\frac{1}{3}pB_{p-3} \pmod{p^2}$$

and

$$H_{p-1}(1,1,1) \equiv 0 \pmod{p}.$$

Then in view of (2.1) we get that

$$\binom{\frac{1}{2}p-1}{p-1} \binom{2p-1}{p-1} \equiv 1 - \frac{7}{12}p^3B_{p-3} \pmod{p^4}. \quad (3.12)$$

Sun [16, Theorem 1.2] proved that

$$\sum_{k=0}^{(p-3)/2} \frac{16^k}{(2k+1)^3 \binom{2k}{k}^2} \equiv \frac{7}{4}B_{p-3} \pmod{p}. \quad (3.13)$$

Substituting (3.12) and (3.13) into (3.11) we immediately obtain Theorem 1.2. \square

Now we can easily obtain Theorem 1.3.

Proof of Theorem 1.3. The case $p > 3$ is the immediate corollary of Theorems 1.1 and 1.2.

Now we consider the case $p = 3$. In view of Theorems 2.1 and 3.1, we only need to prove (1.3) for $r = 1$. In fact, if $p = 3$ and $r = 1$, one may check (1.3) directly. \square

Acknowledgments. The authors are grateful to Prof. Zhi-Wei Sun for his helpful suggestions on this paper. This work was supported by the National Natural Science Foundation of China (grant no. 11971222)

REFERENCES

- [1] H. W. Gould, *Combinatorial Identities*, Morgantown Printing and Binding Co., West Virginia, 1972.
- [2] J. Guillera and W. Zudilin, “Divergent” Ramanujan-type supercongruences, Proc. Amer. Math. Soc. **140** (2012), 765–777.
- [3] V.J.W. Guo, q -Supercongruences modulo the fourth power of a cyclotomic polynomial via creative microscoping, Adv. in Appl. Math. **120** (2020), Art. 102078.
- [4] V.J.W. Guo and M.J. Schlosser, Some q -supercongruences from transformation formulas for basic hypergeometric series, Constr. Approx. (2020), in press.

- [5] V.J.W. Guo and S.-D. Wang, *Some congruences involving fourth powers of central q -binomial coefficients*, Proc. Roy. Soc. Edinburgh Sect. A **150** (2020), 1127–1138.
- [6] Kh. Hessami Pilehrood and T. Hessami Pilehrood, *Congruences arising from Apéry-type series for zeta values*, Adv. in Appl. Math. **49** (2012), 218–238.
- [7] Kh. Hessami Pilehrood, T. Hessami Pilehrood and R. Tauraso, *New properties of multiple harmonic sums modulo p and p -analogues of Leshchiner’s series*, Tran. Amer. Math. Soc. **366** (2014), 3131–3159.
- [8] L. Long, *Hypergeometric evaluation identities and supercongruences*, Pacific J. Math. **249** (2011), 405–418.
- [9] G.-S. Mao and J. Wang, *On some congruences involving Domb numbers and harmonic numbers*, Int. J. Number Theory **15** (2019), 2179–2200.
- [10] G.-S. Mao and T. Zhang, *Proof of Sun’s conjectures on super congruences and the divisibility of certain binomial sums*, Ramanujan J. **50** (2019), 1–11.
- [11] H. Pan and Z.-W. Sun, *Proof of three conjectures on congruences*, Sci. China Math. **57** (2014), 2091–2102.
- [12] M. Petkovšek, H. S. Wilf and D. Zeilberger, *$A = B$* , A K Peters, Wellesley, 1996.
- [13] A. M. Robert, *A Course in p -Adic Analysis*, Graduate Texts in Mathematics, 198. Springer-Verlag, New York, 2000.
- [14] Z.-H. Sun, *Congruences concerning Bernoulli numbers and Bernoulli polynomials*, Discrete Appl. Math. **105** (2000), 193–223.
- [15] Z.-W. Sun, *Super congruences and Euler numbers*, Sci. China Math. **54** (2011), 2509–2535.
- [16] Z.-W. Sun, *p -Adic congruences motivated by series*, J. Number Theory **134** (2014), 181–196.
- [17] Z.-W. Sun, *Open conjectures on congruences*, Nanjing Univ. J. Math. Biquaterly **36** (2019), 1–99.
- [18] Z.-W. Sun and R. Tauraso, *New congruences for central binomial coefficients*, Adv. in Appl. Math. **45** (2010), 125–148.
- [19] R. Tauraso, *A bivariate generating function for zeta values and related supercongruences*, preprint, arXiv:1806.00846v4.
- [20] L. Van Hamme, *Some conjectures concerning partial sums of generalized hypergeometric series*, p -adic functional analysis (Nijmegen, 1996), Lecture Notes in Pure and Appl. Math. 192, Dekker, New York (1997), 223–236.
- [21] C. Wang, *Symbolic summation methods and hypergeometric supercongruences*, J. Math. Anal. Appl. **488** (2020), Art. 124068.
- [22] S.-D. Wang, *Some supercongruences involving $\binom{2k}{k}^4$* , J. Difference Equ. Appl. **24** (2018), 1375–1383.
- [23] W. Zudilin, *Ramanujan-type supercongruences*, J. Number Theory **129** (2009), 1848–1857.

(Chen Wang) DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE’S REPUBLIC OF CHINA

Email address: cwang@smail.nju.edu.cn

(Dian-Wang Hu) SCHOOL OF MATHEMATICS AND STATISTICS, NANYANG NORMAL UNIVERSITY, NANYANG 473061, HENAN, PEOPLE’S REPUBLIC OF CHINA

Email address: hudiwa@163.com