

Consumer Theory with Non-Parametric Taste Uncertainty and Individual Heterogeneity*

Christopher Dobronyi[†] and Christian Gouriéroux[‡]

September 3, 2022

*This paper is to be presented at the [14th International Conference on Computational and Financial Econometrics \(CFE 2020\)](#). The second author gratefully acknowledges financial support from the ACPR Chair (Regulation and Systemic Risks), and the ERC DYSMOIA. We also thank Joann Jasiak and Kyoo il Kim for many useful comments. This research was enabled by support from Sci-Net (www.scinethpc.ca) and Compute Canada (www.computecanada.ca). Our analyses are calculated using data from The Nielsen Company (US), and marketing databases provided by the Kilts Center for Marketing Data Center at The University of Chicago Booth School of Business. The conclusions in this paper do not reflect the views of The Nielsen Company or the ACPR. Neither are responsible for any of our results. Reference: [arXiv:2010.13937](#).

[†]Dobronyi: University of Toronto; christopher.dobronyi@mail.utoronto.ca.

[‡]Gouriéroux: University of Toronto, Toulouse School of Economics, and CREST.

Consumer Theory with Non-Parametric Taste Uncertainty and Individual Heterogeneity

Abstract

We introduce two models of non-parametric random utility for demand systems: the stochastic absolute risk aversion (SARA) model, and the stochastic safety-first (SSF) model. In each model, individual-level heterogeneity is characterized by a distribution $\pi \in \Pi$ of taste parameters, and heterogeneity across consumers is introduced using a distribution F over the distributions in Π . Demand is non-separable and heterogeneity is infinite-dimensional. Both models admit corner solutions. We consider two frameworks for estimation: a Bayesian framework in which F is known, and a hyperparametric framework in which F is a member of a parametric family. Our methods are illustrated by an application to a large panel of scanner data on alcohol consumption.

Keywords: Consumer Theory, Scanner Data, Stochastic Demand, Taste Heterogeneity, Non-Parametric Model, Bayesian Approach.

1 Introduction

The recent availability of databases containing all dated purchases made by a large number of consumers (28,036 in our application) presents a modern challenge for the econometrics of demand systems, requiring new models and estimation approaches (see, for example, [Burda et al., 2008, 2012](#), for discrete choice, and [Guha and Ng, 2019](#), [Chernozhukov et al., 2020](#), and [Dobronyi and Gouriéroux, 2020](#), for the first analyses of such data in the demand literature). This type of data is commonly called *scanner data* because the collection of such data involves retailers or households scanning the UPC code of each purchased good on the date of the purchase. This paper introduces two models of random utility for scanner data: the stochastic absolute risk aversion (SARA) model, and the stochastic safety-first (SSF) model. These models have the following advantages in comparison with the existing literature:

- (i) Both models are *consistent with consumer theory*: Every consumer maximizes a strictly increasing and strictly quasi-concave utility function. The latter property is not accommodated by existing approximations of the utility function like the quadratic approximation of the utility function ([Theil and Neudecker, 1958](#); [Barten, 1968](#)), the translog utility model ([Johansen, 1969](#); [Christensen et al., 1975](#)), or the Almost Ideal Demand System ([Deaton and Muellbauer, 1980](#)) and its extensions ([Banks et al., 1997](#); [Moschini, 1998](#)).
- (ii) Both models are non-parametric. In each model, the utility function is indexed by a functional parameter characterizing the individual heterogeneity, allowing for *infinite-dimensional heterogeneity*. In this respect, our paper differs from the existing literature when finite-dimensional heterogeneity is considered (see [Beckert and Blundell, 2008](#), [Blomquist et al., 2015](#), [Blundell, Horowitz, and Parey, 2017](#), and [Blundell, Kristensen, and Matzkin, 2017](#), for some examples of finite-dimensional restrictions). Our approach is in line with [Dette et al. \(2016\)](#) who write, “in general the multivariate demand function is a non-monotonic function of an infinite-dimensional unobservable—the individual’s preference ordering.”
- (iii) Both models yield demand functions with *non-separable heterogeneity* (see the discussions in [Brown and Walker, 1989](#), [Beckert and Blundell, 2008](#), and [Dette et al., 2016](#)). They are also endowed with precise *structural interpretations*, as heterogeneity is introduced by means of a distribution π of taste parameters, so that we can imagine consumers facing *taste uncertainty*, which they eliminate using expected utility.
- (iv) Both models are *identified* under weak restrictions. Identification follows from the use of panel data. Without such data, we lose identification ([Hausman and Newey, 2016](#)). Of course, the structure of scanner data is extremely important.

Each model is characterized by a basis of functions. This basis is used to generate a family of utility functions. A distribution is, then, placed over this family. To be

precise, we start with a basis of increasing and concave functions. Let $U(x; a)$ denote an element of this basis, where x is a bundle and $a \in \mathcal{A}$ is a finite-dimensional vector of taste parameters. A family of utility functions is generated by taking the convex hull of the basis. Let $U(x; \pi) = \mathbb{E}_\pi[U(x; a)]$ denote an element of this family, where $\pi \in \Pi$ is a distribution on \mathcal{A} . This family is indexed by a functional parameter π , which can be structurally interpreted as taste uncertainty. The heterogeneity across consumers is introduced using a distribution F on the set Π of probability distributions π on \mathcal{A} . Therefore, each model combines uncertainty and heterogeneity: the uncertainty in taste for a given consumer is represented by π and the heterogeneity across consumers is captured by F .

The paper considers a two-good framework and is organized as follows: Section 2 introduces the stochastic absolute risk aversion (SARA) model and Section 3 introduces the stochastic safety-first (SSF) model. For each model, we derive conditions on Π under which there exists a unique demand system, for each $\pi \in \Pi$. In Section 4, the distribution of heterogeneity F is introduced. If F is known, we obtain a Bayesian model in which the functional parameter $\pi \in \Pi$ has to be estimated. If F is a member of a known parametric family, indexed by θ , we obtain a model with a hyperparameter θ in which θ has to be estimated and π has to be filtered. In Section 5, we consider the identification of the taste distribution π *within* each model. Next, we examine if it is possible to distinguish *between* stochastic risk aversion and stochastic safety-first. In Section 6, we use Nielsen scanner data to illustrate our methodology in an application to the consumption of alcohol. Section 7 concludes. The details of the Dirichlet process are in Appendix A; integrability is discussed in Appendix B; an optimization procedure for filtering the taste distributions π after estimating F is in Appendix C; details of the Nielsen Homescan Consumer Panel (NHCP) are placed in Appendix D.

2 A Model with Stochastic Risk Aversion

This section introduces the first utility specification that we consider. It first describes the set of utility functions, then derives conditions under which there exists a unique demand system. The uncertainty is introduced by means of risk aversion parameters.

2.1 The Set of Utility Functions

There are two goods, denoted 1 and 2. Let $\bar{R} = \mathbb{R}_+^2$ denote the non-negative orthant with interior R . A consumer has preferences over the bundles in \bar{R} . Her preferences are summarized by a utility function of the form:

$$U(x; \pi) = -\mathbb{E}_\pi[\exp(-A'x)], \quad (2.1)$$

for every x such that $x_1, x_2 \geq 0$, where $A = (A_1, A_2)$ is a positive stochastic parameter characterizing the consumer's degrees of absolute risk aversion with respect to goods 1

and 2, and π is a joint distribution for this pair of stochastic taste parameters. Her preferences are, as a result, contained in a broad family of utility functions, indexed by a functional parameter π . There are two interpretations of specification (2.1): (i) the preferences are summarized by a deterministic utility function in the convex hull generated by a parametric family, or (ii) the consumer faces “taste uncertainty” and she resolves this uncertainty by using expected utility. We call these preferences *stochastic absolute risk aversion* (SARA) preferences.¹

If π is a point mass at $a = (a_1, a_2)$ such that $a_1, a_2 > 0$, the stochastic parameters are constant, and $U(x; \pi)$ reduces to $U(x; a) = -\exp(-a'x)$. This function is strictly increasing because we have:²

$$\frac{\partial U(x; a)}{\partial x} = \begin{bmatrix} a_1 \exp(-a'x) \\ a_2 \exp(-a'x) \end{bmatrix} > 0, \quad (2.2)$$

at each x such that $x_1, x_2 > 0$, and concave (although not necessarily strictly concave) because the Hessian associated with the utility function:

$$\frac{\partial^2 U(x; a)}{\partial x \partial x'} = -\exp(-a'x) \begin{pmatrix} a_1^2 & a_1 a_2 \\ a_1 a_2 & a_2^2 \end{pmatrix}, \quad (2.3)$$

is negative semi-definite, at each x such that $x_1, x_2 > 0$. This matrix is related to a bivariate measure of absolute risk aversion³ (Richard, 1975; Karni, 1979, 1983; Grant, 1995).

Proposition 1. If preferences are SARA and the consumer’s taste distribution π is not the mixture of point masses $a, a' \in R$ such that a is proportional to a' , then the utility function $U(x; \pi)$ is both strictly increasing and strictly concave given π on \bar{R} .

Proof. The utility function $U(x; \pi)$ is strictly increasing on \bar{R} because:

$$\frac{\partial}{\partial x} \mathbb{E}_\pi [U(x; A)] = \mathbb{E}_\pi \left[\frac{\partial U(x; A)}{\partial x} \right] > 0, \quad (2.4)$$

at every x such that $x_1, x_2 > 0$. It is strictly concave on \bar{R} because a function is strictly concave on \bar{R} , if its Hessian is negative definite on R , and the sum of two 2-by-2 matrices of rank 1, whose columns are not proportional, is a matrix with full rank. \square

¹These preferences differ from those used to describe consumer behaviour in the face of ambiguity or uncertainty, as in, say, Halevy and Feltkamp (2005).

²Here, > 0 means each component is strictly larger than 0.

³Such a measure can be defined as:

$$-\left(\text{diag} \frac{\partial U(x; \pi)}{\partial x} \right)^{-1/2} \frac{\partial^2 U(x; a)}{\partial x \partial x'} \left(\text{diag} \frac{\partial U(x; \pi)}{\partial x} \right)^{-1/2},$$

where $\text{diag} \frac{\partial U(x; \pi)}{\partial x}$ is the diagonal matrix whose diagonal elements are the first derivatives of $U(x; \pi)$.

Proposition 1 implies that we have effectively constructed a family of well-behaved utility functions $\{U(x; \pi) : \pi \in \Pi\}$ indexed by a functional parameter π , describing the taste uncertainty, instead of the standard finite-dimensional parameter usually considered in the literature.

Note that, after integrating out the taste uncertainty, the absolute risk aversions will depend on the consumption level. For instance, when A_1 and A_2 are independent with distributions π_1 and π_2 , then the risk aversion for good 1 becomes:

$$A_1(x_1) = -\frac{d^2 U_1(x_1; \pi_1)/dx_1^2}{dU_1(x_1; \pi_1)/dx_1} = \frac{\mathbb{E}_{\pi_1}[A_1^2 \exp(-A_1 x_1)]}{\mathbb{E}_{\pi_1}[A_1 \exp(-A_1 x_1)]}, \quad (2.5)$$

where $U_1(x_1; \pi_1)$ denotes $\mathbb{E}_{\pi_1}[\exp(-A_1 x_1)]$, the portion of the utility function $U(x; \pi)$ corresponding to good 1. Clearly, $A_1(x_1)$ depends on x_1 . Indeed, it is the average of A_1 given the following modified density:

$$\frac{A_1 \exp(-A_1 x_1)}{\mathbb{E}_{\pi_1}[A_1 \exp(-A_1 x_1)]}, \quad (2.6)$$

with respect to π_1 .

Let $g_\pi(\cdot)$ denote the function defined by the implicit equation:

$$U(x_1, g_\pi(x_1, u); \pi) = u, \quad (2.7)$$

for every $x_1 \geq 0$, and each (attainable) level of utility $u < 0$. This implicit equation has a unique solution because $U(x; \pi)$ is strictly increasing on \bar{R} . The function $g_\pi(\cdot, u)$ is the *indifference curve* associated with the functional parameter π and a utility level of u — $g_\pi(\cdot, u)$ maps every value of x_1 to a value of x_2 for which (x_1, x_2) attains a utility level of u given π . The implicit function theorem implies that $g_\pi(\cdot)$ is twice-continuously-differentiable with respect to x_1 and:

$$\frac{\partial g_\pi(x_1, u)}{\partial x_1} = -\text{MRS}(x_1, g_\pi(x_1, u); \pi), \quad (2.8)$$

on R where $\text{MRS}(x; \pi) \equiv \frac{\partial U(x; \pi)/\partial x_1}{\partial U(x; \pi)/\partial x_2}$ denotes the *marginal rate of substitution* at x —the rate at which the consumer is willing to exchange good 1 for good 2 given x and π .

2.2 The Demand Function

Let $z \in R$ denote a pair $z = (y, p)$ in which y denotes expenditure and p denotes the price of good 1, both normalized by the price of good 2. The consumer can purchase a bundle $x \in \bar{R}$ if, and only if, $px_1 + x_2 \leq y$. She chooses a bundle $x \in \bar{R}$ that solves:

$$\max_{x \in \bar{R}} U(x; \pi) \text{ subject to } px_1 + x_2 \leq y. \quad (2.9)$$

Let $X^*(z; \pi)$ denote the solution to:

$$\max_{x \in \mathbb{R}^2} -\mathbb{E}_\pi[\exp(-A'x)] \quad \text{subject to} \quad px_1 + x_2 \leq y. \quad (2.10)$$

While (2.9) is restricted to bundles in the non-negative orthant, (2.10) allows for negative values. The solution to (2.10) is characterized by a system of first-order conditions:

$$\text{MRS}(x; \pi) \equiv \frac{\mathbb{E}_\pi[A_1 \exp(-A'x)]}{\mathbb{E}_\pi[A_2 \exp(-A'x)]} = p \quad \text{and} \quad px_1 + x_2 - y = 0. \quad (2.11)$$

The first equality says that the marginal rate of substitution equals the relative price p . The second equality says that the budget constraint holds with equality. Equivalently, we can solve the following equality:

$$\mathbb{E}_\pi[(A_1 - pA_2) \exp(-(A_1 - pA_2)x_1) \exp(-A_2y)] = 0, \quad (2.12)$$

for the first component $X_1^*(z; \pi)$, and then use the budget constraint in (2.11) to solve for $X_2^*(z; \pi)$. As long as $A_1 - pA_2$ is not almost surely equal to zero, the first-order partial derivative of the left side of this equality with respect to x_1 is strictly negative:

$$-\mathbb{E}_\pi[(A_1 - pA_2)^2 \exp(-(A_1 - pA_2)x_1) \exp(-A_2y)] < 0. \quad (2.13)$$

The function on the left side of (2.12) is, therefore, strictly decreasing in x_1 , implying that there exists a unique solution $X_1^*(z; \pi)$ to (2.12), and a unique solution $X^*(z; \pi)$ to (2.10). If $X^*(z; \pi)$ is in \bar{R} , then $X^*(z; \pi)$ coincides with the solution to (2.9). Else, the solution to (2.9) is on the boundary of \bar{R} . Let $X(z; \pi)$ denote the solution to (2.9) given z and π . There are three *regimes* of demand in the design space:

$$X(z; \pi) = \begin{cases} (0, y)', & \text{if } X_1^*(z; \pi) \leq 0, \\ X^*(z; \pi), & \text{if } 0 \leq X_1^*(z; \pi) \leq y/p, \\ (y/p, 0)', & \text{if } y/p \leq X_1^*(z; \pi). \end{cases} \quad (2.14)$$

Therefore, we obtain:

Proposition 2. If preferences are SARA and the consumer's taste distribution π is not the mixture of point masses $a, a' \in R$ such that a is proportional to a' , then there exists a unique solution $X(z; \pi)$ to the maximization problem in (2.9) given z and π , for every $z \in R$, almost surely, for every π . There are three regimes defined by (2.14).

In general, the demand function $X(\cdot; \pi)$ is not necessarily invertible. For demand to be invertible (in the second regime), we require the indifference curves of the utility function to be strictly convex, a restriction that is stronger than the single-valuedness of demand $X(\cdot; \pi)$ (see Proposition 2 in Dobronyi and Gouriéroux, 2020).

2.3 Gamma Taste Uncertainty

As an illustration, let us assume that A_1 and A_2 are independent and that A_j has a Gamma distribution $\gamma(\nu_j, \alpha_j)$ with *degree of freedom* $\nu_j > 0$ and *scale factor* $\alpha_j > 0$, for $j = 1, 2$. Under this specification, $\pi = \gamma(\nu_1, \alpha_1) \otimes \gamma(\nu_2, \alpha_2)$, where \otimes denotes the tensor product of distributions. By the Laplace transform of the Gamma distribution:

$$U(x; \pi) = - \left(\frac{\alpha_1}{\alpha_1 + x_1} \right)^{\nu_1} \left(\frac{\alpha_2}{\alpha_2 + x_2} \right)^{\nu_2}. \quad (2.15)$$

Under this specification, the absolute risk aversion for good 1 in (2.5) becomes:

$$A_1(x_1) = \frac{\nu_1}{\alpha_1 + x_1}, \quad (2.16)$$

which is hyperbolic in x_1 . The indifference curve $g_\pi(\cdot)$ associated with utility level u is:

$$x_2 = g_\pi(x_1, u) \equiv \alpha_2 \left\{ \left[-\frac{1}{u} \left(\frac{\alpha_1}{\alpha_1 + x_1} \right)^{\nu_1} \right]^{\frac{1}{\nu_2}} - 1 \right\}, \quad (2.17)$$

for every $x_1 \geq 0$ and $u \in (-1, 0)$ such that:

$$x_1 < \alpha_1 \left[\left(-\frac{1}{u} \right)^{\frac{1}{\nu_1}} - 1 \right]. \quad (2.18)$$

It is easily shown that the second derivative of the indifference curve $g_\pi(\cdot, u)$ equals:

$$\frac{d^2 g_\pi(x_1, u)}{dx_1^2} = c \left(\frac{\alpha_1}{\alpha_1 + x_1} \right)^{\frac{\nu_1}{\nu_2} + 2} > 0, \quad (2.19)$$

for some $c > 0$. Therefore, the indifference curve $g_\pi(\cdot, u)$ is strictly convex. Furthermore, the MRS is equal to:

$$\text{MRS}(x; \pi) = \frac{\nu_1}{\nu_2} \frac{\alpha_2 + x_2}{\alpha_1 + x_1}. \quad (2.20)$$

The unconstrained solution $X_1^*(z; \pi)$ to the first-order condition in (2.11) is equal to:

$$X_1^*(z; \pi) = \frac{\nu_1}{\nu_1 + \nu_2} \cdot \frac{y}{p} + \frac{\nu_1 \alpha_2}{\nu_1 + \nu_2} \cdot \frac{1}{p} - \frac{\nu_2 \alpha_1}{\nu_1 + \nu_2}. \quad (2.21)$$

The second component $X_2^*(z; \pi)$ is deduced from the budget constraint in (2.11). By equation (2.14), the demand function $X(z; \pi)$ coincides with $X^*(z; \pi)$ over the set \mathcal{Z} of pairs z such that:

$$\min \{ \nu_1 y - p \nu_2 \alpha_1 + \nu_1 \alpha_2, \nu_2 y + p \nu_2 \alpha_1 - \nu_1 \alpha_2 \} > 0. \quad (2.22)$$

The three regimes of demand are illustrated in Figure 1 in the design space. The strict convexity of the indifference curve $g_\pi(\cdot, u)$ on \mathcal{Z} implies that the demand function $X(\cdot; \pi)$ associated with this utility function is invertible on \mathcal{Z} .




Figure 1. Regimes for Gamma Taste Uncertainty. The red region contains all designs z for which $X_1(z; \pi) > 0$ and $X_2(z; \pi) = 0$; the blue region contains all designs z for which $X_1(z; \pi) = 0$ and $X_2(z; \pi) > 0$; the green region contains all designs z for which $X_1(z; \pi) > 0$ and $X_2(z; \pi) > 0$.

3 A Model with Stochastic Safety-First

We now consider a model with taste parameters that have a *safety-first* interpretation.

3.1 The Set of Utility Functions

In Section 2, we constructed a family of well-behaved utility functions by taking the convex hull generated by a particular basis. In this section, we consider another basis, consisting of functions with the form:

$$U(x; a) = (x_1 + a_1 x_2) - (x_1 + a_1 x_2 - a_2)^+ = \min \{x_1 + a_1 x_2, a_2\}, \quad (3.1)$$

for every $x_1, x_2 \geq 0$, where $x^+ = \max\{0, x\}$ and $a_1, a_2 > 0$. This function corresponds to the “safety-first” criterion, introduced into the literature on portfolio management by Roy (1952). In order to illustrate, let us consider the consumption of alcohol, as in Dobronyi and Gouriéroux (2020). Suppose that there are two groups of goods: group 1 consisting of drinks with low alcohol by volume such as beers and ciders, and group 2 consisting of drinks with high alcohol by volume such as wines and liquors. Assume that the quantities are measured in identical units such as volume of alcohol—that is, the total volume of the drink in litres multiplied by the alcohol by volume of the drink.⁴ We can, then, add these volumes to aggregate two drinks with different sizes and/or percentages of alcohol. Here, a_1 is the consumer’s relative preference between the two groups of drinks, and a_2 is a “control” parameter, specifying her attempt to limit her intake of alcohol.

Now, let us introduce a distribution π such that $\mathbb{E}_\pi[A_j] < \infty$, $j = 1, 2$, and define:

$$U(x; \pi) = \mathbb{E}_\pi[(x_1 + A_1 x_2) - (x_1 + A_1 x_2 - A_2)^+]. \quad (3.2)$$

⁴Quantities could be, alternatively, measured in calories.

By the law of iterated expectations, we obtain:

$$U(x; \pi) = x_1 + \mathbb{E}_\pi[A_1]x_2 - \mathbb{E}_\pi\mathbb{E}_\pi[(x_1 + A_1x_2 - A_2)^+ | A_1]. \quad (3.3)$$

We call these preferences *stochastic safety-first* (SSF) preferences.

Under mild regularity conditions:

$$\frac{\partial U(x; \pi)}{\partial x_1} = 1 - \mathbb{E}_\pi\mathbb{E}_\pi[\mathbb{1}\{x_1 + A_1x_2 - A_2 > 0\} | A_1], \quad (3.4)$$

$$= \mathbb{E}_\pi[\mathbb{1}\{x_1 + A_1x_2 - A_2 < 0\}], \quad (3.5)$$

$$\frac{\partial U(x; \pi)}{\partial x_2} = \mathbb{E}_\pi[A_1] - \mathbb{E}_\pi[A_1\mathbb{E}_\pi[\mathbb{1}\{x_1 + A_1x_2 - A_2 > 0\} | A_1]], \quad (3.6)$$

$$= \mathbb{E}_\pi[A_1\mathbb{1}\{x_1 + A_1x_2 - A_2 < 0\}], \quad (3.7)$$

for every x such that $x_1, x_2 > 0$. These partial derivatives are strictly positive when π has full support: $\pi(a_1, a_2) > 0$, for $a_1, a_2 > 0$. By taking the second-order derivatives:

$$\frac{\partial^2 U(x; \pi)}{\partial x \partial x'} = - \begin{pmatrix} \mathbb{E}_\pi[\pi_0] & \mathbb{E}_\pi[A_1\pi_0] \\ \mathbb{E}_\pi[A_1\pi_0] & \mathbb{E}_\pi[A_1^2\pi_0] \end{pmatrix}, \quad (3.8)$$

for every x such that $x_1, x_2 > 0$, where $\pi_0 \equiv \pi(x_1 + A_1x_2 | A_1)$ in which $\pi(\cdot | A_1)$ denotes the conditional density of A_2 given A_1 , assuming that such a density exists. This matrix is both symmetric and negative definite when $\pi(\cdot | A_1)$ is continuous and A_1 is not constant. This result follows from the positivity of $\mathbb{E}_\pi[\pi_0]$ and the following equality:

$$\det \frac{\partial^2 U(x; \pi)}{\partial x \partial x'} = \mathbb{E}_\pi[\pi_0]V_{\tilde{\pi}}(A_1) > 0, \quad (3.9)$$

which holds for every x such that $x_1, x_2 > 0$, in which $\tilde{\pi}$ denotes the modified density:

$$\tilde{\pi}(a) = \frac{\pi(x_1 + a_2x_2 | a_1)\pi(a)}{\mathbb{E}_\pi[\pi_0]}. \quad (3.10)$$

Proposition 3. If preferences are SSF and the consumer's taste distribution π is continuous with full support given A_1 , then the utility function $U(x; \pi)$ is strictly increasing and strictly concave given π on \bar{R} .

Consequently, we have constructed another family of well-behaved utility functions $\{U(x; \pi) : \pi \in \Pi\}$ indexed by a functional parameter π , describing taste uncertainty.

3.2 The Demand Function

Let us revisit the utility maximization problem in (2.9). Under the safety-first specification, the analogue of the unconstrained first-order condition in (2.12) is given by:

$$\mathbb{E}_\pi[(1 - pA_1)\mathbb{1}\{x_1(1 - pA_1) + A_1y - A_2 < 0\}] = 0. \quad (3.11)$$

We obtain this equality by equating the marginal rate of substitution with the relative price p , and then using the budget constraint to replace x_2 with $y - px_1$. Under the regularity conditions from above, the left-hand side is strictly monotone in x_1 given π , so that there exists a unique solution to the first-order condition. As in Section 2, we let $X_1^*(z; \pi)$ denote this solution, and let $X_2^*(z; \pi)$ denote the quantity $y - pX_1^*(z; \pi)$.

Proposition 4. If preferences are SSF and the consumer's taste distribution π is continuous with full support given A_1 , then there exists a unique solution $X(z; \pi)$ to the maximization problem in (2.9) given z and π , for every $z \in R$, almost surely, for every π . There are three regimes defined by (2.14).

When the consumer's preferences are SSF, the MRS has the form:

$$\text{MRS}(z; \pi) \equiv \frac{\mathbb{E}_\pi[\mathbf{1}\{x_1 + A_1x_2 - A_2 < 0\}]}{\mathbb{E}_\pi[A_1\mathbf{1}\{x_1 + A_1x_2 - A_2 < 0\}]} = \frac{1}{\mathbb{E}_\pi[A_1|x_1 + A_1x_2 - A_2 < 0]}. \quad (3.12)$$

Thus, the rate at which the consumer is willing to exchange good 1 for good 2 given x and π is equal to the inverse of the expectation of her relative preference between goods A_1 , conditional on not surpassing her control parameter A_2 .

Some functionals of the distribution π can be especially interesting. For instance, in an application to the consumption of alcohol, we might expect the conditional distribution of A_2 given $A_1 = a_1$ to be concentrated around a single mode, characterizing an implicit alcohol limit for this consumer. Then, we can ask the following questions:

- (i) Is this limit positively correlated with A_1 ? In other words, is there a positive relationship between this limit and a preference for *strong* alcoholic beverages?
- (ii) Does a change in the maximum blood alcohol level for driving affect this limit?

These are questions that cannot be answered using classical demand systems like the Almost Ideal Demand System (Deaton and Muellbauer, 1980). In fact, tests based on the Almost Ideal Demand System have rejected rationality in applications to alcohol consumption (Alley et al., 1992). Clearly, it is possible that the Almost Ideal Demand System is misspecified.

3.3 Exponential Threshold Taste Uncertainty

In general, the first-order condition in (3.11) has no closed-form solution. However, its expression can be simplified for some taste distributions π . As an illustration, let us assume that:

- (i) A_1 and A_2 are independent.
- (ii) A_2 follows an exponential distribution $\gamma(1, \lambda)$ with survival function:

$$P(A_2 > a_2) = \exp(-\lambda a_2). \quad (3.13)$$

(iii) A_1 follows a distribution with Laplace transform: $\Psi(v) = \mathbb{E}[\exp(-vA_1)]$, $v \geq 0$.

Under this specification, we can first integrate with respect to A_2 within the expectation in (3.11) in order to obtain the following condition:

$$\mathbb{E}_\pi[(1 - pA_1) \exp\{-\lambda(x_1 + (y - x_1p)A_1)\}] = 0. \quad (3.14)$$

Equivalently, we obtain:

$$\mathbb{E}_\pi[\exp\{-\lambda(y - x_1p)A_1\}] - p\mathbb{E}_\pi[A_1 \exp\{-\lambda(y - x_1p)A_1\}] = 0. \quad (3.15)$$

This equation can be written in terms of the Laplace transform Ψ for A_1 . This yields:

$$\Psi[\lambda(y - x_1p)] + p \frac{d\Psi}{dv}[\lambda(y - x_1p)] = 0, \quad (3.16)$$

which can also be written as:

$$\frac{d \log \Psi}{dv}[\lambda(y - x_1p)] = -\frac{1}{p}. \quad (3.17)$$

Finally, by inverting this expression and rearranging the terms, we get:

$$X_1^*(z; \pi) = \frac{1}{p} \left[y - \frac{1}{\lambda} \left(\frac{d \log \Psi}{dv} \right)^{-1} \left(-\frac{1}{p} \right) \right], \quad (3.18)$$

The second component $X_2^*(z; \pi)$ of the unconstrained solution in (3.11) is deduced from the budget constraint. It follows from equation (2.14) that the demand function $X(z; \pi)$ coincides with $X^*(z; \pi)$ if, and only if:

$$0 \leq \frac{1}{\lambda} \left(\frac{d \log \Psi}{dv} \right)^{-1} \left(-\frac{1}{p} \right) \leq y. \quad (3.19)$$

For instance, if A_1 follows a gamma distribution $\gamma(\nu, \alpha)$, then $\log \Psi(v) = -\nu \log(1 + v/\alpha)$, and we obtain:

$$\frac{d \log \Psi(v)}{dv} = -\frac{\nu}{\alpha + v}, \quad (3.20)$$

for $v \geq 0$. Moreover, by inverting this function, we get:

$$\left(\frac{d \log \Psi}{dv} \right)^{-1}(\xi) = -\left(\frac{\nu}{\xi} + \alpha \right). \quad (3.21)$$

Therefore, the solution $X_1^*(z; \pi)$ has the form:

$$X_1^*(z; \pi) = \frac{1}{p} \left[y + \frac{1}{\lambda}(\alpha - \nu p) \right], \quad (3.22)$$




Figure 2. Regimes for Exponential Threshold Taste Uncertainty.

The red region contains all designs z for which $X_1(z; \pi) > 0$ and $X_2(z; \pi) = 0$; the blue region contains all designs z for which $X_1(z; \pi) = 0$ and $X_2(z; \pi) > 0$; the green region contains all designs z for which $X_1(z; \pi) > 0$ and $X_2(z; \pi) > 0$.

and demand $X(z; \pi)$ coincides with $X^*(z; \pi)$ if, and only if:

$$0 \leq \frac{\nu p - \alpha}{\lambda} \leq y. \quad (3.23)$$

The regimes of demand are illustrated in Figure 2 in the design space. Note, we can also verify that the *Slutsky coefficient* is strictly negative⁵ such that:

$$\Delta_x(z) \equiv \frac{\partial X_1(z; \pi)}{\partial p} + X_1(z; \pi) \frac{\partial X_1(z; \pi)}{\partial y} = -\frac{\nu}{\lambda p} < 0, \quad (3.24)$$

ensuring that the demand function $X(\cdot; \pi)$ is invertible over the set \mathcal{Z} of pairs z on which demand is strictly positive (see Section 2 in Dobronyi and Gouriéroux, 2020).

4 Individual Heterogeneity

Sections 2 and 3 introduced two utility specifications, both indexed by the functional parameter π . Of course, different consumers can have different functional parameters. This individual heterogeneity is introduced in a second layer, by specifying a distribution F over the set Π of distributions on R , such as the Dirichlet process (see, for example, Navarro et al., 2006, for an application of the Dirichlet process in modelling individual differences). More precisely, we make the following theoretical assumption:

Assumption A1 (Latent Stochastic Model).

- (i) There are $n \geq 1$ consumers.

⁵This property holds for any Laplace transform Ψ of A_1 (see Appendix B).

- (ii) Consumers are segmented into M homogeneous groups.
- (iii) Consumers in group m have the utility function $U(x; \pi_m)$, for all $m = 1, \dots, M$.
- (iv) The taste parameters (π_m) are independently drawn from a Dirichlet process F .

Assumption A1 introduces a distribution F over the functional taste parameter π . This distribution F characterizes the heterogeneity across homogeneous groups. It can encompass, for example, regional or demographic differences in preferences. This infinite-dimensional heterogeneity is non-separable in the stochastic demand equation.

The Dirichlet process can be constructed in three steps:

Step 1: Consider the set of (Bernoulli) distributions on $\{0, 1\}$. This set is characterized by $q \in \bar{R}$ such that $q_1 + q_2 = 1$. A distribution defined on this set of distributions is a distribution defined on this parameter set. We can, for instance, introduce a *beta distribution*, denoted $B(\alpha_1, \alpha_2)$. The distribution $B(\alpha_1, \alpha_2)$ has a continuous density:

$$f(q) = \frac{\Gamma(\alpha_1 + \alpha_2) q_1^{\alpha_1} q_2^{\alpha_2}}{\Gamma(\alpha_1) \Gamma(\alpha_2)}, \quad (4.1)$$

with respect to the Lebesgue measure over the simplex $\{(q_1, q_2) \geq 0 : q_1 + q_2 = 1\}$, where Γ denotes the gamma function,⁶ and $\alpha_1, \alpha_2 > 0$ are positive scalar parameters.

Step 2: The beta distribution can be extended to define a distribution on the set of discrete distributions with weights $q_j \geq 0$, $j = 1, \dots, J$, such that $\sum_{j=1}^J q_j = 1$. This procedure leads to the *Dirichlet distribution*, denoted $D(\alpha)$. The resulting distribution $D(\alpha)$ has continuous density:

$$f(q) = \frac{\Gamma(\sum_{j=1}^J \alpha_j) \prod_{j=1}^J q_j^{\alpha_j}}{\prod_{j=1}^J \Gamma(\alpha_j)}, \quad (4.2)$$

with respect to the Lebesgue measure over the simplex:

$$\left\{ q \in \mathbb{R}_+^J : \sum_{j=1}^J q_j = 1 \text{ and } q_j \geq 0, \forall j \right\}, \quad (4.3)$$

(see, for example, Kotz et al., 2000, page 485, and Lin, 2016, for details).

Step 3: Then, the Dirichlet distribution can be extended to define a distribution on a large set of distributions⁷ defined on \bar{R} (see Appendix A). This procedure

⁶The gamma function Γ is defined by $\Gamma(\alpha) = \int_0^\infty \exp(-x) x^{\alpha-1} dx$, for each $\alpha > 0$.

⁷The realizations of a Dirichlet process are, almost surely, discrete distributions. Although we assumed continuity to prove the existence of a unique demand system in Section 3, these realizations can approximate any continuous distribution. This discrepancy has no practical implications.

leads to the *Dirichlet process*. The Dirichlet process is characterized by a distribution μ on \bar{R} and a scaling parameter $c > 0$. The distribution μ can be thought of as the mean of the Dirichlet process, while the parameter c manages its degree of discretization (see Appendix A). This extension of the Dirichlet distribution is much more complicated than the Dirichlet distribution, especially because the notion of the Lebesgue measure on the set of distributions, and the notion of a density, no longer exist (see Ferguson, 1974, Rolin, 1992, and Sethuraman, 1994).

Let us now discuss implications of Assumption A1: If the functional and scaling parameters of the Dirichlet process are known, then we are in a Bayesian framework (see, for example, Geweke, 2012, for a Bayesian analysis of revealed preference) in which the taste distribution $\pi \in \Pi$ has to be estimated. Otherwise, we can assume that the mean μ of our process F is characterized by a finite-dimensional hyperparameter θ . Naturally, the hyperparametric model has two types of parameters: the hyperparameter θ to be estimated, and the functional parameters (π_m) to be filtered.

5 Non-Parametric Identification

In this section, we consider the identification of the functional parameter π *within* each model from the observation of a demand function. Then, we examine if we can distinguish *between* the SARA and SSF models.

Intuitively, a consumer's demand function is identified if we observe her making a lot of consumption decisions at a variety of designs z . Clearly, we can identify her demand function if (i) her preferences are constant over time and we observe a large panel or experiment,⁸ or (ii) she belongs to a large homogeneous segment of consumers with identical preferences. This explains the form of Assumption A1 (as it allows for either interpretation). Later, we apply the segmented approach to scanner data in the application to the consumption of alcohol in Section 6.

With panel data, one no longer requires the assumption that demand is monotonic with respect to unobserved heterogeneity, as commonly assumed in models with finite-dimensional heterogeneity (see Brown and Matzkin, 1995, as well as Matzkin, 2003).

5.1 Within Model Identification

In the models introduced in Sections 2 and 3, and for any π such that demand is invertible (as in Sections 2.3 and 3.3), we can derive the inverse demand function, whose second component coincides with the MRS which can be integrated to obtain a unique preference ordering. Indeed, by construction, the integrability conditions (needed to recover a *unique* well-behaved preference ordering) are satisfied for each individual, implying that preferences are recoverable (see Samuelson, 1948, for the seminal discus-

⁸In this case, when the number of dates T is large, we can have a segment m for each consumer i .

ssion of integrability in the case of two goods, as well as [Samuelson, 1950](#), [Hurwicz and Uzawa, 1971](#), and [Hosoya, 2016](#), for more general approaches). However, the possibility to recover preferences from a consumer's demand function does not imply that the distribution of taste uncertainty π is identified. Indeed, two distinct taste distributions could produce an identical MRS.

For identification, we only consider the information contained in the demand function $X(\cdot; \pi)$ on the set \mathcal{Z} of designs z for which the components of the demand function are strictly positive. This restriction disregards some information that may be available in the first or third regimes of (2.14). However, in most datasets, when a component of the demand function equals zero, the price p is not observed.

5.1.1 Stochastic Absolute Risk Aversion

In the stochastic absolute risk aversion (SARA) model, the identification condition is:

$$\left\{ \frac{\mathbb{E}_\pi[A_1 \exp(-A'x)]}{\mathbb{E}_\pi[A_2 \exp(-A'x)]} = \frac{\mathbb{E}_{\pi'}[A_1 \exp(-A'x)]}{\mathbb{E}_{\pi'}[A_2 \exp(-A'x)]}, \forall x \in R \right\} \Rightarrow \pi = \pi'. \quad (5.1)$$

In the degenerate case in which A is deterministic and equal to (a_1, a_2) , the MRS reduces to a_1/a_2 . Thus, in this special case, the two-dimensional parameter $a = (a_1, a_2)$ is identified up to a positive factor. This reasoning leads us to a question: Does this lack of identification also exist in an extended setting?

Let us first remark that the utility function $U(x; \pi)$ is equal to the moment generating function for π with a negative sign $\Phi(x; \pi) = -U(x; \pi)$. Because this moment generating function characterizes π when the stochastic parameter A is non-negative (see Theorem 1a in Chapter 13 on Tauberian Theorems in [Feller, 1968](#)), it is equivalent to consider the identification of either π , or $\Phi(x; \pi)$.⁹ As mentioned, we can always integrate the MRS to recover a unique preference ordering. In other words, we can recover $U(x; \pi)$ up to a monotonic transformation. We still need to discern the conditions on π under which we can recover $\Phi(x; \pi)$. Indeed, moment generating functions have properties that are not necessarily preserved under monotonic transformations.

We obtain the following result:

Proposition 5. If preferences are SARA, then $\Phi(x; \pi)$ and $\Phi(x; \pi)^\nu$ lead to the same preference ordering, for all positive scalars $\nu > 0$.

Proof. Let $U(x; \pi) = -\Phi(x; \pi)$ and $\tilde{U}(x; \pi) = -\Phi(x; \pi)^\nu$ denote the utility functions associated with $\Phi(x; \pi)$ and $\Phi(x; \pi)^\nu$, respectively. Then, by definition, we must have:

$$\tilde{U}(x; \pi) = -\Phi(x; \pi)^\nu = -(-U(x; \pi))^\nu = \phi_\nu(U(x; \pi)), \quad (5.2)$$

⁹Note, the existence of the moment generating function does not imply the existence of all power moments and, even if all power moments exist, they do not necessarily characterize the distribution. A known example is the log-normal distribution used in the application ([Heyde, 1963](#)).

where $\phi_\nu(u) = -(-u)^\nu$ is strictly increasing for $u < 0$. Since $\tilde{U}(x; \pi)$ is a monotonic transformation of $U(x; \pi)$, these utility functions yield the same preference ordering. \square

This means that we can, at most, identify the class of moment generating functions $\mathcal{C}(\Phi) = \{\Phi^\nu : \nu > 0\}$. Note that, for any moment generating function Φ , the transformed function Φ^ν is also a moment generating function.

Let us now consider identification when A_1 and A_2 are independent:

Proposition 6. Let Φ_j denote the marginal moment generating function for A_j , for $j = 1, 2$. If preferences are SARA, and A_1 and A_2 are independent, then (Φ_1, Φ_2) and (Φ_1^*, Φ_2^*) lead to the same preference ordering if, and only if, for some $\nu > 0$, we have:

$$\Phi_1^* = \Phi_1^\nu \text{ and } \Phi_2^* = \Phi_2^\nu.$$

Proof. The identification criterion becomes:

$$\left(\frac{\partial \Phi_1(x_1)}{\partial x_1} \Phi_2(x_2) \right) \left(\Phi_1(x_1) \frac{\partial \Phi_2(x_2)}{\partial x_2} \right)^{-1} = \left(\frac{\partial \Phi_1^*(x_1)}{\partial x_1} \Phi_2^*(x_2) \right) \left(\Phi_1^*(x_1) \frac{\partial \Phi_2^*(x_2)}{\partial x_2} \right)^{-1},$$

for all $x \in R$. This criterion can, then, be written as:

$$\frac{\partial \log \Phi_1(x_1)}{\partial x_1} \left(\frac{\partial \log \Phi_1^*(x_1)}{\partial x_1} \right)^{-1} = \frac{\partial \log \Phi_2(x_2)}{\partial x_2} \left(\frac{\partial \log \Phi_2^*(x_2)}{\partial x_2} \right)^{-1},$$

for all $x \in R$. Thus, we deduce that, if these distributions yield the same MRS, then:

$$\frac{\partial \log \Phi_j^*(x_j)}{\partial x_j} = \nu \frac{\partial \log \Phi_j(x_j)}{\partial x_j},$$

for some $\nu > 0$, at every $x_j \geq 0$, for both $j = 1, 2$. Because the log-transform of the moment generating function at zero equals zero, by integrating this equation, we get:

$$\log \Phi_j^*(x_j) = \nu \log \Phi_j(x_j), \quad (5.3)$$

at every $x_j \geq 0$, for both $j = 1, 2$. Equivalently, $\Phi_1^* = \Phi_1^\nu$ and $\Phi_2^* = \Phi_2^\nu$. \square

Proposition 6 implies that $\mathcal{C}(\Phi)$ is identified under the independence of A_1 and A_2 . Indeed, we can recover the consumer's preference ordering by using traditional methods, and use the fact that all admissible preference orderings map into a unique class $\mathcal{C}(\Phi)$.

5.1.2 Stochastic Safety-First

In the SSF model, the identification condition is:

$$\left\{ \frac{\mathbb{E}_\pi[A_1 | x_1 + A_1 x_2 - A_2 < 0]}{\mathbb{E}_{\pi'}[A_1 | x_1 + A_1 x_2 - A_2 < 0]} = 1, \forall x \in R \right\} \Rightarrow \pi = \pi'. \quad (5.4)$$

Let us now consider the validity of this condition under an independence assumption.

Proposition 7. If preferences are SSF, A_1 and A_2 are independent, and the marginal distribution of A_2 is continuous, then $\mathbb{E}[A_1]$ is identified, and the marginal distribution of A_2 is identified up to some positive power transformation of its survival function.

Proof. In the SSF model, the MRS is identified, and it satisfies:

$$\mathbb{E}_\pi[A_1 S(x_1 + A_1 x_2)] = \text{MRS}(x; \pi) \mathbb{E}_\pi[S(x_1 + A_1 x_2)], \quad (5.5)$$

where $S(\cdot)$ denotes the survival function of A_2 .

- (i) The expectation $\mathbb{E}_\pi[A_1]$ is identified because $\text{MRS}(x_1, 0; \pi) = \mathbb{E}_\pi[A_1]$.
- (ii) By differentiating (5.5) with respect to x_2 , we get:

$$\begin{aligned} \mathbb{E}_\pi[A_1^2 S'(x_1 + A_1 x_2)] &= \text{MRS}(x; \pi) \mathbb{E}_\pi[A_1 S'(x_1 + A_1 x_2)] \\ &\quad + \frac{\partial \text{MRS}}{\partial x_2}(x; \pi) \mathbb{E}_\pi[S(x_1 + A_1 x_2)]. \end{aligned}$$

When $x_2 = 0$, this equation becomes:

$$S'(x_1) \mathbb{E}_\pi[A_1^2] = \text{MRS}(x_1, 0; \pi) S'(x_1) \mathbb{E}_\pi[A_1] + \frac{\partial \text{MRS}}{\partial x_2}(x_1, 0; \pi) S(x_1).$$

By rearranging, we get:

$$\frac{\partial \text{MRS}}{\partial x_2}(x_1, 0; \pi) = \frac{S'(x_1)}{S(x_1)} (\mathbb{E}_\pi[A_1^2] - \text{MRS}(x_1, 0; \pi) \mathbb{E}_\pi[A_1]) = \frac{S'(x_1)}{S(x_1)} V(A_1).$$

Because the partial derivative of the MRS with respect to x_2 is identified, the hazard function $\lambda(x_1) = -S'(x_1)/S(x_1)$ of the distribution of A_2 is identified up to a positive factor. Since $S(x_1) = \exp\{-\Lambda(x_1)\}$, where $\Lambda(x_1) = \int_0^{x_1} \lambda(t) dt$ is the cumulative hazard function of the distribution of A_2 , we can identify $S(\cdot)$ up to a positive power transformation.

□

Proposition 7 provides no information on the identifiability of the distribution of A_1 beyond its first moment. It seems difficult to obtain a general identification result, but insights into our identification problem can be obtained by considering the two primary families of distributions that are invariant to positive power transformations, that are, the exponential family and the Pareto family.

- (i) **Exponential family:** Suppose that the marginal distribution of A_2 belongs to the exponential family, and that we have identified its survival function up to a positive power transformation such that $S(x) = \exp\{-cx\}$, for some unknown $c > 0$. The MRS in (5.5) becomes:

$$\text{MRS}(x; \pi) = \frac{\mathbb{E}_\pi[A_1 \exp\{-cx_2 A_1\}]}{\mathbb{E}_\pi[\exp\{-cx_2 A_1\}]} \equiv G_0(x_2). \quad (5.6)$$

This expression does not depend on x_1 . Now, let $\Psi(u) = \mathbb{E}_\pi[\exp\{-uA_1\}]$ denote the Laplace transform of A_1 . Under this notation, the equality in (5.6) implies:

$$G_0(x_2) = \frac{d \log \Psi}{du}(cx_2).$$

Or, equivalently, $G_0(u/c) = d \log \Psi(u)/du$. By integrating, we obtain:

$$\log \Psi(u) = c[H(u/c) - H(0)],$$

where $H(\cdot)$ is a primitive of the MRS. Therefore:

Corollary 1. Under the conditions of Proposition 7, if the marginal distribution of A_2 belongs to the exponential family, the following results hold:

- (a) The power transform c is not identified.
- (b) The distribution of A_1 is identified under an identification restriction on c .

We conclude that, under the conditions of Corollary 1, the distributions of A_1 and A_2 are non-parametrically identified up to a single scalar parameter $c > 0$.

- (ii) **Pareto family:** Let us now examine whether a similar result can be obtained for the Pareto family, in which $S(x) = x^{-\alpha}$, for some $\alpha > 0$. The parameter α characterizes the fat tails of the distribution of A_2 and the power transformation on the MRS. This survival function produces:

$$\text{MRS}(x; \pi) = \frac{\mathbb{E}_\pi[A_1(x_1 + A_1 x_2)^{-\alpha}]}{\mathbb{E}_\pi[(x_1 + A_1 x_2)^{-\alpha}]} = \frac{\mathbb{E}_\pi[A_1(x_0 + A_1)^{-\alpha}]}{\mathbb{E}_\pi[(x_0 + A_1)^{-\alpha}]},$$

where $x_0 \equiv x_1/x_2$ denotes a ratio of quantities. Equivalently, we get:

$$\text{MRS}(x; \pi) = \frac{\mathbb{E}_\pi[(x_0 + A_1)^{-\alpha+1}]}{\mathbb{E}_\pi[(x_0 + A_1)^{-\alpha}]} - x_0 \equiv G_0(x_0), \quad (5.7)$$

which only depends on the ratio x_0 . Therefore, we have constructed homothetic preferences. By equation (5.7):

$$e(x) \equiv \frac{d}{dx} \log \mathbb{E}_\pi[(x + A_1)^{-\alpha+1}],$$

is identified up to a multiplicative constant. Therefore, by integration, $\mathbb{E}_\pi[(x + A_1)^{-\alpha+1}]$ is identified up to α and a multiplicative constant κ . However, as x tends to infinity, this expression is equivalent to $\kappa x^{-\alpha+1} \exp E(x)$, where $E(\cdot)$ is a primitive of $e(\cdot)$. This tail behaviour provides both the identification of α and κ . This analysis is summarized by the following result:

Corollary 2. Under the conditions of Proposition 7, if the marginal distribution of A_2 belongs to the Pareto family, the distributions of A_1 and A_2 are both non-parametrically identified.

5.2 Between Model Identification

Once the identification of the consumer's taste distribution π within each model is solved, we still need to consider the identification between the models. This analysis is needed to test whether preferences are consistent with SARA, or SSF, or both. It is important to know whether these two classes of preferences are nested or non-nested. If they are non-nested, we need to characterize their intersection and define a general class encompassing both types of preferences.

To illustrate, suppose that the consumer has SSF preferences:

$$U(x; \pi) = \mathbb{E}_\pi[\min\{x_1 + A_1 x_2, A_2\}], \quad (5.8)$$

where (i) A_1 and A_2 are independent, (ii) A_1 has distribution π_2 , and (iii) A_2 follows an exponential distribution (with unit intensity). Under this specification, we obtain:

$$U(x; \pi) = 1 - \mathbb{E}_\pi[\exp(-x_1 - A_1 x_2)]. \quad (5.9)$$

To clarify this result, observe that, by conditioning on A_1 , we are left with the expectation of the minimum of a set containing a constant and a random variable with an exponential distribution. This utility function is a strictly increasing transformation of a SARA utility function:

$$\tilde{U}(x; \pi) = -\mathbb{E}_\pi[\exp(-B'x)], \quad (5.10)$$

where (i) B_1 follows a point mass at 1, and (ii) B_2 has distribution π_2 . Consequently, these utility functions, one SARA, and the other SSF, induce the same preference ordering over the consumption set.

5.3 Discussion

The possible lack of identification of each consumer's taste distribution π_m has to be taken into account in the economic interpretation of the results. However, it has to be noted that it does not create difficulties for structural inference, where the (scalar or functional) parameters of interest are the parameters characterizing the MRS, rather than the parameters characterizing the utility function.

The lack of identification is due to the special structure of the cone of increasing and concave functions defined on R , and of the extremal elements of this cone. For finite increasing concave functions defined on \mathbb{R}_+ , it is well-known that the extremal functions are of the type:

$$h_0(x) = \min \{ \alpha_1 x + \beta_1, \alpha_2 x + \beta_2 \}, \quad (5.11)$$

in which $(\alpha_j, \beta_j) \in \bar{R}$, for $j = 1, 2$ (see [Blaschke and Pick, 1916](#)), and that any finite positive increasing concave function can be written as:

$$b + \mathbb{E}_\pi [\min(A, x)], \quad (5.12)$$

where b is a positive scalar and π is the distribution of A . Such functions are characterized by b and π . The set of extremal functions in (5.12) is a minimal set of extremal points generating the cone.

Such a property no longer holds for finite positive increasing concave functions defined on \bar{R} . [Johansen \(1974\)](#) has described a large set of extremal points of the type:

$$h_1(x) = \min \{ \alpha_1 x + \beta_1, \dots, \alpha_n x + \beta_n \}, \quad (5.13)$$

for which $h_1(\cdot)$ induces a covering with vertices of order 3 (see page 62 in [Johansen, 1974](#)), and has shown that this set is dense in the cone of finite continuous convex functions defined on a convex set in R (see Theorem 2 in [Johansen, 1974](#)). A minimal set of extremal points generating this cone does not exist. This argument explains why, in Sections 2 and 3, we consider specific subcones and convex subsets generated by parametric functions, and show, ex post, that these generating sets are not minimal.

While we restrict our attention to the SARA and SSF models, other convex hulls could have been considered. For example:

- (i) The convex hull generated by the union of the SARA and SSF models—that is, the smallest structural model containing both of the models in Sections 2 and 3.
- (ii) The convex hull generated by a basis of the form:

$$U(x; a, \nu) = \frac{a_1}{\nu_1} x_1^{\nu_1} + \frac{a_2}{\nu_2} x_2^{\nu_2}, \quad (5.14)$$

for every $x \in \bar{R}$ in which $a \in R$ and $\nu \in (0, 1)^2$. This basis corresponds to a first-order expansion of a utility function (see [Johansen, 1969](#)), and contains the Stone-Geary utility function as a limiting case. Indeed, as ν tends to zero, we obtain: $u(x; a) = a_1 \log x_1 + a_2 \log x_2$. However, the convex hull associated with this basis is not flexible enough, since it only contains additively separable utility functions.

6 An Illustration

This section shows how to use the SARA and SSF models in a non-parametric framework. First, we specify the statistical model by introducing an assumption on the observations, and then we discuss statistical inference. The methodology is illustrated in an application to alcohol consumption using scanner data concerning individual purchase histories.

6.1 Assumptions on Observations

The behavioural models introduced in the previous sections can be completed with an assumption on the available observations. We consider panel data, indexed by the consumer i and date t . After a preliminary treatment of the purchase histories, we have a large number n of consumers and a fixed number T of observed dates. In the preliminary treatment, the goods are aggregated into two groups using a common quantity unit and the dated purchases are aggregated by month (see Section 6.3). Recall that, under Assumption A1, we have M segments of homogeneous consumers.

We introduce the following assumption on the observations:

Assumption A2 (Observations).

- (i) We jointly observe (x_{it}, z_{it}) , for all $i = 1, \dots, n$ and $t = 1, \dots, T$, when $x_{it} > 0$.
- (ii) The individual histories $(x_{it}, z_{it})_{t=1}^T$ are independent given all π_m , $m = 1, \dots, M$.
- (iii) Designs (z_{it}) are exogenous (independent of taste distributions π_m).

Assumption A2 describes the structure of the observations. It implies that we can imagine taste parameters (π_m) being independently drawn from a Dirichlet process F , designs (z_{it}) being independently drawn from some distribution, and consumption x_{it} satisfying $x_{it} = X(z_{it}; \pi_{m_i})$, where m_i is the group of consumer i . Many papers assume that consumption x_{it} is positive (see Section IV.A in [Blundell, Horowitz, and Parey, 2017](#), for this assumption in an application to gasoline demand, as well as Assumption A5 in [Dobronyi and Gouriéroux, 2020](#), for this assumption in an application to the consumption of alcohol); the SARA and SSF models allow for corner solutions. However, in many datasets (including the dataset used in the application in Section 6), there is a problem of partial observability. Let \tilde{y} denote the expenditure (prior to normalization), and let \tilde{p}_j denote the price of good j (prior to normalization). Usually, we only observe the price \tilde{p}_j of a good j when the consumer buys a positive quantity of good j . Then, we only observe (normalized) expenditure y when the consumer buys a positive quantity of good 2, and we only observe the (normalized) price p when the consumer buys a positive quantity of both goods (see [Crawford and Polisson, 2016](#), for an approach to revealed preference that deals with this partial observability problem). This problem explains the specific form of Assumption 2(i).

For deriving the asymptotic properties of estimators, it is also necessary to specify the type of asymptotics to be considered:

Assumption A3. Let n_m denote the size of the m^{th} homogeneous group.

- (i) $n_m T \rightarrow \infty$, as $n \rightarrow \infty$, for all $m = 1, \dots, M$.
- (ii) $n_m T \sim \lambda_m n$, for some $\lambda_m \in (\lambda_\ell, \lambda_h)$, where $0 < \lambda_\ell < \lambda_h < 1$, for $m = 1, \dots, M$.
- (iii) $M \rightarrow \infty$, as $n \rightarrow \infty$.

Assumptions A3(i) and A3(ii) ensure that there are enough observations to non-parametrically estimate the demand function associated with the functional parameter π_m on a sufficiently large subset \mathcal{Z}_m of designs z . Assumption A3(iii) guarantees enough filtered parameters $\hat{\pi}_m$ to estimate the underlying Dirichlet process F . In some special circumstances, T is large, and Assumption A3 can be used with $m = i$ and $M = n$ —that is, a single consumer per group. Otherwise, grouping of homogeneous consumers is needed to identify the demand functions on sufficiently large subsets \mathcal{Z}_m .

6.2 Estimation Method

The Dirichlet process is common in Bayesian estimation (see, for instance, Ferguson, 1974, and Li et al., 2019). This process is useful because it is flexible and, if observations are independently and identically drawn from an unknown distribution, the posterior distribution of this distribution has a closed-form expression. However, our framework is much more complicated for two reasons:

- (i) The observed consumption choices (X_{ijt}) are not identically distributed because consumers make decisions at *different expenditures and prices*.
- (ii) It is difficult to derive a closed-form expression for the demand, as a function of the expenditure, the price, and the functional parameter characterizing taste uncertainty π . It is, therefore, difficult to derive a closed-form expression for the distribution of X_{ijt} conditional on Z_{it} .

These features of our model explain why estimation requires specific numerical algorithms. These specific algorithms have to be able to deal with the non-linear and high-dimensional features of the models. In the Bayesian framework, the Dirichlet process is fixed. In the hyperparametric framework, it is parameterized by a vector θ . These parameters have to be estimated and the functional parameters (π_m) have to be filtered. These estimation approaches are described below.

6.2.1 Bayesian Framework

In a pure Bayesian framework, a Dirichlet process is fixed by selecting a mean distribution μ and a scaling parameter c (see Appendix A). This distribution defines the common prior for the functional taste parameters (π_m) . After, the data are used to compute the posterior distribution for the functional parameters (π_m) . Under Assumption A2, the posterior distribution can be computed separately for each homogeneous group of consumers:

$$\ell(\pi_m | x_{it}, z_{it}, x_{it} > 0, i \in \Lambda_m, t = 1, \dots, T),$$

where Λ_m denotes the group of consumers with preferences characterized by the taste parameter π_m . This approach does not have to account for the potential identification problem discussed in Section 5. If a specific characteristic of π_m is weakly identified, its posterior distribution will be close to the prior distribution.

In our framework, the observations (x_{it}, z_{it}) , conditional on $x_{it} > 0$, must satisfy the deterministic first-order conditions implied by the model. These conditions have the following form:

$$\text{MRS}(x_{it}; \pi) = p_{it}, \quad (6.1)$$

for any observed pair (x_{it}, z_{it}) . Equivalently:

$$\mathbb{E}_\pi \left[\frac{\partial U(x_{it}; A)}{\partial x_1} \right] = p_{it} \mathbb{E}_\pi \left[\frac{\partial U(x_{it}; A)}{\partial x_2} \right], \quad (6.2)$$

for any observed pair (x_{it}, z_{it}) . These conditions are moment restrictions, called *MRS restrictions*. In our big data framework, the number of MRS restrictions is very large, typically several hundred to a thousand. The posterior of π_m is simply the distribution of π_m given these deterministic restrictions on π_m . If the taste distributions, A_1 and A_2 , are independent with marginal distributions, π_1 and π_2 , respectively, then the MRS restrictions are bilinear in π_1 and π_2 —specifically, these restrictions are linear in π_1 given π_2 , and linear in π_2 given π_1 . Later, this property is used to construct a numerically efficient optimization algorithm for filtering all the π_m (see Appendix C).

6.2.2 Hyperparametric Framework

The hyperparametric framework is a complicated non-linear state-space model with two layers of latent state variables. Such a framework can be characterized as follows:

- (i) **Deep layer:** Taste parameters π_m drawn from F (which is parameterized by θ);
- (ii) **Surface layer:** Demand functions $X(\cdot; \pi_m)$ deduced from π_m ;
- (iii) **Measurement equations:** Observed pairs (x_{it}, z_{it}) , given $x_{it} > 0$.

We have partial observability of the demand function because the value of demand $X(z; \pi_m)$ is observed at a finite number of designs z . Moreover, unlike most state-space models, the state variables are infinite-dimensional.

6.2.3 Estimating the Hyperparameter

While it is difficult to derive analytically the distribution of X_{it} given Z_{it} , it is easy to simulate its distribution for a given value of θ (see Appendix A for simulations from the Dirichlet distribution). Therefore, θ can be estimated by the method of simulated moments (MSM), or indirect inference (see [McFadden, 1989](#), [Pakes and Pollard, 1989](#), and [Gouriéroux and Monfort, 1996](#)). That is, θ is estimated by matching some sample and simulated moments of the pair (X_{it}, Z_{it}) .

To illustrate, consider a pure panel such that $M = n$.¹⁰ The steps are the following:

Step 1: Simulate $s = 1, \dots, n$ draws from a Dirichlet process given the parameter θ . Each draw $\pi^s(\theta)$ is associated with an individual consumer i such that $s = i$.

Step 2: Compute simulated consumption $x_{it}^s(\theta)$ by solving the first-order condition in (2.11) with respect to x_1 and applying the transformation in (2.14) given $z_{it} = (y_{it}, p_{it})$ and $\pi^i(\theta)$.

Step 3: Construct a collection of K moments from the observed and simulated data:

$$m \equiv \left[\frac{1}{nT} \sum_{i=1}^n \sum_{t=1}^T m_k(x_{it}, z_{it}) \right]_k \quad \text{and} \quad m(\theta) \equiv \left[\frac{1}{nT} \sum_{i=1}^n \sum_{t=1}^T m_k(x_{it}^s(\theta), z_{it}) \right]_k.$$

Then, numerically solve the following problem:

$$\underset{\theta}{\operatorname{argmin}} \quad \|m - m(\theta)\|, \quad (6.3)$$

in which $\|\cdot\|$ is a Euclidean norm with the form $\|m\| = m' \Omega m$, for some positive-definite $K \times K$ matrix Ω .

Given the estimated hyperparameter $\hat{\theta}$, the taste distributions (π_m) must be filtered. This step is equivalent to applying the Bayesian approach with the estimated Dirichlet distribution as the prior distribution (see Appendix C).

Under Assumptions A1 to A3, the estimator for θ is consistent and asymptotically normal, and it converges at a speed of $1/\sqrt{nT}$. The derivation of the asymptotic properties of the filtered functional parameter $\hat{\pi}_m$ is out of the scope of this paper and left for future research.

6.2.4 Filtering the Taste Distributions

Once the hyperparameter θ is estimated, we can filter π_m by using the following steps:

Step 1: Draw a taste distribution $\tilde{\pi}_m$ from the Dirichlet process given $\hat{\theta}$. Then, by construction, the taste distribution $\tilde{\pi}_m$ is a draw from the prior distribution.

¹⁰When $M < n$, we simulate $n_m T$ observations for the m^{th} draw from the Dirichlet process.

Step 2: Discretize $\tilde{\pi}_m$ on a grid of values for the taste parameters, A_1 and A_2 . Let $\bar{\pi}_m$ denote the result. The aim of this step is to put $\tilde{\pi}_m$ on a grid for optimization.

Step 3: Solve the minimization problem:

$$\min_{\pi} \|\pi - \bar{\pi}_m\| \text{ s.t. MRS restrictions (6.1) and unit mass restrictions.}$$

Let $\hat{\pi}_m^*$ denote the solution. This solution approximates a drawing from the posterior.

Step 4: Replicate these steps to obtain a sequence of solutions: $\hat{\pi}_{m,s}^*$, $s = 1, \dots, S$, where S is the number of replications. The filtered $\hat{\pi}_m$ is obtained by averaging over all simulations such that:

$$\hat{\pi}_m = \frac{1}{S} \sum_{s=1}^S \hat{\pi}_{m,s}^*$$

This procedure involves a high-dimensional argument π_m , and a very large number of MRS restrictions. Indeed, we need several hundred grid points for π_m , and, in the application, we have about one-thousand MRS restrictions, for each $m = 1, \dots, M$. If the taste parameters, A_1 and A_2 , are independent, this procedure can be numerically simplified by using the fact that these restrictions are bilinear (see Section 6.2.1 and Appendix C).

6.3 The Data

We use the Nielsen Homescan Consumer Panel (NHCP). Nielsen provides a sample of households with barcode scanners. Households are asked to scan all purchased goods on the date of each purchase. The prices are entered by the households or linked to retailer data by The Nielsen Company. The households that agree to participate are compensated through benefits and lotteries.

We focus on the consumption of alcoholic drinks (see Manning et al., 1995, for an application to alcohol consumption in economics). We classify drinks by type. Good 1 contains beers and ciders.¹¹ Good 2 contains wines and liquors. We disregard all non-alcoholic beers, ciders, and wines. We are left with 30,635 beers and ciders, and 108,439 wines and liquors, for a total of 139,074 drinks. We convert all measurement units to litres of alcohol by first converting all units to litres and then multiplying by the standard alcohol by volume (ABV) in each subgroup—specifically, 4.5% for beer and cider, 11.6% for wine, and 37% for liquor. For example, if a household buys two packs of six bottles of beer and each bottle contains 355 millilitres of beer, then the household buys 4.26 litres of this beer, or $4.26 \times 0.045 = 0.231$ litres of alcohol. We use

¹¹The NHCP classifies ciders as wine, by default. We reclassify these beverages using UPC product descriptions because most ciders have a low alcohol by volume (ABV).

Table 1. Mean m , standard deviation σ , the ratio $\frac{\sigma}{m}$, and quantiles for expenditure \tilde{y} , prices \tilde{p}_j , normalized expenditure y , and normalized price p . Normalized expenditures y and prices p are conditional on being defined.

Var.	m	σ	σ/m	Quantiles					N
				0%	25%	50%	75%	100%	
\tilde{y}	52.29	70.62	1.35	0.00	13.28	28.17	62.97	2,767.76	63,972
\tilde{p}_1	70.36	44.19	0.62	0.05	46.41	62.55	83.41	2,893.65	33,077
\tilde{p}_2	61.33	326.93	5.33	0.03	29.29	48.56	75.14	39,900.85	45,518
y	1.60	3.48	2.17	0.00	0.27	0.74	1.83	228.57	45,518
p	2.20	4.06	1.54	0.00	0.89	1.37	2.29	139.76	14,659

the standard ABV in each subgroup as a result of data limitations. Our sample only contains purchases made at stores, not purchases made at bars, or restaurants.

We restrict our sample to purchases made from August to November in 2016. This relatively short window is used to diminish the impact of changing tastes and product availability, and to avoid most federal holidays in the United States that are often associated with alcohol consumption such as Independence Day, Christmas Day, and New Year’s Eve. Our sample contains 28,036 households. Some additional details of this restricted sample are placed in Appendix D.

The dated purchases are aggregated by month. For each household and month, the prices are constructed by dividing the total expenditure for each aggregate good (after accounting for the value of coupons) by the amount of alcohol of that aggregate good purchased by the household, when this amount is strictly positive. Then, we normalize by the price of good 2. This procedure yields four monthly observations per household for a total of 112,144. A total of 63,936 observations have positive consumption such that $x_{it} > 0$. Table 1 gives summary statistics conditional on $x_{it} > 0$. The prices (p_{it}) are conditional on being well-defined (see the discussion of partial observability on page 30). For the interpretation of the results, recall that \tilde{y} denotes the expenditure (prior to normalization), and that \tilde{p}_j denotes the price of good j (prior to normalization).

Table 2. Proportion of observations by type.

	$x_2 = 0$	$x_2 > 0$
$x_1 = 0$	0.4298	0.2751
$x_1 > 0$	0.1642	0.1307

There are four regimes of observations: (i) zero expenditure on all goods, (ii) zero




Figure 3. Sample Distributions of Expenditure \tilde{y} by Regime. On the left, we illustrate the sample distribution conditional on $x_1 > 0$ and $x_2 > 0$; on the right, the light histogram illustrates the sample distribution conditional on $x_1 = 0$ and $x_2 > 0$, and the dark histogram illustrates the sample distribution conditional on $x_1 > 0$ and $x_2 = 0$.

expenditure on good 1 and strictly positive expenditure on good 2, (iii) strictly positive expenditure on good 1 and zero expenditure on good 2, and (iv) strictly positive expenditure on all goods. Table 2 provides the proportion of observations in each regime, and shows a large proportion of observations with zero expenditure. Recall, under Assumption A2, designs z_{it} are drawn from a distribution. Therefore, we can interpret this result as a mass at zero in the marginal distribution of expenditure.

Figure 3 displays the sample distribution of expenditure \tilde{y}_{it} by regime: the distribution of expenditure \tilde{y}_{it} conditional on $x_{it} > 0$ is on the left; the sample distributions of expenditure \tilde{y}_{it} for the two other regimes with positive expenditure are on the right. The shape of the sample distribution of expenditure \tilde{y}_{it} does not appear to vary all that much with the regime. That being said, the sample distribution conditional on $x_{it} > 0$ has more probability attributed to higher expenditures.

Figure 4 compares the sample distributions of prices \tilde{p}_j by regime: the sample distributions of \tilde{p}_1 are on the left; the sample distributions of \tilde{p}_2 are on the right. Although the sample distribution of \tilde{p}_1 differs from the sample distribution of \tilde{p}_2 , these distributions do not seem to be affected by the regime.

Figure 5 displays the sample distributions of (normalized) designs $z_{it} = (y_{it}, p_{it})$ and the components of consumption x_{it} given $x_{it} > 0$. As expected, the components of consumption x_{it} are increasing in expenditure y_{it} . Furthermore, the first component of consumption x_{it} is more affected by changes in the price p_{it} than the second component.

Since we consider a rather short window of time, we follow the segmented population approach. We segment the population by state. Large states (e.g. California)




Figure 4. Sample Distributions of Prices \tilde{p}_j by Regime. On the left, the light histogram illustrates the sample distribution of \tilde{p}_1 conditional on $x_1 > 0$ and $x_2 = 0$; on the right, the light histogram illustrates the sample distribution of \tilde{p}_2 conditional on $x_1 = 0$ and $x_2 > 0$; in each plot, the dark histogram illustrates the sample distribution conditional on $x_1 > 0$ and $x_2 > 0$.




Figure 5. Sample Distributions of Designs z_{it} and Consumption. These figures are conditional on $x_{it} > 0$. On the left, colour describes the quantity of good 1; on the right, colour describes the quantity of good 2.

are segmented again by county. Specifically, a county is given its own segment if it has more than 70 observations with positive consumption and it is in a state with more than 1,000 observations with positive consumption. We are left with a total of 65 segments, each corresponding to a state or county. The smallest segment is Wyoming, containing 15 observations with positive consumption; the largest state is Florida (after removing Broward, Hillsborough, Palm Beach, Pinellas, and Miami-Dade counties), containing 880 observations with positive expenditure; the mean number of observations with positive consumption per segment is approximately 226.

Figure 6 displays the sample distributions of (normalized) designs $z_{it} = (y_{it}, p_{it})$ and the first component of consumption x_{it} given $x_{it} > 0$ in two of the larger segments: California (after removing Alameda, Los Angeles, Orange, Riverside, Sacramento, San Bernardino, and San Diego counties), and Florida (after removing Broward, Hillsborough, Palm Beach, Pinellas, and Miami-Dade counties).

Figure 7 displays the Nadaraya-Watson (kernel) estimates of the demand function for beer conditional on $x_{it} > 0$ in California and Florida over a subset of the domain of designs. Demand for beer in California is lower and less responsive to price changes than in Florida.

6.4 Estimation Results

As an illustration, we consider the SARA model in the hyperparametric framework. We assume that the taste parameters, A_1 and A_2 , are independent. Under this assumption, the taste uncertainty is characterized by the marginal distributions, π_1 and π_2 . The marginal distribution π_j of A_j is independently drawn from a Dirichlet process F_j , $j = 1, 2$. The mean of F_j is a log-normal distribution with parameters μ_j and σ_j , and the scale parameter of F_j is c_j . The utility function corresponding to this log-normal mean distribution, say $\bar{\pi}_j$, has a quasi closed-form expression. Indeed, under this distribution, we can write:

$$\log(A_j) = \mu_j + \sigma_j \varepsilon_j, \quad \forall j = 1, 2,$$

where ε_j is distributed with respect to a standard normal distribution. Then:

$$\begin{aligned} \mathbb{E}_{\bar{\pi}_j}[\exp(-A_j x_j)] &= \mathbb{E}_{\bar{\pi}_j}[\exp(-\exp(\mu_j + \sigma_j \varepsilon_j) x_j)], \\ &= \frac{1}{\sqrt{1 + w(x_j \exp(\mu_j) \sigma_j^2)}} \exp \left\{ -\frac{1}{2\sigma_j^2} w(x_j \exp(\mu_j) \sigma_j^2)^2 - \frac{1}{\sigma_j^2} w(x_j \exp(\mu_j) \sigma_j^2) \right\}, \end{aligned}$$

where $w(x)$ is the Lambert function, defined by the implicit equation:

$$w(x) \exp(w(x)) = x,$$

[see equation (1.3) in [Asmussen et al., 2016](#)]. By drawing from the Dirichlet process, we will draw a stochastic utility function around the closed-form expression above. The hyperparameter θ has six components such that:

$$\theta = (\mu_1, \sigma_1, c_1, \mu_2, \sigma_2, c_2).$$




Figure 6. Sample Distributions by State. These figures are conditional on $x_{it} > 0$. California is shown on the top, and Florida is shown on the bottom. On the left, colour describes the quantity of good 1; on the right, colour describes the quantity of good 2.




Figure 7. Demand. Nadaraya-Watson estimates of the demand function for good 1 conditional on $x_{it} > 0$ in California (left) and Florida (right).

6.4.1 The Hyperparameter

As described in Section 6.2.2, the first step involves estimating the hyperparameter θ using the Method of Simulated Moments (MSM). The hyperparameter θ is calibrated by using the following (sample and simulated) moments computed for all of the 63,936 observations with positive consumption:

- (i) marginal moments of (X_{it}) ;
- (ii) cross-moments of $(\log X_{it}, \log P_{it})$ and $(\log X_{it}, \log Y_{it})$;
- (iii) cross-moments of (X_{it}, P_{it}) , (X_{it}, Y_{it}) , $(X_{it}, \log P_{it})$, and $(X_{it}, \log Y_{it})$.

The moments in (ii) are the moments used in the Almost Ideal Demand System (see Deaton and Muellbauer, 1980); the moments in (iii) are introduced in order to capture risk effects, by comparison with the moments in (ii). The optimum is found using a random search algorithm over a sufficiently big support.¹²

To apply MSM, it is necessary to compute simulated consumption $x_{it}^s(\theta)$ for every observation, at each step of the optimization algorithm. This procedure is computationally costly. Note that, the number of simulated observations with positive consumption is stochastic, and not necessarily equal to the number of observations with positive consumption in the sample. This aspect has no impact on the consistency of the MSM estimator.

The estimated hyperparameter is:

$$\hat{\theta} = (0.7987, 3.5516, 45.0951, 0.1201, 3.6597, 3.5544). \quad (6.4)$$

¹²Random search is more efficient than grid search in hyperparameter optimization (Bergstra and Yoshua, 2012).




Figure 8. On the left, black shows the density of the mean of the Dirichlet process for A_1 , and blue shows the density of the mean of the Dirichlet process for A_2 . The x -axis is in log-scale. For scale: $\exp(-5) \simeq 0.0067$ and $\exp(5) \simeq 148.4131$. The figure on the right displays indifference curves associated with these distributions.

Therefore, the median level of risk aversion for the mean of the Dirichlet process¹³ for A_1 is $\exp(0.5495) \simeq 2.2226$, and the median level of risk aversion for the mean of the Dirichlet process for A_2 is $\exp(0.8738) \simeq 1.1276$. The fact that μ_1 is smaller than μ_2 is expected: Since quantities are measured in terms of volume of alcohol, this result is consistent with the faster overall intake of alcohol when consuming drinks with a higher ABV. Moreover, the distribution π_1 of A_1 is much more concentrated around its mean than the distribution π_2 of A_2 , as the scaling parameter $c_1 = 45.0951$ for π_1 is much larger than the scaling parameter $c_2 = 3.5544$ for π_2 .

We do not report any standard errors because they are automatically small from the large number of observations. Indeed, the standard significance test procedures (such as comparing a t -statistic to the critical value of a standard normal at the 5% significance level) are not relevant in this big data framework. The highest degree of uncertainty concerns the filtered parameters ($\hat{\pi}_m$) because π_m is a high-dimensional parameter and the number of observations in each segment Λ_m is much smaller.

The means of these Dirichlet processes are displayed in the left panel in Figure 8. The right panel displays the indifference curves associated with utility levels -0.1000 , -0.0800 , and -0.0680 for a draw from the Dirichlet process given $\hat{\theta}$.

Figure 9 displays the Q-Q plots for two draws (π_1^s, π_2^s) , $s = 1, 2$, from the Dirichlet process given $\hat{\theta}$. In particular, we plot the quantiles of the realization of the distribution π_j of $\log(A_j)$ against the quantiles of the normal distribution given the estimated

¹³This is not the absolute risk aversion of the utility function for the log-normal mean distribution which depends on the consumption level and has to be computed with a modified density.




Figure 9. The Q-Q plots for two draws from the Dirichlet process given $\hat{\theta}$: On the left, the quantiles of $\log(A_1)$ are plotted against the quantiles of the normal distribution given $(\hat{\mu}_1, \hat{\sigma}_1)$; on the right, the quantiles of $\log(A_2)$ are plotted against the quantiles of the normal distribution given $(\hat{\mu}_2, \hat{\sigma}_2)$. In each figure, the green line is the 45-degree line.

hyperparameters $(\hat{\mu}_j, \hat{\sigma}_j)$, for $j = 1, 2$. If these quantiles coincide exactly, they will lie on the 45-degree line. As expected, these Q-Q plots lie approximately around the 45-degree line. The draws (π_1^s) , $s = 1, 2$, for π_1 are closer the 45-degree line and “more continuous” than the draws (π_2^s) , $s = 1, 2$, for π_2 since $c_1 > c_2$.

6.5 Taste Distributions

This section uses the filtering approach described in Section 6.2.4 to recover π_m . In the SARA model, the MRS restriction in (6.2) is:

$$\mathbb{E}_\pi[A_1 \exp(-A'x_{it})] = p_{it} \mathbb{E}_\pi[A_2 \exp(-A'x_{it})].$$

When A_1 and A_2 are independent, this expression becomes:

$$\begin{aligned} & \mathbb{E}_{\pi_1}[A_1 \exp(-A_1 x_{i1t})] \mathbb{E}_{\pi_2}[\exp(-A_2 x_{i2t})] \\ &= p_{it} \mathbb{E}_{\pi_1}[\exp(-A_1 x_{i1t})] \mathbb{E}_{\pi_2}[A_2 \exp(-A_2 x_{i2t})]. \end{aligned} \tag{6.5}$$

To filter π_m , these restrictions have to be imposed for every observation with positive consumption x_{it} associated with segment Λ_m . In California, there are 688 MRS restrictions, and, in Florida, there are 880. Appendix C describes how to numerically solve the resulting optimization problem given the bilinearity of the MRS restrictions under independence.

Figure 10 displays the Q-Q plots for the filtered taste parameter $\hat{\pi}_m$ for California and Florida. In both states, the slope on the left is steeper than the 45-degree line,




Figure 10. The Q-Q plots for the filtered taste distributions for California (black) and Florida (blue): On the left, the quantiles of $\log(A_1)$ are plotted against the quantiles of the normal distribution given $(\hat{\mu}_1, \hat{\sigma}_1)$; on the right, the quantiles of $\log(A_2)$ are plotted against the quantiles of the normal distribution given $(\hat{\mu}_2, \hat{\sigma}_2)$. In each figure, the green line is the 45-degree line.

suggesting that the posterior mean distribution for A_1 is more “dispersed” than its (prior) mean distribution. The convexity of the curves also suggests fatter tails.

For the structural interpretation of these plots, assume that (i) the preferences are SARA, (ii) the taste parameters are independent, (iii) the marginal distribution of A_1 is the same in both states, and (iv) the marginal distribution of A_2 “shifts” such that $\pi'_2(A_2) = \pi_2(cA_2)$, where π_2 and π'_2 denote the marginal distributions of A_2 in these states. Under these assumptions:

$$U(x_1, x_2; \pi') = U(x_1, cx_2; \pi), \quad (6.6)$$

and solving the utility maximization problem in (2.12) yields:

$$X_1(z; \pi') = X_1(cz; \pi) \quad \text{and} \quad X_2(z; \pi) = \left(\frac{1}{c}\right) X_2(cz; \pi). \quad (6.7)$$

Similarly, if there is a “shift” in the marginal distribution of A_1 and the marginal distribution of A_2 is the same in both states, we obtain:

$$X_1(z; \pi') = \left(\frac{1}{c}\right) X_1\left(y, \frac{p}{c}; \pi\right) \quad \text{and} \quad X_2(z; \pi) = X_2\left(y, \frac{p}{c}; \pi\right). \quad (6.8)$$

The relationships given in (6.7) and (6.8) suggest that there exists a complicated non-linear relationship between such demand functions. Therefore, we cannot immediately deduce from Figure 10 which state has a higher demand for beer. For a more formal

analysis, the utility functions associated with each posterior mean taste distribution must be used to derive a posterior MRS, or a posterior demand function.

This analysis has to be completed with a discussion of accuracy. In this framework, the posterior distributions of π_1 and π_2 are infinite-dimensional and cannot be represented. However, posterior distributions of any scalar transformation of π_1 and π_2 can be derived using simulation. In this respect, it is important to know which scalars are of interest. Typically, we are most interested in the MRS, or counterfactual demand, corresponding to a particular design $z_0 = (y_0, p_0)$.

7 Concluding Remarks

This paper is one among pioneering papers attempting to tackle the challenges of performing structural demand analysis with scanner data (see also [Burda et al., 2008, 2012](#), [Crawford and Polisson, 2016](#), [Guha and Ng, 2019](#), [Chernozhukov et al., 2020](#), and [Dobronyi and Gouriéroux, 2020](#)). The recent availability of scanner data permits new developments in the analysis of consumer behaviour. Here, we have shown that, by introducing homogeneous segments of consumers, we can consider a model of consumption with non-parametric preferences and infinite-dimensional heterogeneity, not only from a theoretical point-of-view, but also from a practical one. The distribution of individual heterogeneity in the population can be estimated, and the underlying non-parametric preferences can be filtered by using appropriate algorithms.

We developed an analysis for *two goods* for exposition. This feature of our analysis leaves the question: Can the methods developed in this paper be extended to a framework with, say, 100 goods? A completely *unconstrained* non-parametric analysis would encounter the curse of dimensionality. Specifically, we would need to estimate the distribution of the utility function (a non-parametric function with, in this scenario, 100 arguments). This task would be infeasible, even in our big data framework. But, the SARA model with independent taste parameters is a *constrained* non-parametric model. The structure of the SARA model reduces the non-parametric dimension of the problem, making it feasible. Indeed, when taste parameters are independent, we only need to estimate 100 one-dimensional distributions. A similar remark applies to the algorithm used to filter the taste distributions: The two steps based on the bilinear form of the MRS restrictions in a two good setting can be replaced with 100 successive steps based on the multilinear form of MRS restrictions in a 100 good setting, without increasing the numerical complexity.

In this paper, the consumers are assumed to be rational and divided into homogeneous segments. Since, in each homogeneous segment, the demand function can be non-parametrically estimated over a subset of its domain, the analysis can be continued in order to develop:

- (i) A test for the homogeneity of each segment, or, more generally, a non-parametric method for constructing homogeneous segments.

- (ii) A test of the rationality of the consumers in each segment. That would be, a test of the negative semi-definiteness of the Slutsky matrix (as symmetry is automatic in the case of two goods). In our framework, the integrability condition can be written explicitly in terms of the taste distribution π , or its Laplace transform. A test of this condition can be performed directly, without considering only necessary implications of rationality (see [Dette et al., 2016](#)).

References

- ALLEY, A., D. FERGUSON, AND K. STEWART (1992): “An Almost Ideal Demand System for Alcoholic Beverages in British Columbia,” *Empirical Economics*, 17, 401–418.
- ASMUSSEN, S., J. JENSEN, AND L. ROJAS-NANDAYAPA (2016): “On the Laplace Transform of the Lognormal Distribution,” *Methodology and Computing in Applied Probability*, 18, 441–458.
- BANKS, J., R. BLUNDELL, AND A. LEWBEL (1997): “Quadratic Engel Curves and Consumer Demand,” *The Review of Economics and Statistics*, 79, 527–539.
- BARTEN, A. (1968): “Estimating Demand Equations,” *Econometrica*, 36, 213–251.
- BECKERT, W. AND R. BLUNDELL (2008): “Heterogeneity and the Non-Parametric Analysis of Consumer Choice: Conditions for Invertibility,” *The Review of Economic Studies*, 75, 1069–1080.
- BERGSTRA, J. AND Y. YOSHUA (2012): “Random Search for Hyper-Parameter Optimization,” *Journal of Machine Learning Research*, 13, 281–305.
- BLASCHKE, W. AND G. PICK (1916): “Distanzschätzungen im Funktionenraum II,” *Mathematische Annalen*, 77, 277–302.
- BLOMQUIST, S., A. KUMAR, C. LIANG, AND W. NEWHEY (2015): “Individual Heterogeneity, Nonlinear Budget Sets, and Taxable Income,” *CESifo Working Paper*.
- BLUNDELL, R., D. KRISTENSEN, AND R. MATZKIN (2017): “Individual Counterfactuals with Multidimensional Unobserved Heterogeneity,” *CeMMAP Working Paper*.
- BROWN, B. AND M. WALKER (1989): “The Random Utility Hypothesis and Inference in Demand Systems,” *Econometrica*, 57, 815–829.
- BROWN, D. AND R. MATZKIN (1995): “The Random Utility Model from Data on Consumer Demand,” *Discussion Paper, Yale University*.

- BURDA, M., M. HARDING, AND J. HAUSMAN (2008): “A Bayesian Mixed Logit-Probit Model for Multinomial Choice,” *Journal of Econometrics*, 147, 232–246.
- (2012): “A Poisson Mixture Model of Discrete Choice,” *Journal of Econometrics*, 166, 184–203.
- CHERNOZHUKOV, V., J. HAUSMAN, AND W. NEWHEY (2020): “Demand Analysis with Many Prices,” *Forthcoming in the Journal of Econometrics*.
- CHRISTENSEN, L., D. JORGENSON, AND L. LAU (1975): “Transcendental Logarithmic Utility Functions,” *The American Economic Review*, 65, 367–383.
- CRAWFORD, I. AND M. POLISSON (2016): “Demand Analysis with Partially Observed Prices,” *University of Leicester Working Paper*, 15/12.
- DEATON, A. AND J. MUELLBAUER (1980): “An Almost Ideal Demand System,” *The American Economic Review*, 70, 312–326.
- DETTE, H., S. HODERLEIN, AND N. NEUMEYER (2016): “Testing Multivariate Economic Restrictions Using Quantiles: The Example of Slutsky Negative Semidefiniteness,” *Journal of Econometrics*, 191, 129–144.
- DOBRONYI, C. AND C. GOURIÉROUX (2020): “Stochastic Revealed Preference: A Non-Parametric Analysis,” *Working Paper*.
- FELLER, W. (1968): *An Introduction to Probability Theory and its Applications*, vol. 2, Wiley.
- FERGUSON, T. (1974): “Prior Distributions on Spaces of Probability Measures,” *The Annals of Statistics*, 2, 615–629.
- GEWEKE, J. (2012): “Nonparametric Bayesian Modelling of Monotone Preferences for Discrete Choice Experiments,” *The Journal of Econometrics*, 171, 185–204.
- GOURIÉROUX, C. AND A. MONFORT (1996): *Simulation-Based Econometric Methods*, New York: Oxford University Press.
- GOURIÉROUX, C., A. MONFORT, AND E. RENAULT (1990): “Bilinear Constraints: Estimation and Tests,” *Essays in Honor of Edmond Malinvaud, Empirical Economics*, MIT Press, 166–191.
- GRANT, S. (1995): “A Strong (Ross) Characterization of Multivariate Risk Aversion,” *Theory and Decision*, 38, 131–152.
- GUHA, R. AND S. NG (2019): “A Machine Learning Analysis of Seasonal and Cyclical Sales in Weekly Scanner Data,” *NBER 25899*.

- HALEVY, Y. AND V. FELTKAMP (2005): “A Bayesian Approach to Uncertainty Aversion,” *Review of Economic Studies*, 72, 449–466.
- HAUSMAN, J. AND W. NEWEY (2016): “Individual Heterogeneity and Average Welfare,” *Econometrica*, 84, 1225–1248.
- HEYDE, C. (1963): “On a Property of the Lognormal Distribution,” *Journal of the Royal Statistical Society Series B: Methodological*, 29, 16–18.
- HOSOYA, Y. (2016): “On First-Order Partial Differential Equations: An Existence Theorem and its Applications,” *Advances in Mathematical Economics*, 20, 77–87.
- HURWICZ, L. AND H. UZAWA (1971): “On the Integrability of Demand Functions,” in *Preferences, Utility, and Demand: A Minnesota symposium*, New York, chap. 6, 114–148.
- JOHANSEN, L. (1969): “On the Relationships Between Some Systems of Demand Functions,” *Liiketaloudellinen Aikakauskirja*, 1, 30–41.
- JOHANSEN, S. (1974): “The Extremal Convex Functions,” *Mathematica Scandinavica*, 34, 61–68.
- KARNI, E. (1979): “On Multivariate Risk Aversion,” *Econometrica*, 47, 1391–1401.
- (1983): “On the Correspondence Between Multivariate Risk Aversion and Risk Aversion with State-Dependent Preferences,” *Journal of Economic Theory*, 30, 230–242.
- KITAMURA, Y. AND M. STUTZER (1997): “An Information-Theoretic Alternative to Generalized Method of Moments Estimation,” *Econometrica*, 65, 861–874.
- KOTZ, S., N. BALAKRISHNAN, AND N. JOHNSON (2000): *Continuous Multivariate Distributions, Volume 1: Models and Applications*, New York: Wiley, 2nd ed.
- LEDOIT, O. AND M. WOLF (2004): “A Well-Conditioned Estimator for Large-Dimensional Covariance Matrices,” *Journal of Multivariate Analysis*, 88, 365–411.
- LI, Y., E. SCHOFIELD, AND M. GÖNEN (2019): “A Tutorial on Dirichlet Process Mixture Modeling,” *Journal of Mathematical Psychology*, 91, 128–144.
- LIN, J. (2016): “On the Dirichlet Distribution,” *Queens University, Kingston*.
- MANNING, W., L. BLUMBERG, AND L. MOULTON (1995): “The Demand for Alcohol: The Differential Response to Price,” *Journal of Health Economics*, 14, 123–148.
- MATZKIN, R. (2003): “Nonparametric Estimation of Nonadditive Random Functions,” *Econometrica*, 71, 1339–1376.

- McFADDEN, D. (1989): “A Method of Simulated Moments for Estimation of Discrete Response Models Without Numerical Integration,” *Econometrica*, 57, 995–1026.
- MOSCHINI, G. (1998): “The Semiflexible Almost Ideal Demand System,” *European Economic Review*, 42, 349–364.
- NAVARRO, D., T. GRIFFITHS, M. STEYVERS, AND M. LEE (2006): “Modeling Individual Differences Using Dirichlet Processes,” *Journal of Mathematical Psychology*, 50, 101–122.
- NG, S. (2017): “Opportunities and Challenges: Lessons from Analyzing Terabytes of Scanner Data,” in *Advances in Economics and Econometrics*, Cambridge, vol. 2.
- PAKES, A. AND D. POLLARD (1989): “Simulation and the Asymptotics of Optimization Estimators,” *Econometrica*, 57, 1027–1057.
- RICHARD, S. (1975): “Multivariate Risk Aversion, Utility Independence and Separable Utility Functions,” *Management Science*, 22, 12–21.
- ROCKAFELLAR, R. (1970): *Convex Analysis*, Princeton: Princeton University Press.
- ROLIN, J. (1992): “Some Useful Properties of the Dirichlet Process,” *CORE Discussion Paper, Université Catholique de Louvain*.
- ROY, A. (1952): “Safety First and the Holding of Assets,” *Econometrica*, 20, 431–449.
- SAMUELSON, P. (1948): “Consumption Theory in Terms of Revealed Preference,” *Economica*, 15, 243–253.
- (1950): “The Problem of Integrability in Utility Theory,” *Economica*, 17, 355–385.
- SETHURAMAN, J. (1994): “A Constructive Definition of Dirichlet Priors,” *Statistica Sinica*, 4, 639–650.
- THEIL, H. AND H. NEUDECKER (1958): “Substitution, Complementarity, and the Residual Variation around Engel Curves,” *The Review of Economic Studies*, 25, 114–123.
- VAN ROSEN, D. (2018): *Bilinear Regression Analysis: An Introduction*, Springer.

A The Dirichlet Process

In this appendix, we briefly review the definition and properties of the Dirichlet process, and then describe how to simulate from the Dirichlet process (see [Ferguson, 1974](#), [Rolin, 1992](#), [Sethuraman, 1994](#), [Lin, 2016](#), and [Li et al., 2019](#)).

A.1 Definition and Properties of the Dirichlet Process

For exposition, let us describe the Dirichlet distribution, then the Dirichlet process:

(i) Dirichlet Distribution:

Let $D_J(\alpha)$ denote the J -dimensional Dirichlet distribution with density:

$$f_\alpha(q) = \frac{\Gamma(\sum_{j=1}^J \alpha_j) \prod_{j=1}^J q_j^{\alpha_j}}{\prod_{j=1}^J \Gamma(\alpha_j)}, \quad (\text{A.1})$$

for every $q \in [0, 1]^J$ such that $\sum_{j=1}^J q_j = 1$, where $\alpha \in \mathbb{R}_{++}^J$ denotes a J -dimensional vector of positive parameters. If a random vector (Q_1, \dots, Q_J) has a Dirichlet distribution $D_J(\alpha)$, then:

$$\mathbb{E}[Q_j] = \bar{\alpha}_j \quad \text{and} \quad V(Q_j) = \frac{\bar{\alpha}_j(1 - \bar{\alpha}_j)}{1 + \sum_{j=1}^J \alpha_j}, \quad (\text{A.2})$$

where $\bar{\alpha}_j = \alpha_j / \sum_{j=1}^J \alpha_j$.

(ii) Dirichlet Process:

In the SARA and SSF models, there are two taste parameters, A_1 and A_2 . The probability distribution π of (A_1, A_2) is defined on \mathbb{R}_+^2 . Therefore, in this section, we describe the Dirichlet process in this special case. Let \mathcal{B}_0 denote the Borel sets associated with \mathbb{R}_+^2 , \mathcal{F} denote the set of probability measures defined on $(\mathbb{R}_+^2, \mathcal{B}_0)$, and \mathcal{B}_1 denote the σ -algebra consisting of the Borel sets associated with the topology of weak convergence on \mathcal{F} . Let μ denote a (deterministic) probability measure defined on $(\mathbb{R}_+^2, \mathcal{B}_0)$, and let c denote a strictly positive scalar. A process G with values in \mathcal{F} is a Dirichlet process with functional parameter μ and scaling parameter c if, for every finite and measurable partition $\{C_1, \dots, C_J\}$ of \mathbb{R}_+^2 , the random vector $[G(C_1), \dots, G(C_J)]'$ has a J -dimensional Dirichlet distribution given $\alpha = [c\mu(C_1), \dots, c\mu(C_J)]'$. There exists a Dirichlet process for every probability measure μ defined on $(\mathbb{R}_+^2, \mathcal{B}_0)$ and scaling parameter c . The distribution of the Dirichlet process is a probability measure defined on $(\mathcal{F}, \mathcal{B}_1)$, whose realizations are almost surely *discrete* probability measures defined on $(\mathbb{R}_+^2, \mathcal{B}_0)$, assigning probability one to the set of all discrete probability measures defined on $(\mathbb{R}_+^2, \mathcal{B}_0)$. The support of the distribution of the Dirichlet process is a set of distributions with support contained in the support of $c\mu$ (Ferguson, 1974). The functional parameter μ (sometimes called the base distribution) can be thought of as the mean of the Dirichlet process—indeed, for any measurable set C in \mathbb{R}_+^2 , the mean of the Dirichlet distribution in (A.2) yields $\mathbb{E}[G(C)] = \mu(C)$. Therefore, in our framework, μ represents the expected uncertainty on taste parameters. Intuitively, the scaling parameter c describes the “strength” of discretization: When c is large, the realizations of the Dirichlet process are concentrated around μ ; loosely, as c tends to infinity, the realizations become “more continuous.”

A.2 Simulating a Dirichlet Process

A Dirichlet process is easy to simulate given μ and c . There are a number of ways to simulate a realization—this section outlines the *stick-breaking method*, appropriate for drawing under the independence of A_1 and A_2 , based on the construction of the Dirichlet process in [Sethuraman \(1994\)](#).

Let $B(\alpha_1, \alpha_2)$ denote the beta distribution with continuous density:

$$f(q) = \frac{\Gamma(\alpha_1 + \alpha_2) q_1^{\alpha_1} q_2^{\alpha_2}}{\Gamma(\alpha_1) \Gamma(\alpha_2)}, \quad (\text{A.3})$$

on the simplex $\{(q_1, q_2) \geq 0 : q_1 + q_2 = 1\}$, in which Γ denotes the gamma function, and $\alpha_1, \alpha_2 > 0$ are positive scalar parameters. Under the independence of A_1 and A_2 , it is sufficient to be able to make a draw from a Dirichlet process whose realizations are distributions on $[0, \infty)$. Let μ^* and c^* denote the mean and scaling parameter of this Dirichlet process. We can simulate from this process by using the following steps:

Step 1: For large $L \geq 1$, independently simulate $W_1, \dots, W_L \sim B(1, c^*)$.

Step 2: Compute $W_1^* = W_1$, and:

$$W_\ell^* = W_\ell \prod_{j=1}^{\ell-1} (1 - W_j), \quad \forall \ell = 2, \dots, L. \quad (\text{A.4})$$

Step 3: Independently simulate $V_1, \dots, V_L \sim \mu^*$.

Step 4: Define:

$$G(C) = \sum_{\ell=1}^L W_\ell^* \delta_{V_\ell}(C), \quad \forall C \subseteq \mathbb{R}_+^2, \quad (\text{A.5})$$

where δ_v denotes a point mass at $v \in \mathbb{R}_+^2$.

Theoretically, if we could simulate an infinite number of draws, then this procedure would produce a realization of the Dirichlet process associated with functional parameter μ^* and scaling parameter c^* . Since L is finite, the resulting probability measure G is a *truncated* approximation of a realization of such a process. Figure 11 displays a simulated realization from the Dirichlet process given log-normal μ^* with mean 0 and standard deviation 1 (where these parameters are interpreted on the log-scale) and scaling parameter $c = 100$. This realization was simulated using the stick-breaking method given $L = 100$.

B Integrability

In this appendix, we consider the conditions that are needed for demand to be invertible in the second regime. Here, we first consider SARA preferences, and then SSF preferences. See Section 2 in [Dobronyi and Gouriéroux \(2020\)](#) for a broad overview.




Figure 11. A simulated realization from the Dirichlet process given log-normal μ^* with mean 0 and standard deviation 1 (where these parameters are interpreted on the log-scale) and scaling parameter $c = 100$. This realization was simulated using the stick-breaking method given $L = 100$.

B.1 SARA Model

Suppose A_1 and A_2 are independent. Let Ψ_j denote the Laplace transform of A_j , for $j = 1, 2$. With this notation, we can write:

$$\log U(x; \pi) = \log \Psi_1(x_1) + \log \Psi_2(x_2). \quad (\text{B.1})$$

The indifference curve $g_\pi(\cdot, u)$ associated with $U(x; \pi)$ is obtained by solving:

$$\log \Psi_1(x_1) + \log \Psi_2(x_2) = \log u, \quad (\text{B.2})$$

for x_2 . This procedure leads to:

$$g_\pi(x_1, u) = (\log \Psi_2)^{-1} (\log u - \log \Psi_1(x_1)). \quad (\text{B.3})$$

In general, demand is invertible if the indifference curves are strictly convex such that:

$$\frac{\partial^2 g_\pi(x_1, u)}{\partial x_1^2} > 0. \quad (\text{B.4})$$

When preferences are SARA, this condition becomes:

$$\begin{aligned} & \frac{d^2}{dv^2} [(\log \Psi_2)^{-1}] (\log u - \log \Psi_1(x_1)) \left(\frac{d \log \Psi_1(x_1)}{dx_1} \right)^2 \\ & - \frac{d}{dv} [(\log \Psi_2)^{-1}] (\log u - \log \Psi_1(x_1)) \frac{d^2 \log \Psi_1(x_1)}{dx_1^2} > 0, \end{aligned} \quad (\text{B.5})$$

for every $x_1 > 0$. Of course, an example of a particular specification for $U(\cdot; \pi)$ that satisfies this condition is provided in Section 2.3. Note that, this condition does not appear to be automatic because:

$$\frac{d^2}{dx_1^2} \log \Psi_1(x_1) > 0, \quad \frac{d}{dv} [(\log \Psi_2)^{-1}] (v) < 0, \quad \text{and} \quad \frac{d^2}{dv^2} [(\log \Psi_2)^{-1}] (v) < 0. \quad (\text{B.6})$$

B.2 SSF Model

Now, let us consider SSF preferences. If preferences are SSF, it is rather challenging to derive a closed-form expression for the indifference curve. We can, instead, write the integrability condition using the condition on the bordered Hessian in Lemma 1 in Dobronyi and Gouriéroux (2020), but, for both brevity and exposition, let us restrict our attention to the general specification of utility in the example in Section 3.3, and show that integrability holds for any Laplace transform. Since the strict convexity of the indifference curves is equivalent to the strict negativity of the Slutsky coefficient $\Delta_x(z)$, it is sufficient to check whether $\Delta_x(z)$ is strictly negative. We obtain:

$$\Delta_x(z) = \frac{\partial X_1(z; \pi)}{\partial p} + X_1(z; \pi) \frac{\partial X_1(z; \pi)}{\partial y} = -\frac{1}{\lambda p^3} \frac{d}{dv} \left(\frac{d \log \Psi}{dv} \right)^{-1} \left(-\frac{1}{p} \right). \quad (\text{B.7})$$

It is sufficient to show that:

$$\frac{d}{dv} \left(\frac{d \log \Psi}{dv} \right)^{-1} \left(-\frac{1}{p} \right), \quad (\text{B.8})$$

is strictly positive. To do this, consider the following derivatives:

$$\begin{aligned} \frac{d \log \Psi(v)}{dv} &= -\frac{\mathbb{E}[A_1 \exp(-A_1 v)]}{\mathbb{E}[\exp(-A_1 v)]} \\ \text{and } \frac{d^2 \log \Psi(v)}{dv^2} &= \frac{\mathbb{E}[A_1^2 \exp(-A_1 v)]}{\mathbb{E}[\exp(-A_1 v)]} - \left(\frac{\mathbb{E}[A_1 \exp(-A_1 v)]}{\mathbb{E}[\exp(-A_1 v)]} \right)^2 = V_{\tilde{\pi}}(A_1) > 0, \end{aligned} \quad (\text{B.9})$$

where the variance is with respect to the transformed density:

$$\frac{\exp(-A_1 v)}{\mathbb{E}[\exp(-A_1 v)]} \pi(v). \quad (\text{B.10})$$

Therefore, $\frac{d \log \Psi}{dv}$ is increasing, and so is its inverse $\left(\frac{d \log \Psi}{dv} \right)^{-1}$. Consequently, we can deduce that $\Delta_x(z)$ is strictly negative.

C Numerical Optimization

The optimization problem for filtering can be written as:

$$\begin{aligned} \min_{\pi_1, \pi_2} & (\pi_1 - \hat{\pi}_1)'(\pi_1 - \hat{\pi}_1) + (\pi_2 - \hat{\pi}_2)'(\pi_2 - \hat{\pi}_2) \\ \text{s.t.} & \text{ MRS restrictions (6.2), } e' \pi_1 = 1, \text{ and } e' \pi_2 = 1, \end{aligned} \quad (\text{C.1})$$

where π_1 and π_2 are written on a sufficiently large discrete grid for A_1 and A_2 , and $e = (1, \dots, 1)'$. This optimization problem can be difficult due to the dimension of the problem. The objective function is minimized with respect to the total number $2J$ of grid points in π_1 and π_2 , which is intentionally chosen to be very large (at least several hundred), and the number of constraints is N_m , where N_m denotes the number of observations with positive consumption x_{it} in segment Λ_m , which is typically around 1,000. Note, $2J$ has to be larger than N_m for identification. Therefore, it is important to find a tractable algorithm for such a problem.

We can use the fact that the MRS restrictions are bilinear in π_1 and π_2 . Indeed, these constraints can be written as:

$$A_1(\pi_2)\pi_1 = b_1(\pi_2) \text{ or } A_2(\pi_1)\pi_2 = b_2(\pi_1). \quad (\text{C.2})$$

To illustrate, consider the SARA model, and let a_{1j} and a_{2j} , $j = 1, \dots, J$, denote the locations of the points in the grids for A_1 and A_2 , respectively. Moreover, let

$\pi_1 = (\pi_{1j})$ and $\pi_2 = (\pi_{2j})$ denote the elementary probabilities on (a_{1j}) and (a_{2j}) , respectively. Under the independence of A_1 and A_2 , the MRS restrictions have the form:

$$\sum_{j=1}^J [\pi_{1j} a_{1j} \exp(-a_{1j} x_{i1t})] \sum_{j=1}^J [\pi_{2j} \exp(-a_{2j} x_{i2t})] - p_{it} \sum_{j=1}^J [\pi_{1j} \exp(-a_{1j} x_{i1t})] \sum_{j=1}^J [\pi_{2j} a_{2j} \exp(-a_{2j} x_{i2t})] = 0,$$

for every $i \in \Lambda_m$ and every t with $x_{it} > 0$. The closed-form expressions for $A_1(\pi_2)$, $b_1(\pi_2)$, $A_2(\pi_1)$, and $b_2(\pi_1)$ can be easily deduced. The unit mass restrictions can also be explicitly written as:

$$\sum_{j=1}^J \pi_{1j} = 1 \quad \text{and} \quad \sum_{j=1}^J \pi_{2j} = 1.$$

The equivalent expressions in (C.2) can be used to solve the optimization problem in (C.1) by using a succession of optimization problems with smaller dimensions (see [Gouriéroux et al., 1990](#), and [Van Rosen, 2018](#)). Precisely, let $\pi_1(k)$ and $\pi_2(k)$ denote the solutions for π_1 and π_2 at the k^{th} step of the optimization algorithm. Given $\pi_2(k)$, $\pi_1(k+1)$ is defined as the solution to:

$$\min_{\pi_1} (\pi_1 - \hat{\pi}_1)'(\pi_1 - \hat{\pi}_1) \quad \text{s.t.} \quad A_1[\pi_2(k)]\pi_1 = b_1[\pi_2(k)] \quad \text{and} \quad e'\pi_1 = 1, \quad (\text{C.3})$$

and, similarly, $\pi_2(k+1)$ is defined as the solution to:

$$\min_{\pi_2} (\pi_2 - \hat{\pi}_2)'(\pi_2 - \hat{\pi}_2) \quad \text{s.t.} \quad A_2[\pi_1(k+1)]\pi_2 = b_2[\pi_1(k+1)] \quad \text{and} \quad e'\pi_2 = 1. \quad (\text{C.4})$$

If this algorithm numerically converges, then the limit is the solution to the original optimization problem in (C.1). Moreover, $\pi_1(k)$ and $\pi_2(k)$ have closed-form solutions:

Proposition 8. The solution to (C.3) is equal to:

$$\pi_1(k+1) = \hat{\pi}_1 + A_1^*[\pi_2(k)]' \{A_1^*[\pi_2(k)]A_1^*[\pi_2(k)]'\}^{-1} \{b_1^*[\pi_2(k)] - A_1^*[\pi_2(k)]\hat{\pi}_1\}, \quad (\text{C.5})$$

where A_j^* and b_j^* encompass the MRS constraint and the unit mass constraint together.

Proof. The optimization problem in (C.1) is of the following type:

$$\min_w (w - w_0)'(w - w_0) \quad \text{s.t.} \quad Aw = b.$$

Let us introduce a Lagrange multiplier λ . The first-order conditions are, then:

$$2(w - w_0) - A'\lambda = 0 \quad \text{and} \quad Aw = b. \quad (\text{C.6})$$

The first condition can be written as:

$$w = w_0 + \frac{1}{2}A'\lambda. \quad (\text{C.7})$$

By plugging this expression for w into the second condition, we obtain:

$$\frac{\lambda}{2} = (AA')^{-1}(b - Aw_0). \quad (\text{C.8})$$

Together, (C.7) and (C.8) imply:

$$w = w_0 + A'(AA')^{-1}(b - Aw_0).$$

This expression is exactly the form of the solution in the statement of this proposition. \square

Remark 1. Instead of minimizing the ℓ_2 -distance between π_j and $\hat{\pi}_j$, we could use an information criterion, as in [Kitamura and Stutzer \(1997\)](#). However, we would no longer obtain a closed-form solution for $\pi_1(k)$ and $\pi_2(k)$, and we would have to solve a non-linear system in λ with dimension N_m .

Remark 2. The inversion of AA' is numerically feasible, but can be made more robust numerically by including a regularization. In particular, it can be replaced with the inversion of $AA' + \varepsilon I$, where $\varepsilon > 0$ is a small regularization parameter. This regularization by shrinkage (see, for example, [Ledoit and Wolf, 2004](#)) is preferable to the machine learning practice which replaces AA' with the diagonal matrix made up of the diagonal elements of AA' . In practice, it can also be easier to solve the system in (C.6), instead of using (C.5).

Remark 3. The optimization problem in (C.1) has not explicitly accounted for the positivity of π_1 and π_2 . We can incorporate positivity by adjusting after each step of the algorithm.

D The Nielsen Database

In this appendix, we provide more information about the Nielsen Homescan Consumer Panel (NHCP). First, we describe the individual records, then the representativeness of our restricted sample.

D.1 Individual Records

As mentioned in the text, all purchases are continuously recorded by each consumer. The left panel in Figure 12 displays the daily (total and alcohol-specific) expenditures of a given consumer in October of 2016. During this month, this consumer purchased 166 units of 97 distinct goods (prior to aggregation). The right panel in Figure 12 displays the daily number of units purchased by this consumer. These purchases were all made at three distinct retailers.




Figure 12. Daily Purchases. On the left, we illustrate daily expenditure for a single consumer in October of 2016. On the right, we illustrate the number of units purchased by this consumer. Light shaded bars represent all purchases and dark shaded bars represent alcohol-related purchases.

D.2 Representativeness of Sample

Let us now report the demographics of the households in our data and compare the demographics in our restricted sample to the Current Population Survey (CPS). See [Guha and Ng \(2019\)](#) and [Dobronyi and Gouriéroux \(2020\)](#) for additional summary statistics.

Table 3 gives the distribution of household size in our sample and the CPS. These distributions are similar. Our sample has a slightly smaller proportion of households with single-member and a slightly larger proportion of households with two members. This difference can be explained by one member households simply buying less alcohol.

Table 4 describes the distribution of household income in our sample and the CPS. Once again, these two distributions are quite similar, but our sample has a higher proportion of households earning between \$70,000 and \$99,999.

Tables 5 gives the distribution of the age of the eldest head of the household in our sample and the age of the householder in the CPS. There is no direct comparison between these samples, as the eldest head may differ from the householder. This aspect of the data can explain why our sample seems to be older than the general population.

Table 3. Household size in our sample and in the 2017 Annual Social and Economic Supplement (ASEC) of the CPS. CPS numbers are in thousands.

Size	Sample		CPS	
	Number	Proportion	Number	Proportion
1	5,862	0.2090	35,388	0.2812
2	12,768	0.4554	42,785	0.3400
3	4,121	0.1469	19,423	0.1543
4	3,395	0.1210	16,267	0.1292
5	1,288	0.0459	7,548	0.0599
6	422	0.0150	2,813	0.0223
7+	180	0.0064	1,596	0.0126
Total	28,036	1.0000	125,819	1.0000

Table 4. Annual household income in our sample and in the 2017 Annual Social and Economic Supplement (ASEC) of the CPS. CPS numbers are in thousands.

Income	Sample		CPS	
	Number	Proportion	Number	Proportion
Under \$5,000	265	0.0094	4,138	0.0327
\$5,000 to \$9,999	274	0.0097	3,878	0.0307
\$10,000 to \$14,999	638	0.0227	6,122	0.0485
\$15,000 to \$19,999	694	0.0247	5,838	0.0462
\$20,000 to \$24,999	1,147	0.0409	6,245	0.0494
\$25,000 to \$29,999	1,282	0.0457	5,939	0.0470
\$30,000 to \$34,999	1,480	0.0527	5,919	0.0468
\$35,000 to \$39,999	1,432	0.0510	5,727	0.0453
\$40,000 to \$44,999	1,449	0.0516	5,487	0.0434
\$45,000 to \$49,999	1,637	0.0583	5,089	0.0403
\$50,000 to \$59,999	2,878	0.1026	9,417	0.0746
\$60,000 to \$69,999	2,380	0.0848	8,213	0.0650
\$70,000 to \$99,999	6,459	0.2303	19,249	0.1524
\$100,000+	6,021	0.2147	34,963	0.2769
Total	28,036	1.0000	126,224	1.0000

Table 5. Age of eldest household head in our sample and the householder in the 2017 Annual Social and Economic Supplement (ASEC) of the CPS. CPS numbers are in thousands.

Age	Sample		CPS	
	Number	Proportion	Number	Proportion
Under 20	4	0.0001	753	0.0059
20 to 24	54	0.0019	5,608	0.0445
25 to 29	476	0.0169	9,453	0.0751
30 to 34	1,201	0.0428	10,594	0.0842
35 to 39	1,817	0.0648	10,651	0.0846
40 to 44	1,893	0.0675	10,571	0.0840
45 to 49	2,398	0.0855	11,115	0.0883
50 to 54	3,058	0.1090	12,180	0.0968
55 to 64	7,869	0.2806	23,896	0.1899
65 to 74	6,507	0.2320	17,551	0.1394
75+	2,759	0.0984	13,448	0.1068
Total	28,036	1.0000	125,819	1.0000