

**CLOSE-TO-CONVEXITY PROPERTIES OF CLAUSEN'S
HYPERGEOMETRIC FUNCTION ${}_3F_2(a, b, c; d, e; z)$**

K.CHANDRASEKRAN AND D.J. PRABHAKARAN

ABSTRACT. In this paper, we obtain various conditions on the parameters a, b, c, d and e for which the hypergeometric functions ${}_3F_2(a, b, c; d, e; z)$ to be in the class of all close-to-convex function with respect to some well known convex functions.

1. INTRODUCTION AND PRELIMINARIES

Let \mathcal{A} denote the family of functions f that are analytic in the unit disk $\mathbb{D} = \{z : |z| < 1\}$ and normalized by $f(0) = f'(0) - 1 = 0$ and let $\mathcal{S} \subset \mathcal{A}$ be the class of all univalent functions. In [1, 2], Various subclass of univalent functions are characterised by its geometrical properties and the important subclasses are convex, starlike and close-to-convex functions.

Definition 1.1. A function f in \mathcal{S} is a convex function in \mathbb{D} if and only if

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) > 0$$

Let us denote by \mathcal{C} the class of all convex univalent functions in \mathbb{D} .

Definition 1.2. A function f in \mathcal{S} is a starlike function with respect to the origin in \mathbb{D} if and only if

$$\operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) > 0$$

Let us denote by \mathcal{S}^* , the class of all starlike univalent functions in \mathbb{D} .

Definition 1.3. A function f in \mathcal{S} is a close to convex in \mathbb{D} if and only if

$$\operatorname{Re} \left(\frac{f'(z)}{g'(z)} \right) > 0$$

for some $g \in \mathcal{C}$.

The class of all close to convex univalent functions in \mathbb{D} is denoted by \mathcal{K} . It is well known that the chain of inclusion relations $\mathcal{C} \subset \mathcal{S}^* \subset \mathcal{K} \subset \mathcal{S}$.

2000 *Mathematics Subject Classification.* 30C45.

Key words and phrases. Hypergeometric Function, Starlike, Univalent, Convex, and Close-to-Convex Functions.

Final Version after Correction as on 31.10.2020.

The family of functions in \mathcal{A} which are close-to-convex with respect to $-\log(1-z)$, and also, starlike in \mathbb{D} is denoted by KS^* .

Definition 1.4. Let $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ and $g(z) = z + \sum_{n=2}^{\infty} b_n z^n$ be analytic in \mathbb{D} . Then the Hadamard product or convolution of $f(z)$ and $g(z)$ is defined by

$$f(z) * g(z) = z + \sum_{n=2}^{\infty} a_n b_n z^n, \quad |z| < 1.$$

Definition 1.5. The Alexander transform, is defined as

$$(1) \quad \Lambda_f(z) = \int_0^z \frac{f(t)}{t} dt, \quad f \in \mathcal{S}, z \in \mathbb{D}$$

Using convolution technique the above transform is given by $F(z) = f(z) * h(z)$ where $h(z) = -\log(1-z)$, is not univalent in \mathbb{D} . It is known that the function $h(z)$ is well known and is a convex function of order $1/2$.

For any complex variable $a \neq 0$, the shifted factorial (or Pochhammer symbol) is defined as

$$(a)_0 = 1, \quad (a)_n = a(a+1) \cdots (a+n-1), \quad n = 1, 2, 3, \dots$$

In terms of Euler gamma function, the Pochhammer symbol can also be defined as

$$(a)_n = \frac{\Gamma(n+a)}{\Gamma(a)}, \quad n = 0, 1, 2, \dots$$

where a is neither zero nor a negative integer.

Definition 1.6. The Hypergeometric Function ${}_3F_2(a, b, c; d, e; z)$ is defined by

$$(2) \quad {}_3F_2(a, b, c; d, e; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n (c)_n}{(d)_n (e)_n (1)_n} z^n; \quad a, b, c, d, e \in \mathbb{C}$$

where $a, b, c, d, e \in \mathbb{C}$ and $d, e \neq 0, -1, -2, -3, \dots$ which is analytic and convergent in unit disc \mathbb{D} .

Using (2), the normalized hypergeometric function be written as $z {}_3F_2(a, b, c; d, e; z)$ and defined by

$$(3) \quad f(z) = z {}_3F_2(a, b, c; d, e; z) = z + \sum_{n=2}^{\infty} A_n z^n,$$

where

$$(4) \quad A_n = \frac{(a)_{n-1} (b)_{n-1} (c)_{n-1}}{(d)_{n-1} (e)_{n-1} (1)_{n-1}}, \quad \text{for } n \geq 2.$$

and $A_1 = 1$.

Lemma 5. [7] If $A_n \geq 0$, $\{nA_n\}$ and $\{nA_n - (n+1)A_{n+1}\}$ both are non-increasing, i.e., $\{nA_n\}$ is monotone of order 2, then f defined by (3) is in \mathcal{S}^* .

Lemma 6. [8] Suppose that

$$1 \geq 2A_2 \geq \cdots \geq nA_n \geq \cdots \geq 0$$

or

$$1 \leq 2A_2 \leq \cdots \leq nA_n \leq \cdots \leq 2$$

Then $f(z)$ is defined by (3) is close-to-convex with respect to $-\log(1-z)$.

Lemma 7. [8] Suppose that f is an odd function (i.e., the value of A_{2n} in (3) is zero for each $n \geq 1$) such that

$$1 \geq 3A_3 \geq \cdots \geq (2n+1)A_{2n+1} \geq \cdots \geq 0$$

or

$$1 \leq 3A_3 \leq \cdots \leq (2n+1)A_{2n+1} \leq \cdots \leq 2$$

Then $f \in \mathcal{S}$. In fact, $f(z)$ is close-to-convex with respect to the convex function $\frac{1}{2} \log((1+z)/(1-z))$.

In 1986, Ruscheweyh and Singh [3] obtained the sufficient conditions on the parameters a, b and c for $z {}_2F_1(a, b; c; z)$ to be starlike of order $\beta < 1$. Further the year 1995, Ponnusamy and Vourinen [4] have established the univalence and convexity properties of Gaussian hypergeometric functions. Subsequently Ponnusamy derived the conditions on the parameters a, b and c for univalence and starlike properties of Alexander transform and also the close-to-convexity properties of the Gaussian hypergeometric functions [5, 6].

Inspired by the above results, in this paper, we find conditions on a, b, c, d and e such that the function $z {}_3F_2(a, b, c; d, e; z)$ to be close-to-convex with respect to the functions $-\log(1-z)$, $\frac{1}{2} \log\left(\frac{1+z}{1-z}\right)$ and is in the class KS^* . We also find similar conditions such that the Alexander transform is in the class KS^* .

2. MAIN RESULTS AND PROOFS:

Theorem 8. If $a, b, c > 0$, $de \geq 2abc$ and

$$d + e \geq \text{Max} \left\{ a + b + c, \frac{1}{2}(ab + bc + ac + 2(a + b + c) - 1 - 2abc), 2[ab + bc + ac] - 3abc \right\},$$

then $z {}_3F_2(a, b, c; d, e; z)$ is close-to-convex with respect to $-\log(1-z)$.

Proof. Consider the function $f(z)$ is defined in (3). Replace n by $n+1$ in the equation (4) and Using the Pochhammer symbol, we have

$$(9) \quad A_{n+1} = \frac{(a)_n(b)_n(c)_n}{(d)_n(e)_n(1)_n} = \frac{(a+n-1)(b+n-1)(c+n-1)}{(d+n-1)(e+n-1)n} A_n$$

and observe that $A_n > 0$ for all $n \geq 1$.

To prove f is close-to-convex with respect to $-\log(1-z)$. It is enough to show that $\{nA_n\}$ is non-increasing sequence.

From (4) and (9), we have the following after some manipulation

$$\begin{aligned} nA_n - (n+1)A_{n+1} &= nA_n - \frac{(n+1)(a+n-1)(b+n-1)(c+n-1)}{(d+n-1)(e+n-1)n}A_n \\ &= \frac{A_n U(n)}{(d+n-1)(e+n-1)n} \end{aligned}$$

where

$$\begin{aligned} U(n) &= n^2(e+n-1)(d+n-1) - (n+1)(a+n-1)(b+n-1)(c+n-1) \\ &= (e+d-c-b-a)n^3 + (de - (e+d) - (ab+bc+ac) + c+b+a+1)n^2 \\ &\quad - (abc - c - b - a + 2)n - (a-1)(b-1)(c-1) \end{aligned}$$

For every $n \geq 1$, we have $n^3 \geq 3n^2 - 3n + 1$,

$$\begin{aligned} V(n) &\geq (de + 2(e+d) - (ab+bc+ac) - 2(a+b+c) + 1)n^2 \\ &\quad + (-3(e+d) - abc + 4(c+b+a) - 2)n \\ &\quad + e + d - abc + bc + ac - 2c + ab - 2b - 2a + 1 \end{aligned}$$

Using the fact that $n^2 \geq 2n - 1$, for all $n \geq 1$.

$$\begin{aligned} W(n) &\geq (e+d + 2de - abc - 2(ab+bc+ac))n \\ &\quad - (de + e + d + abc - 2(ab+bc+ac)) \end{aligned}$$

Put $n = 1$ in above the equation, we find

$$W(1) = de - 2abc \geq 0$$

Since $de \geq 2abc$, The above equation implies that

$$U(n) \geq V(n) \geq W(n) \geq W(1) \geq 0.$$

Hence $\{nA_n\}$ is a non-increasing sequence). Hence by Lemma 6, the function $z {}_3F_2(a, b, c; d, e; z)$ is close-to-convex with respect to $-\log(1-z)$. \square

Theorem 10. *If $a, b, c > 0$, $de \geq 3abc$ and*

$$(11) \quad d + e \geq \text{Max} \{a + b + c, \alpha(a, b, c), 3(ab + bc + ac) - 7abc\}$$

where

$$\alpha(a, b, c) = \frac{1}{3} ((2(ab + bc + ac) + 3(a + b + c) - 6abc - 1))$$

then $z {}_3F_2(a, b, c; d, e; z^2)$ is close-to-convex with respect to $\frac{1}{2} \log((1+z)/(1-z))$.

Proof. Consider the function defined as follows by replacing z by z^2 in the equation (3)

$$f(z) = z {}_3F_2(a, b, c; d, e; z^2) = z + \sum_{n=2}^{\infty} A_{2n-1} z^{2n-1},$$

where

$$(12) \quad A_{2n-1} = \frac{(a)_{n-1}(b)_{n-1}(c)_{n-1}}{(d)_{n-1}(e)_{n-1}(1)_{n-1}}, \text{ for } n \geq 2.$$

and $A_1 = 1$. Then, we have the following from the equation (12) by replacing n by $n + 1$.

$$A_{2n+1} = \frac{(a+n-1)(b+n-1)(c+n-1)}{(d+n-1)(e+n-1)n} A_{2n-1}$$

and therefore we obtain

$$(2n-1)A_{2n-1} - (2n+1)A_{2n+1} = \frac{A_{2n-1}X(n)}{(e+n-1)(d+n-1)n}$$

where

$$\begin{aligned} X(n) = & 2(e+d-c-b-a)n^3 \\ & + (2de-3(e+d)-2(ab+bc+ac)+3(c+b+a)+1)n^2 \\ & - (de-(e+d)+2abc-(ab+bc+ac)+2)n \\ & - (a-1)(b-1)(c-1) \end{aligned}$$

Replace $n^3 \geq 3n^2 - 3n + 1$, for every $n \geq 1$, we have

$$\begin{aligned} Y(n) = & (2de+3(e+d)-2(ab+bc+ac)-3(c+b+a)+1)n^2 \\ & - (de+5(e+d)+2abc-(bc+ac+ab)-6(c+b+a)+2)n \\ & + 2(e+d)-abc+ab+bc+ac-3(c+b+a)+1 \end{aligned}$$

Using $n^2 \geq 2n - 1$, for all $n \geq 1$, implies that

$$\begin{aligned} Z(n) = & (3de+e+d-2abc-3(ab+bc+ac))n \\ & - (2de+e+d+abc-3(ab+bc+ac)) \end{aligned}$$

Replace n by 1 in above, we get

$$Z(1) = de - 3abc \geq 0$$

Since $de \geq 3abc$, The above equation implies that

$$X(n) \geq Y(n) \geq Z(n) \geq Z(1) \geq 0$$

Hence by the condition on $d + e$ in (11), we have $X(n)$ is non-negative, for all $n \geq 1$. Thus $\{(2n-1)A_{2n-1}\}$ is non-increasing sequence. The function $z {}_3F_2(a, b, c; d, e; z^2)$ is close-to-convex with respect to $\frac{1}{2} \log((1+z)/(1-z))$ by Lemma 7. \square

Theorem 13. Let a, b , and $c > 0$,

$$d + e \geq \text{Max} \{T_1(a, b, c), T_2(a, b, c), T_3(a, b, c), T_4(a, b, c)\}$$

where

$$\begin{aligned}
T_1(a, b, c) &= (e + d - c - b - a) (e + d - c - b - a + 1) \\
T_2(a, b, c) &= (e + d - c - b - a + 1) (2de + 5(e + d) - 2(ab + bc + ac) - 5(c + b + a) + 2) \\
T_3(a, b, c) &= ((d^2 + 9d + 9) e^2 + (9d^2 + ((-2b - 2a - 8) c + (-2a - 8) b - 8a + 29) d \\
&\quad + ((-2a - 10) b - 10a - 16) c + (-10a - 16) b - 16a + 15) e + 9d^2 \\
&\quad + (((-2a - 10) b - 10a - 16) c + (-10a - 16) b - 16a + 15) d \\
&\quad + (b^2 + (4a + 9) b + a^2 + 9a + 7) c^2 + ((4a + 9) b^2 \\
&\quad + (4a^2 + 24a + 3) b + 9a^2 + 3a - 11) c + (a^2 + 9a + 7) b^2 \\
&\quad + (9a^2 + 3a - 11) b + 7a^2 - 11a + 4) \\
T_4(a, b, c) &= ((4d^2 + 14d + 7) e^2 + (14d^2 + (((-2a - 8) b - 8a - 8) c + (-8a - 8) b \\
&\quad - 8a + 32) d + ((-10a - 16) b - 16a - 8) c + (-16a - 8) b - 8a + 11) e \\
&\quad + 7d^2 + (((-10a - 16) b - 16a - 8) c + (-16a - 8) b - 8a + 11) d \\
&\quad + ((2a + 4) b^2 + (2a^2 + 16a + 10) b + 4a^2 + 10a + 3) c^2 \\
&\quad + ((2a^2 + 16a + 10) b^2 + (16a^2 + 16a - 8) b + 10a^2 - 8a \\
&\quad - 5) c + (4a^2 + 10a + 3) b^2 + (10a^2 - 8a - 5) b \\
&\quad + 3a^2 - 5a + 2)
\end{aligned}$$

and

$$\begin{aligned}
T(a, b, c, d, e) &= (2d^2 + 2d) e^2 + (2d^2 + (2 - 8abc) d - 8abc) e - 8abcd \\
&\quad + ((3a^2 + 3a) b^2 + (3a^2 + 3a) b) c^2 + ((3a^2 + 3a) b^2 + (3a^2 - 5a) b) c \geq 0
\end{aligned}$$

Then $z {}_3F_2(a, b, c; d, e; z)$ is in KS^* .

Proof. The function $f(z) = z {}_3F_2(a, b, c; d, e; z)$ is defined by (3), where A_n is as in (4).

For $a, b, c > 0$, we observe that $de \geq 2abc$ and

$$d+e \geq \text{Max} \left\{ a + b + c, \frac{1}{2}(ab + bc + ac + 2(a + b + c) - 1 - 2abc), 2[ab + bc + ac] - 3abc \right\}.$$

By Theorem 8, this condition implies that the sequence $\{nA_n\}$ is non-increasing. To prove f is starlike. We need to show that the sequence $\{nA_n - (n + 1)A_{n+1}\}$ is also non-increasing using Lemma 5. Let

$$B_n = nA_n - (n + 1)A_{n+1} \quad \text{and} \quad B_{n+1} = (n + 1)A_{n+1} - (n + 2)A_{n+2}$$

Using A_n , we find that

$$\begin{aligned}
(14) \quad B_n - B_{n+1} &= nA_n - 2(n + 1)A_{n+1} + (n + 2)A_{n+2} \\
&= A_n \left[n - 2(n + 1) \left(\frac{A_{n+1}}{A_n} \right) + (n + 2) \left(\frac{A_{n+2}}{A_n} \right) \right]
\end{aligned}$$

Where

$$\frac{A_{n+1}}{A_n} = \frac{(a+n-1)(b+n-1)(c+n-1)}{(d+n-1)(e+n-1)n}$$

and

$$\frac{A_{n+2}}{A_n} = \frac{(a+n)(a+n-1)(b+n)(b+n-1)(c+n)(c+n-1)}{(d+n)(d+n-1)(e+n)(e+n-1)n(n+1)}$$

After some simplification the equation (14) implies

$$(15) \quad B_n - B_{n+1} = \frac{A_n P(n)}{n(n+1)(d+n)(d+n-1)(e+n)(e+n-1)}$$

$$\begin{aligned} P(n) &= n^2(n+1)(d+n)(d+n-1)(e+n)(e+n-1) \\ &\quad - 2(n+1)^2(d+n)(e+n)(a+n-1)(b+n-1)(c+n-1) \\ &\quad + (n+2)(a+n)(a+n-1)(b+n)(b+n-1)(c+n)(c+n-1) \\ &= (e+d-c-b-a)(e+d-c-b-a+1)n^5 \\ &\quad + 2(e+d-c-b-a+1)(de-bc-ac-ab+1)n^4 \\ &\quad + ((d^2+d-1)e^2 + (d^2+((-2b-2a)c-2ab+1)d \\ &\quad + ((-2a-2)b-2a+4)c + (4-2a)b+4a-3)e - d^2 \\ &\quad + (((-2a-2)b-2a+4)c + (4-2a)b+4a-3)d \\ &\quad + (b^2 + (4a+1)b+a^2+a-3)c^2 + ((4a+1)b^2 + (4a^2-9)b \\ &\quad + a^2-9a+7)c + (a^2+a-3)b^2 + (a^2-9a+7)b - 3a^2 \\ &\quad + 7a-4)n^3 \\ &\quad + ((d^2-d)e^2 + (-d^2 + (((-2a-2)b-2a+4)c + (4-2a)b+4a-3)d \\ &\quad + ((2-4a)b+2a)c + 2ab-2)e + (((2-4a)b+2a)c + 2ab-2)d \\ &\quad + ((2a+1)b^2 + (2a^2+4a-5)b+a^2-5a+2)c^2 + ((2a^2+4a-5)b^2 \\ &\quad + (4a^2-20a+11)b - 5a^2+11a-4)c + (a^2-5a+2)b^2 \\ &\quad + (-5a^2+11a-4)b + 2a^2-4a+2)n^2 \\ &\quad + (((((2-4a)b+2a)c + 2ab-2)d + ((2-2a)b+2a-2)c + (2a-2)b \\ &\quad - 2a+2)e + (((2-2a)b+2a-2)c + (2a-2)b-2a+2)d \\ &\quad + ((a^2+3a-2)b^2 + (3a^2-7a+2)b - 2a^2+2a)c^2 \\ &\quad + ((3a^2-7a+2)b^2 + (-7a^2+11a-2)b + 2a^2-2a)c \\ &\quad + (2a-2a^2)b^2 + (2a^2-2a)b)n \\ &\quad - 2(a-1)(b-1)(c-1)(de-abc) \end{aligned}$$

Our aim is to check that $P(n)$ is non-negative for all $n \geq 1$. After few steps of calculation, we get

$$\begin{aligned}
T(n) \geq & ((5d^2 + 9d + 2)e^2 + (9d^2 + (((-8a - 8)b - 8a)c - 8ab + 13)d \\
& + ((-16a - 8)b - 8a)c - 8ab + 2)e + 2d^2 + (((-16a - 8)b - 8a)c \\
& - 8ab + 2)d + ((a^2 + 7a + 3)b^2 + (7a^2 + 13a + 3)b + 3a^2 + 3a)c^2 \\
& + ((7a^2 + 13a + 3)b^2 + (13a^2 - 5a - 5)b + 3a^2 - 5a)c \\
& + (3a^2 + 3a)b^2 + (3a^2 - 5a)b)n \\
& + (-3d^2 - 7d - 2)e^2 + (-7d^2 + ((8b + 8a)c + 8ab - 11)d + ((8a + 8)b + 8a)c \\
& + 8ab - 2)e - 2d^2 + (((8a + 8)b + 8a)c + 8ab - 2)d + ((2a^2 - 4a - 3)b^2 \\
& + (-4a^2 - 10a - 3)b - 3a^2 - 3a)c^2 + ((-4a^2 - 10a - 3)b^2 \\
& + (5 - 10a^2)b - 3a^2 + 5a)c + (-3a^2 - 3a)b^2 + (5a - 3a^2)b
\end{aligned}$$

Put $n = 1$, we have

$$\begin{aligned}
T(1) \geq & (2d^2 + 2d)e^2 + (2d^2 + (2 - 8abc)d - 8abc)e - 8abcd \\
& + ((3a^2 + 3a)b^2 + (3a^2 + 3a)b)c^2 + ((3a^2 + 3a)b^2 + (3a^2 - 5a)b)c \geq 0
\end{aligned}$$

Hence by hypothesis, the following inequalities holds true

$$P(n) \geq Q(n) \geq R(n) \geq S(n) \geq T(n) \geq T(1) \geq 0.$$

Therefore sequences $\{B_n\}$ and $\{nA_n - (n+1)A_{n+1}\}$, is non-increasing. We deduce that f is starlike by Lemma 5. Also the function f is close-to-convex with respect to $-\log(1-z)$. Since the conditions of Lemma 6 are verified. \square

Theorem 16. *Let a, b , and $c > 0$. Suppose that*

$$d + e \geq \max\{T_1(a, b, c), T_2(a, b, c), T_3(a, b, c)\}$$

where

$$\begin{aligned}
T_1(a, b, c) &= (e + d - c - b - a + 1)(e + d - c - b - a + 2), \\
T_2(a, b, c) &= 2(e + d - c - b - a + 2)(de + 2e + 2d - bc - ac - c - ab - b - a + 1), \\
T_3(a, b, c) &= ((d^2 + 7d + 5)e^2 + (7d^2 + ((-2b - 2a - 4)c + (-2a - 4)b - 4a + 21)d \\
& + ((-2a - 6)b - 6a - 4)c + (-6a - 4)b - 4a + 9)e + 5d^2 \\
& + (((-2a - 6)b - 6a - 4)c + (-6a - 4)b - 4a + 9)d \\
& + (b^2 + (4a + 3)b + a^2 + 3a + 1)c^2 + ((4a + 3)b^2 \\
& + (4a^2 + 4a - 5)b + 3a^2 - 5a - 3)c + (a^2 + 3a + 1)b^2 \\
& + (3a^2 - 5a - 3)b + a^2 - 3a + 2)
\end{aligned}$$

and satisfies

$$\begin{aligned}
T(a, b, c, d, e) &= (2d^2 + 2d)e^2 + (2d^2 + (2 - 4abc)d - 4abc)e - 4abcd \\
& + ((a^2 + a)b^2 + (a^2 + a)b)c^2 + ((a^2 + a)b^2 + (a^2 - 3a)b)c \geq 0
\end{aligned}$$

then the Alexander transform is defined by (1) is in KS^* .

Proof. Let $f(z) = z {}_3F_2(a, b, c; d, e; z)$, then from the definition of hypergeometric function ${}_3F_2$, we have

$$f(z) = \sum_{n=1}^{\infty} \frac{(a)_{n-1}(b)_{n-1}(c)_{n-1}}{(d)_{n-1}(e)_{n-1}(1)_{n-1}} z^n$$

so that the corresponding Alexander transform defined by (1) takes the form

$$\Lambda_f(z) = \sum_{n=1}^{\infty} A_n z^n,$$

with $A_1 = 1$ and

$$(17) \quad A_n = \frac{(a)_{n-1}(b)_{n-1}(c)_{n-1}}{n(d)_{n-1}(e)_{n-1}(1)_{n-1}}, \text{ for } n \geq 2.$$

Using the definition of the shifted factorial notation, we have

$$A_{n+1} = \frac{(a)_n(b)_n(c)_n}{(n+1)(d)_n(e)_n(1)_n}$$

and

$$(n+1)A_{n+1} = \frac{(a+n-1)(b+n-1)(c+n-1)}{(d+n-1)(e+n-1)} A_n$$

After some simplification,

$$\begin{aligned} nA_n - (n+1)A_{n+1} &= nA_n - \frac{(a+n-1)(b+n-1)(c+n-1)}{(d+n-1)(e+n-1)} A_n \\ &= A_n \left[\frac{n(d+n-1)(e+n-1) - (a+n-1)(b+n-1)(c+n-1)}{(d+n-1)(e+n-1)} \right] \\ &= \frac{A_n U(n)}{(d+n-1)(e+n-1)} \end{aligned}$$

where

$$\begin{aligned} U(n) &= n(d+n-1)(e+n-1) - (a+n-1)(b+n-1)(c+n-1) \\ &\geq (e+d-c-b-a+1)n^2 \\ &\quad + ((d-1)e-d+(-b-a+2)c+(2-a)b+2a-2)n \\ &\quad + ((1-a)b+a-1)c+(a-1)b-a+1 \\ &= (de+e+d-bc-ac-ab)n - e-d+((1-a)b+a)c+ab \end{aligned}$$

By hypothesis, $d+e \geq ab+bc+ac-abc$ and so for every $n \geq 1$,

$$U(n) \geq U(1) = de - abc \geq 0$$

and since $de \geq abc$, we proved that the above condition satisfied. Thus the sequence $\{nA_n\}$ is decreasing. Next, we prove that $\{nA_n - (n+1)A_{n+1}\}$ is also decreasing.

Let $B_n = nA_n - (n+1)A_{n+1}$. Then after some manipulation, we get

$$\begin{aligned}
B_n - B_{n+1} &= \frac{A_n U(n)}{(d+n-1)(e+n-1)} - \frac{A_{n+1} U(n+1)}{(d+n)(e+n)} \\
&= \frac{A_n U(n)}{(d+n-1)(e+n-1)} - \frac{A_n U(n+1)}{(d+n)(e+n)(n+1)} \\
&\quad \times \left(\frac{(a+n-1)(b+n-1)(c+n-1)}{(d+n-1)(e+n-1)} \right) \\
&= \frac{A_n C(n)}{(d+n-1)(c+n-1)(d+n)(e+n)(n+1)}
\end{aligned}$$

where

$$\begin{aligned}
C(n) &= U(n)(d+n)(e+n)(n+1) - U(n+1)(a+n-1)(b+n-1)(c+n-1) \\
&= (e+d-c-b-a+1)(e+d-c-b-a+2)n^4 \\
&\quad + (e+d-c-b-a+2)(de-bc-ac+c-ab+b+a-1)n^3 \\
&\quad + 2(e+d-c-b-a+2)(de-bc-ac+c-ab+b+a-1)n^2 \\
&\quad + ((d^2-d)e^2 + (-d^2 + ((2-2ab)c + 2b + 2a - 3)d \\
&\quad \quad + ((2-2a)b + 2a - 2)c + (2a-2)b - 2a + 2)e \\
&\quad \quad + (((2-2a)b + 2a - 2)c + (2a-2)b - 2a + 2)d \\
&\quad \quad + ((2a-1)b^2 + (2a^2 - 4a + 1)b - a^2 + a)c^2 \\
&\quad \quad + ((2a^2 - 4a + 1)b^2 + (-4a^2 + 6a - 1)b + a^2 - a)c \\
&\quad \quad + (a - a^2)b^2 + (a^2 - a)b)n \\
&\quad \quad - (a-1)(b-1)(c-1)(2de - abc)
\end{aligned}$$

Our aim is to check that $C(n)$ is non-negative for all $n \geq 1$. After manipulation, we get

$$\begin{aligned}
F(n) &\geq ((3d^2 + 7d + 2)e^2 + (7d^2 + (((-2a-4)b - 4a)c - 4ab + 11)d \\
&\quad + ((-6a-4)b - 4a)c - 4ab + 2)e + 2d^2 + (((-6a-4)b - 4a)c \\
&\quad - 4ab + 2)d + ((2a+1)b^2 + (2a^2 + 4a + 1)b + a^2 + a)c^2 \\
&\quad + ((2a^2 + 4a + 1)b^2 + (4a^2 - 4a - 3)b + a^2 - 3a)c \\
&\quad + (a^2 + a)b^2 + (a^2 - 3a)b)n \\
&\quad + (-d^2 - 5d - 2)e^2 + (-5d^2 + (((4-2a)b + 4a)c + 4ab - 9)d \\
&\quad + ((2a+4)b + 4a)c + 4ab - 2)e - 2d^2 + (((2a+4)b + 4a)c \\
&\quad + 4ab - 2)d + ((a^2 - a - 1)b^2 + (-a^2 - 3a - 1)b - a^2 - a)c^2 \\
&\quad + ((-a^2 - 3a - 1)b^2 + (-3a^2 + a + 3)b - a^2 + 3a)c \\
&\quad + (-a^2 - a)b^2 + (3a - a^2)b
\end{aligned}$$

Put $n = 1$ in the above, we get

$$\begin{aligned}
F(1) &= (2d^2 + 2d)e^2 + (2d^2 + (2 - 4abc)d - 4abc)e - 4abcd \\
&\quad + ((a^2 + a)b^2 + (a^2 + a)b)c^2 + ((a^2 + a)b^2 + (a^2 - 3a)b)c \geq 0
\end{aligned}$$

For every $n \geq 1$, $C(n) \geq D(n) \geq E(n) \geq F(n) \geq F(1)$. Which is true

Therefore, the sequences $\{B_n\}$ and $\{nA_n - (n+1)A_{n+1}\}$ are non-increasing. The Alexander transform $\Lambda_f(z)$ is starlike in unit disc \mathbb{D} by Lemma 5 and already verified $\{nA_n\}$ is non-increasing. Therefore, the Alexander transform $\Lambda_f(z)$ is close-to-convex with respect to $-\log(1-z)$ using Lemma 6. Thus we have $\Lambda_f(z) \in KS^*$. Which completes the proof of Theorem. \square

REFERENCES

1. P. L. Duren, *Univalent functions*, Grundlehren der Mathematischen Wissenschaften, 259, Springer-Verlag, New York, 1983.
2. A. W. Goodman, *Univalent functions*, Vol.I and Vol.II, Tampa Florida Mariner Publishing Company, (1983).
3. St. Ruscheweyh and V. Singh, *On the order of starlikeness of hypergeometric functions*, J. Math. Anal. Appl. **113** (1986), no. 1, 1–11. MR0826655
4. S. Ponnusamy, M. Vuorinen, *Univalence and convexity properties of Gaussian hypergeometric functions*, Preprint 82, Department of Mathematics, University of Helsinki, 1995, 34 pp.
5. S. Ponnusamy, *Univalence of Alexander transform under new mapping properties*, Complex Variables Theory Appl. **30** (1996), no. 1, 55–68. MR1395231
6. S. Ponnusamy, *Close-to-convexity properties of Gaussian hypergeometric functions*, J. Comput. Appl. Math. **88** (1998), no. 2, 327–337. MR1613250
7. L. Fejér, *Trigonometrische Reihen und Potenzreihen mit mehrfach monotoner Koeffizientenfolge*, Trans. Amer. Math. Soc. **39** (1936), no. 1, 18–59.
8. S. Ozaki, *On the theory mmultivalent functions*, Sci. Rep. Tokyo Bunrika Daigaku A **2**(1935), 167–188.

K. CHANDRASEKRAN, DEPARTMENT OF MATHEMATICS, JEPPIAAR SRR ENGINEERING COLLEGE, PADUR, CHENNAI- 603 103, INDIA.

Email address: kchandru2014@gmail.com

D.J. PRABHAKARAN, DEPARTMENT OF MATHEMATICS, ANNA UNIVERSITY, MIT CAMPUS , CHENNAI- 600 005, INDIA.

Email address: asirprabha@yahoo.com