FUNCTION VALUES ARE ENOUGH FOR L₂-APPROXIMATION: PART II

DAVID KRIEG¹ AND MARIO ULLRICH^{1,2}

ABSTRACT. In the first part we have shown that, for L_2 -approximation of functions from a separable Hilbert space in the worst-case setting, linear algorithms based on function values are almost as powerful as arbitrary linear algorithms if the linear widths are square-summable. That is, they achieve the same polynomial rate of convergence. In this sequel, we prove a similar result for separable Banach spaces and other classes of functions.

Let F be a set of complex-valued functions on a set D such that, for all $x \in D$, point evaluation

$$\delta_x \colon F \to \mathbb{C}, \quad f \mapsto f(x)$$

is continuous with respect to some metric d_F on F. We consider numerical approximation of functions from such classes, using only function values, and measure the error in the space $L_2 = L_2(D, \mathcal{A}, \mu)$ of square-integrable functions with respect to an arbitrary measure μ such that F is embedded into L_2 . We are interested in the *n*-th minimal worst-case error

(1)
$$e_n(F, L_2) := \inf_{\substack{x_1, \dots, x_n \in D\\\varphi_1, \dots, \varphi_n \in L_2}} \sup_{f \in F} \left\| f - \sum_{i=1}^n f(x_i) \varphi_i \right\|_{L_2},$$

which is the worst-case error of an optimal linear algorithm that uses at most n function values. These numbers are sometimes called *sampling widths* of F. We

¹INSTITUT FÜR ANALYSIS, JOHANNES KEPLER UNIVERSITÄT LINZ, AUSTRIA

E-mail address: david.krieg@jku.at, mario.ullrich@jku.at. *Date*: March 12, 2021.

²Moscow Center for Fundamental and Applied Mathematics, Lomonosov Moscow State University, Moscow, Russia

²⁰¹⁰ Mathematics Subject Classification. 41A25, 41A45, 41A65, 60B20, 41A63.

Key words and phrases. L_2 -approximation, information-based complexity, least squares, rate of convergence, random matrices, Kadison-Singer.

want to compare the sampling widths with the *linear widths* of F, defined by

(2)
$$a_n(F, L_2) := \inf_{\substack{T: L_2 \to L_2 \\ \operatorname{rank}(T) \le n}} \sup_{f \in F} \left\| f - Tf \right\|_{L_2}.$$

This is the worst-case error of an optimal linear algorithm that uses at most n linear functionals as information. In other words, we want to compare the power of function values (also called standard information) with the power of arbitrary linear information for L_2 -approximation with linear algorithms.

The numbers e_n and a_n are well studied for many particular classes of functions. For an exposition of known results and history on these and related quantities, we refer to the books [18, 19, 20], especially [20, Chapter 26 & 29], as well as [4, 23, 25] and references therein. Note that the linear widths coincide with the Kolmogorov widths in our setting. Here, we want to relate e_n and a_n for general function classes F.

Clearly, we always have $a_n \leq e_n$. On the other hand, an example of Hinrichs, Novak and Vybiral from [8] shows that it is not possible to give any general upper bound for the sampling widths in terms of the linear widths if the latter are not square-summable. We therefore ask for such a relation in the case that the linear widths are square-summable. Let us formulate one particularly interesting open question.

Open Problem. Is it true for any class F and any measure μ as above that there exists a constant $c \in \mathbb{N}$ (possibly depending on F and μ) such that

(3)
$$e_{cn}(F, L_2) \leq \sqrt{\frac{c}{n} \sum_{k \geq n} a_k(F, L_2)^2} \quad for \ all \ n \in \mathbb{N} \ ?$$

Note that (3) is of no use if the linear widths are not square-summable, but would lead to a quite tight bound otherwise. The relation (3) is true for all examples of sufficiently studied function classes F that are known to the authors. On the other hand, general results are only known in the case that F is the unit ball of a reproducing kernel Hilbert space. Even in this case, the problem is open, but the gap is quite small already. Namely, it was shown in [11] that (3) is true with the index cn replaced by $cn \log(n + 1)$, showing that the polynomial order of approximation and sampling widths is the same if the linear widths are square-summable. This was then improved by Nagel/Schäfer/T. Ullrich in [16], who showed (3) with an additional factor of $\sqrt{\log(n+1)}$ on the right hand side. In all these papers, the constant c is independent of F and μ . The aim of this paper is to prove a similar result for general function classes F.

Before we come to our result, let us mention that, recently, another beautiful upper bound for the sampling widths of general classes F has been shown by Temlyakov in [26]. This will be discussed in Section 1.2.

Our main result reads as follows.

Theorem 1. Let (D, \mathcal{A}, μ) be a measure space and let F be a separable metric space of complex-valued functions on D that is continuously embedded into $L_2(D, \mathcal{A}, \mu)$ such that function evaluation is continuous on F. Assume that $(a_n(F, L_2)) \in \ell_p$ for some $0 . There is a universal constant <math>c \in \mathbb{N}$ and a constant $c_p > 0$, depending only on p, such that, for all $n \geq 2$, we have

$$e_{cn}(F, L_2) \leq c_p \sqrt{\log n} \left(\frac{1}{n} \sum_{k \geq n} a_k(F, L_2)^p\right)^{1/p}$$

In particular, the linear widths and the sampling widths have the same polynomial order of convergence: If we assume that $a_n(F, L_2) \leq n^{-\alpha} \log^{\beta}(n+1)$ for some $\alpha > 1/2$ and $\beta \in \mathbb{R}$, then we obtain

(4)
$$e_n(F, L_2) \lesssim n^{-\alpha} \log^{\beta + 1/2} (n+1).$$

Here and in the following, the symbol \leq means that the left hand side is bounded by a constant multiple of the right hand side for all $n \in \mathbb{N}$, whereas the symbol \approx means that this relation holds in both directions. We will also present a bound for the case $\alpha = 1/2$ and $\beta < -3/2$ which is off by an additional log-factor, see Section 4. It is still open if the result of Theorem 1 also holds for p = 2, as it does for Hilbert spaces, see [16].

Our proof of Theorem 1 is not constructive. However, if we know the operators T that achieve the infimum in (2), possibly up to a multiplicative constant, then we can provide an explicit weighted least squares estimator that achieves the stated upper bound up to a further logarithmic factor with high probability, see Theorem 8. Note that these operators are known for a huge variety of smoothness spaces, e.g., on the *d*-dimensional torus, see [4].

FUNCTION VALUES ARE ENOUGH – PART II

1. DISCUSSION AND EXAMPLES

In this section, we give some additional comments on our main result, together with a few illustrative examples and a comparison with existing results.

1.1. The condition on F. The natural condition appearing in the proof of Theorem 1 is that F is a countable subset of L_2 , see Theorem 2 for a precise statement. We then need some kind of continuity in order to extend our result to uncountable sets F. Here, we employ that F is a separable metric space with continuous function evaluation and continuous embedding in L_2 . These assumptions are satisfied, for example, if

- F is the unit ball of a separable normed space on which function evaluation at each point is a continuous functional, or
- the measure μ is finite and F is a compact subset of the space of bounded functions on D.

To see that the conditions of Theorem 1 are matched, we equip F with the metric induced by the normed space in the first case and with the supremum metric in the second case. Note that the theorem might also be applied if F is the unit ball of a *non-separable* normed space, since we might have separability with respect to a weaker norm. For instance, the unit ball F of the non-separable Sobolev space $W^s_{\infty}(0, 1)$ with smoothness $s \in \mathbb{N}$ clearly satisfies the conditions of Theorem 1 when equipped with the supremum metric.

1.2. Temlyakov's L_{∞} -bound. Let us shortly compare our result with the recent result of Temlyakov [26] (see also [3] for a related result). He proved that there are universal constants $c, C \in \mathbb{N}$ such that for any compact domain $D \subset \mathbb{R}^d$, any probability measure μ on D, and any compact subset F of the space $\mathcal{C}(D)$ of continuous functions on D, we have

(5)
$$e_{cn}(F, L_2) \leq C d_n(F, L_\infty)$$

Here, $d_n(F, L_{\infty})$ is the *n*th Kolmogorov width of F in L_{∞} .

First, we observe that the assumptions of Theorem 1 and (5) are quite different. The result (5) does not require the square-summability of the linear widths. For example, it can be applied for classes of functions with small mixed smoothness as considered in [27, 28], where Theorem 1 fails to be of use. On the other hand, only Theorem 1 can be applied for classes of unbounded functions on unbounded domains, like Hermite spaces on \mathbb{R}^d or spaces of functions with singularities. But also for classes of bounded functions on compact domains, it is not possible to say that one result yields better estimates than the other. This is illustrated by the following example, which was kindly provided to us by Erich Novak.

Example 1. Let μ be the Lebesgue measure on [0, 1] and let $(\ell_i)_{i \in \mathbb{N}}$ and $(h_i)_{i \in \mathbb{N}}$ be decreasing zero-sequences with $\sum_{i=1}^{\infty} \ell_i = 1$. Let b_i be the hat function with height one, supported on the interval $I_i = [\sum_{k=1}^{i-1} \ell_k, \sum_{k=1}^{i} \ell_k]$ of length ℓ_i . We consider

$$F = \bigg\{ \sum_{i=1}^{\infty} \lambda_i b_i \, \big| \, |\lambda_i| \le h_i \text{ for all } i \bigg\}.$$

Note that F is a compact subset of $\mathcal{C}([0,1])$ which follows from the theorem of Arzelà and Ascoli. For this example, one can compute that

$$e_n(F, L_2) = \left(\int_0^1 \left(\sum_{i>n} h_i b_i(x)\right)^2 dx\right)^{1/2} = \left(\frac{1}{3}\sum_{i>n} h_i^2 \ell_i\right)^{1/2}$$

and that the Kolmogorov widths in the uniform norm are given by

$$d_n(F, L_\infty) = h_{n+1}.$$

By choosing $\ell_i = i^{-\alpha} / \sum_{k \in \mathbb{N}} k^{-\alpha}$ with some $\alpha > 1$ and $h_i = i^{-\beta}$ with some $\beta > 0$, we obtain that the sampling widths are of order $n^{-(\beta+(\alpha-1)/2)}$ while the Kolmogorov widths are of order $n^{-\beta}$. If α is close to 1 and $\beta < 1/2$, then [26] yields an almost optimal bound while our result yields nothing. If $\alpha > 2$ and β is close to zero, then our results yields an almost optimal bound, while [26] yields almost nothing.

In contrast to this example, it is quite remarkable that Theorem 1 and (5) actually lead to the same upper bounds for many multivariate function classes of mixed smoothness as considered in [4]. We note that the optimal order of the numbers $d_n(F, L_{\infty})$ is often not known, in contrast to the linear widths $a_n(F, L_2)$.

1.3. Korobov classes. As a further example, let us consider the Korobov classes

$$E_d^r = \left\{ f \in L_1(\mathbb{T}^d) \mid |\hat{f}(\mathbf{k})| \le \prod_{j=1}^d \max\{1, |k_j|\}^{-r} \text{ for all } \mathbf{k} \in \mathbb{Z}^d \right\},\$$

for given r > 1 and $d \in \mathbb{N}$, see e.g. [4, Section 3.3]. Here, \mathbb{T}^d is the *d*-dimensional torus and $\hat{f}(\mathbf{k})$ denotes the (classical) Fourier coefficient. It is well known that the

non-increasing rearrangement $(c_n)_{n\in\mathbb{N}}$ of the sequence $(\prod_{j=1}^d \max\{1, |k_j|\}^{-r})_{\mathbf{k}\in\mathbb{Z}^d}$ satisfies $c_n \simeq n^{-r} \log^{r(d-1)}(n+1)$. This can be derived from [1, 15], see also [10] for a direct formulation. We easily obtain

$$a_n(E_d^r, L_2) \le \left(\sum_{k>n} c_k^2\right)^{1/2} \lesssim n^{-r+1/2} \log^{r(d-1)}(n+1)$$

and it follows from Lemma 3.4.5 and Theorem 4.3.5 in [4] that this bound is optimal. Moreover, E_d^r is a bounded subset of the *mixed smoothness* Sobolev space $\mathbf{W}_2^s(\mathbb{T}^d)$ for all s < r - 1/2 and therefore a compact subset of $\mathcal{C}(\mathbb{T}^d)$. Theorem 1 yields

$$e_n(E_d^r, L_2) \lesssim n^{-r+1/2} \log^{r(d-1)+1/2} (n+1).$$

To the best of our knowledge, this bound is new.

2. The result behind Theorem 1

Our main result is based on the following apparently more general theorem.

Theorem 2. Let (D, \mathcal{A}, μ) be a measure space and let F_0 be a countable set of functions in $L_2(D, \mathcal{A}, \mu)$. Assume that $(a_n(F_0, L_2)) \in \ell_p$ for some $0 . Then there is a universal constant <math>c \in \mathbb{N}$ and a constant $c_p > 0$, depending only on p, such that, for all $n \geq 2$, we have

$$e_{cn}(F_0, L_2) \leq c_p \sqrt{\log n} \left(\frac{1}{n} \sum_{k \geq n} a_k(F_0, L_2)^p\right)^{1/p}.$$

Before we prove this theorem, let us show how it implies Theorem 1.

Proof of Theorem 1. Since F is a separable metric space, it contains a countable dense subset F_0 . Now, let $x_1, \ldots, x_n \in D$ and $\varphi_1, \ldots, \varphi_n \in L_2$ be arbitrary. We obtain for every $f \in F$ and $g \in F_0$ that

$$\left\| f - \sum_{i=1}^{n} f(x_i)\varphi_i \right\|_{L_2} \le \left\| f - g \right\|_{L_2} + \left\| g - \sum_{i=1}^{n} g(x_i)\varphi_i \right\|_{L_2} + \left\| \sum_{i=1}^{n} \left(f(x_i) - g(x_i) \right)\varphi_i \right\|_{L_2} \right\|_{L_2}$$

To bound the first and the last term, first note that $U_{\delta}(f) \cap F_0 \neq \emptyset$ for every $\delta > 0$, where

$$U_{\delta}(f) := \{g \in F \colon d_F(f,g) < \delta\}.$$

The continuity of the embedding into L_2 and of function evaluation now implies that for any $\varepsilon > 0$ there is some $\delta > 0$ such that $||f-g||_{L_2} < \varepsilon$ and $|f(y_i)-g(y_i)| < \varepsilon$ for all i = 1, ..., n and all $g \in U_{\delta}(f)$. Therefore, for every $\varepsilon > 0$ and every $f \in F$, we have

$$\left\|f-\sum_{i=1}^n f(x_i)\varphi_i\right\|_{L_2} < \varepsilon + \sup_{g\in F_0} \left\|g-\sum_{i=1}^n g(x_i)\varphi_i\right\|_{L_2} + \varepsilon \sum_{i=1}^n \|\varphi_i\|_{L_2}.$$

We obtain that

$$\sup_{f \in F} \left\| f - \sum_{i=1}^{n} f(x_i) \varphi_i \right\|_{L_2} = \sup_{g \in F_0} \left\| g - \sum_{i=1}^{n} g(x_i) \varphi_i \right\|_{L_2}$$

for all $x_1, \ldots, x_n \in D$ and $\varphi_1, \ldots, \varphi_n \in L_2$. Therefore, an error bound of an algorithm on F_0 carries over to F and so, together with $a_k(F_0, L_2) \leq a_k(F, L_2)$ Theorem 2 implies Theorem 1.

We will now prove Theorem 2 by proving an error bound for a specific algorithm on F_0 . Recall that we have just proven that the same algorithm works for the class F from Theorem 1 if we choose F_0 as a countable dense subset.

3. Algorithm and Proof of Theorem 2

In this whole section, we work under the assumptions of Theorem 2. We start with the following simple observation.

Lemma 3. There is an orthonormal system $\{b_k : k \in \mathbb{N}\}$ in L_2 such that the orthogonal projection P_n onto the span $V_n = \operatorname{span}\{b_1, \ldots, b_n\}$ satisfies

(6)
$$\sup_{f \in F_0} \|f - P_n f\|_{L_2} \le 2 a_{n/4}(F_0, L_2), \qquad n \in \mathbb{N}.$$

Note that the definition of a_n in (2) makes perfect sense for $n \notin \mathbb{N}$.

Proof. Clearly it is enough to find an increasing sequence of subspaces of L_2 ,

$$U_1 \subseteq U_2 \subseteq U_3 \subseteq \dots, \qquad \dim(U_n) \le n,$$

such that the projection P_n onto U_n satisfies (6). By the definition of $a_m, m \in \mathbb{N}$, there is a subspace $W_m \subset L_2$ of dimension m and a linear operator $T_m \colon L_2 \to W_m$ such that

$$\sup_{f \in F_0} \|f - T_m f\|_{L_2} \le 2 a_m(F_0, L_2).$$

We let U_n be the space that is spanned by the union of the spaces W_{2^k} over all $k \in \mathbb{N}_0$ with $2^k \leq n/2$. Note that U_n contains a subspace W_m with $m \geq n/4$. Therefore, $P_n f$ is at least as close to f as $T_m f$ for some $m \geq n/4$, which implies (6).

In what follows $\{b_k : k \in \mathbb{N}\}$ will always be the orthonormal system from Lemma 3. Note that we will consider b_k as a function, where we take an arbitrary representer from the equivalence class in L_2 . We will denote

$$\varepsilon_n := \sup_{f \in F_0} \|f - P_n f\|_{L_2}$$

to ease the notation, keeping in mind that $\varepsilon_n \leq 2 a_{n/4}(F_0, L_2)$. We have almost sure convergence of the (abstract) Fourier series on F_0 .

Lemma 4. There is a measurable subset D_0 of D with $\mu(D \setminus D_0) = 0$ such that for all $x \in D_0$ and $f \in F_0$ we have

$$f(x) = \sum_{k \in \mathbb{N}} \hat{f}(k) b_k(x), \quad \text{where} \quad \hat{f}(k) := \langle f, b_k \rangle_{L_2}.$$

Proof. For all $f \in F_0$, we have

$$\sum_{k\in\mathbb{N}} |\hat{f}(k)|^2 k = \sum_{n\in\mathbb{N}_0} \sum_{k>n} |\hat{f}(k)|^2 \le \sum_{n\in\mathbb{N}_0} \varepsilon_n^2 < \infty.$$

The Rademacher-Menchov Theorem, see e.g. [22], now implies that the Fourier series of f converges to f almost everywhere. Since F_0 is countable, the almost everywhere convergence holds simultaneously for all $f \in F_0$.

Remark 5. Note that the Rademacher-Menchov Theorem holds under much weaker assumptions. However, this is not needed here, because we anyhow require at least the square-summability of the ε_n .

The proof of our bound on the sampling widths is based on an error bound for a specific algorithm. This is a *weighted least squares estimator* of the form

(7)
$$A_{m,n}(f) = \operatorname*{argmin}_{g \in V_n} \sum_{i=1}^m \frac{|g(x_i) - f(x_i)|^2}{\varrho(x_i)},$$

for some $x_1, \ldots, x_m \in D$ and $m \ge n$. We give an explicit formula for the weight function $\varrho: D \to \mathbb{R}$ later. The points x_1, \ldots, x_m will be obtained via a probabilistic argument. They will satisfy $\varrho(x_i) > 0$. The algorithm $A_{m,n}$ may be written as

$$A_{m,n}: F_0 \to L_2, \qquad A_{m,n}(f) := \sum_{k=1}^n (G^+ N f)_k b_k$$

where $N: F_0 \to \mathbb{R}^m$ with $N(f) := \left(\varrho(x_i)^{-1/2} f(x_i)\right)_{i \le m}$ is the weighted information mapping and $G^+ \in \mathbb{R}^{n \times m}$ is the Moore-Penrose inverse of the matrix

(8)
$$G := \left(\varrho(x_i)^{-1/2} b_k(x_i)\right)_{i \le m, k \le n} \in \mathbb{R}^{m \times n}$$

This description of $A_{m,n}$ is actually more precise since it also specifies $A_{m,n}(f)$ in the case that the argmin in (7) is not unique (which is equivalent to G not having full rank). For the state of the art on (weighted) least squares methods for the approximation of individual functions, or in a randomized setting, we refer to [2, 3] and references therein. Here, we consider such methods in the *worst-case setting*, i.e., we measure the error via

$$e(A_{m,n}, F_0, L_2) := \sup_{f \in F_0} \left\| f - A_{m,n}(f) \right\|_{L_2}.$$

Clearly, we have $e_m(F_0, L_2) \leq e(A_{m,n}, F_0, L_2)$ for every choice of x_1, \ldots, x_m and ϱ .

The proof of our upper bound uses the following simple lemma, see [11], which we prove for the reader's convenience. Note that our systematic study of the "power of random information" was initiated in [7], see also [6], and therefore some of the basic ideas behind, like a version of the following lemma, already appeared there.

Lemma 6. Assume that G has full rank, then

$$e(A_{m,n}, F_0, L_2)^2 \leq \varepsilon_n^2 + s_{\min}(G: \ell_2^n \to \ell_2^m)^{-2} \sup_{f \in F_0} \|N(f - P_n f)\|_{\ell_2^m}^2,$$

where $s_{\min}(G: \ell_2^n \to \ell_2^m)$ is the smallest singular value of the matrix G.

Proof. Since G has full rank we obtain from (7) that $A_{m,n}$ satisfies $A_{m,n}(f) = f$ for all $f \in V_n$. Using Lemma 3, we obtain for any $f \in F_0$ that

$$\begin{aligned} \|f - A_{m,n}(f)\|_{L_{2}}^{2} &= \|f - P_{n}(f)\|_{L_{2}}^{2} + \|P_{n}f - A_{m,n}(f)\|_{L_{2}}^{2} \\ &\leq \varepsilon_{n}^{2} + \|A_{m,n}(f - P_{n}f)\|_{L_{2}}^{2} = \varepsilon_{n}^{2} + \|G^{+}N(f - P_{n}f)\|_{\ell_{2}^{n}}^{2} \\ &\leq \varepsilon_{n}^{2} + \|G^{+} \colon \ell_{2}^{m} \to \ell_{2}^{n}\|^{2} \cdot \sup_{f \in F_{0}} \|N(f - P_{n}f)\|_{\ell_{2}^{m}}^{2}. \end{aligned}$$

It only remains to note that the norm of G^+ is the inverse of the smallest singular value of the matrix G.

We note that

$$\varepsilon_n^2 = (\varepsilon_n^p)^{2/p} \le \left(\frac{2}{n} \sum_{k \ge n/2} \varepsilon_k^p\right)^{2/p},$$

due to the monotonicity of (ε_n) . The main result of this paper therefore follows once we prove

$$s_{\min}(G: \ell_2^n \to \ell_2^m)^2 \gtrsim m$$

and

$$\sup_{f \in F_0} \|N(f - P_n f)\|_{\ell_2^m}^2 \lesssim n \log n \left(\frac{1}{n} \sum_{k \ge n/2} \varepsilon_k^p\right)^{2/p}$$

for an instance of $A_{m,n}$ with $n \leq m \leq Cn$. Note that the first bound implies that G has full rank. We divide the proof of this into two parts:

- (1) We show these bounds with high probability for $m \simeq n \log n$ i.i.d. random points and then,
- (2) based on the famous solution to the Kadison-Singer problem, we extract $m \simeq n$ points that fulfill the same bounds.

3.1. Random points. We first observe the result of this paper holds with high probability if we allow a logarithmic oversampling. For this, we now introduce the sampling density $\rho: D \to \mathbb{R}$, which will also specify the weights in the algorithm. For $I_{\ell} := \{n2^{\ell}+1, \ldots, n2^{\ell+1}\}$ and some monotonically decreasing sequence $(v_{\ell})_{\ell \in \mathbb{N}_0}$ with $\sum_{\ell>0} v_{\ell}^2 = 1$ (to be specified later) we define

$$\varrho(x) := \frac{1}{2} \left(\frac{1}{n} \sum_{k \le n} |b_k(x)|^2 + \sum_{\ell \ge 0} \frac{v_\ell^2}{|I_\ell|} \sum_{k \in I_\ell} |b_k(x)|^2 \right).$$

Observe that ρ is indeed a μ -density. Under the assumption $(a_n) \in \ell_p$ with p < 2 that is considered in Theorem 1, we may just use $v_{\ell} \simeq 2^{-\delta \ell}$ for some $0 < \delta < 1/p - 1/2$. If we want to get closer to the condition $(a_n) \in \ell_2$, we need to consider sequences (v_{ℓ}) with polynomial decay, see Section 4.

Remark 7. The form of the density ρ is very much inspired by the density invented in [11], which was already applied in [9, 16, 29]. The density in these papers was needed to prove the result for Hilbert spaces in greatest generality, i.e., for all sequences $(a_n) \in \ell_2$. The density used here is different. It is not clear to us whether one can use the density from [11] for arbitrary classes F and prove a result like Theorem 1 for all $(a_n) \in \ell_2$. Presently, we do not know what happens, e.g., in the case that $a_n \simeq n^{-1/2} \log^{-3/2} (n+1)$, see Section 4. As the result of this part of the proof might be of independent interest, we formulate it as a theorem.

Theorem 8. Let (D, \mathcal{A}, μ) be a measure space and let F_0 be a countable set of functions in $L_2(D, \mathcal{A}, \mu)$. Assume that $(a_n(F_0, L_2)) \in \ell_p$ for some 0 . $Then there is a universal constant <math>C_1 > 0$ and a constant $c_p > 0$, depending only on p, such that for all $n \geq 2$ the algorithm $A_{m,n}$ from (7) with $m = \lceil C_1 n \log n \rceil$ and i.i.d. random variables x_1, \ldots, x_m with μ -density ϱ , satisfies

$$e(A_{m,n}, F_0, L_2) \leq c_p \left(\frac{1}{n} \sum_{k \geq n/8} a_k(F, L_2)^p\right)^{1/p}$$

with probability at least $1 - \frac{5}{n^2}$.

The proof of this result follows a similar reasoning as the original proof in [11], with the improvements from [29] that show the result from [11] with high probability. The crucial difference is that we show an upper bound on the "norm" of the information mapping N on the set

$$F_1 := \left\{ \sum_{k>n} c_k b_k \, \Big| \, \sum_{k>m} c_k^2 \le \varepsilon_m^2 \text{ for all } m \ge n \right\},$$

related to full approximation spaces, instead of the smaller set

$$F_2 := \left\{ \sum_{k>n} c_k b_k \, \Big| \, \sum_{k>n} \left(\frac{c_k}{\varepsilon_k} \right)^2 \le 1 \right\},\,$$

related to Hilbert spaces and considered in [11, 29]. We do this with the help of a dyadic decomposition of the index set $\{k \in \mathbb{N} \mid k > n\}$, together with suitable bounds on the norms of the corresponding random matrices. For this, we use again the matrix concentration result from [21], see also [14].

Proposition 9 ([21, Lemma 1]). Let X be a random vector in \mathbb{C}^k with $||X||_2 \leq R$ with probability 1, and let X_1, X_2, \ldots be independent copies of X. Additionally, let $E := \mathbb{E}(XX^*)$ satisfy $||E|| \leq 1$, where ||E|| denotes the spectral norm of E. Then,

$$\mathbb{P}\left(\left\|\sum_{i=1}^{m} X_i X_i^* - mE\right\| \ge m \cdot t\right) \le 4m^2 \exp\left(-\frac{m}{16R^2}s_t\right),$$

where $s_t = t^2$ for $t \le 2$, and $s_t = 4(t-1)$ for t > 2.

Note that [21, Lemma 1] wrongly states $s_t = \min\{t^2, 4t - 4\}$, which can easily be corrected by looking into the proof. Let us now prove the norm bounds that we need. Namely, we prove the existence of constants C_1 and $C_2 = C_2(p)$ such that the following holds for $m = \lceil C_1 n \log n \rceil$ and i.i.d. points x_1, \ldots, x_m with density ρ .

Fact 1:
$$\mathbb{P}\left[s_{\min}(G:\ell_{2}^{n} \to \ell_{2}^{m})^{2} < \frac{m}{2}\right] \leq \frac{4}{n^{2}}.$$

Fact 2: $\mathbb{P}\left[\sup_{f \in F_{0}} \|N(f - P_{n}f)\|_{\ell_{2}^{m}}^{2} > C_{2}n\log n\left(\frac{1}{n}\sum_{k \geq n/2}\varepsilon_{k}^{p}\right)^{2/p}\right] \leq \frac{1}{n^{2}}$

Together with Lemma 6 and $\varepsilon_k \leq 2a_{n/4}$ these bounds clearly imply Theorem 8.

Proof of Fact 1. Let $X_i := \varrho(x_i)^{-1/2} (b_1(x_i), \dots, b_n(x_i))^\top$. Then we have $\sum_{i=1}^m X_i X_i^* = G^*G$ with G from (8). First observe

(9)
$$||X_i||_2^2 = \varrho(x_i)^{-1} \sum_{k \le n} b_k(x_i)^2 \le 2n =: R^2.$$

Since $E = \mathbb{E}(XX^*) = \text{diag}(1, ..., 1)$ we have ||E|| = 1. Therefore, Proposition 9 with $t = \frac{1}{2}$ and $m = \lceil C_1 n \log n \rceil$ implies

(10)
$$\mathbb{P}\left(\|G^*G - mE\| \ge \frac{m}{2}\right) \le \frac{4}{n^2}$$

if the constant $C_1 > 0$ is large enough. We obtain

$$s_{\min}(G: \ell_2^n \to \ell_2^m)^2 = s_{\min}(G^*G) \ge s_{\min}(mE) - \|G^*G - mE\| \ge m/2$$

with probability at least $1 - \frac{4}{n^2}$.

Proof of Fact 2. Note that we almost surely have for all i = 1, ..., m that $\varrho(x_i)$ is positive and finite and x_i is contained in the set D_0 from Lemma 4 such that we have $f(x_i) = \sum_{k=1}^{\infty} \hat{f}(k)b_k(x_i)$ for every $f \in F_0$. In this certain event, each entry of $N(f - P_n f) \in \mathbb{R}^m$ can be written as

$$\varrho(x_i)^{-1/2} \left(f - P_n f \right)(x_i) = \sum_{k>n} \hat{f}(k) \, \varrho(x_i)^{-1/2} \, b_k(x_i).$$

If we now define $I_{\ell} := \{n2^{\ell} + 1, \dots, n2^{\ell+1}\}$ and the random matrices

$$\Gamma_{\ell} := \left(\varrho(x_i)^{-1/2} b_k(x_i) \right)_{i \le m, k \in I_{\ell}} \in \mathbb{R}^{m \times n2^{\ell}},$$

and set $\hat{f}_{\ell} := (\hat{f}(k))_{k \in I_{\ell}}$, we obtain that

$$\begin{aligned} \|N(f - P_n f)\|_{\ell_2^m} &= \left\|\sum_{\ell=0}^{\infty} \Gamma_\ell \hat{f}_\ell\right\|_{\ell_2^m} \\ &\leq \sum_{\ell=0}^{\infty} \|\Gamma_\ell \colon \ell_2(I_\ell) \to \ell_2^m\| \, \|\hat{f}_\ell\|_{\ell_2(I_\ell)} \leq \sum_{\ell=0}^{\infty} \|\Gamma_\ell \colon \ell_2(I_\ell) \to \ell_2^m\| \, \varepsilon_{n2^\ell}. \end{aligned}$$

It remains to bound the norms of Γ_{ℓ} with high probability, simultaneously for all ℓ . Let us start with an individual ℓ , and consider $X_i := \varrho(x_i)^{-1/2} (b_k(x_i))_{k \in I_{\ell}}^{\top}$ with x_i distributed according to ϱ . Hence, $\sum_{i=1}^m X_i X_i^* = \Gamma_{\ell}^* \Gamma_{\ell}$. We see that

$$||X_i||_2^2 = \varrho(x_i)^{-1} \sum_{k \in I_\ell} |b_k(x_i)|^2 \le \frac{n 2^{\ell+1} \sum_{k \in I_\ell} |b_k(x_i)|^2}{v_\ell^2 \sum_{k \in I_\ell} |b_k(x_i)|^2} \le n 2^{\ell+1} v_\ell^{-2} =: R^2,$$

where we just use the definition of ρ . Moreover, $E = \mathbb{E}(XX^*) = \text{diag}(1, \ldots, 1)$ and so ||E|| = 1. Therefore, Proposition 9 with $m = \lceil C_1 n \log n \rceil$ as above and

$$t = 1 + C_3 \frac{2^{\ell} \log((\ell+1)n)}{v_{\ell}^2 \log n} \ge 2,$$

together with

$$\|\Gamma_{\ell}\|^{2} := \|\Gamma_{\ell} \colon \ell_{2}(I_{\ell}) \to \ell_{2}^{m}\|^{2} = \|\Gamma_{\ell}^{*}\Gamma_{\ell}\| \le m + \|\Gamma_{\ell}^{*}\Gamma_{\ell} - mE\|,$$

implies

$$\mathbb{P}\left(\|\Gamma_{\ell}\|^{2} \geq \frac{C_{4} n 2^{\ell} \log((\ell+1)n)}{v_{\ell}^{2}}\right) \leq \frac{4}{n^{2}(\ell+1)^{2} \pi^{2}}$$

for some constants $C_3, C_4 > 0$ (depending only on C_1). Note that these probabilities are summable, and so a union bound shows

$$\mathbb{P}\left(\exists \ell \in \mathbb{N}_0 \colon \|\Gamma_\ell\|^2 \ge \frac{C_4 n 2^\ell \log((\ell+1)n)}{v_\ell^2}\right) \le \frac{1}{n^2}$$

We therefore obtain, with probability at least $1 - \frac{1}{n^2}$ that

(11)
$$\begin{aligned} \|N(f - P_n f)\|_{\ell_2^m} &\leq C_4 \sum_{\ell=0}^{\infty} \sqrt{n 2^\ell \log\left((\ell+1)n\right)} \cdot \frac{\varepsilon_{n2^\ell}}{v_\ell} \\ &\leq C_5 \sqrt{\log n} \sum_{\ell=0}^{\infty} \sqrt{n 2^\ell \log\left(\ell+1\right)} \cdot \frac{\varepsilon_{n2^\ell}}{v_\ell}. \end{aligned}$$

Clearly, the monotonicity of (ε_n) gives

$$\sum_{k \ge n/2} \varepsilon_k^p \ge n(2^\ell - 1/2)\varepsilon_{n2^\ell}^p$$

and thus

$$\varepsilon_{n2^{\ell}} \leq C_6 2^{-\ell/p} \left(\frac{1}{n} \sum_{k \geq n/2} \varepsilon_k^p\right)^{1/p}$$

for all $\ell \geq 0$. Inserting this in (11) and using $v_{\ell} := c_{\delta} 2^{-\delta \ell}$ for some $0 < \delta < \delta$ 1/p - 1/2, with $c_{\delta} > 0$ such that $\sum_{\ell \ge 0} v_{\ell}^2 = 1$, yields

$$\|N(f - P_n f)\|_{\ell_2^m} \leq C_7 \sqrt{n \log n} \left(\frac{1}{n} \sum_{k \geq n/2} \varepsilon_k^p\right)^{1/p} \sum_{\ell=0}^{\infty} \sqrt{\log(\ell+1)} 2^{-(1/p-1/2-\delta)\ell}.$$

early, the latter series is finite.

Clearly, the latter series is finite.

3.2. Kadison-Singer to reduce the number of points. We now employ the powerful solution to the Kadison-Singer problem due to Marcus, Spielman and Srivastava [13] to show that we can reduce the number of points in our algorithm to $m \simeq n$, without losing the error bound. In detail, we need an equivalent version of the KS problem due to [30], which was already brought into a form that is very useful for us in [16]. Note that the authors of [16] seem to be the first to use this approach in the context of sampling widths. By this, they improved upon [11] and proved the result of Theorem 1 for F being the unit ball of a separable reproducing kernel Hilbert space. For applications of KS to the discretization of the L_2 -norm, which was used to prove the result in [26], see e.g. [12, 24]. Here, we use a special case of [16, Theorem 2.3], see also [12, Lemma 2.2] and [17, Lemma 2].

Proposition 10 ([16, Theorem 2.3]). There exist constants $c_1, c_2, c_3 > 0$ such that, for all $u_1, \ldots, u_m \in \mathbb{C}^n$ such that $||u_i||_2^2 \leq \frac{2n}{m}$ for all $i = 1, \ldots, m$ and

$$\frac{1}{2} \|w\|_2^2 \le \sum_{k=1}^m |\langle w, u_i \rangle|^2 \le \frac{3}{2} \|w\|_2^2, \qquad w \in \mathbb{C}^n,$$

there is a $J \subset \{1, \ldots, m\}$ with $\#J \leq c_1 n$ and

$$c_2 \frac{n}{m} \|w\|_2^2 \le \sum_{k \in J} |\langle w, u_i \rangle|^2 \le c_3 \frac{n}{m} \|w\|_2^2, \qquad w \in \mathbb{C}^n.$$

We now let $u_i = \frac{1}{\sqrt{m}} \left(\varrho(x_i)^{-1/2} b_j(x_i) \right)_{j \le n} \in \mathbb{C}^n$ with random x_1, \ldots, x_m as in the previous section. We clearly have

$$\sum_{k=1}^{n} |\langle w, u_i \rangle|^2 = \frac{1}{m} ||Gw||_2^2, \qquad w \in \mathbb{C}^n.$$

Moreover, from (9), (10) and (11) we see that u_1, \ldots, u_m satisfy with high probability the conditions of the proposition and

$$\sup_{f \in F_0} \|N(f - P_n f)\|_{\ell_2^m} \le c_4 \sqrt{n \log n} \left(\frac{1}{n} \sum_{k \ge n/8} a_k(F_0, L_2)^p\right)^{1/p}$$

for some constant $c_4 > 0$, depending only on p. The proposition yields $J \subset \{1, \ldots, m\}$ with $\#J \leq c_1 n$ such that the matrix

$$G_J := \left(\varrho(x_i)^{-1/2} b_k(x_i)\right)_{i \in J, k \le n}$$

satisfies

$$c_2 \|w\|_2^2 \le \frac{1}{n} \|G_J w\|_2^2 \le c_3 \|w\|_2^2$$

and hence

$$s_{\min}(G_J \colon \ell_2^n \to \ell_2(J))^2 \ge c_2 n.$$

Moreover, we clearly have

$$\sup_{f \in F_0} \|N_J(f - P_n f)\|_{\ell_2(J)} \le c_4 \sqrt{n \log n} \left(\frac{1}{n} \sum_{k \ge n/8} a_k(F_0, L_2)^p\right)^{1/p}$$

for every $J \subset \{1, \ldots, m\}$, where $N_J(f) := (\varrho(x_i)^{-1/2} f(x_i))_{i \in J}$. Using Lemma 6, we see that the algorithm

$$A_{J,n}(f) := \sum_{k=1}^{n} (G_J^+ N_J f)_k b_k = \underset{g \in V_n}{\operatorname{argmin}} \sum_{i \in J} \frac{|g(x_i) - f(x_i)|^2}{\varrho(x_i)}$$

satisfies

$$e(A_{J,n}, F_0, L_2)^2 \leq \varepsilon_n^2 + s_{\min}(G_J \colon \ell_2^n \to \ell_2(J))^{-2} \cdot \sup_{f \in F_0} \|N_J(f - P_n f)\|_{\ell_2(J)}^2$$

$$\leq c_5 \log n \left(\frac{1}{n} \sum_{k \geq n/8} a_k(F_0, L_2)^p\right)^{2/p}$$

for some $c_5 > 0$ that only depends on p. Clearly, the same upper bound holds for $e_{c_1n}(F_0, L_2)^2$ which proves Theorem 2, and thereby Theorem 1.

4. The limiting case p = 2

With our techniques, we were not able to prove a result for arbitrary spaces with $(a_n) \in \ell_2$, and it is not clear if this is possible, even if we allow weaker bounds. However, the condition $(a_n) \in \ell_p$ with p < 2 may be weakened to $((\log n)^q a_n) \in \ell_2$ for any q > 1. A second look at our proof quickly reveals that we actually showed the upper bound

(12)
$$e_{cn}(F, L_2) \leq C \sqrt{\frac{\log n}{n}} \sum_{\ell=0}^{\infty} \sqrt{n2^{\ell} \log (\ell+1)} \cdot \frac{a_{n2^{\ell}}(F, L_2)}{v_{\ell}}$$

for any sequence (v_{ℓ}) with $\sum_{\ell \geq 0} v_{\ell}^2 = 1$, all $n \geq 2$, and with universal constants c and C. We simply skip the additional estimates after equation (11). Theorem 1 is obtained from this estimate if we choose v_{ℓ} with exponential decay. The following is obtained if we choose v_{ℓ} with polynomial decay.

Theorem 11. Let (D, \mathcal{A}, μ) be a measure space and let F be a separable metric space of complex-valued functions on D that is continuously embedded into $L_2(D, \mathcal{A}, \mu)$ such that function evaluation is continuous on F. Then there are universal constants $c, C \in \mathbb{N}$ such that, for all $n \geq 3$,

$$e_{cn}(F, L_2) \le C \cdot \left(\frac{\log n}{n} \sum_{k \ge n} (\log k)^2 (\log \log k)^5 a_k(F, L_2)^2\right)^{1/2}$$

If we apply this estimate to a sequence satisfying $a_n(F, L_2) \leq n^{-1/2} \log^{\beta}(n+1)$ for some $\beta < -3/2$, we obtain for any $\varepsilon > 0$ that

$$e_n(F, L_2) \lesssim n^{-1/2} \log^{\beta+2+\varepsilon}(n+1).$$

Proof. We apply Cauchy Schwarz to (12), choosing the square summable sequence $(\ell^{1/2} \log(\ell + 1))^{-1}$ as the first factor, and obtain

$$e_{2cn}(F, L_2)^2 \leq \tilde{C} \frac{\log n}{n} \sum_{\ell=0}^{\infty} n 2^{\ell} \cdot \ell \, \log^3 \left(\ell + 1\right) \cdot \frac{a_{n2^{\ell+1}}(F, L_2)^2}{v_{\ell}^2}$$
$$\leq \tilde{C} \frac{\log n}{n} \sum_{\ell=0}^{\infty} \frac{\ell \log^3 \left(\ell + 1\right)}{v_{\ell}^2} \sum_{k \in I_{\ell}} a_k(F, L_2)^2.$$

Recall $I_{\ell} = \{n2^{\ell} + 1, \dots, n2^{\ell+1}\}$. The statement is obtained by setting $v_{\ell}^{-2} = \tilde{c}\ell \log^2(\ell+1)$, noting that $\ell \leq \log(k)$ for $k \in I_{\ell}$.

Acknowledgement. We thank Aicke Hinrichs, Erich Novak, Winfried Sickel, Vladimir Temlyakov and Tino Ullrich for various helpful comments. In particular, we thank Erich Novak for providing us with Example 1, and Tino Ullrich for pointing out the present (more elegant) formulation of the upper bound in Theorem 1 and for coming up with the example of Korobov classes. David Krieg is supported by the Austrian Science Fund (FWF) Project F5513-N26, which is a part of the Special Research Program *Quasi-Monte Carlo Methods: Theory and Applications*.

References

- K. I. Babenko. About the approximation of periodic functions of many variable trigonometric polynomials. *Dokl. Akad. Nauk SSR* 32, 247–250, 1960.
- [2] A. Cohen, M. A. Davenport, D. Leviatan. On the Stability and Accuracy of Least Squares Approximations. *Found. Comput. Math.* 13, 819–834, 2013.
- [3] A. Cohen, G. Migliorati. Optimal weighted least-squares methods. SMAI-Journal of Computational Mathematics, 3:181–203, 2017.
- [4] D. Dũng, V.N. Temlyakov, and T. Ullrich. Hyperbolic Cross Approximation. Advanced Courses in Mathematics - CRM Barcelona. Springer International Publishing, 2018.
- [5] D.E. Edmunds, J. Lang. Gelfand numbers and width. J. Approx. Theory, 166:78-84, 2013.
- [6] A. Hinrichs, D. Krieg, E. Novak, J. Prochno, and M. Ullrich. On the power of random information. In F. J. Hickernell and P. Kritzer, editors, *Multivariate Algorithms and Information-Based Complexity*, pages 43–64. De Gruyter, Berlin/Boston, 2020.
- [7] A. Hinrichs, D. Krieg, E. Novak, J. Prochno, and M. Ullrich. Random sections of ellipsoids and the power of random information. arXiv:1901.06639, 2019.
- [8] A. Hinrichs, E. Novak, and J. Vybíral. Linear information versus function evaluations for *l*₂-approximation. J. Approx. Theory, 153(1):97–107, 2008.
- [9] L. Kämmerer, T. Ullrich, T. Volkmer. Worst case recovery guarantees for least squares approximation using random samples. arXiv:1911.10111, 2019.
- [10] D. Krieg. Tensor power sequences and the approximation of tensor product operators. J. Complexity, 44:30–51, 2018.
- [11] D. Krieg, and M. Ullrich. Function values are enough for L₂-approximation. Found. Comput. Math., 2020. https://doi.org/10.1007/s10208-020-09481-w
- [12] I. Limonova and V. Temlyakov. On sampling discretization in L₂. arXiv:2009.10789v1, 2020.
- [13] A. W. Marcus, D. A. Spielman, and N. Srivastava. Interlacing families II: Mixed characteristic polynomials and the Kadison-Singer problem. Ann. of Math. (2), 182(1):327–350, 2015.

- [14] S. Mendelson and A. Pajor. On singular values of matrices with independent rows. *Bernoulli*, 12(5):761–773, 2006.
- [15] B. S. Mityagin. Approximation of functions in L^p and C on the torus. *Math. Notes* 58, 397–414, 1962.
- [16] N. Nagel, M. Schäfer, T. Ullrich. A new upper bound for sampling numbers. to appear in Found. Comput. Math., 2020. arXiv:2010.00327
- [17] S. Nitzan, A. Olevskii, A. Ulanovskii. Exponential frames on unbounded sets. Proc. Amer. Math. Soc., 144(1):109–118, 2016.
- [18] E. Novak and H. Woźniakowski. Tractability of multivariate problems. Vol. 1: Linear information, volume 6 of EMS Tracts in Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [19] E. Novak and H. Woźniakowski. Tractability of multivariate problems. Volume II: Standard information for functionals, volume 12 of EMS Tracts in Mathematics. European Mathematical Society (EMS), Zürich, 2010.
- [20] E. Novak and H. Woźniakowski. Tractability of multivariate problems. Volume III: Standard information for operators, volume 18 of EMS Tracts in Mathematics. European Mathematical Society (EMS), Zürich, 2012.
- [21] R. I. Oliveira. Sums of random Hermitian matrices and an inequality by Rudelson. *Electr. Comm. Probab.*, 15:203–212, 2010.
- [22] R. Salem. A new proof of a theorem of Menchoff. Duke Math. J., 8(2):269–272, 1941.
- [23] V. N. Temlyakov. Approximation of periodic functions, Computational Mathematics and Analysis Series, Nova Science Publishers, Inc., Commack, NY, 1993.
- [24] V. N. Temlyakov. The Marcinkiewicz-type discretization theorems for the hyperbolic cross polynomials. Jaen Journal on Approximation, 9(1):37–63, 2017.
- [25] V. N. Temlyakov. Multivariate Approximation, volume 32 of Cambridge Monographs on Applied and Computational Mathematics. Cambridge University Press, 2018.
- [26] V. Temlyakov. On optimal recovery in L_2 . J. Complexity, 2020. https://doi.org/10.1016/j.jco.2020.101545
- [27] V. Temlyakov, T. Ullrich. Approximation of functions with small mixed smoothness in the uniform norm. arXiv:2012.11983, 2021.
- [28] V. Temlyakov, T. Ullrich. Bounds on Kolmogorov widths and sampling recovery for classes with small mixed smoothness. arXiv:2012.09925, 2021.
- [29] M. Ullrich. On the worst-case error of least squares algorithms for L_2 -approximation with high probability. J. Complexity 60, 2020.
- [30] N. Weaver. The Kadison-Singer problem in discrepancy theory. Discrete Mathematics, 278(1-3):227-239, 2004.