Maximum Matchings and Popularity

Telikepalli Kavitha

Tata Institute of Fundamental Research, Mumbai, India kavitha@tifr.res.in

Abstract. Let $G = (A \cup B, E)$ be a bipartite graph where every node has a strict ranking of its neighbors. For every node, its preferences over neighbors extend naturally to preferences over matchings. Matching N is "more popular" than matching M if the number of nodes that prefer N to M is more than the number that prefer M to N. A maximum matching M in G is a *popular max-matching* if there is no maximum matching in G that is more popular than M. Such matchings are relevant in applications where the set of admissible solutions is the set of maximum matchings and we wish to find a *best* maximum matching as per node preferences.

It is known that a popular max-matching always exists in G. Here we show a compact extended formulation for the popular max-matching polytope. So when there are edge costs, a min-cost popular max-matching can be computed in polynomial time. This is in contrast to the min-cost popular matching problem which is known to be NP-hard. We also consider *Pareto-optimality*, which is a relaxation of popularity, and show that computing a min-cost Pareto-optimal matching/max-matching is NP-hard.

1 Introduction

Consider a bipartite graph $G = (A \cup B, E)$ on n nodes and m edges where every node ranks its neighbors in a strict order of preference. The goal is to find an optimal matching in G and the typical notion of optimality in such an instance is *stability*. A matching M is stable if no edge blocks M; an edge (a, b) blocks M if a and b prefer each other to their respective assignments in M. Gale and Shapley [14] in 1962 showed that stable matchings always exist in G and gave a linear time algorithm to find one.

Stable matchings and their variants are used in many real-world applications such as matching students to schools and colleges [1,4]. Observe that stability empowers every edge with the power to block a matching – thus this is a rather strict notion. Popularity is a natural relaxation of stability that captures collective decision. Preferences of a node over its neighbors extend naturally to preferences over matchings as follows. We say node u prefers matching M to matching N if (i) u is matched in M and unmatched in N or (ii) u is matched in both M and N and it prefers its partner in M to its partner in N.

Let $\phi(M, N)$ be the number of nodes that prefer M to N and similarly, let $\phi(N, M)$ be the number of nodes that prefer N to M. We say N is more popular than M if $\phi(N, M) > \phi(M, N)$.

Definition 1. A matching M is popular if there is no matching in G that is more popular than M, i.e., $\Delta(M, N) \ge 0$ for all matchings N in G, where $\Delta(M, N) = \phi(M, N) - \phi(N, M)$.

The notion of popularity was introduced by Gärdenfors [16] in 1975 and he observed that every stable matching is popular. One of the main merits of popularity is that it allows larger matchings compared to stable matchings. All stable matchings have the same size [15] which may be as low as $|M_{\text{max}}|/2$ (where M_{max} is a maximum matching in G) and it is known that there is always a popular matching of size at least $2|M_{\text{max}}|/3$ and there are simple instances with no larger popular matching [20].

Maximum Matchings. In the domain of *matchings under preferences*, there are applications where the size of the matching computed is of utmost importance, say, in assigning doctors to hospitals during a pandemic. For instance, during the covid-19 pandemic in Mumbai, public hospitals were overwhelmed with rising number of patients and were severely short-staffed; so doctors associated with private clinics were asked to also work in public hospitals [3,19]. This problem can be modeled as described below.¹

¹ In the doctors-hospitals setting, one typically seeks a many-to-one matching since a hospital may need several doctors. Here we focus on one-to-one matchings.

Let A be the set of doctors and B be the set of hospitals: edge (a, b) denotes that doctor a and hospital b are interested in each other. Every doctor has a strict ranking of possible hospitals and similarly, every hospital has a strict ranking of possible doctors. We want the maximum number of doctors to get assigned to hospitals, i.e., we do not wish to compromise at all on the size of our matching. So the set of admissible solutions is the set of maximum matchings and among all maximum matchings, we wish to find a *best* matching as per node preferences. In the context of popularity, this means no maximum matching is "more popular" than the maximum matching we find. So what we seek is a *popular max-matching*, which is defined below.

Definition 2. Call a maximum matching M in $G = (A \cup B, E)$ a popular max-matching if $\Delta(M, N) \ge 0$ for all maximum matchings N in G.

Note that a popular max-matching M need not be popular in the set of *all* matchings—so infeasible matchings (i.e., those that are not maximum) may be more popular than M, however M is popular *within* the set of maximum matchings in G. The relation "more popular than" is not transitive and since stable matchings need not be maximum matchings, it is not clear if every instance admits a popular max-matching. This question was considered in [20] where it was shown that popular max-matchings always exist in $G = (A \cup B, E)$ and one such matching can be computed in O(mn) time.

Suppose there is a cost function $c: E \to \mathbb{R}$ where c(e) is the cost of including the edge e in our matching; for any matching M, $c(M) = \sum_{e \in M} c(e)$. We consider the problem of computing a min-cost popular maxmatching in G. Solving the min-cost popular max-matching problem efficiently implies efficient algorithms for a whole host of popular max-matching problems such as finding one with forced/forbidden edges or one with max-utility or one with min-regret. In general, a cost function allows us to "access" the entire set of popular max-matchings; note that G may have more than 2^n such matchings [27].

While it is easy to find a min-size/max-size popular matching in G [14,17,20], it is NP-hard to compute a min-cost popular matching [10]. On the other hand, there are several polynomial time algorithms to compute a min-cost stable matching and some variants of this problem [11,12,13,18,25,26,28]. Moreover, the stable matching polytope of G has an elegant linear-size description in \mathbb{R}^m [25].

Let \mathcal{M}_G denote the popular max-matching polytope of G. That is, the extreme points of the polytope \mathcal{M}_G are all and only the edge incidence vectors of popular max-matchings in G. A compact description of \mathcal{M}_G (or some extension² of it) implies a polynomial time algorithm to compute a min-cost popular maxmatching. Our main theorem is that the polytope \mathcal{M}_G has a compact extended formulation. So unlike the min-cost popular max-matching problem, interestingly and quite surprisingly, the min-cost popular max-matching problem is tractable.

Theorem 3. Given $G = (A \cup B, E)$ where nodes have strict preferences and $c : E \to \mathbb{R}$, a min-cost popular max-matching can be found in polynomial time.

A relaxation of popularity. Pareto-optimality is a natural relaxation of popularity that any reasonable matching in this domain should satisfy. If M is a matching that is not Pareto-optimal then there is a *better* matching N, i.e., no node is worse-off in N than in M and at least one node is better-off. The unpopularity factor of M is defined as follows [24]: $u(M) = \max_{N \neq M} \phi(N, M) / \phi(M, N)$.

A matching M is Pareto-optimal if $u(M) < \infty$. So there is no matching N such that $\phi(N, M) > 0$ and $\phi(M, N) = 0$. Observe that a popular matching M satisfies $u(M) \leq 1$. A maximum matching M that satisfies $u(M) < \infty$ is a *Pareto-optimal max-matching*. We show that it is NP-hard to find a min-cost Pareto-optimal matching/max-matching (see Theorem 12).

Background. Many algorithmic questions in popular matchings have been investigated in the last 10-15 years; we refer to [6] for a survey. In the domain of popular matchings with two-sided preferences (every node has a preference order ranking its neighbors), other than a handful of positive results [8,17,20,22], most

² A polytope Q that linearly projects to a polytope P is an *extension* of P and a linear description of Q is an *extended* formulation for P. The minimum size of an extension of P is the *extension complexity* of P.

optimization problems (incl. finding a popular matching that is not a min-size/max-size popular matching) have turned out to be NP-hard [10]. Computing a min-cost quasi-popular matching M, i.e., $u(M) \leq 2$, is also NP-hard [9].

Compact extended formulations for the *dominant* matching³ polytope [8,9] and the popular fractional matching polytope [21,23] are known but the popular matching polytope has near-exponential extension complexity [9]. Thus in contrast to stable matchings, the landscape of popular matchings has only a few positive results. However, within the set of maximum matchings, the notion of popularity seems more suitable than stability for algorithmic results. In the context of stability, a *best* maximum matching would be one with the least number of blocking edges. The problem of finding a maximum matching with the least number of blocking edges was considered in [5] and was shown to be NP-hard.

Though an O(mn) time algorithm to find a popular max-matching in G is known [20], there are no previous results on the *optimization* variant of this problem, i.e., to find a min-cost popular max-matching, which is a natural problem here. It is common in this domain to have an efficient algorithm to find a max-size matching in some class – say, Pareto-optimal matchings for one-sided preferences (only nodes in A have preferences) studied in [2] or popular matchings [17,20], however finding a *min-cost* matching in such a class is NP-hard [2,10]. Thus the existence of an efficient algorithm to find some popular max-matching was no guarantee on the tractability of the min-cost popular max-matching problem.

Our Techniques. An algorithm called the "|A|-level Gale-Shapley algorithm" was given in [20] to find a popular max-matching in $G = (A \cup B, E)$. As we show in Section 2, this algorithm is equivalent to running Gale-Shapley algorithm in an auxiliary instance G^* with |A| copies of each node in A and the proof in [20] can be easily adapted to show that there is a linear map from the set of stable matchings in G^* to the set of popular max-matchings in G (see Theorem 5). Our novel contribution here is to show that this map is *surjective*, i.e., every popular max-matching in G is the image of a stable matching in G^*

Our proof that every popular max-matching in G has a stable matching in G^* as its preimage is based on LP-duality. Here we introduce the notion of dual certificates for popular max-matchings—dual certificates for popular matchings are well-understood: these are elements in $\{0, \pm 1\}^n$ [21]. We show in Section 3 that every popular max-matching in G has a dual certificate $\vec{\alpha} \in \mathbb{Z}^n$ where $\alpha_a \in \{0, -2, -4, \ldots\}$ for $a \in A$ and $\alpha_b \in \{0, 2, 4, \ldots\}$ for $b \in B$ and α -values for some nodes can be restricted to fixed values (see Theorem 8). Given a popular max-matching M in G and such a dual certificate $\vec{\alpha}$, we use $\vec{\alpha}$ to construct a preimage S for M and prove that S is a stable matching in G^* .

2 Popular Max-Matchings

In this section we first show a simple characterization of popular max-matchings. Then we show a method to construct matchings that satisfy this characterization.

Let M be any matching in $G = (A \cup B, E)$. The following edge weight function wt_M will be useful here. For any $(a, b) \in E$:

let $\mathsf{wt}_M(a,b) = \begin{cases} 2 & \text{if } (a,b) \text{ blocks } M; \\ -2 & \text{if } a \text{ and } b \text{ prefer their assignments in } M \text{ to each other}; \\ 0 & \text{otherwise.} \end{cases}$

So $\mathsf{wt}_M(e) = 0$ for every $e \in M$. For any cycle/path ρ in G, let $\mathsf{wt}_M(\rho) = \sum_{e \in \rho} \mathsf{wt}_M(e)$. Theorem 4 uses this edge weight function to characterize popular max-matchings.

Theorem 4. For any maximum matching M in G, M is a popular max-matching if and only if (1) there is no alternating cycle C with respect to M such that $wt_M(C) > 0$ and (2) there is no alternating path p with an unmatched node as an endpoint such that $wt_M(p) > 0$.

³ These are popular matchings that are more popular than all larger matchings.

Proof. Let M be a popular max-matching. We need to show that conditions (1) and (2) given in the theorem statement hold. Suppose not. Then there exists either an alternating path with an unmatched node as an endpoint or an alternating cycle wrt M (call this path/cycle ρ) such that $\mathsf{wt}_M(\rho) > 0$. Since $\mathsf{wt}_M(e) \in \{0, \pm 2\}$, $\mathsf{wt}_M(\rho) \ge 2$.

Consider $N = M \oplus \rho$. This is a maximum matching in G and observe that $\Delta(N, M) \ge \operatorname{wt}_M(\rho) - 1$. We are subtracting 1 here to count for that endpoint of ρ (when ρ is a path) that is matched in M but will become unmatched in N. Since $\operatorname{wt}_M(\rho) \ge 2$, $\Delta(N, M) \ge 1$. So N is more popular than M: this is a contradiction to M's popularity within the set of maximum matchings. Thus conditions (1) and (2) have to hold.

To show the converse, suppose M is a maximum matching that obeys conditions (1) and (2). Consider the symmetric difference $M \oplus N$, where N is any maximum matching in G and let C be any alternating cycle here. We know from (1) that $wt_M(C) \leq 0$. Let p be any alternating path in $M \oplus N$. Since M and Nare maximum matchings, p is an alternating path with exactly one node not matched in M as an endpoint. We know from (2) that $wt_M(p) \leq 0$. So we have:

$$\Delta(N,M) \leq \sum_{\rho \in M \oplus N} \operatorname{wt}_M(\rho) \leq 0.$$

Thus M is popular within the set of maximum matchings in G.

An auxiliary instance G^* . We will now construct a new instance $G^* = (A^* \cup B^*, E^*)$ such that every stable matching in G^* maps to a maximum matching in G that satisfies properties (1) and (2) given in Theorem 4. The structure of the instance G^* is inspired by an instance used in [8] whose stable matchings map to dominant matchings in G.

We describe the node sets A^* and B^* . Let $n_0 = |A|$. For every $a \in A$, the set A^* has n_0 copies of a: call them a_0, \ldots, a_{n_0-1} . So $A^* = \bigcup_{a \in A} \{a_0, \ldots, a_{n_0-1}\}$.

Let $B^* = \bigcup_{a \in A} \{ d_1(a), \dots, d_{n_0-1}(a) \} \cup \{ \tilde{b} : b \in B \}$, where $\tilde{B} = \{ \tilde{b} : b \in B \}$ is a copy of the set B. So along with nodes in \tilde{B} , the set B^* also contains $n_0 - 1$ nodes $d_1(a), \dots, d_{n_0-1}(a)$ for each $a \in A$. These will be called *dummy* nodes.

The edge set. For each $(a, b) \in E$, the edge set E^* contains n_0 edges (a_i, \tilde{b}) for $0 \le i \le n_0 - 1$. For each $a \in A$ and $i \in \{1, \ldots, n_0 - 1\}$, E^* also has $(a_{i-1}, d_i(a))$ and $(a_i, d_i(a))$. The purpose of $d_1(a), \ldots, d_{n_0-1}(a)$ is to ensure that in any stable matching in G^* , at most one node among a_0, \ldots, a_{n_0-1} is matched to a neighbor in \tilde{B} . For each $i \in \{1, \ldots, n_0 - 1\}$, the preference order of $d_i(a)$ is $a_{i-1} \succ a_i$.

Preference orders. Let a's preference order in G be $b_1 \succ \cdots \succ b_k$. Then a_0 's preference order in G^* is $\tilde{b}_1 \succ \cdots \succ \tilde{b}_k \succ d_1(a)$, i.e., it is analogous to a's preference order in G with $d_1(a)$ added as a_0 's last choice.

- For $i \in \{1, \ldots, n_0 2\}$, the preference order of a_i in G^* is as follows: $d_i(a) \succ \tilde{b}_1 \succ \cdots \succ \tilde{b}_k \succ d_{i+1}(a)$. So a_i 's top choice is $d_i(a)$ and last choice is $d_{i+1}(a)$.
- The preference order of a_{n_0-1} is $d_{n_0-1}(a) \succ \tilde{b}_1 \succ \cdots \succ \tilde{b}_k$.

Since each of a_0, \ldots, a_{n_0-2} and $d_1(a), \ldots, d_{n_0-1}(a)$ is a top choice neighbor for some node, every stable matching in G^* has to match all these nodes. So the only node among $a_0, \ldots, a_{n_0-1}, d_1(a), \ldots, d_{n_0-1}(a)$ that can possibly be left unmatched in a stable matching in G^* is a_{n_0-1} .

Consider any $b \in B$ and let its preference order in G be $a \succ \cdots \succ z$. Then the preference order of \tilde{b} in G^* is

$$\underbrace{a_{n_0-1} \succ \cdots \succ z_{n_0-1}}_{\text{all subscript } n_0 - 1 \text{ neighbors}} \succ \underbrace{a_{n_0-2} \succ \cdots \succ z_{n_0-2}}_{\text{all subscript } n_0 - 2 \text{ neighbors}} \succ \cdots \succ \underbrace{a_0 \succ \cdots \succ z_0}_{\text{all subscript } 0 \text{ neighbors}}$$

That is, \tilde{b} 's preference order in G^* is all its subscript $n_0 - 1$ neighbors, followed by all its subscript $n_0 - 2$ neighbors, so on, and finally, all its subscript 0 neighbors. For each $i \in \{0, \ldots, n_0 - 1\}$: within all subscript i neighbors, the order of preference for \tilde{b} in G^* is b's order of preference in G.

The set S'. For any stable matching S in G^* , define $S' \subseteq E$ to be the set of edges obtained by deleting edges in S that are incident to dummy nodes and replacing any edge $(a_i, \tilde{b}) \in S$ with the original edge $(a, b) \in E$. Since S matches at most one node among a_0, \ldots, a_{n_0-1} to a neighbor in \tilde{B} , the set S' is a valid matching in G. The proof of Theorem 5 is based on the proof of correctness of the |A|-level Gale-Shapley algorithm (from [20]) in the original instance G.

Theorem 5. Let S be a stable matching in G^* . Then S' is a popular max-matching in G.

Proof. Partition the set A into $A_0 \cup \cdots \cup A_{n_0-1}$ where for $0 \le i \le n_0 - 2$: $A_i = \{a \in A : (a_i, \tilde{b}) \in S \text{ for some } \tilde{b} \in \tilde{B}\}$, i.e., a_i is matched in S to a neighbor in \tilde{B} . The left-out nodes in A, i.e., those in $A \setminus (A_0 \cup \cdots \cup A_{n_0-2})$, form the set A_{n_0-1} (see Fig. 1).

Similarly, partition B into $B_0 \cup \cdots \cup B_{n_0-1}$ where for $1 \leq i \leq n_0 - 1$: $B_i = \{b : (a_i, \tilde{b}) \in S \text{ for some } a \in A_i\}$, i.e., \tilde{b} 's partner in S is a subscript i node. Let $B_0 = B \setminus (B_1 \cup \cdots \cup B_{n_0-1})$ be the set of left-out nodes in B.



Fig. 1. $A = A_0 \cup \cdots \cup A_{n_0-1}$ and $B = B_0 \cup \cdots \cup B_{n_0-1}$. All nodes unmatched in S' are in $A_{n_0-1} \cup B_0$. The dashed edges are blocking edges to S'.

The following properties hold: (these are proved in the appendix)

- 1. $S' \subseteq \bigcup_{i=0}^{n_0-1} (A_i \times B_i)$. Moreover, S' restricted to each set $A_i \cup B_i$ is stable.
- 2. For any i and edge (a, b) where $a \in A_{i+1}, b \in B_i$: we have $\mathsf{wt}_{S'}(a, b) = -2$.
- 3. G has no edge in $A_i \times B_j$ where $i \ge j+2$.
- 4. Any blocking edge to S' has to be in $A_i \times B_j$ where $i \leq j 1$.
- 5. All nodes that are unmatched in S' are in $A_{n_0-1} \cup B_0$.
- 6. S' is a maximum matching in G.

Properties 1-4 imply that for any alternating cycle C wrt S', we have $\mathsf{wt}_{S'}(C) \leq 0$. Properties 1-5 imply that for any alternating path p with one unmatched node as an endpoint, we have $\mathsf{wt}_{S'}(p) \leq 0$. We refer to [20, Theorem 2] for more details. Property 6 states that S' is a maximum matching in G. Hence S' is a popular max-matching in G (by Theorem 4).

Thus every stable matching in G^* maps to a popular max-matching in G. We will show in Section 3 that every popular max-matching in G has to be realized in this manner. This is the tough part and we will use dual certificates here. We will see that dual certificates capture a useful profile of popular max-matchings.

3 Dual Certificates

We first consider the simple case when G admits a perfect matching. Let M be a popular perfect matching in G. So no perfect matching is more popular than M. Consider the following linear program (LP1) that computes a max-weight (with respect to wt_M) perfect matching in G.

maximize
$$\sum_{e \in E} \operatorname{wt}_M(e) \cdot x_e$$
 (LP1)

subject to

$$\sum_{e \in \delta(u)} x_e = 1 \quad \forall u \in A \cup B \quad \text{and} \quad x_e \ge 0 \quad \forall e \in E.$$

It follows from the definition of the function wt_M that the optimal value of (LP1) is $\max_N \Delta(N, M)$ where N is a perfect matching in G. So if M is a popular perfect matching then the optimal value of (LP1) is 0, which is $\Delta(M, M)$, i.e., the edge incidence vector of M is an optimal solution to (LP1).

The linear program (LP2) is the dual of (LP1). Hence if M is a popular perfect matching then there exists a dual feasible $\vec{\alpha}$ such that $\sum_{u \in A \cup B} \alpha_u = 0$.

minimize
$$\sum_{u \in A \cup B} y_u$$
 (LP2)

subject to

$$y_a + y_b \ge \operatorname{wt}_M(a, b) \quad \forall (a, b) \in E.$$

Let $\vec{\alpha}$ be an optimal dual solution. Observe that there exists an integral optimal solution to (LP2) since the constraint matrix is totally unimodular. Thus we can assume that $\vec{\alpha} \in \mathbb{Z}^{2n_0}$, where $n_0 = |A| = |B|$.

Lemma 6. If M is a popular perfect matching in G then there exists an optimal solution $\vec{\alpha}$ to (LP2) such that $\alpha_a \in \{0, -2, -4, \dots, -2(n_0 - 1)\}$ for all $a \in A$ and $\alpha_b \in \{0, 2, 4, \dots, 2(n_0 - 1)\}$ for all $b \in B$.

Proof. The dual feasibility constraints are $\alpha_a + \alpha_b \geq \mathsf{wt}_M(a, b)$ for all $(a, b) \in E$. For each edge $(a, b) \in M$: $\alpha_a + \alpha_b = \mathsf{wt}_M(a, b) = 0$ by complementary slackness. Since $\alpha_b = -\alpha_a$ for $(a, b) \in M$ and because $\mathsf{wt}_M(e) \in \{0, \pm 2\}$ for each edge e, we can assume that in the sorted order of distinct α -values taken by nodes in A, for any two consecutive values $\alpha_{a'}, \alpha_{a''}$, where $\alpha_{a'} > \alpha_{a''}$, we have $\alpha_{a'} - \alpha_{a''} = 2$. Thus we can assume that $\alpha_a \in \{k, k - 2, k - 4, \dots, k - 2(n_0 - 1)\}$ for all $a \in A$ and $\alpha_b \in \{-k, -k + 2, -k + 4, \dots, -k + 2(n_0 - 1)\}$ for all $b \in B$, for some $k \in \mathbb{Z}$.

Observe that k has no impact on the objective function since |A| = |B|: so k's and -k's cancel each other out. Let us update $\vec{\alpha}$ as follows: $\alpha_a = \alpha_a - k$ for every $a \in A$ and $\alpha_b = \alpha_b + k$ for every $b \in B$. The updated vector $\vec{\alpha}$ continues to be dual feasible since $\alpha_a + \alpha_b$, for any edge (a, b), is unchanged by this update. Thus there is an optimal solution $\vec{\alpha}$ to (LP2) such that $\alpha_a \in \{0, -2, -4, \dots, -2(n_0 - 1)\}$ for all $a \in A$ and $\alpha_b \in \{0, 2, 4, \dots, 2(n_0 - 1)\}$ for all $b \in B$.

Let M be a popular perfect matching in G. In order to define a stable matching S in G^* such that M = S' (the set S' is defined above Theorem 5), we will use the vector $\vec{\alpha}$ described in Lemma 6. Since M is perfect, we know that for any $a \in A$, there is an edge $(a, b) \in M$ for some neighbor b of a. Recall that $\alpha_a + \alpha_b = \operatorname{wt}_M(a, b) = 0$ by complementary slackness. We will include the edge (a_i, \tilde{b}) in S where $\alpha_a = -2i$ and $\alpha_b = 2i$. Thus we define S as follows:

$$S = \bigcup_{i=0}^{n_0-1} \{ (a_i, b) : (a, b) \in M \text{ and } \alpha_a = -2i, \alpha_b = 2i \} \cup \{ \text{necessary edges incident to dummy nodes in } G^* \}.$$

In more detail, the edges incident to dummy nodes that are present in S are as follows: if $\alpha_a = -2i$ then these edges are $(a_j, d_{j+1}(a))$ for $0 \le j \le i-1$ and $(a_j, d_j(a))$ for $i+1 \le j \le n_0-1$. Since $(a_i, \tilde{b}) \in S$, all the n_0 nodes a_0, \ldots, a_{n_0-1} and the dummy nodes $d_1(a), \ldots, d_{n_0-1}(a)$ corresponding to a in G^* are matched in S. This holds for every $a \in A$. Also every $\tilde{b} \in \tilde{B}$ is matched in S since M is a perfect matching in G. Thus S is a perfect matching in G^* . It is easy to check that S' = M. What we need to prove is the stability of S in G^* .

Theorem 7. The matching S is stable in G^* .

Proof. We need to show there is no edge in G^* that blocks S. There is no blocking edge incident to a dummy node: this is because a dummy node $d_i(a)$ has only two neighbors and when $d_i(a)$ is matched in S to its second choice neighbor a_i , its top choice neighbor a_{i-1} prefers its partner in S to $d_i(a)$.

Let us now show that no node in a_0, \ldots, a_{n_0-1} has a blocking edge incident to it, for any $a \in A$. Let $(a_i, \tilde{b}) \in S$ where $(a, b) \in M$. All of $a_{i+1}, \ldots, a_{n_0-1}$ are matched to their respective top choice neighbors $d_{i+1}(a), \ldots, d_{n_0-1}(a)$. So there is no blocking edge incident to any of $a_{i+1}, \ldots, a_{n_0-1}$.

All of a_0, \ldots, a_{i-1} are matched to their last choice neighbors—these are the dummy nodes $d_1(a), \ldots, d_i(a)$, respectively. Consider any neighbor $w \in B$ of a. We need to show that $\tilde{w} \in \tilde{B}$ is matched in S to a neighbor preferred to all of a_0, \ldots, a_{i-1} . We have $\alpha_a + \alpha_w \ge \operatorname{wt}_M(a, w)$. Since $\alpha_a = -2i$ and $\operatorname{wt}_M(e) \ge -2$ for every edge e, it follows that $\alpha_w \ge 2i - 2$.

So $(z, w) \in M$ for some neighbor z of w such that $\alpha_z = -\alpha_w \leq -(2i-2)$. Equivalently, $(z_j, \tilde{w}) \in S$ where $j \geq i-1$. Thus there is no blocking edge between \tilde{w} and any of a_0, \ldots, a_{i-2} by \tilde{w} 's preference order in G^* . We will now show that (a_{i-1}, \tilde{w}) cannot be a blocking edge.

- If $j \ge i$ then by \tilde{w} 's preference order in G^* , \tilde{w} prefers z_j to a_{i-1} and so (a_{i-1}, \tilde{w}) does not block S.
- If j = i 1 then $wt_M(a, w) \leq \alpha_a + \alpha_w = -2i + 2i 2 = -2$. So both a and w prefer their respective partners in M to each other. Thus \tilde{w} prefers z_{i-1} to a_{i-1} . So (a_{i-1}, \tilde{w}) does not block S.

Finally, we need to show that there is no blocking edge incident to a_i . By the above argument, we only need to consider edges (a_i, \tilde{w}) where $(z_i, \tilde{w}) \in S$. So $wt_M(a, w) \leq \alpha_a + \alpha_w = -2i + 2i = 0$. Hence either $(a, w) \in M$ or one of a, w prefers its partner in M to the other. So either $(a_i, \tilde{w}) \in S$ or one of a_i, \tilde{w} prefers its partner in S to the other; thus the edge (a_i, \tilde{w}) does not block S. Hence no edge in G^* blocks S. \Box

3.1 The general case

We showed that when G has a perfect matching, our mapping from the set of stable matchings in G^* to the set of popular max-matchings in G is surjective. Now we look at the general case, i.e., G need not have a perfect matching.

Let M be a popular max-matching in G. Let $U \subseteq A \cup B$ be the set of nodes left unmatched in M. Consider (LP3) that computes a max-weight perfect matching (with respect to wt_M) in the subgraph G' induced on $(A \cup B) \setminus U$.

Let E' be the edge set of G'. For any $v \in (A \cup B) \setminus U$, let $\delta'(v) = \delta(v) \cap E'$.

maximize
$$\sum_{e \in E'} \operatorname{wt}_M(e) \cdot x_e$$
 (LP3)

subject to

$$\sum_{e \in \delta'(v)} x_e = 1 \quad \forall v \in (A \cup B) \setminus U \quad \text{and} \quad x_e \ge 0 \quad \forall e \in E'$$

The optimal value of (LP3) is $\max_N \Delta(N, M)$ where N is a perfect matching in G'. Any perfect matching in G' is a maximum matching in G and since M is a popular max-matching in G, $\Delta(N, M) \leq 0$ for any perfect matching N in G'. Since $\Delta(M, M) = 0$, the edge incidence vector of M is an optimal solution to (LP3). The linear program (LP4) is the dual of (LP3).

minimize
$$\sum_{u \in (A \cup B) \setminus U} y_u$$
 (LP4)

subject to

$$y_a + y_b \ge \operatorname{wt}_M(a, b) \quad \forall (a, b) \in E'.$$

Let $n' = |(A \cup B) \setminus U|$. There exists an optimal solution $\vec{\alpha} \in \mathbb{Z}^{n'}$ to (LP4) such that $\sum_{u \in (A \cup B) \setminus U} \alpha_u = 0$. Moreover, we can assume the following (see Lemma 6) where $A' = A \setminus U$, $B' = B \setminus U$, and $|A'| = |B'| = n_0$:

(1) $\alpha_a \in \{0, -2, -4, \dots, -2(n_0 - 1)\}$ for all $a \in A'$ (2) $\alpha_b \in \{0, 2, 4, \dots, 2(n_0 - 1)\}$ for all $b \in B'$.

For any $T \subseteq A \cup B$, let Nbr(T) be the set of neighbors in G of nodes in T. Theorem 8 is our main technical result here. Let $U_A = U \cap A$ and $U_B = U \cap B$.

Theorem 8. Let M be a popular max-matching in G and let U be the set of nodes left unmatched in M. There exists an optimal solution $\vec{\alpha}$ to (LP4), where $\alpha_a \in \{0, -2, \ldots, -2(n_0 - 1)\}$ for $a \in A', \alpha_b \in \{0, 2, \ldots, 2(n_0 - 1)\}$ for $b \in B'$ such that (i) $\alpha_a = 0$ for $a \in Nbr(U_B)$ and (ii) $\alpha_b = 2(n_0 - 1)$ for $b \in Nbr(U_A)$.

Proof. We know there exists an optimal solution $\vec{\alpha}$ to (LP4) such that $\alpha_a \in \{0, -2, \dots, -2(n_0 - 1)\}$ for $a \in A'$ and $\alpha_b \in \{0, 2, \dots, 2(n_0 - 1)\}$ for $b \in B'$. Now we will update $\vec{\alpha}$ so that it remains an optimal solution to (LP4) in the above format and it also satisfies properties (i) and (ii) given in the theorem statement.

We will first update $\vec{\alpha}$ so that $\alpha_b = 2(n_0 - 1)$ for all $b \in Nbr(U_A)$, i.e., property (ii) is obeyed. We will then update $\vec{\alpha}$ so that it satisfies $\alpha_a = 0$ for all $a \in Nbr(U_B)$, i.e., property (i) is obeyed. We then use the fact that M is a maximum matching to show that the second update did not undo the "good" caused by the first update, i.e., $\vec{\alpha}$ satisfies property (i) and property (ii).

Property (ii). Suppose the original vector $\vec{\alpha} \in \{0, \ldots, \pm 2(n_0 - 1)\}^{n'}$ does not satisfy property (ii). Then we will update $\vec{\alpha}$ so that property (ii) is satisfied. First, we increase the α -values of nodes in Nbr(U_A) to $2(n_0 - 1)$ and decrease the α -values of their partners in M to $-2(n_0 - 1)$. Now $\vec{\alpha}$ may no longer be a feasible solution to (LP4). We will use the following three update rules for all $a \in A'$ to make $\vec{\alpha}$ feasible again. Let $\alpha_a = -2i$ where $i \in \{0, \ldots, n_0 - 1\}$. Suppose there is some $(a, b) \in E'$ such that $\alpha_a + \alpha_b < \operatorname{wt}_M(a, b)$. Let M(b) be b's partner in M.

- Rule 1. If wt_M(a, b) = 0 then update $\alpha_b = 2i$ and $\alpha_{M(b)} = -2i$.
- Rule 2. If wt_M(a, b) = -2 then update $\alpha_b = 2(i-1)$ and $\alpha_{M(b)} = -2(i-1)$.
- Rule 3. If $\operatorname{wt}_M(a,b) = 2$ then update $\alpha_b = 2(i+1)$ and $\alpha_{M(b)} = -2(i+1)$.

At the onset, $\vec{\alpha}$ was a feasible solution to (LP4): so $\alpha_a + \alpha_b \ge \mathsf{wt}_M(a, b)$ for $(a, b) \in E'$. Then we moved nodes in $\mathsf{Nbr}(U_A)$ and their partners in M to sets B_{n_0-1} and A_{n_0-1} , resp., where $A_i = \{a \in A' : \alpha_a = -2i\}$ and $B_i = \{b \in B' : \alpha_b = 2i\}$ for all i. The subscript i will be called the *level* of nodes in $A_i \cup B_i$.

The nodes that moved to A_{n_0-1} have a lower α -value than earlier and it is these nodes that "pull" their neighbors upwards to higher levels as given by rules 1-3. Let a be a new node in level i and let b be a neighbor of a such that $\alpha_a + \alpha_b < \mathsf{wt}_M(a, b)$. Then b and M(b) move to: (1) level i if $\mathsf{wt}_M(a, b) = 0$, (2) level i - 1 if $\mathsf{wt}_M(a, b) = -2$, else (3) level i + 1, i.e., if $\mathsf{wt}_M(a, b) = 2$.

In turn, the nodes in A' that have moved to these higher levels by rules 1-3 pull their neighbors and the partners of these neighbors upwards to higher levels by these rules. Thus we may get further new nodes in B_{n_0-1} , A_{n_0-1} and so on. While any of rules 1-3 is applicable, we apply that rule. So a rule may be applied many times to the same edge in E'.

Claim 1 (proved in the appendix) shows a useful property. We show in its proof that such a blocking edge creates a *forbidden* alternating cycle/path wrt M (as given in Theorem 4).

Claim 1 By applying the above rules, suppose a node $v_0 \in A'$ moves to A_{n_0-1} . Then there is no blocking edge (e such that $wt_M(e) = 2$) incident to v_0 .

Applying rules 1-3 increases the α -values of some nodes in B' and it never decreases the α -value of any node in B'. The nodes in B' with increased α -values and their partners have moved to higher levels (see Fig. 1). This upwards movement of nodes has to terminate at level $n_0 - 1$. For the α -value of any $b \in B'$ to be increased beyond $2(n_0 - 1)$, we need a blocking edge (a, b) where $a \in A_{n_0-1}$: this would cause rule 3 to be applied which would increase α_b to $2n_0$. However there is no such blocking edge (by Claim 1).

Since there are n_0 levels and because $|B'| = n_0$, there can be at most n_0^2 applications of these rules. When no rule is applicable, $\vec{\alpha}$ is a feasible solution to (LP4). Moreover, $\sum_u \alpha_u$ is invariant under this update of α -values, since we maintain $\alpha_a + \alpha_b = 0$ for every $(a, b) \in M$. Hence $\vec{\alpha}$ is an optimal solution to (LP4). Thus for every popular max-matching M, there is an optimal solution $\vec{\alpha}$ to (LP4) in the desired format that satisfies property (ii).

Property (i). We now have an optimal solution $\vec{\alpha} \in \{0, \pm 2, \ldots, \pm 2(n_0 - 1)\}^{n'}$ to (LP4), where $\alpha_a \leq 0$ for all $a \in A'$ and $\alpha_b \geq 0$ for all $b \in B'$, such that $\alpha_b = 2(n_0 - 1)$ for all $b \in \mathsf{Nbr}(U_A)$. Suppose property (i) is not satisfied.

Then we increase α -values of certain nodes in A' – this moves several nodes downwards with respect to their level (see Fig. 1) and ensures that property (i) holds. First, we increase the α -values of nodes in Nbr (U_B) to 0 and their partners also have α -values updated to 0. Then we will use the following three update rules for all $b \in B'$. Let $\alpha_b = 2i$ where $i \in \{0, \ldots, n_0 - 1\}$. Suppose there is an edge $(a, b) \in E'$ such that $\alpha_a + \alpha_b < wt_M(a, b)$. Let M(a) be a's partner in M.

- Rule 4. If $wt_M(a,b) = 0$ then update $\alpha_a = -2i$ and $\alpha_{M(a)} = 2i$.
- Rule 5. If $\operatorname{wt}_M(a,b) = -2$ then update $\alpha_a = -2(i+1)$ and $\alpha_{M(a)} = 2(i+1)$.
- Rule 6. If $\operatorname{wt}_M(a,b) = 2$ then update $\alpha_a = -2(i-1)$ and $\alpha_{M(a)} = 2(i-1)$.

Applying rules 4-6 increases the α -values of some nodes in A' and it never decreases the α -value of any node in A'. The nodes in A' with increased α -values and their partners have moved to lower levels. Moreover, the movement of nodes downwards has to stop at level 0 since no blocking edge can be incident to any node that moves to B_0 (analogous to Claim 1). While any of the above three rules is applicable, we apply that rule. When no rule is applicable, $\vec{\alpha}$ is a feasible solution to (LP4). Since $\sum_u \alpha_u = 0$, $\vec{\alpha}$ is an optimal solution to (LP4). So there is an optimal solution $\vec{\alpha}$ to (LP4) in the desired format that satisfies property (i).

Properties (i) and (ii). Note that we cannot claim straightaway that the above $\vec{\alpha}$ satisfies both property (i) and property (ii). This is because rules 4-6 pull nodes downwards and this may have caused $\alpha_b < 2(n_0 - 1)$ for some $b \in Nbr(U_A)$. We will now show this is not possible. Suppose there is such a node $w_1 \in Nbr(U_A)$.

So there is some $z \in U_A$ such that $(z, w_1) \in E$ and though $\alpha_{w_1} = 2(n_0 - 1)$ just before we started applying rules 4-6, the application of these rules caused w_1 to move downwards, i.e., $\alpha_{w_1} < 2(n_0-1)$ at the end. Initially we added nodes in $Nbr(U_B)$ and their partners in M to A_0 and B_0 , respectively. Then we repeatedly applied rules 4-6 and this resulted in w_1 moving downwards from level $n_0 - 1$. Corresponding to what caused w_1 to be "pulled" downwards, we will construct an alternating path $p = w_1 - M(w_1) - w_2 - M(w_2) - \cdots - w_r - M(w_r) - u$ between w_1 and some node $u \in U_B$.

The path p will consist of r pairs of edges for some $r \ge 1$. For $1 \le i \le r-1$: the *i*-th pair of edges is $(w_i, M(w_i))$ and $(M(w_i), w_{i+1})$ where w_{i+1} is the node that pulled $M(w_i)$ and w_i downwards to their level due to the application of one of rules 4-6. The last pair of edges in p is $(w_r, M(w_r))$ and $(M(w_r), u)$, where $w_r \in B_0, M(w_r) \in A_0$, and $u \in U_B$. Thus we have an alternating path p wrt M between w_1 and $u \in U_B$.

By adding the edge (z, w_1) as a prefix to the path p, we get an augmenting path $z - w_1 - \cdots - M(w_r) - u$ with respect to M. However there cannot be any augmenting path wrt M since M is a maximum matching in G. Thus $\vec{\alpha}$ satisfies property (ii). So we have an optimal solution $\vec{\alpha}$ to (LP4) in the desired format that satisfies both property (i) and property (ii).

Theorem 9. Let M be any popular max-matching in G. Then there is a stable matching S in the graph G^* such that M = S'.

Proof. Let U be the set of nodes left unmatched in M. A useful observation is that every node v in Nbr(U) has to be matched in M to some neighbor that v prefers to all its neighbors in U. Otherwise by replacing $(v, w) \in M$ with (v, u), where v prefers $u \in U$ to w, we get a maximum matching more popular than M.

In order to construct the matching S in G^* , we will use the vector $\vec{\alpha}$ described in Theorem 8. Let $S = \bigcup_{i=0}^{n_0-1} \{(a_i, \tilde{b}) : (a, b) \in M \text{ and } \alpha_a = -2i, \alpha_b = 2i\} \cup \{\text{necessary edges incident to dummy nodes in } G^*\}$ where for each $a \in A'$: if $\alpha_a = -2i$ then the necessary edges incident to dummy nodes corresponding to a are $(a_i, d_{i+1}(a))$ for $0 \le j \le i-1$ and $(a_i, d_i(a))$ for $i+1 \le j \le n_0-1$. It is easy to check that S' = M.

We will now show that S is a stable matching in G^* . Consider any edge (a_j, \tilde{w}) in G^* where $(a, w) \in E'$ and $0 \leq j \leq n_0 - 1$. The proof of Theorem 7 shows that (a_j, \tilde{w}) does not block S. Consider any edge (a, w)in $E \setminus E'$.

- Suppose $a \in U_A$. So $w \in Nbr(U_A)$: hence $\alpha_w = 2(n_0 1)$ by property (ii). So $(z_{n_0-1}, \tilde{w}) \in S$ for some neighbor z that w prefers to a. Thus (a_{n_0-1}, \tilde{w}) does not block S. For $i \in \{0, \ldots, n_0 2\}$, none of the edges (a_i, \tilde{w}) can block S (by \tilde{w} 's preference order).
- Suppose $w \in U_B$. So $a \in \mathsf{Nbr}(U_B)$: hence $\alpha_a = 0$ by property (i). Thus $(a_0, \tilde{b}) \in S$ for some neighbor b that a prefers to w. Hence (a_0, \tilde{w}) does not block S. Moreover, none of the edges (a_i, \tilde{w}) for $i \in \{1, \ldots, n_0 1\}$ can block S since a_1, \ldots, a_{n_0-1} are matched to their respective top choice neighbors $d_1(a), \ldots, d_{n_0-1}(a)$.

Finally, no edge incident to a dummy node blocks S. Hence S is stable in G^* .

Thus there is a linear map that is surjective from the set of stable matchings in G^* to the set of popular max-matchings in G (by Theorem 5 and Theorem 9). Hence the stable matching polytope of G^* is an extension of the popular max-matching polytope of G.

3.2 A compact extended formulation for the popular max-matching polytope

The formulation of the stable matching polytope of G^* (from [25]) is given below. For any node u in G^* , the set $\{v' \succ_u v\}$ is the set of all neighbors of u in G^* that it prefers to v. Let $\delta^*(u)$ denote the set of edges incident to u in G^* . Let $T = \bigcup_{a \in A} (\{a_0, \ldots, a_{n_0-2}\} \cup \{d_1(a), \ldots, d_{n_0-1}(a)\})$: every node in T has to be matched in all stable matchings in G^* .

$$\sum_{w \succ_{a_i} \tilde{b}} x_{(a_i,w)} + \sum_{z \succ_{\tilde{b}} a_i} x_{(z,\tilde{b})} + x_{(a_i,\tilde{b})} \ge 1 \qquad \forall (a_i,\tilde{b}) \in E^*$$
$$\sum_{e \in \delta^*(u)} x_e \le 1 \quad \forall u \in A^* \cup B^* \qquad \text{and} \qquad x_e \ge 0 \qquad \forall e \in E^*$$
$$\sum_{e \in \delta^*(u)} x_e = 1 \quad \forall u \in T \quad \text{and} \qquad x_{(a,b)} = \sum_{i=0}^{n_0-1} x_{(a_i,\tilde{b})} \quad \forall (a,b) \in E.$$

- The topmost constraint captures the *stability* constraint for edge $(a_i, \tilde{b}) \in E^*$ where $(a, b) \in E$ and $0 \le i \le n_0 1$.
- The constraints in the second line capture that \vec{x} belongs to the matching polytope of G^* .
- The constraints in the third line (on the left) capture the stability constraint for the edges $(a_{i-1}, d_i(a))$ and $(a_i, d_i(a))$ for $1 \le i \le n_0 - 1$.
- The constraint $x_{(a,b)} = \sum_{i=0}^{n_0-1} x_{(a_i,\tilde{b})}$ for every $(a,b) \in E$ captures the linear map from the edge set E^* to the edge set E.

We showed that the polytope described by the above constraints is an extension of the popular maxmatching polytope of G. Linear programming on the above formulation with $\min \sum_{e \in E} c(e) \cdot x_e$ as the objective function computes a min-cost popular max-matching in G in polynomial time.

4 Min-cost Pareto-optimal (max-)matching

In this section we consider the problems of computing a min-cost Pareto-optimal matching and a min-cost Pareto-optimal max-matching in a marriage instance with edge costs in $\{0, 1\}$. We will show these problems are NP-hard.

Given a 3SAT formula ψ , we will build an instance G_{ψ} with edge costs in $\{0, 1\}$ such that G_{ψ} admits a Pareto-optimal matching of cost 0 if and only if ψ is satisfiable. Any Pareto-optimal matching of cost 0 would have to be a perfect matching in G_{ψ} . Hence this will prove the NP-hardness of both the min-cost Pareto-optimal matching problem and the min-cost Pareto-optimal max-matching problem.

Our reduction resembles a hardness reduction from [7] that showed the NP-hardness of deciding if an instance G has a stable matching M that is also *dominant*, i.e., M is more popular than every larger matching. As done in this reduction, we will first transform ψ so that every clause contains either only positive literals or only negative literals; moreover, there will be a single occurrence of each negative literal in the transformed ψ . This is easy to achieve:

- let X_1, \ldots, X_n be the starting variables. For $i \in [n]$: replace all occurrences of $\neg X_i$ with the same variable X_{n+i} (a new one) and add the two clauses $(X_i \lor X_{n+i}) \land (\neg X_i \lor \neg X_{n+i})$ to capture $\neg X_i \equiv X_{n+i}$. Thus there are 2n variables in the transformed ψ .

We build the graph G_{ψ} as follows. There are two types of gadgets: those that correspond to positive clauses and those that correspond to negative clauses. Fig. 2 shows how a positive clause gadget with 3 literals looks like and Fig. 3 shows how a negative clause gadget looks like.



Fig. 2. The clause gadget for a positive clause $C_{\ell} = x \lor y \lor z$. Every occurrence of a literal in ψ has a separate gadget. So we ought to use labels such as $a_{x,\ell}, b_{x,\ell}, \ldots$ here; for the sake of simplicity, we used the labels a_x, b_x, \ldots here.

We now describe the preference lists of nodes in a positive clause $C_{\ell} = x \lor y \lor z$ (see Fig. 2). The nodes a_x, a'_x, b_x, b'_x occur in x's gadget and a_y, a'_y, b_y, b'_y occur in y's gadget and a_z, a'_z, b_z, b'_z occur in z's gadget: these gadgets are in the ℓ -th clause gadget C_{ℓ} . Every occurrence of a literal has a separate gadget.

a_x	a'_x	a_y	a'_y	a_z	a'_z	<i>b</i>	b'	<i>b.</i> .	b'	<i>b.</i>	b'
	b_r	b_r	b_u	b_{n}	<i>b</i> ~	v_x		v_y	^v y		
$\frac{\tilde{b}_{\pi}}{h_{\pi}}$	b'	<i>b</i>	b'	h_{z}	$\tilde{h'}$	a_y	a'_x	a_z	a'_y	a_x	a'_z
$\frac{\partial x}{\partial t}$		$\frac{\partial y}{\partial t}$	v_y	$\frac{\partial z}{\partial t}$		a_x	c_x	a_y	c_y	a_z	c_z
u_x	_	u_y	_	u_z		- a'	<i>a</i>	a'	а.	a'	a
b'_x	—	b'_y	-	b'_z	-	$^{\omega}x$	ux	^{a}y	uy	uz	$ $ ω_z

Here a_x 's top choice is b_z , second choice b_x , third choice d'_x , fourth choice b'_x , and similarly for other nodes. For every occurrence of a positive literal x: there will be a pair of *consistency edges* – the pair (a_x, d'_x) and (b'_x, c_x) in Fig. 4 – between this gadget of x and the unique gadget of $\neg x$. In our preferences, the neighbors on consistency edges are marked in red.

The preference lists of nodes that occur in a clause gadget with 2 positive literals will be totally analogous to the preference lists of nodes in a clause gadget with 3 positive literals. We will now describe the preference lists of nodes in a negative clause k—the overall picture here is given in Fig. 3.



Fig. 3. A clause gadget corresponding to a negative clause $D_k = \neg x \lor \neg y$; due to our transformation of ψ , every negative clause has only 2 literals.

c_x	c'_x	c_y	c'_y		d	d'	<i>d</i>	d'
d_y	d_x	d_x	d_y		a_x	^u x	uy	
$\overline{d_x}$	d'_x	d_y	d'_u	•	c_y	c_x	c_x	c_y
$b'_{x,i}$	_	$b'_{a',i'}$		-	c_x	$a_{x,i}$	c_y	$a_{y,i'}$
	_	y,ι	_		c'_x	•••	c'_y	• • •
b'.	_	h'	_	-	—	$a_{x,j}$	_	$a_{y,j'}$
$\frac{d'_{x,j}}{d'_{x}}$		d'_{u}			_	c_x	—	c_y

The nodes c_x, c'_x, d_x, d'_x and c_y, c'_y, d_y, d'_y occur in the gadgets of $\neg x$ and $\neg y$, respectively. The nodes $b'_{x,i}, \ldots, b'_{x,j}$ (resp., $b'_{y,i'}, \ldots, b'_{y,j'}$) in the preference lists above are the b'-nodes in the x-gadgets (resp., y-gadgets) in the various clauses that x (resp., y) occurs in. Similarly, $a_{x,i}, \ldots, a_{x,j}$ (resp., $a_{y,i'}, \ldots, a_{y,j'}$) are the *a*-nodes in the x-gadgets (resp., y-gadgets) in the various clauses that x (resp., y) occurs in. The preference order among the b'-nodes and among the *a*-nodes in these lists is not important. The consistency edges between a gadget of x and the gadget of $\neg x$ are shown in Fig. 4.



Fig. 4. For the sake of simplicity, we use a_x, b_x, a'_x, b'_x to denote the 4 nodes in the gadget of x in the ℓ -th clause; c_x, d_x, c'_x, d'_x are the 4 nodes in the unique gadget of $\neg x$. The consistency edges are the red dashed edges.

Edge costs. For each edge e in G_{ψ} , we will set $cost(e) \in \{0, 1\}$ as follows.

- For each variable $r \in \{X_1, \ldots, X_{2n}\}$: set cost(e) = 0 where e is any of the 4 edges in any literal gadget $\langle a_r, b_r, a'_r, b'_r \rangle$ of r or any of the 4 edges in the gadget $\langle c_r, d_r, c'_r, d'_r \rangle$ of $\neg r$.
- For all other edges e, set cost(e) = 1.

In particular, for any edge e in the consistency pair for any variable, we have cost(e) = 1. In our figures, all dashed edges have cost 1 and all solid edges have cost 0. Let M be a Pareto-optimal matching in G_{ψ} with cost(M) = 0. So M has to use only cost 0 edges. Thus M is forbidden to use any edge other than the 4 edges in the gadget of any literal. Moreover, since M is Pareto-optimal, M cannot leave two adjacent nodes unmatched. Thus for $r \in \{X_1, \ldots, X_{2n}\}$:

- 1. From a gadget of r (say, on nodes a_r, b_r, a'_r, b'_r), either (i) $(a_r, b_r), (a'_r, b'_r)$ are in M or (ii) $(a_r, b'_r), (a'_r, b_r)$ are in M.
- 2. From the gadget of $\neg r$ (the nodes are c_r, d_r, c'_r, d'_r), either (i) $(c_r, d_r), (c'_r, d'_r)$ are in M or (ii) $(c_r, d'_r), (c'_r, d_r)$ are in M.

Thus any Pareto-optimal matching in G_{ψ} of cost 0 is a perfect matching. It is easy to show Lemma 10.

Lemma 10. Let M be a Pareto-optimal matching in G_{ψ} . For any $r \in \{X_1, \ldots, X_{2n}\}$, both (a_r, b'_r) and (c_r, d'_r) cannot simultaneously be in M.

Proof. The preferences of the nodes are set such that if both (a_r, b'_r) and (c_r, d'_r) are in M then both the non-matching edges (a_r, d'_r) and (b'_r, c_r) in the alternating cycle $\rho = a_r - d'_r - c_r - b'_r - a_r$ are blocking edges to M. Consider $M \oplus \rho$ versus M. All the 4 nodes a_r, b'_r, c_r, d'_r prefer $M \oplus \rho$ to M while the other nodes are indifferent between $M \oplus \rho$ and M. Thus $\phi(M \oplus \rho, M) = 4$ and $\phi(M, M \oplus \rho) = 0$, so $u(M) = \infty$. This means M is not Pareto-optimal, a contradiction. Thus for any $r \in \{X_1, \ldots, X_{2n}\}$, we cannot have both (a_r, b'_r) and (c_r, d'_r) in M.

Theorem 11 is our main result in this section.

Theorem 11. G_{ψ} has a Pareto-optimal matching M with cost(M) = 0 if and only if ψ is satisfiable.

Proof. Suppose G_{ψ} has a Pareto-optimal matching M with cost(M) = 0. Lemma 10 indicates a natural way of defining an assignment for ψ . For any variable $r \in \{X_1, \ldots, X_{2n}\}$, consider the edges in $\neg r$'s gadget that are in M. If $(c_r, d'_r), (c'_r, d_r)$ are in M then set r = false else set r = true.

Lemma 10 tells us that when r = false, the edges $(a_{r,i}, b_{r,i}), (a'_{r,i}, b'_{r,i})$ from r's gadget in the *i*-th clause are in M (where $a_{r,i}, b_{r,i}, a'_{r,i}, b'_{r,i}$ are the 4 nodes from r's gadget in the *i*-th clause).

Claim 2 The above assignment satisfies ψ .

Claim 2 uses the Pareto-optimality of M to show that every clause has at least one literal set to true. Its proof is given after the proof of Theorem 11. Hence if G_{ψ} admits a Pareto-optimal matching M with cost(M) = 0, then ψ is satisfiable.

The converse. We will now show that if ψ is satisfiable then there is a Pareto-optimal matching M in G_{ψ} such that cost(M) = 0. There is a natural way of constructing the matching M—we will use the satisfying assignment for ψ to choose edges from each literal gadget. For any variable r, include the following edges in the matching M:

- if r = true then take the pair of edges $(c_r, d_r), (c'_r, d'_r)$ from $\neg r$'s gadget and the pair of edges $(a_{r,i}, b'_{r,i}), (a'_{r,i}, b_{r,i})$ from r's gadget in clause *i* (for every clause *i* that *r* belongs to).
- if r = false then take the pair of edges $(c_r, d'_r), (c'_r, d_r)$ from $\neg r$'s gadget and the pair of edges $(a_{r,i}, b_{r,i}), (a'_{r,i}, b'_{r,i})$ from r's gadget in clause *i* (for every clause *i* that *r* belongs to).

It is easy to see that cost(M) = 0. Since M is a perfect matching, there is no alternating path ρ wrt M such that $\phi(M, M \oplus \rho) = 0$. This is because for every alternating path ρ wrt M, we have $|M \oplus \rho| < |M|$ and the nodes matched in M and unmatched in $M \oplus \rho$ prefer M to $M \oplus \rho$, so $\phi(M, M \oplus \rho) > 0$. Hence in order to prove M's Pareto-optimality, what we need to show is Claim 3.

Claim 3 There is no alternating cycle ρ with respect to M such that $\phi(M \oplus \rho, M) > 0$ and $\phi(M, M \oplus \rho) = 0$.

The proof of Claim 3 is given below. This finishes the proof of Theorem 11.

Proof of Claim 2. Suppose this assignment does not satisfy ψ . We have 3 cases here.

1. Let $C_i = x \vee y \vee z$. Suppose all the three variables x, y, z are in false state. Consider the following alternating cycle ρ wrt M:

$$b_{z,i} - (a_{x,i}, b_{x,i}) - (a_{y,i}, b_{y,i}) - (a_{z,i}, b_{z,i}) - a_{x,i}.$$

All non-matching edges in this alternating cycle, i.e., the edges $(b_{z,i}, a_{x,i}), (b_{x,i}, a_{y,i}), (b_{y,i}, a_{z,i})$, are blocking edges with respect to M. In the $M \oplus \rho$ versus M comparison, these 6 nodes $a_{x,i}, b_{x,i}, a_{y,i}, b_{y,i}, a_{z,i}, b_{z,i}$ prefer $M \oplus \rho$ to M while all other nodes in G_{ψ} are indifferent between $M \oplus \rho$ and M. Thus we have $\phi(M \oplus \rho, M) = 6$ and $\phi(M, M \oplus \rho) = 0$. Hence $u(M) = \infty$, contradicting the Pareto-optimality of M.

2. Let $C_j = x \vee y$, i.e., this is a positive clause with 2 literals. Suppose both x and y are in false state. Consider the following alternating cycle ρ wrt M:

$$b_{y,j} - (a_{x,j}, b_{x,j}) - (a_{y,j}, b_{y,j}) - a_{x,j}$$

In the $M \oplus \rho$ versus M comparison, the 4 nodes $a_{x,j}, b_{x,j}, a_{y,j}, b_{y,j}$ prefer $M \oplus \rho$ to M while all the other nodes in G_{ψ} are indifferent between $M \oplus \rho$ and M. Thus $\phi(M \oplus \rho, M) = 4$ and $\phi(M, M \oplus \rho) = 0$. Hence $u(M) = \infty$, contradicting the Pareto-optimality of M.

3. Let $D_k = \neg x \lor \neg y$. Suppose both $\neg x$ and $\neg y$ are in false state. Consider the following alternating cycle ρ wrt M:

$$d_y - (c_x, d_x) - (c_y, d_y) - c_x.$$

In the $M \oplus \rho$ versus M comparison, the 4 nodes c_x, d_x, c_y, d_y prefer $M \oplus \rho$ to M while all the other nodes in G_{ψ} are indifferent between $M \oplus \rho$ and M. So $\phi(M \oplus \rho, M) = 4$ and $\phi(M, M \oplus \rho) = 0$. Hence $u(M) = \infty$, contradicting the Pareto-optimality of M.

Thus every clause in ψ has at least one literal in true state.

Proof of Claim 3. We need to show there is no alternating cycle ρ wrt M such that $\phi(M \oplus \rho, M) > 0$ and $\phi(M, M \oplus \rho) = 0$. Every non-matching edge in such an alternating cycle ρ has to be a blocking edge wrt M. First, we argue that every consistency edge is a non-blocking edge to M; say, this is a consistency edge corresponding to variable r in clause i. It follows from our construction of M that M contains either:

- 1. $(a_{r,i}, b'_{r,i}), (a'_{r,i}, b_{r,i})$ and $(c_r, d_r), (c'_r, d'_r)$ or 2. $(a_{r,i}, b_{r,i}), (a'_{r,i}, b'_{r,i})$ and $(c_r, d'_r), (c'_r, d_r)$.
- In case 1: the node d'_r prefers $M(d'_r) = c'_r$ to $a_{r,i}$ and the node c_r prefers $M(c_r) = d_r$ to $b'_{r,i}$.
- In case 2: the node $a_{r,i}$ prefers $M(a_{r,i}) = b_{r,i}$ to d'_r and the node $b'_{r,i}$ prefers $M(b'_{r,i}) = a'_{r,i}$ to c_r .

Thus in both cases, the consistency edges $(a_{r,i}, d'_r)$ and $(b'_{r,i}, c_r)$ are non-blocking edges to M. Let H be the subgraph of G_{ψ} obtained by preserving only edges in M and blocking edges wrt M. Thus no non-blocking edge (other than edges in M) is included in H – so no consistency edge belongs to H.

Since there are no consistency edges in H, any alternating cycle in H has to be contained within a single clause. We will now show there is no such cycle in H by using the fact that we constructed M using a satisfying assignment for ψ : thus every clause has at least one literal set to true.

Let $C = x \vee y \vee z$ and suppose $y = \text{true in } \psi$. Then (a_y, b_y) and (a_y, b_y) are in M, however the edge (a'_u, b'_u) is non-blocking wrt M and hence it is missing in H. Thus there is no alternating cycle in H that is contained within the clause C. Now consider a negative clause $D = \neg x \lor \neg y$ and suppose $x = \mathsf{false}$ in ψ . Then (c_x, d'_x) and (c'_x, d_x) are in M, however the edge (c'_x, d'_x) is non-blocking wrt M and it is missing in H. Thus there is no alternating cycle in H that is contained within the clause D.

Consider the 4 edges of any literal gadget (say, r) in G_{ψ} : if $(a_r, b_r) \in M$ then (a_r, b'_r) is a non-blocking edge wrt M and if $(a'_r, b_r) \in M$ then (a'_r, b'_r) is a non-blocking edge wrt M. Similarly, in the gadget of $\neg r$: if $(c_r, d_r) \in M$ then (c_r, d'_r) is a non-blocking edge wrt M and if $(c'_r, d_r) \in M$ then (c'_r, d'_r) is a non-blocking edge wrt M. Thus there is no alternating cycle wrt M in H. So there is no alternating cycle ρ in G_{ψ} such that $\phi(M \oplus \rho, M) > 0$ and $\phi(M, M \oplus \rho) = 0$.

Observe that any Pareto-optimal matching in G_{ψ} of cost 0 is a perfect matching. Thus Theorem 11 shows that the min-cost Pareto-optimal matching problem and the min-cost Pareto-optimal max-matching problem are NP-hard. Moreover, these problems are NP-hard to approximate to any multiplicative factor. Thus we have shown the following theorem.

Theorem 12. Given $G = (A \cup B, E)$ with strict preferences and edge costs in $\{0, 1\}$, it is NP-hard to compute a min-cost Pareto-optimal matching in G; moreover, it is NP-hard to approximate this within any multiplicative factor.

Acknowledgments. Thanks to Yuri Faenza for helpful comments that improved the presentation.

References

- 1. A. Abdulkadiroğlu and T. Sönmez. School choice: a mechanism design approach. American Economic Review, 93(3): 729-747, 2003.
- D. J. Abraham, K. Cechl´rová, D. F. Manlove, and K. Mehlhorn. Pareto optimality in house allocation problems. In the 15th International Symposium on Algorithms and Computation (ISAAC): 3-15, 2004.
- 3. T. Barnagarwala. 1172 private doctors sign up to treat patients at govt hospitals in Mumbai. Indian Express, May 11, 2020.
- 4. S. Baswana, P. P. Chakrabarti, S. Chandran, Y. Kanoria, and U. Patange. *Centralized admissions for engineering colleges in India.* INFORMS Journal on Applied Analytics: 49(5): 338-354, 2019.
- 5. P. Biro, D. F. Manlove, and S. Mittal. Size versus stability in the marriage problem. Theoretical Computer Science, 411:1828–1841, 2010.
- 6. Å. Cseh. Popular matchings. Trends in Computational Social Choice, Ulle Endriss (ed.), pp. 105-121, 2017.
- 7. Á. Cseh, Y. Faenza, T. Kavitha, and V. Powers. Understanding popular matchings via stable matchings. http: //arxiv.org/abs/1811.06897
- A. Cseh and T. Kavitha. Popular edges and dominant matchings. Mathematical Programming, 172(1):209–229, Nov 2018.
- 9. Y. Faenza and T. Kavitha. *Quasi-popular matchings, optimality, and extended formulations.* Proceedings of the 31st Annual ACM-SIAM Symposium on Discrete Algorithms (SODA), 325-344, 2020.
- Y. Faenza, T. Kavitha, V. Powers, and X. Zhang. *Popular matchings and limits to tractability*. Proceedings of the 31st Annual ACM-SIAM Symposium on Discrete Algorithms (SODA), 2790-2809, 2019.
- T. Feder. A new fixed point approach for stable networks and stable marriages. Journal of Computer and System Sciences, 45:233–284, 1992.
- 12. T. Feder. Network flow and 2-satisfiability. Algorithmica, 11:291–319, 1994.
- T. Fleiner. A fixed-point approach to stable matchings and some applications. Mathematics of Operations Research, 28(1): 103-126, 2003.
- 14. D. Gale and L. S. Shapley. *College admissions and the stability of marriage*. American Mathematical Monthly, 69:9–15, 1962.
- 15. D. Gale and M. Sotomayor. Some remarks on the stable matching problem. Discrete Applied Mathematics, 11: 223-232, 1985.
- 16. P. Gärdenfors. Match making: assignments based on bilateral preferences. Behavioural Science, 20:166–173, 1975.
- C.-C. Huang and T. Kavitha. Popular matchings in the stable marriage problem. Information and Computation, 222: 180–194, 2013.
- 18. R. W. Irving, P. Leather, and D. Gusfield. An efficient algorithm for the "optimal" stable marriage. Journal of the ACM, 34:532–543, 1987.
- 19. P. B. Jaiswal. Mumbai struggles to tackle increasing number of COVID-19 cases. The Week, May 06, 2020.
- T. Kavitha. A size-popularity tradeoff in the stable marriage problem. SIAM Journal on Computing, 43(1): 52–71, 2014.
- 21. T. Kavitha. *Popular half-integral matchings*. In the 43rd International Colloquium on Automata, Languages, and Programming (ICALP), 22:1–22:13, 2016.
- 22. T. Kavitha. *Popular matchings with one-sided bias.* In the 47th International Colloquium on Automata, Languages, and Programming (ICALP), 70:1–70:18, 2020.
- T. Kavitha, J. Mestre, and M. Nasre. *Popular mixed matchings*. Theoretical Computer Science, 412:2679–2690, 2011.
- 24. M. McCutchen. The least-unpopularity-factor and least-unpopularity-margin criteria for matching problems with one-sided preferences. In the 8th Latin American Symposium on Theoretical Informatics (LATIN), pages 593–604, 2008.
- U. G. Rothblum. Characterization of stable matchings as extreme points of a polytope. Mathematical Programming, 54:57–67, 1992.
- C.-P. Teo and J. Sethuraman. The geometry of fractional stable matchings and its applications. Mathematics of Operations Research, 23:874–891, 1998.
- E. G. Thurber. Concerning the maximum number of stable matchings in the stable marriage problem. Discrete Mathematics, 248(1-3): 195–219, 2002.
- 28. J. H. Vande Vate. Linear programming brings marital bliss. Operations Research Letters, 8(3):147–153, 1989.

Appendix

Missing details from the proof of Theorem 5. The proofs of properties 1-6 are given below.

Proof of property 1. The inclusion $S' \subseteq \bigcup_{i=0}^{n_0-1} (A_i \times B_i)$ follows from the definition of the sets B_i . Recall that for $1 \leq i \leq n_0 - 1$, B_i is the set of nodes b such that $(a_i, \tilde{b}) \in S$ for some $a \in A_i$. Also, B_0 contains all nodes b such that $(a_0, \tilde{b}) \in S$ for some $a \in A_0$. Thus $S' \subseteq \bigcup_{i=0}^{n_0-1} (A_i \times B_i)$.

The stability of S' restricted to each set $A_i \cup B_i$ is by \tilde{b} 's preference order in G^* . Recall that within subscript *i* neighbors, the order of preference for \tilde{b} in G^* is b's order of preference in G. Thus the stability of S in G^* implies the stability of S' restricted to $A_i \cup B_i$ for each *i*.

Proof of property 2. Let $a \in A_{i+1}$. Then $(a_{i+1}, d_{i+1}(a)) \notin S$. So it has to be the case that $(a_i, d_{i+1}(a)) \in S$. Recall that $d_{i+1}(a)$ is a_i 's least preferred neighbor in G^* . So a_i prefers \tilde{b} to its partner in S. Hence it follows from the stability of S in G^* that \tilde{b} prefers its partner in S (this is a subscript i node z_i) to a_i , i.e., b prefers z to a.

Since \tilde{b} prefers subscript i + 1 nodes to subscript i nodes, \tilde{b} prefers a_{i+1} to its partner z_i in S. It follows from the stability of S in G^* that a_{i+1} has to prefer its partner \tilde{w} in S to \tilde{b} , otherwise (a_{i+1}, \tilde{b}) would block S. Hence a prefers w to b. Thus $\mathsf{wt}_{S'}(a, b) = -2$.

Proof of property 3. Suppose $a \in A_i$ where $i \ge j + 2$ and $b \in B_j$. So the edge $(a_{j+1}, d_{j+2}(a)) \in S$. Since $d_{j+2}(a)$ is a_{j+1} 's least preferred neighbor in G^* , the stability of S implies that \tilde{b} prefers its partner in S to a_{j+1} . However $b \in B_j$ and so \tilde{b} 's partner in S is a subscript j node z_j . This contradicts \tilde{b} 's preference order that it prefers any subscript j + 1 neighbor to a subscript j neighbor. Thus there is no edge (a, b) in G with $a \in A_i$ and $b \in B_j$ where $i \ge j + 2$.

Proof of property 4. This follows from properties 1, 2, and 3 given above. Properties 2 and 3 tell us that there is no blocking edge in $A_i \times B_j$ where $i \ge j + 1$. Property 1 tells us that there is no blocking edge in $A_i \times B_j$ for any *i*. So any blocking edge to S' has to be in $A_i \times B_j$ where $i \le j - 1$.

Proof of property 5. This follows from the definitions of the sets A_0, \ldots, A_{n_0-2} and B_1, \ldots, B_{n_0-1} . For each $a \in A_i$ where $0 \le i \le n_0 - 2$: we have $(a_i, \tilde{b}) \in S$ for some $\tilde{b} \in \tilde{B}$ and thus $(a, b) \in S'$. Similarly, for each $b \in B_j$ where $1 \le j \le n_0 - 1$: we have $(a_j, \tilde{b}) \in S$ for some $a \in A_j$ and thus $(a, b) \in S'$. Hence all nodes that are unmatched in S' are in $A_{n_0-1} \cup B_0$.

Proof of property 6. Suppose S' is not a maximum matching in G. Then there is an augmenting path ρ with respect to S'. Let us refer to an edge e that satisfies $\operatorname{wt}_{S'}(e) = -2$ as a negative edge. The endpoints of a negative edge prefer their respective assignments in S' to each other. We know from property 5 above that all nodes in A that are unmatched in S' are in A_{n_0-1} and all nodes in B that are unmatched in S' are in B_0 . We also know that $S' \subseteq \bigcup_i (A_i \times B_i)$ (by prop. 1 above). Moreover, all edges e in $A_{j+1} \times B_j$ are negative edges (by prop. 2 above) and there is no edge in $A_i \times B_j$ where $i \ge j+2$ (by prop. 3 above).

Thus the path ρ starts in A_{n_0-1} at an unmatched node *a* and since there cannot be any negative edge incident to an unmatched node, all of *a*'s neighbors have to be in B_{n_0-1} : this is because every edge *e* in $A_{n_0-1} \times B_{n_0-2}$ is a negative edge. The matched partners of *a*'s neighbors are in A_{n_0-1} . Then the next node can be in B_{n_0-2} (this is by prop. 3 above) and its partner is in A_{n_0-2} and so on. Finally, there is no edge from A_1 to an unmatched node in B_0 : this is because there is no negative edge incident to an unmatched node and we know all edges in $A_1 \times B_0$ are negative edges (by prop. 2 above).

So the *shortest* alternating path ρ from an unmatched $a \in A_{n_0-1}$ to an unmatched $b \in B_0$ moves across sets as follows: $A_{n_0-1} - B_{n_0-1} - A_{n_0-1} - B_{n_0-2} - A_{n_0-2} - B_{n_0-3} - \cdots - A_1 - B_0 - A_0 - B_0$. This implies there are at least $n_0 + 1$ nodes in A. However $|A| = n_0$. So there is no such alternating path, i.e., there is no augmenting path with respect to S'. In other words, S' is a maximum matching in G.

Proof of Claim 1. Let v_0 be the first node that moves to A_{n_0-1} with a blocking edge incident to it. Recall our update procedure – we initially added nodes in Nbr (U_A) and their partners in M to B_{n_0-1} and A_{n_0-1} , respectively. Then we applied rules 1-3 in some order and this resulted in the node v_0 moving to A_{n_0-1} . Corresponding to these rules, we will construct an alternating path $p = v_0 - M(v_0) - v_1 - M(v_1) - v_2 - \cdots - v_k - M(v_k) - u$ between v_0 and some node $u \in U_A$. A useful observation is that all the nodes v_0, \ldots, v_k have moved upwards to their respective levels due to our update of $\vec{\alpha}$ to ensure property (ii) and to maintain $\vec{\alpha}$ as an optimal solution to (LP4).

The path p can be partitioned into k + 1 pairs of edges for some $k \ge 0$. For $0 \le i \le k - 1$: the *i*-th pair consists of the matching edge $(v_i, M(v_i))$ of weight 0 and the non-matching edge $e_i = (M(v_i), v_{i+1})$ where v_{i+1} is the node that pulled $M(v_i)$ and v_i to their current level due to the application of one of the above three rules. Rule 1 implies wt_M $(e_i) = 0$ while rule 2 implies wt_M $(e_i) = -2$ and rule 3 implies wt_M $(e_i) = 2$.

Observe that rule 1 places $M(v_i)$ in the same level as v_{i+1} while rule 2 places $M(v_i)$ one level lower than v_{i+1} and rule 3 places $M(v_i)$ one level higher than v_{i+1} . The last pair of edges in p are $(v_k, M(v_k)) \in A_{n_0-1} \times B_{n_0-1}$ and $(M(v_k), u)$, where the latter edge has weight 0. The node v_0 is in level $n_0 - 1$ and the node $M(v_k)$ is also in level $n_0 - 1$. So v_0 and v_k are at the same level, hence the number of edges in p of weight -2 is exactly the same as the number of edges of weight 2, thus wt_M(p) = 0.

Suppose there is a blocking edge (v_0, w) . If w belongs to p, then it is easy to see that the alternating cycle C obtained by joining the endpoints of the v_0 -w subpath in p with the edge (v_0, w) satisfies $wt_M(C) \ge 2$. This contradicts Theorem 4 since M is a popular max-matching. Hence w does not belong to path p. So let us add the 2-edge path $M(w) - w - v_0$ as a prefix to the v_0 -u path p and call this alternating path q: we have $wt_M(q) = wt_M(p) + wt_M(v_0, w) = 2$. Since $wt_M(q) > 0$ and the unmatched node u is an endpoint of q, this again contradicts Theorem 4. Hence there is no blocking edge incident to v_0 .