# ON THE HARDNESS OF CODE EQUIVALENCE PROBLEMS IN RANK METRIC

by

Alain Couvreur, Thomas Debris–Alazard & Philippe Gaborit

**Abstract.** — In the recent years, the notion of rank metric in the context of coding theory has known many interesting developments in terms of applications such as space time coding, network coding or public key cryptography. These applications raised the interest of the community for theoretical properties of this type of codes, such as the hardness of decoding in rank metric. Among classical problems associated to codes for a given metric, the notion of code equivalence (to decide if two codes are isometric) has always been of the greatest interest, for its cryptographic applications or its deep connexions to the graph isomorphism problem.

In this article, we discuss the hardness of the code equivalence problem in rank metric for  $\mathbb{F}_{q^m}$ -linear and general rank metric codes. In the  $\mathbb{F}_{q^m}$ -linear case, we reduce the underlying problem to another one called *Matrix Codes Right Equivalence Problem*. We prove the latter problem to be either in  $\mathcal{P}$  or in  $\mathcal{ZPP}$  depending of the ground field size. This is obtained by designing an algorithm whose principal routines are linear algebra and factoring polynomials over finite fields. It turns out that the most difficult instances involve codes with non trivial *stabilizer algebras*. The resolution of the latter case will involve tools related to finite dimensional algebras and Wedderburn-Artin theory. It is interesting to note that 30 years ago, an important trend in theoretical computer science consisted to design algorithms making effective major results of this theory. These algorithmic results turn out to be particularly useful in the present article.

Finally, for general matrix codes, we prove that the equivalence problem (both left and right) is at least as hard as the well–studied *Monomial Equivalence Problem* for codes endowed with the Hamming metric.

#### Contents

Introduction	2
1. Basic objects and tools	3
2. About the code equivalence problem	4
3. Main results	7
4. Solving the Matrix Codes Right Equivalence Problem (MCRE)	11
5. The Code Equivalence Problem for $\mathbb{F}_{q^m}$ -linear codes : an Easy Problem	16
6. Reduction from the Code Equivalence Problem in Hamming Metric	20
Conclusion	22
References	23

### Introduction

The code equivalence problem. — Given two codes over a finite field, the Code Equivalence problem (CE) asks if they are isometric when embedded with the Hamming metric, namely if they are image of each other by a permutation. Code equivalence is a longstanding problem in computer science with many applications to cryptography and strong connections with theoretical computer science problems such as the graph isomorphism problem. In particular, it has been proved in [**PR97**] that the graph isomorphism problem reduces to the permutation equivalence problem. Therefore any solver of CE in polynomial time would show that the graph isomorphism problem lies in  $\mathcal{P}$ , statement for which we do not know if it is true or not despite many efforts. This result tends to show that we cannot hope to get a polynomial time algorithm for solving the code equivalence problem. On the other hand, [**PR97**] also proved that this problem is not  $\mathcal{NP}$ -complete unless the polynomial-time hierarchy collapses. Therefore CE lies in an intermediary situation for which its hardness is not clear.

In another line of works, algorithms have been considered to solve the code equivalence problem. The first one, proposed by Leon [Leo82], computes a list of minimum weight codewords of both codes and try to match them (usually with graph techniques) to recover the permutation. However such approach leads to an exponential time algorithm in average over the inputs. Latter Sendrier [Sen00] proposed a new approach based on the computation of the hulls (the intersection of a code and its dual) of both codes. Interestingly enough Sendrier gave an algorithm (when codes are over the binary field) whose "practical" complexity is exponential in the dimension of the hull but since their average dimension for random codes is very low [Sen97] it shows that, in average (and not in the worst case), that for binary codes (and also over  $\mathbb{F}_3$  and  $\mathbb{F}_4$ ), equivalence is easy to decide. On the other hand, the code equivalence problem remains difficult for  $q \ge 5$  [SS13]. We emphasize that the efficiency of the aforementioned algorithms rests on heuristics. They may require an exponential time in some rare worst cases.

Our contribution in this article is to look at the difficulty of the code equivalence problem but when codes are embedded with the rank metric instead of the Hamming weight. Note that these kind of equivalence problem exists also with the Euclidean metric where there is a notion of lattice isomorphism. It has been studied by Haviv and Regev in [HR14].

**Rank metric and its applications.** — Besides the well known notions of Hamming distance for error-correcting codes and Euclidean distance for lattices, there is also the concept of rank metric which was introduced in 1951 by Loo-Keng Hu [Hua51] as "arithmetic distance" for matrices over a field  $\mathbb{F}_q$ . Given two  $n \times n$  matrices **A** and **B** over a finite field  $\mathbb{F}_q$ , the rank distance between **A** and **B** is defined as  $|\mathbf{A} - \mathbf{B}| = \text{Rank}(\mathbf{A} - \mathbf{B})$ . Later, in 1978, Delsarte defined [Del78] the notion of rank distance on the space of bilinear forms and proposed a construction of optimal matrix codes in bilinear form representation. A matrix code over  $\mathbb{F}_q$  is defined as an  $\mathbb{F}_q$ -linear subspace of the space of  $m \times n$  matrices over  $\mathbb{F}_q$  endowed with the rank metric. Later, in 1985, Gabidulin introduced in [Gab85] the notion of rank metric codes in vector representation (as opposed to the matrix representation) over a finite extension field  $\mathbb{F}_{q^m}$  of  $\mathbb{F}_q$ . These codes are known as  $\mathbb{F}_{q^m}$ -linear codes and they form a particular subclass of matrix codes. In the same paper, Gabidulin also introduced an optimal class of  $\mathbb{F}_{q^m}$ -linear codes: the so-called Gabidulin codes, which can be regarded as analogues of Reed-Solomon codes but in a rank metric context, where polynomials are replaced by so-called *linearized* polynomials as introduced by Ore in 1933 in [Ore33]. Interestingly, Gabidulin codes are almost the only matrix codes for which we known an efficient decoding algorithm. All matrix codes with an efficient decoding algorithm are  $\mathbb{F}_{q^m}$ -linear **[Gab85, GMRZ13]** (there is also simple codes **[SKK10]** which are not  $\mathbb{F}_{q^m}$ -linear but they have a trivial structure).

The rank metric and the code equivalence problem. — The notion of code equivalence with the rank metric has been introduced by Berger in [Ber03] (see also [Mor14]) and invariants with respect to this code equivalence are considered in [NPHT20]. However, contrary to the Hamming metric case, to our knowledge, neither the algorithmic resolution of the code equivalence

problem in rank metric nor its theoretical hardness have ever been discussed in the literature. This is the purpose of the present paper.

**Our contributions.** — In this article, we discuss the code equivalence problems in rank metric from a theoretical and algorithmic perspective. Our contributions in this article are three–fold

- 1°) We show that the right equivalence problem is easy *in the worst case*. Interestingly, this proof involves many theoretical and algorithmic developments from Wedderburn-Artin theory,
- 2°) The  $\mathbb{F}_{q^m}$ -linear case is proved to be reduced to the previous case and hence is easy in the worst case,
- 3°) Finally, the general case is proved to be harder than the code equivalence problem in Hamming metric by providing a polynomial time reduction.

It is striking to observe that in rank metric we can get worst case polynomial-time algorithms (possibly Las Vegas) to solve some set of sub-instances of the equivalence problem (points 1 and 2) while no such Hamming counterpart seems reachable. In addition these sub-instances are the most considered one in the rank metric based literature.

**Outline of the article.** — The present article is organised as follows. Section 1 recalls main objects and tools that we consider. Section 2 gives background on the code equivalence problem in Hamming metric and established various code equivalence problems in rank metric. Our main results are stated in Section 3 and their proofs are sketched; detailed on proofs are given in the following sections. In Section 4, we show how to solve the *right equivalence problem*. Section 5 considers the code equivalence problem for  $\mathbb{F}_{q^m}$ -linear codes and finally, in Section 6, we present a polynomial-time reduction from the code equivalence problem in Hamming metric to the (both left and right) equivalence problem in rank metric.

Acknowledgements. — The authors express their deep gratitude to Hugues Randriambololona for his very relevant remarks and to Xavier Caruso for pointing out references from representation theory.

#### 1. Basic objects and tools

Before explaining the main ideas behind our results, we need to recall basic objects and tools that we consider. In all the article, we will consider two different kinds of objects which we will refer to *codes*. Namely:

- vector codes are usual codes, *i.e.* vector subspaces of  $\mathbb{F}_q^n$ . Such codes are usually endowed with the Hamming metric. These codes are denoted with calligraphic letter and with the subscript "Vec" such as  $\mathscr{C}_{\text{Vec}}$ ;
- matrix codes, are subspaces of  $\mathcal{M}_{m,n}(\mathbb{F}_q)$   $(m \times n \text{ matrices whose coefficients belong to } \mathbb{F}_q)$ endowed with the rank metric. They are denoted as  $\mathscr{C}_{Mat}$ .

Nevertheless, there is class of vector codes that we can equip with the rank distance. There are subspaces of  $\mathbb{F}_{q^m}^n$ . Such codes are referred to as rank metric  $\mathbb{F}_{q^m}$ -linear codes and there are a particular class of matrix codes which turn out to be the most commonly studied in the literature. Let us describe them more precisely. Given a vector  $\mathbf{v} \in \mathbb{F}_{q^m}^n$  its rank weight is defined as

(1) 
$$|\mathbf{v}| \stackrel{def}{=} \operatorname{Span}_{\mathbb{F}_q} \left\{ \lambda_1 v_1 + \dots + \lambda_n v_n : \lambda_i \in \mathbb{F}_q \right\}.$$

In other words,  $|\mathbf{v}|$  is the dimension of the  $\mathbb{F}_q$ -linear subspace of  $\mathbb{F}_{q^m}$  spanned by the entries of  $\mathbf{v}$ . It is well-known that any  $\mathbb{F}_{q^m}$ -linear code is isometric to a matrix code as follows. Choose an  $\mathbb{F}_q$ -basis  $\mathcal{B} = (b_1, \ldots, b_m)$  of  $\mathbb{F}_{q^m}$ , then consider the map

(2) 
$$M_{\mathcal{B}}: \begin{cases} \mathbb{F}_{q^m}^n \longrightarrow \mathcal{M}_{m,n}(\mathbb{F}_q) \\ (v_1, \dots, v_n) \longmapsto \begin{pmatrix} v_{11} \cdots v_{1n} \\ \vdots & \vdots \\ v_{m1} \cdots v_{mn} \end{pmatrix}$$

where for any  $j, v_{1j}, \ldots, v_{mj} \in \mathbb{F}_q$  denote the coefficients of  $v_j$  in the basis  $\mathcal{B}$ . That is :  $v_j = v_{1j}b_1 + \cdots + v_{mj}b_m$ . One can easily prove that for any choice of  $\mathbb{F}_q$ -basis of  $\mathbb{F}_{q^m}$ , the above map preserves the metric.

An  $\mathbb{F}_{q^m}$ -linear code of dimension k is through  $M_{\mathcal{B}}$  a km-dimensional matrix code of  $\mathcal{M}_{m,n}(\mathbb{F}_q)$ . However, it requires only k vectors in  $\mathbb{F}_{q^m}^n$  or equivalently k matrices in  $\mathcal{M}_{m,n}(\mathbb{F}_q)$  to be represented while a general km-dimensional matrix code of  $\mathcal{M}_{m,n}(\mathbb{F}_q)$  is represented by a basis of km matrices of size  $m \times n$ . Thus, the representation of an  $\mathbb{F}_{q^m}$ -linear code requires m times less memory size compared to that of a general matrix code of the same  $\mathbb{F}_q$ -dimension. This gain is particularly interesting for cryptographic applications. This is basically what explains why in general McEliece cryptosystems based on rank metric matrix codes have a smaller key size than McEliece cryptosystems based on the Hamming metric. All of these proposals (see for instance [GPT91, GO01, Gab08, GMRZ13, GRSZ14, ABD+19, AAB+17, ABG+19]) are actually built from matrix codes over  $\mathbb{F}_q$  obtained from  $\mathbb{F}_{q^m}$ -linear codes.

On the other hand, vector codes which are  $\mathbb{F}_{q^m}$ -linear can be viewed as structured matrix codes with some "extra" algebraic structure in the same way as, for instance, cyclic linear codes can be viewed as structured versions of linear codes. In the latter case, the code is globally invariant by a linear isometric transform on the codewords corresponding to shifts of a certain length.

Let  $P = \sum_{i=0}^{m-1} a_i X^i + X^m \in \mathbb{F}_q[X]$  be a monic irreducible polynomial of degree m and  $x \in \mathbb{F}_{q^m}$ be a root of P. Then  $\mathcal{B} \stackrel{def}{=} (1, x, \dots, x^{m-1})$  is an  $\mathbb{F}_q$ -basis of  $\mathbb{F}_{q^m}$ . Let  $\mathscr{C}_{\text{Vec}}$  be an  $\mathbb{F}_{q^m}$ -linear code. The  $\mathbb{F}_{q^m}$ -linearity with the definition of  $\mathcal{B}$  means that:

$$\forall \mathbf{c} \in \mathscr{C}_{\mathrm{Vec}}, \forall Q \in \mathbb{F}_q[X], \ Q(x) \cdot \mathbf{c} \in \mathscr{C}_{\mathrm{Vec}}.$$

In terms of matrices this stability can be expressed as follows. It is readily seen that (see (2) for the meaning of  $M_{\mathcal{B}}(\cdot)$ )

$$M_{\mathcal{B}}(P(x)\mathbf{c}) = P(\mathbf{C}_x)M_{\mathcal{B}}(\mathbf{c}) \quad \text{where} \quad \mathbf{C}_x \stackrel{def}{=} \begin{pmatrix} 0 & 0 & \cdots & \cdots & -a_0 \\ 1 & \ddots & \cdots & \cdots & -a_1 \\ 0 & \ddots & \ddots & & \vdots \\ \vdots & & \ddots & \ddots & & \vdots \\ 0 & \cdots & \cdots & 1 & -a_{m-1} \end{pmatrix} \in \mathbb{F}_q^{m \times m}$$

*i.e.*  $\mathbf{C}_x$  is the *companion matrix* of P which represents the multiplication by x in the basis  $\mathcal{B}$ . In other words, the matrix code  $M_{\mathcal{B}}(\mathscr{C}_{\text{Vec}}) \stackrel{def}{=} \{M_{\mathcal{B}}(\mathbf{c}) : \mathbf{c} \in \mathscr{C}_{\text{Vec}}\}$  associated to  $\mathscr{C}_{\text{Vec}}$  is stable by left multiplication by an element of the algebra generated by the companion matrix  $\mathbf{C}_x$  of x.

This property of  $\mathbb{F}_{q^m}$ -linear codes make them as special a case of matrix codes: they have a *large left stabilizer algebra* (see Definition 9). As we will see in Section 5, the easiness of code equivalence problems for  $\mathbb{F}_{q^m}$ -linear codes comes, roughly speaking, from the fact that they have a particular stabilizer algebra.

### 2. About the code equivalence problem

We arrive now to the core of our paper: the code equivalence problem. We first recall here the definition of this problem (and some of its variants) for vector codes endowed with the Hamming metric.

## 2.1. In Hamming Metric. —

**2.1.1.** Statement of the problems. — In Hamming metric, the group of linear isometries of  $\mathbb{F}_q^n$  is the subgroup of  $\mathbf{GL}_n(\mathbb{F}_q)$  (invertible  $n \times n$  matrices over  $\mathbb{F}_q$ ) spanned by permutation matrices and nonsingular diagonal matrices. Any element of this group can be represented as a product **DP** where **D** is diagonal and **P** is a permutation matrix. Such a matrix is usually called a *monomial matrix*. It is a matrix having exactly one non-zero entry per row and per column.

In Hamming metric, one usually considers two code equivalence problems. Given two codes endowed with the Hamming metric, there are two natural questions:

- are they permutation equivalent (image by each other under a permutation)?
- are they monomially equivalent (image by each other under a monomial matrix)?

Some comments. –

- 1. Note that we only consider linear isometries here. Some references in the literature discuss semi-linear isometries, *i.e.* the composition of a monomial transform and some iteration of a component wise Frobenius map (assuming that the ground field  $\mathbb{F}_q$  is not prime). This would lead to an alternative problem but that can be solved from any solver of the monomial equivalence problem in essentially the same time. Indeed if one can solve this problem in polynomial time, then it is enough to brute-force any iterate of the Frobenius.
- 2. Note that MacWilliams equivalence theorem [Mac62, HP03] asserts that if there is a linear map  $\mathscr{C}_{\text{Vec}} \rightarrow \mathscr{D}_{\text{Vec}}$  which is an isometry, then it extends to the whole ambient space. Therefore, solving the monomial equivalence problem is equivalent to decide whether two codes are image of each other by a linear isometry with respect to the Hamming distance.
- **2.1.2.** Algorithms. Several algorithms appeared in the literature to solve these problems.
  - The first work is probably due to Leon [Leo82]. His algorithm computes the full list of minimum weight codewords of both codes, which requires an exponential running time for almost any code.
  - Later, Sendrier [Sen00] introduced the so-called support-splitting algorithm to decide if two codes  $\mathscr{C}_{\text{Vec}}$  and  $\mathscr{D}_{\text{Vec}}$  are permutation equivalent. Its approach consists in comparing the weight enumerators of the hulls of the codes, *i.e.* the codes  $\mathscr{C}_{\text{Vec}} \cap \mathscr{C}_{\text{Vec}}^{\perp}$  and  $\mathscr{D}_{\text{Vec}} \cap \mathscr{D}_{\text{Vec}}^{\perp}$ . The cost of this algorithm is heuristically  $O(2^{\dim \mathscr{C}_{\text{Vec}}} \cap \mathscr{C}_{\text{Vec}}^{\perp} n^{\omega})$  where  $\omega$  denotes the complexity exponent of linear algebraic operations. In particular, the algorithm solves easily the problem if the hulls of the codes have small dimensions, which typically holds [Sen97]. However Sendrier's algorithm does not extend to solve the monomial equivalence unless q = 3and q = 4 if one replaces the hull by its counterpart with respect to the Hermitian inner product.
  - In [Feu09] is proposed a notion of normal form of a class of codes under the action of the monomial group. No mention of complexity appears in this reference but the computation of this normal form seems to require an exponential running time. In addition, a significant speed-up is possible for some input codes using tree–cutting dynamical programming methods.
  - More recently, [ST17, §3.2.4] proposed to decide if two codes are permutation equivalent to solve a polynomial system with Gröbner bases techniques. Actually this approach can be extended to the resolution of the monomial equivalence. However, with this use of Gröbner bases, the average-time complexity analysis is unknown.

**2.1.3.** Theoretical hardness and known reductions. — From a more theoretical point of view, what can we say about the hardness of these problems? Their hardness is not so clear, here is what is known:

- It has been proved in [**PR97**] that the graph isomorphism problem reduces polynomially to the permutation equivalence problem. It is also proved in the same reference that this problem is not  $\mathcal{NP}$ -complete unless the polynomial-time hierarchy collapses;
- Recently, it was proved [**BOS19**] that deciding if two codes with zero hull are permutation equivalent reduces to the graph isomorphism problem;
- Finally, from [SS13], the monomial equivalence problem reduces polynomially to the permutation equivalence problem when the field cardinality q is polynomial in n. However, it should be noticed that the reduction sends any instance into an instance with a large hull, hence in the set of instances which seem to be the hardest ones.

The reductions are summarized in Figure 1.

**2.2.** In Rank Metric. — In rank metric, the following linear automorphisms of  $\mathcal{M}_{m,n}(\mathbb{F}_q)$  are well-known to preserve the rank:

- left multiplication by an element  $\mathbf{P}$  of  $\mathbf{GL}_m(\mathbb{F}_q)$  :  $\mathbf{X} \mapsto \mathbf{PX}$ ;



FIGURE 1. Reductions between various problems. The notation " $A \longrightarrow B$ " means that "Problem A reduces to Problem B in polynomial time". It can be translated as "If one can solve Problem B in polynomial time, then one can solve Problem A in polynomial time.

- right multiplication by an element  $\mathbf{Q}$  of  $\mathbf{GL}_n(\mathbb{F}_q) : \mathbf{X} \mapsto \mathbf{XQ};$
- the transposition map  $\mathbf{X} \mapsto \mathbf{X}^{\top}$  (only when m = n).

A classical result [Hua51] of linear algebra asserts that any rank-preserving linear automorphism of  $\mathcal{M}_{m,n}(\mathbb{F}_q)$  is a composition of these three kinds of maps.

**Definition 1.** — Two matrix codes  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  are said to be *equivalent* if there are matrices  $\mathbf{P} \in \mathbf{GL}_m(\mathbb{F}_q)$  and  $\mathbf{Q} \in \mathbf{GL}_n(\mathbb{F}_q)$  such that  $\mathscr{C}_{Mat} = \mathbf{P}\mathscr{D}_{Mat}\mathbf{Q}$ . When  $\mathbf{P} = \mathbf{I}_m$  (resp.  $\mathbf{Q} = \mathbf{I}_m$ ) codes are said right (resp. left) equivalent.

**Remark 2.** — Note that the previous definition does not involve the possibility of a transposition. Hence two square matrix codes  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_n(\mathbb{F}_q)$  are *equivalent* according to our definition if and only if  $\mathscr{C}_{Mat}$  is the image by a rank preserving linear map of either  $\mathscr{D}_{Mat}$  or  $\mathscr{D}_{Mat}^{\top}$ . In practice, if we benefit from an algorithm which decides whether two square matrix codes are equivalent (according to our definition), then by applying the algorithm successively on the pairs  $(\mathscr{C}_{Mat}, \mathscr{D}_{Mat})$  and  $(\mathscr{C}_{Mat}, \mathscr{D}_{Mat}^{\top})$ , we can decide whether the codes are image of each other by a rank preserving automorphism. In short, considering a possible transposition will only multiply by 2 the algorithm complexity.

**Remark 3.** — Contrary to the Hamming case there does not seem to exist a rank metric McWilliams equivalence theorem [**BG13**, Ex. 2.9]. Two codes may be image of each other by a linear isometry without being equivalent.

In this way, the code equivalence problem for matrix codes endowed with the rank metric can be stated as follows.

# Problem 1 (Matrix Code Equivalence Problem (MCE)). —

- Instance: two matrix codes  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q);$
- Decision: there exist matrices  $\mathbf{P} \in \mathbf{GL}_m(\mathbb{F}_q)$  and  $\mathbf{Q} \in \mathbf{GL}_m(\mathbb{F}_q)$  such that

$$\mathscr{C}_{Mat} = \mathbf{P}\mathscr{D}_{Mat}\mathbf{Q}.$$

In addition to this problem, one can be interested in the equivalence of the most commonly used rank metric codes, namely  $\mathbb{F}_{q^m}$ -linear codes. If codes are represented with vectors, there is no longer equivalence by left multiplying by a non singular matrix but right multiplication is still possible. Therefore, when regarding vector codes as matrix codes, using the expansion operation  $M_{\mathcal{B}}(\cdot)$  defined in (2), equivalence of  $\mathbb{F}_{q^m}$ -linear codes can be regarded as a restriction of the following problem to instances corresponding to matrix representations of  $\mathbb{F}_{q^m}$ -linear codes.

### Problem 2 (Matrix Codes Right Equivalence Problem (MCRE))

- Instance: two matrix codes  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q);$
- Decision: there exists a matrix  $\mathbf{Q} \in \mathbf{GL}_m(\mathbb{F}_q)$  such that

$$\mathscr{C}_{Mat} = \mathscr{D}_{Mat} \mathbf{Q}.$$

Finally, let us introduce a last problem which could be more of cryptographic nature. Suppose given two vector codes  $\mathscr{C}_{\text{Vec}}$ ,  $\mathscr{D}_{\text{Vec}} \subseteq \mathbb{F}_{q^m}^n$  and two  $\mathbb{F}_q$ -bases  $\mathcal{B}, \mathcal{B}'$  of  $\mathbb{F}_{q^m}$  and consider the matrix codes  $M_{\mathcal{B}}(\mathscr{C}_{\text{Vec}})$  and  $M_{\mathcal{B}'}(\mathscr{D}_{\text{Vec}})$ . From the data of these matrix codes and without knowing the bases  $\mathcal{B}, \mathcal{B}'$ , is deciding equivalence easier? This leads to the following problem which is nothing but a restriction of MCE to the subset of instances of matrix codes arising from  $\mathbb{F}_{q^m}$ -linear codes.

### Problem 3 (Hidden Vector Matrix Code Equivalence Problem (HV-MCE))

- Instance:  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  be two spaces of matrices representing  $\mathbb{F}_{q^m}$ -linear codes
- Decision: It exists  $(\mathbf{S}, \mathbf{P}) \in \mathbf{GL}_k(\mathbb{F}_q) \times \mathbf{GL}_n(\mathbb{F}_q)$  such that

$$\mathscr{C}_{Mat} = \mathbf{S}\mathscr{D}_{Mat}\mathbf{P}.$$

# 3. Main results

Our contributions in this article are three-fold

- 1°) We propose a polynomial time algorithm to solve MCRE. This algorithm is deterministic if q is polynomial in mn and Las Vegas for a larger q.
- $2^{\circ}$ ) We prove that HV-MCE is easier than MCE and in particular that it naturally reduces to MCRE in polynomial time when q is polynomial in mn and via a Las Vegas algorithm for larger q.
- 3°) Finally, we prove that the general problem MCE is at least as hard as ME in Hamming metric by providing a polynomial time reduction.

This leads to the following statements.

**Theorem 4.** — MCRE and HV-MCE are in  $\mathcal{P}$  if  $q = mn^{O(1)}$  and in  $\mathcal{ZPP}$  in the general case.

**Theorem 5.** — The monomial equivalence problem reduces in polynomial time to MCE.

The why of  $\mathcal{P}$  v.s.  $\mathcal{ZPP}$ . — In the sequel, we first propose an algorithm to solve MCRE. Then, we give a second algorithm which proves that HV-MCE reduces to MCRE. The major tools of these algorithms are the resolution of linear systems and factorization of univariate polynomials over a finite field. Linear systems are solvable in polynomial time while for factorization of univariate polynomials, Berlekamp algorithm [Ber68] is deterministic polynomial only when the ground field cardinality q is polynomial in the degree. For larger q, one should use a Las Vegas algorithm such as Cantor Zassenhaus algorithm [Ber70, CZ81]. This is the reason why, for a large q, we cannot assert that the problem is in  $\mathcal{P}$  but it is in  $\mathcal{ZPP}$ .

In order to avoid statements making every time this difference between small and large q we introduce the notion of F-algorithm.

**Definition 6.** — An F-algorithm is an algorithm which uses an oracle (subroutine) to factor polynomials over finite fields. The cost of a call of this oracle is the length of the input of the call.

In the sequel we sketch the proofs of these theorems. The detailed proofs appear in the following sections as follows. In Section 4 we show how to solve MCRE efficiently. Section 5 gives an efficient algorithm which reduces HV-MCE to MCRE and therefore proves Theorem 4. Finally, Section 6 is devoted to our reduction, namely the proof of Theorem 5.

**3.1.** A polynomial-time algorithm to solve MCRE. — The resolution of this problem starts with the computation of an object called the *conductor* of  $\mathscr{C}_{Mat}$  into  $\mathscr{D}_{Mat}$  and

**Definition 7** (Conductor of matrix codes). — Let  $\mathscr{C}_{Mat}$ ,  $\mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  and suppose they have the same  $\mathbb{F}_q$ -dimension<sup>(1)</sup> and define the *conductor of*  $\mathscr{C}_{Mat}$  *into*  $\mathscr{D}_{Mat}$  as the  $n \times n$ matrix space:

$$\mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}},\mathscr{D}_{\mathrm{Mat}}) \stackrel{def}{=} \{ \mathbf{M} \in \mathcal{M}_n(\mathbb{F}_q) \mid \mathscr{C}_{\mathrm{Mat}} \mathbf{M} \subseteq \mathscr{D}_{\mathrm{Mat}} \}.$$

**Proposition 8.** — If  $\mathscr{C}_{Mat}\mathbf{Q} = \mathscr{D}_{Mat}$  for some  $\mathbf{Q} \in \mathbf{GL}_n(\mathbb{F}_q)$ , that is to say, if  $\mathscr{C}_{Mat}$  and  $\mathscr{D}_{Mat}$  are right equivalent, then  $\mathbf{Q} \in \mathbf{Cond}(\mathscr{C}_{Mat}, \mathscr{D}_{Mat})$ . Conversely, if  $\mathbf{Cond}(\mathscr{C}_{Mat}, \mathscr{D}_{Mat})$  contains a nonsingular matrix  $\mathbf{Q}$ , then  $\mathscr{C}_{Mat}$  and  $\mathscr{D}_{Mat}$  are right equivalent.

The conductor can be computed by solving a linear system with  $n^2$  unknowns and K(mn - K) equations, where  $K = \dim(\mathscr{C}_{Mat}) = \dim(\mathscr{D}_{Mat})$ . Indeed, our unknowns are matrices  $\mathbf{M} \in \mathbf{Cond}(\mathscr{C}_{Mat}, \mathscr{D}_{Mat})$  whose entries provide  $n^2$  unknowns in  $\mathbb{F}_q$ . For the equations, let  $\mathbf{C}_1, \ldots, \mathbf{C}_K$  be a basis of  $\mathscr{C}_{Mat}$  and  $\mathbf{D}_1, \ldots, \mathbf{D}_{mn-K}$  be a basis of the space

 $\{\mathbf{M} \in \mathcal{M}_{m,n}(\mathbb{F}_q) \mid \forall \mathbf{N} \in \mathscr{D}_{\mathrm{Mat}}, \ \mathrm{Tr}(\mathbf{M}\mathbf{N}^{\top}) = 0\}.$ 

Then, our equations with unknown  $\mathbf{M}$  are

$$\operatorname{Tr}(\mathbf{C}_{i}\mathbf{M}\mathbf{D}_{i}^{\top}) = 0, \quad i \in \{1, \dots, K\}, \ j \in \{1, \dots, N-K\}.$$

In many classical situations,  $m \approx n$ , K is linear in mn and then the number of equations is of  $O(n^4)$  while that of variables is of  $O(n^2)$ . Thus, the system is over constraint and one can expect that one of the two following situations appear:

- either  $\operatorname{Cond}(\mathscr{C}_{\operatorname{Mat}}, \mathscr{D}_{\operatorname{Mat}}) = \{0\}$  and one can assert that the codes are not right equivalent; - or dim( $\operatorname{Cond}(\mathscr{C}_{\operatorname{Mat}}, \mathscr{D}_{\operatorname{Mat}})$ ) = 1 and, it suffices to pick out one non-zero matrix **M** in the conductor and check if it is nonsingular. If it is non singular, we conclude that they are right equivalent, and provide a matrix **M** which realizes the equivalence. If **M** is singular, since the conductor has dimension 1, any other element of the conductor will be singular too and we conclude that the two codes are not equivalent.

In both situations, which are the ones that will "typically" happen, MCRE can be solved in polynomial time.

There remains to treat the case where  $\mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}}, \mathscr{D}_{\mathrm{Mat}})$  has a larger dimension and may contain both singular and non singular elements. In this situation, how to find non singular elements? In the other direction, how to assert that any element in this space is singular? Answering these questions requires the introduction of a fundamental notion: the *stabilizer algebras* of a matrix code.

**Definition 9** (Stabilizer Algebra). — Let  $\mathscr{C}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  be a matrix code. Its right stabilizer algebra is defined as:

$$\mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}}) \stackrel{def}{=} \{ \mathbf{M} \in \mathcal{M}_n(\mathbb{F}_q) \mid \mathscr{C}_{\mathrm{Mat}} \mathbf{M} \subseteq \mathscr{C}_{\mathrm{Mat}} \}$$

while its *left stabilizer* algebra is defined as:

$$\mathrm{Stab}_{\mathrm{left}}(\mathscr{C}_{\mathrm{Mat}}) \stackrel{aej}{=} \{ \mathbf{M} \in \mathcal{M}_m(\mathbb{F}_q) \mid \mathbf{M}\mathscr{C}_{\mathrm{Mat}} \subseteq \mathscr{C}_{\mathrm{Mat}} \}.$$

**Remark 10.** — In terms of terminology, these algebra are also referred to as *idealisers* or *nuclei*. See for instance [LN16, LTZ17].

**Remark 11.** — The right stabilizer algebra is nothing but the conductor of  $\mathscr{C}_{Mat}$  into itself, namely  $Cond(\mathscr{C}_{Mat}, \mathscr{C}_{Mat})$ . In particular, it can be computed by solving a system of linear equations.

<sup>&</sup>lt;sup>(1)</sup>if they do not, then they cannot be equivalent.

It is easily verified that  $\operatorname{Stab}_{\operatorname{left}}(\mathscr{C}_{\operatorname{Mat}})$  and  $\operatorname{Stab}_{\operatorname{right}}(\mathscr{C}_{\operatorname{Mat}})$  are sub-algebras of  $\mathcal{M}_m(\mathbb{F}_q)$  and  $\mathcal{M}_n(\mathbb{F}_q)$  respectively. In particular, for a given code  $\mathscr{C}_{\operatorname{Mat}}$ , its right stabilizer algebra contains scalar matrices, i.e. matrices of the form  $\lambda \mathbf{I}_n$ . If the stabilizer does not contain other matrices, it is said to be *trivial*. Note that most of the matrix codes have trivial stabilizer algebras. The following statement gives a first motivation for the introduction of these algebras.

**Proposition 12.** — Let  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  be two right equivalent codes; i.e. there exists  $\mathbf{Q} \in \mathbf{GL}_n(\mathbb{F}_q)$  such that  $\mathscr{C}_{Mat}\mathbf{Q} = \mathscr{D}_{Mat}$ . Suppose also that  $\dim(\mathbf{Cond}(\mathscr{C}_{Mat}, \mathscr{D}_{Mat})) \ge 2$ . Then,  $\mathscr{C}_{Mat}$  and  $\mathscr{D}_{Mat}$  have non trivial right stabilizer algebra.

*Proof.* — By assumption, there exists  $\mathbf{M} \in \mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}}, \mathscr{D}_{\mathrm{Mat}})$  which is not a scalar multiple of  $\mathbf{Q}$  and  $\mathbf{M}\mathbf{Q}^{-1} \in \mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}}) \setminus \{\lambda \mathbf{I}_n \mid \lambda \in \mathbb{F}_q\}$  and  $\mathbf{Q}^{-1}\mathbf{M} \in \mathrm{Stab}_{\mathrm{right}}(\mathscr{D}_{\mathrm{Mat}}) \setminus \{\lambda \mathbf{I}_n \mid \lambda \in \mathbb{F}_q\}$ .  $\Box$ 

While we observed that, when the conductor has dimension  $\leq 1$ , then solving MCRE is easy to solve. The previous statement asserts that, the cases where the conductor has a higher dimension while the codes are right equivalent are precisely the cases where stabilizer algebras are non-trivial

In such case, the problem is more technical to prove and requires a further study of the stabilizer algebra using Wedderburn–Artin Theory. We refer the reader to **[DK94]** for the mathematical aspects of this theory and to the works of Friedl and Rónyai **[FR85, Rón90]** for the computational aspects. The results of the previous references assert the existence of decompositions

$$\mathscr{C}_{\mathrm{Mat}} = \mathscr{C}_{\mathrm{Mat}} \mathbf{E}_1 \oplus \cdots \oplus \mathscr{C}_{\mathrm{Mat}} \mathbf{E}_s, \qquad \mathscr{D}_{\mathrm{Mat}} = \mathscr{C}_{\mathrm{Mat}} \mathbf{F}_1 \oplus \cdots \oplus \mathscr{C}_{\mathrm{Mat}} \mathbf{F}_t$$

where the  $\mathbf{E}_i$ 's (resp. the  $\mathbf{F}_i$ 's) are projectors matrices onto subspaces in direct sum. Moreover, these decompositions are unique up to conjugation and deciding the right equivalence reduces to decide equivalences of these small pieces as detailed in Section 4.2.3.

**3.2. Equivalence of**  $\mathbb{F}_{q^m}$ -linear codes. — Clearly, an instance of V-MCE is a particular instance of MCRE and hence, according to the previous discussion, can be solved in polynomial time. On the other hand, instances of HV-MCE are particular instances of MCE. Thus, solving HV-MCE requires à *priori* to look for left and right equivalence at the same time. However, it turns out, that the  $\mathbb{F}_{q^m}$ -linear structure permits to treat these problems separately.

Basically, we proceed as follows. We start from a pair of matrix codes  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$ obtained by expanding vector codes  $\mathscr{C}_{Vec}, \mathscr{D}_{Vec} \subseteq \mathbb{F}_{q^m}^n$  in (possibly distinct)  $\mathbb{F}_q$ -bases of  $\mathbb{F}_{q^m}$ . The left stabilizer algebras of these codes can be computed and both are either isomorphic to  $\mathbb{F}_{q^m}$  or contain a subalgebra isomorphic to  $\mathbb{F}_{q^m}$ . The latter case is rather more technical to study (see Section 5.3), thus let us suppose we are in the former case. Classical results of linear algebra permit to prove that two sub-algebras  $\mathrm{Stab}_{\mathrm{left}}(\mathscr{C}_{\mathrm{Mat}}), \mathrm{Stab}_{\mathrm{left}}(\mathscr{D}_{\mathrm{Mat}}) \subseteq \mathcal{M}_m(\mathbb{F}_q)$  which are both isomorphic to  $\mathbb{F}_{q^m}$  are conjugated and that a matrix  $\mathbf{P} \in \mathbf{GL}_m(\mathbb{F}_q)$ , that can be efficiently computed, such that

$$\operatorname{Stab}_{\operatorname{left}}(\mathscr{D}_{\operatorname{Mat}}) = \mathbf{P}^{-1} \operatorname{Stab}_{\operatorname{left}}(\mathscr{C}_{\operatorname{Mat}}) \mathbf{P}.$$

Next, replacing  $\mathscr{D}_{Mat}$  by  $\mathbf{P}\mathscr{D}_{Mat}$ , the two codes turn out to have the same left stabilizer matrix and then, up to some technical details related to the action of the Frobenius map (see Section 5.2), we are then reduced to decide right equivalence on the two codes.

**3.3. Reduction from MCE to ME.** — We prove that the general equivalence problem for matrix codes, MCE, is at least as hard as the monomial equivalence problem ME. We obtain such a statement by providing a polynomial time reduction from ME to MCE. This reduction was suggested by Hugues Randriambololona. Another reduction for the search versions of the problems is given in Section 6.

Consider the map

(3) 
$$\Phi: \left\{ \begin{array}{ccc} \mathbb{F}_{q}^{n} & \longrightarrow & \mathcal{M}_{n}(\mathbb{F}_{q}) \\ (x_{1}, \dots, x_{n}) & \longmapsto & \begin{pmatrix} x_{1} & \\ & \ddots & \\ & & & x_{n} \end{pmatrix} \right\}.$$

An important feature of this map is that it sends vectors with Hamming weight t on matrices with rank t, hence it is an isometry. The reduction rests on the following statement.

**Proposition 13.** — The map  $\Phi$  of (3) sends positive (resp. negative) instances of ME into positive (resp. negative) instances of MCE.

*Proof.* — Suppose two vector codes  $\mathscr{C}_{\text{Vec}}$ ,  $\mathscr{D}_{\text{Vec}} \subseteq \mathbb{F}_q^n$  are monomially equivalent, *i.e.*  $\mathscr{C}_{\text{Vec}}\mathbf{DP} = \mathscr{D}_{\text{Vec}}$  where  $\mathbf{D}$  is a nonsingular diagonal matrix and  $\mathbf{P}$  is a permutation matrix. Then the codes  $\mathscr{C}_{\text{Mat}} \stackrel{def}{=} \Phi(\mathscr{C}_{\text{Vec}})$  and  $\mathscr{D}_{\text{Mat}} \stackrel{def}{=} \Phi(\mathscr{D}_{\text{Vec}})$  satisfy

$$\mathbf{P}^{-1}\mathscr{C}_{\mathrm{Mat}}\mathbf{D}\mathbf{P}=\mathscr{D}_{\mathrm{Mat}}.$$

Hence, the matrix codes are equivalent.

Conversely, suppose that  $\mathscr{C}_{\text{Vec}}$ ,  $\mathscr{D}_{\text{Vec}}$  are not monomially equivalent while  $\mathscr{C}_{\text{Mat}}$ ,  $\mathscr{D}_{\text{Mat}}$  are equivalent. This entails that  $\mathscr{C}_{\text{Mat}}$ ,  $\mathscr{D}_{\text{Mat}}$  are isometric with respect to the rank metric, and, since  $\Phi$  is an isometry too, the codes  $\mathscr{C}_{\text{Vec}}$ ,  $\mathscr{D}_{\text{Vec}}$  are isometric with respect to the Hamming metric. But from McWilliams equivalence theorem [Mac62, HP03], this entails the monomial equivalence, a contradiction.

### 4. Solving the Matrix Codes Right Equivalence Problem (MCRE)

This section is devoted to the resolution of MCRE. As we already noticed in Section 3.1, most of the time, the resolution of the problem is simple since the conductor  $\mathbf{Cond}(\mathscr{C}_{Mat}, \mathscr{D}_{Mat})$  has typically dimension  $\leq 1$ . If it does not, while the codes are right equivalent, then, according to Proposition 12, the codes  $\mathscr{C}_{Mat}$  and  $\mathscr{D}_{Mat}$  should have non trivial stabilizer algebras. The following statements provide further relations between the structure of the conductor with that of the stabilizer algebras, which will be helpful in what follows.

**Proposition 14.** — If two codes  $\mathscr{C}_{Mat}$  and  $\mathscr{D}_{Mat}$  are right equivalent, i.e.  $\mathscr{C}_{Mat}\mathbf{Q} = \mathscr{D}_{Mat}$  for some  $\mathbf{Q} \in \mathbf{GL}_n(\mathbb{F}_q)$ , then their right stabilizer algebras are conjugated under  $\mathbf{Q}$ :

$$Stab_{right}(\mathscr{C}_{Mat}) = \mathbf{Q} \cdot Stab_{right}(\mathscr{D}_{Mat}) \cdot \mathbf{Q}^{-1}$$

*Proof.* — Let  $\mathbf{D} \in \text{Stab}_{\text{right}}(\mathscr{D}_{\text{Mat}})$ . By definition we have the following computation,

$$\mathscr{C}_{\mathrm{Mat}}\mathbf{Q}\mathbf{D}\mathbf{Q}^{-1} = \mathscr{D}_{\mathrm{Mat}}\mathbf{D}\mathbf{Q}^{-1} \subseteq \mathscr{D}_{\mathrm{Mat}}\mathbf{Q}^{-1} = \mathscr{C}_{\mathrm{Mat}}$$

which shows that  $\mathbf{QDQ}^{-1} \in \mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}})$  and easily concludes the proof.

This proposition enables to relate the conductor of right equivalent codes to their stabilizer algebras. It asserts in particular that  $\mathbf{Cond}(\mathscr{C}_{Mat}, \mathscr{D}_{Mat})$  is a left module on  $\mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{Mat})$  and a right module on  $\mathrm{Stab}_{\mathrm{right}}(\mathscr{D}_{Mat})$ .

**Proposition 15.** — Let  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  be matrix codes and  $\mathbf{Q} \in \mathbf{GL}_n(\mathbb{F}_q)$  such that  $\mathscr{C}_{Mat}\mathbf{Q} = \mathscr{D}_{Mat}$ . Then,

$$\mathbf{Cond}(\mathscr{C}_{Mat},\mathscr{D}_{Mat}) = Stab_{right}(\mathscr{C}_{Mat}) \cdot \mathbf{Q} = \mathbf{Q} \cdot Stab_{right}(\mathscr{D}_{Mat}).$$

*Proof.* — Let  $\mathbf{S} \in \text{Stab}_{\text{right}}(\mathscr{C}_{\text{Mat}})$ , then

$$\mathscr{C}_{\mathrm{Mat}}\mathbf{SQ} \subseteq \mathscr{C}_{\mathrm{Mat}}\mathbf{Q} = \mathscr{D}_{\mathrm{Mat}}.$$

Therefore,  $\mathbf{SQ} \in \mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}}, \mathscr{D}_{\mathrm{Mat}})$  and hence  $\mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}})\mathbf{Q} \subseteq \mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}}, \mathscr{D}_{\mathrm{Mat}})$ . Conversely, let  $\mathbf{C} \in \mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}}, \mathscr{D}_{\mathrm{Mat}})$ , then

$$\mathscr{C}_{\mathrm{Mat}}\mathbf{C}\mathbf{Q}^{-1} \subseteq \mathscr{D}_{\mathrm{Mat}}\mathbf{Q}^{-1} = \mathscr{C}_{\mathrm{Mat}}.$$

Therefore,  $\mathbf{CQ}^{-1} \in \mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}})$  or equivalently  $\mathbf{C} \in \mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}})\mathbf{Q}$ . This yields

 $\mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}}, \mathscr{D}_{\mathrm{Mat}}) \subseteq \mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}})\mathbf{Q}.$ 

The equality  $\operatorname{Cond}(\mathscr{C}_{\operatorname{Mat}}, \mathscr{D}_{\operatorname{Mat}}) = \operatorname{\mathbf{Q}Stab}_{\operatorname{right}}(\mathscr{D}_{\operatorname{Mat}})$  is a consequence of the previous equality together with Proposition 14.

**4.1. Finite dimensional algebras.** — As said earlier, the general resolution of MCRE involves results on finite dimensional algebras and the Artin–Wedderburn theory. Here we recall some known results in the literature that we will apply to the right stabilizer algebra of our codes. We refer the reader to [**DK94**] for further details on the theoretical notions and to [**FR85**, **Rón90**] for the algorithmic aspects.

A finite dimensional algebra  $\mathcal{A}$  is finite dimensional vector space over  $\mathbb{F}_q$  which is also a ring (possibly non commutative) with unit. A way to represent such an algebra is to use a basis  $a_1, \ldots, a_\ell$  together with the collection of so-called structure constants with respect to this basis, which are a sequence  $(\lambda_{ijk})_{i,j,k \in \{1,\ldots,\ell\}}$  such that:

$$\forall i, j \in \{1, \dots, \ell\}, \quad a_i a_j = \sum_{k=1}^{\ell} \lambda_{ijk} a_k.$$

Any such algebra can be represented as a subring of  $\mathcal{M}_{\ell}(\mathbb{F}_q)$  ([**DK94**, Thm. 1.3.1]).

A finite dimensional algebra is *simple* if its only two-sided ideals are  $\{0\}$  and the whole algebra itself. Artin–Wedderburn theory asserts that any simple algebra over a finite field  $\mathbb{F}_q$  is isomorphic to  $\mathcal{M}_r(\mathbb{F}_{q^\ell})$  for some positive  $r, \ell$  (see [**DK94**, Cor. 2.4.5 & Thm. 5.2.4]). A *semi-simple* algebra

is a cartesian product of simple algebras. Another fundamental object is the *Jacobson radical* of an algebra  $\mathcal{A}$  which is defined as:

$$\operatorname{Rad}(\mathcal{A}) \stackrel{def}{=} \{ x \in \mathcal{A} : \forall y \in \mathcal{A}, xy \text{ is nilpotent} \}.$$

This is a two-sided ideal and, from [**DK94**, Thm. 3.1.1],  $\mathcal{A}_{\text{Rad}\mathcal{A}}$  is semi-simple. If this last quotient is a field,  $\mathcal{A}$  is said to be *local*.

Finally, another fundamental notion is that of *idempotents*. An element  $e \in \mathcal{A}$  is *idempotent* if  $e^2 = e$ . In matrix algebras, idempotent are nothing but projection matrices. Two idempotents  $e_1$  and  $e_2$  are said to be *orthogonal* if  $e_1e_2 = e_2e_1 = 0$ . An idempotent e is said to be *minimal* if e cannot be decomposed as a sum of two non zero orthogonal idempotents  $e = e_1 + e_2$ . An algebra is local if and only if its only idempotent is the unit.

### 4.2. General resolution of MCRE. -

**4.2.1.** Context. — Let  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  be two matrix codes. Once  $Cond(\mathscr{C}_{Mat}, \mathscr{D}_{Mat})$  is computed, our aim is to decide, and find if exists, a matrix  $\mathbf{Q} \in \mathbf{GL}_n(\mathbb{F}_q)$  such that

$$\mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}},\mathscr{D}_{\mathrm{Mat}}) = \mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}}) \cdot \mathbf{Q} = \mathbf{Q} \cdot \mathrm{Stab}_{\mathrm{right}}(\mathscr{D}_{\mathrm{Mat}})$$

To solve this problem, we first treat the case where  $\operatorname{Stab}_{right}(\mathscr{C}_{Mat})$  is local. Then, we show how to reduce to that case using idempotents of the stabilizer algebra.

**4.2.2.** When the stabilizer algebras are local. — It is easy to decide if  $\mathscr{C}_{Mat}$  and  $\mathscr{D}_{Mat}$  are right equivalent when  $\operatorname{Stab}_{right}(\mathscr{C}_{Mat})$  is a local algebra. Basically it relies on the following lemma.

**Lemma 16.** — Let  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  be two matrix codes that are right equivalent and such that  $Stab_{right}(\mathscr{C}_{Mat})$  is local. Let  $\mathcal{R} \stackrel{def}{=} Rad(Stab_{right}(\mathscr{C}_{Mat}))$ . Then any element of

$$\mathbf{Cond}(\mathscr{C}_{Mat},\mathscr{D}_{Mat}) \smallsetminus (\mathcal{R}\mathbf{Cond}(\mathscr{C}_{Mat},\mathscr{D}_{Mat}))$$

is nonsingular.

*Proof.* — By hypothesis,  $\mathscr{C}_{Mat}\mathbf{Q} = \mathscr{D}_{Mat}$  for some  $\mathbf{Q} \in \mathbf{GL}_n(\mathbb{F}_q)$ . From Proposition 15,  $\mathbf{Cond}(\mathscr{C}_{Mat}, \mathscr{D}_{Mat}) = \mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}})\mathbf{Q}$  and hence any element of  $\mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}}, \mathscr{D}_{\mathrm{Mat}})$  can be written  $\mathbf{CQ}$  where  $\mathbf{C} \in \mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}})$ . Since  $\mathbf{Q}$  is nonsingular,  $\mathbf{CQ}$  is singular if and only if  $\mathbf{C}$  is singular. Finally, since  $\mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}})$  is local, by [**DK94**, Thm. 3.2.2],  $\mathbf{C}$  is singular if and only if it is in  $\mathcal{R}$ .

This situation yields Algorithm 1 which decides the right-equivalence problem in polynomial time when right stabilizer algebras are local.

**Algorithm 1:** An algorithm to decide the right equivalence problem when stabilizer algebras are local

**Input** : Two codes  $\mathscr{C}_{Mat}$ ,  $\mathscr{D}_{Mat} \subseteq Mat_{m,n}(\mathbb{F}_q)$ **Output:** Codes are right-equivalent or not

Compute  $\mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}}, \mathscr{D}_{\mathrm{Mat}}), \mathcal{R} \stackrel{def}{=} \mathrm{Rad}(\mathrm{Stab}_{\mathrm{right}}(\mathscr{C}_{\mathrm{Mat}})) \text{ and } \mathcal{R}\mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}}, \mathscr{D}_{\mathrm{Mat}})$ Pick  $\mathbf{A} \in \mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}}, \mathscr{D}_{\mathrm{Mat}}) \smallsetminus \mathcal{R}\mathbf{Cond}(\mathscr{C}_{\mathrm{Mat}}, \mathscr{D}_{\mathrm{Mat}})$ 

if A non singular then ∟ return Codes are right-equivalent else ∟ return Codes are not right-equivalent

**Theorem 17.** — Let  $\mathscr{C}_{Mat}$ ,  $\mathscr{D}_{Mat}$  be matrix codes with local stabilizer algebras. Algorithm 1 succeeds to decide if they are right equivalent in polynomial time.

*Proof.* — The correctness of the algorithm is a direct consequence of Lemma 16. For the complexity, the computation of the conductor reduces to the resolution of a linear system. The computation of the radical can be done in polynomial time thanks to [FR85].

**4.2.3.** The general case. — In the general case, we aim to reduce the case where stabilizer algebras of codes are local. For this sake, we are going "to decompose" the unit of the right stabilizer algebras thanks to the following proposition.

**Proposition 18.** — Let  $\mathcal{A}$  be a n-dimensional algebra over  $\mathbb{F}_q$ . Then, there exists a polynomial time  $\mathcal{F}$ -algorithm computing a decomposition of the identity of  $\mathcal{A}$  as a sum of minimal orthogonal idempotents:

$$1_{\mathcal{A}} = a_1 + \dots + a_s.$$

Furthermore this conjugation is unique to conjugation, namely if:

$$1_{\mathcal{A}} = b_1 + \dots + b_t$$

we have s = t and there exists  $g \in \mathcal{A}^{\times}$  such that, up to re-indexing the  $b_i$ 's, we have  $a_i = gb_ig^{-1}$ for any  $i \in \{1, \ldots, s\}$ .

*Proof.* — We proceed as follows.

**Step 1.** Using Frield–Rónyai algorithm [**FR85**, Thm 5.7], compute the Jacobson Radical Rad( $\mathcal{A}$ ) of  $\mathcal{A}$ . This is done in polynomial time;

Step 2. Compute the semi-simple algebra  $\mathcal{S} \stackrel{def}{=} \mathcal{A}/\text{Rad}(\mathcal{A})$ . From [Rón90, Theorem 3.1] there is a polynomial time F-algorithm computing bases of simple algebras  $\mathcal{S}_i$  such that

$$\mathcal{S} = \mathcal{S}_1 \oplus \cdots \oplus \mathcal{S}_r.$$

This permits to compute the canonical decomposition of the identity into central idempotents:

$$1_{\mathcal{S}} = 1_{\mathcal{S}_1} + \dots + 1_{\mathcal{S}_r},$$

but these idempotents are not always minimal (unless the  $S_i$ 's are all fields) and should be decomposed into minimal ones. As mentioned in Section 4.1, since the  $S_i$ 's are simple, they are isomorphic to  $\mathcal{M}_{u_i}(\mathbb{F}_{q^{v_i}})$  for some positive  $u_i, v_i$ . Moreover, [**Rón90**, Sec. 5.1] provides a polynomial time F-algorithm to compute these isomorphisms. In the algebra  $\mathcal{M}_{u_i}(\mathbb{F}_{q^{v_i}})$  a minimal decomposition of the identity is obtained as the sum of the matrices with only one non-zero element which is on the diagonal. Using our explicit isomorphism between  $S_i$  and  $\mathcal{M}_{u_i}(\mathbb{F}_{q^{v_i}})$ , we get a decomposition of of  $1_{S_i}$  as a sum of minimal orthogonal idempotents and deduce such a minimal decomposition for S:

$$1_{\mathcal{S}} = e_1 + \dots + e_s$$
 where  $s \ge r$ .

**Step 3.** Since the ground field is finite, from Wedderburn–Malcev Theorem [**DK94**, Thm. 6.2.1], there exists an injective morphism of algebras  $S \hookrightarrow A$  and the image of this morphism is unique up to conjugation. Moreover, such a lift of S can be computed by linear algebra as explained in [**Bre11**, § 1.12]. Using this lift, we deduce a decomposition of the identity of A by minimal orthogonal idempotents:

$$1_{\mathcal{A}} = a_1 + \dots + a_s.$$

According to [DK94, Thm 3.4.1], this decomposition is unique up to conjugation.

In order to decide the right equivalence of two given matrix codes  $\mathscr{C}_{Mat}$ ,  $\mathscr{D}_{Mat}$ , we start by computing their right stabilizer algebras  $\operatorname{Stab}_{right}(\mathscr{C}_{Mat})$  and  $\operatorname{Stab}_{right}(\mathscr{D}_{Mat})$ . Thanks to Proposition 18, one can compute minimal idempotent decompositions of the identity of these algebras

(4) 
$$\mathbf{I}_n = \mathbf{E}_1 + \dots + \mathbf{E}_\ell$$
 and  $\mathbf{I}_n = \mathbf{F}_1 + \dots + \mathbf{F}_{\ell'}$ 

From Proposition 18 the stabilizer algebras are conjugated if an only if  $\ell = \ell'$  and after a suitable re-indexing, there is a matrix **Q** such that for any i,  $\mathbf{QE}_i\mathbf{Q}^{-1} = \mathbf{F}_i$ . In particular, if  $\ell \neq \ell'$  one can stop the process, the codes are not equivalent. Hence, from now on, we suppose  $\ell = \ell'$ . The idea

of the general case is to apply "piecewise" the algorithm designed for codes with local algebras to the terms of the following codes decompositions:

$$\mathscr{C}_{\mathrm{Mat}} = \mathscr{C}_{\mathrm{Mat}} \mathbf{E}_1 \oplus \cdots \oplus \mathscr{C}_{\mathrm{Mat}} \mathbf{E}_\ell \qquad \text{and} \qquad \mathscr{D}_{\mathrm{Mat}} = \mathscr{D}_{\mathrm{Mat}} \mathbf{F}_1 \oplus \cdots \oplus \mathscr{D}_{\mathrm{Mat}} \mathbf{F}_\ell.$$

One technical difficulty to tackle is that terms of the decompositions, are matrix codes whose span of row spaces are not the full space  $\mathbb{F}_q^n$ . Indeed, considering for instance the case of  $\mathscr{C}_{Mat}$ , the matrices  $\mathbf{E}_i$ 's are projectors whose images are in direct sum and any matrix in the space  $\mathscr{C}_{Mat}\mathbf{E}_i$ has a row space contained in that of  $\mathbf{E}_i$ .

To circumvent this issue, we will use the following statement whose proof is left to the reader. Lemma 19. — Let  $\mathbf{E} \in \mathcal{M}_n(\mathbb{F}_q)$  be a projector of rank r then there exist two full-rank matrices

$$\mathbf{E} = \mathbf{A}\mathbf{B}^{\top}$$
 and  $\mathbf{B}^{\top}\mathbf{A} = \mathbf{I}_r$ .

Using this lemma, we decompose the  $\mathbf{E}_i$ 's into  $\mathbf{A}_i \mathbf{B}_i^{\top}$  and the  $\mathbf{F}_i$ 's into  $\mathbf{U}_i \mathbf{V}_i^{\top}$  and will focus on the matrix codes  $\mathscr{C}_{\text{Mat}} \mathbf{A}_i$  (resp.  $\mathscr{D}_{\text{Mat}} \mathbf{U}_i$ ) which are contained in  $\mathcal{M}_{n,\text{Rk}(\mathbf{A}_i)}(\mathbb{F}_q)$  (resp.  $\mathcal{M}_{n,\text{Rk}(\mathbf{U}_i)}(\mathbb{F}_q)$ ).

**Remark 20.** — Note that replacing  $\mathscr{C}_{Mat}$  be a right equivalent code  $\mathscr{C}_{Mat}\mathbf{Q}_0$  for some  $\mathbf{Q}_0 \in \mathbf{GL}_n(\mathbb{F}_q)$  and the  $\mathbf{E}_i$ 's by their conjugates under  $\mathbf{Q}_0$ , one could get  $\mathbf{E}_i$ 's of the form



and the code  $\mathscr{C}_{Mat}\mathbf{A}_1$  would be the code  $\mathscr{C}_{Mat}$  but keeping only the  $Rk(\mathbf{E}_1)$  first columns of each element, the code  $\mathscr{C}_{Mat}\mathbf{A}_2$  would be  $\mathscr{C}_{Mat}$  but only keeping the  $Rk(\mathbf{E}_2)$  next columns of each element, and so on.

Toward the computation of a "picewise" right equivalence, we need the following crucial statements. The first one shows that right equivalence entails picewise right equivalence.

**Proposition 21.** — Let  $\mathscr{C}_{Mat}$ ,  $\mathscr{D}_{Mat}$  be two matrix codes in  $\mathcal{M}_{m,n}(\mathbb{F}_q)$  such that  $\mathscr{C}_{Mat}\mathbf{Q} = \mathscr{D}_{Mat}$  for some  $\mathbf{Q} \in \mathbf{GL}_n(\mathbb{F}_q)$ . Let  $\mathbf{E}_1, \ldots, \mathbf{E}_\ell$  and  $\mathbf{F}_1, \ldots, \mathbf{F}_\ell$  be minimal idempotents of their stabiliser algebras as in (4) together with their respective decompositions  $\mathbf{A}_i \mathbf{B}_i^{\top}$  and  $\mathbf{U}_i \mathbf{V}_i^{\top}$  according to Lemma 19. Then, after a suitable re-indexing, for any *i* the codes  $\mathscr{C}_{Mat}\mathbf{A}_i$  and  $\mathscr{D}_{Mat}\mathbf{A}_i$  are right equivalent.

*Proof.* — Since the  $\mathbf{F}_i$ 's are unique up to conjugation, after possibly replacing  $\mathbf{Q}$  by  $\mathbf{QS}$  for some  $\mathbf{S} \in \mathrm{Stab}_{\mathrm{right}}(\mathscr{D}_{\mathrm{Mat}})$  one may assume that for any  $i, \mathbf{F}_i = \mathbf{Q}^{-1} \mathbf{E}_i \mathbf{Q}$ . Therefore,

(5) 
$$\forall i \in \{1, \dots, \ell\}, \quad \mathscr{C}_{\mathrm{Mat}} \mathbf{E}_i \mathbf{Q} = \mathscr{C}_{\mathrm{Mat}} \mathbf{Q} \mathbf{Q}^{-1} \mathbf{E}_i \mathbf{Q} = \mathscr{D}_{\mathrm{Mat}} \mathbf{F}_i.$$

Next, recall that  $\mathbf{E}_i = \mathbf{A}_i \mathbf{B}_i^{\top}$ ,  $\mathbf{F}_i = \mathbf{U}_i \mathbf{V}_i^{\top}$  with  $\mathbf{V}_i^{\top} \mathbf{U}_i = \mathbf{B}_i^{\top} \mathbf{A}_i = 0$  and consider (5) again

$$\forall i, \quad \mathscr{C}_{\mathrm{Mat}} \mathbf{A}_i \mathbf{B}_i^\top \mathbf{Q} = \mathscr{D}_{\mathrm{Mat}} \mathbf{U}_i \mathbf{V}_i^\top \implies \qquad \mathscr{C}_{\mathrm{Mat}} \mathbf{A}_i \left( \mathbf{B}_i^\top \mathbf{Q} \mathbf{U}_i \right) = \mathscr{D}_{\mathrm{Mat}} \mathbf{U}_i \underbrace{\mathbf{V}_i^\top \mathbf{U}_i}_{=\mathbf{I}_{r_i}} = \mathscr{D}_{\mathrm{Mat}} \mathbf{U}_i,$$

where  $r_i$  denotes the rank of  $\mathbf{E}_i$ .

There remains to prove that  $\mathbf{B}_i^{\top} \mathbf{Q} \mathbf{U}_i \in \mathcal{M}_{r_i}(\mathbb{F}_q)$  is nonsingular. First, note that  $\operatorname{Rk}(\mathbf{B}_i^{\top} \mathbf{Q} \mathbf{U}_i) \geq \operatorname{Rk}(\mathbf{A}_i \mathbf{B}_i^{\top} \mathbf{Q} \mathbf{U}_i \mathbf{V}_i^{\top}) = \operatorname{Rk}(\mathbf{E}_i \mathbf{Q} \mathbf{F}_i).$ 

 $\mathbf{A}, \mathbf{B} \in \mathcal{M}_{n,r}(\mathbb{F}_q)$  such that

Since  $\mathbf{F}_i = \mathbf{Q}^{-1} \mathbf{E}_i \mathbf{Q}$ , the right hand side becomes  $\operatorname{Rk}(\mathbf{E}_i^2 \mathbf{Q}) = \operatorname{Rk}(\mathbf{E}_i \mathbf{Q}) = r_i$ . Therefore  $\mathbf{B}_i^\top \mathbf{Q} \mathbf{U}_i$  is a square matrix with full rank, it is invertible.

Conversely, the following statement asserts that picewise right equivalence entails right equivalence.

**Proposition 22.** — Let  $\mathscr{C}_{Mat}$ ,  $\mathscr{D}_{Mat}$  and the  $\mathbf{E}_i = \mathbf{A}_i \mathbf{B}_i^{\top}$  and  $\mathbf{F}_i = \mathbf{U}_i \mathbf{V}_i^{\top}$  as in Proposition 21. Suppose that for any  $i \in \{1, \ldots, \ell\}$  there exists  $\mathbf{Q}_i \in \mathbf{GL}_{r_i}(\mathbb{F}_q)$ , where  $r_i = \mathrm{Rk}(\mathbf{E}_i)$ , such that  $\mathscr{C}_{Mat}\mathbf{A}_i\mathbf{Q}_i = \mathscr{D}_{Mat}\mathbf{U}_i$ . Then  $\mathscr{C}_{Mat}$  and  $\mathscr{D}_{Mat}$  are right equivalent. In particular,

$$\mathscr{C}_{Mat} \mathbf{Q} = \mathscr{D}_{Mat} \qquad where \qquad \mathbf{Q} \stackrel{def}{=} \sum_{i=1}^{\iota} \mathbf{A}_i \mathbf{Q}_i \mathbf{V}_i^{\top}$$

and the latter matrix  $\mathbf{Q}$  is nonsingular.

*Proof.* — We have,

$$\mathscr{C}_{\mathrm{Mat}}\mathbf{Q} = \sum_{i=1}^{\ell} \mathscr{C}_{\mathrm{Mat}} \mathbf{A}_i \mathbf{Q}_i \mathbf{V}_i^{\top} = \sum_{i=1}^{\ell} \mathscr{D}_{\mathrm{Mat}} \mathbf{U}_i \mathbf{V}_i^{\top} = \mathscr{D}_{\mathrm{Mat}} \sum_{i=1}^{\ell} \mathbf{F}_i = \mathscr{D}_{\mathrm{Mat}}.$$

There remains to prove that  $\mathbf{Q}$  is nonsingular. For this, we will prove that

$$\mathbf{Q}' \stackrel{def}{=} \sum_{i=1}^{\ell} \mathbf{U}_i \mathbf{Q}_i^{-1} \mathbf{B}_i^{ op}$$

is its inverse. Indeed, consider their product:

(6) 
$$\mathbf{Q}\mathbf{Q}' = \sum_{i=1}^{t} \mathbf{A}_{i}\mathbf{Q}_{i}\mathbf{V}_{i}^{\top}\mathbf{U}_{i}\mathbf{Q}_{i}^{-1}\mathbf{B}_{i}^{\top} + \sum_{i\neq j} \mathbf{A}_{i}\mathbf{Q}_{i}\mathbf{V}_{i}^{\top}\mathbf{U}_{j}\mathbf{Q}_{j}^{-1}\mathbf{B}_{j}^{\top}.$$

From Lemma 19, the left-hand sum of (6) is

$$\sum_{i=1}^{\ell} \mathbf{A}_i \mathbf{B}_i^{\top} = \sum_{i=1}^{\ell} \mathbf{E}_i = \mathbf{I}_n.$$

For the right-hand sum of (6), note that, again from Lemma 19,

$$\forall i \neq j, \quad \mathbf{V}_i^\top \mathbf{U}_j = \mathbf{V}_i^\top \underbrace{\mathbf{U}_i \mathbf{V}_i^\top}_{=\mathbf{E}_i} \underbrace{\mathbf{U}_j \mathbf{V}_j^\top}_{=\mathbf{E}_j} \mathbf{U}_j = 0.$$

Hence, all the terms of the right-hand sum of (6) are zero. This concludes the proof.

Finally, we prove that "pieces"  $\mathscr{C}_{Mat}\mathbf{A}_i$  have local right stabiliser algebras. Hence, equivalence can be decided using Algorithm 1.

**Proposition 23.** — Let  $\mathscr{C}_{Mat}$ ,  $\mathbf{E}_i = \mathbf{A}_i \mathbf{B}_i^{\top}$  as in Proposition 21. Then, the code  $\mathscr{C}_{Mat} \mathbf{A}_i$  has a local right stabiliser algebra.

*Proof.* — Suppose that  $\operatorname{Stab}_{\operatorname{right}}(\mathscr{C}_{\operatorname{Mat}}\mathbf{A}_i)$  is not local, then, there is a decomposition of its unit by orthogonal idempotents  $\mathbf{I}_{r_i} = \mathbf{E}_{i_1} + \mathbf{E}_{i_2}$ , where  $\mathbf{E}_{i_1}, \mathbf{E}_{i_2} \in \operatorname{Stab}_{\operatorname{right}}(\mathscr{C}_{\operatorname{Mat}}\mathbf{A}_i)$  Then,

$$\forall j \in \{1, 2\}, \qquad \mathscr{C}_{\mathrm{Mat}} \mathbf{A}_i \mathbf{E}_{i_j} \mathbf{B}_i^\top \subseteq \mathscr{C}_{\mathrm{Mat}} \mathbf{A}_i \mathbf{B}_i^\top = \mathscr{C}_{\mathrm{Mat}} \mathbf{E}_i \subseteq \mathscr{C}_{\mathrm{Mat}}.$$

Therefore,  $\mathbf{A}_i \mathbf{E}_{i_1} \mathbf{B}_i^{\top}$  and  $\mathbf{A}_i \mathbf{E}_{i_2} \mathbf{B}_i^{\top}$  both lie in  $\operatorname{Stab}_{\operatorname{right}}(\mathscr{C}_{\operatorname{Mat}})$ . Next, using Lemma 19, one can prove that they are orthogonal idempotents whose sum equals  $\mathbf{E}_i$ , which contradicts the minimality of  $\mathbf{E}_i$  as an idempotent of  $\operatorname{Stab}_{\operatorname{right}}(\mathscr{C}_{\operatorname{Mat}})$ .

The previous material permits to solve MCRE with Algorithm 2.

Theorem 24. — Algorithm 2 is a polynomial time F-algorithm solving MCRE.

Algorithm 2: An algorithm to solve MCRE
<b>Input</b> : Two codes $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq Mat_{m,n}(\mathbb{F}_q)$
<b>Output:</b> Codes are right-equivalent or not

Compute their right stabiliser algebras and minimal decompositions of their units by orthogonal idempotents:

 $\mathbf{I}_n = \mathbf{E}_1 + \dots + \mathbf{E}_\ell$  and  $\mathbf{I}_n = \mathbf{F}_1 + \dots + \mathbf{F}_{\ell'}$ .

 $\begin{array}{l} \mbox{if } \ell \neq \ell' \mbox{ then } \\ \mbox{ return Codes are not right equivalent } \\ \mbox{else } \\ \mbox{ Find a permutation } \sigma \in \mathfrak{S}_\ell \mbox{ such that } \forall i \in \{1, \ldots, \ell\} \mbox{ it exists a non-singular } \mathbf{Q}_i \mbox{ with } \\ \mbox{ } \mathscr{C}_{\mathrm{Mat}} \mathbf{A}_i \mathbf{Q}_i = \mathscr{D}_{\mathrm{Mat}} \mathbf{U}_{\sigma(i)} \mbox{ (using notation of Proposition 21 ). } \\ \mbox{ if Computation fails then } \\ \mbox{ } \mbox{ } \mbox{ return Codes are not right equivalent } \\ \mbox{ else } \\ \mbox{ } \mbox{ } \mbox{ } \mbox{ equivalent } \\ \mbox{ } \mbox{ } \mbox{ } \mbox{ return Codes are right Equivalent } \end{array}$ 

*Proof.* — The computation of right stabilizer algebras is done in polynomial time. According to Proposition 18 the computation of minimal decompositions of their units by orthogonal idempotents can be done with a polynomial time F-algorithm.

Next step (clearly if  $\ell \neq \ell'$  codes are not right equivalent) of the computation can be done in polynomial time using a search version of Algorithm 1. Furthermore, if there was a permutation  $\sigma \in \mathfrak{S}_{\ell}$  and nonsingular matrices  $\mathbf{Q}_i$  such that

$$orall i, \quad \mathscr{C}_{\mathrm{Mat}} \mathbf{A}_i \mathbf{Q}_i = \mathscr{D}_{\mathrm{Mat}} \mathbf{U}_{\sigma(i)},$$

then according to Proposition 22 the codes are equivalent (and reciprocally) and the equivalence is asserted by the matrix  $\mathbf{Q} \stackrel{def}{=} \sum_{i} \mathbf{A}_{i} \mathbf{Q}_{i} \mathbf{V}_{i}^{\top}$  which concludes the proof.

### 5. The Code Equivalence Problem for $\mathbb{F}_{q^m}$ -linear codes : an Easy Problem

When described as matrix codes,  $\mathbb{F}_{q^m}$ -linear codes are nothing but codes  $\mathscr{C}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_{q^m})$ whose left stabilizer algebra  $\operatorname{Stab}_{left}(\mathscr{C}_{Mat}) \subseteq \mathcal{M}_m(\mathbb{F}_q)$  contains a subalgebra isomorphic to  $\mathbb{F}_{q^m}$ as in the following definition.

**Definition 25** (Representation of  $\mathbb{F}_{q^m}$ ). — A sub-algebra  $\mathcal{F} \subseteq \mathcal{M}_m(\mathbb{F}_q)$  is a representation of  $\mathbb{F}_{q^m}$  if it satisfies one of the following equivalent conditions:

- It is isomorphic to  $\mathbb{F}_{q^m}$  as an  $\mathbb{F}_q$ -algebra;
- It spanned (as an algebra) by a matrix  $\mathbf{A} \in \mathcal{M}_m(\mathbb{F}_q)$  whose characteristic polynomial is  $\mathbb{F}_q$ -irreducible.

In particular, taking any monic irreducible polynomial  $P \in \mathbb{F}_q[X]$  and  $\mathbb{C}_P$  be its companion matrix, then  $\mathbb{F}_q[\mathbb{C}_P]$  is a representation of  $\mathbb{F}_{q^m}$ .

**5.1. Statement of equivalence problems for**  $\mathbb{F}_{q^m}$ -linear codes. — Consider an instance of Problem 3 (HV-MCE) *i.e.*, two matrix codes  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  of which are known to be matrix representations of  $\mathbb{F}_{q^m}$ -linear codes but possibly represented in two distinct and unknown bases. To reduce to an instance of MCRE, our objective is to find a matrix  $\mathbf{P} \in \mathbf{GL}_m(\mathbb{F}_q)$  such that  $\mathbf{P}\mathscr{C}_{Mat}$  is right equivalent to  $\mathscr{D}_{Mat}$ . Such a  $\mathbf{P}$  will be deduced from the left stabilizer algebras of the codes. Recall that these left stabilizer algebras  $\mathrm{Stab}_{\mathrm{left}}(\mathscr{C}_{\mathrm{Mat}}), \mathrm{Stab}_{\mathrm{left}}(\mathscr{D}_{\mathrm{Mat}})$  can be computed

by solving a linear system in a very similar manner as the computation of right stabilizer algebras detailed in Section 4 (see Remark 11). In addition, similarly to Proposition 14, we have that

$$\operatorname{Stab}_{\operatorname{left}}(\mathscr{C}_{\operatorname{Mat}}) = \mathbf{P}^{-1} \operatorname{Stab}_{\operatorname{left}}(\mathscr{D}_{\operatorname{Mat}}) \mathbf{P}.$$

Thus our objective is, from the knowledge of the stabilizer algebras  $\operatorname{Stab}_{\operatorname{left}}(\mathscr{C}_{\operatorname{Mat}})$  and  $\operatorname{Stab}_{\operatorname{left}}(\mathscr{D}_{\operatorname{Mat}})$ , to solve an algebra conjugacy problem. It will be possible since these algebras contain a representation of  $\mathbb{F}_{q^m}$ , and hence are non-trivial We treat the problem by considering separately two cases:

- (i) The left stabilizer algebras of  $\mathscr{C}_{Mat}$ ,  $\mathscr{D}_{Mat}$  are representations of  $\mathbb{F}_{q^m}$ ;
- (ii) The left stabilizer algebras of  $\mathscr{C}_{Mat}$ ,  $\mathscr{D}_{Mat}$  contain representations of  $\mathbb{F}_{q^m}$  as proper subalgebras

**5.2.** When the left stabilizer algebras are representations of  $\mathbb{F}_{q^m}$ . — In the sequel, we need some very classical results in linear algebra and representations of  $\mathbb{F}_{q^m}$  which are summarized in the following technical statement.

**Lemma 26.** — Let  $\mathbf{A} \in \mathcal{M}_m(\mathbb{F}_q)$  be a matrix whose characteristic polynomial  $\chi_{\mathbf{A}}$  is irreducible (for instance the companion matrix of any monic irreducible polynomial of degree m). Then  $\mathbb{F}_q[\mathbf{A}] \simeq \mathbb{F}_{q^m}$  and

- (i) a matrix  $\mathbf{C} \in \mathcal{M}_m(\mathbb{F}_q)$  commutes with any element of  $\mathbb{F}_q[\mathbf{A}]$  if and only if it is an element of  $\mathbb{F}_q[\mathbf{A}]$ ;
- (ii) for any  $\mathbf{x}, \mathbf{y} \in \mathbb{F}_q^m \setminus \{0\}$  there exists  $P(\mathbf{A}) \in \mathbb{F}_q[\mathbf{A}]$  such that  $P(\mathbf{A})\mathbf{x}^\top = \mathbf{y}^\top$ ;
- (iii) There exists  $\Theta \in \mathbf{GL}_m(\mathbb{F}_q)$  such that

$$\Theta^m = \mathbf{I}_m$$
 and  $\forall P(\mathbf{A}) \in \mathbb{F}_q[\mathbf{A}], \quad \Theta P(\mathbf{A})\Theta^{-1} = P(\mathbf{A})^q.$ 

- (iv) Any matrix  $\Gamma \in \mathbf{GL}_m(\mathbb{F}_q)$  satisfying  $\Gamma \mathbb{F}_q[\mathbf{A}]\Gamma^{-1} = \mathbb{F}_q[\mathbf{A}]$  is of the form  $\Gamma = \Theta^j P(\mathbf{A})$  for some  $P(\mathbf{A}) \in \mathbb{F}_q[\mathbf{A}]^{\times}$  and some  $j \in \{0, \dots, m-1\}$ .
- (v) any matrix  $\mathbf{B} \in \mathcal{M}_m(\mathbb{F}_q)$  with the same characteristic polynomial as  $\mathbf{A}$ , then  $\mathbf{A}$  and  $\mathbf{B}$  are conjugated, i.e. there exists  $\mathbf{P} \in \mathbf{GL}_m(\mathbb{F}_q)$  such that  $\mathbf{A} = \mathbf{P}^{-1}\mathbf{B}\mathbf{P}$  and  $\mathbf{P}$  can be computed in polynomial time.

*Proof.* — Clearly  $\mathbb{F}_q[\mathbf{A}]$  is commutative. Conversely, if **C** commutes with any element of  $\mathbb{F}_q[\mathbf{A}]$  then it commutes with **A**. Since the characteristic polynomial  $\chi_{\mathbf{A}}$  of **A** is irreducible and  $\mathbb{F}_q$  is perfect, then **A** has distinct eigenvalues and it is a well–known fact in linear algebra that this latter features entails that **C** should be a polynomial in **A**. This proves (i).

Consider the space  $F \stackrel{def}{=} \{P(\mathbf{A})\mathbf{x}^{\top} \mid P \in \mathbb{F}_q[X]\} \subseteq \mathbb{F}_q^m$ . This space is isomorphic to  $\mathbb{F}_q[X]/(\pi_{\mathbf{A},\mathbf{x}})$ where  $\pi_{\mathbf{A},\mathbf{x}}$  is the monic polynomial P of lowest degree satisfying  $P(\mathbf{A})\mathbf{x}^{\top} = 0$ . This polynomial divides the minimal polynomial of  $\mathbf{A}$  which itself is known to divide  $\chi_{\mathbf{A}}$ . Since, by hypothesis  $\chi_{\mathbf{A}}$  is irreducible, then  $\pi_{\mathbf{A},\mathbf{x}} = \chi_{\mathbf{A}}$  and  $\dim_{\mathbb{F}_q} F = \dim_{\mathbb{F}_q} \mathbb{F}_q[X]/(\chi_{\mathbf{A}}) = m$ . Therefore,  $F = \mathbb{F}_q^m$  and hence contains  $\mathbf{y}$ , which proves (ii).

Using the fact that  $\mathbb{F}_q^m$  is isomorphic as a vector space with  $\mathbb{F}_{q^m}$ , the matrices in  $\mathbb{F}_q[A]$  represent the multiplications by elements of  $\mathbb{F}_{q^m}$  in some given basis. The matrix  $\Theta$  is nothing but a matrix representation of the Frobenius map  $x \mapsto x^q$  in this basis. This proves (iii).

Let  $\Gamma$  be a matrix such that  $\mathbb{F}_q[\mathbf{A}]$  is globally invariant by conjugation under  $\Gamma$ . This conjugation map induces a non-trivial automorphism of the field  $\mathbb{F}_q[\mathbf{A}] = \mathbb{F}_{q^m}$ , which is nothing but an iterate of the Frobenius. Therefore,

$$\forall P(\mathbf{A}) \in \mathbb{F}_q[\mathbf{A}], \quad \mathbf{\Gamma} P(\mathbf{A}) \mathbf{\Gamma}^{-1} = P(\mathbf{A})^{q^j}$$

for some  $j \in \{0, ..., m-1\}$ . Now,  $\Gamma^{-1}\Theta^j$  commutes with any element of  $\mathbb{F}_q[\mathbf{A}]$  and then (iv) can be deduced from (i).

Let  $\mathbf{x} \in \mathbb{F}_q^m \setminus \{0\}$ , reasoning as in the proof of (ii) we see that  $(\mathbf{x}^{\top}, \mathbf{A}\mathbf{x}^{\top}, \dots, \mathbf{A}^{m-1}\mathbf{x}^{\top})$  is a basis of  $\mathbb{F}_q^m$ . Let  $\mathbf{P}_0$  be the transition matrix from the canonical basis to this basis. This matrix can be computed in polynomial time and Then,  $\mathbf{P}_0^{-1}\mathbf{A}\mathbf{P}_0$  is nothing but the companion matrix of

 $\chi_{\mathbf{A}}$ . Similarly, one can compute  $\mathbf{P}_1$  such that  $\mathbf{P}_1^{-1}\mathbf{B}\mathbf{P}_1$  equals this companion matrix of  $\chi_{\mathbf{A}}$ . The matrix  $\mathbf{P} \stackrel{def}{=} \mathbf{P}_1 \mathbf{P}_0^{-1}$  yields (v).

This technical lemma has the following consequence on the matrix representations of  $\mathbb{F}_{q^m}$ .

**Corollary 27.** — Let  $S_1, S_2 \subseteq \mathcal{M}_m(\mathbb{F}_q)$  be two matrix representations of  $\mathbb{F}_{q^m}$ . Then there exists a matrix  $\mathbf{P} \in \mathbf{GL}_m(\mathbb{F}_q)$  such that  $S_1 = \mathbf{P}^{-1}S_2\mathbf{P}$ . In addition,  $\mathbf{P}$  can be computed using a polynomial time *F*-algorithm.

*Proof.* — First, compute matrices  $\mathbf{A}_1, \mathbf{A}_2 \in \mathcal{M}_m(\mathbb{F}_q)$  such that  $\mathcal{S}_1 = \mathbb{F}_q[\mathbf{A}_1]$  and  $\mathcal{S}_2 = \mathbb{F}_q[\mathbf{A}_2]$ . Such matrices can be computed in polynomial time. Denoting by  $\chi_{\mathbf{A}_1}, \chi_{\mathbf{A}_2}$  their respective characteristic polynomials we have

$$\mathcal{S}_1 = \mathbb{F}_q[\mathbf{A}_1] \simeq \mathbb{F}_q[X]/(\chi_{\mathbf{A}_1}) \quad \text{and} \quad \mathcal{S}_2 = \mathbb{F}_q[\mathbf{A}_2] \simeq \mathbb{F}_q[X]/(\chi_{\mathbf{A}_2}).$$

Using an F-algorithm, one can compute a root of  $\chi_{\mathbf{A}_2}$  in  $\mathbb{F}_q[\mathbf{A}_1] \simeq \mathbb{F}_q[X]/\chi_{\mathbf{A}_1}$ , and denote it by  $P(\mathbf{A}_1)$ . Then  $P(\mathbf{A}_1)$  has the same characteristic polynomial as  $\mathbf{A}_2$  and, from Lemma 26(v), there is a matrix  $\mathbf{P}$  which can be computed in polynomial time such that,  $\mathbf{A}_2 = \mathbf{P}^{-1}P(\mathbf{A}_1)\mathbf{P}$ . Moreover, for dimensional reasons,  $\mathbb{F}_q[\mathbf{A}_1] = \mathbb{F}_q[P(\mathbf{A}_1)]$  and hence

$$\mathcal{S}_2 = \mathbb{F}_q[\mathbf{A}_2] = \mathbf{P}^{-1}\mathbb{F}_q[\mathbf{A}_1]\mathbf{P} = \mathbf{P}^{-1}\mathcal{S}_1\mathbf{P}.$$

So, the first part of our algorithm will consist in computing the left stabilizer algebras of  $\mathscr{C}_{Mat}$ and  $\mathscr{D}_{Mat}$  (which are assumed to be representations of  $\mathbb{F}_{q^m}$ ). Next, according to Corollary 27, the stabilizer algebras are conjugated and one can compute (in polynomial time if  $q = m^{O(1)}$  otherwise with a Las Vegas method)  $\mathbf{P} \in \mathbf{GL}_m(\mathbb{F}_q)$  such that:

$$\operatorname{Stab}_{\operatorname{left}}(\mathscr{D}_{\operatorname{Mat}}) = \mathbf{P}^{-1} \operatorname{Stab}_{\operatorname{left}}(\mathscr{C}_{\operatorname{Mat}}) \mathbf{P}.$$

Replacing  $\mathscr{D}_{Mat}$  by the left equivalent code  $\mathbf{P}\mathscr{D}_{Mat}$ , then the codes turn out to have the same left stabilizer algebra. Once we are reduced to the case where the two codes have the same left stabilizer algebra, the following statement asserts that we are almost reduced to solving and instance of MCRE.

**Proposition 28.** — Let  $\mathscr{C}_{Mat}$ ,  $\mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  be two left equivalent matrix codes codes whose left stabilizer algebras are equal and are representations of  $\mathbb{F}_{q^m}$ , then there exists  $j \in \{0, \ldots, m-1\}$  such that

$$\mathscr{C}_{Mat} = \mathbf{\Theta}^{\jmath} \mathscr{D}_{Mat}$$

where  $\Theta$  is a matrix defined in 26 (iii).

*Proof.* — The codes are suppose to be left equivalent, then  $\mathscr{C}_{Mat} = \mathbf{S}\mathscr{D}_{Mat}$  for some  $\mathbf{S} \in \mathbf{GL}_n(\mathbb{F}_q)$ . In addition, if they have the same left stabilizer algebra  $\mathcal{S}$ , which is supposed to be of the form  $\mathbb{F}_q[\mathbf{A}] \simeq \mathbb{F}_{q^m}$  for some  $\mathbf{A} \in \mathcal{M}_m(\mathbb{F}_q)$  with an irreducible characteristic polynomial. Then, from a "left version" of Proposition 14,  $\mathcal{S}$  is stable by conjugation by  $\mathbf{S}$ . Thus, from Lemma 26(*iv*), we have

$$\mathbf{S} = \mathbf{\Theta}^j P(\mathbf{A})$$

for some  $j \in \{0, ..., m-1\}$  and  $P(\mathbf{A}) \in \mathbb{F}_q[\mathbf{A}]^{\times}$ . Therefore, since  $P(\mathbf{A}) \in \mathcal{S} = \text{Stab}_{\text{left}}(\mathscr{D}_{\text{Mat}})$ , then  $\mathbf{S}\mathscr{D}_{\text{Mat}} = \mathbf{\Theta}^j \mathscr{D}_{\text{Mat}}$ , which concludes the proof.

**Corollary 29.** — Let  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  which are equivalent, i.e. there exists  $\mathbf{S} \in \mathbf{GL}_m(\mathbb{F}_q)$  and  $\mathbf{T} \in \mathbf{GL}_n(\mathbb{F}_q)$  such that  $\mathscr{C}_{Mat} = \mathbf{S}\mathscr{D}_{Mat}\mathbf{T}$ . Suppose that the two codes have the same left stabilizer algebra which is a representation of  $\mathbb{F}_{q^m}$ . Then, denoting by  $\Theta$  the matrix of Lemma 26 (iii),  $\mathscr{C}_{Mat}$  is right equivalent to  $\Theta^j \mathscr{D}_{Mat}$  for some  $j \in \{0, \ldots, m-1\}$ .

*Proof.* — Apply Proposition 28 to the pair ( $\mathscr{C}_{Mat}, \mathscr{D}_{Mat}\mathbf{T}$ ).

These results yield to Algorithm 3 which decides HV-MCE in polynomial efficiently when left stabilizer algebras are equal to  $\mathbb{F}_{q^m}$ .

**Algorithm 3:** An algorithm to decide the equivalence problem for  $\mathbb{F}_{q^m}$ -linear code with left stabilizer algebras equal to  $\mathbb{F}_{q^m}$ .

**Input** : Two codes  $\mathscr{C}_{Mat}$ ,  $\mathscr{D}_{Mat} \subseteq Mat_{m,n}(\mathbb{F}_q)$ **Output:** Codes are equivalent or not

**Theorem 30.** — Let  $\mathscr{C}_{Mat}$ ,  $\mathscr{D}_{Mat}$  be two spaces of matrices representing  $\mathbb{F}_{q^m}$ -linear codes and whose left stabilizer algebra is a representation of  $\mathbb{F}_{q^m}$ . Algorithm 3 is a polynomial time F-algorithm which succeeds to decide if they are equivalent.

*Proof.* — The **for** loop is made at most m times, which is polynomial in the size of the input. According to Proposition 27, there is a polynomial time F-algorithm computing **P**. Next, using the polynomial time F-algorithm described in Section 4.2 one can decide the right equivalence of  $\mathscr{C}_{Mat}$  and  $\Theta^t \mathscr{D}_{Mat}$ .

5.3. When the left stabilizer algebras strictly contain representations of  $\mathbb{F}_{q^m}$ . — In the general case, it is possible that the left stabilizer algebras contain representations of  $\mathbb{F}_{q^m}$  as proper subspaces. To understand the way to proceed, we first have to classify sub-algebras of  $\mathcal{M}_m(\mathbb{F}_q)$  containing a representation of  $\mathbb{F}_{q^m}$ , which is the point of the next statement.

**Proposition 31.** — Let  $S \subseteq \mathcal{M}_m(\mathbb{F}_q)$  be an algebra such that

$$\mathbb{F}_q[\mathbf{A}] \simeq \mathbb{F}_{q^m} \subsetneq \mathcal{S} \subseteq \mathcal{M}_m(\mathbb{F}_q),$$

then, S is isomorphic to a matrix algebra of the form  $\mathcal{M}_{\frac{m}{\ell}}(\mathbb{F}_{q^{\ell}})$  for some  $\ell$  dividing m. In addition, any  $\mathbf{C} \in \mathcal{M}_m(\mathbb{F}_q)$  which commutes with any element of the centre of S, then  $\mathbf{C} \in S$ .

*Proof.* — First, let us prove that S is semi-simple, namely that its Jacobson radical is trivial. Indeed, let  $\mathbf{N} \in S \setminus \{0\}$  be such that for any  $\mathbf{M} \in S$ ,  $\mathbf{N}\mathbf{M}$  is nilpotent. Then, there exists  $\mathbf{x} \in \mathbb{F}_{q^m}$  such that  $\mathbf{N}\mathbf{x}^\top \neq 0$  and, from Lemma 26 (ii), there exists  $P(\mathbf{A}) \in \mathbb{F}_q[\mathbf{A}]$  such that  $P(\mathbf{A})\mathbf{N}\mathbf{x}^\top = \mathbf{x}^\top$ . Thus,  $P(\mathbf{A})\mathbf{N} \in S$  and has a non-zero fixed point which contradicts its nilpotence. Therefore,  $\operatorname{Rad}(S) = \{0\}$  and hence S is semi-simple.

Now let us prove that S is simple. For that, using [**DK94**, Cor. 1.7.8], we only need to prove that its centre is simple. But, since S contains  $\mathbb{F}_q[\mathbf{A}]$ , from Lemma 26 (i), its centre is contained in  $\mathbb{F}_q[\mathbf{A}] \simeq \mathbb{F}_{q^m}$  and hence is isomorphic to  $\mathbb{F}_{q^\ell}$  for some  $\ell$  dividing m. In particular, it is simple and so is S. Therefore, S is isomorphic to  $\mathcal{M}_r(\mathbb{F}_{q^\ell})$  for some positive r and there remains to prove that  $r = \frac{m}{\ell}$ .

From Lemma 26(*i*) any element  $\mathbb{F}_q[\mathbf{A}] \simeq \mathbb{F}_{q^m}$  in  $\mathcal{S}$  commuting with any element of  $\mathbb{F}_q[\mathbf{A}]$  is in  $\mathbb{F}_q[\mathbf{A}]$ . Using [**DK94**, Thm. 4.4.6(3)] we deduce that the  $\mathbb{F}_{q^\ell}$ -dimension of  $\mathcal{S}$  is  $(\frac{m}{\ell})^2$ . This proves that  $r = \frac{m}{\ell}$ . The former reference permits also to prove that a matrix in  $\mathcal{M}_m(\mathbb{F}_q)$  commuting with any element of  $\mathcal{S}$  is in  $\mathcal{S}$ .

Before describing our algorithm to decide equivalence when the left stabilizer algebras strictly contain a representation of  $\mathbb{F}_{q^m}$ , we need the following statement, which is an analogue of 29 in the current situation. Note that the current situation is in some sense more favorable since we no longer need to take care of a possible action of the Frobenius morphism.

**Proposition 32.** — Let  $\mathscr{C}_{Mat}, \mathscr{D}_{Mat} \subseteq \mathcal{M}_{m,n}(\mathbb{F}_q)$  be two left equalvalent matrix codes, i.e. such that

$$\exists \mathbf{P} \in \mathbf{GL}_m(\mathbb{F}_q), \ \mathbf{P}\mathscr{C}_{Mat} = \mathscr{D}_{Mat}$$

 $Suppose \ also \ that$ 

$$tab_{left}(\mathscr{C}_{Mat}) = Stab_{left}(\mathscr{D}_{Mat}) \simeq \mathcal{M}_{\frac{m}{4}}(\mathbb{F}_{q^{\ell}}),$$

for some positive  $\ell$  dividing m. Then,  $\mathscr{C}_{Mat} = \mathscr{D}_{Mat}$ .

S

*Proof.* — Let  $S \stackrel{def}{=} \text{Stab}_{\text{left}}(\mathscr{C}_{\text{Mat}}) = \text{Stab}_{\text{left}}(\mathscr{D}_{\text{Mat}})$  Using a left version of 14, we deduce that S is stable by conjugation by **P**. Since S is a central simple algebra over  $\mathbb{F}_{q^{\ell}}$ , from Skolem–Noether Theorem [**DK94**, Cor. 4.4.3], its automorphisms are inner and hence there exists  $\mathbf{R} \in S$  such that

$$orall \mathbf{S} \in \mathcal{S}, \; \mathbf{P}^{-1}\mathbf{S}\mathbf{P} = \mathbf{R}^{-1}\mathbf{S}\mathbf{R}.$$

Therefore, the matrix  $\mathbf{RP}^{-1}$  is in the *centraliser* of S in  $\mathcal{M}_m(\mathbb{F}_q)$ , (*i.e.* the subalgebra of  $\mathcal{M}_m(\mathbb{F}_q)$  of matrices commuting with any element of S). The centraliser contains the centre of S, which is isomorphic to  $\mathbb{F}_{q^\ell}$ . Using [**DK94**, Thm. 4.4.6(3)], we deduce that the centraliser and the centre have the same dimension and hence are equal. Consequently  $\mathbf{RP}^{-1}$  is in  $S = \text{Stab}_{\text{left}}(\mathscr{C}_{\text{Mat}})$  and hence so is **P**. Thus,  $\mathscr{C}_{\text{Mat}} = \mathbf{P}\mathscr{C}_{\text{Mat}} = \mathscr{D}_{\text{Mat}}$ .

With Proposition 31 in hand, we can deduce an algorithm for solving this general case. We describe it as follows.

- 1. Compute the left stabilizer algebras  $\operatorname{Stab}_{\operatorname{left}}(\mathscr{C}_{\operatorname{Mat}}), \operatorname{Stab}_{\operatorname{left}}(\mathscr{D}_{\operatorname{Mat}});$
- 2. Compute their centres  $Z(\operatorname{Stab}_{\operatorname{left}}(\mathscr{C}_{\operatorname{Mat}})), Z(\operatorname{Stab}_{\operatorname{left}}(\mathscr{D}_{\operatorname{Mat}}))$  which are isometric to a same  $\mathbb{F}_{q^{\ell}}$  for some  $\ell > 0$ , otherwise codes are not equivalent;
- 3. These algebras are respectively isomorphic to  $\mathbb{F}_q[\mathbf{A}_1]$ ,  $\mathbb{F}_q[\mathbf{A}_2]$  for some  $\mathbf{A}_1, \mathbf{A}_2 \in \mathcal{M}_m(\mathbb{F}_q)$ whose minimal polynomial is irreducible of degree  $\ell$ . A very similar argument as Corollary 29 permits to assert that the algebras are conjugated and one can compute a matrix  $\mathbf{P} \in \mathbf{GL}_m(\mathbb{F}_q)$  such that  $\mathbb{F}_q[\mathbf{A}_1] = \mathbf{P}\mathbb{F}_q[\mathbf{A}_2]\mathbf{P}^{-1}$ .
- 4. Replacing  $\mathscr{D}_{Mat}$  by  $\mathbf{P}\mathscr{D}_{Mat}$  so that the stabilizer algebras of the codes have the same centre Z. But, from Proposition 31, the stabilizer algebras are then the same since they both equal to the commutator of Z in  $\mathcal{M}_m(\mathbb{F}_q)$ .
- 5. Now, since the two codes have the same left stabiliser algebra, from Proposition 32, they are equivalent if and only if they are right equivalent and we can decide this right equivalence using the algorithm of Section 4.

This enables to prove the following theorem.

**Theorem 33.** — HV-MCE is in  $\mathcal{P}$  if  $q = (mn)^{O(1)}$  and in  $\mathcal{ZPP}$  in the general case.

### 6. Reduction from the Code Equivalence Problem in Hamming Metric

The objective of the present section is to prove that the Matrix Code Equivalence Problem MCE (Problem 1) for rank metric codes is at least as hard as the Monomial Equivalence Problem. Compared to the reduction presented in Section 3.3, the following one treats search versions of the problem and is somehow more explicit.

From the generator matrices point of view, to decide if two codes are monomially equivalent may be formulated as:

- Instance : Two matrices  $\mathbf{A}, \mathbf{B} \in \mathcal{M}_{k,n}(\mathbb{F}_q)$ .
- Decision : it exists  $\mathbf{S} \in \mathbf{GL}_k(\mathbb{F}_q)$ ,  $\mathbf{P} \in \mathcal{M}_n(\mathbb{F}_q)$  be a permutation matrix and a diagonal matrix  $\mathbf{D} \in \mathbf{GL}_n(\mathbb{F}_q)$  such that:

#### $\mathbf{A} = \mathbf{SBDP}.$

Without loss of generality we can assume that the columns of  $\mathbf{A}$  (resp.  $\mathbf{B}$ ) are pairwise linearly independent. If not, we can remove the redundant columns without changing the answer of the problem.

Let  $\mathbf{a}_1^{\top}, \dots, \mathbf{a}_n^{\top}, \mathbf{b}_1^{\top}, \dots, \mathbf{b}_n^{\top} \in \mathbb{F}_q^k$  denote the columns of **A** and **B** respectively. Similarly to the previous section, we keep on considering that vectors are row matrices, which explains why we apply the transposition operator to these column vectors. For any vector  $\mathbf{x} \in \mathbb{F}_q^k$  and any  $1 \leq i \leq n$ , we define the  $n \times k$  matrix  $\mathbf{Row}_i(\mathbf{x})$  over  $\mathbb{F}_q$  whose only non-zero row is the *i*-th one which equals  $\mathbf{x}$ :

$$\mathbf{Row}_i(\mathbf{x}) \stackrel{def}{=} \begin{pmatrix} 0 & \cdots & 0 \\ & \mathbf{x} & \overleftarrow{0} \\ 0 & \cdots & 0 \end{pmatrix} - i.$$

We now build in polynomial time in the size of **A** and **B** the following matrix codes of  $\mathcal{M}_{k+n,k}(\mathbb{F}_q)$ :

$$\mathscr{C}_{\text{Mat}} \stackrel{def}{=} \left\{ \sum_{i=1}^{n} \lambda_i \begin{pmatrix} \mathbf{a}_i^{\top} \mathbf{a}_i \\ \mathbf{Row}_i(\mathbf{a}_i) \end{pmatrix} : \lambda_i \in \mathbb{F}_q \right\},$$
$$\mathscr{D}_{\text{Mat}} \stackrel{def}{=} \left\{ \sum_{i=1}^{n} \lambda_i \begin{pmatrix} \mathbf{b}_i^{\top} \mathbf{b}_i \\ \mathbf{Row}_i(\mathbf{b}_i) \end{pmatrix} : \lambda_i \in \mathbb{F}_q \right\}.$$

The following lemma is crucial for our reduction, it justifies our construction of  $\mathscr{C}_{Mat}$  and  $\mathscr{D}_{Mat}$ .

Lemma 34. — Let  $\mathbf{U} \in \mathbf{GL}_{k+n}(\mathbb{F}_q)$  and  $\mathbf{V} \in \mathbf{GL}_k(\mathbb{F}_q)$  which verify:  $\mathscr{C}_{Mat} = \mathbf{U}\mathscr{D}_{Mat}\mathbf{V}.$ 

Then, there exists a permutation 
$$\sigma \in \mathfrak{S}_n$$
 and  $\alpha_1, \ldots, \alpha_n \in \mathbb{F}_a^{\times}$  such that

$$\begin{pmatrix} \mathbf{a}_i^{\top} \mathbf{a}_i \\ \mathbf{Row}_i(\mathbf{a}_i) \end{pmatrix} = \alpha_{\sigma(i)} \mathbf{U} \begin{pmatrix} \mathbf{b}_{\sigma(i)}^{\top} \mathbf{b}_{\sigma(i)} \\ \mathbf{Row}_{\sigma(i)}(\mathbf{b}_{\sigma(i)}) \end{pmatrix} \mathbf{V}.$$

*Proof.* — By definition, for any  $1 \leq i \leq n$  there exist  $\alpha_1, \ldots, \alpha_n \in \mathbb{F}_q$  such that:

$$\begin{pmatrix} \mathbf{a}_i^{\top} \mathbf{a}_i \\ \mathbf{Row}_i(\mathbf{a}_i) \end{pmatrix} = \sum_{j=1}^n \alpha_j \mathbf{U} \begin{pmatrix} \mathbf{b}_j^{\top} \mathbf{b}_j \\ \mathbf{Row}_j(\mathbf{b}_j) \end{pmatrix} \mathbf{V} = \mathbf{U} \begin{pmatrix} \sum_{j=1}^n \alpha_j \begin{pmatrix} \mathbf{b}_j^{\top} \mathbf{b}_j \\ \mathbf{Row}_j(\mathbf{b}_j) \end{pmatrix} \mathbf{V}.$$

But now, since  $\mathbf{U}$  and  $\mathbf{V}$  are non-singular,

$$\operatorname{Rk}\left(\mathbf{U}\left(\sum_{j=1}^{n}\alpha_{j}\begin{pmatrix}\mathbf{b}_{i}^{\top}\mathbf{b}_{i}\\\mathbf{Row}_{i}(\mathbf{b}_{i})\end{pmatrix}\right)\mathbf{V}\right) = \operatorname{Rk}\left(\sum_{j=1}^{n}\alpha_{j}\begin{pmatrix}\mathbf{b}_{j}^{\top}\mathbf{b}_{j}\\\mathbf{Row}_{j}(\mathbf{b}_{j})\end{pmatrix}\right).$$

Recall that the rows of  $\mathbf{Row}_j(\mathbf{b}_j)$  are all zero but the *j*-th one which equals  $\mathbf{b}_j$ . Here, by hypothesis, the  $\mathbf{b}_j$ 's are pairwise linearly independent. Therefore, if there were at least two non zero  $\alpha_j$ 's, then we would have:

$$\operatorname{Rk}\left(\sum_{j=1}^{n} \alpha_{j} \begin{pmatrix} \mathbf{b}_{j} \mathbf{b}_{j}^{\top} \\ \mathbf{Row}_{j}(\mathbf{b}_{j}^{\top}) \end{pmatrix}\right) \geq 2.$$

Indeed, the considered matrix would have two independent rows in its lower part. A contradiction since  $\begin{pmatrix} \mathbf{a}_i \mathbf{a}_i^\top \\ \mathbf{Row}_i(\mathbf{a}_i^\top) \end{pmatrix}$  has rank 1, which proves the first part of the lemma.

There remains to prove that the map  $i \mapsto j$  is a permutation of  $\{1, \ldots, n\}$ . Suppose that

$$\begin{pmatrix} \mathbf{a}_i^{\top} \mathbf{a}_i \\ \mathbf{Row}_i(\mathbf{a}_i) \end{pmatrix} = \alpha_u \mathbf{U} \begin{pmatrix} \mathbf{b}_u^{\top} \mathbf{b}_u \\ \mathbf{Row}_u(\mathbf{b}_u) \end{pmatrix} \mathbf{V} = \alpha_v \mathbf{U} \begin{pmatrix} \mathbf{b}_v^{\top} \mathbf{b}_v \\ \mathbf{Row}_v(\mathbf{b}_v) \end{pmatrix} \mathbf{V}$$

for two distinct non-zero  $\alpha_u$  and  $\alpha_v$ . Since **U** and **V** are non-singular, the previous equality implies that:

$$\alpha_u \begin{pmatrix} \mathbf{b}_u^\top \mathbf{b}_u \\ \mathbf{Row}_u(\mathbf{b}_u) \end{pmatrix} = \alpha_v \begin{pmatrix} \mathbf{b}_v^\top \mathbf{b}_v \\ \mathbf{Row}_v(\mathbf{b}_v) \end{pmatrix}.$$

This contradicts the definition of  $\mathbf{Row}_u(\mathbf{b}_u)$  and  $\mathbf{Row}_v(\mathbf{b}_v)$ 

Let us now prove that  $(\mathscr{C}_{\text{Vec}}, \mathscr{D}_{\text{Vec}})$  (codes of generator matrices **A** and **B**) is a positive instance of the monomial equivalence problem if and only if  $(\mathscr{C}_{\text{Mat}}, \mathscr{D}_{\text{Mat}})$  is a positive instance of MCE.

6.1. If  $(\mathscr{C}_{\text{Vec}}, \mathscr{D}_{\text{Vec}})$  is a positive instance of the monomial equivalence problem then  $(\mathscr{C}_{\text{Mat}}, \mathscr{D}_{\text{Mat}})$  is a positive instance of MCE.. — It exists  $\mathbf{S} \in \text{GL}_k(\mathbb{F}_q)$ ,  $\mathbf{P} \in \text{GL}_n(\mathbb{F}_q)$  a permutation matrix and  $\mathbf{D} \in \text{GL}_n(\mathbb{F}_q)$  a diagonal matrix such that:

### $\mathbf{A} = \mathbf{SBDP}.$

Denote by  $\alpha_1, \ldots, \alpha_n \in \mathbb{F}_q^{\times}$  the diagonal entries of **D**. Then, for any  $1 \leq i \leq n$ , it exists  $1 \leq j \leq n$  (image of *i* by the permutation given by **P**) such that,

$$\mathbf{a}_i^{\top} = \alpha_j \mathbf{S} \mathbf{b}_j^{\top}$$
 and  $\mathbf{a}_i^{\top} \mathbf{a}_i = \alpha_j^2 \mathbf{S} \mathbf{b}_j^{\top} \mathbf{b}_j \mathbf{S}^{\top}$ 

This gives:

$$\mathscr{C}_{\mathrm{Mat}} = \mathbf{U}\mathscr{D}_{\mathrm{Mat}}\mathbf{V}$$

where

$$\mathbf{U} \stackrel{def}{=} \begin{pmatrix} \mathbf{S} & \mathbf{0} \\ \mathbf{0} & \mathbf{P} \end{pmatrix} \in \mathcal{M}_{k+n}(\mathbb{F}_q) \quad ext{and} \quad \mathbf{V} \stackrel{def}{=} \mathbf{S}^{ op} \in \mathcal{M}_k(\mathbb{F}_q),$$

which are nonsingular matrices. Therefore,  $(\mathscr{C}_{Mat}, \mathscr{D}_{Mat})$  is a positive instance of MCE.

6.2. If  $(\mathscr{C}_{Vec}, \mathscr{D}_{Vec})$  is a positive instance of MCE then  $(\mathscr{C}_{Mat}, \mathscr{D}_{Mat})$  is a positive instance of the monomial equivalence problem. — By definition, there exist two non-singular matrices U and V such that:

$$\mathscr{C}_{Mat} = U \mathscr{D}_{Mat} V.$$

By Lemma 34, for any  $1 \leq i \leq n$ , it exists  $1 \leq \sigma(i) \leq n$  and  $\alpha_{\sigma(i)} \in \mathbb{F}_q^{\times}$  such that:

(7)  
$$\begin{pmatrix} \mathbf{a}_{i}^{\top}\mathbf{a}_{i} \\ \mathbf{Row}_{i}(\mathbf{a}_{i}) \end{pmatrix} = \alpha_{\sigma(i)} \mathbf{U} \begin{pmatrix} \mathbf{b}_{\sigma(i)}^{\top}\mathbf{b}_{\sigma(i)} \\ \mathbf{Row}_{\sigma(i)}(\mathbf{b}_{\sigma(i)}) \end{pmatrix} \mathbf{V}$$
$$= \alpha_{\sigma(i)} \mathbf{U} \begin{pmatrix} \mathbf{b}_{\sigma(i)}^{\top}\mathbf{b}_{\sigma(i)}\mathbf{V} \\ \mathbf{Row}_{\sigma(i)}(\mathbf{b}_{\sigma(i)}\mathbf{V}) \end{pmatrix}.$$

The above matrices are of rank 1 and their row spaces are generated by  $\mathbf{a}_i$  and  $\mathbf{b}_i$ . Therefore as they are equal,  $\mathbf{a}_i$  and  $\mathbf{b}_{\sigma(i)}\mathbf{V}$  are collinear. This yields the monomial equivalence of  $\mathscr{C}_{\text{Vec}}$  and  $\mathscr{D}_{\text{Vec}}$ .

#### Conclusion

This work has presented the equivalence problem of matrix codes endowed with the rank metric as well as some of its natural variants. Our contribution is threefold. First, as summarized in Figure 2, we have shown that the code equivalence problem for matrix codes (MCE) is harder than the monomial equivalence problem. This last problem has been pursued for many years and works on it tend to show its hardness. Thus, the equivalence problem for matrix codes is not likely to be easy and could for instance be considered for cryptographic applications. On the other hand, if one restricts instances in the equivalence problem to  $\mathbb{F}_{q^m}$ -linear codes (particular matrix codes that are appealing for cryptographic applications) or only considers the right-equivalence then it leads to an easy problem *in the worst case*. Using algorithms from the mathematical field of algebras we have shown that these problems fall down in  $\mathcal{P}$  or  $\mathcal{ZPP}$  depending of the ground field size q. To our opinion, this work may have extensions that are of interest.

**Classification of matrix codes.** Our algorithm to decide if two matrix codes are right equivalent takes advantage of the fact that right-equivalence only shifts (multiplicatively) right stabilizer algebras of codes by a non-singular matrix. In this way, stabilizer algebras share, roughly speaking, the same algebraic structure and as we have shown, by decomposing them we can decide if two codes are right equivalent. Interestingly, our algorithm decomposes right stabilizer algebras as "minimal atoms". Then it punctures each code according to these atoms and tries to put them in correspondence.



FIGURE 2. Reductions around the code equivalence problem with our contribution where notation " $A \longrightarrow B$ " means that "Problem A reduces to Problem B in polynomial time".

This may remind techniques that are used in *Support Splitting* approaches [Leo82, Sen00]. It consists for two codes endowed with the Hamming metric to puncture them and to look if they both satisfy the same "invariants". In [Leo82] is proposed to look at the weight distribution of the punctured codes while in [Sen00] it is this distribution but for hulls. In this light, decomposition of right stabilizer algebras can be seen as a good invariant for matrix codes. By *good* we mean that it is an efficient tool to solve equivalence problems. On the other hand, such invariant remains weakly discriminant since for arbitrary codes, the stabilizer algebra will be trivial.

This decomposition of stabilizer algebras could for instance be used to classify matrix codes. Furthermore, stabilizer algebras may help us to identify some properties that could be useful for decoding algorithms. This would be particularly interesting as we know very few families of matrix codes that we can decode efficiently. Except simple codes [SKK10] (that have a trivial structure), all matrix codes with an efficient decoding algorithm are  $\mathbb{F}_{q^m}$ -linear [Gab85, GMRZ13] which exactly corresponds to the case where codes have a rich left stabilizer algebraic structure (that we use to decide the equivalence of these codes).

Finally, it should be emphasized that in the present article, we made the choice to work only over finite fields, which permits a simpler treatment since any central simple algebra is isomorphic to a matrix algebra in this setting (trivial Brauer group). However, some literature exists on codes over infinite fields [Aug14, ALR13, ALR18, ACLN20] and the similar question of solving code equivalence over arbitrary fields makes sense and is left as an open question. Such a treatment will probably represent a slightly harder task while involving non trivial central simple algebras.

**Our Reduction.** We can deduce from our reduction that any algorithm solving a "particular" instance of the matrix code equivalence problem (codes are generated by matrices of rank one) provides an algorithm solving the monomial equivalence problem. Therefore, this reduction gives a new manner to solve the monomial equivalence problem. For instance we could model the matrix equivalence problem into a system of polynomial equations that we would solve with Gröbner bases techniques. This approach, thanks to our reduction, would give a new modelling of the monomial equivalence problem. It may be interesting to study the complexity of this approach as Gröbner and to compare with the results of [**ST17**] where permutation equivalence of codes is treating by solving a system of polynomial equations.

### References

- [AAB<sup>+</sup>17] Carlos Aguilar Melchor, Nicolas Aragon, Slim Bettaieb, Loïc Bidoux, Olivier Blazy, Jean-Christophe Deneuville, Philippe Gaborit, and Gilles Zémor. Rank quasi cyclic (RQC). First round submission to the NIST post-quantum cryptography call, November 2017.
- [ABD<sup>+</sup>19] Nicolas Aragon, Olivier Blazy, Jean-Christophe Deneuville, Philippe Gaborit, Adrien Hauteville, Olivier Ruatta, Jean-Pierre Tillich, Gilles Zémor, Carlos Aguilar Melchor, Slim

Bettaieb, Loïc Bidoux, Bardet Magali, and Ayoub Otmani. ROLLO (merger of Rank-Ouroboros, LAKE and LOCKER). Second round submission to the NIST post-quantum cryptography call, March 2019.

- [ABG<sup>+</sup>19] Nicolas Aragon, Olivier Blazy, Philippe Gaborit, Adrien Hauteville, and Gilles Zémor. Durandal: a rank metric based signature scheme. In Advances in Cryptology - EUROCRYPT 2019 -38th Annual International Conference on the Theory and Applications of Cryptographic Techniques, Darmstadt, Germany, May 19-23, 2019, Proceedings, Part III, volume 11478 of LNCS, pages 728–758. Springer, 2019.
- [ACLN20] Daniel Augot, Alain Couvreur, Julien Lavauzelle, and Alessandro Neri. Rank-metric codes over arbitrary Galois extensions and rank analogues of Reed–Muller codes. arXiv:2006.14489, June 2020.
- [ALR13] Daniel Augot, Pierre Loidreau, and Gwezheneg Robert. Rank metric and Gabidulin codes in characteristic zero. In 2013 IEEE International Symposium on Information Theory, pages 509–513. IEEE, 2013.
- [ALR18] Daniel Augot, Pierre Loidreau, and Gwezheneg Robert. Generalized Gabidulin codes over fields of any characteristic. *Des., Codes Cryptogr.*, 86(8):1807–1848, 2018.
- [Aug14] Daniel Augot. Generalization of Gabidulin codes over fields of rational functions. In 21st International Symposium on Mathematical Theory of Networks and Systems (MTNS 2014), 2014.
- [Ber68] Elwyn R. Berlekamp. Factoring polynomials over finite fields. In E. R. Berlekamp, editor, Algebraic Coding Theory, chapter 6. McGraw-Hill, 1968.
- [Ber70] Elwyn R. Berlekamp. Factoring polynomials over large finite fields. *Math. Comp.*, 24:713–735, 1970.
- [Ber03] Thierry P. Berger. Isometries for rank distance and permutation group of Gabidulin codes. *IEEE Trans. Inform. Theory*, 49(11):3016–3019, 2003.
- [BG13] Aleams Barra and Heide Gluesing-Luerssen. MacWilliams extension theorems and the localglobal property for codes over rings. *CoRR*, abs/1307.7159, 2013.
- [BOS19] Magali Bardet, Ayoub Otmani, and Mohamed Saeed-Taha. Permutation code equivalence is not harder than graph isomorphism when hulls are trivial. In Proc. IEEE Int. Symposium Inf. Theory - ISIT 2019, pages 2464–2468, July 2019.
- [Bre11] Murray R. Bremner. How to compute the Wedderburn decomposition of a finite-dimensional associative algebra. *Groups Complex. Cryptol.*, 3(1):47–66, 2011.
- [CZ81] David G. Cantor and Hans Zassenhaus. A new algorithm for factoring polynomials over finite fields. Math. Comp., pages 587–592, 1981.
- [Del78] Philippe Delsarte. Bilinear forms over a finite field, with applications to coding theory. J. Comb. Theory, Ser. A, 25(3):226–241, 1978.
- [DK94] Yurj A. Drodz and Vladimir V. Kirichenko. *Finite dimensional algebras*. Springer–Verlag Berlin Heidelberg, 1994. Original Russian edition published by: Publisher of Kiev State University, Kiev 1980, Translated by V. Dlab.
- [Feu09] Thomas Feulner. The automorphism groups of linear codes and canonical representatives of their semilinear isometry classes. *Adv. Math. Commun.*, 3(4):363–383, 2009.
- [FR85] Katalin Friedl and Lajos Rónyai. Polynomial time solutions of some problems of computational algebra. In Proceedings of the Seventeenth Annual ACM Symposium on Theory of Computing, STOC '85, pages 153–162, New York, NY, USA, 1985. Association for Computing Machinery.
- [Gab85] Ernest M. Gabidulin. Theory of codes with maximum rank distance. Problemy Peredachi Informatsii, 21(1):3–16, 1985.
- [Gab08] Ernst M. Gabidulin. Attacks and counter-attacks on the GPT public key cryptosystem. Des. Codes Cryptogr., 48(2):171–177, 2008.
- [GMRZ13] Philippe Gaborit, Gaétan Murat, Olivier Ruatta, and Gilles Zémor. Low rank parity check codes and their application to cryptography. In Proceedings of the Workshop on Coding and Cryptography WCC'2013, Bergen, Norway, 2013.
- [GO01] Ernst M. Gabidulin and Alexei V. Ourivski. Modified GPT PKC with right scrambler. Electron. Notes Discrete Math., 6:168–177, 2001.
- [GPT91] Ernst M. Gabidulin, A. V. Paramonov, and O. V. Tretjakov. Ideals over a non-commutative ring and their applications to cryptography. In Advances in Cryptology - EUROCRYPT'91, number 547 in LNCS, pages 482–489, Brighton, April 1991.

- [GRSZ14] Philippe Gaborit, Olivier Ruatta, Julien Schrek, and Gilles Zémor. Ranksign: An efficient signature algorithm based on the rank metric (extended version on arxiv). In *Post-Quantum Cryptography 2014*, volume 8772 of *LNCS*, pages 88–107. Springer, 2014.
- [HP03] W. Cary Huffman and Vera Pless. *Fundamentals of error-correcting codes*. Cambridge University Press, Cambridge, 2003.
- [HR14] Ishay Haviv and Oded Regev. On the lattice isomorphism problem. In Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA '14, pages 391– 404, USA, 2014. Society for Industrial and Applied Mathematics.
- [Hua51] Loo-Keng Hua. A theorem on matrices over a field and its applications. J. Chinese Math. Soc., 1(2):109–163, 1951.
- [Leo82] Jeffrey Leon. Computing automorphism groups of error-correcting codes. *IEEE Trans. Inform. Theory*, 28(3):496–511, 1982.
- [LN16] Dirk Liebhold and Gabriele Nebe. Automorphism groups of gabidulin-like codes. *CoRR*, abs/1603.09565, 2016.
- [LTZ17] Guglielmo Lunardon, Rocco Trombetti, and Yue Zhou. On kernels and nuclei of rank metric codes. Journal of Algebraic Combinatorics, 46, 2017.
- [Mac62] Florence J. MacWilliams. Combinatorial properties of elementary abelian groups. PhD thesis, Radcliffe College, 1962.
- [Mor14] Katherine Morrison. Equivalence for rank-metric and matrix codes and automorphism groups of gabidulin codes. *IEEE Trans. Inform. Theory*, 60(11):7035–7046, 2014.
- [NPHT20] Alessandro Neri, Sven Puchinger, and Anna-Lenna Horlemann-Trautmann. Equivalence and characterizations of linear rank-metric codes based on invariants. *Linear Algebra and its Applications*, pages 418–469, 2020.
- [Ore33] Oystein Ore. On a special class of polynomials. Trans. Amer. Math. Soc., 35(3):559–584, 1933.
- [PR97] Erez Petrank and Ron. Roth. Is code equivalence easy to decide? IEEE Trans. Inform. Theory, 43(5):1602–1604, 1997.
- [Rón90] Lajos Rónyai. Computing the structure of finite algebras. J. Symbolic Comput., 9(3):355–373, 1990.
- [Sen97] Nicolas Sendrier. On the dimension of the hull. SIAM J. Discrete Math., 10(2):282–293, 1997.
- [Sen00] Nicolas Sendrier. Finding the permutation between equivalent linear codes: The support splitting algorithm. *IEEE Trans. Inform. Theory*, 46(4):1193–1203, 2000.
- [SKK10] Danilo Silva, Frank R. Kschischang, and Ralf Kötter. Communication over finite-field matrix channels. *IEEE Trans. Information Theory*, 56(3):1296–1305, 2010.
- [SS13] Nicolas Sendrier and Dimitris E. Simos. The hardness of code equivalence over and its application to code-based cryptography. In *Post-Quantum Cryptography 2013*, volume 7932 of *LNCS*, pages 203–216. Springer, 2013.
- [ST17] Mohamed Ahmed Saeed-Taha. Algebraic Approach for Code Equivalence. PhD thesis, Normandy University, France, 2017.

ALAIN COUVREUR, Inria • LIX, CNRS UMR 7161, École Polytechnique, 1 rue Honoré d'Estienne d'Orves, 91120 PALAISEAU CEDEX • *E-mail* : alain.couvreur@inria.fr

THOMAS DEBRIS-ALAZARD, INIA • LIX, CNRS UMR 7161, École Polytechnique, 1 rue Honoré d'Estienne d'Orves, 91120 PALAISEAU CEDEX • *E-mail* : thomas.debris@inria.fr

PHILIPPE GABORIT, Xlim, CNRS, UMR 7252, Université de Limoges, 123, avenue Albert Thomas, 87060 Limoges Cedex • *E-mail* : gaborit@unilim.fr