

INSTABILITY OF SMALL-AMPLITUDE PERIODIC WAVES FROM FOLD-HOPF BIFURCATION

SHUANG CHEN

Center for Mathematical Sciences
Huazhong University of Sciences and Technology
Wuhan, Hubei 430074, P. R. China
Email: schen@hust.edu.cn; matschen@163.com

JINQIAO DUAN¹

Department of Applied Mathematics
Illinois Institute of Technology
Chicago, IL 60616, USA
Email: duan@iit.edu

ABSTRACT. We study the existence and stability of small-amplitude periodic waves emerging from fold-Hopf equilibria in a system of one reaction-diffusion equation coupled with one ordinary differential equation. This coupled system includes the FitzHugh-Nagumo system, caricature calcium models, consumer-resource models and other models in the real-world applications. Based on the recent results on the averaging theory, we solve periodic solutions in related three-dimensional systems and then prove the existence of periodic waves arising from fold-Hopf bifurcations. Numerical computation in [J. Tsai, W. Zhang, V. Kirk, and J. Sneyd, *SIAM J. Appl. Dyn. Syst.* 11 (2012), 1149–1199] once suggested that the periodic waves from fold-Hopf bifurcations in a caricature calcium model are spectrally unstable, yet without a proof. After analyzing the linearization about periodic waves by the relatively bounded perturbation, we prove the instability of small-amplitude periodic waves through a perturbation of the unstable spectra for the linearizations about the fold-Hopf equilibria. As an application, we prove the existence and stability of small-amplitude periodic waves from fold-Hopf bifurcations in the FitzHugh-Nagumo system with an applied current.

1. INTRODUCTION

Traveling waves play a pivotal role in the study of wave propagation in biological, chemical and physical systems. For example, the traveling waves in FitzHugh-Nagumo equation [20, 40]

Date: December 22, 2024.

2020 Mathematics Subject Classification. Primary 35B10, 35B35; Secondary 34C23, 35B32.

Key words and phrases. Fold-Hopf bifurcation, three-dimensional dynamical system, periodic traveling wave, stability theory, FitzHugh-Nagumo system.

¹The corresponding author

indicate the propagation of nerve impulses in axons, and in the caricature calcium models [34] describe the changes of the free cytoplasmic calcium concentration and the free calcium concentration in the endoplasmic reticulum. Many works have been devoted to homoclinic waves (pulses), heteroclinic waves (fronts and backs) and large-amplitude periodic waves in reaction-diffusion equations coupled with ordinary differential equations in the past.

In the present paper, we focus on the small-amplitude periodic waves emerging from the fold-Hopf bifurcation (also called the Hopf-zero or Gavrilov-Guckenheimer bifurcation) [22, 36]. More precisely, we investigate the small-amplitude periodic traveling waves from fold-Hopf bifurcations in a system of one reaction-diffusion equation coupled with one ordinary differential equation

$$(1.1) \quad \begin{aligned} \frac{\partial u}{\partial t} &= u_{xx} + f(u, w, \alpha), \\ \frac{\partial w}{\partial t} &= g(u, w, \alpha). \end{aligned}$$

where $u(x, t)$ and $w(x, t)$ are system states, x is a one-dimensional spatial variable, t is time, the parameter vector α is in \mathbb{R}^m with $m \geq 1$, and f and g are sufficiently smooth functions. When the diffusion rate of u in (1.1) is a nonzero constant instead of one, we can change it to be one by a simple rescaling. The systems of the form (1.1) are widely used to understand the mechanisms for various phenomena in numerous experiences. For example, the coupled system (1.1) includes the FitzHugh-Nagumo systems with or without an applied current [11, 20, 40], caricature calcium models [47, 49] and consumer-resource models [26, 48].

A solution $(u(x, t), w(x, t))$ of (1.1) is said to be a *traveling wave* if there exists a real number c and the function $(\phi(\cdot), \psi(\cdot)) : \mathbb{R} \rightarrow \mathbb{R}^2$ such that

$$(u(x, t), w(x, t)) = (\phi(x + ct), \psi(x + ct)),$$

where c is the wave speed and $(\phi(\cdot), \psi(\cdot))$ is the wave profile. Additionally, if $(\phi(\cdot), \psi(\cdot))$ is a periodic function with minimal period $A > 0$, that is,

$$(\phi(\xi + A), \psi(\xi + A)) = (\phi(\xi), \psi(\xi)), \quad \xi \in \mathbb{R},$$

then $(u(x, t), w(x, t)) = (\phi(x + ct), \psi(x + ct))$ is called a *periodic traveling wave* of (1.1). In case $c = 0$ the traveling wave is called a *standing wave*. Throughout this paper we focus on small-amplitude traveling waves with nonzero wave speed in (1.1). In the moving coordinate frame $(\xi, t) = (x + ct, t)$, a wave profile $(\phi(\cdot), \psi(\cdot))$ is a steady state of the following system

$$(1.2) \quad \begin{aligned} u_t &= -cu_\xi + u_{\xi\xi} + f(u, w, \alpha), \\ w_t &= -cw_\xi + g(u, w, \alpha). \end{aligned}$$

Set $v = u_\xi$. Then (ϕ, ϕ_ξ, ψ) is a solution of the three-dimensional (3D) ordinary differential system

$$(1.3) \quad \begin{aligned} \frac{du}{d\xi} &= \dot{u} = v, \\ \frac{dv}{d\xi} &= \dot{v} = cv - f(u, w, \alpha), \\ \frac{dw}{d\xi} &= \dot{w} = \frac{1}{c}g(u, w, \alpha). \end{aligned}$$

Concerning the existence of periodic traveling waves in the coupled system (1.1), it can be proved by detecting periodic orbits in the 3D system (1.3). However, the classical Poincaré-Bendixson Theorem for determining periodic orbits in planar systems is not applicable for the 3D system (1.3). This causes a major obstacle to the existence of periodic waves. Some efforts have been made to overcome it. For example, periodic solutions in (1.3) can be constructed by the geometric singular perturbation theory [8, 10, 45], homoclinic bifurcation [37, 42] and the topological methods [25, 39]. These works obtained large-amplitude periodic solutions in the 3D system (1.3). It is worth mentioning that small-amplitude periodic solutions can be obtained by perturbing the fold-Hopf equilibria, at which the Jacobian matrices of the 3D system (1.3) have one zero eigenvalue and a pair of purely imaginary eigenvalues (see [22, 36]). As we know, the 3D system (1.3) near fold-Hopf equilibria could exhibit complex dynamical behaviors by various perturbations. In order to detect periodic solutions bifurcating from fold-Hopf equilibria, we adopt the averaging theory. It is one of efficient methods to solve periodic solutions in nonautonomous differential systems [22, 41]. The recent works such as [6, 7, 38] successfully applied it to prove the existence of periodic orbits for 3D ordinary differential systems. See Section 2.2 for more information on the averaging theory.

The spectral stability of a traveling wave is determined by the spectrum of the linearization about it. Regarding pulses and fronts in the coupled system (1.1), the essential spectra for their linearizations are obtained by analyzing the corresponding asymptotic operators (see [27]), and the point spectra are determined by the zeros of the Evans function [1, 18, 30, 32, 43]. In contrast to pulses or fronts, the linearizations about periodic traveling waves are linear operators with periodic coefficients, whose spectra have no point spectra and consist of spectral bands. In order to analyze the spectra of periodic traveling waves in reaction-diffusion systems, [21] generalized the construction of the Evans function to periodic waves. However, it is still not easy to determine the spectra for periodic waves in multi-component systems. Some works have been contributed to the stability of periodic waves bifurcating from homoclinic orbits, slow-fast periodic waves in systems of the form (1.1) and so on. See, for instance, [16, 19, 39, 42] and the references therein.

It is also interesting to investigate the spectral stability of small-amplitude periodic waves from fold-Hopf bifurcations in the coupled system (1.1). Along this direction, [47, p.1185]

once mentioned that the small-amplitude periodic traveling waves arising from fold-Hopf bifurcations in a simplified model of calcium dynamics are spectrally unstable, yet without a rigorous proof. The recent works [35, 46] gave detailed studies of the spectral stability of small-amplitude periodic waves in scalar Hamiltonian partial differential equations as a perturbation of the zero-amplitude case. Motivated by these works, we prove the spectral instability of small-amplitude periodic waves from fold-Hopf bifurcations in the coupled system (1.1) via the perturbation theory of linear operators. Indeed, by the fold-Hopf conditions (see Lemma 2.1), we obtain that the corresponding background homogeneous rest state is spectrally unstable (see Lemma 3.7). Then this instability leads to the instability in the associated periodic waves via the perturbation of the linearized operator (see Theorem 3.8). In order to prove this result, we first transform the spectral problem for a periodic wave into the Sturm-Liouville problem with periodic boundary condition. This makes the spectrum discrete. Then we decompose it into a linear operator with constant coefficients and a perturbation. The key step to prove the instability is to check that the perturbation is relatively bounded with respect to the linear operator with constant coefficients, which can be viewed as the linearization about the fold-Hopf equilibrium (the zero-amplitude case). After that, we give the proof for the spectral instability of small-amplitude periodic waves by the spectra for the zero-amplitude case and the perturbation theory.

This paper is organized as follows. We first prove the existence of small-amplitude periodic waves emerging from fold-Hopf equilibria via the averaging theory in Section 2. Based on the results on relatively bounded perturbation, we prove the spectral instability of small-amplitude periodic waves in Section 3. As an application, we study the existence and stability of small-amplitude periodic waves for the FitzHugh-Nagumo system with an applied current in Section 4. We end with some concluding remarks in the final section.

2. EXISTENCE OF PERIODIC WAVES VIA AVERAGING THEORY

2.1. Fold-Hopf equilibrium. In this section, we recall that an equilibrium of the 3D system (1.3) is said to be a *fold-Hopf equilibrium* if the Jacobian matrix for the 3D system (1.3) at this point has one zero eigenvalue $\lambda_1 = 0$ and a pair of purely imaginary eigenvalues $\lambda_{2,3} = \pm \mathbf{i} \mu_0$ with $\mu_0 > 0$ (see [36, p.330]), where $\mathbf{i} = \sqrt{-1}$. By perturbing fold-Hopf equilibria, the 3D system (1.3) could undergo fold-Hopf bifurcations. We refer to [22, 36] for more information on the fold-Hopf bifurcation and fold-Hopf equilibrium.

In what follows, we always use T to denote the transpose of a vector in the usual Euclidean inner product. As a preparation, we first provide some conditions under which the 3D system (1.3) has a fold-Hopf equilibrium. More precisely, we have the following results.

Lemma 2.1. *Assume that $P_0 := (u_0, v_0, w_0)^T$ in \mathbb{R}^3 is a fold-Hopf equilibrium of (1.3) with $(\alpha, c) = (\alpha_0, c_0)$ and $c_0 \neq 0$. Let $\lambda_1 = 0$ and $\lambda_{2,3} = \pm \mathbf{i} \mu_0$ for some $\mu_0 > 0$ be the eigenvalues*

of the Jacobian matrix $J(u_0, v_0, w_0)$ at P_0 . Then the following conditions hold:

$$(2.1) \quad v_0 = 0, \quad f(u_0, w_0, \alpha_0) = 0, \quad g(u_0, w_0, \alpha_0) = 0.$$

$$(2.2) \quad c_0 + \frac{1}{c_0}g_w(P_0) = f_u(P_0)g_w(P_0) - f_w(P_0)g_u(P_0) = 0, \quad f_u(P_0) + g_w(P_0) = \mu_0^2 > 0.$$

Proof. Since $P_0 = (u_0, v_0, w_0)^T$ is a fold-Hopf equilibrium of (1.3) with $(\alpha, c) = (\alpha_0, c_0)$, the condition (2.1) holds. It is clear that the Jacobian matrix $J(u_0, v_0, w_0)$ of (1.3) at this point is in the form

$$J(u_0, v_0, w_0) = \begin{pmatrix} 0 & 1 & 0 \\ -f_u(P_0) & c_0 & -f_w(P_0) \\ \frac{1}{c_0}g_u(P_0) & 0 & \frac{1}{c_0}g_w(P_0) \end{pmatrix},$$

and the eigenvalues of $J(u_0, v_0, w_0)$ are determined by the zeros of its characteristic polynomial

$$\begin{aligned} & \det(\lambda I_3 - J(u_0, v_0, w_0)) \\ &= \lambda^3 - \left(c_0 + \frac{1}{c_0}g_w(P_0) \right) \lambda^2 + (f_u(P_0) + g_w(P_0)) \lambda + \frac{1}{c_0} (f_w(P_0)g_u(P_0) - f_u(P_0)g_w(P_0)), \end{aligned}$$

where I_3 is the 3×3 identity matrix. By the definition of the fold-Hopf equilibrium, we have

$$\begin{aligned} c_0 + \frac{1}{c_0}g_w(P_0) &= \lambda_1 + \lambda_2 + \lambda_3 = 0, \\ f_u(P_0) + g_w(P_0) &= \lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3 = \mu_0^2 > 0, \\ \frac{1}{c_0} (f_w(P_0)g_u(P_0) - f_u(P_0)g_w(P_0)) &= \lambda_1\lambda_2\lambda_3 = 0. \end{aligned}$$

Thus, (2.2) holds. This finishes the proof. \square

2.2. Averaging theory. In this section, we take advantage of the averaging theory to find periodic solutions bifurcating from fold-Hopf equilibria in the 3D system (1.3). The averaging theory is one of the efficient methods to find periodic solutions in nonautonomous differential systems (see more details in [6, 22, 38, 41]). In the following we introduce some results on the averaging theory. Consider a nonautonomous differential system in the form

$$(2.3) \quad \dot{X}(t) = \epsilon F(t, X) + \epsilon^2 K(t, X, \epsilon),$$

where $X = (x, y)^T$ is in \mathbb{R}^2 , small parameter ϵ satisfies $0 \leq \epsilon < |\epsilon_1|$ for a sufficiently small $\epsilon_1 > 0$, and the functions F and K are sufficiently smooth and are A -periodic in the variable $t \in \mathbb{R}$. Then (2.3) can be defined in the extended phase space $\mathbb{S}^1 \times \mathbb{R}^2$ by setting $\dot{t} = 1 \pmod{A}$. Let $X(t, Z, \epsilon)$ denote the solution of (2.3) with the initial value $X(0) = Z$ in \mathbb{R}^2 . Then by [6, Lemma 5, p. 3570], the solution $X(t, Z, \epsilon)$ has the expansion

$$X(t, Z, \epsilon) = Z + \epsilon Y(t, Z) + O(\epsilon^2),$$

where Y is given by

$$Y(t, Z) = \int_0^t F(\tau, Z) d\tau.$$

Consequently, the Poincaré map $\Pi(Z, \epsilon) = X(A, Z, \epsilon)$ is in the form

$$\Pi(Z, \epsilon) = Z + \epsilon G(Z) + O(\epsilon^2),$$

where $G(Z)$ is given by

$$G(Z) = Y(A, Z) = \int_0^A F(\tau, Z) d\tau,$$

and is called the *averaged function of order one*. Hence, in order to detect the periodic solutions in (2.3), we turn to solve the equation $\Pi(Z, \epsilon) = Z$. By [38, Theorem A, P. 566] we have the next lemma.

Lemma 2.2. *Assume that $G \neq 0$, and there exists a point Z_0 in \mathbb{R}^2 such that $G(Z_0) = 0$ and the differential $DG(Z_0)$ of $G(Z)$ is an invertible matrix. Then for a sufficiently small $|\epsilon| > 0$, the perturbed system (2.3) has an isolated A -periodic solution $\varphi(t, \epsilon)$ satisfying that $\varphi(0, \epsilon) \rightarrow Z_0$ as $\epsilon \rightarrow 0$.*

In the preceding lemma, we apply the averaged function of order one to find the periodic solutions in (2.3). We also refer to [6, 7, 38] for more recent results on the applications of the averaged function of high order.

2.3. Existence of periodic waves. Now we study the existence of small-amplitude periodic traveling waves emerging from fold-Hopf equilibria via the averaging theory.

Without loss of generality, we assume that for $(\alpha, c) = (\alpha_0, c_0)$, the 3D system (1.3) has a fold-Hopf equilibrium at the origin, which is denoted by 0 for short. Let $X(u, w, v, \epsilon)$ denote the vector field defined by (1.3) with (α, c) in the form

$$(2.4) \quad \alpha = \alpha_0 + \alpha_1(\epsilon) := \alpha_0 + \sum_{k=1}^{+\infty} \tilde{\alpha}_k \epsilon^k, \quad c = c_0 + c_1(\epsilon) := c_0 + \sum_{k=1}^{+\infty} \tilde{c}_k \epsilon^k,$$

where all coefficients will be fixed according to our need such that the radii of convergence of these two series are nonzero. It is clear that by a change

$$(2.5) \quad (u, v, w)^T \rightarrow \epsilon Q(u, v, w)^T, \quad Q = \begin{pmatrix} -\frac{1}{c_0} g_w(0, \alpha_0) & \mu_0 & g_w(0, \alpha_0) \\ -\mu_0^2 & c_0 \mu_0 & 0 \\ \frac{1}{c_0} g_u(0, \alpha_0) & 0 & -g_u(0, \alpha_0) \end{pmatrix},$$

where the matrix Q consists of the eigenvectors corresponding to the Jacobian matrix of (1.3) at the origin, the 3D system (1.3) is transformed into

$$(2.6) \quad \begin{aligned} \dot{u} &= \mu_0 v + X_1(u, v, w, \epsilon), \\ \dot{v} &= -\mu_0 u + X_2(u, v, w, \epsilon), \\ \dot{w} &= X_3(u, v, w, \epsilon), \end{aligned}$$

where

$$\begin{aligned} &(X_1(u, v, w, \epsilon), X_2(u, v, w, \epsilon), X_3(u, v, w, \epsilon))^T \\ &= \epsilon^{-1} Q^{-1} X(\epsilon(u, v, w) Q^T, \epsilon) - (\mu_0 v, -\mu_0 u, 0)^T. \end{aligned}$$

For sufficiently small $|\epsilon|$, the expansion of $(X_1(u, v, w, \epsilon), X_2(u, v, w, \epsilon), X_3(u, v, w, \epsilon))^T$ with respect to ϵ can be written as

$$\begin{aligned} &(X_1(u, v, w, \epsilon), X_2(u, v, w, \epsilon), X_3(u, v, w, \epsilon))^T \\ &= (X_{10} + \epsilon X_{11}(u, v, w), X_{20} + \epsilon X_{21}(u, v, w), X_{30} + \epsilon X_{31}(u, v, w))^T + O(\epsilon^2), \end{aligned}$$

where $X_{j1}(u, v, w) = O(|(u, v, w)|)$ for sufficiently small $|(u, v, w)|$. Note that X_{j0} and X_{j1} depend on the matrix Q , which is not unique. Then for simplicity, we omit their lengthy expressions, which are determined by the first and second partial derivatives of the functions f and g with respect to u, v, w and α . As a concrete example, we give the explicit expressions of X_{j0} and X_{j1} for the FitzHugh-Nagumo system in Section 4.3. Consider system (2.6) in the cylindrical coordinate

$$(u, v, w) = (r \cos \theta, r \sin \theta, w), \quad r \geq 0, \quad 2\pi \geq \theta \geq 0, \quad w \in \mathbb{R}.$$

Then system (2.6) is converted into

$$(2.7) \quad \begin{aligned} \dot{r} &= X_1 \cos \theta + X_2 \sin \theta, \\ r \dot{\theta} &= -\mu_0 r - X_1 \sin \theta + X_2 \cos \theta, \\ \dot{w} &= X_3, \end{aligned}$$

where we write X_j with $(r \cos \theta, r \sin \theta, w, \epsilon)$ being suppressed for simplicity. Then for sufficiently small $|r|$ and $|\epsilon|$,

$$(2.8) \quad \begin{aligned} \frac{dr}{d\theta} &= \frac{r(X_1 \cos \theta + X_2 \sin \theta)}{-\mu_0 r - X_1 \sin \theta + X_2 \cos \theta}, \\ \frac{dw}{d\theta} &= \frac{r X_3}{-\mu_0 r - X_1 \sin \theta + X_2 \cos \theta}. \end{aligned}$$

Clearly, the right-hand side of system (2.8) is 2π -periodic in the variable $\theta \in \mathbb{R}$. Then we have the following results.

Theorem 2.3. (Existence of periodic traveling waves) *Define two functions R_1 and R_2 by*

$$R_1(r, w) = \int_0^{2\pi} (\cos \theta \cdot X_{11}(r \cos \theta, r \sin \theta, w) + \sin \theta \cdot X_{21}(r \cos \theta, r \sin \theta, w)) d\theta,$$

$$R_2(r, w) = \int_0^{2\pi} X_{31}(r \cos \theta, r \sin \theta, w) d\theta.$$

Assume that there exist $\alpha_1(\epsilon)$ and $c_1(\epsilon)$ defined by (2.4) such that $X_{10} = X_{20} = X_{30} = 0$, $R_1(r_, w_*) = R_2(r_*, w_*) = 0$, and*

$$(2.9) \quad \det \begin{pmatrix} \frac{\partial R_1}{\partial r} & \frac{\partial R_1}{\partial w} \\ \frac{\partial R_2}{\partial r} & \frac{\partial R_2}{\partial w} \end{pmatrix} \Big|_{(r_*, w_*)} \neq 0,$$

for some $r_ > 0$ and $w_* \in \mathbb{R}$. Then there exists a sufficiently small $\epsilon_0 > 0$ such that a small-amplitude periodic wave (ϕ, ψ) with period A_ϵ and wave speed $c_0 + c_1(\epsilon)$ emerges from the origin in the coupled system (1.1) with $\alpha = \alpha_0 + \alpha_1(\epsilon)$ for each $\epsilon \in (0, \epsilon_0]$, where*

$$|(\phi, \psi)| = O(\epsilon), \quad A_\epsilon = 2\pi/\mu_0 + O(\epsilon), \quad \epsilon \in (0, \epsilon_0].$$

Proof. Note that the periodic traveling waves of the coupled system (1.1) correspond to the periodic solutions of the 3D system (1.3). Then it suffices to prove that there exists a sufficiently small $\epsilon_0 > 0$ such that a small-amplitude periodic solution (ϕ, ϕ_ξ, ψ) with period A_ϵ bifurcates from the origin in the 3D system (1.3) with $(\alpha, c) = (\alpha_0 + \alpha_1(\epsilon), c_0 + c_1(\epsilon))$ for each $\epsilon \in (0, \epsilon_0]$, where $|(\phi, \phi_\xi, \psi)| = O(\epsilon)$ and $A_\epsilon = 2\pi/\mu_0 + O(\epsilon)$ for $\epsilon \in (0, \epsilon_0]$.

Since $X_{10} = X_{20} = X_{30} = 0$, system (2.8) has the expansion

$$(2.10) \quad \begin{aligned} \frac{dr}{d\theta} &= -\frac{\epsilon}{\mu_0} (\cos \theta X_{11}(r \cos \theta, r \sin \theta, w) + \sin \theta X_{21}(r \cos \theta, r \sin \theta, w)) + O(\epsilon^2), \\ \frac{dw}{d\theta} &= -\frac{\epsilon}{\mu_0} X_{31}(r \cos \theta, r \sin \theta, w) + O(\epsilon^2). \end{aligned}$$

Consequently, by $R_1(r_*, w_*) = R_2(r_*, w_*) = 0$, (2.9) and Lemma 2.2, we have that for sufficiently small ϵ , system (2.10) has an isolated periodic solution $\varphi(\theta, \epsilon)$ of period 2π in θ such that $\varphi(0, \epsilon) \rightarrow (r_*, w_*)$ as $\epsilon \rightarrow 0$. This together with (2.5) yields that the 3D system (1.3) has a periodic solution (ϕ, ϕ_ξ, ψ) with $|(\phi, \phi_\xi, \psi)| = O(\epsilon)$ for sufficiently small $|\epsilon|$. By the second equation in (2.7), we have that for sufficiently small $|\epsilon|$,

$$A_\epsilon = \left| \int_0^{2\pi} \frac{r}{-\mu_0 r - X_1 \sin \theta + X_2 \cos \theta} \Big|_{(r,w)=\varphi(\theta,\epsilon)} d\theta \right| = \frac{2\pi}{\mu_0} + O(\epsilon).$$

Therefore, the proof is now complete. \square

3. INSTABILITY OF PERIODIC WAVES VIA PERTURBATION THEORY

As for small-amplitude periodic waves arising from the fold-Hopf bifurcations, we are interested in the stability of this class of periodic waves in the coupled system (1.1).

3.1. Spectral theory for periodic waves. We first introduce basic notions on the spectral theory for traveling waves in the coupled system (1.1). Assume that the coupled system (1.1) has a traveling wave (ϕ, ψ) with wave speed c . The linearization of (1.2) about the traveling wave (ϕ, ψ) is in the form

$$(3.1) \quad \mathcal{L} \begin{pmatrix} u \\ w \end{pmatrix} := \begin{pmatrix} u_{\xi\xi} - cu_{\xi} + f_u(\phi, \psi, \alpha)u + f_w(\phi, \psi, \alpha)w \\ -cw_{\xi} + g_u(\phi, \psi, \alpha)u + g_w(\phi, \psi, \alpha)w \end{pmatrix},$$

where

$$\mathcal{L} : H^2(\mathbb{R}) \times H^1(\mathbb{R}) \subset L^2(\mathbb{R}) \times L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}) \times L^2(\mathbb{R}).$$

The operator \mathcal{L} is a closed operator on $L^2(\mathbb{R}) \times L^2(\mathbb{R})$ with the domain $\mathcal{D}(\mathcal{L}) = H^2(\mathbb{R}) \times H^1(\mathbb{R})$ and its eigenvalue problem is determined by

$$\mathcal{L}(u, w)^T = \lambda(u, w)^T,$$

where $(u, w)^T$ is in $\mathcal{D}(\mathcal{L})$, and λ is in the complex plane \mathbb{C} . Let $\rho(\mathcal{L})$ denote the *resolvent set* of \mathcal{L} , consisting of all points $\lambda \in \mathbb{C}$ such that $\lambda I - \mathcal{L}$ is invertible and $(\lambda I - \mathcal{L})^{-1}$ is a continuous linear operator, where I is the identity. The complement of $\rho(\mathcal{L})$ in \mathbb{C} is called the *spectrum* of \mathcal{L} , which is denoted by $\sigma(\mathcal{L})$. Then

$$\mathbb{C} = \rho(\mathcal{L}) \cup \sigma(\mathcal{L}).$$

If the dimension of the kernel $\text{Ker}(\mathcal{L})$ for \mathcal{L} and the codimension of the the range $\mathcal{R}(\mathcal{L})$ for \mathcal{L} are both finite, then the operator \mathcal{L} is called a *Fredholm operator* with the *Fredholm index*

$$\text{Ind}(\mathcal{L}) = \dim(\text{Ker}(\mathcal{L})) - \text{codim}(\mathcal{R}(\mathcal{L})).$$

According to the definitions of the Fredholm operator and the Fredholm index, the spectrum $\sigma(\mathcal{L})$ is classified into two different parts, that is,

$$\sigma(\mathcal{L}) = \sigma_{\text{ess}}(\mathcal{L}) \cup \sigma_{\text{pt}}(\mathcal{L}),$$

where $\sigma_{\text{ess}}(\mathcal{L})$ (called the *essential spectrum*) consists of all $\lambda \in \mathbb{C}$ such that $\lambda I - \mathcal{L}$ is not Fredholm or $\lambda I - \mathcal{L}$ is Fredholm with $\text{Ind}(\lambda I - \mathcal{L}) \neq 0$, and $\sigma_{\text{pt}}(\mathcal{L})$ (called the *point spectrum*) consists of all $\lambda \in \mathbb{C}$ such that $\lambda I - \mathcal{L}$ is not invertible and $\text{Ind}(\lambda I - \mathcal{L})$ is zero. Every $\lambda \in \sigma_{\text{pt}}(\mathcal{L})$ is called an *eigenvalue* of \mathcal{L} and every nontrivial $(u, w)^T$ in $\mathcal{D}(\mathcal{L})$ satisfying $\mathcal{L}(u, w)^T = \lambda(u, w)^T$ is called an *eigenfunction* corresponding to λ .

We next introduce the spectral projections associated with $\sigma(\mathcal{L})$, which are useful in the study of the spectral perturbation problem (see Section 3.2). Assume that $\sigma(\mathcal{L})$ is separated into two disjoint parts σ_1 and σ_2 by a simple closed positively oriented cycle Γ , which encloses a bounded open set containing σ_1 in its interior and σ_2 in its exterior. Then by [33, Theorem

6.17, p.178], we can define the spectral projection $P_{\sigma_1}(\mathcal{L})$ associated with σ_1 by the Dunford integral formula

$$P_{\sigma_1}(\mathcal{L}) = \frac{1}{2\pi\mathbf{i}} \oint_{\Gamma} (\lambda I - \mathcal{L}) d\lambda.$$

In particular, if σ_1 only contains an isolated eigenvalue λ_0 of \mathcal{L} , then the dimension of the range of $P_{\lambda_0}(\mathcal{L})$ is called the *algebraic multiplicity* of λ_0 , and the dimension of $\text{Ker}(\lambda_0 I - \mathcal{L}_0)$ is called the *geometric multiplicity* of λ_0 . We refer to [32, 33] for more details on the spectral theory of \mathcal{L} . For notational convenience, these notations are similarly defined for the linear operators which appear in the present paper.

Our goal is to study the stability of periodic traveling waves emerging from the fold-Hopf points. For this reason, we give more details on the spectral theory for this class of periodic waves. Assume that (ϕ, ϕ_ξ, ψ) is a periodic solution with period A_ϵ arising from the fold-Hopf bifurcation in the 3D system (1.3) with $(\alpha, c) = (\alpha_0 + \alpha_1(\epsilon), c_0 + c_1(\epsilon))$ for $\epsilon \in (0, \epsilon_0]$, where α_1 and c_1 are defined as in (2.4). Then (ϕ, ψ) is periodic wave profile with period A_ϵ in system (1.1). So $(\phi(\cdot), \psi(\cdot))$ satisfies (1.2). The spectral stability of the profile $(\phi(\xi), \psi(\xi))$ is determined by the spectra of the operator \mathcal{L} in the form (3.1), where all coefficients are periodic functions with period A_ϵ . Let $Y = (u, v, w)^T$ in \mathbb{R}^3 . Then the spectral problem $\mathcal{L}(u, w)^T = \lambda(u, w)^T$ becomes a first-order system of ordinary differential equations

$$(3.2) \quad \frac{dY}{d\xi} = A(\xi, \lambda)Y, \quad A(\xi, \lambda) = \begin{pmatrix} 0 & 1 & 0 \\ \lambda - f_u(\phi, \psi, \alpha) & c & -f_w(\phi, \psi, \alpha) \\ \frac{1}{c}g_u(\phi, \psi, \alpha) & 0 & \frac{1}{c}(g_w(\phi, \psi, \alpha) - \lambda) \end{pmatrix},$$

where $A(\xi + A_\epsilon) = A(\xi, \lambda)$ for every $\lambda \in \mathbb{C}$. We can check that the operator \mathcal{L} has no point spectrum, and $\lambda = \lambda_0$ is an essential spectrum of \mathcal{L} if and only if there exists a μ in $(-\pi/A_\epsilon, \pi/A_\epsilon]$ such that (3.2) has a nonzero solution Y satisfying

$$Y(A_\epsilon) = e^{\mathbf{i}\mu A_\epsilon} Y(0).$$

By the Bloch-wave decomposition

$$(u, w) \rightarrow (e^{\mathbf{i}\mu\xi}u, e^{\mathbf{i}\mu\xi}w),$$

the operator \mathcal{L} is converted into

$$(3.3) \quad \mathcal{L}_\mu \begin{pmatrix} u \\ w \end{pmatrix} := \begin{pmatrix} (\partial_\xi + \mathbf{i}\mu)^2 u - c(\partial_\xi + \mathbf{i}\mu)u + f_u(\phi, \psi, \alpha)u + f_w(\phi, \psi, \alpha)w \\ -c(\partial_\xi + \mathbf{i}\mu)w + g_u(\phi, \psi, \alpha)u + g_w(\phi, \psi, \alpha)w \end{pmatrix},$$

where

$$\mathcal{L}_\mu : L_{\text{per}}^2([0, A_\epsilon], \mathbb{C}) \times L_{\text{per}}^2([0, A_\epsilon], \mathbb{C}) \rightarrow L_{\text{per}}^2([0, A_\epsilon], \mathbb{C}) \times L_{\text{per}}^2([0, A_\epsilon], \mathbb{C})$$

with the domain $\mathcal{D}(\mathcal{L}_\mu) = H_{\text{per}}^2([0, A_\epsilon], \mathbb{C}) \times H_{\text{per}}^1([0, A_\epsilon], \mathbb{C})$. Then for each μ in $(-\pi/A_\epsilon, \pi/A_\epsilon]$, the point spectrum of \mathcal{L}_μ is an essential spectrum of \mathcal{L} . We refer to [32, pp. 68-70] for more

details on the relation between the spectra of \mathcal{L} and those of \mathcal{L}_μ for μ in $(-\pi/A_\epsilon, \pi/A_\epsilon]$. We summarize the above statements as follows:

Lemma 3.1. *The spectrum $\sigma(\mathcal{L})$ of the operator \mathcal{L} for periodic traveling wave (ϕ, ψ) in (3.1) is in the form*

$$(3.4) \quad \sigma(\mathcal{L}) = \bigcup_{-\frac{\pi}{A_\epsilon} < \mu \leq \frac{\pi}{A_\epsilon}} \sigma_{\text{pt}}(\mathcal{L}_\mu).$$

In order to analyze the spectrum of \mathcal{L}_μ , we also change the spectral problem for \mathcal{L}_μ into a first-order system of ordinary differential equations

$$(3.5) \quad \frac{dY}{d\xi} = (A(\xi, \lambda) - \mathbf{i}\mu I_3)Y, \quad Y(A_\epsilon) = Y(0),$$

where $\mathbf{i} = \sqrt{-1}$ and I_3 is the 3×3 identity matrix. Let $\Phi(\xi, \lambda, \mu)$ denote the fundamental matrix solution of (3.5). Then (3.5) has a nontrivial solution if and only if

$$(3.6) \quad E(\lambda, \mu) := \det(\Phi(A_\epsilon, \lambda, \mu) - I_3) = 0, \quad \lambda \in \mathbb{C}, \quad -\frac{\pi}{A_\epsilon} < \mu \leq \frac{\pi}{A_\epsilon},$$

where $E(\lambda, \mu)$ is called the *Evans function* (see, for instance, [1, 2, 18, 21, 30, 32, 43]). Then the spectrum of \mathcal{L}_μ is determined by the zeros of $E(\lambda, \mu)$.

3.2. Relatively bounded perturbation. By (3.4) we observe that $\sigma(\mathcal{L})$ is spectrally unstable if \mathcal{L}_μ is spectrally unstable for a certain μ in $(-\pi/A_\epsilon, \pi/A_\epsilon]$. Particularly, we consider the case $\mu = 0$ in (3.3). Then \mathcal{L}_μ is reduced to $\mathcal{L}_0 = \mathcal{L}$ with the domain $\mathcal{D}(\mathcal{L}_0) = H_{\text{per}}^2([0, A_\epsilon], \mathbb{C}) \times H_{\text{per}}^1([0, A_\epsilon], \mathbb{C})$. In order to remove the ϵ -dependence associated with the domain $\mathcal{D}(\mathcal{L}_0)$, we make a change

$$\xi \rightarrow A_\epsilon \xi, \quad \epsilon \in (0, \epsilon_0].$$

Then the spectral problem associated with the periodic wave (ϕ, ψ) is converted into the spectral problem for the following operator

$$(3.7) \quad \mathcal{F}(\epsilon) \begin{pmatrix} u \\ w \end{pmatrix} := \begin{pmatrix} u_{\xi\xi} - cA_\epsilon u_\xi + A_\epsilon^2 f_u(\phi, \psi, \alpha)u + A_\epsilon^2 f_w(\phi, \psi, \alpha)w \\ -cA_\epsilon w_\xi + A_\epsilon^2 g_u(\phi, \psi, \alpha)u + A_\epsilon^2 g_w(\phi, \psi, \alpha)w \end{pmatrix} = \lambda \begin{pmatrix} u \\ w \end{pmatrix},$$

where

$$\mathcal{F}(\epsilon) : L_{\text{per}}^2([0, 1], \mathbb{C}) \times L_{\text{per}}^2([0, 1], \mathbb{C}) \rightarrow L_{\text{per}}^2([0, 1], \mathbb{C}) \times L_{\text{per}}^2([0, 1], \mathbb{C})$$

with the domain $\mathcal{D}(\mathcal{F}(\epsilon)) = H_{\text{per}}^2([0, 1], \mathbb{C}) \times H_{\text{per}}^1([0, 1], \mathbb{C})$ for each $\epsilon \in (0, \epsilon_0]$. For each $(u, w)^T$ in $L_{\text{per}}^2([0, 1], \mathbb{C}) \times L_{\text{per}}^2([0, 1], \mathbb{C})$, let the norm of $(u, w)^T$ be defined by

$$|(u, w)^T|_2 = |u|_2 + |w|_2,$$

where $|u|_2$ and $|w|_2$ are the L^2 -norms of u and w in $L^2([0, 1], \mathbb{C})$, respectively.

On the spectra $\sigma(\mathcal{L}_0)$ and $\sigma(\mathcal{F}(\epsilon))$, we have the next lemma.

Lemma 3.2. *The spectra $\sigma(\mathcal{L}_0)$ and $\sigma(\mathcal{F}(\epsilon))$ for each $\epsilon \in (0, \epsilon_0]$ are both composed by a countable number of eigenvalues, which have finite algebraic multiplicities. Furthermore, if λ is an eigenvalue of $\mathcal{F}(\epsilon)$, then λ/A_ϵ^2 is an eigenvalue of \mathcal{L}_0 .*

Proof. The first statement can be obtained by [32, Lemma 8.4.1, p. 242] and the second one can be obtained by a direct computation. This finishes the proof. \square

In order to analyze the spectrum $\sigma(\mathcal{F}(\epsilon))$ by the perturbation theory for linear operators, we decompose $\mathcal{F}(\epsilon)$ into two parts:

$$\mathcal{F}(\epsilon) = \mathcal{F}_0 + \mathcal{F}_1(\epsilon),$$

where

$$(3.8) \quad \mathcal{F}_0 \begin{pmatrix} u \\ w \end{pmatrix} := \begin{pmatrix} u_{\xi\xi} - c_0 A u_\xi + A^2 f_u(P_0, \alpha_0)u + A^2 f_w(P_0, \alpha_0)w \\ -c_0 A w_\xi + A^2 g_u(P_0, \alpha_0)u + A^2 g_w(P_0, \alpha_0)w \end{pmatrix},$$

and $\mathcal{F}_1(\epsilon) = \mathcal{F}(\epsilon) - \mathcal{F}_0$, that is,

$$(3.9) \quad \mathcal{F}_1(\epsilon) \begin{pmatrix} u \\ w \end{pmatrix} := \begin{pmatrix} (c_0 A - c A_\epsilon)u_\xi + \beta_u u + \beta_w w \\ (c_0 A - c A_\epsilon)w_\xi + \gamma_u u + \gamma_w w \end{pmatrix}.$$

Recall that for each $\epsilon \in (0, \epsilon_0]$,

$$\alpha_1(\epsilon) = O(\epsilon), \quad c_1(\epsilon) = O(\epsilon), \quad |(\phi, \phi_\xi, \psi)| = O(\epsilon), \quad |A_\epsilon - A| = O(\epsilon),$$

then we have that $c_0 A - c A_\epsilon = O(\epsilon)$,

$$(3.10) \quad \begin{aligned} \beta_u(\xi, \epsilon) &= A_\epsilon^2 f_u(\phi, \psi, \alpha) - A^2 f_u(P_0, \alpha_0) = O(\epsilon), \\ \beta_w(\xi, \epsilon) &= A_\epsilon^2 f_w(\phi, \psi, \alpha) - A^2 f_w(P_0, \alpha_0) = O(\epsilon), \\ \gamma_u(\xi, \epsilon) &= A_\epsilon^2 g_u(\phi, \psi, \alpha) - A^2 g_u(P_0, \alpha_0) = O(\epsilon), \\ \gamma_w(\xi, \epsilon) &= A_\epsilon^2 g_w(\phi, \psi, \alpha) - A^2 g_w(P_0, \alpha_0) = O(\epsilon), \end{aligned}$$

for $\epsilon \in (0, \epsilon_0]$. As a consequence, the operators $\mathcal{F}(\epsilon)$ for $\epsilon \in (0, \epsilon]$ can be viewed as the perturbations of \mathcal{F}_0 . Next we give a further study on the relation between \mathcal{F}_0 and $\mathcal{F}_1(\epsilon)$.

Lemma 3.3. *Let the operators \mathcal{F}_0 and $\mathcal{F}_1(\epsilon)$ for $\epsilon \in (0, \epsilon_0]$ be defined by (3.8) and (3.9), respectively. Then for each $\epsilon \in (0, \epsilon_0]$ and each $\lambda \in \mathbb{C}$, the operator $\mathcal{F}_1(\epsilon)$ is relatively bounded with respect to $\mathcal{F}_0 - \lambda I$, that is, $\mathcal{D}(\mathcal{F}_0 - \lambda I) \subset \mathcal{D}(\mathcal{F}_1(\epsilon))$ and there exists two nonnegative constants $a(\epsilon)$ and $b(\epsilon)$ such that*

$$(3.11) \quad |\mathcal{F}_1(\epsilon)(u, w)^T|_2 \leq a(\epsilon)|(u, w)^T|_2 + b(\epsilon)|(\mathcal{F}_0 - \lambda I)(u, w)^T|_2,$$

for each $(u, w)^T$ in $\mathcal{D}(\mathcal{F}_0)$, where the constants $a(\epsilon)$ and $b(\epsilon)$ can be in the form

$$\begin{aligned} a(\epsilon) &= \kappa_1(\epsilon) \left\{ 1 + \frac{2\kappa_2(\lambda) + 2n(n+1)}{n-1-|c_0A|} + \frac{2\kappa_2(\lambda)}{|c_0A|} \right\}, \\ b(\epsilon) &= \kappa_1(\epsilon) \left\{ \frac{1}{n-1-|c_0A|} + \frac{1}{|c_0A|} \right\}, \end{aligned}$$

where n is an integer with $n > 1 + |c_0A|$, and $\kappa_1(\epsilon)$ and $\kappa_2(\lambda)$ are in the form

$$\begin{aligned} \kappa_1(\epsilon) &= 3 \cdot \max\{|c_0A - cA_\epsilon|, |\beta_u(\cdot, \epsilon)|_\infty + |\gamma_w(\cdot, \epsilon)|_\infty, |\beta_w(\cdot, \epsilon)|_\infty + |\gamma_u(\cdot, \epsilon)|_\infty\}, \\ \kappa_2(\lambda) &= \max\{A^2(|f_u(P_0, \alpha_0)| + |g_w(P_0, \alpha_0)|), A^2(|f_w(P_0, \alpha_0)| + |g_u(P_0, \alpha_0)|)\} + \frac{|\lambda|}{2}. \end{aligned}$$

Furthermore, the following limits hold:

$$(3.12) \quad a(\epsilon) \rightarrow 0, \quad b(\epsilon) \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0.$$

Proof. Noting that

$$\mathcal{F}_0 : L^2_{\text{per}}([0, 1], \mathbb{C}) \times L^2_{\text{per}}([0, 1], \mathbb{C}) \rightarrow L^2_{\text{per}}([0, 1], \mathbb{C}) \times L^2_{\text{per}}([0, 1], \mathbb{C})$$

has the domain $\mathcal{D}(\mathcal{F}_0) = H^2_{\text{per}}([0, 1], \mathbb{C}) \times H^1_{\text{per}}([0, 1], \mathbb{C})$, and

$$\mathcal{F}_1(\epsilon) : L^2_{\text{per}}([0, 1], \mathbb{C}) \times L^2_{\text{per}}([0, 1], \mathbb{C}) \rightarrow L^2_{\text{per}}([0, 1], \mathbb{C}) \times L^2_{\text{per}}([0, 1], \mathbb{C})$$

has the domain $\mathcal{D}(\mathcal{F}_1(\epsilon)) = H^1_{\text{per}}([0, 1], \mathbb{C}) \times H^1_{\text{per}}([0, 1], \mathbb{C})$, we then conclude that $\mathcal{D}(\mathcal{F}_0) \subset \mathcal{D}(\mathcal{F}_1(\epsilon))$ for each $\epsilon \in (0, \epsilon_0]$. This finishes the proof for the first statement.

Next we prove (3.11). For each $\epsilon \in (0, \epsilon_0]$ and $(u, w)^T$ in $\mathcal{D}(\mathcal{F}_0)$,

$$(3.13) \quad \begin{aligned} |\mathcal{F}_1(\epsilon)(u, w)^T|_2 &= |((c_0A - cA_\epsilon)u_\xi + \beta_u u + \beta_w w, (c_0A - cA_\epsilon)w_\xi + \gamma_u u + \gamma_w w)^T|_2 \\ &\leq \kappa_1(\epsilon) (|u_\xi|_2 + |w_\xi|_2 + |(u, w)^T|_2), \end{aligned}$$

where κ_1 is defined as in this lemma. By (3.8), we have

$$\begin{aligned} &|(u_{\xi\xi} - c_0Au_\xi, -c_0Aw_\xi)^T|_2 \\ &= |\mathcal{F}_0(u, w)^T - (A^2f_u(P_0, \alpha_0)u + A^2f_w(P_0, \alpha_0)w, A^2g_u(P_0, \alpha_0)u + A^2g_w(P_0, \alpha_0)w)^T|_2. \end{aligned}$$

Then

$$\begin{aligned} |c_0Aw_\xi|_2 &\leq |(\mathcal{F}_0 - \lambda I)(u, w)^T|_2 + 2\kappa_2(\lambda)|(u, w)^T|_2, \\ |u_{\xi\xi} - c_0Au_\xi|_2 &\leq |(\mathcal{F}_0 - \lambda I)(u, w)^T|_2 + 2\kappa_2(\lambda)|(u, w)^T|_2, \end{aligned}$$

where $\kappa_2(\lambda)$ is defined as in this lemma. These yield that

$$(3.14) \quad |w_\xi|_2 \leq |c_0A|^{-1} (|(\mathcal{F}_0 - \lambda I)(u, w)^T|_2 + 2\kappa_2(\lambda)|(u, w)^T|_2),$$

$$(3.15) \quad |u_{\xi\xi}|_2 \leq |(\mathcal{F}_0 - \lambda I)(u, w)^T|_2 + 2\kappa_2(\lambda)|(u, w)^T|_2 + |c_0Au_\xi|_2.$$

By the Inequality (1.12) in [33, p.192], we have that for each $n > 1$,

$$|u_\xi|_2 \leq \frac{1}{n-1} |u_{\xi\xi}|_2 + \frac{2n(n+1)}{n-1} |u|_2.$$

This together with (3.15) yields that for $n > 1$,

$$\begin{aligned} \left(1 - \frac{|c_0 A|}{n-1}\right) |u_\xi|_2 &\leq \frac{1}{n-1} (|(\mathcal{F}_0 - \lambda I)(u, w)^T|_2 + 2\kappa_2(\lambda) |(u, w)^T|_2) \\ &\quad + \frac{2n(n+1)}{n-1} |u|_2. \end{aligned}$$

Hence, for every $n > 1 + |c_0 A|$, we have

$$(3.16) \quad \begin{aligned} |u_\xi|_2 &\leq \frac{1}{n-1-|c_0 A|} (|(\mathcal{F}_0 - \lambda I)(u, w)^T|_2 + 2\kappa_2(\lambda) |(u, w)^T|_2) \\ &\quad + \frac{2n(n+1)}{n-1-|c_0 A|} |u|_2. \end{aligned}$$

Substituting (3.14) and (3.16) into (3.13) yields that (3.11) holds.

Finally, we prove (3.12). By (3.10) we have $\kappa_1(\epsilon) = O(\epsilon)$ for $\epsilon \in (0, \epsilon_0]$. Then by the definitions of $a(\epsilon)$ and $b(\epsilon)$, we obtain (3.12). This finishes the proof. \square

Letting $\lambda = 0$ in Lemma 3.3, we have the next corollary.

Corollary 3.1. *For each $\epsilon \in (0, \epsilon_0]$, the operator $\mathcal{F}_1(\epsilon)$ is relatively bounded with respect to \mathcal{F}_0 , and*

$$|\mathcal{F}_1(\epsilon)(u, w)^T|_2 \leq a(\epsilon) |(u, w)^T|_2 + b(\epsilon) |\mathcal{F}_0(u, w)^T|_2,$$

for each $(u, w)^T$ in $\mathcal{D}(\mathcal{F}_0)$, where the constants $a(\epsilon)$ and $b(\epsilon)$ are defined as in Lemma 3.3.

We remark that in order to study the spectral stability of periodic waves arising from Hopf bifurcation in a class of systems of scalar viscous balance laws in one space dimension, the recent work [3] also obtained the similar result as Corollary 3.1. We next study the spectral perturbation problem for \mathcal{F}_0 . By [28, Proposition 15.3, p. 151] we have the following lemma.

Lemma 3.4. *Assume that $\lambda = \lambda_0$ is an isolated point spectrum of \mathcal{F}_0 and has finite algebraic multiplicity $m(\lambda_0)$. Let $\Gamma \subset \rho(\mathcal{F}_0)$ be a small cycle surrounding λ_0 such that λ_0 is the only spectrum point inside the bounded open set surrounded by Γ . Assume that the following statements hold:*

- (i) *there exists a small constant $\tilde{\epsilon}_0 > 0$ such that $\Gamma \subset \rho(\mathcal{F}(\epsilon))$ for all ϵ in $(0, \tilde{\epsilon}_0]$.*
- (ii) *$|P_\Gamma(\mathcal{F}(\epsilon)) - P_\Gamma(\mathcal{F}_0)| \rightarrow 0$ as $\epsilon \rightarrow 0$.*

Then for sufficiently small ϵ , there are \tilde{m}_ϵ eigenvalues $\lambda_1(\epsilon), \dots, \lambda_{\tilde{m}_\epsilon}(\epsilon)$ of $\mathcal{F}(\epsilon)$ inside Γ and $\lambda_j(\epsilon) \rightarrow \lambda_0$ as $\epsilon \rightarrow 0$, and the total algebraic multiplicity of $\lambda_j(\epsilon)$ is equal to $m(\lambda_0)$.

By applying Lemmas 3.3 and 3.4, we have the following results on the spectral perturbation.

Lemma 3.5. *Assume that $\lambda = \lambda_0$ is an isolated point spectrum of \mathcal{F}_0 and has finite algebraic multiplicity $m(\lambda_0)$. Let $\Gamma \subset \rho(\mathcal{F}_0)$ be a small cycle surrounding λ_0 such that λ_0 is the only spectrum point inside the bounded open set surrounded by Γ . Then there exists a constant $\tilde{\epsilon}_0$ with $0 < \tilde{\epsilon}_0 \leq \epsilon_0$ such that for each ϵ in $(0, \tilde{\epsilon}_0]$, there are \tilde{m}_ϵ eigenvalues $\lambda_1(\epsilon), \dots, \lambda_{\tilde{m}_\epsilon}(\epsilon)$ of $\mathcal{F}(\epsilon)$ inside Γ and $\lambda_j(\epsilon) \rightarrow \lambda_0$ as $\epsilon \rightarrow 0$, and the total algebraic multiplicity of $\lambda_j(\epsilon)$ is equal to $m(\lambda_0)$.*

Proof. In order to prove this lemma, it suffices to check the conditions **(i)** and **(ii)** in Lemma 3.4 hold. Since $(\lambda I - \mathcal{F}_0)^{-1}$ is analytic for $\lambda \in \rho(\mathcal{F}_0)$, there exists a constant $M(\Gamma) > 0$, which is dependent of Γ , such that

$$(3.17) \quad |(\lambda I - \mathcal{F}_0)^{-1}| \leq M(\Gamma), \quad \lambda \in \Gamma.$$

Recall that $a(\epsilon)$ and $b(\epsilon)$ are defined as in Lemma 3.3 and satisfy the limits in (3.12). Then there exists a constant $\tilde{\epsilon}_0$ with $0 < \tilde{\epsilon}_0 \leq \epsilon_0$ such that

$$a(\epsilon)|(\lambda - \mathcal{F}_0)^{-1}| + b(\epsilon) < 1, \quad \lambda \in \Gamma, \quad \epsilon \in (0, \tilde{\epsilon}_0].$$

Hence by Lemma 3.3 and [33, Theorem 1.16, p. 196], we have $\Gamma \subset \rho(\mathcal{F}(\epsilon))$ for all ϵ in $(0, \tilde{\epsilon}_0]$, and

$$(3.18) \quad \begin{aligned} |(\lambda I - \mathcal{F}(\epsilon))^{-1}| &\leq \frac{|(\lambda I - \mathcal{F}_0)^{-1}|}{1 - a(\epsilon)|(\lambda I - \mathcal{F}_0)^{-1}| - b(\epsilon)}, \\ |(\lambda I - \mathcal{F}(\epsilon))^{-1} - (\lambda I - \mathcal{F}_0)^{-1}| &\leq \frac{|(\lambda I - \mathcal{F}_0)^{-1}|(a(\epsilon)|(\lambda I - \mathcal{F}_0)^{-1}| + b(\epsilon))}{1 - a(\epsilon)|(\lambda I - \mathcal{F}_0)^{-1}| - b(\epsilon)}, \end{aligned}$$

for $\lambda \in \Gamma$ and $\epsilon \in (0, \tilde{\epsilon}_0]$. This yields that $\Gamma \subset \rho(\mathcal{F}(\epsilon))$ for all ϵ in $(0, \tilde{\epsilon}_0]$.

Let the spectral projections $P_\Gamma(\mathcal{F}_0)$ and $P_\Gamma(\mathcal{F}(\epsilon))$ for $\epsilon \in (0, \tilde{\epsilon}_0]$ be defined by

$$\begin{aligned} P_\Gamma(\mathcal{F}_0) &= \frac{1}{2\pi i} \oint_\Gamma (\lambda I - \mathcal{F}_0)^{-1} d\lambda, \\ P_\Gamma(\mathcal{F}(\epsilon)) &= \frac{1}{2\pi i} \oint_\Gamma (\lambda I - \mathcal{F}(\epsilon))^{-1} d\lambda, \end{aligned}$$

respectively. Then by (3.17) and (3.18), we have

$$|P_\Gamma(\mathcal{F}_0) - P_\Gamma(\mathcal{F}(\epsilon))| \leq \frac{1}{2\pi} \oint_\Gamma |(\lambda I - \mathcal{F}(\epsilon))^{-1} - (\lambda I - \mathcal{F}_0)^{-1}| |d\lambda|.$$

Consequently, by the limits in (3.12), the inequalities (3.17) and (3.18), we have that

$$|P_\Gamma(\mathcal{F}(\epsilon)) - P_\Gamma(\mathcal{F}_0)| \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0.$$

Thus, the proof for this lemma is finished by applying Lemma 3.4. \square

Remark 3.6. The perturbation theory for a family of linear operators $H_\kappa = H + \kappa W$ was considered in Chapter 15 of [28], where H_κ are required to be analytic in κ . In Lemma 3.5, we do not require the smoothness of $\mathcal{F}(\epsilon)$ in ϵ . From Lemma 3.5, the point spectrum of \mathcal{F}_0 is often said to be *stable under the perturbations* $\mathcal{F}_1(\epsilon)$ for sufficiently small $\epsilon > 0$.

3.3. Instability of periodic waves. In order to study the stability of periodic waves emerging from fold-Hopf equilibria, we first study the spectrum $\sigma(\mathcal{F}_0)$ of \mathcal{F}_0 .

Lemma 3.7. *Let the operator \mathcal{F}_0 be defined by (3.8). Assume that $A > 0$. Then the following statements hold:*

- (i) \mathcal{F}_0 has a positive eigenvalue $\lambda = A^2\mu_0^2$ with an eigenfunction $(f_w(P_0, \alpha_0), g_w(P_0, \alpha_0))^T$.
- (ii) the algebraic multiplicity and the geometric multiplicity of $\lambda = A^2\mu_0^2$ are both equal to one.

Proof. For simplification, we only give the proof for $A = 1$. For a general $A > 0$, the results in this lemma can be obtained by taking a rescaling $\xi \rightarrow \xi/A$ in $\mathcal{F}_0(u, w)^T = A^2\lambda(u, w)^T$. Set $v = u_\xi$ and $Y = (u, v, w)^T$. Then the eigenvalue problem $\mathcal{F}_0(u, w)^T = \lambda(u, w)^T$ is equivalent to the following system

$$(3.19) \quad \frac{dY}{d\xi} = A_0(\lambda)Y, \quad A_0(\lambda) = \begin{pmatrix} 0 & 1 & 0 \\ \lambda - f_u(P_0, \alpha_0) & c_0 & -f_w(P_0, \alpha_0) \\ \frac{1}{c_0}g_u(P_0, \alpha_0) & 0 & \frac{1}{c_0}(g_w(P_0, \alpha_0) - \lambda) \end{pmatrix}$$

subject to the periodic boundary condition $Y(1) = Y(0)$. By Lemma 2.1 we have

$$\begin{aligned} \det(A_0(\lambda)) &= -\frac{1}{c_0} (\lambda^2 - (f_u(P_0, \alpha_0) + g_w(P_0, \alpha_0))\lambda \\ &\quad + f_u(P_0, \alpha_0)g_w(P_0, \alpha_0) - f_w(P_0, \alpha_0)g_u(P_0, \alpha_0)) \\ &= -\frac{1}{c_0} \lambda (\lambda - \mu_0^2). \end{aligned}$$

Then for $\lambda = \mu_0^2 > 0$, system (3.19) has a nonzero constant solution $(f_w(P_0, \alpha_0), 0, g_w(P_0, \alpha_0))^T$. This proves (i).

By (2.2) and substituting $\lambda = \mu_0^2$ into (3.19), we have that the rank of $A_0(\mu_0^2)$ is two. Then $\text{Ker}(\mathcal{F}_0 - \mu_0^2 I)$ is one-dimensional and spanned by its eigenfunction $(f_w(P_0, \alpha_0), 0, g_w(P_0, \alpha_0))^T$. Hence, the geometric multiplicity of $\lambda = \mu_0^2$ is one. Since the characteristic polynomial associated with $A_0(\lambda)$ is in the form

$$\begin{aligned} G(z, \lambda) &:= z \left(z^2 - \left(c_0 + \frac{1}{c_0}(g_w(P_0, \alpha_0) - \lambda) \right) z + f_u(P_0, \alpha_0) + g_w(P_0, \alpha_0) - 2\lambda \right) \\ &\quad - \frac{1}{c_0} (\lambda^2 - (f_u(P_0, \alpha_0) + g_w(P_0, \alpha_0))\lambda + f_u(P_0, \alpha_0)g_w(P_0, \alpha_0) - f_w(P_0, \alpha_0)g_u(P_0, \alpha_0)), \end{aligned}$$

by (2.2) we have

$$G(z, \lambda) = z \left(z^2 + \frac{1}{c_0} \lambda z + \mu_0^2 - 2\lambda \right) - \frac{1}{c_0} (\lambda^2 - \mu_0^2 \lambda).$$

Let $z_j(\lambda)$, $j = 1, 2, 3$, denote the zeros of $G(z, \lambda)$ for each $\lambda \in \mathbb{C}$. Especially, for $\lambda = \mu_0^2$ the zeros of $G(z, \mu_0^2)$ are

$$z_1(\mu_0^2) = 0, \quad z_{2,3}(\mu_0^2) = -\frac{1}{2c_0} \mu_0^2 \pm \sqrt{\mu_0^2 + \frac{1}{4c_0^2} \mu_0^4} \neq 0,$$

and $z_1'(\mu_0^2) = -\frac{1}{c_0}$. Then the Evans function $E(\lambda, \mu)$ in (3.6) satisfies

$$E(\lambda, 0) = \det(e^{A_0(\lambda)} - I_3) = \prod_{j=1}^3 (e^{z_j(\lambda)} - 1).$$

Hence, $E(\mu_0^2, 0) = 0$ and

$$\frac{\partial E}{\partial \lambda}(\mu_0^2, 0) = z_1'(\mu_0^2) \cdot e^{z_1(\mu_0^2)} \cdot (e^{z_2(\mu_0^2)} - 1) \cdot (e^{z_3(\mu_0^2)} - 1) \neq 0.$$

This together with [32, Lemma 8.4.1] yields **(ii)**. Thus, the proof is now complete. \square

By Lemmas 3.5 and 3.7, we have the following theorem.

Theorem 3.8. (Instability of periodic traveling waves) *Assume that there exists a sufficiently small $\epsilon_0 > 0$, and two continuous functions $\alpha_1(\epsilon)$ and $c_1(\epsilon)$ with $\alpha_1(\epsilon) = O(\epsilon)$ and $c_1(\epsilon) = O(\epsilon)$ for $\epsilon \in (0, \epsilon_0]$ such that a small-amplitude periodic solution (ϕ, ϕ_ξ, ψ) with period A_ϵ bifurcates from a fold-Hopf point P_0 in the 3D system (1.3) with $(\alpha, c) = (\alpha_0 + \alpha_1(\epsilon), c_0 + c_1(\epsilon))$ for each $\epsilon \in (0, \epsilon_0]$, where*

$$|(\phi, \phi_\xi, \psi) - P_0| = O(\epsilon), \quad A_\epsilon = A + O(\epsilon), \quad A > 0.$$

Then there exists a constant $\tilde{\epsilon}_0$ with $0 < \tilde{\epsilon}_0 \leq \epsilon_0$ such that for each ϵ with $0 < \epsilon \leq \tilde{\epsilon}_0$, the corresponding periodic traveling wave (ϕ, ψ) with wave speed $c_0 + c_1(\epsilon)$ in the system (1.1) with $\alpha_0 + \alpha_1(\epsilon)$ are spectrally unstable.

Proof. By Lemma 3.7, we have that \mathcal{F}_0 has an positive eigenvalue $\lambda_0 = A^2 \mu_0^2$, whose algebraic multiplicity is one. Let Γ in $\text{Re} \lambda > 0$ be a small cycle surrounding $A^2 \mu_0^2$ such that $\lambda_0 = A^2 \mu_0^2$ is the only spectrum point inside Γ . By Lemma 3.5, there exists a constant $\tilde{\epsilon}_0$ with $0 < \tilde{\epsilon}_0 \leq \epsilon_0$ such that for each ϵ in $(0, \tilde{\epsilon}_0]$, there exists a unique eigenvalue $\lambda_1(\epsilon)$ of $\mathcal{F}(\epsilon)$ inside Γ and $\lambda_1(\epsilon) \rightarrow \lambda_0$ as $\epsilon \rightarrow 0$. Then $\sigma(\mathcal{F}(\epsilon))$ has an eigenvalue $\lambda_1(\epsilon)$ with positive real part for $\epsilon \in (0, \tilde{\epsilon}_0]$. This together with Lemma 3.2 yields that the periodic traveling wave (ϕ, ψ) is spectrally unstable. Therefore, the proof is now complete. \square

As a corollary of Theorem 3.8, we have the stability of periodic waves obtained in Theorem 2.3.

Corollary 3.2. *Assume that (ϕ, ψ) is a periodic traveling wave obtained as in Theorem 2.3. Then this periodic traveling wave is spectrally unstable.*

4. APPLICATIONS TO THE FITZHUGH-NAGUMO SYSTEM

In this section, we apply the obtained results to the FitzHugh-Nagumo system. The FitzHugh-Nagumo system is a simplification of the Hodgkin-Huxley model [29] describing the propagation of nerve impulses in axons. Since the FitzHugh-Nagumo system was proposed by FitzHugh [20] and Nagumo, Arimoto and Yoshizawa [40], many efforts have been devoted to understanding complex oscillations and traveling waves in this system in the past tens of years. See [4, 5, 9, 11, 12, 13, 14, 23, 24, 31, 44] and the references therein.

4.1. Mathematical model. We follow [11] and study the FitzHugh-Nagumo system with an applied current

$$(4.1) \quad \begin{aligned} \frac{\partial u}{\partial t} &= du_{xx} + f(u) - w + p, \\ \frac{\partial w}{\partial t} &= \delta(u - \gamma w), \end{aligned}$$

where x is a one-dimensional spatial variable, t is time, the state u represents the plasma membrane electric potential with the diffusion rate d , the state w describes the combined inactivation effects of potassium and sodium ion channels, p is the applied current, f is a bistable function in the form

$$f(u) = u(u - 1)(\alpha - u),$$

and all system parameters are positive.

In the moving coordinate frame $(\xi, t) = (x + ct, t)$, a periodic wave profile $(\phi(\cdot), \psi(\cdot))$ is a periodic steady state of the following system

$$(4.2) \quad \begin{aligned} u_t &= du_{\xi\xi} - cu_{\xi} + f(u) - w + p, \\ w_t &= -cw_{\xi} + \delta(u - \gamma w). \end{aligned}$$

Set $v = u_{\xi}$. Then the vector (ϕ, ϕ_{ξ}, ψ) is a periodic solution of the three-dimensional ordinary differential system

$$(4.3) \quad \begin{aligned} \frac{du}{d\xi} &= \dot{u} = v, \\ \frac{dv}{d\xi} &= \dot{v} = \frac{1}{d}(-f(u) + cv + w - p), \\ \frac{dw}{d\xi} &= \dot{w} = \frac{\delta}{c}(u - \gamma w). \end{aligned}$$

For convenience, we call (4.3) the *3D-FHN system*. Throughout this paper, the values of the parameters α , d and δ are fixed as they were given in [4, 11, 14, 23], that is,

$$(4.4) \quad \alpha = 0.1, \quad d = 5.0, \quad \delta = 0.01.$$

Then the 3D-FHN system (4.3) depends on three parameters c , γ and p .

4.2. Fold-Hopf equilibrium. If $(u_0, v_0, w_0)^T$ in \mathbb{R}^3 is an equilibrium of the 3D-FHN system (4.3), then $(u_0, v_0, w_0)^T$ satisfies the equations:

$$v_0 = 0, \quad f(u_0) + p = w_0, \quad u_0 = \gamma w_0.$$

This yields that the equilibria of the 3D-FHN system (4.3) are determined by the roots of equation

$$g(u, \gamma) = p, \quad u \in \mathbb{R}, \quad \gamma, p > 0,$$

where

$$(4.5) \quad g(u, \gamma) := \frac{1}{\gamma}u - f(u) = u^3 - 1.1u^2 + \left(0.1 + \frac{1}{\gamma}\right)u.$$

For each $\gamma, p > 0$, the results on the zeros of $g(u, \lambda) = p$ are summarized as follows.

Lemma 4.1. *Let $N(\gamma, p)$ denote the number of the zeros for $g(u, \gamma) = p$ with $\gamma > 0$ and $p > 0$. Then the following statements hold:*

- (i) *If γ satisfies $0 < \gamma \leq \frac{300}{91}$, then $N(\gamma, p) = 1$ for every $p > 0$.*
- (ii) *If γ satisfies $\gamma > \frac{300}{91}$, then $g_u(u, \gamma) = 0$ has precisely two positive roots $u_1(\gamma)$ and $u_2(\gamma)$ with $0 < u_1(\gamma) < u_2(\gamma)$ for each $\gamma > \frac{300}{91}$. Let $u_0(\gamma)$ and $u_3(\gamma)$ be defined by*

$$\begin{aligned} g(u_0(\gamma)) &= g(u_2(\gamma)), & u_0(\gamma) &< u_1(\gamma), \\ g(u_3(\gamma)) &= g(u_1(\gamma)), & u_3(\gamma) &> u_2(\gamma). \end{aligned}$$

Moreover, the following dichotomies hold:

- (ii.1) *If γ satisfies $\frac{300}{91} < \gamma < \frac{400}{81}$, then $N(\gamma, p) = 1$ for $p > g(u_1(\gamma))$ and $0 < p < g(u_2(\gamma))$, $N(\gamma, p) = 2$ for $p = g(u_1(\gamma))$ and $p = g(u_2(\gamma))$, and $N(\gamma, p) = 3$ for $g(u_2(\gamma)) < p < g(u_1(\gamma))$.*
- (ii.2) *If γ satisfies $\gamma \geq \frac{400}{81}$, then $N(\gamma, p) = 1$ for $p > g(u_1(\gamma))$, $N(\gamma, p) = 2$ for $p = g(u_1(\gamma))$, and $N(\gamma, p) = 3$ for $0 < p < g(u_1(\gamma))$.*

The graphs of $g(u, \gamma)$ for different (γ, p) are shown in Figure 1.

Proof. Note that the first-order partial derivative of g with respect to u is in the form

$$g_u(u, \gamma) = 3u^2 - 2.2u + 0.1 + \frac{1}{\gamma}.$$

Thus, $g_u(u, \gamma) \geq 0$ for all $u \geq 0$ and $0 < \gamma \leq \frac{300}{91}$. This yields (i).




FIGURE 1. The graphs of g with respect to u for different values of γ : 1(a) $\gamma \leq \frac{300}{91}$, 1(b) $\frac{300}{91} < \gamma < \frac{400}{81}$, 1(c) $\gamma = \frac{400}{81}$ and 1(d) $\gamma > \frac{400}{81}$.

For each $\gamma > \frac{300}{91}$, equation $g_u(u, \gamma) = 0$ has precisely two zeros $u_1(\gamma)$ and $u_2(\gamma)$ with $u_1(\gamma) < u_2(\gamma)$, and $g(u, \gamma)$ has a zero of multiplicity two for $\gamma = \frac{400}{81}$. Hence $u_0(\gamma)$ and $u_3(\gamma)$ are well-defined, and the statements in **(ii)** can be obtained by a standard analysis. Thus, the proof is complete. \square

We next study the conditions under which an equilibrium is a fold-Hopf equilibrium. If the 3D-FHN system (4.3) with (4.4) has an equilibrium $(u_0, v_0, w_0)^T$ in \mathbb{R}^3 , then $g(u_0, \gamma) = p$. Clearly, the Jacobian matrix $J(u_0, v_0, w_0)$ at this point is in the form

$$J(u_0, v_0, w_0) = \begin{pmatrix} 0 & 1 & 0 \\ -\frac{1}{d}f'(u_0) & \frac{c}{d} & \frac{1}{d} \\ \frac{\delta}{c} & 0 & -\frac{\delta\gamma}{c} \end{pmatrix},$$

and the eigenvalues λ_j , $j = 1, 2, 3$, of $J(u_0, v_0, w_0)$ are determined by the zeros of its characteristic polynomial

$$\begin{aligned} & \det(\lambda I_3 - J(u_0, v_0, w_0)) \\ &= \lambda^3 + \left(\frac{\delta\gamma}{c} - \frac{c}{d} \right) \lambda^2 + \frac{1}{d} (f'(u_0) - \delta\gamma) \lambda + \frac{\delta}{cd} (\gamma f'(u_0) - 1), \end{aligned}$$

where I_3 is the 3×3 identity matrix. Assume that $(u_0, v_0, w_0)^T$ is a fold-Hopf equilibrium. Then without loss of generality, we assume $\lambda_1 = 0$ and $\lambda_{2,3} = \pm \mathbf{i}\mu_0$ for some $\mu_0 > 0$. Then

$$\gamma f'(u_0) - 1 = 0, \quad \sum_{j=1}^3 \lambda_j = \frac{\delta\gamma}{c} - \frac{c}{d} = 0, \quad \frac{1}{d} (f'(u_0) - \delta\gamma) = \mu_0^2 > 0.$$

This yields that if $(u_0, v_0, w_0)^T$ in \mathbb{R}^3 is a fold-Hopf equilibrium, then (c, γ, p) satisfies

$$(4.6) \quad \begin{aligned} g(u_0, \gamma) &= p, & \frac{1}{\gamma} &= f'(u_0) > 0.01\gamma > 0, \\ c &= \sqrt{0.05\gamma}, & \mu_0^2 &= \frac{1}{500\gamma}(100 - \gamma^2) > 0. \end{aligned}$$

More properties on the fold-Hopf equilibria of the 3D-FHN system (4.3) are summarized as follows.

Lemma 4.2. *Let $c = \sqrt{0.05\gamma}$ for each $\gamma > 0$. Then the following statements hold:*

- (i) *For $0 < \gamma < \frac{300}{91}$ and $\gamma \geq 10$, the 3D-FHN system (4.3) has no fold-Hopf equilibria.*
- (ii) *For $\gamma = \frac{300}{91}$, the 3D-FHN system (4.3) has precisely one fold-Hopf equilibrium at $(\frac{11}{30}, 0, \frac{1001}{9000})^T$, and p takes the value $\frac{1331}{27000}$.*
- (iii) *For $\frac{300}{91} < \gamma < \frac{400}{81}$, the 3D-FHN system (4.3) has precisely two fold-Hopf equilibria at $(u_1(\gamma), 0, u_1(\gamma)/\gamma)^T$ for $p = g(u_1(\gamma), \gamma)$ and $(u_2(\gamma), 0, u_2(\gamma)/\gamma)^T$ for $p = g(u_2(\gamma), \gamma)$, respectively, where $u_1(\gamma)$ and $u_2(\gamma)$ are defined as in (ii) of Lemma 4.1.*
- (iv) *For $\frac{400}{81} \leq \gamma < 10$, the 3D-FHN system (4.3) has precisely one fold-Hopf equilibrium at $(u_1(\gamma), 0, u_1(\gamma)/\gamma)^T$ for $p = g(u_1(\gamma), \gamma)$, where $u_1(\gamma)$ is defined as in (ii) of Lemma 4.1.*

Proof. By the proof for (i) of Lemma 4.1, we have that $\frac{1}{\gamma} > f'(u)$ for each $u > 0$ and γ with $0 < \gamma < \frac{300}{91}$. Then (i) holds for $0 < \gamma < \frac{300}{91}$. Since $\frac{1}{\gamma} \leq 0.01\gamma$ for $\gamma \geq 10$, the proof for (i) is finished by (4.6).

Note that $\frac{1}{\gamma} = f'(u)$ implies $g'(u, \gamma) = 0$. Then by Lemma 4.1 and a direct computation, the remaining statements hold. Thus, the proof is complete. \square

4.3. Existence and stability of periodic waves. By Lemma 2.2 and Theorem 3.8, we can obtain the existence and stability of periodic traveling waves emerging fold-Hopf equilibria. More precisely, we have the following theorem.

Theorem 4.3. *Let (α, d, δ) be in the form (4.4). Assume that the 3D-FHN system (4.3) with $(c, \gamma, p) = (c_0, \gamma_0, p_0)$ has a fold-Hopf equilibrium $(u_0, v_0, w_0)^T$ whose linearization has the eigenvalues $\lambda_1 = 0$ and $\lambda_{2,3} = \pm \mathbf{i}\mu_0$ for some $\mu_0 > 0$. Then there is a sufficiently small $\epsilon_0 > 0$, and two continuous functions $\gamma(\epsilon)$ and $p(\epsilon)$ given by*

$$\gamma(\epsilon) = \gamma_0 + \gamma_1\epsilon + O(\epsilon^2), \quad p(\epsilon) = p_0 + O(\epsilon^3),$$

for $\epsilon \in (0, \epsilon_0]$ and $\gamma_1 \neq 0$ with $\gamma_1(11 - 30u_0) > 0$, such that a periodic solution (ϕ, ϕ_ξ, ψ) with period A_ϵ bifurcates from $(u_0, v_0, w_0)^T$ in the 3D-FHN system (4.3) with $(c, \gamma, p) = (c_0 + O(\epsilon^3), \gamma(\epsilon), p(\epsilon))$ for each $\epsilon \in (0, \epsilon_0]$, and

$$|(\phi, \phi_\xi, \psi) - (u_0, v_0, w_0)| = O(\epsilon), \quad A_\epsilon = 2\pi/\mu_0 + O(\epsilon), \quad \epsilon \in (0, \epsilon_0].$$

Furthermore, the corresponding periodic traveling wave (ϕ, ψ) in the FitzHugh-Nagumo system (4.1) is spectrally unstable.

Proof. We begin by proving the first statement by the averaging theory. After translating the equilibrium (u_0, v_0, w_0) of the 3D-FHN system (4.3) to the origin, we change the variables by

$$(4.7) \quad (u, v, w) \rightarrow (\epsilon(u + \gamma_0 w), \epsilon \mu_0 v, \epsilon(\delta_0 \gamma_0 u - c_0 \mu_0 v + w)),$$

then the 3D-FHN system (4.3) is converted into

$$(4.8) \quad \begin{aligned} \dot{u} &= \mu_0 v - \frac{1}{d_0 \mu_0^2} (c_0 g_1(u, w) + g_2(u, v, w)) \epsilon + O(\epsilon^2), \\ \dot{v} &= -\mu_0 u + \frac{1}{\mu_0} g_1(u, w) \epsilon + O(\epsilon^2), \\ \dot{w} &= \frac{1}{d_0 \gamma_0 \mu_0^2} (c_0 g_1(u, w) + g_2(u, v, w)) \epsilon + O(\epsilon^2), \end{aligned}$$

where

$$\begin{aligned} g_1(u, w) &= \frac{1}{d_0} (3u_0 - 1.1)(u + \gamma_0 w)^2, \\ g_2(u, v, w) &= -\delta_0 \gamma_1 (\delta_0 \gamma_0 u - c_0 \mu_0 v + w). \end{aligned}$$

By another change of variables

$$(u, v, w) \rightarrow (r \cos \theta, r \sin \theta, w), \quad r \geq 0, \quad 2\pi \geq \theta \geq 0,$$

we have

$$(4.9) \quad \begin{aligned} \dot{r} &= \left(-\frac{1}{d_0 \mu_0^2} (c_0 g_1 + g_2) \cos \theta + \frac{1}{\mu_0} g_1 \sin \theta \right) \epsilon + O(\epsilon^2), \\ r \dot{\theta} &= -\mu_0 r + \left(\frac{1}{d_0 \mu_0^2} (c_0 g_1 + g_2) \sin \theta + \frac{1}{\mu_0} g_1 \cos \theta \right) \epsilon + O(\epsilon^2), \\ \dot{w} &= \frac{1}{d_0 \gamma_0 \mu_0^2} (c_0 g_1 + g_2) \epsilon + O(\epsilon^2), \end{aligned}$$

where we write $g_1 = g_1(r \cos \theta, w)$ and $g_2 = g_2(r \cos \theta, r \sin \theta, w)$ for short. Then for sufficiently small $|r|$ and $|\epsilon|$, we change system (4.8) into a nonautonomous system

$$(4.10) \quad \begin{aligned} \frac{dr}{d\theta} &= \left(\frac{1}{d_0 \mu_0^3} (c_0 g_1 + g_2) \cos \theta - \frac{1}{\mu_0^2} g_1 \sin \theta \right) \epsilon + O(\epsilon^2) := H_1(\theta, r, w) \epsilon + O(\epsilon^2), \\ \frac{dw}{d\theta} &= -\frac{1}{d_0 \gamma_0 \mu_0^3} (c_0 g_1 + g_2) \epsilon + O(\epsilon^2) := H_2(\theta, r, w) \epsilon + O(\epsilon^2). \end{aligned}$$

Clearly, the functions $H_j(\cdot, r, w)$ are periodic functions of period 2π . In order to find the periodic solutions in system (4.10), we apply the averaging theory and compute the averaged function of order one $(G_1(r, w), G_2(r, w))$, which is obtained by the following integrals

$$G_j(r, w) = \int_0^{2\pi} H_j(\theta, r, w) d\theta, \quad j = 1, 2.$$

These yields

$$\begin{aligned} G_1(r, w) &= \frac{\pi\gamma_0 r}{d_0^2 \mu_0^3} (2(3u_0 - 1.1)w + d_0 \delta_0^2 \gamma_1), \\ G_2(r, w) &= -\frac{2\pi}{d_0 \gamma_0 \mu_0^3} \left(\frac{c_0}{d_0} (3u_0 - 1.1) \left(\frac{1}{2} r^2 + \gamma_0^2 w^2 \right) - \delta_0 \gamma_1 w \right). \end{aligned}$$

Solving $G_1(r, w) = 0$ and $G_2(r, w) = 0$ in the set $r > 0$ yields that

$$r = r_*^+ = \frac{d_0 \gamma_0 |\gamma_1|}{|3u_0 - 1.1|} \sqrt{\left| \frac{c_0 \gamma_0^4 - 2\delta_0}{2c_0} \right|}, \quad w_* = \frac{5d_0 \gamma_0^2 \gamma_1}{11 - 30u_0},$$

where $11 - 30u_0 \neq 0$ can be checked by the proof of Lemma 4.2. Note that $2\delta_0 - c_0 \gamma_0^4 = 0$ if and only if $\gamma_0 = 1/5^{1/3} < 1$. Then by Lemma 4.2, we have $2\delta_0 - c_0 \gamma_0^4 \neq 0$. This implies

$$\det \begin{pmatrix} \frac{\partial G_1}{\partial r} & \frac{\partial G_1}{\partial w} \\ \frac{\partial G_2}{\partial r} & \frac{\partial G_2}{\partial w} \end{pmatrix} \Big|_{(r_*, w_*)} = \frac{2\pi^2 (2\delta_0 - c_0 \gamma_0^4) \gamma_0^2 \gamma_1^2}{d_0^2 \mu_0^6} \neq 0.$$

Thus by Lemma 2.2, for sufficiently small ϵ there exists an isolated periodic solution $\varphi(\theta, \epsilon)$ of period 2π in (4.10) such that $\varphi(0, \epsilon) \rightarrow (r_*^+, w_*)$ as $\epsilon \rightarrow 0$. Consequently, by (4.7) and (4.9), the first statement holds. The second one is obtained by Theorem 3.8. Therefore, the proof is now complete. \square

5. CONCLUDING REMARKS

We have investigated the existence and stability of small-amplitude periodic traveling waves emerging from fold-Hopf equilibria in a system of one reaction-diffusion equation coupled with one ordinary differential equation. This is a complement to the existing works on homoclinic, heteroclinic and periodic waves in systems such as FitzHugh-Nagumo system, caricature calcium models and other models in the real-world applications. Our work has three main contributions:

(i) Applying the recent results on the averaging theory, we not only prove the existence of small-amplitude periodic traveling waves emerging from the fold-Hopf bifurcations, but also give the periods for the obtained periodic waves.

(ii) Tsai et al. [47] studied a simplified model of calcium dynamics

$$(5.1) \quad \begin{aligned} \frac{\partial u}{\partial t} &= Du_{xx} + f(u)w - g(u), \\ \frac{\partial w}{\partial t} &= -\gamma(f(u)w - g(u)), \end{aligned}$$

where

$$f(u) = \alpha + k_f \frac{u^2}{u^2 + \varphi_1^2} \cdot \frac{\varphi_2}{u + \varphi_2}, \quad g(u) = f(u)u + k_s u.$$

We refer to [47] for the biological descriptions of the model parameters. So model (5.1) has exactly the same form as system (1.1). Tsai et al. [47] numerically found that the small-amplitude periodic traveling waves from fold-Hopf bifurcations in (5.1) are spectrally unstable. By a similar method used to the FitzHugh-Nagumo system (4.1) (see Section 4), we can prove the spectral instability of a class of small-amplitude periodic waves from fold-Hopf bifurcation in model (5.1).

(iii) We construct the periodic waves near the fold-Hopf equilibria via the averaging theory and prove the spectral instability of the periodic waves for the FitzHugh-Nagumo system with an applied current. This indicates that unstable periodic wave impulses can arise in axons through the fold-Hopf bifurcations.

Analyzing the spectral stability of periodic traveling waves is not an easy task (see [21] for more details). More recently, much attention has been paid to the stability of small-amplitude periodic traveling waves bifurcating from a homogeneous rest state in, such as, scalar viscous balance laws [3] and Hamiltonian partial differential equations [35]. As shown in [3, 35], these periodic traveling waves appear to inherit the instability of the homogeneous rest state. After proving the instability of the corresponding homogeneous rest state, we apply the perturbation theory to show the similar phenomenon in the coupled system (1.1).

Note that in the present paper we have only investigated a class of periodic waves arising from fold-Hopf bifurcations in the coupled system (1.1). As shown in the references [22, 36], there exist complex dynamical behaviors arising near the fold-Hopf equilibria through the fold-Hopf bifurcations. It is also interesting to further study the properties of other perturbed solutions for the coupled system (1.1). By the similar method used in the present paper, it is also possible to study the stability of small-amplitude periodic traveling waves arising in other equations such as the classical and generalized Ginzburg-Landau equations [15, 17].

ACKNOWLEDGMENTS

The authors would like to thank the editor and the anonymous referees for carefully reading the manuscript and providing valuable comments. This work was partly supported by the Science and Technology Project for Excellent Postdoctors of Hubei.

REFERENCES

1. J. Alexander, R. Gardner, and C. K. R. T. Jones, A topological invariant arising in the stability analysis of traveling waves, *J. reine angew. Math.* **410** (1990), 167–212.
2. J. Alexander, M. G. Grillakis, C. K. R. T. Jones, and B. Sandstede, Stability of pulses on optical fibers with phase-sensitive amplifiers, *Z. Angew. Math. Phys.* **48** (1997), 175–192.
3. E. Álvarez and R. G. Plaza, Existence and spectral instability of bounded spatially periodic traveling waves for scalar viscous balance laws, <https://doi.org/10.1090/qam/1591>.
4. G. Arioli and H. Koch, Existence and stability of traveling pulse solutions of the FitzHugh-Nagumo equation, *Nonlinear Anal.* **113** (2015), 51–70.
5. R. Cai, Z. He, Y. Liu, J. Duan, J. Kurths, and X. Li, Effects of Lévy noise on the FitzHugh-Nagumo model: a perspective on the maximal likely trajectories, *J. Theoret. Biol.* **480** (2019), 166–174.
6. M. Cândido, J. Llibre, and D. Novaes, Persistence of periodic solutions for higher order perturbed differential systems via Lyapunov-Schmidt reduction, *Nonlinearity* **30** (2017), 3560–3586.
7. M. Cândido, D. Novaes, and C. Valls, Periodic solutions and invariant torus in the Rössler system, *Nonlinearity* **33** (2020), 4512–4538.
8. G. A. Carpenter, A geometric approach to singular perturbation problems with applications to nerve impulse equations, *J. Differential Equations* **23** (1977), 335–367.
9. P. Carter and B. Sandstede, Fast pulses with oscillatory tails in the FitzHugh-Nagumo system, *SIAM J. Math. Anal.* **47** (2015), 3393–3441.
10. P. Carter and A. Scheel, Wave train selection by invasion fronts in the FitzHugh-Nagumo equation, *Nonlinearity* **31** (2018), 5536–5572.
11. A. R. Champneys, V. Kirk, E. Knobloch, B. Oldeman, and J. Sneyd, When Shil’nikov meets Hopf in excitable systems, *SIAM J. Appl. Dyn. Syst.* **6** (2007), 663–693.
12. C. Chen and X. Hu, Stability analysis for standing pulse solutions to FitzHugh-Nagumo equations, *Calc. Var. Partial Differential Equations* **49** (2014), 827–845.
13. P. Cornwell and C. K. R. T. Jones, On the existence and stability of fast traveling waves in a doubly diffusive FitzHugh-Nagumo system, *SIAM J. Appl. Dyn. Syst.* **17** (2018), 754–787.
14. A. Czechowski and P. Zgliczyński, Existence of periodic solutions of the FitzHugh-Nagumo equations for an explicit range of the small parameter, *SIAM J. Appl. Dyn. Syst.* **15** (2016), 1615–1655.
15. A. Doelman, *On the Nonlinear Evolution of Patterns (Modulation Equations and Their Solutions)*, PhD thesis, University of Utrecht, 1990.
16. A. Doelman, B. Sandstede, A. Scheel, and G. Schneider, *The Dynamics of Modulated Wave Trains*, Mem. Amer. Math. Soc. **199** (2009).
17. J. Duan, P. Holmes, Fronts, domain walls and pulses in a generalized Ginzburg-Landau equation, *Proc. Edinburgh Math. Soc.* **38** (1995), 77–97.
18. J. Evans, Nerve axon equations. IV. The stable and the unstable impulse, *Indiana Univ. Math. J.* **24** (1974/75), 1169–1190.
19. E. G. Eszter, *Evans Function Analysis of the Stability of Periodic Traveling Wave Solutions Associated with the FitzHugh-Nagumo System*, PhD thesis, University of Massachusetts Amherst, 1999.
20. R. FitzHugh, Thresholds and plateaus in the Hodgkin-Huxley nerve conduction equations, *J. Gen. Physiol.* **43** (1960), 867–896.
21. R. A. Gardner, On the structure of the spectra of periodic travelling waves, *J. Math. Pures Appl.* **72** (1993), 415–439.
22. J. Guckenheimer and P. Holmes, *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, Appl. Math. Sci., Vol. 42, Springer, New York, 1983.

23. J. Guckenheimer and C. Kuehn, Homoclinic orbits of the FitzHugh-Nagumo equation: the singular-limit, *Discrete Contin. Dyn. Syst. Ser. S* **2** (2009), 851–872.
24. J. Guckenheimer and C. Kuehn, Homoclinic orbits of the FitzHugh-Nagumo equation: bifurcations in the full system, *SIAM J. Appl. Dyn. Syst.* **9** (2010), 138–153.
25. S. Hastings, The existence of periodic solutions to Nagumo’s equation, *Quart. J. Math. Oxford* **3** (1974), 369–378.
26. X. He, K. Lam, Y. Lou, and W.-M. Ni, Dynamics of a consumer-resource reaction-diffusion model, *J. Math. Biol.* **78** (2019), 1605–1636.
27. D. Henry, *Geometric Theory of Semilinear Parabolic Equations*, Lecture Notes in Math., Vol. **840**, Springer, Berlin, 1981.
28. P. D. Hislop and I. M. Sigal, *Introduction to Spectral Theory with Applications to Schrödinger Operators*, Vol. **113**, Appl. Math. Sci., Springer-Verlag, New York, 1996.
29. A. L. Hodgkin and A.F. Huxley, A quantitative description of membrane current and its application to conduction and excitation in nerve, *J. Physiol. London* **117** (1952), 500–544.
30. C. K. R. T. Jones, Stability of the travelling wave solution of the FitzHugh-Nagumo system, *Trans. Amer. Math. Soc.* **286** (1984), 431–469.
31. C. K. R. T. Jones, N. Kopell, and R. Langer, Construction of the FitzHugh-Nagumo pulse using differential forms, *Patterns and Dynamics in Reactive Media* (Minneapolis, MN, 1989), 101–115, IMA Vol. Math. Appl., **37**, Springer, New York, 1991.
32. T. Kapitula and K. Promislow, *Spectral and Dynamical Stability of Nonlinear Waves*, Appl. Math. Sci., Vol. 185, Springer, New York, 2013.
33. T. Kato, *Perturbation Theory for Linear Operators*, Reprint of the 1980 edition, Classics in Mathematics, Springer-Verlag, Berlin, 1995.
34. J. Keener and J. Sneyd, *Mathematical Physiology*, Int. Appl. Math., Vol. 8, Springer-Verlag, New York, 1998.
35. R. Kollár, B. Deconinck, and O. Trichtchenko, Direct characterization of spectral stability of small-amplitude periodic waves in scalar Hamiltonian problems via dispersion relation, *SIAM J. Math. Anal.* **51** (2019), 3145–3169.
36. Y. Kuznetsov, *Elements of Applied Bifurcation Theory*, Appl. Math. Sci., Vol. 112, Springer-Verlag, New York, 1998.
37. X.-B. Lin, Using Melnikov’s method to solve Shilnikov’s problems, *Proc. R. Soc. Edinburgh A* **116** (1990), 295–325.
38. J. Llibre, D. Novaes, and M. Teixeira, Higher order averaging theory for finding periodic solutions via Brouwer degree, *Nonlinearity* **27** (2014), 563–583.
39. K. Maginu, Existence and stability of periodic travelling wave solutions to Nagumo’s nerve equation, *J. Math. Biol.* **10** (1980), 133–153.
40. J. Nagumo, S. Arimoto, and S. Yoshizawa, An active pulse transmission line simulating nerve axon, *Proc. IRE.* **50** (1962), 2061–2070.
41. J. A. Sanders, F. Verhulst, and J. Murdock, *Averaging Methods in Nonlinear Dynamical Systems*, Second edition, Appl. Math. Sci., Vol. 59, Springer, New York, 2007.
42. B. Sandstede and A. Scheel, On the stability of periodic travelling waves with large spatial period, *J. Differential Equations* **172** (2001), 134–188.
43. B. Sandstede, Stability of travelling waves, in *Handbook of Dynamical Systems*, Vol. 2, North-Holland, Amsterdam, 2002.

44. W. Schouten-Straatman and H. Hupkes, Traveling waves for spatially discrete systems of FitzHugh-Nagumo type with periodic coefficients, *SIAM J. Math. Anal.* **51** (2019), 3492–3532.
45. C. Soto-Treviño, A geometric method for periodic orbits in singularly-perturbed systems, *Multiple-Time-Scale Dynamical Systems*, IMA Volumes in Mathematics and its Applications, Vol. 122, ed. C. K. R. T. Jones and A Khibnik Springer, New York, pp. 141–202, 2001.
46. O. Trichtchenko, B. Deconinck, and R. Kollár, Stability of periodic traveling wave solutions to the Kawahara equation, *SIAM J. Appl. Dyn. Syst.* **17** (2018), 2761–2783.
47. J. Tsai, W. Zhang, V. Kirk, and J. Sneyd, Traveling waves in a simplified model of calcium dynamics, *SIAM J. Appl. Dyn. Syst.* **11** (2012), 1149–1199.
48. B. Zhang, K. Alex, M. L. Keenan, Z. Lu, L. R. Arrix, W.-M. Ni, D. L. DeAngelis, and J. D. Van Dyken, Carrying capacity in a heterogeneous environment with habitat connectivity, *Ecol. Lett.* **20** (2017), 1118–1128.
49. W. Zhang, J. Sneyd, and J. Tsai, Curvature dependence of propagating velocity for a simplified calcium model, *SIAM J. Appl. Math.* **74** (2014), 1442–1462.