

The anti-vaxx movement and epidemic spreading in the era of social networks: nonmonotonic effects, bistability and network segregation

Marcelo A. Pires^{1,*}, Andre L. Oestereich^{2,†}, Nuno Crokidakis^{3,‡} and Sílvia M. Duarte Queirós^{1,4,5,§}

¹*Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro/RJ, Brazil*

²*Unidade Central De Educação Faem Faculdade, Itapiranga/SC, Brazil*

³*Instituto de Física, Universidade Federal Fluminense, Niterói/RJ, Brazil*

⁴*National Institute of Science and Technology for Complex Systems, Brazil*

⁵*i3N, Universidade de Aveiro, Campus de Santiago, 3810-193 Aveiro, Portugal*

(Dated: February 28, 2022)

In this work, we address a multicoupled dynamics on complex networks with tunable structural segregation. Specifically, we work at a networked epidemic spreading under a vaccination campaign with agents in favor and against the vaccine. Our results show that such coupled dynamics exhibit a myriad of phenomena such as nonequilibrium transitions accompanied by bistability. Besides we observe the emergence of an intermediate optimal segregation level where the community structure can favor the negative opinions about vaccination but counterintuitively hinders - rather than enhance - the global disease spreading. Thus, our results point out that vaccination campaigns should avoid policies that end up segregating excessively anti-vaccine groups.

I. INTRODUCTION

In such a complex contemporary society where elements – people and events – influence one another and feedback at different scales [1], the application of tools set forth in Statistical Physics in order to cope with collective phenomena has gained prominence in other areas such as Biology and Medicine, Social Sciences, and Humanities, which have put quantitative tools in the methodologies they apply [2–7].

Accordingly, phenomena in which there is a change in the collective behavior displayed by a social system have turn into an appropriate field for the application of such techniques [8–10]; among the several different instances we can find important contributions within the spreading of epidemics as well as opinion dynamics (see eg [11] and Sec. II). In spite of the fact that the two subject-matters are not related at first, the dissemination of a causal relationship between neurological disorders and vaccinations [12] has prompted an urban myth that ultimately has jeopardized the elimination of the disease in countries with a very high Human Development Index as the USA [13].

As the world faces a pandemic the quash of which has been pointed to a worldwide vaccination program, the interplay between the spreading of opinion — particularly the diffusion of ‘anti-vaxxers’ ideas — and the dissemination of a contagious disease is a natural focus of attention. Therefore, assuming a complex network of agents we implement paradigmatic models of epidemic and opinion dynamics in order to understand to what extent the existence of a strong anti-vaccine population can imply in the proliferation of the respective disease.

The remaining of this manuscript is organised as follows: in Sec. II, we establish the state-of-the-art of the problem; in Sec. III, we introduce our model for the combined dynamics of opinion and contagion; in Sec. IV, we discuss the results for the model; and in Sec. V, we present our final observations on the work and future perspectives about it.

II. LITERATURE REVIEW

Modular networks [14] are generated by an algorithm that leads to networks with an architecture of communities. A given node in each community can be connected to nodes of the same community (intracommunity links) and/or to nodes of the other community (intercommunity links).

The impact of network modularity in spreading processes has been investigated in recent years. Since the work of Ref. [14], a series of works were published regarding the subject of optimal network modularity; therein, the authors showed that modular structure may have counterintuitive effects on information diffusion. Indeed, it was discussed that the presence of strong communities in modular networks can facilitate global diffusion by enhancing local, intracommunity spreading.

Still in relation to modular networks, it was recently found that an optimal community structure that maximizes spreading dynamics which can pave the way to rich phase diagrams with exhibiting first-order phase transitions [15]. Within the same context, the authors in [16] discussed the impact of social reinforcement in information diffusion. They also found optimal multi-community network modularity for information diffusion, i.e., depending on the range of the parameters the multi-community structure can facilitate information diffusion instead of hindering it.

Regarding biological systems, it was recently found there is a nonlinear relation between modularity and

* piresma@cbpf.br

† andreoestereich@uceff.edu.br

‡ nuno@mail.if.uff.br

§ sdqueiro@cbpf.br

global efficiency in animal networks, with the latter peaking at intermediate values of the former [17]. In addition, in neural networks there exists an optimal modularity for memory performance, where a balance between local cohesion and global connectivity is established, allowing optimally modular networks to remember longer [18].

The authors in [19] studied the importance of close and ordinary social contacts in promoting large-scale contagion and found an optimal fraction of ordinary contacts for outbreaks at a global scale. With respect to correlations in complex networks, it was found that constraining the mean degree and the fraction of initially informed nodes, the optimal structure can be assortative (modular), core-periphery or even disassortative [20]. Other recent works leading with optimal modularity in networks can be found in [21, 22].

In a recent work [23], it was proposed a model of disease spreading in a structural modular complex network and studied how the number of bridge nodes n that connect communities affects disease spreading. It was verified that near the critical point as n increases, the disease reaches most of the communities, but each community has only a small fraction of recovered nodes.

Disease information can spark strong emotions like fear — or even panic — that would affect behavior during an epidemic. The authors in [24] considered an agent-based model that assumes that agents can obtain a complete picture of the epidemic via information from local daily contacts or global news coverage. Those results helped conclude that such model can be used to mimic real-world epidemic situations and explain disease transmission, behavior changes, and distribution of prevalence panic. Game theory was also considered to reproduce the decision-making process of individuals during the evolution of a disease. In [25] a spatial evolutionary game was coupled to a SIR model, and the results showed that protective behaviors decrease the numbers of infected individuals and delay the peak time of infection. The study also concluded that increased numbers of risk-averse individuals and preemptive actions can more effectively mitigate disease transmission; however, changes in human behavior require a high social cost (such as avoidance of crowded places leading to absences in schools, workplaces, or other public places).

A recent work considered a coupled behavior-change and infection in a structured population characterized by homophily and outgroup aversion [26]. It was found that homophily can either increase or decrease the final size of the epidemic depending on its relative strength in the two groups. In addition, homophily and outgroup aversion can also produce a ‘second wave’ in the first group that follows the peak of the epidemic in the second group.

Models of opinion dynamics were applied in the context of opinions about vaccination (pro versus anti-vaccine) without coupling an epidemic process [27]. Later, kinetic opinion dynamics were coupled to classical epidemic models in order to study the feedback among risk perception, opinions about vaccination, and the disease spread-

ing. In [28] it was found that the engagement of the pro-vaccine individuals can be crucial for stopping the epidemic spreading. On the other hand, the work [29] found counterintuitive outcomes like the fact that an increment in the initial fraction of the population that is pro-vaccine can lead to smaller epidemic outbreaks in the short term, but it also contributes to the survival of the chain of infections in the long term.

Recently, the anti-vaccine sentiment was treated as a cultural pathogen. The authors in [30] modeled it as a ‘infection’ dynamics. The authors showed that interventions to increase vaccination can potentially target any of three types of transitions - decreasing sentiment transmission to undecided individuals, increasing pro-vaccine decisions among undecided individuals, or increasing sentiment switching among anti-vaccine individuals.

We previously cited anti-vaccine opinions, thus it is important to mention some recent discussion about the global anti-vaccine movement. Since the online discussions dominate the social interactions in our modern world, the propagation of such anti-vaccine opinions is growing fast. A recent report noted that 31 million people follow anti-vaccine groups on Facebook, with 17 million people subscribing to similar accounts on YouTube [31]. The authors in [32] recently pointed that if the current trends continue, anti-vaccine views will dominate online discussion in 10 years. The importance of anti-vaccine movement is fundamental for the evolution of COVID-19 outbreak. Indeed, the authors in [33] called attention to the fact that it is a key point to qualitatively assess how the administration of a vaccine could affect the COVID-19 outbreak, taking into account of the behavioral changes of individuals in response to the information available on the status of the disease in the community. According to a study published in August 2020, nearly one in four adults would not get a vaccine for COVID-19 [34] and in some countries, more than half of the population would not get it, including Poland and France [35]. In September 2020, it was verified that only 42 percent of Americans said *yes* to receiving a future COVID-19 vaccine, across all political sides. It means that even in a best-case scenario where a future high performing vaccine is 95% effective in an individual, it would only impact $42 \times 95 \approx 40\%$ of the population, which is way below predicted thresholds for herd immunity [36].

III. MODEL

A. I: Opinion dynamics

Based on [29] (and the original model [37]), we consider an agent-based dynamics in which the opinion about vaccination, $o_i \in [-1, 1]$, of each agent, i , evolves with

$$o_i(t+1) = o_i(t) + \epsilon o_j(t) + w I_{neig(i)}(t) \quad (1)$$




FIG. 1. Examples of modular networks with $N = 100$, $\langle k \rangle = 10$ for different values of μ . In these examples we can see the strengthening of the community structure for lower values of μ .

A negative (positive) value of o_i represents an individual i supporting anti-vaccine (pro-vaccine) opinion. Equation (1) takes into account the agent's opinion, $o_i(t+1)$, depends on multiple factors: (i) his previous opinion $o_i(t)$; (ii) the peer pressure exerted by a randomly selected neighbor, j , modulated by a stochastic heterogeneity ϵ , uniformly distributed in the interval $[0, 1]$; (iii) the proportion of infected neighbors, $I_{neig(i)}(t)$, modulated by a risk perception parameter, w .

The opinion dynamics regarding the vaccination campaign is coupled with the epidemic dynamics, due to the factor $I_{neig(i)}(t)$ in Eq. (1).

B. II: epidemics-vaccination dynamics

Based on [28, 29] (and references therein), we define the transitions among the epidemic compartments as follows:

- $S \xrightarrow{\gamma} R$: a Susceptible agent i becomes Vaccinated

with probability γ_i ;

- $S \xrightarrow{(1-\gamma_i)\lambda} I$: a Susceptible agent i becomes Infected with probability $(1-\gamma_i)\lambda$ if he is in contact with an Infected agent;
- $I \xrightarrow{\alpha} S$: an Infected agent i recovers with probability α ;
- $R \xrightarrow{\phi} S$: a immune agent i becomes Susceptible again with the resusceptibility probability ϕ . Based on [38–42] we assume that Vaccinated and Recovered agents are in the same compartment.

The vaccination probability γ_i of an agent i is proportional to his opinion about vaccination $-1 \leq o_i \leq 1$:

$$\gamma_i(t) = \frac{1 + o_i(t)}{2} \in [0, 1] \quad (2)$$

C. Community structure

Based on [43] and related literature, we start by picking the first $N_1 = N/2$ of the N nodes and attaching them to the community 1, and assigning the other $N_2 = N - N_1$ nodes to community 2. We then proceed by randomly assigning $(1-\mu)M$ connections among pairs of nodes from the same community and μM connections are randomly distributed among pairs of nodes that belong to distinct communities.

The parameter μ regulates the community strength: large values of μ means more ties between the two communities consequently a weaker community organization.

D. Initial condition

We consider that the community 1 holds a positive stance on vaccination, whereas the community 2 holds a negative opinion about that. We also assume the chain of infections starts in the community 2, because this case is more important since $o_i < 0$ leads to a low propensity for the agents to get vaccinated. If the epidemic would start in the community 1, the positive opinions $o_i > 0$ would induce a relatively high probability for an agent to get vaccinated which ends up disrupting the chain of contagions.

Let $U(a, b)$ be a single random value from a uniform distribution in the range $[a, b]$.

At $t = 0$, we set:

- For i in $0 \dots N/2 - 1$: (community 1: $o_i > 0$; 0% infected)
 - $o_i \sim U(0, 1)$
 - status(i) = S
- For i in $N/2 \dots N - 1$: (community 2: $o_i < 0$; 1% of infected)




FIG. 2. Steady-state for the spreading measure I_i and collective opinion m_i for each community $i = \{1, 2\}$. Symbols are the steady-state outcome for each sample, i.e., each symbol is the result from each Monte Carlo realization. Results for $\mu = 0.1$. For this high level of segregation, each community ends up preserving the sign of its initial opinion. Besides, the chain of contagion starts in the community 2 and cannot become permanent in the community 1.

- $o_i \sim U(-1, 0)$
- $\text{status}(i) = \text{S}$ with probability 0.99
- $\text{status}(i) = \text{I}$ with probability 0.01

IV. RESULTS AND DISCUSSION

In this section, we present our results come from Monte Carlo simulations of networks with $N = 10^4$ nodes and $k=20$. In all simulations, we set $\alpha = 0.1$ and $\phi = 0.01$, without losing generality. In Figs. 2-5, we show the steady-state density of infected agents in the community u , I_u . We also depict the behavior of the stationary opin-

ion in the community u , m_u . In turn, I_{tot} and m_{tot} refer to the global proportion of infected individuals and global mean opinion, respectively.

The results in Fig. 2 show that in the community 2 — the seed community — there is a transition from the absorbing phase (extinction of the epidemic) to the epidemic survival phase. In the community 1, there is no survival of the chain of infections in the long term. In this setting with $\mu = 0.1$ (weak modular structure) the seed community remains with the negative opinion about vaccination, which weakens the vaccination campaign and thus facilitates the local permanence of the disease. Similarly, there is a persistence of the initial opinion in the community 1, which in this case is pro-vaccine and there-




FIG. 3. Steady-state for the spreading measure I_i and collective opinion m_i for each community $i = \{1, 2\}$. Symbols are the steady-state outcome for each sample, ie, each symbol is the result from each Monte Carlo realization. Results for $\mu = 0.2$. For this intermediate level of segregation, there is the possibility for a switch of opinion in the community 2 (seed). The epidemic spreading does not survive at the global and local levels.

fore favors the vaccine uptake that makes the epidemic spreading unsustainable. This means that a low number of intercommunity ties hinders the change in the community stance over vaccination; that creates a strong distinction in the epidemic spread between both communities with community 1 being unfavorable to epidemic spreading since $m_1 > 0$, and community 2 being favorable since $m_2 < 0$.

In Fig. 3, it is notable that an intermediate community strength leads to the elimination of the epidemic transmission in both communities even when there is a dominance of the negative opinion about vaccination in the community 2. The epidemic contagion spreading is halted in the community 2, even though the agents have

a negative opinion about the vaccination, due to the intermediate number of bridges, $\mu = 0.2$, to the other community. These bridges are just strong enough to drain the infected agents of the community 2, but not strong enough to change its average opinion.

In Fig. 4, with $\mu = 0.3$ there is a high number of intercommunity links. This additional connectivity between communities weakens the initial epidemic spreading in the community 2, but it is sufficient to introduce the possibility of a wide opinion change in the community 1. The opinion change in the community 1 facilitates the epidemic spreading in that community. This effect is limited because we can see for high infection probabilities $\lambda > 0.8$ the epidemic spread vanishes. So, we have




FIG. 4. Steady-state for the spreading measure I_i and collective opinion m_i for each community $i = \{1, 2\}$. Symbols are the steady-state outcome for each sample, i.e., each symbol is the result from each Monte Carlo realization. Results for $\mu = 0.3$. For this low level of segregation, there is the possibility for a switch of opinion in both communities. The epidemic dynamics can survive if the contagion is not too aggressive (intermediate values of λ).

a counterintuitive effect, because for higher transmissibility the epidemic spread vanishes. The reason behind this is the risk perception, wI in Eq. (1), which promotes vaccination, so higher transmissibility leads to a bigger outbreak that in turn results in better opinions about vaccination which ends up stopping the epidemic outbreak.

The emergence of an intermediate range of μ that blocks the local and global epidemic spreading is visible in Fig. 5. Regarding the opinion dynamics, an initial increase in μ leads to a decrease in m_1 and an increase in m_2 , that is the collective opinions tend to be less extremist for an initial rise in the amount of inter-communities routes. Then a further increase in μ promotes a sudden

rise in m_1 and m_2 which means a speed up in the switch of opinions in the community 2. A further rise in μ leads to a bistable behavior in both communities.

This intermediary range of inter-community connectivity that promotes a minimal epidemic spreading seems to also come from a perceived increment in the probability of an infected individual having a vaccinated neighbor. The increment of the bridges between communities the initially infected agents have a bigger probability of having a neighbor that was vaccinated because initially most of the infected people are in the community 2 and most agents with a positive opinion about vaccination are in the community 1. This effect does not persist for higher values of μ because then both communities tend




FIG. 5. Steady-state for the spreading measure I_i and collective opinion m_i for each community $i = \{1, 2\}$. Symbols are the steady-state outcome for each sample, i.e., each symbol is the result from each Monte Carlo realization. Results for $w = 0.1$ and $\lambda = 0.8$.

to adopt the same average opinion about vaccination and this opinion can, in some cases, be negative. A negative global opinion about vaccines does not guarantee that the epidemic spread will persist, as can be seen in some cases for $\mu \approx 0.23$ where all samples had no infected individual but some of them had negative opinions about the vaccination. This can occur due to the fact that the number of infected agents can become zero before the negative global consensus about vaccines is reached.

While in Figs. 2-3 there is a single stable steady-state (either extinction or persistence), Fig. 4 display bistable solutions depending on the randomness 'embedded' in the dynamics. Moreover, the results in Fig. 2 suggest that the absorbing-active epidemic transition is continuous for strong communities (such as $\mu = 0.1$) whereas

the results shown in Fig. 4 signalize this extinction-persistence epidemic transition is discontinuous for weak communities (such as $\mu = 0.3$). Therefore, the structural factors present in the modular networks can induce the emergence of bistability in the epidemic-vaccination-opinion dynamics as well as a change in the nature of the absorbing-active transitions.

An overall look into Figs. 2-5 reveals that a sudden transition can emerge from structural factors (increasing μ) or epidemiological factors (increasing λ). The transitions from the Disease-Free phase to the active phase and vice-versa (epidemic resurgence) highlight the non-monotonic behavior of the full dynamics with the transmissibility λ .

Comparing with other works, we see that while in [14]

there is an optimal modularity for enhancing information spreading, here there is an optimal modularity for hindering epidemic spreading.

V. FINAL REMARKS

In [44] was shown with a binary opinion dynamics that the spread of opinions against vaccination is one of the potential responsible for the large outbreaks of vaccine-preventable diseases in many high-income countries. Here we show with continuous opinion dynamics coupled to a networked SIRS model that the spectrum of scenarios arising from the competition of pro- vs anti-vaccine views during an epidemic spreading is highly complex.

The several outcomes shown in Figs.2-5 point out that our model produces a diverse phenomenology where the social and biological scenarios exhibit a nonmonotonic dependence with spreading rate λ . From the perspective of the dynamical systems, our results provide a new mechanism for bistability in a biological-social setting. From a practical point of view, our work offers new perspectives for the development of novel strategies for halting epidemic spreading based on tuning the modularity to an optimal degree.

Some pro-vaccine strategies can have as side effects the

segregation between individuals with conflicting views about the vaccines. In this regard, it was shown recently [45] that segregation of anti-vaxxers can potentially extend the duration of an epidemic spreading. In [46] it was found that an increase in the contact between vaccine refusers and the rest of the society can lead to a scenario where vaccination alone may not be able to prevent an outbreak. Here we show that too much or too low segregation of anti-vaxxers favors the chain of contagion, but an intermediate level of segregation disfavor the epidemic spreading. Therefore, our results indicate that vaccination campaigns should avoid strategies that have as a side effect too much segregation of anti-vaccine groups.

Our work produces a thought-provoking analogy. In a small-world architecture, there is an intermediate number of long-range bridges that lead the full network to have unusual properties such as high clustering and low path lengths. Here, a structure with an intermediate number of inter-community ties leads the dynamics in the full network to produce an interesting outcome, namely the suppression of the epidemics.

In future works, it would be valuable to consider more sophisticated network architectures. Besides, it will be worthwhile to consider the interplay between several sources of heterogeneity in agent's bias, namely plurality and polarization [47].

-
- [1] G. Parisi, Complex systems: a physicist's viewpoint, *Physica A* **263**, 557 (1999).
- [2] S. Thurner, R. Hanel, and P. Klimek, *Introduction to the Theory of Complex Systems* (Oxford, University Press, Oxford, 2018).
- [3] D. Stauffer, S. M. M. De Oliveira, P. M. C. De Oliveira, and J. S. de Sá Martins, *Biology, sociology, geology by computational physicists* (Elsevier, 2006).
- [4] C. Castellano, S. Fortunato, and V. Loreto, Statistical physics of social dynamics, *Reviews of modern physics* **81**, 591 (2009).
- [5] P. M. C. de Oliveira and D. Stauffer, *Evolution, Money, War, and Computers: Non-Traditional Applications of Computational Statistical Physics*, Vol. 34 (Springer Science & Business Media, 2013).
- [6] S. Galam, Sociophysics: A review of galam models, *International Journal of Modern Physics C* **19**, 409 (2008).
- [7] P. Sen and B. K. Chakrabarti, *Sociophysics: an introduction* (Oxford University Press, 2014).
- [8] R. Pastor-Satorras, P. Van Mieghem, and A. Vespignani, Epidemic processes in complex networks, *Rev. Mod. Phys.* **87**, 925 (2015).
- [9] G. F. de Arruda, F. A. Rodrigues, and Y. Moreno, Fundamentals of spreading processes in single and multilayer complex networks, *Phys. Rep.* **756**, 1 (2018).
- [10] W. Wang, M. Tang, L. A. Braunstein, and E. H. Stanley, Unification of theoretical approaches for epidemic spreading on complex networks, *Rep. Prog. Phys.* **80**, 036603 (2017).
- [11] W. Wang, Q.-H. Liu, J. Liang, Y. Hu, and T. Zhou, Coevolution spreading in complex networks, *Physics Reports* **10.1016/j.physrep.2019.07.001** (2019).
- [12] R. Gasparini, D. Panatto, P. L. Lai, and D. Amicizia, The "urban myth" of the association between neurological disorders and vaccinations, *J. Prev. Med. Hyg.* **56**, e1 (2015).
- [13] K. A. Feemster and C. Szipszky, Resurgence of measles in the united states: how did we get here?, *Curr. Opin. Pediatr.* **32**, 139 (2020).
- [14] A. Nematzadeh, E. Ferrara, A. Flammini, and Y.-Y. Ahn, Optimal network modularity for information diffusion, *Phys. Rev. Lett.* **113**, 088701 (2014).
- [15] Z. Su, W. Wang, L. Li, H. E. Stanley, and L. A. Braunstein, Optimal community structure for social contagions, *New Journal of Physics* **20**, 053053 (2018).
- [16] J. Wu, R. Du, Y. Zheng, and D. Liu, Optimal multi-community network modularity for information diffusion, *International Journal of Modern Physics C* **27**, 1650092 (2016).
- [17] V. Romano, M. Shen, J. Pansanel, A. J. MacIntosh, and C. Sueur, Social transmission in networks: global efficiency peaks with intermediate levels of modularity, *Behavioral ecology and sociobiology* **72**, 154 (2018).
- [18] N. Rodriguez, E. Izquierdo, and Y.-Y. Ahn, Optimal modularity and memory capacity of neural reservoirs, *Network Neuroscience* **3**, 551 (2019).
- [19] P.-B. Cui, W. Wang, S.-M. Cai, T. Zhou, Y.-C. Lai, *et al.*, Close and ordinary social contacts: How important are they in promoting large-scale contagion?, *Physical Review E* **98**, 052311 (2018).

- [20] G. Curato and F. Lillo, Optimal information diffusion in stochastic block models, *Physical Review E* **94**, 032310 (2016).
- [21] A. Nematzadeh, N. Rodriguez, A. Flammini, and Y.-Y. Ahn, Optimal modularity in complex contagion, in *Complex spreading phenomena in social systems* (Springer, 2018) pp. 97–107.
- [22] H. Peng, A. Nematzadeh, D. M. Romero, and E. Ferrara, Network modularity controls the speed of information diffusion, *Phys. Rev. E* **102**, 052316 (2020).
- [23] L. D. Valdez, L. A. Braunstein, and S. Havlin, Epidemic spreading on modular networks: The fear to declare a pandemic, *Physical Review E* **101**, 032309 (2020).
- [24] K. Bi, Y. Chen, S. Zhao, D. Ben-Arieh, and C.-H. J. Wu, Modeling learning and forgetting processes with the corresponding impacts on human behaviors in infectious disease epidemics, *Computers & Industrial Engineering* **129**, 563 (2019).
- [25] S. Zhao, Y. Kuang, C.-H. Wu, K. Bi, and D. Ben-Arieh, Risk perception and human behaviors in epidemics, *IIEE Transactions on Healthcare Systems Engineering* **8**, 315 (2018).
- [26] P. E. Smaldino and J. H. Jones, Coupled dynamics of behavior and disease contagion among antagonistic groups, *bioRxiv* (2020).
- [27] S. Galam, Public debates driven by incomplete scientific data: the cases of evolution theory, global warming and h1n1 pandemic influenza, *Physica A: Statistical Mechanics and its Applications* **389**, 3619 (2010).
- [28] M. A. Pires and N. Crokidakis, Dynamics of epidemic spreading with vaccination: impact of social pressure and engagement, *Physica A* **467**, 167 (2017).
- [29] M. A. Pires, A. L. Oestereich, and N. Crokidakis, Sudden transitions in coupled opinion and epidemic dynamics with vaccination, *J. Stat. Mech.* **2018**, 053407 (2018).
- [30] R. S. Mehta and N. A. Rosenberg, Modelling anti-vaccine sentiment as a cultural pathogen, *Evolutionary Human Sciences* **2** (2020).
- [31] T. Burki, The online anti-vaccine movement in the age of covid-19, *The Lancet Digital Health* **2**, e504 (2020).
- [32] N. F. Johnson, N. Velásquez, N. J. Restrepo, R. Leahy, N. Gabriel, S. El Oud, M. Zheng, P. Manrique, S. Wuchty, and Y. Lupu, The online competition between pro-and anti-vaccination views, *Nature* **582**, 230 (2020).
- [33] B. Buonomo, Effects of information-dependent vaccination behavior on coronavirus outbreak: insights from a siri model, *Ricerche di Matematica* **69**, 483 (2020).
- [34] N. Boyon and K. Silverstein, Three in four adults globally say they would get a vaccine for covid-19, *Ipsos, News & Pools* (2020).
- [35] R. P. Curiel and H. G. Ramírez, Vaccination strategies against covid-19 and the diffusion of anti-vaccination views (2020), [arXiv:2009.13674](https://arxiv.org/abs/2009.13674) [physics.soc-ph].
- [36] N. F. Johnson, N. Velásquez, R. Leahy, O. Jha, and Y. Lupu, Not sure? handling hesitancy of covid-19 vaccines (2020), [arXiv:2009.08413](https://arxiv.org/abs/2009.08413) [physics.soc-ph].
- [37] M. Lallouache, A. S. Chakrabarti, A. Chakraborti, and B. K. Chakrabarti, Opinion formation in kinetic exchange models: Spontaneous symmetry-breaking transition, *Phys. Rev. E* **82**, 056112 (2010).
- [38] G.-Z. Zeng and L.-S. Chen, Complexity and asymptotical behavior of a sirs epidemic model with proportional impulsive vaccination, *Advances in Complex Systems* **8**, 419 (2005).
- [39] F. Rao, P. S. Mandal, and Y. Kang, Complicated epidemics of an sirs model with a generalized incidence under preventive vaccination and treatment controls, *Applied Mathematical Modelling* **67**, 38 (2019).
- [40] I. Moneim and D. Greenhalgh, Threshold and stability results for an sirs epidemic model with a general periodic vaccination strategy, *Journal of biological systems* **13**, 131 (2005).
- [41] A. Lahrouz, L. Omari, D. Kiouach, and A. Belmaâti, Complete global stability for an sirs epidemic model with generalized non-linear incidence and vaccination, *Applied Mathematics and Computation* **218**, 6519 (2012).
- [42] P. Doutor, P. Rodrigues, M. do Céu Soares, and F. A. Chalub, Optimal vaccination strategies and rational behaviour in seasonal epidemics, *Journal of mathematical biology* **73**, 1437 (2016).
- [43] A. L. Oestereich, M. A. Pires, and N. Crokidakis, Three-state opinion dynamics in modular networks, *Phys. Rev. E* **100**, 032312 (2019).
- [44] M. Salathé and S. Bonhoeffer, The effect of opinion clustering on disease outbreaks, *Journal of The Royal Society Interface* **5**, 1505 (2008).
- [45] M. Bizzarri, F. Panebianco, and P. Pin, Is segregating anti-vaxxers a good idea?, *arXiv preprint arXiv:2007.08523* (2020).
- [46] C. M. Saad-Roy, C. E. Wagner, R. E. Baker, S. E. Morris, J. Farrar, A. L. Graham, S. A. Levin, M. J. Mina, C. J. E. Metcalf, and B. T. Grenfell, Immune life history, vaccination, and the dynamics of sars-cov-2 over the next 5 years, *Science* **370**, 811 (2020).
- [47] A. L. Oestereich, M. A. Pires, S. D. Queirós, and N. Crokidakis, Hysteresis and disorder-induced order in continuous kinetic-like opinion dynamics in complex networks, *Chaos, Solitons & Fractals* **137**, 109893 (2020).