

# On a class of integrable Hamiltonian equations in 2+1 dimensions

B. Gormley<sup>1</sup>, E.V. Ferapontov<sup>1,2</sup>, V.S. Novikov<sup>1</sup>

<sup>1</sup>Department of Mathematical Sciences  
Loughborough University  
Loughborough, Leicestershire LE11 3TU  
United Kingdom

<sup>2</sup>Institute of Mathematics, Ufa Federal Research Centre,  
Russian Academy of Sciences, 112, Chernyshevsky Street,  
Ufa 450077, Russia

e-mails:

B.Gormley@lboro.ac.uk  
E.V.Ferapontov@lboro.ac.uk  
V.S.Novikov@lboro.ac.uk

## Abstract

We classify integrable Hamiltonian equations of the form

$$u_t = \partial_x \left( \frac{\delta H}{\delta u} \right), \quad H = \int h(u, w) \, dx dy,$$

where the Hamiltonian density  $h(u, w)$  is a function of two variables: dependent variable  $u$  and the non-locality  $w = \partial_x^{-1} \partial_y u$ . Based on the method of hydrodynamic reductions, the integrability conditions are derived (in the form of an involutive PDE system for the Hamiltonian density  $h$ ). We show that the generic integrable density is expressed in terms of the Weierstrass  $\sigma$ -function:  $h(u, w) = \sigma(u)e^w$ . Dispersionless Lax pairs, commuting flows and dispersive deformations of the resulting equations are also discussed.

MSC: 35Q51, 37K10.

**Keywords:** Hamiltonian PDEs, hydrodynamic reductions, Einstein-Weyl geometry, dispersionless Lax pairs, commuting flows, dispersive deformations, Weierstrass elliptic functions.

*To Allan Fordy on the occasion of his 70th birthday*

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Formulation of the problem . . . . .	2
1.2	Equivalent approaches to dispersionless integrability . . . . .	3
1.3	Summary of the main results . . . . .	5
<b>2</b>	<b>Proofs</b>	<b>6</b>
2.1	The method of hydrodynamic reductions: proof of Theorem 1 . . . . .	6
2.2	Canonical forms of integrable densities: proof of Theorem 2 . . . . .	7
2.3	Dispersionless Lax pairs . . . . .	9
2.4	Commuting flows: proof of Theorem 3 . . . . .	11
2.5	Commuting flows and dispersionless Lax pairs . . . . .	14
<b>3</b>	<b>Dispersive deformations</b>	<b>17</b>
<b>4</b>	<b>Appendix: dispersionless Lax pair for <math>h = \sigma(u)e^w</math></b>	<b>18</b>

## 1 Introduction

### 1.1 Formulation of the problem

In this paper we investigate Hamiltonian systems of the form

$$u_t = \partial_x \left( \frac{\delta H}{\delta u} \right), \quad H = \int h(u, w) \, dx dy. \quad (1)$$

Here  $\partial_x$  is the Hamiltonian operator, and the Hamiltonian density  $h(u, w)$  depends on  $u$  and the nonlocal variable  $w = \partial_x^{-1} \partial_y u$  (equivalently,  $w_x = u_y$ ). Since  $\frac{\delta H}{\delta u} = h_u + \partial_x^{-1} \partial_y (h_w)$  we can rewrite equation (1) in the two-component first-order quasilinear form:

$$u_t = (h_u)_x + (h_w)_y, \quad w_x = u_y. \quad (2)$$

Familiar examples of this type include the dispersionless KP equation ( $h = \frac{1}{2}w^2 + \frac{1}{6}u^3$ ) and the dispersionless Toda (Boyer-Finley) equation ( $h = e^w$ ). Our main goal is to classify integrable systems within class (2) and to construct their dispersionless Lax pairs, commuting flows and dispersive deformations. Before stating our main results, let us begin with a brief description of the existing approaches to dispersionless integrability in 2+1 dimensions.

## 1.2 Equivalent approaches to dispersionless integrability

Here we summarise three existing approaches to integrability of equations of type (2), namely, the method of hydrodynamic reductions, the geometric approach based on integrable conformal geometry (Einstein-Weyl geometry), and the method of dispersionless Lax pairs. Based on seemingly different ideas, these approaches lead to equivalent integrability conditions/classification results [13].

**The method of hydrodynamic reductions**, see e.g. [16, 9], consists of seeking multi-phase solutions to system (2) in the form

$$u = u(R^1, R^2, \dots, R^n), \quad w = w(R^1, R^2, \dots, R^n) \quad (3)$$

where the phases  $R^i(x, y, t)$  (also known as Riemann invariants; note that their number  $n$  can be arbitrary) satisfy a pair of commuting hydrodynamic-type systems:

$$R_y^i = \mu^i(R)R_x^i, \quad R_t^i = \lambda^i(R)R_x^i; \quad (4)$$

we recall that the commutativity conditions are equivalent to the following constraints for the characteristic speeds  $\mu^i, \lambda^i$  [18, 19]:

$$\frac{\partial_j \mu^i}{\mu^j - \mu^i} = \frac{\partial_j \lambda^i}{\lambda^j - \lambda^i}, \quad (5)$$

$i \neq j$ ,  $\partial_j = \partial_{R^j}$ . Substituting ansatz (3) into (2) and using (4), (5) one obtains an overdetermined system for the unknowns  $u, w, \mu^i, \lambda^i$ , viewed as functions of  $R^1, \dots, R^n$  (the so-called generalised Gibbons-Tsarev system, or GT-system). System (2) is said to be integrable by the method of hydrodynamic reductions if it possesses ‘sufficiently many’ multi-phase solutions of type (3), in other words, if the corresponding GT-system is involutive. Note that the coefficients of GT-system will depend on the density  $h(u, w)$  and its partial derivatives. The requirement that GT-system is involutive imposes differential constraints for the Hamiltonian density  $h$ , the so-called integrability conditions. Details of this computation will be given in Section 2.1.

**Integrability via Einstein-Weyl geometry.** Let us first introduce a conformal structure defined by the characteristic variety of system (2). Given a  $2 \times 2$  quasilinear system

$$A(v)v_{x^1} + B(v)v_{x^2} + C(v)v_{x^3} = 0$$

where  $A, B, C$  are  $2 \times 2$  matrices depending on  $v = (u, w)^T$ , the characteristic equation of this system,  $\det(Ap_1 + Bp_2 + Cp_3) = 0$ , defines a conic  $g^{ij}p_i p_j = 0$ . This gives the characteristic conformal structure  $[g] = g_{ij}dx^i dx^j$  where  $g_{ij}$  is the inverse matrix of  $g^{ij}$ . For system (2) direct calculation gives

$$[g] = 4h_{ww}dxdt - dy^2 - 4h_{uw}dydt + 4(h_{ww}h_{uu} - h_{uw}^2)dt^2; \quad (6)$$

here we set  $(x^1, x^2, x^3) = (x, y, t)$ . Note that  $[g]$  depends upon a solution to the system (we will assume  $[g]$  to be non-degenerate, which is equivalent to the condition  $h_{ww} \neq 0$ ). It turns out that integrability of system (2) can be reformulated geometrically as the Einstein-Weyl property of the characteristic conformal structure  $[g]$ . We recall that Einstein-Weyl geometry is a triple  $(\mathbb{D}, [g], \omega)$  where  $[g]$  is a conformal structure,  $\mathbb{D}$  is a symmetric affine connection and  $\omega = \omega_k dx^k$  is a 1-form such that

$$\mathbb{D}_k g_{ij} = \omega_k g_{ij}, \quad R_{(ij)} = \Lambda g_{ij} \quad (7)$$

for some function  $\Lambda$  [2, 3]; here  $R_{(ij)}$  is the symmetrised Ricci tensor of  $\mathbb{D}$ . Note that the first part of equations (7), known as the Weyl equations, uniquely determines  $\mathbb{D}$  once  $[g]$  and  $\omega$  are specified. It was observed in [13] that for broad classes of dispersionless integrable systems (in particular, for systems of type (2)), the one-form  $\omega$  is given in terms of  $[g]$  by a universal explicit formula

$$\omega_k = 2g_{kj} \mathcal{D}_s g^{js} + \mathcal{D}_k \ln(\det g_{ij})$$

where  $\mathcal{D}_k$  denotes the total derivative with respect to  $x^k$ . Applied to  $[g]$  given by (6), this formula implies

$$\begin{aligned} \omega_1 &= 0, \\ \omega_2 &= \frac{2(h_{uuw}v_{xx} + 2h_{uww}v_{xy} + h_{www}v_{yy})}{h_{ww}}, \\ \omega_3 &= \frac{4(h_{uw}(h_{uuw}v_{xx} + 2h_{uww}v_{xy} + h_{www}v_{yy}))}{h_{ww}} - \frac{h_{ww}(h_{www}v_{xx} + 2h_{uuw}v_{xy} + h_{uww}v_{yy})}{h_{ww}}. \end{aligned} \quad (8)$$

To summarise, integrability of system (2) is equivalent to the Einstein-Weyl property of  $[g]$ ,  $\omega$  given by (6), (8) on every solution of system (2). Note that in 3D, Einstein-Weyl equations (7) are themselves integrable by the twistor construction [17], see also [8], and thus constitute ‘integrable conformal geometry’.

**Dispersionless Lax pair** of system (2) consist of two Hamilton-Jacobi type equations for an auxiliary function  $S$ ,

$$S_t = F(S_x, u, w), \quad S_y = G(S_x, u, w),$$

whose compatibility condition,  $S_{ty} = S_{yt}$ , is equivalent to system (2). Dispersionless Lax pairs were introduced in [20] as quasiclassical limits of Lax pairs of integrable soliton equations in 2+1D. It is known that the existence of a dispersionless Lax representation is equivalent to hydrodynamic/geometric integrability discussed above [10, 13]. We refer to Section 2.3 for dispersionless Lax pairs of integrable systems (2).

### 1.3 Summary of the main results

Our first result is the set of integrability conditions for the Hamiltonian density  $h$ .

**Theorem 1.** *The following conditions are equivalent:*

- (a) *System (2) is integrable by the method of hydrodynamic reductions;*
- (b) *Characteristic conformal structure  $[g]$  and covector  $\omega$  given by (6), (8) satisfy Einstein-Weyl equations (7) on every solution of system (2);*
- (c) *System (2) possesses a dispersionless Lax pair;*
- (d) *Hamiltonian density  $h(u, w)$  satisfies the following set of integrability conditions:*

$$\begin{aligned}
 h_{www}^2 - h_{ww}h_{www} &= 0, \\
 h_{uww}h_{www} - h_{ww}h_{uwww} &= 0, \\
 h_{uuw}h_{www} - h_{ww}h_{uuww} &= 0, \\
 h_{uuu}h_{www} - h_{ww}h_{uuuu} &= 0, \\
 -3h_{uuw}^2 + 4h_{uww}h_{uuu} - h_{ww}h_{uuuu} &= 0.
 \end{aligned} \tag{9}$$

Theorem 1 is proved in Section 2.1. The system of integrability conditions (9) is involutive, and modulo natural equivalence transformations its solutions can be reduced to one of the six canonical forms.

**Theorem 2.** *Solutions  $h(u, w)$  of system (9) can be reduced to one of the six canonical forms:*

$$\begin{aligned}
 h(u, w) &= \frac{1}{2}w^2 + \frac{1}{6}u^3, \\
 h(u, w) &= w^2 + u^2w - \frac{1}{4}u^4, \\
 h(u, w) &= uw^2 + \beta u^7, \\
 h(u, w) &= e^w, \\
 h(u, w) &= ue^w, \\
 h(u, w) &= \sigma(u; 0, g_3)e^w;
 \end{aligned}$$

here  $\beta$  and  $g_3$  are constants, and  $\sigma(u; g_2, g_3)$  denotes the Weierstrass sigma function.

Theorem 2 is proved in Section 2.2. Dispersionless Lax pairs for the corresponding systems (2) are constructed in Section 2.3.

It turns out that every integrable system (2) possesses a higher commuting flow of the form

$$\begin{aligned}
 u_\tau &= a(u, w, v)u_x + b(u, w, v)u_y + c(u, w, v)w_y + d(u, w, v)v_y, \\
 w_x &= u_y, \\
 v_x &= (p(u, w))_y,
 \end{aligned} \tag{10}$$

where  $\tau$  is the higher ‘time’, and  $v = \partial_x^{-1} \partial_y p(u, w)$  is an extra nonlocal variable (in contrast to the 1+1 dimensional case, higher commuting flows in 2+1 dimensions require higher nonlocalities). Remarkably, the structure of higher nonlocalities is uniquely determined by the original system (2), in particular, the function  $p(u, w)$  can be expressed in terms of  $h(u, w)$ :  $p = h_w$ . Furthermore, commuting flow (10) is automatically Hamiltonian.

**Theorem 3.** *Every integrable system (2) possesses a higher commuting flow (10) with the nonlocality  $v_x = (h_w)_y$ . Commuting flow (10) is Hamiltonian with the Hamiltonian density  $f(u, w, v)$  of the form*

$$f(u, w, v) = v h_w(u, w) + g(u, w),$$

where  $g(u, w)$  can be recovered from the compatible equations

$$\begin{aligned} g_{ww} &= 4h_{uw}h_{ww} + \alpha w h_{ww}, \\ g_{uuu} &= 8h_{uw}h_{uuu} + \alpha w h_{uuu}, \\ g_{uuw} &= 6h_{uw}h_{uuw} + 2h_{ww}h_{uuu} + \alpha w h_{uuw}. \end{aligned}$$

Here the constant  $\alpha$  is defined by the relation  $\alpha = 2h_{ww} \frac{\partial}{\partial w} \left( \frac{h_{uw}}{h_{ww}} \right)$  which follows from integrability conditions (9).

Theorem 3 is proved in Section 2.4. Dispersionless Lax pairs for commuting flows are constructed in Section 2.5.

## 2 Proofs

In this section we prove Theorems 1-3 and construct Lax pairs for integrable systems (2) and their commuting flows (10).

### 2.1 The method of hydrodynamic reductions: proof of Theorem 1

**Equivalences (a)  $\Leftrightarrow$  (b) and (a)  $\Leftrightarrow$  (c)** of Theorem 1 follow from the results of [13] and [10] which hold for general two-component systems of hydrodynamic type in 2+1 dimensions.

**Equivalence (a)  $\Leftrightarrow$  (d)** can be demonstrated as follows. Let us rewrite system (2) in the form

$$u_t = h_{uu}u_x + 2h_{uw}u_y + h_{ww}w_y, \quad w_x = u_y,$$

and substitute the ansatz  $u = u(R^1, R^2, \dots, R^n)$ ,  $w = w(R^1, R^2, \dots, R^n)$ . Using equations (4) and collecting coefficients at  $R_x^i$  we obtain  $\partial_i w = \mu^i \partial_i u$ , along with the dispersion relation  $\lambda^i = h_{uu} + 2h_{uw}\mu^i + h_{ww}(\mu^i)^2$ . Substituting the last formula into the commutativity conditions (5) we obtain

$$\partial_j \mu^i = \frac{h_{uuu} + h_{uuw}(\mu^j + 2\mu^i) + h_{uw}w(2\mu^i \mu^j + (\mu^i)^2) + h_{ww}w \mu^j (\mu^i)^2}{h_{ww}(\mu^j - \mu^i)} \partial_j u. \quad (11)$$

Finally, the compatibility condition  $\partial_i \partial_j w = \partial_j \partial_i w$  results in

$$\partial_i \partial_j u = \frac{2h_{uuu} + 3h_{uww}(\mu^j + \mu^i) + h_{uw}((\mu^i)^2 + 4\mu^i \mu^j + (\mu^j)^2) + h_{www}(\mu^j(\mu^i)^2 + \mu^i(\mu^j)^2)}{h_{ww}(\mu^j - \mu^i)^2} \partial_i u \partial_j u. \quad (12)$$

Equations (11), (12) constitute the corresponding GT-system. As one can see, it contains partial derivatives of the Hamiltonian density  $h$  in the coefficients. Verifying involutivity of GT-system amounts to checking the compatibility conditions  $\partial_k(\partial_j \mu^i) = \partial_j(\partial_k \mu^i)$  and  $\partial_k(\partial_i \partial_j u) = \partial_j(\partial_i \partial_k u)$ . Direct computation (performed in Mathematica) results in the integrability conditions (9) for  $h(u, w)$ . Note that without any loss of generality one can restrict to the case when the number of Riemann invariants  $R^i$  is equal to three, indeed, all compatibility conditions involve three distinct indices only. This finishes the proof of Theorem 1.

## 2.2 Canonical forms of integrable densities: proof of Theorem 2

We have five integrability conditions, namely

$$h_{www}^2 - h_{ww}h_{www} = 0, \quad (13)$$

$$h_{uww}h_{www} - h_{ww}h_{uww} = 0, \quad (14)$$

$$h_{uuw}h_{www} - h_{ww}h_{uuw} = 0, \quad (15)$$

$$h_{uuu}h_{www} - h_{ww}h_{uuu} = 0, \quad (16)$$

$$-3h_{uuw}^2 + 4h_{uww}h_{uuu} - h_{ww}h_{uuuu} = 0. \quad (17)$$

The classification of solutions will be performed modulo equivalence transformations leaving system (2) form-invariant (and therefore preserving the integrability conditions). These include

$$\tilde{x} = x - 2at, \quad \tilde{y} = y - 2bt, \quad \tilde{h} = h + au^2 + buw + mu + nw + p, \quad (18)$$

as well as

$$\tilde{x} = x - sy, \quad \tilde{w} = w + su; \quad (19)$$

(other variables remain unchanged). We will always assume  $h_{ww} \neq 0$  which is equivalent to the requirement of irreducibility of the dispersion relation. There are two main cases to consider.

**Case 1:**  $h_{www} = 0$ . Then

$$h(u, w) = \alpha(u)w^2 + \beta(u)w + \gamma(u),$$

and the integrability conditions imply

$$\alpha'' = 0, \quad \beta''' = 0, \quad -3\beta''^2 + 8\alpha'\gamma''' - 2\alpha\gamma'''' = 0.$$

There are two further subcases:  $\alpha = 1$  and  $\alpha = u$ .

The subcase  $\alpha = 1$  leads, modulo equivalence transformations (18), to densities of the form

$$h(u, w) = w^2 + \beta_1 u^2 w - \frac{\beta_1^2}{4} u^4 + \gamma_1 u^3,$$

$\beta_1, \gamma_1 = \text{const.}$  For  $\beta_1 = 0$  we obtain the first case of Theorem 2 (after a suitable rescaling). If  $\beta_1 \neq 0$  then we can eliminate the term  $u^3$  by a translation of  $u$ . This gives the second case of Theorem 2 (after rescaling of  $u$  and  $w$ ).

The subcase  $\alpha = u$  leads, modulo equivalence transformations (18), to densities of the form

$$h(u, w) = uw^2 + \beta_1 u^2 w + \gamma_1 u^7 + \frac{\beta_1^2}{4} u^3.$$

$\beta_1, \gamma_1 = \text{const.}$  Note that we can set  $\beta_1 = 0$  using transformation (19) with  $s = \beta_1/2$ . This gives the third case of Theorem 2.

**Case 2:**  $h_{www} \neq 0$ . Then the first two integrability conditions (13) and (14) imply  $h_{www} = ch_{ww}$  for some constant  $c$  (which can be set equal to 1). This gives

$$h(u, w) = a(u)e^w + p(u)w + q(u).$$

The next two integrability conditions (15) and (16) give  $p'' = 0$  and  $q''' = 0$ , respectively. Thus, modulo equivalence transformations (18) we can assume  $h(u, w) = a(u)e^w$ . Finally, equation (17) implies

$$aa'''' - 4a'a''' + 3a''^2 = 0,$$

which is the classical equation for the Weierstrass sigma function (equianharmonic case  $g_2 = 0$ ). Setting  $\wp = -(\ln a)''$  we obtain  $\wp'' = 6\wp^2$ , which integrates to

$$\wp'^2 = 4\wp^3 - g_3, \tag{20}$$

$g_3 = \text{const.}$  There are three subcases.

*Subcase  $g_3 = 0, \wp = 0$ .* Then  $a(u) = e^{\alpha u + \beta}$  and modulo equivalence transformations (19) we obtain Case 4 of Theorem 2.

*Subcase  $g_3 = 0, \wp = \frac{1}{u^2}$ .* Then  $a(u) = ue^{\alpha u + \beta}$  and modulo equivalence transformations (19) we obtain Case 5 of Theorem 2.

*Subcase  $g_3 \neq 0$ .* Then  $a(u) = \sigma(u; 0, g_3)e^{\alpha u + \beta}$  and modulo equivalence transformations (19) we obtain the last case of our classification. This finishes the proof of Theorem 2.

**Remark.** The paper [12] gives a classification of integrable two-component Hamiltonian systems of the form

$$\begin{bmatrix} U_t \\ W_t \end{bmatrix} = \begin{bmatrix} 0 & \partial_x \\ \partial_x & \partial_y \end{bmatrix} \begin{bmatrix} \frac{\delta H}{\delta U} \\ \frac{\delta H}{\delta W} \end{bmatrix} \tag{21}$$

where  $H = \int F(U, W) dx dy$ . Explicitly, we have

$$U_t = (F_W)_x, \quad W_t = (F_U)_x + (F_W)_y.$$

Let us introduce a contact change of variables  $(U, W, F) \rightarrow (u, w, f)$  via partial Legendre transform:

$$w = F_W, \quad u = U, \quad f = F - WF_W, \quad f_w = -W, \quad f_u = F_U.$$

In the new variables the system becomes

$$w_y = -(f_u)_x - (f_w)_t, \quad u_t = w_x.$$

Modulo relabelling  $u \leftrightarrow w$ ,  $f \rightarrow -h$ ,  $y \rightarrow t$ ,  $t \rightarrow x$ ,  $x \rightarrow y$  these equations coincide with (2). Thus, Hamiltonian formalisms (1) and (21) are equivalent. Examples of dKP and Boyer-Finley equations suggest however that Hamiltonian formalism (1) is more natural and convenient, indeed, in the form (1) both equations arise directly in their ‘physical’ variables.

### 2.3 Dispersionless Lax pairs

In this section we provide dispersionless Lax representations for all six canonical forms of Theorem 2. The results are summarised in Table 1 below.

Table 1: Dispersionless Lax pairs for integrable systems (2)

Hamiltonian density $h(u, w)$	Dispersionless Lax pair
$h(u, w) = \frac{1}{2}w^2 + \frac{1}{6}u^3$ System (2) : $u_t = uu_x + w_y$ $w_x = u_y$	$S_t = \frac{1}{3}S_x^3 + uS_x + w$ $S_y = \frac{1}{2}S_x^2 + u$
$h(u, w) = w^2 + u^2w - \frac{1}{4}u^4$ System (2) : $u_t = (2w - 3u^2)u_x + 4uu_y + 2w_y$ $w_x = u_y$	$S_t = (3u^2 + 2w)S_x + 2uS_x^4 + \frac{2}{7}S_x^7$ $S_y = uS_x + \frac{1}{4}S_x^4$
$h(u, w) = uw^2 + \beta u^7$ System (2) : $u_t = 42\beta u^5 u_x + 4uw_y + 2uw_y$ $w_x = u_y$	$S_t = 4u^2 \wp(S_x)(w + \frac{1}{5}u^3 \wp'(S_x))$ $S_y = u^2 \wp(S_x)$ here $\wp'^2 = 4\wp^3 - 35\beta$
$h(u, w) = e^w$ System (2) : $u_t = e^w w_y$ $w_x = u_y$	$S_t = -\frac{e^w}{S_x + u}$ $S_y = -\ln(S_x + u)$
$h(u, w) = ue^w$ System (2) : $u_t = e^w(2u_y + uu_y)$ $w_x = u_y$	$S_t = \frac{3u^2 e^w S_x}{u^3 - S_x^3}$ $S_y = \ln(S_x - u) + \epsilon \ln(S_x - \epsilon u) + \epsilon^2 \ln(S_x - \epsilon^2 u)$ here $\epsilon = \exp\left(\frac{2\pi i}{3}\right)$
$h(u, w) = \sigma(u)e^w$ System (2) : $u_t = e^w(\sigma''u_x + 2\sigma'u_y + \sigma w_y)$ $w_x = u_y$	$S_t = \sigma(u)e^w G_u(S_x, u)$ $S_y = G(S_x, u)$ here $\sigma(u) = \sigma(u; 0, g_3)$

In the last case the function  $G(p, u)$  is defined by the equations

$$G_p = \frac{G_{uu}}{G_u} - \zeta(u), \quad G_{uuu}G_u - 2G_{uu}^2 + 2\wp(u)G_u^2 = 0 \quad (22)$$

where  $\zeta$  and  $\wp$  are the Weierstrass functions (equianharmonic case  $g_2 = 0$ ). The general solution of these equations is given by the formula

$$G(p, u) = \ln \sigma(\lambda(p - u)) + \epsilon \ln \sigma(\lambda(p - \epsilon u)) + \epsilon^2 \ln \sigma(\lambda(p - \epsilon^2 u)) \quad (23)$$

where  $\epsilon = e^{2\pi i/3} = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$  and  $\lambda = \frac{i}{\sqrt{3}}$ . Note that the degeneration  $g_3 \rightarrow 0$ ,  $\sigma(u) \rightarrow u$  takes the Lax pair corresponding to the Hamiltonian density  $h = \sigma(u)e^w$  to the Lax pair for the density  $h = ue^w$ . We refer to the Appendix for a proof that formula (23) indeed solves the equations (22): this requires some non-standard identities for equianharmonic elliptic functions.

## 2.4 Commuting flows: proof of Theorem 3

Our aim is to show that every integrable system (2) possesses a commuting flow of the form (10),

$$\begin{aligned} u_\tau &= a(u, w, v)u_x + b(u, w, v)u_y + c(u, w, v)w_y + d(u, w, v)v_y, \\ w_x &= u_y, \\ v_x &= (p(u, w))_y. \end{aligned}$$

Here  $\tau$  is the higher ‘time’ variable and  $v = \partial_x^{-1}\partial_y p(u, w)$  is a new nonlocality (to be determined). Due to the presence of nonlocal variables, direct computation of compatibility condition  $u_{t\tau} = u_{\tau t}$  is not straightforward. Therefore, we adopt a different approach and require that the combined system (2)  $\cup$  (10),

$$u_t = (h_u)_x + (h_w)_y, \tag{24}$$

$$u_\tau = a(u, w, v)u_x + b(u, w, v)u_y + c(u, w, v)w_y + d(u, w, v)v_y, \tag{25}$$

$$w_x = u_y, \tag{26}$$

$$v_x = (p(u, w))_y, \tag{27}$$

possesses hydrodynamic reductions. Thus, we seek multiphase solutions of the form  $u = u(R^1, \dots, R^n)$ ,  $w = w(R^1, \dots, R^n)$  and  $v = v(R^1, \dots, R^n)$  where the Riemann invariants  $R^i$  satisfy a triple of commuting systems of hydrodynamic type:

$$R_y^i = \mu^i(R)R_x^i, \quad R_t^i = \lambda^i(R)R_x^i, \quad R_\tau^i = \eta^i(R)R_x^i.$$

We recall that the commutativity conditions are equivalent to

$$\frac{\partial_j \mu^i}{\mu^j - \mu^i} = \frac{\partial_j \lambda^i}{\lambda^j - \lambda^i} = \frac{\partial_j \eta^i}{\eta^j - \eta^i}. \tag{28}$$

Following the same procedure as in Section 2.1, from equations (24) and (26) we obtain the relations  $\partial_i w = \mu^i \partial_i u$ , the GT-system (11), (12), and the integrability conditions (9) for the Hamiltonian density  $h(u, w)$ . Similarly, equation (27) implies

$$\partial_i v = (p_u \mu^i + p_w (\mu^i)^2) \partial_i u,$$

and the compatibility condition  $\partial_j \partial_i v = \partial_i \partial_j v$  results in the relations

$$h_{uuw}p_w - h_{ww}p_{uu} = 0, \quad h_{uww}p_w - h_{ww}p_{uw} = 0, \quad h_{www}p_w - h_{ww}p_{ww} = 0.$$

Modulo unessential constants of integration (which can be removed by equivalence transformations) these relations uniquely specify the nonlocality:

$$p(u, w) = h_w(u, w).$$

Finally, equation (25) gives an additional dispersion relation,

$$\eta^i = a + b\mu^i + (c + p_u d)(\mu^i)^2 + p_w d(\mu^i)^3.$$

Substituting  $\eta^i$  into the commutativity conditions (28) we obtain the following set of relations:

$$p_w^2 d_v = 0, \quad (29)$$

$$((p_{ww}d + p_w d_w) + p_w p_u d_v) h_{ww} = 2p_w dh_{www}, \quad (30)$$

$$(p_{uw}d + p_w d_u) h_{ww} = 2p_w dh_{uww}, \quad (31)$$

$$h_{ww}(c_v + p_u d_v) p_w = p_w dh_{www}, \quad (32)$$

$$h_{ww}((c_w + p_{uw}d + p_u d_w) + (c_v + p_u d_v) p_u) = 5h_{uww} dp_w + ch_{www} + p_u dh_{www}, \quad (33)$$

$$h_{ww} b_v p_w = 2p_w dh_{uww}, \quad (34)$$

$$h_{ww}(c_u + p_{uu}d + p_u d_u) = 4p_w dh_{uuw} + ch_{uww} + p_u dh_{uww}, \quad (35)$$

$$h_{ww} a_v p_w = p_w dh_{uuw}, \quad (36)$$

$$h_{ww}(b_w + b_v p_u) = 4p_w dh_{uuw} + 2ch_{uww} + 2p_u dh_{uww}, \quad (37)$$

$$h_{ww} b_u = 2p_w dh_{uuu} + 2ch_{uuw} + 2p_u dh_{uuw}, \quad (38)$$

$$h_{ww}(a_w + a_v p_u) = p_w dh_{uuu} + ch_{uuw} + p_u dh_{uuw}, \quad (39)$$

$$h_{ww} a_u = ch_{uuu} + p_u dh_{uuu}. \quad (40)$$

Using the fact that  $p = h_w$  we solve these relations modulo the integrability conditions (9), recall that  $h_{ww} \neq 0$ . Equation (29) gives  $d_v = 0$ . Equations (30, 31) imply

$$dh_{www} - h_{ww} d_w = 0, \quad dh_{uww} - h_{ww} d_u = 0,$$

which can be solved for  $d$ :

$$d = \delta h_{ww},$$

for some constant  $\delta$  (which will be set equal to 2 in what follows). Equation (32) gives

$$c_v = d \frac{h_{www}}{h_{ww}}.$$

Setting  $c = d \frac{h_{www}}{h_{ww}} v + c_1$  for some  $c_1 = c_1(u, w)$  and substituting into equation (35) we find

$$c_1 = 3dh_{uw} + c_2(w)h_{ww}.$$

Substituting  $c = d \frac{h_{www}}{h_{ww}} v + 3dh_{uw} + c_2(w)h_{ww}$  into equation (33) we find

$$(c_2)_w = d \left( \frac{h_{uww} h_{ww} - h_{www} h_{uw}}{h_{ww}^2} \right) = d \frac{\partial}{\partial w} \left( \frac{h_{uw}}{h_{ww}} \right).$$

It turns out that modulo the integrability conditions  $(c_2)_w$  is a constant. If we set

$$\alpha = d \frac{\partial}{\partial w} \left( \frac{h_{uw}}{h_{ww}} \right),$$

the final formula for  $c$  can be written as

$$c = d \frac{h_{www}}{h_{ww}} v + 3dh_{uw} + \alpha w h_{ww}.$$

The equations for the coefficients  $a$  and  $b$  cannot be integrated explicitly; rearranging the remaining equations gives the following final result:

$$\begin{cases} a_u = \frac{h_{uuu}}{h_{ww}}(c + dh_{uw}), \\ a_w = \frac{h_{uuw}}{h_{ww}}c + dh_{uuu}, \\ a_v = d \frac{h_{uuw}}{h_{ww}}. \end{cases} \quad \begin{cases} b_u = 2(dh_{uuu} + \frac{h_{uuu}}{h_{ww}}(c + dh_{uw})), \\ b_w = 2(\frac{h_{uuw}}{h_{ww}}c + 2dh_{uuu}), \\ b_v = 2d \frac{h_{uuw}}{h_{ww}}. \end{cases}$$

$$c = d \frac{h_{www}}{h_{ww}} v + 3dh_{uw} + \alpha w h_{ww}, \quad d = \delta h_{ww}, \quad \alpha = d \frac{\partial}{\partial w} \left( \frac{h_{uw}}{h_{ww}} \right).$$

The equations for  $a$  and  $b$  are consistent modulo integrability conditions (9). This proves the existence of commuting flows (10).

**Hamiltonian formulation of commuting flows.** Our next goal is to show that the obtained commuting flow can be cast into Hamiltonian form

$$u_\tau = \partial_x \left( \frac{\delta F}{\delta u} \right), \quad F = \int f(u, w, v) \, dx dy, \quad (41)$$

with the nonlocal variables  $w, v$  defined by  $w_x = u_y$ ,  $v_x = (h_w)_y$ . More precisely, we claim that the commuting density  $f$  is given by the formula

$$f(u, w, v) = v h_w + g(u, w)$$

where the function  $g(u, w)$  is yet to be determined. We have

$$\frac{\delta F}{\delta u} = 2v h_{uw} + g_u + \partial_x^{-1} \partial_y (2v h_{ww} + g_w),$$

so that equation (41) takes the form

$$\begin{aligned} u_\tau &= (2v h_{uw} + g_u)_x + (2v h_{ww} + g_w)_y, \\ w_x &= u_y, \\ v_x &= (h_w)_y. \end{aligned} \quad (42)$$

Explicitly, (42) gives

$$\begin{aligned} u_\tau &= (2vh_{uuw} + g_{uu})u_x + (4vh_{uww} + 2g_{uw} + 2(h_{uw})^2)u_y \\ &\quad + (2h_{uw}h_{ww} + 2vh_{www} + g_{ww})w_y + 2h_{ww}v_y. \end{aligned}$$

Comparing this with (25) we thus require

$$\begin{aligned} a &= 2vh_{uuw} + g_{uu}, \\ b &= 4vh_{uww} + 2g_{uw} + 2h_{uw}^2, \\ c &= 2vh_{www} + 2h_{uw}h_{ww} + g_{ww}, \\ d &= 2h_{ww}. \end{aligned}$$

Using the expressions for  $a, b, c, d$  calculated above we obtain the equations for  $g(u, w)$ :

$$\begin{aligned} g_{ww} &= 4h_{uw}h_{ww} + \alpha wh_{ww}, \\ g_{uuu} &= 8h_{uw}h_{uuu} + \alpha wh_{uuu}, \\ g_{uuw} &= 6h_{uw}h_{uuw} + 2h_{ww}h_{uuu} + \alpha wh_{uuw}; \end{aligned}$$

note that these equations are consistent modulo the integrability conditions (9). This finishes the proof of Theorem 3.

## 2.5 Commuting flows and dispersionless Lax pairs

In this section we calculate commuting flows of integrable systems (2) and construct their dispersionless Lax pairs.

**1. Hamiltonian density**  $h(u, w) = \frac{1}{6}u^3 + \frac{1}{2}w^2$ . The commuting density is

$$f(u, w, v) = vw + u^2w.$$

Commuting flow has the form (note that  $\alpha = 0$ ):

$$\begin{aligned} u_\tau &= 2wu_x + 4uu_y + 2v_y, \\ w_x &= u_y, \\ v_x &= w_y. \end{aligned}$$

Dispersionless Lax pair:

$$\begin{aligned} S_y &= \frac{1}{2}S_x^2 + u, \\ S_\tau &= \frac{1}{2}S_x^4 + 2uS_x^2 + 2wS_x + 2u^2 + 2v. \end{aligned}$$

**2. Hamiltonian density**  $h(u, w) = w^2 + u^2w - \frac{1}{4}u^4$ . The commuting density is

$$f(u, w, v) = 2wv + u^2v + 8uw^2 - \frac{8}{5}u^5.$$

Commuting flow has the form (note that  $\alpha = 0$ ):

$$\begin{aligned} u_\tau &= (4v - 32u^3)u_x + (32w + 8u^2)u_y + 24uw_y + 4v_y, \\ w_x &= u_y, \\ v_x &= (2w + u^2)_y. \end{aligned}$$

Dispersionless Lax pair:

$$\begin{aligned} S_y &= uS_x + \frac{1}{4}S_x^4, \\ S_\tau &= (4v + 32uw + 16u^3)S_x + (8w + 24w^2)S_x^4 + 8uS_x^7 + \frac{4}{5}S_x^{10}. \end{aligned}$$

**3. Hamiltonian density**  $h(u, w) = uw^2 + \beta u^7$ . The commuting density is

$$f(u, w, v) = 2uvw + 4uw^3 + 20\beta u^7w.$$

Commuting flow has the form (note that  $\alpha = 4$ ):

$$\begin{aligned} u_\tau &= 840\beta u^5 w u_x + (8v + 32w^2 + 280\beta u^6)u_y + 32u w w_y + 4v_y, \\ w_x &= u_y, \\ v_x &= (2uw)_y. \end{aligned}$$

Dispersionless Lax pair:

$$\begin{aligned} S_y &= u^2\wp(S_x), \\ S_\tau &= (8u^2v + 32u^2w^2 + 16u^5w\wp'(S_x) + 8u^8\wp^3(S_x))\wp(S_x), \end{aligned}$$

where  $\wp'^2 = 4\wp^3 - 35\beta$ .

**4. Hamiltonian density**  $h(u, w) = e^w$ . The commuting density is

$$f(u, w, v) = ve^w.$$

Commuting flow has the form (note that  $\alpha = 0$ ):

$$\begin{aligned} u_\tau &= 2ve^w w_y + 2e^w v_y, \\ w_x &= u_y, \\ v_x &= (e^w)_y. \end{aligned}$$

Dispersionless Lax pair:

$$\begin{aligned} S_y &= -\ln(S_x + u), \\ S_\tau &= \frac{-2ve^w}{S_x + u} + \frac{e^{2w}}{(S_x + u)^2}. \end{aligned}$$

**5. Hamiltonian density**  $h(u, w) = ue^w$ . The commuting density is

$$f(u, w, v) = uve^w + ue^{2w}.$$

Commuting flow has the form (note that  $\alpha = 0$ ):

$$\begin{aligned} u_\tau &= (4ve^w + 6e^{2w})u_y + (2uve^w + 6ue^{2w})w_y + 2ue^w v_y, \\ w_x &= u_y, \\ v_x &= (ue^w)_y. \end{aligned}$$

Dispersionless Lax pair:

$$\begin{aligned} S_y &= \ln(S_x - u) + \varepsilon \ln(S_x - \varepsilon u) + \varepsilon^2 \ln(S_x - \varepsilon^2 u), \\ S_\tau &= \frac{3u^2 e^w S_x (2u^3 v - 2v S_x^3 - 3e^w S_x^3)}{(S_x^3 - u^3)^2}. \end{aligned}$$

**6. Hamiltonian density**  $h(u, w) = \sigma(u)e^w$ . The commuting density is

$$f(u, w, v) = v\sigma(u)e^w + \sigma(u)\sigma'(u)e^{2w}.$$

Commuting flow has the form (note that  $\alpha = 0$ ):

$$\begin{aligned} u_\tau &= (2v\sigma''e^w + (\sigma\sigma')''e^{2w})u_x + (4v\sigma'e^w + (4\sigma\sigma'' + 6\sigma'^2)e^{2w})u_y + (2v\sigma e^w + 6\sigma\sigma'e^{2w})w_y + 2\sigma e^w v_y, \\ w_x &= u_y, \\ v_x &= (\sigma e^w)_y. \end{aligned}$$

Dispersionless Lax pair:

$$\begin{aligned} S_y &= G(S_x, u) \\ S_\tau &= 2[ve^w \sigma(u) + e^{2w} \sigma(u)\sigma'(u)]G_u(S_x, u) - e^{2w} \sigma(u)^2 G_{uu}(S_x, u), \end{aligned}$$

here  $G(S_x, u)$  is defined by equations (22).

### 3 Dispersive deformations

Dispersive deformations of hydrodynamic type systems in  $1 + 1$  dimensions were thoroughly investigated in [4, 5, 6, 7] based on deformations of the corresponding hydrodynamic symmetries. In  $2 + 1$  dimensions, an alternative approach based on deformations of hydrodynamic reductions was proposed in [14, 15].

It still remains a challenging problem to construct dispersive deformations of all Hamiltonian systems (2) obtained in this paper. In general, all three ingredients of the construction may need to be deformed, namely, the Hamiltonian operator  $\partial_x$ , the Hamiltonian density  $h(u, w)$  and the nonlocality  $w$ . Here we give just two examples.

**Example 1: dKP equation.** The Hamiltonian density  $h = \frac{1}{2}w^2 + \frac{1}{6}u^3$  results in the dKP equation:

$$u_t = uu_x + w_y, \quad w_x = u_y.$$

It possesses an integrable dispersive deformation

$$u_t = uu_x + w_y - \epsilon^2 u_{xxx}, \quad w_x = u_y,$$

which is the full KP equation (in this section  $\epsilon$  denotes an arbitrary deformation parameter). The KP equation corresponds to the deformed Hamiltonian density

$$h(u, w) = \frac{1}{2}w^2 + \frac{1}{6}u^3 + \frac{\epsilon^2}{2}u_x^2,$$

while the Hamiltonian operator  $\partial_x$  and the nonlocality  $w = \partial_x^{-1}\partial_y u$  stay the same. Indeed, we have

$$u_t = \partial_x \frac{\delta H}{\delta u} = \partial_x (\partial_x^{-1}\partial_y w + \frac{1}{2}u^2 - \epsilon^2 u_{xx}) = uu_x + w_y - \epsilon^2 u_{xxx}.$$

**Example 2: Boyer-Finley equation.** The Hamiltonian density  $h = e^w$  results in the dispersionless Toda (Boyer-Finley) equation:

$$u_t = e^w w_y, \quad w_x = u_y.$$

It possesses an integrable dispersive deformation

$$u_t = \left( \frac{1 - T^{-1}}{\epsilon} \right) e^w, \quad w_x = \left( \frac{T - 1}{\epsilon} \right) u,$$

which is the full Toda equation. Here  $T$  and  $T^{-1}$  denote the forward/backward  $\epsilon$ -shifts in the  $y$ -direction, so that  $\frac{T-1}{\epsilon}$  and  $\frac{1-T^{-1}}{\epsilon}$  are the forward/backward discrete  $y$ -derivatives. The Toda equation corresponds to the deformed nonlocality  $w = \partial_x^{-1} \frac{T-1}{\epsilon} u$ , while the

Hamiltonian operator  $\partial_x$  and the Hamiltonian density  $h = e^w$  stay the same. Indeed, we have

$$\frac{\delta H}{\delta u} = \partial_x^{-1} \left( \frac{1 - T^{-1}}{\epsilon} \right) e^w,$$

so that

$$u_t = \partial_x \frac{\delta H}{\delta u} = \left( \frac{1 - T^{-1}}{\epsilon} \right) e^w,$$

as required.

## 4 Appendix: dispersionless Lax pair for $h = \sigma(u)e^w$

Here we prove that expression (23),

$$G(p, u) = \ln \sigma(\lambda(p - u)) + \epsilon \ln \sigma(\lambda(p - \epsilon u)) + \epsilon^2 \ln \sigma(\lambda(p - \epsilon^2 u)),$$

where  $\epsilon = e^{2\pi i/3} = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$  and  $\lambda = \frac{i}{\sqrt{3}}$ , solves the equations (22),

$$G_p = \frac{G_{uu}}{G_u} - \zeta(u), \quad G_{uuu}G_u - 2G_{uu}^2 + 2\wp(u)G_u^2 = 0.$$

In what follows we will use the addition formula

$$\zeta(u + v) = \zeta(u) + \zeta(v) + \frac{1}{2} \frac{\wp'(u) - \wp'(v)}{\wp(u) - \wp(v)}. \quad (43)$$

We will also need the following identity:

**Proposition 1.** *In the equianharmonic case, the Weierstrass functions satisfy the identity*

$$\lambda \frac{\wp'(\lambda u)}{\wp(\lambda u)} + 3\lambda\zeta(\lambda u) - \zeta(u) = 0, \quad \lambda = \frac{i}{\sqrt{3}}. \quad (44)$$

**Proof:**

Using the standard expansions

$$\zeta(z) = \frac{1}{z} - \frac{g_3}{140}z^5 - \dots, \quad \wp(z) = \frac{1}{z^2} + \frac{g_3}{28}z^4 + \dots,$$

one can show that formula (44) holds to high order in  $z$  for the specific parameter value  $\lambda = \frac{i}{\sqrt{3}}$ . Therefore, it is sufficient to establish the differentiated (by  $u$ ) identity (44), namely,

$$-\lambda^2\wp(\lambda u) + \frac{\lambda^2 g_3}{\wp^2(\lambda u)} + \wp(u) = 0, \quad (45)$$

where we have used  $\wp'' = 6\wp^2$  and  $\wp'^2 = 4\wp^3 - g_3$ . Explicitly, (45) reads

$$\wp(iu/\sqrt{3}) - \frac{g_3}{\wp^2(iu/\sqrt{3})} + 3\wp(u) = 0.$$

Setting  $u = i\sqrt{3}v$  we obtain

$$\wp(v) - \frac{g_3}{\wp^2(v)} + 3\wp(i\sqrt{3}v) = 0. \quad (46)$$

Thus, it is sufficient to establish (46). Formulae of this kind appear in the context of complex multiplication for elliptic curves with extra symmetry. Let us begin with the standard invariance properties of the equianharmonic  $\zeta$ -function:

$$\zeta(\epsilon z) = \epsilon^2 \zeta(z), \quad \zeta(\epsilon^2 z) = \epsilon \zeta(z);$$

here  $\epsilon = e^{2\pi i/3} = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$  is the cubic root of unity. Setting  $z = 2v$  this gives

$$\zeta(-v + i\sqrt{3}v) = \epsilon^2 \zeta(2v), \quad \zeta(-v - i\sqrt{3}v) = \epsilon \zeta(2v).$$

Using the addition formula (43) one can rewrite these relations in the form

$$-\zeta(v) + \zeta(i\sqrt{3}v) + \frac{1 - \wp'(v) - \wp'(i\sqrt{3}v)}{2(\wp(v) - \wp(i\sqrt{3}v))} = \epsilon^2 \zeta(2v)$$

and

$$-\zeta(v) - \zeta(i\sqrt{3}v) + \frac{1 - \wp'(v) + \wp'(i\sqrt{3}v)}{2(\wp(v) - \wp(i\sqrt{3}v))} = \epsilon \zeta(2v),$$

respectively. Adding these relations together (and keeping in mind that  $1 + \epsilon + \epsilon^2 = 0$ ) we obtain

$$-2\zeta(v) - \frac{\wp'(v)}{\wp(v) - \wp(i\sqrt{3}v)} + \zeta(2v) = 0.$$

Using the duplication formula  $\zeta(2v) = 2\zeta(v) + \frac{3\wp^2(v)}{\wp'(v)}$  this simplifies to

$$-\frac{\wp'(v)}{\wp(v) - \wp(i\sqrt{3}v)} + \frac{3\wp^2(v)}{\wp'(v)} = 0,$$

which is equivalent to (46) via  $\wp'^2 = 4\wp^3 - g_3$ . □

**Proposition 2.** *Expression (23) solves the equations (22).*

**Proof:**

Computation of partial derivatives of  $G(p, u)$  gives

$$G_p = \lambda\zeta(\lambda(p-u)) + \lambda\epsilon\zeta(\lambda(p-\epsilon u)) + \lambda\epsilon^2\zeta(\lambda(p-\epsilon^2 u)),$$

$$G_u = -\lambda\zeta(\lambda(p-u)) - \lambda\epsilon^2\zeta(\lambda(p-\epsilon u)) - \lambda\epsilon\zeta(\lambda(p-\epsilon^2 u)).$$

Using the addition formula (43), the identity  $1 + \epsilon + \epsilon^2 = 0$ , and the invariance

$$\begin{aligned}\zeta(\epsilon z) &= \epsilon^2\zeta(z), & \zeta(\epsilon^2 z) &= \epsilon\zeta(z), \\ \wp(\epsilon z) &= \epsilon\wp(z), & \wp(\epsilon^2 z) &= \epsilon^2\wp(z), \\ \wp'(\epsilon z) &= \wp'(z), & \wp'(\epsilon^2 z) &= \wp'(z),\end{aligned}$$

we obtain:

$$\begin{aligned}\frac{1}{\lambda}G_p &= \zeta(\lambda(p-u)) + \epsilon\zeta(\lambda(p-\epsilon u)) + \epsilon^2\zeta(\lambda(p-\epsilon^2 u)) \\ &= \zeta(\lambda p) - \zeta(\lambda u) + \frac{1}{2} \frac{\wp'(\lambda p) + \wp'(\lambda u)}{\wp(\lambda p) - \wp(\lambda u)} \\ &\quad + \epsilon \left( \zeta(\lambda p) - \epsilon^2\zeta(\lambda u) + \frac{1}{2} \frac{\wp'(\lambda p) + \wp'(\lambda u)}{\wp(\lambda p) - \epsilon\wp(\lambda u)} \right) \\ &\quad + \epsilon^2 \left( \zeta(\lambda p) - \epsilon\zeta(\lambda u) + \frac{1}{2} \frac{\wp'(\lambda p) + \wp'(\lambda u)}{\wp(\lambda p) - \epsilon^2\wp(\lambda u)} \right) \\ &= -3\zeta(\lambda u) + \frac{\wp'(\lambda p) + \wp'(\lambda u)}{2} \left( \frac{1}{\wp(\lambda p) - \wp(\lambda u)} + \frac{\epsilon}{\wp(\lambda p) - \epsilon\wp(\lambda u)} + \frac{\epsilon^2}{\wp(\lambda p) - \epsilon^2\wp(\lambda u)} \right) \\ &= -3\zeta(\lambda u) + \frac{\wp'(\lambda p) + \wp'(\lambda u)}{2} \frac{3\wp^2(\lambda u)}{\wp^3(\lambda p) - \wp^3(\lambda u)} \\ &= -3\zeta(\lambda u) + \frac{\wp'(\lambda p) + \wp'(\lambda u)}{2} \frac{12\wp^2(\lambda u)}{\wp'^2(\lambda p) - \wp'^2(\lambda u)} \\ &= -3\zeta(\lambda u) + \frac{6\wp^2(\lambda u)}{\wp'(\lambda p) - \wp'(\lambda u)}.\end{aligned}$$

A similar calculation gives:

$$\begin{aligned}
-\frac{1}{\lambda}G_u &= \zeta(\lambda(p-u)) + \epsilon^2\zeta(\lambda(p-\epsilon u)) + \epsilon\zeta(\lambda(p-\epsilon^2u)) \\
&= \zeta(\lambda p) - \zeta(\lambda u) + \frac{1}{2}\frac{\wp'(\lambda p)+\wp'(\lambda u)}{\wp(\lambda p)-\wp(\lambda u)} \\
&+ \epsilon^2\left(\zeta(\lambda p) - \epsilon^2\zeta(\lambda u) + \frac{1}{2}\frac{\wp'(\lambda p)+\wp'(\lambda u)}{\wp(\lambda p)-\epsilon\wp(\lambda u)}\right) \\
&+ \epsilon\left(\zeta(\lambda p) - \epsilon\zeta(\lambda u) + \frac{1}{2}\frac{\wp'(\lambda p)+\wp'(\lambda u)}{\wp(\lambda p)-\epsilon^2\wp(\lambda u)}\right) \\
&= \frac{\wp'(\lambda p)+\wp'(\lambda u)}{2}\left(\frac{1}{\wp(\lambda p)-\wp(\lambda u)} + \frac{\epsilon^2}{\wp(\lambda p)-\epsilon\wp(\lambda u)} + \frac{\epsilon}{\wp(\lambda p)-\epsilon^2\wp(\lambda u)}\right) \\
&= \frac{\wp'(\lambda p)+\wp'(\lambda u)}{2}\frac{3\wp(\lambda p)\wp(\lambda u)}{\wp^3(\lambda p)-\wp^3(\lambda u)} \\
&= \frac{\wp'(\lambda p)+\wp'(\lambda u)}{2}\frac{12\wp(\lambda p)\wp(\lambda u)}{\wp'^2(\lambda p)-\wp'^2(\lambda u)} \\
&= \frac{6\wp(\lambda p)\wp(\lambda u)}{\wp'(\lambda p)-\wp'(\lambda u)}.
\end{aligned}$$

To summarise, we have:

$$G_p = -3\lambda\zeta(\lambda u) + \frac{6\lambda\wp^2(\lambda u)}{\wp'(\lambda p) - \wp'(\lambda u)}, \quad G_u = -\frac{6\lambda\wp(\lambda p)\wp(\lambda u)}{\wp'(\lambda p) - \wp'(\lambda u)}.$$

This gives

$$\frac{G_{uu}}{G_u} = (\ln G_u)_u = \lambda\frac{\wp'(\lambda u)}{\wp(\lambda u)} + \frac{6\lambda\wp^2(\lambda u)}{\wp'(\lambda p) - \wp'(\lambda u)}, \quad (47)$$

and the first equation (22),  $G_p = \frac{G_{uu}}{G_u} - \zeta(u)$ , is satisfied identically due to (44). Finally, the second equation (22),  $G_{uuu}G_u - 2G_{uu}^2 + 2\wp(u)G_u^2 = 0$ , which can be written in the equivalent form

$$-\left(\frac{G_{uu}}{G_u}\right)_u + \left(\frac{G_{uu}}{G_u}\right)^2 = 2\wp(u),$$

is satisfied identically due to (47) and (45).  $\square$

## Acknowledgements

We thank Yurii Brezhnev and Maxim Pavlov for clarifying discussions. The research of EVF was supported by the EPSRC grant EP/N031369/1. The research of VSN was supported by the EPSRC grant EP/V050451/1.

## References

- [1] N.I. Akhiezer, Elements of the theory of elliptic functions, Translated from the second Russian edition by H. H. McFaden. Translations of Mathematical Monographs, 79. American Mathematical Society, Providence, RI ( 1990) 237 pp.
- [2] E. Cartan, Sur une classe d'espaces de Weyl, Ann. Sci. École Norm. Sup. (3) **60** (1943) 1-16.
- [3] E. Cartan, The geometry of differential equations of third order, Revista Mat. Hisp.-Amer. **4** (1941) 3-33.
- [4] B.A. Dubrovin, Hamiltonian PDEs: deformations, integrability, solutions, J. Phys. A **43**, no. 43 (2010) 434002, 20 pp.
- [5] B.A. Dubrovin, On Hamiltonian perturbations of hyperbolic systems of conservation laws. II. Universality of critical behaviour, Comm. Math. Phys. **267**, no. 1 (2006) 117-139.
- [6] B.A. Dubrovin, Si-Qi Liu and Youjin Zhang, On Hamiltonian perturbations of hyperbolic systems of conservation laws. I. Quasi-triviality of bi-Hamiltonian perturbations, Comm. Pure Appl. Math. **59**, no. 4 (2006) 559-615.
- [7] B.A. Dubrovin and Youjin Zhang, Bi-Hamiltonian hierarchies in 2D topological field theory at one-loop approximation, Comm. Math. Phys. **198**, no. 2 (1998) 311-361.
- [8] M. Dunajski, E.V. Ferapontov and B. Kruglikov, On the Einstein-Weyl and conformal self-duality equations, J. Math. Phys. **56** (2015) 083501.
- [9] E.V. Ferapontov and K.R. Khusnutdinova, On integrability of (2+1)-dimensional quasilinear systems, Comm. Math. Phys. **248** (2004) 187-206.
- [10] E.V. Ferapontov and K.R. Khusnutdinova, The characterization of 2-component (2+1)-dimensional integrable systems of hydrodynamic type, J. Phys. A: Math. Gen. **37**, no. 8 (2004) 2949-2963.
- [11] E.V. Ferapontov, A. Moro and V. V. Sokolov, Hamiltonian systems of hydrodynamic type in 2+1 dimensions, Comm. Math. Phys. **285**, no. 1 (2009) 31-65.
- [12] E.V. Ferapontov, A.V. Odesskii and N.M. Stoilov, Classification of integrable two-component Hamiltonian systems of hydrodynamic type in 2+1 dimensions, Journal of Mathematical Physics, **52** (2011) 073505; Manuscript ID: 10-1152R.
- [13] E.V. Ferapontov and B. Kruglikov, Dispersionless integrable systems in 3D and Einstein-Weyl geometry, J. Diff. Geom. **97** (2014) 215-254.

- [14] E.V. Ferapontov and A. Moro, Dispersive deformations of hydrodynamic reductions of 2D dispersionless integrable systems, *J. Phys. A: Math. Theor.* **42**, no. 3 (2009) 035211, 15 pp.
- [15] E.V. Ferapontov, A. Moro and V.S. Novikov, Integrable equations in 2+1 dimensions: deformations of dispersionless limits, *J. Phys. A: Math. Theor.* **42**, no. 34 (2009) 345205, 18 pp.
- [16] J. Gibbons and S.P. Tsarev, Reductions of the Benney equations, *Phys. Lett. A* **211** (1996) 19-24.
- [17] N.J. Hitchin, Complex manifolds and Einstein's equations, *Twistor geometry and nonlinear systems* (Primorsko, 1980), 73-99, *Lecture Notes in Math.* **970**, Springer, Berlin-New York (1982).
- [18] S.P. Tsarev, Poisson brackets and one-dimensional Hamiltonian systems of hydrodynamic type, *Dokl. Akad. Nauk SSSR* **282**, no. 3 (1985) 534-537.
- [19] S.P. Tsarev, Geometry of Hamiltonian systems of hydrodynamic type. The generalized hodograph method, *Izvestija AN USSR Math.* **54**, no. 5 (1990) 1048-1068.
- [20] E.V. Zakharov, Dispersionless limit of integrable systems in 2 + 1 dimensions, in *Singular Limits of Dispersive Waves*, Ed. N.M. Ercolani et al., Plenum Press, NY, (1994) 165-174.