The membership problem for constant-sized quantum correlations is undecidable

Honghao Fu^{*1}, Carl A. Miller^{+1,2}, and William Slofstra^{‡3}

¹Joint Institute for Quantum Information and Computer Science, University of Maryland, College Park, MD, 20740, USA

²National Institute of Standards and Technology, 100 Bureau Dr., Gaithersburg, MD 20899, USA ³Institute for Quantum Computing and Department of Pure Mathematics, University of Waterloo,

Waterloo, Canada

May 3, 2022

Abstract

When two spatially separated parties make measurements on an unknown entangled quantum state, what correlations can they achieve? How difficult is it to determine whether a given correlation is a quantum correlation? These questions are central to problems in quantum communication and computation. Previous work has shown that the general membership problem for quantum correlations is computationally undecidable. In the current work we show something stronger: there is a family of constant-sized correlations — that is, correlations for which the number of measurements and number of measurement outcomes are fixed — such that solving the quantum membership problem for this family is computationally impossible. Thus, the undecidability that arises in understanding Bell experiments is not dependent on varying the number of measurements in the experiment. This places strong constraints on the types of descriptions that can be given for quantum correlation sets. Our proof is based on a combination of techniques from quantum self-testing and undecidability results for linear system nonlocal games.

^{*}h7fu@umd.edu

[†]camiller@umd.edu

[‡]weslofst@uwaterloo.ca

Contents

1	Introduction	2			
	1.1 Paper overview	7			
	1.2 Acknowledgements	7			
2	Notation and group theory background	7			
3	Quantum correlations	12			
4	Quantum correlations and group theory	17			
	4.1 Solution groups and correlations	18			
	4.2 Embedding groups in solution groups	20			
5	Minsky machines and Kharlampovich-Myasnikov-Sapir groups	22			
	5.1 Minsky machines	22			
	5.2 Kharlampovich-Myasnikov-Sapir groups	25			
	5.3 An extension of the Kharlampovich-Myasnikov-Sapir group	27			
	5.4 Embedding KMS groups in solution groups	33			
6	Constant-sized correlations \mathfrak{C}_p for the dihedral groups	34			
7	Membership problems 4				

1 Introduction

Suppose two spatially separated parties, say Alice and Bob, are each able to perform different measurements on their local system. If Alice can perform n_A different measurements, each with m_A outcomes, and Bob can perform n_B different measurements, each with m_B outcomes, then from the point of view of an outside observer, their behaviour is captured by the collection

$$P = \{P(a, b | x, y) : 0 \le a < m_A, 0 \le b < m_B, 0 \le x < n_A, 0 \le y < n_B\}$$

where P(a, b|x, y) is the probability that Alice measures outcome *a* and Bob measures outcome *b*, given that Alice performs measurement *x* and Bob performs measurement *y*. The collection *P* is called a *correlation (matrix)* or *behaviour* [Tsi93]. Colloquially, the size of a correlation is given by the tuple (n_A, n_B, m_A, m_B) .

It is natural to ask which correlations can occur in nature. Suppose measurement x on Alice's system always gives outcome c_x , and measurement y on Bob's system always gives outcome d_y . Then the corresponding correlation is $P(a, b|x, y) = \delta_{a,c_x} \delta_{b,d_y}$, where δ is the Kronecker delta. Correlations of this form are called *deterministic correlations*. The convex hull of the set of deterministic correlations is denoted by $C_c(n_A, n_B, m_A, m_B)$, or C_c when the tuple (n_A, n_B, m_A, m_B) is clear. Correlations in C_c are called *classical correlations*. All deterministic correlations obviously occur in nature, and if Alice and Bob have access to shared randomness, they can also achieve all correlations in C_c . It is a fundamental fact of

quantum mechanics, first observed theoretically by John Bell and now verified in many experiments, that Alice and Bob can achieve correlations outside of C_c by using quantum entanglement [Bel64].

Bell's theorem leads to the question of which correlations can be achieved in quantum mechanics. To study this question, Tsirelson introduced the set of quantum correlations [Tsi93]. There are actually several ways to define the set of quantum correlations, depending on whether we assume that all Hilbert spaces are finite-dimensional, and whether we use the tensor-product axiom or commuting-operator axiom for joint systems. This leads to four different choices for the set of quantum correlations: the finite-dimensional quantum correlations C_q , the quantum-spatial correlations C_{qs} , the quantum-approximate correlations C_{qa} , and the commuting-operator correlations C_{qc} . We use the same convention as for classical correlations, in that C_t refers to $C_t(n_A, n_B, m_A, m_B)$ when the tuple (n_A, n_B, m_A, m_B) is clear. Tsirelson suggested that all four sets should be equal, but we now know that (for large enough n_A, n_B, m_A, m_B) all four sets are different, and hence give a strictly increasing sequence

$$C_c \subsetneq C_q \subsetneq C_{qs} \subsetneq C_{qa} \subsetneq C_{qc}$$

[Slo19, CS20, JNV⁺20, DPP19, Col20]. The last inequality $C_{qa} \subsetneq C_{qc}$ is a very exciting consequence of the recent proof [JNV⁺20] that MIP^{*} = RE by Ji, Natarajan, Vidick, Wright, and Yuen, and following [Fri12, JNP⁺11], this inequality gives a negative resolution to the Connes embedding problem.

As the convex hull of a finite set, C_c is a polytope in \mathbb{R}^N , where $N = n_A n_B m_A m_B$. The sets C_t , $t \in \{q, qs, qa, qc\}$, are also convex subsets of \mathbb{R}^N (in addition, C_{qa} and C_{qc} are closed), but it follows from a result of Tsirelson [Tsi87] that these sets are not polytopes. Following up on this point in [Tsi93, Problem 2.10], Tsirelson asks whether the sets of quantum correlations might still have nice geometric descriptions, specifically by analytic or even polynomial inequalities. This question is significant for two reasons: practical, in that the quantum correlation set captures what is possible with quantum entanglement, and thus a description of this set tells us what is theoretically achievable in experiments and quantum technologies; and conceptual, in that a nice description of the set of quantum correlations could improve our conceptual understanding of quantum entanglement, similarly to how the description of C_c as the convex hull of deterministic correlations is central to our understanding of classical correlations.

Due to the significance of this question, describing the set of quantum correlations has been a central question in the field. On the geometric side, Tsirelson's original results show that when $m_A = m_B = 2$, a certain linear slice of the quantum correlation set is the elliptope, a convex set described by quadratic inequalities ([Tsi87], see also [Lan88, WW01, Mas03, Pit08] for subsequent work on the special case that $n_A = n_B = 2$, and [TVC19] for a description as the elliptope). The convex geometry of $C_q(2, 2, 2, 2)$ is studied in detail in [GKW⁺18]. The case of $C_q(2, 2, 2, 2)$ benefits from a dimension reduction argument: by Jordan's lemma, any correlation in $C_q(2, 2, 2, 2)$ can be expressed as a convex combination of correlations from two-qubit systems. In general, we might ask whether there is a bound on the dimension of Hilbert spaces needed to realize correlations in $C_q(n_A, n_B, m_A, m_B)$. There are several different proofs that, as the number of questions and outcomes increases, there are correlations which require Hilbert spaces of arbitrarily high dimension [Tsi93, BPA⁺08, Slo11]. If we fix n_A , n_B , m_A , m_B such that $C_q(n_A, n_B, m_A, m_B)$ is not closed (which again, happens if n_A , n_B , m_A , m_B are large enough), then $C_q(n_A, n_B, m_A, m_B)$ contains correlations which require Hilbert spaces of arbitrarily large dimension. The first author gives an explicit family of correlations { $P_d | d \ge 1$ } with n_A , n_B , m_A , m_B fixed requiring maximally entangled states of dimension d in [Fu22]. Thus the methods used to study $C_q(2, 2, 2, 2)$ do not work for more measurements or outcomes. Using a different dimension reduction argument, Russell describes another linear slice of C_q , the synchronous correlations, in $C_q(3, 3, 2, 2)$, but again this description does not extend to other numbers of measurements and outcomes [Rus20].

In another line, a number of authors have considered whether it's possible to give a conceptual, rather than geometric, description of the quantum correlation sets. The first result in this line comes from Tsirelson's original definition of quantum correlations, where he observes that quantum correlations belong to the set of nonsignalling correlations, which are those correlations *P* for which the sums

$$\sum_{b} P(a, b|x, y) \text{ and } \sum_{a} P(a, b|x, y)$$

are independent of *y* and *x* respectively. This condition captures the fact that, when spatially separated, Alice and Bob cannot communicate with each other. Since the set of nonsignalling correlations is strictly larger than the commuting-operator correlations C_{qc} , the fact that Alice and Bob cannot communicate does not identify the set of quantum correlations among all correlations. But it is natural to ask whether there might not be additional principles which would suffice to identify the set of correlations. Some examples of conditions which further restrict the set of nonsignalling correlations and which are satisfied by quantum correlations can be found in [BBL+06, PPK+09, NW09, FSA+13, SGAN18], but so far these do not give a complete description of the set of quantum correlations.

Based on the apparent difficulty of describing the set of quantum correlations, there has also been a line of work studying the computational complexity of problems related to these sets. The main line of inquiry, initiated in [CHTW04], has been to consider the difficulty of determining the quantum and commuting-operator values of a nonlocal game. For example, one can consider the problem of determining whether a given nonlocal game has a perfect strategy.

Problem PerfectStrategy_t. Given a tuple of natural numbers (n_A, n_B, m_A, m_B) and a nonlocal game *G* with n_A and n_B questions and m_A and m_B answers, does *G* have a perfect strategy in C_t ?

From the point of view of convex geometry, the quantum (resp. commuting-operator) value of a nonlocal game is the maximum a certain linear functional on the set C_{qa} (resp. C_{qc}). Asking whether such a nonlocal game has a perfect strategy corresponds to asking whether this maximum is equal to 1. Leading up to [JNV⁺20], there was a series of deep works showing that even the approximate version of this optimization problem is indeed very difficult [IV12, RUV13, Ji17, NV18, NW19]. These results have implications

in computational complexity theory, as they imply lower bounds on the complexity class MIP^{*} of multiprover proofs with entangled provers. In the exact (rather than approximate) case, previous results by the last author of the current paper imply that the problems PerfectStrategy_t are undecidable for $t \in \{q, qs, qa, qc\}$ [Slo19, Slo20, FJVY19]. To understand the difficulty of approximating the quantum and commuting-operator values, we can look at a gapped variant of PerfectStrategy_t:

Problem GappedPerfectStrategy_t. Given a tuple of natural numbers (n_A, n_B, m_A, m_B) and a nonlocal game *G* with n_A and n_B questions and m_A and m_B answers, decide whether *G* has a perfect strategy in C_t , or the quantum value of *G* is $\leq 1/2$, given that one of the two is the case.

The result by Ji, Natarajan, Vidick, Wright, and Yuen mentioned above shows that GappedPerfectStrategy_t is also undecidable for $t \in \{q, qs, qa\}$ [JNV⁺20].

Rather than looking at nonlocal games, a more straightforward way to study the difficulty of describing quantum correlation sets is to look at the membership problem for these sets. Specifically, we can look at the decision problems for $t \in \{q, qs, qa, qc\}$ and subfields $\mathbb{K} \subseteq \mathbb{R}$.

Problem Membership_{*t*,**K**}. Given a tuple (n_A, n_B, m_A, m_B) , and a correlation $P \in \mathbb{K}^{n_A n_B m_A m_B}$, is $P \in C_t(n_A, n_B, m_A, m_B)$?

The point of restricting to correlations in $\mathbb{K}^{n_A n_B m_A m_B}$ rather than $\mathbb{R}^{n_A n_B m_A m_B}$ is that it is not possible to describe all real numbers in a finite fashion. We are primarily interested in subfields of \mathbb{R} such as \mathbb{Q} , where it is practical to work with elements of the field on a computer. For our results we actually need to take a larger field than \mathbb{Q} , so in what follows we'll set $\mathbb{K} = \overline{\mathbb{Q}} \cap \mathbb{R}$ unless otherwise noted, where $\overline{\mathbb{Q}}$ is the algebraic closure of the rationals.¹

The problems Membership_{*t*,K} are a very general way of studying descriptions of the sets C_t for $t \in \{q, qs, qa, qc\}$, since we don't restrict to any particular form of description, but instead just look at a basic functionality that we would hope to have from any nice description, namely a way of being able to distinguish elements inside the set from those outside. The decision problems Membership_{*t*,K} are not equivalent to the problems PerfectStrategy_{*t*} or GappedPerfectStrategy_{*t*}, since nonlocal games do not necessarily have unique perfect strategies in C_t . Nonetheless, the two families of decision problems are closely related. Indeed, the methods used in [Slo20] to show the undecidability of PerfectStrategy_{*qc*} are adapted in [CS19] to show the undecidability of Membership_{*qc*,K} [CS19]. The methods of [Slo19] can be adapted to show the undecidability of Membership_{*t*,K} for $t \in \{q, qs, qa\}$ in similar fashion (although some work is needed for the case t = q). The undecidability of GappedPerfectStrategy_{*t*} can be used (in a blackbox fashion, without referring to the proof methods) to get the stronger result that Membership_{*t*,Q} is undecidable for $t \in \{q, qs, qa\}$ [JNV⁺20].

The above undecidability results put very strong restrictions on what descriptions of the quantum correlation sets are possible. For instance, they imply that there is no

¹Since $\overline{\mathbb{Q}}$ is computable, it is possible to work with $\overline{\mathbb{Q}}$ and $\overline{\mathbb{Q}} \cap \mathbb{R}$ on a computer, and indeed support for this is included in Mathematica and other computer algebra packages.

Turing machine which takes tuples (n_A, n_B, m_A, m_B) as inputs, and outputs a description of $C_t(n_A, n_B, m_A, m_B)$ in terms of a finite list of polynomial inequalities, since such a Turing machine would allow us to decide Membership_{t,K} (such a description is sometimes called a uniform polynomial description). Similarly, these results also imply that there can be no finite set of principles, independent of (n_A, n_B, m_A, m_B) , such that we can decide whether a correlation satisfies every principle, and such that a correlation satisfies all the principles if and only if it belongs to $C_t(n_A, n_B, m_A, m_B)$.

However, we note that the reasoning in the last two paragraphs depends crucially on the fact that the parameters (n_A, n_B, m_A, m_B) can vary. The papers [Slo20, Slo19, JNV⁺20] all involve games with unbounded alphabet size. Hence these results leave open the possibility that every set $C_t(n_A, n_B, m_A, m_B)$ has a nice description, but that it is just not possible to have a Turing machine which outputs these descriptions as a function of (n_A, n_B, m_A, m_B) (in other words, they might have a non-uniform description). Thus it is natural to ask what happens to the complexity of Membership_{*t*,K} when (n_A, n_B, m_A, m_B) is held constant. This question motivates our main result.

Problem Membership $(n_A, n_B, m_A, m_B)_{t,\mathbb{K}}$. Given a correlation $P \in \mathbb{K}^{n_A n_B m_A m_B}$, is $P \in C_t(n_A, n_B, m_A, m_B)$?

Theorem 1.1. There is an integer α such that the decision problem Membership $(n_A, n_B, m_A, m_B)_{t,\mathbb{K}}$ is undecidable for $t \in \{qa, qc\}$ and $n_A, n_B, m_A, m_B > \alpha$.

This result shows that the undecidability of membership in C_{qa} and C_{qc} is not only a consequence of varying the size of the correlation, but is in fact embedded into the shape of a single set $C_t(n_A, n_B, m_A, m_B)$ for large enough (n_A, n_B, m_A, m_B) . As a practical consequence, the result shows that there is no description of the set $C_t(n_A, n_B, m_A, m_B)$ (e.g. by polynomial inequalities) that would allow us to decide membership in that set.

As mentioned above, in this theorem \mathbb{K} is the intersection $\mathbb{Q} \cap \mathbb{R}$. However, the proof of this theorem does not rely on writing down very complicated elements of $\overline{\mathbb{Q}}$. In fact, \mathbb{K} could be replaced with $\mathbb{K}_0 \cap \mathbb{R}$, where \mathbb{K}_0 is the subfield of $\overline{\mathbb{Q}}$ generated by roots of unity. In this way, the theorem is similar to the undecidability results for (Membership_{*t*, \mathbb{K}}) that follow from [Slo19, Slo20, CS19]. However, in those results, if the correlations are defined in terms of observables instead of measurements, then it is possible to take $\mathbb{K} = \mathbb{Q}$. In our case, even if we work with correlations defined in terms of observables, we still need to use roots of unity. We also note that the correlations constructed in the proof of Theorem 1.1 are synchronous (see Definition 3.3), so Theorem 1.1 holds for the subsets of synchronous correlations.

It is interesting to also consider upper bounds on the complexity of the problem Membership $(n_A, n_B, m_A, m_B)_{t,\mathbb{K}}$. When t = qc, this problem is contained in coRE, and Theorem 1.1 actually shows that this problem is coRE-complete (for large enough n_A, n_B, m_A, m_B). When t = q or t = qs, this problem is contained in RE, but when t = qa, the best known upper bound on this decision problem is Π_2^0 . In this case, Theorem 1.1 only shows that Membership $(n_A, n_B, m_A, m_B)_{qa,\mathbb{K}}$ is coRE-hard, so this lower bound is not necessarily tight. Recently, Mousavi, Nezhadi, and Yuen have shown that PerfectStrategy_{qa} is Π_2^0 -complete [MNY21], and it seems reasonable to conjecture that Membership $(n_A, n_B, m_A, m_B)_{qa,\mathbb{K}}$ is also Π_2^0 -complete for large enough n_A, n_B, m_A, m_B . We leave this for future research.

1.1 Paper overview

We summarize the technical content of this paper. The starting point for the proof of Theorem 1.1 is the fact that the halting problem for Minsky machines is undecidable [Min67]. Minsky machines, which we review in Section 5, are a model of universal computation similar to Turing machines. To relate Minsky machines to correlations, we go through group theory: specifically the Kharlampovich-Myasnikov-Sapir (KMS) groups [KMS17], also described in Section 5. To construct correlations from these groups, we use the machinery of [Slo19], described in Section 4. Section 2 and Section 3 contain some basic background on group theory and quantum correlations, respectively.

For the proof of Theorem 1.1, we pick a Minsky machine MM with an undecidable halting problem. For each $n \ge 1$, we then write down a finite set of correlations F_n such that $F_n \cap C_{qa} \ne \emptyset$ if MM does not accept n, and $F_n \cap C_{qc} = \emptyset$ otherwise. The correlations in F_n can be thought of as a combination of two subcorrelations. The first subcorrelation is constructed from MM using the method of [Slo19], and is independent of n. The second subcorrelation encodes the input n using the method of [Fu22] to keep the number of measurements and outcomes fixed. This is described in Section 6. In this way, the number of measurements and measurement outcomes for correlations in F_n will depend only on MM, not on n. The proof of Theorem 1.1 is completed in Section 7.

1.2 Acknowledgements

The authors thank Henry Yuen for helpful conversations about the topics of this paper. CAM thanks Johannes Bausch for a conversation about [BCLPG20] which helped to inspire this project. WS is supported by NSERC DG 2018-03968 and an Alfred P. Sloan Research Fellowship. This paper is partly a contribution of the U. S. National Institute of Standards and Technology, and is not subject to copyright in the United States.

2 Notation and group theory background

In this section we give a brief description of some of the notation and group theory concepts we'll use throughout the paper. For basic notation, we denote the set $\{0, 1, ..., n-1\}$ by [n]and the set $\{x \in \mathbb{R} \mid x \ge c\}$ by $\mathbb{R}_{\ge c}$. We index vectors in \mathbb{C}^n starting from 0, so $\mathbb{C}^n = \mathbb{C}^{[n]}$. The *n*-th root of unity is denoted by $\omega_n := e^{i2\pi/n}$. For a Hilbert space \mathcal{H} , we let $\mathcal{L}(\mathcal{H})$ be the set of all bounded linear operators acting on \mathcal{H} , and $\mathcal{U}(\mathcal{H})$ be the group of unitaries acting on \mathcal{H} . We let $\|\cdot\|_{op}$ denote the operator norm on $\mathcal{L}(\mathcal{H})$. For finite-dimensional Hilbert spaces, we also work with the *normalized Hilbert-Schmidt norm*, which is defined by

$$\|M\| = \sqrt{\frac{\operatorname{Tr}(M^{\dagger}M)}{d}}.$$

for $M \in \mathcal{L}(\mathbb{C}^d)$. Note that we don't use any subscript to distinguish this from other norms, as this will be our default norm. We also let $\tilde{Tr}(M)$ be the *normalized trace* Tr(M)/d of M. When working with a group G, we use e for the identity, and let [g, h] be the commutator

 $g^{-1}h^{-1}gh$ of $g,h \in G$. We let g^h denote the conjugate $h^{-1}gh$ of g by h. This notation matches with [KMS17].

If *S* is a set, we let $\mathcal{F}(S)$ be the free group generated by *S*. If *R* is a subset of $\mathcal{F}(S)$, then we let $\langle S : R \rangle$ be the quotient of $\mathcal{F}(S)$ by the normal subgroup generated by *R*. The pair *S*, *R* is called a *presentation* of $\langle S : R \rangle$, and as usual we use $\langle S : R \rangle$ to refer to the presentation and the group defined by the presentation interchangeably. We also will write $\langle S : r_i = t_i, i \in I \rangle$ to mean the presentation $\langle S : \{r_i t_i^{-1}, i \in I\} \rangle$. If *S* and *R* are finite, then $\langle S : R \rangle$ is said to be *finitely-presented*. If $G = \langle S_G : R_G \rangle$, *S* is disjoint from S_G , and *R* is a subset of $\mathcal{F}(S_G \cup S)$, then we sometimes denote the presentation $\langle S_G \cup S : R_G \cup R \rangle$ by $\langle G, S : R \rangle$. An example of a finitely-presented group that we'll use is the *dihedral group*

$$D_n = \langle t_1, t_2 : t_1^2 = t_2^2 = (t_1 t_2)^n = e \rangle$$

This group has order 2*n*, and the elements are $(t_1t_2)^j$ and $t_2(t_1t_2)^j$ for $j \in [n]$.

The *free product* of a group *G* with a group *H* is denoted by G * H. Note that if $G = \langle S_G : R_G \rangle$ and $H = \langle S_H : R_H \rangle$, then $G * H = \langle S_G \cup S_H : R_G \cup R_H \rangle = \langle G, S_H : R_H \rangle$, where the unions of S_G and S_H are disjoint. A more general notion of the free product of groups is *the free product of groups with amalgamation*. Let G_1 and G_2 be two groups with subgroups H_1 and H_2 respectively such that there exists an isomorphism $\phi : H_1 \to H_2$. Then the free product of G_1 and G_2 with amalgamation is defined by $G_1 *_{\phi} G_2 := G_1 * G_2 / \langle h_1 \phi(h_1)^{-1} | h_1 \in H_1 \rangle$.

Another way to construct new groups from a given group is by Higman-Neumann-Neumann extension (*HNN-extension*) [HNN49]. If *H* is a subgroup of *G* and $\phi : H \to H$ is an injective homomorphism, then the HNN-extension of *G* is $\overline{G} = \langle G, t : t^{-1}ht = \phi(h), h \in H \rangle$. By [Rot12, Theorem 11.70], the natural homomorphism sending $g \in G$ to its image in \overline{G} is injective, meaning that we can regard *G* as a subgroup of \overline{G} . We shall introduce other important properties of the free product with amalgamation and the HNN-extension later when they are needed. For more background on these concepts, we refer to [Rot12].

When ϕ : $H \to H$ is an isomorphism of order n, we similarly define the \mathbb{Z}_n -HNN extension of G by $\hat{G} = \langle G, t : t^n = e, t^{-1}ht = \phi(h)$ for $h \in H \rangle$. As in the case of the ordinary HNN-extension, G is embedded in \hat{G} :

Lemma 2.1. Let *G* be a group, *H* a subgroup of *G*, $\phi : H \to H$ an isomorphism of order *n*, and $\hat{G} := \langle G, t : t^n = e, t^{-1}ht = \phi(h)$ for $h \in H \rangle$ the \mathbb{Z}_n -HNN extension. Then the inclusion

$$G \to \hat{G} : g \mapsto g$$

is injective, and t has order n.

Proof. Let G^{*n} denote the free product of G with itself n times, where we index the factors by elements of \mathbb{Z}_n . Let $i_k : G \hookrightarrow G^{*n}$ be the inclusion of the kth factor, $k \in \mathbb{Z}_n$, and let $\psi : G^{*n} \to G^{*n}$ be the cyclic shift, so $\psi(i_k(g)) = \psi(i_{k+1}(g))$ for $k \in \mathbb{Z}_n$.

Let *N* be the normal subgroup of G^{*n} generated by $i_k(h^{-1})i_{k+1}(\phi(h))$ for all $h \in H$ and $0 \le k \le n-2$. Since we only include these relations for $k \le n-2$, we can check (for instance, by looking at presentations) that G^{*n}/N is an iterated amalgamated product of

G with itself *n* times. But observe that *N* also contains

$$\begin{pmatrix} i_0(\phi(h))^{-1})i_1(\phi^2(h)) \end{pmatrix} \left(i_1(\phi^2(h)^{-1})i_2(\phi^3(h)) \right) \cdots \left(i_{n-2}(\phi^{n-1}(h)^{-1})i_{n-1}(\phi^n(h)) \right)$$

= $i_0(\phi(h)^{-1})i_{n-1}(h),$

and hence contains $i_{n-1}(h^{-1})i_0(\phi(h))$. Consequently $\psi(N) = N$, so ψ induces an automorphism $\tilde{\psi}$ of G^{*n}/N . We claim that \hat{G} is isomorphic to $K := G^{*n}/N \rtimes_{\tilde{\psi}} \mathbb{Z}_n$. Indeed, suppose *t* is the generator of \mathbb{Z}_n , so that

$$t^k \cdot x = \tilde{\psi}^k(x) \cdot t^k$$
 for all $x \in G^{*n}/N$.

Then

$$t^{-1}i_0(h)Nt = \tilde{\psi}^{-1}(i_0(h)N) = i_{n-1}(h)N = i_0(\phi(h))N$$

for all $h \in H$, so there is a homomorphism $\alpha : \hat{G} \to K$ sending $g \mapsto i_0(g)N$, $g \in G$, and $t \mapsto t$.

Going the other way, there is a homomorphism $\beta : G^{*n} \to \hat{G}$ sending $i_k(g) \mapsto t^k g t^{-k}$. This homomorphism sends

$$i_k(h^{-1})i_{k+1}(\phi(h)) \mapsto t^k h^{-1} t^{-k} \cdot t^{k+1} \phi(h) t^{-(k+1)} = t^{k+1} (t^{-1} h^{-1} t) \phi(h) t^{-(k+1)} = e,$$

so β descends to a homomorphism $\tilde{\beta} : G^{*n}/N \to \hat{G}$. If $g \in G$ and $a \in \mathbb{Z}_n$ then

$$t^{a}\tilde{\beta}(i_{k}(g)N)t^{-a} = t^{a+k}gt^{-(a+k)} = \tilde{\beta}(i_{a+k}(g)N) = \tilde{\beta}(\tilde{\psi}^{a}(i_{k}(g)N)).$$

We conclude that $\tilde{\beta}(\tilde{\psi}^a(x)) = t^a \tilde{\beta}(x)t^{-a}$ for all $x \in G^{*n}/N$, and hence there is a homomorphism $K \to \hat{G}$ sending $x \in G^{*n}/N$ to $\tilde{\beta}(x)$ and $t \mapsto t$. Since, in particular, this homomorphism sends $i_0(g)N$ to g, it is an inverse to α , proving the claim that \hat{G} and K are isomorphic.

Since G^{*n}/N is an iterated amalgamated free product, the homomorphism $G \to G^{*n}/N$ sending $g \mapsto i_0(g)N$ is injective. Since G^{*n}/N is a subgroup of K, $g \mapsto i_0(g)N$ is still injective when considered as a homomorphism $G \to K$. Composing with the isomorphism $K \cong \hat{G}$, we get that the homomorphism $G \to \hat{G} : g \mapsto g$ is injective. Finally, as the generator of \mathbb{Z}_n , *t* has order *n*.

A unitary representation ρ of a group G on the Hilbert space \mathcal{H} is a homomorphism $\rho : G \to \mathcal{U}(\mathcal{H})$. For any set X, we let $\ell^2 X$ denote the Hilbert space with Hilbert basis $\{|x\rangle : x \in X\}$. The *left regular representation* $L : G \to \mathcal{U}(\ell^2 G)$ of a group G is defined by $L(g)|h\rangle = |gh\rangle$, and right regular representation $R : G \to \mathcal{U}(\ell^2 G)$ is defined by $R(g)|h\rangle = |hg^{-1}\rangle$ for all $g, h \in G$. Note that L(g) and R(g') commute for all $g, g' \in G$.

To construct correlations in C_{qa} (which, recall from the introduction, are limits of finite-dimensional correlations), we use finite-dimensional approximate representations of groups. The norm we use for these approximate representations is the normalized Hilbert-Schmidt norm.

Definition 2.2 (Definition 5 of [Slo19]). Let $G = \langle S : R \rangle$ be a finitely-presented group, and let \mathcal{H} be a finite-dimensional Hilbert space. A finite-dimensional ϵ -approximate representation of G is a homomorphism $\phi : \mathcal{F}(S) \to \mathcal{U}(\mathcal{H})$ such that $\|\phi(r) - \mathbb{1}\| \leq \epsilon$ for all $r \in R$.

An element $g \in G = \langle S : R \rangle$ represented by a word $w \in \mathcal{F}(S)$ is *nontrivial in approximate representations* of *G* if there exists some $\delta > 0$ such that, for all $\epsilon > 0$, there is an ϵ -approximate representation $\phi : \mathcal{F}(S) \to \mathcal{U}(\mathcal{H})$ such that $\|\phi(w) - \mathbb{1}\| \ge \delta$ (this does not depend on the choice of word w). Otherwise we say that *g* is *trivial in approximate representations*. If *F* is a finite subset of elements which are non-trivial in approximate representations, then we can find ϵ -representations where all the elements of *F* are bounded away from the identity, and in fact we can do this by having the trace of all the elements of *F* be close to zero:

Proposition 2.3. Let $G = \langle S : R \rangle$ and W be a finite subset of $\mathcal{F}(S)$. Then, for every $\epsilon, \zeta > 0$, there is an ϵ -approximate representation ϕ such that for all $w \in W$,

- 1. *if* w *is trivial in approximate representations of* G*, then* $1 \zeta \leq \tilde{Tr}(\phi(w)) \leq 1$ *, and*
- 2. *if* w *is nontrivial in approximate representations of* G*, then* $0 \leq \tilde{Tr}(\phi(w)) \leq \zeta$ *.*

The proof of the proposition above is very similar to the proof of [Slo19, Lemma 12], so we omit it here. The next two well-known lemmas are useful when working with approximate representations.

Lemma 2.4. Let $G = \langle S : R \rangle$ be a finitely presented group. Suppose $f \in \mathbb{C}[\mathcal{F}(S)]$. Then there is a constant c (depending on f) such that for any ϵ -approximate representation $\phi : \mathcal{F}(S) \to \mathcal{U}(\mathcal{H})$, $\|\phi(f)\|_{op} \leq c$. Furthermore, if f = 0 in $\mathbb{C}[G]$, then there is a constant c' (also depending on f) such that for any ϵ -approximate representation $\phi : \mathcal{F}(S) \to \mathcal{U}(\mathcal{H}), \|\phi(f)\| \leq c' \epsilon$.

Proof. Suppose $f = \sum_{i \in [k]} a_i u_i$ where $a_i \in \mathbb{C}$ and $u_i \in \mathcal{F}(S)$. Then

$$\|\phi(f)\|_{op} \le \sum_{i \in [k]} |a_i| \|\phi(u_i)\|_{op} = \sum_{i \in [k]} |a_i|.$$

If f = 0 in $\mathbb{C}[G]$, we can write

$$f = \sum_{i \in [k]} b_i x_i (e - r_i) y_i,$$

where $b_i \in \mathbb{C}$, $x_i, y_i \in \mathcal{F}(S)$ and $r_i \in R$. Then

$$\|\phi(f)\| \le \sum_{i \in [k]} |b_i| \|\phi(x_i)(1 - \phi(r_i))\phi(y_i)\| = \sum_{i \in [k]} |b_i| \|1 - \phi(r_i)\| \le \sum_{i \in [k]} |b_i|\epsilon,$$

by the unitary invariance of the normalized Hilbert-Schmidt norm. The lemma follows from taking $c = \sum_{i \in [k]} |a_i|$ and $c' = \sum_{i \in [k]} |b_i|$.

Lemma 2.5. There is a nondecreasing function $\Delta : \mathbb{R}_{\geq 1} \times \mathbb{N} \to \mathbb{R}_{\geq 1}$ such that if $\{P_i \mid i \in [n]\} \subset \mathcal{L}(\mathbb{C}^d)$ is a set of matrices such that

$$\|P_i\|_{op} \le c, \quad \|P_i^2 - P_i\| \le \epsilon, \quad \|P_i^* - P_i\| \le \epsilon, \quad \|P_iP_j\| \le \epsilon, \text{ and } \quad \|\sum_{k \in [n]} P_k - \mathbb{1}\| \le \epsilon$$

for all $i, j \in [n]$, $i \neq j$ and some $c \in \mathbb{R}$, then there is a projective measurement $\{\Pi_i \mid i \in [n]\} \subset \mathcal{L}(\mathbb{C}^d)$ such that $\|\Pi_i - P_i\| \leq \Delta(c, n)\epsilon$ for all $i \in [n]$.

Proof. When c = 1 and P_i is positive for all $i \in [n]$, this is shown in Lemma 3.5 of [KPS18], with function $\Delta(1, n) = \Delta_{pos}(n)$, where $\Delta_{pos}(n)$ is defined recursively by $\Delta_{pos}(n+1) = (40n+3)\Delta_{pos}(n)$ and $\Delta_{pos}(1) = 2\sqrt{2}$. To reduce to this case, suppose that $\{P_i \mid i \in [n]\}$ satisfy the conditions of the lemma. If $\{Q_i \mid i \in [n]\} \subseteq \mathcal{L}(\mathbb{C}^d)$ are self-adjoint operators such that $\|Q_i\|_{op} \leq c'$ and $\|P_i - Q_i\| \leq \delta$ for all $i \in [n]$, then

$$\begin{aligned} \|Q_i^2 - Q_i\| &\leq \|Q_i(Q_i - P_i)\| + \|(Q_i - P_i)P_i\| + \|Q_i - P_i\| + \|P_i^2 - P_i\| \\ &\leq \|Q_i\|_{op}\|Q_i - P_i\| + \|P_i\|_{op}\|Q_i - P_i\| + \|Q_i - P_i\| + \|P_i^2 - P_i\| \\ &\leq (c + c' + 1)\delta + \epsilon. \end{aligned}$$

Similarly, $||Q_iQ_j|| \le (c+c')\delta + \epsilon$ and $||\sum_{i\in[n]}Q_i - 1|| \le n\delta + \epsilon$.

If we take $Q_i = (P_i + P_i^*)/2$, then Q_i is self-adjoint, $||Q_i||_{op} \le c$ and $||Q_i - P_i|| \le \epsilon/2$. Let $\chi_{[1/2,\infty)}$ be the indicator function of the interval $[1/2,\infty)$. Since $|\chi_{[1/2,\infty)}(t) - t| \le 2|t^2 - t|$ for all $t \in \mathbb{R}$,

$$\|\chi_{[1/2,\infty)}(Q_i) - Q_i\| \le 2\|Q_i^2 - Q_i\| \le (2c+3)\epsilon.$$

Hence the self-adjoint projections $Q'_i = \chi_{[1/2,\infty)}(Q_i)$ satisfy the conditions

$$\|Q_i'Q_j'\| \le (c+1)(2c+3)\epsilon + \|Q_iQ_j\| \le (2c^2 + 7c + 4)\epsilon \text{ and} \\\|\sum_{i\in[n]} Q_i' - 1\| \le (2c+3)n\epsilon + \|\sum_{i\in[n]} Q_i - 1\| \le (2c+7/2)n\epsilon + \epsilon.$$

Applying Lemma 3.5 of [KPS18] to $\{Q'_i \mid i \in [n]\}$ yields $\{\Pi_i \mid i \in [n]\}$ such that

$$\|\Pi_i - Q'_i\| \le \Delta_{pos}(n)(2c^2 + 7c + 5)n\epsilon.$$

Since

$$\|\Pi_i - P_i\| \le \|\Pi_i - Q_i'\| + \|Q_i' - Q_i\| + \|Q_i - P_i\| \le \Delta_{pos}(n)(2c^2 + 7c + 5)n\epsilon + (2c + 3)\epsilon + 1/2\epsilon,$$

the lemma is true with $\Delta(c, n) = \Delta_{pos}(n)(2c^2 + 7c + 5)n + 2c + 4$.

There are variants of Lemma 2.5 that reduce the dependence on *n* (see, e.g., [dlS21]). In this paper, *c* and *n* are fixed, so $\Delta(c, n)$ is a constant.

By the definition of the normalized Hilbert-Schmidt norm, the set of elements of *G* that are trivial in finite-dimensional approximate representations forms a normal subgroup of *G*, denoted by N^{fa} . For a group *G*, we define

$$G^{fa} := G/N^{fa}.$$

If ϕ : $G \to H$ is a homomorphism between finitely-presented groups and $x \in G$ is trivial in approximate representations of G, then $\phi(x)$ is trivial in approximate representations of H, so there is an induced homomorphism $G^{fa} \to H^{fa}$.

Definition 2.6 (Definition 14 of [Slo19]). For finitely-presented groups G and H, a homomorphism $\phi : G \to H$ is an **fa-embedding** if the induced map: $G^{fa} \to H^{fa}$ is injective.

In other words ϕ : $G \rightarrow H$ is an fa-embedding if whenever x is non-trivial in approximate representations of G, then $\phi(x)$ is non-trivial in approximate representations of H.

A finitely-presented group *G* is said to be *hyperlinear* if every non-trivial element is non-trivial in approximate representations. Although we've defined hyperlinearity only for finitely-presented groups, whether a group is hyperlinear is independent of the presentation. To show that groups are hyperlinear, we use the stronger properties of solvability, amenability, and soficity. Recall that a group *G* is *solvable* if it has subgroups $G_0 = \{e\}, G_1, \ldots, G_{k-1}$ and $G_k = G$ such that G_{j-1} is normal in G_j and G_j/G_{j-1} is an abelian group, for $1 \le j \le k$. For the purposes of our paper, the definitions of amenable and sofic groups are irrelevant; we just need the following well-known properties of these classes of groups (see [CLP15, Proposition 2.4.1]):

- 1. Solvable groups are amenable, amenable groups are sofic, and sofic groups are hyperlinear.
- 2. If *H* is an amenable subgroup of a sofic group *G*, and $\alpha : H \to H$ is an injective homomorphism, then the HNN-extension of *G* by α is sofic.
- 3. If H_1 and H_2 are amenable subgroups of sofic groups G_1 and G_2 , and $\alpha : H_1 \to H_2$ is an isomorphism, then the free product of G_1 and G_2 with amalgamation, $G_1 *_{\alpha} G_2$, is sofic.
- 4. If *N* is a normal subgroup of *G* such that *N* is sofic and *G*/*N* is amenable, then *G* is sofic.

To these properties we can add:

Lemma 2.7. If *H* is an amenable subgroup of a sofic group *G*, and ϕ : $H \rightarrow H$ is an isomorphism of order *n*, then the \mathbb{Z}_n -HNN-extension of *G* by ϕ is sofic.

Proof. We continue with the notation from the proof of Lemma 2.1. As G^{*n}/N is an iterated almagamated free product over the amenable group H, G^{*n}/N is hyperlinear. Since $K/(G^{*n}/N) \cong \mathbb{Z}_n$ and \mathbb{Z}_n is amenable, K is sofic.

3 Quantum correlations

In this section, we now introduce our main object of study. Consider a scenario with two parties or players, Alice and Bob, and a referee. The referee chooses questions to send to Alice and Bob from finite sets \mathcal{X} and \mathcal{Y} respectively, and they return answers from finite sets \mathcal{A} and \mathcal{B} . As mentioned in the introduction, this can also be thought of as a scenario in which Alice and Bob perform measurements labelled by the elements of \mathcal{X} and \mathcal{Y} , and

receive outcomes from A and B respectively. Alice and Bob's behaviour in this scenario can be described by the function

$$P: \mathcal{A} \times \mathcal{B} \times \mathcal{X} \times \mathcal{Y} \to \mathbb{R}_{>0}: (k, \ell, i, j) \mapsto P(k, \ell | i, j),$$

where $P(k, \ell | i, j)$ is the probability of answers $(k, \ell) \in \mathcal{A} \times \mathcal{B}$ with questions $(i, j) \in \mathcal{X} \times \mathcal{Y}$. We call a tuple $(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ of finite sets a *nonlocal scenario*, and a function $P : \mathcal{A} \times \mathcal{B} \times \mathcal{X} \times \mathcal{Y} \to \mathbb{R}_{>0}$ such that

$$\sum_{k \in \mathcal{A}, \ell \in \mathcal{B}} P(k, \ell | i, j) = 1$$

for all $i \in \mathcal{X}, j \in \mathcal{Y}$ a *bipartite correlation* for the scenario $(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$.

With quantum correlations, we want to capture what Alice and Bob can do in a nonlocal scenario when they cannot communicate. Even though Alice and Bob cannot communicate, the rules of quantum mechanics do allow them to share entanglement. We can visualize this scenario as in Figure 1.



Figure 1: A nonlocal test between Alice and Bob

Recall that a vector state in a Hilbert space \mathcal{H} is a unit vector, and a projective measurement (with *m* outcomes) is a collection $\{P^{(i)} \mid i \in [m]\}$ of orthogonal projections on \mathcal{H} such that

$$\sum_{i \in [m]} P^{(i)} = \mathbb{1}$$

There are two ways in quantum mechanics to handle the restriction that Alice and Bob cannot communicate. In the first, we require that their joint Hilbert space be a tensor product of subspaces:

Definition 3.1. *A bipartite correlation P for the scenario* $(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ *is a quantum correlation if there are*

- (a) finite-dimensional Hilbert spaces \mathcal{H}_A and \mathcal{H}_B ,
- (b) a vector state $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$,
- (c) a collection of projective measurements $\{M_i^{(k)} | k \in A\}$ on \mathcal{H}_A for every $i \in \mathcal{X}$, and
- (d) a collection of projective measurements $\{N_j^{(\ell)} \mid \ell \in \mathcal{B}\}$ on \mathcal{H}_B for every $j \in \mathcal{Y}$,

such that

$$P(k, \ell | i, j) = \langle \psi | M_i^{(k)} \otimes N_j^{(\ell)} | \psi \rangle$$

for all $i \in X$, $j \in Y$, $k \in A$, $\ell \in B$.

The set $C_q(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ is the set of all quantum correlations for scenario $(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$, and the set $C_{qa}(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ is the closure of $C_q(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ in $\mathbb{R}^{\mathcal{A} \times \mathcal{B} \times \mathcal{X} \times \mathcal{Y}}$. If n_A, n_B, m_A , and m_B are positive integers, and $t \in \{q, qa\}$, then we set $C_t(n_A, n_B, m_A, m_B) := C_t([n_A], [n_B], [m_A], [m_B])$.

If $|\mathcal{X}| = n_A$, $|\mathcal{Y}| = n_B$, $|\mathcal{A}| = m_A$, and $|\mathcal{B}| = m_B$, then $\mathbb{R}^{\mathcal{A} \times \mathcal{B} \times \mathcal{X} \times \mathcal{Y}} \cong \mathbb{R}^{m_A m_B n_A n_B}$, and this is used to define the closure of $C_q(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$. In fact, this isomorphism identifies $C_t(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ with $C_t(n_A, n_B, m_A, m_B)$ for $t \in \{q, qa\}$. So although it's convenient to be able to use arbitrary labels for questions and answers (and we'll use non-integer labels in this paper), we could just work with the sets $C_t(n_A, n_B, m_A, m_B)$ if we wanted.

Some additional terminology we'll use: a collection of Hilbert spaces, vector state, and projective measurements as in parts (a)-(d) is called a *quantum strategy*. Also, although we don't study this set in this paper, the set $C_{qs}(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ of *quantum-spatial correlations* is defined similarly to the set of quantum correlations, but the Hilbert spaces \mathcal{H}_A and \mathcal{H}_B in the strategy are allowed to be infinite-dimensional. The closure of C_{qs} is also equal to C_{qa} [SW08].

Moving on, a more general way to handle the restriction that Alice and Bob cannot communicate is to drop the requirement that Alice and Bob's projective measurements act on different Hilbert spaces, and instead just require that their projective measurements commute:

Definition 3.2. A bipartite correlation P for a scenario $(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ commuting-operator correlation if there is

- (a) a Hilbert space \mathcal{H} ,
- (b) a vector state $|\psi\rangle \in \mathcal{H}$,
- (c) a collection of projective measurements $\{M_i^{(k)} | k \in A\}$ on \mathcal{H} for every $i \in \mathcal{X}$, and

(d) a collection of projective measurements $\{N_i^{(\ell)} \mid \ell \in \mathcal{B}\}$ on \mathcal{H} for every $j \in \mathcal{Y}$,

such that

$$M_i^{(k)} N_j^{(\ell)} = N_j^{(\ell)} M_i^{(k)}$$

and

$$P(k,\ell|i,j) = \langle \psi | M_i^{(k)} \cdot N_j^{(\ell)} | \psi \rangle$$

for all $i \in \mathcal{X}$, $j \in \mathcal{Y}$, $k \in \mathcal{A}$, $\ell \in \mathcal{B}$.

The set $C_{qc}(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ is the set of all commuting-operator correlations for the scenario $(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$, and if n_A , n_B , m_A , and m_B are positive integers, then $C_{qc}(n_A, n_B, m_A, m_B) := C_{qc}([n_A], [n_B], [m_A], [m_B])$.

As with quantum correlations, we refer to a Hilbert space, vector state, and projective measurements as in (a)-(d) as a *commuting-operator strategy*. Note that the Hilbert space \mathcal{H} in a commuting-operator strategy does not have to be finite-dimensional.

We define one more subtype of correlation that we'll use:

Definition 3.3. A bipartite correlation P for scenario $(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ is synchronous if $\mathcal{X} = \mathcal{Y}$, $\mathcal{A} = \mathcal{B}$, and

$$\sum_{k \in \mathcal{A}} P(k, k | i, i) = 1$$

for all $i \in \mathcal{X}$.

Equivalently, a correlation with $\mathcal{X} = \mathcal{Y}$ and $\mathcal{A} = \mathcal{B}$ is synchronous if $P(k, \ell | i, i) = 0$ for all $k \neq \ell$ and *i*, or in other words if Alice and Bob always return the same answer when given the same question. The following fact about synchronous correlations is well-known:

Proposition 3.4 (Theorem 5.5(i) in [PSS⁺16]). Let *P* be a synchronous correlation for $(\mathcal{X}, \mathcal{X}, \mathcal{A}, \mathcal{A})$, and let $\mathcal{H}, |\psi\rangle, \{M_i^{(k)} | k \in \mathcal{A}\}, \{N_i^{(\ell)} | \ell \in \mathcal{A}\}$ be a commuting operator strategy for *P*. Then

$$M_i^{(k)}|\psi
angle=N_i^{(k)}|\psi
angle$$

for all $i \in \mathcal{X}$, $k \in \mathcal{A}$.

Proposition 3.4 is an immediate consequence of the following lemma, which is contained in the proof of [PSS⁺16, Theorem 5.5(i)].

Lemma 3.5. Let $|\psi\rangle \in \mathcal{H}$ be a quantum state, and $\{M_j \mid j \in [n]\}$ and $\{N_j \mid j \in [n]\}$ be two projective measurements on \mathcal{H} for some $n \geq 2$, such that $M_jN_k = N_kM_j$ for all $j,k \in [n]$. If $\langle \psi | M_jN_k | \psi \rangle = 0$ for all $j \neq k \in [n]$, then

$$M_j|\psi
angle=N_j|\psi
angle$$

for each $j \in [n]$.

The next two conclusions of Lemma 3.5 will help us work with correlations.

Lemma 3.6. Let $|\psi\rangle \in \mathcal{H}$ be a quantum state, and let $\{M_0^{(k)} | k \in [m_A]\}$ and $\{M_1^{(k)} | k \in [m_A]\}$ be two projective measurements on \mathcal{H} , both of which commute with the projective measurement $\{N^{(l,l')} | l, l' \in [m_A]\}$ on \mathcal{H} . If

$$\langle \psi | M_0^{(k)} N^{(l,l')} | \psi \rangle = \langle \psi | M_1^{(k')} N^{(l,l')} | \psi \rangle = 0$$

for any $k \neq l$ and $k' \neq l'$, then

$$M_0^{(k)} M_1^{(k')} |\psi\rangle = M_1^{(k')} M_0^{(k)} |\psi\rangle$$

for any $k, k' \in [m_A]$.

Proof. The condition implies that the two measurement pairs

$$\{M_0^{(k)} \mid k \in [m_A]\}, \{\sum_{l' \in [m_A]} N^{(k,l')} \mid k \in [m_A]\}$$

and

$$\{M_1^{(k')} \mid k' \in [m_A]\}, \{\sum_{l \in [m_A]} N^{(l,k')} \mid k' \in [m_A]\}$$

both satisfy the condition of Lemma 3.5 with respect to $|\psi\rangle$. We conclude that

$$egin{aligned} M_0^{(k)}|\psi
angle &= \sum\limits_{l'\in[m_A]} N^{(k,l')}|\psi
angle ext{ and } \ M_1^{(k')}|\psi
angle &= \sum\limits_{l\in[m_A]} N^{(l,k')}|\psi
angle \end{aligned}$$

for each $k, k' \in [m_A]$. Then,

$$\begin{split} M_0^{(k)} M_1^{(k')} |\psi\rangle &= M_0^{(k)} \sum_{l \in [m_A]} N^{(l,k')} |\psi\rangle = \sum_{l \in [m_A]} N^{(l,k')} M_0^{(k)} |\psi\rangle \\ &= \sum_{l \in [m_A]} N^{(l,k')} \sum_{l' \in [m_A]} N^{(k,l')} |\psi\rangle = N^{(k,k')} |\psi\rangle = \sum_{l' \in [m_A]} N^{(l',k)} \sum_{l \in [m_A]} N^{(l,k')} |\psi\rangle \\ &= M_1^{(k')} \sum_{l' \in [m_A]} N^{(l',k)} |\psi\rangle = M_1^{(k')} M_0^{(k)} |\psi\rangle, \end{split}$$

for each $k, k' \in [m_A]$.

Lemma 3.7 (Substitution Lemma). Let $|\psi\rangle \in \mathcal{H}$ be a quantum state. Suppose there exist sequences of unitaries $\{V\}$, $\{V_i \mid i \in [k]\}$ and $\{M_i \mid i \in [n]\}$ on \mathcal{H} commuting with another sequence of unitaries $\{N_i \mid i \in [n]\}$ on \mathcal{H} , such that

$$M_i |\psi\rangle = N_i |\psi\rangle$$

for each $i \in [n]$ *, and*

$$egin{aligned} V|\psi
angle &= \prod_{i\in [k]} V_i |\psi
angle. \end{aligned}$$

Then

$$V\prod_{i\in[n]}M_i|\psi
angle = \left(\prod_{i\in[k]}V_i\right)\left(\prod_{i\in[n]}M_i\right)|\psi
angle.$$

Proof. We prove this lemma by induction on *n*. The n = 0 case follows the condition that $V|\psi\rangle = \prod_{i \in [k]} V_i |\psi\rangle$. Assume the conclusion holds for j = m and consider the case

j = m + 1. Then

$$V \prod_{i \in [m+1]} M_i |\psi\rangle = V \left(\prod_{i \in [m]} M_i\right) M_m |\psi\rangle = V \left(\prod_{i \in [m]} M_i\right) N_m |\psi\rangle$$
$$= N_m V \left(\prod_{i \in [m]} M_i\right) |\psi\rangle = N_m \left(\prod_{i \in [k]} V_i\right) \left(\prod_{i \in [m]} M_i\right) |\psi\rangle$$
$$= \left(\prod_{i \in [k]} V_i\right) \left(\prod_{i \in [m]} M_i\right) N_m |\psi\rangle = \left(\prod_{i \in [k]} V_i\right) \left(\prod_{i \in [m+1]} M_i\right) |\psi\rangle,$$

which completes the proof.

When working with correlations, it sometimes simplifies arguments if we restrict to strategies with the following property:

Definition 3.8. A commuting operator strategy

$$|\psi\rangle \in \mathcal{H}, \{M_i^{(k)} \mid k \in \mathcal{A}\}, i \in \mathcal{X}, \{N_j^{(\ell)} \mid \ell \in \mathcal{B}\}, j \in \mathcal{Y}$$

is good if

(a) for all $i \in \mathcal{X}$ and $k \in \mathcal{A}$, if $\langle \psi | M_i^{(k)} | \psi \rangle = 0$, then $M_i^{(k)} = 0$, and (b) for all $j \in \mathcal{Y}$ and $\ell \in \mathcal{B}$, if $\langle \psi | N_i^{(\ell)} | \psi \rangle = 0$, then $N_i^{(\ell)} = 0$.

Proposition 3.9. *If* $P \in C_{qc}(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B})$ *, then* P *has a good commuting-operator strategy. Proof.* Suppose $(|\psi\rangle, \{M_i^{(k)}\}, \{N_i^{(\ell)}\})$ is a strategy for *P*. If $P(k.\ell \mid i, j) \neq 0$ then

$$\langle \psi | M_i^{(k)} | \psi \rangle = \sum_{\ell' \in \mathcal{B}} P(k, \ell' \mid i, j) \neq 0.$$

If $\langle \psi | M_i^{(k)} | \psi \rangle = 0$, then $\langle \psi | M_i^{(k)} N_i^{(\ell)} | \psi \rangle = P(k, \ell \mid i, j) = 0$ for all $j \in \mathcal{Y}$ and $\ell \in \mathcal{B}$. Since $\sum_{k' \in \mathcal{A}} M_i^{(k')} = \mathbb{1}$, there must be some k' such that $\langle \psi | M_i^{(k')} | \psi \rangle \neq 0$. If we replace $M_i^{(k')}$ with $M_i^{(k')} + M_i^{(k)}$ and $M_i^{(k)}$ with 0, we get another commuting-operator strategy for *P*, and doing this for all $k \in A$, $i \in \mathcal{X}$ with $\langle \psi | M_i^{(k)} | \psi \rangle = 0$, and similarly for all $\ell \in \mathcal{B}$, $j \in \mathcal{Y}$ with $\langle \psi | N_i^{(\ell)} | \psi \rangle = 0$, gives a good strategy for *P*.

Quantum correlations and group theory 4

In this section, we introduce the notion of a perfect correlation associated with a binary linear system. We then recall the notion of a solution group associated to a linear system, and use the solution group to show that every linear system has a perfect correlation. Finally we recall how to embed an arbitrary finitely presented group in a solution group.

4.1 Solution groups and correlations

Definition 4.1. Let Ax = 0 be an $m \times n$ binary linear system, so A is an $m \times n$ matrix over \mathbb{Z}_2 , and $0 \in \mathbb{Z}_2^n$. Suppose that each row of A has κ non-zero entries. For each $i \in [m]$, let

$$I_i = \{ j \in [n] \mid A_{ij} = 1 \}$$

and let $\phi_i : I_i \to [\kappa]$ be the unique order-preserving bijection (so the smallest element of I_i maps to 0, the largest maps to $\kappa - 1$, and so on). Let $\mathcal{X}_{var} := \{x_i \mid i \in [n]\}$, and let $\mathcal{X} := [m] \cup \mathcal{X}_{var}$. Let $S := \{v \in \mathbb{Z}_2^{\kappa} : \sum_{i \in [\kappa]} v_i = 0\}$. A correlation P for the scenario $(\mathcal{X}, \mathcal{X}, \mathbb{Z}_2^{\kappa}, \mathbb{Z}_2^{\kappa})$ is a **perfect correlation for** Ax = 0 if P(a, b|x, y) = 0 whenever

- (1) $x \in [m]$ and $a \notin S$ or $y \in [m]$ and $b \notin S$;
- (2) $x \in \mathcal{X}_{var}$ and $(a_0, \ldots a_{\kappa-2}) \neq (0, 0, \ldots, 0)$, or $y \in \mathcal{X}_{var}$ and $(b_0, \ldots b_{\kappa-2}) \neq (0, 0, \ldots, 0)$;
- (3) $x, y \in [m]$ and $a_{\phi_x(k)} \neq b_{\phi_y(k)}$ for some $k \in I_x \cap I_y$;
- (4) $x \in [m]$, $y = x_i \in \mathcal{X}_{var}$ for some $i \in I_x$, and $a_{\phi_x(i)} \neq b_{\kappa-1}$ in \mathbb{Z}_2 ;
- (5) $x = x_i \in \mathcal{X}_{var}$ for some $i \in I_y$, $y \in [m]$, and $a_{\kappa-1} \neq b_{\phi_y(i)}$ in \mathbb{Z}_2 ; or

(6)
$$x = y \in \mathcal{X}_{var}$$
, and $a_{\kappa-1} \neq b_{\kappa-1}$.

In perfect correlations as defined here, the questions that Alice and Bob receive are either variables in \mathcal{X}_{var} , or indices from [m]. When Alice or Bob gets an index $x \in [m]$, condition (1) requires them to return an element $a \in S$. This element should be thought of as an assignment to the *x*th equation, where variable x_k , $k \in I_x$, receives value $a_{\phi_x(k)}$. When Alice or Bob gets a variable x_i in \mathcal{X}_{var} , condition (2) forces them to return an element $a \in \mathbb{Z}_2^{\kappa}$ with $a_j = 0$ for $0 \le j \le \kappa - 2$ and $a_{\kappa-1} \in \{0, 1\}$. This should be thought of as an assignment to x_i from \mathbb{Z}_2 . The remaining conditions state that if Alice and Bob are asked about the same variable (either as part of an equation or directly) then their assignments to that variable must agree. Note that conditions (3) and (6) imply that every perfect correlation for Ax = 0 is synchronous.

Definition 4.1 is stated for linear systems with a constant number κ of non-zero entries in each row. This allows us to use answer sets \mathbb{Z}_2^{κ} . It is possible to define perfect correlations for systems with a varying number of entries in each row, by either using answer sets which vary with the question, or by using larger answer sets \mathbb{Z}_2^n as in [KPS18]. However, all the linear systems we work with have a constant number of non-zero entries in each row, and making this assumption in Definition 4.1 simplifies later analysis.

The point of perfect correlations is that any strategy for a perfect correlation yields a representation of a certain group associated with Ax = 0.

Definition 4.2 (Definition 17 of [Slo19]). Let Ax = 0 be an $m \times n$ linear system over \mathbb{Z}_2 , so A is an $m \times n$ matrix with entries in \mathbb{Z}_2 and $0 \in \mathbb{Z}_2^n$. For $j \in [m]$, define $I_j = \{k \in [n] \mid A_{jk} = 1\}$.

The homogeneous solution group of Ax = 0 *is*

$$\Gamma(A) := \langle x_0, x_1, \dots x_{n-1} : x_j^2 = e \text{ for all } j \in [n],$$
$$\prod_{k \in I_i} x_k = e \text{ for all } i \in [m],$$
$$[x_j, x_k] = e \text{ if } j, k \in I_i \text{ for some } i \rangle$$

Proposition 4.3. Let Ax = 0 be a binary linear system with κ non-zero entries in each row. Suppose *P* is a perfect correlation for Ax = 0, and that \mathcal{H} , $|\psi\rangle$, $\{M_x^{(a)} \mid a \in \mathbb{Z}_2^{\kappa}\}$, $x \in \mathcal{X}$, $\{N_y^{(b)} \mid b \in \mathbb{Z}_2^{\kappa}\}$, $y \in \mathcal{X}$ is a good commuting-operator strategy for *P*. Let $\mathcal{H}_0 := \overline{\mathcal{A} \cdot |\psi\rangle}$, the closure of $\mathcal{A}|\psi\rangle$ in \mathcal{H} , where \mathcal{A} is the algebra generated by $M_x^{(a)}$ and $N_x^{(a)}$ for $x \in \mathcal{X}$, $a \in \mathbb{Z}_2^{\kappa}$. For $i \in [n]$, let

$$M(x_i) := M_{x_i}^{(0,\dots,0,0)} - M_{x_i}^{(0,\dots,0,1)} \text{ and } N(x_i) := N_{x_i}^{(0,\dots,0,0)} - N_{x_i}^{(0,\dots,0,1)}.$$

Then

- (1) $M(x_i)$ and $N(x_i)$ are binary observables such that $M(x_i)|\psi\rangle = N(x_i)|\psi\rangle$ for all $i \in [n]$,
- (2) there are unitary representations Φ_M and Φ_N of $\Gamma(A)$ on \mathcal{H}_0 sending $x_i \in \Gamma(A)$ to $M(x_i)|_{\mathcal{H}_0}$ and $N(x_i)|_{\mathcal{H}_0}$ respectively, and
- (3) if $\Phi_M(r)|\psi\rangle = |\psi\rangle$ for some $r \in \Gamma(A)$, then $\Phi_M(r) = \mathbb{1}_{\mathcal{H}_0}$.

Proposition 4.3 is similar to [CLS17, Lemma 8]. We include a full proof for completeness.

Proof of Proposition 4.3. Since perfect correlations are synchronous, $M_x^{(a)}|\psi\rangle = N_x^{(a)}|\psi\rangle$ for all $x \in \mathcal{X}$, $a \in \mathbb{Z}_2^{\kappa}$ by Proposition 3.4. As a result, we have $M(x_i)|\psi\rangle = N(x_i)|\psi\rangle$ for all $i \in [n]$. For the second part of the proposition, define projections

$$M_{i,k}^{(c)} := \sum_{\substack{a \in \mathbb{Z}_{2}^{\kappa} \\ a_{\phi_{i}(k)} = c}} M_{i}^{(a)}, i \in [m], k \in I_{i}, c \in \mathbb{Z}_{2}.$$

Observe that by condition (2) of Definition 4.1, if $i \in [n]$ and $b \in \mathbb{Z}_2^{\kappa}$ with $(b_0, \ldots, b_{\kappa-2}) \neq (0, \ldots, 0)$, then

$$\langle \psi | N_{x_i}^{(b)} | \psi \rangle = \sum_{a \in \mathbb{Z}_2^{\kappa}} \langle \psi | M_1^{(a)} N_{x_i}^{(b)} | \psi \rangle = \sum_{a \in \mathbb{Z}_2^{\kappa}} P(a, b | 1, x_i) = 0,$$

and hence $N_{x_i}^{(b)}|\psi\rangle = 0$. Since the strategy is a good strategy, by Definition 3.8, $N_{x_i}^{(b)} = 0$ for $(b_0, \ldots, b_{\kappa-2}) \neq (0, \ldots, 0)$. Thus $\{N_{x_i}^{(0)}, N_{x_i}^{(1)}\}$ is a complete measurement, where for convenience we write $N_{x_i}^{(c)}$ for $N_{x_i}^{(0,\ldots,0,c)}$ for all $i \in [n]$ and $c \in \mathbb{Z}_2$. Then $N(x_i) =$ $N_{x_i}^{(0)} - N_{x_i}^{(1)}$ is a binary observable on \mathcal{H} . Similarly, condition (1) of Definition 4.1 and Definition 3.8 imply that $M_i^{(a)} = 0$ for all $i \in [m]$ and $a \notin S$. By condition (4) of Definition 4.1,

$$\langle \psi | M_{i,k}^{(0)} \cdot N_{x_k}^{(1)} | \psi \rangle = \sum_{\substack{a \in \mathbb{Z}_2^{\kappa} \\ a_{\phi_i(k)} = 0}} \langle \psi | M_i^{(a)} N_{x_k}^{(1)} | \psi \rangle = \sum_{\substack{a,b \in \mathbb{Z}_2^{\kappa} \\ a_{\phi_i(k)} = 0}} P(a, (0, \dots, 0, 1) | i, x_k) = 0,$$

and $\langle \psi | M_{i,k}^{(1)} \cdot N_{x_k}^{(0)} | \psi \rangle = 0$ similarly. Applying Lemma 3.5 to the projective measurements $\{M_{i,k}^{(0)}, M_{i,k}^{(1)}\}$ and $\{N_{x_k}^{(0)}, N_{x_k}^{(1)}\}$, we get that $M_{i,k}^{(c)} | \psi \rangle = N_{x_k}^{(c)} | \psi \rangle$ for all $i \in [m]$, $k \in I_i$, and $c \in \mathbb{Z}_2$.

Now let $M_{i,k} := M_{i,k}^{(0)} - M_{i,k}^{(1)}$ for some $i \in [m]$. Since the projections $M_i^{(a)}$, $a \in \mathbb{Z}_2^{\kappa}$ commute, $[M_{i,k}, M_{i,l}] = 0$ for all $i \in [m]$ and $k, l \in I_i$. If $k_0, \ldots, k_{\ell-1}$ is a sequence in I_i , and σ is a permutation of $[\ell]$, then

$$\begin{split} N(x_{k_0})\cdots N(x_{k_{\ell-1}})|\psi\rangle &= M_{i,k_{\ell-1}}M_{i,k_{\ell-2}}\cdots M_{i,k_0}|\psi\rangle = M_{i,k_{\sigma(\ell-1)}}\cdots M_{i,k_{\sigma(0)}}|\psi\rangle \\ &= N(x_{\sigma(k_0)})\cdots N(x_{\sigma(k_{\ell-1})})|\psi\rangle, \end{split}$$

so the operators $N(x_k)$, $k \in I_i$ commute on $|\psi\rangle$. When we take the product across all of I_i ,

$$\begin{split} \prod_{k \in I_i} N(x_k) |\psi\rangle &= \prod_{k \in I_i} M_{i,k} |\psi\rangle = \prod_{k \in I_i} \left(\sum_{a \in \mathbb{Z}_2^{\kappa}} (-1)^{a_{\phi_i(k)}} M_i^{(a)} \right) |\psi\rangle = \sum_{a \in \mathbb{Z}_2^{\kappa}} (-1)^{\sum_{k \in [\kappa]} a_k} M_i^{(a)} |\psi\rangle \\ &= \sum_{a \in S} M_i^{(a)} |\psi\rangle = \sum_{a \in \mathbb{Z}_2^{\kappa}} M_i^{(a)} |\psi\rangle = |\psi\rangle, \end{split}$$

where we use that $M_i^{(a)} = 0$ for $a \notin S$ and $\sum_k a_k = 0$ for $a \in S$. Finally, $N(x_k)^2 = N_{x_k}^{(0)} + N_{x_k}^{(1)} = \mathbb{1}_{\mathcal{H}}$.

To finish the proof, let \mathcal{A}_0 (resp. \mathcal{A}_1) be the algebra generated by $M_x^{(a)}$ (resp. $N_x^{(a)}$) for $x \in \mathcal{X}$ and $a \in \mathbb{Z}_2^{\kappa}$. A standard result about synchronous correlations, following immediately from the fact that $M_x^{(a)}|\psi\rangle = N_x^{(a)}|\psi\rangle$, is that $\mathcal{A}|\psi\rangle = \mathcal{A}_0|\psi\rangle = \mathcal{A}_1|\psi\rangle$. If $R \in \mathcal{A}_1$ satisfies $R|\psi\rangle = 0$, then $RT|\psi\rangle = TR|\psi\rangle = 0$ for all $T \in \mathcal{A}_0$, and thus Rv = 0for all $v \in \mathcal{H}_0 = \overline{\mathcal{A}}|\psi\rangle = \overline{\mathcal{A}}_0|\psi\rangle$. Define $\Phi_N : \mathcal{F}(x_1, \ldots, x_n) \to U(\mathcal{H}_0) : x_i \mapsto N(x_i)|_{\mathcal{H}_0}$, and suppose $r \in \mathcal{F}(x_1, \ldots, x_n)$ is a defining relation for $\Gamma(A)$ from Definition 4.2. We've shown that $\Phi_N(r)|\psi\rangle = |\psi\rangle$, and hence $\mathbb{1} - \Phi_N(r)$ is 0 on \mathcal{H}_0 . It follows that Φ_N induces a representation of $\Gamma(A)$ on \mathcal{H}_0 sending $x_i \mapsto N(x_i)|_{\mathcal{H}_0}$. Switching M and N in the above argument, we see that there is also a representation of $\Gamma(A)$ on \mathcal{H}_0 sending $x_i \mapsto M(x_i)|_{\mathcal{H}_0}$.

We can also construct perfect correlations for Ax = 0 from representations of the solution group $\Gamma(A)$. We do this for particular representations in Section 7.

4.2 Embedding groups in solution groups

To embed KMS groups into solution groups, we use the embedding results of [Slo19]. Recall the following technical definition:

Definition 4.4 (Definition 32 of [Slo19]). Let A be an $m \times n$ matrix over \mathbb{Z}_2 , $C_0 \subseteq [n] \times [n] \times [n]$, $C_1 \subseteq [\ell] \times [n] \times [n]$ for some $\ell \geq 1$, and L be an $\ell \times \ell$ lower-triangular matrix with non-negative integer entries. Let

$$E\Gamma(A, C_0, C_1, L) := \langle \Gamma(A), y_0, \dots, y_{\ell-1} : x_i x_j x_i = x_k \text{ for all } (i, j, k) \in C_0, y_i^{-1} x_j y_i = x_k \text{ for all } (i, j, k) \in C_1, y_i^{-1} y_j y_i = y_j^{L_{ij}} \text{ for all } i > j \text{ with } L_{ij} > 0 \rangle.$$

We say a group G is an $m \times n \times \ell$ *extended homogeneous-linear-plus-conjugacy group if it has a presentation of this form.*

By [Slo19, Propositions 27 and 33], extended homogenous-linear-plus-conjugacy groups can be *fa*-embedded into solution groups. We use the following version of that result:

Proposition 4.5. Let $G = E\Gamma(A, C_0, C_1, L)$ be an $m \times n \times \ell$ extended homogeneous linear-plusconjugacy group. Then there is an $m' \times n'$ matrix A' over \mathbb{Z}_2 for some $m' \ge m$ and $n' \ge n$, and a homomorphism

$$\phi: \mathcal{F}(x_0,\ldots,x_{n-1},y_0,\ldots,y_{\ell-1}) \to \mathcal{F}(x_0,\ldots,x_{n'-1}),$$

such that:

- (a) Each row in A' has only three non-zero entries.
- (b) $\widetilde{\phi}(x_i) = x_i$ for all $i \in [n]$.
- (c) For all $i \in [\ell]$ there are $j, k \in [n']$ such that $\widetilde{\phi}(y_i) = x_j x_k$.
- (d) If r is a defining relation of G, then $\tilde{\phi}(r)$ is in the normal subgroup generated by the defining relations of $\Gamma(A')$, so $\tilde{\phi}$ induces a homomorphism $\phi : G \to \Gamma(A')$.
- (e) There are integers n_1, n_2, n_3 such that if γ is an ϵ -representation of G, then there is an $O(\epsilon)$ -representation α of Γ with

$$\alpha(\widetilde{\phi}(g)) = \gamma(g)^{\oplus n_1} \oplus \overline{\gamma(g)}^{\oplus n_2} \oplus \mathbb{1}_{n_3}$$

for all $g \in \mathcal{F}(x_0, ..., x_{n-1}, y_0, ..., y_{\ell-1})$. As a result, the homomorphism $\phi : G \to \Gamma(A')$ is an fa-embedding.

Furthermore, both A' and $\tilde{\phi}$ are constructible, in the sense that there's a Turing machine which, given A, C₀, C₁, and L, will output A' and $\tilde{\phi}(y_i)$ for all $i \in [\ell]$.

Although we don't need this fact, A' and $\tilde{\phi}$ can be constructed in polynomial time.

Proof. Parts (b)-(e) follow from the proofs of Propositions 27 and 33 in [Slo19]. Proposition 27 in [Slo19] is actually about non-homogeneous solution groups $\Gamma(A, b)$, which have an additional central element *J* of order 2 representing a scalar in \mathbb{Z}_2 . However, the non-homogeneous solution group $\Gamma(A, b)$ associated with an $m \times n$ linear system Ax = b can

be regarded as a homogeneous solution group by adding variables x_n, \ldots, x_{2n} , replacing J with x_{2n} wherever it occurs in the presentation, and adding linear relations $x_i x_{n+i} x_{2n} = e$ for all $0 \le i < n$ to force x_{2n} to be central. Alternatively, the proof of Proposition 27 can be adapted to the homogeneous case by replacing J with e wherever it occurs in group presentations, and ignoring assignments to J in approximate representations.

For part (a), given an $m_0 \times n_0$ matrix A_0 over \mathbb{Z}_2 , we can find an $m_1 \times n_1$ matrix A_1 , where $m_1 \ge m_0$ and $n_1 \ge n_0$, such that there is an isomorphism $\Gamma(A_0) \to \Gamma(A_1)$ sending $x_i \mapsto x_i$ for all $i \in [n_0]$, and A_1 has exactly three non-zero entries in each row. Indeed, suppose the *i*th row has non-zero entries in columns j_1, \ldots, j_r , where r > 3. Adding variables z_{1t}, z_{2t}, z_3 and equations $z_3 = x_{j_1}x_{j_2}, z_{1t}x_{j_1}x_{j_t} = e$, and $z_{2t}x_{j_2}x_{j_t} = e$ for all $t = 3, \ldots, r$, and replacing the *i*th row of A with the equation $z_3x_{j_3} \cdots x_{j_r} = e$, we get an isomorphic solution group where equation *i* has the number of non-zero entries reduced by one, and all the added equations have length exactly three (the equations $z_{ij}x_{j_i}x_{j_i}$ are needed to force x_{j_i} and x_{j_i} to commute). If row *i* of A_0 has exactly one non-zero entry in column *j*, then we can add variables z_1, z_2, z_3 and replace the *i*th equation with equations $x_j z_1 z_2 = x_j z_1 z_3 = x_j z_2 z_3 = z_1 z_2 z_3 = e$ (which together force $x_j = z_1 = z_2 = z_3 = e$). If row *i* has exactly two non-zero entries in columns *j* and *k*, then we can add variables z_1, z_2 , and replace the *i*th equation with equations $x_j z_1 z_2 = x_k z_1 z_2 = e$. Iterating these steps, we eventually get A_1 as desired.

5 Minsky machines and Kharlampovich-Myasnikov-Sapir groups

5.1 Minsky machines

A *k-glass Minsky Machine* [Min67], denoted by MM, consists of *k* glasses where each glass can hold arbitrarily many coins, a set of states [N], and a finite list of commands. Just like a Turing machine, a configuration of MM describes which state the machine is in and how many coins are in each of the glasses. A computation running on MM is a sequence of commands, where each command maps one configuration to another. Commands can leave a glass unchanged, add a coin to a glass, or remove a coin from a non-empty glass, as well as change the state. Glasses are numbered starting from 1. In formal language, this means that a *configuration of* MM is an element $(i; n_1, n_2, ..., n_k) \in [N] \times (\mathbb{Z}_{\geq 0})^{[k]}$. The state 0 is regarded as a final halt state, and 1 is regarded as a start state. The *accept configuration* is (0; 0, 0, ...0) and the *starting configuration* with input *m* is (1; m, 0, ...0). There are four types of commands:

1. Adding coins: When the state is *i*, add a coin to each of the glasses numbered $j_1, j_2 \dots j_\ell$ where $\ell \le k$, and go to state *j*. This command is encoded as

$$i; \rightarrow j; Add(j_1, j_2 \dots j_\ell).$$

2. Removing coins: When the state is *i*, if the glasses numbered $j_1, j_2 \dots j_\ell$, $\ell \le k$, are all nonempty, then remove a coin from each of the glasses numbered $j_1, j_2 \dots j_\ell$, and go

to state *j*. This command is encoded as

$$i; n_{j_1} > 0, \ldots, n_{j_\ell} > 0 \rightarrow j; Sub(j_1, j_2, \ldots, j_\ell).$$

3. Empty check: When the state is *i*, if the glasses numbered $j_1, j_2 \dots j_\ell$, $\ell \le k$, are empty, go to state *j*. This command is encoded as

$$i; n_{j_1} = 0, n_{j_2} = 0, \dots, n_{j_{\ell}} = 0 \quad \to \quad j.$$

4. Stop: When the state is *i*, change state to 0. This command is encoded as

$$i; \rightarrow 0.$$

In addition, the input state *i* to each command must be non-zero, so there are no commands leaving the halt state. As can be seen from the descriptions, commands can change a configuration, but not every command can be applied to every configuration. Some configurations may not have any applicable commands, while some configurations can have more than one applicable command. If for every configuration, there is at most one command that can be applied, the Minsky machine is *deterministic*. Otherwise, the Minsky machine is *non-deterministic*. A *k*-glass Minsky machine MM *accepts* an input $n \in \mathbb{Z}_{\geq 0}$ if there is a sequence of configurations

$$(1; n, 0, \dots, 0) =: C_0 \to C_1 \to \dots \to C_N := (0; 0, \dots, 0)$$

from the input configuration to the accept configuration such that for every $0 \le i < N$, there is a command of MM that applies to C_i and transforms it to C_{i+1} . In general, we let \equiv_{MM} be the equivalence relation on configurations generated by the relations $C \equiv_{MM} C'$, where C and C' are configurations for which there is a command of MM which applies to Cand transforms it to C'. Note that the equivalence relation generated by these relations also includes the relations $C' \equiv_{MM} C$ whenever there is a command transforming C to C', as well as the transitive closure of both types of relations. So $C \equiv_{MM} C'$ does not necessarily mean that C can be transformed to C' by applying commands from MM, but rather that C can be transformed to C' by applying commands or the inverses of commands. If MM is deterministic, then it's easy to see that $C_1 \equiv_{MM} C_2$ if and only if C_1 and C_2 can both be transformed to the same configuration C_3 by the operations of MM. Since there are no commands with input state 0, a deterministic Minsky machine MM accepts an input n if and only if $(1; n, 0, \ldots, 0) \equiv_{MM} (0; 0, \ldots, 0)$.

Recall that a subset *S* of natural numbers is *recursively enumerable*, or RE, if there is an algorithm such that the algorithm halts on input *s* if and only if $s \in S$. Minsky machines can recognize any recursively enumerable set (or in other words, Minsky machines are Turing complete):

Theorem 5.1 (Theorem 2.7, part (a) of [KMS17]). *X* is recursively enumerable if and only if there exists a 3-glass deterministic Minsky machine MM such that $n \in X$ if and only if MM accepts *n*.

Proof. In Theorem 2.7, part (a) of [KMS17], the theorem is stated for 2-glass Minsky machines with a different encoding of the input. However, the statement above for 3-glass Minsky machines is used as part of the proof of the 2-glass case.

If we change the input encoding, then 2-glass Minsky machines are also sufficiently powerful to recognize all RE sets. However, we use 3-glass Minsky machines as the starting point for our construction, since the simpler input encoding is necessary to connect with other parts of the construction.

Lemma 5.2. Let MM be a k-glass deterministic Minsky machine. Then there is a (k + 1)-glass deterministic Minsky machine MM' such that

- (a) n is accepted by MM' if and only if n is accepted by MM, and
- (b) if n is not accepted by MM, then $(1; n, 0, ..., 0) \equiv_{MM'} (1; m_1, ..., m_{k+1})$ if and only if $(n, 0, ..., 0) = (m_1, ..., m_{k+1})$.

Proof. To construct MM', we add six new states 0', 1', 2', 3', 4', 5' and 6' to MM. We keep the commands of MM the same, except that we replace states 0 and 1 with 0' and 1' wherever they occur in commands. We then add commands

To explain these commands, suppose we start in configuration $(1; m_1, ..., m_{k+1})$ for MM'. We are allowed to move onto state 2' if and only if $m_2 = ... = m_{k+1} = 0$, or in other words we are in an input configuration (1; n, 0, ..., 0). In states 2' and 3', we copy the 1st glass to the 2nd and k + 1th glass. Once the configuration is (2'; 0, n, 0, ..., 0, n), we can move on to state 4', where we copy back the 2nd glass to the 1st glass to get configuration (4'; n, 0, ..., 0, n). At this point, we can move on to state 1', where we can now start applying commands from MM. Note that commands from MM do not affect the k + 1th glass. If input *n* is accepted by MM, then at some point MM' will end up in configuration (0'; 0, ..., 0, n). At this point the state changes to 6', and we empty out the k + 1th glass and then move to the accept configuration (0; 0, ..., 0).

Since 0 is not an allowed input state for commands of MM, the commands coming from MM in MM' do not share input states with any of the added commands above, so MM' is deterministic. The explanation of the commands above shows that if MM accepts

n, then so does MM'. If MM does not accept *n*, then MM' will never reach configuration (0'; 0, ..., 0, n), and thus cannot proceed to state 0, so MM' does not accept *n* either.

Finally, suppose $c_1 \equiv_{MM'} c_2$, where $c_1 := (1; n, 0, ..., 0)$ and $c_2 := (1; m_1, ..., m_{k+1})$, and n is not accepted by MM'. Since MM' is deterministic, this means that MM' will take (1; n, 0, ..., 0) and $(1; m_1, ..., m_{k+1})$ to a common configuration $c_3 = (s; p_1, ..., p_{k+1})$. If s = 1 then $c_1 = c_3 = c_2$. Otherwise $m_2 = ... = m_{k+1} = 0$. If $s \in \{1', 2', 3', 4', 5'\}$ then MM' will send c_1 and c_2 to $(1'; n, 0, ..., 0, n) = (1'; m_1, 0, ..., 0; m_1)$. Since MM does not accept n, we cannot have $s \in \{6', 0\}$. Since commands from MM do not change the k + 1th glass of MM', if $s \notin \{1', 2', 3', 4', 5'\}$ then we must have $n = p_{k+1} = m_1$. In each case, we conclude that $(n, 0, ..., 0) = (m_1, m_2, ..., m_{k+1})$.

5.2 Kharlampovich-Myasnikov-Sapir groups

For a *k*-glass Minsky machine MM, deterministic or non-deterministic, the *Kharlampovich-Myasnikov-Sapir (KMS) group G*(MM) is a finitely presented group defined in [KMS17], based on an earlier construction of Kharlampovich [Kha82]. The definition of this group depends on a parameter *p*, which we always take to be 2. We let *S*(MM) and *R*(MM) denote the generating set and relations for the presentation given in [KMS17, Section 4.1].

The point of KMS groups is the following theorem:

Theorem 5.3 (Properties 3.1 and 3.2 and Theorem 4.3 of [KMS17]). Let MM be a Minsky machine. Then G(MM) is solvable, and there is a computable function w from configurations c of MM to words w(c) in the free group $\mathcal{F}(S(MM))$, such that

$$w(c) = w(c')$$
 in $G(MM)$ if and only if $c \equiv_{MM} c'$.

In particular, if w(n) := w((1; n, 0, ..., 0)) is the word for input configuration and $w_{accept} := w((0; 0, ..., 0))$ is the word for the accept configuration, then

 $w(n) = w_{accept}$ in G(MM) if and only if $(1; n, 0, ..., 0) \equiv_{MM} (0; 0, ..., 0)$.

If MM is deterministic, then we can replace this last condition with the condition that MM accepts *n*.

For our purposes, we need some details of the definition of G(MM) and the function w (we also include some additional details for context). We use the notation from [KMS17] for ease of reference. Suppose MM has *k*-glasses and state set [N + 1]. The generating set S(MM) is divided into subsets L_0 , L_1 , and L_2 , where

$$L_0 = \{x(q_i A_{i_1} \cdots A_{i_m}) \mid i \in [N+1], 0 \le m \le k, 0 \le i_1 < i_2 < \dots < i_m \le k\},\$$

$$L_1 = \{A_i \mid 0 \le i \le k\}, \text{ and }$$

$$L_2 = \{a_i, a'_i, \tilde{a}_i, \tilde{a}'_i \mid 1 \le i \le k\}.$$

Intuitively, the generators x(u) are used to keep track of state, the generators A_i represent the bottom of the *i*th glass, and the generators a_i are used to keep track of coins in the *i*th glass.

The relations R(MM) include a number of relations (marked as (G1)-(G7) in [KMS17, Section 4.1]) which are common to all *k*-glass Minsky machines with state set [N + 1], and then a number of relations (marked as (G8) in the same section) for the commands. For the purpose of discussion, we'll call these *common relations* and *command relations*. The common relations include relations specifying that the elements of L_0 and L_1 have order two, and that [x, y] = e for all $x, y \in L_i$, i = 0, 1, 2. The order of the generators in L_2 is unspecified. The common relations also have the property that if MM' has state set [N' + 1]with $N \leq N'$, then the common relations of R(MM) are common relations of R(MM').

For the command relations, there is one relation for each command. To specify these relations, the following notation is used: if $f \in \mathcal{F}(S(MM))$ and $1 \le j \le k$, let

$$f \circledast a_j := f^{-1} f^{a_j} (f^{-1})^{a_j^{-1}} f^{a_j^{\prime-1}},$$

and

$$f \circledast A_j := [f, A_j]$$

(recall that $f^a := a^{-1}fa$). Also, let

$$t_1 \circledast t_2 \ldots \circledast t_m := (\ldots (t_1 \circledast t_2) \circledast \ldots) \circledast t_m \text{ and } t_1 \circledast t_2^{\circledast n} := t_1 \circledast \underbrace{t_2 \circledast \ldots \circledast t_2}_{\text{n times}}.$$

Then with this notation, the relation for i; \rightarrow j; $Add(j_1, \dots, j_\ell)$ is

 $x(q_iA_0) = x(q_jA_0) \circledast a_{j_1} \circledast \cdots \circledast a_{j_\ell},$

the relation for $i; n_{j_1} > 0, \dots, n_{j_\ell} > 0 \rightarrow j; Sub(j_1, \dots, j_\ell)$ is

$$x(q_iA_0) \circledast a_{j_1} \circledast \cdots \circledast a_{j_\ell} = x(q_jA_0),$$

the relation for $i; n_{j_1} = 0, n_{j_2} = 0, \dots, n_{j_\ell} = 0 \rightarrow j$ is

$$x(q_iA_0) \circledast A_{j_1} \circledast \cdots \circledast A_{j_\ell} = x(q_jA_0) \circledast A_{j_1} \circledast \cdots \circledast A_{j_\ell},$$

and the relation for $i \rightarrow 0$ is

$$x(q_i A_0) = x(q_0 A_0).$$

For $n \in \mathbb{Z}_{\geq 0}$, the word corresponding to the input configuration for *n* is

$$w(n) := x(q_1 A_0) \circledast a_1^{\circledast n} \circledast A_1 \circledast \ldots \circledast A_k.$$

In particular, $w(0) := x(q_1A_0) \circledast A_1 \circledast \ldots \circledast A_k$. For the accept configuration, the group element is

$$w_{accept} := x(q_0 A_0) \circledast A_1 \circledast \ldots \circledast A_k.$$

By [KMS17, Relations (G5a) and (G1)], we can actually reduce these two words to

$$w(0) = x(q_1A_0A_1...A_k)$$
 and $w_{accept} = x(q_0A_0A_1...A_k)$.

In particular, $w_{accept}^2 = w(0)^2 = [w(0), w_{accept}] = e$.

Suppose we start with a *k*-glass Minsky machine MM_0 , and add states and commands to form a *k*-glass Minsky machine MM_1 . Then the generating set $S(MM_0) \subseteq S(MM_1)$. As previously mentioned, all the common relations for $R(MM_0)$ belong to $R(MM_1)$, and since all the commands of MM_0 are commands of MM_1 , the same is true for the command relations. Hence, we also have $R(MM_0) \subseteq R(MM_1)$. This leads immediately to the following lemma:

Lemma 5.4. Let MM_0 and MM_1 be k-glass Minsky machines with state sets $[N_0]$ and $[N_1]$ respectively, where $N_0 \leq N_1$. If every command of MM_0 is a command of MM_1 , then there is a homomorphism $G(MM_0) \rightarrow G(MM_1)$ sending $x \mapsto x$ for all $x \in S(MM_0) \subseteq S(MM_1)$. In particular, this homomorphism sends the elements w(0) and w_{accept} for $G(MM_0)$ to the same elements for $G(MM_1)$.

We finish this section by stating how KMS groups connect to correlations:

Lemma 5.5 (Lemma 42 of [Slo19]). Let MM be a Minsky machine, and let $S(MM) = L_0 \cup L_1 \cup L_2$ be the partition described above. Then the KMS group G(MM) has a presentation as an $m \times n \times \ell$ extended homogeneous linear-plus-conjugacy group $E\Gamma(A, C_0, C_1, L)$, in which:

- (a) the generators in L_0 and L_1 and the elements $x(q_1A_0) \circledast a_1$ and $w(0)w_{accept}$ all belong to the generating set $\{x_0, \ldots, x_{n-1}\}$, and
- (b) the generators in L_2 belong to $\{y_0, \ldots, y_{\ell-1}\}$.

Proof. By Lemma 42 of [Slo19], G(MM) has a presentation as an $m' \times n' \times \ell$ extended homogeneous linear-plus-conjugacy group, in which $x(q_1A_0) \circledast a_1$ belongs to the generating set $\{x_1, \ldots, x_{n'}\}$ (in the notation of [Slo19], $x(q_1A_0) \circledast a_1$ belongs to $\mathcal{N}(L_0, S)$, which is abelian [Slo19, Theorem 40, part (c)]). Looking at the proof of Lemma 42, this presentation is constructed by starting with the generators for G(MM), and then adding additional generators so that all the relations for G(MM) can be rewritten as either linear or conjugacy relations. The generators from L_0 and L_1 end up in the generating set $\{x_1, \ldots, x_{n'}\}$ for the constructed presentation, and the generators L_2 end up in the generating set $\{y_1, \ldots, y_\ell\}$.

The element $w(0)w_{accept}$ also belongs to $\mathcal{N}(L_0, S)$, and hence we can also add additional generators and relations to include this in the generating set of our presentation, following the produre detailed in the proof of Lemma 42. Alternatively, since $w(0) = x(q_1A_0A_1...A_k)$ and $w_{accept} = x(q_0A_0A_1...A_k)$ are already generators in the presentation, and w(0) and w_{accept} commute, we can also just add one additional generator $x_{n'+1}$, along with the linear relation $x_{n'+1}w(0)w_{accept} = e$, to get a presentation with $w(0)w_{accept}$ in the generating set as required.

5.3 An extension of the Kharlampovich-Myasnikov-Sapir group

This section is devoted to proving the following proposition.

Proposition 5.6. Let p(n), $n \ge 1$ be an increasing sequence of prime numbers, where the function $p : \mathbb{Z}_{\ge 1} \to \mathbb{Z}_{\ge 1}$ is computable, let X be a recursively enumerable set of positive

integers, and let *r* be a positive integer which is coprime to p(n) for all $n \ge 1$. Then there exists an $m \times n' \times \ell$ extended homogeneous linear-plus-conjugacy group $H = E\Gamma(A, C_0, C_1, L)$ and generators $x \in \{x_0, ..., x_{n'-1}\}$ and $u, t \in \{y_0, ..., y_{\ell-1}\}$ satisfying the following properties:

- (a) $u^{-1}tu = t^r$ in H,
- (b) $H/\langle t^{p(n)} = e \rangle$ is sofic for all $n \ge 1$,

(c)
$$x = e$$
 in $H/\langle t^{p(n)} = e \rangle$ if and only if $n \in X$, and

(d) t has order p(n) in $H/\langle t^{p(n)} = e \rangle$.

The function of the different generators x, t, and u will be explained in the next section. We prove Proposition 5.6 in a number of steps, starting with:

Lemma 5.7. Let p(n), $n \ge 1$ be an increasing sequence of prime numbers, where the function $p : \mathbb{Z}_{\ge 1} \to \mathbb{Z}_{\ge 1}$ is computable, and let X be a recursively enumerable set of positive integers. Then the set

$$P_X := \{ p(n) \mid n \in X \}$$

is recursively enumerable.

Proof. Let A_X be the Turing machine that accepts $x \in \mathbb{N}$ if and only if $x \in X$. Consider the Turing machine which does the following: Given $q \in \mathbb{N}$, it computes p(n) for all $n \leq q$. If q = p(k) for some $k \leq q$, then it runs A_X on k and accept if A_X accepts. If $q \neq p(k)$ for all $k \leq q$, then it runs forever without accepting. Since p(n) is an increasing sequence, if q = p(k) for some $k \in \mathbb{Z}_{\geq 1}$ then $k \leq q$. As a result, this algorithm accepts q if and only if q = p(k) for $k \in X$.

Let \mathbf{MM}_0 be a 3-glass deterministic Minsky machine that accepts $q \in \mathbb{N}$ if and only if $q \in P_X$ (where P_X is the set from Lemma 5.7), and let \mathbf{MM} be a 4-glass deterministic Minksy machine satisfying parts (a) and (b) of Lemma 5.2 with respect to $\mathbf{MM} = \mathbf{MM}_0$. Let $G(\mathbf{MM}) = \langle S(\mathbf{MM}) : R(\mathbf{MM}) \rangle$ be the KMS group of \mathbf{MM} . Although $w(p(n)) = w_{accept}$ in $G(\mathbf{MM})$ if and only if $n \in X$, for the group H in Proposition 5.6 we need elements x and twhich do not depend on n, such that x = e in $H/\langle t^{p(n)} = e \rangle$ if and only if $n \in X$. To get these elements, we'll start with the group

$$\overline{G(\mathsf{MM})} := \langle G(\mathsf{MM}), t : [t, a_1] = [t, a_1'] = e, t^{-1} x(q_1 A_0) t = x(q_1 A_0) \circledast a_1 \rangle,$$
(1)

where a_1 , a'_1 , and $x(q_1A_0)$ are the generators of G(MM) described above. We'll then construct *H* at the end of the subsection by adding *u* to $\overline{G(MM)}$. For our construction, we also want to consider two other groups

$$G_{p(n)}(\mathsf{M}\mathsf{M}) := \langle G(\mathsf{M}\mathsf{M}) : x(q_1A_0) \circledast a_1^{\circledast p(n)} = x(q_1A_0) \rangle \text{ and}$$

$$\overline{G_{p(n)}(\mathsf{M}\mathsf{M})} := \langle \overline{G(\mathsf{M}\mathsf{M})} : x(q_1A_0) \circledast a_1^{\circledast p(n)} = x(q_1A_0), t^{p(n)} = e \rangle,$$

defined for every $n \ge 1$. The group $\overline{G_{p(n)}(\mathsf{MM})}$ is the quotient of $\overline{G(\mathsf{MM})}$ by $\langle t^{p(n)} = e \rangle$. To show this, we need to explain the definition of $\overline{G(\mathsf{MM})}$ a little more:

Lemma 5.8. Let MM be a Minsky machine, and let K be the subgroup of G(MM) generated by $x(q_1A_0)$, a_1 , and a'_1 . Then there is a homomorphism $\alpha : K \to K$ sending $a_1 \mapsto a_1, a'_1 \mapsto a'_1$, and $x(q_1A_0) \mapsto x(q_1A_0) \circledast a_1$.

Proof. As we've already noted, the generators of *K* satisfy the relations $x(q_1A_0)^2 = [a_1, a'_1] = e$. Let

$$\psi: \mathcal{F}(S(\mathsf{MM})) \to \langle b_1, b_2: [b_1, b_2] = e \rangle = \mathbb{Z} \times \mathbb{Z}$$

be the homomorphism defined by $\psi(a_1) = b_1$, $\psi(a'_1) = b_2$, and $\psi(s) = e$ for all $s \in S(MM) \setminus \{a_1, a'_1\}$. Checking the relations in [KMS17], we see that $\psi(r) = e$ for all $r \in R(MM)$. Hence ψ descends to a homomorphism $G(MM) \to \mathbb{Z} \times \mathbb{Z}$, and this homomorphism restricts to a surjective homomorphism $\psi : K \to \mathbb{Z} \times \mathbb{Z}$. Let $\langle x(q_1A_0) \rangle^K$ be the normal subgroup generated by $x(q_1A_0)$ in K. Since $\psi(x(q_1A_0)) = e$, this normal subgroup is contained in the kernel of ψ , and hence there is a surjective homomorphism $\mathbb{Z} \times \mathbb{Z} \to K$ sending $b_1 \mapsto a_1$ and $b_2 \mapsto a'_1$. Hence $K / \langle x(q_1A_0) \rangle^K \cong \mathbb{Z} \times \mathbb{Z}$, and thus $\langle x(q_1A_0) \rangle^K$ is the kernel of ψ in K. We conclude that $K \cong \langle x(q_1A_0) \rangle^K \rtimes (\mathbb{Z} \times \mathbb{Z})$, and in particular every element of K can be written uniquely as $ta_1^n(a'_1)^m$ for some $t \in \langle x(q_1A_0) \rangle^K$ and $n, m \in \mathbb{Z}$.

By [KMS17, Lemma 4.1], $\langle x(q_1A_0) \rangle^K$ is abelian. Hence the functions $\langle x(q_1A_0) \rangle^K \rightarrow \langle x(q_1A_0) \rangle^K$ sending $f \mapsto f^{-1}$, $f \mapsto f^{a_1}$, $f \mapsto (f^{-1})^{a_1}$, and $f \mapsto f^{(a'_1)^{-1}}$ are all homomorphisms, and so we conclude that

$$\langle x(q_1A_0)\rangle^K \to \langle x(q_1A_0)\rangle^K : f \mapsto f \circledast a_1$$

is a homomorphism. Using the relation $[a_1, a'_1] = e$ again, we also see that

$$f^{a_1^n} \circledast a_1 = (f \circledast a_1)^{a_1^n}$$
 and $f^{(a_1')^m} \circledast a_1 = (f \circledast a_1)^{(a_1')^m}$

for all $n, m \in \mathbb{Z}$. Using the fact that $K \cong \langle x(q_1A_0) \rangle^K \rtimes (\mathbb{Z} \times \mathbb{Z})$, we see that there is a homomorphism $\alpha : K \to K$ sending $a_1 \mapsto a_1, a'_1 \mapsto a'_1$, and $f \in \langle x(q_1A_0) \rangle^K$ to $f \circledast a_1$ as desired.

Let *K* and α be the group and homomorphism from Lemma 5.8, with MM = **MM**. The presentation of $\overline{G(\mathbf{MM})}$ states that $t^{-1}ft = \alpha(f)$ for $f \in \{a_1, a'_1, x(q_1A_0)\}$, so this identity holds for all $f \in K$. In other words, $\overline{G(\mathbf{MM})}$ is analogous to the HNN-extension of $G(\mathbf{MM})$ by α , although strictly speaking we do not know if $\overline{G(\mathbf{MM})}$ is an HNN extension, since we do not know if α is injective. This is enough to show:

Corollary 5.9. $\overline{G(\mathsf{MM})} / \langle t^{p(n)} = e \rangle \cong \overline{G_{p(n)}(\mathsf{MM})}.$

Proof. Let *K* and α be the group and homomorphism from Lemma 5.8, with MM = **MM**. Since $t^{-1}ft = \alpha(f)$ for all $f \in K$, $t^{-k}ft^k = \alpha^k(f)$ for all $k \ge 0$. If $f \in \langle x(q_1A_0) \rangle^K$, then the proof of Lemma 5.8 shows that $\alpha(f) = f \circledast a_1 \in \langle x(q_1A_0) \rangle^K$, so $\alpha^k(f) = f \circledast a_1^{\circledast k}$ for all $k \ge 0$. Thus, in $\overline{G(\mathbf{MM})} / \langle t^{p(n)} = e \rangle$ we have

$$x(q_1A_0) \circledast a_1^{\circledast p(n)} = t^{-p(n)} x(q_1A_0) t^{p(n)} = x(q_1A_0).$$

So
$$\overline{G(\mathsf{MM})} / \langle t^{p(n)} = e \rangle = \langle \overline{G(\mathsf{MM})} : t^{p(n)} = e, x(q_1 A_0) \circledast a_1^{\circledast p(n)} = x(q_1 A_0) \rangle = \overline{G_{p(n)}(\mathsf{MM})}.$$

Although we can't show that G(MM) is an HNN-extension, we can show:

Lemma 5.10. $\overline{G_{p(n)}(\mathsf{MM})}$ is a $\mathbb{Z}_{p(n)}$ -HNN-extension of $G_{p(n)}(\mathsf{MM})$ over the subgroup $K_{p(n)}$ generated by $x(q_1A_0)$, a_1 , and a'_1 in $G_{p(n)}(\mathsf{MM})$. In addition, $K_{p(n)}$ is amenable.

Proof. We continue with the notation from the proof of Lemma 5.8, with MM = MM. Note that the map $\psi : \mathcal{F}(S(MM)) \to \langle b_1, b_2 : [b_1, b_2] = e \rangle$ sending $a_1 \mapsto b_1, a'_1 \mapsto b_2$, and $s \mapsto e$ for all $s \in S(MM) \setminus \{a_1, a'_1\}$ sends $r \mapsto e$ for all r in the normal subgroup of $\mathcal{F}(S(MM))$ generated by $S(MM) \setminus \{a_1, a'_1\}$. If f belongs to this subgroup, then $f \circledast a_1$ also belongs to this subgroup, and hence

$$\psi(x(q_1A_0) \circledast a_1^{\circledast p(n)}) = e = \psi(x(q_1A_0)).$$

We conclude that ψ induces a homomorphism $G_{p(n)}(\mathsf{MM}) \to \mathbb{Z} \times \mathbb{Z}$. We can then follow the proof of Lemma 5.8 exactly to show that $K_{p(n)} = \langle x(q_1A_0) \rangle^{K_{p(n)}} \rtimes (\mathbb{Z} \times \mathbb{Z})$, and that there is a homomorphism $\tilde{\alpha} : K_{p(n)} \to K_{p(n)}$ such that $\tilde{\alpha}(a_1) = a_1$, $\tilde{\alpha}(a'_1) = a'_1$, and $\tilde{\alpha}(f) = f \circledast a_1$ for all $f \in \langle x(q_1A_0) \rangle^{K_{p(n)}}$.

If $f \in \langle x(q_1A_0) \rangle^{K_{p(n)}}$, then $\widetilde{\alpha}(f) = f \circledast a_1 \in \langle x(q_1A_0) \rangle^{K_{p(n)}}$, so $\widetilde{\alpha}^k(f) = f \circledast a_1^{\circledast k}$ for all $k \ge 0$. Hence

$$\widetilde{\alpha}^{p(n)}(x(q_1A_0)) = x(q_1A_0) \circledast a_1^{\circledast p(n)} = x(q_1A_0).$$

Since $\tilde{\alpha}^{p(n)}(a_1) = a_1$ and $\tilde{\alpha}^{p(n)}(a'_1) = a'_1$ as well, we conclude that $\tilde{\alpha}^{p(n)} = 1$ on $K_{p(n)}$. In other words, $\tilde{\alpha}$ is an automorphism of order p(n). Looking at the presentations, we see that $\overline{G_{p(n)}(\mathsf{MM})}$ is the $\mathbb{Z}_{p(n)}$ -HNN extension of $G_{p(n)}(\mathsf{MM})$ by $\tilde{\alpha}$.

To see that $K_{p(n)}$ is amenable, observe that $\langle x(q_1A_0)^{K_{p(n)}}$ is a subgroup of the group *T* from [KMS17, Lemma 4.5], and hence is abelian. As a semidirect product of two abelian groups, $K_{p(n)}$ is solvable, and hence amenable.

Towards proving part (b) of Proposition 5.6, we get:

Corollary 5.11. The group $\overline{G_{p(n)}(\mathsf{MM})}$ is sofic.

Proof. Since $G(\mathsf{MM})$ is solvable and $G_{p(n)}(\mathsf{MM})$ is a quotient of $G(\mathsf{MM})$, $G_{p(n)}(\mathsf{MM})$ is solvable. Since $\overline{G_{p(n)}(\mathsf{MM})}$ is a $\mathbb{Z}_{p(n)}$ -HNN-extension of $G_{p(n)}(\mathsf{MM})$ over the amenable subgroup $K_{p(n)}$, $\overline{G_{p(n)}(\mathsf{MM})}$ is sofic by Lemma 2.7.

For the proof of Proposition 5.6, we'll take $x = w(0)w_{accept}$. We already have the ingredients to show that x = e in $\overline{G_{p(n)}(\mathsf{MM})}$ if $n \in X$. However, we also need to ensure that $x \neq e$ if $n \notin X$. For this, we introduce a non-deterministic modification of MM , denoted by $\mathsf{MM}^{(p(n))}$. To construct $\mathsf{MM}^{(p(n))}$ from MM , we add p(n) - 1 additional states, which we denote by $2', \ldots, p(n)'$. We include all the commands of MM in $\mathsf{MM}^{p(n)}$, and we add p(n) new commands:

1;
$$\rightarrow$$
 2'; Add(1)
i'; \rightarrow (i+1)'; Add(1) for $2 \leq i < p(n)$
 $p(n)'; \rightarrow$ 1; Add(1).

In other words, in configuration (1; k, 0, 0, 0) in **MM**^{p(n)}, we have two choices. We can either apply commands from **MM**, or add a coin to the first glass and proceed to state 2'. After this choice, we are forced to go through the states i' for i = 2, ..., p(n), adding a coin to the first glass each time, until we return to state 1 in configuration (1; k + p(n), 0, 0, 0). Since **MM** was constructed using Lemma 5.2, we can show:

Lemma 5.12. Let $n \ge 1$. Then $(1; 0, 0, 0, 0) \equiv_{MM^{p(n)}} (0; 0, 0, 0, 0)$ if and only if $n \in X$.

Proof. If $n \in X$, then (1; p(n), 0, 0, 0) is accepted by **MM**. So in **MM**^{p(n)} there is a computation path going from (1; 0, 0, 0, 0) to (1; p(n), 0, 0, 0), and from there to the accept configuration. Hence $(1; 0, 0, 0, 0) \equiv_{\mathbf{MM}^{p(n)}} (0; 0, 0, 0, 0)$.

For the other direction, suppose $(1; 0, 0, 0, 0) \equiv_{\mathsf{MM}^{p(n)}} (0; 0, 0, 0, 0)$. Let *N* be the smallest integer such that there is a sequence of configurations

$$(1;0,0,0,0) =: C_0 \to C_1 \to \ldots \to C_N := (0;0,0,0,0),$$

where for all i = 1, ..., N, either C_{i-1} can be transformed to C_i by a command of $\mathsf{MM}^{p(n)}$, or C_i can be transformed to C_{i-1} . Let k be the largest integer such that $C_0 \to C_1 \to ... \to C_k$ does not use any commands from MM . The states of the configurations C_i for $0 \le i \le k$ must belong to $\{1, 2', 3', ..., p(n)'\}$, so in particular k < N. In addition, since $C_k \to C_{k+1}$ must use a command from MM , C_k must be in in state 1. The sequence $C_0 \to ... \to C_k$ shows that $(1; 0, 0, 0, 0) \equiv_{\mathsf{MM}'} C_k$ in the Minsky machine MM' with states $\{1, 2', ..., p(n)'\}$ and commands

1;
$$\rightarrow$$
 2'; Add(1)
i'; \rightarrow (i+1)'; Add(1) for $2 \leq i < p(n)$
 $p(n)'; \rightarrow$ 1; Add(1).

MM' is deterministic and sends (1; a, 0, 0, 0) to (1; a + mp(n), 0, 0, 0) for $m \ge 0$. Thus $(1; a, 0, 0, 0) \equiv_{MM'} (1; b, 0, 0, 0)$ if and only if $a = b \mod p(n)$. We conclude that $C_k = (1; mp(n), 0, 0, 0)$ for some $m \ge 0$.

Let ℓ be the largest integer such that $C_k \to \ldots \to C_\ell$ involves only commands from **MM**. If $\ell = N$, then $(1; mp(n), 0, 0, 0) \equiv_{\mathsf{MM}} (0; 0, 0, 0, 0)$. Since **MM** is deterministic, that would mean that mp(n) is accepted by **MM**, so m = 1 and $n \in X$ as desired.

Suppose $\ell < N$, so $C_{\ell} \to C_{\ell+1}$ involves a command not in **MM**. The configurations C_i for $k \le i \le \ell$ cannot be in states $\{2', \ldots, p(n)'\}$, so C_{ℓ} must be in state 1. The sequence $C_k \to \ldots \to C_{\ell}$ shows that $C_k \equiv_{\mathsf{MM}} C_{\ell}$. If $n \notin X$, then mp(n) is not accepted by **MM** for all $m \ge 0$, so $C_{\ell} = C_k$ by Lemma 5.2. But then $C_0 \to \ldots \to C_k \to C_{\ell+1} \to \ldots C_N$ is a sequence showing that $(1;0,0,0,0) \equiv_{\mathsf{MM}^{p(n)}} (0;0,0,0,0)$. Since $k < \ell$ by the definition of k, this contradicts the minimality of N. So again, we conclude that $n \in X$ as desired.

We are now ready to show:

Lemma 5.13. In $G_{p(n)}(MM)$, $w(0) = w_{accept}$ if and only if $n \in X$.

Proof. Because $x(q_1A_0) \circledast a_1^{\circledast p(n)} = x(q_1A_0)$ in $G_{p(n)}(\mathsf{MM})$,

$$w(0) = x(q_1A_0) \circledast A_1 \circledast A_2 \circledast A_3 \circledast A_4 = x(q_1A_0) \circledast a_1^{\circledast p(n)} \circledast A_1 \circledast A_2 \circledast A_3 \circledast A_4 = w(p(n)).$$

If $n \in X$, then p(n) is accepted by MM, so $w(p(n)) = w_{accept}$ in G(MM), and hence the same is true in $G_{p(n)}(MM)$. Thus we conclude that $w(0) = w_{accept}$ in $G_{p(n)}(MM)$ when $n \in X$.

For the other direction, we want to show that there is a homomorphism from $G_{p(n)}(\mathsf{MM})$ to $G(\mathsf{MM}^{p(n)})$. By Lemma 5.4, there is a homomorphism $G(\mathsf{MM}) \to G(\mathsf{MM}^{p(n)})$ which is the identity on generators. From the states and commands added to $\mathsf{MM}^{p(n)}$, we have generators $x(q_{i'}A_0)$ for $2 \le i \le p(n)$, and relations

$$\begin{aligned} x(q_1 A_0) &= x(q_{2'} A_0) \circledast a_1, \\ x(q_{i'} A_0) &= x(q_{(i+1)'} A_0) \circledast a_1 \text{ for all } 2 \le i < p(n), \\ x(q_{p(n)'} A_0) &= x(q_1 A_0) \circledast a_1. \end{aligned}$$

Combining these relations, we see that

$$x(q_1A_0) \circledast a_1^{\circledast p(n)} = x(q_1A_0)$$

in $G(\mathsf{MM}^{(p(n))})$, so the homomorphism $G(\mathsf{MM}) \to G(\mathsf{MM}^{p(n)})$ descends to a homomorphism $G_{p(n)}(\mathsf{MM}) \to G(\mathsf{MM}^{p(n)})$. If $n \notin X$, then by Lemma 5.12, $(1;0,0,0,0) \not\equiv_{\mathsf{MM}^{p(n)}}$ (0;0,0,0,0), so $w(0) \neq w_{accept}$ in $G(\mathsf{MM}^{p(n)})$. Hence $w(0) \neq w_{accept}$ in $G_{p(n)}(\mathsf{MM})$. \Box

The relations between $\overline{G(\mathbf{MM})}/\langle t^{p(n)} = e \rangle$, $G_{p(n)}(\mathbf{MM})$, $\overline{G_{p(n)}(\mathbf{MM})}$ and $G(\mathbf{MM}^{(p(n))})$ are summarized in the following equation:

$$G(\mathsf{MM}^{p(n)}) \longleftarrow G_{p(n)}(\mathsf{MM}) \longleftrightarrow \overline{G_{p(n)}(\mathsf{MM})} \cong \overline{G(\mathsf{MM})} / \langle t^{p(n)} = e \rangle.$$

We are finally ready to prove:

Proof of Proposition 5.6. Let

$$H = \langle \overline{G(\mathbf{MM})}, u : u^{-1}tu = t^r \rangle$$

We take $x = w(0)w_{accept} \in H$, and u and t to be the generators already defined in H. Part (a) follows immediately from the definition. For parts (b)-(d), observe that

$$H/\langle t^{p(n)} = e \rangle = \langle \overline{G_{p(n)}(\mathsf{MM})}, u : u^{-1}tu = t^r \rangle$$

by Corollary 5.9. By Lemma 5.10, $\overline{G_{p(n)}}(\mathsf{MM})$ is a $\mathbb{Z}_{p(n)}$ -HNN extension of $G_{p(n)}(\mathsf{MM})$. Hence *t* has order p(n) in $\overline{G_{p(n)}}(\mathsf{MM})$ by Lemma 2.1. Since *r* is coprime to p(n), $t^i \mapsto t^{ri}$ is an automorphism of the subgroup generated by *t* in $\overline{G_{p(n)}}(\mathsf{MM})$, so $H/\langle t^{p(n)} = e \rangle$ is an HNN-extension of $\overline{G_{p(n)}}(\mathsf{MM})$ over the subgroup generated by *t*. Since the subgroup generated by *t* is finite (and hence amenable), $H/\langle t^{p(n)} = e \rangle$ is sofic by [CLP15, Proposition 2.4.1], proving part (b). We also get that $\overline{G_{p(n)}}(\mathsf{MM})$ is a subgroup of $H/\langle t^{p(n)} = e \rangle$, so *t* has order p(n) in $H/\langle t^{p(n)} = e \rangle$ as well, proving part (d). And since $G_{p(n)}(\mathsf{MM})$ is a subgroup of $\overline{G_{p(n)}}(\mathsf{MM})$ by Lemma 2.1, x = e in $H/\langle t^{p(n)} = e \rangle$ if and only if $n \in X$ by Lemma 5.13, proving part (c).

It remains to show that *H* is an exended homogeneous linear-plus-conjugacy group, with *u*, *t*, and *x* in the generating set. By Lemma 5.5, $G(\mathbf{MM})$ has a presentation as an $m \times n' \times \ell'$ extended homogeneous linear-plus-conjugacy group $E\Gamma(A, C_0, C_1, L)$, in which $x(q_1A_0)$, $x = w(0)w_{accept}$ and $x(q_1A_0) \circledast a_1$ belong to the generating set $\{x_0, \ldots, x_{n'-1}\}$, and a_1 and a'_1 belong to the generating set $\{y_0, \ldots, y_{\ell'-1}\}$. To present *H* as an extended homogeneous linear-plus-conjugacy group, we can add two additional generators $y_{\ell'}$ and $y_{\ell'+1}$ for *t* and *u* respectively, and add the conjugacy relations

$$y_{\ell'}^{-1}x(q_1A_0)y_{\ell'} = x(q_1A_0) \circledast a_1, y_{\ell'}^{-1}a_1y_{\ell'} = a_1, y_{\ell'}^{-1}a_1'y_{\ell'} = a_1', \text{ and } y_{\ell'+1}^{-1}y_{\ell'}y_{\ell'+1} = y_{\ell'}^r.$$

This gives a presentation of *H* as an $m \times n' \times \ell$ extended homogeneous linear-plusconjugacy group, where $\ell = \ell' + 2$.

5.4 Embedding KMS groups in solution groups

Proposition 5.14. Let p(n), $n \ge 1$ be an increasing sequence of prime numbers, where the function $p : \mathbb{Z}_{\ge 1} \to \mathbb{Z}_{\ge 1}$ is computable, let X be a recursively enumerable set of positive integers, and let r be a positive integer which is coprime to p(n) for all $n \ge 1$. Then there is an $m \times n$ solution group $\Gamma(A)$ with generators $x, t_1, t_2, u_1, u_2 \in \{x_0, \dots, x_{n-1}\}$ such that

- (a) $u_2u_1t_1t_2u_1u_2 = (t_1t_2)^r$ in $\Gamma(A)$,
- (b) $x = e \text{ in } \Gamma(A) / \langle (t_1 t_2)^{p(n)} = e \rangle$ if and only if $n \in X$,
- (c) if $n \notin X$, then x is non-trivial in approximate representations of $\Gamma(A) / \langle (t_1 t_2)^{p(n)} = e \rangle$,
- (d) t_1t_2 has order p(n) in $\Gamma(A)/\langle (t_1t_2)^{p(n)} = e \rangle$,
- (e) $w \in \langle t_1, t_2 \rangle \setminus \{e\}$ is nontrivial in approximate representations of $\Gamma(A) / \langle (t_1t_2)^{p(n)} = e \rangle$, and
- (f) each row of A only has three nonzero entries.

Proof. Let *H* be the $m \times n \times \ell$ extended homogeneous linear-plus-conjugacy group from Proposition 5.6, with elements $x, t, u \in H$. Let $\Gamma(A')$ be the $m' \times n'$ solution group corresponding to *H* from Proposition 4.5, and let

$$\widetilde{\phi} : \mathcal{F}(x_0, \dots, x_{n-1}, y_0, \dots, y_{\ell-1}) \to \mathcal{F}(x_0, \dots, x_{n'-1}) \text{ and } \phi : H \to \Gamma(A')$$

be the homomorphisms from that proposition. By parts (b) and (c) of Proposition 4.5, $\tilde{\phi}(x) = x$, $\tilde{\phi}(t) = t_1 t_2$, and $\tilde{\phi}(u) = u_1 u_2$ for some generators x, t_1 , t_2 , u_1 , $u_2 \in \{x_0, \ldots, x_{n'-1}\}$. Since ϕ is a homomorphism,

$$u_2 u_1 t_1 t_2 u_1 u_2 = \phi(u^{-1} t u) = \phi(t^r) = (t_1 t_2)^r,$$

proving part (a). Let $\Gamma_n := \Gamma(A') / \langle (t_1 t_2)^{p(n)} = e \rangle$. If $n \in X$, then x = e in $H / \langle t^{p(n)} = e \rangle$, so x = e in Γ_n .

By part (e) of Proposition 4.5, there are integers n_1 , n_2 , n_3 such that for any ϵ -representation ψ of H, there is an $O(\epsilon)$ -representation α of $\Gamma(A')$ such that

$$\alpha(\widetilde{\phi}(g)) = \psi(g)^{\oplus n_1} \oplus \overline{\psi(g)}^{\oplus n_2} \oplus \mathbb{1}_{n_3} \text{ for all } g \in \mathcal{F}(x_0, \ldots, x_{n-1}, y_0, \ldots, y_{\ell-1}).$$

If ψ is an ϵ -representation of $H/\langle t^{p(n)} = e \rangle$, then $\|\psi(t^{p(n)}) - \mathbb{1}\| \leq \epsilon$, so

$$\|\alpha((t_1t_2)^{p(n)}) - \mathbb{1}\| = \|\alpha(\widetilde{\phi}(t^{p(n)})) - \mathbb{1}\| = \|\psi(t^{p(n)})^{\oplus n_1} \oplus \overline{\psi(t^{p(n)})}^{\oplus n_2} \oplus \mathbb{1}_{n_3} - \mathbb{1}\| \le \epsilon$$

as well, and α is an $O(\epsilon)$ -representation of Γ_n .

Suppose $w \neq e$ in $H/\langle t^{p(n)} = e \rangle$. Since $H/\langle t^{p(n)} = e \rangle$ is sofic, w is non-trivial in approximate representations of $H/\langle t^{p(n)} = e \rangle$, which means that there is $\delta > 0$ such that for all $\epsilon > 0$, there is an ϵ -representation ψ with $\|\psi(w) - \mathbb{1}\| \geq \delta$. If α is an $O(\epsilon)$ -representation of Γ_n as above, then

$$\|\alpha(\widetilde{\phi}(w)) - \mathbb{1}\| \ge \sqrt{\frac{n_1d + n_2d}{n_1d + n_2d + n_3}} \delta \ge \sqrt{\frac{n_1 + n_2}{n_1 + n_2 + n_3}} \delta.$$

Since n_1 , n_2 , and n_3 are independent of ϵ , $\tilde{\phi}(w)$ is non-trivial in approximate representations of Γ_n .

If $n \notin X$, then $x \neq e$ in $H/\langle t^{p(n)} = e \rangle$, so x is non-trivial in approximate representations of Γ_n . In particular, $x \neq e$ in Γ_n , so parts (b) and (c) hold. Similarly, $t^i \neq e$ in $H/\langle t^{p(n)} = e \rangle$ for all 0 < i < p(n), and thus $(t_1t_2)^i \neq e$ in Γ_n , proving part (d). This also shows that $(t_1t_2)^i$ is nontrivial in approximate representations of Γ_n for 0 < i < p(n), or in other words that $(t_1t_2)^i \neq e$ in Γ_n^{fa} . This means that t_1 and t_2 must have order 2 in Γ_n^{fa} . Thus $\langle t_1, t_2 \rangle_{\Gamma_n^{fa}} \cong D_{p(n)}$. Since $\langle t_1, t_2 \rangle_{\Gamma_n} \cong D_{p(n)}$ as well, this proves part (e). Finally, part (f) follows immediately from part (a) of Proposition 4.5.

6 Constant-sized correlations \mathfrak{C}_p for the dihedral groups

In Section 4, we showed that any strategy for a perfect correlation of a solution group must come from a representation of that group. The dihedral group D_n has a presentation as an extended homogeneous linear-plus-conjugacy group, and thus can be embedded in solution groups. However, the size of this solution group will depend on n. In this section we write down a constant-sized correlation \mathfrak{C}_p for the dihedral group D_p , p a prime, such that any commuting-operator strategy (meeting a condition which we will enforce in the next section using perfect correlations for the solution group from Proposition 5.14) induces a representation of the dihedral group. The construction we use comes from [Fu22], although we modify the construction slightly so that the correlation comes from the regular representation, rather than the representation used in [Fu22]. This modification is necessary for the next section, where we use the correlation from this section in conjunction with perfect correlations for the solution from this section in conjunction with perfect correlations for the solution from this section in conjunction with perfect correlations for the solution from this section in conjunction with perfect correlations for the solution group in Proposition 5.14.

To define \mathfrak{C}_p , recall that D_p is generated by t_1 and t_2 , and consists of the elements $(t_1t_2)^j$ and $t_2(t_1t_2)^j$ for $j \in [p]$. As in Section 2, let *L* and *R* denote the left and right regular representations of D_p on

$$\ell^2 D_p = \operatorname{span}(\{|(t_1t_2)^j\rangle, |t_2(t_1t_2)^j\rangle \mid j \in [p]\}).$$

Define elements

$$\begin{aligned} \pi_{0}^{(0)} &= \frac{1}{p} \sum_{j \in [p]} (t_{1}t_{2})^{j}, \\ \pi_{0}^{(1)} &= \frac{2}{p} \sum_{j \in [p]} \cos(\frac{2j\pi}{p})(t_{1}t_{2})^{j}, \\ \pi_{0}^{(2)} &= e - \pi_{0}^{(0)} - \pi_{0}^{(1)}, \\ \pi_{1}^{(0)} &= \frac{1}{2} \pi_{0}^{(1)} + \frac{1}{p} \sum_{j \in [p]} \cos(\frac{(2j+1)\pi}{p})t_{2}(t_{1}t_{2})^{j}, \\ \pi_{1}^{(1)} &= \pi_{0}^{(1)} - \pi_{1}^{(0)}, \\ \pi_{1}^{(2)} &= e - \pi_{0}^{(1)}, \\ \pi_{2}^{(0)} &= \frac{1}{2} \pi_{0}^{(1)} + \frac{1}{p} \sum_{j \in [p]} \sin(\frac{(2j+1)\pi}{p})t_{2}(t_{1}t_{2})^{j}, \\ \pi_{2}^{(1)} &= \pi_{0}^{(1)} - \pi_{2}^{(0)}, \text{ and} \\ \pi_{2}^{(2)} &= e - \pi_{0}^{(1)}. \end{aligned}$$

in the group algebra $\mathbb{C}[D_p]$. These elements are all projections. For instance, $\pi_0^{(0)}$ is the sum of central projections for the trivial and sign representations. $\pi_0^{(1)}$ is the central projection for the 2-dimensional irreducible representation $V^{(1)}$ sending

$$t_1 \mapsto \begin{pmatrix} 0 & \omega_p \\ \omega_p^{-1} & 0 \end{pmatrix}$$
 and $t_2 \mapsto \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

The projections $\pi_1^{(0)}$ and $\pi_2^{(0)}$ are more complicated: they correspond to the rank-one projections

$$\frac{1}{2} \begin{pmatrix} 1 & \omega_{2p} \\ \omega_{2p}^{-1} & 1 \end{pmatrix} \text{ and } \frac{1}{2} \begin{pmatrix} 1 & -i\omega_{2p} \\ i\omega_{2p}^{-1} & 1 \end{pmatrix}$$

in $V^{(1)}$.

Before we define \mathfrak{C}_p , we first consider the correlation \mathfrak{C}'_p for the scenario ([5], [5], [3], [3]) defined by

$$\mathfrak{C}'_{p}(a,b|x,y) = \frac{\langle e|L(e-\pi_{0}^{(0)})\tilde{P}_{x}^{(a)}\tilde{Q}_{y}^{(b)}L(e-\pi_{0}^{(0)})|e\rangle}{\|L(e-\pi_{0}^{(0)})|e\rangle\|^{2}}, x, y \in [5], a, b \in [3], b$$

where

$$\tilde{P}_{x}^{(a)} = \begin{cases} L(\pi_{0}^{(a+1)}) & \text{if } x = 0, a \in \{0,1\} \\ \frac{L(e) + (-1)^{a}L(t_{x})}{2} & \text{if } x \in \{1,2\}, a \in \{0,1\} \\ L(\pi_{1}^{(a)}) & \text{if } x = 3 \\ L(\pi_{2}^{(a)}) & \text{if } x = 4 \\ 0 & \text{otherwise} \end{cases}$$

$$\tilde{Q}_{y}^{(b)} = \begin{cases} R(\pi_{0}^{(b+1)}) & \text{if } y = 0, b \in \{0,1\} \\ \frac{R(e) + (-1)^{b}R(t_{y})}{2} & \text{if } y \in \{1,2\}, b \in \{0,1\} \\ R(\pi_{1}^{(b)}) & \text{if } y = 3 \\ R(\pi_{2}^{(b)}) & \text{if } y = 4 \\ 0 & \text{otherwise} \end{cases}$$

Direct calculation gives us the following.

Lemma 6.1. The correlation \mathfrak{C}'_p is the correlation $\hat{P}_{-\pi/p}$ defined in [Fu22, Definition 3.2].

One important property of $\hat{P}_{-\pi/p}$ is summarized in the following proposition.

Proposition 6.2 (Proposition 3.3 of [Fu22]). Let $(|\psi\rangle, \{\{P_x^{(a)}\}\}, \{\{Q_y^{(b)}\}\})$ be an inducing strategy of $\hat{P}_{-\pi/p}$, and let

$$|\phi_1\rangle = \frac{1}{2} \left(P_3^{(0)} + iP_4 P_3^{(1)} - iP_4 P_3^{(0)} + P_3^{(1)} \right) |\psi\rangle, \tag{3}$$

where $P_4 = P_4^{(0)} - P_4^{(1)}$. Then

$$\||\phi_1\rangle\|^2 = 1/(p-1), P_1P_2|\phi_1\rangle = \omega_p^{-1}|\phi_1\rangle, and Q_1Q_2|\phi_1\rangle = \omega_p|\phi_1\rangle,$$

where $P_x = P_x^{(0)} - P_x^{(1)}$ and $Q_y = Q_y^{(0)} - Q_y^{(1)}$ for $x, y \in \{1, 2\}$.

Note that the proof of Proposition 3.3 of [Fu22] also tells us that

$$|\phi_1\rangle = \frac{1}{2} \left(Q_3^{(0)} - iQ_4 Q_3^{(1)} + iQ_4 Q_3^{(0)} + Q_3^{(1)} \right) |\psi\rangle, \tag{4}$$

where $Q_4 = Q_4^{(0)} - Q_4^{(1)}$. Now we are ready to define \mathfrak{C}_p . Let $I := \{0, 1, 2, t_1, t_2, (0, t_1), (0, t_2)\}$. We define \mathfrak{C}_p as a correlation for the scenario $(I, I, [3] \times [2], [3] \times [2])$ by

$$\mathfrak{C}_p(a,b|x,y) = \langle e|\tilde{M}_x^{(a)}\tilde{N}_y^{(b)}|e\rangle, x,y \in I, a,b \in [3] \times [2],$$

where $|e\rangle \in L_2D_p$,

$$\begin{split} \tilde{M}_{x}^{(a_{0},a_{1})} &= \begin{cases} L(\pi_{x}^{(a_{0})}) & \text{if } x \in \{0,1,2\}, a_{1} = 0, \\ \frac{L(e) + (-1)^{a_{1}}L(x)}{2} & \text{if } x \in \{t_{1},t_{2}\}, a_{0} = 0, \\ \tilde{M}_{0}^{(a_{0},0)}\tilde{M}_{t}^{(0,a_{1})} & \text{if } x = (0,t) \\ 0 & \text{otherwise} \end{cases} \\ \tilde{N}_{y}^{(b_{0},b_{1})} &= \begin{cases} R(\pi_{y}^{(b_{0})}) & \text{if } y \in \{0,1,2\}, b_{1} = 0, \\ \frac{R(e) + (-1)^{b_{1}}R(y)}{2} & \text{if } y \in \{t_{1},t_{2}\}, b_{0} = 0, \\ \tilde{N}_{0}^{(b_{0},0)}\tilde{N}_{t}^{(0,b_{1})} & \text{if } y = (0,t) \\ 0 & \text{otherwise} \end{cases} \end{split}$$

It's easy to see that the families $\{\tilde{M}_x^{(a_0,a_1)}: (a_0,a_1) \in [3] \times [2]\}$ and $\{\tilde{N}_y^{(b_0,b_1)}: (b_0,b_1) \in [3] \times [2]\}$ are projective measurements for all $x, y \in \{0, 1, 2, t_1, t_2\}$. For $x, y \in \{(0, t_1), (0, t_2)\}$, this follows from the fact that $\pi_0^{(a)}$ is a central projection for all $a \in [3]$. So \mathfrak{C}_p is in $C_{qc}(I, I, [3] \times [2], [3] \times [2])$. Some of the most important entries of \mathfrak{C}_p are shown in Table 1. We summarize some additional properties in the following lemma:

Lemma 6.3.

- (a) \mathfrak{C}_p is a synchronous correlation in $C_{qc}(I, I, [3] \times [2], [3] \times [2])$.
- (b) The entries of \mathfrak{C}_p are computable elements of $\overline{\mathbb{Q}}$.

Proof. Let $\iota : \mathbb{C}[D_p] \to \mathbb{C}[D_p]$ be the linear involution sending $g \mapsto g^{-1}$ for all $g \in D_p$. It is not hard to see that $\pi_j^{(i)} = \iota(\pi_j^{(i)})$ for all *i* and *j*. In addition, $R(\alpha)|e\rangle = L(\iota(\alpha))|e\rangle$ for all $\alpha \in \mathbb{C}[D_p]$, so $\tilde{N}_y^{(b)}|e\rangle = \tilde{M}_y^{(b)}|e\rangle$ for all $y \in I$ and $b \in [3] \times [2]$. So

$$\langle e | ilde{M}_x^{(a)} ilde{N}_x^{(b)} | e
angle = \langle e | ilde{M}_x^{(a)} ilde{M}_x^{(b)} | e
angle = 0$$

for all $x \in I$ and $a \neq b \in [3] \times [2]$. We conclude that \mathfrak{C}_p is synchronous. Part (b) follows from the fact that the elements $\pi_i^{(i)}$ belong to $\overline{\mathbb{Q}}[D_p]$.

We now come to the main theorem of this section. To state this theorem, recall that an integer r is a primitive root of a prime p if all the integers between 1 and p - 1 are congruent modulo p to some power of r.

Theorem 6.4. Let $S = (|\psi\rangle, \{M_x^{(a_0,a_1)}\}, \{N_y^{(b_0,b_1)}\})$ be a good strategy for \mathfrak{C}_p and let r be a primitive root of p. Suppose there exist unitaries U_A and U_B such that

$$\begin{aligned} U_A U_B |\psi\rangle &= U_B U_A |\psi\rangle, \\ U_A N_y^{(b_0,b_1)} |\psi\rangle &= N_y^{(b_0,b_1)} U_A |\psi\rangle \text{ for all } y, b_0, b_1, \\ U_B M_x^{(a_0,a_1)} |\psi\rangle &= M_x^{(a_0,a_1)} U_B |\psi\rangle \text{ for all } x, a_0, a_1, \\ U_A U_B |\psi\rangle &= |\psi\rangle, \\ (N_{t_1} N_{t_2}) U_B |\psi\rangle &= U_B (N_{t_1} N_{t_2})^r |\psi\rangle, \text{ and} \\ (M_{t_1} M_{t_2}) U_A |\psi\rangle &= U_A (M_{t_1} M_{t_2})^r |\psi\rangle, \end{aligned}$$

where $M_x = M_x^{(0,0)} - M_x^{(0,1)}$ and $N_y = N_y^{(0,0)} - N_y^{(0,1)}$ for $x, y \in \{t_1, t_2\}$. Then $(M_{t_1}M_{t_2})^p |\psi\rangle = |\psi\rangle.$

When we use this theorem in the next section, the existence of U_A and U_B will be guaranteed by Proposition 5.14.

The rest of the section is devoted to the proof of Theorem 6.4, so for the remainder of the section, we will assume that we have a good strategy *S* and unitaries U_A and U_B satisfying the conditions of the theorem. Note that in a good strategy, $M_x^{(a_0,a_1)} = N_x^{(a_0,a_1)} = 0$ if $x \in \{0,1,2\}$ and $a_1 \neq 0$, or if $x \in \{t_1, t_2\}$ and $a_0 \neq 0$. In particular, M_{t_i} and N_{t_i} are binary observables. The basic idea of the proof is to find a decomposition of $|\psi\rangle$ as $|\psi\rangle = \sum_{j=0}^p |\psi_j\rangle$, where $||\psi_0\rangle||^2 = ||\psi_p\rangle||^2 = 1/2p$, $||\psi_j\rangle||^2 = 1/p$ for $1 \leq j \leq p - 1$, and $|\psi_j\rangle$ is an eigenvector of $M_{t_1}M_{t_2}$ with eigenvalue ω_p^j . Intuitively, $|\psi_0\rangle$ and $|\psi_p\rangle$ are in the 1-dimensional irreducible representation of D_p , and $|\psi_j\rangle$ and $|\psi_{p-j}\rangle$ are in the 2-dimensional irreducible representation of D_p , in which

$$t_1 t_2 \mapsto \begin{pmatrix} \omega_p^j & 0\\ 0 & \omega_p^{-j} \end{pmatrix}$$

for $1 \le j \le (p-1)/2$. The norms of the vectors are chosen because the multiplicity of the 2-dimensional irreducible representations in the regular representation of D_p is 2, and the multiplicity of the 1-dimensional irreducible representations is 1.

The vectors $|\psi_0\rangle$ and $|\psi_p\rangle$ are defined as

$$|\psi_0\rangle = M_{t_1}^{(0,0)} M_0^{(0,0)} |\psi\rangle$$
, and $|\psi_p\rangle = M_{t_1}^{(0,1)} M_0^{(0,0)} |\psi\rangle$. (5)

It follows immediately from the definition of M_{t_1} that

$$M_{t_1}|\psi_0\rangle = |\psi_0\rangle$$
, and $M_{t_1}|\psi_p\rangle = -|\psi_p\rangle$. (6)

To see that

$$\||\psi_0\rangle\|^2 = \||\psi_p\rangle\|^2 = \frac{1}{2p'},\tag{7}$$

		y = 0				
		b = (0,0)	b = (1,0)	b = (2, 0)		
	a = (0, 0)	1/p	0	0		
x = 0	a = (1, 0)	0	2/p	0		
	a = (2, 0)	0	0	(p-3)/p		

(a) \mathfrak{C}_p : the correlation values for x = y = 0.

		y =	= 1	<i>y</i> = 2		
		b = (0,0)	b = (1,0)	b = (0,0)	b = (1,0)	
$x = t_1$	a = (0, 0)	$\frac{\cos^2(\pi/2p)}{p}$	$\frac{\sin^2(\pi/2p)}{p}$	$\frac{1 - \sin(\pi/p)}{2p}$	$\frac{1+\sin(\pi/p)}{2p}$	
	a = (0, 1)	$\frac{\sin^2(\pi/2p)}{p}$	$\frac{\cos^2(\pi/2p)}{p}$	$\frac{1+\sin(\pi/p)}{2p}$	$\frac{1-\sin(\pi/p)}{2p}$	
$x = t_2$	a = (0, 0)	$\frac{\cos^2(\pi/2p)}{p}$	$\frac{\sin^2(\pi/2p)}{p}$	$\frac{1+\sin(\pi/p)}{2p}$	$\frac{1-\sin(\pi/p)}{2p}$	
	a = (0, 1)	$\frac{\sin^2(\pi/2p)}{p}$	$\frac{\cos^2(\pi/2p)}{p}$	$\frac{1-\sin(\pi/p)}{2p}$	$\frac{1+\sin(\pi/p)}{2p}$	

(b) \mathfrak{C}_p : the correlation values for $x \in \{t_1, t_2\}$ and $y \in \{1, 2\}$.

		x = 1		x = 2			x = 0		
		$a_0 = 0$	$a_0 = 1$	$a_0 = 2$	$a_0 = 0$	$a_0 = 1$	$a_0 = 2$	$a_0 = 1$	$a_0 \neq 1$
	$b_0 = 0$	$\frac{1}{p}$	0	0	$\frac{1}{2p}$	$\frac{1}{2p}$	0	$\frac{1}{p}$	0
y = 1	$b_0 = 1$	0	$\frac{1}{p}$	0	$\frac{1}{2p}$	$\frac{1}{2p}$	0	$\frac{1}{p}$	0
	$b_0 = 2$	0	0	$\frac{p-2}{p}$	0	0	$\frac{p-2}{p}$	0	$\frac{p-2}{p}$
	$b_0 = 0$	$\frac{1}{2p}$	$\frac{1}{2p}$	0	$\frac{1}{p}$	0	0	$\frac{1}{p}$	0
y = 2	$b_0 = 1$	$\frac{1}{2p}$	$\frac{1}{2p}$	0	0	$\frac{1}{p}$	0	$\frac{1}{p}$	0
	$b_0 = 2$	0	0	$\frac{p-2}{p}$	0	0	$\frac{p-2}{p}$	0	$\frac{p-2}{p}$
u = 0	$b_0 = 1$	$\frac{1}{p}$	$\frac{1}{p}$	0	$\frac{1}{p}$	$\frac{1}{p}$	0	$\frac{2}{p}$	0
3 0	$b_0 \neq 1$	0	0	$\frac{p-2}{p}$	0	0	$\frac{p-2}{p}$	0	$\frac{p-2}{p}$

(c) \mathfrak{C}_p : the correlation values for $x, y \in \{0, 1, 2\}$.

$y = (0, t_1)$							
		b = (0, 0)	b = (0, 1)	b = (1,0)	b = (1, 1)	b = (2, 0)	b = (2, 1)
	$a_0 = 0$	$\frac{1}{2p}$	$\frac{1}{2p}$	0	0	0	0
x = 0	$a_0 = 1$	0	0	$\frac{1}{p}$	$\frac{1}{p}$	0	0
	$a_0 = 2$	0	0	0	0	$\frac{p-3}{2p}$	$\frac{p-3}{2p}$
$x = t_1$	$a_1 = 0$	$\frac{1}{2p}$	0	$\frac{1}{p}$	0	$\frac{p-3}{2p}$	0
	$a_1 = 1$	0	$\frac{1}{2p}$	0	$\frac{1}{p}$	0	$\frac{p-3}{2p}$

(d) \mathfrak{C}_p : the correlation values for the commutation test for Alice's questions 0 and t_1 .

		y = 0	$(0, t_2)$
		b=(0,0)	b = (0, 1)
$\mathbf{r} = (0, t_1)$	a = (0, 0)	1/p	0
$x = (0, t_1)$	a = (0, 1)	0	1/p

(e) \mathfrak{C}_p : some of the values for $x = (0, t_1), y = (0, t_2), a_0 = b_0 = 0$.

Table 1: Some important values of \mathfrak{C}_p .

we need the following identities. Lemma 3.6 applied to Table 1d implies that

$$M_x^{(0,a_1)} M_0^{(a_0,0)} |\psi\rangle = N_{(0,x)}^{(a_0,a_1)} |\psi\rangle = M_0^{(a_0,0)} M_x^{(0,a_1)} |\psi\rangle$$
(8)

for $a_0 \in [3], a_1 \in [2]$ and $x \in \{t_1, t_2\}$. Then

$$\||\psi_0\rangle\|^2 = \langle \psi|M_0^{(0,0)}M_{t_1}^{(0,0)}M_0^{(0,0)}|\psi\rangle = \langle \psi|M_0^{(0,0)}N_{(0,t_1)}^{(0,0)}|\psi\rangle = \frac{1}{2p}$$

as shown in Table 1d. The derivation of $|||\psi_p\rangle||^2 = \frac{1}{2p}$ is similar. Next, Lemma 3.5 applied to Table 1e implies that

$$M^{(0,a_1)}_{(0,t_1)}|\psi
angle = N^{(0,a_1)}_{(0,t_2)}|\psi
angle$$

for each $a_1 \in [2]$. Hence,

$$|\psi_0\rangle = M_{t_1}^{(0,0)} M_0^{(0,0)} |\psi\rangle = N_{(0,t_2)}^{(0,0)} |\psi\rangle = M_{t_2}^{(0,0)} M_0^{(0,0)} |\psi\rangle,$$

and similarly, $|\psi_p
angle=M_{t_2}^{(0,1)}M_0^{(0,0)}|\psi
angle$. Thus

$$M_{t_2}|\psi_0\rangle = |\psi_0\rangle$$
, and $M_{t_2}|\psi_p\rangle = -|\psi_p\rangle$, (9)

implying that $|\psi_0\rangle$ and $|\psi_p\rangle$ are 1-eigenvectors of $M_{t_1}M_{t_2}$.

The vectors $|\psi_j\rangle$ for $2 \le j \le p-1$ can be constructed from $|\psi_1\rangle$, but the construction of $|\psi_1\rangle$ is more complicated and requires Proposition 6.2. We first show a strategy for \mathfrak{C}'_p can be extracted from any strategy for \mathfrak{C}_p .

Proposition 6.5. Suppose $(|\psi\rangle, \{M_x^{(a)} \mid a \in [3] \times [2]\}, x \in I, \{N_y^{(b)} \mid b \in [3] \times [2]\}, y \in I)$ is a good strategy for \mathfrak{C}_p . Let $|\psi'\rangle = (\mathbb{1} - M_0^{(0,0)})|\psi\rangle/\|(\mathbb{1} - M_0^{(0,0)})|\psi\rangle\|$, and define projective measurements $\{P_x^{(a)} \mid a \in [3]\}, x \in [5], and \{Q_y^{(b)} \mid b \in [3]\}, y \in [5]$ by

$$\begin{array}{ll} P_{0}^{(0)} = M_{0}^{(1,0)}, & Q_{0}^{(0)} = N_{0}^{(1,0)}, \\ P_{0}^{(1)} = M_{0}^{(2,0)}, & Q_{0}^{(1)} = N_{0}^{(2,0)}, \\ P_{0}^{(2)} = 0, & Q_{0}^{(2)} = 0, \\ for \ x = 1,2: & P_{x}^{(a)} = \begin{cases} M_{t_{x}}^{(0,a)} & if \ a = 0,1 \\ 0 & otherwise \end{cases}, & Q_{x}^{(a)} = \begin{cases} N_{t_{x}}^{(0,a)} & if \ a = 0,1 \\ 0 & otherwise \end{cases}, \\ P_{3}^{(a)} = M_{1}^{(a,0)}, & Q_{3}^{(a)} = N_{1}^{(a,0)}, \\ P_{4}^{(a)} = M_{2}^{(a,0)}, \ and & Q_{4}^{(a)} = N_{2}^{(a,0)}. \end{cases}$$

Then $(|\psi'\rangle, \{P_x^{(a)} \mid a \in [3]\}, x \in [5], \{Q_y^{(b)} \mid b \in [3]\}, y \in [5])$ *is a strategy for* \mathfrak{C}'_p .

Proof. For this proof, the first key observation is that

$$\|(\mathbb{1} - M_0^{(0,0)})|\psi\rangle\|^2 = (p-1)/p_0$$

which follows from the fact that $||M_0^{(0,0)}|\psi\rangle||^2 = 1/p$. We also need the following identities:

$$(M_0^{(0,0)} + M_0^{(2,0)})|\psi\rangle = M_1^{(2,0)}|\psi\rangle = M_2^{(2,0)}|\psi\rangle = N_1^{(2,0)}|\psi\rangle = N_2^{(2,0)}|\psi\rangle = (N_0^{(0,0)} + N_0^{(2,0)})|\psi\rangle,$$
(10)

and

$$(M_1^{(0,0)} + M_1^{(1,0)})|\psi\rangle = (M_2^{(0,0)} + M_2^{(1,0)})|\psi\rangle = M_0^{(1,0)}|\psi\rangle = N_0^{(1,0)}|\psi\rangle = (N_1^{(0,0)} + N_1^{(1,0)})|\psi\rangle = (N_2^{(0,0)} + N_2^{(1,0)})|\psi\rangle.$$
(11)

To prove Equation (10), first observe that $\{M_i^{(0,0)}, M_i^{(1,0)}, M_i^{(2,0)}\}$ and $\{N_i^{(0,0)}, N_i^{(1,0)}, N_i^{(2,0)}\}$ are projective measurements for i = 0, 1, 2. By Table 1d, the two projective measurements $\{M_0^{(1,0)}, (M_0^{(0,0)} + M_0^{(2,0)})\}$ and $\{(N_1^{(0,0)} + N_1^{(1,0)}), N_1^{(2,0)}\}$ satisfy the conditions of Lemma 3.5 with respect to $|\psi\rangle$, and hence we have $(M_0^{(0,0)} + M_0^{(2,0)})|\psi\rangle = N_1^{(2,0)}|\psi\rangle$. For the same reason, $N_1^{(2,0)}|\psi\rangle = M_1^{(2,0)}|\psi\rangle = (N_2^{(0,0)} + N_2^{(1,0)})|\psi\rangle$ and so on. Equation (11) follows from Equation (10) as $1 - (M_0^{(0,0)} + M_0^{(2,0)}) = M_0^{(1,0)}, 1 - N_1^{(2,0)} = N_1^{(0,0)} + N_1^{(1,0)}$ and so on.

Since $||M_x^{(a,0)}N_0^{(2,0)}|\psi\rangle||^2 = \langle \psi | M_x^{(a,0)}N_0^{(2,0)}|\psi\rangle = 0$ from Table 1d, we have $M_x^{(a,0)}N_0^{(2,0)}|\psi\rangle = 0$. Using the identities in Equations (10) and (11), we can prove that, for x = 1, 2 and $a \in \{0, 1\}$,

$$\begin{split} M_x^{(a,0)}(\mathbb{1} - M_0^{(0,0)})|\psi\rangle &= M_x^{(a,0)}(N_0^{(1,0)} + N_0^{(2,0)})|\psi\rangle = M_x^{(a,0)}N_0^{(1,0)}|\psi\rangle \\ &= M_x^{(a,0)}(M_x^{(0,0)} + M_x^{(1,0)}|\psi\rangle = M_x^{(a,0)}|\psi\rangle. \end{split}$$

That is,

$$M_x^{(a,0)}|\psi\rangle = M_x^{(a,0)}(\mathbb{1} - M_0^{(0,0)})|\psi\rangle.$$
(12)

The same argument can also give us that, for x = 1, 2 and $a \in \{0, 1\}$,

$$N_x^{(a,0)}|\psi\rangle = N_x^{(a,0)}(\mathbb{1} - M_0^{(0,0)})|\psi\rangle.$$
(13)

Following the definitions of $M_0^{(a,0)}$ and $N_0^{(a,0)}$ and using $N_0^{(0,0)}|\psi\rangle = M_0^{(0,0)}|\psi\rangle$, we can also see that

$$M_0^{(a,0)}|\psi\rangle = M_0^{(a,0)}(\mathbb{1} - M_0^{(0,0)})|\psi\rangle \text{ and } N_0^{(a,0)}|\psi\rangle = N_0^{(a,0)}(\mathbb{1} - M_0^{(0,0)})|\psi\rangle$$
(14)

for *a* = 1, 2.

To see the extracted strategy induces \mathfrak{C}'_p , we need to determine $\langle \psi | (\mathbb{1} - M_0^{(0,0)}) P_x^{(a)} Q_y^{(b)} (\mathbb{1} - M_0^{(0,0)}) | \psi \rangle$ for $x, y \in [5]$ and $a, b \in [3]$ from the values of \mathfrak{C}_p . If $x, y \in \{0, 3, 4\}$, it follows from the definitions of $P_x^{(a)}$ and $Q_y^{(b)}$ and Equations (12) to (14) that

$$\langle \psi | (\mathbb{1} - M_0^{(0,0)}) P_x^{(a)} Q_y^{(b)} (\mathbb{1} - M_0^{(0,0)}) | \psi \rangle = \langle \psi | P_x^{(a)} Q_y^{(b)} | \psi \rangle.$$

If $x \in \{1, 2\}$ and $y \in \{0, 3, 4\}$, Equations (8), (13) and (14) imply that

$$\langle \psi | (\mathbb{1} - M_0^{(0,0)}) P_x^{(a)} Q_y^{(b)} (\mathbb{1} - M_0^{(0,0)}) | \psi \rangle = \langle \psi | P_x^{(a)} Q_y^{(b)} (\mathbb{1} - M_0^{(0,0)})^2 | \psi \rangle = \langle \psi | P_x^{(a)} Q_y^{(b)} | \psi \rangle.$$

The same identity holds for $x \in \{0,3,4\}$ and $y \in \{1,2\}$ by symmetry. Finally, when $x, y \in \{1,2\}$, Equation (8) implies that

$$\begin{aligned} \langle \psi | (\mathbb{1} - M_0^{(0,0)}) P_x^{(a)} Q_y^{(b)} (\mathbb{1} - M_0^{(0,0)}) | \psi \rangle &= \langle \psi | P_x^{(a)} (\mathbb{1} - M_0^{(0,0)})^2 Q_y^{(b)} | \psi \rangle \\ &= \langle \psi | (\mathbb{1} - M_0^{(0,0)}) P_x^{(a)} Q_y^{(b)} | \psi \rangle. \end{aligned}$$

If a = 0,

$$P_x^{(0)}(\mathbb{1}-M_0^{(0,0)})|\psi\rangle = M_{t_x}^{(0,0)}(\mathbb{1}-M_0^{(0,0)})|\psi\rangle = M_{t_x}^{(0,0)}|\psi\rangle - |\psi_0\rangle,$$

where $|\psi_0\rangle = M_{t_x}^{(0,0)} M_0^{(0,0)} |\psi\rangle$. Observe that, because \mathfrak{C}_p is synchronous,

$$|\psi_0\rangle = M_{t_x}^{(0,0)}(M_0^{(0,0)})^2 |\psi\rangle = M_{t_x}^{(0,0)} M_0^{(0,0)} N_0^{(0,0)} |\psi\rangle = N_0^{(0,0)} |\psi_0\rangle.$$

Hence,

$$\begin{split} \langle \psi | (\mathbb{1} - M_0^{(0,0)}) P_x^{(a)} Q_y^{(b)} | \psi \rangle &= \langle \psi | M_{t_x}^{(0,0)} Q_y^{(b)} | \psi \rangle - \langle \psi_0 | Q_y^{(b)} | \psi \rangle \\ &= \langle \psi | M_{t_x}^{(0,0)} Q_y^{(b)} | \psi \rangle - \langle \psi_0 | N_0^{(0,0)} Q_y^{(b)} | \psi \rangle \\ &= \langle \psi | M_{t_x}^{(0,0)} Q_y^{(b)} | \psi \rangle - \langle \psi_0 | \psi_j \rangle, \end{split}$$

where j = 0 if b = 0 and j = p otherwise. Since $|\psi_0\rangle$ and $|\psi_p\rangle$ are 1 and -1-eigenvectors of M_{t_1} by Equation (6), $\langle \psi_0 | \psi_p \rangle = 0$. If a = 1, the same calculation holds except that $\langle \psi_0 |$ is replaced by $\langle \psi_p |$ and $M_{t_x}^{(0,0)}$ is replaced by $M_{t_x}^{(0,1)}$. Thus in all cases, we can calculate

$$\langle \psi | (\mathbb{1} - M_0^{(0,0)}) P_x^{(a)} Q_y^{(b)} (\mathbb{1} - M_0^{(0,0)}) | \psi \rangle$$

from \mathfrak{C}_p , and direct calculation shows that the induced correlation is \mathfrak{C}'_p .

We can now define $|\psi_1\rangle$ based on Propositions 6.2 and 6.5.

Corollary 6.6. Let $(|\psi\rangle, \{M_x^{(a)} | a \in [3] \times [2]\}, x \in I, \{N_y^{(b)} | b \in [3] \times [2]\}, y \in I)$ be a good strategy for \mathfrak{C}_p , and let

$$|\psi_1\rangle = \frac{1}{2} (M_1^{(0,0)} + iM_2 M_1^{(1,0)} - iM_2 M_1^{(0,0)} + M_1^{(1,0)}) |\psi\rangle, \tag{15}$$

where $M_2 = M_2^{(0,0)} - M_2^{(1,0)}$. Then

$$|\psi_1\rangle = \frac{1}{2} (N_1^{(0,0)} - iN_2 N_1^{(1,0)} + iN_2 N_1^{(0,0)} + N_1^{(1,0)}) |\psi\rangle, \tag{16}$$

$$\||\psi_1\rangle\|^2 = \frac{1}{p} = \langle \psi|\psi_1\rangle, \tag{17}$$

$$M_{t_1}M_{t_2}|\psi_1\rangle = \omega_p|\psi_1\rangle$$
, and (18)

$$N_{t_1}N_{t_2}|\psi_1\rangle = \omega_p^{-1}|\psi_1\rangle,\tag{19}$$

where $M_{t_x} = M_{t_x}^{(0,0)} - M_{t_x}^{(0,1)}$ and $N_{t_x} = N_{t_x}^{(0,0)} - N_{t_x}^{(0,1)}$ for x = 1, 2.

Proof. Suppose $(|\psi\rangle, \{M_x^{(a)} \mid a \in [3] \times [2]\}, x \in I, \{N_y^{(b)} \mid b \in [3] \times [2]\}, y \in I)$ is a good strategy for \mathfrak{C}_p . Then

$$\langle \psi | \psi_1 \rangle = \frac{1}{2} \langle \psi | (N_1^{(0,0)} + i(M_2^{(0,0)} - M_2^{(1,0)}) N_1^{(1,0)} - i(M_2^{(0,0)} - M_2^{(1,0)}) N_1^{(0,0)} + N_1^{(1,0)}) | \psi \rangle = \frac{1}{p} \langle \psi | (N_1^{(0,0)} + i(M_2^{(0,0)} - M_2^{(1,0)}) N_1^{(1,0)} - i(M_2^{(0,0)} - M_2^{(1,0)}) N_1^{(0,0)} + N_1^{(1,0)}) | \psi \rangle = \frac{1}{p} \langle \psi | (N_1^{(0,0)} + i(M_2^{(0,0)} - M_2^{(1,0)}) N_1^{(1,0)} - i(M_2^{(0,0)} - M_2^{(1,0)}) N_1^{(0,0)} + N_1^{(1,0)}) | \psi \rangle = \frac{1}{p} \langle \psi | (N_1^{(0,0)} + i(M_2^{(0,0)} - M_2^{(1,0)}) N_1^{(1,0)} - i(M_2^{(0,0)} - M_2^{(1,0)}) N_1^{(1,0)} + N_1^{(1,0)}) | \psi \rangle = \frac{1}{p} \langle \psi | (N_1^{(0,0)} + i(M_2^{(0,0)} - M_2^{(1,0)}) N_1^{(1,0)} - i(M_2^{(0,0)} - M_2^{(1,0)}) N_1^{(1,0)} + N_1^{(1,0)}) | \psi \rangle = \frac{1}{p} \langle \psi | (N_1^{(0,0)} + i(M_2^{(0,0)} - M_2^{(1,0)}) N_1^{(1,0)} + N_1^{(1,0)}) | \psi \rangle$$

by Table 1c. Let $(|\psi'\rangle, \{P_x^{(a)} \mid a \in [3]\}, x \in [5], \{Q_y^{(b)} \mid b \in [3]\}, y \in [5])$ be the strategy for \mathfrak{C}'_p from Proposition 6.5, and let $|\phi_1\rangle = \frac{1}{2} \left(P_3^{(0)} + iP_4P_3^{(1)} - iP_4P_3^{(0)} + P_3^{(1)} \right) |\psi\rangle$ as in Proposition 6.2. Expanding $P_x^{(a)}$, we see that

$$\begin{split} \|(\mathbb{1} - M_0^{(0,0)})|\psi\rangle\| \cdot |\phi_1\rangle &= \frac{1}{2} (M_1^{(0,0)} - iM_2 M_1^{(1,0)} + iM_2 M_1^{(0,0)} + M_1^{(1,0)})(\mathbb{1} - M_0^{(0,0)})|\psi\rangle \\ &= \frac{1}{2} (M_1^{(0,0)} - iM_2 M_1^{(1,0)} + iM_2 M_1^{(0,0)} + M_1^{(1,0)})|\psi\rangle \\ &= |\psi_1\rangle, \end{split}$$

where the second equality follows from Equation (12). Equation (16) follows from expanding Equation (4) similarly. Equations (17) to (19) follow immediately from Proposition 6.2 and the fact that $\|(\mathbb{1} - M_0^{(0,0)})|\psi\rangle\|^2 = (p-1)/p$.

We can now finish the proof of Theorem 6.4.

Proof of Theorem 6.4. We have already defined $|\psi_0\rangle$ and $|\psi_p\rangle$ in Equation (5), and $|\psi_1\rangle$ in Corollary 6.6. Next we define $|\psi_i\rangle$ for $2 \le j \le p - 1$ as

$$|\psi_j\rangle = (U_A U_B)^{\log_r j} |\psi_1\rangle$$

where $\log_r j$ is the discrete log of j modulo p (in other words, $\log_r j = a$ where $r^a \equiv j \pmod{p}$). The discrete log is defined for all $1 \le j \le p - 1$ because r is a primitive root of p. Since U_A and U_B are unitary, $||\psi_j\rangle||^2 = 1/p$. To prove

$$(M_{t_1}M_{t_2})|\psi_j\rangle = \omega_p^j |\psi_j\rangle$$
, and $(N_{t_1}N_{t_2})|\psi_j\rangle = \omega_p^{-j} |\psi_j\rangle$, (20)

observe that since $M_{t_i}|\psi\rangle = N_{t_i}|\psi\rangle$, $(M_{t_1}M_{t_2})^n|\psi\rangle = (N_{t_1}N_{t_2})^{-n}|\psi\rangle$, and similarly $U_A^n|\psi\rangle = U_B^{-n}|\psi\rangle$. Thus

$$(M_{t_1}M_{t_2})^n U_A |\psi\rangle = (M_{t_1}M_{t_2})^{n-1} U_A (M_{t_1}M_{t_2})^r |\psi\rangle$$

= $(N_{t_1}N_{t_2})^{-r} (M_{t_1}M_{t_2})^{n-1} U_A |\psi\rangle$
...
= $(N_{t_1}N_{t_2})^{-nr} U_A |\psi\rangle$
= $U_A (M_{t_1}M_{t_2})^{nr} |\psi\rangle.$

Hence

$$(M_{t_1}M_{t_2})U_A^n|\psi\rangle = (M_{t_1}M_{t_2})U_A(U_B^{\dagger})^{n-1}|\psi\rangle$$

= $(U_B^{\dagger})^{n-1}U_A(M_{t_1}M_{t_2})^r|\psi\rangle$
= $(U_B^{\dagger})^{n-2}U_A(M_{t_1}M_{t_2})^rU_A|\psi\rangle$
= $(U_B^{\dagger})^{n-2}U_A^2(M_{t_1}M_{t_2})^{r^2}|\psi\rangle$
...
= $U_A^n(M_{t_1}M_{t_2})^{r^n}|\psi\rangle.$

Then

$$\begin{split} (M_{t_1}M_{t_2})|\psi_j\rangle &= (M_{t_1}M_{t_2})(U_AU_B)^{\log_r j}|\psi_1\rangle \\ &= \frac{1}{2}U_B^{\log_r j}(N_1^{(0,0)} - iN_2N_1^{(1,0)} + iN_2N_1^{(0,0)} + N_1^{(1,0)})(M_{t_1}M_{t_2})U_A^{\log_r j}|\psi\rangle \\ &= \frac{1}{2}U_B^{\log_r j}(N_1^{(0,0)} - iN_2N_1^{(1,0)} + iN_2N_1^{(0,0)} + N_1^{(1,0)})U_A^{\log_r j}(M_{t_1}M_{t_2})^j|\psi\rangle \\ &= (U_AU_B)^{\log_r j}(M_{t_1}M_{t_2})^j|\psi_1\rangle = \omega_p^j(U_AU_B)^{\log_r j}|\psi_1\rangle = \omega_p^j|\psi_j\rangle, \end{split}$$

as desired.

Let

$$|\psi'
angle = \sum_{j=0}^p |\psi_j
angle.$$

Since eigenvectors with different eigenvalues are orthogonal, $\langle \psi_j | \psi_k \rangle = 0$ for $0 \le j \ne k \le p$. As a result, $|||\psi'\rangle|| = 1$. If j = 0 or p, then $\langle \psi | \psi_j \rangle = |||\psi_j\rangle||^2 = 1/2p$. If $1 \le j \le p - 1$, then $\langle \psi | \psi_j \rangle = \langle \psi | (U_A U_B)^{\log_r j} | \psi_1 \rangle = \langle \psi | \psi_1 \rangle = 1/p$ using Corollary 6.6 and the fact that $U_A U_B | \psi \rangle = |\psi\rangle$. Thus,

$$\langle \psi | \psi'
angle = \langle \psi | \psi_0
angle + \langle \psi | \psi_p
angle + \sum_{j=1}^{p-1} \langle \psi | \psi_j
angle$$

 $= \frac{1}{2p} + \frac{1}{2p} + (p-1)\frac{1}{p} = 1,$

implying that $|\psi\rangle = |\psi'\rangle$. We conclude that

$$(M_{t_1}M_{t_2})^p |\psi\rangle = (M_{t_1}M_{t_2})^p (\sum_{j=0}^p |\psi_j\rangle) = \sum_{j=0}^p \omega_p^{jp} |\psi_j\rangle = |\psi\rangle,$$

which completes the proof.

7 Membership problems

Recall that Membership $(n_A, n_B, m_A, m_B)_{t,\mathbb{K}}$, where $t \in \{q, qs, qa, qc\}$ and \mathbb{K} is a subfield of \mathbb{R} , is defined in Section 1 as the problem of deciding if a correlation $P \in \mathbb{K}^{n_A n_B m_A m_B}$ is in the correlation set $C_t(n_A, n_B, m_A, m_B)$. In this section, we let $\mathbb{K} = \mathbb{K}_0 \cap \mathbb{R}$, where \mathbb{K}_0 is the subfield of \mathbb{C} generated by the roots of unity ω_n^k for $k, n \in \mathbb{Z}$. We then drop the subscript \mathbb{K} when referring to membership problems. The hardness of Membership $(n_A, n_B, m_A, m_B)_t$ is related to the hardness of a more general problem:

Problem Intersection $(n_A, n_B, m_A, m_B)_t$. Given a set of correlations $F \subset \mathbb{K}^{n_A n_B m_A m_B}$ with constants n_A, n_B, m_A and m_B , is $F \cap C_t(n_A, n_B, m_A, m_B) \neq \emptyset$?

Proposition 7.1. *For fixed* n_A , n_B , m_A , $m_B \in \mathbb{N}$ *and* $t \in \{q, qs, qa, qc\}$,

Intersection $(n_A, n_B, m_A, m_B)_t$ and Membership $(n_A, n_B, m_A, m_B)_t$

are equivalent under Cook reduction.

Proof. If D_M is a decider for Membership $(n_A, n_B, m_A, m_B)_t$, we can decide if $F \cap C_t(n_A, n_B, m_A, m_B) = \emptyset$ by running D_M on all the elements of F. If D_I is a decider for Intersection $(n_A, n_B, m_A, m_B)_t$, we can decide if a correlation $P \in C_t(n_A, n_B, m_A, m_B)$ by running D_I on $\{P\}$.

The main result of this section is the following.

Theorem 7.2. For every recursively enumerable set X of positive integers, there exists $N \in \mathbb{N}$ and a computable family of finite sets of correlations $\{F_n \mid n \in \mathbb{N}\}$, where $F_n \subset \mathbb{K}^{N^2 \times 8^2}$, such that

$$F_n \cap C_{qc}(N, N, 8, 8) = \emptyset \text{ if } n \in X, \text{ and}$$

$$F_n \cap C_{aa}(N, N, 8, 8) \neq \emptyset \text{ if } n \notin X.$$

Before proving Theorem 7.2, we first observe that Theorem 1.1 follows directly from Theorem 7.2.

Proof of Theorem 1.1. Let *X* be a RE-complete set of positive integers, and let *N* and $F_n \subset \mathbb{K}^{N^2 \times 8^2}$ be as in Theorem 7.2. Set $\alpha = \max(N, 8)$, and suppose n_A , n_B , m_A , $m_B \ge \alpha$. For any $n \in \mathbb{N}$ and $C \in F_n$, define $C' \in \mathbb{K}^{n_A n_B m_A m_B}$ by

$$C'(a,b \mid x,y) = \begin{cases} C(a,b \mid \delta(x), \delta(y)) & \text{if } a, b < 8\\ 0 & \text{otherwise} \end{cases}$$

where $\delta : \mathbb{N} \to \mathbb{N}$ is defined by

$$\delta(i) = \begin{cases} i & \text{if } i < N \\ N & \text{otherwise} \end{cases}$$

It follows easily from the definitions that

 $C' \in C_t(n_A, n_B, m_A, m_B)$ if and only if $C \in C_t(N, N, 8, 8)$.

Hence if $F'_n = \{C' \mid C \in F_n\}$, then

 $F'_n \cap C_t(n_A, n_B, m_A, m_B) \neq \emptyset$ if and only if $F_n \cap C_t(N, N, 8, 8) \neq \emptyset$.

Since $C_{qa}(n_A, n_B, m_A, m_B) \subseteq C_{qc}(n_A, n_B, m_A, m_B)$, Theorem 7.2 implies that

$$F'_n \cap C_t(n_A, n_B, m_A, m_B) \neq \emptyset$$
 if and only if $n \notin X$

for both t = qa and t = qc. Thus Intersection $(n_A, n_B, m_A, m_B)_t$ is coRE-hard for t = qa, qc. By Proposition 7.1, Membership $(n_A, n_B, m_A, m_B)_t$ is also coRE-hard.

Although we take $m_A, m_B \ge \alpha$ in the proof, note that it is sufficient to choose m_A , $m_B \ge 8$. Also, it has been shown that Membership $(n_A, n_B, m_A, m_B)_{qc}$ is in coRE [NPA08]. Hence, Membership $(n_A, n_B, m_A, m_B)_{qc}$ is coRE-complete for $n_A, n_B \ge N$ and $m_A, m_B \ge 8$.

To prove Theorem 7.2, we first construct F_n from X. Recall that an integer r is a primitive root of a prime p if all the integers between 1 and p - 1 are congruent modulo p to some power of r. By a result of Gupta and Murty [GM84], there are integers r which are primitive roots of infinitely many primes. We use a version of this result due to Heath-Brown.

Lemma 7.3 ([HB86], see also [Mur88]). *There exists* $r \in \{2, 3, 5\}$ *such that r is a primitive root of infinitely many primes.*

For the construction of F_n , fix $r \in \{2,3,5\}$ such that r is a primitive root of infinitely many primes. Let p(n) be the n-th prime greater than r for which r is a primitive root. Since we can decide if r is a primitive root of a given prime, the sequence of primes p(1) < p(2) < ... is computable. Let $\Gamma(A)$ be the solution group from Proposition 5.14 for the function p, set X and integer r. Let m and ℓ be the number of rows and columns of A respectively, and note that each row of A has three nonzero entries. Recall that the generating set $\{x_i \mid i \in [\ell]\}$ has special generators x, t_1, t_2, u_1 and u_2 . By reordering the generators, we can take $x = x_0$, which lets us use x for other things. Recall that for perfect correlations associated with Ax = 0, we use question set $[m] \cup \mathcal{X}_{var}$, where $\mathcal{X}_{var} = \{x_i \mid i \in [\ell]\}$ is the set of variables in the system, and answer set \mathbb{Z}_2^3 . For correlations in F_n , the question set is $\mathcal{X} = \mathcal{X}_{var} \cup [m] \cup \{m, m + 1, m + 2, (m, t_1), (m, t_2)\}$, and the answer set is $\mathcal{A} = \mathbb{Z}_2^3$. The questions $m, m + 1, m + 2, (m, t_1)$ and (m, t_2) will correspond to questions $0, 1, 2, (0, t_1)$ and $(0, t_2)$ from the correlation $\mathfrak{C}_{p(n)}$.

To define the entries of the correlations in F_n , we use the notations from Definition 4.1. In particular, $I_j = \{k \in [\ell] \mid A(j,k) \neq 0\}$ for $j \in [m]$. Let

$$G_n = \langle x_0, x_1, \dots, x_{\ell-1} : x_j^2 = e \text{ for all } j \in [\ell], [x_j, x_k] = e \text{ if } j, k \in I_i \text{ for some } i, (t_1 t_2)^{p(n)} = e \rangle,$$

so that $\Gamma(A)/\langle (t_1t_2)^{p(n)} = e \rangle$ is a quotient of G_n . G_n is a Coxeter group, so its word problem is decidable [Hum90, Chapter 5]. Specifically, two words w_0 and w_1 over the generators of G_n are equal in G_n if they can both be transformed into a third word using

the transformations

$$x_j^2 \to e,$$

$$x_j x_k \to x_k x_j \text{ if } j, k \in I_i \text{ for some } i,$$

$$\underbrace{t_1 t_2 \dots t_1}_{\text{length } p(n)} \to \underbrace{t_2 t_1 \dots t_2}_{\text{length } p(n)}, \text{ and}$$

$$\underbrace{t_2 t_1 \dots t_2}_{\text{length } p(n)} \to \underbrace{t_1 t_2 \dots t_1}_{\text{length } p(n)}.$$

Since the transformations never increase the length of a word, determining if two words are equal is a finite problem.

Recall that a group algebra like $\mathbb{C}[G_n]$ is a *-algebra under the operation $(\sum_g \alpha_g g)^* = \sum_g \overline{\alpha_g} g^{-1}$. We define a mapping $\sigma : \mathcal{X} \times \mathcal{A} \to \mathbb{C}[G_n]$ from question-answer pairs to selfadjoint projections in $\mathbb{C}[G_n]$ as follows. For $(a_0, a_1) \in \mathbb{Z}_2^2$, let $\#(a_0, a_1)$ be the element of [4] with binary representation (a_0, a_1) .

• When $x \in \mathcal{X}_{var}$,

$$\sigma(x,a) = \begin{cases} \frac{e+(-1)^{a_2}x}{2} & \text{if } (a_0,a_1) = (0,0) \\ 0 & \text{otherwise} \end{cases}$$

• When $x = i \in [m]$,

$$\sigma(i,a) = \prod_{k \in I_i} \frac{e + (-1)^{a_{\phi_i(k)}} x_k}{2}$$

• When $x \in \{m, m+1, m+2\}$,

$$\sigma(x,a) = \begin{cases} \pi_0^{(\#(a_0,a_1))} & \text{if } x = m \text{ and } \#(a_0,a_1) \le 2, a_2 = 0\\ \pi_1^{(\#(a_0,a_1))} & \text{if } x = m+1 \text{ and } \#(a_0,a_1) \le 2, a_2 = 0\\ \pi_2^{(\#(a_0,a_1))} & \text{if } x = m+2 \text{ and } \#(a_0,a_1) \le 2, a_2 = 0\\ 0 & \text{otherwise} \end{cases}$$

where $\pi_i^{(a)} \in \mathbb{C}[\langle t_1, t_2 \rangle] \cong \mathbb{C}[D_{p(n)}]$ is defined in Section 6, Equation (2).

• When $x = (m, t_i)$ for i = 1, 2,

$$\sigma((m,t_1),(a_0,a_1,a_2)) = \begin{cases} \pi_0^{(\#(a_0,a_1))} \left(\frac{e+(-1)^{a_2}t_1}{2}\right) & \text{if } \#(a_0,a_1) < 3\\ 0 & \text{otherwise} \end{cases} \text{ and} \\ \sigma((m,t_2),(a_0,a_1,a_2)) = \begin{cases} \pi_0^{(\#(a_0,a_1))} \left(\frac{e+(-1)^{a_2}t_2}{2}\right) & \text{if } \#(a_0,a_1) < 3\\ 0 & \text{otherwise} \end{cases},$$

where again $\pi_0^{(a)} \in \mathbb{C}[\langle t_1, t_2 \rangle] \cong \mathbb{C}[D_{p(n)}]$ is defined in Section 6, Equation (2).

If $z = \sum_{g} \alpha_{g} g$, let

$$\operatorname{supp}(z) = \{g \in G_n \mid \alpha_g \neq 0\}.$$

Define

$$W_n = \bigcup_{x,y \in \mathcal{X}, a, b \in \mathcal{A}} \operatorname{supp}(\sigma(x, a)\sigma(y, b)),$$

and let

$$\mathcal{F}_n = \{ f : W_n \to \{0,1\} \mid f(e) = 1, f(x_0) = 0, f(g) = 0 \text{ for } g \in \langle t_1, t_2 \rangle \setminus \{e\} \}.$$

Functions $f : W_n \to \{0,1\}$ can be regarded as linear functions $\text{span}_{\mathbb{C}}(W_n) \to \mathbb{C}$ by extending linearly. Hence, given a function $f \in \mathcal{F}_n$, we can define a bipartite correlation C_f for the scenario $(\mathcal{X}, \mathcal{X}, \mathcal{A}, \mathcal{A})$ by $C_f(a, b|x, y) = f(\sigma(x, a)\sigma(y, b))$. These correlations contain a copy of the correlation $\mathfrak{C}_{p(n)}$ from Section 6.

Proposition 7.4. Let $Q = \{t_1, t_2, m, m+1, m+2, (m, t_1), (m, t_2)\} \subseteq \mathcal{X}$ and let $I = \{t_1, t_2, 0, 1, 2, (0, t_1), (0, t_2)\}$ as in Section 6. Let $\alpha : Q \to I$ be the bijection $\alpha(t_i) = t_i, \alpha((m, t_i)) = (0, t_i)$ for i = 1, 2, and $\alpha(m + j) = j$ for $j \in [3]$. If $x, y \in Q$ and $a, b \in \mathcal{A}$ such that $\#(a_0, a_1), \#(b_0, b_1) < 3$, then

$$C_f(a,b|x,y) = \mathfrak{C}_{p(n)}((\#(a_0,a_1),a_2),(\#(b_0,b_1),b_2)|\alpha(x),\alpha(y)),$$
(21)

for all $f \in \mathcal{F}_n$. Furthermore, if $S = (|\psi\rangle \in \mathcal{H}, \{M_x^{(a)}, N_x^{(a)} | a \in \mathcal{A}\}, x \in \mathcal{X})$ is a good strategy for C_f , then $S' = (|\psi\rangle \in \mathcal{H}, \{\tilde{M}_x^{(a)}, \tilde{N}_x^{(a)} | a \in [3] \times [2]\}, x \in I)$ is a good strategy for $\mathfrak{C}_{p(n)}$, where

$$\tilde{M}_{x}^{(\#(a_{0},a_{1}),a_{2})} = M_{\alpha^{-1}(x)}^{(a)} \text{ and } \tilde{N}_{x}^{(\#(a_{0},a_{1}),a_{2})} = N_{\alpha^{-1}(x)}^{(a)} \text{ for } x \in I, a \in \mathcal{A}, \#(a_{0},a_{1}) < 3.$$

Proof. Since the subgroup generated by t_1 and t_2 in G_n is a parabolic subgroup, $\langle t_1, t_2 \rangle \subseteq G_n$ is isomorphic to $D_{p(n)}$. By construction, if $f \in \mathcal{F}_n$ and $g \in \langle t_1, t_2 \rangle$ then f(g) = 1 if and only if g = e. Hence when $x, y \in Q$ and $\#(a_0, a_1), \#(b_0, b_1) < 3$,

$$f(\sigma(x,a)\sigma(y,b)) = \langle e|L(\sigma(x,a))R(\sigma(y,b))|e\rangle,$$

where $|e\rangle \in \ell^2 D_{p(n)}$ and $L : \mathbb{C}[D_{p(n)}] \to \mathcal{U}(\ell^2 D_{p(n)})$ and $R : \mathbb{C}[D_{p(n)}] \to \mathcal{U}(\ell^2 D_{p(n)})$ are the left and right regular representations of $\mathbb{C}[D_{p(n)}]$. Since $L(\sigma(x, a)) = \tilde{M}_{\alpha(x)}^{(\#(a_0, a_1), a_2)}$ and $R(\sigma(y, b)) = \tilde{N}_{\alpha(y)}^{(\#(b_0, b_1), b_2)}$ from the definition of \mathfrak{C}_p in Section 6, Equation (21) follows.

If $a \in \mathcal{A}$ with $\#(a_0, a_1) = 3$, then $C_f(a, b | x, y) = C_f(b, a | y, x) = 0$ for all $x \in Q, y \in \mathcal{X}$ and $b \in \mathcal{A}$. Hence $\langle \psi | M_x^{(a)} | \psi \rangle = \langle \psi | N_x^{(a)} | \psi \rangle = 0$ for all $x \in Q$. Since *S* is a good strategy, $M_x^{(a)} = N_x^{(a)} = 0$. We conclude that $\{\tilde{M}_x^{(a)} \mid a \in [3] \times [2]\}$ and $\{\tilde{N}_x^{(a)} \mid a \in [3] \times [2]\}$ are projective measurements for all $x \in I$, and thus *S'* is a good strategy for $\mathfrak{C}_{p(n)}$. \Box Finally, we are ready to define *F*_{*n*}:

$$F_n = \{C_f, f \in \mathcal{F}_n \mid C_f |_{\mathcal{A} \times \mathcal{A} \times (\mathcal{X}_{var} \cup [m]) \times (\mathcal{X}_{var} \cup [m])} \text{ is a perfect correlation for } Ax = 0\},\$$

where $C_f|_{\mathcal{A}\times\mathcal{A}\times(\mathcal{X}_{var}\cup[m])\times(\mathcal{X}_{var}\cup[m])}$ is the restriction of C_f to the question set $\mathcal{X}\cup[m]$, and A is the matrix fixed above. Since support sets are finite, W_n and \mathcal{F}_n are finite, and hence F_n is finite. Since the word problem of G_n is decidable, the sets W_n and \mathcal{F}_n are computable from n. Whether $C_f|_{\mathcal{A}\times\mathcal{A}\times(\mathcal{X}_{var}\cup[m])\times(\mathcal{X}_{var}\cup[m])}$ is a perfect correlation for Ax = 0 is also decidable, and therefore F_n is computable from n.

Proposition 7.5. If $n \in X$, $F_n \cap C_{qc}(\mathcal{X}, \mathcal{X}, \mathcal{A}, \mathcal{A}) = \emptyset$.

Proof. Assume $C_f \in F_n \cap C_{qc}(\mathcal{X}, \mathcal{X}, \mathcal{A}, \mathcal{A})$ for some *f*. By Proposition 3.9, there is a good commuting-operator strategy

$$S = (|\psi\rangle \in \mathcal{H}, \{M_x^{(a)}, N_x^{(a)} \mid a \in \mathcal{A}\}, x \in \mathcal{X})$$

for C_f . Since $C_f|_{\mathcal{A}\times\mathcal{A}\times(\mathcal{X}_{var}\cup[m])\times(\mathcal{X}_{var}\cup[m])}$ is a perfect correlation for Ax = 0, and $S' = (|\psi\rangle, \{M_x^{(a)}, N_x^{(a)} | a \in \mathcal{A}\}, x \in \mathcal{X}_{var} \cup [m])$ is a strategy for this perfect correlation, Proposition 4.3 states that there exists a subspace \mathcal{H}_0 of \mathcal{H} containing $|\psi\rangle$ and unitary operators $M(x_i)$ and $N(x_i)$ for $x_i \in \mathcal{X}_{var}$ on \mathcal{H} inducing commuting representations Φ_M and Φ_N of $\Gamma(A)$ on \mathcal{H}_0 . The operators $M(x_i)$ and $N(x_i)$ are defined as

$$M(x_i) = M_{x_i}^{(0,0,0)} - M_{x_i}^{(0,0,1)}$$
 and $N(x_i) = N_{x_i}^{(0,0,0)} - N_{x_i}^{(0,0,1)}$.

By Proposition 5.14, the generators t_1 , t_2 , u_1 and u_2 of $\Gamma(A)$ satisfy the relation $u_2u_1t_1t_2u_1u_2 = (t_1t_2)^r$, so

$$(M(t_1)M(t_2))(M(u_1)M(u_2))|\psi\rangle = (M(u_1)M(u_2))(M(t_1)M(t_2))^r|\psi\rangle \text{ and } (N(t_1)N(t_2))(N(u_1)N(u_2))|\psi\rangle = (N(u_1)N(u_2))(N(t_1)N(t_2))^r|\psi\rangle.$$

Let $S' = (|\psi\rangle \in \mathcal{H}, \{\tilde{M}_x^{(a)}, \tilde{N}_x^{(a)} | a \in [3] \times [2]\}, x \in I)$ be the strategy for $\mathfrak{C}_{p(n)}$ from Proposition 7.4. Notice that $\tilde{M}_{t_i} := \tilde{M}_{t_i}^{(0,0)} - \tilde{M}_{t_i}^{(0,1)} = M_{t_i}^{(0,0,0)} - M_{t_i}^{(0,0,1)} = M(t_i)$ for i = 1, 2. Similarly $\tilde{N}_{t_i} := \tilde{N}_{t_i}^{(0,0)} - \tilde{N}_{t_i}^{(0,1)} = N(t_i)$ for i = 1, 2. Let $U_A = M(u_1)M(u_2)$ and $U_B = N(u_1)N(u_2)$. By Proposition 4.3, $M(x_i)N(x_i)|\psi\rangle = |\psi\rangle$, so $U_A U_B |\psi\rangle = |\psi\rangle$. Hence U_A and U_B satisfy the conditions of Theorem 6.4 with the strategy S'. We conclude that

$$\Phi_M((t_1t_2)^{p(n)})|\psi\rangle = (M(t_1)M(t_2))^{p(n)}|\psi\rangle = |\psi\rangle.$$

By part (3) of Proposition 4.3, Φ_M descends to a representation of $\Gamma(A) / \langle (t_1 t_2)^{p(n)} = e \rangle$ on \mathcal{H}_0 . On the other hand,

$$\langle \psi | M(x_0) | \psi \rangle = \langle \psi | M_{x_0}^{(0,0,0)} - M_{x_0}^{(0,0,1)} | \psi \rangle = f(\sigma(x_0, (0,0,0)) - \sigma(x_0, (0,0,1))) = f(x_0) = 0$$

by the definition of \mathcal{F}_n . Hence $\Phi_M(x_0) \neq \mathbb{1}_{\mathcal{H}_0}$, which implies that $x \neq e$ in $\Gamma(A) / \langle (t_1 t_2)^{p(n)} = e \rangle$. By part (b) of Proposition 5.14, $n \notin X$.

Proposition 7.6. If $n \notin X$, $F_n \cap C_{qa}(\mathcal{X}, \mathcal{X}, \mathcal{A}, \mathcal{A}) \neq \emptyset$.

Proof. For this proof, let $\Gamma_n := \Gamma(A)/\langle (t_1t_2)^{p(n)} = e \rangle$ and let $W_n^{fa} = \{g \in W_n \mid g \neq e \text{ in } \Gamma_n^{fa}\}$. Suppose $n \notin X$, and define $f : W_n \to \mathbb{C}$ by f(g) = 0 if $g \in W_n^{fa}$ and f(g) = 1 if $g \in W_n \setminus W_n^{fa}$. (The function f is the pullback of the canonical trace on Γ_n^{fa} to G_n and restricted to W_n .) Since $n \notin X$, $x_0 \in W_n^{fa}$ by part (c) of Proposition 5.14, so $f(x_0) = 0$. By parts (d) and (e) of Proposition 5.14, the set $\langle t_1, t_2 \rangle \setminus \{e\} \subseteq W_n^{fa}$ as well. Thus $f \in \mathcal{F}_n$.

In the rest of proof, we show that $C_f \in C_{qa}(\tilde{\mathcal{X}}, \mathcal{X}, \mathcal{A}, \tilde{\mathcal{A}})$. Let $\pi : \mathcal{F}(\mathcal{X}_{var}) \to G_n$ be the quotient homomorphism. For each $x \in \mathcal{X}$ and $a \in \mathcal{A}$, let $\tilde{\sigma}(x, a) \in \mathbb{C}[\mathcal{F}(\mathcal{X}_{var})]$ be a lift of $\sigma(x, a)$, i.e. an element such that $\pi(\tilde{\sigma}(x, a)) = \sigma(x, a)$. Let

$$\tilde{W}_n = \bigcup_{x,y \in \mathcal{X}, b \in \mathcal{A}} \operatorname{supp}(\tilde{\sigma}(x,a)\tilde{\sigma}(y,b))$$

and

$$\tilde{W}_n^{fa} = \{ g \in \tilde{W}_n \mid g \neq e \text{ in } \Gamma_n^{fa} \}.$$

Note that if $w \in \tilde{W}_n$, then $\pi(w) \in W_n$, and $w \in \tilde{W}_n^{fa}$ if and only if $\pi(w) \in W_n^{fa}$. The polynomial $\tilde{\sigma}(x,a)^2 - \tilde{\sigma}(x,a)$ is not necessarily 0 in $\mathbb{C}[\mathcal{F}(\mathcal{X}_{var})]$, but is 0 in $\mathbb{C}[\Gamma_n]$, and the same is true for the polynomials $\tilde{\sigma}(x,a)^* - \tilde{\sigma}(x,a)$, $a \in \mathcal{A}$, $x \in \mathcal{X}$, $\tilde{\sigma}(x,a)\tilde{\sigma}(x,b)$, $a \neq b$, $x \in \mathcal{X}$, and $\sum_{a \in \mathcal{A}} \tilde{\sigma}(x,a) - e$, $x \in \mathcal{X}$. By Lemma 2.4, there is a constant *c* such that for all ϵ -approximate representation $\rho : \mathcal{F}(\mathcal{X}_{var}) \to \mathcal{U}(\mathbb{C}^d)$ of Γ_n

$$\begin{aligned} \|\rho(\tilde{\sigma}(x,a))\|_{op} &\leq c, \quad \|\rho(\tilde{\sigma}(x,a)^2 - \tilde{\sigma}(x,a))\| \leq c\epsilon, \quad \|\rho(\tilde{\sigma}(x,a)^* - \tilde{\sigma}(x,a))\| \leq c\epsilon, \\ \|\rho(\tilde{\sigma}(x,a)\tilde{\sigma}(x,b))\| &\leq c\epsilon, \text{ and } \|\rho(\sum_{a' \in \mathcal{A}} \tilde{\sigma}(x,a') - e)\| \leq c\epsilon \end{aligned}$$

for all $x \in \mathcal{X}$ and $a \neq b \in \mathcal{A}$. By Proposition 2.3, for any $\epsilon, \zeta > 0$ there is an ϵ -approximate representation $\rho : \mathcal{F}(\mathcal{X}_{var}) \to \mathcal{U}(\mathbb{C}^d)$ of Γ_n , where d depends on ϵ and ζ , such that $0 \leq \tilde{\mathrm{Tr}}(\rho(w)) \leq \zeta$ for each $w \in W_n^{fa}$, and $1 - \zeta \leq \tilde{\mathrm{Tr}}(\rho(w)) \leq 1$ for each $w \in W_n \setminus W_n^{fa}$. For $x \in \mathcal{X}$ and $a \in \mathcal{A}$, let

$$\tilde{M}_x^a = \rho(\tilde{\sigma}(x,a)).$$

For all $x, y \in \mathcal{X}$ and $a, b \in \mathcal{A}$, we can write $\tilde{\sigma}(x, a)\tilde{\sigma}(y, b) = \sum_{g \in \tilde{W}_n} u_g g$ for some $u_g \in \mathbb{R}$, where $|u_g| < 1$ for all $g \in \tilde{W}_n$ by definition of $\tilde{\sigma}(x, a)$. Thus

$$\begin{aligned} |C_f(a,b \mid x,y) - \tilde{\mathrm{Tr}}(\tilde{M}_x^a \tilde{M}_y^b)| &\leq \sum_{g \in \tilde{W}_n} |u_g| |f(\pi(g)) - \tilde{\mathrm{Tr}}(\rho(g))| \\ &= \sum_{g \in \tilde{W}_n^{fa}} |u_g| |0 - \tilde{\mathrm{Tr}}(\rho(g))| + \sum_{g \in \tilde{W}_n \setminus \tilde{W}_n^{fa}} |u_g| |1 - \tilde{\mathrm{Tr}}(\rho(g))| \\ &\leq |\tilde{W}_n| \zeta. \end{aligned}$$

Unfortunately, $\{\tilde{M}_x^a \mid a \in A\}$ may not be a measurement. However, by Lemma 2.5 there are projective measurements $\{M_x^a \mid a \in A\}$, $x \in \mathcal{X}$, such that

$$\|M_x^a - \tilde{M}_x^a\| \leq \Delta(c, 8)c\epsilon$$

for all $x \in \mathcal{X}$ and $a \in \mathcal{A}$ (where 8 comes from the size of \mathcal{A}). Then

$$\begin{split} |\tilde{\mathrm{Tr}}(M^a_x M^b_y) - \tilde{\mathrm{Tr}}(\tilde{M}^a_x \tilde{M}^b_y)| &\leq \|M^a_x M^b_y - \tilde{M}^a_x \tilde{M}^b_y\| \\ &\leq \|M^a_x\|_{op} \|M^b_y - \tilde{M}^b_y\| + \|\tilde{M}^b_y\|_{op} \|M^a_x - \tilde{M}^a_x\| \\ &\leq (1+c) \cdot \Delta(c,8)c\epsilon. \end{split}$$

Let $|\psi\rangle = 1/\sqrt{d} \sum_{i=1}^{d} |i\rangle \otimes |i\rangle \in \mathbb{C}^{d} \otimes \mathbb{C}^{d}$, where $|i\rangle$ is the i-th standard basis of \mathbb{C}^{d} , and let $N_{x}^{a} = (M_{x}^{a})^{T}$. Since $\langle \psi | M_{x}^{a} \otimes N_{y}^{b} | \psi \rangle = \tilde{\mathrm{Tr}}(M_{x}^{a}M_{y}^{b})$, we conclude that

$$\begin{aligned} |\langle \psi | M_x^a \otimes N_y^b | \psi \rangle - C_f(a, b \mid x, y)| &\leq |\tilde{\mathrm{Tr}}(M_x^a M_y^b) - \tilde{\mathrm{Tr}}(\tilde{M}_x^a \tilde{M}_y^b)| + |\tilde{\mathrm{Tr}}(\tilde{M}_x^a \tilde{M}_y^b) - C_f(a, b \mid x, y)| \\ &\leq (1+c) \cdot \Delta(c, 8)c\epsilon + |\tilde{W}_n|\zeta. \end{aligned}$$

The correlation defined by $(|\psi\rangle, \{M_x^a \mid a \in \mathcal{A}\}, \{N_x^a \mid a \in \mathcal{A}\}, x \in \mathcal{X})$ belongs to $C_q(\mathcal{X}, \mathcal{X}, \mathcal{A}, \mathcal{A})$. Since ϵ and ζ can be arbitrarily small, we see that $C_f \in \overline{C_q(\mathcal{X}, \mathcal{X}, \mathcal{A}, \mathcal{A})} = C_{qa}(\mathcal{X}, \mathcal{X}, \mathcal{A}, \mathcal{A})$.

The proof of Theorem 7.2 follows immediately from Propositions 7.5 and 7.6.

References

- [BBL⁺06] G. Brassard, H. Buhrman, N. Linden, A. A. Methot, A. Tapp, and F. Unger. A limit on nonlocality in any world in which communication complexity is not trivial. *Phys. Rev. Lett.*, 96(250401), 2006. URL: https://link.aps.org/doi/ 10.1103/PhysRevLett.96.250401, doi:10.1103/PhysRevLett.96.250401.
- [BCLPG20] Johannes Bausch, Toby S Cubitt, Angelo Lucia, and David Perez-Garcia. Undecidability of the spectral gap in one dimension. *Physical Review X*, 10(3):031038, 2020. URL: https://link.aps.org/doi/10.1103/PhysRevX.10.031038, doi: 10.1103/PhysRevX.10.031038.
- [Bel64] John S. Bell. On the Einstein Podolsky Rosen paradox. *Physics*, 1(3):195–200, 1964. URL: https://link.aps.org/doi/10.1103/PhysicsPhysiqueFizika. 1.195, doi:10.1103/PhysicsPhysiqueFizika.1.195.
- [BPA⁺08] Nicolas Brunner, Stefano Pironio, Antonio Acin, Nicolas Gisin, André Allan Méthot, and Valerio Scarani. Testing the dimension of Hilbert spaces. *Physical review letters*, 100(21):210503, 2008. doi:10.1103/PhysRevLett.100.210503.

- [CHTW04] Richard Cleve, Peter Hoyer, Ben Toner, and John Watrous. Consequences and limits of nonlocal strategies. In *Proceedings*. 19th IEEE Annual Conference on Computational Complexity, 2004., pages 236–249, June 2004. doi:10.1109/CCC. 2004.1313847.
- [CLP15] Valerio Capraro, Martino Lupini, and Vladimir Pestov. *Introduction to sofic and hyperlinear groups and Connes' embedding conjecture*, volume 2136. Springer, 2015. doi:10.1007/978-3-319-19333-5.
- [CLS17] Richard Cleve, Li Liu, and William Slofstra. Perfect commuting-operator strategies for linear system games. *Journal of Mathematical Physics*, 58(1):012202, 2017. doi:10.1063/1.4973422.
- [Col20] Andrea Coladangelo. A two-player dimension witness based on embezzlement, and an elementary proof of the non-closure of the set of quantum correlations. *Quantum*, 4:282, 2020. doi:10.22331/q-2020-06-18-282.
- [CS19] Matthew Coudron and William Slofstra. Complexity lower bounds for computing the approximately-commuting operator value of non-local games to high precision. *arXiv preprint arXiv:1905.11635*, 2019.
- [CS20] Andrea Coladangelo and Jalex Stark. An inherently infinite-dimensional quantum correlation. *Nature Communications*, 11(1):3335, 2020. URL: https://doi. org/10.1038/s41467-020-17077-9, doi:10.1038/s41467-020-17077-9.
- [dlS21] Mikael de la Salle. Orthogonalization of positive operator valued measures. *arXiv preprint arXiv:2103.14126*, 2021.
- [DPP19] Ken Dykema, Vern I Paulsen, and Jitendra Prakash. Non-closure of the set of quantum correlations via graphs. *Communications in Mathematical Physics*, 365(3):1125–1142, 2019. doi:10.1007/s00220-019-03301-1.
- [FJVY19] Joseph Fitzsimons, Zhengfeng Ji, Thomas Vidick, and Henry Yuen. Quantum proof systems for iterated exponential time, and beyond. In *Proceedings of the* 51st Annual ACM SIGACT Symposium on Theory of Computing, pages 473–480. ACM, 2019. doi:10.1145/3313276.3316343.
- [Fri12] Tobias Fritz. Tsirelson's problem and Kirchberg's conjecture. *Reviews in Mathematical Physics*, 24(05):1250012, 2012. doi:10.1142/S0129055X12500122.
- [FSA⁺13] T. Fritz, A. B. Sainz, R. Augusiak, J. B. Brask, R. Chaves, A. Leverrier, and A. Acin. Local orthogonality as a multipartite principle for quantum correlations. *Nature Communications*, 4(2263), 2013. doi:10.1038/ncomms3263.
- [Fu22] Honghao Fu. Constant-sized correlations are sufficient to self-test maximally entangled states with unbounded dimension. *Quantum*, 6:614, January 2022. URL: https://doi.org/10.22331/q-2022-01-03-614, doi:10.22331/ q-2022-01-03-614.

- [GKW⁺18] Koon Tong Goh, Jedrzej Kaniewski, Elie Wolfe, Tamás Vértesi, Xingyao Wu, Yu Cai, Yeong-Cherng Liang, and Valerio Scarani. Geometry of the set of quantum correlations. *Phys. Rev. A*, 97:022104, Feb 2018. URL: https://link.aps. org/doi/10.1103/PhysRevA.97.022104, doi:10.1103/PhysRevA.97.022104.
- [GM84] Rajiv Gupta and M. Ram Murty. A remark on Artin's conjecture. *Inventiones mathematicae*, 78(1):127–130, 1984. URL: https://doi.org/10.1007/ BF01388719, doi:10.1007/BF01388719.
- [HB86] D. R. Heath-Brown. Artin's conjecture for primitive roots. The Quarterly Journal of Mathematics, 37(1):27–38, 03 1986. URL: https://doi.org/10.1093/ qmath/37.1.27, arXiv:https://academic.oup.com/qjmath/article-pdf/ 37/1/27/4354561/37-1-27.pdf, doi:10.1093/qmath/37.1.27.
- [HNN49] Graham Higmann, Bernhard Neumann, and Hanna Neumann. Embedding theorems for groups. *Journal of the London Mathematical Society*, 1(4):247–254, 1949. doi:10.1112/jlms/s1-24.4.247.
- [Hum90] James E. Humphreys. *Reflection Groups and Coxeter Groups*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1990. doi:10.1017/ CB09780511623646.
- [IV12] Tsuyoshi Ito and Thomas Vidick. A multi-prover interactive proof for NEXP sound against entangled provers. In 2012 IEEE 53rd Annual Symposium on Foundations of Computer Science, pages 243–252. IEEE, 2012. doi:10.1109/F0CS. 2012.11.
- [Ji17] Zhengfeng Ji. Compression of quantum multi-prover interactive proofs. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*, pages 289–302, 2017. URL: https://doi.org/10.1145/3055399.3055441, doi:10.1145/3055399.3055441.
- [JNP⁺11] M. Junge, M. Navascués, C. Palazuelos, D. Pérez-García, V. B. Scholz, and R. F. Werner. Connes' embedding problem and Tsirelson's problem. *Journal of Mathematical Physics*, 52(1):012102, 2011. doi:10.1063/1.3514538.
- [JNV⁺20] Zhengfeng Ji, Anand Natarajan, Thomas Vidick, John Wright, and Henry Yuen. $MIP^* = RE. arXiv preprint arXiv:2001.04383, 2020.$
- [Kha82] Olga Kharlampovich. A finitely presented solvable group with unsolvable word problem. *Mathematics of the USSR-Izvestiya*, 19(1):151, 1982.
- [KMS17] Olga Kharlampovich, Alexei Myasnikov, and Mark Sapir. Algorithmically complex residually finite groups. *Bulletin of Mathematical Sciences*, 7(2):309–352, 2017. doi:10.1007/s13373-017-0103-z.
- [KPS18] Se-Jin Kim, Vern Paulsen, and Christopher Schafhauser. A synchronous game for binary constraint systems. *Journal of Mathematical Physics*, 59(3):032201, 2018. doi:10.1063/1.4996867.

- [Lan88] L. J. Landau. Empirical two-point correlation functions. *Foundations of Physics*, 18:449–460, 1988. doi:10.1007/BF00732549.
- [Mas03] Ll. Masanes. Necessary and sufficient condition for quantum-generated correlations. 2003. arXiv:quant-ph/0309137.
- [Min67] Marvin Lee Minsky. *Computation: Finite and Infinite Machines*. Prentice-Hall Englewood Cliffs, 1967. doi:10.5555/1095587.
- [MNY21] Hamoon Mousavi, Seyed Sajjad Nezhadi, and Henry Yuen. Nonlocal Games, Compression Theorems, and the Arithmetical Hierarchy. *arXiv preprint arXiv:2110.04651*, 2021.
- [Mur88] M. Ram Murty. Artin's conjecture for primitive roots. *The Mathematical Intelligencer*, 10(4):59–67, 1988. doi:10.1007/BF03023749.
- [NPA08] Miguel Navascués, Stefano Pironio, and Antonio Acín. A convergent hierarchy of semidefinite programs characterizing the set of quantum correlations. *New Journal of Physics*, 10(7):073013, 2008. doi:10.1088/1367-2630/10/7/073013.
- [NV18] Anand Natarajan and Thomas Vidick. Low-degree testing for quantum states, and a quantum entangled games PCP for QMA. In 2018 IEEE 59th Annual Symposium on Foundations of Computer Science (FOCS), pages 731–742. IEEE, 2018. doi:10.1109/F0CS.2018.00075.
- [NW09] Miguel Navascues and Harald Wunderlich. A glance beyond the quantum model. *Proc. Royal Soc. A*, 466:881–890, 2009. doi:10.1098/rspa.2009.0453.
- [NW19] Anand Natarajan and John Wright. *NEEXP* is Contained in *MIP**. In 2019 *IEEE 60th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 510–518. IEEE, 2019. doi:10.1109/F0CS.2019.00039.
- [Pit08] I. Pitowsky. Geometry of quantum correlations. *Physical Review A*, 77:062109, 2008. URL: https://link.aps.org/doi/10.1103/PhysRevA.97.022104, doi: 10.1103/PhysRevA.97.022104.
- [PPK⁺09] M. Pawlowski, T. Paterek, D. Kaszlikowski, V. Scarani, A. Winter, and M. Zukowski. Information causality as a physical principle. *Nature*, 461:1101– 1104, 2009. doi:10.1038/nature08400.
- [PSS⁺16] Vern I Paulsen, Simone Severini, Daniel Stahlke, Ivan G Todorov, and Andreas Winter. Estimating quantum chromatic numbers. *Journal of Functional Analysis*, 270(6):2188–2222, 2016. doi:10.1016/j.jfa.2016.01.010.
- [Rot12] Joseph J Rotman. *An introduction to the theory of groups*, volume 148. Springer Science & Business Media, 2012. doi:/10.1007/978-1-4612-4176-8.
- [Rus20] Travis B. Russell. Geometry of the set of synchronous quantum correlations. *Journal of Mathematical Physics*, 61:052201, 2020. doi:10.1063/1.5115010.

- [RUV13] Ben W Reichardt, Falk Unger, and Umesh Vazirani. Classical command of quantum systems. *Nature*, 496(7446):456, 2013. doi:10.1038/nature12035.
- [SGAN18] Ana Belén Sainz, Yelena Guryanova, Antonio Acín, and Miguel Navascués. Almost-quantum correlations violate the no-restriction hypothesis. *Phys. Rev. Lett.*, 120:200402, May 2018. URL: https://link.aps.org/doi/10.1103/ PhysRevLett.120.200402, doi:10.1103/PhysRevLett.120.200402.
- [Slo11] William Slofstra. Lower bounds on the entanglement needed to play XOR non-local games. *Journal of Mathematical Physics*, 52(10):102202, 2011. doi: 10.1063/1.3652924.
- [Slo19] William Slofstra. The set of quantum correlations is not closed. In *Forum of Mathematics, Pi*, volume 7. Cambridge University Press, 2019. doi:10.1017/ fmp.2018.3.
- [Slo20] William Slofstra. Tsirelson's problem and an embedding theorem for groups arising from non-local games. *Journal of the American Mathematical Society*, 33(1):1–56, 2020. doi:10.1090/jams/929.
- [SW08] V. B. Scholz and R. F. Werner. Tsirelson's Problem. *arXiv preprint arXiv:0812.4305, 2008.*
- [Tsi87] B.S. Tsirelson. Quantum analogues of the Bell inequalities. The case of two spatially separated domains. *Journal of Soviet Mathematics*, 36(4):557–570, 1987. doi:10.1007/BF01663472.
- [Tsi93] Boris S. Tsirelson. Some results and problems on quantum Bell-type inequalities. *Hadronic Journal Supplement*, 8:329–345, 1993.
- [TVC19] Le Phuc Thinh, Antonios Varvitsiotis, and Yu Cai. Geometric structure of quantum correlators via semidefinite programming. *Phys. Rev. A*, 99:052108, May 2019. URL: https://link.aps.org/doi/10.1103/PhysRevA.99.052108, doi:10.1103/PhysRevA.99.052108.
- [WW01] R. F. Werner and M. M. Wolf. All-multipartite Bell-correlation inequalities for two dichotomic observables per site. *Phys. Rev. A*, 64:032112, Aug 2001. URL: https://link.aps.org/doi/10.1103/PhysRevA.64.032112, doi:10.1103/PhysRevA.64.032112.