

TRANSITION SPACE FOR THE CONTINUITY OF THE LYAPUNOV EXPONENT OF QUASIPERIODIC SCHRÖDINGER COCYCLES

LINGRUI GE, YIQIAN WANG, JIANGONG YOU, AND XIN ZHAO

ABSTRACT. We construct discontinuous point of the Lyapunov exponent of quasiperiodic Schrödinger cocycles in the Gevrey space G^s with $s > 2$. In contrast, the Lyapunov exponent has been proved to be continuous in the Gevrey space G^s with $s < 2$ [17, 37]. This shows that G^2 is the transition space for the continuity of the Lyapunov exponent.

1. INTRODUCTION

Let X be a C^r compact manifold, $A(x)$ be a $SL(2, \mathbb{R})$ -valued function on X and (X, T, μ) be ergodic with μ a normalized T -invariant measure. Dynamical systems on $X \times \mathbb{R}^2$ given by

$$(x, w) \rightarrow (T(x), A(x)w)$$

are called a $SL(2, \mathbb{R})$ -cocycle and denoted by (T, A) . In particular, if $X = \mathbb{T}^n = \mathbb{R}^n/2\pi\mathbb{Z}^n$ and $T = T_\alpha : x \rightarrow x + 2\pi\alpha$ with α independent over \mathbb{Q} , we call (T_α, A) a quasiperiodic $SL(2, \mathbb{R})$ -cocycle, which is simply denoted by (α, A) . If moreover,

$$A(x) = S_{E,v}(x) = \begin{pmatrix} E - v(x) & -1 \\ 1 & 0 \end{pmatrix}$$

with $v(x)$ a 2π -periodic function in each variable, we call $(\alpha, S_{E,v})$ a quasiperiodic Schrödinger cocycle.

The n -th iteration of the cocycle (T, A) is denoted by $(T, A)^n = (T^n, A_n)$ where

$$A_n(x) = \begin{cases} A(T^{n-1}x) \cdots A(x), & n \geq 1 \\ I_2, & n = 0 \\ A_{-n}(T^n x)^{-1}, & n \leq -1 \end{cases}.$$

The (maximum) Lyapunov exponent $L(A)$ of the cocycle is defined as

$$L(A) = \lim_{n \rightarrow \infty} \frac{1}{n} \int_X \ln \|A_n(x)\| d\mu = \inf_n \frac{1}{n} \int_X \ln \|A_n(x)\| d\mu \geq 0.$$

The limit exists and is equal to the infimum since $\{\int_X \ln \|A_n(x)\| d\mu\}_{n \geq 1}$ is a subadditive sequence. Moreover, by Kingman's subadditive ergodic theorem, we also have

$$L(A) = \lim_{n \rightarrow \infty} \frac{1}{n} \ln \|A_n(x)\|$$

for μ -almost every $x \in X$.

Regularity of the Lyapunov exponent (LE) is one of the central subjects in smooth dynamical systems, which depends subtly on the base dynamics T and the smoothness of the matrix A . In the present paper, we are mainly interested in how the regularity of A affects the continuity of LE for quasiperiodic $SL(2, \mathbb{R})$ /Schrödinger cocycles. Our motivation comes from the pioneering (opposite) results on the continuity of the Lyapunov exponent in C^ω and C^∞ spaces.

- For any quasiperiodic $SL(2, \mathbb{R})$ -cocycle, the Lyapunov exponent is always continuous with respect to A in C^ω topology [2, 15].
- the Lyapunov exponent is not always continuous with respect to A in C^∞ topology [47, 48].

It is well-known that the Gevrey spaces G^s , $1 < s < \infty$ are between the C^∞ and analytic spaces. Roughly speaking, a 2π -periodic cocycle map $A \in G^s$ means that $\widehat{A}(k)$, the Fourier coefficients of A , decay sub-exponentially like $\mathcal{O}(e^{-k^{1/s}h})$, while the Fourier coefficients of an analytic function decay exponentially and the Fourier coefficients of a smooth function decay faster than any polynomials. In this paper, we are interested in finding the optimal Gevrey space to ensure the continuity of the Lyapunov exponent. More concretely, we prove that the Lyapunov exponent of quasiperiodic Schrödinger cocycles is discontinuous in the Gevrey space G^s with $s > 2$. In contrast, the Lyapunov exponent is continuous in G^s with $s < 2$ [17, 37]. This shows that G^2 is the transition space for the continuity of the Lyapunov exponent.

It is known that a powerful tool to prove the continuity of LE is the large deviation theorem (LDT) and avalanche principle (AP). Our results in some sense show that LDT breaks down for general $G^s(\mathbb{S}, SL(2, \mathbb{R}))$ -cocycles with $s > 2$. One can also compare our result with the result in [37] where Klein showed LE is continuous with respect to the energies E if the potential is in an open and dense subspace of $G^s(\mathbb{S}^1)$ with $s > 2$ where certain non-degeneracy condition is satisfied. Our result shows that such non-degeneracy condition is necessary as the LE would be discontinuous if the potential is sufficiently “degenerate”.

We finally remark that transition phenomenon seems to be a common phenomenon in quasiperiodic dynamical systems and always attracts people’s attention. For example, it was shown in [16] that any Lagrangian torus with a given unique rotation vector of an integrable Hamiltonian can be destructed by an arbitrarily $C^{2d-\delta}$ -small perturbation. In contrast, it was shown that KAM torus with constant type frequency persists for all $C^{2d+\delta}$ -small perturbations [43]. Thus C^{2d} is the transition space for the persistence of KAM torus.

1.1. Transition phenomena for quasiperiodic Schrödinger operators. The discrete one dimensional quasiperiodic Schrödinger operators on $\ell^2(\mathbb{Z})$ are given by

$$(1.1) \quad (H_{\lambda v, x, \alpha} u)_n = u_{n+1} + u_{n-1} + \lambda v(x + n\alpha)u_n, \quad n \in \mathbb{Z},$$

where $\alpha \in \mathbb{R}^d$ is the frequency, $x \in \mathbb{S}^d$ is the phase, $\lambda \in \mathbb{R}$ is the coupling constant and $v \in C^r(\mathbb{S}^d, \mathbb{R})$ ($r = 0, 1, \dots, \infty, \omega$) is called the potential. The spectral properties of operator (1.1) is closely related to the Schrödinger cocycle $(\alpha, S_{E, \lambda v}) \in \mathbb{S}^d \times C^r(\mathbb{S}^d, SL(2, \mathbb{R}))$. Quasiperiodic Schrödinger operators naturally arise in solid-state physics, describing the influence of an external magnetic field on the electrons of a crystal.

Different from random Schrödinger operators, an important feature of one dimensional quasiperiodic operators is that the family $\{H_{\lambda v, x, \alpha}\}_{\lambda \in \mathbb{R}}$ undergoes a so called metal-insulator transition when $|\lambda|$ changes from small to large. Indeed, besides the metal-insulator transition, various spectral transition phenomena take place for quasiperiodic operators. Here we give some perfect examples.

Example 1.1 (Metal-insulator transition). Assume α is Diophantine¹ and $v(x) = 2 \cos 2\pi x$, the following results were given by Jitomirskaya [29] in 1999,

- $|\lambda| > 1$, $H_{\lambda v, x, \alpha}$ has Anderson localization for a.e. x ,
- $|\lambda| = 1$, $H_{\lambda v, x, \alpha}$ has purely singular continuous for a.e. x ,
- $|\lambda| < 1$, $H_{\lambda v, x, \alpha}$ has purely absolutely continuous spectrum for a.e. x .

Example 1.2 (Sharp spectral transition in frequency). We denote by

$$\beta(\alpha) = \limsup_{k \rightarrow \infty} -\frac{\ln \|k\alpha\|_{\mathbb{R}/\mathbb{Z}}}{|k|},$$

¹ $\alpha \in \mathbb{R}$ is called *Diophantine*, denoted by $\alpha \in \text{DC}(\kappa, \tau)$, if there exist $\kappa > 0$ and $\tau > 1$ such that

$$(1.2) \quad \text{DC}(\kappa, \tau) := \left\{ \alpha \in \mathbb{R} : \inf_{j \in \mathbb{Z}} |n\alpha - j| > \frac{\kappa}{|n|^\tau}, \quad \forall n \in \mathbb{Z} \setminus \{0\} \right\}.$$

where $\|x\|_{\mathbb{R}/\mathbb{Z}} = \text{dist}(x, \mathbb{Z})$. Let $v(x) = 2 \cos 2\pi x$, i.e., the famous almost Mathieu operators,

- $|\lambda| > e^{\beta(\alpha)}$, $H_{\lambda v, x, \alpha}$ has Anderson localization for a.e. x [4, 33]²,
- $1 \leq |\lambda| < e^{\beta(\alpha)}$, $H_{\lambda v, x, \alpha}$ has purely singular continuous for all x [4, 30]³,
- $|\lambda| < 1$, $H_{\lambda v, x, \alpha}$ has purely absolutely continuous spectrum for all x [1].

Example 1.3 (Sharp spectral transition in phase). We denote by

$$\delta(\alpha, x) = \limsup_{k \rightarrow \infty} -\frac{\ln \|2x + k\alpha\|_{\mathbb{R}/\mathbb{Z}}}{|k|}.$$

Let $v(x) = 2 \cos 2\pi x$, i.e., the famous almost Mathieu operators,

- $|\lambda| > e^{\delta(\alpha, x)}$, $H_{\lambda v, x, \alpha}$ has Anderson localization for Diophantine α [34],
- $1 \leq |\lambda| < e^{\delta(\alpha, x)}$, $H_{\lambda v, x, \alpha}$ has purely singular continuous for all α [30, 34].
- $|\lambda| < 1$, $H_{\lambda v, x, \alpha}$ has purely absolutely continuous spectrum for all α [1].

Although the transition phenomenon is common for quasiperiodic Schrödinger operators, however, the exact transition points are usually difficult to obtain as it depends sensitively on the arithmetic properties of the frequency and phase. This paper will give explicit transition space for the continuity of the Lyapunov exponent. For our purpose, we introduce the following space of Gevrey functions and its topology.

For any smooth function f defined on \mathbb{S}^1 , let

$$|f|_{s, K} := \frac{4\pi^2}{3} \sup_k \frac{(1 + |k|)^2}{K^k (k!)^s} |\partial^k f|_{C^0(\mathbb{S}^1)},$$

$G^{s, K}(\mathbb{S}^1) = \{f \in C^\infty(\mathbb{S}^1, \mathbb{R}) \mid |f|_{s, K} < \infty\}$ and $G^s(\mathbb{S}^1) = \cup_{K > 0} G^{s, K}(\mathbb{S}^1)$. Note that $G^{s, K}(\mathbb{S}^1)$ is a Banach space. Obviously, $G^1(\mathbb{S}^1)$ is the space of analytic functions and for any $s \geq 1$, $G^s(\mathbb{S}^1)$ is a subspace of the space of smooth functions. We equip $G^s(\mathbb{S}^1)$ with the usual inductive limit topology. That is, f_n converges to f in $G^s(\mathbb{S}^1)$ -topology if and only if $|f_n - f|_{s, K} \rightarrow 0$ as $n \rightarrow \infty$ for some $K > 0$.

We say $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ is of bounded type if there exists $M > 0$, such that the continued fraction expansion of α , denoted by p_n/q_n satisfying

$$q_{n+1} \leq M q_n, \quad \forall n \in \mathbb{N}.$$

Theorem 1.1. *Assume α is of bounded type, for quasiperiodic Schrödinger cocycle $(\alpha, S_{E, v})$, we have*

- (1) *For any $v \in G^s(\mathbb{S}^1)$ with $s < 2$, the Lyapunov exponent is continuous with respect to v in G^s -topology.*
- (2) *There exists $v_0 \in G^s(\mathbb{S}^1)$ with $s > 2$, such that the Lyapunov exponent is discontinuous at v_0 in G^s -topology.*

Remark 1.1. Part (1) of Theorem 1.1 was recently proved in [17] (Theorem 6.3 of [17]), we list here for completeness. The main result of the present paper is part (2).

Remark 1.2. The bounded type α is dense in \mathbb{R} .

Part (2) of Theorem 1.1 can be obtained in the same way as in [47] (See page 2367, proof of Theorem 2 in [47]) from the following examples in $SL(2, \mathbb{R})$ -cocycles.

Theorem 1.2. *Consider quasiperiodic $SL(2, \mathbb{R})$ -cocycles over \mathbb{S}^1 with α being a fixed irrational number of bounded-type. For any $s > 2$, there exists a cocycle $D_s \in G^s(\mathbb{S}^1, SL(2, \mathbb{R}))$ such that the Lyapunov exponent is discontinuous at D_s in G^s -topology.*

²[4] proved the measure version and [33] proved the arithmetic version, actually [33] proved Anderson localization holds for Diophantine phases.

³[4] proved the case $\beta > 0$, [30] proved the case α irrational and $|\lambda| = 1$.

1.2. A Brief review on the continuity of the Lyapunov exponent. As we mentioned above, both the base dynamics T and the smoothness of the matrix A affect the regularity of the Lyapunov exponent. This has been the object of considerable recent interests, see Viana [45], Wilkinson [51] and the references therein.

If the base dynamics has some hyperbolicity, then the Lyapunov exponent is continuous. For example, Furstenberg-Kifer [23] and Hennion [28] proved the continuity of the largest LE of i.i.d random matrices under a condition of almost irreducibility. Bocker and Viana [10] proved continuity of Lyapunov exponents with respect to the cocycle and the invariant probability for random products of $SL(2, \mathbb{R})$ matrices in the Bernoulli setting. In higher dimensions, continuous dependence with respect to A of all Lyapunov exponents for i.i.d. random products of matrices in $GL(d, \mathbb{R})$ was proved by Avila et al. [3]. If the base dynamics is a subshift of finite type or, more generally, a hyperbolic set, then Backes-Brown-Butler [5] proved that the Lyapunov exponents are always continuous among Hölder continuous fiber-bunched $SL(2, \mathbb{R})$ -cocycles.

If $A \in C^r(X, SL(2, \mathbb{R}))$, it is known that $L(A)$ is upper semicontinuous; thus, it is continuous at generic A . Especially, it is continuous at A with $L(A) = 0$ and at uniformly hyperbolic cocycles. The most interesting issue is the continuity of $L(A)$ at the nonuniformly hyperbolic cocycles, which is found to depend on the class of cocycles under consideration including its topology. LE was proved to be discontinuous at any nonuniformly hyperbolic cocycles in C^0 -topology by Furman [22] (Continuity at uniformly hyperbolic cocycles is well-known). Motivated by Mane [40, 41], Bochi [9] further proved that any nonuniformly hyperbolic $SL(2, \mathbb{R})$ -cocycle over a fixed ergodic system on a compact space, can be arbitrarily approximated by cocycles with zero LE in the C^0 -topology.

In this paper, we are interested in the quasiperiodic cocycles. The base system is a rotation on the torus in this case, things are very complicated: it will depend on the smoothness of A in a very sensitive way. If the cocycle is analytic, the Hölder continuity of the Lyapunov exponent in the positive Lyapunov exponent regime was proved by Goldstein and Schlag [25] assuming that α is strong Diophantine. Similar results were proved in [14] by Bourgain, Goldstein, and Schlag when the underlying dynamics is a shift or skew-shift of a higher-dimensional torus. For more results of this favor, here is a partial list [12, 19–21, 24–27, 39, 42, 44, 49, 52, 53]. Later, it was proved by Bourgain-Jitomirskaya in [15] that the LE is joint continuous for $SL(2, \mathbb{R})$ cocycles, in frequency and cocycle map, at any irrational frequencies. Jitomirskaya-Koslover-Schulteis [31] got the continuity of LE with respect to potentials for a class of analytic quasiperiodic $M(2, \mathbb{C})$ -cocycles. Bourgain [13] extended the results in [15] to multi-frequency case. Jitomirskaya-Marx [35] extended the results in [15] to all (including singular) $M(2, \mathbb{C})$ -cocycles. More recently, continuity of the Lyapunov exponents for one-frequency analytic $M(m, \mathbb{C})$ cocycles was given by Avila-Jitomirskaya-Sadel [2]. Weak Hölder continuity of the Lyapunov exponents for multi-frequency $GL(m, \mathbb{C})$ -cocycles, $m \geq 2$, was recently obtained by Schlag [44] and Duarte-Klein [20]. For the lower regularity case, Klein [37] proved that for Schrödinger operators with potentials in a Gevrey class G^s with $1 \leq s < 2$, the LE is weak Hölder continuous on any compact interval of the energy provided that the frequency is strong Diophantine and the LE is large than 0. While if we further lower the regularity of the potential, Wang-You [47] constructed examples to show that the LE of quasiperiodic Schrödinger cocycles can be discontinuous with respect to the potential even in the C^∞ -topology. Jitomirskaya-Marx [35] obtained similar results in the complex category $M(2, \mathbb{C})$ by the tools of harmonic analysis. Recently, Wang-You [48] improved the result in [47] by showing that in C^r -topology, $1 \leq r \leq +\infty$, there exists Schrödinger cocycles with a positive LE that can be approximated by ones with zero LE. For other results about results on discontinuity of LE, one can see [18, 46].

1.3. Outline of the proof and the structure of this paper. The main results of this paper are based on several improvements of the results in [47] where the authors constructed examples of discontinuity of LE in C^∞ -topology. We first give a quick review of the main ideas. We construct $D_s \in G^s(\mathbb{S}^1, SL(2, \mathbb{R}))$ ($s > 2$) as the limit of a sequence of cocycles $\{A_n, n = N, N + 1, \dots\}$

in $G^s(\mathbb{S}^1, SL(2, \mathbb{R}))$ ($s > 2$), the sequence $\{A_n, n = N, N + 1, \dots\}$ possesses some kind of finite hyperbolic property, that is, $\|A_n^{r_n^+}(x)\| \approx \lambda^{r_n^+}$ for most $x \in \mathbb{S}^1$ and $\lambda \gg 1$ with $r_n^+ \rightarrow \infty$ as $n \rightarrow \infty$, which gives a lower bound estimate $(1 - \varepsilon) \ln \lambda$ of the Lyapunov exponent of the limit cocycle (α, D_s) . Then we modify $\{A_n, n = N, N + 1, \dots\}$, and construct another sequence of cocycles $\{\tilde{A}_n, n = N, N + 1, \dots\}$ with some kind of degenerate property such that $\tilde{A}_n \rightarrow D_s$ in G^s -topology as $n \rightarrow \infty$. Moreover, for each n , the Lyapunov exponent of (α, \tilde{A}_n) is less than $(1 - \delta) \ln \lambda$ with $\delta \gg \varepsilon$, which implies the discontinuity of the Lyapunov exponent at D_s .

Compared to [47], the main technical improvements of the present paper are the following two aspects:

- (1) Since we need to construct examples in Gevrey space, we need explicit examples of Gevrey functions. We find the C^∞ -bump functions are all Gevrey functions based on an optimal estimate of the upper bound of its derivatives. Surprisingly, this easy but important observation makes it possible for us to construct a counterexample in Gevrey space.
- (2) Another technical difficulty (the most difficult part) is to prove the sequences $\{A_n\}_{n=N}^\infty$ and $\{\tilde{A}_n\}_{n=N}^\infty$ converge in G^s -topology ($s > 2$). It is much more difficult than to prove the convergence in C^∞ -topology since one needs very delicate control of the derivatives, and it is out of reach by the methods in [47]. We overcome this difficulty by developing a G^s version of the concatenation of finitely many hyperbolic matrices, i.e., Lemma 4.1 and Lemma 4.2 in our paper. Our new Lemmas enable us to not only greatly simplify the proofs in [47], but also optimize almost all estimates in [47]. Our construction is optimal since [17] has shown that for the case of $s < 2$, the Lyapunov exponent is continuous.

A key technique in the construction of $A_n(x)$ comes from Young [54], which was derived from Benedicks and Carleson [6]. Based on this technique, Wang-Zhang [49] developed a new iteration scheme to prove LDT for Schrödinger cocycles with a class of finitely differential potential. They proved that for C^2 cos-like (Morse) potential with a large coupling, LE is weak-Hölder continuous. In this aspect, we also give an improvement of the non-resonance lemma proved in [49], which plays an important role in our proof. For more applications of Benedicks-Carleson-Young's method to quasiperiodic Schrödinger operators, we refer readers to [7, 8, 50].

The structure of this paper is as follows. In Section 2, we give some basic concepts and preparations for $SL(2, \mathbb{R})$ matrices and Gevrey functions. The main idea of the proof will be sketched in Section 3. In Section 4, we give the details of the construction, which is the key part of this paper. Finally, We give the proof of some basic properties of Gevrey functions in Section 5.

2. PREPARATIONS AND SOME TECHNICAL LEMMAS

For $\theta \in \mathbb{S}^1$, let

$$R_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \in SO(2, \mathbb{R}).$$

Define the map

$$s : SL(2, \mathbb{R}) \rightarrow \mathbb{RP}^1 = \mathbb{R}/(\pi\mathbb{Z})$$

so that $s(A)$ is the most contraction direction of $A \in SL(2, \mathbb{R})$. That is, for a unit vector $\hat{s}(A) \in s(A)$, it holds that $\|A \cdot \hat{s}(A)\| = \|A\|^{-1}$. Abusing the notation a little, let

$$u : SL(2, \mathbb{R}) \rightarrow \mathbb{RP}^1 = \mathbb{R}/(\pi\mathbb{Z})$$

be determined by $u(A) = s(A^{-1})$ and $\hat{u}(A) \in u(A)$. Then for $A \in SL(2, \mathbb{R})$, it is clear that

$$A = R_u \cdot \begin{pmatrix} \|A\| & 0 \\ 0 & \|A\|^{-1} \end{pmatrix} \cdot R_{\frac{\pi}{2}-s},$$

where $s, u \in [0, \pi)$ are angles corresponding to the directions $s(A), u(A) \in \mathbb{R}/(\pi\mathbb{Z})$.

2.1. Hyperbolic sequences of $SL(2, \mathbb{R})$ -matrices. For a sequence of matrices $\{\cdots A_{-1}, A_0, A_1, \cdots\}$, we denote

$$A^n = A_{n-1} \cdots A_1 A_0$$

and

$$A^{-n} = A_{-n}^{-1} \cdots A_{-1}^{-1}.$$

Definition 2.1. For any $1 < \mu \leq \lambda$, we say the block of matrices $\{A_0, A_1, \cdots, A_{n-1}\}$ is μ -hyperbolic if

- (1) $\|A_i\| \leq \lambda, \quad \forall i,$
- (2) $\|A^i\| \geq \mu^{i(1-\varepsilon)}, \quad \forall i,$

and (1) and (2) hold if A_0, \cdots, A_{n-1} are replaced by $A_{n-1}^{-1}, \cdots, A_0^{-1}$.

The following lemma is due to Young [54] which tells us when the concatenation of two hyperbolic blocks is still a hyperbolic block.

Lemma 2.1 (Lemma 5 of [54]). *Suppose that C satisfies $\|C\| \geq \mu^m$ with $\mu \gg 1$. Assume that $\{A_0, A_1, \cdots, A_{n-1}\}$ is a μ -hyperbolic sequence, and assume that $\angle(s(C^{-1}), s(A^n)) = 2\theta \ll 1$. Then $\|A^n \cdot C\| \geq \mu^{(m+n)(1-\varepsilon)} \cdot \theta$.*

2.2. The Gevrey functions.

2.2.1. Basic properties. In the following, $s \geq 1, K > 0$ will always be some fixed constants. We give some basic properties on the product and composition of Gevrey functions whose proofs will be given in Section 5. Abusing the notations a little bit, for any $f \in G^{s,K}(I)$, we denote

$$|f|_{s,K} = |f|_{G^{s,K}(I)} := \frac{4\pi^2}{3} \sup_k \frac{(1+|k|)^2}{K^k (k!)^s} |\partial^k f|_{C^0(I)}.$$

Proposition 2.1. *Assume $f, g \in G^{s,K}(I)$ and $\varepsilon > 0$ is sufficiently small, we have*

- (1) $|fg|_{s,K} \leq |f|_{s,K} |g|_{s,K}.$
- (2) *For any $\varepsilon > 0$, $|\partial f|_{s, (1+\varepsilon \frac{1}{s})K} \leq \frac{K}{\varepsilon}.$*
- (3) *Assume $|f - 1|_{s,K} \leq \varepsilon$, then*

$$\left| \frac{1}{f} - 1 \right|_{s, (1+\varepsilon \frac{1}{s+8})K} \leq \varepsilon^{\frac{1}{12}}.$$

- (4) *Assume $|f - 1|_{s,K} \leq \varepsilon$, then*

$$\left| \sqrt{f} - 1 \right|_{s, (1+\varepsilon \frac{1}{s+16})K} \leq \varepsilon^{\frac{1}{12}}.$$

- (5) *Assume $|f|_{s,K} \leq \varepsilon$, then $\arcsin(f) \in G^{s,4K}(I)$ and*

$$|\sin f|_{s, (1+\varepsilon \frac{1}{s+8})K}, \quad |\cos f - 1|_{s, (1+\varepsilon \frac{1}{s+8})K} \leq \varepsilon^{\frac{1}{12}}.$$

2.2.2. Explicit examples. Given a C^∞ -bump function, an interesting question is to investigate the decay rate of its Fourier coefficients (equivalently, the growth rate of its derivatives). In this part, we investigate the Gevrey exponent of various C^∞ -bump functions, the proofs will be also postponed to Section 5. We remark that our estimates of the upper bound of the derivatives of C^∞ bump functions are even optimal, we refer readers to [36] for more details.

Lemma 2.2. *Assume that $0 < \nu < \infty$ and*

$$f(x) = \begin{cases} e^{-\frac{1}{|x|^\nu}} & x \neq 0 \\ 0 & x = 0, \end{cases}$$

then there is some $C > 0$ such that

$$|f^{(n)}(x)| \leq \begin{cases} C^n e^{-\frac{1}{2|x|^\nu}} (n!)^{1+\frac{1}{\nu}} & x \neq 0 \\ 0 & x = 0 \end{cases}, \quad \forall n \in \mathbb{N}.$$

As a corollary, $f \in G^{1+\frac{1}{\nu}}(\mathbb{R})$.

Corollary 2.1. Assume that $0 < \nu < \infty$ and

$$f(x) = e^{\frac{1}{|x|^\nu}}, \quad x \neq 0.$$

Then there is some $C > 0$ such that for any $x \neq 0$, we have

$$|f^{(n)}(x)| \leq C^n e^{\frac{2}{|x|^\nu}} (n!)^{1+\frac{1}{\nu}}, \quad \forall n \in \mathbb{N}.$$

Corollary 2.2. We define a 2π -periodic function as follows

$$g(x) = \begin{cases} ce^{-\left(\frac{1}{(x-c_1-k\pi)^\nu} + \frac{1}{(c_1+(k+1)\pi-x)^\nu}\right)} & x \in (c_1 + k\pi, c_1 + (k+1)\pi) \\ 0 & x \in \{c_1 + k\pi, c_1 + (k+1)\pi\} \end{cases},$$

then there is some $C > 0$ such that for any $x \in \mathbb{S}^1$, we have

$$|g^{(n)}(x)| \leq C^n e^{-\left(\frac{1}{2|x-c_1|^\nu} + \frac{1}{2|x-c_1-\pi|^\nu}\right)} (n!)^{1+\frac{1}{\nu}}, \quad \forall n \in \mathbb{N}.$$

As a corollary, $g \in G^{1+\frac{1}{\nu}}(\mathbb{S}^1)$.

$$\text{Let } I_{n,1} = \left[c_1 - \frac{1}{q_n^\beta}, c_1 + \frac{1}{q_n^\beta} \right], I_{n,2} = \left[c_2 - \frac{1}{q_n^\beta}, c_2 + \frac{1}{q_n^\beta} \right] \text{ and } I_n = I_{n,1} \cup I_{n,2}.$$

Lemma 2.3. Assume $0 < \nu < 1$, $\beta > 1$ and $\delta > 0$ satisfy $0 < \frac{\beta}{1-\delta} < 1$. For any $n \geq N$, there exist an absolute constant C and a 2π -periodic function $f_n \in G^{1+\frac{1}{\nu}, C}(\mathbb{S}^1)$ such that

$$f_n(x) \begin{cases} = 1 & x \in \frac{I_n}{10} \\ \in (0, 1] & x \in I_n \setminus \frac{I_n}{10} \\ = 0 & x \in \mathbb{S}^1 \setminus I_n \end{cases}$$

and

$$|f_n|_{1+\frac{1}{\nu}, C} \leq (Cq_n^\beta)^{q_n^{\frac{\nu\beta}{1-\delta}}}.$$

3. PROOF OF THEOREM 1.2

We first introduce some notations. Let p_n/q_n be the continued fraction expansion of α . The general settings of the cocycle (α, A) are

- $q_{n+1} \leq Mq_n$, $n \in \mathbb{N}$,
- $A \in G^{1+\frac{1}{\nu}}(\mathbb{S}^1)$ with $0 < \nu < 1$.

Let $M, N > 0$ be sufficiently large such that

$$\varepsilon = M^{-100} \ll \delta = \frac{1}{4}M^{-20}, \quad \lambda = e^{q_N^\beta} \gg 1, \quad 1 < \beta < \frac{1}{\nu}.$$

Denote by $\gamma = \nu\beta$. For $n \geq N$, we inductively define

$$\begin{aligned} \ln \lambda_{n+1} &= \ln \lambda_n - 10^4 q_{n+1}^{\gamma-1}, & \lambda_N &= \lambda^{1-\varepsilon}. \\ \ln \widetilde{\lambda}_{n+1} &= \ln \widetilde{\lambda}_n + 10^4 q_{n+1}^{\gamma-1}, & \widetilde{\lambda}_N &= \lambda^{1+\varepsilon}. \end{aligned}$$

Choose q_N sufficiently large such that $\sum_{i=N+1}^{\infty} 10^4 q_i^{\gamma-1} < \varepsilon$, then $\lambda_{\infty} \geq \lambda^{1-2\varepsilon}$ and $\widetilde{\lambda}_{\infty} \leq \lambda^{1+2\varepsilon}$.

We define

- The critical set: $C_0 = \{c_1, c_2\}$ where $c_1 \in [0, \pi)$ and $c_2 = c_1 + \pi$.
- The critical interval: $I_{n,1} = \left[c_1 - \frac{1}{q_n^{\beta}}, c_1 + \frac{1}{q_n^{\beta}}\right]$, $I_{n,2} = \left[c_2 - \frac{1}{q_n^{\beta}}, c_2 + \frac{1}{q_n^{\beta}}\right]$ and $I_n = I_{n,1} \cup I_{n,2}$.
- The first return time: For $x \in I_n$, we denote the smallest positive integer i with $T^i x \in I_n$ (respectively $T^{-i} x \in I_n$) by $r_n^+(x)$ (respectively $r_n^-(x)$), and define $r_n^{\pm} = \min_{x \in I_n} r_n^{\pm}(x)$. Obviously, $r_n^{\pm} \geq \frac{q_n}{2}$.
- The sample function: The 2π -periodic smooth function ϕ_0 is defined as

$$\sin \phi_0(x) = ce^{-\left(\frac{1}{(x-c_1-k\pi)^{\nu}} + \frac{1}{(c_1+(k+1)\pi-x)^{\nu}}\right)}, \quad x \in [c_1 + k\pi, c_1 + (k+1)\pi), \quad k \in \mathbb{Z},$$

where c is sufficiently small.

Remark 3.1. To ensure $r_n^{\pm} \geq \frac{q_n}{2}$, we must require $\beta > 1$. To ensure $\sum_{i=N+1}^{\infty} 10^4 q_i^{\gamma-1} < \varepsilon$, we must require $\beta\nu < 1$. Thus our construction is possible only if $\nu < 1$. Indeed, it is essential since if $\nu > 1$, the LE is continuous [17, 37].

Remark 3.2. By (5) in Proposition 2.1 and Corollary 2.2, we have $\phi_0 \in G^{1+\frac{1}{\nu}}(\mathbb{S}^1)$.

For $C \geq 1$, we denote by $\frac{I_{n,i}}{C}$ the sets $\left[c_i - \frac{1}{Cq_n^{\beta}}, c_i + \frac{1}{Cq_n^{\beta}}\right]$, $i = 1, 2$ and by $\frac{I_n}{C}$ the set $\frac{I_{n,1}}{C} \cup \frac{I_{n,2}}{C}$.

Let $\Lambda = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$. Theorem 1.2 follows from the following two propositions whose proofs will be given in Section 4.

Proposition 3.1. *There exist functions $\phi_n(x)$ on \mathbb{S}^1 ($n = N, N+1, \dots$) such that*

$$(1)_n \quad |\phi_n - \phi_{n-1}|_{1+\frac{1}{\nu}, K} \leq \lambda_n^{-\frac{q_n-1}{100}} \text{ for some } K > 0, \text{ if } n > N,$$

$$(2)_n \quad A_n(x), A_n(Tx), \dots, A_n(T^{r_n^+(x)-1}(x)) \text{ is } \lambda_n\text{-hyperbolic for } x \in I_n \text{ where } A_n(x) = \Lambda R_{\frac{\pi}{2} - \phi_n(x)},$$

(3)_n It holds

$$(a)_n \quad s_n(x) - s'_n(x) = \phi_0(x) \quad x \in \frac{I_n}{10},$$

$$(b)_n \quad |s_n(x) - s'_n(x)| \geq \frac{1}{2} |\phi_0(x)| \geq ce^{-10^{\nu} q_n^{\gamma}}, \quad x \in I_n \setminus \frac{I_n}{10},$$

$$\text{where } s_n(x) = s(A_n^{r_n^+}(x)), s'_n(x) = s(A_n^{-r_n^-}(x)),$$

(4)_n It holds

$$\|A_{r_n^{\pm}}\|_{G^{1+\frac{1}{\nu}, K}(I_n)} \leq \widetilde{\lambda}_n^{r_n^{\pm}}, \quad \left| \frac{1}{\|A_{r_n^{\pm}}\|} \right|_{G^{1+\frac{1}{\nu}, K}(I_n)} \leq \lambda_n^{-r_n^{\pm}}.$$

Proposition 3.2. *There exist functions $\tilde{\phi}_n(x)$ on \mathbb{S}^1 ($n = N, N+1, \dots$) such that*

$$(1)_n \quad |\phi_n(x) - \tilde{\phi}_n(x)|_{1+\frac{1}{\nu}, K} \leq Cq_n^{-2} \text{ for some } K > 0, \text{ if } n > N,$$

$$(2)_n \quad \tilde{A}_n(x), \tilde{A}_n(Tx), \dots, \tilde{A}_n(T^{r_n^+(x)-1}(x)) \text{ is } \lambda_n\text{-hyperbolic for each } x \in I_n \text{ where } \tilde{A}_n(x) = \Lambda R_{\frac{\pi}{2} - \tilde{\phi}_n(x)},$$

(3)_n It holds

$$\tilde{s}_n(x) = \tilde{s}'_n(x), \quad x \in \frac{I_n}{10},$$

where $\tilde{s}_n(x) = \overline{s(\tilde{A}_n^{r_n^+}(x))}$, $\tilde{s}'_n(x) = \overline{s(\tilde{A}_n^{-r_n^-}(x))}$.

Proof of Theorem 1.2: By (1)'s in Proposition 3.1 and Proposition 3.2, there exists $D_\nu \in G^{1+\frac{1}{\nu}, K}$ such that $\|A_n - D_\nu\|_{1+\frac{1}{\nu}, K} \rightarrow 0$ and $\|\tilde{A}_n - D_\nu\|_{1+\frac{1}{\nu}, K} \rightarrow 0$. Then Theorem 1.2 is a direct conclusion of the followings:

- (a) $L(D_\nu) \geq (1 - 4\varepsilon) \ln \lambda$,
- (b) $L(\tilde{A}_n) \leq (1 - \delta) \ln \lambda, \quad \forall n > N$.

Step 1: Proof of (a). We say $x \in \mathbb{S}^1$ is nonresonant for $A_n(x)$ if

$$(3.1) \quad \begin{cases} \text{dist}(T^i x, C_0) > \frac{1}{q_N^\beta}, & 0 \leq i < q_N, \\ \text{dist}(T^i x, C_0) > \frac{1}{q_k^\beta}, & q_{k-1} \leq i < q_k, N < k \leq n. \end{cases}$$

The Lebesgue measure of the set of all nonresonant points $x \in \mathbb{S}^1$ is at least $2\pi(1 - \sum_{N \leq k < n} 1/q_k^{\beta-1})$, which is larger than $2\pi(1 - \varepsilon/2C\pi)$ for $N \gg 1$. For any x satisfying the nonresonant property (3.1), let j_0 be the first time such that $T^{j_0}x \in I_N$ and let n_0 be such that $T^{j_0}x \in I_{n_0} \setminus I_{n_0+1}$. In general, let j_i and n_i be defined so that $T^{j_i}x \in I_{n_i} \setminus I_{n_i+1}$ and let $T^{j_{i+1}}x$ be the next return of $T^{j_i}x$ to I_{n_i} . It is obvious that $j_{i+1} - j_i \geq \frac{q_{n_i}}{2}$. By condition (2), we have $\{A_n(T^{j_i}x), \dots, A_n(T^{j_{i+1}-1}x)\}$ is λ_∞ -hyperbolic (See also [54] for similar arguments).

Since $T^{j_i}x \notin I_{n_i+1}$, by (3)_n of Proposition 3.1 and the definition of ϕ_0 , we have

$$\angle(s_n(T^{j_i}x), s'_n(T^{j_i}x)) \geq ce^{-10^\nu q_{n_i+1}^\gamma}.$$

On the other hand, it holds that

$$\angle\left(s(A_n^{-j_i}(T^{j_i}x)), s(A_n^{j_{i+1}-j_i}(T^{j_i}x))\right) > \frac{1}{2} \angle\left(s_n(T^{j_i}x), s'_n(T^{j_i}x)\right) \geq ce^{-10^\nu q_{n_i+1}^\gamma}.$$

By Lemma 2.1, we have

$$\begin{aligned} \|A_n^{j_{i+1}}(x)\| &\geq \|A_n^{j_i}(x)\| \cdot \|A_n^{j_{i+1}-j_i}(T^{j_i}x)\| \cdot \angle\left(s(A_n^{-j_i}(T^{j_i}x)), s(A_n^{j_{i+1}-j_i}(T^{j_i}x))\right) \\ &\geq \|A_n^{j_i}(x)\| \cdot \lambda_\infty^{(j_{i+1}-j_i)(1-\varepsilon)} ce^{-10^\nu q_{n_i+1}^\gamma}. \end{aligned}$$

Inductively

$$\|A_n^{j_s}(x)\| \geq \|A_n^{j_0}(x)\| \cdot \lambda_\infty^{j_s-j_0} \cdot \prod_{i=0}^{s-1} ce^{-10^\nu q_{n_i+1}^\gamma}.$$

Notice that $j_s - j_0 = \sum_{i=1}^s j_i - j_{i-1} \geq \sum_{i=0}^{s-1} \frac{q_{n_i}}{2}$, thus if s is sufficiently large, we have

$$j_0 \leq q_N^C \leq \frac{\varepsilon}{4} j_s, \quad \prod_{i=0}^{s-1} ce^{-10^\nu q_{n_i+1}^\gamma} \geq \lambda^{-\frac{\varepsilon}{4}(j_s-j_0)}.$$

It follows that

$$\|A_n^{j_s}(x)\| \geq \lambda_\infty^{(1-\frac{3\varepsilon}{2})j_s} \geq \lambda_\infty^{(1-2\varepsilon)j_s}.$$

Now we are ready to prove the main result. From the subadditivity of the cocycle, the finite Lyapunov exponent of a cocycle converges to the Lyapunov exponent. Thus there exists a large $s_0 \geq N_0$ such that

$$\left| \frac{1}{j_{s_0}} \int_{\mathbb{T}} \ln \|D_\nu^{j_{s_0}}(x)\| dx - L(\alpha, D_\nu) \right| \leq \varepsilon.$$

By (1)_n of Proposition 3.1, there exists $N_1 > N_0$, such that for any $n > N_1$, it holds that

$$\left| \frac{1}{j_{s_0}} \int_{\mathbb{T}} \ln \|D_\nu^{j_{s_0}}(x)\| dx - \frac{1}{j_{s_0}} \int_{\mathbb{T}} \ln \|A_n^{j_{s_0}}(x)\| dx \right| \leq \varepsilon.$$

On the other hand

$$\frac{1}{j_{s_0}} \int_{\mathbb{T}} \ln \|A_n^{j_{s_0}}(x)\| dx \geq (1 - 2\varepsilon) \ln \lambda_\infty - C \frac{\varepsilon}{C} \geq (1 - 3\varepsilon) \ln \lambda.$$

Thus we finish the proof of (a).

Step 2: Proof of (b). The following two lemmas have been proved in [47].

Lemma 3.1 (Lemma 4.1 of [47]). *Suppose A and B are two hyperbolic matrices such that $\|A\| = \lambda_1^m$ and $\|B\| = \lambda_2^n$ with $m, n > 0$ and $\lambda_1, \lambda_2 \gg 1$. If $A(s(A)) \parallel u(B)$, then $\|BA\| \leq 2 \max\{\lambda_1^m \lambda_2^{-n}, \lambda_2^n \lambda_1^{-m}\}$.*

Lemma 3.2 (Corollary 4.1 of [47]). *Let $\min r_n = \min_{x \in I_n} \min\{i > 0 | T^i x \pmod{2\pi} \in I_n\}$, and let $\max r_n = \max_{x \in \frac{1}{10} I_n} \min\{i > 0 | T^i x \pmod{2\pi} \in \frac{1}{10} I_n\}$, Then $M^{-20} \leq \frac{\min r_n}{\max r_n} \leq 1$.*

Let $\dots < n_{j-1} < n_j < n_{j+1} < \dots$ be the returning times of $x \in \frac{I_n}{10}$ to $\frac{I_n}{10}$. Moreover, we let n_{j+} be the first returning time of $x \in I_n$ to I_n after n_j . Similarly, we denote by n_{j-} the last returning time of $x \in I_n$ to I_n before n_j . Obviously, it holds that $n_{j-1} \leq n_{j-} < n_j$ and $n_j < n_{j+} \leq n_{j+1}$. By Lemma 3.2, we have

$$n_{j+} - n_j, n_j - n_{j-} \leq (1 - \frac{1}{2} M^{-20})(n_{j+} - n_{j-}).$$

Since $T^{n_j} x \in I_n$, by Proposition 3.2 and Lemma 3.1, we have

$$\begin{aligned} \left\| \tilde{A}_n(T^{n_{j+}x}) \dots \tilde{A}_n(T^{n_jx}) \dots \tilde{A}_n(T^{n_{j-}x}) \right\| &\leq 2 \max \left\{ \tilde{A}_n^{n_{j+} - n_j}(T^{n_jx}), \tilde{A}_n^{n_j - n_{j-}}(T^{n_{j-}x}) \right\} \\ &\leq 2 \lambda^{\max\{n_{j+} - n_j, n_j - n_{j-}\}} \leq \lambda^{(1 - \frac{1}{2} M^{-20})(n_{j+} - n_{j-})}. \end{aligned}$$

It follows that

$$\left\| \tilde{A}_n(T^{n_{j+1}x}) \dots \tilde{A}_n(T^{n_{j-1}x}) \right\| \leq \lambda^{n_{j+1} - n_{j-1} - \frac{1}{2} M^{-20}(n_{j+} - n_{j-})} \leq \lambda^{(n_{j+1} - n_{j-1})(1 - \frac{1}{4} M^{-40})}.$$

Thus for any even k ,

$$\left\| \tilde{A}_n(T^{n_k x}) \dots \tilde{A}_n(x) \right\| \leq \lambda^{(1 - \frac{1}{4} M^{-40}) \sum_{j=0}^{\frac{k}{2}-1} (n_{2j+2} - n_{2j})} \leq \lambda^{n_k (1 - \frac{1}{4} M^{-40})},$$

which implies that $L(\tilde{A}_n) \leq (1 - \delta) \ln \lambda$.

4. PROOF OF PROPOSITION 3.1 AND 3.2

In this section, we aim to prove Proposition 3.1 and Proposition 3.2 which are the main technical parts of this paper. The proof is split into the following three subsections.

4.1. Key Lemmas. Let $2 < s < \infty$, $0 < \gamma < 1$, $K > 0$, $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ be bounded, p_n/q_n be the continued fraction expansion of α and $\lambda > 1$ be sufficiently large. Without loss of generality, we may assume $2 < s < 3$ since we are mainly concerned with the case s is sufficiently close to 2. Recall that for any $f \in C^\infty(I)$, we denote

$$|f|_{s,K} := \frac{4\pi^2}{3} \sup_k \frac{(1 + |k|)^2}{K^k (k!)^s} |\partial^k f|_{C^0(I)},$$

the Gevrey norm of f restricting to I .

In the following, we prove a Gevrey version of the concatenation of hyperbolic matrices. It greatly simplifies and improves the proofs in [47].

Lemma 4.1. *Let*

$$E(x) = \begin{pmatrix} e_2(x) & 0 \\ 0 & e_2^{-1}(x) \end{pmatrix} R_{\theta(x)} \begin{pmatrix} e_1(x) & 0 \\ 0 & e_1^{-1}(x) \end{pmatrix},$$

where $e_1, e_2, \theta \in G^{s,K}(I)$ satisfying

$$(4.1) \quad \inf_{x \in I} \left| \theta(x) - \frac{\pi}{2} \right| \geq ce^{-q_n^\gamma} \gg \min \left\{ \inf_{x \in I} e_1(x), \inf_{x \in I} e_2(x) \right\}^{-\frac{1}{100}},$$

$$(4.2) \quad \left| \frac{1}{\cos \theta} \right|_{s,K}, |\tan \theta|_{s,K} \leq Ce^{q_n^\gamma}, \quad |\cos \theta|_{s,K}, |\cot \theta|_{s,K} \leq C,$$

$$(4.3) \quad |e_i^{-1}|_{s,K} \leq C\lambda^{-\frac{1}{3}q_n-1}, \quad i = 1, 2.$$

Then, for $e(x) := \|E(x)\|$, it holds that

$$(4.4) \quad \inf_{x \in I} e(x) \geq c \inf_{x \in I} e_1(x) \cdot \inf_{x \in I} e_2(x) \cdot e^{-q_n^\gamma},$$

$$(4.5) \quad |e|_{s,(1+\eta)K} \leq C^2 |e_1|_{s,K} |e_2|_{s,K},$$

$$(4.6) \quad |e^{-1}|_{s,(1+\eta)K} \leq C^2 |e_1^{-1}|_{s,K} |e_2^{-1}|_{s,K} e^{q_n^\gamma}.$$

Let $s(x) = s(E(x))$ and $u(x) = u(E(x))$, we further have

$$(4.7) \quad \left| \frac{\pi}{2} - s \right|_{s,(1+\eta)K} \leq |e_1^{-1}|_{s,K}^3 |e_1|_{s,K}^2, \quad |u|_{s,(1+\eta)K} \leq |e_2^{-1}|_{s,K}^3 |e_2|_{s,K}^2,$$

where $\eta = \lambda^{-\frac{1}{4000}q_n-1}$.

Proof. For simplicity, let us omit the dependence on x in the following computation. Direct computations show that

$$E^t E = \begin{pmatrix} e_1^2 e_2^2 \cos^2 \theta + e_1^2 e_2^{-2} \sin^2 \theta & (e_2^{-2} - e_2^2) \sin \theta \cos \theta \\ (e_2^{-2} - e_2^2) \sin \theta \cos \theta & e_1^{-2} e_2^{-2} \cos^2 \theta + e_2^2 e_1^{-2} \sin^2 \theta \end{pmatrix}.$$

It is obvious that

$$\begin{aligned} e^2 + e^{-2} &= e_1^2 e_2^2 \cos^2 \theta + e_1^2 e_2^{-2} \sin^2 \theta + e_1^{-2} e_2^{-2} \cos^2 \theta + e_2^2 e_1^{-2} \sin^2 \theta \\ &= e_1^2 e_2^2 \cos^2 \theta (1 + e_2^{-4} \tan^2 \theta + e_1^{-4} e_2^{-4} + e_1^{-4} \tan^2 \theta) := b. \end{aligned}$$

Thus

$$(4.8) \quad e = \sqrt{\frac{b + \sqrt{b^2 - 4}}{2}} = \sqrt{b} \sqrt{\frac{1 + \sqrt{1 - 4b^{-2}}}{2}},$$

$$(4.9) \quad e^{-1} = \sqrt{\frac{2}{b + \sqrt{b^2 - 4}}} = \sqrt{\frac{1}{b}} \sqrt{\frac{2}{1 + \sqrt{1 - 4b^{-2}}}}.$$

By (4.1) and (4.8), we have

$$(4.10) \quad \inf_{x \in I} e(x) \geq c \inf_{x \in I} e_1(x) \cdot \inf_{x \in I} e_2(x) \cdot e^{-q_n^\gamma}.$$

By (1) in Proposition 2.1, (4.3) and (4.2), we have

$$(4.11) \quad \begin{aligned} &|e_2^{-4} \tan^2 \theta + e_1^{-4} e_2^{-4} + e_1^{-4} \tan^2 \theta|_{s,K} \\ &\leq |e_2^{-1}|_{s,K}^4 |\tan \theta|_{s,K}^2 + |e_1^{-1}|_{s,K}^4 |e_2^{-1}|_{s,K}^4 + |e_1^{-1}|_{s,K}^4 |\tan \theta|_{s,K}^2 \\ &\leq C\lambda^{-\frac{4}{3}q_n-1} e^{2q_n^\gamma} + C\lambda^{-\frac{8}{3}q_n-1} + C\lambda^{-\frac{4}{3}q_n-1} e^{2q_n^\gamma} \leq \lambda^{-q_n-1}. \end{aligned}$$

The last inequality holds since α is bounded and λ is sufficiently large.

Let $\eta_1 = \lambda^{-\frac{1}{20}q_{n-1}}$ and $\eta_2 = \lambda^{-\frac{1}{240}q_{n-1}}$, by (4.11) and (3)-(4) in Proposition 2.1, we have

$$(4.12) \quad \left| \frac{1}{1 + e_2^{-4} \tan^2 \theta + e_1^{-4} e_2^{-4} + e_1^{-4} \tan^2 \theta} - 1 \right|_{s, (1+\eta_1)K} \leq \lambda^{-\frac{1}{12}q_{n-1}}.$$

$$(4.13) \quad \left| \sqrt{1 + e_2^{-4} \tan^2 \theta + e_1^{-4} e_2^{-4} + e_1^{-4} \tan^2 \theta} - 1 \right|_{s, (1+\eta_1)K} \leq \lambda^{-\frac{1}{12}q_{n-1}}.$$

$$(4.14) \quad \left| \sqrt{\frac{1}{1 + e_2^{-4} \tan^2 \theta + e_1^{-4} e_2^{-4} + e_1^{-4} \tan^2 \theta}} - 1 \right|_{s, (1+\eta_2)(1+\eta_1)K} \leq \lambda^{-\frac{1}{144}q_{n-1}}.$$

By (1) in Proposition 2.1, (4.3), (4.2), (4.12), (4.13) and (4.14), we have

$$(4.15) \quad \begin{aligned} \left| \frac{1}{b} \right|_{s, (1+\eta_1)K} &\leq |e_1^{-1}|_{s,K}^2 |e_2^{-1}|_{s,K}^2 \left| \frac{1}{\cos^2 \theta} \right|_{s,K} \left| \frac{1}{1 + e_2^{-4} \tan^2 \theta + e_1^{-4} e_2^{-4} + e_1^{-4} \tan^2 \theta} \right|_{s, (1+\eta_1)K} \\ &\leq C^2 e^{2q_n^\gamma} \lambda^{-\frac{4}{3}q_{n-1}} (1 + \lambda^{-\frac{1}{12}q_{n-1}}) \leq \lambda^{-q_{n-1}}. \end{aligned}$$

$$(4.16) \quad \begin{aligned} \left| \sqrt{b} \right|_{s, (1+\eta_1)K} &\leq |e_1|_{s,K} |e_2|_{s,K} |\cos \theta|_{s,K} \left| \sqrt{1 + e_2^{-4} \tan^2 \theta + e_1^{-4} e_2^{-4} + e_1^{-4} \tan^2 \theta} \right|_{s, (1+\eta_1)K} \\ &\leq 2C |e_1|_{s,K} |e_2|_{s,K}. \end{aligned}$$

$$(4.17) \quad \begin{aligned} \left| \sqrt{\frac{1}{b}} \right|_{s, (1+\eta_2)(1+\eta_1)K} &\leq \left| \frac{1}{e_1} \right|_{s,K} \left| \frac{1}{e_2} \right|_{s,K} \left| \frac{1}{\cos \theta} \right|_{s,K} \left| \sqrt{\frac{1}{1 + e_2^{-4} \tan^2 \theta + e_1^{-4} e_2^{-4} + e_1^{-4} \tan^2 \theta}} \right|_{s, (1+\eta_2)(1+\eta_1)K} \\ &\leq 2C |e_1^{-1}|_{s,K} |e_2^{-1}|_{s,K} e^{q_n^\gamma}. \end{aligned}$$

By (4.15), (3) and (4) in Proposition 2.1, we have

$$(4.18) \quad \begin{aligned} \left| \sqrt{1 - 4b^{-2}} - 1 \right|_{s, (1+\eta_1)^2 K} &\leq \lambda^{-\frac{1}{12}q_{n-1}}, \\ \left| \sqrt{\frac{1 + \sqrt{1 - 4b^{-2}}}{2}} - 1 \right|_{s, (1+\eta_2)(1+\eta_1)^2 K} &\leq \lambda^{-\frac{1}{12^2}q_{n-1}}, \end{aligned}$$

$$(4.19) \quad \left| \sqrt{\frac{2}{1 + \sqrt{1 - 4b^{-2}}}} - 1 \right|_{s, (1+\eta_3)(1+\eta_2)(1+\eta_1)^2 K} \leq \lambda^{-\frac{1}{12^3}q_{n-1}},$$

where $\eta_2 = \lambda^{-\frac{1}{2880}q_{n-1}}$.

By (4.16), (4.17), (4.18) and (4.19), we have

$$\begin{aligned} |e|_{s, (1+\eta)K} &= \left| \sqrt{b} \right|_{s, (1+\eta)K} \left| \sqrt{\frac{1 + \sqrt{1 - 4b^{-2}}}{2}} \right|_{s, (1+\eta)K} \leq C^2 |e_1|_{s,K} |e_2|_{s,K}. \\ |e^{-1}|_{s, (1+\eta)K} &= \left| \sqrt{\frac{1}{b}} \right|_{s, (1+\eta)K} \left| \sqrt{\frac{2}{1 + \sqrt{1 - 4b^{-2}}}} \right|_{s, (1+\eta)K} \leq C^2 |e_1^{-1}|_{s,K} |e_2^{-1}|_{s,K} e^{q_n^\gamma}. \end{aligned}$$

We finish the proofs of (4.4)-(4.6).

Now we prove (4.7). By polar decomposition procedure, we have $s(x) = \frac{\pi}{2} + \theta(E(x))$ where $s(x)$ is the most contraction direction of $E(x)$ and $\theta(E(x))$ is the eigen-direction of $E^t(x)E(x)$

corresponding to the eigenvalue $\|E(x)\|^2$. Let $a = e_1 e_2$, $c = \frac{e_1}{e_2}$, $u = 2(e_2^2 - e_2^{-2}) \sin \theta \cos \theta$ and $U = (a^2 - a^{-2}) \cos^2 \theta + (c^2 - c^{-2}) \sin^2 \theta$. It's easy to calculate that

$$(4.20) \quad \tan s(x) = \tan\left(\frac{\pi}{2} + \theta(E(x))\right) = \frac{u}{\sqrt{U^2 + u^2} - U} = \frac{\sqrt{U^2 + u^2} + U}{u}.$$

Since $|\theta - \pi/2| > ce^{-q_n^\gamma} \gg \min\left\{\inf_{x \in I} e_1(x), \inf_{x \in I} e_2(x)\right\}^{-\frac{1}{100}}$, we have

$$(4.21) \quad U \geq ce_1^2 e_2^2 e^{-2q_n^\gamma} - e_1^2 e_2^{-2} - e_1^{-2} e_2^2 > 0.$$

(4.20) and (4.21) imply that

$$(4.22) \quad s(x) = \arctan\left(\operatorname{sgn}(u) \left(\sqrt{\frac{U^2}{u^2} + 1} + \frac{U}{|u|}\right)\right).$$

A direct calculation shows that

$$(4.23) \quad g := \frac{U}{u} = \frac{(e_1 e_2)^2 - (e_1 e_2)^{-2}}{2(e_2^2 - e_2^{-2})} \cot \theta + \frac{(e_1 e_2^{-1})^2 - (e_1 e_2^{-1})^{-2}}{2(e_2^2 - e_2^{-2})} \tan \theta.$$

Without loss of generality, we only consider the case $g(x) > 0$. A direct computation shows

$$\begin{aligned} \frac{ds}{dx} &= \frac{1}{2} \frac{1}{1+g^2} \frac{dg}{dx}, \\ g &= \frac{e_1^2 \cot \theta}{2} \frac{1 - e_1^{-2} e_2^{-4} + (e_2^{-4} - e_1^{-4}) \tan^2 \theta}{1 - e_2^{-4}} := \frac{e_1^2 \cot \theta}{2} \cdot h, \\ \frac{1}{1+g^2} &= 4e_1^{-4} \tan^2 \theta \frac{1}{4e_1^{-4} \tan^2 \theta + h^2}. \end{aligned}$$

By (4.3), (4.2), (1) and (3) in Proposition 2.1, we have

$$\begin{aligned} |h - 1|_{s, (1+\eta_1)K} &= \left| \frac{1 - e_1^{-2} e_2^{-4} + (e_2^{-4} - e_1^{-4}) \tan^2 \theta}{1 - e_2^{-4}} - 1 \right|_{s, (1+\eta_1)K} \leq \lambda^{-q_n-1}, \\ |h^2 - 1|_{s, (1+\eta_1)K} &\leq |h - 1|_{s, (1+\eta_1)K} |h + 1|_{s, (1+\eta_1)K} \leq 4\lambda^{-q_n-1}, \\ \left| \frac{1}{4e_1^{-4} \tan^2 \theta + h^2} - 1 \right|_{s, (1+\eta_1)^2 K} &\leq 4\lambda^{-\frac{q_n-1}{12}}. \end{aligned}$$

It follows that

$$\begin{aligned} |g|_{s, (1+\eta_1)K} &\leq C |e_1|_{s, K}^2 |\cot \theta|_{s, K} \left| \frac{1 - e_1^{-2} e_2^{-4} + (e_2^{-4} - e_1^{-4}) \tan^2 \theta}{1 - e_2^{-4}} \right|_{s, (1+\eta_1)K} \leq C |e_1|_{s, K}^2, \\ \left| \frac{1}{1+g^2} \right|_{s, (1+\eta_1)^2 K} &\leq 2 |\tan \theta|_{s, K}^2 |e_1^{-1}|_{s, K}^4 \leq C |e_1^{-1}|_{s, K}^4 e^{2q_n^\gamma}. \end{aligned}$$

By (2) in Proposition 2.1 and (4.3), we have

$$\left| \frac{\pi}{2} - s \right|_{s, (1+\eta)K} \leq \left| \frac{1}{1+g^2} \right|_{s, (1+\eta_1)^2 K} |\partial g|_{s, (1+\eta)K} \leq C^2 |e_1^{-1}|_{s, K}^4 e^{2q_n^\gamma} |e_1|_{s, K}^2 K \lambda^{\frac{1}{20}q_n-1} \leq |e_1^{-1}|_{s, K}^3 |e_1|_{s, K}^2.$$

Similar results hold for u . Thus we finish the whole proof. \square

Consider a sequence of maps

$$E^\ell \in G^{s,K}(I, SL(2, \mathbb{R})), \quad 0 \leq \ell \leq n-1.$$

Let $s^\ell(x) = s[E^\ell(x)]$, $u^\ell(x) = u[E^\ell(x)]$, $e^\ell(x) = \|E^\ell(x)\|$ and $\Lambda^\ell(x) = \begin{pmatrix} e^\ell(x) & 0 \\ 0 & (e^\ell(x))^{-1} \end{pmatrix}$. By polar decomposition, it holds that

$$E^\ell(x) = R_{u^\ell(x)} \Lambda^\ell(x) R_{\frac{\pi}{2} - s^\ell(x)}.$$

Set for each $0 \leq \ell \leq n-1$,

$$E_k^\ell(x) = \begin{cases} E^{k-1+\ell}(x) \cdots E^\ell(x), & 1 \leq k \leq n-\ell \\ I_2, & k=0 \\ \left(E_{-k}^{\ell+k}(x)\right)^{-1}, & -\ell \leq k \leq -1 \end{cases}.$$

For $k \geq 1$, let $s_k^\ell(x) = s[E_k^\ell(x)]$, $u_k^\ell(x) = s[E_{-k}^\ell(x)]$, $e_k^\ell(x) = \|E_k^\ell(x)\|$ and $\Lambda_k^\ell(x) = \begin{pmatrix} e_k^\ell(x) & 0 \\ 0 & (e_k^\ell(x))^{-1} \end{pmatrix}$. Again by polar decomposition, it holds that

$$E_k^\ell(x) = R_{u_k^{\ell+k}(x)} \Lambda_k^\ell(x) R_{\frac{\pi}{2} - s_k^\ell(x)}.$$

Lemma 4.2. *Let $0 \leq \ell < n-1 \leq q_m^C$, $1 < \xi < \frac{1}{1000}$ and $\theta_\ell(x) =: u^{\ell-1}(x) - s^\ell(x) + \frac{\pi}{2}$. Assume that*

$$(4.24) \quad \inf_{x \in I} \left| \frac{\pi}{2} - \theta_\ell(x) \right| = \inf_{x \in I} \left| s^\ell(x) - u^{\ell-1}(x) \right| \geq c e^{-q_m^\gamma} \gg \min_{0 \leq \ell \leq n-1} \left\{ \inf_{x \in I} e^\ell(x) \right\}^{-\frac{1}{100}},$$

$$(4.25) \quad \left| \frac{1}{\cos \theta_\ell} \right|_{s,K}, |\tan \theta_\ell|_{s,K} \leq C e^{q_m^\gamma}, \quad |\cos \theta_\ell|_{s,K}, |\cot \theta_\ell|_{s,K} \leq C,$$

$$(4.26) \quad |(e^\ell)^{-1}|_{s,K} \leq \left(|e^\ell|_{s,K} \right)^{-1+\xi} \leq C \lambda^{-\frac{1}{3}q_m-1}.$$

Then it holds that

$$(4.27) \quad \inf_{x \in I} e_n^0(x) \geq c^n e^{-2nq_m^\gamma} \prod_{\ell=0}^{n-1} \inf_{x \in I} e^\ell(x),$$

$$(4.28) \quad |e_n^0|_{s,(1+\eta)K} \leq C^{2n} \prod_{\ell=0}^{n-1} |e^\ell|_{s,K},$$

$$(4.29) \quad |(e_n^0)^{-1}|_{s,(1+\eta)K} \leq C^{2n} \prod_{\ell=0}^{n-1} |(e^\ell)^{-1}|_{s,K} e^{4nq_m^\gamma},$$

$$(4.30) \quad |s^0 - s_n^0|_{s,(1+\eta)K} \leq \lambda^{-\frac{1}{10}q_m-1}, \quad |u^{n-1} - u_n^n|_{s,(1+\eta)K} \leq \lambda^{-\frac{1}{10}q_m-1},$$

where $\eta = \lambda^{-\frac{1}{8000}q_m-1}$.

Proof. We prove it by induction. In the case of the product of two matrices, i.e. $n=2$, it follows from Lemma 4.2. Now, we assume for $k \leq n-1$ and for all possible ℓ , we have

$$(4.31) \quad \inf_{x \in I} e_k^\ell(x) \geq c^k e^{-2kq_m^\gamma} \prod_{j=0}^{k-1} \inf_{x \in I} e^{\ell+j}(x),$$

$$(4.32) \quad |e_k^\ell|_{s,(1+\eta)^{2k}K} \leq C^{2k} \prod_{j=0}^{k-1} |e^{\ell+j}|_{s,K},$$

$$(4.33) \quad |(e_k^\ell)^{-1}|_{s,(1+\eta)^{2k}K} \leq C^{2k} \prod_{j=0}^{k-1} |(e^{\ell+j})^{-1}|_{s,K} e^{4kq_m^\gamma},$$

$$(4.34) \quad \left| s_{k-1}^\ell - s_k^\ell \right|_{s,(1+\eta)^{2k}K} \leq \lambda^{-\frac{1}{9}kq_{m-1}},$$

$$(4.35) \quad \left| u_{k-1}^{n-\ell} - u_k^{n-\ell} \right|_{s,(1+\eta)^{2k}K} \leq \lambda^{-\frac{1}{9}kq_{m-1}},$$

where $\eta = \lambda^{-\frac{1}{4000}q_{m-1}}$.

Clearly, (4.34) implies that for $\ell = 0, 1$,

$$\left| s^\ell - s_{n-1}^\ell \right|_{s,(1+\eta)^{2n-2}K} \leq \sum_{j=1}^{n-1} \lambda^{-\frac{1}{9}jq_{m-1}} \leq 2\lambda^{-\frac{1}{9}q_{m-1}},$$

$$\left| u^{n-\ell-1} - u_{n-1}^{n-\ell} \right|_{s,(1+\eta)^{2n-2}K} \leq \sum_{j=1}^{n-1} \lambda^{-\frac{1}{9}jq_{m-1}} \leq 2\lambda^{-\frac{1}{9}q_{m-1}}.$$

Combining the above with (4.24), we have

$$(4.36) \quad \inf_{x \in I} |u_{n-1}^{n-1}(x) - s^{n-1}(x)| \geq ce^{-q_m^\gamma},$$

$$\inf_{x \in I} |s_{n-1}^1(x) - u^0(x)| \geq ce^{-q_m^\gamma}.$$

Let $\tilde{\theta}_{n-1}(x) = u_{n-1}^{n-1}(x) - s^{n-1}(x)$, then

$$\left| \tilde{\theta}_{n-1} - \theta_{n-1} \right|_{G^{s,(1+\eta)^{2n-2}K}(I)} \leq |u_{n-1}^{n-1} - u^n|_{G^{s,(1+\eta)^{2n-2}K}(I)} \leq 2\lambda^{-\frac{1}{9}q_{m-1}}.$$

Note that

$$\frac{1}{\cos \tilde{\theta}_{n-1}} = \frac{1}{\cos \theta_{n-1} \cos(\tilde{\theta}_{n-1} - \theta_{n-1}) - \tan \theta_{n-1} \sin(\tilde{\theta}_{n-1} - \theta_{n-1})}.$$

By (5) in Proposition 2.1 and (4.25), we have

$$\left| \cos(\tilde{\theta}_{n-1} - \theta_{n-1}) - \tan \theta_{n-1} \sin(\tilde{\theta}_{n-1} - \theta_{n-1}) - 1 \right|_{G^{s,(1+\eta)^{2n-1}K}} \leq \lambda^{-\frac{1}{120}q_{m-1}}.$$

By (3) in Proposition 2.1, we have

$$(4.37) \quad \left| \frac{1}{\cos(\tilde{\theta}_{n-1} - \theta_{n-1}) - \tan \theta_{n-1} \sin(\tilde{\theta}_{n-1} - \theta_{n-1})} \right|_{G^{s,(1+\eta)^{2n-1}K}} \leq 2.$$

By (4.25) and (4.37), we have

$$(4.38) \quad \left| \frac{1}{\cos \tilde{\theta}_{n-1}} \right|_{G^{s,(1+\eta)^{2n-1}K}(I)} \leq 2 \left| \frac{1}{\cos \theta_{n-1}} \right|_{G^{s,(1+\eta)^{2n-1}K}(I)} \leq 2Ce^{q_m^\gamma}.$$

Similarly

$$(4.39) \quad \left| \tan \tilde{\theta}_{n-1} \right|_{G^{s,(1+\eta)^{2n-1}K}(I)} \leq 2Ce^{q_m^\gamma}, \quad \left| \cos \tilde{\theta}_{n-1} \right|_{G^{s,(1+\eta)^{2n-1}K}(I)}, \quad \left| \cot \tilde{\theta}_{n-1} \right|_{G^{s,(1+\eta)^{2n-1}K}(I)} \leq 2C.$$

(4.32), (4.33), (4.36), (4.38) and (4.39) imply that we can apply Lemma 4.2 to the product

$$E_n^0(x) = E^{n-1}(x)E_{n-1}^0(x) = E_{n-1}^1(x)E^0(x),$$

which implies that

$$\inf_{x \in I} e_n^0(x) \geq c \inf_{x \in I} e_{n-1}^0(x) \inf_{x \in I} e^{n-1}(x) e^{-q_m^\gamma} \geq c^n e^{-2nq_m^\gamma} \prod_{\ell=0}^{n-1} \inf_{x \in I} e^\ell(x),$$

$$|e_n^0|_{s, (1+\eta)^{2n}K} \leq C^2 |e^{n-1}|_{s, (1+\eta)^{2(n-1)}K} |e_{n-1}^0|_{s, (1+\eta)^{2(n-1)}K} \leq C^{2n} \prod_{\ell=0}^{n-1} |e^\ell|_{s, K},$$

$$|(e_n^0)^{-1}|_{s, (1+\eta)^{2n}K} \leq C^2 e^{2q_m^\gamma} |(e^{n-1})^{-1}|_{s, (1+\eta)^{2(n-1)}K} |(e_{n-1}^0)^{-1}|_{s, (1+\eta)^{2(n-1)}K} \leq C^{2n} e^{4nq_m^\gamma} \prod_{\ell=0}^{n-1} |(e^\ell)^{-1}|_{s, K}.$$

By Lemma 4.2 and (4.26), we have

$$\begin{aligned} |s_{n-1}^0 - s_n^0|_{s, (1+\eta)^{2n}K} &\leq |(e_{n-1}^0)^{-1}|_{s, (1+\eta)^{2n-2}K}^3 |e_{n-1}^0|_{s, (1+\eta)^{2n-2}K}^2 \\ &\leq C^{6n-6} e^{12nq_m^\gamma} \prod_{\ell=0}^{n-2} |(e^\ell)^{-1}|_{s, (1+\eta)^{2n-2}K}^3 C^{4n-4} \prod_{\ell=0}^{n-2} |e^\ell|_{s, (1+\eta)^{2n-2}K}^2 \\ &\leq C^{10n} e^{12nq_m^\gamma} \prod_{j=0}^{n-2} |(e^j)^{-1}|_{s, (1+\eta)^{2n-2}K}^{\frac{1}{2}} \leq \lambda^{-\frac{1}{9}nq_m-1}, \end{aligned}$$

thus

$$|s^0 - s_n^0|_{s, (1+\eta)^{2n}K} \leq |s^0 - s_{n-1}^0|_{s, (1+\eta)^{2n}K} + |s_{n-1}^0 - s_n^0|_{s, (1+\eta)^{2n}K} \leq \lambda^{-\frac{1}{10}nq_m-1}.$$

Similar results hold for u_n^n , we finish the proof since $(1+\eta)^{2n} \leq 1 + \lambda^{-\frac{1}{8000}nq_m-1}$ for $n < q_m^C$. \square

In the following, we will fix $0 < \nu_2 < \nu_1 < 1$, $s_2 = 1 + \frac{1}{\nu_2} > s_1 = 1 + \frac{1}{\nu_1} > 2$, $\beta > 1$ such that $0 < \gamma_2 = \beta\nu_2 < \gamma_1 = \beta\nu_1 < 1$. Let $\delta_1 > 0$ be sufficiently small such that $0 < \frac{\nu_1\beta}{1-\delta_1\nu_1} < 1$. Recall that

- The critical set: $C_0 = \{c_1, c_2\}$ where $c_1 \in [0, \pi)$ and $c_2 = c_1 + \pi$.
- The critical interval: $I_{n,1} = \left[c_1 - \frac{1}{q_n^\beta}, c_1 + \frac{1}{q_n^\beta}\right]$, $I_{n,2} = \left[c_2 - \frac{1}{q_n^\beta}, c_2 + \frac{1}{q_n^\beta}\right]$ and $I_n = I_{n,1} \cup I_{n,2}$.
- The first return time: For $x \in I_n$, we denote the smallest positive integer i with $T^i x \in I_n$ (respectively $T^{-i} x \in I_n$) by $r_n^+(x)$ (respectively $r_n^-(x)$), and define $r_n^\pm = \min_{x \in I_n} r_n^\pm(x)$. Obviously, $r_n^\pm \geq \frac{q_n}{2}$.

Remark 4.1. If α is bounded, we have $r_n^\pm \leq q_n^C$ for some C only depending on α . See [32] for the proof.

4.2. Proof of Proposition 3.1. We prove Proposition 3.1 by induction. Instead of $r_n^\pm(x)$, sometimes, we use r_n^\pm for short when the difference between $r_n^\pm(x)$ and r_n^\pm are negligible. Recall that

$$\begin{aligned} \ln \lambda_{n+1} &= \ln \lambda_n - 10^4 q_{n+1}^{\gamma_1-1}, \quad \gamma_1 = \nu_1\beta, \quad \lambda_N = \lambda^{1-\varepsilon}. \\ \ln \widetilde{\lambda}_{n+1} &= \ln \widetilde{\lambda}_n + 10^4 q_{n+1}^{\gamma_1-1}, \quad \gamma_1 = \nu_1\beta, \quad \widetilde{\lambda}_N = \lambda^{1+\varepsilon}. \end{aligned}$$

We first construct $\phi_N(x)$ and $A_N(x)$ such that (1)_N – (4)_N hold.

Construction of $\phi_N(x)$ and $A_N(x)$: Let $c_1, c_2 \in \mathbb{T}$ with $c_1 \in [0, \pi)$ and $c_2 = c_1 + \pi$. We define a 2π -periodic smooth function ϕ_0 by

$$\sin(\phi_0(x)) = ce^{-\left(\frac{1}{(x-c_1-k\pi)^{\nu_1}} + \frac{1}{(c_1+(k+1)\pi-x)^{\nu_1}}\right)}, \quad x \in [c_1 + k\pi, c_1 + (k+1)\pi),$$

for some $0 < c < \frac{1}{1000}$. In view of Proposition 2.1 and Corollary 2.2, it's easy to see

- (1) ϕ_0 is a $G^{1+\frac{1}{\nu_1}, C}$ - 2π periodic function for some $C > 0$.
- (2) $|\phi_0|_{C^0(\mathbb{S}^1)} \leq \frac{\pi}{6}$ and for $i = 1, 2$, $|\phi_0(x)| \geq ce^{-|x-c_i|^{-\nu_1}}$ for some $c > 0$.

Let $A(x) = \Lambda \cdot R_{\frac{\pi}{2} - \phi_0(x)} = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix} \cdot R_{\frac{\pi}{2} - \phi_0(x)}$, by [54], there exists a large $\lambda_0 > 0$ depending on ϕ_0 , ν_1 and ε such that if $\lambda > \lambda_0$,

$$\{A(x), \dots, A(T^{r_N^+}(x)^{-1}x)\} \text{ is } \lambda_N - \text{hyperbolic}, \forall x \in I_N.$$

Let $\bar{s}_N(x) = \overline{s(A^{r_N^+}(x))}$, $\bar{s}'_N(x) = \overline{s(A^{-r_N^-}(x))}$ for $x \in I_N$. Let $e_N(x)$ be a 2π -periodic C^∞ -function such that $e_N(x) = \phi_0(x) - (\bar{s}'_N(x) - \bar{s}_N(x))$ for $x \in I_N$. Let $\hat{e}_N(x) = e_N(x) \cdot f_N(x)$ where f_N is defined in Lemma 2.3 and $\phi_N(x) = \phi(x) + \hat{e}_N(x)$ for $x \in \mathbb{S}^1$.

Verifying (1)_N and (4)_N of Proposition 3.1: Let $\eta_N = \lambda_N^{-\frac{1}{8000}}$, $n = r_N^+ \leq q_N^C$, $e^\ell(x) = \|A(x + \ell\alpha)\|$ and $I = I_N$,

$$E^\ell(x) = A(x + \ell\alpha) = \Lambda \cdot R_{\frac{\pi}{2} - \phi_0(x + \ell\alpha)}.$$

For $0 \leq \ell < n - 1$, since $x + \ell\alpha \notin I_N$, one can easily verify that

$$\inf_{x \in I} \|A(x + \ell\alpha)\| = \lambda,$$

$$(4.40) \quad \inf_{x \in I} \left| \frac{\pi}{2} - \theta_\ell(x) \right| := \inf_{x \in I} |\phi_0(x + \ell\alpha)| \geq ce^{-q_N^{\beta\nu_1}} = ce^{-q_N^{\gamma_1}},$$

$$(4.41) \quad |(e^\ell)^{-1}|_{G^{s_1, C}(I)} = |e^\ell|_{G^{s_1, C}(I)}^{-1} = \lambda^{-1}.$$

By Corollary 2.1 and Corollary 2.2, there is some $C > 0$ such that

$$(4.42) \quad |\cos \theta_\ell|_{G^{s_1, C}(I)} = |\sin(\phi_0(x + \ell\alpha))|_{G^{s_1, C}(I)}, \quad |\cot \theta_\ell|_{G^{s_1, C}(I)} = |\tan(\phi_0(x + \ell\alpha))|_{G^{s_1, C}(I)} \leq C,$$

$$(4.43) \quad \left| \frac{1}{\cos \theta_\ell} \right|_{G^{s_1, C}(I)} = \left| \frac{1}{\sin(\phi_0(x + \ell\alpha))} \right|_{G^{s_1, C}(I)}, \quad |\tan \theta_\ell|_{G^{s_1, C}(I)} = |\cot(\phi_0(x + \ell\alpha))|_{G^{s_1, C}(I)} \leq Ce^{q_N^{\gamma_1}}.$$

Set $q_{N-1} = 1$, (4.40)-(4.43) imply that all the assumptions in Lemma 4.2 are satisfied. It follows

$$\begin{aligned} \inf_{x \in I_N} \|A_{r_N^+}(x)\| &\geq \lambda^{r_N^+} c^{r_N^+} e^{-r_N^+ q_N^{\gamma_1}} \geq \lambda_N^{r_N^+}, \quad \|A_{r_N^+}\|_{G^{s_1, (1+\eta_N)C}(I_N)} \leq C^{2r_N^+} \lambda^{r_N^+} \leq \widetilde{\lambda}_N^{r_N^+}, \\ \left| \frac{1}{\|A_{r_N^+}\|} \right|_{G^{s_1, (1+\eta_N)C}(I_N)} &\leq C^{2r_N^+} \lambda^{-r_N^+} e^{4r_N^+ q_N^{\gamma_1}} \leq \lambda_N^{-r_N^+}, \quad |e_N|_{G^{s_1, (1+\eta_N)C}(I)} \leq \lambda_N^{-\frac{1}{10}}. \end{aligned}$$

Similar results hold for r_N^- , we omit the proof.

By the definition and Lemma 2.3, we have

$$\begin{aligned} |\phi_N - \phi|_{G^{s_1, (1+\eta_N)C}(\mathbb{S}^1)} &= |\hat{e}_N|_{G^{s_1, (1+\eta_N)C}(\mathbb{S}^1)} \leq |e_N|_{G^{s_1, (1+\eta_N)C}(I_N)} |f_N|_{G^{s_1, (1+\eta_N)C}(\mathbb{S}^1)} \\ &\leq \lambda_N^{-\frac{1}{10}} (Cq_N^\beta)^{q_N^{\frac{\nu_1\beta}{1-\delta_1\nu_1}}} \leq \lambda_N^{-\frac{1}{20}}. \end{aligned}$$

The last inequality holds since $\frac{\nu_1\beta}{1-\delta_1\nu_1} < 1$ and $\lambda \gg e^{q_N^{\gamma_1}}$.

Verifying (2)_N of Proposition 3.1: Let $A_N(x) = \Lambda \cdot R_{\frac{\pi}{2} - \phi_N(x)}$. Obviously, $A_N(x) = A(x) \cdot R_{-\hat{e}_N(x)}$.

Lemma 4.3 ([47]). *For $x \in I_N$, it holds that*

$$A_N^{r_N^+}(x) = A^{r_N^+}(x) \cdot R_{-\hat{e}_N(x)}$$

and

$$A_N^{-r_N^-}(x) = R_{\hat{e}_N(T^{-r_N^-}x)} \cdot A^{-r_N^-}(x).$$

Thus, for any $x \in I_N$, $\{A_N(x), \dots, A_N(T_N^{r_N^+}(x)^{-1}x)\}$ is a λ_N -hyperbolic sequence.

Verifying (3)_N of Proposition 3.1: $(s_N - s'_N)(x) = (\bar{s}_N - \bar{s}'_N)(x) + \hat{e}_N(x)$ which implies that $(s_N - s'_N)(x) = \phi_0(x)$ on $\frac{I_N}{10}$, since $|e_N(x)| \leq \lambda_N^{-\frac{1}{20}}$ in I_N . Thus we have

$$|(s_N - s'_N)(x)| \geq |\phi_0(x)| - \lambda_N^{-\frac{1}{20}} \geq ce^{-10\nu_1 q_N^{\gamma_1}},$$

on $I_N \setminus \frac{I_N}{10}$ since $\lambda > e^{q_N^{\gamma_1}}$.

Inductively, we assume that $\phi_N(x), \dots, \phi_{n-1}(x)$ have been constructed such that Proposition 3.1 holds for $N \leq i \leq n-1$, i.e.,

$$(1)_i \quad |\phi_i(x) - \phi_{i-1}(x)|_{\mathcal{S}^{1, (1+\eta_i)C}} \leq \lambda_i^{-\frac{q_i-1}{100}} \quad \text{where } \eta_i = \prod_{j=N}^{i-1} (1 + \lambda_j^{-\frac{1}{8000}q_j}) - 1.$$

(2)_i For each $x \in I_i$, $A_i(x), A_i(Tx), \dots, A_i(T_N^{r_i^+}(x)^{-1}(x))$ is λ_i -hyperbolic.

(3)_i We have

$$(a)_i \quad s_i(x) - s'_i(x) = \phi_0(x) \quad x \in \frac{I_i}{10};$$

$$(b)_i \quad |s_i(x) - s'_i(x)| \geq \frac{1}{2} |\phi_0(x)| \geq \frac{1}{2} e^{-10\nu_1 q_i^{\nu_1 \beta}}, \quad x \in I_i \setminus \frac{I_i}{10}.$$

(4)_i It holds

$$\|A_{r_i^\pm}\|_{G^{\mathcal{S}^{1, (1+\eta_i)C}(I_n)}} \leq \tilde{\lambda}_i^{r_i^\pm}, \quad \left| \frac{1}{\|A_{r_i^\pm}\|} \right|_{G^{\mathcal{S}^{1, (1+\eta_i)C}(I_n)}} \leq \lambda_i^{-r_i^\pm}.$$

Now we construct $\phi_n(x)$ and verify (1)_n – (4)_n.

Constructing $\phi_n(x)$: From (2)_{n-1}, we have that

$$\left\| A_{n-1}^{r_{n-1}^+}(x) \right\| \cdot e^{-(10q_{n-1}^\beta)\nu_1} \geq \lambda_{n-1}^{q_{n-1}} \cdot e^{-(10q_{n-1}^\beta)\nu_1} \geq \lambda_n^{(1-\epsilon)q_{n-1}}, \quad x \in I_{n-1}.$$

Combing the above with (3)_{n-1}, for each $x \in I_n$, $A_{n-1}(x), A_{n-1}(Tx), \dots, A_{n-1}(T_N^{r_n^+}(x)^{-1}(x))$ is λ_n -hyperbolic. Let $\bar{s}_n(x) = s(A_N^{r_n^+}(x))$, $\bar{s}'_n(x) = s(A_N^{-r_n^-}(x))$. Let $e_n(x)$ be a 2π -periodic C^∞ -function such that $e_n(x) = \phi_0(x) - (\bar{s}'_n(x) - \bar{s}_n(x))$ for $x \in I_n$. Let $\hat{e}_n(x) = e_n(x) \cdot f_n(x)$ and $\phi_n(x) = \phi(x) + \hat{e}_n(x)$ for $x \in \mathbb{S}^1$.

Verifying (1)_n and (4)_n of Proposition 3.1: For any $x \in I_n$, let j_i be defined so that $T^{j_i}x \in I_{n-1} \setminus I_n$ and let $T^{j_{i+1}}x$ be the next return of $T^{j_i}x$ to I_{n-1} . Let $\eta_n = \prod_{j=N}^{n-1} (1 + \lambda_j^{-\frac{1}{8000}q_j}) - 1$, $n_0 \leq q_n^C$, $e^\ell(x) = \|A_{j_{\ell+1}-j_\ell}(x + j_\ell\alpha)\|$ and $I = I_n$.

$$E^\ell(x) = A_{j_{\ell+1}-j_\ell}(x + j_\ell\alpha) = R_{u_{j_{\ell+1}-j_\ell}(x+j_\ell\alpha)} \begin{pmatrix} e_\ell(x) & 0 \\ 0 & (e_\ell(x))^{-1} \end{pmatrix} R_{\frac{\pi}{2}-s_{j_{\ell+1}-j_\ell}(x+j_\ell\alpha)}.$$

By (2)_{n-1}, we have

$$\inf_{x \in I} \|A_{j_{\ell+1}-j_\ell}(x + j_\ell\alpha)\| \geq \lambda_{n-1}^{j_{\ell+1}-j_\ell} \geq \lambda_{n-1}^{\frac{q_{n-1}}{2}}, \quad 0 \leq \ell \leq n_0 - 1.$$

By (3)_{n-1}, for $0 \leq \ell < n_0 - 1$,

$$\begin{aligned} \inf_{x \in I} \left| \frac{\pi}{2} - \theta_\ell(x) \right| &:= |s_{j_{\ell+1}-j_\ell}(x + j_\ell \alpha) - u_{j_\ell-j_{\ell-1}}(x + j_\ell \alpha)| \\ &\geq \frac{1}{2} |s_{n-1}(x + j_\ell \alpha) - s'_{n-1}(x + j_\ell \alpha)| \geq ce^{-q_n^{\beta\nu_1}} = ce^{-q_n^{\gamma_1}}. \end{aligned}$$

By (4)_{n-1}, we have

$$\begin{aligned} |e^\ell|_{G^{s,(1+\eta_{n-1})}C} \left| \frac{1}{e^\ell} \right|_{G^{s,(1+\eta_{n-1})}C} &\leq \left(\frac{\widetilde{\lambda_{n-1}}}{\lambda_{n-1}} \right)^{j_{\ell+1}-j_\ell} \leq \lambda^{4\varepsilon(j_{\ell+1}-j_\ell)} \leq |e_\ell|_{G^{s,(1+\eta_{n-1})}C}^\xi, \\ |e^\ell|_{G^{s,(1+\eta_{n-1})}C}^{-1+\xi} &\leq \lambda_{n-1}^{-\frac{j_{\ell+1}-j_\ell}{2}} \leq \lambda_{n-1}^{-\frac{q_{n-1}}{3}}. \end{aligned}$$

By (1)_{n-1}, we have $|\phi_{n-1} - \phi_0|_{G^{s,(1+\eta_{n-1})}C} \leq 2\lambda^{-\frac{1}{100}}$. By Proposition 2.1 and similar arguments as above, we have

$$\begin{aligned} |\cos \theta_\ell|_{G^{s_1,(1+\eta_{n-1})}C(I)}, \quad |\tan \theta_\ell|_{G^{s_1,(1+\eta_{n-1})}C(I)} &\leq C, \\ \left| \frac{1}{\cos \theta_\ell} \right|_{G^{s_1,(1+\eta_{n-1})}C(I)}, \quad |\cot \theta|_{G^{s_1,(1+\eta_{n-1})}C(I)} &\leq Ce^{q_n^{\gamma_1}}. \end{aligned}$$

Thus all the assumptions in Lemma 4.2 are satisfied, it follows

$$\begin{aligned} \|A_{r_n^+}\|_{G^{s_1,(1+\eta_n)}C} &\leq C^{2n_0} \prod_{\ell=0}^{n_0-1} |e^\ell|_{G^{s_1,(1+\eta_{n-1})}K} \leq C^{2n_0} \widetilde{\lambda_{n-1}}^{\sum_{\ell=0}^{n_0-1} (j_{\ell+1}-j_\ell)} \leq \widetilde{\lambda}_n^{r_n^+}, \\ \left| \frac{1}{\|A_{r_n^+}\|} \right|_{G^{s_1,(1+\eta_i)}C} &\leq C^{2n_0} e^{4n_0 q_n^{\gamma_1}} \prod_{\ell=0}^{n_0-1} |(e^\ell)^{-1}|_{G^{s_1,(1+\eta_{n-1})}K} \leq \lambda_n^{-r_n^+}, \\ |e_n|_{G^{s_1,(1+\eta_n)}C(I_n)} &\leq \lambda_n^{-\frac{1}{20}q_{n-1}}. \end{aligned}$$

Similar results hold for r_n^- .

By the definition, we have

$$\begin{aligned} |\phi_n - \phi_{n-1}|_{G^{s_1,(1+\eta_n)}C(\mathbb{S}^1)} &= |\hat{e}_n|_{G^{s_1,(1+\eta_n)}C(\mathbb{S}^1)} \leq |e_n|_{G^{s_1,(1+\eta_n)}C(I_n)} |f_n|_{G^{s_1,(1+\eta_n)}C(\mathbb{S}^1)} \\ &\leq \lambda_{n-1}^{-\frac{q_{n-1}}{2}} (C_0 q_n^\beta)^{q_n^{\frac{\nu_1 \beta}{1-\delta_1 \nu_1}}} \leq \lambda_n^{-\frac{1}{40}q_{n-1}}. \end{aligned}$$

The last inequality holds because α is bounded.

Verifying (2)_n of Proposition 3.1: Define $A_n(x) = \Lambda \cdot R_{\frac{\pi}{2}-\phi_n(x)}$. Obviously, $A_n(x) = A_{n-1}(x) \cdot R_{-\hat{e}_n(x)}$.

Lemma 4.4 ([47]). *For $x \in I_n$, it holds that*

$$A_n^{r_n^+}(x) = A_{n-1}^{r_n^+}(x) \cdot R_{-\hat{e}_n(x)}$$

and

$$A_n^{-r_n^-}(x) = R_{\hat{e}_n(T^{-r_n^-}x)} \cdot A_{n-1}^{-r_n^-}(x).$$

Thus, for any $x \in I_n$, $\{A_n(x), \dots, A_n(T^{r_n^+}(x)^{-1}x)\}$ is a λ_n -hyperbolic sequence.

Verifying (3)_n of Proposition 3.1: $(s_n - s'_n)(x) = (\bar{s}_n - \bar{s}'_n)(x) + \hat{e}_n(x)$ which implies that $(s_n - s'_n)(x) = \phi_0(x)$ on $\frac{I_n}{10}$, since $|e_n(x)|_{G^{s_1, (1+\eta_n)C}} \leq \lambda_{n-1}^{-q_{n-1}}$ in I_n . Thus we have

$$|(s_n - s'_n)(x)| \geq |\phi_0(x)| - \lambda_{n-1}^{-q_{n-1}} \geq \frac{1}{2} e^{-(10q_n^\beta)^{\nu_1}},$$

on $I_n \setminus \frac{I_n}{10}$.

Thus we finish the proof by letting $K_1 = \lim_{n \rightarrow \infty} (1 + \eta_n)C$.

4.3. Proof of Proposition 3.2. For any $n \geq N$, let $\tilde{e}_n(x) = -(s_n(x) - s'_n(x)) \cdot f_n(x)$ be a 2π -periodic smooth function such that it is $-(s_n(x) - s'_n(x))$ on $\frac{I_n}{10}$ and vanishes outside I_n . From (3)_n in Proposition 3.1, we have that $\tilde{e}_n(x) = -\phi_0(x) \cdot f_n(x)$. Then we define $\tilde{\phi}_n(x) = \phi_n(x) + \tilde{e}_n(x)$ and $\tilde{A}_n(x) = \Lambda \cdot R_{\frac{\pi}{2} - \tilde{\phi}_n(x)}$.

Verifying of (1)_n of Proposition 3.2: It follows from the following Lemma.

Lemma 4.5. $|\tilde{e}_n|_{G^{s_2, C}} \leq e^{-\frac{1}{4}q_n^\gamma}$ for $n > N$.

Proof. By (5.7) in Lemma 2.2, we have $\|\phi_0\|_{G^{s_1, C}(I_n)} \leq e^{-\frac{1}{2}q_n^{\beta\nu_1}}$ for some $C > 0$. On the other hand, we choose δ_2 sufficiently small such that $\frac{\beta}{s_2-1-\delta_2} = \frac{\beta}{\nu_2-\delta_2} < \beta\nu_1$, by Lemma 2.3, we have

$$|f_n|_{s_2, C} \leq (Cq_n^\beta)^{q_n^{\frac{\beta}{s_2-1-\delta_2}}}.$$

Thus

$$|\tilde{e}_n|_{G^{s_2, C}(\mathbb{S}^1)} \leq |\phi_0|_{G^{s_1, C}(I_n)} |f_n|_{G^{s_2, C}(\mathbb{S}^1)} \leq e^{-\frac{1}{2}q_n^{\beta\nu_1}} (Cq_n^\beta)^{q_n^{\frac{\beta}{s_2-1-\delta_2}}} \leq e^{-\frac{1}{4}q_n^{\gamma_1}}.$$

□

Verifying (2)_n of Proposition 3.2: Since for each $x \in I_n$, $\{A_n(x), A_n(Tx), \dots, A_n(T^{r_n^+(x)-1}x)\}$ is λ_n -hyperbolic and $\tilde{\phi}_n(x) = \phi_n(x)$ on $\mathbb{S}^1 \setminus I_n$, we see that $\{\tilde{A}_n(x), \tilde{A}_n(Tx), \dots, \tilde{A}_n(T^{r_n^+(x)-1}x)\}$ is λ_n -hyperbolic. Thus $\tilde{s}_n(x) = s(\tilde{A}_n^{r_n^+}(x))$ and $\tilde{s}'_n(x) = s(\tilde{A}_n^{-r_n^+}(x))$ are well defined.

Verifying (3)_n of Proposition 3.2: Notice that $\tilde{s}_n(x) - \tilde{s}'_n(x) = s_n(x) - s'_n(x) - \tilde{e}_n(x)$. Thus from the definition of $\tilde{e}_n(x)$, it holds that

$$\tilde{s}_n(x) = \tilde{s}'_n(x) \quad x \in \frac{I_n}{10}.$$

Thus we finish the whole proof by choosing $\nu = \frac{1}{s_2-1}$ and $K = \max\{K_1, C\}$.

5. THE PROOFS OF TECHNICAL LEMMAS

5.1. Proof of Proposition 2.1. The following two Lemmas will be used frequently.

Lemma 5.1 (The Formula of Faa di Bruno, see Theorem 1.3.2 in [38]). *Assume f and g are two smooth functions in an open interval (a, b) , let $h = g \circ f$, then*

$$h^{(n)}(x) = \sum_{k_1+2k_2+\dots+nk_n=n} \frac{n!}{k_1!k_2!\dots k_n!} g^{(k)}(f(x)) \left(\frac{f^{(1)}}{1!}\right)^{k_1} \left(\frac{f^{(2)}}{2!}\right)^{k_2} \dots \left(\frac{f^{(n)}}{n!}\right)^{k_n},$$

where $k = k_1 + k_2 + \dots + k_n$.

Lemma 5.2 (Lemma 1.4.1 in [38]).

$$\sum_{k_1+2k_2+\dots+nk_n=n} \frac{k!}{k_1!k_2!\dots k_n!} R^k = R(1+R)^{n-1},$$

where $k = k_1 + k_2 + \dots + k_n$.

By Stirling formula, one has

$$(5.1) \quad \left(\frac{n}{e}\right)^n \leq n! \leq C \left(\frac{n}{e}\right)^n \sqrt{n}.$$

Lemma 5.3. For any $0 < \varepsilon \leq \frac{1}{2}$ and any $\sigma > 0$, we have

$$n^\sigma \leq (2\sigma)^\sigma \varepsilon^{-\sigma} (1+\varepsilon)^n.$$

Proof. Note that $\frac{x^\sigma}{(1+\varepsilon)^x} = e^{-x \ln(1+\varepsilon) + \sigma \ln x}$. Let $f(x) = -x \ln(1+\varepsilon) + \sigma \ln x$, then

$$f'(x) = -\ln(1+\varepsilon) + \frac{\sigma}{x}.$$

It follows that $\max |f(x)| = f\left(\frac{\sigma}{\ln(1+\varepsilon)}\right) = -\sigma + \sigma \ln \frac{\sigma}{\ln(1+\varepsilon)}$. Hence

$$\frac{x^\sigma}{(1+\varepsilon)^x} \leq \left(\frac{\sigma}{\ln(1+\varepsilon)}\right)^\sigma \leq \left(\frac{2\sigma}{\varepsilon}\right)^\sigma.$$

We finish the proof. \square

Proof of Proposition 2.1: The proof of (1) and (2) can be found in [11]. Now we prove (3), note that $\left(\frac{1}{x}\right)^{(n)} = \frac{(-1)^n n!}{x^{n+1}}$. By Lemma 5.1, we have

$$(5.2) \quad \left(\frac{1}{f}\right)^{(n)} = \sum_{k_1+2k_2+\dots+nk_n=n} \frac{n!}{k_1!k_2!\dots k_n!} \frac{(-1)^k k!}{f^{k+1}} \left(\frac{f^{(1)}}{1!}\right)^{k_1} \left(\frac{f^{(2)}}{2!}\right)^{k_2} \dots \left(\frac{f^{(n)}}{n!}\right)^{k_n},$$

where $k = k_1 + k_2 + \dots + k_n$. Recall that

$$|f|_{s,K} := \frac{4\pi^2}{3} \sup_n \frac{(1+|n|)^2}{K^n (n!)^s} |\partial^n f|_{C^0(I)},$$

it follows that

$$(5.3) \quad \inf_{x \in I} |f(x)| \geq \frac{3}{4\pi^2} (1-\varepsilon), \quad \sup_{x \in I} |f^{(n)}(x)| \leq \begin{cases} \frac{3}{4\pi^2} (1+\varepsilon) & n=0 \\ \frac{3\varepsilon}{4\pi^2} \frac{K^n (n!)^s}{(1+n)^2} & n \geq 1. \end{cases}$$

Let $c = \frac{3}{4\pi^2} (1-\varepsilon)$. By (5.2) and (5.3), for $n \geq 1$, we have

$$\begin{aligned} \left| \left(\frac{1}{f}\right)^{(n)} \right| &\leq c^{-1} \left| \sum_{k_1+2k_2+\dots+nk_n=n} \frac{n!}{k_1!k_2!\dots k_n!} \frac{k!}{c^k} \left(\frac{f^{(1)}}{1!}\right)^{k_1} \left(\frac{f^{(2)}}{2!}\right)^{k_2} \dots \left(\frac{f^{(n)}}{n!}\right)^{k_n} \right| \\ &\leq c^{-1} \left| \sum_{k_1+2k_2+\dots+nk_n=n} \frac{n!}{k_1!k_2!\dots k_n!} \frac{k!}{c^k} \varepsilon^k K^n \left(\frac{(2!)^s}{2!}\right)^{k_2} \dots \left(\frac{(n!)^s}{n!}\right)^{k_n} \right|. \end{aligned}$$

By (5.1) and Lemma 5.3, we have

$$n! \leq C \left(\frac{n}{e}\right)^n \sqrt{n} \leq C \left(\frac{n}{e}\right)^n (1+\varepsilon^{\frac{1}{s+1}})^n \varepsilon^{-\frac{1}{2(s+1)}}.$$

Thus

$$\left| \left(\frac{(2!)^s}{2!}\right)^{k_2} \dots \left(\frac{(n!)^s}{n!}\right)^{k_n} \right| \leq C^{(s-1)k} \varepsilon^{-\frac{1}{2}k} \left(\frac{n}{e}\right)^{(s-1)n} (1+\varepsilon^{\frac{1}{s+1}})^{(s-1)n}.$$

It follows that

$$\begin{aligned} \left| \left(\frac{1}{f} \right)^{(n)} \right| &\leq c^{-1} n! \binom{n}{e}^{(s-1)n} (1 + \varepsilon^{\frac{1}{s+1}})^{(s-1)n} K^n \left| \sum_{k_1+2k_2+\dots+nk_n=n} \frac{k!}{k_1!k_2!\dots k_n!} (\varepsilon^{\frac{1}{2}} c^{-1} C^s)^k \right| \\ &\leq \varepsilon^{\frac{1}{2}} c^{-2} C^s (n!)^s \left[K(1 + \varepsilon^{\frac{1}{2}} c^{-1} C^s) (1 + \varepsilon^{\frac{1}{s+1}})^{(s-1)} \right]^n, \end{aligned}$$

where $k = k_1 + k_2 + \dots + k_n$ and the last inequality follows from Lemma 5.2 and (5.1). For sufficiently small ε depending on s , we have

$$(1 + \varepsilon^{\frac{1}{2}} c^{-1} C^s) (1 + \varepsilon^{\frac{1}{s-1}})^{(s-1)} \leq 1 + \frac{1}{4} \varepsilon^{\frac{1}{s}}, \quad \varepsilon^{\frac{1}{2}} c^{-2} C^s \leq \varepsilon^{\frac{1}{3}}.$$

Hence by Lemma 5.3 again,

$$\frac{4\pi^2}{3} \sup_n \frac{(1 + |n|)^2}{\left(K(1 + \varepsilon^{\frac{1}{s+8}}) \right)^n (n!)^s} \left| \partial^n \frac{1}{f} \right|_{C^0(I)} \leq \varepsilon^{\frac{1}{3}} |n|^2 \left(1 + \frac{1}{2} \varepsilon^{\frac{1}{s+8}} \right)^{-n} \leq \varepsilon^{\frac{1}{12}}.$$

By the definition, we have

$$\left| \frac{1}{f} - 1 \right|_{s, (1 + \varepsilon^{\frac{1}{s+8}})K} \leq \varepsilon^{\frac{1}{12}}.$$

For (4), note that $|(\sqrt{x})^n| = |\frac{1}{2} \cdots (\frac{1}{2} - n + 1) x^{\frac{1}{2}-n}| \leq (n+2)! \sqrt{|x|} |x|^{-n}$. Similar to the proof of (2), we have

$$\left| (\sqrt{f})^{(n)} \right| \leq \varepsilon^{\frac{1}{2}} c^{-1} C^s (n+2)^2 (n!)^s \left[K(1 + \varepsilon^{\frac{1}{2}} c^{-1} C^s) (1 + \varepsilon^{\frac{1}{s-1}})^{(s+1)} \right]^n.$$

By Lemma 5.3, we have

$$\frac{4\pi^2}{3} \sup_n \frac{(1 + |n|)^2}{\left(K(1 + \varepsilon^{\frac{1}{s+16}}) \right)^n (n!)^s} \left| \partial^n \sqrt{f} \right|_{C^0(I)} \leq \varepsilon^{\frac{1}{3}} |n|^4 \left(1 + \frac{1}{2} \varepsilon^{\frac{1}{s+16}} \right)^{-n} \leq \varepsilon^{\frac{1}{12}}.$$

The proof of (5) is exactly the same as (2) since $|\arcsin^{(n)} x| \leq 2^n n!$, $|\sin^{(n)} x|$, $|\cos^{(n)} x| \leq 1 \leq n!$ for any $n \in \mathbb{N}$. Thus we finish the proof.

5.2. Proof of Lemma 2.2. We inductively prove for $x > 0$,

$$(5.4) \quad f^{(n)}(x) = \sum_{i=1}^n \frac{a_i^n(\nu)}{x^{i\nu+n}} e^{-\frac{1}{x^\nu}},$$

$$(5.5) \quad |a_i^n(\nu)| \leq (2\nu + 2)^{n+i} (\nu + n)^{n-i}, \quad 1 \leq i \leq n.$$

Assume for $k \leq n$, (5.4) and (5.5) hold, then for $k = n + 1$, we have

$$f^{(n+1)}(x) = \sum_{i=1}^n \frac{\nu a_i^n(\nu)}{x^{i\nu+n+\nu+1}} e^{-\frac{1}{x^\nu}} - \sum_{i=1}^n \frac{a_i^n(\nu)(i\nu + n)}{x^{i\nu+n+1}} e^{-\frac{1}{x^\nu}} := \sum_{i=1}^{n+1} \frac{a_i^{n+1}(\nu)}{x^{i\nu+n+1}} e^{-\frac{1}{x^\nu}},$$

where

$$a_i^{n+1} = \begin{cases} -a_1^n(\nu)(\nu + n) & i = 1 \\ a_{i-1}^n(\nu)\nu - a_i^n(\nu)(i\nu + n) & 2 \leq i \leq n \\ a_n^n(\nu)\nu & i = n + 1 \end{cases}.$$

By (5.5), we have

$$|a_i^{n+1}| \leq \begin{cases} |a_i^n(\nu)|(\nu+n) \leq (2\nu+2)^{n+i}(\nu+n)^{n+1-i} \leq (2\nu+2)^{n+1+i}(\nu+n+1)^{n+1-i} & i=1 \\ |a_{i-1}^n(\nu)\nu| + |a_i^n(\nu)(i\nu+n)| \leq (2\nu+2)^{n+i+1}(\nu+n+1)^{n+1-i} & 2 \leq i \leq n. \\ |a_i^n(\nu)\nu| \leq (2\nu+2)^{n+1+i}(\nu+n)^{n-i} \leq (2\nu+2)^{n+1+i}(\nu+n+1)^{n+1-i} & i=n+1 \end{cases}$$

(5.4) and (5.5) imply that

$$(5.6) \quad |f^{(n)}(x)| \leq \sum_{i=1}^n \frac{(2\nu+2)^{n+i}(\nu+n)^{n-i}}{|x|^{i\nu+n}} e^{-\frac{1}{|x|^\nu}}.$$

Notice that

$$\sup_{x \in \mathbb{R}} \frac{1}{|x|^{i\nu+n}} e^{-\frac{1}{2|x|^\nu}} \leq \left(\frac{2(i\nu+n)}{\nu} \right)^{i+n/\nu},$$

it follows that

$$(5.7) \quad \begin{aligned} |f^{(n)}(x)| &\leq e^{-\frac{1}{2|x|^\nu}} \sum_{i=1}^n (2\nu+2)^{n+i}(\nu+n)^{n-i} \left(\frac{2(i\nu+n)}{\nu} \right)^{i+n/\nu} \\ &\leq e^{-\frac{1}{2|x|^\nu}} C^n (\nu+n)^{n(1+\frac{1}{\nu})} \leq e^{-\frac{1}{2|x|^\nu}} C^n (n!)^{1+\frac{1}{\nu}}. \end{aligned}$$

5.3. Proof of Corollary 2.1. By the same argument as in Lemma 2.2, we have for any $x > 0$,

$$(5.8) \quad f^{(n)}(x) = \sum_{i=1}^n \frac{a_i^n(\nu)}{x^{i\nu+n}} e^{\frac{1}{x^\nu}},$$

$$(5.9) \quad |a_i^n(\nu)| \leq (2\nu+2)^{n+i}(\nu+n)^{n-i}, \quad 1 \leq i \leq n.$$

(5.8) and (5.9) imply that

$$(5.10) \quad |f^{(n)}(x)| \leq \sum_{i=1}^n \frac{(2\nu+2)^{n+i}(\nu+n)^{n-i}}{|x|^{i\nu+n}} e^{\frac{1}{|x|^\nu}}.$$

Notice that

$$\sup_{x \in \mathbb{R}} \frac{1}{|x|^{i\nu+n}} e^{-\frac{1}{|x|^\nu}} \leq \left(\frac{i\nu+n}{\nu} \right)^{i+n/\nu},$$

it follows that

$$|f^{(n)}(x)| \leq e^{\frac{2}{|x|^\nu}} \sum_{i=1}^n (2\nu+2)^{n+i}(\nu+n)^{n-i} \left(\frac{i\nu+n}{\nu} \right)^{i+n/\nu} \leq e^{\frac{2}{|x|^\nu}} C^n (n!)^{1+\frac{1}{\nu}}.$$

5.4. Proof of Corollary 2.2. For any $k \in \mathbb{Z}$ and $x \in [c_1 + k\pi, c_1 + (k+1)\pi)$, we have

$$g(x) = cf(x - c_1 - k\pi)f(c_1 + (k+1)\pi - x).$$

Let us firstly show that $g \in C^\infty(\mathbb{S}^1)$, for which we only need to verify the derivative exists for $x = c_1 + k\pi$. By a direct calculation, we have

$$g^{(n)}(x) = \begin{cases} \sum_{k=0}^n \binom{n}{k} f^{(n-k)}(x - c_1 - k\pi) (-1)^k f^{(k)}(c_1 + (k+1)\pi - x) & c_1 + k\pi < x < c_1 + (k+1)\pi \\ \sum_{k=0}^n \binom{n}{k} f^{(n-k)}(x - c_1 - (k-1)\pi) (-1)^k f^{(k)}(c_1 + k\pi - x) & c_1 + (k-1)\pi < x < c_1 + k\pi \end{cases}.$$

We inductively prove that

$$(5.11) \quad g_+^{(n)}(c_1 + k\pi) = g_-^{(n)}(c_1 + k\pi) = 0.$$

Assume (5.11) holds for all $k \leq n$. For $k = n + 1$, by Lemma 2.2, we have

$$\begin{aligned} & \lim_{x \searrow c_1 + k\pi} \left| \frac{g^{(n)}(x) - g^{(n)}(c_1 + k\pi)}{x - c_1 - k\pi} \right| \\ & \leq \lim_{x \searrow c_1 + k\pi} \frac{\sum_{k=0}^n \binom{n}{k} e^{-\frac{1}{2|x-c_1-k\pi|^\nu}} C^{n-k} ((n-k)!)^{1+\frac{1}{\nu}} e^{-\frac{1}{2|c_1+(k+1)\pi-x|^\nu}} C^k ((k)!)^{1+\frac{1}{\nu}}}{x - c_1 - k\pi} = 0. \\ & \lim_{x \nearrow c_1 + k\pi} \left| \frac{g^{(n)}(x) - g^{(n)}(c_1 + k\pi)}{x - c_1 - k\pi} \right| \\ & \leq \lim_{x \nearrow c_1 + k\pi} \frac{\sum_{k=0}^n \binom{n}{k} e^{-\frac{1}{2|x-c_1-(k-1)\pi|^\nu}} C^{n-k} ((n-k)!)^{1+\frac{1}{\nu}} e^{-\frac{1}{2|c_1+k\pi-x|^\nu}} C^k ((k)!)^{1+\frac{1}{\nu}}}{|x - c_1 - k\pi|} = 0. \end{aligned}$$

Thus

$$g_+^{(n+1)}(c_1 + k\pi) = g_-^{(n+1)}(c_1 + k\pi) = 0.$$

By (1) in Proposition 2.1 and Lemma 2.2, we have

$$|g|_{s,C} \leq |f(\cdot - c_1)|_{s,C} |f(c_1 + \pi - \cdot)|_{s,C} \leq ce^{-\frac{1}{2|x-c_1|^\nu}} e^{-\frac{1}{2|x-c_1-\pi|^\nu}}.$$

We thus finish the proof.

5.5. Proof of Lemma 2.3. Define

$$\phi(x) = \begin{cases} e^{-\frac{1}{x^{1/\delta}}} & x > 0 \\ 0 & x \leq 0 \end{cases}.$$

Let

$$w_1(x) = \begin{cases} w_0(-x) & x > 0 \\ w_0(x) & x \leq 0 \end{cases},$$

where $w_0(x) = \frac{\phi(x+2)}{\phi(x+2) + \phi(-x-1)}$. It's easy to verify that

$$w_1(x) = \begin{cases} 0 & x \geq 2 \\ \frac{e^{-\frac{1}{(-x+2)^{1/\delta}}}}{e^{-\frac{1}{(-x+2)^{1/\delta}}} + e^{-\frac{1}{(x-1)^{1/\delta}}} & 1 < x < 2 \\ 1 & -1 \leq x \leq 1 \\ \frac{e^{-\frac{1}{(x+2)^{1/\delta}}}}{e^{-\frac{1}{(x+2)^{1/\delta}}} + e^{-\frac{1}{(-x-1)^{1/\delta}}} & -2 < x < -1 \\ 0 & x \leq -2 \end{cases}.$$

By similar arguments as Corollary 2.2, we have $w_1 \in G^{1+\delta}(\mathbb{R})$.

Then we define f_n to be a π -periodic function such that

$$f_n(x) = w_1(10q_n^\beta(x - c_1)), \quad x \in \left[c_1 - \frac{\pi}{2}, c_1 + \frac{\pi}{2} \right].$$

From the definition, we have that $f_n^{(r)}(x) = (10q_n)^{\beta r} \cdot w_1^{(r)}(y)$ where $y = 10q_n^2(x - c_1)$. By the definition of $G^{1+\delta}$ -norm, there exists $C > 0$ such that

$$\sup_{x \in I_n} |f_n^{(r)}(x)| \leq \frac{(Cq_n)^{\beta r} (r!)^{1+\delta}}{1 + r^2}.$$

Thus

$$\sup_{x \in I_n} \frac{|f_n^{(r)}(x)|(1+r^2)}{C^r(r!)^{1+\frac{1}{\nu}}} \leq q_n^{\beta r} (r!)^{\delta - \frac{1}{\nu}} \leq (Cq_n^\beta) q_n^{\frac{\nu\beta}{1-\delta\nu}}.$$

ACKNOWLEDGEMENT

We would like to thank Svetlana Jitomirskaya for many valuable discussions. L. Ge and X. Zhao were partially supported by NSF DMS-1901462. L. Ge was partially supported by AMS-Simons Travel Grant 2020-2022. Y. Wang was supported by NSF of China (11771205). J. You was partially supported by National Key R&D Program of China (2020YFA0713300), NNSF of China (11871286). X. Zhao was partially supported by China Scholarship Council (No. 201906190072).

REFERENCES

- [1] A. Avila. The absolutely continuous spectrum of the almost Mathieu operator, preprint. arXiv:0810.2965 (2008).
- [2] A. Avila, S. Jitomirskaya and C. Sadel. Complex one-frequency cocycles. *J. Eur. Math. Soc* **16(9)** (2014), 1915-1935.
- [3] A. Avila, M. Viana and A. Eskin. Continuity of Lyapunov exponents of random matrix products. In preparation
- [4] A. Avila, J. You and Q. Zhou. Sharp phase transitions for the almost Mathieu operator. *Duke Math. J.* **166(14)** (2017), 2697-2718.
- [5] L. Backes, A.W. Brown and C. Butler. Continuity of Lyapunov exponents for cocycles with invariant holonomies. Preprint. arXiv:1507.08978v2 (2015).
- [6] M. Benedicks and L. Carleson. The dynamics of the Hénon map. *Ann. of Math.* **133(1)** (1991), 73-169.
- [7] K. Bjerklöv. Positive Lyapunov exponent and minimality for a class of one-dimensional quasi-periodic Schrödinger equations. *Ergodic Theory Dynam. Systems* **25(4)** (2005), 1015-1045.
- [8] K. Bjerklöv. The dynamics of a class of quasi-periodic Schrödinger cocycles. *Ann. Henri Poincaré* **16(4)** (2015), 961-1031.
- [9] J. Bochi. Genericity of zero Lyapunov exponents. *Ergod. Theory Dyn. Syst.* **22(6)** (2002), 1667-1696.
- [10] C. Bocker-Neto and M. Viana. Continuity of Lyapunov exponents for random two-dimensional matrices. *Ergodic Theory Dyn. Syst.* **37** (2017), 1413-1442.
- [11] A. Bounemoura and J. Féjoz. Hamiltonian perturbation theory for ultra-differentiable functions. arXiv:1710.01156 (2017).
- [12] J. Bourgain. Hölder regularity of integrated density of states for the almost Mathieu operator in a perturbative regime. *Lett. Math. Phys.* **51** (2000), 83-118.
- [13] J. Bourgain. Positivity and continuity of the Lyapunov exponent for shifts on \mathbb{T}^d with arbitrary frequency vector and real analytic potential. *J. Anal. Math.* **96** (2005), 313-355.
- [14] J. Bourgain, M. Goldstein and W. Schlag. Anderson localization for Schrödinger operators on \mathbb{Z} with potentials given by skew-shift. *Comm. Math. Phys.* **220** (2001), 583-621.
- [15] J. Bourgain and S. Jitomirskaya. Continuity of the Lyapunov exponent for quasiperiodic operators with analytic potential. *J. Stat. Phys.* **108** (2002) 1203-1218.
- [16] C. Cheng, L. Wang. Destruction of Lagrangian torus for positive definite Hamiltonian systems. *Geom. Funct. Anal.* **23(3)** (2013) 848-866.
- [17] H. Cheng, L. Ge, J. You, Q. Zhou. Global rigidity results for ultra-differentiable quasiperiodic cocycles and its spectral applications. Preprint. arXiv:2101.11150v1 (2021).
- [18] D. Damanik, Z Gan and H. Krüger. Limit-periodic Schrödinger operators with a discontinuous Lyapunov exponent. *J. Funct. Anal.* **279(4)** (2020) 108565, 16 pp.

- [19] P. Duarte and S. Klein. Continuity of the Lyapunov exponents for quasiperiodic cocycles. *Commun. Math. Phys.* **332(3)** (2014), 1113-1166.
- [20] P. Duarte and S. Klein. Continuity, positivity, and simplicity of the Lyapunov exponents for quasi-periodic cocycles. *J. Eur. Math. Soc.* **21(7)** (2019), 2051-2106.
- [21] J. Figueras and T. Timoudas, Sharp $\frac{1}{2}$ -Hölder continuity of the Lyapunov exponent at the bottom of the spectrum for a class of Schrödinger cocycles. *Discrete and continuous dynamical systems* **40(7)** (2020) 4519-4531.
- [22] A. Furman. On the multiplicative ergodic theorem for uniquely ergodic systems. *Ann. Inst. Henri. Poincaré Prob. Stat.* **33** (1997), 797-815.
- [23] H. Furstenberg and Y. Kifer. Random matrix products and measures in projective spaces. *Isr. J. Math.* **46** (1983), 12-32.
- [24] L. Ge, J. You and X. Zhao. Hölder regularity of the integrated density of states for quasi-periodic long-range operators on $\ell^2(\mathbb{Z}^d)$. arXiv:2009.08004.
- [25] M. Goldstein and W. Schlag. Hölder continuity of the integrated density of states for quasi-periodic Schrödinger equations and averages of shifts of subharmonic functions. *Ann of Math.* **154** (2001), 155-203.
- [26] M. Goldstein and W. Schlag. Fine properties of the integrated density of states and a quantitative separation property of the Dirichlet eigenvalues. *Geom. Funct. Anal.* **18** (2008), 755-869.
- [27] R. Han and S. Zhang. Large deviation estimates and Hölder regularity of the Lyapunov exponents for quasi-periodic Schrödinger cocycles. *Int. Math. Res. Not.* **rnz319** (2020).
- [28] H. Hennion. Loi des grands nombres et perturbations pour des produits réductibles de matrices aléatoires indépendantes. *Z. Wahrsch. Verw. Gebiete* **67** (1984), 265-278.
- [29] S. Jitomirskaya. Metal-Insulator Transition for the almost Mathieu operator. *Ann. of Math.* **150** (1999), 1159-1175.
- [30] S. Jitomirskaya. On point spectrum of critical almost Mathieu operators. To appear in *Advances in Mathematics*.
- [31] S. Jitomirskaya, D. Koslover and M. Schulteis. Continuity of the Lyapunov exponent for analytic quasiperiodic cocycles. *Ergod. Th. & Dynam. Sys.* **29(6)** (2009), 1881-1905.
- [32] S. Jitomirskaya and I. Krasovsky. Continuity of the measure of the spectrum for discrete quasiperiodic operators. *Math. Res. Lett.* **9(4)** (2002), 413-421.
- [33] S. Jitomirskaya and W. Liu. Universal hierarchical structure of quasi-periodic eigenfunctions. *Ann. of Math.* **187(3)** (2018), 721-776.
- [34] S. Jitomirskaya and W. Liu. Universal reflective-hierarchical structure of quasiperiodic eigenfunctions and sharp spectral transition in phase. arXiv:1802.00781.
- [35] S. Jitomirskaya and C.A. Marx. Analytic quasi-periodic cocycles with singularities and the Lyapunov exponent of extended Harper's model. *Commun. Math. Phys.* **316** (2012), 237-267.
- [36] S. G. Johnson. Saddle-point integration of C^∞ bump functions, arXiv:1508.04376 (2015).
- [37] S. Klein. Anderson localization for the discrete one-dimensional quasi-periodic Schrödinger operator with potential defined by a Gevrey-class function. *J. Funct. Anal.* **218(2)** (2005), 255-292.
- [38] S.G. Krantz and H. R. Parks. *A Primer of Real Analytic Functions*. Birkhäuser Adv. Texts Basler Lehrbücher, Birkhäuser, Boston, MA, 2002.
- [39] J. Liang, Y. Wang and J. You. Hölder continuity of Lyapunov exponent for a class of C^2 Schrödinger cocycles, arxiv:806.03284 (2018).
- [40] R. Mane. Oseledec's theorem from the generic viewpoint. *Proc. ICM* **1** (1983), 1269-1276. (Warsaw), PWN, Warsaw, (1984).
- [41] R. Mane. The Lyapunov exponents of generic area preserving diffeomorphisms, In *International Conference on Dynamical Systems (Montevideo, 1995)*. *Pitman Res. Notes Math. Longman* **362** (1996), 110-119.

- [42] M. Poletti and M. Viana. Simple Lyapunov spectrum for certain linear cocycles over partially hyperbolic maps. *Nonlinearity* **32**(1) (2018), 238-284.
- [43] J. Pöschel. Integrability of Hamiltonian systems on Cantor sets. *Commun. Pure Appl. Math.* **35**(5) (1982), 653-696.
- [44] W. Schlag. Regularity and convergence rates for the Lyapunov exponents of linear cocycles. *J. Mod. Dyn.* **7**(4) (2013), 619-637.
- [45] M. Viana. Lectures on Lyapunov Exponents. *Cambridge Studies in Advanced Mathematics* **145** (2014) Cambridge University Press, Cambridge.
- [46] M. Viana. (Dis)continuity of Lyapunov exponents. *Ergodic Theory Dynam. Systems* **40**(3) (2020), 577-611.
- [47] Y. Wang and J. You. Examples of discontinuity of Lyapunov exponent in smooth quasiperiodic cocycles. *Duke Math. J.* **162**(13) (2013), 2363-2412.
- [48] Y. Wang and J. You. The set of smooth quasi-periodic Schrödinger cocycles with positive Lyapunov exponent is not open. *Commun. Math. Phys.* **362** (2018), 801-826.
- [49] Y. Wang and Z. Zhang. Uniform positivity and continuity of Lyapunov exponents for a class of C^2 quasiperiodic Schrödinger cocycles. *J. Funct. Anal.* **268**(9) (2015), 2525-2585.
- [50] Y. Wang and Z. Zhang. Cantor spectrum for a class of C^2 quasiperiodic Schrödinger operators. *Int. Math. Res. Not.* **8** (2017), 2300-2336.
- [51] A. Wilkinson. What are Lyapunov exponents, and why are they important? *Bull. Amer. Math. Soc.* **54** (2017), 79-105.
- [52] J. Xu, L. Ge and Y. Wang. The Hölder continuity of Lyapunov exponents for a class of Cos-type quasiperiodic Schrödinger cocycles. arXiv:2006.03381.
- [53] J. You and S. Zhang. Hölder continuity of the Lyapunov exponent for analytic quasiperiodic Schrödinger cocycle with weak Liouville frequency. *Ergod. Th. & Dynam. Sys.* **34**(4) (2014), 1395-1408.
- [54] L.S. Young. Lyapunov exponents for some quasi-periodic cocycles. *Ergod. Th. & Dynam. Sys.* **17**(2) (1997), 483-504.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA IRVINE, CA, 92697-3875, USA
Email address: `lingruig@uci.edu`

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, CHINA
Email address: `yiqianw@nju.edu.cn`

CHERN INSTITUTE OF MATHEMATICS AND LPMC, NANKAI UNIVERSITY, TIANJIN 300071, CHINA
Email address: `jyou@nankai.edu.cn`

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, CHINA AND DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA IRVINE, CA, 92697-3875, USA
Email address: `njuzhaox@126.com`