

A HIGHER-ORDER GENERALIZED SINGULAR VALUE DECOMPOSITION FOR RANK DEFICIENT MATRICES*

IDRIS KEMPF[†], PAUL J. GOULART[†], AND STEPHEN R. DUNCAN[†]

Abstract. The higher-order generalized singular value decomposition (HO-GSVD) is a matrix factorization technique that extends the GSVD to $N \geq 2$ data matrices, and can be used to identify shared subspaces in multiple large-scale datasets with different row dimensions. The standard HO-GSVD factors N matrices $A_i \in \mathbb{R}^{m_i \times n}$ as $A_i = U_i \Sigma_i V^T$, but requires that each of the matrices A_i has full column rank. We propose a reformulation of the HO-GSVD that extends its applicability to rank-deficient data matrices A_i . If the matrix of stacked A_i has full rank, we show that the properties of the original HO-GSVD extend to our reformulation. The HO-GSVD captures shared right singular vectors of the matrices A_i , and we show that our method also identifies directions that are unique to the image of a single matrix. We also extend our results to the higher-order cosine-sine decomposition (HO-CSD), which is closely related to the HO-GSVD. Our extension of the standard HO-GSVD allows its application to datasets with $m_i < n$, such as are encountered in bioinformatics, neuroscience, control theory or classification problems.

Key words. Higher-Order Generalized Singular Value Decomposition, Higher-Order Cosine-Sine Decomposition, Diagonalization, Multimodal Data Fusion.

AMS subject classifications. 65F15, 65F55.

1. Introduction. The *generalized singular value decomposition* (GSVD) [12] is an extension of the well-known *singular value decomposition* (SVD) to $N = 2$ matrices. The GSVD is concerned with the decomposition of two matrices, $A_1 \in \mathbb{R}^{m_1 \times n}$ and $A_2 \in \mathbb{R}^{m_2 \times n}$ with $m_1 \geq n$, and factorizes each of the matrices as $A_i = U_i \Sigma_i V^T$. The matrix of right basis vectors $V \in \mathbb{R}^{n \times n}$, with $\det V \neq 0$, is shared among the decompositions but is not an orthogonal matrix, in contrast to the standard SVD. The columns of the matrices $U_i \in \mathbb{R}^{m_i \times m_i}$ are commonly referred to as *left generalized singular vectors*, each satisfying $U_i^T U_i = I$. The matrices $\Sigma_i = \text{diag}(\sigma_{i,1}, \dots, \sigma_{i,r})$, with $\sigma_{i,k} \geq 0$ and $r = \text{rank}[A_1^T, A_2^T]^T$, contain the *generalized singular values*. In contrast to an SVD, where the singular values represent an amplification in a particular direction, the generalized singular values measure the relative significance of the right basis vectors in the image of each A_i . If $\sigma_{1,k} = \sigma_{2,k}$, then v_k contributes equally to the images of A_1 and A_2 .

The *higher-order GSVD* (HO-GSVD) [11] is an extension of the GSVD to $N \geq 2$ matrices. Given N matrices A_1, \dots, A_N , the HO-GSVD decomposes each A_i as

$$(1.1) \quad A_i = U_i \Sigma_i V^T,$$

where $U_i \in \mathbb{R}^{m_i \times n}$, $\Sigma_i \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{n \times n}$ with $\det V \neq 0$ is shared among all factorizations. The matrix V is obtained from the eigensystem $S_\pi V = V \Sigma$, where $\Sigma := \text{diag}(\varsigma_1, \dots, \varsigma_n)$ and S_π is the arithmetic mean of all pairwise quotients $D_{i,\pi} D_{j,\pi}^{-1}$,

$$(1.2) \quad S_\pi := \frac{1}{N(N-1)} \sum_{i=1}^N \sum_{j=i+1}^N (D_{i,\pi} D_{j,\pi}^{-1} + D_{j,\pi} D_{i,\pi}^{-1}),$$

with $D_{i,\pi}$ defined as

$$(1.3) \quad D_{i,\pi} := A_i^T A_i + \pi A^T A,$$

where $A := [A_1^T, \dots, A_N^T]^T$. The standard HO-GSVD framework [11] corresponds to the case $\pi = 0$. We introduce the term $\pi A^T A$ with $\pi > 0$ and will show that it allows us to accommodate rank-deficient matrices A_i .

As in the case of the GSVD, the columns of the matrices U_i are the left generalized singular vectors and the diagonal matrices Σ_i contain the generalized singular values. It can be shown that the GSVD is a special case of the HO-GSVD with $N = 2$ and that the standard SVD of A_j can be obtained from setting $A_i = I$ for $i \neq j$ and $N \geq 2$ [11]. For the case $\pi = 0$ it was shown in [11] that the subspace associated with the unit eigenvalues of S_π forms a *common HO-GSVD subspace*, and we show that this property is preserved for $\pi > 0$. This subspace

*Submitted February 22, 2021.

Funding: The research leading to these results is supported by Diamond Light Source and the Engineering and Physical Sciences Research Council (EPSRC) with a CASE studentship.

[†]Department of Engineering Science, University of Oxford, Parks Road, Oxford, OX1 3PJ, UK (idris.kempf@eng.ox.ac.uk, paul.goulart@eng.ox.ac.uk, stephen.duncan@eng.ox.ac.uk).

is spanned by the right basis vectors v_k that are of equal significance in all A_i and A_j and therefore associated with the generalized singular values for which $\sigma_{i,k} = \sigma_{j,k}$.

The HO-GSVD is a technique that can be attributed to the field of multimodal data fusion [10], which aims to indentify common features across multiple data sets that describe related phenomena. The HO-GSVD can be compared to other matrix or tensor decompositions, such as the parallel factor analysis (PARAFAC [6] or PARAFAC2 [7]), multilinear SVDs [2] or multilinear principal component analysis [4]. On one hand, some of these decompositions allow constraints to be imposed on the factor matrices U_i , Σ_i and V in (1.1), such as orthogonality or rank conditions, but the matrices A_i are often approximated as $A_i \approx U_i \Sigma_i V^T$ by minimizing $\sum_{i=1}^N \|A_i - U_i \Sigma_i V^T\|_F$. On the other, tensor decompositions require that the matrices A_i share the same dimensions, e.g. a third-order tensor $\mathcal{A} = A_1 \times A_2 \times \dots \times A_N$ requires that all matrices A_i have dimensions $m \times n$, which imposes constraints on the data acquisition. In contrast, the HO-GSVD is an exact matrix factorization so that $A_i = U_i \Sigma_i V^T$ for $i = 1, \dots, N$, and it can accommodate $A_i \in \mathbb{R}^{m_i \times n}$ with different row dimensions m_i , although no constraints are imposed on the factor matrices.

One shortcoming of the original HO-GSVD framework [11] is that the arithmetic mean (1.2) is only well defined for matrices A_i that have full column rank. If $\text{rank } A_i < n$ or $m_i < n$ for some i , then the inverse $(A_i^T A_i)^{-1}$ does not exist and so S_0 in (1.2) is not well defined. In addition, computing (1.2) may be inaccurate when one or more of the A_i have small singular values. In Section 7, we describe several applications in which the problem data generally has $m_i < n$.

To extend the HO-GSVD framework to rank-deficient A_i , we introduce the term $\pi A^T A$ with parameter $\pi > 0$ in (1.3). Provided that the matrix A of stacked A_i has full rank, this has the effect of shifting the eigenvalues of each $D_{i,\pi}$, so that the terms $D_{i,\pi}$ are guaranteed to be invertible and the HO-GSVD can be computed for A_i with arbitrary rank. When all A_i have full column rank, we show that S_π with $\pi > 0$ and S_0 both capture the common subspaces of A_1, \dots, A_N , so that our reformulation is equivalent to the original HO-GSVD [11]. We introduce the notion of an *isolated HO-GSVD subspace* that accounts for the fact that a rank deficient A_i can have a non-empty (right) nullspace. An isolated HO-GSVD subspace captures directions that are in the kernel of all but one A_i .

The GSVD is closely related to the (thin) *cosine-sine decomposition* (CSD) [3, Ch. 2.5.4]. In essence, the CSD states that the SVDs of $Q_1 \in \mathbb{R}^{m_1 \times n}$ and $Q_2 \in \mathbb{R}^{m_2 \times n}$ satisfying $Q_1^T Q_1 + Q_2^T Q_2 = I$ share the same matrix of right singular vectors. The term CSD originates from the fact that the singular values $\mu_{i,k}$ of Q_1 and Q_2 satisfy $\mu_{1,k}^2 + \mu_{2,k}^2 = 1$. The GSVD can be obtained from applying a CSD to the matrices Q_1 and Q_2 that are obtained from the thin QR factorization of the stacked matrices $[A_1^T, A_2^T]^T = QR$, where Q is conformably partitioned such that $A_i = Q_i R$. Analogous to the GSVD and the CSD, the HO-GSVD is closely related to the *higher-order CSD* (HO-CSD) [13]. The HO-GSVD of N matrices A_i can be obtained from the HO-CSD of Q_1, \dots, Q_N that are obtained from the thin QR factorization of the stacked matrices $[A_1^T, \dots, A_N^T]^T$. As in the case of the HO-GSVD, the computation of the HO-CSD proposed in [13] is limited to the case that all Q_i have full rank. In this paper, we also propose to compute the HO-CSD in a different way, which allows for the factorization of rank-deficient Q_i satisfying $Q_1^T Q_1 + \dots + Q_N^T Q_N = I$.

The paper is organized as follows. Section 2 presents the HO-CSD and the HO-GSVD that are applicable to rank-deficient matrices. In Section 3, we extend the notion of common HO-CSD and HO-GSVD subspaces to rank-deficient matrices. In Section 4, the canonical decompositions are derived by applying the results from Section 3 to the decompositions from Section 2. Finally, we relate our findings to existing methods in Section 5 and provide a numerical example in Section 6.

2. Main results. Given N matrices $A_i \in \mathbb{R}^{m_i \times n}$, let A denote the matrix of stacked A_i and $QR = A$ be its thin QR factorization,

$$(2.1) \quad A = \begin{bmatrix} A_1 \\ \vdots \\ A_N \end{bmatrix} = QR = \begin{bmatrix} Q_1 \\ \vdots \\ Q_N \end{bmatrix} R, \quad Q_i \in \mathbb{R}^{m_i \times n}, \quad R \in \mathbb{R}^{n \times n}.$$

The matrices $A_i = Q_i R$ can individually have arbitrary rank, but we assume that $\text{rank } A = n$ so that $\det R \neq 0$. Because $Q^T Q = I$, it holds that $\sum_{i=1}^N Q_i^T Q_i = I$ and $\|Q_i\|_2 \leq 1$. The quotient terms $D_{i,\pi}$ (2.2) of the arithmetic mean S_π (1.2) can be rewritten as

$$(2.2) \quad D_{i,\pi} = A_i^T A_i + \pi A^T A = R^T (Q_i^T Q_i + \pi I) R,$$

with parameter $\pi > 0$. Since $A_i^T A_i \succeq 0$ and $\pi A^T A \succ 0$, the terms $D_{i,\pi}$ are guaranteed to be invertible.

The majority of our developments on the HO-GSVD are based on the HO-CSD. Define the mean of reciprocal inverses, T_π , as

$$(2.3) \quad T_\pi := \frac{1}{N} \sum_{i=1}^N (Q_i^T Q_i + \pi I)^{-1}.$$

The eigensystem of T_π will lead us to the HO-CSD of the matrices Q_i . It can be shown (Appendix A) that S_π and T_π are related by:

$$(2.4) \quad R^{-T} S_\pi R^T = \frac{1}{N-1} ((1 + \pi N) T_\pi - I).$$

THEOREM 2.1. *The matrix T_π is non-defective. There exists $Z \in \mathbb{R}^{n \times n}$ such that $Z^T T_\pi Z = \text{diag}(\tau_1, \dots, \tau_n)$ with $Z^T Z = I$, where the columns of Z are eigenvectors of T_π and the eigenvalues τ_i of T_π satisfy*

$$\tau_i \in \left[\frac{N}{1 + \pi N}, \frac{\pi N + N - 1}{\pi N(1 + \pi)} \right].$$

Proof. From (2.3), the matrix T_π is real and symmetric, hence non-defective with an orthonormal eigenbasis [3, Ch. 8.1]. For the bounds on τ_i , see Appendix B. \square

THEOREM 2.2. *The matrix S_π is non-defective. There exists an invertible $V \in \mathbb{R}^{n \times n}$ such that $V^{-1} S_\pi V = \text{diag}(\varsigma_1, \dots, \varsigma_n)$, where the columns of V are eigenvectors of S_π and the eigenvalues ς_i satisfy*

$$\varsigma_i \in \left[1, 1 + \frac{1}{\pi N(1 + \pi)} \right].$$

Proof. Pre- and post-multiplying (2.4) with Z^T and Z from Thm. 2.1 yields

$$Z^T R^{-T} S_\pi R^T Z = \frac{1}{N-1} ((1 + \pi N) Z^T T_\pi Z - I).$$

Since $Z^T T_\pi Z = \text{diag}(\tau_1, \dots, \tau_n)$, the matrix $Z^T R^{-T} S_\pi R^T Z$ is diagonal. Set $V := R^T Z$, which is invertible because $\det R \neq 0$ and $Z^T Z = I$, then the columns of V are eigenvectors for S_π associated with eigenvalues $\varsigma_i = ((1 + \pi N)\tau_i - 1)/(N - 1)$. The bounds on ς_i are obtained from the bounds on τ_i . \square

The significance of Thm. 2.1 and 2.2 is that the diagonalizable matrices T_π and S_π have eigenvalues that are both bounded away from zero and contained in finite intervals, in contrast to the formulation [11] that requires the full rank condition, which corresponds to $\pi = 0$. More precisely, the range of eigenvalues of S_π is contracted from $[1, \infty)$ for the full-rank case to $[1, 1 + 1/(\pi N(1 + \pi))]$ in our case, which bounds the condition number as $\kappa(S_\pi) \leq 1 + 1/(\pi N(1 + \pi))$.

Before examining the eigenvalues of S_π and T_π further, we state our version of the HO-CSD and HO-GSVD. The HO-CSD and HO-GSVD have already been described in [13] and [11], respectively, but our modified $D_{i,\pi}$ from (2.2) allows us to omit the requirements that A_i and Q_i be full rank.

DEFINITION 2.3 (HO-CSD). *If $\sum_{i=1}^N Q_i^T Q_i = I$, then the HO-CSD of $Q_i \in \mathbb{R}^{m_i \times n}$ is given by $Q_i = U_i \Sigma_i Z^T$, $i = 1, \dots, N$, where $Z \in \mathbb{R}^{n \times n}$ is such that $Z^T Z = I$ and $Z^T T_\pi Z = \text{diag}(\tau_1, \dots, \tau_n)$, with T_π defined as in (2.3). The matrices $\Sigma_i \in \mathbb{R}^{n \times n}$ with $\Sigma_i = \text{diag}(\sigma_{i,1}, \dots, \sigma_{i,n}) \succeq 0$ are obtained from*

$$B_i := Q_i Z, \quad B_i = [b_{i,1}, \dots, b_{i,n}], \quad \sigma_{i,k} = \|b_{i,k}\|_2,$$

and $U_i \in \mathbb{R}^{m_i \times n}$ with $U_i = [u_{i,1}, \dots, u_{i,n}]$ from

$$u_{i,k} = \begin{cases} b_{i,k}/\sigma_{i,k} & \text{if } \sigma_{i,k} > 0 \\ u \in \mathbb{R}^{m_i} \text{ with } \|u\|_2 = 1 & \text{if } \sigma_{i,k} = 0. \end{cases}$$

Because we allow for $\text{rank } Q_i < n$, it is possible that $Q_i z_k = 0$ for some eigenvector z_k of T_π , consequently making the corresponding generalized singular value $\sigma_{i,k} = 0$. In these cases, the column $u_{i,k}$ can be chosen freely or the corresponding row of Σ_i can be dropped. The columns of U_i have unit 2-norm and are, under certain circumstances, orthogonal (see Section 3).

DEFINITION 2.4 (HO-GSVD). *If the matrix A in (2.1) has full column-rank, then the HO-GSVD of the N matrices A_1, \dots, A_N is given by $A_i = U_i \Sigma_i V^T$, where $V \in \mathbb{R}^{n \times n}$ is such that $V^{-1} S_\pi V = \text{diag}(\varsigma_1, \dots, \varsigma_n)$ where S_π is defined in (1.2) using (2.2). The matrices $\Sigma_i \in \mathbb{R}^{n \times n}$ with $\Sigma_i = \text{diag}(\sigma_{i,1}, \dots, \sigma_{i,n}) \succeq 0$ are obtained from*

$$V B_i^T := A_i^T, \quad B_i = [b_{i,1}, \dots, b_{i,n}], \quad \sigma_{i,k} = \|b_{i,k}\|_2,$$

and $U_i \in \mathbb{R}^{m_i \times n}$ with $U_i = [u_{i,1}, \dots, u_{i,n}]$ from

$$u_{i,k} = \begin{cases} b_{i,k}/\sigma_{i,k} & \text{if } \sigma_{i,k} > 0 \\ u \in \mathbb{R}^{m_i} \text{ with } \|u\|_2 = 1 & \text{if } \sigma_{i,k} = 0. \end{cases}$$

3. Common and isolated subspaces. The HO-CSD and HO-GSVD identify directions, corresponding to columns of Z and V , that contribute equally to the right images of each Q_i and A_i , respectively. These directions are the right singular vectors that are shared across each Q_i and A_i and associated with identical singular values. The shared right singular vectors form subspaces [11, 13], which are referred to as the common HO-CSD and HO-GSVD subspaces. However, these have been defined for rank $Q_i = n$ and rank $A_i = n$. The following two theorems extend the notion of common subspaces to consider rank-deficient matrices.

THEOREM 3.1 (Common and isolated HO-CSD subspaces). *Suppose the assumptions on Q_i from Def. 2.3 hold. The common HO-CSD subspace is defined as*

$$\mathcal{T}_N \{Q_1, \dots, Q_N\} := \{z \in \mathbb{R}^n \mid T_\pi z = \tau_{\min} z\},$$

and the isolated HO-CSD subspace as

$$\mathcal{T}_1 \{Q_1, \dots, Q_N\} := \{z \in \mathbb{R}^n \mid T_\pi z = \tau_{\max} z\},$$

where τ_{\min} and τ_{\max} correspond to the left and right boundaries on the eigenvalues of T_π defined in Thm. 2.1. A vector $z \in \mathcal{T}_N$ is a right singular vector for each Q_i and associated with a singular value equal to $1/\sqrt{N}$. A vector $z \in \mathcal{T}_1$ is in the nullspace of all but one Q_i .

Proof. See Appendix B.

Note that for an arbitrary set of matrices Q_1, \dots, Q_N , the subspaces defined in Thm. 3.1 are likely to be empty. However, as conjectured in [11], the eigenvectors associated with eigenvalues for T_π , which lie close to the boundaries τ_{\min} or τ_{\max} , are likely to be directions that are equally amplified across all Q_i or singular to one particular Q_i .

THEOREM 3.2 (Common and isolated HO-GSVD subspaces). *Suppose the assumptions on A_i from Def. 2.4 hold. The common HO-GSVD subspace is defined as*

$$\mathcal{S}_N \{A_1, \dots, A_N\} := \{v \in \mathbb{R}^n \mid S_\pi v = \varsigma_{\min} v\},$$

and the isolated HO-GSVD subspace as

$$\mathcal{S}_1 \{A_1, \dots, A_N\} := \{v \in \mathbb{R}^n \mid S_\pi v = \varsigma_{\max} v\},$$

where ς_{\min} and ς_{\max} correspond to the left and right boundaries on the eigenvalues of S_π defined in Thm. 2.2. A vector $v \in \mathcal{S}_N$ is a right singular vector for each A_i and associated with an identical singular value. A vector $v \in \mathcal{S}_1$ is in the nullspace of all but one A_i .

Proof. Follows from Thm. 2.2 and 3.1, since if z is an eigenvector for T_π , then $v = R^T z$ is an eigenvector of S_π . Therefore if z is a right singular vector for each Q_i associated with an identical singular value, then v is a right singular vector for each A_i associated with an identical singular value. \square

4. Canonical Forms. The common subspaces capture the right singular vectors that are shared among matrices and have equal amplifications. For the HO-CSD and HO-GSVD, these directions are given by the eigenvectors of T_π and S_π , respectively, associated with a particular eigenvalue. It turns out that these directions produce the left basis vectors, given by the columns of U_i in Def. 2.3 and 2.4, that are orthogonal to all other columns of U_i . The following corollaries consider this additional property.

COROLLARY 4.1 (Canonical HO-CSD). *Suppose that the assumptions on Q_i from Def. 2.3 hold and let $Z^T T_\pi Z = \text{diag}(\tau_1, \dots, \tau_n)$ be the Schur decomposition of T_π with $Z = [z_1, \dots, z_n]$ and $\mathcal{T}_N \{Q_1, \dots, Q_N\} = \text{span}\{z_1, \dots, z_p\}$ the common HO-CSD subspace. Then, using Def. 2.3, $Q_i = U_i \Sigma_i Z^T$ with*

$$U_i = \begin{bmatrix} U_i^c & | & U_i^u \\ \hline p & & n-p \end{bmatrix}, \quad Z = \begin{bmatrix} Z^c & | & Z^u \\ \hline p & & n-p \end{bmatrix}, \quad \Sigma_i = \begin{bmatrix} I_p/\sqrt{N} & 0 \\ 0 & \Sigma_i^u \end{bmatrix},$$

where Σ_i^u is diagonal. The columns of each U_i^c are orthonormal and $(U_i^c)^T U_i^u = 0$. If, in addition, one of the $n - p$ column vectors z_k of Z^u is such that $z_k \in \mathcal{T}_1\{Q_1, \dots, Q_N\}$, then the corresponding column $u_{i,k}$ of each U_i^u can be chosen to be orthogonal to all other columns of U_i and the corresponding generalized singular value $\sigma_{i,k}$ is either zero or equal to 1.

Proof. From Thm. 3.1, a vector $z_k \in \mathcal{T}_N\{Q_1, \dots, Q_N\}$ is a right singular vector for each Q_i associated with a (generalized) singular value $\sigma_{i,k} = 1/\sqrt{N}$, whereas a vector $z_k \in \mathcal{T}_1\{Q_1, \dots, Q_N\}$ is a right singular vector for each Q_i associated with a (generalized) singular value $\sigma_{i,k}$ that is either zero or equal to 1. For $z_k \in \mathcal{T}_N\{Q_1, \dots, Q_N\}$, let $b_{i,k} = Q_i z_k = \sigma_{i,k} z_k$ denote the columns of the matrix B_i from Def. 2.3. Then for $k \neq j$ and each Q_i

$$b_{i,k}^T b_{i,j} = (Q_i z_k)^T Q_i z_j = (Q_i^T Q_i z_k)^T z_j = \sigma_{i,k} z_k^T z_j = 0.$$

The property $(U_i^c)^T U_i^u = 0$ follows from Def. 2.3. For $z_k \in \mathcal{T}_1\{Q_1, \dots, Q_N\}$, Def. 2.3 allows for a free choice of the corresponding column $u_{i,k}$. Choose this particular column to be the (standard) left singular vector associated with a singular value that is either zero or 1. The orthogonality property follows. \square

COROLLARY 4.2 (Canonical HO-GSVD). *Suppose that the assumptions on A_i from Def. 2.4 hold and let $V^{-1} S_\pi V = \text{diag}(\varsigma_1, \dots, \varsigma_n)$ be the Schur decomposition of S_π with $V = [v_1, \dots, v_n]$ and $\mathcal{S}_N\{A_1, \dots, A_N\} = \text{span}\{v_1, \dots, v_p\}$ the common HO-CSD subspace. Then, using Def. 2.4, $A_i = U_i \Sigma_i V^T$ with*

$$U_i = \begin{bmatrix} U_i^c & | & U_i^u \\ p & & n-p \end{bmatrix}, \quad V = \begin{bmatrix} V^c & | & V^u \\ p & & n-p \end{bmatrix}, \quad \Sigma_i = \begin{bmatrix} I_p / \sqrt{N} & 0 \\ 0 & \Sigma_i^u \end{bmatrix},$$

where Σ_i^u is diagonal. The columns of each U_i^c are orthonormal and $(U_i^c)^T U_i^u = 0$. If, in addition, one of the $n - p$ columns vectors v_k of V^u is such that $v_k \in \mathcal{S}_1\{A_1, \dots, A_N\}$, then the corresponding column $u_{i,k}$ of each U_i^u is orthogonal to all other columns of U_i and the corresponding generalized singular value $\sigma_{i,k}$ is either zero or equal to 1.

Proof. Follows directly from Corollary 4.1 and $v_k = R^T z_k$. \square

5. Comparison with standard HO-GSVD, GSVD and SVD. When one out of two matrices is the identity matrix, the GSVD reduces to the standard SVD [12]. The same has been shown for the full-rank HO-GSVD [11]. When $N - 1$ matrices A_i are identity matrices, then the full-rank HO-GSVD reverts to the standard SVD of A_j , $j \neq i$. Here, this fact is demonstrated for our HO-GSVD as given in Def. 2.4.

LEMMA 5.1. *Suppose that $N = 2$, $A_1 \in \mathbb{R}^{m_1 \times n}$, $A_2 = I_n$ and $\text{rank}[A_1^T, A_2^T]^T = n$. Then the HO-GSVD of A_1 and A_2 yields the standard SVD of A_1 .*

Proof. See Appendix C.

THEOREM 5.2. *Suppose that $N > 2$. The HO-GSVD of $N - 1$ matrices satisfying $A_i = I$ yields the SVD of A_j , $i \neq j$.*

Proof. See Appendix D.

The HO-GSVD from Def. 2.4 can also be related to the GSVD. For the special case that $N = 2$, $A_1 \in \mathbb{R}^{m_1 \times n}$ with $m_1 \geq n$ and $\text{rank} A_1 = n$ and an arbitrary $A_2 \in \mathbb{R}^{m_2 \times n}$, it can be shown that the HO-GSVD yields Σ_i with $\Sigma_1^T \Sigma_1 + \Sigma_2^T \Sigma_2 = I$ and orthonormal U_1 and U_2 .

LEMMA 5.3. *Let $N = 2$, then T_π from Def. 2.3, $(Q_i^T Q_i + \pi I)^{-1}$ and $Q_i^T Q_i$ with $i = 1, 2$ and $Q_1^T Q_1 + Q_2^T Q_2 = I$ share the same eigenspace for any $\pi \in \mathbb{R}_+$.*

Proof. See Appendix E.

THEOREM 5.4. *For $N = 2$, the HO-CSD from Def. 2.3 yields the CSD.*

Proof. Using Lemma 5.3, the eigenvectors z_k for T_π can be chosen such that they are right singular vectors for Q_1 and Q_2 . Let $b_{i,k}$ denote the columns of $B_i = Q_i Z$, then for $j \neq k$,

$$b_{i,k}^T b_{i,j} = (Q_i z_k)^T Q_i z_j = (\bar{\sigma}_{i,k} \bar{u}_{i,k})^T \bar{\sigma}_{i,j} \bar{u}_{i,j} = 0,$$

where $\bar{\sigma}_\times$ and \bar{u}_\times denote standard singular values and left singular vectors, respectively. Hence, from $U_i \Sigma_i = B_i$, the columns of U_i are either zero or orthonormal. Substituting $Q_i = U_i \Sigma_i V^T$ in $Q_1^T Q_1 + Q_2^T Q_2 = I$ yields $Z \Sigma_1^T \Sigma_1 Z^T + Z \Sigma_2^T \Sigma_2 Z^T = I$, and from $Z^T Z = I$, follows $\Sigma_1^T \Sigma_1 + \Sigma_2^T \Sigma_2 = I$. \square

COROLLARY 5.5. *For $N = 2$, $A_1 \in \mathbb{R}^{m_1 \times n}$ with $m_1 \geq n$ and $\text{rank} A_1 = n$ and an arbitrary $A_2 \in \mathbb{R}^{m_2 \times n}$, the HO-GSVD from Def. 2.4 yields the GSVD [12].*

Proof. Follows from Thm. 5.4 with $V = R^T Z$. \square

Remark 5.6. Note that Thm. 5.4 and Corollary 5.5 must be handled with caution. Lemma 5.3 shows that the three matrices, T_π , $Q_1^T Q_1$ and $Q_2^T Q_2$, share the same eigenspace, but not every eigendecomposition of T_π yields eigenvectors that are parallel to those of $Q_1^T Q_1$ and $Q_2^T Q_2$. For example, suppose that $\dim(\ker(Q_i)) = 1$ and that $q_i \in \ker(Q_i)$, $i = 1, 2$, are linearly independent. Note that from $Q_1^T Q_1 + Q_2^T Q_2 = I$, it holds that $q_1^T q_2 = 0$. It follows that $\dim(\mathcal{T}_1) = 2$, so that T_π has an eigenvalue with algebraic and geometric multiplicity of two. When the associated eigenvectors are computed using numerical software, these will not necessarily be parallel to q_1 and q_2 , and the HO-CSD will not necessarily yield orthonormal matrices U_i .

The HO-GSVD from Def. 2.4 can also be compared with the full-rank HO-GSVD [11]. For $N = 2$ and full-rank matrices A_i , both HO GSVDs have been shown to be equivalent to the GSVD. Both HO GSVDs have also been shown to yield the SVD of A_j when $A_i = I$ for $i \neq j$.

For $N > 2$, however, the HO-GSVD from Def. 2.4 and [11] will in general *not* yield identical factorizations $A_i = U_i \Sigma_i V^T$, even when $\text{rank } A_i = n$. This can be seen by comparing the eigenspaces of T_π from (2.3) for varying π , where $\pi = 0$ corresponds to the standard HO-CSD [13]. For $N = 2$, Lemma 5.3 shows that the eigenvectors of T_π are independent of the value of π because its eigenvectors are fixed by the orthogonality property $Q_1^T Q_1 + Q_2^T Q_2 = I$, while for $N > 2$ this property is lost. From $V = R^T Z$, it follows that the same holds for the HO-GSVD. However, it can be shown that in case the matrices A_i and Q_i have full rank, then the common HO-CSD and HO-GSVD subspaces will be the same for any value of π .

LEMMA 5.7. *The common and isolated subspaces from Thm. 3.1 and 3.2 are independent of the value of π .*

Proof. Follows from the definitions of the common and isolated subspaces (see Appendix B), which require that for $z \in \mathcal{T}_N$ and $v \in \mathcal{S}_N$, z and v must be right singular vectors for each Q_i and A_i , respectively. This requirement is independent of the value of π . \square

6. Numerical Example. To illustrate the common subspaces of order $N - P$, we consider the following example with $N = 3$ matrices,

$$(6.1) \quad A = \begin{bmatrix} -\frac{A_1}{-} \\ -\frac{A_2}{-} \\ -\frac{A_3}{-} \end{bmatrix} = \begin{bmatrix} -\frac{a_1}{-} & -\frac{a_2}{-} \\ -\frac{0}{-} & -\frac{1}{-} \\ -\frac{1}{-} & -\frac{0}{-} \end{bmatrix}, \quad (a_1, a_2) = \begin{cases} (1, 0) & \text{for case 1} \\ (1, 1) & \text{for case 2} \end{cases}$$

where $\text{rank } A = 2$. The matrix A_2 (A_3) has right singular vectors $[1, 0]^T$ and $[0, 1]^T$ associated with singular values 0 and 1 (1 and 0), respectively. For each of the cases, the matrix S_π from (1.2) is obtained as

$$(6.2) \quad S_\pi \Big|_{\text{case 1}} = \begin{bmatrix} \frac{1+6\pi+12\pi^2}{6\pi+12\pi^2} & 0 \\ 0 & \frac{1+3\pi+3\pi^2}{3\pi+3\pi^2} \end{bmatrix}, \quad S_\pi \Big|_{\text{case 2}} = \begin{bmatrix} \frac{1+4\pi+6\pi^2}{4\pi+6\pi^2} & 0 \\ 0 & \frac{1+4\pi+6\pi^2}{4\pi+6\pi^2} \end{bmatrix},$$

with eigenvalues that can be read from the diagonal.

For case 1, $A_1 = A_3$ and comparison with Thm. 3.2 shows that the eigenvector $[0, 1]^T$ spans \mathcal{S}_1 and is a common right singular vector for each A_i associated with a zero singular value for A_1 and A_3 and with a singular value equal to 1 for A_2 .

For case 2, none of the A_i have common right singular vectors associated with identical singular values. Comparison with Thm. 3.2 shows that none of the eigenvalues of S_π correspond to a value that would yield a non-empty \mathcal{S}_3 or \mathcal{S}_1 .

7. Applications. The HO-GSVD has already been applied in various fields, and the following paragraphs outline applications for which the rank-deficient case is relevant.

Bioinformatics: The (HO)-GSVD has been used for genomic signal processing [11, 15, 9]. In [11], the $N > 2$ matrices A_i represent measurements taken from different organism with the rows of A_i representing organism specific genes and the columns time instants. Because the study includes several thousands of genes, but only few time instants, the matrices have full-column rank. In other studies, such as [15] or [17], the n columns of A_i are genes that are shared across different organism. In these cases, the standard HO-GSVD cannot be applied because the matrices typically satisfy $m_i < n$. In [9], $N = 2$ vaccines are analyzed by comparing datasets A_i that represents n genes and measurements across m_i patients. These matrices also have dimensions $m_i < n$ and our HO-GSVD extension could be used to extend the study to $N > 2$ vaccines.

Neuroscience: In [16], a tensor decomposition has been used to factorize matrices obtained from $N > 2$ trials, where the n columns of A_i model a large number of neurons, while the m_i rows represent the time instants at which the measurements are taken. These matrices typically have more columns than rows and the tensor

decomposition used in [16] requires that $m_i = m_j$ for $i, j = 1, \dots, N$. Our HO-GSVD extension could be used for such a factorization and, in contrast to the tensor decomposition, our decomposition would allow for varying m_i across the trials.

Artificial Intelligence: Tensor decompositions are widely used in artificial intelligence applications [4]. The decompositions serve to classify data or extract useful information from a vast amount of data. In a classification problem, for example, the data is composed out of m_i classified examples with n features that are shared among N data classes. For the case that $m_i < n$, our HO-GSVD extension could be used to identify shared and unshared features of the different data classes.

Control Theory: In [8], the GSVD has been used to diagonalize the orbital dynamics of electrons in a synchrotron with $N = 2$ actuator (magnet) arrays. The actuator matrices typically have full row-rank and fewer rows than columns. Our HO-GSVD extension could be used to diagonalize a system with $N > 2$ actuator arrays, such as encountered in the cross-directional control of paper machines [14]. In [1], the GSVD has been used to design an observer. Our HO-GSVD extension could be used to simultaneously design the observer and the controller.

8. Conclusion. In this paper, we have extended the standard HO-GSVD [11] to accommodate column rank-deficient matrices. By adding the term $\pi A^T A$ to each of the quotient terms $D_{i,\pi} = A_i^T A_i + \pi A^T A$, we shifted their eigenvalues and bounded them away from zero. This allowed for omitting the full-rank requirement on each A_i and extending the notion of common subspaces to consider the right nullspace of each A_i .

The choice of adding a multiple of $A^T A$ was motivated by the relationship between the arithmetic mean of pairwise quotients S_π and the mean of reciprocal inverses T_π , which yielded the same relationship than in [11] for $\pi = 0$. The parameter π was assumed to be positive, but otherwise left unspecified. The common HO-CSD and HO-GSVD subspaces are identified irrespective of the value of π . We could have chosen to define $D_{i,\pi}$ as $D_{i,\pi_i} = A_i^T A_i + \pi_i A^T A$ with a parameter π_i that is different for each term. The resulting relationship between S_π and T_π would have been similar to (2.4). An indexed parameter π_i , for example, could account for the case that the matrices A_i share a right singular vector v associated with different singular values and an appropriate choice of π_i could make v an eigenvector for S_π . Future research could investigate the role of the weight π or the application of indexed parameter π_i .

The majority of our developments were based on the HO-CSD. Using the QR factorization of $A = [A_1^T, \dots, A_N^T]^T$, each A_i was represented as $A_i = Q_i R$ and our findings were developed for Q_1, \dots, Q_N , from which we inferred the corresponding properties for A_1, \dots, A_N . This procedure allowed us to exploit the orthogonality property $\sum_{i=1}^N Q_i^T Q_i = I$ and ensure the invertibility of the factor matrix V in $A_i = U_i \Sigma_i V^T$, but required a full column-rank A , so that $\det R \neq 0$. For a column rank-deficient A , the HO-GSVD remains applicable iff the $D_{i,\pi}$ are invertible, but it is unclear whether the matrix V remains invertible. The invertibility of V is a key requirement for the application of matrix factorizations, as it allows one to switch from the original space to the modal space. A further investigation could focus on elaborating properties for the arithmetic mean of pairwise quotients S_π that are not based on the HO-CSD.

We bounded the eigenvalues of T_π (S_π) and showed that the extremal eigenvalues are attained iff the corresponding eigenvectors are right singular vectors for each Q_i (A_i) associated with a particular singular value. This led to the definition of the common and isolated HO-CSD (HO-GSVD) subspaces. In Appendix B, we also showed that if the Q_i share a right singular vector v associated with a zero singular value for P matrices Q_i and with an identical singular value for the other $N - P$ matrices Q_j , then T_π will have a particular eigenvalue $\tau(P)$ associated with the eigenvector v . Future research could investigate whether a biconditional (“iff”) connection holds.

REFERENCES

- [1] G. P. DADA AND A. ARMAOU, *110th anniversary: Generalized singular value decomposition reduced-order observers for linear time-invariant systems with noisy measurements*, Ind. Eng. Chem. Process Des. Dev., 58 (2019), pp. 23201–23210.
- [2] L. DE LATHAUWER, B. DE MOOR, AND J. VANDEWALLE, *A multilinear singular value decomposition*, SIAM J. Matrix Anal. Appl., 21 (2000), pp. 1253–1278.
- [3] G. H. GOLUB AND C. F. VAN LOAN, *Matrix Computations*, JHU Press, Baltimore, MD, USA, 4 ed., 2013.
- [4] HAIPING LU, K. N. PLATANIOTIS, AND A. N. VENETSANOPOULOS, *A survey of multilinear subspace learning for tensor data*, Pattern Recognit., 44 (2011), pp. 1540–1551.
- [5] G. H. HARDY, J. E. LITTLEWOOD, AND G. PÓLYA, *Inequalities*, CUP, Cambridge, UK, 1 ed., 1934.
- [6] R. A. HARSHMAN, *Foundations of the PARAFAC procedure: Models and conditions for an “explanatory” multimodal factor analysis*, UCLA Work. Pap. Phon., 16 (1970), pp. 1–84.
- [7] R. A. HARSHMAN, *PARAFAC2: Mathematical and technical notes*, UCLA Work. Pap. Phon., 22 (1972), pp. 30–44.
- [8] I. KEMPF, S. R. DUNCAN, P. J. GOULART, AND G. REHM, *Multi-array electron beam stabilization using block-circulant transformation and generalized singular value decomposition*, in Proc. 59th CDC, Jeju Island, Republic of Korea, Dec. 2020.

- [9] K. VAN DEUN, A. K. SMILDE, L. THORREZ, H. A. L. KIERS, AND I. VAN MECHELEN, *Identifying common and distinctive processes underlying multiset data*, Chemom. Intell. Lab. Syst., 129 (2013), pp. 40–51.
- [10] D. LAHAT, T. ADALI, AND C. JUTTEN, *Multimodal data fusion: An overview of methods, challenges, and prospects*, Proc. IEEE, 103 (2015), pp. 1449–1477.
- [11] S. P. PONNAPALLI, M. A. SAUNDERS, C. F. VAN LOAN, AND O. ALTER, *A higher-order generalized singular value decomposition for comparison of global mRNA expression from multiple organisms*, PLOS ONE, 6 (2011), pp. 1–11.
- [12] C. F. VAN LOAN, *Generalizing the singular value decomposition*, SIAM J. Numer. Anal., 13 (1976), pp. 76–83.
- [13] C. F. VAN LOAN, *Lecture 6. The higher-order generalized singular value decomposition*, June 2015, <http://www.dm.unibo.it/%7Esimoncin/CIME/Vanloan.Lec6.pdf>.
- [14] J. G. VANANTWERP, A. P. FEATHERSTONE, R. D. BRAATZ, AND B. A. OGUNNAIKE, *Cross-directional control of sheet and film processes*, Automatica, 43 (2007), pp. 191–211.
- [15] L. J. VAN'T VEER, D. HONGYUE, M. J. VAN DE VIJVER, Y. D. HE, A. A. M. HART, M. MAO, H. L. PETERSE, K. VAN DER KOOY, M. J. MARTON, A. T. WITTEVEEN, G. J. SCHREIBER, R. M. KERKHOVEN, C. ROBERTS, P. S. LINSLEY, R. BERNARDS, AND S. H. FRIEND, *Gene expression profiling predicts clinical outcome of breast cancer*, Nature, 415 (2002), pp. 530–536.
- [16] A. H. WILLIAMS, T. H. KIM, F. WANG, S. VYAS, S. I. RYU, K. V. SHENOY, M. SCHNITZER, T. G. KOLDA, AND S. GANGULI, *Unsupervised discovery of demixed, low-dimensional neural dynamics across multiple timescales through tensor component analysis*, Neuron, 98 (2018), pp. 1099–1115.
- [17] B. ZHANG AND S. HORVATH, *A general framework for weighted gene co-expression network analysis*, Stat. Appl. Genet. Mol. Biol., 4 (2005).

Appendix A. Relation between S_π and T_π . Let $D_{i,\pi} = Q_i^T Q_i + \pi I$ and define $K_i := Q_i^T Q_i + \pi I$. Using (2.2), the matrix S_π is written as

$$\begin{aligned} S_\pi &= \frac{1}{N(N-1)} \sum_{i=1}^N \sum_{j=i+1}^N (D_{i,\pi} D_{j,\pi}^{-1} + D_{j,\pi} D_{i,\pi}^{-1}) \\ &= \frac{1}{N(N-1)} R^T \left(\sum_{i=1}^N \sum_{j=i+1}^N K_i K_j^{-1} + K_j K_i^{-1} \right) R^{-T}, \end{aligned}$$

so that by considering $\sum_i (K_i + \pi I) = (1 + \pi N)I$

$$\begin{aligned} R^{-T} S_\pi R^T &= \frac{1}{N(N-1)} \sum_{i=1}^N \sum_{j=i+1}^N K_i K_j^{-1} + K_j K_i^{-1} = \frac{1}{N(N-1)} \sum_{i=1}^N K_i \sum_{j=1}^N K_j^{-1} - \frac{1}{N-1} I \\ &= \frac{1}{N-1} ((1 + \pi N)T_\pi - I). \end{aligned}$$

Appendix B. Characterizing T_π . The eigenspace of T_π defined in (2.3) with $\sum_{i=1}^N Q_i^T Q_i = I$ and $\|Q_i\|_2 \leq 1$ has properties that can be used to characterize common subspaces of Q_1, \dots, Q_N , i.e. common features of the matrices Q_i . By setting $u = (Q_i^T Q_i + \pi I)^{\frac{1}{2}} t$ and $v = (Q_i^T Q_i + \pi I)^{-\frac{1}{2}} t$ with $\|t\|_2 = 1$, the Cauchy-Schwarz inequality $(u^T v)^2 \leq \|u\|_2^2 \|v\|_2^2$ becomes

$$(B.1) \quad t^T (Q_i^T Q_i + \pi I)^{-1} t \geq (t^T (Q_i^T Q_i + \pi I) t)^{-1},$$

with equality iff t is an eigenvector for $Q_i^T Q_i$ [5, Thm. 7]. Using (B.1) and the harmonic-mean arithmetic-mean (HM-AM) inequality [5, Thm. 16], a lower bound on $t^T T_\pi t$, $\|t\|_2 = 1$, can be established as

$$\begin{aligned} (B.2a) \quad t^T T_\pi t &= \frac{1}{N} \sum_{i=1}^N t^T (Q_i^T Q_i + \pi I)^{-1} t \geq \frac{1}{N} \sum_{i=1}^N \frac{1}{t^T (Q_i^T Q_i + \pi I) t} \\ (B.2b) \quad &\geq \frac{N}{\pi N + \sum_{i=1}^N t^T (Q_i^T Q_i) t} = \frac{N}{\pi N + 1}. \end{aligned}$$

From the Cauchy-Schwarz inequality, (B.2a) becomes an equality iff t is a common eigenvector for each $Q_i^T Q_i + \pi I$ or equivalently for each $Q_i^T Q_i$. From the HM-AM inequality, inequality (B.2b) becomes an equality iff $t^T (Q_i^T Q_i + \pi I) t = t^T (Q_j^T Q_j + \pi I) t$ for $i, j = 1, \dots, N$. In this case, it follows that

$$1 = t^T Q^T Q t = \sum_{i=1}^N t^T (Q_i^T Q_i) t = \sum_{i=1}^N \mu_Q = N \mu_Q,$$

so that the common eigenvalue of $Q_i^T Q_i$ associated with the common eigenvector y is $\mu_Q = 1/N$. The common vector t is therefore a common right singular vector for each Q_i with singular value $\sigma_{i,k} = 1/\sqrt{N}$. According to (B.2), t is an eigenvector for T_π associated with the smallest eigenvalue $\tau_i = N/(1 + \pi N)$.

An upper bound on the eigenvalues of T_π can be found by noticing that for each term, $t^\top (Q_i^\top Q_i + \pi I)^{-1} t \leq 1/\pi$ for $\|t\|_2 = 1$, which results in

$$(B.3) \quad t^\top T_\pi t < 1/\pi,$$

so that the maximum eigenvalue of T_π is bounded above by $1/\pi$. Note that the strict inequality in (B.3) holds because $\sum_{i=1}^N Q_i^\top Q_i = I$ and t can not simultaneously be in the nullspace of each Q_i . The orthogonality condition $\sum_{i=1}^N Q_i^\top Q_i = I$ leads to a tighter bound that is established in the following theorem.

THEOREM B.1. *It holds that $\rho(T_\pi) \leq (\pi N + N - 1)/(\pi N(1 + \pi))$, where T_π is defined in (2.3) and $\rho(T_\pi)$ denotes the maximum eigenvalue of T_π . Equality holds iff there exists a vector that lies in the kernel of all but one $Q_i^\top Q_i$.*

Proof. Let $\bar{V}_i \bar{\Sigma}_i^2 \bar{V}_i^\top = Q_i^\top Q_i$ be the eigendecomposition of $Q_i^\top Q_i$ and set $\bar{\Sigma}_i^2 = \text{diag}(\bar{\sigma}_{i,j}^2)$, $j = 1, \dots, n$, so that $(\bar{\Sigma}_i^2 + \pi I)^{-1} = \text{diag}(\bar{\sigma}_{i,j}^2 + \pi)^{-1}$. Let t with $\|t\|_2 = 1$ be an eigenvector of T_π associated with $\rho(T_\pi)$, then

$$\rho(T_\pi) = t^\top T_\pi t = \frac{1}{N} \sum_{i=1}^N t^\top \bar{V}_i \text{diag}\left(\frac{1}{\bar{\sigma}_{i,1}^2 + \pi}, \dots, \frac{1}{\bar{\sigma}_{i,n}^2 + \pi}\right) \bar{V}_i^\top t.$$

For each i , express t as $t = \sum_{j=1}^n a_{i,j} \bar{v}_{i,j}$, where $\bar{v}_{i,j}$ are the columns of the orthonormal matrices \bar{V}_i and $a_{i,j} \in \mathbb{R}$ with $\sum_{j=1}^n a_{i,j}^2 = 1$ because $\|t\|_2 = 1$. In addition, define $y_{i,j} := \bar{\sigma}_{i,j}^2 + \pi \in [\pi, 1 + \pi]$ and rewrite $\rho(T_\pi)$ as

$$\rho(T_\pi) = t^\top T_\pi t = \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^n \frac{a_{i,j}^2}{y_{i,j}}, \quad \sum_{j=1}^n a_{i,j}^2 = 1, \quad y_{i,j} \in [\pi, 1 + \pi].$$

Next, bound each of the terms $1/y_{i,j}$ by a function that is linear in $y_{i,j}$, i.e. $f(y_{i,j}) = m y_{i,j} + c$ with $1/y_{i,j} \leq f(y_{i,j})$ for $y_{i,j} \in [\pi, 1 + \pi]$. Note that the function $1/y_{i,j}$ is convex on $y_{i,j} \in [\pi, 1 + \pi]$ and m and c can be chosen such that equality holds at $y_{i,j} = \pi$ and $y_{i,j} = 1 + \pi$, i.e. by selecting m and c as

$$m = \frac{-1}{\pi(1 + \pi)}, \quad c = \frac{1 + 2\pi}{\pi(1 + \pi)}.$$

We obtain a bound on $\rho(T_\pi)$ as

$$\rho(T_\pi) \leq \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^n a_{i,j}^2 \frac{-y_{i,j} + 1 + 2\pi}{\pi(1 + \pi)} = \left(\frac{1}{N} \sum_{i=1}^N \sum_{j=1}^n \frac{-a_{i,j}^2 y_{i,j}}{\pi(1 + \pi)} \right) + \frac{1 + 2\pi}{\pi(1 + \pi)},$$

where we used that $\sum_{j=1}^n a_{i,j}^2 = 1$. Clearly, equality holds iff the $y_{i,j}$ associated with non-zero $a_{i,j}^2$ are at the boundaries of $[\pi, 1 + \pi]$. Finally, exploit the orthogonality condition $\sum_{i=1}^N Q_i^\top Q_i = \sum_{i=1}^N \bar{V}_i \bar{\Sigma}_i^2 \bar{V}_i^\top = I$. By adding $N\pi I$, we can express the same condition using $y_{i,j}$ as $\sum_{i=1}^N \bar{V}_i \text{diag}(y_{i,j}) \bar{V}_i^\top = (1 + N\pi)I$. Left- and right-multiplying the modified orthogonality condition by t^\top and t , respectively, and expressing t using the columns $\bar{v}_{i,j}$ as above yields

$$\sum_{i=1}^N t^\top \bar{V}_i \text{diag}(y_{i,j}) \bar{V}_i^\top t = \sum_{i=1}^N \sum_{j=1}^n a_{i,j}^2 y_{i,j} = 1 + N\pi.$$

Substituting in the upper bound on $\rho(T_\pi)$ yields

$$\rho(T_\pi) \leq -\frac{1 + N\pi}{\pi(1 + \pi)} + \frac{1 + 2\pi}{\pi(1 + \pi)} = \frac{1}{N} \left(\frac{N-1}{\pi} + \frac{1}{1 + \pi} \right) = \frac{\pi N + N - 1}{\pi N(1 + \pi)}.$$

Equality holds iff each $y_{i,j} = \bar{\sigma}_{i,j}^2 + \pi$ is at the boundaries of $[\pi, 1 + \pi]$, i.e. either $y_{i,j} = \pi$, in which case $\bar{\sigma}_{i,j}^2 = 0$, or $y_{i,j} = 1 + \pi$, in which case $\bar{\sigma}_{i,j}^2 = 1$. Note that the orthogonality condition requires at least one non-zero $\bar{\sigma}_{i,j}^2$. In addition, if $\bar{\sigma}_{i,j}^2 = 1$ associated with $\bar{v}_{i,j}$ for some i and j , then the vector $\bar{v}_{i,j}$ must be shared among all \bar{V}_i and associated with $\bar{\sigma}_{k,j}^2 = 0$ for $k \neq i$, because $\bar{v}_{i,j}^\top (\sum_{i=1}^N \bar{V}_i \bar{\Sigma}_i^2 \bar{V}_i^\top) \bar{v}_{i,j} > 1$ otherwise. It therefore holds that $\rho(T_\pi) = (\pi N + N - 1)/(\pi N(1 + \pi))$ iff the $Q_i^\top Q_i$ have a shared eigenvector associated with a zero eigenvalue for $N - 1$ matrices and a unit eigenvalue for one matrix. \square

Finally, note that if there exists a vector t with $\|t\|_2 = 1$ in the nullspace of P matrices Q_j , but in the range of all other Q_i with index $i \in \mathcal{R}$, then the inequalities (B.2a)-(B.2b) can be reformulated as

$$(B.4a) \quad \begin{aligned} t^T T_\pi t &= \frac{1}{N} \sum_{i=1}^N t^T (Q_i^T Q_i + \pi I)^{-1} t = \frac{P}{\pi N} + \frac{1}{N} \sum_{i \in \mathcal{R}} t^T (Q_i^T Q_i + \pi I)^{-1} t \\ &\geq \frac{P}{\pi N} + \frac{1}{N} \sum_{i \in \mathcal{R}} \frac{1}{t^T (Q_i^T Q_i + \pi I) t} \end{aligned}$$

$$(B.4b) \quad \geq \frac{P}{\pi N} + \frac{N-P}{N} \frac{N-P}{\pi(N-P) + \underbrace{\sum_{i \in \mathcal{R}} t^T (Q_i^T Q_i) t}_{=1}} = \frac{P(1-\pi N) + \pi N^2}{\pi N(1 + \pi(N-P))}.$$

The term on the right-hand side of (B.4b) corresponds to the minimum and maximum eigenvalues of T_π for $P = 0$ and $P = N - 1$, respectively. If there exists a shared vector t in the nullspace of P matrices Q_j , but in the range of all other Q_i , then an eigenvalue of T_π will be equal to the corresponding value on the right-hand side of (B.4b). Note that (B.4) does *not* prove the converse.

Appendix C. Proof of Lemma 5.1. Let $\bar{U}_1 \bar{\Sigma}_1 \bar{V}_1^T = A_1$ be the standard SVD of A_1 with $\bar{\Sigma}_1 \in \mathbb{R}^{m_1 \times n} \succeq 0$. Substitute the SVD and $A_2 = I_n$ in (1.2) using (2.2):

$$(C.1) \quad S_\pi = \frac{1}{2} \bar{V}_1 \left((\tilde{\pi} \bar{\Sigma}_1^T \bar{\Sigma}_1 + I) (\tilde{\pi} I + \bar{\Sigma}_1^T \bar{\Sigma}_1)^{-1} + (\tilde{\pi} I + \bar{\Sigma}_1^T \bar{\Sigma}_1) (\tilde{\pi} \bar{\Sigma}_1^T \bar{\Sigma}_1 + I)^{-1} \right) \bar{V}_1^T,$$

where $\tilde{\pi} = (1 + \pi)/\pi$. Therefore, $S_\pi = S_\pi^T \succ 0$, hence non-defective with an orthonormal eigenbasis $V = \bar{V}_1$. According to Def. 2.4, the HO-GSVD $A_1 = U_1 \Sigma_1 V^T$ is obtained from $V B_1^T = A_1^T$ and $U_1 \Sigma_1 = B_1$. We show that B_1 has orthogonal columns. From the QR factorization (2.1), $A_2 = I = Q_2 R$ so that $Q_2 = R^{-1}$,

$$I = Q_1^T Q_1 + Q_2^T Q_2 = Q_1^T Q_1 + R^{-T} R^{-1},$$

and

$$(C.2) \quad R^T R - I = R^T Q_1^T Q_1 R = A_1^T A_1.$$

Therefore,

$$(C.3) \quad B_1^T B_1 = V^T A_1^T A_1 V = V^T (R^T R - I) V \stackrel{Z=RV}{=} Z^T Z - V^T V,$$

where the columns of Z are the orthogonal eigenvectors of $T_\pi = T_\pi^T$, thus the matrix on the right-hand side is diagonal. We follow that B_1 has orthogonal columns. If A_1 has $r = \text{rank } A_1 < n$, then $n - r$ columns of B_1 will be zero and, from $U_1 \Sigma_1 = B_1$, the associated columns of U_1 can be chosen freely. These can always be chosen to have unit length and such that U_1 has at least m_1 orthogonal columns. We obtain a factorization $A_1 = U_1 \Sigma_1 V^T$, where $\Sigma_1 \succeq 0$ is diagonal and V is orthonormal. If $U_1 \in \mathbb{R}^{m_1 \times n}$ has more columns than rows, i.e. $m_1 < n$, we remove $n - m_1$ of the linear dependent columns of U_1 and the corresponding rows of Σ_1 , which are necessarily zero-rows. The remaining part of U_1 is orthonormal, thus the HO-GSVD of A_1 must be the standard SVD of A_1 [11, Thm. S5]. \square

Appendix D. Proof of Lemma 5.2. Follows from Lemma 5.1 and from the fact that the matrices M and $\tilde{M} := aM + bI$, where $a, b \in \mathbb{R}$, share the same eigenspace so that all right singular vectors for $A_j \neq I$ are eigenvectors for S_π as in (C.1). For $N - 1$ matrices $A_i = I$, $i \neq j$, equation (C.2) becomes $A_j^T A_j = R^T R - (N - 1)I$. From substituting $A_j^T A_j$ in equation C.3, it can be seen that $B_j^T B_j$ is orthogonal. \square

Appendix E. Proof of Lemma 5.3. Let t be an eigenvector for $Q_1^T Q_1$ associated with eigenvalue λ . Then, using $Q_1^T Q_1 + Q_2^T Q_2 = I$,

$$Q_2^T Q_2 t = (I - Q_1^T Q_1) t = (1 - \lambda) t,$$

so that t is an eigenvector for $Q_2^T Q_2$ associated with eigenvalue $1 - \lambda$. Note that t is also an eigenvector for $(Q_1^T Q_1 + \pi I)^{-1}$ and $(Q_2^T Q_2 + \pi I)^{-1}$ associated with eigenvalues $1/(\pi + \lambda)$ and $1/(1 + \pi - \lambda)$, respectively. From 2.3,

$$T_\pi t = \frac{1}{2} \left((Q_1^T Q_1 + \pi I)^{-1} + (Q_2^T Q_2 + \pi I)^{-1} \right) t = \frac{1}{2} \left(\frac{1}{\pi + \lambda} + \frac{1}{1 + \pi - \lambda} \right) t,$$

i.e. t is an eigenvector for T_π associated with eigenvalue $\frac{1/2 + \pi}{(\pi + \lambda)(1 + \pi - \lambda)}$. \square