

Finite-size scaling analysis of eigenstate thermalization

Yichen Huang (黄溢辰)*

Center for Theoretical Physics, Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139, USA

April 6, 2021

Abstract

We study the fluctuations of eigenstate expectation values in a microcanonical ensemble. Assuming the eigenstate thermalization hypothesis, an analytical formula for the finite-size scaling of the fluctuations is derived. The same problem was studied by Beugeling et al. [Phys. Rev. E 89, 042112 (2014)]. We compare our results with theirs.

1 Introduction

The eigenstate thermalization hypothesis (ETH) [5, 14, 13, 4, 6] is a sufficient condition for the emergence of statistical mechanics from the unitary evolution of isolated quantum many-body systems. It states that eigenstates that are close in energy have similar local expectation values, i.e., the fluctuations of eigenstate expectation values (EEV) in a microcanonical ensemble vanish in the thermodynamic limit.

What is the asymptotic behavior of EEV fluctuations as the system size diverges? The answer to this question depends on how EEV fluctuations are defined. Assuming the ETH, we rigorously derive an analytical formula for the finite-size scaling of some definitions (including the one in Ref. [2]) of EEV fluctuations.

The rest of this paper is organized as follows. Section 2 sets the stage and introduces some definitions of EEV fluctuations. Section 3 presents the main results. Section 4 compares our results with those of Beugeling et al. [2] for the same problem. The main text of this paper should be easy to read, for most of the technical details are deferred to Appendix A.

2 Definitions

Throughout this paper, standard asymptotic notations are used extensively. Let $f, g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be two functions. One writes $f(x) = O(g(x))$ if and only if there exist constants $M, x_0 > 0$

*yichuang@mit.edu

such that $f(x) \leq Mg(x)$ for all $x > x_0$; $f(x) = \Omega(g(x))$ if and only if there exist constants $M, x_0 > 0$ such that $f(x) \geq Mg(x)$ for all $x > x_0$; $f(x) = \Theta(g(x))$ if and only if there exist constants $M_1, M_2, x_0 > 0$ such that $M_1g(x) \leq f(x) \leq M_2g(x)$ for all $x > x_0$.

Consider a system of N spins on a hypercubic lattice in $D = \Theta(1)$ spatial dimensions, where each lattice site has a spin. The dimension of the Hilbert space is $d = d_{\text{loc}}^N$, where $d_{\text{loc}} = \Theta(1)$ is the local dimension of each spin. The system is governed by a (not necessarily translationally invariant) local Hamiltonian $H = \sum_i H_i$. The sum is over $\Theta(N)$ lattice sites. Each term H_i has operator norm $\|H_i\| = \Theta(1)$ and is supported in a small neighborhood of site i . Assume without loss of generality that $\text{tr } H_i = 0$ (traceless) so that the mean energy of H is $\text{tr } H/d = 0$.

Let $\{|j\rangle\}_{j=1}^d$ be a complete set of eigenstates of H with corresponding energies $\{E_j\}$. Let $J := \{j : -N\delta_1 \leq E_j \leq N\delta_2\}$ be a microcanonical ensemble in the middle of the energy spectrum, where $\delta_1, \delta_2 = \Theta(1)$ are arbitrary positive constants. Let A be a traceless local operator with $\|A\| = 1$, and $A_{jj} := \langle j|A|j\rangle$ be the EEV so that $\sum_{j=1}^d A_{jj} = \text{tr } A = 0$.

In this paper, we consider three definitions of EEV fluctuations. The first

$$\Delta A := \frac{1}{d} \sum_{j=1}^d |A_{jj}|^2 \quad (1)$$

is simply the variance of A_{jj} in all eigenstates. The second

$$\Delta A_J := \frac{1}{|J|} \sum_{j \in J} \left| A_{jj} - \frac{1}{|J|} \sum_{k \in J} A_{kk} \right|^2 = \frac{1}{|J|} \sum_{j \in J} |A_{jj}|^2 - \frac{1}{|J|^2} \left| \sum_{j \in J} A_{jj} \right|^2 \quad (2)$$

is the variance of A_{jj} in J . The third definition [2] is slightly more complicated. Let $K_j = \{k : |E_j - E_k| \leq N\delta_3\}$, where $\delta_3 = \Theta(1)$ is an arbitrary positive constant. Let

$$\Delta' A_J := \frac{1}{|J|} \sum_{j \in J} \left| A_{jj} - \frac{1}{|K_j|} \sum_{k \in K_j} A_{kk} \right|^2. \quad (3)$$

3 Results

In the thermodynamic limit $N \rightarrow +\infty$, the fluctuations $\Delta A_J, \Delta' A_J$ depend weakly on the hyperparameters $\delta_1, \delta_2, \delta_3$, and are approximately equal to ΔA up to exponentially small additive errors.

Lemma 1. *For any traceless local operator A with $\|A\| = 1$,*

$$|\Delta A_J - \Delta A| = e^{-\Omega(N)}, \quad (4)$$

$$|\Delta' A_J - \Delta A| = e^{-\Omega(N)}. \quad (5)$$

Proof. See Appendix A.1. □

It suffices to assume the ETH for eigenstates in the middle of the energy spectrum.

Assumption 1 (eigenstate thermalization hypothesis in the middle of the spectrum [9, 8]). Let ϵ be an arbitrarily small positive constant. For any traceless local operator A with $\|A\| = 1$, there is a function $f_A : [-\epsilon, \epsilon] \rightarrow \{z \in \mathbb{C} : |z| \leq 1\}$ such that

$$|A_{jj} - f_A(E_j/N)| \leq 1/\text{poly}(N) \quad (6)$$

for all j with $|E_j| \leq N\epsilon$, where $\text{poly}(N)$ denotes a polynomial of sufficiently high degree in N . We assume that $f_A(x)$ is smooth in the sense of having a Taylor expansion to some low order around $x = 0$.

It was proposed analytically [15] and supported by numerical simulations [11] that the right-hand side of (6) can be improved to $e^{-\Omega(N)}$. For our purposes, however, a (much weaker) inverse polynomial upper bound suffices.

Lemma 2. *For a traceless local operator A with $\|A\| = 1$, Assumption 1 implies that*

$$\Delta A = \frac{|\text{tr}(HA)|^2}{d \text{tr}(H^2)} + O(1/N^2). \quad (7)$$

For a generic A , $\text{tr}(HA)$ is nonzero, and hence the first term on the right-hand side of Eq. (7) is $\Theta(1/N)$ (see Eq. (10)).

Lemma 2 appears in Refs. [9, 12], neither of which has rigorously bounded the approximation error in the derivation of Eq. (7). In Appendix A.2, we follow Ref. [9] and present a complete proof of Lemma 2 with rigorous error analysis.

Combining Lemmas 1 and 2, we obtain

Theorem 1. *For a traceless local operator A with $\|A\| = 1$, Assumption 1 implies that*

$$\Delta A_J = \frac{|\text{tr}(HA)|^2}{d \text{tr}(H^2)} + O(1/N^2), \quad \Delta' A_J = \frac{|\text{tr}(HA)|^2}{d \text{tr}(H^2)} + O(1/N^2). \quad (8)$$

4 Discussion

The finite-size scaling of the EEV fluctuation $\Delta' A_J$ was studied by Beugeling et al. [2]. For $\delta_3 = 0.025$ and J being the middle 20% of the spectral range, they presented numerical evidence that $\Delta' A_J$ scales as $d^{-1/2} = e^{-\Theta(N)}$ in generic non-integrable systems, where d is the dimension of the Hilbert space. Furthermore, a heuristic ‘‘typicality argument’’ was provided to explain the $d^{-1/2}$ behavior. This result is different from ours. Recall that Eq. (8) states that for a generic A , $\Delta' A_J$ scales as $\Theta(1/N)$.

In fact, the typicality argument has already been proved to be problematic in a different but related setting [7]. Using the equivalence of ΔA_J and $\Delta' A_J$ (Lemma 1), the proof can be extended to the quantity $\Delta' A_J$. This requires some amount of work, and we will not present the extension here.

For a constant δ_3 and J being the middle 10% of the spectral range, Sugimoto et al. [16] found numerically that

$$\Delta'' A_J = e^{-\Omega(N)}, \quad \Delta'' A_J := \max_{j \in J} \left| A_{jj} - \frac{1}{|K_j|} \sum_{k \in K_j} A_{kk} \right|^2 \quad (9)$$

in generic translationally invariant and generic disordered spin chains. Since $\Delta'' A_J \geq \Delta' A_J$, Eq. (9) is inconsistent with Eq. (8).

We suspect that the discrepancy between Theorem 1 and the numerical results of Refs. [2, 16] is due to finite-size effects in the simulations.

Acknowledgments

We would like to thank Fernando G.S.L. Brandão, Xie Chen, and Yong-Liang Zhang for collaboration on a related project [9]. This work was supported by NSF grant PHY-1818914.

A Proofs

Lemma 3 (moments [9]). *For any integer $m \geq 0$,*

$$\frac{1}{d} \sum_j E_j^{2m} = \frac{1}{d} \text{tr}(H^{2m}) = \Theta(N^m), \quad (10)$$

$$\frac{1}{d} \left| \sum_j E_j^3 \right| = \frac{1}{d} |\text{tr}(H^3)| = O(N). \quad (11)$$

Proof. Expanding H in the generalized Pauli basis, we count the number of terms that do not vanish upon taking the trace in the expansion of H^{2m} . There are $\Theta(N^m)$ such terms, the trace of each of which is $\Theta(d)$. Therefore, we obtain Eq. (10). Equation (11) can be proved in the same way. \square

Almost all eigenstates have vanishing energy density:

Lemma 4 (concentration of eigenvalues [1]). *For any $\epsilon > 0$,*

$$|\{j : |E_j| \geq N\epsilon\}|/d = e^{-\Omega(N\epsilon^2)}. \quad (12)$$

This lemma allows us to upper bound the total contribution of all eigenstates away from the middle of the spectrum. Let $C = O(1)$ be a sufficiently large constant such that

$$\frac{1}{d} \sum_{j:|E_j| \geq \Lambda} 1 \leq \frac{1}{d} \sum_{j:|E_j| \geq \Lambda} |E_j| \leq q, \quad \Lambda := C\sqrt{N \log N}, \quad q := 1/\text{poly}(N), \quad (13)$$

where $\text{poly}(N)$ denotes a polynomial of sufficiently high degree in N .

Lemma 4 and Eqs. (10), (13) are related to the fact that E_j 's approach a normal distribution in the thermodynamic limit $N \rightarrow +\infty$ [10, 3]. Indeed, $|E_j| = \Theta(\sqrt{N})$ for almost all j .

A.1 Proof of Lemma 1

Proof of Eq. (4). By definition,

$$\begin{aligned}\Delta A_J - \Delta A &= \frac{1}{|J|} \sum_{j \in J} |A_{jj}|^2 - \frac{1}{|J|^2} \left| \sum_{j \in J} A_{jj} \right|^2 - \frac{1}{d} \sum_j |A_{jj}|^2 \\ &= \left(\frac{1}{|J|} - \frac{1}{d} \right) \sum_{j \in J} |A_{jj}|^2 - \frac{1}{|J|^2} \left| \sum_{j \notin J} A_{jj} \right|^2 - \frac{1}{d} \sum_{j \notin J} |A_{jj}|^2.\end{aligned}\quad (14)$$

Since $\delta_1, \delta_2 = \Theta(1)$ are positive constants, Lemma 4 implies that $1 - |J|/d = e^{-\Omega(N)}$. Therefore,

$$|\Delta A_J - \Delta A| \leq (1/|J| - 1/d)|J| + (d - |J|)^2/|J|^2 + (d - |J|)/d = e^{-\Omega(N)}.\quad (15)$$

□

Proof of Eq. (5). Let $\delta' := \min\{\delta_1, \delta_2, \delta_3/2\} = \Theta(1)$ and $J' := \{j : |E_j| \leq N\delta'\} \subseteq J$. By definition,

$$\begin{aligned}\Delta' A_J - \Delta A &= \frac{1}{|J|} \sum_{j \in J} \left| A_{jj} - \frac{1}{|K_j|} \sum_{k \in K_j} A_{kk} \right|^2 - \frac{1}{d} \sum_j |A_{jj}|^2 \\ &= \frac{1}{|J|} \sum_{j \in J'} \left(\frac{A_{jj}^*}{|K_j|} \sum_{k \notin K_j} A_{kk} + \frac{A_{jj}}{|K_j|} \sum_{k \notin K_j} A_{kk}^* + \frac{1}{|K_j|^2} \left| \sum_{k \notin K_j} A_{kk} \right|^2 \right) \\ &\quad + \left(\frac{1}{|J|} - \frac{1}{d} \right) \sum_{j \in J'} |A_{jj}|^2 + \frac{1}{|J|} \sum_{j \in J \setminus J'} \left| A_{jj} - \frac{1}{|K_j|} \sum_{k \in K_j} A_{kk} \right|^2 - \frac{1}{d} \sum_{j \notin J'} |A_{jj}|^2.\end{aligned}\quad (16)$$

For any $j \in J'$,

$$K_j = \{k : |E_j - E_k| \leq N\delta_3\} \supseteq \{k : |E_k| \leq N\delta_3/2\}.\quad (17)$$

For such j , Lemma 4 implies that $1 - |K_j|/d = e^{-\Omega(N)}$. Therefore,

$$|\Delta' A_J - \Delta A| \leq \frac{3}{|J|} \sum_{j \in J'} \frac{d - |K_j|}{|K_j|} + \left(\frac{1}{|J|} - \frac{1}{d} \right) |J'| + \frac{4(|J| - |J'|)}{|J|} + \frac{d - |J'|}{d} = e^{-\Omega(N)}.\quad (18)$$

□

A.2 Proof of Lemma 2

Lemma 5 ([8]). *For a traceless local operator A with $\|A\| = 1$, Assumption 1 implies that*

$$|f_A(0)| = O(1/N),\quad (19)$$

$$|f'_A(0) - N \operatorname{tr}(HA) / \operatorname{tr}(H^2)| = O(1/N).\quad (20)$$

We include the proof of this lemma for completeness. For notational simplicity, let $x \stackrel{\delta}{=} y$ denote $|x - y| \leq \delta$.

Proof of Eq. (19).

$$\begin{aligned} 0 &= \frac{1}{d} \operatorname{tr} A = \frac{1}{d} \sum_j A_{jj} \stackrel{O(q)}{=} \frac{1}{d} \sum_{j:|E_j|<\Lambda} A_{jj} \stackrel{1/\text{poly}(N)}{=} \frac{1}{d} \sum_{j:|E_j|<\Lambda} f_A(E_j/N) \\ &\approx \frac{1}{d} \sum_{j:|E_j|<\Lambda} \left(f_A(0) + \frac{f'_A(0)E_j}{N} \right) \stackrel{O(q)}{=} \frac{1}{d} \sum_j \left(f_A(0) + \frac{f'_A(0)E_j}{N} \right) = f_A(0), \end{aligned} \quad (21)$$

where we used the inequality (13), the ETH (6), and the Taylor expansion

$$f_A(E_j/N) = f_A(0) + f'_A(0)E_j/N + f''_A(0)E_j^2/(2N^2) + O(|E_j|^3/N^3) \quad (22)$$

in the steps marked with “ $O(q)$,” “ $1/\text{poly}(N)$,” and “ \approx ,” respectively. The approximation error in the “ \approx ” step is

$$\frac{1}{d} \sum_{j:|E_j|<\Lambda} O(E_j^2/N^2) \leq \frac{1}{d} \sum_j O(E_j^2/N^2) = O(1/N), \quad (23)$$

where we used Eq. (10) with $m = 1$. We obtain Eq. (19) by combining (21), (23). \square

Proof of Eq. (20).

$$\begin{aligned} \frac{1}{d} \operatorname{tr}(HA) &= \frac{1}{d} \sum_j E_j A_{jj} \stackrel{O(q)}{=} \frac{1}{d} \sum_{j:|E_j|<\Lambda} E_j A_{jj} \stackrel{1/\text{poly}(N)}{=} \frac{1}{d} \sum_{j:|E_j|<\Lambda} E_j f_A(E_j/N) \\ &\approx \frac{1}{d} \sum_{j:|E_j|<\Lambda} \left(E_j f_A(0) + \frac{f'_A(0)E_j^2}{N} + \frac{f''_A(0)E_j^3}{2N^2} \right) \\ &\stackrel{O(q)}{=} \frac{1}{d} \sum_j \left(E_j f_A(0) + \frac{f'_A(0)E_j^2}{N} + \frac{f''_A(0)E_j^3}{2N^2} \right) \stackrel{O(1/N)}{=} \frac{f'_A(0) \operatorname{tr}(H^2)}{Nd}, \end{aligned} \quad (24)$$

where we used (13), (6), the Taylor expansion (22), and Eq. (11) in the steps marked with “ $O(q)$,” “ $1/\text{poly}(N)$,” “ \approx ,” and “ $O(1/N)$,” respectively. The approximation error in the “ \approx ” step is

$$\frac{1}{d} \sum_{j:|E_j|<\Lambda} O(E_j^4/N^3) \leq \frac{1}{d} \sum_j O(E_j^4/N^3) = O(1/N), \quad (25)$$

where we used Eq. (10) with $m = 2$. We obtain Eq. (20) by combining (24), (25). \square

We are ready to prove Lemma 2:

$$\begin{aligned} \Delta_A &= \frac{1}{d} \sum_j |A_{jj}|^2 \stackrel{O(q)}{=} \frac{1}{d} \sum_{j:|E_j|<\Lambda} |A_{jj}|^2 \stackrel{1/\text{poly}(N)}{=} \frac{1}{d} \sum_{j:|E_j|<\Lambda} |f_A(E_j/N)|^2 \\ &\approx \frac{1}{d} \sum_{j:|E_j|<\Lambda} \left| f_A(0) + \frac{f'_A(0)E_j}{N} \right|^2 \stackrel{O(q)}{=} \frac{1}{d} \sum_j \left| f_A(0) + \frac{f'_A(0)E_j}{N} \right|^2 \\ &= |f_A(0)|^2 + \frac{|f'_A(0)|^2}{d} \sum_j \frac{E_j^2}{N^2} = |f_A(0)|^2 + \frac{|f'_A(0)|^2 \operatorname{tr}(H^2)}{N^2 d} \stackrel{O(1/N^2)}{=} \frac{|\operatorname{tr}(HA)|^2}{d \operatorname{tr}(H^2)}, \end{aligned} \quad (26)$$

where we used (13), (6), and Lemma 5 in the steps marked with “ $O(q)$,” “ $1/\text{poly}(N)$,” and “ $O(1/N^2)$,” respectively. In the “ \approx ” step, we used (22) with the approximation error upper bounded by

$$\begin{aligned} O(1/d) \left| \sum_{j:|E_j|<\Lambda} \frac{E_j^3}{N^3} \right| + O(1/d) \sum_{j:|E_j|<\Lambda} \left(\frac{|f_A(0)|E_j^2}{N^2} + \frac{E_j^4}{N^4} \right) \\ \leq O(1/d) \left| \sum_j \frac{E_j^3}{N^3} \right| + O(q) + O(1/d) \sum_j \left(\frac{E_j^2}{N^3} + \frac{E_j^4}{N^4} \right) = O(1/N^2), \quad (27) \end{aligned}$$

where we used Lemma 3. We complete the proof of Lemma 2 by combining (26), (27).

References

- [1] A. Anshu. Concentration bounds for quantum states with finite correlation length on quantum spin lattice systems. *New Journal of Physics*, 18(8):083011, 2016.
- [2] W. Beugeling, R. Moessner, and M. Haque. Finite-size scaling of eigenstate thermalization. *Physical Review E*, 89(4):042112, 2014.
- [3] F. G. S. L. Brandao and M. Cramer. Equivalence of statistical mechanical ensembles for non-critical quantum systems. arXiv:1502.03263, 2015.
- [4] L. D’Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol. From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics. *Advances in Physics*, 65(3):239–362, 2016.
- [5] J. M. Deutsch. Quantum statistical mechanics in a closed system. *Physical Review A*, 43(4):2046–2049, 1991.
- [6] J. M. Deutsch. Eigenstate thermalization hypothesis. *Reports on Progress in Physics*, 81(8):082001, 2018.
- [7] R. Hamazaki and M. Ueda. Atypicality of most few-body observables. *Physical Review Letters*, 120(8):080603, 2018.
- [8] Y. Huang. Convergence of eigenstate expectation values with system size. arXiv:2009.05095, 2020.
- [9] Y. Huang, F. G. S. L. Brandão, and Y.-L. Zhang. Finite-size scaling of out-of-time-ordered correlators at late times. *Physical Review Letters*, 123(1):010601, 2019.
- [10] J. P. Keating, N. Linden, and H. J. Wells. Spectra and eigenstates of spin chain Hamiltonians. *Communications in Mathematical Physics*, 338(1):81–102, 2015.
- [11] H. Kim, T. N. Ikeda, and D. A. Huse. Testing whether all eigenstates obey the eigenstate thermalization hypothesis. *Physical Review E*, 90(5):052105, 2014.

- [12] M. Mierzejewski and L. Vidmar. Quantitative impact of integrals of motion on the eigenstate thermalization hypothesis. *Physical Review Letters*, 124(4):040603, 2020.
- [13] M. Rigol, V. Dunjko, and M. Olshanii. Thermalization and its mechanism for generic isolated quantum systems. *Nature*, 452(7189):854–858, 2008.
- [14] M. Srednicki. Chaos and quantum thermalization. *Physical Review E*, 50(2):888–901, 1994.
- [15] M. Srednicki. The approach to thermal equilibrium in quantized chaotic systems. *Journal of Physics A: Mathematical and General*, 32(7):1163–1175, 1999.
- [16] S. Sugimoto, R. Hamazaki, and M. Ueda. Test of the eigenstate thermalization hypothesis based on local random matrix theory. *Physical Review Letters*, 126(12):120602, 2021.