

Assessing the effect of sample bias correction in species distribution models

Nicolas Dubos,^{1,*} Clémentine Préau,¹ Maxime Lenormand,^{1,*} Guillaume Papuga,² Sophie Monsarrat,^{3,4}
Pierre Denelle,⁵ Marine Le Louarn,¹ Stien Heremans,⁶ Roel May,⁷ Philip Roche,⁸ and Sandra Luque¹

¹*TETIS, Univ Montpellier, AgroParisTech, Cirad, CNRS, INRAE, Montpellier, France*

²*AMAP, Univ Montpellier, CIRAD, CNRS, INRAE, IRD, Montpellier, France*

³*Center for Biodiversity Dynamics in a Changing World (BIOCHANGE),
Department of Biology, Aarhus University, Ny Munkegade 114, DK-8000 Aarhus C, Denmark*

⁴*Section for Ecoinformatics and Biodiversity, Department of Biology,
Aarhus University, Ny Munkegade 114, DK-8000 Aarhus C, Denmark*

⁵*Biodiversity, Macroecology & Biogeography, University of Goettingen, Göttingen, Germany*

⁶*Research Institute for Nature and Forest (INBO), Brussels, Belgium*

⁷*Norwegian institute for nature research (NINA),*

P.O. Box 5685 Torgarden, NO-7485 Trondheim, Norway

⁸*RECOVER, INRAE, 3275 Route de Cézanne,*

CS 40061, 13182, Aix en Provence, Cedex 5, France

ABSTRACT

1. Open-source biodiversity databases contain a large amount of species occurrence records, but these are often spatially biased, which affects the reliability of species distribution models based on these records. Sample bias correction techniques include data filtering at the cost of record numbers or require considerable additional sampling effort. However, independent data are rarely available and assessment of the correction technique must rely on performance metrics computed with subsets of the only available (biased) data, which may be misleading.

2. Here we assess the extent to which an acknowledged sample bias correction technique is likely to improve models' ability to predict species distributions in the absence of independent data. We assessed the variation in model predictions induced by the correction and model stochasticity, i.e. the variability between model replicates which is related to a random component (pseudo-absences sets and cross-validation subsets). We present an index of the effect of correction relative to model stochasticity, the Relative Overlap Index (ROI). We tested whether the ROI better represented the effect of correction than classic performance metrics (Boyce index, cAUC, AUC and TSS) and absolute overlap metrics (Shoener's D, Pearson's and Spearman's correlation coefficients) using 70 vertebrate species and 18 virtual species with a generated sample bias.

3. When based on absolute overlaps and cross-validation performance metrics, we found no effect of correction, except for cAUC. When considering its effect relative to model stochasticity, the effect of correction depended on the site and the species. Virtual species enabled us to verify that the correction actually improved distribution predictions and the biological relevance of the selected variables at the sites with a clear gradient of sample bias, and when species distribution predictors are not correlated with sample bias patterns.

4. In absence of additional independent data, the assessment of sample bias correction based on subsample data may be misleading. Along with the investigation of the biological relevance of environmental variables selected and the use of discrimination metrics calibrated with null geographic models (e.g. cAUC), we propose to assess the effect of sample bias correction on the basis of its effect relative to model stochasticity.

Keywords. Accessibility maps, cross-validation, performance metrics, overlap, pseudo-absence selection, terrestrial vertebrates, variable selection, virtual species.

INTRODUCTION

There is a growing demand for species data in a view of producing robust statistical models and evidence-based conservation actions, but the availability of standardised data remain limited. The recent development of extensive biodiversity databases is supported mainly by opportunistic presence-only data from citizen science programs and naturalists

associations. Despite its limitations, opportunistic, non-standardised data still constitute a promising venue for improving biodiversity assessments (McKinley *et al.*, 2017). Such data are often limited by the heterogeneity of their sources and spatial biases due to uneven sampling effort (Beck *et al.*, 2014; Bird *et al.*, 2014; Botella *et al.*, 2021; Johnston *et al.*, 2020; Otegui *et al.*, 2013). Specifically, sampling may be biased by field accessibility such as the number of observations that is influenced by the proximity to urban areas and roads, which often leads to spatial autocorrelation among observations (Phillips *et al.*, 2009; Stolar & Nielsen, 2015). As a result, environ-

* Corresponding authors: nicolas.dubos@inrae.fr & maxime.lenormand@inrae.fr

mental bias may arise and models tend to overestimate/underestimate the environmental suitability in the zone with higher/lower density of occurrence data. This may be problematic in studies aiming to provide guidelines for management (Yackulic *et al.*, 2013). Accounting for spatial bias in opportunistic data from heterogeneous sources (e.g., citizen science, naturalists and experts associations) would increase their potential use in ecological studies and enable the inclusion of a broader range of species in Species Distribution Models (SDMs).

SDM is one of the most commonly used tool to test ecological hypotheses (Anderson *et al.*, 2009), assess alien species invasion risks (Bellard *et al.*, 2013; Briscoe Runquist *et al.*, 2019), forecast the potential effect of environmental change (Araújo *et al.*, 2005), and support conservation management (Leroy *et al.*, 2014; Mikolajczak *et al.*, 2015; Schwartz, 2012). While presence-only biodiversity databases are increasingly used in SDMs, there is still a persistent blur around central methodological questions such as sample bias corrections (Johnston *et al.*, 2020; Meynard *et al.*, 2019).

Spatial sampling bias is a major factor affecting the predictive performance of SDMs (Araújo & Guisan, 2006; Barbet–Massin *et al.*, 2012; Kramer–Schadt *et al.*, 2013; Meynard *et al.*, 2019). A number of procedures have been developed in order to account for sampling bias, including spatial filtering of presence points (Boria *et al.*, 2014; Edrén *et al.*, 2010), environmental filtering (Gábor *et al.*, 2020; Varela *et al.*, 2014) and production of a similar sampling bias in non-presence background data/pseudo-absences (Phillips *et al.*, 2009). However, presence points and environmental filtering consist in removing occurrence data, inducing a loss of information and statistical power, which can be critical when dealing with rare, or poorly detected species (Kramer–Schadt *et al.*, 2013; Lobo & Tognelli, 2011; Robinson *et al.*, 2018; Vollering *et al.*, 2019). In presence/background or presence/pseudo-absence models, a range of pseudo-absence selection techniques were recently developed (e.g. Fourcade *et al.* (2014); Hertzog *et al.* (2014); Iturbide *et al.* (2015); Senay *et al.* (2013)), enabling to reduce the effect of sampling bias and improve model performance without removing any occurrence points. For instance, pseudo-absence selection based on sampling bias reference maps has been acknowledged as an efficient method to account for spatially biased occurrence data (Hertzog *et al.*, 2014; Phillips *et al.*, 2009). The use of reference maps such as the target-group (TG) approach or accessibility maps can be used to represent a sampling bias map specific to a given study area and offer a promising approach to improve the predictive performance of SDMs (Monsarrat *et al.*, 2019; Ranc *et al.*, 2017). The TG approach relies on the hypothesis that the study species share the same sampling pattern as the target group. Accessibility maps rely on the hypothesis that constraining features are identified (e.g. geographical barriers, social conflicts, long distances). These do not provide

explicit information on sampling efforts but might be more appropriate when species richness patterns are heterogeneous (Ranc *et al.*, 2017) and when treating data from heterogeneous sources with different sampling patterns (Monsarrat *et al.*, 2019). The ability of accessibility maps to predict sampling bias was shown for historical data, and remains to be tested on modern data (see Sicacha–Parada *et al.* (2020)). Besides, the effectiveness of correction techniques can vary depending on the species and the environmental context (e.g. Gábor *et al.* (2020); Varela *et al.* (2014)), and pseudo-absence selection techniques were mostly tested on specific case-studies (one or two species, or one taxonomic group). This prevents from drawing general conclusions and calls the need for their assessment at broader taxonomic scales in a variety of locations.

The efficiency of a given sample bias correction technique is often measured on the basis of comparisons of performance metrics between corrected and uncorrected models. In both corrected and uncorrected groups, performance metrics are based on a comparison of the ability of models built with a subset of the original data (i.e. training/calibration dataset) to accurately predict the remaining data (test/evaluation dataset), a process often called “cross-validation” (e.g. Boria *et al.* (2014); Senay *et al.* (2013)). The most common model performance metrics such as the Area under the operating curve (AUC), the True Skill Statistic (TSS), the Boyce index, and Similarity indices provide quantitative measures of discrimination ability between models built with training data and the full dataset (Fourcade *et al.*, 2018). However, a large majority of studies test the efficiency of sample bias correction by performing internal cross-validation (see point 3B of the Standard for SDMs on data sharing the same bias in Araújo *et al.* (2019)), an approach that has shown strong limitations when the purpose of the SDM is to extrapolate predictions to a different time/region (Araújo *et al.*, 2019 2005; Beck *et al.*, 2014; Fourcade *et al.*, 2018; Hertzog *et al.*, 2014). Ideally, the improvement conferred by a correction technique should be evaluated with an independent, unbiased dataset (Hertzog *et al.*, 2014; Johnston *et al.*, 2020; Norberg *et al.*, 2019; Phillips *et al.*, 2009). Field validation or evaluations relative to independent unbiased datasets represents the best standard practice to assess models’ ability to predict species distributions (Araújo *et al.*, 2019). Nevertheless, field validation is highly demanding in terms of prospecting effort and sometimes unfeasible (e.g. taxonomically or geographically extensive scale study sites), and independent standardised datasets are rarely available (Hao *et al.*, 2019). A possible method to assess sample bias correction techniques, when relying on a partitioned dataset, is to select data subsets that are subject to different types of bias (e.g. Bean *et al.* (2012)). However, this method may also be highly demanding in terms of spatio-temporal coverage and may not be feasible for most species (Johnston *et al.*, 2020). A cost-effective method has been proposed by Hijmans




Figure 1. Location of the three study sites in Europe and accessibility maps for each site. The accessibility index is inversely proportional to the Euclidean distance between cities and roads and represents the degree to which a geographic point is easily reached by an observer. Black dots represent occurrence records of all species pooled together ($n = 54$, 11 and 8 species in Trondheim, Grote Nete and Thau, respectively).

(2012) to assess the potential efficiency of sample bias correction techniques with an AUC calibrated with a null geographic model. The efficiency of this method may vary through space and between species (Hijmans, 2012), calling for the need to characterise those sites and species. Virtual species can be used to assess bias correction techniques (e.g. Fourcade *et al.* (2014); Phillips *et al.* (2009); Ranc *et al.* (2017); Varela *et al.* (2014)), by simulating a sampling bias and producing performance metrics that are relative to a perfectly known distribution. The projection of a range of virtual species on multiple real regions may represent a cost-effective approach to test whether a correction technique is likely to actually improve the accuracy of SDMs, provided virtual and real species are comparable.

Species distribution models can be calibrated with a range of model parameters generated with a random component (e.g. pseudo absence selection, cross-validation subsets) inducing a stochasticity among models that are otherwise identical. These model parameters can be sources of uncertainty in model projections (Buisson *et al.*, 2010; Thibaud *et al.*, 2014). Sample bias correction should induce variation in the

predicted values (and subsequently species range projections), but may be negligible if the variation is of the same magnitude as the sources of uncertainty. Therefore, the effect of sample bias correction may be assessed on the basis of its effect between corrected and uncorrected modalities relative to intra-modality variation.

We hypothesise that a sample bias correction technique improves model predictions if its relative effect is stronger than that of the remaining model input parameters. We tested this assumption using virtual species with a generated sample bias, modelled at the same sites and with the same range of model parameters as the real species. This study aims (1) at assessing the effect of a sampling bias correction technique on distribution projections in absence of independent data over a range of terrestrial vertebrate species ($n = 70$) in three contrasted regions. We measured the effect of correction by computing the degree of overlap between uncorrected and corrected models. We predicted that the effect of correction on projections differs between sites and species. We further evaluated (2) whether the effect of correction can be assessed with a range of validation metrics. We also

tested whether this effect can be represented by an index of overlap between correction modalities relative to intra-modality variation. We tested (3) whether sample bias correction actually improved model predictions using virtual species with a simulated sample bias. Finally, we provide recommendations for the assessment of sample bias correction when independent data are unavailable.

MATERIALS AND METHODS

Study sites and accessibility

We focused on three regions, in the surroundings of Thau (southern France), Grote Nete (northern Belgium) and Trondheim (central Norway). The sites are characterised by contrasted distributions of roads and cities, with a homogeneous distribution overall in Thau and in Grote Nete while there was a strong gradient in road and town density in Trondheim. At each site, we produced accessibility maps (Figure 1) by computing an Accessibility Index (AI). Accessibility Indices represent the degree to which a geographic point is easily reached by an observer and is context specific (Monsarrat *et al.*, 2019). Here we use occurrence data from heterogeneous sources, hence we assumed that accessibility depends mostly on distance from cities and from roads (e.g. Sicacha-Parada *et al.* (2020)). The Accessibility Index was computed as follows:

$$AI_i = \frac{1}{2} \left(e^{-\frac{1}{2} \left(\frac{dist_C}{\sigma_C} \right)^2} + e^{-\frac{1}{2} \left(\frac{dist_R}{\sigma_R} \right)^2} \right) \quad (1)$$

where AI_i is the accessibility index at pixel i , $dist_C$ is the Euclidean distance from the closest city centre (≥ 200 inhabitants), $dist_R$ is the Euclidean distance from the closest primary and secondary road. σ_C and σ_R are the standard deviations of the distances distributions to the nearest city and road, respectively.

Environmental data

We used land use variables retrieved from Corine Land Cover habitat classes, a European biophysical dataset derived from remote sensing. Land use variables are more relevant than climatic predictors in species distribution models at the local scale (Ficetola *et al.*, 2014; Soberón & Nakamura, 2009). For each $200 \times 200 \text{ m}^2$ pixel, we measured the distance from nearest habitat features using 8 habitat classes (artificial surface, forest edge, intensive farmland, non-intensive farmland, scrubland/herbaceous areas, coastal areas, water courses, water bodies). We also computed the proportion of a given habitat type within a range of buffer zones around occurrence points (200m, 500m and 1000m) corresponding to

species habitat use at the landscape scale according to previous studies (e.g. in reptiles, amphibians and bats; Azam *et al.* (2016); Jeliaskov *et al.* (2014)). In birds, the landscape may be influential at larger scales (e.g. 5000m; Dubos *et al.* (2018)). However, given the scale of our study sites, the use of larger buffer zones would cause a lack of variability in environmental conditions at occurrence points, so we limited the extent of our buffer zones to 1000m.




Figure 2. Boxplots of measures of sample bias in the spatial distribution of the 70 real species according to the case study site. The bias is assessed with Boyce indices measuring how occurrence data are accurately predicted by the Accessibility Index. Each boxplot is composed of the first decile, the lower hinge, the median, the upper hinge and the ninth decile.

Occurrence records

We used occurrence records obtained from biodiversity databases (*Ligue pour la Protection des Oiseaux*¹, the Global Biodiversity Information Facility² and Natuurpunt Studie Association³) at the three sites for 79 terrestrial vertebrate species (58 birds, 10 mammals, 6 amphibians and 5 reptiles). More details in Table S1 in Appendix. Species with fewer than 15 occurrence points per site were discarded ($n = 9$). We selected one occurrence point per $200 \times 200 \text{ m}^2$ pixel (i.e. the resolution of our environmental variables). This process is used as a rule of thumb to limit sampling bias driven by multiple observations within the pixel, but does not account for sampling bias driven

¹ <https://www.lpo.fr>, last accessed 17/02/2021

² <https://www.gbif.org>, last accessed 17/02/2021

³ <https://www.natuurpunt.be/afdelingen/natuurpunt-studie>, last accessed 17/02/2021




Figure 3. Steps of the methodology developed to study the effect of sample bias correction applied on the species distribution modelling of 70 real and 18 virtual study species.

by aggregated observations in the surrounding pixels and at larger scales. The level of sample bias can be estimated using the Boyce index, usually used as a evaluation metric for presence-only data to assess the extent to which a spatial layer correctly predicts presence points. In our case the spatial layer is the accessibility map (Figure 1). Overall, as can be observed in Figure 2, the spatial bias due to accessibility in occurrence data was negative in Grote Nete (Boyce index = -0.936), slightly positive in Thau (Boyce index = 0.733) and strongly positive in Trondheim (Boyce index = 0.948).

Distribution modelling

We built species distribution models using the Biomod2 R package (Thuiller *et al.*, 2009), and an ensemble of eight modelling techniques: generalised linear models (GLM), generalised boosting model (GBM), classification tree analysis (CTA), artificial neural network (ANN), surface range envelop (SRE), flexible discriminant analysis (FDA), general additive model (GAM) and random forest (RF). The modelling procedure included (1) a method for pseudo-absence selection, (2) an environmental variable selection process, (3) a final model calibration and (4) a model evaluation process, and is

summarised in Figure 3.

Pseudo-absence selection. For each group (uncorrected and corrected groups as defined below), we included 10 different sets of generated pseudo-absences with equal number to that of presence points (Meynard *et al.*, 2019). For models which did not account for field accessibility (and subsequent sampling bias), hereafter referred as the “uncorrected group”, we randomly and evenly selected a number of pseudo-absences in the study area, equal to the number of occurrence data within the background (Barbet-Massin *et al.*, 2012; Liu *et al.*, 2019). For models which accounted for sampling bias, we randomly selected pseudo-absences with a sampling probability weighted by AIs after excluding presence pixels. This enabled pseudo-absences to share the same bias than presence points, following the original concept proposed in Phillips *et al.* (2009). Thus, for species that were negatively biased by accessibility (i.e. more found in inaccessible areas), we weighted sampling probability by the negative AIs. Treating each site separately, we selected one variable per group of inter-correlated variables to avoid collinearity (Pearson’s $r \geq 0.7$). For each species individually (Hawkins *et al.*, 2017), we assessed the relative importance of each variable kept (calculated as the Pearson’s coefficient between initial model

predictions and model predictions made when the assessed variable is randomly permuted) with 10 permutations. The final set of variables included in the final models were those with a relative importance ≥ 0.05 for across at least 50% of model runs (Bellard *et al.*, 2016).

Final models. We used the eight aforementioned modelling techniques, with 10 sets of pseudo absence, and three runs of calibration with 70% of the data (30% for evaluation).

Effect of sample bias correction

Effect on model predictions. We measured the “absolute” effect of sample bias correction using indices of similarity and correlation coefficients between uncorrected and corrected predictions. We computed the Shoener’s D as a measure of projection overlap (computed with the ENMTool R package (Rödger & Engler, 2011; Warren *et al.*, 2010)), the Pearson’s correlation coefficient (Li & Guo, 2013) and the Spearman’s rank coefficient (Phillips *et al.*, 2009). Shoener’s D was computed as follows:

$$D(p_x, p_y) = 1 - \frac{1}{2} \sum_i |p_{x_i} - p_{y_i}| \quad (2)$$

where p_{x_i} and p_{y_i} are the normalized suitability scores for uncorrected x and corrected y prediction in grid cell i , for each species, modelling technique, cross-validation run and pseudo-absence run individually.

For comparison, we also assessed model performance using four classical evaluation metrics based on data subsets, i.e. the Boyce index (computed with the ecospat R package (Cola *et al.*, 2017), the true skill statistic (TSS), the area under the relative operating characteristic curve (AUC), and a calibrated AUC (cAUC). The cAUC was computed following Hijmans (2012) as the AUC calibrated on the AUC of a null geographic model (the null geographic model was computed with the geoDist function of the dismo R package (Hijmans *et al.*, 2015)). The Boyce index is a reliability metric, indicating the extent to which a spatial layer correctly predicts presence points. The TSS, AUC and cAUC are discrimination metrics, indicating the ability to distinguish between occupied and unoccupied sites. For a given performance metric X , we define the δ performance as follows:

$$\delta = \frac{X_{corrected} - X_{uncorrected}}{X_{uncorrected}} \quad (3)$$

where $X_{corrected}$ and $X_{uncorrected}$ stand for the performance metric obtained with the corrected and uncorrected prediction, respectively.

Effect relative to model stochasticity. For each species and modelling technique, we assessed the extent to which the correction technique affected predictions relative to the sources of stochastic variation between

models of the corrected group (i.e. cross-validation runs, pseudo-absence set runs, for each modelling technique). Model stochasticity was quantified using the aforementioned overlap metrics (i.e. Shoener’s D, Pearson’s and Spearman’s coefficient) between all pairwise combinations of model projections (i.e. 10 pseudo-absence datasets and 3 cross validation subsets, individually for the 70 species and the 8 modelling techniques, resulting in $n = 252\,000$ measures for each metric). We present the Relative Overlap Index (ROI), an index of mean overlap between predictions of the uncorrected and the corrected groups relative to the average overlap between pairwise model projections of the corrected group. The two overlap components of the ROI can be assessed either with similarity metrics or correlation coefficients. When based on Schoener’s D, the ROI was computed as follows:

$$ROI = \frac{\bar{D}_0 - \bar{D}(p_x, p_y)}{\bar{D}_0} \quad (4)$$

where \bar{D}_0 is the mean overlap between model runs of the corrected group and $\bar{D}(p_x, p_y)$ is the mean overlap between runs of the uncorrected and the corrected models. A value close to 0 represents a perfect match between predictions (i.e. no effect of sample bias correction). The overlaps between uncorrected and corrected groups tend to be significantly smaller than the overlaps between runs when the ROI approaches 1 (i.e. effect of sample bias correction). The formula was similar when based on Pearson’s and Spearman’s rank coefficient, but values were transformed in order to range between 0 and 1 (by adding 1 and dividing by 2).

Effect on variable selection. We estimated the degree of similarity in the selected variables between the uncorrected and the corrected group. We used the Jaccard index (Jaccard, 1912) when only considering whether the variable is selected or not. We also computed the Bray-Curtis index (Bray & Curtis, 1957) when accounting for variable importance.

Factors influencing the effect of correction. We investigated whether the effect of correction differed between sites and modelling techniques, and depended on the level of sample bias (inferred from Boyce indices computed with species occurrences and accessibility maps) and sample size (number of occurrence points). The latter two factors were slightly correlated (Pearson’s $r = 0.38$).

Testing whether changes correspond to actual improvements using virtual species

For each of the three study sites individually, we generated a set of 6 virtual species with different ecological niches generated at each site individually (18 species in total) using the Virtualspecies R package




Figure 4. Site-specific variation in the effect of sample bias correction. (a) Boxplots of the three absolute overlap measures between corrected and uncorrected predictions. (b) Boxplots of the δ performance between corrected and uncorrected predictions. (c) Boxplots of the variable selection similarity. Each boxplot is composed of the first decile, the lower hinge, the median, the upper hinge and the ninth decile.

(Leroy *et al.*, 2016). Their probability of presence was generated according to a relationship with a single environmental variable (parameters and shape of the relationships are provided in Table 1). We produced presence-absence maps using a probability threshold of 0.6, and sampled 300 presence points, and 300 absence points (except for species which we associated with the grassland index and the proportion of herbaceous areas within 500m, for which the number of pixels represented was lower than 300, respectively 100 and 80 presence-absence points were sampled). For the 18 virtual species, we repeated the sample procedure with a sample probability weighted by the Accessibility Index to simulate a sample bias. The simulated bias was overall well represented by an increase in the Boyce index based on accessibility maps (Figure 2).

Did the correction effect correspond to an improvement? The virtual species generated and their simulated sample bias enabled us to test whether the correction technique actually improved the models' ability to predict the "true" distribution (Meynard & Kaplan, 2012). To quantify the degree to which model predictions were improved by the correction technique, we compared the predicted probability of occurrence of the corrected group with the "true" probability of occurrence using the root-mean-square error (RMSE), the Pearson's correlation coefficient and Shoener's D overlap.

Did the correction improve the biological relevance of variable selection? We further verified whether variable selection was relevant by comparing the variables used to generate the virtual species and the variables selected before and after correction.

RESULTS

Effect of sample bias correction

Effect on model predictions. The effect of correction was rather consistent between sites, as shown by the "absolute" overlap between models built from uncorrected versus corrected pseudo-absences plotted in Figure 4a. Model projections shared about 75% of common information (Shoener's D) between uncorrected and corrected models overall. Maps of the projection are available in Appendix (Figures S1 to S17). There was no difference in Boyce, AUC and TSS indices between groups (Figure 4b). Only the cAUC showed a significant effect of correction in Thau and Trondheim.

Effect on variable selection. We found significant differences in variable selection (Jaccard indices) and variable importance (Bray-Curtis indices) between corrected and uncorrected models at Trondheim (Figure 4c). Variables importance show evident differences in Trondheim.

Relative effect of correction

The effect of correction was high compared to model stochasticity in Trondheim, as shown by the lower overlaps between treatments compared to overlaps between model replicates (Figure 5). At the two remaining sites, the effect of correction was of similar magnitude than that of model replicates. This result was consistent when using Pearson's and Spearman's coefficients as a measure of overlap (Figures S18 and S19 in Appendix).




Figure 5. Site-specific variation in the effect of sample bias correction Relationship between the effect of sample bias correction (Schoener's D overlap between uncorrected and corrected predictions) and the effect of model stochasticity (Schoener's D overlap between model runs). Points located below the $y = x$ red line represent models for which the relative effect of sample bias correction exceeds that of model stochasticity. The inset shows the boxplots of the Relative Overlap Index (ROI) according to the case study site. Each boxplot is composed of the first decile, the lower hinge, the median, the upper hinge and the ninth decile. Similar plots obtained with the Pearson's and Spearman's coefficients are available in Figures S18 and S19 in Appendix.

Factors influencing the relative effect of correction

The Relative Overlap Index (ROI) showed a higher effect of correction for the largest sample sizes and the most positively biased samples overall (Figure 6a and 6b). As can be observed in Figure 6c, the different modelling techniques showed consistent results overall (the coefficient of variation of the ROI between modelling techniques was ≤ 1 for 66% of the species). The ROI was more variable with SRE (as known as BIOCLIM) than the other seven techniques.

Did the correction actually improve predictions?

The correction enabled to remarkably improve distribution predictions in Trondheim and to a lesser extent in Grote Nete, but not in Thau (Figure 7). Maps of the projection are available in Appendix (Figures S20 to S25).

Did the correction improve the biological relevance of the selected variables?

The generated sample bias induced the selection of irrelevant variables for 10 virtual species (i.e. variables which were not used to generate the virtual species; Table 1). For six of those, the correction improved the biological relevance of variable selection (i.e. the irrelevant variables were not selected after correction). In contrast, the correction decreased the relevance of variable selection for one species. This species was the only case showing a poor performance of the correction in Trondheim, and was generated using a variable which was strongly correlated with the accessibility index (i.e. Distance from intensive agriculture; Pearson's $r = -0.80$).

DISCUSSION

The efficiency of sample bias correction based on accessibility varies differently between sites and species. Correction effect depends on the landscape complexity of the site and on the degree of spatial bias in oc-




Figure 6. Variation of the Relative Overlap Index (ROI) across quartiles of sample size and sample bias for 70 species, and across modelling techniques. (a) Boxplots of the ROI according to the sample size (divided into quartiles). (b) Boxplots of the ROI according to the sample bias measured with the Boyce (divided into quartiles). (c) Probability Density Function (PDF) of the coefficient of variations of the ROI between modelling techniques for each species individually. The inset shows the boxplots of the ROI according to the modelling technique. Each boxplot is composed of the first decile, the lower hinge, the median, the upper hinge and the ninth decile.

currence data and in predictor variables. In absence of independent data, the effect of the correction can be assessed using the cAUC and the Relative Overlap Index that we introduced in this study.

Effect of sample bias correction

In this study, the impact of sample bias correction primarily depends on the study site, and to a lower extent on the species. The correction mostly affected the predictions of species distributions in Trondheim, which is consistent with the results obtained from virtual species. Comparisons with the “true” probability of occurrence of virtual species suggested that the correction was likely to improve model predictions at this site. The differences in the impact of the correction between sites may be explained by the spatial scale, and the landscape complexity of the sites. At the two sites with a low impact of correction, characterised by a smaller extent and an accessibility more scattered, model predictions were already acceptable in absence of correction (Figure 5). The accessibility index is built on the basis of site-specific characteristics, and reflects a relative measure of distance. At small spatial scales, the

less accessible zones were geographically closer to the more accessible ones, and the absolute distance may not be problematic in terms of accessibility. This suggests that sampling bias correction based on accessibility is unnecessary at small spatial scales in areas with homogeneous distributions of roads and towns. Accessibility maps may also not reflect sample bias equally for all species and sites. Depending on the species or the site, accessibility maps can be built on the basis of topography, land use and property to better represent species or site-specific sample bias.

Effect of sample bias. The impact of correction on predictions was greater when sample bias was highly positive but remained highly heterogeneous, even among species with strong sample biases. The impact of correction may be limited for species for which the habitat is small, or not represented in inaccessible areas. For instance, two of the most biased species for which the correction did not change predictions (i.e. *Accipiter nisus* and *Actitis hypoleucos* in Figure S6 in Appendix) were dependent on coastal areas and water bodies, respectively. These two habitats were mostly distributed in urbanised areas, and the species were absent from other habitats. This is also true for species that are dependent on habitats




Figure 7. Overlap with the “true” probability of occurrence (virtual species) for the uncorrected and the corrected groups at three sites. An improvement in model performance is indicated by a higher overlap (Shoener’s D). Each boxplot is composed of the first decile, the lower hinge, the median, the upper hinge and the ninth decile. Similar plots obtained with the Pearson’s coefficient and the RMSE are available in Figures S26 in Appendix.

that are negatively correlated with accessibility, as shown by the virtual species distributed away from intensive agriculture (see *Fugiagrís norwegensis* in Figure S25 in Appendix). This calls the need to carefully investigate the selected variables and their distribution relative to factors of accessibility.

Effect of sample size. Correction affected the most species with the largest sample sizes, presumably because the model stochasticity may be lower as a result of higher accuracy (Stockwell & Peterson, 2002). A data subset based on a smaller number of occurrences is more likely to produce varying predictions between model runs. In our case, the largest samples –corresponding to common species frequently observed in accessible areas– were also among the most spatially biased ones. Despite sample bias and size were not collinear, a high number of occurrences is more likely to be provided in accessible areas, which leads to a spatial bias and may explain the strong effect of correction for those species.

Biological relevance of variable selection

Variable selection differed the most at the site where the correction was effective. The majority of the models of the uncorrected group often selected additional

variables that were not biologically meaningful, presumably because they were locally correlated with the accessibility index (e.g. Intensive agriculture, water courses and distance to the coast in Trondheim; Figures S1 to S17 in Appendix). Models tended to explain the absence of occurrence data in the inaccessible area by the environmental variable which was the most represented there. Virtual species showed that the correction enabled the selection of variables which were biologically meaningful since they were used to generate the species distribution most of the time. Consistent with the effect of correction on predictions (see “Effect of sample bias”), variable selection did not differ when the species habitat variable was correlated with accessibility. In some cases, the correction could induce the selection of variables which were biologically irrelevant. For instance, the correction decreased the biological relevance of variable selection for one virtual species, and subsequently the effect of the correction (Figures S1 to S17 in Appendix). Along with the degree of correlation with accessibility factors, this stresses the need to carefully investigate the biological relevance of the variable selected before and after correction in accordance with the recommendations from Hijmans (2012) and Fourcade *et al.* (2018).

Consistent with the target-group approach, the benefit conveyed by sample bias correction with accessibility maps may be limited when the geographic sampling bias translates into a bias in the environ-

Table 1. Variables and response functions used to generate virtual species, and selected variables during the modelling process for the uncorrected and the corrected group. We give the distribution family and the parameters of the response function. Variables that are biologically irrelevant are shown in bold.

Variable used for generation	Response function	Site	Random group Selected variable	Variable importance	Corrected group Selected variable	Variable importance	
Distance from non-intensive agriculture	Logistic (beta = -500, alpha = -5000)	Grote Nete Trondheim Thau	Proportion of non-intensive agriculture within	0.968	Distance from non-intensive agriculture	0.971	
			Distance from non-intensive agriculture	0.968	Distance from non-intensive agriculture	0.969	
			Distance from non-intensive agriculture	0.972	Distance from non-intensive agriculture	0.969	
Distance from intensive agriculture	Logistic (beta = 500, alpha = -5000)	Grote Nete Trondheim	Distance from intensive agriculture	0.983	Distance from intensive agriculture	0.984	
			Distance from intensive agriculture	0.925	Distance from intensive agriculture	0.9	
			Distance from forest edges	0.111	Distance from forest edges	0.096	
		Thau	Distance from open natural areas		0.049	Proportion of coniferous forests within 1000m	0.067
			Proportion of water bodies within 1000m		0.035	Proportion of water bodies within 1000m	0.035
			Distance from intensive agriculture	0.984	Distance from intensive agriculture	0.983	
Distance from water courses	Gaussian (mean = 100, SD = 100)	Grote Nete Trondheim Thau	Distance from water courses	0.982	Distance from water courses	0.983	
			Distance from water courses	0.903	Distance from water courses	0.983	
			Distance from coastal areas	0.151			
Distance from water courses	Gaussian (mean = 100, SD = 500)	Grote Nete Trondheim Thau	Distance from water courses	0.984	Distance from water courses	0.984	
			Distance from non-intensive agriculture	0.185			
			Distance from water courses	0.711	Distance from water courses	0.984	
Distance from water courses	Gaussian (mean = 100, SD = 500)	Grote Nete Trondheim Thau	Distance from coastal areas	0.435			
			Distance from water courses	0.985	Distance from water courses	0.985	
Grassland Index	Logistic (beta = -1000, alpha = -50)	Grote Nete	Grassland Index	0.837	Grassland Index	0.734	
			Proportion of non-intensive agriculture within	0.17	Proportion of non-intensive agriculture within 200m	0.189	
		Trondheim Thau	Distance from open natural areas		0.136	Distance from open natural areas	
			Grassland Index	0.972	Grassland Index	0.971	
Distance from open natural areas	Logistic (beta = -500, alpha = -5000)	Grote Nete Trondheim	Distance from open natural areas	0.341	Proportion of open natural areas within 200m	0.684	
			Proportion of open natural areas within 200m	0.524	Distance from open natural areas	0.196	
			Distance from open natural areas	0.417	Proportion of open natural areas within 200m	0.525	
		Thau	Proportion of open natural areas within 200m	0.39	Distance from open natural areas	0.337	
			Distance from forest edges	0.126			
			Distance from open natural areas	0.52	Distance from open natural areas	0.556	
Distance from open natural areas	Logistic (beta = -500, alpha = -2000)	Grote Nete Trondheim	Proportion of open natural areas within 200m	0.335	Proportion of open natural areas within 200m	0.306	
		Thau	Distance from open natural areas	0.968	Distance from open natural areas	0.965	
			Distance from open natural areas	0.619	Distance from open natural areas	0.967	
		Distance from intensive agriculture	0.479				
		Distance from open natural areas	0.971	Distance from open natural areas	0.965		

mental predictors (Ranc *et al.*, 2017). In other terms, sample bias issues cannot be fixed on the basis of accessibility maps when the distribution of environmental variables matches that of accessibility. In our study regions, this might be the case for species that are the most impacted by urbanisation (Geslin *et al.*, 2013) and agricultural intensification (Jeliazkov *et al.*, 2016; Olivier *et al.*, 2020).

Correction effect relative to model stochasticity

In absence of independent, standardised data, the performance of SDMs and correction methods cannot be assessed properly. Here we propose a measure of the effect of the correction technique relative to the within-model stochasticity (between runs of varying input parameters, individually for each modelling technique) to inform the potential benefit conveyed by the correction. We show that the Relative Overlap Index was in better agreement with changes between corrected and uncorrected predictions of virtual species than the classic cross-validation performance metrics. This is also true for changes in the selected environmental variables across sites and species. The use of this index may be generalisable to species for which habitat is not restricted to the same section of the accessibility gradient. This metric can be used to indicate whether species distribution models are likely to be improved by sample bias correction.

Cross-validation metrics

The performance metrics based on cross-validation failed to detect an improvement in species range predictions, except for cAUC. This is consistent with Hertzog *et al.* (2014), who assessed the performance of a variety of bias correction techniques on the model's ability to predict the range of a beetle and found a striking difference between the evaluations based on partitioned datasets and field validation. In their study, cAUC was in agreement with field validation. In our case, cAUC was not correlated with the Relative overlap index (Figure S27 in Appendix). As specified by Hijmans (2012), a null geographic model may not be relevant when a species occurs in a single continuous range, which may explain why it did not detect an improvement of the correction in some cases. Another recently developed evaluation method consists in taking into account the accumulation curve of occurrences with the area predicted as suitable, and the amount of uninformative niche space predicted (Jiménez & Soberón, 2020). This method is appropriate when absence data are unavailable and is similar to our approach in that the evaluation is relative to a random component. The report of multiple metrics describing different aspects of model performance is recommended to improve the understanding and transparency of SDMs (Araújo *et al.*, 2019). We therefore recommend the use of Relative Overlap

Index and alternative metrics such as the cAUC as complementary measures to assess the performance of sample bias correction. The cAUC indicates the level of discrimination (the ability to distinguish between occupied and unoccupied sites) accounting for sorting sample bias. The ROI is to be interpreted as the degree of change in spatial predictions while accounting for variation between model replicates.

Increasing the potential use of biodiversity databases

Despite the increasing amount of open source, publicly available data, and EU policies to promote open access and data sharing, high quality data availability is still a major hinder to effectively inform decision-making. Here we presented some of the limitations inherent to existing database's to be able to advance in data mobilisation. Accounting for sample bias is challenging, especially for rare species or species for which the distribution range is small, with a subsequent low number of occurrence points. Pseudo-absence selection weighted by accessibility maps enable to account for spatial sample bias without requiring the use of filtering techniques and hence, without reducing the amount of data available. Sample bias correction is also relevant for any species that is broadly distributed when occurrence data are spatially biased (e.g. Hertzog *et al.* (2014)), provided the effect of correction is assessed. A proper assessment of the efficiency of sample bias correction techniques remains challenging when standardised data are unavailable. However, the use of a relative measure of its effect, along with virtual species represents a promising tool to increase the use of large, heterogeneous, biased datasets in species distribution models and biodiversity assessments.

ACKNOWLEDGEMENTS

This study was partly supported by the IMAGINE project (ERANET BIODIVERSA). Natuurpunt Studie Association provided the data for Grote Nete, Region in Belgium, LPO (*Ligue pour la Protection des Oiseaux*) Occitanie, provided the data for Thau Region, the French site. Data from Norway are part of GBIF. A special thank goes to Boris Leroy for helpful discussions and wise advices.

AUTHOR CONTRIBUTIONS

ND, ML and SL designed the analyses. ND performed the analyses. CP built the environmental variables and the map. GP, SM and PD aided in interpreting the results and worked on the manuscript. MLL and FR, SH and RM helped to organize data collected in France, Belgium and Norway, respectively. ND led the writing of the manuscript. All authors

read, commented and validated the final version of the manuscript.

REFERENCES

- Anderson, B. J., Arroyo, B. E., Collingham, Y. C., Etheridge, B., Fernandez–De–Simon, J., Gillings, S., Gregory, R. D., Leckie, F. M., Sim, I. M. W., Thomas, C. D., Travis, J. & Redpath, S. M. (2009) Using distribution models to test alternative hypotheses about a species’ environmental limits and recovery prospects. *Biological Conservation*, **142**, 488–499.
- Araújo, M. B., Anderson, R. P., Barbosa, A. M., Beale, C. M., Dormann, C. F., Early, R., Garcia, R. A., Guisan, A., Maiorano, L., Naimi, B., O’Hara, R. B., Zimmermann, N. E. & Rahbek, C. (2019) Standards for distribution models in biodiversity assessments. *Science Advances*, **5**, eaat4858.
- Araújo, M. B. & Guisan, A. (2006) Five (or so) challenges for species distribution modelling. *Journal of Biogeography*, **33**, 1677–1688.
- Araújo, M. B., Pearson, R. G., Thuiller, W. & Erhard, M. (2005) Validation of species–climate impact models under climate change. *Global Change Biology*, **11**, 1504–1513.
- Azam, C., Le Viol, I., Julien, J.-F., Bas, Y. & Kerbiriou, C. (2016) Disentangling the relative effect of light pollution, impervious surfaces and intensive agriculture on bat activity with a national-scale monitoring program. *Landscape Ecology*, **31**, 2471–2483.
- Barbet–Massin, M., Jiguet, F., Albert, C. H. & Thuiller, W. (2012) Selecting pseudo-absences for species distribution models: how, where and how many? *Methods in Ecology and Evolution*, **3**, 327–338.
- Bean, W. T., Stafford, R. & Brashares, J. S. (2012) The effects of small sample size and sample bias on threshold selection and accuracy assessment of species distribution models. *Ecography*, **35**, 250–258.
- Beck, J., Böller, M., Erhardt, A. & Schwanghart, W. (2014) Spatial bias in the GBIF database and its effect on modeling species’ geographic distributions. *Ecological Informatics*, **19**, 10–15.
- Bellard, C., Leroy, B., Thuiller, W., Rysman, J.-F. & Courchamp, F. (2016) Major drivers of invasion risks throughout the world. *Ecosphere*, **7**, e01241.
- Bellard, C., Thuiller, W., Leroy, B., Genovesi, P., Bakkenes, M. & Courchamp, F. (2013) Will climate change promote future invasions? *Global Change Biology*, **19**, 3740–3748.
- Bird, T. J., Bates, A. E., Lefcheck, J. S., Hill, N. A., Thomson, R. J., Edgar, G. J., Stuart–Smith, R. D., Wotherspoon, S., Krkosek, M., Stuart–Smith, J. F., Pecl, G. T., Barrett, N. & Frusher, S. (2014) Statistical solutions for error and bias in global citizen science datasets. *Biological Conservation*, **173**, 144–154.
- Boria, R. A., Olson, L. E., Goodman, S. M. & Anderson, R. P. (2014) Spatial filtering to reduce sampling bias can improve the performance of ecological niche models. *Ecological Modelling*, **275**, 73–77.
- Botella, C., Joly, A., Bonnet, P., Munoz, F. & Monestiez, P. (2021) Jointly estimating spatial sampling effort and habitat suitability for multiple species from opportunistic presence-only data. *Methods in Ecology and Evolution*.
- Bray, J. R. & Curtis, J. T. (1957) An Ordination of the Upland Forest Communities of Southern Wisconsin. *Ecological Monographs*, **27**, 325–349.
- Briscoe Runquist, R. D., Lake, T., Tiffin, P. & Moeller, D. A. (2019) Species distribution models throughout the invasion history of Palmer amaranth predict regions at risk of future invasion and reveal challenges with modeling rapidly shifting geographic ranges. *Scientific Reports*, **9**, 2426.
- Buisson, L., Thuiller, W., Casajus, N., Lek, S. & Grenouillet, G. (2010) Uncertainty in ensemble forecasting of species distribution. *Global Change Biology*, **16**, 1145–1157.
- Cola, V. D., Broennimann, O., Petitpierre, B., Breiner, F. T., D’Amen, M., Randin, C., Engler, R., Pottier, J., Pio, D., Dubuis, A., Pellissier, L., Mateo, R. G., Hordijk, W., Salamin, N. & Guisan, A. (2017) ecospat: an R package to support spatial analyses and modeling of species niches and distributions. *Ecography*, **40**, 774–787.
- Dubos, N., Viol, I. L., Robert, A., Teplitsky, C., Ghislain, M., Dehorter, O., Julliard, R. & Henry, P. Y. (2018) Disentangling the effects of spring anomalies in climate and net primary production on body size of temperate songbirds. *Ecography*, **41**, 1319–1330.
- Edrén, S. M. C., Wisz, M. S., Teilmann, J., Dietz, R. & Söderkvist, J. (2010) Modelling spatial patterns in harbour porpoise satellite telemetry data using maximum entropy. *Ecography*, **33**, 698–708.
- Ficetola, G. F., Bonardi, A., Múcher, C. A., Gilissen, N. L. M. & Padoa–Schioppa, E. (2014) How many predictors in species distribution models at the landscape scale? Land use versus LiDAR-derived canopy height. *International Journal of Geographical Information Science*, **28**, 1723–1739.

- Fourcade, Y., Besnard, A. G. & Secondi, J. (2018) Paintings predict the distribution of species, or the challenge of selecting environmental predictors and evaluation statistics. *Global Ecology and Biogeography*, **27**, 245–256.
- Fourcade, Y., Engler, J. O., Rödder, D. & Secondi, J. (2014) Mapping Species Distributions with MAXENT Using a Geographically Biased Sample of Presence Data: A Performance Assessment of Methods for Correcting Sampling Bias. *PLOS ONE*, **9**, e97122.
- Gábor, L., Moudrý, V., Barták, V. & Lecours, V. (2020) How do species and data characteristics affect species distribution models and when to use environmental filtering? *International Journal of Geographical Information Science*, 1567–1584.
- Geslin, B., Gauzens, B., Thébault, E. & Dajoz, I. (2013) Plant Pollinator Networks along a Gradient of Urbanisation. *PLOS ONE*, **8**, e63421.
- Hao, T., Elith, J., Guillera-Aroita, G. & Lahoz-Monfort, J. J. (2019) A review of evidence about use and performance of species distribution modelling ensembles like BIOMOD. *Diversity and Distributions*, **25**, 839–852.
- Hawkins, B. A., Leroy, B., Rodríguez, M., Singer, A., Vilela, B., Villalobos, F., Wang, X. & Zelený, D. (2017) Structural bias in aggregated species-level variables driven by repeated species co-occurrences: a pervasive problem in community and assemblage data. *Journal of Biogeography*, **44**, 1199–1211.
- Hertzog, L. R., Besnard, A. & Jay-Robert, P. (2014) Field validation shows bias-corrected pseudo-absence selection is the best method for predictive species-distribution modelling. *Diversity and Distributions*, **20**, 1403–1413.
- Hijmans, R., Phillips, S., Leathwick, J. & Elith, J. (2015) Dismo: Species distribution modeling. *R Package Version 1.0-12*.
- Hijmans, R. J. (2012) Cross-validation of species distribution models: removing spatial sorting bias and calibration with a null model. *Ecology*, **93**, 679–688.
- Iturbide, M., Bedia, J., Herrera, S., del Hierro, O., Pinto, M. & Gutiérrez, J. M. (2015) A framework for species distribution modelling with improved pseudo-absence generation. *Ecological Modelling*, **312**, 166–174.
- Jaccard, P. (1912) The distribution of the flora in the alpine zone. *New Phytologist*, **11**, 37–50.
- Jeliazkov, A., Chiron, F., Garnier, J., Besnard, A. & Silvestre, M. and Jiguet, F. (2014) Level-dependence of the relationships between amphibian biodiversity and environment in pond systems within an intensive agricultural landscape. *Hydrobiologia*, **723**, 7–23.
- Jeliazkov, A., Mimet, A., Chargé, R., Jiguet, F., Devictor, V. & Chiron, F. (2016) Impacts of agricultural intensification on bird communities: New insights from a multi-level and multi-facet approach of biodiversity. *Agriculture, Ecosystems & Environment*, **216**, 9–22.
- Jiménez, L. & Soberón, J. (2020) Leaving the area under the receiving operating characteristic curve behind: An evaluation method for species distribution modelling applications based on presence-only data. *Methods in Ecology and Evolution*, **11**, 1571–1586.
- Johnston, A., Moran, N., Musgrove, A., Fink, D. & Baillie, S. R. (2020) Estimating species distributions from spatially biased citizen science data. *Ecological Modelling*, **422**, 108927.
- Kramer-Schadt, S., Niedballa, J., Pilgrim, J. D., Schröder, B., Lindenborn, J., Reinfelder, V., Stillfried, M., Heckmann, I., Scharf, A. K., Augeri, D. M., Cheyne, S. M., Hearn, A. J., Ross, J., Macdonald, D. W., Mathai, J., Eaton, J., Marshall, A. J., Semiadi, G., Rustam, R., Bernard, H., Alfred, R., Samejima, H., Duckworth, J. W., Breitenmoser-Wuersten, C., Belant, J. L., Hofer, H. & Wilting, A. (2013) The importance of correcting for sampling bias in MaxEnt species distribution models. *Diversity and Distributions*, **19**, 1366–1379.
- Leroy, B., Bellard, C., Dubos, N., Colliot, A., Vasseur, M., Courtial, C., Bakkenes, M., Canard, A. & Ysnel, F. (2014) Forecasted climate and land use changes, and protected areas: the contrasting case of spiders. *Diversity and Distributions*, **20**, 686–697.
- Leroy, B., Meynard, C. N., Bellard, C. & Courchamp, F. (2016) virtualspecies, an R package to generate virtual species distributions. *Ecography*, **39**, 599–607.
- Li, W. & Guo, Q. (2013) How to assess the prediction accuracy of species presence-absence models without absence data? *Ecography*, **36**, 788–799.
- Liu, C., Newell, G. & White, M. (2019) The effect of sample size on the accuracy of species distribution models: considering both presences and pseudo-absences or background sites. *Ecography*, **42**, 535–548.
- Lobo, J. M. & Tognelli, M. F. (2011) Exploring the effects of quantity and location of pseudo-absences and sampling biases on the performance of distribution models with limited point occurrence data. *Journal for Nature Conservation*, **19**, 1–7.

- McKinley, D. C., Miller-Rushing, A. J., Ballard, H. L., Bonney, R., Brown, H., Cook-Patton, S. C., Evans, D. M., French, R. A., Parrish, J. K., Phillips, T. B., Ryan, S. F., Shanley, L. A., Shirk, J. L., Stepenuck, K. F., Weltzin, J. F., Wiggins, A., Boyle, O. D., Briggs, R. D., Chapin, S. F., Hewitt, D. A., Preuss, P. W. & Soukup, M. A. (2017) Citizen science can improve conservation science, natural resource management, and environmental protection. *Biological Conservation*, **208**, 15–28.
- Meynard, C. N. & Kaplan, D. M. (2012) The effect of a gradual response to the environment on species distribution modeling performance. *Ecography*, **35**, 499–509.
- Meynard, C. N., Leroy, B. & Kaplan, D. M. (2019) Testing methods in species distribution modelling using virtual species: what have we learnt and what are we missing? *Ecography*, **42**, 2021–2036.
- Mikolajczak, A., Maréchal, D., Sanz, T., Isenmann, M., Thierion, V. & Luque, S. (2015) Modelling spatial distributions of alpine vegetation: A graph theory approach to delineate ecologically-consistent species assemblages. *Ecological Informatics*, **30**, 196–202.
- Monsarrat, S., Boshoff, A. F. & Kerley, G. I. H. (2019) Accessibility maps as a tool to predict sampling bias in historical biodiversity occurrence records. *Ecography*, **42**, 125–136.
- Norberg, A., Abrego, N., Blanchet, F. G., Adler, F. R., Anderson, B. J., Anttila, J., Araújo, M. B., Dallas, T., Dunson, D., Elith, J., Foster, S. D., Fox, R., Franklin, J., Godsoe, W., Guisan, A., O'Hara, B., Hill, N. A., Holt, R. D., Hui, F. K. C., Husby, M., Kålås, J. A., Lehtikoinen, A., Luoto, M., Mod, H. K., Newell, G., Renner, I., Roslin, T., Soininen, J., Thuiller, W., Vanhatalo, J., Warton, D., White, M., Zimmermann, N. E., Gravel, D. & Ovaskainen, O. (2019) A comprehensive evaluation of predictive performance of 33 species distribution models at species and community levels. *Ecological Monographs*, **89**, e01370.
- Olivier, T., Thébaud, E., Elias, M., Fontaine, B. & Fontaine, C. (2020) Urbanization and agricultural intensification destabilize animal communities differently than diversity loss. *Nature Communications*, **11**, 2686.
- Otegui, J., Ariño, A. H., Encinas, M. A. & Pando, F. (2013) Assessing the Primary Data Hosted by the Spanish Node of the Global Biodiversity Information Facility (GBIF). *PLOS ONE*, **8**, e55144.
- Phillips, S. J., Dudík, M., Elith, J., Graham, C. H., Lehmann, A., Leathwick, J. & Ferrier, S. (2009) Sample selection bias and presence-only distribution models: implications for background and pseudo-absence data. *Ecological Applications*, **19**, 181–197.
- Ranc, N., Santini, L., Rondinini, C., Boitani, L., Poitevin, F., Angerbjörn, A. & Maiorano, L. (2017) Performance tradeoffs in target-group bias correction for species distribution models. *Ecography*, **40**, 1076–1087.
- Robinson, O. J., Ruiz-Gutierrez, V. & Fink, D. (2018) Correcting for bias in distribution modelling for rare species using citizen science data. *Diversity and Distributions*, **24**, 460–472.
- Rödger, D. & Engler, J. O. (2011) Quantitative metrics of overlaps in Grinnellian niches: advances and possible drawbacks. *Global Ecology and Biogeography*, **20**, 915–927.
- Schwartz, M. W. (2012) Using niche models with climate projections to inform conservation management decisions. *Biological Conservation*, **155**, 149–156.
- Senay, S. D., Worner, S. P. & Ikeda, T. (2013) Novel Three-Step Pseudo-Absence Selection Technique for Improved Species Distribution Modelling. *PLOS ONE*, **8**, e71218.
- Sicacha-Parada, J., Steinsland, I., Cretois, B. & Borgelt, J. (2020) Accounting for spatial varying sampling effort due to accessibility in Citizen Science data: A case study of moose in Norway. *Spatial Statistics*, 100446.
- Soberón, J. & Nakamura, M. (2009) Niches and distributional areas: Concepts, methods, and assumptions. *Proceedings of the National Academy of Sciences*, **106**, 19644–19650.
- Stockwell, D. R. B. & Peterson, A. T. (2002) Effects of sample size on accuracy of species distribution models. *Ecological Modelling*, **148**, 1–13.
- Stolar, J. & Nielsen, S. E. (2015) Accounting for spatially biased sampling effort in presence-only species distribution modelling. *Diversity and Distributions*, **21**, 595–608.
- Thibaud, E., Petitpierre, B., Broennimann, O., Davison, A. C. & Guisan, A. (2014) Measuring the relative effect of factors affecting species distribution model predictions. *Methods in Ecology and Evolution*, **5**.
- Thuiller, W., Lafourcade, B., Engler, R. & Araújo, M. B. (2009) BIOMOD – a platform for ensemble forecasting of species distributions. *Ecography*, **32**, 369–373.
- Varela, S., Anderson, R. P., García-Valdés, R. & Fernández-González, F. (2014) Environmental filters reduce the effects of sampling bias and improve predictions of ecological niche models. *Ecography*, **37**, 1084–1091.

- Vollering, J., Halvorsen, R., Auestad, I. & Rydgren, K. (2019) Bunching up the background betters bias in species distribution models. *Ecography*, **42**, 1717–1727.
- Warren, D. L., Glor, R. E. & Turelli, M. (2010) ENM-Tools: a toolbox for comparative studies of environmental niche models. *Ecography*, **33**, 607–611.
- Yackulic, C. B., Chandler, R., Zipkin, E. F., Royle, J. A., Nichols, J. D., Grant, E. H. C. & Veran, S. (2013) Presence-only modelling using MAXENT: when can we trust the inferences? *Methods in Ecology and Evolution*, **4**, 236–243.

APPENDIX




Figure S1. Species projections in Grote Nete (uncorrected versus corrected group). From top to bottom, *Alcedo atthis*, *Anthus trivialis*, *Castor fiber* and *Dryocopus martius*.




Figure S2. Species projections in Grote Nete (uncorrected versus corrected group). From top to bottom, *Lullula arborea*, *Luscinia megarhynchos*, *Luscinia svecica* and *Oriolus oriolus*.




Figure S3. Species projections in Grote Nete (uncorrected versus corrected group). From top to bottom, *Phoenicurus hoenicurus*, *Poecile montanus* and *Sciurus vulgaris*.




Figure S4. Species projections in Thau (uncorrected versus corrected group). From top to bottom, *Epidalea calamita*, *Natrix natrix*, *Plecotus austriacus* and *Rhinolophus ferrumequinum*.




Figure S5. Species projections in Thau (uncorrected versus corrected group). From top to bottom, *Tadarida teniotis*, *Timon lepidus* and *Triturus marmoratus*.




Figure S6. Species projections in Trondheim (uncorrected versus corrected group). From top to bottom, *Accipiter gentilis*, *Accipiter nisus*, *Actitis hypoleucos* and *Aegithalos caudatus*.




Figure S7. Species projections in Trondheim (random versus corrected group). From top to bottom, *Alauda arvensis*, *Anas acuta*, *Anas penelope* and *Anthus petrosus*.




Figure S8. Species projections in Trondheim (uncorrected versus corrected group). From top to bottom, *Anthus pratensis*, *Anthus rivialis*, *Asio lammeus* and *Aythya fuligula*.




Figure S9. Species projections in Trondheim (uncorrected versus corrected group). From top to bottom, *Bombycilla garrulus*, *Bucephala clangula*, *Calidris alpina* and *Calidris pugnax*.




Figure S10. Species projections in Trondheim (uncorrected versus corrected group). From top to bottom, *Calidris temminckii*, *Carduelis flammea*, *Cepphus grylle* and *Certhia familiaris*.




Figure S11. Species projections in Trondheim (uncorrected versus corrected group). From top to bottom, *Charadrius hiaticula*, *Cinclus cinclus*, *Coccythraustes coccythraustes* and *Cygnus cygnus*.




Figure S12. Species projections in Trondheim (uncorrected versus corrected group). From top to bottom, *Delichon urbicum*, *Dendrocopos major*, *Dendrocopos minor* and *Dryocopus martius*.




Figure S13. Species projections in Trondheim (uncorrected versus corrected group). From top to bottom, *Emberiza citrinella*, *Emberiza schoeniclus*, *Erithacus rubecula* and *Ficedula hypoleuca*.




Figure S14. Species projections in Trondheim (uncorrected versus corrected group). From top to bottom, *Gallinago gallinago*, *Garrulus glandarius*, *Gavia arctica* and *Gavia stellata*.




Figure S15. Species projections in Trondheim (uncorrected versus corrected group). From top to bottom, *Glaucidium passerinum*, *Grus grus*, *Haematopus ostralegus* and *Hippolais icterina*.




Figure S16. Species projections in Trondheim (uncorrected versus corrected group). From top to bottom, *Hirundo rustica*, *Lanius excubitor*, *Lissotriton vulgaris* and *Loxia curvirostra*.




Figure S17. Species projections in Trondheim (uncorrected versus corrected group). From top to bottom, *Loxia pytyopsittacus*, *Luscinia svecica*, *Mergus serrator* and *Motacilla alba*.




Figure S18. Site-specific variation in the effect of sample bias correction. Relationship between the effect of sample bias correction (Pearson's overlap between uncorrected and corrected predictions) and the effect of model stochasticity (Pearson's overlap between model runs). Points located below the $y = x$ red line represent models for which the relative effect of sample bias correction exceeds that of model stochasticity. The inset shows the boxplots of the Relative Overlap Index (ROI) according to the case study site. Each boxplot is composed of the first decile, the lower hinge, the median, the upper hinge and the ninth decile.




Figure S19. Site-specific variation in the effect of sample bias correction. Relationship between the effect of sample bias correction (Spearman's overlap between uncorrected and corrected predictions) and the effect of model stochasticity (Spearman's overlap between model runs). Points located below the $y = x$ red line represent models for which the relative effect of sample bias correction exceeds that of model stochasticity. The inset shows the boxplots of the Relative Overlap Index (ROI) according to the case study site. Each boxplot is composed of the first decile, the lower hinge, the median, the upper hinge and the ninth decile.




Figure S20. Virtual species projections in Grote Note (true, uncorrected and corrected group). From top to bottom, *Philocourdos belgiumensis*, *Philocourdogenos belgiumensis* and *Philoherbas belgiumensis*.




Figure S21. Virtual species projections in Grote Note (true, uncorrected and corrected group). From top to bottom, *Philoherbagenos belgiumensis*, *Fugiagris belgiumensis* and *Alteragris belgiumensis*.




Figure S22. Virtual species projections in Thau (true, uncorrected and corrected group). From top to bottom, *Philocourdos thauensis*, *Philocourdogenos thauensis* and *Philoherbas thauensis*.




Figure S23. Virtual species projections in Thau (true, uncorrected and corrected group). From top to bottom, *Philoherbagenos thauensis*, *Fugiagrish thauensis* and *Alteragris thauensis*.




Figure S24. Virtual species projections in Trondheim (true, uncorrected and corrected group). From top to bottom, *Philocourdos norwegensis*, *Philocourdogenos norwegensis* and *Philoherbas norwegensis*.




Figure S25. Virtual species projections in Trondheim (true, uncorrected and corrected group). From top to bottom, *Philoherbagenos norwayensis*, *Fugiagris norwayensis* and *Alteragris norwayensis*.




Figure S26. Overlap with the “true” probability of occurrence (virtual species) for the uncorrected and the corrected group at three sites. An improvement in model performance is indicated by a higher overlap Pearson's r (a) and a lower RMSE (b). Each boxplot is composed of the first decile, the lower hinge, the median, the upper hinge and the ninth decile.




Figure S27. Relationship between delta cAUC (difference between corrected and uncorrected group) and the Relative Overlap Index (Pearson's $r = 0.03$).

Table S1. Study species, corresponding site, sample size and sample bias. Sample bias was estimated from Boyce indices based on occurrence points and accessibility maps.

Species	Site	Sample size	Sample bias
<i>Alcedo.atthis</i>	Grote Nete	612	-0.95
<i>Anthus.trivialis</i>	Grote Nete	223	-0.94
<i>Anthus.trivialis</i>	Grote Nete	223	0.48
<i>Castor.fiber</i>	Grote Nete	72	-0.67
<i>Dryocopus.martius</i>	Grote Nete	443	0.71
<i>Dryocopus.martius</i>	Grote Nete	443	-0.94
<i>Lullula.arborea</i>	Grote Nete	169	-0.94
<i>Luscinia.megarhynchos</i>	Grote Nete	268	-0.89
<i>Luscinia.svecica</i>	Grote Nete	177	-0.14
<i>Luscinia.svecica</i>	Grote Nete	177	-0.73
<i>Oriolus.oriolus</i>	Grote Nete	129	-0.85
<i>Phoenicurus.phoenicurus</i>	Grote Nete	89	-0.59
<i>Poecile.montanus</i>	Grote Nete	303	-0.80
<i>Sciurus.vulgaris</i>	Grote Nete	380	0.10
<i>Epidalea.calamita</i>	Thau	82	0.20
<i>Natrix.natrix</i>	Thau	25	-0.46
<i>Plecotus.austriacus</i>	Thau	129	0.05
<i>Rhinolophus.ferrumequinum</i>	Thau	110	-0.24
<i>Tadarida.teniotis</i>	Thau	32	-0.60
<i>Tarentola.mauritanica</i>	Thau	293	0.74
<i>Timon.lepidus</i>	Thau	54	0.55
<i>Triturus.marmoratus</i>	Thau	63	0.30
<i>Anthus.trivialis</i>	Trondheim	398	0.48
<i>Anthus.trivialis</i>	Trondheim	398	-0.94
<i>Dryocopus.martius</i>	Trondheim	587	0.71
<i>Dryocopus.martius</i>	Trondheim	587	-0.94
<i>Luscinia.svecica</i>	Trondheim	53	-0.14
<i>Luscinia.svecica</i>	Trondheim	53	-0.73
<i>Accipiter.gentilis</i>	Trondheim	522	0.83
<i>Accipiter.nisus</i>	Trondheim	876	0.94
<i>Actitis.hypoleucos</i>	Trondheim	1285	0.74
<i>Aegithalos.caudatus</i>	Trondheim	390	0.36
<i>Alauda.arvensis</i>	Trondheim	219	0.78
<i>Anas.acuta</i>	Trondheim	20	-0.43
<i>Anas.penelope</i>	Trondheim	1472	0.74
<i>Anthus.petrosus</i>	Trondheim	119	0.70
<i>Anthus.pratensis</i>	Trondheim	979	0.89
<i>Asio.flammeus</i>	Trondheim	22	0.10
<i>Aythya.fuligula</i>	Trondheim	2713	0.29
<i>Bombus.garrulus</i>	Trondheim	2028	0.89
<i>Bucephala clangula</i>	Trondheim	9213	0.69
<i>Calidris.alpina</i>	Trondheim	421	0.39
<i>Calidris.canutus</i>	Trondheim	77	
<i>Calidris.minuta</i>	Trondheim	21	0.52
<i>Calidris.pugnax</i>	Trondheim	189	0.15
<i>Calidris.temminckii</i>	Trondheim	56	0.68
<i>Carduelis.flammea</i>	Trondheim	3226	0.89
<i>Cephus.grylle</i>	Trondheim	576	-0.30
<i>Certhia.familiaris</i>	Trondheim	1327	0.88
<i>Charadrius.hiaticula</i>	Trondheim	1095	0.84
<i>Cinclus.cinclus</i>	Trondheim	1119	0.73
<i>Coccothraustes.coccothraustes</i>	Trondheim	614	0.50
<i>Cygnus.cygnus</i>	Trondheim	1825	0.94
<i>Delichon.urbicum</i>	Trondheim	404	0.84
<i>Dendrocopos.major</i>	Trondheim	3521	0.83
<i>Dendrocopos.minor</i>	Trondheim	207	0.74
<i>Emberiza.citrinella</i>	Trondheim	5509	0.86
<i>Emberiza.schoeniclus</i>	Trondheim	635	0.73
<i>Eptesicus.nilssonii</i>	Trondheim	21	0.23
<i>Erithacus.rubecula</i>	Trondheim	5342	0.92
<i>Ficedula.hypoleuca</i>	Trondheim	607	0.84
<i>Gallinago.gallinago</i>	Trondheim	299	0.83
<i>Garrulus.glandarius</i>	Trondheim	1372	0.89
<i>Gavia.arctica</i>	Trondheim	197	0.06
<i>Gavia.stellata</i>	Trondheim	1587	0.41
<i>Glaucidium.passerinum</i>	Trondheim	60	-0.29
<i>Grus.grus</i>	Trondheim	501	0.66
<i>Haematopus.ostralegus</i>	Trondheim	2477	0.51
<i>Hippolais.icterina</i>	Trondheim	513	0.78
<i>Hirundo.rustica</i>	Trondheim	1637	0.80
<i>Lanius.excubitor</i>	Trondheim	111	0.81
<i>Lissotriton.vulgaris</i>	Trondheim	22	-0.06
<i>Loxia.curvirostra</i>	Trondheim	590	0.89
<i>Loxia.leucoptera</i>	Trondheim	84	-0.20
<i>Loxia.pytyopsittacus</i>	Trondheim	86	-0.59
<i>Mergus.merganser</i>	Trondheim	2818	0.58
<i>Mergus.serrator</i>	Trondheim	2295	0.17
<i>Motacilla.alba</i>	Trondheim	3118	0.88
<i>Rana.temporaria</i>	Trondheim	34	0.34