

Species-area relationships emerge from the assembly of complex ecological communities

Ankit Vikrant^{1*} | ankitv@chalmers.se

Martin Nilsson Jacobi¹ | mjacobi@chalmers.se

¹ *Department of Space, Earth and Environment, Chalmers University of Technology, Maskingränd 2, 412 58 Gothenburg, Sweden*

* Corresponding Author: Ankit Vikrant | Physical Resource Theory, Department of Space, Earth and Environment, Chalmers University of Technology, Maskingränd 2, 412 58 Gothenburg, Sweden | Telephone: +46761145991 | E-mail: ankitv@chalmers.se

Running Title: SARs from assembly of large ecological communities.

Keywords: Community assembly, macroecology, species-area relationship, complex ecological communities

Type of article: Letter

Number of words: 157 (abstract) and 3672 (main text)

Number of figures: 6 (main text) + 4 (supporting information)

Number of references: 38

Authorship: AV and MNJ conceived and designed the study, AV performed the simulations and wrote the paper, AV and MNJ analyzed the results.

Data accessibility statement: No new data were used in the preparation of this manuscript.

Abstract

It has been a century since the species-area relationship (SAR) was first proposed as a power law to explain how species richness scales with area. There have been many attempts to explain the origin of this predominant form. Apart from the power law, numerous empirical studies also report a semi-log form of the SAR, but very few have addressed its incidence. In this work, we test whether these relationships could emerge from the assembly of large random communities. To this end, we reformulate the generalized Lotka-Volterra model replacing the interaction strength parameter by the inverse of area. This framework allows us to track the species richness values resulting from the assembly of large competitive communities over a range of areas. Our analysis demonstrates that the two most widely reported relationship forms can emerge due to differences in immigration rates and skewness towards weak interactions. We support our results regarding immigration rates through two related studies from the literature.

Introduction

The species-area relationship (SAR) is arguably the most widely studied scaling law in ecology, which has received empirical support from numerous studies spanning different geographical regions and taxa (Drakare et al., 2006; Lomolino and Weiser, 2001). The predominant power law form of the SAR was first described by O. Arrhenius in 1921 (Arrhenius, 1921). It related the number of species S to the area of a habitat A as $S \sim A^z$, where the exponent z varies between 0 and 1. A quantitative meta-analysis of a large number of SAR studies estimated its average value as 0.27 (Drakare et al., 2006). The power law was contested by a semi-log relationship in 1922, that advocated the form $S \sim \log(A^z)$ (Gleason, 1922). While the power law relationship is more widely reported, the semi-log SAR has also found support from numerous studies (Drakare et al., 2006; Lomolino and Weiser, 2001). There have been attempts to explain the power law form based on species distributions (Šizling and Storch, 2004; Picard et al., 2004; Leitner and Rosenzweig, 1997; Coleman, 1981), abundance distributions (Preston, 1948) or population dynamics through constraints on immigration (Bastolla et al., 2001; Durrett and Levin, 1996). The prevalence of SARs has also been attributed to the combined effects of widely observed abundance distributions and the fact that individuals from the same species cluster (Martín and Goldenfeld, 2006). The semi-log relationship can be recovered from the power law SAR in some limit using species-incidence functions that depend on colonization and extinction rates (?). However, there is no unified framework to explain the emergence of these competing SARs.

These scaling relationships are emergent in that they could be described by coarse-grained dynamics of large communities at the species level without reference to finer details and properties of individual organisms. Understanding the assembly of large communities could therefore underpin mechanisms

that shape these scaling laws. The analysis of large systems has benefitted from many emerging approaches in the recent decades. In 1972, P.W. Anderson influenced the philosophy of science by suggesting that ‘more is different’ (Anderson, 1972), based on accumulating evidence from various disciplines. This means that the properties of a collective composed of many parts could be drastically different from the parts themselves. In the same year, May used random matrix theory to show that large ecosystems become unstable when their complexity increases beyond a threshold (May, 1972), which contradicted the prevailing notion that diversity increases stability. May’s analytical results showed that one cannot have indefinite stability in large and complex ecosystems with many interactions. There is a limit beyond which an ecosystem is not resilient to small perturbations and can exhibit large fluctuations in the population abundances of the constituent species. He defined complexity in terms of connectance and interaction strength of the random matrix that encodes species interactions.

The complex dynamics of such random interaction networks can be modelled using the Generalized Lotka-Volterra (GLV) equations. This model has been employed to uncover theoretical results ranging from identification of structural properties that affect coexistence (Serván et al., 2018) to the study of generic assembly patterns that are consistent across network structures (Barbier et al., 2018; Bunin, 2017). The tractability of these equations has also invited their widespread use in modelling microbial communities based on experimental data (Bashan et al., 2016; Buffie et al., 2015; Marino et al., 2014; Mounier et al., 2008). Some recent studies have investigated the distribution of number of coexisting species that results from GLV dynamics of much larger species pools (Serván et al., 2018). Others have even explored the progression and boundaries of extinction in large ecosystems (Pettersson et al., 2020). These studies depart from identifying constraints on parameters that result in complete co-existence of all species.

When one increases the interaction strength beyond the regime where all species co-exist, the system wades through a phase characterized by single-species extinctions that was termed as the extinction continuum in Pettersson et al. (2020). May’s stability limit marks the end of this phase beyond which no stable equilibria exist.

We hypothesize that a modified GLV model accounting for spatial scaling could exhibit SARs through the assembly of random communities of different sizes. Our argument relies on interpreting the inverse of interaction strength parameter as area to test these questions. We explore a large part of this area parameter space to recover different number of surviving species beyond the regime of complete co-existence. By further allowing demographic immigration in the modified GLV model, we demonstrate that the two popular forms of the SAR stem from differences in immigration rates and the skewness towards weak interactions.

Material and Methods

Generalized Lotka-Volterra with spatial scaling

In its usual form, the GLV model describes the dynamics of species with densities y_i through the following equations:

$$\frac{dy_i}{dt} = r_i y_i \left(1 - \frac{y_i}{K_i}\right) + \sigma y_i \sum_j B_{ij} y_j \quad (1)$$

where K_i and r_i denote the carrying capacity and growth rate of i^{th} species. B_{ij} expresses pairwise inter-specific interactions between species i and any other species j . The full matrix B contains information about all possible pairwise interactions between species. Equation 1 implies that in the absence of interactions, each species grows to its carrying capacity K_i .

For a given value of σ below May’s limit, the system eventually relaxes to a stable equilibrium that represents an assembled community where species densities are resilient

to small perturbations (Fig. 1). The equilibrium densities of species at this stable fixed point could either be zero or positive. Fixed points without any extinctions are called feasible solutions. Though many studies have looked at feasibility in large ecosystems (Dougoud et al., 2018; Stone, 2018; Goh and Jennings, 1977), we focus on the relationship between species richness of assembled communities and σ along the extinction continuum (Pettersson et al., 2020).

A natural question is whether there is a physical interpretation of the parameter σ . First note that σ operates by scaling all the interactions within the ecosystem by the same magnitude. This implies that an increase in this parameter reflects increased encounter rates between various species. We posit that σ behaves like the inverse of area such that an increase in this parameter is emulated by a corresponding decrease in area. This especially holds for a community of competitive species since they are more likely to encounter each other when contesting for common resources.

To make this argument clearer, we replace densities in the GLV model by absolute abundances x_i and area. Without the interaction strength, the equations are:

$$\frac{dx_i}{dt} = r_i x_i \left(1 - \frac{x_i}{K_i}\right) + \frac{x_i A_0}{A} \sum_j B_{ij} x_j \quad (2)$$

where A_0 parameterises this model for a given ecosystem. We set this parameter equal to 1 from hereon.

This set of equations is of the same form as the GLV model but now the interaction strength parameter σ is replaced by the inverse of area. Increase in interaction strength is analogous to decrease in area, which increases the encounter rates between species. Tuning the interaction strength also simultaneously scales the variance of the interaction matrix. One could instead directly tune the variance of the interaction matrix ignoring the interaction strength parameter. How is the change in variance in the two cases different? The variance in

interactions could partially arise from habitat heterogeneity. Increased heterogeneity in an ecosystem with a fixed area reduces the effective area available to each species as per the area-heterogeneity tradeoff (Ben-Hur and Kadmon, 2020; Allouche et al., 2012; Kadmon and Allouche, 2007). This scenario would better correspond to maintaining a constant interaction strength (or area as per our interpretation) and only scaling the variance of the interaction matrix. In the absence of colonisation by new species, the number of species would effectively decrease as heterogeneity increases. This decrease in the species richness is almost identical whether one scales the variance or the interaction strength.

We investigate the dynamics of large ecosystems from a spatial scaling perspective using the modified GLV model with an area parameter. We use our model to simulate ecosystem dynamics as follows:

1. We pick entries of the interaction matrix B_{ij} from a normal distribution that is symmetric around a negative mean. Our specific choice is a normal distribution with mean -1 and variance 0.2. This particular choice allows for a large range of area values for which the system relaxes to stable equilibria.
2. The growth rates r_i are drawn from a normal distribution with mean +1. The constraints on interactions and growth rates describe a community of competitive species.
3. Starting from an initial area, the number of surviving species is plotted against decreasing area A .

Analogous to the scenario of increasing σ , the system relaxes to a unique stable fixed point when the area parameter is above a certain threshold. We obtain the number of surviving species from the fixed point for each value of the area parameter (Fig. 1). We hypothesize that the different number of surviving species

obtained by varying the area parameter result in widely reported SARs. These relationships are usually studied for one type of species or species that are placed in the same trophic level. This is congenial to our choice of a competitive interaction matrix. A competitive system could represent functional groups such as pollinators that compete for some common resources. A competitive GLV model with demographic noise has been shown to reproduce neutral island theories of Wilson-MacArthur and Hubbell (Kessler and Shnerb, 2015). The power-law SAR has also been recovered from a spatially-explicit extension of the Lotka-Volterra competition model that allowed migration between patches (O’Sullivan et al., 2019).

Our analysis describes ecosystems shrinking in area, where equilibrium abundances obtained for a given area parameter are used as initial abundances for the subsequent area parameter value. An alternative but equivalent interpretation is a scenario where a regional pool of species is available to colonise different islands in a region (Kessler and Shnerb, 2015). For an island defined by its area, the dynamics resulting from our model culminates in a final community where some species from the regional pool might not be feasible. Islands of different sizes yield communities with different compositions as a consequence. This interpretation results in spatial patterns identical to the area-reduction scenario described here.

Spatial scaling patterns with immigration

Immigration slows down the decline in number of surviving species through introduction of new species (MacArthur and Wilson, 1963) or by delaying extinctions through incoming individuals of existing species (demographic immigration) (Brown and Kodric-Brown, 1977). What effects do different levels of immigration have on spatial scaling patterns in our model ecosystem? To address this, we redefine our GLV model with an additional term for demographic immigration:

$$\frac{dx_i}{dt} = r_i x_i \left(1 - \frac{x_i}{K_i}\right) + \frac{x_i}{A} \sum_j B_{ij} x_j + \lambda e^{-\beta/\sqrt{A}} \quad (3)$$

The last term represents the immigration rate. This term has a negligible contribution for smaller values of area, where a species may go extinct without support from the growth and interaction terms. As the area of an ecosystem shrinks, its distance from other patches also increases. This reduces the possibility of immigrants entering the ecosystem. The immigration term in the above equation has an exponential function that represents varying levels of demographic rescue (Brown and Kodric-Brown, 1977). For large values of area, λ is the effective immigration rate. β is a constant in the exponential function analogous to the characteristic length scale in the spatially extended GLV model described in (O’Sullivan et al., 2019). We fixed $\beta = 1000$ and compared the results for different values of λ .

We also consider interaction networks with more realistic connectances and distributions of interactions between species. Real food-webs are characterized by many weak and few strong interactions that also endow stability (?). We study how the preponderance of weak interactions influences SARs. We use interactions drawn from exponential distributions that represent communities with varying skewness towards weak interactions. The rate parameter of the exponential distribution serves as a measure of this skew.

Results

Fig. 2 corresponds to the simplest case of an ecosystem with full connectance and no immigration. Starting with 100 species, we plot the number of surviving species against the area parameter after the first extinction occurs. The SAR is best represented by a semi-log function through our model. The curve saturates at an upper asymptote for very high values of the area parameter.

The slope of the semi-log SAR varies with changes in the means of interactions and growth rates. It is also worth noting that for an intermediate range of areas, even the log-log plot could show a misleadingly good fit for a power law SAR (Fig. 2).

What determines a log-log or a semi-log SAR?

Our model – in its simplest form – supports the semi-log relationship, that is also widely reported in literature (Drakare et al., 2006). Our analysis suggests that varying levels of immigration lead to different functional forms of the SAR. We start with a very low value of λ and progressively increase it to check the resulting SAR. For very low immigration rates, the semi-log relationship is supported (see Fig. S1 in Supporting Information) as seen in the scenario without immigration (Fig. 2). However, there exists an intermediate regime best characterised by a power law (Fig. 3). This form of the SAR also lacks the upper asymptote that we observed in the semi-log fit (Fig. 2). Interestingly, using area in the immigration term instead of its square root does not change the above results (see Fig. S2).

The level of skew towards weak interactions strongly influences SAR shape. Given the same immigration level, a higher skew towards weak interactions favours a semi-log relationship (see Fig. S3 in Supporting Information, also Fig. 4). This result does not change for fat-tailed distributions such as the Pareto distribution in the regime where stable solutions exist (preferably shape parameter ≥ 2 , see Fig. S3 in Supporting Information). When studied for communities with low connectances, the SARs thus obtained have lower slopes as in most natural communities (Fig. 4).

Discussion

Immigration shapes SARs: A case of bird studies from the Southwest Pacific Islands

An important study from the Solomon Archipelago (Diamond and Mayr, 1976) demonstrates some of the predictions from our theoretical model. The dataset used in that study has islands with areas spanning over six orders of magnitude, conclusively differentiating between competing forms of the SAR. The authors assume that intra-archipelago immigration rates are much higher than the immigration rates from the ‘source’ island of New Guinea, that is at least 600 kilometres away from the nearest island.

They further plot the SAR for three groups of islands within the Solomon Archipelago, which supports a semi-log form. The slope of the SAR is nearly the same across these three groups of islands (Fig. 5). We surmise that the immigration of birds into an island is also balanced by emigration to other islands within the same group. In other words, the system is at a steady-state of zero or very low immigration within the archipelago. Thus, any effective immigration should emanate from the source island or from islands in other distant archipelagos. The large distance from these other sources implies that the net immigration rates to the Solomon islands are very low. In fact, the authors state that with increasing isolation of an archipelago, the SAR may shift in form from a power function to an exponential (semi-log). This is consistent with what we find from our theoretical model that low immigration rates lead to semi-log SARs. The authors exclude ‘isolated’ islands from this analysis, that are far from large islands within the archipelago. This is congenial to our model because many of these islands have endemic species that are not found elsewhere in the archipelago such as *Zosterops rennellianus* and *Pachycephala feminina*, both restricted to the Rennell island. Speciation might influence the assembled communities especially on islands with fewer species.

This dataset also has many islands smaller than a few square kilometres, which are usually absent in many SAR studies (Lomolino and Weiser, 2001). Both forms of the SAR could show a very good (and similar) fit to data for larger island sizes (Fig. 5). As the authors point out, really small islands should be included in SAR analyses to conclusively identify the correct form of the relationship (Diamond and Mayr, 1976).

Another study from islands that lie 5 to 300 miles from New Guinea, found a power-law SAR (Diamond, 1972). Considering that these islands lie closer to the ‘source’ island of New Guinea, the immigration rates are likely to be higher than those for the Solomon Archipelago. This lends support to our theoretical results on the incidence of power-law SARs for higher immigration levels. The power-law SAR in Diamond (1972) excludes islands whose avifaunas have not reached equilibrium. These are recolonized volcanic islands and islands that have undergone overall size contraction or modification of connecting land-bridges in the past c. 10,000 years. This highlights the importance of selecting only those areas where assembled communities have reached stable equilibrium.

Limits to inferring SAR forms and the relevance of community data

Much attention has been devoted to explain the power law form of the SAR. Our results show the emergence of two most widely observed forms of SAR through differences in immigration rates and skewness towards weak interactions (Fig. 6). Our GLV model-based approach supports the view that community dynamics brings about such spatial patterns. The area form of the GLV model works more effectively for competitive communities – which are relevant to study of SARs – because similar species compete for common resources. This scales the interactions more uniformly with change in area. In general, since we relate area to the interaction strength, the SARs seem to stem from the scaling of interactions for species of the same

trophic level.

Our analysis also alludes to the debate around existence of an upper asymptote in SARs. Many empirical studies are also constrained by the lack of very large islands in their dataset (Tjørve, 2003) and fail to address this issue about existence of an upper asymptote. Our simulations track SARs corresponding to areas over 2 orders of magnitude. We obtain these asymptotes for very large values of area especially for highly connected communities with no immigration (Fig. 2). These features are almost absent for cases that support the power-law relationship (Fig. 3, Fig. 4). We re-emphasize that in the absence of immigration, even the power law relationship shows a decent fit for intermediate range of areas suggesting that empirical studies might miss the correct form of SARs if they only have access to areas over a small range. Some of the claimed power laws may in fact alternatively be explained by a semi-log SAR. In general, we expect areas spanning more than one order of magnitude to discern whether a power law SAR exists or not. In case the power law shows a good fit, it is imperative to check if the data might be better explained by an alternative form. This prescription is akin to that for deducing power-law distributions from data (Clauset et al., 2009).

In addition to immigration rates and skewness towards weak interactions, connectance also influences SAR slopes. If all other parameters are kept the same, then systems with lower connectance result in lower SAR slopes (compare Fig. S3 in Supplementary Information versus Fig. 4). We also find that higher immigration rates correspond to higher SAR slopes, such that the number of surviving species fall off much more sharply with area for large areas. The SAR slopes we obtain are much more reasonable for choices based on realistic interaction networks (Fig. 4). We expect that some network structures could result in even lower slopes, but without much effect on the SAR form. An interesting direction is to study these spatial patterns in a multi-trophic community, and possibly see how the

relationships vary across trophic levels. Our framework could be used to study the spatially explicit dynamics of a mosaic of connected ecosystems, especially those with competitive communities. The migration of species between patches could be viewed as a diffusion process where the local dynamics of each patch is described by our model.

It is interesting that given a random configuration of competitive species, we recover many known features of SARs while also identifying factors that might best explain the variation in these relationships. Our model allows us to track a large range of areas for which stable communities exist, enabling a robust deduction of the correct SAR. While broadly studying the emergence of SARs, we also propose cautionary prescriptions for empirical studies to conclusively identify the shapes of these relationships. Given that many studies predict extinction scenarios based on assumed SARs, it is indispensable to reject possibilities that cannot be supported with limited data. We hope that our theoretical model prompts empirical studies to systematically evaluate the effects of immigration and community structure on species-area relationships.

References

- Omri Allouche, Michael Kalyuzhny, Gregorio Moreno-Rueda, Manuel Pizarro, and Ronen Kadmon. Area-heterogeneity tradeoff and the diversity of ecological communities. *Proceedings of the National Academy of Sciences*, 109(43):17495–17500, 2012.
- Philip W Anderson. More is different. *Science*, 177(4047):393–396, 1972.
- Olof Arrhenius. Species and area. *Journal of Ecology*, 9(1):95–99, 1921.
- Matthieu Barbier, Jean-François Arnoldi, Guy Bunin, and Michel Loreau. Generic assembly patterns in complex

- ecological communities. *Proceedings of the National Academy of Sciences*, 115(9):2156–2161, 2018.
- Amir Bashan, Travis E Gibson, Jonathan Friedman, Vincent J Carey, Scott T Weiss, Elizabeth L Hohmann, and Yang-Yu Liu. Universality of human microbial dynamics. *Nature*, 534(7606):259–262, 2016.
- U Bastolla, M Lässig, SC Manrubia, and A Valleriani. Diversity patterns from ecological models at dynamical equilibrium. *Journal of Theoretical Biology*, 212(1):11–34, 2001.
- Eyal Ben-Hur and Ronen Kadmon. An experimental test of the area–heterogeneity tradeoff. *Proceedings of the National Academy of Sciences*, 117(9):4815–4822, 2020.
- James H Brown and Astrid Kodric-Brown. Turnover rates in insular biogeography: effect of immigration on extinction. *Ecology*, 58(2):445–449, 1977.
- Charlie G Buffie, Vanni Bucci, Richard R Stein, Peter T McKenney, Lilan Ling, Asia Gobourne, Daniel No, Hui Liu, Melissa Kinnebrew, Agnes Viale, et al. Precision microbiome reconstitution restores bile acid mediated resistance to *clostridium difficile*. *Nature*, 517(7533):205–208, 2015.
- Guy Bunin. Ecological communities with lotka-volterra dynamics. *Physical Review E*, 95(4):042414, 2017.
- Aaron Clauset, Cosma Rohilla Shalizi, and Mark EJ Newman. Power-law distributions in empirical data. *SIAM review*, 51(4):661–703, 2009.
- Bernard D Coleman. On random placement and species-area relations. *Mathematical Biosciences*, 54(3-4):191–215, 1981.
- Jared M Diamond. Biogeographic kinetics: estimation of relaxation times for avifaunas of southwest pacific islands. *Proceedings of the National Academy of Sciences*, 69(11):3199–3203, 1972.
- Jared M Diamond and Ernst Mayr. Species-area relation for birds of the solomon archipelago. *Proceedings of the National Academy of Sciences*, 73(1):262–266, 1976.
- Michaël Dougoud, Laura Vinckenbosch, Rudolf P Rohr, Louis-Félix Bersier, and Christian Mazza. The feasibility of equilibria in large ecosystems: A primary but neglected concept in the complexity-stability debate. *PLoS computational biology*, 14(2):e1005988, 2018.
- Stina Drakare, Jack J Lennon, and Helmut Hillebrand. The imprint of the geographical, evolutionary and ecological context on species–area relationships. *Ecology letters*, 9(2):215–227, 2006.
- Rick Durrett and Simon Levin. Spatial models for species–area curves. *Journal of Theoretical Biology*, 179(2):119–127, 1996.
- Henry Allan Gleason. On the relation between species and area. *Ecology*, 3(2):158–162, 1922.
- BS Goh and LS Jennings. Feasibility and stability in randomly assembled lotka-volterra models. *Ecological Modelling*, 3(1):63–71, 1977.
- Ronen Kadmon and Omri Allouche. Integrating the effects of area, isolation, and habitat heterogeneity on species diversity: a unification of island biogeography and niche theory. *The American Naturalist*, 170(3):443–454, 2007.
- David A Kessler and Nadav M Shnerb. Generalized model of island biodiversity. *Physical Review E*, 91(4):042705, 2015.
- Wade A Leitner and Michael L Rosenzweig. Nested species-area curves and stochastic sampling: a new theory. *Oikos*, pages 503–512, 1997.
- MV Lomolino and MD Weiser. Towards a more general species–area relationship: diversity on all islands, great and small. *Journal of biogeography*, pages 431–445, 2001.

- Robert H MacArthur and Edward O Wilson. An equilibrium theory of insular zoogeography. *Evolution*, pages 373–387, 1963.
- Simeone Marino, Nielson T Baxter, Gary B Huffnagle, Joseph F Petrosino, and Patrick D Schloss. Mathematical modeling of primary succession of murine intestinal microbiota. *Proceedings of the National Academy of Sciences*, 111(1): 439–444, 2014.
- Héctor García Martín and Nigel Goldenfeld. On the origin and robustness of power-law species–area relationships in ecology. *Proceedings of the National Academy of Sciences*, 103(27):10310–10315, 2006.
- Robert M May. Will a large complex system be stable? *Nature*, 238(5364):413–414, 1972.
- Jérôme Mounier, Christophe Monnet, Tatiana Vallaeys, Roger Arditi, Anne-Sophie Sarthou, Arnaud Hélias, and Françoise Irlinger. Microbial interactions within a cheese microbial community. *Applied and environmental microbiology*, 74(1): 172–181, 2008.
- Jacob D O’Sullivan, Robert J Knell, and Axel G Rossberg. Metacommunity-scale biodiversity regulation and the self-organised emergence of macroecological patterns. *Ecology Letters*, 22(9):1428–1438, 2019.
- Susanne Pettersson, Van M Savage, and Martin Nilsson Jacobi. Predicting collapse of complex ecological systems: quantifying the stability–complexity continuum. *Journal of the Royal Society Interface*, 17(166):20190391, 2020.
- Nicolas Picard, Moussa Karembé, and Philippe Birnbaum. Species-area curve and spatial pattern. *Ecoscience*, 11(1):45–54, 2004.
- Frank W Preston. The commonness, and rarity, of species. *Ecology*, 29(3):254–283, 1948.
- Carlos A Serván, José A Capitán, Jacopo Grilli, Kent E Morrison, and Stefano Allesina. Coexistence of many species in random ecosystems. *Nature ecology & evolution*, 2(8): 1237–1242, 2018.
- Arnošt L Šizling and David Storch. Power-law species–area relationships and self-similar species distributions within finite areas. *Ecology letters*, 7(1):60–68, 2004.
- Lewi Stone. The feasibility and stability of large complex biological networks: a random matrix approach. *Scientific reports*, 8(1):1–12, 2018.
- Even Tjørve. Shapes and functions of species–area curves: a review of possible models. *Journal of Biogeography*, 30(6): 827–835, 2003.
- Kristian Trøjelsgaard, Marcos Báez, Xavier Espadaler, Manuel Nogales, Pedro Oromí, Francisco La Roche, and Jens M Olesen. Island biogeography of mutualistic interaction networks. *Journal of Biogeography*, 40(11):2020–2031, 2013.

Figures




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure Captions

Figure 1. Plot showing different assembled communities over time starting from a species pool of 100 species. The clustered lines represent species abundances (left y-axis) for different species as a function of time. The time series is divided into slices of 400 time steps each, where every slice shows dynamics corresponding to a fixed value of the interaction strength parameter (σ). Starting from a value of 2.67×10^{-5} for the first slice, σ increases progressively between consecutive slices by a factor of 1.33. The bold black line traces the number of surviving species at different time slices as shown by the right y-axis. For a given slice, the brief transients eventually settle down at stable equilibrium values, that serve as starting population abundances for the subsequent slice. Note that species go extinct – represented by the corresponding lines converging to zero abundance – as σ increases progressively along the extinction continuum (Pettersson et al., 2020). The interactions, growth rates and carrying capacities are chosen from normal distributions.

Figure 2. Species-area plots generated through 50 realizations of interaction matrix with mean = -1 and variance = 0.2. The semi-log plot (A) shows a better fit with an obvious upper asymptote.

Figure 3. Species area plots demonstrating the better fit of power law SAR for intermediate values of immigration rates. Panels A and B show the fits when $\lambda = 1.5$ for 50 instances of the interaction matrix with mean -1 and variance 0.2. C and D correspond to $\lambda = 0.5$ with the same conditions on the interaction matrix. The solid line illustrates the better fit in both cases.

Figure 4. SAR plots for exponentially distributed interactions with two different rate parameters. All plots correspond to $\lambda = 0.1$ and connectance = 0.05, where the entries of the interaction matrix are chosen randomly as an Erdős-Rényi graph. The semi-log form is better supported for rate parameter = 0.5, as demonstrated by the fits in A and B. Plots C and D show the fits for rate parameter = 0.33, where the power law performs better.

Figure 5. SAR plots for three groups of non-isolated islands within the Solomon Archipelago. These groups differ in how the islands within them were connected during the Pleistocene period. The islands in Group 3 did not have any history of connections. The semi-log relationship shows a good fit to data (A). The R^2 values for the regression lines are 0.989, 0.991 and 0.977 for Group 1, 2 and 3 respectively. The slopes for the different groups are very similar. Panel B shows a clear departure from a power-law relationship for smaller areas. The linear regression lines indicate a good fit for islands larger than one square mile. In particular, the R^2 value for such islands in group 1 is 0.988 from the power-law SAR. This demonstrates that a naive inference could support a power law, in spite of the islands spanning over four orders of magnitude in area (> 1 square mile).

Figure 6. Immigration rates and skewness towards weak interactions determine SAR forms. Semi-log relationship dominates in the absence of immigration. Higher immigration rates from a source pool result in power law relationships but these could shift to semi-log SARs if the relative proportion of weak interactions is increased. S , A and z represent the number of species, area and the scaling law exponent respectively.