Evaluating distributional regression strategies for modelling self-reported sexual age-mixing

Timothy M Wolock^{*,a}, Seth R Flaxman^a, Kathryn A Risher^{b,c}, Tawanda Dadirai^d, Simon Gregson^{b,d}, Jeffrey W Eaton^b

^aDepartment of Mathematics, Imperial College London, London, UK ^bMRC Centre for Global Infectious Disease Analysis, School of Public Health, Imperial College London, London, UK ^cLondon School of Hygiene & Tropical Medicine, London, UK ^dManicaland Centre for Public Health Research, Biomedical Research and Training Institute, Harare,

Zimbabwe

Abstract

The age dynamics of sexual partnership formation determine patterns of sexually transmitted disease transmission and have long been a focus of researchers studying human immunodeficiency virus. Data on self-reported sexual partner age distributions are available from a variety of sources. We sought to explore statistical models that accurately predict the distribution of sexual partner ages over age and sex. We identified which probability distributions and outcome specifications best captured variation in partner age and quantified the benefits of modelling these data using distributional regression. We found that distributional regression with a sinh-arcsinh distribution replicated observed partner age distributions most accurately across three geographically diverse data sets. This framework can be extended with well-known hierarchical modelling tools and can help improve estimates of sexual age-mixing dynamics.

Introduction

Patterns in sexual mixing across ages determine patterns of transmission of sexually transmitted infections (STIs). Consequently, sexual age-mixing has been of great interest to researchers studying the human immunodeficiency virus (HIV) since the beginning of the global epidemic. Anderson et al. (1992) used a model

^{*}Corresponding Author

Email address: t.wolock18@imperial.ac.uk (Timothy M Wolock)

of partnership formation to predict that mixing between young women and older men would amplify the already-substantial effect on HIV on population growth. Garnett & Anderson (1994) used a mathematical model to show that patterns of age-mixing could substantially influence the magnitude and timing of hypothetical epidemic trajectories, while Hallett et al. (2007) demonstrated that delaying sexual debut and increasing age-similar partnerships could reduce an individual's risk of HIV infection in a highly endemic setting.

These modelling studies have been complemented by analyses of survey and population cohort data on age-mixing patterns. Gregson et al. (2002) observed that individuals with older partners were at greater risk of HIV infection. Ritchwood et al. (2016) and Maughan-Brown et al. (2016) found that larger age differences were associated with more risky sexual behaviour in surveys of young South African people. On the other hand, Harling et al. (2014) found that age-disparate relationships were not associated greater risk of HIV acquisition in young women in South Africa.

These results underscore the importance of considering age-mixing dynamics when designing and evaluating HIV prevention strategies, and, consequently, the importance of measuring them accurately. For example, an intervention aiming to prevent new HIV infections among young women could be undermined by high prevalence among older men. Identifying changes in sexual partner age distributions and attributing them to interventions might even be a valuable end by itself, in which case accurate measurement must be complemented by an effective modelling strategy.

Data about sexual partner age-mixing are routinely collected by long-term cohort studies (such as those that comprise the ALPHA Network) and large-scale household surveys (such as the Demographic and Health Surveys) (Reniers et al., 2016; *The DHS Program*, 2021). Typically, these data consist of the respondent's age and sex and the ages of their sexual partners in the last 12 months. These data are highly variable, skewed, and often deviate substantially from conventional parametric distributions, such as the normal distribution or the gamma distribution (Beauclair et al., 2018).

One may consider statistical modelling approaches for the distribution of partner age as a function of respondent age and sex. Some notable previous approaches to modelling partner age distributions include Hallett et al., who used a log-logistic distribution to model partner age differences for women aged 15 to 45 years, assuming that the partner age difference distributions did not vary over respondent age. More recently, as an input to a model of *Chlamydia trachomatis*, Smid et al. (2018) fit skew normal distributions to each age-/sex-specific partner age distribution and used a secondary regression model to smooth the estimated skew normal parameters across respondent age. They observed substantial changes in the estimated skew normal parameters with respect to respondent age. Although this method allows for non-linear variation across respondent age, their two-stage estimation process makes uncertainty propagation complex. Replacing this process with a single "distributional" regression model, in which all distributional parameters (e.g. the location, scale, skewness, etc.) are modelled as functions of data (Kneib & Umlauf, 2017), allows for complex variation across respondent age while still robustly incorporating uncertainty.

More broadly, no previous work has systematically evaluated the wide variety of distributions potentially available to model partner age distributions. These data are skewed, heavy-tailed, and otherwise dissimilar to conventional statistical distributions due to personal preferences, social dynamics, demographic change, and any number of other factors. We were specifically interested in distributions that introduce parameters to control tail weight, which may capture intergenerational mixing that could sustain endemic HIV and STI transmission. This led us to test the ability of the four-parameter "sinh-arcsinh" distribution originally proposed by Jones & Pewsey (2009) to fit to these data.

We hypothesized that integrating the sinh-arcsinh distribution into a distributional modelling framework would allow us to replicate observed partner age distributions more accurately than prior modelling strategies. We tested this theory by comparing a variety candidate strategies, which varied along three dimensions: the parametrisation of the dependent variable, the choice of distribution, and the method for incorporating variability across respondent age and sex.

Methods

We conducted two model comparison experiments to identify which of a set of strategies best replicated partner age distributions. First, in our probability distribution comparison, we identified which of a set of distribution-dependent variable combinations fit best to age-/sex-specific data subsets, and then, in our distributional regression evaluation, we tested whether distributional regression methods could be used to estimate age-/sex-specific partner age distributions by sharing strength across observations. We divided the model comparison into two separate experiments to make the probability distribution comparison as fair as possible (accounting for the possibility that certain distributions would perform particularly well under certain regression specification).

Data

We analysed data on sexual partner age distributions from three sources: the Africa Centre Demographic Information System, a health and demographic surveillance site in uMkhanyakude district, South Africa collected by the African Health Research Institute (AHRI) (Gareta et al., 2021; Gareta et al., 2020a, 2020b), the Manicaland General Population Cohort in Zimbabwe (Gregson et al., 2017), and the 2016-2017 Demographic and Health Survey (DHS) in Haiti (Institut Haïtien de l'Enfance - IHE/Haiti & ICF, 2018).

The AHRI and Manicaland studies are multi-round, open, general population cohort studies designed to measure the dynamics of HIV, sexual risk behaviour, and demographic change in sub-Saharan African settings. We used rounds 1 through 6 of the Manicaland study, collected between 1998 and 2013. The AHRI data we used were collected annually between 2004 and 2018. The 2016-17 Haiti DHS was a large, nationally representative household survey conducted in 2016 and 2017. We did not incorporate the weights associated with the survey into this analysis because our primary interest was in statistical modelling of partner age distribution as a function of respondent age, not producing population representative statistics for the Haitian population.

These data sets consisted of individuals' reports of their own age and sex and the ages of each of their sexual partners from the last year. Let $i \in (1, ..., N)$ index reported partnerships, $a_i \in [15, 64]$ and $s_i \in \{0, 1\}$ be the age and sex

of the respondent in partnership *i* with s = 1 indicating female, and p_i be the age of non-respondent partner in partnership *i*. These questionnaires do not ask specifically about partner sex, but self-reporting of non-heterosexual partnerships in these populations is thought to be low.

Respondents in each of these data sets are disproportionately likely to report that their partners' ages are multiples of five or multiples of five away from their own age, leading to distinct "heaping" in the empirical partner age (or age difference) distributions at multiples of five. We tested the sensitivity of our results to heaping by developing a simple "deheaping" algorithm, applying it to the AHRI data, and running each analysis on the deheaped AHRI data. We present these results in Appendix section "Age heaping."

Probability distribution comparison

To identify the best probability distribution for modelling sexual partner age distributions, we split each data set into 12 subsets by sex and five-year age bin ranging from 20 to 50, resulting in 36 subsets, and fit a number of distribution-dependent variable combinations to each subset.

Distributions. We tested five candidate probability distributions: normal, skew normal, beta, gamma, and sinh-arcsinh. Table 1 summarises the domains, parameters, and probability density functions (PDFs) of these distributions. Because the gamma distribution is always right-skewed and men typically partner with women who are younger than them, we transformed data among male respondents to be right-skewed when using the gamma distribution. Specifically, we multiplied the men's partners' ages by -1 to reflect the distribution horizontally across the y-axis, and added 150 to the reflected ages to ensure that all resulting values were positive. Similarly, the beta distribution is only defined on the interval (0, 1), so, only when using a beta distribution, we scaled all partner ages to be between zero and one using upper and lower bounds of 0 and 150.

The sinh-arcsinh distribution, presented by Jones & Pewsey (2009), is an extension of Johnson's S_U distribution (Johnson, 1949). It has four parameters: location, scale, skewness, and tail weight (denoted, μ , σ , ϵ , and δ respectively), and it can deviate substantially from the normal distribution. Figure 1 plots

Distribution	Parameters	Domain	PDF
Normal	μ (location) $\sigma > 0$ (scale)	R	$\frac{1}{\sigma\sqrt{2\pi}}\exp\left[\frac{-x_z}{2}\right]$
Skew normal	$\mu \text{ (location)} \\ \sigma > 0 \text{ (scale)} \\ \epsilon \text{ (skewness)} $	R	$\frac{2}{\sigma}p(x_z)\Phi(\epsilon x_z)$
Gamma	k > 0 (shape) $\theta > 0$ (scale)	\mathbb{R}^+	$\frac{1}{\Gamma(k)\theta^k} x^{k-1} \exp\left[\frac{-x}{\theta}\right]$
Beta	$\alpha > 0$ (left) $\beta > 0$ (right)	$\mathbb{R}^{(0,1)}$	$\frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha,\beta)}$
Sinh- arcinh	$\mu \text{ (location)} \\ \sigma > 0 \text{ (scale)} \\ \epsilon \text{ (skewness)} \\ \delta > 0 \text{ (tail weight)} $	R	$\frac{1}{\sigma\sqrt{2\pi}}\frac{\delta C_{\epsilon,\delta}(x_z)}{\sqrt{1+x_z^2}}\exp\left[-\frac{S_{\epsilon,\delta}(x_z)^2}{2}\right]$

Table 1: Details of the five distributions tested in this analysis. We define $x_z = (x - \mu)/\sigma$, p(x) to be the standard normal PDF, $\Phi(x)$ to be the standard normal cumulative density function, $S_{\epsilon,\delta}(x) = \sinh(\epsilon + \delta \sinh(x))$, and $C_{\epsilon,\delta} = \cosh(\epsilon + \delta \sinh(x))$.

the density of this distribution with $\mu = 0$ and $\sigma = 1$ for a variety of values of skewness and tail weight.

Dependent variable transformations. We considered the possibility that certain distributions could interact well with particular transformations of the dependent variable (partner age) by testing a set of four potential outcome parametrisations. For example, if *X* is a positive-valued, right-skewed random variable, then assuming log *X* is normally distributed might be more effective than assuming that *X* itself is normal.

Let y_i be the dependent variable value for partnership *i*, and let a_i and p_i be the respondent age and partner age of partnership *i*, respectively. We tested the following dependent variables:

- 1. **Linear age**: $y_i = p_i$. This is untransformed partner age, included as a baseline. It has the undesirable quality of being able to predict negative ages.
- 2. Age difference: $y_i = p_i a_i$. If changes in expected partner age are consistent across respondent age then this variable would be more consistent across respondent age than the linear age. This parametrisation also allows for negative partner age predictions.



Figure 1: The sinh-arcsinh density with $\mu = 0$, $\sigma = 1$, and a variety of assumptions about ϵ and δ

- 3. **Log-age**: $y_i = \log p_i$. We can use a log link function to ensure that our predictions will be positive-valued.
- 4. Log-ratio: $y_i = \log(p_i/a_i)$. Finally, we can combine the link function and differencing approaches by modelling the log of the ratio of partner to respondent age. This variable will only produce positive predictions and should exhibit the same relative stability as the age difference variable.

Because the gamma and beta distributions are not defined on the entire real line, we only fit them with the linear age dependent variable with the previously discussed transformations.

To identify which distribution-dependent variable combination best modelled the characteristics of sexual partner age distributions, we stratified each of our three data sets by sex and five-year age bin from 20-24 through 45-49. We fit every viable distribution-dependent variable combination to all 36 subsets independently. Given that we fit only the linear age dependent variable to the gamma and beta distributions, comprising a total of 504 models (14 per data set). We fit each model using the brms R package (Bürkner, 2018), defining custom families as necessary.

Distributional regression evaluation

Given a probability distribution that accurately replicated the non-Gaussian characteristics of partner age distributions, we tested whether or not distributional regression would allow us to pool data across age and sex without sacrificing fit. In distributional regression, we make all of our distributional parameters, not just the mean, functions of data (Kneib & Umlauf, 2017). Taking conventional Bayesian regression as an example, we have

$$y_i \sim \mathbf{N}(\mu_i, \sigma)$$

 $\mu_i = \beta \mathbf{X}_i,$

where β and log σ are free parameters. There is an explicit assumption in this model that the standard deviation of the generating distribution is constant across all observations. We can use distributional regression to relax this assumption, making σ a function of data:

$$\begin{array}{rcl} y_i & \sim & \mathrm{N}(\mu_i, \sigma_i) \\ \mu_i & = & \beta^{\mu} \mathbf{X}_i^{\mu} \\ \log \sigma_i & = & \beta^{\sigma} \mathbf{X}_i^{\sigma}, \end{array}$$

where β^{μ} and β^{σ} are now our free parameters. Note that we have not assumed that $\mathbf{X}^{\mu} = \mathbf{X}^{\sigma}$. If \mathbf{X}^{σ} is a column of ones, this model is identical to the conventional case. This approach increases the complexity of the model and requires more data, but, based on previously described characteristics of how the distribution of partnership age distribution changes with age, even a simple model for our distributional parameters could yield large improvements.

In this case, we used a sinh-arcsinh distribution and specified a model for each of its four parameters. We fit a series of increasingly complex distributional regression specifications to the three data sets using brms (Bürkner, 2018), which has deep support for distributional regression.

- 1. **Conventional**: linear age-sex interaction for location and constants for all three higher-order parameters
- 2. **Distributional 1**: linear age-sex interaction for location and independent age and sex effects for all other parameters
- 3. **Distributional 2**: linear age-sex interactions for all four parameters
- 4. **Distributional 3**: sex-specific spline with respect to age for location and linear age-sex interactions for all other parameters
- 5. Distributional 4: sex-specific splines with respect to age for all four pa-

Table 2: Summary	of five n	nodels fit in	ı this a	nalysis.
------------------	-----------	---------------	----------	----------

Model	Distributional?	Location	Other parameters
Conventional	No	Age-sex interaction	Constant
Distributional 1	Yes	Age-sex interaction	Age and sex effects
Distributional 2	Yes	Age-sex interaction	Age-sex interaction
Distributional 3	Yes	Sex-specific splines	Age-sex interaction
Distributional 4	Yes	Sex-specific splines	Sex-specific splines

rameters

Table 2 describes all five models. By fitting a wide set of specifications, we hoped to assess whether the additional complexity incurred by distributional regression was valuable. More detailed descriptions of each model are available in the "Model specification details" section of the Appendix.

Model comparison

Across both analyses, we used two metrics to measure model fit. First, we calculated the expected log posterior density (ELPD), which estimates the density of the model at a new, unobserved data point (Vehtari et al., 2017). In cases where we wanted to compare across dependent variables, we multiplied the posterior densities of any variables resulting from non-linear transformations of observed partner ages by the Jacobians of the transformations. For example, if our observation model was defined on the log-age dependent variable $y_i = \log p_i$, we divided the posterior density by p_i . We used the loo R package (Vehtari et al., 2020) to calculate ELPD values.

To measure the ability of our models to replicate partner age distributions in an objective and interpretable way, we found the root mean squared error (RMSE) between the observed and posterior predictive quantiles. We calculated quantiles from 10 to 90 in increments of 10 by age bin and sex in the data and in the posterior predictions, and found the error in model prediction of each quantile. This measure tells how well our model predicts the entire distribution in the same units as our predictions. It is equivalent to finding the mean squared or median absolute distance from the line of equality in a quantile-quantile (QQ) plot.

Software

We conducted all of these analysis using the R programming language (R Core Team, 2020) and the brms library (Bürkner, 2018). We used the loo library to estimate all ELPDs (Vehtari et al., 2020), and produced all plots in this paper with the ggplot2 library (Wickham, 2016). We cannot provide the data we used for this analysis, but we do provide code and data for an simulated case on GitHub (https://github.com/twolock/distreg-illustration).

Results

The AHRI data included 77,619 partnerships, Manicaland had 58,676, and the Haiti DHS had 12,447. As an illustrative example of the distribution of partner ages, Figure 2 presents histograms of reported partner ages among women aged 35-39 for each of our three data sets. Figure 3 shows the sex- and age bin-specific empirical moments for the three data sets. Mean partner age increased with respondent age consistently for both sexes across all three data sets: among women, mean partner age increased by 26.0, 22.7, and 23.7 years in the AHRI data, Haiti DHS data, and Manicaland data, respectively, between age bins 20-24 and 45-49. However, higher order moments were less consistent: the standard deviation of women's partners' ages changed by 2.3, 0.5, and 3.5 years in the AHRI data, Haiti DHS data, and Manicaland data, respectively.

Within each data set, there is systematic variation across sex. For example, the standard deviation of partner ages in the Haiti DHS increased by 2.5 years among men and only by 0.5 years among women. These summary statistics illustrate the heterogeneity of partner age distributions across age and sex.

Probability distribution comparison

To identify the probability distribution that most accurately described the variation in sexual partner age distributions, we first determined the dependent variable with the highest ELPD for each distribution-dependent variable combination. Figure 4 illustrates each probability distribution's best fit to AHRI data among women aged 35-39 with each of the best distribution-specific dependent



Figure 2: Observed partner age distributions among women aged 34 years in all three data sets.

Table 3: Share of subsets in which each dependent variable yields the highest ELPD given each probability distribution (excluding deheaped AHRI data).

Variable	Normal	Skew normal	Sinh-arcsinh
Age difference	22.2%	25.0%	16.7%
Linear age	8.3%	5.6%	16.7%
Log-age	19.4%	41.7%	30.6%
Log-ratio	50.0%	27.8%	36.1%

variables. Results for all 36 data subsets and the 12 deheaped subsets are in Appendix section "Full results."

The best dependent variable varied across data subset and probability distribution. Table 3 provides the share of data sets for which each dependent variable has the highest ELPD given each distribution. The log-ratio dependent variable was best in 50.0% of subsets with a normal distribution, but it was best in only 27.8% of subsets with a skew normal distribution. The dependent variable that was best in a plurality of subsets in each probability distribution (i.e. the variable with the highest percentage in each column of Table 3) used a log link function. We restricted all remaining comparisons to each distribution-subset combination's best dependent variable.

The sinh-arcsinh distribution had the highest ELPD in 35 of 36 data subsets (98%). In 29 of the 35 (83%) cases in which the sinh-arcsinh provided the highest



Figure 3: Observed means, variances, skewnesses, and kurtoses of partner age by five-year age bin and sex in all three data sets



Figure 4: Observed partner age distributions (grey bars) and posterior predictive partner age distributions (lines) for each probability distribution among women aged 35-39 in the AHRI data set. Posterior predictive distributions come from fitting each age bin/sex combination independently.

Distribution	AHRI	Haiti 2016-17 DHS	Manicaland
ELPD			
Gamma	-14847.2	-2917.9	-13152.8
Beta	-14748.0	-2896.5	-13003.5
Normal	-14593.7	-2868.4	-12856.8
Skew normal	-14505.1	-2854.0	-12778.5
Sinh-arcsinh	-14312.5	-2839.5	-12625.8
QQ RMSE			
Gamma	0.83	0.82	0.95
Beta	0.99	0.82	1.11
Normal	0.82	0.68	0.97
Skew normal	0.77	0.65	0.85
Sinh-arcsinh	0.36	0.37	0.44

Table 4: Model comparison metrics averaged across all data subsets for all three data sets. Higher ELPD values indicate better fit. Lower QQ RMSE values indicate more accurate prediction of empirical quantiles. Bolded rows are best across all three data sets.

ELPD, the absolute value of the ratio of the difference between the two best ELPDs and the estimated standard error of the difference was greater than 2, indicating that the sinh-arcsinh distribution was significantly better than the alternatives in the majority of cases. In one case, men aged 20-24 in the Haiti DHS, the skew normal distribution resulted in a slightly higher ELPD than the sinh-arcsinh distribution, but the standard error of the difference was greater than the difference. These results were not affected by deheaping the data (Appendix section "Age heaping").

To summarise each distribution's performance, we calculated the average ELPD and QQ RMSE across the three data sets (Table 4). The sinh-arcsinh distribution had the highest average ELPD and lowest average QQ RMSE in all three data sets. The sinh-arcsinh distribution was, on average, able to predict the empirical quantiles of each data set within half a year of accuracy (0.36, 0.37, and 0.44 years for the AHRI, Haiti DHS, and Manicaland data, respectively). Overlaid QQ plots and non-aggregated tables of ELPD differences and QQ RMSEs are presented in the "Full results" section of the Appendix.

Table 5: ELPD and QQ RMSE values for all five distributional regression models fit to each data set. The models increase in complexity from Conventional Regression to Distributional Model 4. Bolded ELPD values are more than two standard errors higher than the next best value in the column. Bolded QQ RMSE values are lowest in their column.

Model	AHRI	Haiti 2016-17 DHS	Manicaland
ELPD			
Conventional	52689.2	4777.8	21011.3
Distributional 1	54335.2	5140.8	23192.5
Distributional 2	54794.8	5138.7	23472.1
Distributional 3	55534.2	5196.7	24313.7
Distributional 4	55841.9	5207.6	24516.1
QQ RMSE			
Conventional	1.30	1.33	2.05
Distributional 1	1.15	0.98	1.89
Distributional 2	1.21	0.99	1.80
Distributional 3	0.93	0.91	1.34
Distributional 4	0.66	0.84	1.04

Distributional regression evaluation

We fit all five distributional regression specifications to all three of our data sets with sinh-arcsinh distributions and log-ratio dependent variables and compared the ELPDs and QQ RMSEs as before (provided in Table 5). Across all three data sets, the most complex distributional model (Distributional 4) had the highest ELPD and lowest QQ RMSE. When fit to the AHRI and Manicaland data sets (but not for the Haiti DHS), the most complex distributional model was a least two standard errors better than the next best model. Notably, the largest ELPD improvements came from moving from conventional regression (Conventional) to the simplest distributional model (improvements of 1646.0 units, 361.0 units, and 2181.2 units in the AHRI, Haiti DHS, and Manicaland data, respectively). Full tables are available in the "Full results" section of the Appendix.

Figure 5 shows the posterior predictive distributions from the conventional regression model and the most complex distributional model among men aged 16 years, 24 years, and 37 years in the AHRI data to illustrate the effect of distributional regression. Not only does the distributional model capture the high peak in the youngest age more accurately, but it also allows the variance of the distributions to change appropriately (beyond the change that naturally results from the log link function).



Figure 5: Observed partner age distributions (grey bars) and posterior predictive partner age distributions (lines) for conventional regression and the most complex distributional model among men aged 16, 24, and 37 years in the AHRI data set. Posterior predictive distributions come from regression models fit to the entire AHRI data set.

Figure 6 illustrates posterior summaries among men and women in the AHRI data for all four distributional parameters for the conventional regression model, the simplest distributional model, and the most complex distributional model. The red estimates (Conventional Regression) of the three higher order parameters were constant across age and sex, whereas the blue estimates (Distributional Model 1) included independent, linear age and sex effects. The orange estimates (Distributional Model 4) were generated sex-specific splines with respect to age, allowing for flexible variation across age and sex.

The third row of plots in Figure 6, which corresponds to the skewness parameter, illustrates the impact of incorporating sex and age effects into the model. The conventional regression model estimated that neither the distrbution for men nor women exhibited much skewness; the estimated parameter value was -0.05 (95% UI: -0.06 to -0.05) regardless of age, with 0.0 corresponding to perfect symmetry. However, when we allowed independent age and sex effects in Distributional Model 1, we estimated that at age 15, women's skewness was -0.26 (95% UI: -0.27 to -0.25) and men's was 0.11 (95% UI: 0.10 to 0.12).

The most complex model (Distributional Model 4) inferred sex-specific, nonlinear variation with respect to age in all four distributional parameters. The non-linearity was particularly dramatic in the scale parameter among men. The scale value began at 0.05 (95% UI: 0.05 to 0.06) among 15-year-olds, peaked among 37-year-olds at 0.11 (95% UI: 0.10 to 0.11), and decreased back down to 0.05 (95% UI: 0.04 to 0.06) at age 64.

Finally, Figure 7 presents inferred distributional parameters from Distributional Model 4 for both men and women for all three data sets. Based on those plots, the flexible model was justified for most distributional parameters in all three data sets. Were we to continue developing these models, this plot suggests that skewness might only need linear, sex-specific effects with respect to age. Interestingly, the 2016-2017 Haiti DHS and Manicaland estimates exhibit similar patterns across all four parameters, despite the different socio-cultural contexts surrounding partnerships in the two populations. We also note that the DHS does not collect data on adults aged 50 years and older, so our estimates in Haiti from age 50 to age 64 are purely extrapolated.



Figure 6: Estimated sinh-arcsinh distributional parameters from the conventional regression model, and distributional models 1 and 4 fit to the AHRI data. "Conventional" assumes no variation across age and sex, "Distributional 1" allows for independent age and sex effects, and "Distributional 4" includes sex-specific splines with respect to age.



Figure 7: Estimated sinh-arcsinh distributional parameters for Distributional Model 4 fit to the three main data sets.

Discussion

We found that the sinh-arcsinh distribution reproduced observed sexual partner age distributions better than a number of other possible distributional assumptions across age and sex in three distinct data sets. We integrated this finding into a distributional regression framework using existing statistical modelling software. Even the simplest distributional regression in our set of candidate models far outperformed conventional regression, in which all moments except the first are estimated as constants. Our most complex distributional model fit better than all other models in all three data sets, suggesting that modelling these data benefits from the additional complexity.

These results indicate that distributional regression models with sinh-arcsinh distributions can accurately replicate age-/sex-specific sexual partner age distributions. This approach presents a number of advantages over previous methods. First, like Smid et al., it allows a unique distribution for every age-sex combination. As Figure 3 illustrates, partner age distributions can exhibit substantial, systematic variation across age and sex in any of the first four moments, so we must consider modelling strategies that allow for such variation. Second, distributional regression offers a principled method to propagate uncertainty through this estimation process.

Finally, distributional regression implemented through brms provides access to a deep set of hierarchical modelling tools that could enable estimation in a variety of low-data settings. We evaluated a small set of relatively simple distributional models in this work, but, theoretically, each distributional parameter could have its own, arbitrarily complex hierarchical regression model. Using these tools, one could estimate unique partner age distributions across levels of stratification that are substantively interesting but do not provide sufficient sample size for independent estimation (e.g. study sites or geographic areas).

We have identified several limitations in this approach. First, the amount of data required to produce usefully precise estimates is not tested. Each additional distributional parameter introduces model parameters, so this method is more complex than conventional regression. The sinh-arcsinh distribution did fail to produce the highest ELPD in our smallest data subset (N = 170), but it was not significantly worse than the best distribution. More importantly, by integrating

these data into a distributional modelling framework, we gain the ability to impose structure on these parameters, which could easily offset the cost of any additional model parameters.

Interpreting the inferred model parameters in sinh-arcsinh regression can also be difficult. Whereas conventional regression estimates the effects of covariates on expected values, the sinh-arcsinh distribution is parametrised in terms of a location parameter. This parameter correlates closely with the central tendency of the distribution, but it is not strictly equal to the mean. We can reparametrise the distribution so that we estimate a mean (and therefore effects of covariates on the expected value), but it is not currently possible in the probabilistic programming software that underlies brms.

Third, our analysis assumed that we were operating at a level of stratification at which partnerships are basically comparable, but any number of factors could lead to fundamentally different partner age distributions. For example, we did not control for whether the partnership was same-sex or the type of the partnership (married, casual, etc.). That said, our distributional framework would allow us to incorporate data on any of those factors directly into the model.

Despite these limitations, we believe that the strategy we present will work well in future projects that require estimates of partner age distributions. We plan to use these methods to produce age-mixing matrices to inform epidemic models of HIV, but there are a number of additional directions that could be explored. We are specifically interested in leveraging the spatio-temporal structure of the survey data used here. Hierarchical mapping exercises with household survey data are increasingly common in epidemiology, but estimating spatially varying partner age distributions would require an evaluation of how best to model higher order moments over space. We would, for example, need to consider how the variance of partner age distributions varies by urbanicity.

Similarly, population-based studies typically collect far more detailed information on partnerships than we took advantage of here. Relationship type is a key confounder of the association between respondent age and partner age (that we ignored for the purposes of our experiments). We might expect the age distributions of casual partners to vary substantially from those of long-term cohabiting partners. Because we have built our model in a pre-existing regression framework, incorporating new covariates into any of the distributional regression specifications should be simple.

We believe that our framework offers a flexible, accurate, and robust method for smoothing and interpolating sexual partner age distributions, but these methods are not specific to partner age distributions. The sinh-arcsinh distribution is relatively easy to implement without incurring high computational cost, so it could be applied in many settings. Even without the distributional regression framework we have used here, allowing the third and fourth moments of the distribution to vary from the "default" normal values could be valuable across a variety of applications.

Distributional regression is also underutilised in social science applications. We often work with large surveys that would comfortably support models for higher order parameters. Data requirements will vary by application and model, but, as we have shown here, even a simple distributional model can improve fit and avoid biasing estimates.

Acknowledgements

JWE, SG, and KAR acknowledge funding support from the Bill & Melinda Gates Foundation. JWE and SG acknowledge funding support from the MRC Centre for Global Infectious Disease Analysis (reference MR/R015600/1), jointly funded by the UK Medical Research Council (MRC) and the UK Foreign, Commonwealth & Development Office (FCDO), under the MRC/FCDO Concordat agreement and is also part of the EDCTP2 programme supported by the European Union. JWE also acknowledges funding support from National Institute of Allergy and Infectious Disease of the National Institutes of Health under award number R01AI136664. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. SRF acknowledges funding support from the EPSRC (EP/V002910/1). TMW's work is funded by the Imperial College President's PhD Scholarship. We thank all of the people who participated in the three studies that shared their data with us, as well as the survey and data management teams, without whom this work would not be possible.

Competing interests

None declared.

References

- Anderson, R. M., May, R. M., Ng, T. W., & Rowley, J. T. (1992). Age-dependent choice of sexual partners and the transmission dynamics of HIV in subsaharan africa. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 336(1277), 135–155. https://doi.org/10.1098/rstb.1992. 0052
- Beauclair, R., Hens, N., & Delva, W. (2018). The role of age-mixing patterns in HIV transmission dynamics: Novel hypotheses from a field study in cape town, south africa. *Epidemics*, 25, 61–71. https://doi.org/10.1016/j.epidem. 2018.05.006
- Bürkner, P.-C. (2018). Advanced bayesian multilevel modeling with the r package brms. *The R Journal*, *10*(1), 395–411. https://doi.org/10.32614/RJ-2018-017
- Gareta, D., Baisley, K., Mngomezulu, T., Smit, T., Khoza, T., Nxumalo, S., Dreyer, J., Dube, S., Majozi, N., Ording-Jesperson, G., Ehlers, E., Harling, G., Shahmanesh, M., Siedner, M., Hanekom, W., & Herbst, K. (2021). Cohort profile update: Africa centre demographic information system (ACDIS) and population-based HIV survey. *International Journal of Epidemiology*, 50(1), 33–34. https://doi.org/10.1093/ije/dyaa264
- Gareta, D., Dube, S., & Herbst, K. (2020a). AHRI.PIP.men's general health.all.release 2020-07. Africa Health Research Institute (AHRI). https://doi.org/10.23664/ AHRI.PIP.RD04-99.MGH.ALL.202007
- Gareta, D., Dube, S., & Herbst, K. (2020b). AHRI.PIP.women's general health.all.release 2020-07. Africa Health Research Institute (AHRI). https://doi.org/10.23664/ AHRI.PIP.RD03-99.WGH.ALL.202007
- Garnett, G. P., & Anderson, R. M. (1994). Balancing sexual partnerships in an age and activity stratified model of HIV transmission in heterosexual populations. *IMA Journal of Mathematics Applied in Medicine and Biology*, 11(3), 161–192. https://doi.org/10.1093/imammb/11.3.161
- Gregson, S., Mugurungi, O., Eaton, J., Takaruza, A., Rhead, R., Maswera, R., Mutsvangwa, J., Mayini, J., Skovdal, M., Schaefer, R., Hallett, T., Sherr, L.,

Munyati, S., Mason, P., Campbell, C., Garnett, G. P., & Nyamukapa, C. A. (2017). Documenting and explaining the HIV decline in east zimbabwe: The manicaland general population cohort. *BMJ Open*, 7(10), e015898. https://doi.org/10.1136/bmjopen-2017-015898

- Gregson, S., Nyamukapa, C. A., Garnett, G. P., Mason, P. R., Zhuwau, T., Caraël, M., Chandiwana, S. K., & Anderson, R. M. (2002). Sexual mixing patterns and sex-differentials in teenage exposure to HIV infection in rural zimbabwe. *Lancet* (*London, England*), 359(9321), 1896–1903. https://doi.org/10.1016/ S0140-6736(02)08780-9
- Hallett, T. B., Gregson, S., Lewis, J. J. C., Lopman, B. A., & Garnett, G. P. (2007). Behaviour change in generalised HIV epidemics: Impact of reducing crossgenerational sex and delaying age at sexual debut. *Sexually Transmitted Infections*, 83 Suppl 1, i50–54. https://doi.org/10.1136/sti.2006.023606
- Harling, G., Newell, M.-L., Tanser, F., Kawachi, I., Subramanian, S., & Bärnighausen, T. (2014). Do age-disparate relationships drive HIV incidence in young women? Evidence from a population cohort in rural KwaZulu-natal, south africa. *Journal of Acquired Immune Deficiency Syndromes* (1999), 66(4), 443–451. https://doi.org/10.1097/QAI.00000000000198
- Institut Haïtien de l'Enfance IHE/Haiti, & ICF. (2018). Haiti enquête mortalité, morbidité et utilisation des services 2016-2017 - EMMUS-VI [dataset]. IHE/Haiti, ICF [Producers]. http://dhsprogram.com/pubs/pdf/FR326/FR326.pdf
- Johnson, N. L. (1949). Systems of frequency curves generated by methods of translation. *Biometrika*, 36(1), 149. https://doi.org/10.2307/2332539
- Jones, M. C., & Pewsey, A. (2009). Sinh-arcsinh distributions. *Biometrika*, 96(4), 761–780. https://doi.org/10.1093/biomet/asp053
- Kneib, T., & Umlauf, N. (2017). A primer on bayesian distributional regression (Working Paper No. 2017-13). Working Papers in Economics; Statistics. https://www.econstor.eu/handle/10419/180164
- Maughan-Brown, B., Evans, M., & George, G. (2016). Sexual behaviour of men and women within age-disparate partnerships in south africa: Implications for young women's HIV risk. *PLOS ONE*, 11(8), e0159162. https://doi.org/ 10.1371/journal.pone.0159162
- R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. https://www.R-project.org
- Reniers, G., Wamukoya, M., Urassa, M., Nyaguara, A., Nakiyingi-Miiro, J., Lutalo, T., Hosegood, V., Gregson, S., Gómez-Olivé, X., Geubbels, E., Crampin, A. C., Wringe, A., Waswa, L., Tollman, S., Todd, J., Slaymaker, E., Ser-

wadda, D., Price, A., Oti, S., ... Zaba, B. (2016). Data resource profile: Network for analysing longitudinal population-based HIV/AIDS data on africa (ALPHA network). *International Journal of Epidemiology*, 45(1), 83–93. https://doi.org/10.1093/ije/dyv343

- Ritchwood, T. D., Hughes, J. P., Jennings, L., MacPhail, C., Williamson, B., Selin, A., Kathleen, K., Gómez-Olivé, F. X., & Pettifor, A. (2016). Characteristics of age-discordant partnerships associated with HIV risk among young south african women (HPTN 068). *Journal of Acquired Immune Deficiency Syndromes* (1999), 72(4), 423–429. https://doi.org/10.1097/QAI.00000000000088
- Smid, J. H., Garcia, V., Low, N., Mercer, C. H., & Althaus, C. L. (2018). Age difference between heterosexual partners in britain: Implications for the spread of chlamydia trachomatis. *Epidemics*, 24, 60–66. https://doi.org/10. 1016/j.epidem.2018.03.004
- *The DHS program.* (2021). USAID. https://dhsprogram.com/data/
- Vehtari, A., Gabry, J., Magnusson, M., Yao, Y., Bürkner, P.-C., Paananen, T., & Gelman, A. (2020). *Loo: Efficient leave-one-out cross-validation and WAIC for bayesian models*. https://mc-stan.org/loo
- Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, 27(5), 1413–1432. https://doi.org/10.1007/s11222-016-9696-4
- Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York. https://ggplot2.tidyverse.org



Figure 8: Illustration of the effect of the deheaping algorithm on women aged exactly 24 years in the AHRI data. Dark grey bars correspond to ages identified as potentially heaped (multiples of five away from 24). The red line is the expected count of observations estimated by excluding any potentially heaped ages.

Appendix

Age heaping

Respondents in each of these data sets are disproportionately likely to report that their partners' ages are multiples of five or multiples of five away from their own age, leading to distinct "spikes" in the empirical partner age (or age difference) distributions at multiples of five. The left panel of Figure 8 illustrates this phenomenon among women aged 24 years in the AHRI data. These spikes, widely referred to as "heaping," could bias our results towards certain probability distributions, so we developed a simple deheaping algorithm, applied it to the AHRI data.

To account for the possibility that heaping affected the results, we developed a simple deheaping algorithm and treated the deheaped AHRI data as a fourth data set. Due to the structure of the questionnaire ("how many years older or younger is your partner than you?"), the AHRI partner age data exhibit strong heaping on partner ages that are multiples of five years from the respondent's age. For example, among women aged 24 years, we observe far more partners aged exactly 29 years than expected.

Let $n_{s,a,p}$ be the number of observed partnerships with $s_i = s$, $a_i = a$, and $p_i = p$.

Fixing age to be *a* and sex to be *s*, we can find the expected count at partner age p, $\hat{n}_{s,a,p}$ by fitting a Nadaraya-Watson estimator to all ordered pairs $(p, n_{s,a,p})$ such that p - a is *not* a multiple of five. We can then find the positive-valued excess counts at all p such that p - a is a multiple of five:

$$e_{s,a,p} = \max(n_{s,a,p} - \hat{n}_{s,a,p}, 0).$$

This quantity, $e_{s,a,p}$, is what the Nadaraya-Watson estimator has identified as number of heaped observations. Fixing p^* to be a partner age such that $(p^* - a) \mod 5 \equiv 0$, we assume that all of the excess mass at p^* will be allocated to the four partner ages on either side of p^* . We find the share of e_{s,a,p^*} to be allocated to each of $(p^* - 2, ..., p^* + 2)$, denoted $b_{s,a,p}$, as

$$b_{s,a,p} = \frac{n_{s,a,p}}{\sum_{i=-2}^{2} n_{s,a,p^{\star}+i}},$$

substituting in \hat{n}_{s,a,p^*} for $n_{s,a,p}$ wherever applicable. Finally, we find the number of individuals to be reassigned from p^* to each p within two years of y^* as $d_{s,a,p} = b_{s,a,p} \cdot e_{s,a,p^*}$. Note that each partner age can only "receive" partnerships from its nearest multiple of five and that each multiple of five can only "send" partnerships to itself and the four partner ages on either side of it. For each ywithin two years of y^* , we randomly select $\lfloor d_{s,a,p} \rceil$ individuals to move from p^* to p. We apply this method for both sexes and all respondent ages with at least two observations separately.

Figure 8 illustrates the effect of this process on data among women aged 24 in the AHRI data. Despite its simplicity, the deheaping algorithm seems to produce distributions that should be sufficiently plausible for the purposes of this experiment. If our results differed substantially between the original and deheaped AHRI data, we would need to consider the possibility that our results could be an artefact of heaping.

This method is quite simple, but it seems to work reasonably well on the AHRI data. Regardless, we do not need a perfect deheaping algorithm for this application; we just need one that will give us a *plausibly* deheaped version of the AHRI data. If the results differ drastically between the heaped and



Figure 9: Observed sexual partner age distributions among women in the AHRI data. The left panel is original data, and the right panel is the same data set after deheaping age differences from multiples of five.

deheaped data sets (*i.e.* if one probability distribution works perfectly only on the deheaped data), then we will know that our results are sensitive to irregularities in the data.

Results

Figure 9 shows the presence of age heaping among women in the AHRI data, as well as the effects of our deheaping algorithm. Visible diagonal lines indicate that women were disproportionately likely to report that the difference between their partner's age and their own age was a multiple of five. Heaping to partner ages (not partner age differences) would manifest as horizontal lines. As we can see in the right panel, the deheaping procedure resolves the majority of the heaping. We cannot validate the algorithm, but for the purposes of this experiment, simply producing plausibly deheaped age distributions should be sufficient.

Table 6 provides ELPD and QQ RMSE values for all five regression models fit to

Table 6: ELPD and QQ RMSE values for all five models fit to deheaped AHRI data The models increase in complexity from Conventional Regression to Distributional Model 4. Bolded ELPD values are more than two standard errors higher than the next best value in the column. Bolded QQ RMSE values are lowest in their column.

Model	AHRI Deheaped
ELPD	
Conventional	55296.2
Distributional 1	57097.4
Distributional 2	57503.7
Distributional 3	58219.2
Distributional 4	58504.0
QQ RMSE	
Conventional	1.26
Distributional 1	1.06
Distributional 2	1.14
Distributional 3	0.92
Distributional 4	0.62

the deheaped AHRI data. As with the heaped AHRI data, the most complex distributional model had the highest ELPD (58504.0). From these results, we conclude that the presence of heaping in the three main data sets is unlikely to have substantially altered the results of this analysis.

Model specification details

We modelled the log-ratio dependent variable using the four-parameter sinharcsinh distribution:

$$\begin{array}{rcl} y_i &\sim & \sinh(\mu_i, \sigma_i, \epsilon_i, \delta_i) \\ \mu_i &= & \beta^{\mu} \mathbf{X}_i^{\mu} \\ \log \sigma_i^{\star} &= & \beta^{\sigma} \mathbf{X}_i^{\sigma} \\ \epsilon_i &= & \beta^{\epsilon} \mathbf{X}_i^{\epsilon} \\ \log \delta_i &= & \beta^{\delta} \mathbf{X}_i^{\delta} \\ \sigma_i &= & \sigma_i^{\star} \delta_i, \end{array}$$

where β^{μ} , β^{σ} , β^{ϵ} , and β^{δ} are free parameters. We placed essentially arbitrary shrinkage priors on all coefficients:

$$\beta^{\mu}, \beta^{\sigma}, \beta^{\epsilon}, \beta^{\delta} \sim \mathrm{N}(0, 5).$$

First, we fit a conventional regression, in which only the location parameter, μ , is a function of data. Specifically, we allowed for linear sex and age effects and a linear interaction between respondent sex and age (s_i and a_i , respectively) in the model of μ :

$$\begin{aligned} \mathbf{X}_i^{\mu} &= (1, s_i, a_i, s_i \cdot a_i) \\ \mathbf{X}_i^{\sigma}, \mathbf{X}_i^{\epsilon}, \mathbf{X}_i^{\delta} &= (1). \end{aligned}$$

In the second model, we allowed the three higher order distributional parameters to vary by age and sex:

$$\begin{aligned} \mathbf{X}_{i}^{\mu} &= (1, s_{i}, a_{i}, s_{i} \cdot a_{i}) \\ \mathbf{X}_{i}^{\sigma}, \mathbf{X}_{i}^{\varepsilon}, \mathbf{X}_{i}^{\delta} &= (1, s_{i}, a_{i}). \end{aligned}$$

In the third model, all four distributional parameters had age, sex, and age-sex interaction effects:

$$\mathbf{X}_{i}^{\mu}, \mathbf{X}_{i}^{\sigma}, \mathbf{X}_{i}^{\epsilon}, \mathbf{X}_{i}^{\delta} = (1, s_{i}, a_{i}, s_{i} \cdot a_{i})$$

To allow for the possibility of non-linear variation with respect to age in the fourth model, we modelled the location parameter using sex-specific natural splines on age:

$$\begin{aligned} \mathbf{X}_{i}^{\mu} &= (1, s_{i}, \phi_{1}(a_{i}), ..., \phi_{K}(a_{i}), s_{i} \cdot \phi_{1}(a_{i}), ..., s_{i} \cdot \phi_{K}(a_{i})) \\ \mathbf{X}_{i}^{\sigma}, \mathbf{X}_{i}^{\epsilon}, \mathbf{X}_{i}^{\delta} &= (1, s_{i}, a_{i}, s_{i} \cdot a_{i}), \end{aligned}$$

where *K* is the number of columns in the spline design matrix. By including a second set of basis function values that are multiplied by s_i , we are estimating an additional, female-specific trend with respect to age.

Finally, we fit a fifth model, in which all four distributional parameters were

modelled as sex-specific splines with respect to age:

$$\mathbf{X}_{i}^{\mu}, \mathbf{X}_{i}^{\sigma}, \mathbf{X}_{i}^{\epsilon}, \mathbf{X}_{i}^{\delta} = (1, s_{i}, \phi_{1}(a_{i}), ..., \phi_{K}(a_{i}), s_{i} \cdot \phi_{1}(a_{i}), ..., s_{i} \cdot \phi_{K}(a_{i})).$$

Full Results

Supplementary Figures



Figure 10: Overlaid quantile-quantile (QQ) plots for each probability distribution's best fit to data in all three main data sets. Presented quantiles range from 10th to 90th in increments of 10. Lines closer to the line of equality indicate better fit to empirical quantiles



Figure 11: Observed partner age distributions (grey bars) and posterior predictive partner age distributions (lines) for each probability distribution among women in the AHRI data set. Here, we plot the posterior predicitve distribution associated with each distribution's highest-ELPD dependent variable.



Figure 12: Observed partner age distributions (grey bars) and posterior predictive partner age distributions (lines) for each probability distribution among men in the AHRI data set. Here, we plot the posterior predicitve distribution associated with each distribution's highest-ELPD dependent variable.



Figure 13: Observed partner age distributions (grey bars) and posterior predictive partner age distributions (lines) for each probability distribution among women in the AHRI Deheaped data set. Here, we plot the posterior predicitve distribution associated with each distribution's highest-ELPD dependent variable.



Figure 14: Observed partner age distributions (grey bars) and posterior predictive partner age distributions (lines) for each probability distribution among men in the AHRI Deheaped data set. Here, we plot the posterior predicitve distribution associated with each distribution's highest-ELPD dependent variable.



Figure 15: Observed partner age distributions (grey bars) and posterior predictive partner age distributions (lines) for each probability distribution among women in the Haiti 2016-17 DHS data set. Here, we plot the posterior predicitve distribution associated with each distribution's highest-ELPD dependent variable.



Figure 16: Observed partner age distributions (grey bars) and posterior predictive partner age distributions (lines) for each probability distribution among men in the Haiti 2016-17 DHS data set. Here, we plot the posterior predicitve distribution associated with each distribution's highest-ELPD dependent variable.



Figure 17: Observed partner age distributions (grey bars) and posterior predictive partner age distributions (lines) for each probability distribution among women in the Manicaland data set. Here, we plot the posterior predicitve distribution associated with each distribution's highest-ELPD dependent variable.



Figure 18: Observed partner age distributions (grey bars) and posterior predictive partner age distributions (lines) for each probability distribution among men in the Manicaland data set. Here, we plot the posterior predicitve distribution associated with each distribution's highest-ELPD dependent variable.

Supplementary Tables

Table 7: Full ELPD and QQ RMSE table for women in the AHRI data set. Higher ELPD values and lower QQ RMSE values are better.

Papl	Distribution	רים די		SE of Diff	OO PMSE
Канк	Distribution	ELFD	ELLUDIII		QQ KIVISE
AHRI	Female 20-24			2.22	
1	Sinh-arcsinh	-31750.94	0.00	0.00	0.32
2	Skew normal	-32056.39	-305.46	48.63	0.47
3	Normal	-32414.54	-663.61	60.54	0.62
4	Beta	-32953.92	-1202.98	112.08	0.77
5	Gamma	-33461.85	-1710.92	148.15	0.80
AHRI	Female 25-29				
1	Sinh-arcsinh	-24647.65	0.00	0.00	0.28
2	Skew normal	-24906.22	-258.57	43.27	0.52
3	Normal	-25238.71	-591.06	54.82	0.68
4	Beta	-25701.13	-1053.48	114.84	0.89
5	Gamma	-25995.81	-1348.16	132.15	0.90
AHRI	Female 30-34				
1	Sinh-arcsinh	-19831.53	0.00	0.00	0.44
2	Skew normal	-20200.44	-368.91	69.40	0.51
3	Normal	-20314.79	-483.26	52.24	0.80
4	Beta	-20575.61	-744.08	67.46	0.93
5	Gamma	-20708.35	-876.82	73.89	0.91
AHRI	Female 35-39				
1	Sinh-arcsinh	-15469.18	0.00	0.00	0.31
2	Skew normal	-15749.79	-280.61	53.04	0.77
3	Normal	-15834.32	-365.14	41.23	0.80
4	Beta	-16026.51	-557.33	53.99	1.18
5	Gamma	-16087.40	-618.22	57.06	1.04
AHRI	Female 40-44				
1	Sinh-arcsinh	-12556.61	0.00	0.00	0.45
2	Skew normal	-12876.71	-320.10	45.85	1.27
3	Normal	-12935.34	-378.73	52.38	0.92
4	Beta	-13137.69	-581.08	69.18	1.38
5	Gamma	-13150.66	-594.05	62.73	1.19
AHRI	Female 45-49				
1	Sinh-arcsinh	-10059.21	0.00	0.00	0.59
2	Skew normal	-10391.95	-332.74	42.75	1.36
3	Normal	-10433.64	-374.43	48.91	1.53
4	Gamma	-10527.00	-467.79	50.72	1.35
5	Beta	-10545.33	-486.12	56.02	1.58

1	Distribution	ELDD			
Kank	Distribution	ELPD	ELPD Diff	SE of Diff	QQ KMSE
AHRI	Male 20-24				
1	Sinh-arcsinh	-20428.11	0.00	0.00	0.23
2	Skew normal	-20499.86	-71.75	17.12	0.25
3	Normal	-20503.89	-75.79	16.85	0.22
4	Beta	-20545.59	-117.49	23.21	0.22
5	Gamma	-20700.24	-272.13	43.53	0.29
AHRI	Male 25-29				
1	Sinh-arcsinh	-12664.21	0.00	0.00	0.26
2	Skew normal	-12727.03	-62.82	17.86	0.28
3	Beta	-12739.03	-74.82	18.65	0.31
4	Normal	-12753.25	-89.04	19.35	0.29
5	Gamma	-12788.26	-124.05	35.07	0.38
AHRI	Male 30-34				
1	Sinh-arcsinh	-9301.03	0.00	0.00	0.29
2	Skew normal	-9357.18	-56.15	14.08	0.43
3	Beta	-9371.86	-70.83	16.48	0.37
4	Normal	-9385.63	-84.60	14.67	0.46
5	Gamma	-9419.34	-118.31	35.11	0.27
AHRI	Male 35-39				
1	Sinh-arcsinh	-6746.89	0.00	0.00	0.30
2	Skew normal	-6812.77	-65.88	17.73	0.64
3	Normal	-6817.86	-70.97	23.24	0.70
4	Beta	-6830.95	-84.06	17.95	0.71
5	Gamma	-6832.47	-85.58	32.03	0.44
AHRI	Male 40-44				
1	Sinh-arcsinh	-4610.95	0.00	0.00	0.35
2	Skew normal	-4711.78	-100.84	18.66	0.92
3	Normal	-4713.78	-102.83	18.82	0.78
4	Gamma	-4718.28	-107.33	24.83	0.63
5	Beta	-4742.70	-131.75	17.28	1.07
AHRI	Male 45-49				
1	Sinh-arcsinh	-3683.47	0.00	0.00	0.34
2	Skew normal	-3770.59	-87.12	16.56	0.81
3	Gamma	-3776.33	-92.86	15.56	0.87
4	Normal	-3778.84	-95.37	14.50	1.17
5	Beta	-3805.78	-122.31	17.40	1.36

Table 8: Full ELPD and QQ RMSE table for men in the AHRI data set. Higher ELPD values and lower QQ RMSE values are better.

Rank	Distribution	ELPD	ELPD Diff	SE of Diff	QQ RMSE
AHRI	Deheaped Fem	ale 20-24			
1	Sinh-arcsinh	-31411.24	0.00	0.00	0.26
2	Skew normal	-31797.37	-386.13	53.15	0.59
3	Normal	-32179.29	-768.05	65.50	0.56
4	Beta	-32737.57	-1326.32	118.47	0.76
5	Gamma	-33254.17	-1842.92	155.14	0.78
AHRI	Deheaped Fem	ale 25-29			
1	Sinh-arcsinh	-24439.47	0.00	0.00	0.27
2	Skew normal	-24768.06	-328.59	46.71	0.65
3	Normal	-25104.46	-664.99	58.32	0.82
4	Beta	-25574.33	-1134.86	119.65	1.03
5	Gamma	-25870.30	-1430.83	137.51	1.05
AHRI	Deheaped Fem	ale 30-34			
1	Sinh-arcsinh	-19680.77	0.00	0.00	0.41
2	Skew normal	-20112.70	-431.94	72.95	0.55
3	Normal	-20228.52	-547.76	56.19	0.81
4	Beta	-20492.23	-811.46	70.53	0.92
5	Gamma	-20624.82	-944.06	76.98	0.80
AHRI	Deheaped Fem	ale 35-39			
1	Sinh-arcsinh	-15381.68	0.00	0.00	0.26
2	Skew normal	-15703.77	-322.09	55.30	0.68
3	Normal	-15788.73	-407.05	43.67	0.82
4	Beta	-15983.57	-601.90	56.31	1.13
5	Gamma	-16044.22	-662.54	59.17	1.04
AHRI	Deheaped Fem	ale 40-44			
1	Sinh-arcsinh	-12491.91	0.00	0.00	0.25
2	Skew normal	-12846.63	-354.72	47.38	0.99
3	Normal	-12905.04	-413.12	54.14	0.89
4	Beta	-13109.82	-617.91	70.96	1.31
5	Gamma	-13121.45	-629.53	64.12	1.13
AHRI	Deheaped Fem	ale 45-49			
1	Sinh-arcsinh	-9981.83	0.00	0.00	0.53
2	Skew normal	-10357.85	-376.01	45.08	1.43
3	Normal	-10401.64	-419.80	51.57	1.46
4	Gamma	-10493.73	-511.90	52.90	1.37
5	Beta	-10513.46	-531.63	58.21	1.61

Table 9: Full ELPD and QQ RMSE table for women in the AHRI Deheaped data set. Higher ELPD values and lower QQ RMSE values are better.

Rank	Distribution	ELPD	ELPD Diff	SE of Diff	QQ RMSE
AHRI	Deheaped Male	e 20-24			
1	Sinh-arcsinh	-20310.35	0.00	0.00	0.27
2	Skew normal	-20429.90	-119.55	27.09	0.22
3	Normal	-20459.73	-149.38	35.78	0.29
4	Beta	-20574.15	-263.80	75.13	0.22
5	Gamma	-20899.52	-589.17	175.99	0.27
AHRI	Deheaped Male	e 25-29			
1	Sinh-arcsinh	-12585.54	0.00	0.00	0.28
2	Skew normal	-12680.59	-95.05	21.53	0.44
3	Beta	-12697.00	-111.46	23.31	0.37
4	Normal	-12701.76	-116.23	22.96	0.41
5	Gamma	-12763.81	-178.27	41.24	0.39
AHRI	Deheaped Male	e 30-34			
1	Sinh-arcsinh	-9227.26	0.00	0.00	0.37
2	Skew normal	-9302.42	-75.16	16.15	0.41
3	Beta	-9318.24	-90.97	19.07	0.39
4	Normal	-9327.58	-100.31	16.18	0.41
5	Gamma	-9372.32	-145.06	38.27	0.27
AHRI	Deheaped Male	e 35-39			
1	Sinh-arcsinh	-6694.86	0.00	0.00	0.30
2	Skew normal	-6774.11	-79.26	19.32	0.61
3	Normal	-6780.69	-85.84	25.42	0.44
4	Beta	-6791.95	-97.10	19.81	0.69
5	Gamma	-6796.41	-101.55	34.45	0.40
AHRI	Deheaped Male	e 40-44			
1	Sinh-arcsinh	-4591.04	0.00	0.00	0.49
2	Skew normal	-4700.54	-109.51	19.38	1.16
3	Normal	-4703.52	-112.49	19.93	1.00
4	Gamma	-4708.43	-117.40	25.94	0.89
5	Beta	-4731.41	-140.37	17.84	1.30
AHRI	Deheaped Male	e 45-49			
1	Sinh-arcsinh	-3680.18	0.00	0.00	0.30
2	Normal	-3796.06	-115.88	19.24	1.15
3	Skew normal	-3797.14	-116.95	23.48	1.02
4	Gamma	-3801.02	-120.83	24.51	0.98
5	Beta	-3817.97	-137.79	19.37	1.39

Table 10: Full ELPD and QQ RMSE table for men in the AHRI Deheaped data set. Higher ELPD values and lower QQ RMSE values are better.

Rank	Distribution	ELPD	ELPD Diff	SE of Diff	QQ RMSE
Haiti 2	2016-17 DHS Fe	male 20-24			
1	Sinh-arcsinh	-3259.31	0.00	0.00	0.49
2	Skew normal	-3263.46	-4.15	4.95	0.53
3	Normal	-3338.23	-78.92	19.54	0.91
4	Beta	-3441.91	-182.60	45.77	1.24
5	Gamma	-3504.85	-245.54	53.90	1.29
Haiti 2	2016-17 DHS Fer	male 25-29			
1	Sinh-arcsinh	-4447.43	0.00	0.00	0.26
2	Skew normal	-4471.22	-23.78	8.41	0.57
3	Normal	-4527.25	-79.82	18.72	0.86
4	Beta	-4625.97	-178.54	40.88	1.23
5	Gamma	-4678.20	-230.77	45.81	1.22
Haiti 2	2016-17 DHS Fea	male 30-34			
1	Sinh-arcsinh	-4720.12	0.00	0.00	0.44
2	Skew normal	-4749.57	-29.45	9.06	0.68
3	Normal	-4763.78	-43.66	10.51	0.62
4	Beta	-4809.11	-88.99	17.19	0.85
5	Gamma	-4836.82	-116.70	20.32	0.83
Haiti 2	2016-17 DHS Fe	male 35-39			
1	Sinh-arcsinh	-4490.82	0.00	0.00	0.33
2	Skew normal	-4518.58	-27.75	8.14	0.57
3	Normal	-4526.55	-35.73	8.59	0.73
4	Beta	-4561.27	-70.45	13.60	0.94
5	Gamma	-4577.84	-87.01	15.27	0.86
Haiti 2	2016-17 DHS Fer	male 40-44			
1	Sinh-arcsinh	-3601.02	0.00	0.00	0.35
2	Skew normal	-3629.45	-28.43	7.51	0.83
3	Normal	-3633.14	-32.11	7.96	0.71
4	Beta	-3641.61	-40.59	9.76	0.73
5	Gamma	-3644.89	-43.86	10.47	0.64
Haiti 2	2016-17 DHS Fer	male 45-49			
1	Sinh-arcsinh	-3106.27	0.00	0.00	0.39
2	Skew normal	-3133.10	-26.82	7.68	0.88
3	Gamma	-3133.61	-27.33	7.50	0.68
4	Normal	-3134.62	-28.35	7.46	0.81
5	Beta	-3136.89	-30.62	8.61	0.88

Table 11: Full ELPD and QQ RMSE table for women in the Haiti 2016-17 DHS data set. Higher ELPD values and lower QQ RMSE values are better.

Rank	Distribution	ELPD	ELPD Diff	SE of Diff	QQ RMSE			
Haiti 2016-17 DHS Male 20-24								
1	Skew normal	-468.98	0.00	0.00	0.43			
2	Sinh-arcsinh	-469.60	-0.62	1.12	0.41			
3	Normal	-475.31	-6.33	4.28	0.67			
4	Beta	-483.53	-14.55	7.33	0.65			
5	Gamma	-500.53	-31.55	13.35	0.94			
Haiti 2016-17 DHS Male 25-29								
1	Sinh-arcsinh	-1386.13	0.00	0.00	0.38			
2	Skew normal	-1390.54	-4.41	3.19	0.49			
3	Normal	-1395.47	-9.34	4.79	0.60			
4	Beta	-1407.46	-21.32	7.31	0.62			
5	Gamma	-1434.18	-48.04	11.75	0.79			
Haiti 2	Haiti 2016-17 DHS Male 30-34							
1	Sinh-arcsinh	-2217.20	0.00	0.00	0.44			
2	Skew normal	-2222.10	-4.89	3.42	0.69			
3	Normal	-2223.97	-6.76	4.48	0.45			
4	Beta	-2240.58	-23.37	9.32	0.52			
5	Gamma	-2281.18	-63.98	17.91	0.73			
Haiti 2	Haiti 2016-17 DHS Male 35-39							
1	Sinh-arcsinh	-2185.96	0.00	0.00	0.28			
2	Skew normal	-2189.87	-3.91	2.67	0.69			
3	Beta	-2191.05	-5.10	3.68	0.48			
4	Normal	-2191.11	-5.16	3.55	0.49			
5	Gamma	-2205.69	-19.73	9.57	0.52			
Haiti 2	Haiti 2016-17 DHS Male 40-44							
1	Sinh-arcsinh	-2051.62	0.00	0.00	0.39			
2	Skew normal	-2060.16	-8.54	4.21	0.72			
3	Normal	-2060.38	-8.75	4.57	0.69			
4	Beta	-2062.00	-10.37	4.87	0.70			
5	Gamma	-2063.79	-12.17	5.73	0.47			
Haiti 2	Haiti 2016-17 DHS Male 45-49							
1	Sinh-arcsinh	-2138.34	0.00	0.00	0.23			
2	Normal	-2150.53	-12.19	6.38	0.35			
3	Skew normal	-2151.51	-13.17	5.97	0.56			
4	Gamma	-2152.88	-14.54	8.93	0.25			
5	Beta	-2156.13	-17.79	6.14	0.48			

Table 12: Full ELPD and QQ RMSE table for men in the Haiti 2016-17 DHS data set. Higher ELPD values and lower QQ RMSE values are better.

Rank	Distribution	ELPD	ELPD Diff	SE of Diff	QQ RMSE		
Manic	Manicaland Female 20-24						
1	Sinh-arcsinh	-16390.77	0.00	0.00	0.31		
2	Skew normal	-16502.01	-111.25	21.22	0.44		
3	Normal	-16779.93	-389.16	37.05	0.67		
4	Beta	-17111.57	-720.80	62.02	0.86		
5	Gamma	-17387.38	-996.61	76.80	1.02		
Manicaland Female 25-29							
1	Sinh-arcsinh	-18702.50	0.00	0.00	0.53		
2	Skew normal	-18923.04	-220.53	25.27	0.94		
3	Normal	-19080.66	-378.16	36.05	0.83		
4	Beta	-19405.80	-703.30	64.97	1.05		
5	Gamma	-19615.53	-913.03	76.38	1.09		
Manicaland Female 30-34							
1	Sinh-arcsinh	-16523.81	0.00	0.00	0.48		
2	Skew normal	-16877.96	-354.15	40.36	0.87		
3	Normal	-16886.62	-362.80	36.41	0.99		
4	Beta	-17021.26	-497.44	43.60	1.12		
5	Gamma	-17094.58	-570.76	49.53	0.93		
Manic	aland Female 3	5-39					
1	Sinh-arcsinh	-14397.76	0.00	0.00	0.48		
2	Skew normal	-14736.64	-338.88	28.35	1.25		
3	Normal	-14798.55	-400.79	36.87	1.39		
4	Beta	-14824.80	-427.04	33.02	1.47		
5	Gamma	-14835.11	-437.35	34.49	1.14		
Manic	Manicaland Female 40-44						
1	Sinh-arcsinh	-12293.13	0.00	0.00	0.68		
2	Skew normal	-12488.28	-195.15	21.36	1.03		
3	Gamma	-12500.93	-207.80	22.18	1.03		
4	Normal	-12508.91	-215.78	23.29	1.28		
5	Beta	-12537.14	-244.01	25.41	1.22		
Manic	Manicaland Female 45-49						
1	Sinh-arcsinh	-9183.03	0.00	0.00	0.56		
2	Skew normal	-9455.87	-272.83	23.57	1.68		
3	Normal	-9477.33	-294.30	23.55	1.62		
4	Gamma	-9497.31	-314.27	25.08	1.44		
5	Beta	-9576.44	-393.40	32.07	1.94		

Table 13: Full ELPD and QQ RMSE table for women in the Manicaland data set. Higher ELPD values and lower QQ RMSE values are better.

Rank	Distribution	ELPD	ELPD Diff	SE of Diff	QQ RMSE			
Manicaland Male 20-24								
1	Sinh-arcsinh	-9770.00	0.00	0.00	0.30			
2	Skew normal	-9895.82	-125.83	33.35	0.40			
3	Normal	-10139.11	-369.11	79.13	0.49			
4	Beta	-10587.64	-817.64	181.23	0.56			
5	Gamma	-11594.58	-1824.59	388.26	1.15			
Manicaland Male 25-29								
1	Sinh-arcsinh	-13978.59	0.00	0.00	0.40			
2	Skew normal	-13990.39	-11.80	8.51	0.48			
3	Normal	-14018.60	-40.00	17.48	0.45			
4	Beta	-14152.35	-173.76	48.77	0.40			
5	Gamma	-14500.47	-521.87	117.58	0.55			
Manic	Manicaland Male 30-34							
1	Sinh-arcsinh	-12949.24	0.00	0.00	0.31			
2	Skew normal	-13016.44	-67.21	25.01	0.37			
3	Normal	-13037.46	-88.22	16.57	0.49			
4	Beta	-13070.31	-121.07	54.74	0.41			
5	Gamma	-13285.47	-336.23	171.92	0.42			
Manic	Manicaland Male 35-39							
1	Sinh-arcsinh	-11496.14	0.00	0.00	0.27			
2	Skew normal	-11528.36	-32.22	9.83	0.39			
3	Normal	-11530.43	-34.29	9.72	0.26			
4	Gamma	-11531.75	-35.61	12.47	0.24			
5	Beta	-11582.63	-86.49	12.97	0.48			
Manic	Manicaland Male 40-44							
1	Sinh-arcsinh	-8714.06	0.00	0.00	0.35			
2	Skew normal	-8749.78	-35.72	10.11	0.51			
3	Gamma	-8777.08	-63.02	10.38	0.55			
4	Normal	-8791.45	-77.38	12.23	0.76			
5	Beta	-8860.22	-146.16	18.15	0.93			
Manic	Manicaland Male 45-49							
1	Sinh-arcsinh	-7110.27	0.00	0.00	0.42			
2	Skew normal	-7177.03	-66.75	25.08	0.76			
3	Gamma	-7213.99	-103.72	13.02	1.07			
4	Normal	-7232.04	-121.77	13.09	1.28			
5	Beta	-7312.35	-202.08	18.61	1.61			

Table 14: Full ELPD and QQ RMSE table for men in the Manicaland data set. Higher ELPD values and lower QQ RMSE values are better.

Table 15: LOO-CV estimated ELPD values, differences, and standard errors of differences, as well as QQ RMSE values, for all five regression models fit to all four data sets. The "difference" value of a row is the difference between that row's ELPD value and data set-specific best ELPD value. Higher ELPD values and lower QQ RMSE values are better.

Rank	Model	ELPD	ELPD Diff	SE of Diff	QQ RMSE		
AHRI							
1	Distributional 4	55841.91	0.00	0.00	0.66		
2	Distributional 3	55534.16	-307.75	32.36	0.93		
3	Distributional 2	54794.79	-1047.12	51.69	1.21		
4	Distributional 1	54335.19	-1506.72	72.32	1.15		
5	Conventional	52689.21	-3152.70	100.59	1.30		
AHRI Deaheaped							
1	Distributional 4	58503.98	0.00	0.00	0.62		
2	Distributional 3	58219.23	-284.75	28.64	0.92		
3	Distributional 2	57503.68	-1000.30	47.14	1.14		
4	Distributional 1	57097.39	-1406.59	64.48	1.06		
5	Conventional	55296.25	-3207.73	99.42	1.26		
Haiti 2	Haiti 2016-17 DHS						
1	Distributional 4	5207.57	0.00	0.00	0.84		
2	Distributional 3	5196.69	-10.89	6.54	0.91		
3	Distributional 1	5140.77	-66.80	12.27	0.98		
4	Distributional 2	5138.75	-68.83	12.24	0.99		
5	Conventional	4777.78	-429.80	30.54	1.33		
Manic	aland						
1	Distributional 4	24516.15	0.00	0.00	1.04		
2	Distributional 3	24313.74	-202.40	20.52	1.34		
3	Distributional 2	23472.07	-1044.08	47.77	1.80		
4	Distributional 1	23192.49	-1323.66	54.97	1.89		
5	Conventional	21011.29	-3504.86	89.01	2.05		