

# A compendium of covariances and correlation coefficients of coalescent tree properties

Egor Alimpiev\* and Noah A Rosenberg\*

October 10, 2022

**Abstract.** Gene genealogies are frequently studied by measuring properties such as their height ( $H$ ), length ( $L$ ), sum of external branches ( $E$ ), sum of internal branches ( $I$ ), and mean of their two basal branches ( $B$ ), and the coalescence times that contribute to the other genealogical features ( $T$ ). These tree properties and their relationships can provide insight into the effects of population-genetic processes on genealogies and genetic sequences. Here, under the coalescent model, we study the 15 correlations among pairs of features of genealogical trees:  $H_n$ ,  $L_n$ ,  $E_n$ ,  $I_n$ ,  $B_n$ , and  $T_k$  for a sample of size  $n$ , with  $2 \leq k \leq n$ . We report high correlations among  $H_n$ ,  $L_n$ ,  $I_n$ , and  $B_n$ , with all pairwise correlations of these quantities having values greater than or equal to  $\sqrt{6}[6\zeta(3) + 6 - \pi^2]/(\pi\sqrt{18 + 9\pi^2 - \pi^4}) \approx 0.84930$  in the limit as  $n \rightarrow \infty$ . Although  $E_n$  has an expectation of 2 for all  $n$  and  $H_n$  has expectation 2 in the limit as  $n \rightarrow \infty$ , their limiting correlation is 0. The results contribute toward understanding features of the shapes of coalescent trees.

---

\*Department of Biology, Stanford University, Stanford, CA 94305.

# 1 Introduction

In coalescent theory, gene genealogies are investigated in relation to the evolutionary processes that are included in population-genetic models (Hein *et al.*, 2005; Wakeley, 2009). In analyzing genetic sequences in populations, features of gene genealogies that underlie the sequences are often used to understand the evolutionary processes that have given rise to the data.

The use of genealogies for interpreting genetic sequences relies on an understanding of the ways in which different features of the genealogies behave when affected by population-genetic processes. For example, comparing a constant-sized and an exponentially growing population, exponential growth increases the total length of the branches of a gene genealogy in relation to its height (Slatkin & Hudson, 1991; Slatkin, 1996; Sano & Tachida, 2005). Coalescences are rare in recent generations, when the population is large, and they occur primarily in the period deep in the past when the population was small.

Several tree features have been used for measuring effects of population-genetic processes on gene genealogies (Slatkin, 1996; Uyenoyama, 1997; Schierup & Hein, 2000; Rosenberg, 2006). For a binary ultrametric tree of  $n$  lineages, these features (Figure 1) include the tree height from the tips to the root ( $H_n$ ), the total length of all the branches ( $L_n$ ), the total length of external branches connecting tips to the nearest internal node ( $E_n$ ), the total length of internal branches connecting internal nodes to other internal nodes ( $I_n$ ), and the mean length of the two basal branches incident to the root node ( $B_n$ ).

These tree features are correlated, as increases or decreases in a coalescence time—the time  $T_k$  for coalescence of  $k$  to  $k - 1$  lineages, for example, with  $2 \leq k \leq n$ —can affect more than one of the quantities. For example, an increase in the total length  $L_n$  necessarily increases the length  $E_n$  of the external branches, the length  $I_n$  of the internal branches, or both.

Analyses of coalescent models have examined some of the correlations between tree features, notably the relationship between  $H_n$  and  $L_n$  (Fu, 1996; Griffiths & Tavaré, 1996; Rosenberg & Hirsh, 2003; Arbisser *et al.*, 2018). Here, we perform a detailed investigation of correlations and ratios among  $H_n$ ,  $L_n$ ,  $E_n$ ,  $I_n$ , and  $B_n$ . For each pair, under the coalescent, assuming a constant-sized population, we evaluate their covariance and correlation. We explore limiting values as  $n \rightarrow \infty$ . The approach follows Arbisser *et al.* (2018), who obtained the covariance and correlation of  $H_n$  and  $L_n$ ; we perform analogous calculations for all 10 pairs among  $\{H_n, L_n, E_n, I_n, B_n\}$ , as well as for the five pairs involving one of  $\{H_n, L_n, E_n, I_n, B_n\}$  and  $T_k$ .

## 2 Tree properties

We consider the standard coalescent model of a constant-sized population of size  $N$  haploids. Time is measured in units of the population size, with one time unit representing  $N$  generations. For sample size  $n \geq 2$ , we examine tree properties  $H_n$ ,  $L_n$ ,  $E_n$ ,  $I_n$ , and  $B_n$ , as well as the coalescence time  $T_k$ ,  $2 \leq k \leq n$ . In this section, we recall basic features of the various quantities.

For convenience, for a mathematical expression we will use frequently, we write

$$S_{p,n} = \sum_{k=1}^n \frac{1}{k^p}. \quad (1)$$

Recall that the harmonic sum  $S_{1,\infty}$  diverges, and the sum of the reciprocals of squares is  $S_{2,\infty} = \pi^2/6 \approx 1.64493$ .

The sum of the reciprocals of cubes is Apéry's constant,  $S_{3,\infty} = \zeta(3) \approx 1.20206$ .

### 2.1 $T_k$

$T_k$  is a random variable representing the time during which  $k$  lineages coalesce to  $k-1$  lineages. The  $T_k$ ,  $2 \leq k \leq n$ , are independent and exponentially distributed with probability density function  $f_{T_k}(t_k) = \binom{k}{2} e^{-\binom{k}{2} t_k}$  (Wakeley, 2009, p. 60). The expectation and variance of  $T_k$  are then (Wakeley, 2009, p. 61)

$$\mathbb{E}[T_k] = \frac{2}{k(k-1)}, \quad (2)$$

$$\text{Var}[T_k] = \frac{4}{k^2(k-1)^2}. \quad (3)$$

As  $n, k \rightarrow \infty$  with  $k \leq n$ , both  $\mathbb{E}[T_k]$  and  $\text{Var}[T_k]$  have limit 0.

### 2.2 $H_n$

For  $n \geq 2$ , the height  $H_n$  of a tree from root to leaves can be written

$$H_n = \sum_{k=2}^n T_k. \quad (4)$$

The expectation and variance of  $H_n$  are then found using the expectation and variance of  $T_k$  (eqs. 2 and 3), noting that the  $T_k$  are independent (Wakeley, 2009, p. 76):

$$\mathbb{E}[H_n] = \sum_{k=2}^n \mathbb{E}[T_k] = \frac{2(n-1)}{n}, \quad (5)$$

$$\text{Var}[H_n] = 8 \left( \sum_{k=2}^n \frac{1}{k^2} \right) - 4 \left( \frac{n-1}{n} \right)^2. \quad (6)$$

The variance can be written  $\text{Var}[H_n] = 4(2S_{2,n}n^2 - 3n^2 + 2n - 1)/n^2$ . The limits are  $\lim_{n \rightarrow \infty} \mathbb{E}[H_n] = 2$  and  $\lim_{n \rightarrow \infty} \text{Var}[H_n] = 4\pi^2/3 - 12 \approx 1.15947$  (Wakeley, 2009, p. 76).

### 2.3 $L_n$

For  $n \geq 2$ , the total tree length, summing the lengths of all branches of a tree, is

$$L_n = \sum_{k=2}^n kT_k. \quad (7)$$

By eqs. 2 and 3 and the independence of the  $T_k$ , we have (Wakeley, 2009, p. 76)

$$\mathbb{E}[L_n] = \sum_{k=2}^n k\mathbb{E}[T_k] = 2 \sum_{k=1}^{n-1} \frac{1}{k}, \quad (8)$$

$$\text{Var}[L_n] = 4 \sum_{k=1}^{n-1} \frac{1}{k^2}. \quad (9)$$

In terms of  $S_{p,n}$  (eq. 1), these expressions are  $\mathbb{E}[L_n] = 2S_{1,n-1}$  and  $\text{Var}[L_n] = 4S_{2,n-1}$ . The limits are  $\lim_{n \rightarrow \infty} \mathbb{E}[L_n] = \infty$  and  $\lim_{n \rightarrow \infty} \text{Var}[L_n] = 2\pi^2/3 \approx 6.57974$  (Wakeley, 2009, p. 76).

### 2.4 $E_n$

The external branches of a tree are the branches that connect leaves to their nearest internal nodes. Denoting the individual external branch lengths  $e_1^{(n)}, e_2^{(n)}, \dots, e_n^{(n)}$ , the sum of external branch lengths is

$$E_n = e_1^{(n)} + e_2^{(n)} + \dots + e_n^{(n)}.$$

The  $e_k^{(n)}$  are identically distributed, and we write  $e_n$  for the length of a randomly chosen external branch of a tree of  $n$  lineages. The sum of the external branches has expectation

$$\mathbb{E}[E_n] = n\mathbb{E}[e_n]. \quad (10)$$

The random variable  $e_n$  can be written recursively as (Fu & Li, 1993, eq. 7)

$$e_n = \begin{cases} e_{n-1} + T_n, & \text{probability } \frac{n-2}{n}, \\ T_n, & \text{probability } \frac{2}{n}. \end{cases} \quad (11)$$

Expressions for  $\mathbb{E}[e_n]$ ,  $\mathbb{E}[E_n]$ , and  $\text{Var}[E_n]$  can then be obtained by solving recurrence equations (Fu & Li, 1993). We have

$$\mathbb{E}[e_n] = \frac{2}{n}. \quad (12)$$

For the mean and variance of  $E_n$ , we obtain (Fu & Li, 1993, eqs. 10 and 14)

$$\mathbb{E}[E_n] = 2, \quad (13)$$

$$\text{Var}[E_n] = \begin{cases} 4, & n = 2, \\ \frac{8}{(n-1)(n-2)}[n(\sum_{k=1}^{n-1} \frac{1}{k}) - 2(n-1)], & n > 2. \end{cases} \quad (14)$$

$\mathbb{E}[E_n]$  is equal to 2 irrespective of the choice of  $n$ , so that  $\lim_{n \rightarrow \infty} \mathbb{E}[E_n] = 2$ . The limit of the variance is  $\lim_{n \rightarrow \infty} \text{Var}[E_n] = 0$  (Fu & Li, 1993).

## 2.5 $I_n$

The internal branches connect internal nodes to other internal nodes. Their total length is  $I_n$ , with

$$I_n = L_n - E_n. \quad (15)$$

The mean and variance of the sum of internal branches are (Fu & Li, 1993, eqs. 12 and 17)

$$\mathbb{E}[I_n] = \mathbb{E}[L_n] - \mathbb{E}[E_n] = 2 \left( \sum_{k=1}^{n-1} \frac{1}{k} \right) - 2, \quad (16)$$

$$\text{Var}[I_n] = 4 \left[ \frac{2[S_{1,n-1}n - 2(n-1)]}{(n-1)(n-2)} - \frac{2S_{1,n-1}}{n-1} + S_{2,n-1} \right]. \quad (17)$$

The limits are  $\lim_{n \rightarrow \infty} \mathbb{E}[I_n] = \infty$  and  $\lim_{n \rightarrow \infty} \text{Var}[I_n] = 2\pi^2/3 \approx 6.57974$ , the same as for  $L_n$  (Section 2.3).

## 2.6 $B_n$

Finally, we consider the basal branches, the two branches that extend from the root. We define  $B_n$  as the mean of the two branch lengths. One of the branches has length  $T_2$ , and we denote the other length  $b_n$ . We assume here that  $n \geq 4$  for calculations involving  $B_n$ . The appendix of Uyenoyama (1997) gives

$$B_n = \frac{T_2 + b_n}{2}, \quad (18)$$

with

$$b_n = \left[ \sum_{j=3}^{n-1} \sum_{k=2}^j \frac{T_k}{j} \prod_{i=3}^{j-1} \left(1 - \frac{1}{i}\right) \right] + \left[ \sum_{k=2}^n T_k \prod_{i=3}^{n-1} \left(1 - \frac{1}{i}\right) \right]$$

for  $n \geq 4$ . A convenient form for  $b_n$  encodes the fact that with probability  $2/[j(j-1)]$ ,  $b_n = H_j$  for  $j = 3, 4, \dots, n-1$ , and with probability  $2/(n-1)$ ,  $b_n = H_n$ :

$$b_n = \left[ \sum_{j=3}^{n-1} \sum_{k=2}^j \frac{2}{j(j-1)} T_k \right] + \left( \sum_{k=2}^n \frac{2}{n-1} T_k \right). \quad (19)$$

Assuming  $n \geq 4$ , the branch length  $b_n$  has expectation (Uyenoyama, 1997):

$$\mathbb{E}[b_n] = \frac{4}{n} + 4 \sum_{k=3}^{n-1} \frac{1}{k^2}. \quad (20)$$

The expectation and variance of  $B_n$  then equal

$$\mathbb{E}[B_n] = \frac{2}{n} + 2 \sum_{k=2}^{n-1} \frac{1}{k^2}, \quad (21)$$

$$\text{Var}[B_n] = \frac{2(3S_{2,n-1}n^2 - 2S_{2,n-1}^2n^2 + n^2 - 4S_{2,n-1}n + 3n - 4)}{n^2}. \quad (22)$$

The expectation appears in the appendix of Uyenoyama (1997). We calculate the expression for the variance in Section 3.11. Taking limits of these equations, we obtain  $\lim_{n \rightarrow \infty} \mathbb{E}[B_n] = \pi^2/3 - 2 \approx 1.28987$  and  $\lim_{n \rightarrow \infty} \text{Var}[B_n] = 2 + \pi^2 - \pi^4/9 \approx 1.04637$ .

### 3 Theoretical results

For pairs of variables among  $\{H_n, L_n, E_n, I_n, B_n, T_k\}$ , we apply results from Section 2 to compute covariances and correlations. First, for each pair, we compute their covariance. The covariance together with the variances of the two quantities from Section 2 provides their correlation. We obtain the limiting correlation for large trees by taking  $n \rightarrow \infty$ . Among the 15 pairs, our analyses for 13 are exact; for  $(E_n, B_n)$  and  $(I_n, B_n)$ , we offer approximate covariances and correlations. We also provide the derivation of eq. 22 for  $\text{Var}[B_n]$ .

Note that correlations in pairs involving  $E_n$  have distinct forms for  $n = 2$  and  $n \geq 3$ , owing to the piecewise definition of  $\text{Var}[E_n]$  in eq. 14. We exclude the case of  $n = 2$  for pairs involving  $I_n$ , as  $I_2 = 0$  with  $\text{Var}[I_2] = 0$ . We also assume that  $B_n$  is defined only for  $n \geq 4$ .

We present a summary of our mathematical results in Tables 1 and 2. Table 1 shows covariances of pairs of variables and their limits as  $n \rightarrow \infty$ . Table 2 shows correlations and their  $n \rightarrow \infty$  limits.

#### 3.1 $H_n$ and $T_k$

We calculate the covariance of  $H_n$  and  $T_k$  using  $\text{Cov}[H_n, T_k] = \mathbb{E}[H_n T_k] - \mathbb{E}[H_n] \mathbb{E}[T_k]$ . Recalling that  $T_i$  and  $T_j$  are independent for  $i \neq j$  (Section 2.1), we have  $\mathbb{E}[T_i T_j] = \mathbb{E}[T_i] \mathbb{E}[T_j]$  for  $i \neq j$ . Hence, inserting

eq. 4 for  $H_n$  and eq. 2 for  $\mathbb{E}[T_i]$ , we have

$$\begin{aligned}
\text{Cov}[H_n, T_k] &= \mathbb{E} \left[ T_k \sum_{i=2}^n T_i \right] - \mathbb{E} \left[ \sum_{i=2}^n T_i \right] \mathbb{E}[T_k] \\
&= \sum_{i=2}^n \mathbb{E}[T_i T_k] - \sum_{i=2}^n \mathbb{E}[T_i] \mathbb{E}[T_k] \\
&= \text{Var}[T_k] + \sum_{i=2, i \neq k}^n \mathbb{E}[T_i] \mathbb{E}[T_k] - \sum_{i=2, i \neq k}^n \mathbb{E}[T_i] \mathbb{E}[T_k] = \text{Var}[T_k] = \frac{4}{k^2(k-1)^2}, \quad (23)
\end{aligned}$$

where the last step uses  $\text{Var}[T_k]$  from eq. 3. We observe that the covariance is independent of  $n$ .

For the correlation coefficient  $\text{Corr}[H_n, T_k] = \text{Cov}[H_n, T_k] / \sqrt{\text{Var}[H_n] \text{Var}[T_k]}$ , applying eq. 6 for  $\text{Var}[H_n]$ , eq. 3 for  $\text{Var}[T_k]$ , and eq. 23 for  $\text{Cov}[H_n, T_k]$ , we have

$$\text{Corr}[H_n, T_k] = \frac{n}{\sqrt{2S_{2,n}n^2 - 3n^2 + 2n - 1}} \frac{1}{k(k-1)}. \quad (24)$$

Taking a limit as  $n \rightarrow \infty$ , we obtain

$$\lim_{n \rightarrow \infty} \text{Corr}[H_n, T_k] = \frac{\sqrt{3}}{\sqrt{\pi^2 - 9}} \frac{1}{k(k-1)}. \quad (25)$$

The limiting correlation decreases to 0 with  $k$  from an initial value of  $\frac{1}{2}[\sqrt{3/(\pi^2 - 9)}] \approx 0.92869$  at  $k = 2$ .

### 3.2 $L_n$ and $T_k$

For  $L_n$  and  $T_k$ , applying eqs. 7, 8, and 2, we have for the covariance

$$\begin{aligned}
\text{Cov}[L_n, T_k] &= \mathbb{E}[L_n T_k] - \mathbb{E}[L_n] \mathbb{E}[T_k] \\
&= \mathbb{E} \left[ T_k \sum_{i=2}^n iT_i \right] - \mathbb{E} \left[ \sum_{i=2}^n iT_i \right] \mathbb{E}[T_k] \\
&= \mathbb{E}[kT_k^2] - k\mathbb{E}[T_k]^2 = k \text{Var}[T_k] = \frac{4}{k(k-1)^2}, \quad (26)
\end{aligned}$$

where the last step uses eq. 3. The covariance of  $L_n$  and  $T_k$ , like  $\text{Cov}[H_n, T_k]$  (eq. 23), is independent of  $n$ .

Now we calculate the correlation coefficient from eqs. 26, 9, and 3:

$$\text{Corr}[L_n, T_k] = \frac{1}{\sqrt{S_{2,n-1}}} \frac{1}{k-1}. \quad (27)$$

If we let  $n \rightarrow \infty$ , then this quantity becomes

$$\lim_{n \rightarrow \infty} \text{Corr}[L_n, T_k] = \frac{\sqrt{6}}{\pi} \frac{1}{k-1}. \quad (28)$$

The limiting correlation decreases to 0 with  $k$ , starting for  $k = 2$  at  $\sqrt{6}/\pi \approx 0.77970$ .

### 3.3 $H_n$ and $L_n$

Arbisser *et al.* (2018) reported the covariance and correlation of  $H_n$  and  $L_n$ . By eq. 4 and the linearity of the covariance,

$$\text{Cov}[H_n, L_n] = \sum_{k=2}^n \text{Cov}[L_n, T_k].$$

Applying eq. 26, we obtain

$$\text{Cov}[H_n, L_n] = 4S_{2,n-1} - 4 + \frac{4}{n}. \quad (29)$$

The limit of the covariance is

$$\lim_{n \rightarrow \infty} \text{Cov}[H_n, L_n] = \frac{2\pi^2}{3} - 4 \approx 2.57974. \quad (30)$$

Dividing the covariance in eq. 29 by the square root of the product of eqs. 6 and 9, we obtain

$$\text{Corr}[H_n, L_n] = \frac{S_{2,n-1}n - n + 1}{\sqrt{S_{2,n-1}(2S_{2,n}n^2 - 3n^2 + 2n - 1)}}. \quad (31)$$

The limit is

$$\lim_{n \rightarrow \infty} \text{Corr}[H_n, L_n] = \frac{\pi^2 - 6}{\pi\sqrt{2(\pi^2 - 9)}} \approx 0.93399. \quad (32)$$

### 3.4 $H_n$ and $E_n$

For the covariance  $\text{Cov}[H_n, E_n] = \mathbb{E}[H_n E_n] - \mathbb{E}[H_n] \mathbb{E}[E_n]$ , we first note that by eqs. 5 and 13, the second term is simply  $4(1 - \frac{1}{n})$ . Expanding  $\mathbb{E}[H_n E_n]$  by using the definition of  $H_n$  (eq. 4) gives us

$$\mathbb{E}[H_n E_n] = \sum_{i=2}^n \mathbb{E}[E_n T_i] = n \sum_{i=2}^n \mathbb{E}[e_n T_i],$$

as all external branch lengths are identically distributed (eq. 10).

For integers  $k, i$  with  $2 \leq k, i \leq n$ , the external branch length  $e_k$ , representing the length of a randomly chosen external branch for a tree with  $k$  leaves, and the coalescence time  $T_i$ , satisfy (eq. 11)

$$e_k T_i = \begin{cases} e_{k-1} T_i + T_k T_i, & \text{probability } \frac{k-2}{k}, \\ T_k T_i, & \text{probability } \frac{2}{k}, \end{cases} \quad (33)$$

where for convenience, we write  $e_1 = 0$ .

Note that  $e_k$  and  $T_i$  are independent for  $i > k$ , as the recurrence for  $e_k$  is constructed only using coalescence times  $T_2, T_3, \dots, T_k$  (eq. 11); each of these times is independent of  $T_i$  for  $i > k$  (Section 2.1).



We solve to find  $\mathbb{E}[e_n T_i]$  by computing  $\mathbb{E}[e_k T_i]$ , incrementing  $k$  from 2 to  $n$ . The calculations are similar to those of the Appendix of Fu & Li (1993).

$\mathbb{E}[e_2 T_2]$  is trivial, with  $e_2 = T_2$ , and  $\mathbb{E}[e_2 T_2] = \mathbb{E}[T_2^2] = 2$  by eqs. 2 and 3. By eqs. 12 and 2 and the independence of  $e_k$  and  $T_i$  for  $i > k$ , for  $i \geq 3$ ,

$$\mathbb{E}[e_{i-1} T_i] = \mathbb{E}[e_{i-1}] \mathbb{E}[T_i] = \frac{4}{i(i-1)^2}.$$

Noting  $\mathbb{E}[T_i^2] = \text{Var}[T_i] + \mathbb{E}[T_i]^2 = 2\mathbb{E}[T_i]^2$  by eqs. 2 and 3, we use eq. 33 to write an expression for  $\mathbb{E}[e_i T_i]$ :

$$\mathbb{E}[e_i T_i] = \frac{i-2}{i} \mathbb{E}[e_{i-1} T_i] + \mathbb{E}[T_i^2] = \frac{4}{i(i-1)^2}. \quad (34)$$

Next, incrementing eq. 34, we have

$$\mathbb{E}[e_{i+1} T_i] = \frac{i-1}{i+1} \mathbb{E}[e_i T_i] + \mathbb{E}[T_{i+1}] \mathbb{E}[T_i] = \frac{4}{i^2(i-1)}, \quad (35)$$

by eqs. 2 and 12.

The final step is to solve the recurrence equation

$$\mathbb{E}[e_n T_i] = \frac{n-2}{n} \mathbb{E}[e_{n-1} T_i] + \mathbb{E}[T_n] \mathbb{E}[T_i],$$

with initial condition eq. 35. Recalling the case of  $i = n = 2$ , with  $2 \leq i \leq n$ , we obtain solution

$$\mathbb{E}[e_n T_i] = \frac{4}{i(i-1)(n-1)}. \quad (36)$$

Applying eq. 10, the expression for  $\text{Cov}[H_n, E_n]$  becomes

$$\text{Cov}[H_n, E_n] = n \sum_{i=2}^n \frac{4}{i(i-1)(n-1)} - 4 \left(1 - \frac{1}{n}\right) = \frac{4}{n}. \quad (37)$$

The limit of the covariance as  $n \rightarrow \infty$  is

$$\lim_{n \rightarrow \infty} \text{Cov}[H_n, E_n] = 0. \quad (38)$$

Dividing eq. 37 by the square root of the product of variances from eqs. 6 and 14, the correlation is

$$\text{Corr}[H_n, E_n] = \begin{cases} 1, & n = 2, \\ \frac{\sqrt{(n-1)(n-2)}}{\sqrt{2(2S_{2,n}n^2 - 3n^2 + 2n - 1)(S_{1,n-1}n - 2n + 2)}}, & n > 2. \end{cases} \quad (39)$$

The limit of the correlation is

$$\lim_{n \rightarrow \infty} \text{Corr}[H_n, E_n] = 0. \quad (40)$$

### 3.5 $E_n$ and $T_k$

In the process of computing  $\text{Cov}[H_n, E_n]$ , we have obtained an expression for  $\mathbb{E}[e_n T_i]$  (eq. 36), from which we can obtain  $\text{Cov}[E_n, T_k] = n\mathbb{E}[e_n T_k] - \mathbb{E}[E_n]\mathbb{E}[T_k]$ . Applying eqs. 13 and 2, we have

$$\text{Cov}[E_n, T_k] = \frac{4}{k(k-1)(n-1)}. \quad (41)$$

Irrespective of the value of  $k$ , we have

$$\lim_{n \rightarrow \infty} \text{Cov}[E_n, T_k] = 0. \quad (42)$$

Applying eqs. 3 and 14, the correlation coefficient is

$$\text{Corr}[E_n, T_k] = \begin{cases} 1, & n = 2, \\ \frac{\sqrt{n-2}}{\sqrt{2(n-1)(S_{1,n-1}n-2n+2)}}, & n \geq 3. \end{cases} \quad (43)$$

The correlation coefficient is independent of  $k$ , and it has limit

$$\lim_{n \rightarrow \infty} \text{Corr}[E_n, T_k] = 0. \quad (44)$$

### 3.6 $L_n$ and $E_n$

Fu & Li (1993) provided the expression for  $\mathbb{E}[L_n E_n]$  (see also p. 167 of Durrett (2008), with all values scaled by  $2N_e$ ). The main result is the following expression, obtained by solving recurrence equations:

$$\mathbb{E}[L_n E_n] = \frac{4S_{1,n-1}n}{n-1}.$$

We can use this result to calculate the covariance of  $L_n$  and  $E_n$  by  $\text{Cov}[L_n, E_n] = \mathbb{E}[L_n E_n] - \mathbb{E}[L_n]\mathbb{E}[E_n]$  with eqs. 8 and 13. The covariance can also be quickly obtained from eqs. 7 and 41,

$$\text{Cov}[L_n, E_n] = \sum_{k=2}^n k \text{Cov}[E_n, T_k] = \frac{4S_{1,n-1}}{n-1}. \quad (45)$$

The limit is

$$\lim_{n \rightarrow \infty} \text{Cov}[L_n, E_n] = 0. \quad (46)$$

Applying eqs. 45, 9, and 14, the correlation coefficient of  $L_n$  and  $E_n$  is

$$\text{Corr}[L_n, E_n] = \begin{cases} 1, & n = 2, \\ \frac{S_{1,n-1}\sqrt{n-2}}{\sqrt{2S_{2,n-1}(n-1)(S_{1,n-1}n-2n+2)}}, & n \geq 3, \end{cases} \quad (47)$$

with the limit

$$\lim_{n \rightarrow \infty} \text{Corr}[L_n, E_n] = 0. \quad (48)$$

### 3.7 $H_n$ and $I_n$

For the pair  $H_n$  and  $I_n$ , we exploit results obtained for other pairs to quickly obtain the covariance.

Remembering that  $I_n = L_n - E_n$  (eq. 15), we use eqs. 29 and 37 to obtain for  $n \geq 3$

$$\begin{aligned} \text{Cov}[H_n, I_n] &= \text{Cov}[H_n, L_n] - \text{Cov}[H_n, E_n] \\ &= 4S_{2,n-1} - 4. \end{aligned} \quad (49)$$

For this covariance, we have

$$\lim_{n \rightarrow \infty} \text{Cov}[H_n, I_n] = \frac{2\pi^2}{3} - 4 \approx 2.57974. \quad (50)$$

From the covariance in eq. 49 and variances in eqs. 6 and 17, we compute the correlation coefficient:

$$\text{Corr}[H_n, I_n] = \frac{(S_{2,n-1} - 1) n \sqrt{(n-1)(n-2)}}{\sqrt{(2S_{2,n}n^2 - 3n^2 + 2n - 1) [4S_{1,n-1} + S_{2,n-1}(n-1)(n-2) - 4(n-1)]}}. \quad (51)$$

The limit is the same as that of  $\text{Corr}[H_n, L_n]$ , or (eq. 32)

$$\lim_{n \rightarrow \infty} \text{Corr}[H_n, I_n] = \frac{\pi^2 - 6}{\pi \sqrt{2(\pi^2 - 9)}} \approx 0.93399. \quad (52)$$

### 3.8 $I_n$ and $T_k$

By eqs. 15, 26, and 41, assuming  $n \geq 3$ , we have

$$\begin{aligned} \text{Cov}[I_n, T_k] &= \text{Cov}[L_n, T_k] - \text{Cov}[E_n, T_k] \\ &= \frac{4(n-k)}{k(k-1)^2(n-1)}. \end{aligned} \quad (53)$$

The limit of this expression is a rapidly decreasing function of  $k$ ,

$$\lim_{n \rightarrow \infty} \text{Cov}[I_n, T_k] = \frac{4}{k(k-1)^2}. \quad (54)$$

Using the variances in eqs. 17 and 3, the correlation coefficient is

$$\text{Corr}[I_n, T_k] = \frac{(n-k)\sqrt{n-2}}{(k-1)\sqrt{(n-1) [4S_{1,n-1} + S_{2,n-1}(n-1)(n-2) - 4(n-1)]}}, \quad (55)$$

with limit

$$\lim_{n \rightarrow \infty} \text{Corr}[I_n, T_k] = \frac{\sqrt{6}}{\pi} \frac{1}{k-1}. \quad (56)$$

The limit follows that of  $\text{Corr}[L_n, T_k]$  (eq. 28),

### 3.9 $L_n$ and $I_n$

By eq. 15, we can apply eqs. 9 and 45 to obtain for  $n \geq 3$

$$\begin{aligned} \text{Cov}[L_n, I_n] &= \text{Var}[L_n] - \text{Cov}[L_n, E_n] \\ &= 4S_{2,n-1} - \frac{4S_{1,n-1}}{n-1}. \end{aligned} \quad (57)$$

The limit as  $n \rightarrow \infty$  is

$$\lim_{n \rightarrow \infty} \text{Cov}[L_n, I_n] = \frac{2\pi^2}{3} \approx 6.57974. \quad (58)$$

For the correlation coefficient, applying eqs. 57, 9, and 17, we get

$$\text{Corr}[L_n, I_n] = \frac{[S_{2,n-1}(n-1) - S_{1,n-1}]\sqrt{n-2}}{(n-1)\sqrt{S_{2,n-1}[4S_{1,n-1} + S_{2,n-1}(n-1)(n-2) - 4(n-1)]}}, \quad (59)$$

with

$$\lim_{n \rightarrow \infty} \text{Corr}[L_n, I_n] = 1. \quad (60)$$

### 3.10 $E_n$ and $I_n$

For this pair, with  $n \geq 3$ , the covariance was reported by Fu & Li (1993):

$$\text{Cov}[E_n, I_n] = \frac{4S_{1,n-1}}{n-1} - \frac{8S_{1,n-1}n}{(n-1)(n-2)} + \frac{16}{n-2}. \quad (61)$$

We can also obtain this result quickly from eqs. 15, 57, and 17, as  $\text{Cov}[E_n, I_n] = \text{Cov}[L_n - I_n, I_n] = \text{Cov}[L_n, I_n] - \text{Var}[I_n]$ . In the limit, we have

$$\lim_{n \rightarrow \infty} \text{Cov}[E_n, I_n] = 0. \quad (62)$$

For the correlation coefficient, we divide eq. 61 by the product of the square roots of eqs. 14 and 17:

$$\text{Corr}[E_n, I_n] = \frac{4(n-1) - S_{1,n-1}(n+2)}{\sqrt{2(S_{1,n-1}n - 2n + 2)[4S_{1,n-1} + S_{2,n-1}(n-1)(n-2) - 4(n-1)]}}, \quad (63)$$

with the limit

$$\lim_{n \rightarrow \infty} \text{Corr}[E_n, I_n] = 0. \quad (64)$$

This result matches the limit for  $\text{Corr}[L_n, E_n]$  (eq. 48).

### 3.11 Var $[B_n]$

To obtain correlation coefficients involving  $B_n$ , assuming  $n \geq 4$ , we first verify the expression for  $\text{Var}[B_n]$  in eq. 22. By definition of  $B_n$  in eq. 18, we have

$$\text{Var}[B_n] = \mathbb{E} \left[ \frac{1}{4}(T_2 + b_n)^2 \right] - \mathbb{E} \left[ \frac{1}{2}(T_2 + b_n) \right]^2 = \frac{1}{4} \text{Var}[b_n] + \frac{1}{2} \text{Cov}[b_n, T_2] + \frac{1}{4}, \quad (65)$$

where we have used  $\text{Var}[T_2] = 1$  (eq. 3).

To calculate  $\text{Var}[b_n]$ , we first recall that for  $j = 3, 4, \dots, n-1$ , with probability  $2/[j(j-1)]$ , we have  $b_n = H_j$ ; with probability  $2/(n-1)$  we have  $b_n = H_n$ . Hence, applying eq. 19 and  $\mathbb{E}[H_k^2] = \text{Var}[H_k] + \mathbb{E}[H_k]^2$  with eqs. 5 and 6, we have

$$\begin{aligned} \mathbb{E}[b_n^2] &= \left[ \sum_{j=3}^{n-1} \frac{2}{j(j-1)} \mathbb{E}[H_j^2] \right] + \frac{2}{n-2} \mathbb{E}[H_n^2] \\ &= \frac{-16S_{2,n-1}n^2 + 30n^2 - 16n - 16}{n^2}. \end{aligned}$$

Using the expression for  $\mathbb{E}[b_n]$  from eq. 20, we obtain

$$\text{Var}[b_n] = \frac{24S_{2,n-1}n^2 - 16S_{2,n-1}^2n^2 + 5n^2 - 32S_{2,n-1}n + 24n - 32}{n^2}. \quad (66)$$

Next, we compute  $\text{Cov}[b_n, T_k]$  and insert  $k = 2$ . By eq. 19, applying the independence of the  $T_i$  (Section 2.1) and inserting eq. 3, we have

$$\begin{aligned} \text{Cov}[b_n, T_k] &= \left[ \sum_{j=3}^{n-1} \sum_{i=2}^j \frac{2}{j(j-1)} \text{Cov}[T_i, T_k] \right] + \left( \sum_{i=2}^n \frac{2}{n-1} \text{Cov}[T_i, T_k] \right) \\ &= \left[ \sum_{i=2}^{n-1} \sum_{\substack{j=i \text{ if } i \geq 3 \\ j=3 \text{ if } i=2}}^{n-1} \frac{2}{j(j-1)} \text{Cov}[T_i, T_k] \right] + \left( \frac{2}{n-1} \text{Var}[T_k] \right) \\ &= \begin{cases} \left[ \sum_{j=3}^{n-1} \frac{2}{j(j-1)} \text{Var}[T_k] \right] + \left( \frac{2}{n-1} \text{Var}[T_k] \right), & k = 2, \\ \left[ \sum_{j=k}^{n-1} \frac{2}{j(j-1)} \text{Var}[T_k] \right] + \left( \frac{2}{n-1} \text{Var}[T_k] \right), & k = 3, 4, \dots, n-1, \\ \frac{2}{n-1} \text{Var}[T_k], & k = n, \end{cases} \\ &= \begin{cases} 1, & k = 2, \\ \frac{8}{k^2(k-1)^3}, & k = 3, 4, \dots, n. \end{cases} \end{aligned} \quad (67)$$

Inserting  $\text{Var}[b_n]$  from eq. 66 and  $\text{Cov}[b_n, T_2]$  from eq. 67 into eq. 65, we confirm eq. 22.

### 3.12 $B_n$ and $T_k$

We extract  $\text{Cov}[B_n, T_k]$  from Section 3.11, as  $\text{Cov}[B_n, T_k] = \text{Cov}[b_n, T_k]/2 + \text{Cov}[T_2, T_k]/2$  by the definition in eq. 18, and  $\text{Cov}[T_2, T_k] = \delta_{k,2}$ , where  $\delta$  is the Kronecker delta (Section 2.1). By eq. 67, recalling  $n \geq 4$ ,

$$\text{Cov}[B_n, T_k] = \frac{4}{k^2(k-1)^3}. \quad (68)$$

The covariance is independent of  $n$ .

For the correlation coefficient, using eqs. 68, 22, and 3, we have

$$\text{Corr}[B_n, T_k] = \frac{\sqrt{2}n}{\sqrt{3S_{2,n-1}n^2 - 2S_{2,n-1}^2n^2 + n^2 - 4S_{2,n-1}n + 3n - 4}} \frac{1}{k(k-1)^2}, \quad (69)$$

with the limit

$$\lim_{n \rightarrow \infty} \text{Corr}[B_n, T_k] = \frac{6}{\sqrt{18 + 9\pi^2 - \pi^4}} \frac{1}{k(k-1)^2}. \quad (70)$$

The limit begins at  $3/\sqrt{18 + 9\pi^2 - \pi^4} \approx 0.97759$  for  $k = 2$  and quickly decreases to 0 as  $k$  increases.

### 3.13 $H_n$ and $B_n$

To obtain  $\text{Cov}[H_n, B_n]$  with  $n \geq 4$ , we begin from eq. 18:

$$\text{Cov}[H_n, B_n] = \frac{1}{2} \text{Cov}[H_n, b_n] + \frac{1}{2} \text{Cov}[H_n, T_2].$$

The second term was computed in eq. 23,  $\text{Cov}[H_n, T_2] = 1$ .

For the first term,  $\text{Cov}[H_n, b_n]$ , we decompose  $H_n$  (eq. 4) and apply eq. 67 to obtain

$$\begin{aligned} \text{Cov}[H_n, b_n] &= \sum_{k=2}^n \text{Cov}[b_n, T_k] \\ &= 1 + \sum_{k=3}^n \frac{8}{k^2(k-1)^3}. \end{aligned}$$

We use a partial fraction decomposition to sum the series, obtaining

$$\begin{aligned} \text{Cov}[H_n, B_n] &= 1 + \sum_{k=3}^n \frac{4}{k^2(k-1)^3} \\ &= \frac{4[S_{3,n-1}n^2 - 3S_{2,n-1}n^2 + (n-1)(4n+1)]}{n^2}. \end{aligned} \quad (71)$$

The asymptotic limit of  $\text{Cov}[H_n, B_n]$  is

$$\lim_{n \rightarrow \infty} \text{Cov}[H_n, B_n] = 4\zeta(3) + 16 - 2\pi^2 \approx 1.06902. \quad (72)$$

The correlation coefficient is then equal to:

$$\text{Corr}[H_n, B_n] = \frac{\sqrt{2}[S_{3,n-1}n^2 - 3S_{2,n-1}n^2 + (n-1)(4n+1)]}{\sqrt{(2S_{2,n}n^2 - 3n^2 + 2n - 1)(3S_{2,n-1}n^2 - 2S_{2,n-1}^2n^2 + n^2 - 4S_{2,n-1}n + 3n - 4)}}. \quad (73)$$

The limit of the correlation coefficient is quite close to 1:

$$\lim_{n \rightarrow \infty} \text{Corr}[H_n, B_n] = \frac{3\sqrt{3}[2\zeta(3) + 8 - \pi^2]}{\sqrt{(\pi^2 - 9)(18 + 9\pi^2 - \pi^4)}} \approx 0.97054. \quad (74)$$

### 3.14 $L_n$ and $B_n$

In a manner similar to that used in Section 3.13, with  $n \geq 4$ , we expand  $\text{Cov}[L_n, B_n]$  using eq. 18:

$$\text{Cov}[L_n, B_n] = \frac{1}{2} \text{Cov}[L_n, b_n] + \frac{1}{2} \text{Cov}[L_n, T_2].$$

The second term is  $\text{Cov}[L_n, T_2] = 2$  by eq. 26. The first term is decomposable by eq. 7; applying eq. 67,

$$\begin{aligned} \text{Cov}[L_n, b_n] &= \sum_{k=2}^n k \text{Cov}[b_n, T_k] \\ &= 2 + \sum_{k=3}^n \frac{8}{k(k-1)^3}. \end{aligned}$$

Summing the series, we have

$$\begin{aligned} \text{Cov}[L_n, B_n] &= 2 + \sum_{k=3}^n \frac{4}{k(k-1)^3} \\ &= \frac{4[S_{3,n-1}n - S_{2,n-1}n + n - 1]}{n}. \end{aligned} \quad (75)$$

The limiting covariance is

$$\lim_{n \rightarrow \infty} \text{Cov}[L_n, B_n] = 4\zeta(3) + 4 - \frac{2\pi^2}{3} \approx 2.22849. \quad (76)$$

Using eqs. 75, 9, and 22, we now obtain an expression for the correlation coefficient:

$$\text{Corr}[L_n, B_n] = \frac{\sqrt{2}(S_{3,n-1}n - S_{2,n-1}n + n - 1)}{\sqrt{S_{2,n-1}(3S_{2,n-1}n^2 - 2S_{2,n-1}^2n^2 + n^2 - 4S_{2,n-1}n + 3n - 4)}}. \quad (77)$$

The limit is

$$\lim_{n \rightarrow \infty} \text{Corr}[L_n, B_n] = \frac{\sqrt{6}[6\zeta(3) + 6 - \pi^2]}{\pi\sqrt{18 + 9\pi^2 - \pi^4}} \approx 0.84930. \quad (78)$$

### 3.15 $E_n$ and $B_n$

For  $\text{Cov}[E_n, B_n]$ , we obtain an approximate rather than exact answer. Decomposing  $B_n$  by eq. 18, we have

$$\text{Cov}[E_n, B_n] = \frac{1}{2} \text{Cov}[E_n, T_2] + \frac{1}{2} \text{Cov}[E_n, b_n]. \quad (79)$$

Recall that  $b_n$  can be defined conditionally (Section 2.6). For a random variable  $J$ ,  $b_n = H_j$  with probability  $p_j$ , where  $p_j = \mathbb{P}[J = j] = 2/[j(j-1)]$  for  $J = 3, 4, \dots, n-1$  and  $p_j = 2/(j-1)$  for  $J = n$ . We can then decompose the covariance  $\text{Cov}[E_n, b_n]$  by the conditional covariance formula, conditioning on  $J$ :

$$\text{Cov}[E_n, b_n] = \mathbb{E} \left[ \text{Cov}[E_n, b_n | J] \right] + \text{Cov} \left[ \mathbb{E}[E_n | J], \mathbb{E}[B_n | J] \right]. \quad (80)$$

We next perform an approximation by ignoring the second term in the covariance decomposition. Noting that  $\text{Cov}[E_n, T_2] = \frac{2}{n-1}$  by eq. 41, we use eq. 79 together with eq. 80 to write approximations

$$\widetilde{\text{Cov}}[E_n, b_n] = \mathbb{E} \left[ \text{Cov}[E_n, b_n | J] \right] \quad (81)$$

$$\widetilde{\text{Cov}}[E_n, B_n] = \frac{1}{n-1} + \frac{1}{2} \widetilde{\text{Cov}}[E_n, b_n]. \quad (82)$$

Weighting each  $\text{Cov}[H_j, E_n]$  by the associated probability  $p_j$ , and decomposing  $H_j$  by eq. 4, eq. 81 gives

$$\begin{aligned} \mathbb{E} \left[ \text{Cov}[E_n, b_n | J] \right] &= \sum_{j=3}^n p_j \text{Cov}[E_n, b_n | J = j] \\ &= \left[ \sum_{j=3}^{n-1} \frac{2}{j(j-1)} \text{Cov}[H_j, E_n] \right] + \frac{2}{n-1} \text{Cov}[H_n, E_n] \\ &= \left[ \sum_{j=3}^{n-1} \frac{2}{j(j-1)} \sum_{k=2}^j \text{Cov}[E_n, T_k] \right] + \frac{2}{n-1} \sum_{k=2}^j \text{Cov}[E_n, T_k]. \end{aligned}$$

We can then insert the result of eq. 41 and simplify to obtain

$$\widetilde{\text{Cov}}[E_n, b_n] = \frac{2(4S_{2,n-1}n - 5n + 4)}{n(n-1)}. \quad (83)$$

Finally, inserting eq. 83 into eq. 82, we have

$$\widetilde{\text{Cov}}[E_n, B_n] = \frac{4(S_{2,n-1}n - n + 1)}{n(n-1)}. \quad (84)$$

The limit is

$$\lim_{n \rightarrow \infty} \widetilde{\text{Cov}}[E_n, B_n] = 0. \quad (85)$$



For the approximate correlation coefficient  $\widetilde{\text{Corr}}[E_n, B_n] = \widetilde{\text{Cov}}[E_n, B_n] / \sqrt{\text{Var}[E_n] \text{Var}[B_n]}$ , we use eqs. 84, 14, and 22 to obtain

$$\widetilde{\text{Corr}}[E_n, B_n] = \frac{(S_{2,n-1}n - n + 1)\sqrt{n-2}}{\sqrt{(n-1)(S_{1,n-1}n - 2n + 2)(3S_{2,n-1}n^2 - 2S_{2,n-1}^2n^2 + n^2 - 4S_{2,n-1}n + 3n - 4)}}, \quad (86)$$

with limit

$$\lim_{n \rightarrow \infty} \widetilde{\text{Corr}}[E_n, B_n] = 0. \quad (87)$$

### 3.16 $I_n$ and $B_n$

We use eq. 15 and results involving  $L_n$  (eq. 75) and  $E_n$  (eq. 84) to obtain

$$\begin{aligned} \widetilde{\text{Cov}}[I_n, B_n] &= \text{Cov}[L_n, B_n] - \widetilde{\text{Cov}}[E_n, B_n] \\ &= \frac{4(S_{3,n-1}n - S_{2,n-1}n + n - S_{3,n-1} - 1)}{n-1}, \end{aligned} \quad (88)$$

with limit

$$\lim_{n \rightarrow \infty} \widetilde{\text{Cov}}[I_n, B_n] = 4\zeta(3) + 4 - \frac{2\pi^2}{3} \approx 2.22849. \quad (89)$$

Dividing eq. 88 by the product of the square roots of eqs. 17 and 22, the approximate correlation is

$$\widetilde{\text{Corr}}[I_n, B_n] = \frac{\sqrt{2}(S_{3,n-1}n - S_{2,n-1}n + n - S_{3,n-1} - 1)n\sqrt{n-2}}{\sqrt{(n-1)[4S_{1,n-1} + S_{2,n-1}(n-1)(n-2) - 4(n-1)](3S_{2,n-1}n^2 - 2S_{2,n-1}^2n^2 + n^2 - 4S_{2,n-1}n + 3n - 4)}}, \quad (90)$$

with limit

$$\lim_{n \rightarrow \infty} \widetilde{\text{Corr}}[I_n, B_n] = \frac{\sqrt{6}[6\zeta(3) + 6 - \pi^2]}{\pi\sqrt{18 + 9\pi^2 - \pi^4}} \approx 0.84930. \quad (91)$$

## 4 Numerical and simulation-based analysis

### 4.1 Analysis methods

We examine the results of Section 3 summarized in Tables 1 and 2 numerically and by coalescent simulation. For 13 of 15 covariances and correlations, the theoretical results are exact, and simulations merely verify that the mathematics has proceeded without error. For the covariances and correlations involving  $(E_n, B_n)$  and  $(I_n, B_n)$ , the theoretical results are approximate, and the simulations assess the accuracy of the approximations.

We simulated the coalescent process for a series of values of  $n$  beginning with  $n = 2$ , at each value of  $n$  performing 100,000 replicate simulations. To generate the simulated replicates, we employed `ms` (Hudson, 2002), using the command `ms n 100000 -T`, with  $n$  taken from  $\{2, 3, \dots, 50\}$ . In the set of simulated replicates, we evaluated simulated covariances and correlation coefficients for pairs of quantities.

We plot the mathematical results of Tables 1 and 2 together with simulation values in Figures 2-5. Figures 2 and 3 show covariances of pairs of variables; Figures 4 and 5 show correlations.

## 4.2 Accuracy of approximations

Figure 2 shows the analytical and simulated covariances, and Figure 4 shows the analytical and simulated correlations, for pairs of variables among  $\{H_n, L_n, I_n, E_n, B_n\}$ . For pairs of variables for which no approximations were needed in obtaining covariances—all except  $(E_n, B_n)$  and  $(I_n, B_n)$ —the simulated and analytical values produce plots that are nearly indistinguishable.

For  $(E_n, B_n)$  and  $(I_n, B_n)$ , the approximate and simulated correlations are close, but noticeably different (Figure 4); the mean absolute difference between the analytical and simulated values across choices of  $n$  from 4 to 30 is 0.02458 for  $(E_n, B_n)$  and 0.01089 for  $(I_n, B_n)$ . For covariance, which unlike the correlation coefficient is not standardized to lie in  $[-1, 1]$ , the approximate and simulated values are quite close, with corresponding mean absolute deviations of 0.02372 for  $(E_n, B_n)$  and 0.03101 for  $(I_n, B_n)$ .

## 4.3 Properties of correlations

We observe that  $H_n$ ,  $L_n$ ,  $I_n$ , and  $B_n$  all remain strongly correlated as  $n$  increases, with the six limiting correlations among these four quantities lying between 0.84930 for  $\text{Corr}[L_n, B_n]$  and  $\text{Corr}[I_n, B_n]$  and 1 for  $\text{Corr}[L_n, I_n]$  (Table 2). The high limiting  $\text{Corr}[H_n, L_n]$  of approximately 0.93399 reflects the strong influence of times  $T_k$  with small  $k$  on both  $H_n$  and  $L_n$  (Figures 3 and 5). As  $n$  increases,  $\mathbb{E}[I_n]$  increases without bound (eq. 16), whereas  $\mathbb{E}[E_n]$  remains constant (eq. 13); the contribution of  $E_n$  to the total tree length  $L_n$  becomes negligible, and  $\text{Corr}[L_n, I_n]$  approaches 1.  $\text{Corr}[H_n, I_n]$  has the same limiting value as  $\text{Corr}[H_n, L_n]$ , and  $H_n$ ,  $L_n$ , and  $I_n$  all have limiting correlation 0 with  $E_n$ . Interestingly, although  $H_n$  and  $E_n$  have the same limiting expectation of 2, the limit of their correlation  $\text{Corr}[H_n, E_n]$  is 0.

The correlations of  $H_n$ ,  $L_n$ , and  $I_n$  with  $B_n$ , like their correlations with each other, are relatively high.

$\text{Corr}[H_n, B_n]$  is nearly constant in  $n$ , with limit approximately 0.97054; both  $H_n$  and  $B_n$  are determined in large part by the  $T_k$  with small  $k$  (eqs. 5 and 21), so that little change occurs in the correlation as  $n$  increases. Because  $\text{Corr}[H_n, B_n]$  is high and  $\text{Corr}[H_n, L_n]$  is also high, the constraint on a correlation  $\text{Corr}[Y, Z]$  given  $\text{Corr}[X, Y]$  and  $\text{Corr}[X, Z]$ , or (Wickens, 2014, eq. 7.1),

$$\text{Corr}[Y, Z] \geq \text{Corr}[X, Y] \text{Corr}[X, Z] - \sqrt{1 - \text{Corr}[X, Y]^2} \sqrt{1 - \text{Corr}[X, Z]^2} \quad (92)$$

$$\text{Corr}[Y, Z] \leq \text{Corr}[X, Y] \text{Corr}[X, Z] + \sqrt{1 - \text{Corr}[X, Y]^2} \sqrt{1 - \text{Corr}[X, Z]^2}, \quad (93)$$

forces a high value for  $\text{Corr}[L_n, B_n]$  as well. In particular, placing  $H_n, L_n, B_n$  in the roles of  $X, Y, Z$ , with  $\lim_{n \rightarrow \infty} \text{Corr}[H_n, L_n] \approx 0.93399$  and  $\lim_{n \rightarrow \infty} \text{Corr}[H_n, B_n] \approx 0.97054$ , we obtain an interval  $0.82037 \leq \lim_{n \rightarrow \infty} \text{Corr}[L_n, B_n] \leq 0.99256$  from eqs. 92 and 93;  $\lim_{n \rightarrow \infty} \text{Corr}[L_n, B_n] \approx 0.84930$  lies near its lower end. Eqs. 92 and 93 similarly force a high value for  $\lim_{n \rightarrow \infty} \text{Corr}[I_n, B_n]$ , using  $H_n, I_n, B_n$  as  $X, Y, Z$ .

Next, for correlations involving the  $T_k$ , we observe that for fixed  $n$ , as  $k$  increases from 2 to  $n$ ,  $\text{Corr}[H_n, T_k]$  decreases (Figure 5). At fixed  $n$  and  $k$ ,  $\text{Corr}[L_n, T_k]$  generally exceeds  $\text{Corr}[H_n, T_k]$ ;  $k$  copies of the branch length  $T_k$  contribute to tree length  $L_n$  (eq. 7), whereas only one copy contributes to the tree height  $H_n$  (eq. 4), giving rise to a greater value for the correlation of  $T_k$  with  $L_n$  than with  $H_n$ . For  $k > 2$ ,  $\text{Corr}[B_n, T_k]$  is generally smaller than  $\text{Corr}[H_n, T_k]$ ; because  $B_n$  is determined to a larger extent by  $T_2$  than is  $H_n$ , the correlations of  $B_n$  with  $T_k$  for  $k > 2$  are generally smaller. Finally, because tree length  $L_n$  consists primarily of internal branches for large  $n$ , the correlation  $\text{Corr}[I_n, T_k]$  is similar to  $\text{Corr}[L_n, T_k]$  (Figure 5), approaching the same limit as  $n \rightarrow \infty$  (Table 2); the correlation of  $E_n$  and  $T_k$  is a constant that does not depend on  $k$ .

## 5 Discussion

We have examined relationships between pairs of tree features under the coalescent model by deriving expressions for their covariances and correlation coefficients (Tables 1 and 2). For 13 of 15 pairs examined, we obtained exact expressions for the covariances and correlation coefficients, and for the remaining two pairs, we obtained quantities observed in simulations to closely approximate the desired quantities (Figures 2 and 4). The results provide a compendium of basic relationships among coalescent tree features, contributing to a more precise understanding of the way in which the properties of coalescent trees relate to each other.

In most cases, the covariances have relatively simple expressions, comparable to the simplicity of most

expressions for expectations and variances (Table 1). Expressions for the correlation coefficients are somewhat more complex, in many cases with  $n \rightarrow \infty$  limits that contain terms resulting from the limit  $\sum_{k=1}^{\infty} 1/k^2 = \pi^2/6$ .

Numerically, we obtain tight correlations between  $H_n$ ,  $L_n$ ,  $I_n$ , and  $B_n$  as  $n$  grows large, with all of these quantities possessing limiting correlations of 0.84930 or greater (Table 2). In the limit,  $L_n$  and  $I_n$  are perfectly correlated, and all limiting correlations of other quantities with  $I_n$  are equal to their corresponding correlations with  $L_n$ . Decreasing correlations are observed for  $H_n$ ,  $L_n$ ,  $I_n$ , and  $B_n$  with  $E_n$ , with limits of 0 observed in all cases (Table 2). Although  $H_n$  and  $E_n$  both have limiting expectation 2 (eqs. 5 and 13), their limiting correlation coefficient is 0. The correlations among  $H_n$ ,  $L_n$ , and  $B_n$  are all large; however, the limiting correlation for  $(L_n, B_n)$  is near the lower end of the interval suggested by the larger limiting correlations for  $(H_n, L_n)$  and  $(H_n, B_n)$  (eqs. 92 and 93). This result suggests that  $L_n$  and  $B_n$  capture relatively distinct features of coalescent trees in relation to the constraints placed on a pair of correlated variables that are each highly correlated with a third variable ( $H_n$ ). A similar observation can be made concerning  $I_n$  and  $B_n$ , as  $L_n$  and  $I_n$  are asymptotically fully correlated.

Interest in tree properties such as  $H_n$ ,  $L_n$ ,  $E_n$ ,  $I_n$ , and  $B_n$  arises in part from their relationship to statistical tests that assess the fit of the coalescent model to data on genetic variation. Features of tree shape underlie predictions of the coalescent regarding allele frequencies; in particular, tree properties contribute to predictions for the unfolded site-frequency spectrum (SFS) of a genomic region, the vector that for a sample of size  $n$  tabulates how many variable (biallelic) sites in the region possess allele frequencies  $1/n, 2/n, \dots, (n-1)/n$  for the derived allele (e.g. Fu, 1995; Ferretti *et al.*, 2017). Test statistics then assess agreement of site-frequency spectra with the predictions (e.g. Achaz, 2009; Ferretti *et al.*, 2010; Ronen *et al.*, 2013), so that correlations among statistics emphasizing different aspects of site-frequency spectra emerge from dependence on correlated tree features. In this context, further understanding of correlations among tree properties can assist in understanding the behavior of SFS-based tests of the coalescent model.

The computations augment earlier calculations concerning quantities associated with coalescent trees. The pairs  $(H_n, L_n)$  (Arbisser *et al.*, 2018) and  $(L_n, E_n)$  and  $(E_n, I_n)$  (Fu & Li, 1993) have been studied in some detail; we have added consideration of many additional pairs of features. For the variable  $B_n$ , used by Uyenoyama (1997) but not commonly examined in studies of coalescent quantities, we derived an expression for its variance (eq. 22), as well as exact covariances and correlations with  $H_n$ ,  $L_n$ , and  $T_k$  and approximate

covariances and correlations with  $E_n$  and  $I_n$ . Several studies have extended the work of Fu & Li (1993) on features of the external and internal branch lengths (Blum & François, 2005; Caliebe *et al.*, 2007; Janson & Kersting, 2011; Dahmer & Kersting, 2015, 2017; Disanto & Wiehe, 2020); it may be possible to seek exact rather than approximate covariances and correlations for  $(E_n, B_n)$  and  $(I_n, B_n)$  by building on these studies.

When examining joint distributions of  $H_n$  and  $L_n$ , Arbisser *et al.* (2018) obtained not only the covariance and correlation under the coalescent, but also approximations for the expectation and variance of the ratio  $H_n/L_n$ . These approximations relied on computations of the means of  $H_n$  and  $L_n$ , the variances of  $H_n$  and  $L_n$ , and the covariance of  $H_n$  and  $L_n$ . If we rely on the same approach as Arbisser *et al.* (2018), because we have computed the exact or approximate covariances for pairs of variables among  $\{H_n, L_n, E_n, I_n, B_n, T_k\}$ , similar approximations for expectations and variances of ratios are now possible to obtain for additional pairs.

**Acknowledgments.** We acknowledge support from NIH grants R01 GM131404 and R01 HG005855.

## References

- Achaz, G. 2009. Frequency spectrum neutrality tests: one for all and all for one, *Genetics* **183** (1), 249–258.
- Arbisser, I. M., Jewett, E. M., and Rosenberg, N. A. 2018. On the joint distribution of tree height and tree length under the coalescent, *Theoretical Population Biology* **122**, 46–56.
- Blum, M. G. B. and François, O. 2005. Minimal clade size and external branch length under the neutral coalescent, *Advances in Applied Probability* **37**, 647–662.
- Caliebe, A., Neininger, R., Krawczak, M., and Rösler, U. 2007. On the length distribution of external branches in coalescence trees: genetic diversity within species, *Theoretical Population Biology* **72**, 245–252.
- Dahmer, I. and Kersting, G. 2015. The internal branch lengths of the Kingman coalescent, *Annals of Applied Probability* **25** (3), 1325–1348.
- Dahmer, I. and Kersting, G. 2017. The total external length of the evolving Kingman coalescent, *Probability Theory and Related Fields* **167** (3-4), 1165–1214.
- Disanto, F. and Wiehe, T. 2020. Mapping the external branches of a Kingman tree: a discrete approach, *Theoretical Population Biology* **134**, 92–105.

- Durrett, R. 2008. “Probability Models for DNA Sequence Evolution”, Springer-Verlag, New York, second edition.
- Ferretti, L., Ledda, A., Wiehe, T., Achaz, G., and Ramos-Onsins, S. E. 2017. Decomposing the site frequency spectrum: the impact of tree topology on neutrality tests, *Genetics* **207** (1), 229–240.
- Ferretti, L., Perez-Enciso, M., and Ramos-Onsins, S. 2010. Optimal neutrality tests based on the frequency spectrum, *Genetics* **186** (1), 353–365.
- Fu, Y.-X. 1995. Statistical properties of segregating sites, *Theoretical Population Biology* **48** (2), 172–197.
- Fu, Y.-X. 1996. Estimating the age of the common ancestor of a DNA sample using the number of segregating sites, *Genetics* **144** (2), 829–838.
- Fu, Y.-X. and Li, W.-H. 1993. Statistical tests of neutrality of mutations, *Genetics* **133** (3), 693–709.
- Griffiths, R. and Tavaré, S. 1996. Monte Carlo inference methods in population genetics, *Mathematical and Computer Modelling* **23** (8-9), 141–158.
- Hein, J., Schierup, M., and Wiuf, C. 2005. “Gene Genealogies, Variation and Evolution”, Oxford University Press, Oxford.
- Hudson, R. R. 2002. Generating samples under a Wright–Fisher neutral model of genetic variation, *Bioinformatics* **18** (2), 337–338.
- Janson, S. and Kersting, G. 2011. On the total external length of the Kingman coalescent, *Electronic Journal of Probability* **16**, 2203–2218.
- Ronen, R., Udpa, N., Halperin, E., and Bafna, V. 2013. Learning natural selection from the site frequency spectrum, *Genetics* **195** (1), 181–193.
- Rosenberg, N. A. 2006. Gene genealogies, in “Evolutionary Genetics: Concepts and Case Studies” (C. W. Fox and J. B. Wolf, eds), pp. 173–189, Oxford University Press, Oxford.
- Rosenberg, N. A. and Hirsh, A. E. 2003. On the use of star-shaped genealogies in inference of coalescence times, *Genetics* **164** (4), 1677–1682.
- Sano, A. and Tachida, H. 2005. Gene genealogy and properties of test statistics of neutrality under population growth, *Genetics* **169** (3), 1687–1697.

- Schierup, M. H. and Hein, J. 2000. Consequences of recombination on traditional phylogenetic analysis, *Genetics* **156** (2), 879–891.
- Slatkin, M. 1996. Gene genealogies within mutant allelic classes, *Genetics* **143** (1), 579–587.
- Slatkin, M. and Hudson, R. R. 1991. Pairwise comparisons of mitochondrial DNA sequences in stable and exponentially growing populations, *Genetics* **129** (2), 555–562.
- Uyenoyama, M. K. 1997. Genealogical structure among alleles regulating self-incompatibility in natural populations of flowering plants, *Genetics* **147** (3), 1389–1400.
- Wakeley, J. 2009. “Coalescent Theory”, Roberts & Company, Greenwood Village, CO.
- Wickens, T. D. 2014. “The Geometry of Multivariate Statistics”, Psychology Press, New York.

$(X_n, Y_n)$	$\text{Cov}[X_n, Y_n]$	$\lim_{n \rightarrow \infty} \text{Cov}[X_n, Y_n]$	Reference
$H_n, T_k$	$\frac{4}{k^2(k-1)^2}$	$\frac{4}{k^2(k-1)^2}$	3.1, eq. 23
$H_n, L_n$	$4S_{2,n-1} - 4 + \frac{4}{n}$	$\frac{2\pi^2}{3} - 4 \approx 2.57974$	3.3, eqs. 29, 30
$H_n, E_n$	$\frac{4}{n}$	0	3.4, eqs. 37, 38
$H_n, I_n$	$4S_{2,n-1} - 4$	$\frac{2\pi^2}{3} - 4 \approx 2.57974$	3.7, eqs. 49, 50
$H_n, B_n$	$\frac{4[S_{3,n-1}n^2 - 3S_{2,n-1}n^2 + (n-1)(4n+1)]}{n^2}$	$4\zeta(3) + 16 - 2\pi^2 \approx 1.06902$	3.13, eqs. 71, 72
$L_n, T_k$	$\frac{4}{k(k-1)^2}$	$\frac{4}{k(k-1)^2}$	3.2, eq. 26
$L_n, E_n$	$\frac{4S_{1,n-1}}{n-1}$	0	3.6, eqs. 45, 46
$L_n, I_n$	$4S_{2,n-1} - \frac{4S_{1,n-1}}{n-1}$	$\frac{2\pi^2}{3} \approx 6.57974$	3.9, eqs. 57, 58
$L_n, B_n$	$\frac{4[S_{3,n-1}n - S_{2,n-1}n + n-1]}{n}$	$4\zeta(3) + 4 - \frac{2\pi^2}{3} \approx 2.22849$	3.14, eqs. 75, 76
$E_n, T_k$	$\frac{4}{k(k-1)(n-1)}$	0	3.5, eqs. 41, 42
$E_n, I_n$	$\frac{4S_{1,n-1}}{n-1} - \frac{8S_{1,n-1}n}{(n-1)(n-2)} + \frac{16}{n-2}$	0	3.10, eqs. 61, 62
$E_n, B_n$	$\frac{4(S_{2,n-1}n - n + 1)}{n(n-1)}$	0	3.15, eqs. 84, 85
$I_n, T_k$	$\frac{4(n-k)}{k(k-1)^2(n-1)}$	$\frac{4}{k(k-1)^2}$	3.8, eqs. 53, 54
$I_n, B_n$	$\frac{4(S_{3,n-1}n - S_{2,n-1}n + n - S_{3,n-1} - 1)}{n-1}$	$4\zeta(3) + 4 - \frac{2\pi^2}{3} \approx 2.22849$	3.16, eqs. 88, 89
$B_n, T_k$	$\frac{4}{k^2(k-1)^3}$	$\frac{4}{k^2(k-1)^3}$	3.12, eq. 68

Table 1: Covariances of pairs of variables. Expressions involving  $E_n$  or  $I_n$  apply for  $n \geq 3$  and expressions involving  $B_n$  apply for  $n \geq 4$ .



$(X_n, Y_n)$	$\text{Corr}[X_n, Y_n]$	$\lim_{n \rightarrow \infty} \text{Corr}[X_n, Y_n]$	Reference
$H_n, T_k$	$\frac{1}{\sqrt{2n^2 S_{2,n} - 3n^2 + 2n - 1}} \frac{1}{k(k-1)}$	$\frac{\sqrt{3}}{\sqrt{\pi^2 - 9}} \frac{1}{k(k-1)} \approx \frac{1.85738}{k(k-1)}$	3.1, eqs. 24, 25
$H_n, L_n$	$\frac{S_{2,n-1} n - n + 1}{\sqrt{S_{2,n-1} (2S_{2,n} n^2 - 3n^2 + 2n - 1)}}$	$\frac{\pi^2 - 6}{\pi \sqrt{2(\pi^2 - 9)}} \approx 0.93399$	3.3, eqs. 31, 32
$H_n, E_n$	$\frac{\sqrt{(n-1)(n-2)}}{\sqrt{2(2S_{2,n} n^2 - 3n^2 + 2n - 1)(S_{1,n-1} n - 2n + 2)}}$	0	3.4, eqs. 39, 40
$H_n, I_n$	$\frac{(S_{2,n-1} - 1)n \sqrt{(n-1)(n-2)}}{(2S_{2,n} n^2 - 3n^2 + 2n - 1)[4S_{1,n-1} + S_{2,n-1}(n-1)(n-2) - 4(n-1)]}$	$\frac{\pi^2 - 6}{\pi \sqrt{2(\pi^2 - 9)}} \approx 0.93399$	3.7, eqs. 51, 52
$H_n, B_n$	$\frac{\sqrt{2}[S_{3,n-1} n^2 - 3S_{2,n-1} n^2 + (n-1)(4n+1)]}{\sqrt{(2S_{2,n} n^2 - 3n^2 + 2n - 1)(3S_{2,n-1} n^2 - 2S_{2,n-1}^2 n^2 + n^2 - 4S_{2,n-1} n + 3n - 4)}}$	$\frac{3\sqrt{3}[2\zeta(3) + 8 - \pi^2]}{\sqrt{(\pi^2 - 9)(18 + 9\pi^2 - \pi^4)}} \approx 0.97054$	3.13, eqs. 73, 74
$L_n, T_k$	$\frac{1}{\sqrt{S_{2,n-1}}} \frac{1}{k-1}$	$\frac{\sqrt{6}}{\pi} \frac{1}{k-1} \approx \frac{0.77970}{k-1}$	3.2, eqs. 27, 28
$L_n, E_n$	$\frac{S_{1,n-1} \sqrt{n-2}}{\sqrt{2S_{2,n-1}(n-1)(S_{1,n-1} n - 2n + 2)}}$	0	3.6, eqs. 47, 48
$L_n, I_n$	$\frac{[S_{2,n-1}(n-1) - S_{1,n-1}] \sqrt{n-2}}{(n-1) \sqrt{S_{2,n-1} [4S_{1,n-1} + S_{2,n-1}(n-1)(n-2) - 4(n-1)]}}$	1	3.9, eqs. 59, 60
$L_n, B_n$	$\frac{\sqrt{2}(S_{3,n-1} n - S_{2,n-1} n + n - 1)}{\sqrt{S_{2,n-1} (3S_{2,n-1} n^2 - 2S_{2,n-1}^2 n^2 + n^2 - 4S_{2,n-1} n + 3n - 4)}}$	$\frac{\sqrt{6}[6\zeta(3) + 6 - \pi^2]}{\pi \sqrt{18 + 9\pi^2 - \pi^4}} \approx 0.84930$	3.14, eqs. 77, 78
$E_n, T_k$	$\frac{\sqrt{n-2}}{\sqrt{2(n-1)(S_{1,n-1} n - 2n + 2)}}$	0	3.5, eqs. 43, 44
$E_n, I_n$	$\frac{4(n-1) - S_{1,n-1}(n+2)}{\sqrt{2(S_{1,n-1} n - 2n + 2)[4S_{1,n-1} + S_{2,n-1}(n-1)(n-2) - 4(n-1)]}}$	0	3.10, eqs. 63, 64
$E_n, B_n$	$\frac{(S_{2,n-1} n - n + 1) \sqrt{n-2}}{\sqrt{(n-1)(S_{1,n-1} n - 2n + 2)(3S_{2,n-1} n^2 - 2S_{2,n-1}^2 n^2 + n^2 - 4S_{2,n-1} n + 3n - 4)}}$	0	3.15, eqs. 86, 87
$I_n, T_k$	$\frac{(n-k) \sqrt{n-2}}{(k-1) \sqrt{(n-1)[4S_{1,n-1} + S_{2,n-1}(n-1)(n-2) - 4(n-1)]}}$	$\frac{\sqrt{6}}{\pi} \frac{1}{k-1} \approx \frac{0.77970}{k-1}$	3.8, eqs. 55, 56
$I_n, B_n$	$\frac{\sqrt{2}(S_{3,n-1} n - S_{2,n-1} n + n - S_{3,n-1} - 1)n \sqrt{n-2}}{\sqrt{(n-1)[4S_{1,n-1} + S_{2,n-1}(n-1)(n-2) - 4(n-1)](3S_{2,n-1} n^2 - 2S_{2,n-1}^2 n^2 + n^2 - 4S_{2,n-1} n + 3n - 4)}}$	$\frac{\sqrt{6}[6\zeta(3) + 6 - \pi^2]}{\pi \sqrt{18 + 9\pi^2 - \pi^4}} \approx 0.84930$	3.16, eqs. 90, 91
$B_n, T_k$	$\frac{\sqrt{2n}}{\sqrt{3S_{2,n-1} n^2 - 2S_{2,n-1}^2 n^2 + n^2 - 4S_{2,n-1} n + 3n - 4}} \frac{1}{k(k-1)^2}$	$\frac{6}{\sqrt{18 + 9\pi^2 - \pi^4}} \frac{1}{k(k-1)^2} \approx \frac{1.95518}{k(k-1)^2}$	3.12, eqs. 69, 70

Table 2: Correlation coefficients of pairs of variables. Expressions involving  $E_n$  or  $I_n$  apply for  $n \geq 3$  and expressions involving  $B_n$  apply for  $n \geq 4$ .




Figure 1: Tree features. The tree height is  $H_n$  and the sum of the lengths of all tree branches is  $L_n$ . External branches, with total length  $E_n$ , appear in green; internal branches, with total length  $I_n$ , appear in red; basal branches, with mean length  $B_n$ , appear in purple.




Figure 2: Simulated and theoretical values of covariances of pairs of variables, plotted as functions of sample size  $n$ . Expressions for theoretical values are taken from Table 1.




Figure 3: Theoretical values of covariances  $\text{Cov}[X, T_k]$  for variables  $X$  in  $\{H_n, L_n, E_n, I_n, B_n\}$ , plotted as functions of  $k$  for  $n = 10$ ,  $n = 20$ , and  $n = 50$ . The plots appear on a logarithmic scale.




Figure 4: Simulated and theoretical values of correlation coefficients of pairs of variables, plotted as functions of sample size  $n$ . Expressions for theoretical values are taken from Table 2.




Figure 5: Theoretical values of correlation coefficients  $\text{Corr}[X, T_k]$  for variables  $X$  in  $\{H_n, L_n, E_n, I_n, B_n\}$ , plotted as functions of  $k$  for  $n = 10$ ,  $n = 20$ , and  $n = 50$ .