ON NON ALMOST-FIBERED KNOTS

MARIO EUDAVE-MUÑOZ, ARACELI GUZMÁN-TRISTÁN, ENRIQUE RAMÍREZ-LOSADA

ABSTRACT. An almost-fibered knot is a knot whose complement possesses a circular thin position in which there is one and only one weakly incompressible Seifert surface and one incompressible Seifert surface. Infinite examples of almost-fibered knots are known. In this article, we show the existence of infinitely many hyperbolic genus one knots that are not almost-fibered.

1. Introduction

Let K be a knot in S^3 . A regular circle-valued Morse function on the knot complement $f: S^3 \setminus K \to S^1$ induces a handle decomposition on the knot exterior. In [3], F. Manjarrez-Gutiérrez adapted the ideas of [8] to re-order the handles in such a way that the regular level surfaces are as simple as possible, giving rise to the definition of circular width and circular thin position of the knot exterior. Circular thin position of the knot exterior gives a sequence of Seifert surfaces which are alternately incompressible and weakly incompressible. In this context a fibered knot is a knot whose exterior has a circular thin position with one and only one incompressible level surface and none weakly incompressible level surface. This is the unique circular thin position for a fibered knot (see [1] or [11]). The circular width of a fibered knot is defined to be zero. An almost-fibered knot is a knot whose complement possesses a circular thin position in which there is one and only one weakly incompressible Seifert surface and one incompressible Seifert surface. Goda [2] showed that all prime non-fibered knots up to ten crossings are almost-fibered knots. On the other hand, F. Manjarrez-Gutiérrez, V. Núñez and the third author [5] showed that all non-fibered knots which are free genus one, are almost-fibered knots. It is worth mentioning that any non-fibered knot has a circular decomposition with only one thick level (see Subsection 2.2 for a definition), that is, any knot has a decomposition that looks like an almost-fibered knot, except that the thick surface may not be weakly incompresible, then the decomposition is not thin.

Until now, the only known examples of non almost-fibered knots are some connected sum of knots, for instance, the connected sum of asmall knots [4]. In 2017, at the Mathematical Congress of Americas, Hans Boden asked whether there are non almost-fibered, hyperbolic knots. In this paper we answer this question in the affirmative.

Theorem 1. There exists an infinite family of genus one hyperbolic knots which are not almost-fibered.

The paper is organized as follows. In Section 2 we review definitions about circular thin position of knots as well as definitions concerning to companion annuli in genus two handlebodies. In Section 3 we give an explicit construction of the desired family of knots, and in Section 4 we develop the proofs of the main results.

2. Preliminaries

2.1. **Knots and surfaces.** Let K be a knot in S^3 . The knot complement will be denoted by $C_K = S^3 \setminus K$. A closed regular neighborhood of K will be denoted by $\mathcal{N}(K)$ and the exterior of the knot K by $E(K) = S^3 \setminus \mathcal{N}(K)^\circ$.

A Seifert surface for a knot K is an oriented compact surface whose boundary is the knot K. It should be noted that such a surface always exists for any knot in S^3 .

If F is a two-sided surface, we say that F is weakly incompressible if any two compressing disks for F on opposite sides of F intersect along their boundary.

2.2. Circular thin position for knots. In this section, much of the notation and definitions follow from F. Manjarrez-Gutiérrez [3].

Let $K \subset S^3$ be a knot. Let $F: S^3 \backslash K \to S^1$ be a Morse function and define $f: E(K) \to S^1$ to be the restriction of F to E(K). A fundamental cobordism of f can be isotoped to have no local maxima or minima [7], we assume that all critical points of f have index 1 or 2.

We construct a handle decomposition of E(K) from f as follows: Choose a Seifert surface R at a regular level of f between an index-1 and an index-2 critical points. There are many such choices: We assume R is chosen to have smallest genus among all choices. If f has no critical points, then K is fibered and there is only one choice for R. Otherwise, we see that the critical points of f define collections $N = \{N_1, N_2, \ldots, N_k\}$ and $B = \{B_1, B_2, \ldots, B_k\}$ of 1- and 2-handles, respectively. We assume the handles in N_1 appear first after R and, moreover, that the handles in N_i appear before the handles in B_i and that the handles in B_i appear before the handles in N_{i+1} (taking all indices modulo k where necessary). Construct the compact manifold

$$H = (R \times I) \cup (N_1 \cup B_1) \cup \ldots \cup (N_k \cup B_k)$$

by flowing R along E(K) via the gradient of f, attaching handles from N and B as prescribed by the critical points of f. Choose a regular level S_i separating N_i from B_i ; that is,

$$S_i \cong \overline{\partial[(R \times I) \cup (N_1 \cup B_1) \cup \ldots \cup N_i] \setminus [\partial E(K) \cup (R \times \{0\})]}$$

Call S_i a thick level of f and set $S = \bigcup_{i=1}^k S_i$, similarly choose a regular level F_i separating B_i from N_{i+1} ; that is,

$$F_i = \overline{\partial[(R \times I) \cup (N_1 \cup B_1) \cup \ldots \cup (N_i \cup B_i)] \setminus [\partial E(K) \cup (R \times \{0\})]}$$

Call F_i a thin level of f and set $\mathcal{F} = \bigcup_{i=1}^k F_i$. We also define

$$W_i = (\text{collar of } F_i) \cup (N_i \cup B_i)$$

which is a 3-manifold with boundary $\partial W_i = F_i \cup F_{i+1} \cup (W_i \cap \partial E(K))$. The thick level of S_i defines a compact (but not closed) Heegaard surface for W_i , dividing it into compression bodies

$$A_i = (\text{collar of } F_i) \cup N_i \quad \text{and} \quad G_i = (\text{collar of } S_i) \cup B_i$$

The boundary ∂A_i can be seen as the union of three components; that is, $\partial A_i = S_i \cup F_i \cup \partial_v A_i$, where $\partial_v A_i = \partial A_i \cap \partial E(K)$. We call $\partial_v A_i$ the vertical boundary of A_i . We can similarly define $\partial_v G_i = \partial G_i \cap \partial E(K)$ to be the vertical boundary of G_i . Note that the vertical boundary is an annulus.

Observe that F_k is diffeomorphic to R. The function f defines a diffeomorphism $\phi: R \to R$. When K is fibered, ϕ is called the *monodromy* of K. In this case, we see that

$$E(K) = H/(R \times \{0\}) = \phi(R \times \{1\})$$

The collection $\mathcal{D} = \{(W_i; A_i, G_i)\}_{i=1}^k$ will be called the *circular handle decomposition* of E(K) induced by f.

For a closed, connected surface $S \neq S^2$ define its complexity $c(S) = 1 - \chi(S)$. If S has nonempty boundary, define $c(S) = 1 - \chi(\overline{S})$, where \overline{S} denotes S with its boundary components capped off with disks. We define $c(S^2) = 0$ and $c(D^2) = 0$. If f is disconnected, define $c(S) = \sum c(S_i)$, where $S = \bigcup_i S_i$ and each S_i is connected.




FIGURE 1.

For a knot $K \subset S^3$ with circular handle decomposition \mathcal{D} for E(K), the circular width $cw(\mathcal{D})$ of the decomposition \mathcal{D} is the multiset of integers $\{c(S_i)\}_{i=1}^k$, ordered in a non-increasing way, and $|cw(\mathcal{D})| = k$ is the number of thick levels in \mathcal{D} . The circular width cw(E(K)) of the knot exterior E(K) is defined to be the minimal circular width among all circular handle decompositions of E(K). The minimum is taken using the lexicographic ordering of multisets of integers.

The pair $(E(K), \mathcal{D})$ is in *circular thin position* if \mathcal{D} realizes the circular width of E(K). When K is a fibered knot, we define $cw(E(K)) = \emptyset$, so |cw(E(K))| = 0. If |cw(E(K))| = 1, we say that K is almost-fibered.

F. Manjarrez-Gutiérrez [3] examined knot exteriors and showed the following results:

Theorem 2 (F. Manjarrez-Gutiérrez). Let $K \subset S^3$ be a knot. At least one of the following holds:

- K is fibered;
- *K* is almost-fibered;
- K contains a closed essential surface in its complement. Moreover, this closed essential surface is in the complement of an incompressible Seifert surface of K;
- K has at least two non-isotopic, incompressible Seifert surfaces.

Theorem 3 (F. Manjarrez-Gutiérrez). If $(E(K), \mathcal{D})$ is in circular thin position, then

- Each Heegaard splitting S_i of W_i is strongly irreducible.
- Each F_i is incompressible in E(K).
- Each S_i is weakly incompressible surface in E(K).

The converse of this theorem is not true in general. That is, a circular handle decomposition satisfying the three properties above need not be thin.

2.3. Companion annuli in genus two handlebodies. In this section much of the definitions and notations follow from L. Valdez-Sánchez [10].

Let M a 3-manifold with boundary and $\gamma \subset \partial M$ a circle which is non-trivial in M. We say that a separating annulus A properly embedded in M is a companion annulus of γ if A is not parallel into ∂M and the circle components of ∂A cobound an annulus $A_{\gamma} \subset \partial M$ with core isotopic to γ in ∂M . If the region cobounded by A and A_{γ} in M is a solid torus V, we say that V is a companion solid torus of γ in M and denote the components of $M \setminus A$ by M_A and V.

Let H be a genus two handlebody and $\gamma, \gamma' \subset \partial H$ be mutually disjoint and non-parallel circles. We say that

- γ is a *primitive circle* in H if γ represents a primitive element in the free group $\pi_1(H)$; geometrically, this is equivalent to the presence of a disk in H which intersects γ minimally in one point.
- γ is a power circle in H if γ represents a non-trivial power in $\pi_1(H)$, that is, if γ represents a power $p \geq 2$ of some non-trivial element in $\pi_1(H)$.

In the part 2 of Lemma 3.3 of [10], Valdez-Sánchez prove the next

Lemma 1. Let $\gamma \subset \partial H$ a circle which is non-trivial in H. Then γ has a companion annulus in H iff γ is a power circle in H; more precisely,

- the companion annulus A of γ is unique up to isotopy and cobounds with ∂H a companion solid torus of γ , of whose core γ represents a non-trivial power in $\pi_1(H)$.
- $H \setminus A$ consists of a genus two handlebody H_A and a solid torus, and the core of A is a primitive circle in H_A .

3. Knots with four genus one Seifert surfaces

Suppose that $K \subset S^3$ is a genus one knot and $\mathbb{T} = T_1 \sqcup T_2 \sqcup T_3 \sqcup T_4$ is a collection of four mutually disjoint and non-parallel genus one Seifert surfaces for K, where the T_i 's are labeled consecutively around $\partial \mathcal{N}(K)$ following some fixed orientation of the meridian slope $\mu \subset \partial \mathcal{N}(K)$. For $i \in \{1, \ldots, 4\}$, denote by H_i the region in E(K) cobounded by T_i and T_{i+1} . For each $i = 1, \ldots, 4$, set $\partial H_i = T_i \cup J_i \cup T_{i+1} \pmod{4}$ with $J_i \subset \partial \mathcal{N}(K)$ an annulus whose core is parallel to K. See Figure 2.

Assume that H_i satisfy the following conditions:

- Each of the regions H_i is a genus two handlebody.
- There is a power circle $\omega_i \subset T_i$ in H_i with companion annulus $C_i \subset H_i$, such that the circles ∂C_i cobound an annulus $\overline{C}_i \subset T_i$,




Figure 2.

and such that the once-punctured torus obtained by isotoping $(T_i \setminus \overline{C}_i) \cup C_i$ slightly off T_i is parallel to T_{i+1} .

- ω_i represents a power $p \geq 3$ of some non-trivial element in $\pi_1(H)$.
- Let ω'_i be a parallel copy of the core of the annulus C_i in T_{i+1} . Then ω'_i and ω_{i+1} intersect transversely in one point.

Note that the curve ω'_i in T_{i+1} is a power circle in H_i , with companion annulus C'_i , such that $C_i \cap C'_i$ intersect in two simple closed curves.

Theorem 4. There is an infinite family of genus one knots $K(\ell, m, n)$ that satisfy all the above properties.

Proof. Let $K' \cup L$ be the link shown in Figure 3, where ℓ, m, n are integers. The box labelled by m stands for a sequence of m horizontal crossings. Assume that $|m|, |n|, |\ell| \geq 3$. Note that both K' and L are trivial knots.

Let D_1 , D_2 , D_3 and D_4 be the disks shown in Figure 4, that is, disks properly embedded in E(K'), their boundary lying in $\partial \mathcal{N}(K')$. Note that $\partial D_i = K'$, and that D_i intersects L transversely in 3 points.

Let $p: S^3 \to S^3$ be the double cover of S^3 branched along L. The lifting $K = K(\ell, m, n) = p^{-1}(K')$ is a knot in S^3 since the linking number between K' and L is an odd number. Let $T_i = p^{-1}(D_i)$, i = 1, 2, 3, 4. Then T_i is a genus one Seifert surface for K. Let H'_i be the region bounded by D_i and D_{i+1} . This is a 3-ball that intersects L in 3 arcs. Note that considered as a 3-tangle, it is a trivial tangle. Therefore, $H_i = p^{-1}(H'_i)$ is a genus two handlebody. Let E'_i be a disk in D_i that contains two points of intersection with L, and such that there is another disk E_i in H'_i , with $\partial E'_i = \partial E_i$, and $E'_i \cup E_i$ bounds a 3-ball B_i contained in H'_i which encloses the crossings labelled ℓ , m, or n. Let C'_i and C_i be the liftings of E'_i and E_i , respectively, these are annuli. The lifting of B_i is a solid torus R_i whose boundary is




FIGURE 3.




FIGURE 4.

the union of C'_i and C_i . Let ω_i be the core of C'_i , this is a curve that wraps around R_i at least thrice (in fact it wraps around R_i , ℓ , m or n times). Then ω_i is a power curve in H_i , and C_i is a companion annulus. Clearly, $(T_i \setminus C'_i) \cup C_i$ is parallel to T_{i+1} . The curve ω_i is the lift of a curve on E'_i parallel to its boundary. Note that a copy of this curve, and the corresponding curve in E_{i+1} intersect in two points. However,




FIGURE 5.

these curves lift to a pair of curves, and in fact, each pair of curves intersect in one point. \Box

Theorem 5. The exterior of the knots $K(\ell, m, n)$ admits a circular decomposition of width $\{3, 3\}$.

Proof. The link $K' \cup L$ shown in Figure 3 is isotopic to the link shown in Figure 5. Let D_2 , F_1 , D_4 and F_2 be the disks shown in Figure 5, these are disks properly embedded in E(K'), with their boundary lying in $\partial \mathcal{N}(K')$. Note that F_1 and F_2 intersect L transversely in 5 points. As before, $T_i = p^{-1}(D_i)$, i = 2, 4, is a genus one Seifert surface for $K = K(\ell, m, n)$, while $\Sigma_i = p^{-1}(F_i)$, i = 1, 2, is a genus two Seifert surface for K. By taking a collection of parallel disks between D_2 and F_1 , we note that there is collection of disks intersecting L in three points, then there is a single disk intersecting L in three points and in a local maximum point and then a collection of disks intersecting Lin five points. In the double branched cover, this implies that we pass from T_2 to Σ_1 by adding a 1-handle. Similarly, when passing from F_1 to D_4 , there is a disk that intersects L in a local minimum. In the double branched cover this implies that we pass from Σ_1 to T_4 by adding a 2-handle. Something similar can be done for D_4 , Σ_2 and D_2 . Then T_2 , Σ_1 , T_4 and Σ_2 define a circular decomposition for the exterior of K, which has width $\{3,3\}$.

4. Knots without a decomposition of width {3}

Suppose that $K \subset S^3$ is a genus one knot and $\mathbb{T} = T_1 \sqcup T_2 \sqcup T_3 \sqcup T_4$ is a collection of four mutually disjoint and non-parallel genus one Seifert surfaces for K. Denote by H_1 , H_2 , H_3 , and H_4 the complementary regions of the surfaces in the exterior of K. Suppose that all conditions for the surfaces and the regions stated just before Theorem 4 are satisfied.

Lemma 2. The knot K is hyperbolic.

Proof. The given conditions imply that for each T_i (i = 1, ..., 4) a power circle in H_i cannot have companion annuli on both sides of T_i . Then part 1 of Lemma 8.1 of [10] implies that H is hyperbolic.

Lemma 3. For each i = 1, ..., 4, there are up to isotopy five incompressible annuli properly embedded in H_i , whose boundaries lie in $T_i \sqcup T_{i+1}$, and which are not parallel onto T_i or T_{i+1} .

Proof. First we note that H_i contains five incompressible annuli properly embedded associated to the power circle in H_i . These annuli are:

- The companion annulus C_i for the power circle in H_i such that $\partial C_i \subset T_i$.
- The companion annulus C'_i for the power circle in H_i such that $\partial C'_i \subset T_{i+1}$.
- The spanning annulus A_i for the power circle in H_i with one boundary component in T_i and the other in T_{i+1} .
- A second spanning annulus A'_i for the power circle in H_i with one boundary component in T_i and the other in T_{i+1} .
- An annulus \widehat{A}_i boundary parallel into $\partial \mathcal{N}(K)$.

The spanning annuli A_i and A'_i are obtained by doing one of the two possible oriented double curve sums of C_i and C'_i . In the complementary region H_i we represent these five annuli as in Figure 6. That is, we represent by a blue circle the companion torus, by green arcs the companion annuli, by red arcs the spanning annuli, and by a purple arc the annulus parallel into $\partial \mathcal{N}(K)$.

Note that each one of these annuli is not boundary parallel into T_i or T_{i+1} , and that any ∂ -compression disk for these annuli must intersect the annulus J_i contained in $\partial \mathcal{N} K$.

Now suppose that B is a properly embedded and incompressible annulus in H_i , whose boundary lies in $T_i \sqcup T_{i+1}$. If there is a ∂ -compression disk for B disjoint from J_i , then B is boundary parallel, for otherwise T_i or T_{i+1} would be compressible. Isotope B to intersect




Figure 6.

 $C_i \sqcup C_i' \sqcup A_i \sqcup A_i' \sqcup \widehat{A_i}$ minimally. Trivial closed intersection curves on B or $C_i \sqcup C_i' \sqcup A_i \sqcup A_i' \sqcup \widehat{A_i}$ can be eliminated since the surfaces are incompressible and H_i is irreducible. Suppose first that ∂B is not disjoint from the boundary curves of all the annuli $C_i \sqcup C_i' \sqcup A_i \sqcup A_i' \sqcup \widehat{A_i}$. Then there must be intersection arcs between them. If there is an arc of intersection which is trivial in one of the annuli, then either it can be removed by an isotopy or it would imply that one of the annuli is ∂ -compressible, which is not possible. If ∂B has both components on T_i , then there must be trivial intersection arcs on A_i , and if ∂B has one component in T_i and the other in T_{i+1} , then there is a trivial intersection arc in B.

Then suppose that ∂B is disjoint from the boundary curves of all the annuli $C_i \sqcup C'_i \sqcup A_i \sqcup A'_i \sqcup \widehat{A_i}$. If there are closed essential intersection curves on B or $C_i \sqcup C'_i \sqcup A_i \sqcup A'_i \sqcup \widehat{A_i}$, then the boundary curves of B would be parallel to the boundary curves of one of the annuli $C_i, C'_i, A_i, A'_i, A_i$. By taking an outermost curve of intersection lying in B, an isotopy can be done to reduce the number of curves of intersection. Thus we have that $B \cap (C_i \sqcup C'_i \sqcup A_i \sqcup A'_i \sqcup \widehat{A_i}) = \emptyset$. Suppose B is not parallel into ∂H_i . By symmetry we may assume only two cases: $\partial B \subset T_i$ (or T_{i+1}) and $\partial_1 B \subset T_i, \, \partial_2 B \subset T_{i+1}$. If $\partial B \subset T_i$, then ∂B cobound an annulus on T_i and B is separating on H_i . Then B is a companion annulus, but there is up to isotopy only one companion annulus in H_i (Lemma 1), so B is parallel to C_i . Similarly, if $\partial B \subset T_{i+1}$, then B is parallel to C'_i . Now, suppose $\partial_1 B \subset T_i$, $\partial_2 B \subset T_{i+1}$. Since $B \cap (C_i \sqcup C'_i \sqcup A_i \sqcup A'_i \sqcup \widehat{A_i}) = \emptyset$, $\partial_1 B \subset T_i$ must be parallel to either ∂C_i or $\partial_1 A_i$ and $\partial_2 B \subset T_{i+1}$ must be parallel to either $\partial C'_i$ or $\partial_2 A_i$. Then B is parallel to one of the spanning annuli A_i, A'_i , or $\widehat{A_i}$.

Lemma 4. If A is an incompressible annulus properly embedded in $H_i \cup H_{i+1}$, whose boundary lies on $T_i \cup T_{i+2}$, then either it is parallel to an annulus in $\partial \mathcal{N}(K)$, or it can be isotoped to lie in H_i or H_{i+1} .

Proof. This follows from Lemma 3 and from the fact that there are no companion annuli on both sides of T_{i+1} with the same boundary slope (Lemma 5.1 of [10]).

Lemma 5. For each i = 1, ..., 4, H_i cannot contain a properly embedded pair of pants P with all its boundaries parallel to K, that is, such that each component of ∂P lies on T_i (or T_{i+1}) and it is parallel to ∂T_i (or ∂T_{i+1}), or it is an essential curve on $J_i \subset \partial \mathcal{N}(K)$.

Proof. Suppose H_i contains a properly embedded pair of pants P, with boundaries all parallel to K. Up to isotopy, we can assume that ∂P is contained in J_i . Then there is planar surface with three boundary components properly embedded in E(K), which is not possible by [6], unless P is compressible. But if this happens, by compressing P we get that K is the trivial knot, which is not possible.

Lemma 6. Suppose that P is an incompressible pair of pants properly embedded in H_i , whose boundary is contained in $T_i \sqcup T_{i+1}$. Suppose that two of its boundary components are essential, non-boundary parallel, and non power curves on T_{i+1} . Suppose that the third boundary component lies on T_i . Then this last curve is parallel on T_i to ∂T_i .

Proof. Consider the intersection between P and the companion C'_i . Note that P and C'_i must intersect, and the intersection consists of arcs, for simple closed curves of intersection can be removed by the incompressibility of the surfaces. Suppose that γ is an arc of intersection in C'_i , with endpoints lying in the same boundary component of $\partial C'_i$, and that γ is outermost. So, γ cuts off a disk D in C'_i which is a ∂ -compression disk for P. By performing the boundary compression, we get an annulus, with one boundary component in T_{i+1} which is parallel to ∂T_{i+1} . Then by Lemma 5, the other boundary component of the annulus, that is, the boundary component of P lying in T_i , must be parallel to ∂T_i .

The remaining possibility is that all the arcs of intersection in C'_i are essential, that is, spanning arcs in C'_i . There are at least two arcs of intersection, for there are at least four points of intersection between ∂P and $\partial C'_i$. Take two consecutive arcs of intersection in C'_i , note that when considered in P, there are two possibilities for the arcs. A possible case is that each arc has endpoints on the same boundary component, but a different component from the other arc. In this case one of the arcs is trivial, and then there is a ∂ -compression disk for

 C'_i , which is not possible. The other case is when both arcs join two different boundary components. Then there is a disk D in P formed by these two arcs plus one arc in each of the boundary components. Let $\overline{C'_i}$ be the annulus in T_{i+1} bounded by $\partial C'_i$. One possibility is that D is a compression disk for the torus $(T_{i+1} \setminus \overline{C'_i}) \cup C'_i$, which is not possible. The remaining possibility is that D is a meridian disk of the companion solid torus of H_i , but this implies that the power curve in H_i is a power of order two, which by hypothesis is not possible.

Lemma 7. The exterior of K, E(K), cannot contain another genus one Seifert surface for K non isotopic to any T_i ; i = 1, ..., 4.

Proof. Suppose that E(K) contains a fifth genus one Seifert surface for K, S, non isotopic to any T_i (i = 1, ..., 4). Isotope S in order to make ∂S and ∂T_i disjoint for every i = 1, ..., 4.

If $S \cap \mathbb{T} = \emptyset$, then $S \subset H_i$ for some $i \in \{1, ..., 4\}$. By Lemma 6.8 of [10], S is parallel to T_i or T_{i+1} .

Now, if $S \cap \mathbb{T} \neq \emptyset$, by the incompressibility of S and T_i (i = 1, ..., 4) we can isotope S to intersect transversely \mathbb{T} in a minimal set of essential closed curves (in both S and T_i ; i = 1, ..., 4). \mathbb{T} divide S into a collection of annuli, pair of pants, and once punctured tori. Suppose $\partial S \subset J_i \subset H_i$ for some $i \in \{1, ..., 4\}$.

If there is an annulus A in S with one boundary ∂S and the other, $\partial_2 A$ in T_{i+1} , then $\partial_2 A$ is parallel to K and we can isotope S to eliminate such intersection. Then the curves of intersection in S, are not parallel to ∂S . If there is a curve of intersection between T_i and S, which is parallel to ∂T_i in T_i , then by doing an annulus compression to S, we get a pair of pants with its boundaries contained in $\partial \mathcal{N}(K)$, which is not possible, by the same argument as in Lemma 5. We can assume that \mathbb{T} divide S into a collection of annuli and a pair of pants. Note that in this collection, we can not have adjacent annuli, by Lemma 4.

In this way, \mathbb{T} divide S into an annulus A and a pair of pants P. Since the annulus A is parallel to the companion annulus, say in H_i , its boundary curves are the power circle in H_i and assuming that the pair of pants P is contained in H_{i-1} , we have that in H_{i-1} , P has two boundary curves in T_i which are not power circles H_{i-1} and one boundary curve parallel to K. It follows that P is parallel to T_i minus an annulus, then S can be isotoped to be contained in H_i and to be parallel to T_{i+1} , a contradiction.

Lemma 8. Suppose that P is an incompressible pair of pants properly embedded in H_i , whose boundary is contained in T_i . Suppose that two of its boundary components are essential, non-boundary parallel on T_i ,

and that the third boundary component is parallel on T_i to ∂T_i . Then either P is parallel into T_i , or two boundary components of ∂P are isotopic to the power curve of H_i , and P union an annulus in T_i is parallel into T_{i+1} .

Proof. If two boundary components of P are not the power curve of H_i , then P must intersect the annulus A_i . By looking at an outermost arc of intersection in A_i we get that P is ∂ -compressible. By ∂ -compressing P we get an annulus parallel to T_i , and hence P must be parallel into T_i . If two boundary components of P are isotopic to the power curve of H_i . These curves bound an annulus C on T_i , such that $P \cup C$ is a once-puncured tori, which by Lemma 7 is either parallel to T_i , or to T_{i+1} .

Theorem 6. E(K) cannot admit a circular decomposition of width $\{3\}$.

Proof. We are assuming that K is a knot whose exterior contains 4 mutually disjoint and non-parallel genus one Seifert surfaces T_i ; $i = 1, \ldots, 4$, and that $\mathbb{T} = \bigsqcup_{i=1}^4 T_i$. Suppose E(K) has a circular decomposition of width $\{3\}$. So,

$$E(K) = (T_1 \times I) \cup N \cup B/T_1 \times 0 \sim T_1 \times 1$$

where N and B stand for a one handle and a two handle, respectively. By Lemma 7, T_i $(i=1,\ldots,4)$ are the only genus one Seifert surfaces for K.

Let $\Sigma := \partial((T_1 \times I) \cup N)$. Then Σ is a genus two Seifert surface for K. By Theorem 3, Σ is weakly incompressible.

We note that $\Sigma \cap T_i \neq \emptyset$ for $i \in \{2, 3, 4\}$. To show this, suppose that $\Sigma \cap T_4 = \emptyset$. Let D the core disk of B. By the incompressibility of T_4 , we can isotope D to be disjoint of T_4 . By compressing Σ along D we obtain a genus one surface which does not intersect the region H_4 and then cannot be isotopic to T_1 , a contradiction.

Without lost of generality, suppose that $\partial \Sigma \subset H_1$ or $\partial \Sigma \subset H_2$. By general position, $\Sigma \cap T_i$ for $i \in \{2,3,4\}$ consists of simple closed curves. Moreover, as Σ is weakly incompressible, $\Sigma \cap T_i$ consists of simple closed curves which are essential on both Σ and T_i , by Lemma 2.6 of [9]. We assume that Σ has been isotoped such that the number of connected components of $\Sigma \setminus \mathbb{T}$, denoted $c(\Sigma)$, is minimal. \mathbb{T} divides Σ into a collection of annuli, pair of pants, four punctured spheres, once punctured tori, and twice punctured tori. Let $\Sigma_i = \Sigma \cap H_i$ for $i \in \{1, 2, 3, 4\}$.

If there is an annulus in Σ with one boundary in $\partial \mathcal{N}(K)$ and the other in T_2 or T_3 , isotope Σ to eliminate such intersection. Also, by

Lemma 4, there cannot be adjacent annuli on Σ with boundary slope a power circle in H_i .

Lemma 9. No component of Σ_4 is a 4-punctured sphere.

Proof. Suppose that a component of Σ_4 is a 4-punctured sphere S_4 . Then $\partial S_4 \subset T_4$. As Σ is a genus 2 surface, the complementary regions of Σ_4 in Σ are an annulus A and a pair of pants P. Note that A must be contained in H_3 , and it is a companion annulus there, otherwise we can push it into H_4 , reducing $c(\Sigma)$. P is contained in the union of H_3 , H_2 and H_1 . ∂P consists of $\partial \Sigma$ and two curves α_1 , α_2 . No curve of intersection of P with T_3 and T_2 is parallel to $\partial \Sigma$. So $P \cap (T_3 \cup T_2)$ consists of curves parallel to α_1 or α_2 . This implies that $P \cap H_3$ consists of annuli, and also $P \cap H_2$ consists of annuli. As we have two adjacent annuli these must be parallel to a neighborhood of the knot. That is, α_1 is a curve parallel to K. If one of these annuli have its boundary components in the same T_i , an isotopy removes the intersection. If all annuli are spanning, then in H_1 there is a pair of pants, whose all boundary components are parallel to K, which contradicts Lemma 5.

Lemma 10. If Σ_4 contains a once punctured torus T or a pair of pants, then part of it can be isotoped through T_4 , changing it into an annulus, without changing $c(\Sigma)$.

Proof. If Σ_4 contains a once punctured torus T, then it has its boundary on T_4 , which is parallel to ∂T_4 , and then it is parallel to T_1 . Part of T can be pushed into H_3 , except for an annulus, which remains in H_4 . This is done without changing $c(\Sigma)$.

If Σ_4 contains a pair of pants P, then its boundary lies on T_4 . If only one or the three boundary components of P were essential curves on T_4 , then there would be a simple closed curve in T_4 intersecting P and then Σ an odd number of times, which is not possible for any such closed curve has linking number 0 with K. Then two boundary components of P are essential curves on T_4 and the third one is parallel to ∂T_4 . Then by Lemma 8, either P can be pushed towards H_3 , or P union an annulus in T_4 is parallel into T_1 . Then, as in the previous case, part of the pair of pants can be pushed into H_3 , except for an annulus, which remains in H_4 .

We can assume that Σ_4 contains an annulus A_4 , which is then a companion annulus in H_4 .

Lemma 11. The annulus A_4 is a non-separating annulus in Σ .




Figure 7.

Proof. Suppose that A_4 is a separating annulus in Σ . Then it separates Σ into a once-punctured torus T' and a twice punctured torus. If T' is contained in H_3 , then by looking at the intersections between T' and the annulus C_3 , we see that there is a boundary compression disk for T' contained in C_3 , but this contradicts the so called parity rule. The other possibility is that $T' \cap H_3$ is a pair of pants and $T' \cap H_2$ is an annulus. But this annulus has to be a companion annulus in H_2 , and then the pair of pants contradicts Lemma 6.

We use the symbols A, P, S, T' to denote an annulus, a pair of pants, a four punctured sphere and a twice punctured torus, respectively. By a symbol Q_i we mean a surface of type Q, so that $Q_i = \Sigma \cap H_i$. Then by a sequence $A_4.P_3.S_2.A_1$ we mean that Σ intersects the handlebodies H_i in an annulus, a pair of pants, a four punctured sphere, and an annulus, respectively. If we have a sequence $Q_4.Q_3.Q_2.Q_1$, we also mean that $Q_i \cap T_i = Q_{i-1} \cap T_i$.

Lemma 12. There are the following possibilities for the intersections of Σ with the H_i .

- $(1) A_4.P_3.P_2.P_1$
- (2) $A_4.P_3.A_2.P_1.P_2$
- (3) $A_4.S_3.\{A_2,A_2'\}.P_1$
- $(4) A_4.P_3.A_2.S_1.A_2'$
- (5) $A_4.P_3.S_2.A_1$
- (6) $A_4.P_3.A_2.T_1'$




Figure 8.

Proof. Note that $\chi(\Sigma) = \sum_{i=1}^{i=4} \chi(\Sigma_i) = -3$, and then the pieces of Σ consist of 3 pair of pants, or a 4-punctured sphere and a pair of pants, or a twice punctured torus and a pair of pants, plus several annuli in each case. It is shown by inspection that the cases listed above are all possible cases. A schematic figure of all possible cases is shown in Figure 7.

We have that Σ_4 is a non-separating annulus, which is a companion annulus in H_4 , the region adjacent to it and contained in H_3 , can not contain an annulus, by Lemma 4. So, Σ_3 is either a pair of pants P_3 or a 4-punctured sphere S_3 . Suppose first that $\Sigma_3 = S_3$. What is left of Σ is a pair of pants; note that Σ_1 can not contain an annulus containing $\partial \Sigma$, and then the only possibility is that Σ_2 consists of two annuli A_2 and A'_2 , and Σ_1 consists of a pair of pants P_1 . So we have Case 3.

Suppose now that $\Sigma_3 = P_3$. What is left of Σ is a twice punctured torus. Note that by Lemma 6, the boundary of P_3 lying in T_3 is parallel to ∂T_3 . If Σ_2 contains a pair of pants P_2 , then Σ_1 must be a pair of pants P_1 , so we have Case 1. If Σ_2 contains a 4-punctured sphere S_2 , then it must contain $\partial \Sigma$, and Σ_1 is an annulus A_1 . So, we have Case 5. If Σ_2 contains an annulus A_2 adjacent to P_3 , what is left of Σ is a twice punctured torus, and there are three possibilities. First, Σ_1 is a twice punctured torus T'_1 , this is Case 6. Second, Σ_1 consists of a pair of pants P_1 and Σ_2 contains a pair of pants P_2 , this is Case 2. And third, Σ_1 consists of a 4-punctured sphere S_1 and Σ_2 contains one more annulus A'_2 , this is Case 4.

To finish the proof of Theorem 6, we analyze each one of the cases and show that it can not arise.

Case 1: $A_4.P_3.P_2.P_1$.

In this case A_4 is a companion annulus in H_4 , two of the boundary curves of P_3 are essential in T_4 , and the other lies in T_3 , and is boundary parallel, by Lemma 6. One boundary curve of P_2 lies in T_3 , and is boundary parallel there, the other two are essential curves in T_2 . Finally, two of the boundary curves of P_1 are essential in T_2 and parallel to the boundary curves of the companion annulus in H_1 (for otherwise P_1 could be isotoped to lie in H_2) and the last one lies in I_3 . This is case is schematically shown in Figure 8. In this and the subsequent figures, the blue circles represent the companion tori, the green arcs the companion annuli, and the surface Σ is represented by a red curve.

As Σ is bicompressible, let D be a compressing disk lying in the external side, that is, the side intersecting $\partial \mathcal{N}(K)$ in a single annulus lying in H_1 . Consider $D \cap \mathbb{T}$. Trivial closed intersection curves on D can be eliminated by an isotopy. Then we can assume $D \cap \mathbb{T}$ consists only of arcs.

Take an innermost intersection arc t on D, then there exists an arc $\sigma \subset \Sigma$ such that $\sigma \cup t$ cobound a disk $E \subset D$ and $int E \cap \mathbb{T} = \emptyset$.

• If $t \subset T_4$, we have two options: $\sigma \subset H_4$ or $\sigma \subset H_3$.

Suppose $\sigma \subset H_4$. In H_4 , the annulus A_4 is parallel to the companion annulus of H_4 and its boundary components lie in T_4 . If σ joins one boundary component of A_4 to itself, we can eliminate such intersection by an isotopy. Otherwise, σ joins the two boundary components of A_4 , giving rise to a boundary compression of this annulus, a contradiction.

Suppose now $\sigma \subset H_3$. In H_3 , the pair of pants P_3 is parallel to $T_4 \backslash A$ for an annulus $A \subset T_4$. Here E is a compression disk for $P_3 \cup A$, which is a surface parallel to T_4 , a contradiction.

• If $t \subset T_3$, we have two options: $\sigma \subset H_3$ or $\sigma \subset H_2$.

Suppose $\sigma \subset H_3$. As in the precedent case, we can think of P_3 in H_3 as $T_4 \backslash A$ for an annulus $A \subset T_4$. Consider the region in P_3 parallel to $\partial E(K)$ between ∂T_3 and ∂T_4 which contains an annulus A'. In A', two subarcs of σ cobound a disk F such that $E \cup F$ is an annulus. It means that σ and t can be completed to coannular closed curves $\hat{\sigma}$ and \hat{t} , $\hat{\sigma} \subset T_4$ and $\hat{t} \subset T_3$. If the curve $\hat{\sigma}$ is parallel to the core of the companion annulus in H_4 so is \hat{t} , but this is not possible. Also, $\hat{\sigma}$ can not be parallel to the core of the companion annulus in H_3 . Then $\hat{\sigma}$ and \hat{t} are parallel to K. This implies that we can eliminate the innermost arc in D by an isotopy.




Figure 9.

Now if $\sigma \subset H_2$, we follow the same argument as before, since P_2 is isotopic to $T_2 \setminus A$ where a core of the annulus A has the slope of the companion annulus in H_1 .

• If $t \subset T_2$, we have two options: $\sigma \subset H_2$ or $\sigma \subset H_1$. If $\sigma \subset H_2$ we proceed as in the first case when $t \subset T_4$ and $\sigma \subset H_3$. Suppose now $\sigma \subset H_1$. P_1 is a pair of pants parallel to $T_1 \setminus A$, where A is an annulus parallel to an annulus in T_2 having t as an essential arc. Then E is a compression disk for $P_1 \cup A$ which is parallel to T_1 , a contradiction.

Case 2: $A_4.P_3.A_2.P_1.P_2$.

This case is schematically shown in Figure 9. In this case Σ_2 consists of an spanning annulus A_2 parallel to an annulus in $\partial \mathcal{N}(K)$ and a pair of pants P_2 , whose boundary consists of two curves on T_2 and $\partial \Sigma$. Then Σ would have a self-intersection, which is impossible.

Case 3: $A_4.S_3.\{A_2, A_2'\}.P_1.$

This case is schematically shown in Figure 10. In this case Σ_2 consists of the annuli A_2 and A'_2 , which are spanning annuli in H_2 , whose boundary curves are power circles in H_2 . It follows that P_1 is a pair of pants whose boundaries are two curves in T_2 , which are primitive circles in H_1 and one curve parallel to K, then it is possible to push P_1 into H_2 to eliminate $\Sigma \cap H_1$, a contradiction.

Case 4: $A_4.P_3.A_2.S_1.A'_2$.




FIGURE 10.




FIGURE 11. Case 4

This case is schematically shown in Figure 11. In this case Σ_2 two consists of two annuli, A_2 and A'_2 , where A_2 is a boundary parallel annulus and A'_2 is parallel to the companion annulus in H_2 with boundary on T_2 , and then S_1 is a 4-punctured sphere.

As in Case 1, let D be a compressing disk for Σ lying in the external side, such that $D \cap \mathbb{T}$ consists only of arcs. As before, take an innermost intersection arc t on D, and an arc $\sigma \subset \Sigma$ such that $\sigma \cup t$ cobound a disk $E \subset D$ and $intE \cap \mathbb{T} = \emptyset$. Note that the cases where $t \subset T_4$ or $t \subset T_3$ and $\sigma \subset H_3$ can be eliminated as in case 1.

Now suppose that $t \subset T_3$ and $\sigma \subset H_2$. Remember that Σ_2 is a disjoint union of $\Sigma_2 = A_2 \cup A_2$, where A_2 is parallel to the annulus J_2 and A_2 is parallel to the companion annulus in H_2 , with its boundary lying in T_2 . Then $\sigma \subset A_2$, but this is implies that σ is not essential, a contradiction.

Now suppose that $t \subset T_2$ and $\sigma \subset H_2$. Then $\sigma \subset A'_2$. Let A be an annulus in T_2 with $\partial A = \partial A'_2$. Then $t \cup \sigma$ lies in $(T_2 \setminus A) \cup A'_2$, which implies that E is a compression disk for a surface parallel to T_3 , which is not possible.

So we have that $t \subset T_2$ and $\sigma \subset H_1$. The four-punctured sphere S_1 has two boundary slopes $\partial_1 S_1$, $\partial_2 S_1$ which are primitive curves in H_1 and two boundary slopes parallel to K, one lying in T_2 ($\partial_3 S_1$) and the other in $\partial \mathcal{N}(K)$ ($\partial_4 S_1$). By making a ∂ -compression in S_1 with the disk E we obtain a pair of pants or an annulus and a pair of pants. We have the following cases:

- If t joins $\partial_1 S_1$ and $\partial_2 S_1$. By making a ∂ -compression with E, we obtain a pair of pants whose boundaries are all parallel to K, which is not possible, by Lemma 5.
- If t joins $\partial_1 S_1$ (or $\partial_2 S_1$) with $\partial_3 S_1$. By making a ∂ -compression with E, we obtain a pair of pants with boundary components $\partial_2 S_1$, $\partial_4 S_1$ and one more curve parallel to $\partial_2 S_1$. As $\partial_2 S_1$ is a primitive curve in H_1 , then the pair of pants is parallel into T_2 . It follows that it is possible to isotope S_1 to lie in H_2 , a contradiction.
- If t joins $\partial_1 S_1$ or $\partial_2 S_1$ with itself. By making a ∂ -compression with E, we obtain an annulus and a pair of pants which are parallel into T_2 and which are possible to isotope to lie in H_2 , a contradiction.
- If t joins $\partial_3 S_1$ with itself. By making a ∂ -compression with E, we obtain an annulus and a pair of pants which are parallel into T_2 and which are possible to isotope to lie in H_2 , a contradiction.

Case 5: $A_4.P_3.S_2.A_1$.

This case is schematically shown in Figure 12. Led D, E, t and σ be as in Case 1. Note that the cases where $t \subset T_4$ or $t \subset T_3$ and $\sigma \subset H_3$ can be eliminated as in Case 1.

The four-punctured sphere S_2 has two boundary curves $\partial_1 S_2$, $\partial_2 S_2$, which lie in T_2 and are primitive curves in H_2 and two boundary curves parallel to K, one lying in T_3 ($\partial_3 S_2$) and the other in $\partial \mathcal{N}(K)$ ($\partial_4 S_2$).

Now we assume $t \subset T_3$ and $\sigma \subset H_2$. Then t is an arc joining $\partial_3 S$ with itself. By making a ∂ -compression in S_2 with E, we obtain an annulus and a pair of pants. The boundary of the annulus consists of an essential curve in T_3 and either $\partial_1 S_2$, $\partial_2 S_2$ or $\partial_4 S_2$. In any case, this is in contradiction with Lemma 3.




Figure 12.

Now suppose, $t \subset T_2$ and $\sigma \subset H_2$. If t joins $\partial_1 S_2$ and $\partial_2 S_2$, then by doing a ∂ -compression on S_2 with E, we obtain a pair of pants whose all boundaries are parallel to K, a contradiction to Lemma 5. Now, if t joins $\partial_1 S_2$ (or $\partial_2 S_2$) to itself, by doing a ∂ -compression to S_2 we obtain an annulus A and a pair of pants P. The curve $\partial_1 S_2$ is split into two curves, say ∂_{11} which is parallel to ∂T_2 , and ∂_{12} which is parallel to $\partial_2 S_2$. If the annulus A has ∂_{12} as a boundary component, then the other boundary component can not be $\partial_3 S_2$ or $\partial_4 S_2$, and then it has to be $\partial_2 S_2$. But in this case the three boundary components of P are parallel to K, which is not possible. Then the annulus A has ∂_{11} as a boundary component. The other boundary component can not be $\partial_2 S_2$. It can not be $\partial_3 S_2$, for in this case A and P will intersect. Then the other boundary component of A is $\partial_4 S_2$. In this case A can be isotoped into A, we get a configuration as in Case 1.

Finally, suppose that $t \subset T_2$ and $\sigma \subset H_1$. We note that A_1 is parallel to the companion annulus in H_1 . Let A be an annulus in T_2 with the same boundary as A_1 . Then E is a compression disk for the surface $(T_2 - A) \cup A_1$, which is parallel to T_1 , which is not possible.

Case 6: $A_4.P_3.A_2.T_1'$.

This case is schematically shown in Figure 13. Note that one boundary component of T_1' lies in J_1 , which is precisely $\partial \Sigma$, and the other lies in T_2 and it is parallel to ∂T_2 . Led D, E, t and σ be as in Case 1. Note that the cases where $t \subset T_4$ or $t \subset T_3$ can be eliminated as in Case 1. The only remaining case is when $t \subset T_2$ and $\sigma \subset H_1$. In this




Figure 13.

case the arc t joins points in the same boundary component of T_1' . By pushing T_1' along the disk E, we get a new configuration for Σ , $\Sigma \cap H_3$ and $\Sigma \cap H_4$ are both a pair of pants, that is, we have a configuration as in Case 1, which is already solved.

Theorem 7. There exists an infinite family of genus one hyperbolic knots which are not almost-fibered.

Proof. Let $K(\ell, m, n)$ be the family of knots constructed in Theorem 4. These knots have genus one and by Theorem 5 their exterior admits a circular decomposition of width $\{3,3\}$. Also, by Lemma 2 the knots $K(\ell, m, n)$ are hyperbolic. If they are almost-fibered, then their exterior would admit a circular decomposition of width $\{3\}$ which is impossible by Theorem 6.

Acknowledgements. The first author is partially supported by grant PAPIIT-UNAM IN116720. The second author's research is supported by a CONACYT postdoctoral fellowship. The third author made this research on a sabbatical year at IMATE-UNAM, Unidad Juriquilla.

References

- [1] G. Burde, H. Zieschang, Neuwirth Knoten und Flachenabbildungen, Abh. Math. Sem. Univ. Hamburg 31 (1967), 239–246.
- [2] H. Goda, On handle number of Seifert surfaces in S³, Osaka J. Math. 30 (1993), 63–80.

- [3] F. Manjarrez-Gutiérrez, Circular thin position for knots in S³, Algebr. Geom. Topol. **9** (2009), 429–454.
- [4] F. Manjarrez-Gutiérrez, Additivity of handle number and Morse-Novikov number of a-small knots. Topology Appl. 160 (2013), no. 1, 117–125.
- [5] F. Manjarrez-Gutiérrez, V. Núñez, E. Ramírez-Losada, Circular handle decompositions of free genus one knots, Pac. J. Math. 275 (2015), 361–407.
- [6] D. Gabai, Foliations and the topology of 3-manifolds, J. Differ. Geom. 18 (1983), 445–503.
- [7] J. Milnor, Lectures on the h-cobordism theorem, Princeton University Press. Princeton, N.J., 1963.
- [8] M. Scharlemann, A. Thompson, *Thin position for 3-manifolds*, Contemp. Math., **164**, AMS, (1994), 231–238.
- [9] M. Scharlemann, M. Tomova, Alternate Heegaard genus bounds distance, Geom. Topol. 10 (2006), 593–617.
- [10] L. Valdez-Sánchez, Seifert surfaces for genus one hyperbolic knots in the 3-sphere, Algebr. Geom. Topol. 19 (2019), 2151–2231.
- [11] W. Whitten, *Isotopy types of knot spanning surfaces*, Topology **12** (1973), 373–380.

Instituto de Matemáticas, Universidad Nacional Autónoma de México, Campus Juriquilla, Querétaro, México

Email address: mario@matem.unam.mx

CENTRO DE INVESTIGACIÓN EN MATEMÁTICAS, A.C. 402, 36000 GUANAJUATO, GTO., MÉXICO

Email address: araceli.guzman@cimat.mx

CENTRO DE INVESTIGACIÓN EN MATEMÁTICAS, A.C. 402, 36000 GUANAJUATO, GTO., MÉXICO

Email address: kikis@cimat.mx