On the completeness of generalized eigenfunctions of the Hamiltonian of reggeon field theory in Bargmann space

Abdelkader Intissar^(1,2)

(1) Equipe d'Analyse spectrale, UMR-CNRS n:6134, Université de Corse, Quartier Grossetti, 20 250 Corté-France.

E.mail: abdelkader.intissar@orange.fr

(2) Le Prador, 129, rue du commandant Rolland, 13008 Marseille-France.

Abstract

The aim of the present paper is to review and study some new spectral properties of the operator $H_{\mu,\lambda} = \mu A^*A + i\lambda A^*(A+A^*)A$ which characterizes the Reggon field theory, where μ , λ are real parameters and $i^2 = -1$. $A \longrightarrow \frac{d}{dz}$ and $A^* \longrightarrow z = x + iy; (x,y) \in \mathbb{R}^2$ respectively are boson annihilation and creation operators, satisfying the commutation relation $[A,A^*] = I$ in Bargmann space $\mathbb{B} = \{\varphi : \mathbb{C} \longrightarrow \mathbb{C} \ entire \ ; \int_{\mathbb{C}} |\varphi(z)|^2 \ e^{-|z|^2} dx dy < \infty \}$ with standard scalar product $\langle \varphi, \psi \rangle = \frac{1}{\pi} \int_{\mathbb{C}} \varphi(z) \overline{\psi(z)} e^{-|z|^2} dx dy$

The Hamiltonian takes the form $H_{\mu,\lambda} = \mu z \frac{d}{dz} + i\lambda(z \frac{d^2}{dz^2} + z^2 \frac{d}{dz}) := \mu \mathcal{N} + i\lambda H_I$, where $\mathcal{N} = z \frac{d}{dz}$ and $H_I = z \frac{d^2}{dz^2} + z^2 \frac{d}{dz}$.

The Hamiltonian $H_{\mu,\lambda}$ is non-Hermitian with respect to the above standard scalar product. Hence the question arises, whether the eigenfunctions by the finite norm condition form a complete basis.

The main new results of this article are the determination of the boundary conditions for the eigenvalue problem associated to $H_{\mu,\lambda}$ and the proof of the completeness of the basis mentioned in the question above by transforming $H_{\mu,\lambda}$ in the Hermitian form.

This article also provides an explicit integral form of the inverse of $H_{\mu,\lambda}$ on the negative imaginary axis z = -iy; y > 0 and some spectral properties of $H_{\mu,\lambda}^{-1}$; $\mu > 0$.

 $\textbf{Keywords:} \ Spectral \ theory: Gribov \ operators: semigroups: Non-self-adjoint \ operators: Bargmann \ space: Reggeon \ field \ theory.$

§ 1. Introduction

Le \mathbb{B} be the Bargmann space [4] defined by :

$$\mathbb{B} = \{ \varphi : \mathbb{C} \longrightarrow \mathbb{C} \ entire \ ; \int_{\mathbb{C}} |\varphi(z)|^2 \ e^{-|z|^2} dx dy < \infty \}$$
 (1.1)

with standard scalar product

$$<\varphi,\psi> = \frac{1}{\pi} \int_{\mathbb{C}} \varphi(z) \overline{\psi(z)} e^{-|z|^2} dx dy$$
 (1.2)

and with usual basis

$$e_n(z) = \frac{z^n}{\sqrt{n!}}; \ n \in \mathbb{N}$$
 (1.3)

and

 \mathbb{P} is the set of polynomials which is dense in \mathbb{B} .

The constant $\frac{1}{\pi}$ is chosen in scalar product such that the norm of the constant function $\varphi(z) = 1$ is one

Let
$$\mathbb{B}_s = \{(a_n)_{n=0}^{\infty} \in \mathbb{C}; \sum_{n=0}^{\infty} n! \mid a_n \mid^2 < \infty \}$$
 (1.4)

 \mathbb{B}_s is also a Hilbert space. The scalar product on \mathbb{B}_s is defined by

$$\langle (a_n), (b_n) \rangle_s = \sum_{n=0}^{\infty} n! a_n \overline{b_n}$$
(1.5)

and the associated norm is denoted by $|| . ||_s$.

It is well known that \mathbb{B} is related to \mathbb{B}_s by unitary transform of \mathbb{B} onto \mathbb{B}_s , given by the following transform :

$$I: \mathbb{B} \longrightarrow \mathbb{B}_s, \varphi(z) = \sum_{n=0}^{\infty} a_n z^n \longrightarrow I(\varphi) = \left(\frac{1}{n!} \varphi^{(n)}(0)\right)_{n=0}^{\infty} = (a_n)_{n=0}^{\infty}$$
(1.6)

In Bargmann's representation, we have :

the annihilation operator A is :

$$A = \frac{d}{dz} \text{ with domain } D(A) = \{ \varphi \in \mathbb{B}; \frac{d}{dz} \varphi \in \mathbb{B} \}$$
 (1.7)

the creation operator A^* is :

the multiplication by z with domain
$$D(A^*) = \{ \varphi \in \mathbb{B}; z\varphi \in \mathbb{B} \}$$
 (1.8)

It is well known that A^* is adjoint of A (with respect above standard scalar product), D(A) is dense in \mathbb{B} , $D(A) = D(A^*)$ and the injection $D(A) \longrightarrow \mathbb{B}$ is a compact mapping.(see [13] or lemma 0.2 in [15] for an elementary proof).

and the Hamiltonian of reggeon field theory in zero transverse dimensions takes the following form :

$$(\bullet_1) H_{\mu,\lambda} = \mu z \frac{d}{dz} + i\lambda \left(z \frac{d^2}{dz^2} + z^2 \frac{d}{dz}\right) := \mu \mathcal{N} + i\lambda H_I; \ (\mu,\lambda) \in \mathbb{R}^2 \text{ and } i^2 = -1$$

$$(1.9)$$

with maximal domain $D(H_{\mu,\lambda}) = \{ \varphi \in \mathbb{B}; H_{\mu,\lambda} \varphi \in \mathbb{B} \}$.

where

$$\mathcal{N} = z \frac{d}{dz}$$
 with domain $D(\mathcal{N}) = \{ \varphi \in \mathbb{B}; \mathcal{N}\varphi \in \mathbb{B} \}$ (1.10)

and

$$H_I = z \frac{d^2}{dz^2} + z^2 \frac{d}{dz} \text{ with maximal domain } D(H_I) = \{ \varphi \in \mathbb{B}; H_I \varphi \in \mathbb{B} \}.$$
 (1.11)

(•2) If we call generalized Laguerre-type derivatives the operators of the form $\mathcal{D}(r,s) = A^r \mathcal{N}^s$, with r = 1, 2, ..., s = 0, 1, ... then as \mathcal{N}^s conserves the number of bosons, we observe that these operators act as monomials in boson operators which annihilate r bosons and that the operator $H_{\mu,\lambda}$ can be written in the form

$$H_{\mu,\lambda} = \mu A^* A + i\lambda (\mathcal{D}(1,1) + \mathcal{D}^*(1,1) - (A + A^*)) \tag{1.12}$$

(•3) In 1987, we have given in [13] many spectral properties of $H_{\mu,\lambda}$ for $\mu > 0$ in particular the minimal domain of $H_{\mu,\lambda}$ coincides with its maximal domain, the positiveness of its eigenvalues, the existence of the smallest eigenvalue $\sigma_0 \neq 0$ and an asymptotic expansion of its semigroup $e^{-tH_{\lambda,\mu}}$ as $t \longrightarrow +\infty$.

Where the minimal domain of $H_{\mu,\lambda}$ is:

$$D(H_{\mu,\lambda}^{min}) = \{ \varphi \in \mathbb{B}, \exists p_n \in \mathbb{P}, p_n \longrightarrow \varphi, \exists \psi \in \mathbb{B}; H_{\mu,\lambda}p_n \longrightarrow \psi \} . \tag{1.13}$$

- (•4) In 1984, Ando and Zerner have schown in [3] that the smallest eigenvalue σ_0 of $H_{\mu,\lambda}$ can be analytically continued in μ on entire real axis.
- (•5) In 1998, we have given in [14] the boundary conditions at infinity for a description of all maximal dissipative extensions in Bargmann space of the minimal Heun's operator $H_I = z \frac{d^2}{dz^2} + z^2 \frac{d}{dz}$. The characteristic functions of the dissipative extensions have computed and some completeness theorems have obtained for the system of generalized eigenvectors. it is well known that the restriction H_I^{min} of the closure of H_I on the polynomials set \mathbb{P} is symmetric.

But the minimal domain $D(H_I^{min}) = \{ \varphi \in \mathbb{B}, \exists \, p_n \in \mathbb{P}, p_n \longrightarrow \varphi, \exists \, \psi \in \mathbb{B}; H_I p_n \longrightarrow \psi \} \text{ of } H_I \text{ is different of its maximal domain } D(H_I) = \{ \varphi \in \mathbb{B}; H_I \varphi \in \mathbb{B} \}.$

- (\bullet_6) It is also well known that H_I is chaotic operator in Devaney's sense see [8] or [16], in particular its spectrum is $\sigma(H_I) = \mathbb{C}$.
- (\bullet_7) The Hamiltonian $H_{\mu,\lambda}$ is non-Hermitian with respect to above standard scalar product and the domain of the adjoint and anti-adjoint parts are not included in each other, nor is the domain of their commutator.
- (•8) Hence the question arises, whether the eigenfunctions by the finite norm condition form the complete basis. The aim new results of this article is the determination of boundary conditions of eigenvalue problem associated to $H_{\mu,\lambda}$ and to give a answer to above question by transform $H_{\mu,\lambda}$ to the Hermitian form. Also, in this article we give an explicit integral form of the inverse of $H_{\mu,\lambda}$ on the negative imaginary axis z = -iy, y > 0 and some spectral properties of $H_{\mu,\lambda}$; $\mu > 0$ on a weighted space.

Now we give an outline of the content of this paper :

The Section 2 deals the action of some elementary operators on Bargmann space usefull to study of eigenvalue problem associated to the Hamiltonian $H_{\mu,\lambda}$.

In section 3, we give an asymptotic analysis of the solutions of bi-confluent Heun equation associated to eigenvalue problem of the Hamiltonian $H_{\mu,\lambda}$.

In section 4, we give on the negative imaginary axis z = -iy, y > 0 a transformation procedure of $H_{\mu,\lambda}$ to a symmetric operator with compact resolvent and we show the main result of this paper that the eigenfunctions of $H_{\mu,\lambda}$ form a complete basis.

And

In section 5, we give an explicit integral form of the inverse of $H_{\mu,\lambda}$ on the negative imaginary axis z = -iy; y > 0 and some spectral properties of $H_{\mu,\lambda}^{-1}$; $\mu > 0$ as integral operator on a weighted Hilbert space.

\S 2 Action of some elementary operaors on Bargmann space usefull to study of eigenvalue problem associated to the Hamiltonian $H_{\mu,\lambda}$

• We introduce the operator P (changing the sign of z) defined by

$$PzP^{-1} = -z$$
 and $P\frac{d}{dz}P^{-1} = -\frac{d}{dz}$ then we deduce that
$$PHP^{-1} = \mu z \frac{d}{dz} - i\lambda(z\frac{d^2}{dz^2} + z^2\frac{d}{dz}) := H_{\mu,-\lambda} \text{ (the formal adjoint of } H_{\mu,\lambda})$$
(2.1)

The operator P is unitary and Hermitian in Bargmann space, since

$$< P\varphi, P\psi > = < \varphi, \psi >$$
 and $< \varphi, P\psi > = < P\varphi, \psi >$

Now, let the following eigenvalue problem :

$$H_{\mu,\lambda}\varphi_n = \sigma_n\varphi_n \text{ with real } \sigma_n$$
 (2.2)

and

$$\varphi_n^*(z) = -P\varphi_n(z) = -\varphi_n(-z) \tag{2.3}$$

then we get:

$$\langle \varphi_m^*, H_{\mu,\lambda} \varphi_n \rangle = \sigma_n \langle \varphi_n^*, \varphi_n \rangle$$
 (2.4)

and

$$H_{\mu,-\lambda}\varphi_m^* = -H_{\mu,-\lambda}P\varphi_m = -PH_{\mu,\lambda}\varphi_m = \sigma_m\varphi_m^*$$
(2.5)

and

$$\langle H_{\mu,-\lambda}\varphi_m^*;\varphi_n\rangle = \sigma_m \langle \varphi_m^*;\varphi_n\rangle$$
 (2.6)

it follows that

$$(\sigma_n - \sigma_m) < \varphi_m^*, \varphi_n > = < \varphi_m^*, H_{\mu,\lambda}\varphi_n > - < H_{\mu,-\lambda}\varphi_m^*, \varphi_n > - < \varphi_m^*, \varphi_$$

$$= \langle \varphi_m^*, H_{\mu,\lambda} \varphi_n \rangle - \langle \varphi_m^*, H_{\mu,\lambda} \varphi_n \rangle = 0$$

so that

$$\langle \varphi_m^*, \varphi_n \rangle = 0 \text{ when } \sigma_n \neq \sigma_m$$
 (2.7)

 Determination of an orthogonal scalar product in Bargmann space associated to the operator P.

We introduce the operator ν (the Hermitian operator of the sign) on the eigenfunctions $\{\varphi_n\}$, n = 1, 2, ...; of $H_{\mu,\lambda}$ defined by

$$\nu(\varphi_n) = \nu_n \varphi_n$$

where

$$\left\{ \begin{array}{l} \nu_n = +1, \quad if < \varphi_n, P\varphi_n >> 0 \\ \\ \nu_n = -1, \quad if < \varphi_n, P\varphi_n >< 0 \end{array} \right.$$

We introduce the positively defined scalar product $<<\varphi,\psi>>=<\varphi,\nu P\psi>$ with respect to which the eigenstates of $H_{\mu,\lambda}$ are orthogonal.

\S 3 Asymptotic analysis of the solutions of bi-confluent Heun equation associated to eigenvalue problem of the Hamiltonian $H_{\mu,\lambda}$

We recall that $H_{\mu,\lambda}$ has the form $p(z)\frac{\partial^2}{\partial z^2} + q(z)\frac{\partial}{\partial z}$ with $p(z) = i\lambda z$ and $q(z) = i\lambda z^2 + \mu z$ of degree 2 and $H_{\mu,\lambda}$ is of Heun operator type.

The eigenvalue problem associated to $H_{\mu,\lambda}$ does not satisfy the classical ordinary differential equations of the form :

$$p(z)\frac{\partial^2 \varphi}{\partial z^2} + q(z))\frac{\partial \varphi}{\partial z} = \sigma \varphi \tag{3.1}$$

where p(z) is a polynomial of degree at most two, q(z) is a polynomial of the degree exactly one and σ is a constant.

• If $\lambda \neq 0$, the eigenvalue problem associated to $H_{\mu,\lambda}$ can be written as follows:

$$z\frac{\partial^2 \varphi}{\partial z^2} + (z^2 - i\rho z)\frac{\partial \varphi}{\partial z} = -i\frac{\sigma}{\lambda}\varphi; \quad \rho = \frac{\mu}{\lambda}$$
(3.2)

Let $z=i\sqrt{2}\xi$ and $\varphi(\sqrt{2}\xi)=\psi(\xi)$ then (3.2) can be transformed to :

$$\xi \frac{\partial^2 \psi}{\partial \xi^2} + (-2\xi^2 + \rho\sqrt{2}\xi) \frac{\partial \psi}{\partial \xi} - \frac{\sigma\sqrt{2}}{\lambda}\psi = 0$$
(3.3)

the above equation belongs to bi-confluent Heun equations, denoted by $BHE(\alpha, \beta, \gamma, \delta)$ which are in the form :

$$\xi \frac{d^2 u}{d\xi^2} + (1 + \alpha - \beta \xi - 2\xi^2) \frac{du}{d\xi} + \{ (\gamma - \alpha - 2)\xi - \frac{1}{2} (\delta + (\alpha + 1)\beta) \} u = 0 \; ; \; (\alpha, \beta, \gamma, \delta) \in \mathbb{C}^4 \; (3.4)$$

It is well known that

- the singular points of the bi-confluent Heun equation (3.4) are 0 and ∞ .
- The singularity is regular at 0 and irregular at ∞ .
- At 0, if $(\alpha, \beta, \gamma, \delta) \in (\mathbb{C} \mathbb{Z}) \times \mathbb{C}^3$ then a basis of $BHE(\alpha, \beta, \gamma, \delta)$ is given by

$$\{N(\alpha, \beta, \gamma, \delta, \xi), \xi^{-\alpha}N(-\alpha, \beta, \gamma, \delta, \xi)\}\$$
where $N(\alpha, \beta, \gamma, \delta, \xi) = \Gamma(\alpha)\sum_{n=0}^{\infty} \frac{A_n(\alpha, \beta, \gamma, \delta)}{\Gamma(\alpha + 1 + n)n!}\xi^n,$

the A_n are polynomials in $\alpha, \beta, \gamma, \delta$ defined by the following relation

$$A_{n+2} = [\beta(n+1) + \frac{1}{2}(\delta + \beta(1+\alpha))]A_{n+1} - (\gamma - 2 - \alpha - 2n)(n+1)(n+1+\alpha)A_n \text{ for } n \ge 0$$

and $A_{-1} = 0, A_0 = 1$, (see proposition 5 of ref. [22] or ref. [5]) (\star_1)

- If $\alpha \in \mathbb{Z}$ there are logarithmic terms in the bases of solutions of $BHE(\alpha, \beta, \gamma, \delta)$ at 0 which is a regular singularity (except for some values see [20]). (\star_2)
- If $(\alpha, \beta, \gamma, \delta) \in \mathbb{C}^4$ then $BHE(\alpha, \beta, \gamma, \delta)$ admits as basis of formal solutions at ∞ ,

$$\{\xi^{-\frac{1}{2}(\gamma-\alpha-2)} \sum_{n=0}^{\infty} a_n(\alpha,\beta,\gamma,\delta) \xi^{-n}, e^{-\frac{1}{2}(\gamma+\alpha+2} e^{\xi^2+\beta\xi} \sum_{n=0}^{\infty} a_n(\alpha,i\beta,-\gamma,-i\delta) (i\xi)^{-n} \}.$$

where the complex numbers an are defined by

$$2(n+2)a_{n+2} = \left[\frac{1}{2}(\delta + \beta(\gamma - 1) - \beta(n+1)\right]a_{n+1} - (\frac{\gamma - \alpha - 2}{2} - n)(\frac{\gamma + \alpha - 2}{2} - n)a_n \text{ for } n \ge 0 \text{ and } a_{-1} = 0, a_0 = 1. \text{ (see proposition 8 of ref [22])}$$

These equations are more complicated than the equations for classical special functions such as Bessel functions, hypergeometric functions, and confluent hypergeometric functions (see [7] and [1]).

Remark 3.1

(i) If we take
$$\alpha = -1, \beta = -\rho\sqrt{2}, \gamma = 1$$
 and $\delta = \frac{2\sigma\sqrt{2}}{\lambda}$ in (3.4) then we get (3.3), so it is $BHE(-1, -\frac{\mu\sqrt{2}}{\lambda}, 1, -\frac{2\sigma\sqrt{2}}{\lambda})$.

- (ii) As $\alpha = -1 \in \mathbb{Z}$ then the proposition 5 of [22] (see \star_1) is not applicable.
- (iii) We will show there are logarithmic terms in the bases of solutions of $BHE(-1, -\frac{\mu\sqrt{2}}{\lambda}, 1, \frac{2\sigma\sqrt{2}}{\lambda})$ at 0 which is a regular singularity by using the classical Frobenius-Fuch's method.

In (3.2) denoting $\frac{d}{dz}\varphi(z)=\varphi^{'}(z)$ and $\frac{d^{2}}{dz^{2}}\varphi(z)=\varphi^{"}(z)$ where a prime throughout denotes a derivative with respect to z, then the eigenvalue problem associated to $H_{\mu,\lambda}$; $\lambda\neq 0$ can be written as follows

$$\varphi''(z) + p(z)\varphi'(z) + q(z)\varphi(z) = 0; \quad p(z) = (z - i\rho) \text{ and } q(z) = \frac{i\sigma}{\lambda z}$$
 (3.5)

Eq. (3.5) is bi-confluent equation designated in the Ince classification [11] as $[0, 1, 1_4]$ which has, as in the case of the confluent hypergeometric equation, two singularities located, respectively, at the origin and infinity: one regular, and the second irregular with a singularity rank higher by unity than that for the confluent hypergeometric equation.

We recall that z_0 is regular singular point of (3.5) if

• z_0 is singular;

•
$$(z-z_0)p(z)$$
 and $(z-z_0)^2q(z)$ are analytic at z_0 . (3.6)

As $\lim zp(z) = \lim z^2q(z) = 0$ as $z \longrightarrow 0$ then $z_0 = 0$ is regular singular point of (3.5).

The usual power series method, that is setting $\varphi(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n$, breaks down if z_0 is a singular point. Here "breaks down" means "cannot find all solutions".

The solution of the our bi-confluent Heun equation in a power series of z can be easily constructed by applying the Method of Frobenius to (3.5) whose the basic idea is to look for solutions

of the form
$$\varphi(z) = z^s \sum_{n=0}^{+\infty} a_n z^n$$
:

Substitute the expansion $\varphi(z) = z^s \sum_{n=0}^{+\infty} a_n z^n$ into the equation (3.5), we have

$$(z^{s} \sum_{n=0}^{+\infty} a_{n} z^{n})^{"} + (z - i\rho)(z^{s} \sum_{n=0}^{+\infty} a_{n} z^{n})' + \frac{i\sigma}{\lambda z} z^{s} \sum_{n=0}^{+\infty} a_{n} z^{n+s} = 0$$
 (3.7)

Carrying out the differentiation, we reach

$$\sum_{n=0}^{+\infty} (n+s)(n+s-1)a_n z^{n+s-2} + \sum_{n=0}^{+\infty} (n+s)a_n z^{n+s} + i(\frac{\sigma}{\lambda} - \rho) \sum_{n=0}^{+\infty} (n+s)a_n z^{n+s-1} = 0$$
(3.8)

Shifting index:

$$\bullet \sum_{n=0}^{+\infty} (n+s)a_n z^{n+s-1} = \sum_{n=1}^{+\infty} (n+s-1)a_{n-1} z^{n+s-2}$$
(3.9)

$$\bullet \sum_{n=0}^{+\infty} (n+s)a_n z^{n+s} = \sum_{n=1}^{+\infty} (n+s-3)a_{n-2} z^{n+s-2}; a_{-1} = 0$$
(3.10)

Now if we put $\alpha = i(\frac{\sigma}{\lambda} - \rho)$ then the equation becomes

$$s(s-1)a_0z^{s-2} + \sum_{n=1}^{+\infty} [(n+s)(n+s-1)a_n + \alpha(n+s-1)a_{n-1} + (n+s-3)a_{n-2}]z^{n+s-2} = 0$$
(3.11)

and the indicial equation is

$$s(s-1) = 0. (3.12)$$

by using the theorem of Fuchs:

Theorem 3.2 (theorem of Fuchs)

Consider the equation $\varphi''(z) + p(z)\varphi' + q(z)\varphi(z) = 0$

Let z_0 be a regular singular point. That is $(z-z_0)p(z)=\sum_{n=0}^{+\infty}p_n(z-z_0)^n$ and

$$(z-z_0)^2 q(z) = \sum_{n=0}^{+\infty} q_n (z-z_0)^n$$
 with certain radii of convergence.

Let r be no bigger than the radius of convergence of either $(z - z_0)p(z)$ or $(z - z_0)^2q(z)$ and s_1, s_2 solve the indicial equation

$$s(s-1) + p_0 s + q_0 = 0. (3.13)$$

Then

1. If $s_1 \neq s_2$ and $s_1 - s_2$ is not an integer, then the two linearly independent solutions are given by

$$\varphi_1(z) = (z - z_0)^{s_1} \sum_{n=0}^{+\infty} a_n (z - z_0)^n, \, \varphi_2(z) = (z - z_0)^{s_1} \sum_{n=0}^{+\infty} b_n (z - z_0)^n$$
(3.14)

2. If $s_1 = s_2$ then $\varphi_1(z)$ is given by the same formula as above, and $\varphi_2(z)$ is of the form

$$\varphi_2(z) = \varphi_1(z)\ln(z - z_0) + (z - z_0)^{s_1} \sum_{n=0}^{+\infty} d_n(z - z_0)^n$$
(3.15)

3. If $s_1 - s_2$ is an integer, then take s_1 to be the larger root (More precisely, when s_1 , s_2 are both complex, take s_1 to be the one with larger real part, that is $\Re e s_1 > \Re e s_2$). Then $\varphi_1(z)$ is still the same, while

$$\varphi_2(z) = c\varphi_1(z)ln(z - z_0) + (z - z_0)^{s_2} \sum_{n=0}^{+\infty} e_n(z - z_0)^n$$
(3.16)

(Note that c may be 0).

All the solutions constructed above converge at least for $0 < |z - z_0| < r$.

The proof of this theorem is through careful estimate of the size of an using the recurrence relation, see [2]:

In fact the converse of this theorem is also true. That is if all solutions of the equation satisfies $Lim(z-z_0)^s\varphi(z)=0$ as $z\longrightarrow z_0$ for some s, then $(z-z_0)p(z)$ and $(z-z_0)^2q(z)$ are analytic at z_0 . This is called FuchsTheorem. Its proof is a tour deforce of complex analysis and can be found in [23] see also, the following references about this topic [10] or [21]

We deduce that (3.5) has tow linearly independent solutions given by

$$\begin{cases}
\varphi_1(z) = z\eta(z); & \eta \quad is \quad analytic, \eta'(0) \neq 0 \\
\varphi_2(z) = czln(z)\xi(z) + \sum_{n=0}^{+\infty} c_n z^n; & \xi(z) \quad is \quad analytic, \xi'(0) \neq 0
\end{cases}$$
(3.17)

As $\varphi_2(z)$ contains non-analytic terms then it is not acceptable in Bargmann space and thus it is rejected then

$$\varphi(z) \sim z \quad (z \longrightarrow 0) \tag{3.18}$$

Now, we turn to the point $z=\infty$, which is the second possible singular point for the solution of (3.5)). To analyse the point $z=\infty$, we can first perform the change of independent variable from z to ξ , $z=\frac{1}{\xi}$ and $\varphi(z)=\varphi(\frac{1}{\xi})=\phi(\xi)$ and study the behaviour of the transformed equation at $\xi=0$ to get :

$$\begin{cases} \frac{d}{dz} = -\xi^2 \frac{d}{d\xi} \, and \, \frac{d^2}{dz^2} = \xi^4 \frac{d^2}{d\xi^2} + 2\xi^3 \frac{d}{d\xi} \\ \phi''(\xi) + p(\xi)\phi'(\xi) + q(\xi)\phi(\xi) = 0 \\ where \\ p(\xi) = -\frac{2\xi^2 - i\rho\xi + 1}{\xi^3} \, and \, q(\xi) = \frac{i\sigma}{\lambda\xi^3} \end{cases}$$
(3.19)

• The origin is regular singular point of (3.19) if $\xi p(\xi)$ and $\xi^2 q(\xi)$ are analytic functions in the neighborhood of the origin. As these latter conditions are not fulfilled, then the origin is an irregular singular point and it is of order 3.

Consequently, the theorem of Fauchs is not applicable to (3.19) and this equation cannot be approximated by an Euler equation in the neighborhood of the origin.

If we like to explore some of these techniques, a good starting point would be Chapter 3 of [6] or the book of Kristensson [19].

The case of an irregular singular point is much more difficult to address, although some techniques do exist for obtaining the solution to a homogeneous second-order linear differential equation in the vicinity of an irregular singular point.

The applications of the concepts on irregular singular points in the current literature to our second order linear differential equation does not inform us about the solutions belonging to the domains of our operators, in particular to the Bargmann space and what are the boundary

conditions verified by them at infinity?

For example, if we look for a solution of (3.19) in the form $\phi(\xi) = \sum_{k=0}^{+\infty} a_k \xi^{k+s}$ then we get the following recurrence relation:

$$k(k-2)a_k + i[\rho(k+1) + \frac{\sigma}{\lambda}]a_{k+1} - 2(k+2)a_{k+2} = 0; \ k \ge 2$$
(3.20)

- It is difficult to know if the sequence $(a_k)_{k\in\mathbb{N}}$ belongs to \mathbb{B}_s .
- In the vicinity of $z = i\sqrt{2}\xi = \infty$, the difficulty remains even if we use the techniques of the Thomé series in equation (3.3) by using the proposition 8 of [22] where one can choose as two linearly independent solutions of (3.3) the functions ψ_1 and ψ_2 , defined by the power series:

$$\psi_1(\xi) = e^{-\frac{1}{2}(\gamma - \alpha - 2)} \sum_{n=0}^{\infty} a_n(\alpha, \beta, \gamma, \delta) \xi^{-n}$$
(3.21)

$$\psi_2(\xi) = e^{-\frac{1}{2}(\gamma + \alpha + 2)} e^{\xi^2 + \beta \xi} \sum_{n=0}^{\infty} a_n(\alpha, i\beta, -\gamma, -i\delta) (-i\xi)^{-n}$$
(3.22)

where $\alpha = -1, \beta = -\frac{\mu\sqrt{2}}{\lambda}, \gamma = 1, \delta = \frac{2\sigma\sqrt{2}}{\lambda}$ and the complex numbers a_n are defined by (\star_3) .

Now as $\varphi(z) = \psi(\xi)$ where $z = i\sqrt{\xi}$, then we deduce from (3.21) and (3.22) that (3.2) at ∞ , satisfies:

$$\varphi_1(z) \sim const. \text{ as } z \longrightarrow \infty$$
 (3.23)

$$\varphi_2(z) \sim \frac{1}{z} e^{-\frac{1}{2}z^2 + i\rho z} \text{ as } z \longrightarrow \infty$$
 (3.24)

So that for any given direction in the complex z-plane an eigenfunction will be asymptotically a linear combination of (3.23) and (3.24)

In the following, we will show that Eq (3.24) is not acceptable in Bargmann space and thus it can be rejected.

• As there are many important applications of the closedness of the range of linear operator in the spectral study of differential operators and also in the context of perturbation theory (see e.g. [Goldberg]) [9] then to require an asymptotic behavior of the solutions of the eigenvalue problem associated with $H_{\mu,\lambda}$, we begin by showing that the image of $H_{\mu,\lambda}$ is closed that permit us to deduce a natural asymptotic behavior of the solutions of the eigenvalue problem associated to $H_{\mu,\lambda}$ in vicinity of $z=\infty$

Proposition 3.3

Let $\mathbb{B}_0 = \{ \varphi \in \mathbb{B}; \varphi(0) = 0 \}$ and $H_{\mu,\lambda}^{mim} = H_{\mu,\lambda}$ with domain $D(H_{\mu,\lambda}^{mim})$ and $\mathcal{R}(H_{\mu,\lambda}^{mim})$ denotes the range of $H_{\mu,\lambda}^{mim}$ then

i) In Bargmann space
$$\mathbb{B}$$
, we have $\varphi \in \mathbb{B} \iff z \longrightarrow g(z) = \frac{\varphi^{'}(z) - \varphi^{'}(0)}{z} \in \mathbb{B}$.

ii) For
$$\mu \neq 0$$
, $||H_{\mu,\lambda}^{mim}\varphi|| \geq |\mu|||\varphi||$, $\forall \varphi \in D(H_{\mu,\lambda}^{mim})$.

- iii) $\mathcal{R}(H_{\mu,\lambda}^{mim})$ is closed in Bargmann space.
- iv) $\mathcal{R}(H_{\mu,\lambda}^{mim})$ is dense in Bargmann space.

v) For
$$\mu \neq 0$$
, $H_{\mu,\lambda}^{mim}$ is invertible.

Remark 3.4

- 1) The properties of this proposition are well known. In particular (i) is the lemma 1 page 127 in [3] see also [12] or lemma 0.3 page 338 in [15].
- 2) It is convenient to reproduce the proofs of these properties which serve as a starting point of our study on the completeness of eigenfunctions of $H_{\mu,\lambda}$.

Proof

i) For
$$\phi(z) = \sum_{n=0}^{\infty} a_n z^n$$
, we have $\phi(z) = \sum_{n=0}^{\infty} a_{n+1} z^{n+1}$ and $\frac{\phi'(z) - \varphi'(0)}{z} = \sum_{n=0}^{\infty} a_{n+2} z^{n+2}$.

By using the isometry between spaces \mathbb{B} and \mathbb{B}_s , we deduce that

$$||\frac{\phi'(z) - \varphi'(0)}{z}||^2 = \sum_{n=0}^{\infty} n!(n+2)^2 |a_{n+2}|^2.$$

i.e.

$$\left\| \frac{\phi'(z) - \varphi'(0)}{z} \right\|^2 = \sum_{n=0}^{\infty} \frac{n+2}{n+1} (n+2)! \mid a_{n+2} \mid^2.$$
 (3.23)

As
$$\frac{n+2}{n+1} \le 2$$
 then $||\frac{\phi'(z) - \varphi'(0)}{z}||^2 \le 2\sum_{n=0}^{\infty} (n+2)! |a_{n+2}|^2$ Then

if
$$\varphi \in \mathbb{B}$$
, we have $z \longrightarrow \frac{\varphi'(z) - \varphi'(0)}{z} \in \mathbb{B}$.

For the reciprocity, we remark that $\frac{n+2}{n+1} > 1$ and from (3.23) we deduce that $||\varphi|| \le ||g||$ where $g(z) = \frac{\varphi'(z) - \varphi'(0)}{z}$.

ii) For $\lambda \in \mathbb{R}$ we observe that

$$|\Re e < H_{\mu,\lambda}^{min} \varphi, \varphi > |=|\mu| ||A\varphi||^2 = |\mu| \sum_{n=0}^{\infty} (n+1) |a_n|^2 \ge |\mu| \sum_{n=0}^{\infty} |a_n|^2 = |\mu| ||\varphi||^2$$
 (3.24)

Now by using the Cauchy's inequality, we deduce that

$$(\alpha) \mid\mid H_{\mu,\lambda}^{min} \varphi \mid\mid \geq \mid \mu \mid\mid\mid A\varphi \mid\mid, \forall \varphi \in D(H_{\mu,\lambda}^{min})$$
(3.25)

In particular

(
$$\beta$$
) the injection of $D(H_{\mu,\lambda}^{min})$ in $D(A)$ is continuous. (3.26)

and

$$(\gamma) || H_{\mu,\lambda}^{min} \varphi || \ge |\mu| || \varphi ||, \forall \varphi \in D(H_{\mu,\lambda}^{min})$$

$$(3.27)$$

(iii) From (3.25) and as the injection of D(A) in \mathbb{B} is a compact mapping, we deduce that :

The injection
$$D(H_{\mu,\lambda}^{min}) \longrightarrow \mathbb{B}$$
 is a compact mapping. (3.28)

(iv) From (3.27), it follows that $\mathcal{N}(H_{\mu,\lambda}^{min}) = \{0\}$ and $\mathcal{R}(H_{\mu,\lambda}^{min})$ is closed, where $\mathcal{N}(H_{\mu,\lambda}^{min})$ and $\mathcal{R}(H_{\mu,\lambda}^{min})$ denote the kernel and the range of $H_{\mu,\lambda}^{min}$ respectively.

(v) In fact,
$$\langle H_{\mu,\lambda}^{min}\varphi,\psi\rangle = 0 \iff -i\lambda z\psi^{''}(z) + (-i\lambda z^2 + \mu z)\psi^{'}(z) = 0 \iff H_{\mu,-\lambda}\psi = 0.$$

for
$$\lambda \neq 0$$
, we have $\psi'(z) = e^{-\frac{1}{2}z^2 + i\frac{\mu}{\lambda}z}$.

Now we show that the normalizability requirement for $\psi(z)$ is not verified in some direction of the z-plane.

In fact, we will use the fundamental property (i) for $z \longrightarrow -i\infty$.

$$\text{Let } z = \Re ez + i\Im mz \text{, then } e^{-|z|^2} \mid \frac{\psi'(z)}{z} \mid = e^{-|z|^2} \mid \frac{e^{-\frac{1}{2}z^2 + i\frac{\mu}{\lambda}z}}{z} \mid^2 = \frac{1}{\mid z\mid^2} e^{-2(\Re ez)^2} e^{-2\frac{\mu}{\lambda}\Im mz}.$$

Given μ and λ , there exists a direction of z-plane where $\psi(z) = \int_0^z e^{-\frac{1}{2}z\xi^2 + i\frac{\mu}{\lambda}\xi} d\xi$ cannot be considered in Bargmann space.

Remark 3.5

1) To sum up in $\mathbb{B}_0 = \{ \varphi \in \mathbb{B}; \varphi(0) = 0 \}$ the Hilbert space orthogonal to the vacuum (the only state of exactly zero energy), the eigenvalue conditions of $H_{\mu,\lambda}\varphi(z) = \sigma\varphi(z)$ are :

$$\varphi(z) \sim z \text{ for } z \sim 0$$
 (3.29_a)

and

$$\varphi(z) \sim const. \text{ for } z \sim -i\infty$$
 (3.29_b)

2) From iv) and v) we deduce that $H_{\mu,\lambda}^{min}$ is invertible i.e.,

$$0 \in \rho(H_{\mu,\lambda}^{min})$$
 where $\rho(H_{\mu,\lambda}^{min})$ denotes the resolvent set of $H_{\mu,\lambda}^{min}$. (3.30)

- 4) From (3.28) and as $\rho(H_{\mu,\lambda}^{min}) \neq \emptyset$ we deduce that the resolvent of $H_{\mu,\lambda}^{min}$ is compact.
- 5) From [15] (see theorem 1.2), it is well known that for $\mu \neq 0$ we have $D(H_{\mu,\lambda}^{min}) = D(H_{\mu,\lambda})$.
- 6) From lemmas 9, 10 and 11 of the ref. [13] (see page 280), it is well known that the eigenvalues of $H_{\mu,\lambda}$ are real.
- \S 4 A transformation procedure of $H_{\mu,\lambda}$ to a symmetric operator with compact resolvent on the negative imaginary axis

We begin by recalling an original lemma on Bargmann space with its proof was established in [13]

Lemma 4.1

Let \mathbb{B} be Bargmann space, if $\varphi \in \mathbb{B}$ then its restriction on $x + i\mathbb{R}$ is square integrable function with measure $e^{-|y|^2}dy, y \in \mathbb{R}$ for all fixed $x \in \mathbb{R}$.

Proof

Bargmann has built an isometry between the space \mathbb{B} and $L_2(\mathbb{R})$ so that, $\forall \varphi \in \mathbb{B}$ is uniquely represented by $f \in L_2(\mathbb{R})$ by means of the following integral:

$$\varphi(z) = c \int_{\mathbb{D}} e^{-\frac{z^2}{2} - \frac{q^2}{2} + \sqrt{2}zq} f(q) dq \tag{4.1}$$

such that
$$||\varphi|| = ||f||_{L_2(\mathbb{R})}$$
 (4.2)

Now, we put $g_z(q) = e^{-\frac{q^2}{2} + \sqrt{2}zq}$, which is function belonging to $L_2(\mathbb{R})$. Then we can write (4.1) in the following form:

$$e^{\frac{z^2}{2}}\phi(z) = c\int_{\mathbb{D}} g_z(q)f(q)dq \tag{4.3}$$

As $g_z(q)$ and f are in $L_2(\mathbb{R})$, we can apply the Parseval's identity at (4.3) to get :

$$e^{\frac{z^2}{2}}\varphi(z) = c_1 \int_{\mathbb{D}} \hat{g}_z(p)\hat{f}(p)dp \tag{4.4}$$

As $g_z(q)$ is a Gaussian's function, one knows how to calculate its Fourier transform:

$$\hat{g}_z(p) = \int_{\mathbb{R}} e^{ipq} g_z(q) dq = \int_{\mathbb{R}} e^{-\frac{q^2}{2} + (\sqrt{2}z - ip)q} dq = (2\pi)^{\frac{1}{2}} e^{\frac{1}{2}(\sqrt{2}z - ip)^2}$$

i.e.

$$\hat{g}_z(p) = (2\pi)^{\frac{1}{2}} e^{\frac{1}{2}(\sqrt{2}z - ip)^2} \tag{4.5}$$

Let z=x+iy, then for all fixed $x\in\mathbb{R}$, the function $\hat{g}_z(p)$ can be written under following form :

$$\hat{g}_z(p) = (2\pi)^{\frac{1}{2}} e^{\frac{1}{2}(\sqrt{2}x - i(p - \sqrt{2}y)^2}$$
(4.6)

If we put $h_x(p) = e^{x^2} e^{-\frac{1}{2}(p^2 + i\frac{\sqrt{2}}{2}px)}$ then $e^{\frac{z^2}{2}}\varphi(z)$ is the convolution product of h_x with \hat{f} evaluated in $\sqrt{2}y$, i.e.,

$$e^{\frac{(x+iy)^2}{2}}\varphi(x+iy) = C_2 h_x * \hat{f}(\sqrt{2}y)$$
(4.7)

It follows that $y \to e^{\frac{(x+iy)^2}{2}} \varphi(x+iy)$ is in $L_2(\mathbb{R})$ for all $x \in \mathbb{R}$

By applying the Young's inequality we deduce that :

$$||e^{\frac{(x+iy)^2}{2}}\varphi(x+iy)||_{L_2(\mathbb{R})}^2 \le ||h_x||_{L_1(\mathbb{R}^N)}^2 \cdot ||\hat{f}||_{L_2(\mathbb{R})}^2$$
(4.8)

where C_1, C_2 and C are constants.

As $||\ h_x\ ||^2_{L_1(\mathbb{R})} = C_3 e^{2|x|^2}$ et $||\ \hat{f}\ ||^2_{L_2(\mathbb{R})} = C_4\ ||\ f\ ||^2_{L_2(\mathbb{R})},$ then we deduce that :

$$\int_{\mathbb{P}} e^{-|y|^2} |\varphi(x+iy)|^2 dy \le C_5 e^{-|x|^2} ||\varphi||^2 \tag{4.9}$$

where C_5 depends of all the previous constants.

• From the above lemma, it is possible to restrict ourselves to the negative imaginary axis, z = -iy, y > 0, where we impose the boundary conditions (3.29).

Let us write therefore $u(y) = \varphi(-iz)$ then we have

$$H_{R}^{\mu,\lambda}u(y) = -\lambda y u''(y) + (\lambda y^{2} + \mu y)u'(y) = \lambda \left[-y\frac{d^{2}}{dy^{2}} + (y^{2} + \rho y)\frac{d}{dy}\right]u(y);$$

$$\rho = \frac{\mu}{\lambda}$$
(4.10)

The eigenfunction equation $H_{\scriptscriptstyle R}\mu, \lambda u(y) = \sigma u(y)$ may be rewritten (for $\lambda \neq 0$) in the form :

$$\begin{cases} \lambda y \left[-\frac{d^2}{dy^2} + (y + \rho) \frac{d}{dy} \right] u(y) = \sigma u(y) \\ u(y) \sim y, \quad y \longrightarrow 0 \\ u(y) \sim const., \quad y \longrightarrow +\infty \end{cases}$$
(4.11)

The term proportional to u'(y) can be eliminated by the transformation

$$u(y) = T(y)v(y)$$
 where $T(y) = e^{\frac{1}{4}(y+\rho)^2}$ (4.11)

that gives the eigenvalue equation

$$\begin{cases} \lambda y \left[-\frac{d^2}{dy^2} + \frac{1}{4}(y+\rho)^2 - \frac{1}{2} \right] v(y) = \sigma v(y) \\ v(y) \sim y, \quad y \longrightarrow 0 \\ v(y) \sim e^{-\frac{y}{2}(\frac{y}{2}+\rho)}, \quad y \longrightarrow +\infty \end{cases}$$

$$(4.12)$$

As in vicinity of y=0 we have $v(y) \sim y$ and in vicinity of $y=+\infty$ we have $|v(y)|^2 \sim e^{-(\frac{y^2}{2}+\rho y)}$ then v vanishes at origin and $v \in L^2[0,\infty[$.

We denote by $L_R^2[0,\infty[$ the space of functions in $L^2[0,\infty[$ satisfying both contraints.

$$\text{Let } \underline{H}^{\mu,\lambda} = e^{-T(y)} H_{\scriptscriptstyle R}^{\mu,\lambda} e^{T(y)}, \text{ then } \underline{H}\mu, \lambda = \lambda y [-\frac{d^2}{dy^2} + \frac{1}{4}(y+\rho)^2 - \frac{1}{2}].$$

Now, if we consider the change of variable $y=x^2$ and the similarity transform given by $y^{\frac{1}{4}}=x^{\frac{1}{2}}$ then we get

On completeness of eigenfunctions of $H_{\mu,\lambda}$

A. Intissar

$$\tilde{H}_{\mu,\lambda} = x^{-\frac{1}{2}} \underline{H}^{\mu,\lambda} x^{\frac{1}{2}}$$

$$= x^{-\frac{1}{2}} e^{-T(x^2)} H_{{\scriptscriptstyle R}} \mu, \lambda e^{T(x^2)} x^{\frac{1}{2}}$$

$$= \frac{\lambda}{4} \left[-\frac{d^2}{dx^2} + \frac{3/4}{x^2} + x^2((x^2 + \rho)^2 - 2) \right] \tag{4.13}$$

$$v(y) = (x^2)^{\frac{1}{4}} w(x) = x^{\frac{1}{2}} w(x), \quad y = x^2.$$

In particular

$$w(x) = x^{-\frac{1}{2}}u(y) \sim x^{\frac{3}{2}}, \ x \longrightarrow 0$$
 (4.14)

and

$$w(x) \sim x^{-\frac{1}{2}} e^{-\frac{1}{4}(x^2 + \rho)^2}, \ x \longrightarrow \infty.$$
 (4.15)

Remark 4.2

- 1) The similarity transformation from $H_{\mu,\lambda}$ to $\tilde{H}_{\mu,\lambda}$ is non-unitary, but bijective.
- 2) On $L_2[0, +\infty[, \tilde{H}_{\mu,\lambda}]$ with domain $D(\tilde{H}_{\mu,\lambda}) = \{w \in L_2[0, +\infty[; \tilde{H}_{\mu,\lambda}] w \in L_2[0, +\infty[]\}$ is a symmetric operator.
- 3) As the injection of maximal domain of $H_{\mu,\lambda}$ in Bargmann space $\mathbb B$ is compact then we deduce that :
- (i) the injection of maximal domain of $\tilde{H}_R^{\mu,\lambda}$ in

$$\mathbb{B}_R = \{ u : [0, +\infty[\longrightarrow \mathbb{C} \text{ analytic}; \int_0^\infty |u(y)|^2 e^{-y^2} dy < +\infty \}$$

is compact.

and

- (ii) the injection of maximal domain of $\tilde{H}_{\mu,\lambda}$ in $L_2[0,+\infty[$ is also compact, in particular $\tilde{H}_{\mu,\lambda}$ is a symmetric operator with compact resolvent.
- 4) The equation $H_{\mu,\lambda}u = \sigma u$ is equivalent to $\tilde{H}w = \sigma w$, where

$$w(x) = x^{-\frac{1}{2}} e^{-\frac{1}{4}(x^2 + \rho)^2} u(y = x^2). \tag{4.14}$$

is an analytical function in x.

We end this section by the main result of this paper

Proposition 4.3

The eigenfunctions of $H_{\mu,\lambda}, \mu > 0$ form a complete basis in Bargmann space.

Proof

The above formulated problem of finding the spectrum of the Hamiltonian $H_{\mu,\lambda}$ or of $H_R^{\mu,\lambda}$ acting on Bargmann space or on its restriction functions respecti to negative imaginary axis is equivalent to the problem of finding the spectrum of the symmetric operator $\tilde{H}_{\mu,\lambda}$ in the space $L^2([0,+\infty[)$.

As $\tilde{H}_{\mu,\lambda}$ is symmetric with compact resolvent, then by using the spectral theorem for symmetric operators with compact resolvent, we deduce that the eigenfunctions of $\tilde{H}_{\mu,\lambda}$ form a complete basis:

Theorem 4.4 (spectral theorem for symmetric operators with compact resolvent)

- 1) Let T be a densely defined linear operator in a Hilbert space \mathcal{H} and let $\sigma \in \rho(T) \cap \mathbb{R}$ then T is symmetric iff $R_{\sigma} = (T \sigma I)^{-1}$ is self-adjoint.
- 2) Let T be a symmetric operator in a Hilbert space \mathcal{H} such that $(T \sigma_0 I)^{-1}$ is compact, then $\sigma_p(T) = \{\sigma_0 + \frac{1}{\gamma}; \gamma \in \sigma_p(T \sigma_0 I)^{-1}\}$ where $\sigma_p(T)$ denotes the ponctual spectrum of T.
- 3) If T is a symmetric operator with compact resolvent then there exists $\sigma_1 \in \rho(T) \cap \mathbb{R}$ such that $(T \sigma_1 I)^{-1}$ is a compact self-adjoint operator.
- 4) Let T be a symmetric operator with compact resolvent in a Hilbert space \mathcal{H} .

For each $\sigma \in \sigma_p(T)$, define $n(\sigma) = \dim \ker(T - \sigma I)$, and choose an orthonormal basis

$$\{\varphi_{\sigma,1},\varphi_{\sigma,2},...,\varphi_{\sigma,n(\sigma)}\}\$$
of $ker(T-\sigma I)$. Then $\{\varphi_{\sigma,i};\sigma\in\sigma_p(T),i=1,2,...,n(\sigma)\}$ forms

a complete orthonormal set for \mathcal{H} . : Moreover, we have the following :

i)
$$\varphi \in D(T) \iff \sum_{\sigma \in \sigma_n(T)} \sum_{i=1}^{n(\sigma)} |\sigma|^2 |\langle \varphi, \varphi_{\sigma,i} \rangle|^2 < \infty.$$

ii) For any
$$\varphi \in D(T)$$
, $T\varphi = \sum_{\sigma \in \sigma_p(T)} \sum_{i=1}^{n(\sigma)} \sigma < \varphi$, $\varphi_{\sigma,i} > \varphi_{\sigma,i}$.

\S 5 An explicit integral form of the inverse of $H_{\mu,\lambda}$ on negative imaginary axis and some spectral properties of $H_{\mu,\lambda}^{-1}$

A) Implementation of integral equation associated to $H_{\mu,\lambda}$

On Bargmann space, we consider the equation $H_{\mu,\lambda}\varphi(z) = \tilde{\sigma}\varphi(z)$. If we put $\rho = \frac{\mu}{\lambda}$ and $\sigma = \frac{\tilde{\sigma}}{\lambda}$ with $\lambda \neq 0$ then this eigenvalue equation can be written in the following form:

$$\varphi''(z) + p(z)\varphi'(z) + q(z)\varphi(z) = 0; p(z) = (z - i\rho) \text{ and } q(z) = \frac{i\sigma}{\lambda z}$$

$$(5.1)$$

We restrict the study of the equation (5.1) on the lines parallel to the imaginary axis:

Let z = a + iy and $\varphi(a + iy) = \varphi_a(y)$ where a is fixed. Then we get

$$\begin{cases}
\varphi_a^{"}(y) - (y - \gamma)\varphi_a^{'}(y) - \frac{i\sigma}{a + iy}\varphi_a(y) = 0 \\
\gamma = \rho + ia
\end{cases}$$
(5.2)

We put

$$\varphi_a'(y) = K_a(y)e^{\frac{1}{2}(y-\gamma)^2} \tag{5.3}$$

and substitute it in (5.2) then we get

$$K'_{a}(y) = i\sigma e^{-\frac{1}{2}(y-\gamma)^{2}} \frac{\varphi_{a}(y)}{a+iy}$$
(5.4)

We choose a primitive representation of the function $y \longrightarrow K_a^{'}(y)$ as follows

$$K_a(y) = i\sigma \int_{-\infty}^{y} e^{-\frac{1}{2}(u-\gamma)^2} \frac{\varphi_a(u)}{a+iu} du + c_a; y \in]-\infty, 0]$$
(5.5)

where c_a is a constant.

Consequently, equation (5.3) can be written as follows

$$\varphi_{a}'(y) = i\sigma e^{\frac{1}{2}(y-\gamma)^{2}} \int_{-\infty}^{y} e^{-\frac{1}{2}(u-\gamma)^{2}} \frac{\varphi_{a}(u)}{a+iu} du + c_{a} e^{\frac{1}{2}(y-\gamma)^{2}}$$
(5.6)

Lemma 5.1

For $\rho > 0$ the representation (5.5) has a sense.

Proof

For $\varphi \in \mathbb{B}$ we have $\int_{\mathbb{R}^2} e^{-(x^2+y^2)} |\varphi(x+iy)|^2 dxdy < +\infty$ then we apply the Fubini theorem to deduce that

$$\int_{\mathbb{R}} e^{-y^2} \mid \varphi_x(y) \mid^2 dy < +\infty \tag{5.7}$$

Now, as $\int_{-\infty}^{y} e^{-\frac{1}{2}(u-\gamma)^2} \frac{\varphi_a(u)}{a+iu} du = e^{-\frac{1}{2}\gamma^2} \int_{-\infty}^{y} e^{-\frac{1}{2}u^2} \varphi_a(u) \frac{e^{\gamma u}}{a+iu} du$ and we look for φ in Bargmann space \mathbb{B} , we have on the one hand $u \longrightarrow e^{-\frac{1}{2}u^2} \varphi_a(u)$ is square integrable and on the other hand for $\rho > 0$ the function $u \longrightarrow \frac{e^{\gamma u}}{a+iu}$ is also square integrable on $]-\infty,0]$. Consequently, $\int_{-\infty}^{y} e^{-\frac{1}{2}(u-\gamma)^2} \frac{\varphi_a(u)}{a+iu} du \text{ has a sense.}$

Remark 5.2

(i) If $\rho < 0$, we choose a primitive representation of the function $y \longrightarrow K_a^{'}(y)$ as follows

On completeness of eigenfunctions of $H_{\mu,\lambda}$

A. Intissar

$$K_a(y) = i\sigma \int_y^{+\infty} e^{-\frac{1}{2}(u-\gamma)^2} \frac{\varphi_a(u)}{a+iu} du + c_a, c_a \text{ is constant}$$

$$(5.8)$$

This representation has a sense.

(ii) From (5.3) and (5.5) we deduce that

$$\varphi_{a}'(y) = i\sigma e^{\frac{1}{2}(y-\gamma)^{2}} \int_{u}^{+\infty} e^{-\frac{1}{2}(u-\gamma)^{2}} \frac{\varphi_{a}(u)}{a+iu} du + c_{a} e^{\frac{1}{2}(u-\gamma)^{2}}$$
(5.9)

Theorem 5.3

(i) In the representation (5.5) we have $c_a = 0$

(ii)
$$\lim e^{-\frac{1}{2}(y-\gamma)^2} \varphi_a'(y) = 0$$
 as $y \longrightarrow -\infty$ (boundary condition) (5.10)

Proof

(i) By multiplying the two members of the equation (5.9) by $e^{-\frac{1}{2}(u-\gamma)^2}$, we get

$$e^{-\frac{1}{2}(u-\gamma)^{2}}\varphi_{a}'(y) = i\sigma \int_{y}^{+\infty} e^{-\frac{1}{2}(u-\gamma)^{2}} \frac{\varphi_{a}(u)}{a+iu} du + c_{a}$$
(5.11)

As the function $u \longrightarrow e^{-\frac{1}{2}(u-\gamma)^2} \frac{\varphi_a(u)}{a+iu}$ is integrable we deduce that

$$Lim e^{-\frac{1}{2}(y-\gamma)^{2}} \varphi_{a}^{'}(y) = c_{a} \text{ as } y \longrightarrow -\infty$$

$$(5.12)$$

Now, if $c_a \neq 0$ then we get that

$$\varphi_{a}^{'}(y) \sim c_{a} e^{\frac{1}{2}(y-\gamma)^{2}} \text{ and } \frac{e^{-\frac{1}{2}y^{2}}}{y} \varphi_{a}^{'}(y) \sim c_{a} \frac{e^{-\rho y}}{y} e^{\frac{\gamma^{2}}{2} - ia}$$
 (5.13)

We consider the space

$$\mathbb{B}_a = \{ \varphi_a : \mathbb{R} \longrightarrow \mathbb{C} \text{ entire}; \int_{\mathbb{R}} e^{-\frac{1}{2}y^2} \mid \varphi_a(y) \mid^2 dy < +\infty \}$$
 (5.14)

Then

- (α) From lemma (4.1), we deduce that $\forall \varphi \in \mathbb{B}$ then $\varphi_a \in \mathbb{B}_a$
- (β) From (i) of proposition (3.3), we deduce that $\forall \varphi_a \in \mathbb{B}_a$ then the function

$$y \longrightarrow \frac{\varphi_a'(y) - \varphi_a'(0)}{y} \in \mathbb{B}_a$$

As in (5.13) the function $y \longrightarrow \frac{e^{-\rho y}}{y} e^{\frac{\gamma^2}{2} - ia}$ is not square integrable then $\varphi_a \notin \mathbb{B}_a$

which is contradictory with the property (β) and therefore $c_a = 0$.

(ii) we apply (i) to (5.12).

Remark 5.4

By applying the above theorem, equation (5.6) can be written as follows

$$\varphi_{a}'(y) = i\sigma e^{\frac{1}{2}(y-\gamma)^{2}} \int_{-\infty}^{y} e^{-\frac{1}{2}(u-\gamma)^{2}} \frac{\varphi_{a}(u)}{a+iu} du; \ y \in]-\infty, 0]$$
(5.15)

B) Justification for crossing to the limit in the equation (5.15) as $a \longrightarrow 0$

Proposition 5.5

If $a \longrightarrow 0$ in (5.15) then we get

$$\varphi_0'(y) = \sigma e^{\frac{1}{2}(y-\rho)^2} \int_{-\infty}^y e^{-\frac{1}{2}(u-\rho)^2} \frac{\varphi_0(u)}{u} du; \ y \in]-\infty, 0]$$
(5.16)

Proof

As the functions
$$a \longrightarrow \varphi'_a(y)$$
 and $a \longrightarrow \frac{e^{-\frac{1}{2}(u-\rho-ia)^2}}{a+iu}\varphi_a(u)$ are continuous then $Lim \varphi'_a(y) = \varphi'_0(y)$ and $Lim \frac{e^{-\frac{1}{2}(u-\rho-ia)^2}}{a+iu}\varphi_a(u) = \frac{e^{-\frac{1}{2}(u-\rho)^2}}{iu}\varphi_0(u)$ as $a \longrightarrow 0$. (5.17)

We have to show that $u \longrightarrow \frac{e^{-\frac{1}{2}(u-\rho)^2}}{iu} \varphi_0(u)$ is integrable. For this purpose we consider the following function

$$f_a(u) = \frac{e^{-\frac{1}{2}(u-\rho-ia)^2}}{a+iu} \varphi_a(u)$$
 (5.18)

As for $\epsilon > 0$ enough small we have $\varphi_a(a + iu) \sim a + iu; u \in]-\epsilon, 0]$ (by using lemma 4.1), we deduce that

$$|f_a(u)| \le e^{\rho u}. \tag{5.19}$$

Now, for $u \in]-\infty, -\epsilon[$, we observe that

$$|f_a(u)| \le e^{-\frac{\rho^2}{2}} \frac{e^{\rho u}}{u}.$$
 (5.20)

As the function

$$g(u) = \begin{cases} e^{-\frac{\rho^2}{2}} \frac{e^{\rho u}}{u}; u \in]-\infty, -\epsilon[\\ e^{\rho u}; u \in]-\epsilon, 0] \end{cases}$$

$$(5.21)$$

is integrable and the the sequence $f_a(u)$ converges pointwise to $\frac{e^{-\frac{1}{2}(u-\rho^2)}}{iu}\varphi_0(u)$ as $a\longrightarrow 0$ then by applying the Lebesgue's Dominated Convergence Theorem, we deduce that

$$Lim \int_{-\infty}^{y} \frac{e^{-\frac{1}{2}(u-\rho-ia)^{2}}}{a+iu} \varphi_{a}(u)du = \int_{-\infty}^{y} \frac{e^{-\frac{1}{2}(u-\rho^{2})}}{iu} \varphi_{0}(u) \text{ as } a \longrightarrow 0$$

$$(5.22)$$

Consequently, we get

$$\varphi_0'(y) = \sigma e^{\frac{1}{2}(y-\rho)^2} \int_{-\infty}^y e^{-\frac{1}{2}(u-\rho)^2} \frac{\varphi_0(u)}{u} du; \ y \in]-\infty, 0]$$
(5.23)

Theorem 5.6

(i) An operator integral associated to $H_{\mu,\lambda}$ is given as follows

$$K\psi(y) = \int_{-\infty}^{0} \mathcal{N}(y, s)\psi(s)ds$$
 (5.24)

where

$$(\alpha) \ \psi(y) = \frac{\varphi_0'(y)}{y} e^{-\frac{y^2}{2}} \theta(y)$$
with $\theta(y) = \begin{cases} y; & y \in [-1, 0] \\ -1; & y \in [-\infty, -1] \end{cases}$ (5.25)

$$(\beta) \mathcal{N}(y,s) = e^{-\rho y} \frac{\theta(y)}{y} \cdot \frac{s}{\theta(s)} e^{\frac{s^2}{2}} \int_{-\infty}^{\min(y,s)} e^{-\frac{1}{2}(u-\rho)^2} u du$$

$$\text{where } \min(y,s) = \begin{cases} y; & y \le s \\ s; & s \le y \end{cases}$$

$$(5.26)$$

$$(\gamma) \ \psi \in L_2(]-\infty,0], \theta(y)dy) \tag{5.27}$$

(ii) For $\psi \in \mathbb{B}_0$, an explicit inverse of $H_{\mu,\lambda}$ restricted on imaginary axis; $y \in [0, +\infty[$ is given by

$$H_{\mu,\lambda}^{-1}\psi(-iy) = \int_0^\infty \mathcal{N}_{\mu,\lambda}(y,s)\psi(-is)ds$$
where $\mathcal{N}_{\mu,\lambda}(y,s) = \frac{1}{\lambda s}e^{-\frac{s^2}{2} - \frac{\mu}{\lambda}s} \int_0^{\min(y,s)} e^{\frac{u^2}{2} + \frac{\mu}{\lambda}u} du$

$$(5.28)$$

Proof

(i) We substitue $\varphi_0(u) = -\int_u^0 \varphi_0'(s) ds$ in (5.23) to obtain $\varphi_0'(y) = -\sigma e^{\frac{1}{2}(y-\rho)^2} \int_{-\infty}^y \frac{e^{-\frac{1}{2}(u-\rho)^2}}{u} \int_u^0 \varphi_0'(s) ds; \ y \in]-\infty, 0]$ (5.29)

By applying the Fubini theorem we get

$$\int_{-\infty}^{y} \frac{e^{-\frac{1}{2}(u-\rho)^{2}}}{u} \int_{u}^{0} \varphi_{0}^{'}(s)ds = \int_{-\infty}^{0} \left[\int_{-\infty}^{\min(y,s)} \frac{e^{-\frac{1}{2}(u-\rho)^{2}}}{u} du\right] \varphi_{0}^{'}(s)ds \tag{5.30}$$

and

$$\varphi_0'(y) = -\sigma e^{\frac{1}{2}(y-\rho)^2} \int_{-\infty}^0 \left[\int_{-\infty}^{\min(y,s)} \frac{e^{-\frac{1}{2}(u-\rho)^2}}{u} du \right] \varphi_0'(s) ds \tag{5.31}$$

or

On completeness of eigenfunctions of $H_{\mu,\lambda}$

$$e^{-\frac{y^{2}}{2}}\varphi_{0}'(y) = -\sigma e^{\frac{\rho^{2}}{2}} \int_{-\infty}^{0} \left[e^{\rho y} \int_{-\infty}^{\min(y,s)} \frac{e^{-\frac{1}{2}(u-\rho)^{2}}}{u} du\right] \varphi_{0}'(s) ds$$
 (5.32)

Now, if we put
$$\psi(y) = \frac{\varphi_0'(y)}{y} e^{-\frac{y^2}{2}} \theta(y)$$
 (5.33)

with
$$\theta(y) = \begin{cases} y; & y \in [-1, 0] \\ -1; & y \in]-\infty, -1] \end{cases}$$

and if we substitute ψ in (5.32)) then we get

$$\psi(y) = -\sigma e^{\frac{\rho^2}{2}} \int_{-\infty}^{0} \mathcal{N}(y, s) \psi(s) ds \tag{5.34}$$

Where $\mathcal{N}(y,s)$ is given by (5.27).

(ii) Let $\varphi \in \mathbb{B}_0$ and $\psi \in \mathbb{B}_0$, we consider the equation $H_{\mu,\lambda}\varphi = \psi(z)$ i.e.

$$i\lambda z\varphi''(z) + (i\lambda z^2 + \mu z)\varphi'(z) = \psi(z)$$
(5.35)

Let z = -iy, $u(y) = \varphi(-iy)$ and $f(y) = \psi(-iy)$ with $y \in [0, \infty[$ then (5.34) can be

written in the following form:

$$-\lambda y u''(y) + (\lambda y^2 + \mu y) u'(y) = f(y)$$
(5.36)

or

$$-u''(y) + (y + \rho)u'(y) = \frac{f(y)}{\lambda y}$$
 (5.37)

with help lemma (4.1), the property i) of proposition (3.3) and a similar technique used in proof of (i) of this theorem we get (5.28).

In the following, we show that the integral operator K is an operator of Hilbert-Schmidt on $L_2(]-\infty,0], \theta(y)dy)$ and we give some properties of kernel $\mathcal{N}(y,s)$ (see (5.28)) in $L^2(]-\infty,0]\times]-\infty,0], \theta(y)\theta(s)dyds)$.

Theorem 5.7

Let
$$\mathcal{N}(y,s) = e^{-\rho y} \frac{\theta(y)}{y} \frac{s}{\theta(s)} e^{\frac{1}{2}s^2} \int_{-\infty}^{\min(y,s)} \frac{e^{-\frac{1}{2}(u-\rho)^2}}{u} du$$
 be the kernel of the equation :

$$-\lambda y u''(y) + (\lambda y^2 + \mu y) u'(y) = \psi(-iy) , \psi \in \mathbb{B}_0$$
(5.38)

then:

- (i) $\mathcal{N}(y,s)$ is Hilbert-Shmidt on $\mathcal{N}(y,s)$ in $L^2(]-\infty,0]\times]-\infty,0]$, $\theta(y)\theta(s)dyds)$.
- (ii) $\mathcal{N}(y,s)$ in $L^2(]-\infty,0]\times]-\infty,0],\theta(y)\theta(s)dyds)$.

Proof

(i) To apply the theorem of dominated convergence, we will increase $\mathcal{N}(y,s)$ by a function $\tilde{\mathcal{N}}(y,s) \in L^2(]-\infty,0]\times]-\infty,0],\theta(y)\theta(s)dyds)$

For this we pose min(y, s) = m then we have :

$$\left| \frac{e^{-\frac{1}{2}(u-\rho)^2}}{u} \right| \le e^{-\frac{1}{2}\rho^2} \frac{1}{m(m-\rho)} \left| (\rho-u)e^{-\frac{1}{2}u^2+\rho u} \right|$$

The function $u \longrightarrow \frac{1}{u(u-\rho)}$ is bounded for $-\infty < u \le m < 0$

Hence a first increase :

$$|\mathcal{N}(y,s)| \le e^{-\frac{1}{2}\rho^2} |\frac{e^{-\rho y}}{y} \frac{\theta(y)}{\theta(s)} s e^{\frac{1}{2}s^2} \frac{e^{-\frac{1}{2}m^2 + \rho^m}}{m(m-\rho)} |$$

• If y < s, we have :

$$m = y$$
 et $e^{\frac{1}{2}s^2 - \frac{1}{2}y^2} < 1$.

Then:

$$\mid \mathcal{N}(y,s) \mid \leq e^{-\frac{1}{2}\rho^2} \mid \frac{\theta(y)}{y} \frac{s}{\theta(s)} \frac{1}{y(y-\rho)}$$

• If $s \leq y$, we have :

$$m = s \text{ et } e^{\rho(s-y)} < 1.$$

Therefore

$$|\mathcal{N}(y,s)| \le e^{-\frac{1}{2}\rho^2} |\frac{\theta(y)}{y} \frac{s}{\theta(s)} \frac{1}{s(s-\rho)}|$$

Finally we put:

$$\tilde{\mathcal{N}}(y,s) = \left\{ \begin{array}{ll} e^{-\frac{1}{2}\rho^2} \mid \frac{\theta(y)}{y} \frac{s}{\theta(s)} \frac{1}{y(y-\rho)} & pour \quad y \leq s \\ \\ e^{-\frac{1}{2}\rho^2} \mid \frac{\theta(y)}{y} \frac{s}{\theta(s)} \frac{1}{s(s-\rho)} & pour \quad s \leq y \end{array} \right.$$

to get
$$|\mathcal{N}(y,s)| \leq |\tilde{\mathcal{N}}(y,s)|$$

It remains to show that $\tilde{\mathcal{N}}(y,s) \in L^2(]-\infty,0]\times]-\infty,0],\theta(y)\theta(s)dyds$

Let
$$\Delta =]-\infty, 0] \times]-\infty, 0]-]-1, 0] \times]-1, 0], \Delta_1 = \{(y, s) \in \Delta; y \leq s\}$$
 and

$$\Delta_2 = \{ (y, s) \in \Delta; s \le y \}.$$

Then

$$\int_{-\infty}^{0} \theta(y) dy \int_{-\infty}^{0} \theta(s) \mid \tilde{\mathcal{N}}(y,s) \mid^{2} ds =$$

$$\int_{\Delta_{1}} \theta(y) \theta(s) \mid \frac{\theta(y)}{y} \frac{s}{\theta(s)} \frac{1}{y(y-\rho)} \mid^{2} dy ds + \int_{\Delta_{2}} \theta(y) \theta(s) \mid \frac{\theta(y)}{y} \frac{s}{\theta(s)} \frac{1}{s(s-\rho)} \mid^{2} dy ds$$

As $\rho > 0$ and the function $u \to \frac{1}{(u-\rho)^2}$ is integrable at the origin, we deduce that :

$$\int_{-\infty}^{0} \theta(y) dy \int_{-\infty}^{0} \theta(s) |\tilde{\mathcal{N}}(y,s)|^{2} ds < +\infty$$

then $\mathcal{N}(y,s)$ is a kernel of Hilbert-Shmidt.

Corollary 5.8 (some spectral properties of K)

I) We note that the functions:

$$y \longrightarrow \int_{-\infty}^{y} \frac{e^{-\frac{1}{2}(u-\rho)^2}}{u} du; \quad y \le s$$

and

$$s \longrightarrow \int_{-\infty}^{s} \frac{e^{-\frac{1}{2}(u-\rho)^2}}{u} du; \quad s \le y.$$

are decreasing and tend to zero respectively when $y \longrightarrow -\infty$ and $s \longrightarrow -\infty$.

- ii) The kernel $-\mathcal{N}(y,s)$ is non negative.
- iii) the operator $K\psi(y) = \int_{-\infty}^{0} \mathcal{N}(y,s)\psi(s)ds$ is an operator of Hilbert-Shmidt.
- (iv) As K is a Hilbert-Schmidt operator then its spectrum is discrete and as its kernel is non negative then by using Jentzsch theorem [17] or Krein-Rutman theorem [18], we deduce the existence of real eigenvalue of K.

In particular, we deduce the existence of real eigenvalue of $H_{\mu,\lambda}$; $\mu > 0$ and that $H_{\mu,\lambda}^{-1}$ is a positive operator on Bargmann space \mathbb{B} in the following sense:

Let $\mathcal{C} \subset \mathbb{B}$ be a closed cone with non empty interior $int(\mathcal{C})$ then $H^{-1}_{\mu,\lambda}(\mathcal{C} - \{0\}) \subset int(\mathcal{C})$.

(v) But the question of the existence of $\sigma > 0$ such that $(H_{\mu,\lambda} - \sigma I)^{-1}$ is a positive operator is always an open question.

References

- [1] Abramovitz, M. and Stegun, I. A.: Handbook of Mathematical Functions, New York (1968).
- [2] Agarwal, R.P. and Regan, D.O.: Ordinary and Partial Differential Equations: With Special Functions, Fourier Series, and Boundary Value Problems, Lecture 5.
- [3] Ando, T. and Zerner, M. :. Sur une valeur propre d'un opérateur, Comm. Math. Phys. 93 (1984).
- [4] Bargmann, V. :. On a Hilbert space of analytic functions and an associated integral transform I, Comm. Pure Appl. Math. 14 (1962) 187-214.
- [5] Batola, F. :. Une généralisation d'une formule de Meixner-tricomi, Can. J. Math., Vol. XXXIV, No. 2, 1982, pp. 411- 422

- [6] Bender, C. M., Steven A. and Orszag, S. A.: Advanced Mathematical Methods for Scientists and Engineers (McGraw-Hill Book Company, New York, 1978).
- [7] Bateman, H. and Erdelyi, A.:. Higher Transcendental Functions, Vol. 1, New York (1953).
- [8] Decarreau, A. Emamirad, H. and Intissar, A. :. Chaoticité de l'opérateur de Gribov dans l'espace de Bargmann, C. R. Acad. Sci. Paris Sér. I Math. 331 (2000) 751-756.
- [9] Goldberg, S.: Unbounded Linear Operators. Mc Graw Hill, New York, 1966.
- [10] Hille, E. :. Ordinary Differential Equations in the Complex Domain, Dover Publications Inc., New edition (1997)
- [11] Ince, E.L.: Ordinary Differential Equations, Dover, New York, 1956.
- [12] Intissar, A., Le Bellac, M. and Zerner, M. .. Properties of the Hamiltonian of Reggeon field theory, Phys. Lett. B 113 (1982) 487-489.
- [13] Intissar, A. :. Etude spectrale d'une famille d'opérateurs non-symétriques intervenant dans la théorie des champs de Reggeons, Comm. Math. Phys. 113 (1987) 263-297.
- [14] Intissar, A. :. Analyse de Scattering d'un opérateur cubique de Heun dans l'espace de Bargmann, Comm. Math. Phys. 199 (1998) 243-256.
- [15] Intissar, A. :. Spectral analysis of non-selfadjoint Jacobi-Gribov operator and asymptotic analysis of its generalized eigenvectors, Advances in Mathematics (China), Vol.44, no 3, (2015), 335-353, doi: 10.11845/sxjz.2013117b
- [16] Intissar, A. and Intissar, J. K.:. On Chaoticity of the Sum of Chaotic Shifts with Their Adjoints in Hilbert Space and Applications to Some Weighted Shifts Acting on Some Fock-Bargmann Spaces, Complex Anal. Oper. Theory, Volume 11, issue 3, (2017), 491-505
- [17] Jentzsch, R.:. Uber Integralgleichungen mit positiven Kern, J. Reine Angew. Math. 141 (1912), 235-244.
- [18] Krein, M. G. and M. A. Rutman, M. A.:. Linear operators leaving invariant a cone in a Banach space, Amer. Math. Soc. Transl. Ser. 1 10 (1950), 199-325 [originally Uspekhi Mat. Nauk 3 (1948), 3-95]
- [19] Kristensson, G. :. Second Order Differential Equations : Special Functions and Their Classification (Springer Science, New York, 2010).
- [20] Maroni, P. :. Biconfluent Heun equation, In : A. Ronveau (Ed.), Heun's diff erential equations, 191-249, Oxford University Press, Oxford 1995
- [21] Morse, P.M. and H Feshbach, H.: Methods of Theoretical Physics, I, http://people.physics.tamu.edu/pope/methods
- [22] Roseau, A.:. On the solutions of the biconfluent Heun equations, Bull. Belg. Math. Soc. Simon Stevin, Volume 9, Number 3 (2002), 321-342. doi:10.36045/bbms/1102715058.

On completeness of eigenfunctions of $H_{\mu,\lambda}$

 $A.\ Intissar$

 $\rm https://projecteuclid.org/euclid.bbms/1102715058$

[23] Yosida, K. :. Lectures on Differential and Integral Equations, pp. 37-40.