

ON THE EXISTENCE, UNIQUENESS AND STABILITY OF PERIODIC WAVES FOR THE FRACTIONAL BENJAMIN-BONA-MAHONY EQUATION

SABRINA AMARAL, HANDAN BORLUK, GULCIN M. MUSLU, FÁBIO NATALI*, AND GOKSU ORUC

ABSTRACT. The existence, uniqueness and stability of periodic traveling waves for the fractional Benjamin-Bona-Mahony equation is considered. In our approach, we give sufficient conditions to prove a uniqueness result for the single-lobe solution obtained by a constrained minimization problem. The spectral stability is then shown by determining that the associated linearized operator around the wave restricted to the orthogonal of the tangent space related to the momentum and mass at the periodic wave has no negative eigenvalues. We propose the Petviashvili's method to investigate the spectral stability of the periodic waves for the fractional Benjamin-Bona-Mahony equation, numerically. Some remarks concerning the orbital stability of periodic traveling waves are also presented.

1. INTRODUCTION.

In this paper, we show the spectral/orbital stability of zero mean periodic traveling wave solutions associated to the fractional Benjamin-Bona-Mahony (fBBM) type equation

$$u_t + u_x + uu_x + (D^\alpha u)_t = 0. \quad (1.1)$$

Here $u = u(x, t) : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is a 2π -periodic function at the x -variable and D^α represents the fractional differential operator defined as a Fourier multiplier by

$$\widehat{D^\alpha g}(\xi) = |\xi|^\alpha \widehat{g}(\xi), \quad \xi \in \mathbb{Z}, \quad (1.2)$$

where $0 < \alpha \leq 2$.

For the case $\alpha = 2$, we have the well known Benjamin-Bona-Mahony (BBM) equation arising as an improvement of the Korteweg-de Vries equation for modeling long surface gravity waves of small amplitude which propagate unidirectionally. It also describes the propagation of long waves with a balance of nonlinear and dissipative effects. In addition, the model appears in the analysis of the surface waves of long wavelength in liquids, hydromagnetic waves in cold plasma, acoustic-gravity waves in compressible fluids, and acoustic waves in harmonic crystals (see [7] and related references).

If $\alpha = 1$, we obtain the regularized Benjamin-Ono equation (rBO) which is a regularized version of the standard Benjamin-Ono equation. The rBO equation is a model for the time evolution of long-crested waves at the interface between two immiscible fluids. Some situations in which the equation

2000 *Mathematics Subject Classification.* 76B25, 35Q51, 35Q53.

Key words and phrases. BBM type-equations, existence and uniqueness of minimizers, spectral stability, orbital stability, Petviashvili's method.

* *Corresponding author*

Date: June, 2021.

is useful are the pycnocline in the deep ocean, and the two-layer system created by the inflow of fresh water from a river into the sea [25].

The fBBM equation (1.1) admits the following conserved quantities, smooth in their domains, as

$$E(u) = \frac{1}{2} \int_{-\pi}^{\pi} [(D^{\frac{\alpha}{2}} u)^2 - \frac{1}{3} u^3] dx \quad (1.3)$$

$$P(u) = \frac{1}{2} \int_{-\pi}^{\pi} [(D^{\frac{\alpha}{2}} u)^2 + u^2] dx, \quad (1.4)$$

and

$$M(u) = \int_{-\pi}^{\pi} u dx. \quad (1.5)$$

A traveling wave solution for (1.1) is of the form $u(x, t) = \phi(x - ct)$, where $\phi : \mathbb{T} \rightarrow \mathbb{R}$, with $\mathbb{T} = [-\pi, \pi]$, is a smooth 2π -periodic function and c represents the wave speed. Substituting this form into the fBBM equation (1.1), we obtain

$$cD^\alpha \phi + (c - 1)\phi - \frac{1}{2}\phi^2 + A = 0, \quad (1.6)$$

where A is a constant of integration. If we suppose that $\phi : \mathbb{T} \rightarrow \mathbb{R}$ is a periodic function with the zero mean property, then $A = A(c)$ is defined by

$$A(c) := \frac{1}{4\pi} \int_{-\pi}^{\pi} \phi^2(x) dx. \quad (1.7)$$

Combining (1.6) and (1.7), we see that the equation (1.6) can be expressed by the following boundary-value problem

$$cD^\alpha \phi + (c - 1)\phi = \frac{1}{2}\Pi_0 \phi^2, \quad \phi \in H_{per}^\alpha(\mathbb{T}), \quad (1.8)$$

where $\Pi_0 f := f - \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx$ is the projection operator restricted to the mean zero 2π -periodic functions.

The conserved quantities (1.3), (1.4) and (1.5) play an important role in our spectral stability analysis. In fact, they allow us to consider the augmented Lyapunov functional

$$G(u) = E(u) + (c - 1)P(u) + AM(u), \quad (1.9)$$

where $G'(\phi) = 0$, that is, ϕ is a critical point of G . Moreover, from (1.9) we obtain the Hessian operator around the wave ϕ , which is commonly called in the current literature, as linearized operator around ϕ and given by

$$\mathcal{L} := G''(\phi) = cD^\alpha + c - 1 - \phi. \quad (1.10)$$

It is easy to see that \mathcal{L} is a self-adjoint operator defined in $L_{per}^2(\mathbb{T})$ with dense domain $H_{per}^\alpha(\mathbb{T})$.

Next, we establish our linearized spectral problem for the fBBM equation. Indeed, substituting $u(x, t) = \phi(x - ct) + v(x - ct, t)$ into (1.1) and using that (1.8) is satisfied by ϕ , we obtain that v is a solution of the nonlinear equation

$$(\partial_t - c\partial_x)(v + D^\alpha v) + \partial_x(v + \phi v) + v\partial_x v = 0. \quad (1.11)$$

Replacing (1.11) by its linearization about ϕ we obtain the basic spectral stability problem

$$\partial_t(v + D^\alpha v) = \partial_x \mathcal{L}v \quad (1.12)$$

where \mathcal{L} is given by (1.10). Since ϕ depends on x and but not t , the equation (1.12) has a separation of variables of the form $v(x, t) = e^{\lambda t} \eta(x)$ with some $\lambda \in \mathbb{C}$ and $\eta : \mathbb{T} \rightarrow \mathbb{C}$. These arguments allow us to consider the spectral problem

$$\partial_x \mathcal{L} \eta = \lambda(1 + D^\alpha) \eta. \quad (1.13)$$

We can rewrite the spectral problem in (1.13) as

$$J\mathcal{L} \eta = \lambda \eta, \quad (1.14)$$

where $J := (1 + D^\alpha)^{-1} \partial_x$. Let us denote the spectrum of $J\mathcal{L}$ by $\sigma(J\mathcal{L})$. In a general setting, the periodic wave $\phi \in H_{per}^\alpha(\mathbb{T}) \cap X_0$ is said to be spectrally stable if $\sigma(J\mathcal{L}) \subset i\mathbb{R}$ in $L_{per}^2(\mathbb{T})$. Otherwise, that is, if $\sigma(J\mathcal{L})$ in $L_{per}^2(\mathbb{T})$ contains a point λ with $Re(\lambda) > 0$, the periodic wave ϕ is said to be spectrally unstable.

According to the classical spectral stability results as in [18], the problem given by (1.14) can not be handled in periodic context since $J = (1 + D^\alpha)^{-1} \partial_x$ is not a one-to-one operator. To overcome this difficulty, we can consider the linearization of (1.1) around ϕ restricted to the space of zero mean periodic function to obtain the modified problem (see [12])

$$J\mathcal{L}|_{X_0} \chi = \lambda \chi, \quad (1.15)$$

where $\mathcal{L}|_{X_0} = \mathcal{L} + \frac{1}{2\pi} \langle \phi, \cdot \rangle$ is a restriction of \mathcal{L} on the closed subspace X_0 of periodic functions with zero mean,

$$X_0 = \left\{ f \in L_{per}^2(\mathbb{T}) : \int_{-\pi}^{\pi} f(x) dx = 0 \right\}. \quad (1.16)$$

Clearly, operator $\mathcal{L}|_{X_0}$ is defined in the space X_0 with domain $H_{per}^\alpha(\mathbb{T}) \cap X_0$. The new spectral problem (1.15) allows to consider a new definition for the spectral stability restricted to the periodic space X_0 as:

Definition 1.1. *The periodic wave $\phi \in H_{per}^\alpha(\mathbb{T}) \cap X_0$ is said to be spectrally stable if $\sigma(J\mathcal{L}|_{X_0}) \subset i\mathbb{R}$ in X_0 . Otherwise, that is, if $\sigma(J\mathcal{L}|_{X_0})$ in X_0 contains a point λ with $Re(\lambda) > 0$, the periodic wave ϕ is said to be spectrally unstable.*

Remark 1.2. *In Definition 1.1, we consider a smooth zero mean periodic wave ϕ . However, it is possible to obtain the spectral stability/instability in the space X_0 using the same definition for non-zero mean periodic waves as reported, for instance, in [12] and [23].*

In [23], the existence of minimizers for the energy functional $V := P - E$ with fixed momentum P and mass M has been established. The periodic wave obtained by this minimization problem is smooth in terms of the independent parameters A and c in (1.6) and it has been determined that they are spectrally stable in the sense of Definition 1.1. The orbital stability of the periodic minimizers is then established by assuming that the 2-by-2 determinant $\{P, M\}_{A,c}$ is nonzero (see the precise definition of orbital stability in the last section of this paper). We also refer the reader to see [11]

for a different approach. However, in the referred work, a different minimization problem compared to [23] has been established and the authors obtained a different criterium for the orbital stability in the energy space.

For the cases $\alpha = 1$ and $\alpha = 2$, precise results of orbital stability for the equation (1.6) with $A = 0$ were determined in [5] combined that \mathcal{L} has only one negative eigenvalue which is simple and zero is a simple eigenvalue whose associated eigenfunction is ϕ' . In order to show that \mathcal{L} has only one negative eigenvalue which is simple and zero is a simple eigenvalue with associated eigenfunction ϕ' they have used the Fourier expansion of the explicit solutions together with the main theorem in [6]. In addition, the explicit solutions were also used to calculate the positiveness of $\frac{d}{dc}P(\phi)$.

Concerning the case $\alpha = 2$ in (1.1) with a general power nonlinearity as

$$u_t + u_x + (p + 1)u^p u_x - u_{xxt} = 0, \quad (1.17)$$

where $p \geq 1$ is an integer and traveling ϕ waves satisfying the equation

$$-c\phi'' + (c - 1)\phi - \phi^{p+1} = 0, \quad (1.18)$$

existence and stability results of small amplitude periodic waves near of the equilibrium solution $(c - 1)^{1/p}$ for $c > 1$ have been considered in [21]. These waves are spectrally stable for all speeds $c > 1$ when $1 \leq p \leq 2$ in both $L^2(\mathbb{R})$ and $C_b(\mathbb{R})$. Here, $C_b(\mathbb{R})$ stands the space of bounded uniformly continuous functions. For $p \geq 3$, there exists a critical speed $c_p = \frac{p(p+4)}{p^2+p+6}$, $1 < c_p < \frac{p}{p-3}$, where the waves are stable for $c \in \left(c_p, \frac{p}{p-3}\right)$ and spectrally unstable for $c \in (1, c_p) \cup \left(\frac{p}{p-3}, \infty\right)$ by considering the same perturbations. For perturbations which have the same period L as the wave, all small amplitude periodic waves are spectrally stable for all values of p .

For the model (1.1) posed on unbounded domains, the author in [3] showed the existence of solitary waves employing the arguments in [15]. He used the abstract approach in [18] ([28]) and determined the orbital (spectral) stability of solitary waves if:

i) $c > 1$ and $\alpha \in \left[\frac{1}{2}, 1\right)$,

ii) $c > c_0 := \frac{2 + \sqrt{2(3\alpha - 1)}}{6\alpha} > 1$ and $\alpha \in \left(\frac{1}{3}, \frac{1}{2}\right)$.

If $\alpha \in \left(\frac{1}{3}, \frac{1}{2}\right)$ and $c \in (1, c_0)$, the solitary wave is said to be spectrally unstable using the arguments in [28]. His proof relies on the scaling argument for the solitary wave ϕ which gives good features about the Hessian matrix $\frac{d}{dc}P(\phi)$. The same method in [3] can be used to establish the orbital (spectral) stability of solitary waves for the case $\alpha \in [1, 2]$ and $c > 1$.

We can cite some additional contributions concerning the spectral/orbital stability of periodic waves for related equations to (1.1) and other dispersive equations (see [1], [4], [6], [10], [12], [16], [17], [19], [24], [30], [31], and references therein). In some cases, the study of the stability was considered in relation to perturbations of the same period of the wave and other types of perturbations (in the latter case, we can consider: anti-periodic, bounded and so on).

The main results of our work can be presented according to the following theorem:

Theorem 1.3. *Let $\alpha \in (\frac{1}{3}, 2]$ and $\tau > 0$ be fixed. For every $c > \frac{1}{2}$, there exists an even and periodic single-lobe profile $\phi \in H_{per}^\alpha(\mathbb{T})$ which is solution of the constrained minimization problem,*

$$\inf_{u \in H_{per}^{\frac{\alpha}{2}}(\mathbb{T})} \left\{ \int_{-\pi}^{\pi} c(D^{\frac{\alpha}{2}}u)^2 + (c-1)u^2 dx; \int_{-\pi}^{\pi} u^3 dx = \tau, \int_{-\pi}^{\pi} u dx = 0 \right\}, \quad (1.19)$$

and a solution of (1.8). If $\ker(\mathcal{L}|_{X_0}) = [\phi']$ for all $c > \frac{1}{2}$, we obtain a smooth curve $c \in (\frac{1}{2}, +\infty) \mapsto \phi \in H_{per}^\alpha(\mathbb{T}) \cap X_0$ of even periodic waves with fixed period and furthermore, denoting $d := 1 + 2A(c) - c - cA'(c)$, we have,

- i) for the linearized operator \mathcal{L} : a simple negative eigenvalue if $d < 0$ and two negative eigenvalues if $d \geq 0$,*
- ii) zero is a simple eigenvalue of \mathcal{L} if and only if $d \neq 0$,*
- iii) ϕ is the unique solution of the problem (1.19) for the case $\alpha \in (\frac{1}{2}, 2]$,*
- iv) the periodic wave ϕ is spectrally stable if:*

- $d \neq 0$ and $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi) > 0$,
- $\alpha \in (\frac{1}{2}, 2]$ and $d \neq 0$ for all $c > \frac{1}{2}$,

- v) the periodic wave ϕ is orbitally stable if $\alpha \in (1, 2]$ and $d \neq 0$ for all $c > \frac{1}{2}$.*

The first two parts in Theorem 1.3 concern the existence of periodic minimizers for the problem (1.19) and the existence of smooth curve of solutions with fixed period. To prove the first part, we need to use the Poincaré-Wirtinger inequality to find the bifurcation point $c = \frac{1}{2}$ and compactness tools to establish that the periodic waves exist for all $c > \frac{1}{2}$. The second part is similar to [31, Lemma 3.8] and it can be determined by using the crucial hypothesis $\ker(\mathcal{L}|_{X_0}) = [\phi']$ and the implicit function theorem.

Assumption $\ker(\mathcal{L}|_{X_0}) = [\phi']$ is also important to give precise informations about items i)-v) in Theorem 1.3. In fact, item i) is determined since we show that the our periodic waves solve the minimization problem (1.19). Thus, we combine this information with convenient index formulas (see [33, Theorem 4.1]) to obtain that $n(\mathcal{L}) = 1$, if $d < 0$ and $n(\mathcal{L}) = 2$, if $d \geq 0$. Here, $n(\mathcal{L})$ stands for the number of negative eigenvalues of \mathcal{L} . The value of d is crucial in our analysis since it determines the existence of *fold points*, that is, values of c (and depending on α) such that $d = 0$. Folds points are related with the existence of additional elements in $\ker(\mathcal{L})$, besides ϕ' , and they give the exact value of c where the number of negative eigenvalues change. It is worth mentioning that they were first studied in [31] for the case of fractional Korteweg-de Vries (fKdV) equation.

Second item can be established since the operator \mathcal{L} in (1.10) satisfies an *Oscillation Theorem* for fractional linear operators with smooth periodic potentials according to [15], [23] and [31]. In fact, we obtain that $\dim(\ker(\mathcal{L})) \in \{1, 2\}$ and $\dim(\ker(\mathcal{L})) = 1$ if and only if $d \neq 0$.

Regarding item iii). Using the arguments in [15], our work establishes sufficient conditions for the *uniqueness of periodic minimizers* associated to the problem (1.19). In what follows, let $\alpha_0 \in (\frac{1}{2}, 2]$ be fixed. Assume that $(\phi_0, c_0) \in V \times (\frac{1}{2}, +\infty)$ with ϕ_0 being a non-zero solution of (1.8) and satisfying assumption (3.24) with $\alpha = \alpha_0$ and $c = c_0$. The implicit function theorem guarantees the existence

of a $\delta > 0$ and an C^1 -map $\alpha \in I = [\alpha_0, \alpha_0 + \delta) \mapsto (\phi_\alpha, c_\alpha)$ such that (ϕ_α, c_α) is the unique local solution of (1.8) in a convenient neighbourhood around (ϕ_0, c_0) . In addition, it is possible to obtain for all $\alpha \in I$ that

$$\int_{-\pi}^{\pi} \phi_\alpha^3 dx = \int_{-\pi}^{\pi} \phi_0^3 dx. \quad (1.20)$$

Using compactness arguments, we prove that the maximal branch (ϕ_α, c_α) and equality (1.20) extends to the interval $[\alpha_0, 2)$. Next, since it is well known that for $\alpha = 2$ and a fixed $c > \frac{1}{2}$ the solution with dnoidal profile (see (3.42)) is unique, we obtain the uniqueness of minimizers for the problem (1.19) using the similar arguments as in [15, Theorem 2.4].

We determine item iv) employing again the index formula to get

$$n(\mathcal{L}|_{\{1, (D^{\alpha+1}\phi)^\perp\}}) = n(\mathcal{L}) - n_0 - z_0,$$

where n_0 is the number of negative eigenvalues determined by the Hessian symmetric matrix formed by momentum and mass and z_0 indicates the dimension of the kernel of the same Hessian matrix. To simplify the analysis, n_0 and z_0 can be both precisely determined by analysing the following quantity for the case $d \neq 0$ (see Section 5).

$$\det S(0) := \frac{4\pi^2 d^{-1}}{c} \left(A'(c) + \frac{1}{\pi} \mathcal{B}_c(\phi) \right). \quad (1.21)$$

To obtain the precise statements for this item, we need to discuss the existence (and spectral stability) of periodic small amplitude waves associated with the equation (1.1) for the case $\alpha \in (0, 2]$. For a small amplitude parameter a , we put forward the explicit solutions of (1.8) as

$$\phi(x) = a \cos(x) + \frac{a^2}{2(2^\alpha - 1)} \cos(2x) + O(a^3). \quad (1.22)$$

The wave speed c and parameter A are given respectively by

$$c = \frac{1}{2} + \frac{a^2}{4(2^\alpha + 1)(2^\alpha - 1)} + O(a^4), \quad \text{and} \quad A(c) = \frac{a^2}{4} + O(a^4). \quad (1.23)$$

Orbital/spectral stability and related topics of small amplitude periodic waves associated to several evolution models have been exhaustively studied (see [20], [22], [24], [27], [31], and references therein). In our paper, we show the existence of two fold points: $\alpha = \tilde{\alpha} \approx 0.2924$ and $\alpha = \frac{1}{2}$. We prove that $n(\mathcal{L}) = 1$, if $\alpha \in (0, \tilde{\alpha}) \cup (\frac{1}{2}, 2]$ and $n(\mathcal{L}) = 2$, if $\alpha \in (\tilde{\alpha}, \frac{1}{2})$. To determine $A'(c)$ in (1.21), we can use both expressions in (1.23) to get

$$A'(c) = (2^\alpha + 1)(2^\alpha - 1) + O(a^3). \quad (1.24)$$

Since $A'(c) > 0$ for $\alpha \in (0, 2]$, it follows that n_0 and z_0 are determined only by the sign of d . In fact, if $d < 0$, one has $n_0 = 1$ and $z_0 = 0$, while $d > 0$ gives us that $n_0 = 2$ and $z_0 = 0$. In both cases, we obtain the spectral stability of the small amplitude periodic wave ϕ (1.22). The degenerate case $d = 0$ is also established and the wave is said to be spectrally stable. For the case $\alpha = 2$ in (1.1), our results are compatible with those ones in [21] when $p = 1$ in equation (1.17) but, in our case, the small amplitude periodic waves are near of the equilibrium solution $\phi = 0$.

Using the spectral information obtained by the small amplitude periodic waves, we establish that the number of negative eigenvalues of the linearized operator \mathcal{L} is always equal to one if and only if $\dim(\ker(\mathcal{L})) = 1$ for all $c > \frac{1}{2}$. In addition, since $d < 0$ for the periodic wave obtained by the problem (1.19), we conclude $A(c) > c - \frac{1}{2}$ and this bound allows us to deduce that $\det S(0)$ in (1.21) is always negative. This fact gives us the spectral stability and moreover, it recovers the same result as in [3] without using scaling argument. In addition, we obtain periodic travelling wave solutions of the equation (1.1) numerically by using a Petviashvili's iteration method. The numerical results also confirm the analytical results stating that the periodic wave is spectrally stable for $\alpha \in (\frac{1}{2}, 2]$.

Next, we discuss the case $\alpha \in (\frac{1}{3}, \frac{1}{2}]$. In fact, numerical experiments will be used to decide the exact sign of the quantities d and $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi)$ in (1.21) in terms of $c > \frac{1}{2}$. Our numerical results point out that there exists a critical wave speed $c^* \in (\frac{1}{2}, +\infty)$ at which the sign of d changes. For $c \in (c^*, +\infty)$, we observe numerically that $d < 0$ and $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi) > 0$, therefore the periodic wave is spectrally stable. On the other hand, for $c \in (\frac{1}{2}, c^*)$, we have $d > 0$ and $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi) > 0$. Thus $\det S(0) > 0$ which implies $n_0 = 0$ or $n_0 = 2$. Since the first entry of the matrix $S(0)$ is negative, we obtain $n_0 = 2$ and the spectral stability of periodic wave is observed numerically for $c \in (\frac{1}{2}, c^*)$.

In addition to the spectral stability, we prove the orbital stability for $\alpha \in (1, 2]$ by employing an adaptation of the recent arguments in [11] and new ingredients. Indeed, in our paper we are not considering, for the orbital stability, the old method which minimizes the energy E in (1.3) with fixed momentum P in (1.4) and mass M in (1.5). When this kind of approach is considered, we need to prove, besides good spectral properties for the linearized operator \mathcal{L} in (1.10), the positiveness of the Hessian matrix $\frac{d}{dc}P(\phi)$ (see [1], [5], [6], [11], [18], and references therein). As far as we can see, even in the case $\alpha = 2$, when explicit solutions are known in terms of the Jacobi elliptic functions, the calculation of the derivative of $P(\phi)$ in terms of c becomes a hard task (see [5]). Instead of this, we use the conservation law $V(u) = P(u) - E(u)$ as a constrained manifold associated to the augmented Lyapunov functional G in (1.9). This consideration sheds new light in the stability theory since we obtain our results without using any additional information of the wave, only the basic bound $A(c) > c - \frac{1}{2}$ determined to prove the spectral stability. This fact enables us to conclude item v) in Theorem 1.3.

Our paper is organized as follows. In Section 2, we establish local and global well posedness results for the Cauchy problem associated to the equation (1.1). In Section 3 we show the existence of periodic minimizers for the problem (1.19). and a precise information about the number of negative eigenvalues of (1.10). Also in this section, we show the uniqueness of periodic minimizers for the problem (1.19) in the interval $\alpha \in (\frac{1}{2}, 2]$. In Section 4, we present our spectral stability result for the waves obtained in Section 3. Section 5 is devoted to the numerical investigation of the spectral stability of periodic waves. Finally, in Section 6 we give some important remarks concerning the orbital stability of periodic waves.

2. WELL-POSEDNESS RESULTS

In this section, we present a brief comment concerning the global well-posedness for the Cauchy problem associated to the equation (1.1)

$$\begin{cases} u_t + u_x + uu_x + (D^\alpha u)_t = 0, \\ u(x, 0) = u_0(x). \end{cases} \quad (2.1)$$

In the whole real line, the best result of local well posedness for the Cauchy problem (2.1) has been determined in [29] for initial data $u_0 \in H^s(\mathbb{R})$, $s > \frac{3}{2} - \frac{3}{8}\alpha$ and $\alpha \in (0, 1)$. For initial data $u_0 \in H_{per}^{\alpha/2}(\mathbb{T})$ and $\alpha > 1$, we are going to use fixed point arguments applied to the representation of (2.1) in an integral form as

$$u(x, t) = u_0(x) + \int_0^t \left[\mathcal{K} * \left(u + \frac{u^2}{2} \right) \right] (x, s) ds, \quad (2.2)$$

where \mathcal{K} is defined using the periodic Fourier transform

$$\widehat{\mathcal{K}v}(k) = \frac{-ik}{1 + |k|^\alpha} \widehat{v}(k).$$

Proposition 2.1. *Let $\alpha > 1$ be fixed. For each $s \geq \frac{\alpha}{2}$ and $u_0 \in H_{per}^s(\mathbb{T})$, there exist $T > 0$ and a unique solution u of (2.1) such that $u \in C([-T, T]; H_{per}^s(\mathbb{T}))$. Moreover, for all $T^* < T$, there exists a neighborhood V of u_0 in $H_{per}^s(\mathbb{T})$ such that the data-solution map*

$$w_0 \in V \subset H_{per}^s(\mathbb{T}) \mapsto w \in C([-T^*, T^*]; H_{per}^s(\mathbb{T})),$$

is continuous.

Proof. The proof of this result is classical because of the integral representation in (2.2) (for the cases $\alpha = 1, 2$, see [5]). Define, $Y = C([-T, T]; H_{per}^s(\mathbb{T}))$ with the norm $\|w\|_Y = \sup_{t \in [-T, T]} \|w(t)\|_{H_{per}^s}$. Consider the map $\Upsilon : B_r \rightarrow B_r$ given by

$$\Upsilon(u) = u_0(x) + \int_0^t \mathcal{K} * \left(u + \frac{u^2}{2} \right) ds, \quad (2.3)$$

where $r > 0$ will be chosen later and $B_r = \{w \in Y; \|w\|_Y \leq r\}$. We show that Υ is well defined, in the sense that $\Upsilon(u) \in B_r$ for $u \in B_r$, and Υ is a strict contraction.

In fact, consider $v \in H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$. Since $\alpha > 1$, we have

$$\|\mathcal{K} * v\|_{H_{per}^s}^2 \leq 2\pi \sum_{k \in \mathbb{Z}} (1 + k^2)^s |\widehat{v}(k)|^2 = \|v\|_{H_{per}^s}^2. \quad (2.4)$$

In addition, if $v \in L_{per}^2(\mathbb{T})$ we obtain the basic smoothing effect given by

$$\|\mathcal{K} * v\|_{H_{per}^{\frac{\alpha}{2}}}^2 \leq 2\pi C_1 \sum_{k \in \mathbb{Z}} |\widehat{v}(k)|^2 = C_1 \|v\|_{L_{per}^2}^2, \quad (2.5)$$

where $C_1 > 0$ does not depend on v .

On the other hand $\alpha > 1$ implies that $H_{per}^s(\mathbb{T})$ is a Banach algebra for all $s \geq \frac{\alpha}{2}$, and thus

$$\|u^2\|_{H_{per}^s} \leq C_2 \|u\|_{H_{per}^s}^2, \quad (2.6)$$

where $C_2 > 0$ does not depend on u .

Let $u, v \in Y$ be fixed. Gathering (2.4), (2.6) and the definition of Υ , we obtain

$$\|\Upsilon u\|_Y \leq \|u_0\|_{H_{per}^s} + T (\|u\|_Y + C_3 \|u\|_Y^2), \quad (2.7)$$

where $C_3 > 0$ is positive constant depending on $C_2 > 0$.

Next,

$$\|\mathcal{K} * (u^2 - v^2)\|_{H_{per}^s} \leq (\|u\|_Y + \|v\|_Y) \|u - v\|_Y. \quad (2.8)$$

Therefore, by using a similar procedure as in (2.7) we obtain that

$$\|\Upsilon u - \Upsilon v\|_Y \leq T (1 + \|u\|_Y + \|v\|_Y) \|u - v\|_Y. \quad (2.9)$$

By (2.7), (2.9) and the fact that $u, v \in Y$, we have

$$\|\Upsilon u\|_Y \leq \|u_0\|_{H_{per}^s} + T (r + C_3 r^2) \quad \text{and} \quad \|\Upsilon u - \Upsilon v\|_Y \leq T (1 + 2r) \|u - v\|_Y. \quad (2.10)$$

Using (2.10), we can choose $r = 2\|u_0\|_{H_{per}^s}$ and $T = \frac{1}{2(1+C_4r)}$, where $C_4 = \max\{2, C_3\} > 0$, to obtain

$$\|\Upsilon u\|_Y \leq r \quad \text{and} \quad \|\Upsilon u - \Upsilon v\|_Y \leq \frac{1}{2} \|u - v\|_Y. \quad (2.11)$$

Inequalities in (2.11) give us that $\Upsilon : B_r \rightarrow B_r$ is well defined and a strict contraction. From the Banach Fixed Point Theorem, we obtain the existence of a unique $u \in B_r$ such that $\Upsilon u(t) = u(t)$ for all $t \in [-T, T]$. The uniqueness in the whole space and the continuous dependence are determined by a direct application of standard arguments. \square

Proposition 2.2. *Let $\alpha > 1$ be fixed. The quantities E , P and M defined in (1.3), (1.4) and (1.5), respectively are conservation laws.*

Proof. The proof of this result is standard and we skip the details. \square

Propositions 2.1 and 2.2 give us the following result.

Proposition 2.1. *Let $\alpha > 1$ be fixed. For each $u_0 \in H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$, the Cauchy problem (2.1) is globally well posed in $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$ with $u \in C(\mathbb{R}; H_{per}^{\frac{\alpha}{2}}(\mathbb{T}))$.* \square

3. EXISTENCE AND UNIQUENESS OF MINIMIZERS - SPECTRAL PROPERTIES.

3.1. Existence of Periodic Minimizers. In this subsection, we first give a sufficient condition for the existence of periodic waves for the equation (1.6) by showing the existence of minimizers associated to a convenient variational problem. After that, we use the minimizers to obtain good spectral properties for the linearized operator \mathcal{L} in (1.10).

Before starting, we need an important definition which characterizes the solutions of the equation (1.8).

Definition 3.1. *We say that the periodic traveling wave solutions satisfying the equation (1.8) has a single-lobe profile ϕ if there exist only one maximum and minimum of ϕ on \mathbb{T} . Without the loss of generality, the maximum of ϕ is placed at $x = 0$.*

For a fixed $\tau > 0$, let us consider the set

$$Y_0 = \left\{ u \in H_{per}^{\frac{\alpha}{2}}(\mathbb{T}); \int_{-\pi}^{\pi} u^3 = \tau, \int_{-\pi}^{\pi} u = 0 \right\}. \quad (3.1)$$

Our goal is to find a minimizer of the constrained minimization problem

$$q_c = \inf_{u \in Y_0} \mathcal{B}_c(u), \quad (3.2)$$

where

$$\mathcal{B}_c(u) = \frac{1}{2} \int_{-\pi}^{\pi} c(D^{\frac{\alpha}{2}}u)^2 + (c-1)u^2 dx. \quad (3.3)$$

The next lemma establishes the first part of Theorem 1.3.

Lemma 3.2. *Let $\alpha > 1/3$ and $c > \frac{1}{2}$ be fixed. The minimization problem (3.2) has at least one solution, that is, there exists a $\phi \in Y_0$ satisfying*

$$\mathcal{B}_c(\phi) = \inf_{u \in Y_0} \mathcal{B}_c(u). \quad (3.4)$$

If $\alpha \in (\frac{1}{3}, 2]$, the periodic minimizer of (3.2) has an even, single-lobe profile in the sense of Definition 3.1.

Proof. Let $c > \frac{1}{2}$ be fixed. By Poincaré-Wirtinger inequality, we see that

$$2\mathcal{B}_c(u) = \int_{-\pi}^{\pi} c(D^{\frac{\alpha}{2}}u)^2 + (c-1)u^2 dx \geq (2c-1) \int_{-\pi}^{\pi} u^2 dx \geq 0. \quad (3.5)$$

Combining (3.5) with the Garding inequality, we obtain that \mathcal{B}_c is an equivalent norm in $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$ yielding $q_c \geq 0$. Since \mathcal{B}_c is a smooth functional in $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$, let $\{u_n\}$ be a minimizing sequence for (3.2), that is, a sequence in Y_0 satisfying

$$\mathcal{B}_c(u_n) \rightarrow \inf_{u \in Y_0} \mathcal{B}_c(u), \quad \text{as } n \rightarrow +\infty.$$

There exist positive constants M_0 and M_1 depending on c satisfying

$$M_0 \|u_n\|_{H_{per}^{\frac{\alpha}{2}}}^2 \leq \mathcal{B}_c(u_n) \leq M_1 \|u_n\|_{H_{per}^{\frac{\alpha}{2}}}^2, \quad \forall n \in \mathbb{N},$$

that is, $\{u_n\}$ is bounded in $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$. Thus, there is $\phi \in H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$ such that, up to a subsequence,

$$u_n \rightharpoonup \phi \text{ weakly in } H_{per}^{\frac{\alpha}{2}}(\mathbb{T}), \quad \text{as } n \rightarrow +\infty.$$

On other hand, since $\alpha > 1/3$ one sees that the energy space $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$ is compactly embedded in $L_{per}^3(\mathbb{T})$ (see [2, Theorem 4.2]), and $u_n \rightarrow \phi$ in $L_{per}^3(\mathbb{T})$, as $n \rightarrow +\infty$. In addition, using the fact

$$\left| \int_{-\pi}^{\pi} (u_n^3 - \phi^3) dx \right| \leq \|u_n - \phi\|_{L_{per}^3} + 3\|u_n - \phi\|_{L_{per}^3} \|\phi\|_{L_{per}^3} \|u_n\|_{L_{per}^3},$$

one has that $\int_{-\pi}^{\pi} \phi^3 dx = \tau$. A similar argument as above and using the fact that $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$ is compactly embedded in $L_{per}^1(\mathbb{T})$, give us $\int_{-\pi}^{\pi} \phi dx = 0$.

Moreover, since the weak lower semi-continuity of \mathcal{B}_c , we have $\mathcal{B}_c(\phi) \leq \liminf_{n \rightarrow +\infty} \mathcal{B}_c(u_n) = q_c$.

The symmetric rearrangements $\phi^{\#} \in Y_0$ associated to the solution ϕ^1 leave the L_{per}^2 -norm of $\phi^{\#}$ and ϕ invariants and $\int_{-\pi}^{\pi} (D^{\alpha/2} \phi^{\#})^2 dx$ does not increase in comparison with $\int_{-\pi}^{\pi} (D^{\alpha/2} \phi)^2 dx$ thanks to the fractional Polya–Szegő inequality (for further details and similar applications, see [10, Lemma A.1], [23, Proposition 2.1] and [31, Theorem 2.1]). Therefore, for $c > \frac{1}{2}$ we have $\mathcal{B}_c(\phi^{\#}) = q_c$. Invoking the original notation ϕ instead of $\phi^{\#}$ to simplify the comprehension of the reader, we see that the minimizer $\phi \in Y_0$ of $\mathcal{B}_c(u)$ must decrease away symmetrically from the maximum point. Using the translational invariance, the maximum point can be placed at $x = 0$, which yields an even single-lobe profile for ϕ . \square

From Lemma (3.2) and Lagranges’s Multiplier Theorem, there exists C_1 and C_2 such that

$$cD^{\alpha} \phi + (c - 1)\phi = C_1 \phi^2 + C_2. \quad (3.6)$$

We see that ϕ is a nontrivial single-lobe because $\phi \in Y_0$. Since $\int_{-\pi}^{\pi} \phi dx = 0$, we deduce from (3.6) that $C_2 = -\frac{C_1}{2\pi} \int_{-\pi}^{\pi} \phi^2 dx$. In addition, multiplying equation (3.6) by ϕ and integrating the result over $[0, L]$, we obtain by the Poincaré–Wirtinger inequality, the fact $c > \frac{1}{2}$ and since $\tau > 0$ that $C_1 > 0$. By the homogeneity of C_1 ($\phi^2 - \frac{1}{2\pi} \int_{-\pi}^{\pi} \phi^2 dx$), we see that C_1 can be chosen as $C_1 = \frac{1}{2}$. Indeed, for all $s > 0$, we obtain

$$\begin{aligned} \mathcal{B}_c(s\phi) &= s^2 \mathcal{B}_c(\phi) \\ &= \inf \left\{ \mathcal{B}_c(su); u \in H_{per}^{\frac{\alpha}{2}}(\mathbb{T}), \int_{-\pi}^{\pi} u^3 dx = \tau, \int_{-\pi}^{\pi} u dx = 0 \right\} \\ &= \inf \left\{ \mathcal{B}_c(u); u \in H_{per}^{\frac{\alpha}{2}}(\mathbb{T}), \int_{-\pi}^{\pi} u^3 dx = s^3 \tau, \int_{-\pi}^{\pi} u dx = 0 \right\}. \end{aligned} \quad (3.7)$$

Thus, if ϕ solves (3.2) we have that $\psi = s\phi$ is a solution of the minimization problem (3.7). As above and by Lagrange’s Multiplier Theorem, there exists $C_3 > 0$ and $C_4 = -\frac{C_3}{2\pi} \int_{-\pi}^{\pi} \psi^2 dx$ such that

$$s(cD^{\alpha} \phi + (c - 1)\phi) = s^2 C_3 \left(\phi^2 - \frac{1}{2\pi} \int_{-\pi}^{\pi} \phi^2 dx \right). \quad (3.8)$$

¹We can take $\phi^{\#}$ as a symmetric rearrangement in Y_0 because it leaves $\int_{-\pi}^{\pi} u^3 dx$ and $\int_{-\pi}^{\pi} u dx$ invariant (see [23, Proposition 2.1])

Since $s > 0$ is arbitrary and ϕ is non-trivial, we can choose $s = \frac{1}{2C_3}$ to obtain that ϕ solves the equation

$$cD^\alpha\phi + (c-1)\phi - \frac{1}{2}\phi^2 + A = 0, \quad (3.9)$$

so that C_1 can be chosen as $C_1 = \frac{1}{2}$. Moreover, by a standard bootstrap argument (see [11]), we obtain that ϕ is smooth.

Remark 3.3. *Let $\alpha \in (\frac{1}{3}, 2]$ and $L > 0$ be fixed. The periodic wave ϕ obtained in Lemma 3.2 can be considered with a general period L instead of the normalized period $L = 2\pi$. In the general case, the wave speed needs to satisfy (by using Poincaré-Wirtinger inequality for general periods) the basic bound $c > \frac{1}{1+(\frac{2\pi}{L})^\alpha}$ for the existence of periodic waves in the space $X_{0,L} = \{f \in L^2_{per}(\mathbb{T}_L); \int_{-L/2}^{L/2} f dx = 0\}$, where $\mathbb{T}_L = [-L/2, L/2]$. All results in this section, Sections 4 and 6 can be obtained with a general period by replacing the normalized bifurcation point $c = \frac{1}{2}$ by $c = \frac{1}{1+(\frac{2\pi}{L})^\alpha}$ and slight modifications in the arguments when necessary. In Section 5, where we present our numerical experiments, it makes necessary to fix the period in order to obtain the plots which give us, for example, the behaviour of ϕ , d , $A'(c)$ and etc. However, the restriction to consider normalized periodic solutions does not affect the generality of the proposed results in a general context.*

Next, let us denote $\tilde{\mathcal{L}} := \frac{1}{c}\mathcal{L} = D^\alpha + (1 - \frac{1}{c}) - \frac{1}{c}\phi$ the linearized and self-adjoint operator around the periodic wave ϕ and suppose that $\ker(\tilde{\mathcal{L}}|_{X_0}) = [\phi']$. Thus, a direct application of the implicit function theorem as in [31, Lemma 3.8] gives us that $c \in (\frac{1}{2}, +\infty) \mapsto \phi \in H^\infty_{per}(\mathbb{T})$ defines a smooth curve of periodic waves all of them with the same period 2π . Thus, one has that $\frac{d}{dc}\phi \in D(\tilde{\mathcal{L}}) = H^\alpha_{per}(\mathbb{T})$,

$$\tilde{\mathcal{L}}\left(\frac{d}{dc}\phi\right) = \frac{1}{c^2}\left(A(c) - \phi - \frac{1}{2}\phi^2 - cA'(c)\right), \quad \tilde{\mathcal{L}}\left(\frac{1}{c}\right) = \frac{1}{c^2}(c-1-\phi) \quad (3.10)$$

and

$$\tilde{\mathcal{L}}\left(\frac{1}{c}\phi\right) = -\frac{1}{c^2}\left(\frac{1}{2}\phi^2 + A(c)\right). \quad (3.11)$$

Equations in (3.10) and (3.11) give us an important relation

$$c^2\tilde{\mathcal{L}}\left(\frac{d}{dc}\phi - \frac{1}{c} - \frac{1}{c}\phi\right) = (1 + 2A(c) - c - cA'(c)) = d. \quad (3.12)$$

Since $\tilde{\mathcal{L}}\phi = -\frac{1}{c}\left(\frac{1}{2}\phi^2 + A(c)\right)$, we have $\langle \tilde{\mathcal{L}}\phi, \phi \rangle < 0$ and we deduce that $\tilde{\mathcal{L}}$ has at least one negative eigenvalue. Next result establishes items i) and ii) of Theorem 1.3 by giving the precise behaviour of the first eigenvalues associated with the linearized operator $\tilde{\mathcal{L}}$.

Proposition 3.1. *Let $\alpha \in (\frac{1}{3}, 2]$ be fixed and assume that $\ker(\tilde{\mathcal{L}}|_{X_0}) = [\phi']$. For $c > \frac{1}{2}$ consider $d = 1 + 2A(c) - c - cA'(c)$. The linearized operator $\tilde{\mathcal{L}} = D^\alpha + (1 - \frac{1}{c}) - \frac{1}{c}\phi$ around the periodic*

solution ϕ satisfies

$$n(\tilde{\mathcal{L}}) = \begin{cases} 1, & \text{if } d < 0, \\ 2, & \text{if } d \geq 0 \end{cases} \quad (3.13)$$

and

$$z(\tilde{\mathcal{L}}) = \begin{cases} 1, & \text{if } d \neq 0, \\ 2, & \text{if } d = 0. \end{cases} \quad (3.14)$$

Proof. In view of (3.2), since ϕ locally minimizes \mathcal{B}_c , we have $\tilde{\mathcal{L}}|_{\{1, \phi^2\}^\perp} \geq 0$, that is, $n(\tilde{\mathcal{L}}|_{\{1, \phi^2\}^\perp}) = 0$ and $n(\tilde{\mathcal{L}}) \in \{1, 2\}$, since $\langle \tilde{\mathcal{L}}\phi, \phi \rangle = -2\mathcal{B}_c(\phi) < 0$.

For $\lambda \notin \sigma(\tilde{\mathcal{L}})$, let $R(\lambda)$ be the following matrix given by

$$R(\lambda) := \begin{bmatrix} \langle (\tilde{\mathcal{L}} - \lambda)^{-1}1, 1 \rangle & \langle (\tilde{\mathcal{L}} - \lambda)^{-1}\phi^2, 1 \rangle \\ \langle (\tilde{\mathcal{L}} - \lambda)^{-1}\phi^2, 1 \rangle & \langle (\tilde{\mathcal{L}} - \lambda)^{-1}\phi^2, \phi^2 \rangle \end{bmatrix}. \quad (3.15)$$

For $d \neq 0$, we compute $R(\lambda)$ at $\lambda = 0$. In fact,

$$\langle \tilde{\mathcal{L}}^{-1}\phi^2, \phi^2 \rangle = -4c^2 \int_{-\pi}^{\pi} (D^{\frac{\alpha}{2}}\phi)^2 dx - 8\pi A(c) \left(\frac{A(c)}{c} d^{-1} + 2(c^2 - c) \right) \quad (3.16)$$

$$\langle \tilde{\mathcal{L}}^{-1}1, \phi^2 \rangle = \langle \tilde{\mathcal{L}}^{-1}\phi^2, 1 \rangle = \frac{4\pi A(c)}{c} d^{-1}, \quad (3.17)$$

and

$$\langle \tilde{\mathcal{L}}^{-1}1, 1 \rangle = -\frac{2\pi}{c} d^{-1}. \quad (3.18)$$

In (3.16), (3.17) and (3.18), $\tilde{\mathcal{L}}^{-1}\phi^2$ and $\tilde{\mathcal{L}}^{-1}1$ are calculated by combining (3.10) and (3.11) to obtain

$$\tilde{\mathcal{L}} \left(d^{-1} \left(\frac{d}{dc}\phi - \frac{1}{c} - \frac{1}{c}\phi \right) \right) = 1, \quad (3.19)$$

and

$$\tilde{\mathcal{L}} \left(-2A(c) \left(d^{-1} \left(\frac{d}{dc}\phi - \frac{1}{c} - \frac{1}{c}\phi \right) \right) - 2c\phi \right) = \phi^2. \quad (3.20)$$

The determinant of $R(0)$ is now given by

$$\det R(0) = 8\pi d^{-1} \left(c \int_{-\pi}^{\pi} (D^{\frac{\alpha}{2}}\phi)^2 dx + (c-1) \int_{-\pi}^{\pi} \phi^2 dx \right) = 16\pi d^{-1} \mathcal{B}_c(\phi). \quad (3.21)$$

Since $c > \frac{1}{2}$, we obtain that $\mathcal{B}_c(\phi) > 0$ and the sign of $R(0)$ is obtained only by d^{-1} .

On the other hand, by [33, Theorem 4.1] we have the following identities

$$n(\tilde{\mathcal{L}}|_{\{1, \phi^2\}^\perp}) = n(\tilde{\mathcal{L}}) - n_0 - z_0, \quad (3.22)$$

and

$$z(\tilde{\mathcal{L}}|_{\{1, \phi^2\}^\perp}) = z(\tilde{\mathcal{L}}) + z_0 - z_\infty \quad (3.23)$$

where n_0 is the number of negative eigenvalues of $R(0)$, z_0 denotes the dimension of the kernel of $R(0)$ and z_∞ corresponds the number of diverging eigenvalues, that is, $z_\infty = 1$ if $d = 0$, and $z_\infty = 0$ otherwise. We then deduce the result (3.22) from the equation (3.13).

Finally, since ϕ is a single-lobe solution, we see that $\phi' \in \ker(\tilde{\mathcal{L}})$ has only two zeroes in the interval $[-\pi, \pi)$. This means, from the Oscillation Theorem in [23], that $z(\tilde{\mathcal{L}}) \in \{1, 2\}$. If $d \neq 0$, we deduce from the second equality in (3.10), (3.19) and (3.20) that $\{1, \phi, \phi^2\} \subset \text{range}(\tilde{\mathcal{L}})$, so that $z(\tilde{\mathcal{L}}) = 1$ by using [23, Proposition 3.1]. If $d = 0$, we have $z_\infty = 1$ and since $\phi' \in \{1, \phi^2\}^\perp$, one has $z(\tilde{\mathcal{L}}|_{\{1, \phi^2\}^\perp}) \geq 1$. Thus, $z(\tilde{\mathcal{L}}) = 2$ as requested. \square

Remark 3.1. Let $\alpha \in (\frac{1}{3}, 2]$ be fixed. If $\ker(\mathcal{L}|_{X_0}) = [\phi']$, by [31, Lemma 3.8] we obtain the existence of a smooth curve $\omega \in I \mapsto \psi_\omega \in H_{per}^\infty(\mathbb{T}) \cap X_{0,e}$ of periodic waves for the equation (1.8), where $I \subset (\frac{1}{2}, +\infty)$ is a convenient open interval containing c . Important to mention that we can not assure if ψ_ω solves the minimization problem (3.2), neither that it is a single-lobe, except at $\omega = c$, where $\psi_\omega = \phi$.

3.2. Uniqueness of periodic minimizers. We give sufficient conditions to determine a result of uniqueness for the periodic minimizer ϕ obtained in Lemma 3.2. To do so, we will follow similar arguments in [15, Section 5].

Let us consider $\mathcal{L}|_{X_0} = \mathcal{L} + \frac{1}{2\pi}\langle \phi, \cdot \rangle$. Throughout this section we assume that

$$z(\mathcal{L}|_{X_0}) = 1, \quad (3.24)$$

for every $c > \frac{1}{2}$ and $\alpha \in (\frac{1}{2}, 2]$. Using that $\phi \in X_0$ we obtain $\langle \mathcal{L}|_{X_0}\phi, \phi \rangle = \langle \mathcal{L}\phi, \phi \rangle = -\frac{1}{2}\int_{-\pi}^{\pi} \phi^3 dx = -2\mathcal{B}_c(\phi) < 0$, so that $n(\mathcal{L}|_{X_0}) \geq 1$. By [31, Corollary 4.5], we can obtain automatically the condition (3.24) by assuming the additional hypothesis $n(\mathcal{L}) = 1$ for every $c > \frac{1}{2}$ and $\alpha \in (\frac{1}{2}, 2]$. Equality in (3.24) enables us to conclude an important property for $\mathcal{L}|_{X_0}$ as

$$\ker(\mathcal{L}|_{X_0}) = [\phi']. \quad (3.25)$$

To simplify the notation, we define the real Banach space

$$V = \{f \in L_{per}^3(\mathbb{T}) \cap X_0; f \text{ is even and real-valued}\} \quad (3.26)$$

whose norm, since $L_{per}^3(\mathbb{T}) \hookrightarrow L_{per}^2(\mathbb{T})$, is given by $\|f\|_V := \|f\|_{L_{per}^3(\mathbb{T})}$. For a fixed $\alpha_0 \in (\frac{1}{2}, 2]$, our first goal is to construct a local branch of solutions $(\phi_\alpha, c_\alpha) \in V \times (\frac{1}{2}, +\infty)$ of (1.8) parameterized by the index α in some interval $[\alpha_0, \alpha_0 + \delta)$, for some $\delta > 0$ small enough.

Proposition 3.2. Let $\alpha_0 \in (\frac{1}{2}, 2]$. Suppose that $(\phi_0, c_0) \in V \times (\frac{1}{2}, +\infty)$ with ϕ_0 being a nonzero solution of (1.8) and satisfying assumption (3.24) with $\alpha = \alpha_0$ and $c = c_0$. Then, for some $\delta > 0$, there exists a C^1 -map $\alpha \in I \mapsto (\phi_\alpha, c_\alpha) \in V \times (\frac{1}{2}, +\infty)$, defined in the interval $I = [\alpha_0, \alpha_0 + \delta)$, such that the following holds:

- (i) (ϕ_α, c_α) solves equation (1.8) with $c = c_\alpha$, for all $\alpha \in I$ and the pair (ϕ_α, c_α) satisfies (3.24).

- (ii) *There exists $\varepsilon > 0$ such that (ϕ_α, c_α) is the unique solution of (1.8) for $\alpha \in I$ in the neighborhood $\{(\phi, c) \in V \times (\frac{1}{2}, +\infty); \|\phi - \phi_0\|_V + |c - c_0| < \varepsilon\}$.*
- (iii) *For all $\alpha \in I$, we have $\int_{-\pi}^{\pi} \phi_\alpha^3 dx = \int_{-\pi}^{\pi} \phi_0^3 dx$.*

Proof. The proof of this result relies on the implicit function theorem. Since $c > \frac{1}{2}$, we obtain that (1.8) can be written as $\phi = (cD^\alpha + (c - 1))^{-1} \frac{1}{2} \Pi_0 \phi^2$. For some $\delta > 0$ to be chosen later, let us define the mapping

$$\mathcal{F} : V \times \left(\frac{1}{2}, +\infty\right) \times I \rightarrow V \times \mathbb{R}, \quad (3.27)$$

given by

$$\mathcal{F}(\phi, c, \alpha) = \begin{bmatrix} \phi - (cD^\alpha + (c - 1))^{-1} \frac{1}{2} \Pi_0 \phi^2 \\ \int_{-\pi}^{\pi} \phi^3 dx - \int_{-\pi}^{\pi} \phi_0^3 dx \end{bmatrix}. \quad (3.28)$$

We see that \mathcal{F} is a well defined C^1 map and $\mathcal{F}(\phi_0, c_0, \alpha_0) = 0$. Our intention is to show the invertibility of the Fréchet derivative of \mathcal{F} with respect (ϕ, c) at (ϕ_0, c_0, α_0) . In fact, we see that

$$D_{\phi, c} \mathcal{F}(\phi_0, c_0, \alpha_0) = \begin{bmatrix} 1 - (c_0 D^{\alpha_0} + (c_0 - 1))^{-1} \Pi_0 \phi_0 & (c_0 D^{\alpha_0} + (c_0 - 1))^{-2} (D^{\alpha_0} + 1) \frac{1}{2} \Pi_0 \phi_0^2 \\ 3\langle \phi_0^2, \cdot \rangle & 0 \end{bmatrix}.$$

We claim that $D_{\phi, c} \mathcal{F}$ is invertible at (ϕ_0, c_0, α_0) . To do so, for every $g \in V$ and $\lambda \in \mathbb{R}$ given, we need to show the existence of a unique pair $(f, \beta) \in V \times \mathbb{R}$ such that

$$\begin{bmatrix} f - (c_0 D^{\alpha_0} + (c_0 - 1))^{-1} \Pi_0(\phi_0 f) + \frac{\beta}{2} (c_0 D^{\alpha_0} + (c_0 - 1))^{-2} (D^{\alpha_0} + 1) \Pi_0 \phi_0^2 \\ 3\langle \phi_0^2, f \rangle \end{bmatrix} = \begin{bmatrix} g \\ \lambda \end{bmatrix}. \quad (3.29)$$

In order to simplify the notation, we define $\mathcal{P} = -(c_0 D^{\alpha_0} + (c_0 - 1))^{-1} \Pi_0 \phi_0$ and $h_0 = \frac{1}{2} (c_0 D^{\alpha_0} + (c_0 - 1))^{-2} (D^{\alpha_0} + 1) \Pi_0 \phi_0^2$. By (3.29) we see that

$$(1 + \mathcal{P})f + \beta h_0 = g, \quad (3.30)$$

$$3 \int_{-\pi}^{\pi} \phi_0^2 f dx = \lambda. \quad (3.31)$$

Now, \mathcal{P} is a compact operator on $X_{0,e} = \{f \in X_0; f \text{ is even}\}$ and from (3.24) we obtain that -1 is not an element of the spectrum of \mathcal{P} . Thus, $1 + \mathcal{P}$ is an invertible operator on $X_{0,e}$ and since

$\mathcal{P} : V \rightarrow V$ (this fact follows similarly from [15, Lemma E.1]), we obtain that $(1 + \mathcal{P})^{-1}$ exists on V . Thus, we can express f uniquely as

$$f = -\beta(1 + \mathcal{P})^{-1}h_0 + (1 + \mathcal{P})^{-1}g. \quad (3.32)$$

Gathering results in (3.31) and (3.32), we deduce

$$3\beta \int_{-\pi}^{\pi} \phi_0^2(1 + \mathcal{P})^{-1}h_0 dx = -\lambda + 3 \int_{-\pi}^{\pi} \phi_0^2(1 + \mathcal{P})^{-1}g dx. \quad (3.33)$$

By (3.25), we see that $(\mathcal{L}|_{X_0})^{-1}$ exists on $X_{0,e}$ and therefore, on $H_{per}^{\alpha_0}(\mathbb{T}) \cap X_{0,e}$

$$(1 + \mathcal{P})^{-1} = (\mathcal{L}|_{X_0})^{-1}(c_0 D^{\alpha_0} + (c_0 - 1)) \quad (3.34)$$

is well defined. To show that β can be explicitly expressed in terms of ϕ_0 , \mathcal{P} and g , we need to prove that $\langle \phi_0^2, (1 + \mathcal{P})^{-1}h_0 \rangle \neq 0$. Indeed, using (3.34), the fact that $\mathcal{L}|_{X_0}\phi_0 = -\frac{1}{2}\Pi_0\phi_0^2$ and since $h_0 \in V$, we see from (3.33) that

$$\begin{aligned} \langle \phi_0^2, (1 + \mathcal{P})^{-1}h_0 \rangle &= \langle A(c_0), (1 + \mathcal{P})^{-1}h_0 \rangle - 2\langle \mathcal{L}|_{X_0}\phi_0, (1 + \mathcal{P})^{-1}h_0 \rangle \\ &= -2\langle \mathcal{L}|_{X_0}\phi_0, (\mathcal{L}|_{X_0})^{-1}(c_0 D^{\alpha_0} + (c_0 - 1))h_0 \rangle \\ &= -\langle (c_0 D^{\alpha_0} + (c_0 - 1))\phi_0, (c_0 D^{\alpha_0} + (c_0 - 1))^{-2}(D^{\alpha_0} + 1)\Pi_0\phi_0^2 \rangle \\ &= -\frac{1}{2}\langle \Pi_0\phi_0^2, (c_0 D^{\alpha_0} + (c_0 - 1))^{-2}(D^{\alpha_0} + 1)\Pi_0\phi_0^2 \rangle \\ &= -\pi \sum_{k \neq 0} \frac{|k|^{\alpha_0} + 1}{(c_0|k|^{\alpha_0} + (c_0 - 1))^2} |\widehat{\Pi_0\phi_0^2}(k)|^2 \neq 0, \end{aligned}$$

where \widehat{f} indicates the periodic Fourier transform for a function $f \in L_{per}^2(\mathbb{T})$. The rest of the proof can be done by a direct application of the implicit function theorem. \square

The next step is to follow similar arguments as found in [15, Subsection 5.2]. In fact, we consider the corresponding maximal extension of the branch (ϕ_α, c_α) for $\alpha \in [\alpha_0, \alpha_*)$, where α_* is given by

$$\alpha_* := \sup\{q; \alpha_0 < q < 2, (\phi_\alpha, c_\alpha) \in C^1([\alpha_0, q]; V \times (\frac{1}{2}, +\infty)) \text{ given by Proposition 3.2}$$

$$\text{and } (\phi_\alpha, c_\alpha) \text{ satisfies (1.8) for } \alpha \in [\alpha_0, q)\}.$$

It is clear that $\alpha_* \leq 2$ and we prove that $\alpha_* = 2$.

Proposition 3.3. *Let $\{\alpha_n\}_{n=1}^{n=+\infty} \subset (\frac{1}{2}, \alpha_*)$ be a sequence such that $\alpha_n \rightarrow \alpha_*$. Furthermore, we assume that $\phi_{\alpha_n} \in X_{0,e}$ are the corresponding solutions obtained in Proposition 3.2 with wave speed c_{α_n} . Up to a subsequence, it follows that*

$$\phi_{\alpha_n} \rightarrow \phi_* \text{ in } V \cap L_{per}^4(\mathbb{T}) \text{ and } c_{\alpha_n} \rightarrow c_*$$

where $c_* \in (\frac{1}{2}, +\infty)$ and ϕ_* satisfy (1.8). Moreover, the corresponding maximal branch $(\phi_\alpha, c_\alpha) \in C^1([\alpha_0, \alpha_*]; V \times (\frac{1}{2}, +\infty))$ extends to $\alpha_* = 2$.

Proof. First, let us suppose that $\{c_{\alpha_n}\}_{n=1}^{+\infty} \subset (\frac{1}{2}, +\infty)$ is not a bounded sequence. For all $M > 0$ large enough, there exists an index $m := n_M \in \mathbb{N}$ such that $c_{\alpha_m} > M$. Thus, from (1.8) and the fact that $\int_{-\pi}^{\pi} \phi_{\alpha_n}^3 dx = \int_{-\pi}^{\pi} \phi_0^3 dx > 0$ for all $n \in \mathbb{N}$, we obtain

$$0 \leq \int_{-\pi}^{\pi} (D^{\alpha_m/2} \phi_{\alpha_m})^2 dx \leq \frac{1}{2M} \int_{-\pi}^{\pi} \phi_{\alpha_n}^3 dx \leq \int_{-\pi}^{\pi} \phi_0^3 dx.$$

Since $\alpha_m \geq \alpha_0$, the Sobolev embedding $H_{per}^{\alpha_m/2}(\mathbb{T}) \cap X_{0,e} \hookrightarrow H_{per}^{\alpha_0/2}(\mathbb{T}) \cap X_{0,e}$ for all $\alpha_0 > \frac{1}{2}$ is valid. Now, $f \mapsto \|D^s f\|_{L_{per}^2(\mathbb{T})}$ is an equivalent norm on the space $H_{per}^s(\mathbb{T}) \cap X_{0,e}$ for all $s > 0$, and thus $\{\phi_{\alpha_m}\}_n$ is a bounded sequence on $H_{per}^{\alpha_0/2}(\mathbb{T}) \cap X_{0,e}$. Since $H_{per}^{\alpha_0/2}(\mathbb{T}) \cap X_{0,e}$ is compactly embedded into $V \hookrightarrow X_{0,e}$ and into $L_{per}^4(\mathbb{T})$, there exists $\phi_0 \in V \cap L_{per}^4(\mathbb{T})$ such that, up to a subsequence $\int_{-\pi}^{\pi} \phi_{\alpha_m}^3 dx \rightarrow \int_{-\pi}^{\pi} \phi_0^3 dx > 0$, $\int_{-\pi}^{\pi} \phi_{\alpha_m}^4 dx \rightarrow \int_{-\pi}^{\pi} \phi_0^4 dx > 0$ and $\int_{-\pi}^{\pi} \phi_{\alpha_m}^2 dx \rightarrow \int_{-\pi}^{\pi} \phi_0^2 dx > 0$, as $m \rightarrow +\infty$. By (1.8) and the Poincaré-Wirtinger inequality, we deduce

$$0 \leq (2M - 1) \int_{-\pi}^{\pi} \phi_{\alpha_m}^2 dx \leq (2c_{\alpha_m} - 1) \int_{-\pi}^{\pi} \phi_{\alpha_m}^2 dx \leq \frac{1}{2} \int_{-\pi}^{\pi} \phi_0^3 dx. \quad (3.35)$$

When $m \rightarrow +\infty$, we obtain from (3.35) a contradiction, so that $\{c_{\alpha_n}\}_{n=1}^{+\infty}$ is a bounded sequence on \mathbb{R} . Therefore, there exists a $c_* \in [\frac{1}{2}, +\infty)$ such that, up to a subsequence $c_{\alpha_n} \rightarrow c_*$. On the other hand, the fact that $c_{\alpha_n} \rightarrow c_* \geq \frac{1}{2}$ enables us to obtain,

$$0 \leq (2c_{\alpha_n} - 1) \|\phi_{\alpha_n}\|_{L_{per}^2}^2 \leq 2\mathcal{B}_{c_{\alpha_n}}(\phi_{\alpha_n}) = \frac{1}{2} \int_{-\pi}^{\pi} \phi_{\alpha_n}^3 dx = \frac{1}{2} \int_{-\pi}^{\pi} \phi_0^3 dx,$$

using again the Poincaré-Wirtinger inequality. Similarly as determined above, we guarantee the existence of $\phi_* \in V \cap L_{per}^4(\mathbb{T})$ such that

$$\int_{-\pi}^{\pi} \phi_{\alpha_n}^3 dx \rightarrow \int_{-\pi}^{\pi} \phi_*^3 dx, \quad \int_{-\pi}^{\pi} \phi_{\alpha_n}^2 dx \rightarrow \int_{-\pi}^{\pi} \phi_*^2 dx, \quad \text{and} \quad \int_{-\pi}^{\pi} \phi_{\alpha_n}^4 dx \rightarrow \int_{-\pi}^{\pi} \phi_*^4 dx. \quad (3.36)$$

Next, from the equation (1.8), we obtain

$$D^{\alpha_n} \phi_{\alpha_n} = - \left(1 - \frac{1}{c_{\alpha_n}}\right) \phi_{\alpha_n} + \frac{1}{2c_{\alpha_n}} \phi_{\alpha_n}^2 - \frac{A(c_{\alpha_n})}{c_{\alpha_n}},$$

so that $\{\phi_{\alpha_n}\}_n$ is bounded in $H_{per}^{\alpha_n}(\mathbb{T})$. Since $\alpha_n > \frac{1}{2}$, it follows that $H_{per}^{\alpha_n}(\mathbb{T})$ is a Banach algebra and thus $\{\phi_{\alpha_n}^2\}_n$ is bounded in $H_{per}^{\alpha_n}(\mathbb{T})$. From

$$D^{2\alpha_n} \phi_{\alpha_n} = - \left(1 - \frac{1}{c_{\alpha_n}}\right) D^{\alpha_n} \phi_{\alpha_n} + \frac{1}{2c_{\alpha_n}} D^{\alpha_n} \phi_{\alpha_n}^2, \quad (3.37)$$

one has that $\{\phi_{\alpha_n}\}_n$ is bounded in $H_{per}^{2\alpha_n}(\mathbb{T})$. Following with this inductive process, there is $r \in \mathbb{N}$ large enough such that $r\alpha_n > \alpha_*$ and $\{\phi_{\alpha_n}\}_n$ is bounded in $H_{per}^{r\alpha_n}(\mathbb{T})$. By the compact embedding $H_{per}^{r\alpha_n}(\mathbb{T}) \hookrightarrow H_{per}^{\alpha_*}(\mathbb{T})$, one has (modulus to a subsequence)

$$\phi_{\alpha_n} \rightarrow \phi_* \text{ in } H_{per}^{\alpha_*}(\mathbb{T}), \text{ as } n \rightarrow +\infty. \quad (3.38)$$

Therefore,

$$\|D^{\alpha_n} \phi_{\alpha_n} - D^{\alpha_*} \phi_*\|_{L^2_{per}} \leq \|D^{\alpha_*}(\phi_{\alpha_n} - \phi_*)\|_{L^2_{per}} + \|D^{\alpha_n} \phi_* - D^{\alpha_*} \phi_*\|_{L^2_{per}} \quad (3.39)$$

We obtain from (3.38), (3.39) and the fact that $\alpha_n \rightarrow \alpha_*$,

$$D^{\alpha_n} \phi_{\alpha_n} \rightarrow D^{\alpha_*} \phi_* \text{ in } L^2_{per}(\mathbb{T}), \text{ as } n \rightarrow +\infty. \quad (3.40)$$

Since $c_{\alpha_n} \rightarrow c_*$, the second and third convergences in (3.36) give us by (3.40) that ϕ_* satisfies the equation

$$c_* D^{\alpha_*} \phi_* + (c_* - 1) \phi_* - \frac{1}{2} \phi_*^2 + A(c_*) = 0. \quad (3.41)$$

To prove $\alpha_* = 2$, since ϕ_* satisfies (3.41), Proposition 3.2 allows to extend the branch (ϕ_α, c_α) beyond α_* . This fact contradicts the maximality property of α_* . This fact proves that $\alpha_* = 2$.

It remains to establish $c_* > \frac{1}{2}$. First of all, we obtain the smoothness of ϕ_* by a standard bootstrapping argument. If $c_* = \frac{1}{2}$, the only possibility for the smooth solution $\phi_* \in H^\infty_{per}(\mathbb{T}) \cap X_{0,e}$ of (3.41) with $\alpha_* = 2$ is that $\phi_* \equiv 0$ (see Remark 3.4 below). This fact generates a contradiction since $\int_{-\pi}^{\pi} \phi_*^3 dx > 0$.

□

Remark 3.4. *The explicit solution for the equation (3.41) with $\alpha_* = 2$ depends on the Jacobi elliptic function of dnoidal type and it is given by*

$$\phi_*(x) = a_* \left(\operatorname{dn}^2 \left(\frac{K(\kappa)}{\pi} x, \kappa \right) - \frac{E(\kappa)}{K(\kappa)} \right), \quad (3.42)$$

where

$$a_* = 12c_* \frac{K(\kappa)^2}{\pi^2}.$$

The value of c_* can be expressed by

$$c_* = \frac{\pi^2}{(4\kappa^2 - 8)K(\kappa)^2 + 12K(\kappa)E(\kappa) + \pi^2}, \quad (3.43)$$

where K and E are, respectively, the complete elliptic integral of first and second kind. Both functions depending on the modulus of the Jacobi elliptic function $\kappa \in (0, 1)$. For this solution, the integration constant

$$A(c_*) = \frac{1}{16} \frac{(24\kappa^2 - 24)K(\kappa)^4 + (96 - 48\kappa^2)E(\kappa)K(\kappa)^3 - 72K(\kappa)^2E(\kappa)^2}{((\kappa^2 - 2)K(\kappa)^2 + 3E(\kappa)K(\kappa) + \frac{\pi^2}{4})^2} \quad (3.44)$$

is obtained by using symbolic computations in Maple. By the explicit expressions given by (3.43) and (3.44), respectively, it is easy to see that c_* and $A(c_*)$ are strictly monotonic functions in terms of $\kappa \in (0, \kappa_0)$, where $\kappa_0 \approx 0.994$ is the unique zero of the function $p(\kappa) = (4\kappa^2 - 8)K(\kappa)^2 + 12K(\kappa)E(\kappa) + \pi^2$. Furthermore, we have

$$c_* \rightarrow \frac{1}{2} \text{ and } A(c_*) \rightarrow 0^+, \text{ as } \kappa \rightarrow 0^+.$$

From (3.43), we see that the limit $c_* > \frac{1}{2}$ is independent of the sequence $\{\alpha_n\}_{n=1}^{+\infty}$ presented by the Proposition 3.3. Furthermore, since ϕ_* is unique for each fixed c_* and satisfying $c_* > \frac{1}{2}$, we conclude that the limit ϕ_* is also independent of $\{\alpha_n\}_{n=1}^{+\infty}$.

Collecting all the results enunciated above, we can prove our uniqueness result to establish the precise statement in Theorem 1.3-iii).

Proposition 3.5. *Let $\alpha \in (\frac{1}{2}, 2]$ be fixed. If (3.24) is valid, then the solution obtained in Lemma 3.2 is unique.*

Proof. The proof of this result has the same spirit as determined in [15, Theorem 2.4]. Suppose that ϕ_0 and $\tilde{\phi}_0$ are solutions of the problem (1.19) satisfying $\phi_0 \not\equiv \tilde{\phi}_0$. Since both ϕ_0 and $\tilde{\phi}_0$ are in Y_0 , we obtain that $\int_{-\pi}^{\pi} \phi_0 dx = \int_{-\pi}^{\pi} \tilde{\phi}_0 dx = 0$ and

$$\int_{-\pi}^{\pi} \phi_0^3 dx = \int_{-\pi}^{\pi} \tilde{\phi}_0^3 dx \quad (3.45)$$

For the case of solitary waves, equality (3.45) is one of the crucial parts in the uniqueness proof contained in [15] since they use the classical Pohozaev equality to reach the result. According to our best knowledge, it is well known that we do not have a similar equality in the periodic context, so that equality (3.45) is essential for our purpose. By Proposition 3.3, there exist two smooth branch of solutions $(\phi_\alpha, c_\alpha), (\tilde{\phi}_\alpha, \tilde{c}_\alpha) \in C^1([\alpha_0, 2]; V \times (\frac{1}{2}, +\infty))$ satisfying (1.8) with $c_\alpha, \tilde{c}_\alpha \in (\frac{1}{2}, +\infty)$, $(\phi_{\alpha_0}, c_{\alpha_0}) := (\phi_0, c_0)$ and $(\tilde{\phi}_{\alpha_0}, \tilde{c}_{\alpha_0}) := (\tilde{\phi}_0, c_0)$. Notice that by the local uniqueness obtained from Proposition 3.2, the smooth branches (ϕ_α, c_α) and $(\tilde{\phi}_\alpha, \tilde{c}_\alpha)$ do not have a common point. Thus, for each branch (ϕ_α, c_α) and $(\tilde{\phi}_\alpha, \tilde{c}_\alpha)$, we have by Proposition 3.3 the existence of (ϕ_*, c_*) and $(\tilde{\phi}_*, \tilde{c}_*)$ satisfying the following ordinary differential equations

$$-c_*\phi_*'' + (c_* - 1)\phi_* - \frac{1}{2}\phi_*^2 + A(c_*) = 0 \quad (3.46)$$

and

$$-\tilde{c}_*\tilde{\phi}_*'' + (\tilde{c}_* - 1)\tilde{\phi}_* - \frac{1}{2}\tilde{\phi}_*^2 + A(\tilde{c}_*) = 0, \quad (3.47)$$

respectively. Moreover, by Proposition 3.2-(iii), one has

$$\int_{-\pi}^{\pi} \phi_\alpha^3 dx = \int_{-\pi}^{\pi} \phi_0^3 dx = \int_{-\pi}^{\pi} \tilde{\phi}_0^3 dx = \int_{-\pi}^{\pi} \tilde{\phi}_\alpha^3 dx. \quad (3.48)$$

Again, using compactness arguments as done in the proof of Proposition 3.3, it follows that $\int_{-\pi}^{\pi} \phi_\alpha^3 dx \rightarrow \int_{-\pi}^{\pi} \phi_*^3 dx$ and $\int_{-\pi}^{\pi} \tilde{\phi}_\alpha^3 dx \rightarrow \int_{-\pi}^{\pi} \tilde{\phi}_*^3 dx$, as $\alpha \rightarrow 2$. By (3.48) we have

$$\int_{-\pi}^{\pi} \phi_*^3 dx = \int_{-\pi}^{\pi} \tilde{\phi}_*^3 dx. \quad (3.49)$$

We claim that $c_* = \tilde{c}_*$. Indeed, for $\alpha_* = 2$, let us consider $\varphi := \varphi_\omega$ a general zero mean solution for the equation (3.46) with wave speed

$$\omega := \omega_\kappa = \frac{\pi^2}{(4\kappa^2 - 8)K(\kappa)^2 + 12K(\kappa)E(\kappa) + \pi^2},$$

instead of c_* . Deriving equation (3.46) with respect to ω , we obtain

$$-\omega \frac{d}{d\omega} \varphi'' - \omega \varphi'' + (\omega - 1) \frac{d}{d\omega} \varphi + \varphi - \varphi \frac{d}{d\omega} \varphi + \frac{d}{d\omega} A(\omega) = 0. \quad (3.50)$$

Next, multiplying equation (3.50) by φ , integrating the result over $[-\pi, \pi]$, we have after an integration by parts that

$$\frac{\omega}{2} \frac{d}{d\omega} \left(\int_{-\pi}^{\pi} (\varphi')^2 + \varphi^2 dx \right) - \frac{1}{2} \frac{d}{d\omega} \int_{-\pi}^{\pi} \varphi^2 dx + \int_{-\pi}^{\pi} (\varphi')^2 + \varphi^2 dx - \frac{1}{3} \frac{d}{d\omega} \int_{-\pi}^{\pi} \varphi^3 dx = 0. \quad (3.51)$$

On the other hand, multiplying (3.46) by φ and integrating the final result over $[-\pi, \pi]$, it follows that

$$\omega \left(\int_{-\pi}^{\pi} (\varphi')^2 + \varphi^2 dx \right) - \int_{-\pi}^{\pi} \varphi^2 dx - \frac{1}{2} \int_{-\pi}^{\pi} \varphi^3 dx = 0. \quad (3.52)$$

Deriving expression (3.52) with respect to ω and combining the final result with (3.51), we obtain

$$\frac{d}{d\omega} \int_{-\pi}^{\pi} \varphi^3 dx = 6 \left(\int_{-\pi}^{\pi} (\varphi')^2 + \varphi^2 dx \right) > 0, \quad (3.53)$$

so that, $\omega \mapsto \int_{-\pi}^{\pi} \varphi^3 dx$ is one-to-one, and consequently $c_* = \tilde{c}_*$.

Finally, by Remark 3.4, we have that ω given explicitly by

$$\omega = \frac{\pi^2}{(4\kappa^2 - 8)K(\kappa)^2 + 12K(\kappa)E(\kappa) + \pi^2} \quad (3.54)$$

is positive and strictly increasing in terms of $\kappa \in (0, \kappa_0)$. We have that ϕ_* and $\tilde{\phi}_*$ are 2π -periodic solutions given explicitly by (3.42) with the same wave speed $c_* = \tilde{c}_*$. Since ω is one-to-one according with (3.54), it follows that the corresponding modulus of c_* and \tilde{c}_* must satisfy $\kappa_* = \tilde{\kappa}_*$, so that $\tilde{\phi} \equiv \phi_*$. The remainder of the proof is similar to [15, Theorem 2.4]. \square

Remark 3.6. Suppose that $n(\mathcal{L}) = 1$ for every $c > \frac{1}{2}$ and $\alpha \in (\frac{1}{2}, 2]$. Since (3.24) is valid, let $\{\alpha_n\}_{n=1}^{n=+\infty} \subset (\frac{1}{2}, \alpha_*)$ be a sequence such that $\alpha_n \rightarrow \alpha_*$. The additional assumption on the number of negative eigenvalues gives us immediately the following equality:

$$n(\mathcal{L}_{\alpha_n}) = n(\mathcal{L}_{\alpha_*}) = 1,$$

where $\mathcal{L}_{\alpha_n} = c_{\alpha_n} D^{\alpha_n} + (c_{\alpha_n} - 1) - \phi_{\alpha_n}$ and $\mathcal{L}_{\alpha_*} = c_* D^{\alpha_*} + (c_* - 1) - \phi_*$ indicate the linearized operators around the periodic waves ϕ_{α_n} and ϕ_{α_*} , respectively. It is important to mention that our uniqueness result remains valid if the quantity of negative eigenvalues for \mathcal{L}_{α_n} is not stable in the sense that we may have $2 = n(\mathcal{L}_{\alpha_{n_0}}) > n(\mathcal{L}_{\alpha_*}) = 1$ for some $n_0 \in \mathbb{N}$. However, our numerical approach in Section

5 attests that $d < 0$ for all $c > \frac{1}{2}$, so that the quantity of negative eigenvalues of \mathcal{L}_{α_n} remains stable along $c \in (\frac{1}{2}, +\infty)$ for all $n \in \mathbb{N}$.

Remark 3.7. *It is worth mentioning that Proposition 3.5 agrees in some sense with the uniqueness result for positive and periodic waves associated to the fractional Korteweg-de Vries equation as in [27]. The authors employed Krasnoselskii's fixed-point theorem and three additional facts:*

- i) $n(\mathcal{L}) = 1$.
- ii) $\ker(\mathcal{L}) = [\phi']$.
- iii) $\alpha \in (0.5849, 2]$.

Remark 3.8. *Similarly as reported on page 1962 of [31], if ϕ is a solution of (1.6) obtained from Lemma 3.2, we also conjecture that (3.24) is always valid for all $c > \frac{1}{2}$. Our conjecture and the results in this section allow to conclude that the uniqueness of minimizers is always expected.*

3.3. Existence of the small amplitude solutions. Let $0 < \alpha \leq 2$. In this subsection, we determine the existence of small amplitude periodic wave solutions for the boundary-value problem (1.8) which is an even profile. We show that this kind of solutions are given by the corresponding expansion of the wave for c near $\frac{1}{2}$.

For obtaining the small amplitude periodic waves of the boundary-value problem (1.8), we use the arguments contained in [9, Chapter 8.4], where the authors have determined the existence of the referred waves via bifurcation theory and the Lyapunov-Schmidt reduction.

For $\alpha \in (0, 2]$, let $F : H_{per}^\alpha(\mathbb{T}) \cap X_{0,e} \times (0, +\infty) \rightarrow X_{0,e}$ be the map defined by

$$F(g, \omega) := \omega D^\alpha g + (\omega - 1)g - \frac{1}{2}\Pi_0 g^2. \quad (3.55)$$

We see that F is smooth in all variables. Moreover, $F(g, \omega) = 0$ if and only if, g is a solution of the boundary-value problem (1.8) corresponding to the wave speed $\omega \in (0, +\infty)$. In particular, it is clear that $F(0, \omega) = 0$ for all $\omega \in (0, +\infty)$.

The Fréchet derivative associated to the function F with respect to g is given by

$$D_g F(g, \omega) : H_{per}^\alpha(\mathbb{T}) \cap X_{0,e} \rightarrow X_{0,e}, \quad f \mapsto (\omega D^\alpha + \omega - 1 - \Pi_0 g)f. \quad (3.56)$$

Let $c_0 > 0$ be fixed and consider the corresponding linearized operator \mathcal{L} around the wave ϕ defined as in (1.10). The derivative (3.56) at the point (ϕ, c_0) becomes the well known self-adjoint projector operator $\mathcal{L}|_{X_0}$. Moreover, at the point $(0, c_0)$, we have

$$D_g F(0, c_0) = c_0 D^\alpha + c_0 - 1. \quad (3.57)$$

Notice that the nontrivial kernel of $D_g F(0, c_0)$ is determined by functions $h \in H_{per}^\alpha(\mathbb{T}) \cap X_{0,e}$ satisfying

$$\widehat{h}(k)((c_0 - 1) + c_0 |k|^\alpha) = 0, \quad k \neq 0. \quad (3.58)$$

Then, $D_g F(0, c_0)$ has the one-dimensional kernel if and only if, $c_0 = (1 + |k|^\alpha)^{-1}$ for some $k \in \mathbb{Z}$, in which case it is given by

$$\ker D_g F(0, c_0) = [\tilde{\phi}_k(x)], \quad (3.59)$$

where $\tilde{\phi}_k(x) = \cos(xk)$.

The local bifurcation theory in [9, Chapter 8.4] enables us to guarantee the existence of an open interval \mathcal{J} containing $c_0 = (1 + |k|^\alpha)^{-1}$, an open ball $B_r \in H_{per}^\alpha(\mathbb{T}) \cap X_{0,e}$ of radius $r > 0$ centered at $g = 0$ and a unique smooth mapping $c \in \mathcal{J} \mapsto \phi(c) \in H_{per}^\alpha(\mathbb{T}) \cap X_{0,e}$ such that $F(\phi(c), c) = 0$ for all $c \in \mathcal{J}$ and $\phi(c_0) = 0$. In other words, every solution of $F(g, \omega) = 0$ has the form $(\phi(c), c)$.

For each integer $k \geq 1$, the point $(0, \tilde{c}_k)$, where $\tilde{c}_k := (1 + k^\alpha)^{-1}$, is a bifurcation point. Furthermore, there exists $a_0 > 0$ and a local bifurcation curve $a \in (0, a_0) \mapsto (\phi_k(a), c_k(a))$ through $(0, \tilde{c}_k)$ constituted by even $\frac{2\pi}{k}$ -periodic solutions of the boundary-value problem (1.8). Additionally, one has $c_k(0) = \tilde{c}_k$ and $D_a \phi_k(0) = \tilde{\phi}_k(x) = \cos(xk)$ and all solutions of $F(g, \omega) = 0$ in a neighbourhood of $(0, \tilde{c}_k)$ belongs to the above curve depending on a .

We apply the method of Lyapunov-Schmidt reduction to obtain the explicit expansion of the wave ϕ for c near $\frac{1}{2}$. In addition, the local bifurcation curve extends to a global smooth curve $c \in ((1 + k^\alpha)^{-1}, +\infty) \mapsto (\phi_k(c), c_k(c))$ of solutions for the equation $F(g, \omega) = 0$. For the simple case $k = 1$, both results can be summarized in the next proposition.

Proposition 3.4. *Let $\alpha \in (0, 2]$ be fixed. There exists $a_0 > 0$ such that for all $a \in (0, a_0)$ there is a unique even local periodic solution ϕ for the boundary value problem (1.8) given by the following expansion:*

$$\phi(x) = a \cos(x) + \frac{a^2}{2(2^\alpha - 1)} \cos(2x) + O(a^3) \quad (3.60)$$

and

$$c = \frac{1}{2} + \frac{a^2}{4(2^\alpha + 1)(2^\alpha - 1)} + O(a^4). \quad (3.61)$$

In addition, constant A is given by

$$A(c) = \frac{a^2}{4} + O(a^4). \quad (3.62)$$

The pair $(\phi, c) \in H_{per}^\alpha(\mathbb{T}) \cap X_{0,e} \times (\frac{1}{2}, +\infty)$ is global in terms of the parameter $c > \frac{1}{2}$ and it satisfies (1.8).

Proof. The proof of this result relies on slight modifications of [8, Theorem 5.4 and Theorem 5.6]. \square

In the next result, we see that q_c in (3.2) is continuous in c and for $c > \frac{1}{2}$ and that $q_c \rightarrow 0^+$ as $c \rightarrow \frac{1}{2}^+$. This fact gives us that the small amplitude periodic waves obtained in the Proposition 3.4 satisfy the minimization problem (3.2) for $c \rightarrow \frac{1}{2}^+$.

Proposition 3.5. *Let $\phi \in Y_0$ be the solution of the constrained minimization problem obtained in Lemma 3.2 and $q_c = \mathcal{B}_c(\phi)$. Then q_c is continuous in c for $c > \frac{1}{2}$ and $q_c \rightarrow 0$ as $c \rightarrow \frac{1}{2}$.*

Proof. This result is similar to [31, Lemma 2.3]. \square

Next result establishes good spectral properties for the linearized operator \mathcal{L} in (1.10) around the single-lobe solution ϕ obtained in Lemma 3.2 by knowing the spectral information for the same operator around the small amplitude periodic waves obtained by Proposition 3.4.

Proposition 3.6. *Let $\alpha \in (\frac{1}{2}, 2]$ be fixed. For every $c > \frac{1}{2}$, we have $n(\mathcal{L}) = 1$ if and only if $z(\mathcal{L}) = 1$. Moreover, \mathcal{L} consists in a countable sequence of positive eigenvalues bounded away from zero.*

Proof. The proof has the same spirit as in [27, Lemma 2.3]. According to the Proposition 3.1, the number of negative eigenvalues of \mathcal{L} may change in the parameter continuations in c if and only if, the eigenvalues pass through zero eigenvalue. If $n(\mathcal{L}) = 1$ for every $c > \frac{1}{2}$, we see that $\ker(\mathcal{L}|_{X_0}) = [\phi']$ by [31, Corollary 4.5]. If $z(\mathcal{L}) = 2$ for some $c_1 > \frac{1}{2}$ we see that parameter d defined in 3.1 satisfies $d = 0$ at $c = c_1$. This means by Proposition 3.1 that $n(\mathcal{L}) = 2$ at $c = c_1$ which is a contradiction. Suppose that $z(\mathcal{L}) = 1$ for every $c > \frac{1}{2}$. For the small amplitude periodic waves ϕ obtained by the Proposition 3.4, we see that $c \mapsto \phi \in H_{per}^\infty(\mathbb{T})$ is smooth. Now, from (3.61), we have the expansion

$$d = 1 + 2A(c) - c - cA'(c) = \frac{1 - 16^\alpha + 14 \cdot 2^{-2+2\alpha} - 3}{2 \cdot 4^\alpha - 1} + O(a^3). \quad (3.63)$$

For $\alpha \in (\frac{1}{2}, 2]$, we have $d < 0$ and this means by Proposition 3.1 that $n(\mathcal{L}) = 1$ for the small amplitude periodic waves. By a continuity argument, we obtain that $n(\mathcal{L}) = 1$ for the linearized operator \mathcal{L} around the single-lobe solution ϕ determined by Lemma 3.2 for all $c > \frac{1}{2}$. \square

4. SPECTRAL STABILITY

In this section, we study the spectral stability of (1.13). First, we see that the even solution for the boundary-value problem (1.8) obtained in Lemma 3.2 is smooth by using similar arguments as in [31, Proposition 2.4]. Moreover ϕ has a single-lobe profile, namely, there exists only one maximum (at $x = 0$) and minimum of ϕ on \mathbb{T} . This fact is in accordance with the approach in [23] to decide about the non-degeneracy of the $\ker(\tilde{\mathcal{L}})$.

For $\alpha \in (0, 2]$ and since ϕ is a single-lobe solution, we see by Oscillation Theorem in [23] that the operator $\tilde{\mathcal{L}}$ has at most two negative eigenvalues, that is, $n(\tilde{\mathcal{L}}) \in \{1, 2\}$. Consequently, there may be at most two eigenfunctions of $\tilde{\mathcal{L}}$ for the zero eigenvalue, that is, $z(\tilde{\mathcal{L}}) \in \{1, 2\}$. Recall that $\ker(\tilde{\mathcal{L}}|_{X_0}) = [\phi']$ implies $c \in (\frac{1}{2}, +\infty) \mapsto \phi$ is smooth and d in (3.12) can be used to decide about the non-degeneracy of $\ker(\tilde{\mathcal{L}})$ and the exact quantity of $n(\tilde{\mathcal{L}})$ according to the Proposition 3.1. Indeed, if $d \neq 0$ we conclude by the first equality in (3.10), (3.11), and (3.12) that $\{1, \phi, \phi^2\} \in \text{range}(\tilde{\mathcal{L}})$. Proposition 3.1 in [23] is now used to conclude that $\ker(\tilde{\mathcal{L}}) = [\phi']$. On the other hand, if $d = 0$, we have $\ker(\tilde{\mathcal{L}}) = [\phi', \frac{d}{dc}\phi - \frac{1}{c} - \frac{1}{c}\phi]$. In addition, $d < 0$ implies $n(\tilde{\mathcal{L}}) = 1$ and $d > 0$ gives us that $n(\tilde{\mathcal{L}}) = 2$.

First, we prove the spectral stability for the small amplitude periodic waves associated to the case $\alpha \in (0, 2]$.

Proposition 4.1. *Let $\alpha \in (0, 2]$. The small amplitude periodic waves $\phi \in H_{per}^\infty(\mathbb{T})$ determined by Proposition 3.4 are spectrally stable in the sense of Definition 1.1.*

Proof. Since \mathcal{L} is a Hessian operator for $G(u)$ in (1.10) and $\tilde{\mathcal{L}} = \frac{1}{c}\mathcal{L}$ for all $c > \frac{1}{2}$, the spectral stability holds if $\tilde{\mathcal{L}}|_{\{1, (D^{\alpha+1})\phi\}^\perp} \geq 0$, that is, if $n(\tilde{\mathcal{L}}|_{\{1, (D^{\alpha+1})\phi\}^\perp}) = 0$. On the other hand, the periodic wave ϕ is spectrally unstable if $n(\tilde{\mathcal{L}}|_{\{1, (D^{\alpha+1})\phi\}^\perp}) = 1$.

For $\lambda \notin \sigma(\tilde{\mathcal{L}})$, let us define the following symmetric 2-by-2 matrix given by

$$S(\lambda) := \begin{bmatrix} \langle (\tilde{\mathcal{L}} - \lambda I)^{-1} 1, 1 \rangle & \langle (\tilde{\mathcal{L}} - \lambda I)^{-1} (D^\alpha + 1)\phi, 1 \rangle \\ \langle (\tilde{\mathcal{L}} - \lambda I)^{-1} 1, (D^\alpha + 1)\phi \rangle & \langle (\tilde{\mathcal{L}} - \lambda I)^{-1} (D^\alpha + 1)\phi, (D^\alpha + 1)\phi \rangle \end{bmatrix}.$$

Using (3.63), for $\alpha \in (0, \tilde{\alpha}) \cup (\frac{1}{2}, +\infty)$ we obtain $d < 0$, whereas for $\tilde{\alpha} < \alpha < \frac{1}{2}$, we have $d > 0$. For $\alpha = \tilde{\alpha} \approx 0.2924$ and $\alpha = \frac{1}{2}$, we have the existence of fold points for which $d = 0$.

Assume $d \neq 0$, we compute at $\lambda = 0$. Indeed,

$$\langle \tilde{\mathcal{L}}^{-1} (D^\alpha + 1)\phi, (D^\alpha + 1)\phi \rangle = -\frac{2\pi d^{-1}}{c} (A'(c))^2 - 2\pi A'(c) - \frac{1}{6} \gamma'(c), \quad (4.1)$$

$$\langle \tilde{\mathcal{L}}^{-1} (D^\alpha + 1)\phi, 1 \rangle = \langle \tilde{\mathcal{L}}^{-1} 1, (D^\alpha + 1)\phi \rangle = \frac{2\pi d^{-1}}{c} A'(c), \quad (4.2)$$

and

$$\langle \tilde{\mathcal{L}}^{-1} 1, 1 \rangle = -2\pi \frac{d^{-1}}{c}, \quad (4.3)$$

where $\gamma(c) = \int_{-\pi}^{\pi} \phi^3 dx$ and $\frac{1}{3} \gamma'(c) = \frac{8\pi}{c} A(c) - \frac{1}{c} \gamma(c)$. The determinant of $S(0)$, for $d \neq 0$, is given by

$$\det S(0) = \frac{4\pi^2 d^{-1}}{c} \left(A'(c) + \frac{1}{\pi} \mathcal{B}_c(\phi) \right). \quad (4.4)$$

By (1.24) we have $A'(c) > 0$ for all $\alpha \in (0, 2]$. Since $\mathcal{B}_c(\phi) > 0$, we see that the sign of $\det S(0)$ is determined only by d^{-1} since the terms between parentheses in (4.4) are strictly positive.

We denote by n_0 and z_0 the number of negative and zero eigenvalues of $S(0)$, respectively. If $d = 0$, then $S(0)$ is singular, in which case we denote the number of diverging eigenvalues of $D(\lambda)$ as $\lambda \rightarrow 0$ by z_∞ . Thus, [33, Theorem 4.1] gives us the following relations:

$$\begin{cases} n(\tilde{\mathcal{L}} |_{\{1, (D^\alpha + 1)\phi\}^\perp}) &= n(\tilde{\mathcal{L}}) - n_0 - z_0, \\ z(\tilde{\mathcal{L}} |_{\{1, (D^\alpha + 1)\phi\}^\perp}) &= z(\tilde{\mathcal{L}}) + z_0 - z_\infty \end{cases}.$$

Assume $d \neq 0$, so that $z_\infty = 0$. If $0 < \alpha < \tilde{\alpha}$ or $\alpha \in (\frac{1}{2}, 2]$, we have $z_0 = 0$ and $\det S(0) < 0$ which implies that $n_0 = 1$. On the other hand by Corollary 3.5, we see that the waves in (3.60) solve the minimization problem (3.2) and Proposition 3.1 gives us $n(\mathcal{L}) = 1$. Therefore, we have the spectral stability in this case because $n(\tilde{\mathcal{L}} |_{\{1, (D^\alpha + 1)\phi\}^\perp}) = 1 - 1 - 0 = 0$. For the case $\tilde{\alpha} < \alpha < \frac{1}{2}$, we have $n_0 = 0$ or $n_0 = 2$. Since the trace of $S(0)$ is strictly negative, we conclude that $n_0 = 2$. Hence, we obtain the spectral stability of ϕ since $n(\tilde{\mathcal{L}}) = 2$ and $n(\tilde{\mathcal{L}} |_{\{1, (D^\alpha + 1)\phi\}^\perp}) = 2 - 2 - 0 = 0$.

Finally, for the case $\alpha = \tilde{\alpha}$ or $\alpha = \frac{1}{2}$, we obtain from the relations above and Proposition 3.1 that $z_\infty = 1$, $n(\tilde{\mathcal{L}}) = 2$, so that $n(\tilde{\mathcal{L}} |_{\{1, (D^\alpha + 1)\phi\}^\perp}) = 2 - 2 = 0$. \square

Remark 4.2. *The result contained in Proposition 4.1 can be extended for the case $\alpha > 2$.*

The next result guarantees sufficient conditions for the spectral stability of the periodic minimizer ϕ and it gives the proof of Theorem 1.3-iv).

Proposition 4.3. *Let $\alpha \in (\frac{1}{3}, 2]$ be fixed. If $\ker(\mathcal{L}|_{X_0}) = [\phi']$, the periodic minimizer ϕ obtained in Lemma 3.2 is spectrally stable in the sense of Definition 1.1 if:*

- i) $d \neq 0$ and $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi) > 0$.
- ii) $\alpha \in (\frac{1}{2}, 2]$ and $d \neq 0$ for all $c > \frac{1}{2}$.

Proof. We use the same computations as determined in the proof of Proposition 4.1. To prove i), suppose that $\ker(\mathcal{L}|_{X_0}) = [\phi']$, $d > 0$ and $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi) > 0$. By Proposition 3.1, we have $n(\mathcal{L}) = 2$ and by (4.3) the first entry of the matrix $S(0)$ in Proposition 4.1 is negative. Thus $n_0 = 2$ and we have the spectral stability since $n(\tilde{\mathcal{L}}|_{\{1, (D^{\alpha+1})\phi\}^\perp}) = n(\tilde{\mathcal{L}}) - n_0 - z_0 = 2 - 2 - 0 = 0$. If $d < 0$ and $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi) > 0$, we see that $\det S(0) < 0$ and we have the spectral stability since $n(\tilde{\mathcal{L}}|_{\{1, (D^{\alpha+1})\phi\}^\perp}) = n(\tilde{\mathcal{L}}) - n_0 - z_0 = 1 - 1 - 0 = 0$. To prove ii), since $d \neq 0$ for all $c > \frac{1}{2}$, we obtain by Proposition 3.1 that $z(\mathcal{L}) = 1$, so that $n(\mathcal{L}) = 1$ by Proposition 3.6. We claim that $1 + 2A(c) - c - cA'(c) < 0$ for all $c > \frac{1}{2}$. Indeed, if there exists $c_2 > \frac{1}{2}$ such that $1 + 2A(c_2) - c_2 - c_2A'(c_2) \geq 0$, we deduce by Proposition 3.1 that $n(\mathcal{L}) = 2$ at $c = c_2$. This fact generates a contradiction and the claim is satisfied. Next, we obtain after multiplying this inequality by the factor $\frac{1}{c^3}$ that

$$\frac{d}{dc} \left(A(c) \frac{1}{c^2} \right) > \frac{1}{c^3} - \frac{1}{c^2}. \quad (4.5)$$

Integrating (4.5) over the interval $(\frac{1}{2}, c]$ and using $A(c) \rightarrow 0$ when $c \rightarrow \frac{1}{2}$, we obtain the important inequality

$$A(c) > c - \frac{1}{2}. \quad (4.6)$$

Thus, we have from the Poincaré-Wirtinger inequality, (4.6), inequality $1 + 2A(c) - c - cA'(c) < 0$, and a simple computation

$$A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi) \geq \left(c - \frac{1}{2} \right)^2 > 0. \quad (4.7)$$

Therefore, if $c > \frac{1}{2}$ one has by (4.7) that $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi) > 0$. From (4.4), we obtain

$$\det S(0) = \frac{4\pi^2 d^{-1}}{c} \left(A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi) \right) < 0.$$

This together with $n(\mathcal{L}) = 1$ give us from the index formula contained in Proposition 4.1 the spectral stability of the single-lobe ϕ . □

Remark 4.4. *The numerical approach in the next section will attest for a fixed $\alpha \in (\frac{1}{2}, 2]$ that $d \neq 0$ for all $c > \frac{1}{2}$, so that the periodic waves obtained in Lemma 3.2 are spectrally stable by Proposition 4.3 – ii). For the case $\alpha \in (\frac{1}{3}, \frac{1}{2}]$ we will have the following situation: there exists a unique c^* such that $d > 0$ for all $c \in (\frac{1}{2}, c^*)$ and $d < 0$ for all $c \in (c^*, +\infty)$. Since in both cases we have $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi) > 0$, we obtain from (4.4) that $\det S(0) > 0$ for the case $d > 0$ and $\det S(0) < 0$ if*

$d < 0$. In both cases, the spectral stability is obtained by Proposition 4.3 – i). For the case $d = 0$, we also have the spectral stability of ϕ using a similar procedure as in Proposition 4.1.

5. NUMERICAL EXPERIMENTS

In this section we propose a Petviashvili's iteration method for the numerical generation of periodic travelling wave solutions of the equation (1.1). The method is widely used for the generation of travelling wave solutions [13, 14, 27, 32, 34]. The iteration method for the periodic waves, a modification of standard Petviashvili's algorithm [14, 27], is based on the following solution steps. First, we use the transformation

$$\psi = \frac{1}{2c} \left[\phi - (c-1) + \sqrt{(c-1)^2 + 2A} \right] \quad (5.1)$$

to convert the equation (1.6) into

$$D^\alpha \psi + w\psi - \psi^2 = 0, \quad (5.2)$$

where $w = \frac{1}{c} \sqrt{(c-1)^2 + 2A}$. Thus, we obtain the equation with a zero integration constant. Integrating (5.1) on $[-\pi, \pi]$ and using the fact that $\phi : \mathbb{T} \rightarrow \mathbb{R}$ is a periodic function with the zero mean value yield that

$$c = \left(1 - w + \frac{1}{\pi} \int_{-\pi}^{\pi} \psi dx \right)^{-1}. \quad (5.3)$$

Next, we use the standard Petviashvili's method to solve the equation (5.2). Employing the Fourier transform to the equation (5.2) gives

$$(|\xi|^\alpha + w) \widehat{\psi}(\xi) - \widehat{\psi^2}(\xi) = 0. \quad (5.4)$$

A simple iterative algorithm for numerical calculation of $\widehat{\psi}(\xi)$ for the above equation can be proposed in the form

$$\widehat{\psi}_{n+1}(\xi) = \frac{\widehat{\psi_n^2}(\xi)}{|\xi|^\alpha + w}, \quad n \in \mathbb{N} \quad (5.5)$$

where $\widehat{\psi}_n(\xi)$ is the Fourier transform of ψ_n which is the n^{th} iteration of the numerical solution. Here the solutions are constructed under the assumption

$$|\xi|^\alpha + w \neq 0. \quad (5.6)$$

Since the above algorithm is usually divergent, we finally present the Petviashvili's method as

$$\widehat{\psi}_{n+1}(\xi) = \frac{M_n^\nu}{|\xi|^\alpha + w} \widehat{\psi_n^2}(\xi) \quad (5.7)$$

by introducing the stabilizing factor

$$M_n = \frac{\langle (D^\alpha + w)\psi_n, \psi_n \rangle}{\langle \psi_n^2, \psi_n \rangle}, \quad \psi_n \in H_{per}^\alpha(\mathbb{T}) \quad (5.8)$$

where $\langle \cdot, \cdot \rangle$ denotes the standard inner product in $L^2_{per}(\mathbb{T})$. Here, the free parameter ν is chosen as 2 for the fastest convergence (see [34] for details). The iterative process is controlled by the error between two consecutive iterations given by

$$Error(n) = \|\psi_n - \psi_{n-1}\|_{L^\infty_{per}}$$

and the stabilization factor error given by

$$|1 - M_n|.$$

The residue of the interaction process is determined by $RES(n) = \|\mathcal{S}\psi_n\|_{L^\infty_{per}}$, where

$$\mathcal{S}\psi = D^\alpha\psi + w\psi - \psi^2.$$

The periodic traveling wave solution ϕ of the BBM equation corresponds to boundary value problem (1.6) with $\alpha = 2$ is given in the equation (3.42).




FIGURE 5.1. The exact and the numerical solutions of the BBM equation with the wave speed $c = 1.2181$ and the variation of $Error(n)$, $|1 - M_n|$ and RES with the number of iterations in semi-log scale.

In order to test the accuracy of our scheme, we compare the exact solution (3.42) with the numerical solution obtained by using the expansion corresponding to (5.2) as the initial guess

$$\psi_0(x) = 1 + a \cos(x) + a^2 \left[\frac{1}{2} - \frac{1}{2(2^\alpha - 1)} + \frac{\cos(2x)}{2(2^\alpha - 1)} \right] + a^3 \frac{\cos(3x)}{2(2^\alpha - 1)(3^\alpha - 1)}, \quad (5.9)$$

where $a = 0.2$ and $\alpha = 2$. In this experiment, the space interval is $[-\pi, \pi]$ and number of grid points is chosen as $N = 2^{12}$. In the first panel of Figure 5.1, we depict the exact and numerical solutions for the wave speed $c = 1.2181$. As it is seen from the figure, the exact and the numerical solutions coincide. In the other panels of Figure 5.1, the variations of three different errors with the number of iteration are presented. These results show that our numerical scheme captures the solution remarkably well.




FIGURE 5.2. Variation of exact and numerical values of integration constant A versus c (left panel) and variation of numerical value of d versus c (right panel) for $\alpha = 2$.

The left panel of Figure 5.2 shows the exact and numerical variation of the integration constant A with respect to c . The exact relation between A and c is illustrated by using the eqs. (3.43) and (3.44). From the numerical point of view, first we obtain the numerical solution by using Petviashvili's scheme (5.7) for every $w \in (1, +\infty)$. Finally, the numerical value of the integration constant $A(c)$ is evaluated by

$$A(c) = \frac{1}{2} [c^2 w^2 - (c - 1)^2]. \quad (5.10)$$

As it is seen from the figure, the numerical and the exact values of $A(c)$ coincide. In the right panel of the figure, we present the variation of d with c . We observe that d is always negative. The single-lobe periodic solution to the boundary value problem (5.2) for $\alpha = 1$ is given by

$$\psi(x) = \frac{\sinh \gamma}{\cosh \gamma - \cos x}, \quad (5.11)$$




FIGURE 5.3. The exact and the numerical solutions of rBO equation for the wave speed $c = 1.2192$.




FIGURE 5.4. Variation of exact and numerical values of integration constant A versus c (left panel) and variation of numerical value of d versus c (right panel) for $\alpha = 1$.

where parameter $\gamma \in (0, +\infty)$ is given by $\gamma = \coth^{-1}(w)$. The transformation (5.1) allows one to obtain the single-lobe periodic solution for the rBO equation as

$$\phi(x) = 2c \left(\frac{\sinh \gamma}{\cosh \gamma - \cos x} - 1 \right). \quad (5.12)$$

We compare the exact solution (5.12) with the numerical solution obtained by using the expansion (5.9) with $a = 0.2$ and $\alpha = 1$ as the initial guess. The space interval is $[-\pi, \pi]$ and number of grid points is chosen as $N = 2^9$. We present the exact and numerical solutions for the wave speed $c = 1.2192$ in Figure 5.3. Since we have $\int_0^\pi \psi(x) dx = \pi$ from (5.11), we obtain $c = (3 - w)^{-1}$ by using (5.3). Eliminating w in (5.10), we compute the integration constant $A(c) = 4c^2 - 2c$ explicitly. In the

left panel of Figure 5.4, we compare the exact and numerical variation of the integration constant A with respect to c for $\alpha = 1$. The right panel shows the variation of d with c for $\alpha = 1$. Figures 5.2 and 5.4 show that $A(c)$ is strictly increasing for all values of c and d is always negative. Therefore, numerical results are compatible with Proposition 4.3 stating the spectral stability of single-lobe solution for $\alpha \in (\frac{1}{2}, 2]$.




FIGURE 5.5. Various numerical wave profiles (left panel) and the modulus of the Fourier coefficients of the numerical solution (right panel) for $\alpha = 0.55$.

In the next numerical experiment, we choose $\alpha = 0.55 > \alpha_0$ since the fold point $\alpha_0 = 0.5$ for the fBBM equation. In the left panel of Figure 5.5, we illustrate the periodic wave profiles for several values of c . It can be seen that the amplitude becomes more and more peaked with the increasing wave speed c . The right panel of Figure 5.5 shows that the modulus of the Fourier coefficients computed via a discrete Fourier transformation decreases to machine precision for $N = 2^{12}$ Fourier modes. In Figure 5.6, we depict the variation of $A(c)$ and d with c . The numerical results are again compatible with the Proposition 4.3 for $\alpha > \frac{1}{2}$.

In Figure 5.7, we show the variation of $A(c)$ with c for $\alpha = 0.45 < \alpha_0$. The picture shows a lack of convergence using Petviashvili's scheme in the sense that function $A(c)$ can not reach the bifurcation point $c = \frac{1}{2}$ as determined for the case $\alpha \in (\frac{1}{2}, 2]$. To do best of our knowledge, this phenomena gives us (at least numerically) an indication that the number of negative eigenvalues for the linearized operator is 2 and/or the existence of a fold point, that is, a value of c such that $\dim(\ker(\mathcal{L})) = 2$. In both cases, it is well known that the numerical method does not converge (see [27, 31, 34]) and prevent us to show that ϕ solves the minimization problem (1.19). Due to the lack of numerical data for $c \in (0.5, 1.8)$, we illustrate the curve (dashed line) by using the relation

$$A(c) \approx \left(c - \frac{1}{2}\right) (2^\alpha + 1)(2^\alpha - 1), \quad (5.13)$$




FIGURE 5.6. Variation of A versus c (left panel) and variation of d versus c (right panel) both evaluated numerically for $\alpha = 0.55$.




FIGURE 5.7. Variation of values of A versus c (left panel) and close-up look to the gap (right panel) ($\alpha = 0.45$). The solid line shows the numerical result obtained by Petviashvili's method and the dashed line shows the relation (5.13) from the small amplitude expansion.

obtained by the expressions of c and $A(c)$ in (1.23). The right panel gives a closer look to the gap between the branch of the small amplitude periodic waves and numerical result. In order to fill this gap, we use the Newton's method for $c \in (0.5, 1.8)$ in Figure 5.8.

In Figure 5.9, we present the variation of d and the term $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi)$ with c for $\alpha = 0.45$. We observe that $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi)$ is positive for all values of c . However, d is positive up to a critical speed $c^* \approx 0.953$ and then becomes negative. Therefore, $\det S(0)$ is positive for $c \in (\frac{1}{2}, c^*)$ and negative




FIGURE 5.8. Variation of numerical values of A versus c with the combination of Newton and Petviashvili method ($\alpha = 0.45$)




FIGURE 5.9. Variation of d versus c (top left), close-up look near critical value c^* (top right) and the variation of $A'(c) + \frac{1}{\pi}\mathcal{B}_c(\phi)$ with c (bottom) for $\alpha = 0.45$.




FIGURE 5.10. Variation of d versus c (top left), close-up look near critical value c^* (top right) and the variation of $A'(c) + \frac{1}{\pi}B_c(\phi)$ with c (bottom) for $\alpha = 0.5$.

for $c \in (c^*, +\infty)$. We also illustrate the variation of d and the term $A'(c) + \frac{1}{\pi}B_c(\phi)$ with c in Figure 5.10 for the fold point $\alpha_0 = 0.5$. The numerical results are very similar to the ones for $\alpha = 0.45$ with $c^* \approx 0.67$. As it is seen from the figures, $\det S(0) < 0$ for $c \in (c^*, +\infty)$ which indicates the spectral stability. However, $\det S(0) > 0$ for $c \in (\frac{1}{2}, c^*)$ which yields that $n_0 = 0$ or $n_0 = 2$. Since in this case $d > 0$, we have by Proposition 6.1 that ϕ is spectrally stable for all $c \in (\frac{1}{2}, c^*)$.

6. REMARKS ON THE ORBITAL STABILITY OF PERIODIC WAVES

In this section, we present a brief discussion concerning the orbital stability of the periodic wave ϕ obtained by Lemma 3.2.

Before stating the result, we need some preliminary tools. For functions u and v in $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$, we

define ρ as the “distance” between u and v given by

$$\rho(u, v) = \inf_{y \in \mathbb{R}} \|u - v(\cdot + y)\|_{H_{per}^{\frac{\alpha}{2}}}.$$

Our precise definition of orbital stability is given below.

Definition 6.1. *We say that the periodic solution ϕ is orbitally stable in $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$, by the periodic flow of (1.1), if for any $\varepsilon > 0$ there exists $\delta > 0$ such that for any $u_0 \in H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$ satisfying $\|u_0 - \phi\|_{H_{per}^{\frac{\alpha}{2}}} < \delta$, the solution $u(t)$ of (1.1) with initial data u_0 exists globally and satisfies*

$$\rho(u(t), \phi) < \varepsilon,$$

for all $t \geq 0$.

Remark 6.2. *Our notion of orbital stability prescribes the existence of global solutions. Thus, from Section 2 we need to assume that $\alpha > 1$.*

Next, let us introduce the conserved quantity

$$V(u) = \frac{1}{2} \int_{-\pi}^{\pi} \left(u^2 + \frac{u^3}{3} \right) dx = P(u) - E(u) \quad (6.1)$$

and the auxiliary functional

$$Q(u) := -V(u) + (c - 1 - A(c))M(u).$$

In what follows, we set

$$\Upsilon_0 = \{u \in H_{per}^{\frac{\alpha}{2}}(\mathbb{T}); \langle Q'(\phi), u \rangle = 0\}.$$

Note that Υ_0 is nothing but the tangent space to $\{u \in H_{per}^{\frac{\alpha}{2}}(\mathbb{T}); Q(u) = Q(\phi)\}$ at ϕ . With these notations, the result in [11, Theorem 2.1] (see also [1]) reads as follows.

Proposition 6.1. *Suppose that $n(\mathcal{L}) = 1$ and $z(\mathcal{L}) = 1$. If there exists $\Phi \in H_{per}^{\alpha}(\mathbb{T})$ such that $\langle \mathcal{L}\Phi, \Psi \rangle = 0$ for all $\Psi \in \Upsilon_0$ and $\langle \mathcal{L}\Phi, \Phi \rangle < 0$, then ϕ is orbitally stable in $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$ by the periodic flow of (1.1).*

□

Finally, we can use the estimate (4.6) to obtain the orbital stability of ϕ . This last result gives us the proof of Theorem 1.3-v).

Proposition 6.2. *Let $\alpha \in (1, 2]$ be fixed. If $\ker(\mathcal{L}|_{X_0}) = [\phi']$ and $d \neq 0$ for all $c > \frac{1}{2}$, the periodic wave ϕ obtained in Lemma 3.2 is orbitally stable in $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$ in the sense of Definition 6.1.*

Proof. Since $n(\mathcal{L}) = z(\mathcal{L}) = 1$ by Proposition 3.6, we only need to use Proposition 6.1 for a convenient Φ . Indeed, let us consider $\Phi = 1 + \phi$. For

$$Q(u) = -V(u) + (c - 1 - A(c))M(u),$$

we obtain $\langle Q'(\phi), \phi' \rangle = 0$ and $\langle \mathcal{L}(1 + \phi), \Psi \rangle = 0$ for all $\Psi \in \Upsilon_0$. If $\langle \mathcal{L}(1 + \phi), 1 + \phi \rangle < 0$, one has the orbital stability of ϕ in the energy space $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$. To calculate $\langle \mathcal{L}(1 + \phi), 1 + \phi \rangle$, we employ the Poincaré-Wirtinger inequality and (4.6) to obtain

$$\begin{aligned} \langle \mathcal{L}(1 + \phi), 1 + \phi \rangle &= 2\pi(c - 1) - \int_{-\pi}^{\pi} c(D^{\frac{\alpha}{2}}\phi)^2 + (c + 1)\phi^2 dx \\ &\leq 2\pi(c - 1) - 4\pi(2c + 1)A(c) \\ &\leq 2\pi(c - 1) - 4\pi(2c + 1)\left(c - \frac{1}{2}\right) = -2\pi c(4c - 1) < 0. \end{aligned} \tag{6.2}$$

By Proposition 6.1 one has the orbital stability in the energy space $H_{per}^{\frac{\alpha}{2}}(\mathbb{T})$. \square

ACKNOWLEDGMENTS

The authors are grateful to the two anonymous referees for their valuable suggestions and comments which greatly improved the presentation of the paper. S. Amaral was supported by the regular doctorate scholarship from CAPES. F. Natali is partially supported by CNPq (grant 304240/2018-4), Fundação Araucária (grant 002/2017) and CAPES MathAmSud (grant 88881.520205/2020-01).

REFERENCES

- [1] G. ALVES, F. NATALI AND A. PASTOR, *Sufficient conditions for orbital stability of periodic traveling waves*, J. Diff. Equat., 267 (2019), pp. 879-901.
- [2] V. AMBROSIO, *On some convergence results for fractional periodic Sobolev spaces*, Opuscula Math., 40 (2020), pp. 5-20.
- [3] J. ANGULO, *Stability properties of solitary waves for fractional KdV and BBM equations*, Nonlinearity, 31 (2018), pp. 920-956.
- [4] J. ANGULO, J., E. CARDOSO JR. AND F. NATALI, *Stability properties of periodic traveling waves for the intermediate long wave equation*, Rev. Mat. Iber. 33 (2017), pp. 417-448.
- [5] J. ANGULO, C. BANQUET AND M. SCIALOM, *The regularized Benjamin-Ono and BBM equations: Well-posedness and nonlinear stability*, J. Diff. Equat., 250 (2011), pp. 4011-4036.
- [6] J. ANGULO AND F. NATALI, *Positivity properties of the Fourier transform and the stability of periodic travelling-wave solutions*, SIAM J. Math. Anal., 40 (2008), pp. 1123-1151.
- [7] T.B. BENJAMIN, J.L. BONA AND J.J. MAHONY, *Model Equations for Long Waves in Nonlinear Dispersive Systems*, Phil. Trans. Royal Soc. London. Series A, Math. Phys. Sci., 272 (1972), pp. 47-78
- [8] G. BRUELL AND R.N. DHARA, *Waves of maximal height for a class of nonlocal equations with homogeneous symbol*, Indiana Univ. Math. Journal, to appear, (2020).
- [9] B. BUFFONI AND J. TOLAND, *Analytic Theory of Global Bifurcation*. Princeton Series in Applied Mathematics. Princeton University Press, Princeton, NJ, 2003.
- [10] K. CLAASEN AND M. JOHNSON, *Nondegeneracy and stability of antiperiodic bound states for fractional nonlinear Schrödinger equations*, J. Diff. Eqs., 266 (2019), 5664-5712.
- [11] F. CRISTÓFANI, F. NATALI AND A. PASTOR, *Periodic Traveling-wave solutions for regularized dispersive equations: Sufficient conditions for orbital stability with applications*, Comm. Math. Sci., 18 (2020), pp. 613-634.

- [12] B. DECONINCK AND T. KAPITULA *On the spectral and orbital stability of spatially periodic stationary solutions of generalized Korteweg-de Vries equations*, in Hamiltonian Partial Diff. Eq. Appl., 75 285-322., Fields Inst. Comm., Springer, New York, 2015.
- [13] A. DURAN, *An efficient method to compute solitary wave solutions of fractional Korteweg-de Vries equations*. Int J Comp Math., 95 (2018), pp. 1362-1374.
- [14] A. DURAN, *Numerical generation of periodic traveling wave solutions of some nonlinear dispersive wave systems*. J. Comp. App. Math., 316 (2017), pp. 29-39.
- [15] R.L. FRANK AND E. LENZMANN, *Uniqueness of non-linear ground states for fractional Laplacians in \mathbb{R}* , Acta Math., 210 (2013), pp. 261–318.
- [16] T. GALLAY AND M. HÄRÄĞUŞ, *Stability of small periodic waves for the nonlinear Schrödinger equation*, J. Diff. Equat., 234 (2007), pp. 544-581.
- [17] S. BENZONI-GAVAGE, C. MIETKA AND L.M. RODRIGUES, *Co-periodic stability of periodic waves in some Hamiltonian PDEs*, Nonlinearity, 29 (2016), pp. 3241–3308.
- [18] M. GRILLAKIS, J. SHATAH, AND W. STRAUSS, *Stability theory of solitary waves in the presence of symmetry I*, J. Funct. Anal., 74 (1987), pp. 160-197.
- [19] S. HAKKAEV., *Nonlinear Stability of Periodic Traveling Waves of the BBM System*, Comm. Math. Anal., 15 (2013), pp. 39-51.
- [20] M. HÄRÄĞUŞ AND E. WAHLÉN, *Transverse instability of periodic and generalized solitary waves for a fifth-order KP model*, J. Diff. Equat., 262 (2017), pp. 3235-3249.
- [21] M. HÄRÄĞUŞ, *Stability of periodic waves for the generalized BBM equation*, Rev. Roumaine Math. Pures Appl., 53 (2008), pp. 445–463
- [22] V.M. HUR AND A.K. PANDEY, *Modulational instability in nonlinear non local equations of regularized long wave type*, Phys. D, 325 (2016), pp. 98-112.
- [23] V.M. HUR AND M. JOHNSON, *Stability of periodic traveling waves for nonlinear dispersive equations*, SIAM J. Math. Anal., 47 (2015), 3528–3554.
- [24] M. JOHNSON, *Stability of small periodic waves in fractional KdV-type equations*, SIAM J. Math. Anal., 45 (2013), pp. 3168-3193.
- [25] H. KALISCH, *Error analysis of a spectral projection of the regularized Benjamin–Ono equation*, BIT Numer. Math., 45 (2005), pp. 69–89.
- [26] H. KIELHÖFER, *Bifurcation theory*, Appl. Math. Sci., Springer, New York, 2012.
- [27] U. LE AND D.E. PELINOVSKY, *Convergence of Petviashvili’s method near periodic waves in the fractional Korteweg-de Vries equation*, SIAM J. Math. Anal., 51 (2019), pp. 2850-2583.
- [28] Z. LIN, *Instability of nonlinear dispersive solitary waves*, J. Funct. Anal., 255 (2008), pp. 1091-1124.
- [29] F. LINARES, D. PILOD AND J-C. SAUT *Dispersive perturbations of Burgers and hyperbolic equations I: local theory*, SIAM J. Math. Anal., 46 (2015), pp. 1505–1537.
- [30] F. NATALI, D.E. PELINOVSKY AND U. LE, *Periodic waves in the fractional modified Korteweg–de Vries equation*, to appear in J. Dyn. Diff. Equat. (2022).
- [31] F. NATALI, U. LE AND D.E. PELINOVSKY, *New variational characterization of periodic waves in the fractional Korteweg-de Vries equation*, Nonlinearity, 33 (2020), pp 1956-1986.
- [32] G. ORUC, H. BORLUK, G. M MUSLU, *The generalized fractional Benjamin-Bona-Mahony equation: Analytical and numerical results*, Physica D: Nonlinear Phenomena, (2020), Article number:132499.
- [33] D.E. PELINOVSKY, *Localization in periodic potentials: from Schrödinger operators to the Gross–Pitaevskii equation*, LMS Lecture Note Series, 390 Cambridge University Press, Cambridge, 2011.
- [34] D.E. PELINOVSKI AND Y.A. STEPANYANTS, *Convergence of Petviashvili’s iteration method for numerical approximation of stationary solution of nonlinear wave equations*. SIAM J. Numer. Anal., 42 (2004), pp. 1110-1127.

(S. Amaral) DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF MARINGÁ, MARINGÁ, PR, BRAZIL.

Email address: `sabrinasuelen20@gmail.com`

(H. Borluk) OZYEGIN UNIVERSITY, DEPARTMENT OF NATURAL AND MATHEMATICAL SCIENCES, CEKMEKOY, ISTANBUL, TURKEY.

Email address: `handan.borluk@ozyegin.edu.tr`

(G. M. Muslu) ISTANBUL TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS, MASLAK, ISTANBUL, TURKEY.

Email address: `gulcin@itu.edu.tr`

(F. Natali) DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF MARINGÁ, MARINGÁ, PR, BRAZIL.

Email address: `fmanatali@uem.br`

(G. Oruc) ISTANBUL TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS, MASLAK, ISTANBUL, TURKEY.

Email address: `topkarci@itu.edu.tr`