# A NOTE ON BARKER SEQUENCES OF EVEN LENGTH

#### JÜRGEN WILLMS

ABSTRACT. A Barker sequence is a binary sequence for which all nontrivial aperiodic autocorrelations are either 0, 1 or -1. The only known Barker sequences have length 2, 3, 4, 5, 7, 11 or 13. It is an old conjecture that no longer Barker sequences exist and in fact, there is an overwhelming evidence for this conjecture. For binary sequences of odd length, this conjecture is known to be true, whereas for even length it is still open, whether a Barker sequence of even length greater 4 exists.

Similar to the well-known fact that a Barker sequence of odd length is necessarily skew-symmetric, we show that in the case of even length there is also a form of symmetry albeit weaker. In order to exploit this symmetry, we derive different formulas for the calculation of the aperiodic correlation. We prove by using only elementary methods that there is no Barker sequence of even length n>4 with  $C_1=C_3=\cdots=C_{\frac{n}{2}-1}$ , where  $C_k$  denotes the kth aperiodic autocorrelation of the sequence.

### 1. Introduction

For an integer n > 1 an element of  $\{-1,1\}^n$  is called a binary sequence a of length n. We write  $a_j$  for the j-th entry in a. For  $k = 0, 1, \dots, n-1$  the kth aperiodic autocorrelation of the binary sequence a is defined by

$$C_k(a) = \sum_{j=1}^{n-k} a_j a_{j+k}.$$

The autocorrelation  $C_0(a)$  is called the *peak autocorrelation* of a, whereas all other values  $C_k(a)$  are called the *nontrivial* aperiodic autocorrelations. Note that  $C_0(n) = n$ . In the following a will always denote a binary sequence of length n > 1 and we will abbreviate  $C_k(a)$  as  $C_k$ . A simple parity argument shows that

$$(1.1) C_k \equiv n - k \pmod{2}$$

for each  $k = 1, 2, \dots, n - 1$ . Furthermore, we have

(1.2) 
$$(\sum_{j=1}^{n} a_j)^2 = n + 2\sum_{k=1}^{n-1} C_k.$$

Moreover, use the simple parity argument  $xy \equiv x-y+1 \pmod 4$  for  $x,y \in \{-1,1\}$  to obtain for  $k=1,2,\cdots,n-1$ 

(1.3) 
$$C_k + C_{n-k} = \sum_{j=1}^n a_j a_{j+k} \equiv n \pmod{4},$$

Date: 1.4.2021.

2000 Mathematics Subject Classification. 11B83, 05B10, 94A55.

Key words and phrases. binary sequence, Barker sequence, autocorrelation.

where the index j + k of a is reduced modulo n if j + k > n.

Of particular interest are binary sequences for which the absolute values of all aperiodic autocorrelations are small; for a recent survey see for example [5]. An ideal sequence from this point of view is a Barker sequence. A binary sequence a is called  $Barker\ sequence$  if all nontrivial aperiodic autocorrelations of a are either 0, 1 or -1. Hence by (1.1) all nontrivial aperiodic autocorrelations of a Barker sequence are in magnitude as small as possible.

The only known Barker sequences have length 2, 3, 4, 5, 7, 11 or 13. It is a longstanding conjecture that no longer Barker sequences exist. If n is odd, then a well-known result of Turyn and Storer [7] published in 1961 says, that Barker sequences of odd length n > 13 do not exist. More then 50 years later, it was noted [9] that their elementary but arduous inductive proof is incomplete. A correction of the proof was suggested in [6]. A new proof was given by Borwein and Erdélyi [1] in 2013 and an even simpler and very different proof was presented by Schmidt and Willms [6] in 2016.

The Barker conjecture for binary sequences of even length is still open, but there is an overwhelming evidence for it. For example, results from Borwein and Mossinghoff [2] and Leung and B. Schmidt [4] show that there is no Barker sequence of even length n for  $4 < n \le 4 \cdot 10^{33}$ .

Let us briefly try to explain why in the Barker conjecture the case of even length appears to be considerably harder than the case of odd length. Suppose that a is a Barker sequence of length n and let  $1 \le k < n$ . If the length n is odd then by (1.1) we know that  $C_k = 0$  if k is odd, and (1.3) gives us that  $C_k = (-1)^{\frac{n-1}{2}}$  if k is even; so in this case all aperiodic autocorrelations are fixed. In contrast, if the length n of a Barker sequence is even, then  $C_k = 0$  if k is even. But now (1.3) gives us only that

$$(1.4) C_k = -C_{n-k}.$$

So in this case only half of the aperiodic autocorrelations that are not zero are fixed; no further constraints on the other half (as for example  $C_1, C_3, \dots, C_{\frac{n}{2}-1}$ ) are known (apart of course from the fact that each value is either 1 or -1).

Furthermore, by (1.2) if a is a Barker sequence of even length  $n \ge 4$  then  $|\sum_{j=1}^n a_j| = \sqrt{n}$  and hence  $n = 4r^2$  for some  $r \in \mathbb{N}$ ; in particular, n is a multiple of 4. There are many more stronger and deeper partial results on the non-existence of Barker sequences of even length (see for example [3] or [2]), which however we will not use in the following.

Let a be a Barker sequences of even length  $n \geq 4$  and suppose that the first  $\frac{n}{4}$  odd aperiodic autocorrelations  $C_1, C_3, \dots, C_{\frac{n}{2}-1}$  are all equal. The next result shows that this is only possible if the length n of a equals 4.

**Theorem 1.** Suppose a is a Barker sequences of even length  $n \ge 4$  with  $C_1 = C_3 = \cdots = C_{\frac{n}{2}-1}$ . Then n = 4.

We need a couple of lemmata in order to prove Theorem 1 and proceed as follows. In the Section 2, we consider what we call weak symmetric sequences, and show that a putative Barker sequence of even length  $n \geq 4$  must be weak symmetric. In Section 3, we derive different formulas for the calculation of the aperiodic correlation that are in particular useful in the weak symmetric case. Finally, in Section 4 we apply our results to sequences with a stronger form of symmetry, which enables us to prove Theorem 1.

Let us outline the main idea behind the proof of Theorem 1. Suppose that a is a putative Barker sequence of even length  $n \geq 4$  with  $C_1 = C_3 = \cdots = C_{\frac{n}{2}-1}$  and put  $u = \frac{n}{4}$ . Similar as in [7] we show that

(1.5) 
$$a_j a_{j+1} = a_{2j} a_{2j+1}$$
 for each  $j$  satisfying  $1 \le j < u$ .

Now construct a new sequence p from a by reversing the first half of a and do the same with the second half of a. It follows from Lemma 8 that p is also a Barker sequence of length n. Applying again (1.5) gives us that  $|C_u(b)| = u \neq 0$  where b denotes the binary sequence of length 2u with  $b_j = a_j$  for each  $1 \leq j \leq 2u$ . Since by Lemma 7 we have  $C_k(b) = 0$  for even k, it follows that u must be odd. Now assume that n > 4. Then it is not difficult to show that both  $C_2(b)$  and  $C_{2u-2}(b)$  must be zero. But by (1.3) we have  $C_2(b) + C_{2u-2}(b) \equiv 2u \pmod{4}$  which implies that u is even, a contradiction. Hence n = 4.

## 2. Weak Symmetry

In the following, we will exploit the symmetry of a putative Barker sequence of even length. A very similar approach was used in the case of odd length Barker sequences; all proofs in this case rely heavily on the fact that a Barker sequence of odd length n is skew-symmetric, that is  $a_j a_{n+1-j} = (-1)^{\frac{n+1}{2}+j}$  for each  $1 \leq j \leq n$ . In the case of a Barker sequence of even length we will prove a weaker form of symmetry that leads us to the following definition. We call a binary sequence a weak symmetric if the length n of a is a multiple of 4 and  $a_j a_{j+1} = -a_{n+1-j} a_{n-j}$  holds for each odd integer j with  $1 \leq j < \frac{n}{2}$  (and thus for each odd integer j with  $1 \leq j < n$ ). In Proposition 4 we will show that a Barker sequence of even length  $n \geq 4$  is indeed weak symmetric.

For  $k = 1, 2, \dots, n$  put

$$T_k = \sum_{j=1}^k a_j a_{n+1-j}.$$

The next result shows that weak symmetry,  $T_k$  and  $C_{n-k}$  are strongly related.

**Lemma 2.** Let a be a binary sequence of length n and let  $1 \le k < n$ . Then:

- (i)  $C_{n-k} \equiv T_k \pmod{4}$ .
- (ii) Suppose that a is weak symmetric. Then  $T_k = 0$  if k is even, and  $T_k = a_k a_{n+1-k}$  if k is odd.

*Proof.* (i) Use the simple counting argument that is stated in Lemma 3 (i) to obtain

$$(-1)^{\frac{k-T_k}{2}} = \prod_{j=1}^k a_j a_{n+1-j} = \prod_{j=1}^k a_j a_{j+n-k} = (-1)^{\frac{k-C_{n-k}}{2}}$$

which implies that  $C_{n-k} \equiv T_k \pmod{4}$ .

(ii) Suppose a is weak symmetric. Then  $a_1a_2 = -a_{n-1}a_n$  and thus  $T_2 = 0$ . Since for k > 2 we have  $T_k = T_{k-2} + a_{k-1}a_{n-k} + a_ka_{n+1-k}$ , it follows that  $T_k = 0$  if k is even. Set  $T_0 = 0$ ; if k is odd, then k = 2j + 1 for some  $j \ge 0$  and thus  $T_k = T_{2j} + a_ka_{n+1-k} = a_ka_{n+1-k}$ .

<sup>&</sup>lt;sup>1</sup>The conclusion that u is odd, here derived by merely elementary arguments, follows also from a result of Turyn [8] using fairly deep methods of character theory.

The next lemma uses a simple counting argument. Recall that  $b_j$  denotes the j-th entry in b.

**Lemma 3.** Let b be a binary sequence of length m.

(i) Let  $S = \sum_{j=1}^{m} b_j$ ; then

$$\prod_{j=1}^{m} b_j = (-1)^{\frac{m-S}{2}}$$

(ii) Let  $R = \sum_{j=1}^{m} (-1)^{j} b_{j}$ ; then

$$\prod_{j=1}^{m} b_j = \begin{cases} (-1)^{\frac{R}{2}} & \text{if } m \text{ is even} \\ (-1)^{\frac{R+1}{2}} & \text{if } m \text{ is odd.} \end{cases}$$

*Proof.* (i) Denote by  $\alpha$  resp.  $\beta$  the number of positive resp. negative entries in b. Then  $\alpha + \beta = m$ ,  $S = \alpha - \beta$  and hence  $m - S = 2\beta$ . Since  $\prod_{j=1}^{m} b_j = (-1)^{\beta}$ , the assertion (i) follows directly.

(ii) Let  $\gamma$  denote the cardinality of the set  $\{1 \leq j \leq m : j \text{ odd}\}$ . Then  $\gamma = \frac{m}{2}$  if m is even, and  $\gamma = \frac{m+1}{2}$  if m is odd. Hence we have  $(-1)^{\gamma} \prod_{j=1}^{m} b_j = \prod_{j=1}^{m} (-1)^j b_j$ . Use (i) to obtain  $\prod_{j=1}^{m} (-1)^j b_j = (-1)^{\frac{m-R}{2}}$ . By considering the cases m even and m odd separately, assertion (ii) follows easily.

We have already seen in Section 1 that the length n of a Barker sequence of even length is a multiple of 4 and that its nontrivial aperiodic autocorrelations  $C_k$  equal zero if k is even; moreover, it is also weak symmetric as the next result shows.

**Proposition 4.** Let a be a Barker sequence of even length  $n \ge 4$ . Then a is weak symmetric, and  $C_k = -a_k a_{n+1-k}$  if k is odd with  $1 \le k < n$ .

Proof. Let  $1 \leq j \leq \frac{n}{2}$ . Note that  $T_1 = a_1 a_n = C_{n-1}$  and  $|T_{j+1} - T_j| = 1$ . Suppose that a is a Barker sequence of even length  $n \geq 4$ . Then  $|C_{n-j} - C_{n-j-1}|$ . Since by Lemma 2 (i) we have  $C_{n-j} \equiv T_j \pmod 4$ , it follows that  $C_{n-j} \equiv T_j$ . In particular, we have  $T_j = 0$  if j is even. If j is odd then  $0 = T_{j+1} - T_{j-1} = a_j a_{n+1-j} + a_{j+1} a_{n-j}$ , which shows that a is weak symmetric. If k is odd with  $1 \leq k < n$  then by Lemma 2 (ii) and (1.4) we have  $C_k = -C_{n-k} = -T_k = -a_k a_{n+1-k}$ .

## 3. Correlation Formulas

In order to simplify the notation and to exploit the symmetry we will use the following abbreviation. For a binary sequence a of length n put  $\delta_k = a_k a_{n+1-k}$  for  $k = 1, 2, \dots, n$ .

Note that for  $1 \le k \le n$  we have

$$\delta_k = \delta_{n+1-k}.$$

Moreover, if the binary sequence a of length n is weak symmetric, then it follows directly from the definition that for each odd k (with  $1 \le k < n$ )

$$\delta_k = -\delta_{k+1}.$$

One goal of this section is to express the aperiodic autocorrelations  $C_k$  of a binary sequence a of length n=2m by using only the first m entries  $a_1,a_2,\cdots,a_m$  of a together with  $\delta_1,\delta_2,\cdots,\delta_m$ . The next lemma is a first step towards this goal.

**Lemma 5.** Let a be a binary sequence a of length n and let  $1 \le k < n$ .

(i) If k is even then

$$C_{n-k} = \sum_{j=1}^{\frac{k}{2}} a_j a_{k+1-j} (\delta_j + \delta_{k+1-j}).$$

(ii) If k is odd then

$$C_{n-k} = \delta_{\frac{k+1}{2}} + \sum_{j=1}^{\frac{k-1}{2}} a_j a_{k+1-j} (\delta_j + \delta_{k+1-j}).$$

*Proof.* By definition we have  $C_{n-k} = \sum_{j=1}^k a_j a_{j+n-k} = \sum_{j=1}^k a_j a_{k+1-j} \delta_{k+1-j}$ . In order to prove (i) and (ii) combine in the last sum the first with the last summand, the second with the second last, and so on.

Let a be a binary sequence a of even length n=2m and suppose that  $1 \leq k \leq m$ . Note that the formula for  $C_{n-k}$  in Lemma 5 does not refer to an element  $a_j$  or  $\delta_j$  with j > m. In order to to achieve the same goal when computing  $C_k$  (instead of  $C_{n-k}$ ) we have to split the sum on the right hand side of (i) resp. (ii) of Lemma 5 into two parts as shown in the next lemma.

**Lemma 6.** Let a be a binary sequence a of even length n = 2m and let  $1 \le k < m$ .

(i) If k is even then

$$C_k = \sum_{j=1}^{m-k} a_j a_{j+k} (1 + \delta_j \delta_{j+k}) + \sum_{j=1}^{\frac{k}{2}} a_{j+m-k} a_{m+1-j} (\delta_{j+m-k} + \delta_{m+1-j}).$$

(ii) If k is odd then

$$C_k = \delta_{\frac{n-k+1}{2}} + \sum_{j=1}^{m-k} a_j a_{j+k} (1 + \delta_j \delta_{j+k}) + \sum_{j=1}^{\frac{k-1}{2}} a_{j+m-k} a_{m+1-j} (\delta_{j+m-k} + \delta_{m+1-j}).$$

*Proof.* We show that both (i) and (ii) follow directly from Lemma 5. Suppose first that k is even. Then Lemma 5 (i) gives us that  $C_k = \sum_{j=1}^{\frac{n-k}{2}} a_j a_{n-k+1-j} (\delta_j + \delta_{n-k+1-j})$ . If  $1 \leq j \leq m-k$  then by (3.1) we have  $a_{n-k+1-j} = a_{k+j} \delta_{k+j}$ ,  $\delta_{k+j} = \delta_{n-k+1-j}$ , and hence  $a_j a_{n-k+1-j} (\delta_j + \delta_{n-k+1-j}) = a_j a_{j+k} \delta_{k+j} (\delta_j + \delta_{n-k+1-j}) = a_j a_{j+k} (\delta_j \delta_{j+k} + 1)$ . Since

$$\sum_{j=m-k+1}^{\frac{n-k}{2}} a_j a_{n-k+1-j} (\delta_j + \delta_{n-k+1-j}) = \sum_{j=1}^{\frac{k}{2}} a_{j+m-k} a_{m+1-j} (\delta_{j+m-k} + \delta_{m+1-j})$$

assertion (i) follows. Suppose next that k is odd. Then by Lemma 5 (ii) we have  $C_k = \delta_{\frac{n-k+1}{2}} + \sum_{j=1}^{\frac{n-k-1}{2}} a_j a_{n-k+1-j} (\delta_j + \delta_{n-k+1-j})$ , and the same argument as above can be used to prove (ii).

## 4. Strong Symmetric Binary Sequences

In the this section we will use Lemma 5 and Lemma 6 in order to derive a formula for the aperiodic correlation if the binary sequence exhibits a stronger form of symmetry, which is for example the case if a is a Barker sequence of even length  $n = 2m \ge 4$  with  $C_1 = C_3 = \cdots = C_{m-1} = 1$ .

Suppose that a is a Barker sequence of even length  $n=2m\geq 4$  with  $C_1=C_3=\cdots=C_{m-1}=1$  and let  $1\leq k\leq m$ . By Proposition 4 we have  $a_ka_{n+1-k}=-1$  if k is odd, and  $a_ka_{n+1-k}=1$  if k is even. This motivates the following definition.

A binary sequence a of length n is called *strong symmetric* if the length n of a is a multiple of 4 and and  $a_i a_{n+1-j} = (-1)^j$  for each  $1 \le j \le \frac{n}{2}$ .

Note that if a is a strong symmetric binary sequence of length n=2m then a is weak symmetric and  $\delta_k=(-1)^k$  for each  $1\leq k\leq m$ , and  $\delta_k=(-1)^{k+1}$  for each  $m< k\leq n$  (recall that  $\delta_k$  is defined as  $\delta_k=a_ka_{n+1-k}$  for  $1\leq k\leq n$ ). Note further, that a strong symmetric binary sequence a of length n=2m is already determined by the first m entries  $a_1,a_2,\cdots,a_m$ .

Applying Lemma 5 and Lemma 6 to strong symmetric binary sequences gives us the following result.

**Lemma 7.** Let a be a strong symmetric binary sequence of length n = 2m. Denote by b be the binary sequence of length m with  $b_j = a_j$  for each  $1 \le j \le m$ . If k is even with  $1 \le k < n$ , then

$$C_k = \begin{cases} 2C_k(b) & \text{if } k < m \\ 0 & \text{if } k \ge m. \end{cases}$$

Furthermore, if k is odd with  $1 \le k < m$ , then following two statements hold:

(i)

$$C_{n-k} = (-1)^{\frac{k+1}{2}} + 2\sum_{j=1}^{\frac{k-1}{2}} (-1)^j b_j b_{k+1-j}.$$

(ii)

$$C_k = (-1)^{\frac{k-1}{2}} - 2\sum_{j=1}^{\frac{k-1}{2}} (-1)^j b_{j+m-k} b_{m+1-j}.$$

Note that  $C_k(b)$  denotes the kth aperiodic autocorrelation of the binary sequence b, whereas  $C_k$  as always refers to the sequence a, i.e.  $C_k = C_k(a)$ . Note further that the sequence b corresponds to the first half of the sequence a, so equations (i) and (ii) hold also, if we replace on the right hand side the entries of b with the corresponding entries of a.

Proof. Let  $1 \leq k < n$ . Suppose first that k is even. If  $k \geq m$  then  $C_k = 0$  follows directly from Lemma 5 (i), since in this case  $\delta_j + \delta_{n-k+1-j} = 0$  for each  $1 \leq j \leq \frac{n-k}{2}$ . For the case k < m we use Lemma 6 (i). Note that in this case  $1 + \delta_j \delta_{j+k} = 2$  for each  $1 \leq j \leq m-k$  and  $\delta_{j+m-k} + \delta_{m+1-j} = 0$  for each  $1 \leq j \leq \frac{k}{2}$ ; hence  $C_k = 2 \sum_{j=1}^{m-k} a_j a_{j+k} = 2C_k(b)$ . Let us suppose next that k is odd and  $1 \leq k < m$ . Then by Lemma 5 (ii) we have

Let us suppose next that k is odd and  $1 \le k < m$ . Then by Lemma 5 (ii) we have  $C_{n-k} = \delta_{\frac{k+1}{2}} + \sum_{j=1}^{\frac{k-1}{2}} a_j a_{k+1-j} (\delta_j + \delta_{k+1-j})$ . Note that in this case  $\delta_j + \delta_{k+1-j} = 2(-1)^j$  for all  $1 \le j \le \frac{k-1}{2}$  and that  $\delta_{\frac{k+1}{2}} = (-1)^{\frac{k+1}{2}}$ , which proves (i). In order to

prove (ii) use Lemma 6; for this note that  $\delta_{\frac{n-k+1}{2}} = (-1)^{\frac{k-1}{2}}$ ,  $1 + \delta_j \delta_{j+k} = 0$  for each  $1 \leq j \leq m-k$ , and since m is even we have  $\delta_{j+m-k} + \delta_{m+1-j} = -2(-1)^j$  for each  $1 \leq j \leq \frac{k-1}{2}$ .

Suppose that a and p are both strong symmetric binary sequences of length n=2m. If the first m entries of p are the same as the first m entries of a but in reversed order, then the aperiodic autocorrelations of a and p are strongly related; this shows the next lemma.

**Lemma 8.** Let a and p be strong symmetric binary sequences of length n = 2m and suppose that  $p_j = a_{m+1-j}$  for each  $1 \le j \le m$ . Then for each  $1 \le k < n$ 

$$C_k(p) = \begin{cases} C_k(a) & \text{if } k \text{ is even} \\ -C_{n-k}(a) & \text{if } k \text{ is odd.} \end{cases}$$

Proof. Let  $1 \le k < n$ , and as in Lemma 7 let b denote the binary sequence of length m with  $b_j = a_j$  for each  $1 \le j \le m$ . Consider first the case when k is even. Then by Lemma 7 we have  $C_k(a) = 0 = C_k(p)$  if  $k \ge m$ . Since reversing a binary sequence does not chance its aperiodic autocorrelations, it follows that  $C_k(a) = 2C_k(b) = C_k(p)$  if k < m.

Next consider the case when k is odd. If k < m then by Lemma 7 (ii) and (i) we have  $C_k(p) = (-1)^{\frac{k-1}{2}} - 2\sum_{j=1}^{\frac{k-1}{2}} (-1)^j a_{k+1-j} a_j$  and  $C_{n-k}(a) = (-1)^{\frac{k+1}{2}} + 2\sum_{j=1}^{\frac{k-1}{2}} (-1)^j a_j a_{k+1-j}$ . This shows that  $C_k(p) = -C_{n-k}(a)$  if k < m. The same argument gives us also that  $C_k(a) = -C_{n-k}(p)$  if k < m. Hence  $C_k(p) = -C_{n-k}(a)$  for all odd k satisfying  $1 \le k \le n$ .

Recall that if a is a Barker sequence of even length  $n \geq 4$  then n is a multiple of 4. We now consider the special case of a Barker sequence of even length n = 4u where the first u odd aperiodic autocorrelations all have the same value.

**Lemma 9.** Suppose that a is a Barker sequence of even length  $n = 2m \ge 4$  with  $C_1 = C_3 = \cdots = C_{m-1}$ , and let  $u = \frac{n}{4}$ . Then we have  $a_j a_{j+1} = a_{2j} a_{2j+1} = a_{u+j} a_{u+j+1}$  for each j satisfying  $1 \le j < u$ .

Proof. Recall that by Proposition 4 the Barker sequence a is weak symmetric. Consider first the case that  $C_1=C_3=\cdots=C_{m-1}=1$ , and let  $1\leq k < m$ . Then by Proposition 4 we have  $\delta_k=a_ka_{n+1-k}=-1$  if k is odd; since a is weak symmetric this implies that a is in fact strong symmetric. If k is odd then by (1.4) and Lemma 7 (i) we have  $-1=C_{n-k}=(-1)^{\frac{k+1}{2}}+2R_k$  with  $R_k=\sum_{j=1}^{\frac{k-1}{2}}(-1)^ja_ja_{k+1-j}$ . Depending on whether  $\frac{k+1}{2}$  is odd or even,  $R_k$  is either 0 or -1. In both cases Lemma 3 (ii) gives us that  $\prod_{j=1}^{\frac{k-1}{2}}a_ja_{k+1-j}=1$  and hence  $\prod_{j=1}^ka_j=a_{j+1}=1$  for each odd k with  $1\leq k < m$ . Similar as in [7] multiplying two successive equations of this latter form shows that

$$(4.1) a_j a_{j+1} = a_{2j} a_{2j+1} \text{ for each } j \text{ satisfying } 1 \le j < u.$$

Next consider the case  $C_1 = C_3 = \cdots = C_{m-1} = -1$ . Since the binary sequence s of length n defined by  $s_j = (-1)^j a_j$  is a Barker sequence with  $C_1(s) = C_3(s) = \cdots = C_{m-1}(s) = 1$  Equation (4.1) holds also in this case.

It remains to show that  $a_{2j}a_{2j+1} = a_{u+j}a_{u+j+1}$  for  $1 \leq j < u$ . As in Lemma 8 let p be the (well defined) strong symmetric binary sequence of length n with

 $p_j = a_{m+1-j}$  for each  $1 \le j \le m$ . Lemma 8 shows that p is also a Barker sequence of length n with  $C_1(p) = C_3(p) = \cdots = C_{m-1}(p) = -C_1$ . Thus by (4.1) we have  $a_{m+1-k}a_{m-k} = a_{m+1-2k}a_{m-2k}$  for  $1 \le k < u$ . Putting j = u - k this can be rewritten as  $a_{u+j+1}a_{u+j} = a_{2j+1}a_{2j}$  for  $1 \le j < u$ .

We are now in the position to prove Theorem 1.

Proof of Theorem 1. Suppose that a is a Barker sequence of even length  $n = 2m \ge 4$  with  $C_1 = C_3 = \cdots = C_{m-1}$ . Without loss of generality we can assume that  $C_1 = 1$ . Otherwise, instead of a consider the binary sequence s of length n defined by  $s_j = (-1)^j a_j$ ; then, as mentioned before, s is also a Barker sequence of length n with  $C_1(s) = C_3(s) = \cdots = C_{m-1}(s) = 1$ .

Put  $u = \frac{n}{4}$  and as in Lemma 7 let b denote the binary sequence of length m with  $b_j = a_j$  for each  $1 \le j \le m$ . Then by Lemma 9 we have  $a_j a_{j+u} = a_{j+1} a_{j+1+u}$  for all j satisfying  $1 \le j < u$  and hence  $|C_u(b)| = u \ne 0$ . Thus by Lemma 7 the integer u must be odd.

Now assume that m > 2. Then Lemma 7 gives us that  $C_2(b) = 0$  and  $C_{m-2}(b) = 0$ . But by (1.3) we have  $0 = C_2(b) + C_{m-2}(b) \equiv m \pmod{4}$  which implies that u is even, a contradiction. Hence m = 2.

## 5. Conclusion

We have shown that Barker sequences of even length  $n \geq 4$  exhibit certain symmetry properties. In order to exploit this symmetry, different formulas for the calculation of the aperiodic correlation have been derived. By using only elementary methods we have proven that the special case where  $C_1 = C_3 = \cdots = C_{\frac{n}{2}-1}$  is for Barker sequences of even length  $n \geq 4$  only possible if n = 4. It is an interesting open question whether a similar elementary approach can be used to further reduce the number of cases for which the Barker conjecture for even length can be proved.

# References

- Peter Borwein and Tamás Erdélyi. A note on barker polynomials. International Journal of Number Theory, 9(03):759-767, 2013.
- [2] Peter Borwein and Michael J Mossinghoff. Wieferich pairs and barker sequences, ii. LMS Journal of Computation and Mathematics, 17(1):24–32, 2014.
- [3] Jonathan Jedwab. What can be used instead of a barker sequence? Contemporary Mathematics, 461:153–178, 2008.
- [4] Ka Hin Leung and Bernhard Schmidt. The anti-field-descent method. Journal of Combinatorial Theory, Series A, 139:87–131, 2016.
- [5] Kai-Uwe Schmidt. Sequences with small correlation. Designs, Codes and Cryptography, 78(1):237–267, 2016.
- [6] Kai-Uwe Schmidt and Jürgen Willms. Barker sequences of odd length. Designs, Codes and Cryptography, 80(2):409–414, 2016.
- [7] R Turyn and J Storer. On binary sequences. Proceedings of the American Mathematical Society, 12(3):394–399, 1961.
- [8] Richard Turyn. Character sums and difference sets. Pacific Journal of Mathematics, 15(1):319–346, 1965.
- [9] Jürgen Willms. Counterexamples to Theorem 1 of Turyn's and Storer's paper 'On binary sequences'". arXiv preprint arXiv:1404.4833, 2014.

Email address: willms.juergen@fh-swf.de

Institut für Computer Science, Vision and Computational Intelligence, Fach-Hochschule Südwestfalen, Lindenstrasse 53, D-59872 Meschede, Germany