THE COHOMOLOGY INVARIANT FOR CLASS DIII TOPOLOGICAL INSULATORS

GIUSEPPE DE NITTIS AND KYONORI GOMI

ABSTRACT. This work concerns with the description of the topological phases of band insulators of class DIII by using the equivariant cohomology. The main result is the definition of a cohomology class for general systems of class DIII which generalizes the well-known \mathbb{Z}_2 -invariant given by the Teo-Kane formula in the one-dimension case. In the two-dimensional case this cohomology invariant allows a complete description of the strong and weak phases. The relation with the KR-theory, the Noether-Fredholm index and the classification of "Real" gerbes are also discussed.

MSC 2010: Primary: 14D21; Secondary: 57R22, 55N25, 19L64.

Keywords: Class DIII topological insulators, "Quaternionic" vector bundles, equivariant cohomology, KR-theory.

CONTENTS

1. Introduction	2			
Structure of the paper.				
Acknowledgements	5			
2. Construction of the \mathbb{Z}_2 -invariant	5			
2.1. Underlying "Quaternionic" structure	5			
2.2. The topological invariant associated to the sewing matrix	8			
2.3. The relation with the KR-theory	11			
3. The one-dimensional case	17			
3.1. The Teo-Kane formula	17			
3.2. The role as topological obstruction				
3.3. The role as an index	23			
3.4. The relation with the KR-theory				
3.5. The relation with the classification of "Real" gerbes				
4. The two-dimensional case				
4.1. Strong invariant: the Dirac case				
4.2. Weak-strong invariant: the Bloch case	31			
Appendix A. Equivariant cohomology and homotopy				
Appendix B. Strong equivalence of "Quaternionic" structures				
References	37			

Date: April 2, 2021.

1. Introduction

This work concerns the classification of the protected phases of topological insulators of class DIII from the point of view of the equivariant cohomology theory. In this sense this work complements the research carried out by the same authors for classes AI, AII and AIII [DG1, DG2, DG3, DG4, DG6]. The literature on *topological insulators* is too vast to be reported here without crucial omissions. For this reason, and for the benefit of the reader, we will refer directly to the recent review [CTSR] which provides a global overview of the subject accompanied by a rich list of references. The works [RSFL, TK, BA], among others, deserve to be mentioned for the specific focus on systems of class DIII.

In order to present the logic and the main results of this work we need first to introduce some basic definition. An *involutive space* is a pair (X,τ) made of a compact Hausdorff space X endowed with a continuous *involution* $\tau: X \to X$, $\tau \circ \tau = \mathrm{Id}_X$. The *fixed point set* of (X,τ) is defined as $X^\tau := \{x \in X \mid \tau(x) = x\}$. A special example is provided by the pair (\mathbb{S}^1,ι) made by the unit circle $\mathbb{S}^1 := \mathbb{R}/2\pi\mathbb{Z}$, parametrized by $k \in [0,2\pi)$, and endowed with the involution ι defined by $\iota(k) := -k$. In this specific case the fixed-point set is $(\mathbb{S}^1)^\iota = \{0,\pi\}$.

Let \mathcal{H} be a finite dimensional Hilbert space, $\mathfrak{Herm}(\mathcal{H})$ the set of self-adjoint (or Hermitian) operators on \mathcal{H} and $\mathfrak{Herm}(\mathcal{H})^{\times}$ the subset of invertible elements of $\mathfrak{Herm}(\mathcal{H})$. Let $T: \mathcal{H} \to \mathcal{H}$ and $C: \mathcal{H} \to \mathcal{H}$ be two anti-unitary maps that meet the conditions

$$T^2 = -\mathbf{1}, \qquad C^2 = +\mathbf{1}, \qquad TC = -CT$$
 (1.1)

where $\mathbf{1}$ is the identity operator on \mathcal{H} . In the theory of topological insulators it is common to refer to T as *time-reversal symmetry* (TSR) and to C as *particle-hole symmetry* (PHS). The conditions on the square say that T is an *odd* symmetry while C is an *even* symmetry. The anti-commutation condition corresponds to fixing an arbitrary relative phase between T and C. With this convention, the linear operator $\chi := TC = -CT$, called *chiral symmetry*, satisfies

$$\chi^2 = \mathbf{1}$$
, $T\chi = -\chi T$, $C\chi = -\chi C$. (1.2)

The next definition is borrowed from the standard jargon for topological insulators [AZ, SRFL, Kit, RSFL].

Definition 1.1 (Band insulator of class DIII). A band insulator of class DIII over the involutive space (X, τ) is a continuous map

$$H: X \longrightarrow \mathfrak{Herm}(\mathcal{H})^{\times}$$

such that

$$\begin{cases} C H(x) = -H(\tau(x)) C \\ T H(x) = H(\tau(x)) T \end{cases}, \forall x \in X.$$
 (1.3)

Two band insulators of class DIII H and H' over the same involutive space (X, τ) and subjected to the same system of symmetries T, C are *unitarily equivalent* if and only if there exists a continuous map $V: X \to \mathbb{U}(\mathcal{H})$ such that

$$\begin{cases} H'(x) = V(x) H(x) V(x)^* \\ C V(x) = V(\tau(x)) C \\ T V(x) = V(\tau(x)) T \end{cases}, \forall x \in X.$$
 (1.4)

The unitary equivalence is called *strong* if and only if $V(\tau(x)) = V(x)$ for all $x \in X$. When the involutive space is explicitly given by (\mathbb{S}^1, ι) , then one refers to a *one-dimensional* (1D) band insulators of class DIII.

Definition 1.1 along with the relations (1.2) imply

$$\chi H(x) = -H(x) \chi, \quad \forall x \in X.$$
 (1.5)

An immediate consequence of this relation is that the dimension of the space \mathcal{H} must be even, *i. e.* dim(\mathcal{H}) = 2m for some $\mathfrak{m} \in \mathbb{N}$.

Let us focus for the moment on the one-dimensional case. Let H and H' be two 1-dimensional band insulators of class DIII and consider the *spectrally flattened* band operators $Q_H := H|H|^{-1}$ and $Q_{H'} := H'|H'|^{-1}$. It is known that the triple ($\mathbb{S}^1 \times \mathcal{H}$, Q_H , $Q_{H'}$) defines an element of the KR-group $KR^{-3}(\mathbb{S}^1, \iota) \simeq \mathbb{Z}_2$ [Ati, Kit, FM] (see also Section 3.4). Thus, there exists a \mathbb{Z}_2 -invariant which classifies (the difference of) 1-dimensional band insulators of class DIII. The aim of this work is to interpret this invariant from a cohomology point of view, and possibly to generalize it to more generic situations.

The key (not new) observation is that as a consequence of the commutation relations between H and the symmetries T and C, the topological information carried by H is completely encoded into a continuous map $q_H: \mathbb{S}^1 \to \mathbb{U}(m)$ such that

$$q_{\mathsf{H}}(\iota(k)) = -q_{\mathsf{H}}(k)^{\mathsf{t}}, \quad \forall k \in \mathbb{S}^{1},$$
 (1.6)

where the superscript t denotes the *transpose* of the matrix $q_H(k)$. There are at least two immediate ways in which the matrix q_H can produce a topological invariant.

(I) The map q_H can be interpreted as the *sewing matrix* of a "Quaternionic" vector bundle [Kah, Dup, DG2, DG3, DG4]. More precisely, the map q_H endows the product bundle $\mathbb{S}^1 \times \mathbb{C}^m$ with the structure of a "Quaternionic" vector bundle \mathscr{E}_{q_H} explicitly given by

$$(k, v) \longmapsto (\iota(k), q_H(k) \cdot \overline{v}) , \qquad \forall (k, v) \in \mathbb{S}^1 \times \mathbb{C}^m .$$
 (1.7)

Unfortunately, it is known that "Quaternionic" vector bundles on (\mathbb{S}^1, ι) are automatically trivial [DG2, Theorem 1.2]. Therefore, this is not this vector bundle the good candidate to detect the topology of 1-dimensional insulators of class DIII.

(II) In the presence of a fixed point $k = \iota(k)$, the rank m of $\mathbb{U}(m)$ is forced to be even, *i. e.* m = 2n. In turn, the determinant of q_H provides a map $\det[q_H] : \mathbb{S}^1 \to \mathbb{U}(1)$ which is *invariant* in the sense that

$$det[q_H(\iota(k))] = det[q_H(k)], \quad \forall k \in \mathbb{S}^1.$$

Since the homotopy classes of invariant maps on (\mathbb{S}^1, ι) are classified by the group $H^1_{\mathbb{Z}_2}(\mathbb{S}^1, \mathbb{Z})$ (cf. Appendix A), one gets the topological invariant

$$\left[\, \text{det}[\mathfrak{q}_H] \right] \; \in \; H^1_{\mathbb{Z}_2}(\mathbb{S}^1,\mathbb{Z})$$

associated to H. However, also in this case the invariant turns out to be trivial since $H^1_{\mathbb{Z}_2}(\mathbb{S}^1,\mathbb{Z})=0$ [DG1, Lemma 5.6]. The implication of this fact will be discussed in Remark 2.14.

Thus, to detect the class of $KR^{-3}(\mathbb{S}^1, \iota)$ associated to a one-dimensional band insulator of class DIII we need to extract from q_H some other type of topological invariant finer than (I) and (II). It turns out that a closer inspection to the triviality of the "Quaternionic" vector bundle \mathscr{E}_{q_H} allows to build a suitable non-trivial $\mathbb{Z}/2$ -invariant. The principal result of this work is the construction of such a topological invariant for band insulators of class DIII over a large class of involutive spaces (X, τ) .

More concretely, given a sewing matrix $q: X \to \mathbb{U}(2n)$ on the involutive space (X, τ) subject to certain conditions (*cf.* Assumption 2.5), one can associate to q a homotopy invariant

$$\nu_{q} \in H^{1}(X, \mathbb{Z})/H^{1}_{\mathbb{Z}_{2}}(X|X^{\tau}, \mathbb{Z}(1))$$
 (1.8)

In formula (1.8), $H^1(X, \mathbb{Z})$ is the first cohomology group of X with integer coefficients while $H^1_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))$ is the first equivariant cohomology group X with coefficients in the local system $\mathbb{Z}(1)$ relative to the fixed point set X^τ [DG2, Section 3.1]. In the case of the involutive circle (\mathbb{S}^1,ι) , one has that

$$H^1(\mathbb{S}^1,\mathbb{Z}) \simeq \mathbb{Z} \simeq H^1_{\mathbb{Z}_2}(\mathbb{S}^1|(\mathbb{S}^1)^{\iota},\mathbb{Z}(1)),$$

but the quotient group in (1.8) turns out to be isomorphic to \mathbb{Z}_2 (cf. Lemma 3.1), showing that ν_q is a \mathbb{Z}_2 -invariant as expected.

It is worth to point out, however, that the invariant ν_q of the sewing matrix q cannot serve as an *absolute* invariant for band insulators of class DIII. This is a consequence of the fact that the way to identify the symmetries with a fixed *standard form* is not unique, and can be altered by an automorphism (*cf.* Proposition 2.1). Accordingly, the sewing matrix associated with H is not unique, and if q_H and q'_H are two sewing matrices associated with the same H one can have $\nu_{q_H} \neq \nu_{q'_H}$ in general. In this sense, ν_{q_H} does not produce an invariant of the Hamiltonian H. However, if one fixes a way to identify the symmetries with the standard form, then the sewing matrix can be used to detect the difference of two band insulators H and H' of class DIII. In this sense the sewing matrix can be used to construct a *relative* invariant which provides a well-defined non-trivial

homomorphism

$$\mathsf{KR}^{-3}(X,\tau) \ \supseteq \ \mathsf{Ker}(\kappa) \ \stackrel{\nu}{\longrightarrow} \ \mathsf{H}^1(X,\mathbb{Z})/\mathsf{H}^1_{\mathbb{Z}_2}\big(X,X^\tau,\mathbb{Z}(1)\big) \tag{1.9}$$

(cf. Theorem 2.11 for more details). In the one-dimensional case (\mathbb{S}^1, ι) this construction provides a bijection $\nu : KR^{-3}(\mathbb{S}^1, \iota) \simeq \mathbb{Z}_2$ (Proposition 3.15) and the \mathbb{Z}_2 -value of the class ν_q associated with the sewing matrix q can be computed with the well known *Teo-Kane formula* (Theorem 3.5) or as the \mathbb{Z}_2 -index of an associated Toeplitz operator (Theorem 3.12).

Structure of the paper. In Section 2, we start by fixing the standard form of the symmetries of a band insulator of class DIII. Then, we introduce the invariant ν_q of a sewing matrix q in a general setup. The construction of the homomorphism ν in equation (1.9) and its relation with the KR-theory are also provided in this section. Section 3 is devoted to the study of the one-dimensional case. In this section we provide the equivalence of ν_q with the well-known Teo-Kane formula [CTSR, eq. (3.70)], we clarify the role of v_q as an obstruction class and we establish a bulk-edge-type correspondence as the coincidence of ν_q with a \mathbb{Z}_2 -index for Toeplitz operators. Finally, we will show that the homomorphism (1.9) amounts to an isomorphism in the one-dimensional case and we will shed some light on the role of this invariant in the classification of "Real" gerbes over (\mathbb{S}^1, ι) . In **Section** 4 we investigate the application of the invariant v_q for two-dimensional band insulators of class DIII over the sphere (Dirac case) and the torus (Bloch case). It turns out that ν_q provides a complete description of the weak topological invariants in the two-dimensional case. Appendix A contains a brief summary of the relationship between cohomology and homotopy in the \mathbb{Z}_2 -equivariant category and **Appendix B** deals with the notion of strong equivalence between "Quaternionic" structures.

Acknowledgements. GD's research is supported by the grant Fondecyt Regular - 1190204. KG's research is supported by JSPS KAKENHI Grant Numbers 20K03606 & JP17H06461.

2. Construction of the \mathbb{Z}_2 -invariant

2.1. **Underlying "Quaternionic" structure.** In view of condition (1.5), and the invertibility of H(x), we argued that band insulators of class DIII can be seated only in Hilbert spaces $\mathcal H$ of even dimension. Moreover, the eigenspaces of the symmetry χ , determined by the range of the eigenprojections $\chi_{\pm} := \frac{1}{2}(1 \pm \chi)$, must have same dimension. As a consequence there exists a (non unique) unitary isomorphism $\mathcal H \simeq \mathbb C^{2m}$ which provides the matrix representation

$$\chi = \begin{pmatrix} +\mathbf{1}_{m} & 0\\ 0 & -\mathbf{1}_{m} \end{pmatrix} \tag{2.1}$$

where $\mathbf{1}_m$ is the $m \times m$ identity matrix. For the next result we need to introduce the operator K which implements the standard real structure on \mathbb{C}^m . This is nothing more than the complex conjugation $Kv := \overline{v}$ for every $v \in \mathbb{C}^m$.

Proposition 2.1. (Standard representation) Let H, T, C be the elements which describe a band insulator of class DIII over the involutive space (X, τ) , according to Definition 1.1. Then, there exists a (non unique) unitary isomorphism $\mathcal{H} \simeq \mathbb{C}^{2m}$ which provides the matrix representation

$$T = \begin{pmatrix} 0 & -K \\ +K & 0 \end{pmatrix}, \qquad C = \begin{pmatrix} 0 & -K \\ -K & 0 \end{pmatrix}. \tag{2.2}$$

Moreover, the Hamiltonian $H: X \to \mathfrak{Herm}(\mathcal{H})^{\times}$ *is represented by*

$$H(x) = \begin{pmatrix} 0 & h(x)^{-1} \\ h(x) & 0 \end{pmatrix}, \qquad (2.3)$$

where the map $h: X \to \mathbb{GL}(m)$ meets the condition

$$h(\tau(x)) = -K h(x)^{-1} K, \qquad \forall x \in X.$$
 (2.4)

Proof. Under the unitary transformation $\mathcal{H}\simeq\mathbb{C}^{2m}$ which provides the representation (2.1), the operator T takes the form

$$T = U_T \begin{pmatrix} K & 0 \\ 0 & K \end{pmatrix}$$

where U_T is a unitary operator. Since the anti-linear part induced by K commutes with χ , it follows that the second constraint in (1.2) can be satisfied if and only if $\chi U_T = -U_T \chi$. However, this forces U_T to be off-diagonal with respect to the diagonal representation of χ , and in turn

$$T = \begin{pmatrix} 0 & \widetilde{U}_T K \\ \widetilde{U}_T^* K & 0 \end{pmatrix}$$

with $\widetilde{U}_T \in \mathbb{U}(m)$ a given $m \times m$ unitary matrix. The condition $T^2 = -\mathbf{1}$ implies that $K\widetilde{U}_T K = -\widetilde{U}_T$. Therefore, one gets

$$\begin{pmatrix} \mathbf{1}_{m} & \mathbf{0} \\ \mathbf{0} & \widetilde{\mathbf{U}}_{T} \end{pmatrix} \begin{pmatrix} \mathbf{0} & \widetilde{\mathbf{U}}_{T} \mathbf{K} \\ \widetilde{\mathbf{U}}_{T}^{*} \mathbf{K} & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{1}_{m} & \mathbf{0} \\ \mathbf{0} & \widetilde{\mathbf{U}}_{T}^{*} \end{pmatrix} = \begin{pmatrix} \mathbf{0} & -\mathbf{K} \\ +\mathbf{K} & \mathbf{0} \end{pmatrix}$$

and at the same time

$$\left(\begin{array}{cc} \mathbf{1}_{\mathfrak{m}} & \mathbf{0} \\ \mathbf{0} & \widetilde{\mathbf{U}}_{\mathsf{T}} \end{array} \right) \chi \left(\begin{array}{cc} \mathbf{1}_{\mathfrak{m}} & \mathbf{0} \\ \mathbf{0} & \widetilde{\mathbf{U}}_{\mathsf{T}}^* \end{array} \right) \; = \; \chi \; .$$

Then, up to a modification of the initial unitary transform $\mathcal{H}\simeq\mathbb{C}^{2m}$, one can represent T as in (2.2) without affecting the representation (2.1). The representation of C in (2.2) is obtained by the formula $C=\chi T$. The anti-commutation relation between H(x) and χ implies that H(x) must be off-diagonal. The invertibility of H(x) for every $x\in X$ justifies the representation (2.3). Condition (2.4) follows from the first of (1.3).

A straightforward adaptation of the arguments used in the proof of Proposition 2.1 provides the following result that will be used several times in the following.

Lemma 2.2. Let $V \in \mathbb{U}(2n)$ be a unitary matrix which commutes with χ and T as given by the representations (2.1) and (2.2), respectively. Then there exists a unitary matrix $\varphi_V \in \mathbb{U}(m)$ such that

$$V \,=\, \left(egin{array}{cc} \varphi_V & 0 \ 0 & \overline{\varphi_V} \end{array}
ight) \,, \qquad \overline{\varphi_V} \,:=\, K \varphi_V K \,.$$

Given the invertible matrix $H \in \mathfrak{Herm}(\mathcal{H})^{\times}$ one can define the *flattened* operator

$$Q_H := \frac{H}{|H|}, \qquad |H| := \sqrt{H^*H}.$$
 (2.5)

Evidently $Q_H \in \mathbb{U}(\mathcal{H}) \cap \mathfrak{Herm}(\mathcal{H})^{\times}$ is a unitary and self-adjoint operator, namely an involution $Q_H^2 = \mathbf{1}$. If $H: X \to \mathfrak{Herm}(\mathcal{H})^{\times}$ is a band insulator of class DIII, then $Q_H: X \to \mathfrak{Herm}(\mathcal{H})^{\times}$ is also a band insulator of class DIII, and Proposition 2.1 applies. In particular one has that

$$Q_{H}(x) = \begin{pmatrix} 0 & q_{H}(x)^{-1} \\ q_{H}(x) & 0 \end{pmatrix} = \begin{pmatrix} 0 & q_{H}(x)^{*} \\ q_{H}(x) & 0 \end{pmatrix}, \qquad (2.6)$$

where the map $q_H: X \to \mathbb{U}(n)$ meets the condition

$$q_{H}(\tau(x)) = -Kq_{H}(x)^{*}K = -q_{H}(x)^{t}.$$
 (2.7)

The map q_H allows to endow the trivial product bundle $X \times \mathbb{C}^m$ with the extra structure $\Theta_H: X \times \mathbb{C}^m \to X \times \mathbb{C}^m$ defined by

$$\Theta_{\mathsf{H}} : (x, v) \longmapsto (\tau(x), q_{\mathsf{H}}(k) \cdot \overline{v}) , \qquad \forall (x, v) \in X \times \mathbb{C}^{\mathsf{m}} .$$
 (2.8)

It turns out that $(X \times \mathbb{C}^m, \Theta_H)$ is a "Quaternionic" vector bundle in the sense of [Kah, Dup, DG2, DG3, DG4].

Definition 2.3 (Sewing matrix). Let $H: X \to \mathfrak{Herm}(\mathcal{H})^{\times}$ be a band insulator of class DIII over the involutive space (X, τ) according to Definition 1.1. Then $(X \times \mathbb{C}^m, \Theta_H)$ will be called the *associated* "Quaternionic" vector bundle, and q_H the *sewing matrix* of the associated "Quaternionic" structure.

It is known that the existence of fixed points of the involutive space (X, τ) implies that the rank of the "Quaternionic" vector bundles over (X, τ) must be even [DG2, Proposition 2.1]. This fact has an immediate consequence:

Proposition 2.4. Let $H: X \to \mathfrak{Herm}(\mathfrak{H})^{\times}$ be a band insulator of class DIII over the involutive space (X,τ) , according to Definition 1.1. Assume that X is path-connected and $X^{\tau} \neq \emptyset$. Then the associated "Quaternionic" vector bundle has even rank $\mathfrak{m}=2\mathfrak{n}$ and $\mathfrak{q}_H: X \to \mathbb{U}(2\mathfrak{n})$. Moreover, the standard representation described in Proposition 2.1 is induced by a unitary isomorphism $\mathfrak{H} \simeq \mathbb{C}^{4\mathfrak{n}}$.

2.2. The topological invariant associated to the sewing matrix. In order to define a sufficiently good topological invariant for band insulators of class DIII we need to establish some working hypotheses.

Assumption 2.5. Let (X, τ) be an involution space and $q: X \to \mathbb{U}(2n)$ a sewing matrix, *i. e.* a map such that

$$q\big(\tau(x)\big) \; = \; -q(x)^{\mathtt{t}} \; , \qquad \forall \; x \in X \; .$$

Let $\mathscr{E}_q = X \times \mathbb{C}^{2n}$ be the "Quaternionic" vector bundle with "Quaternionic" Θ_q structure induced by q, i. e.

$$\Theta_{q}: (x,v) \longmapsto (\tau(x),q(k)\cdot \overline{v}), \qquad (x,v) \in X \times \mathbb{C}^{2n}.$$

We will assume that:

- (a) X is a path connected \mathbb{Z}_2 -CW complex;
- (b) The fixed point set is non-empty, *i. e.* $X^{\tau} \neq \emptyset$;
- (c) The FKMM invariant $\kappa(\mathscr{E}_q) \in H^2_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))$ is trivial.

For the definition and the properties of the *FKMM invariant* for "Quaternionic" vector bundles we refer to [DG2, DG3, DG4]. For more details on the relative cohomology groups $H^{\bullet}_{\mathbb{Z}_2}(X|X^{\tau},\mathbb{Z}(1))$ we will refer to [DG2, Section 3.1] and references therein. It is worth recalling that these groups fit into the long exact sequence for the pair (X, X^{τ}) . The relevant part of the latter for the aims of this work reads

where the first term of the sequence corresponds to $H^0_{\mathbb{Z}_2}(X^\tau,\mathbb{Z}(1))=0$ [Gom]. As a matter of fact, the image of $\kappa(\mathscr{E}_q)$ under $H^2_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))\to H^2_{\mathbb{Z}_2}(X,\mathbb{Z}(1))$ is the "Real" Chern class $c_1^R(\det(\mathscr{E}_q))$ that classifies the determinant "Real" line bundle $\det(\mathscr{E}_q)$ [Kah, DG1]. Therefore, $\kappa(\mathscr{E}_q)=0$ implies that the determinant line bundle $\det(\mathscr{E}_q)$ is trivial as "Real" line bundle. Let us also recall the isomorphisms $H^1(X,\mathbb{Z})\simeq [X,\mathbb{U}(1)]$ and $H^1_{\mathbb{Z}_2}(X,\mathbb{Z}(1))\simeq [X,\mathbb{U}(1)]_{\mathbb{Z}_2}$, where the (standard) involution on $\mathbb{U}(1)$ is given by the complex conjugation $z\mapsto \overline{z}$ [Gom, Proposition A.2].

For the next result we need to introduce some notation. Let $C(X, \mathbb{U}(1))$ be the set of continuous functions from X to $\mathbb{U}(1)$ and $q: X \to \mathbb{U}(2n)$ a sewing matrix as in Assumption 2.5. When $x \in X^{\tau}$ is a fixed point under the involution, then $q(x) = -q(x)^{t}$ is a skew-symmetric matrix and one can calculate the associated *Pfaffian* $Pf[q(x)] \in \mathbb{U}(1)$. Let us define the subset

$$C(X,\mathbb{U}(1))_q \; := \; \left\{ p: X \to \mathbb{U}(1) \left| \begin{array}{cc} det[q(x)] = p(\tau(x))p(x) & \forall \; x \in X, \\ p(x) = Pf[q(x)] & \forall \; x \in X^\tau \end{array} \right\} \; .$$

Under the isomorphism $H^1(X, \mathbb{Z}) \simeq [X, \mathbb{U}(1)]$ the subset $C(X, \mathbb{U}(1))_q$ identifies a subset $H^1(X, \mathbb{Z})_q \subset H^1(X, \mathbb{Z})$ which coincides with the set of homotopy classes of maps in $C(X, \mathbb{U}(1))_q$.

Lemma 2.6. *Under Assumption* **2.5**, *the following facts hold true:*

- (1) The set $C(X, \mathbb{U}(1))_q$ is non-empty, and is a torsor under the group of \mathbb{Z}_2 -equivariant maps $r: X \to \mathbb{U}(1)$ which take the value 1 on the fixed point set X^{τ} .
- (2) The set $H^1(X,\mathbb{Z})_q$ is a torsor under the group $H^1_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))$ with action induced by

$$H^1_{\mathbb{Z}_2}\big(X|X^\tau,\mathbb{Z}(1)\big) \ \to \ H^1_{\mathbb{Z}_2}\big(X,\mathbb{Z}(1)\big) \ \to \ H^1(X,\mathbb{Z}) \ \supset \ H^1(X,\mathbb{Z})_q \ .$$

Proof. Let us start with (1). As pointed out above, the triviality of the FKMM invariant $\kappa(\mathscr{E}_q)$ implies the triviality of the determinant "Real" line bundle $\det(\mathscr{E}_q)$. Since the information on $\det(\mathscr{E}_q)$ is completely encoded in $\det[q]:X\to \mathbb{U}(1)$, from the triviality of $\det(\mathscr{E}_q)$ one infers the existence of the a global trivialization $p:X\to \mathbb{U}(1)$ which intertwines between the "Real" structure provided by $\det[q]$ and the *trivial* (or constant) "Real" structure. The latter fact is equivalent to the constraint $\det[q(x)] = p(\tau(x))p(x)$ for all $x \in X$. From [DG3, Proposition 2.10] one gets that the FKMM invariant $\kappa(\mathscr{E}_q) \in H^2_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))$ coincides with the injective image under

$$H^1_{\mathbb{Z}_2}\big(X^\tau,\mathbb{Z}(1)\big)/H^1_{\mathbb{Z}_2}\big(X,\mathbb{Z}(1)\big) \ \longrightarrow \ H^2_{\mathbb{Z}_2}\big(X|X^\tau,\mathbb{Z}(1)\big)$$

of the element whose representative is given by the map $X^{\tau} \ni x \mapsto p(x)/Pf[q(x)] \in \mathbb{Z}_2$ where p is the global trivialization introduced above. Assumption 2.5 (c) says that $\kappa(\mathscr{E}_q)$ is trivial. Hence, one can choose p so that p(x) = Pf[q(x)] for every $x \in X^{\tau}$. This proves that the set $C(X, \mathbb{U}(1))_q$ is non-empty. For a given pair $p', p \in C(X, \mathbb{U}(1))_q$, let $r: X \to \mathbb{U}(1)$ be the map $r(x) := p'(x)\overline{p(x)}$. It turns out that $r(\tau(x)) = \overline{r(x)}$ for every $x \in X$ and r(x) = 1 when $x \in X^{\tau}$. Hence $C(X, \mathbb{U}(1))_q$ is a torsor as stated. For (2) it is enough to consider the the homotopy classes of maps in (1).

Let us recall that by definition a torsor coincides with the orbit generated by any of its points. We are now in position to introduce the main topological invariant of this work.

Definition 2.7 (DIII-invariant). Under Assumption 2.5, one defines ν_q to be the orbit of the subset $H^1(X,\mathbb{Z})_q \subset H^1(X,\mathbb{Z})$ under $H^1_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))$, *i. e.*

$$\nu_q \; := \; \left[H^1(X,\mathbb{Z})_q \right] \; \in \; H^1(X,\mathbb{Z}) / H^1_{\mathbb{Z}_2} \big(X | X^\tau, \mathbb{Z}(1) \big) \; .$$

Let us point out that the notation used in Definition 2.7 is a little redundant since $[H^1(X,\mathbb{Z})_q]$ and $H^1(X,\mathbb{Z})_q$ coincide as a set in view of the torsor property. Anyway, the notation proposed as the advantage of emphasizing that ν_q is an equivalence class (or an orbit). The topological properties of ν_q are specified in the following result.

Theorem 2.8. Under Assumption 2.5 the following facts hold true:

- (1) v_q is an invariant of the homotopy class of q.
- (2) Let $q_0: X \to \mathbb{U}(2n)$ be the constant map given by

$$q_0(x) \; = \; Q \; := \; \left(\begin{array}{cc} 0 & -\mathbf{1}_n \\ +\mathbf{1}_n & 0 \end{array} \right) \; , \label{eq:q0}$$

then $v_{q_0} = 0$.

(3) Let $q: X \to \mathbb{U}(2n)$ and $q': X \to \mathbb{U}(2n')$ be as in Assumption 2.5, and consider $q \oplus q': X \to \mathbb{U}(2n+2n')$. Then

$$\nu_{\mathfrak{q}\oplus\mathfrak{q}'}\,=\,\nu_{\mathfrak{q}}\,+\,\nu_{\mathfrak{q}'}$$

as elements of the quotient group $H^1(X,\mathbb{Z})/H^1_{\mathbb{Z}_2}(X|X^{\tau},\mathbb{Z}(1))$.

(4) Let $q:X\to \mathbb{U}(2n)$ and $q':X\to \mathbb{U}(2n)$ be related by

$$q'(x) = h(\tau(x))^t q(x) h(x)$$

for a given map $h: X \to \mathbb{U}(2n)$ *. Then,*

$$\nu_{\mathfrak{q}'} \,=\, det[h] \cdot \nu_{\mathfrak{q}} \,=\, \left[\, det[h] \cdot H^1(X,\mathbb{Z})_{\mathfrak{q}} \right] \,.$$

Proof. (1) Let q and q' be sewing matrices satisfying Assumption 2.5. Suppose that these two matrices are connected by a homotopy $\tilde{q}: X \times [0,1] \to \mathbb{U}(2n)$ of sewing matrices. The map \tilde{q} defines the "Quaternionic" vector bundles $\mathcal{E}_{\tilde{q}}$ over $X \times [0,1]$. By the homotopy property of "Quaternionic" vector bundles [DG2, Theorem 2.3], $\mathcal{E}_{\tilde{q}}$ is isomorphic to the pull-back of \mathcal{E}_{q} under the projection $X \times [0,1] \to X$. Hence, both the determinant "Real" line bundle of $\mathcal{E}_{\tilde{q}}$ and the FKMM invariant of $\mathcal{E}_{\tilde{q}}$ are trivial. As a result, $C(X \times [0,1],\mathbb{U}(1))_{\tilde{q}}$ is non-empty as a consequence of Lemma 2.6. Let us choose a $\tilde{p} \in C(X \times [0,1],\mathbb{U}(1))_{\tilde{q}}$. Such a \tilde{p} provides a homotopy between the representatives $p := \tilde{p}|_{X \times \{0\}}$ and $p' := \tilde{p}|_{X \times \{1\}}$ of v_q and $v_{q'}$, respectively. Therefore, $v_q = v_{q'}$.

- (2) One can take $p_0 \in C(X, \mathbb{U}(1))_q$ to be the constant map $p_0(x) = 1$. This map represents the trivial element $0 \in H^1(X, \mathbb{Z})$, so that $\nu_{q_0} = 0$.
- (3) If $p \in C(X, \mathbb{U}(1))_q$ and $p' \in C(X, \mathbb{U}(1))_{q'}$, then the map $pp' : X \to \mathbb{U}(1)$ given by

$$(pp')(x) \,:=\, p(x)p'(x)$$

belongs to $C(X,\mathbb{U}(1))_{q\oplus q'}$. In the quotient group $H^1(X,\mathbb{Z})/H^1_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))$, the invariants ν_q and $\nu_{q'}$ are represented by the homotopy classes $[p]\in H^1(X,\mathbb{Z})$ and $[p']\in H^1(X,\mathbb{Z})$, respectively. One has that $[pp']=[p]+[p']\in H^1(X,\mathbb{Z})$, so that $\nu_{q\oplus q'}=\nu_q+\nu_{q'}$.

(4) If $p \in C(X, \mathbb{U}(1))_q$, then one gets $p' \in C(X, \mathbb{U}(1))_{q'}$ by setting

$$p'(x) := \det[h(x)]p(x).$$

Thus, the multiplication by det[h] induces a transformation on $H^1(X, \mathbb{Z})$ which carries $H^1(X, \mathbb{Z})_q$ to $H^1(X, \mathbb{Z})_{q'}$. As a consequence, $\nu_q \mapsto \nu_{q'}$ in the quotient group.

Remark 2.9. Let us recall that the for the equivariant cohomology the concept of reduced cohomology makes sense (see *e. g.* [DG1, Section 5.1]). Therefore, one has the usual decomposition

$$H^1_{\mathbb{Z}_2}\big(X,\mathbb{Z}(1)\big) \ \simeq \ \tilde{H}^1_{\mathbb{Z}_2}\big(X,\mathbb{Z}(1)\big) \ \oplus \ H^1_{\mathbb{Z}_2}\big(\{*\},\mathbb{Z}(1)\big),$$

where $\tilde{H}^1_{\mathbb{Z}_2}(X,\mathbb{Z}(1))$ is the reduced cohomology group and $H^1_{\mathbb{Z}_2}(\{*\},\mathbb{Z}(1))\simeq \mathbb{Z}_2$. By means of a spectral sequence, one can show

$$\tilde{H}^1_{\mathbb{Z}_2}\big(X,\mathbb{Z}(1)\big) \; \simeq \; H^1(X,\mathbb{Z})^{\mathbb{Z}_2}_- \; := \; \{f \in H^1(X;\mathbb{Z}) | \, \tau^*(f) = -f\}.$$

In particular, $\tilde{H}^1_{\mathbb{Z}_2}(X,\mathbb{Z}(1))$ turns out to be a free abelian group. Since the generator of $H^1_{\mathbb{Z}_2}(\{*\},\mathbb{Z}(1)) \subset H^1_{\mathbb{Z}_2}(X,\mathbb{Z}(1))$ can be represented by the constant map $X \ni x \mapsto -1 \in \mathbb{U}(1)$, it follows from the exact sequence

$$0 \to H^1_{\mathbb{Z}_2}\big(X|X^\tau,\mathbb{Z}(1)\big) \to H^1_{\mathbb{Z}_2}\big(X,\mathbb{Z}(1)\big) = \tilde{H}^1_{\mathbb{Z}_2}\big(X,\mathbb{Z}(1)\big) \oplus \mathbb{Z}_2$$
 that $H^1_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))$ injects into the free part $\tilde{H}^1_{\mathbb{Z}_2}(X,\mathbb{Z}(1))$.

2.3. The relation with the KR-theory. The KR-group $KR^{-3}(X,\tau)$ of a space X with involution τ may be regarded as a group which classifies the "difference" of two band insulators of class DIII. In this section we will relate $KR^{-3}(X,\tau)$ with the invariant ν_q of a sewing matrix q, provided that certain assumptions are satisfied.

Let us start by recalling the formulation of the *twisted equivariant K-theory* [FM] (see also [DG5, Section 5] and [Gom2]). Let \mathbb{G} be a finite group acting on a space X endowed with two homomorphisms $\phi: \mathbb{G} \to \mathbb{Z}_2$ and $c: \mathbb{G} \to \mathbb{Z}_2$ and a group 2-cocycle $\sigma: \mathbb{G} \times \mathbb{G} \times X \to \mathbb{U}(1)$, which satisfies

$$\sigma(g_2,g_3,x)^{\varphi(g_1)} \; \sigma(g_1g_2,g_3,x)^{-1} \; \sigma(g_1,g_2g_3,x) \; \sigma(g_1,g_2,g_3(x))^{-1} \; = \; 1$$

for all $g_1, g_2, g_3 \in \mathbb{G}$ and all $x \in X$. A (φ, σ, c) -twisted (ungraded) vector bundle with a $\text{Cl}_{p,q}$ -action is a (finite-rank) Hermitian vector bundle $\pi : \mathscr{E} \to X$ equipped with:

- (a) An isometric bundle map $\rho(g):\mathscr{E}\to\mathscr{E}$ covering the action of $g\in\mathbb{G}$ on X for each $g\in\mathbb{G}$;
- (b) Unitary maps $\gamma_1, \ldots, \gamma_{p+q}$ on \mathscr{E} , which satisfy the Clifford relations

$$\gamma_i\gamma_j+\gamma_j\gamma_i \;=\; \left\{ \begin{array}{ll} 2 & \text{ if } \ i=j=1,\ldots,p \ , \\ -2 & \text{ if } \ i=j=p+1,\ldots,p+q \ , \\ 0 & \text{ otherwise }; \end{array} \right.$$

(c) The compatibility relations

$$\begin{split} i\,\rho(g) \;&=\; \varphi(g)\rho(g)\,i\ ,\\ \rho(g)\rho(h) \;&=\; \sigma(g,h)\rho(gh)\ ,\\ \gamma_j\rho(g) \;&=\; c(g)\rho(g)\gamma_j\ , \end{split}$$

valid for every $q, h \in \mathbb{G}$ and j = 1, ..., p + q.

A *homomorphism* from a twisted bundle $(\mathscr{E}, \rho, \gamma)$ to a twisted bundle $(\mathscr{E}', \rho', \gamma)$ is a map of complex vector bundles $f : \mathscr{E} \to \mathscr{E}'$ such that

$$f \circ \rho(g) = \rho'(g) \circ f$$
, $f \circ \gamma_i = \gamma'_i \circ f$,

for every $g \in \mathbb{G}$ and every j = 1, ..., p + q. The notion of *isomorphism* follows naturally.

Let $(\mathscr{E}, \rho, \gamma)$ be a twisted bundle with a Clifford action. An invertible Hermitian map $H : \mathscr{E} \to \mathscr{E}$ is said to be *compatible* with the twisted action if it holds true that

$$H\rho(g) = c(g)\rho(g)H$$
, $H\gamma_i = -\gamma_iH$.

for every $g \in \mathbb{G}$ and every $j = 1, \ldots, p+q$. If H and H' are invertible Hermitian maps on $(\mathscr{E}, \rho, \gamma)$ and $(\mathscr{E}', \rho', \gamma')$ respectively, such that $f \circ H = H' \circ f$ for a given isomorphism $f : \mathscr{E} \to \mathscr{E}'$, then H and H' will be regarded as *isomorphic*. Note that the self-adjoint involution $Q_H := H/|H|$ is nothing but a gradation, or a \mathbb{Z}_2 -grading of the twisted bundle \mathscr{E} . Since Q_H and H are homotopic, we can generalize Karoubi's formulation of twisted equivariant K-theory [DK, Ros, FHT, FM, Gom2] as follows:

Definition 2.10 (Twisted equivariant K-theory). Let X, ϕ , σ and c be as above.

- (a) We define ${}^{\varphi}\mathcal{M}^{(\sigma,c)+(p,q)}_{\mathbb{G}}(X)$ to be the monoid of isomorphism classes of triples (\mathscr{E},H_0,H_1) consisting of (φ,σ,c) -twisted bundles \mathscr{E} with $Cl_{p,q}$ -action and two invertible Hermitian maps H_0 and H_1 on \mathscr{E} compatible with the twisted actions.
- (b) We define ${}^{\varphi}\mathcal{Z}_{\mathbb{G}}^{(\sigma,c)+(p,q)}(X)$ to be the submonoid of ${}^{\varphi}\mathcal{M}_{\mathbb{G}}^{(\sigma,c)+(p,q)}(X)$ consisting of isomorphism classes of triples (\mathscr{E},H_0,H_1) such that H_0 and H_1 are homotopic within invertible Hermitian maps compatible with the twisted actions.
- (c) We define ${}^{\phi}K_{\mathbb{G}}^{(\sigma,c)+(p,q)}(X) := {}^{\phi}\mathcal{M}_{\mathbb{G}}^{(\sigma,c)+(p,q)}(X)/{}^{\phi}\mathcal{M}_{G}^{(\sigma,c)+(p,q)}(X)$ to be the quotient monoid.

In view of the (1, 1)-periodicity ${}^{\varphi}K_{\mathbb{G}}^{(\sigma,c)+(p+1,q+1)}(X)\cong {}^{\varphi}K_{\mathbb{G}}^{(\sigma,c)+(p,q)}(X)$ [Gom2], we put

$${}^{\phi}K_{\mathbb{G}}^{(\sigma,c)+q-p}(X) := {}^{\phi}K_{\mathbb{G}}^{(\sigma,c)+(p,q)}(X). \tag{2.9}$$

Let us now describe how from a band insulator $H:X\to \mathfrak{Herm}(\mathbb{C}^{2m})^{\times}$ of class DIII on the involutive space (X,τ) one gets an invertible Hermitian map on a twisted bundle. Consider the group $\mathbb{G}=\mathbb{Z}_2=\{\pm 1\}$ as generated by $\tau\equiv -1$, i. e. let \mathbb{G} act on X through the involution. We choose $\varphi:\mathbb{G}\to\mathbb{Z}_2$ to be the identity homomorphism $\varphi(\pm 1)=\pm 1$ and $c:\mathbb{G}\to\mathbb{Z}_2$ to be the trivial homomorphism $c(\pm 1)=0$. These choices can be shortly summarized by $\varphi=1$ and c=0. Let $\sigma:\mathbb{G}\times\mathbb{G}\to\mathbb{U}(1)$ be the 2-cocycle (independent of the points of X) defined by

$$\sigma(1,1) = \sigma(-1,1) = \sigma(1,-1) = 1, \qquad \sigma(-1,-1) = -1.$$

Let $\mathscr E=X\times\mathbb C^{2m}$ be the product Hermitian vector bundle on X and define $\rho(g):\mathscr E\to\mathscr E$ as follows:

$$\rho(1) : (x, v) \longmapsto (x, v), \qquad \rho(-1) : (x, v) \longmapsto (\tau(x), T\bar{v}) \tag{2.10}$$

where T denotes the TRS. We also define $\gamma:\mathscr{E}\to\mathscr{E}$ to be the unitary map defined by

$$\gamma: (x, v) \longmapsto (x, i\chi v).$$
 (2.11)

where χ denotes the chiral symmetry. Then $(\mathscr{E},\rho,\gamma)$ is a (φ,σ) -twisted vector bundle with $Cl_{0,1}$ -action. Finally, from $H:X\to \mathfrak{Herm}(\mathbb{C}^{2m})^{\times}$ one gets the invertible Hermitian map $H:(x,v)\mapsto (x,H(x)v)$ on \mathscr{E} compatible with the twisted action. As a result one has that differences of band insulators of class DIII provide elements of the twisted equivariant K-theory ${}^{\varphi}K^{\sigma+1}_{\mathbb{Z}_2}(X)$ where we used the notation (2.9) and the reference to the trivial map c=0 has been omitted. As a matter of fact a φ -twisted vector bundle is nothing but a "Real" vector bundle, so that ${}^{\varphi}K^0_{\mathbb{Z}_2}(X)=KR^0(X,\tau)$ [Ati]. Since the cocycle σ has the effect of the degree shift by -4 one gets

$${}^{\phi}K_{\mathbb{Z}_2}^{\sigma+1}(X) = KR^{-3}(X,\tau)$$
 (2.12)

The next result provides the link between the group $KR^{-3}(X,\tau)$ which classifies the (differences of) band insulators of class DIII and the invariant of sewing matrices described in Definition 2.7. For that we need to introduce the homomorphism

$$\kappa : KR^{-3}(X,\tau) \longrightarrow H^1_{\mathbb{Z}_2}(X|X^{\tau},\mathbb{Z}(1))$$
 (2.13)

induced by the FKMM invariant [DG2, DG3, DG4]. More precisely, the homomorphism κ assigns to a band insulator of class DIII, thought of as a class in $KR^{-3}(X,\tau)$, the FKMM invariant of the associated "Quaternionic" vector bundle as described in Definition 2.3.

Theorem 2.11. Let (X, τ) be a path connected finite \mathbb{Z}_2 -CW complex which admits a fixed point. Then:

- (1) The map (2.13) induced by the FKMM invariant of sewing matrices provides a well-defined homomorphism;
- (2) The invariant of sewing matrices introduced in Definition 2.7 provides a well-defined homomorphism

$$\nu \,:\, \text{Ker}(\kappa) \,\longrightarrow\, H^1(X,\mathbb{Z})/H^1_{\mathbb{Z}_2}\big(X|X^\tau,\mathbb{Z}(1)\big) \;.$$

Proof. Let (\mathscr{E}, H_0, H_1) be a triple representing an element of $KR^{-3}(X, \tau)$. In general, a twisted graded vector bundle on a compact space equivariant under a finite group admits a complementary twisted bundle \mathscr{F} such that $\mathscr{E} \oplus \mathscr{F} \simeq X \times \mathcal{H}$ is isomorphic to a product bundle for some finite dimensional Hilbert space \mathcal{H} and the twisted actions are independent of the points on X (this fact is a straightforward generalization of a property of the usual equivariant vector bundles). Let $H_F : \mathscr{F} \to \mathscr{F}$ be an invertible Hermitian map. In view of the definition of $KR^{-3}(X,\tau)$ as a quotient monoid, (\mathscr{E}, H_0, H_1) and $(\mathscr{E} \oplus \mathscr{F}, H_0 \oplus H_F, H_1 \oplus H_F)$ provide the same class in the K-theory. Thus, we can assume that $\mathscr{E} = X \times \mathcal{H}$ with twisted actions independent of points on X since the beginning. Therefore, as shown in Proposition 2.1, we can represent the twisted actions on $X \times \mathcal{H}$ in a standard form, and express H_0 and H_1 in terms of sewing matrices $q_0 := q_{H_0}$ and

 $q_1 := q_{H_1}$, respectively.

(1) With the argument above (and the identification $\mathcal{H} \equiv \mathbb{C}^{4n}$ in view of Proposition 2.4) one can specify the action of the map (2.13) as follows

$$\kappa([X \times \mathbb{C}^{4n}, H_0, H_1]) := \kappa(\mathscr{E}_{q_0}) - \kappa(\mathscr{E}_{q_1})$$
 (2.14)

where \mathscr{E}_{q_i} is the "Quaternionic" vector bundle associated with the sewing matrix q_i , with i=0,1. Notice that this definition is independent of the way to identify the standard presentation. In fact if one changes the identification, one obtains new sewing matrices q_i' which are related to q_i via a continuous map $\varphi:X\mapsto \mathbb{U}(2n)$ according to equation (B.1). This map induces an isomorphism of the associated "Quaternionic" vector bundles $\mathscr{E}_{q_i'}\simeq\mathscr{E}_{q_i}$. Since the FKMM invariant takes the same value on isomorphic "Quaternionic" bundles, it follows that equation (2.14) is insensitive on the way of fixing the standard presentation. Furthermore, if H_0 and H_1 are homotopic, then so are q_0 and q_1 . In this case the homotpy invariance of "Quaternionic" bundles imples $\kappa(\mathscr{E}_{q_0})=\kappa(\mathscr{E}_{q_1})$, and in turn one gets $\kappa([X\times\mathbb{C}^{4n},H_0,H_1])=0$ for the trivial class. This fact, along with the additivity of the FKMM invariant under the direct sum of "Quaternionic" bundles, proves that (2.13) is a well-defined homomorphism.

(2) Let us start by showing that when $[X \times \mathbb{C}^{4n}, H_0, H_1] \in Ker(\kappa)$ then it is possible to choose the sewing matrices q_i in such a way that $\kappa(\mathscr{E}_{q_i}) = 0$ for i = 0, 1. To see this observe that $[X \times \mathbb{C}^{4n}, H_0, H_1] \in Ker(\kappa)$ implies $\kappa(\mathscr{E}_{q_0}) = \kappa(\mathscr{E}_{q_1})$. As described in the proof of Lemma 2.6, the FKMM invariant $\kappa(\mathscr{E}_{q_i})$ can be represented by the map $X^\tau \ni x \mapsto p_i(x)/Pf[q_i(x)] \in \mathbb{Z}_2$ where $p_i : X \to \mathbb{U}(1)$ is such that $det[q_i(x)] = p_i(\tau(x))p_i(x)$ for all $x \in X$. Let $q_2 : X \to \mathbb{U}(2n)$ be the sewing matrix defined by $q_2(x) := q_0(x)^{-1}$. This provides $Pf[q_2(x)] = (-1)^n Pf[q_0(x)]^{-1}$ for all $x \in X^\tau$. If we define $p_2 : X \to \mathbb{U}(1)$ by $p_2(x) := (-1)^n p_0(x)^{-1}$ for all $x \in X$, then one gets $det[q_2(x)] = p_2(\tau(x))p_2(x)$ for all $x \in X$ and the FKMM invariant $\kappa(\mathscr{E}_{q_2})$ is represented by the map

$$X^{\tau} \ni x \longmapsto \frac{p_2(x)}{\operatorname{Pf}[q_2(x)]} = \left(\frac{p_0(x)}{\operatorname{Pf}[q_0(x)]}\right)^{-1} \in \mathbb{Z}_2.$$

This means that $\kappa(\mathscr{E}_{q_2}) = -\kappa(\mathscr{E}_{q_0})$. Now, let us introduce the Hamiltonian $H_2: X \to \mathfrak{Herm}(\mathbb{C}^{4n})^{\times}$ given by

$$H_2(x) := \left(egin{array}{cc} 0 & q_2(x)^{-1} \\ q_2(x) & 0 \end{array}
ight) \,, \qquad x \in X \,,$$

and the new triple $(X \times \mathcal{H}', H'_0, H'_1)$ defined by

$$\begin{split} (X\times \mathfrak{H}',H_0',H_1') \; := \; (X\times \mathbb{C}^{4n},H_0,H_1) \; \oplus \; (X\times \mathbb{C}^{4n},H_2,H_2) \\ &= \; \left(X\times (\mathbb{C}^{4n}\oplus \mathbb{C}^{4n}),H_0\oplus H_2,H_1\oplus H_2\right). \end{split}$$

The classes $[X \times \mathbb{C}^{4n}, H_0, H_1]$ and $[X \times \mathcal{H}', H_0', H_1']$ represent the same element in $Ker(\kappa)$ and

$$\begin{split} \kappa(\mathscr{E}_{q_0'}) \; &=\; \kappa(\mathscr{E}_{q_0} \oplus \mathscr{E}_{q_2}) \; = \; \kappa(\mathscr{E}_{q_0}) + \kappa(\mathscr{E}_{q_2}) \; = \; 0 \\ \kappa(\mathscr{E}_{q_1'}) \; &=\; \kappa(\mathscr{E}_{q_1} \oplus \mathscr{E}_{q_2}) \; = \; \kappa(\mathscr{E}_{q_1}) + \kappa(\mathscr{E}_{q_2}) \; = \; \kappa(\mathscr{E}_{q_0}) + \kappa(\mathscr{E}_{q_2}) \; = \; 0 \; . \end{split}$$

As a consequence, we can show that

$$Ker(\kappa) = {}^{\varphi} \mathfrak{N}_{\mathbb{Z}_2}^{\sigma+(0,1)}(X) / \left({}^{\varphi} \mathfrak{N}_{\mathbb{Z}_2}^{\sigma+(0,1)}(X) \ \cap \ {}^{\varphi} \mathfrak{Z}_{\mathbb{Z}_2}^{\sigma+(0,1)}(X) \right) \ ,$$

where the submonoid ${}^{\varphi}\mathcal{N}^{\sigma+(0,1)}_{\mathbb{Z}_2}(X)\subset {}^{\varphi}\mathcal{M}^{\sigma+(0,1)}_{\mathbb{Z}_2}(X)$ is defined by

$${}^\varphi \mathfrak{N}^{\sigma+(0,1)}_{\mathbb{Z}_2}(X) \;:=\; \left\{ (X\times \mathfrak{H},H_0,H_1)\in {}^\varphi \mathfrak{M}^{\sigma+(0,1)}_{\mathbb{Z}_2}(X) \;\middle|\; \kappa(\mathcal{E}_{\mathfrak{q}_0})=\kappa(\mathcal{E}_{\mathfrak{q}_1})=0 \right\} \;.$$

Now, let us define the map ν which associates to $(X \times \mathcal{H}, H_0, H_1) \in {}^{\phi}\mathcal{N}_{\mathbb{Z}_2}^{\sigma+(0,1)}(X)$ the difference of the invariants of the sewing matrices q_0 and q_1 , *i.e.*

$$\nu((X \times \mathcal{H}, H_0, H_1)) := \nu_{q_0} - \nu_{q_1}.$$
 (2.15)

The rest of the proof consists in proving that the map ν as defined by (2.15) provides a well-defined homomorphism on the group $\text{Ker}(\kappa)$. First of all, let us notice that the sewing matrices q_0, q_1 depend on the choice of the isomorphism $X \times \mathcal{H} \simeq X \times \mathbb{C}^{4n}$ which makes the twisted actions in the standard form as discussed in Lemma 2.2. By combining the notation introduced in (2.10) and (2.11) with Proposition 2.1, one gets that the standard actions are given by

$$\rho(-1) : (x,v) \longmapsto (x,Tv) = \left(x, \begin{pmatrix} 0 & -\mathbf{1}_{2n} \\ +\mathbf{1}_{2n} & 0 \end{pmatrix} \overline{v}\right)$$

and

$$\gamma \;:\; (x,v) \;\longmapsto\; (x,\,i\,\chi v) \;=\; \left(x,\left(\begin{array}{cc} +\,i\,\textbf{1}_{2\,n} & \textbf{0} \\ \textbf{0} & -\,i\,\textbf{1}_{2\,n} \end{array}\right)v\right)\;.$$

Moreover, every continuous map $f:X\to \mathbb{U}(2n)$ provides the automorphism ψ_f of $X\times \mathbb{C}^{4n}$ given by

$$\psi_f : (x, v) \longmapsto \left(x, \left(\begin{array}{cc} f(x) & 0 \\ 0 & \overline{f(\tau(x))} \end{array}\right) v\right).$$

Let $\phi: X \times \mathcal{H} \cong X \times \mathbb{C}^{4n}$ and $\phi': X \times \mathcal{H} \cong X \times \mathbb{C}^{4n}$ be two standard representation intertwined by the automorphism ψ_f , *i. e.* $\phi = \psi_f \circ \phi'$. Let q_0, q_1 be the sewing matrices associated to H_0 and H_1 , respectively, via the isomorphism ϕ . Similarly, let q_0', q_1' be the related sewing matrices obtained via the isomorphism ϕ' . A straightforward computation shows that

$$q'_{j}(x) = f(\tau(x))^{t} q_{j}(x) f(x), \qquad j = 0, 1.$$

By using Theorem 2.8 (4) one gets the following fact: If ν_{q_j} is represented by $p_j \in C(X, \mathbb{U}(1))_{q_j}$, then $\nu_{q_i'}$ is represented by $p_j' = p_j \det[f]$. In terms of classes in $H^1(X, \mathbb{Z}) = f(x_j)$

[X, U(1)], one gets

$$\begin{aligned} [p_0'] - [p_1'] &= [p_0 \det[f]] - [p_1 \det[f]] \\ &= ([p_0] + [\det[f]]) - ([p_1] + [\det[f]]) = [p_0] - [p_1] . \end{aligned}$$

Therefore $\nu_{q_0'} - \nu_{q_1'} = \nu_{q_0} - \nu_{q_1}$ in $H^1(X,\mathbb{Z})/H^1_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))$, and the map ν defined by (2.15) turns out to be independent of the choice of the standard representation $X \times \mathcal{H} \simeq X \times \mathbb{C}^{4n}$. This also implies that $\nu(X \times \mathcal{H}, H_0, H_1) = \nu(X \times \mathcal{H}', H_0', H_1')$ whenever the triples $(X \times \mathcal{H}, H_0, H_1)$ and $(X \times \mathcal{H}', H_0', H_1')$ are isomorphic. This leads to a map

$$\nu \;:\; {}^\varphi \mathcal{N}^{\sigma+(0,1)}_{\mathbb{Z}_2}(X) \;\longrightarrow\; H^1(X,\mathbb{Z})/H^1_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))\;,$$

which is a monoid homomorphism by Theorem 2.8 (2) and (3). By Theorem 2.8 (1), the monoid homomorphism is trivial on the submonoid ${}^{\varphi}\mathcal{Z}_{\mathbb{Z}_2}^{\sigma+(0,1)}(X) \cap {}^{\varphi}\mathcal{N}_{\mathbb{Z}_2}^{\sigma+(0,1)}(X)$. Consequently, we get a well-defined homomorphism ν on the group $\operatorname{Ker}(\kappa)$. Equation (2.12) completes the proof.

Remark 2.12. It is well known that the involutive space (\mathbb{S}^1, ι) admits only the trivial "Quaternionic" vector bundle [DG2, Theorem 1.2 (ii)]. As a consequence one gets that $\operatorname{Ker}(\kappa) = K^{-3}(\mathbb{S}^1, \iota)$ in this special case. The properties of the homomorphism ν on $K^{-3}(\mathbb{S}^1, \iota)$ will be studied in Proposition 3.15.

Remark 2.13. As pointed out in the proof of Theorem 2.11, the sewing matrix q depends on the way to express the twisted bundle $X \times \mathcal{H}$ in a standard form. If we change the isomorphism by an automorphism associated to $f: X \to \mathbb{U}(m)$, then the sewing matrix q is changed into a sewing matrix q' and the relation between the two sewing matrices is expressed by the equation $q'(x) = f(\tau(x))^t q(x) f(x)$. The difference $\nu_{q'} - \nu_q$ is represented by $[\det[f]]$ in the group $H^1(X,\mathbb{Z})/H^1_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))$. Since $\det: [X,\mathbb{U}(m)] \to [X,\mathbb{U}(1)]$ is surjective one has that all the elements in $H^1(X,\mathbb{Z})/H^1_{\mathbb{Z}_2}(X|X^\tau,\mathbb{Z}(1))$ arise just by changing the isomorphism $X \times \mathcal{H} \cong X \times \mathbb{C}^{2m}$. This means that we cannot define an *absolute* invariant of a band insulator H of class DIII by using the invariant ν_q of its sewing matrix q. Notice, however, that the invariant is able to detect the *difference* of two band insulators, as established in Theorem 2.11. In other words ν_q works as *relative* invariant for band insulators of class DIII.

Remark 2.14. A map $\phi: X \to \mathbb{U}(1)$ over the involutive space (X,τ) is called *invariant* if $\phi = \phi \circ \tau$. The equivariant cohomology $H^1_{\mathbb{Z}_2}(X;\mathbb{Z})$ is isomorphic to the group $[X,\mathbb{U}(1)]_{inv}$ of homotopy classes of invariant maps (see Appendix A). Thus, in general, $[\det[q]] \in H^1_{\mathbb{Z}_2}(X,\mathbb{Z})$ is an invariant of the homotopy class of a sewing matrix $q: X \to \mathbb{U}(2n)$. By a proof similar to that of Theorem 2.11, one can show that this invariant defines a homomorphism

$$det : KR^{-3}(X,\tau) \longrightarrow H^1_{\mathbb{Z}_2}(X,\mathbb{Z}) .$$

However, this invariant does not detect $KR^{-3}(\mathbb{S}^1,\iota)\simeq\mathbb{Z}_2$ since $H^1_{\mathbb{Z}_2}(\mathbb{S}^1,\mathbb{Z})=0$. This observation is in agreement with the content item (II) presented in Section 1.

3. The one-dimensional case

In this section we will analyze in detail the case of *one-dimensional* band insulators of class DIII. According to Definition 1.1, the latter are band insulators of class DIII defined on the involutive space (\mathbb{S}^1 , ι). This peculiar case is relevant for its application to physical systems. In fact, in this section we will show the coincidence of the topological invariant ν_q described in Definition 2.7 with the topological invariant for topological insulators of class DIII usually used in the physical literature [RSFL, TK, BA, CTSR]. For the benefit of the reader we report here a table of some cohomology groups of the involutive space (\mathbb{S}^1 , ι) which will be useful in the following:

n	$H^{n}(\mathbb{S}^{1},\mathbb{Z})$	$H^{\mathfrak{n}}_{\mathbb{Z}_{2}}(\mathbb{S}^{1},\mathbb{Z})$	$H^{\mathfrak{n}}_{\mathbb{Z}_{2}}(\mathbb{S}^{1},\mathbb{Z}(1))$	$H^{\mathfrak{n}}_{\mathbb{Z}_{2}}((\mathbb{S}^{1})^{\iota},\mathbb{Z}(1))$	$H_{\mathbb{Z}_2}^n(\mathbb{S}^1 (\mathbb{S}^1)^{\iota},\mathbb{Z}(1))$
0	$\mathbb Z$	$\mathbb Z$	0	0	0
1	\mathbb{Z}	0	$\mathbb{Z}\oplus\mathbb{Z}_2$	$\mathbb{Z}_2 \oplus \mathbb{Z}_2$	$\mathbb Z$
2	0	$\mathbb{Z}_2 \oplus \mathbb{Z}_2$	0	0	0
3	0	0	$\mathbb{Z}_2 \oplus \mathbb{Z}_2$	$\mathbb{Z}_2 \oplus \mathbb{Z}_2$	

The proof of the results listed in the table can be found in [Gom, Lemma 2.12], in [DG1, Section 5.3] and in [DG3, Appendix B]. Let us also recall that $H^1(\mathbb{S}^1, \mathbb{Z}) \simeq [\mathbb{S}^1, \mathbb{U}(1)]$ is generated by the identity map under the identification $\mathbb{U}(1) \simeq \mathbb{S}^1$. Similarly, in view of $H^1_{\mathbb{Z}_2}(\mathbb{S}^1, \mathbb{Z}(1)) \simeq [\mathbb{S}^1, \mathbb{U}(1)]_{\mathbb{Z}_2}$ [Gom, Proposition A.2] one gets that the free subgroup \mathbb{Z} is generated by the identity map under the identification of $\mathbb{U}(1)$, endowed with the complex conjugation, with the involutive space (\mathbb{S}^1, ι) . The torsion subgroup \mathbb{Z}_2 is generated by the constant map with value -1. All "Real" and "Quaternionic" vector bundles on (\mathbb{S}^1, ι) are trivial as proved in [DG1, Theorem 1.6] and [DG2, Theorem 1.2]. Finally, let us observe that (\mathbb{S}^1, ι) satisfies the properties (a), (b) and (c) listed in Assumption 2.5.

3.1. **The Teo-Kane formula.** In this section we will provide an explicit formula to compute the invariant v_q in the one-dimensional case. In particular we will establish the connection with the *Teo-Kane formula*, the latter being the topological invariant commonly used in the physical literature for one-dimensional for topological insulators of class DIII (see e. g. [TK, eq. (4.27)] or [BA, eq. (8)] or [CTSR, eq. (3.70)]).

Let us start with a couple of preliminary results about the topology of the maps which define the invariant ν_q . The first result concerns with the analysis of the range of the invariant ν_q .

Lemma 3.1. Consider the involutive space (\mathbb{S}^1, ι) . Let $\delta : C(\mathbb{S}^1, \mathbb{U}(1)) \to \mathbb{Z}_2$ be the map defined by $\delta(\mathfrak{p}) := (-1)^{\deg(\mathfrak{p})}$ where $\deg(\mathfrak{p})$ denotes the degree (or winding number) of \mathfrak{p} as a map $\mathfrak{p} : \mathbb{S}^1 \to \mathbb{U}(1) \simeq \mathbb{S}^1$. Then δ induces the isomorphism of groups

$$H^{1}(\mathbb{S}^{1}, \mathbb{Z})/H^{1}_{\mathbb{Z}_{2}}(\mathbb{S}^{1}|(\mathbb{S}^{1})^{\iota}, \mathbb{Z}(1)) \simeq \mathbb{Z}_{2}. \tag{3.1}$$

Proof. As anticipated in the introductory part of this section, $H^1_{\mathbb{Z}_2}(\mathbb{S}^1, \mathbb{Z}(1)) \simeq \mathbb{Z} \oplus \mathbb{Z}_2$ is generated by the identity map $\mathbb{S}^1 \to \mathbb{U}(1)$ and the constant map with value -1. From the

exact sequence [DG3, eq. (2.6)]

$$\begin{array}{ll} 0 \ = \ H^0_{\mathbb{Z}_2}\big((\mathbb{S}^1)^\iota,\mathbb{Z}(1)\big) \longrightarrow H^1_{\mathbb{Z}_2}\big(\mathbb{S}^1|(\mathbb{S}^1)^\iota,\mathbb{Z}(1)\big) \stackrel{\jmath}{\longrightarrow} H^1_{\mathbb{Z}_2}\big(\mathbb{S}^1,\mathbb{Z}(1)\big) \\ \stackrel{r}{\longrightarrow} H^1_{\mathbb{Z}_2}\big((\mathbb{S}^1)^\iota,\mathbb{Z}(1)\big) \longrightarrow H^2_{\mathbb{Z}_2}\big(\mathbb{S}^1|(\mathbb{S}^1)^\iota,\mathbb{Z}(1)\big) \ = \ 0, \end{array}$$

one gets that the map r, induced by the restriction $\mathbb{S}^1 \to (\mathbb{S}^1)^\iota$, is surjective. Moreover, one infers that the injective image of $H^1_{\mathbb{Z}_2}(\mathbb{S}^1|(\mathbb{S}^1)^\iota,\mathbb{Z}(1))$ under \mathfrak{g} is generated by the equivariant map $e:\mathbb{S}^1\to\mathbb{U}(1)$ given by $e(z):=e^{i\,2k}$. Since the map $f:H^1_{\mathbb{Z}_2}(\mathbb{S}^1,\mathbb{Z}(1))\to H^1(\mathbb{S}^1,\mathbb{Z})$ which forgets the \mathbb{Z}_2 -action is the projection which induces the bijection on the free part of the two groups, it follows that the image of

$$H^1_{\mathbb{Z}_2}\big(\mathbb{S}^1|(\mathbb{S}^1)^\iota,\mathbb{Z}(1)\big) \ \stackrel{\jmath}{\longrightarrow} \ H^1_{\mathbb{Z}_2}\big(\mathbb{S}^1,\mathbb{Z}(1)\big) \ \stackrel{f}{\longrightarrow} \ H^1(\mathbb{S}^1,\mathbb{Z})$$

is generated by maps $\mathbb{S}^1 \to \mathbb{U}(1)$ with even degree. This proves (3.1) with the isomorphism given by the parity of the degree of the maps from \mathbb{S}^1 to $\mathbb{U}(1)$.

The next result provides a formula for the degree of the equivariant maps on the involutive space (\mathbb{S}^1, ι) with value in $\mathbb{U}(1)$. Let us emphasize that a map $r : \mathbb{S}^1 \to \mathbb{U}(1)$ is equivariant if $r(\iota(k)) = \overline{r(k)}$ for every $k \in \mathbb{S}^1$.

Lemma 3.2. Let $r: \mathbb{S}^1 \to \mathbb{U}(1)$ be an equivariant map defined on the involutive space (\mathbb{S}^1, ι) . Then, it holds true that

$$(-1)^{\deg(r)} = \frac{r(\pi)}{r(0)}$$
.

Proof. The equivariance condition implies that $r(k) = \pm 1$ at the fixed points $k = 0, \pi$. In particular, the value at each fixed point is invariant under equivariant homotopy deformations of r. Therefore, the assignment $\Delta: r \mapsto r(\pi)/r(0)$ induces a well-defined homomorphism $\Delta: H^1_{\mathbb{Z}_2}(\mathbb{S}^1, \mathbb{Z}(1)) \to \mathbb{Z}_2$ (still denoted with the same symbol). By using the explicit basis of $H^1_{\mathbb{Z}_2}(\mathbb{S}^1, \mathbb{Z}(1))$, one can verify the equality $\Delta(r) = (-1)^{\deg(r)}$.

With the help of Lemma 3.1 and Lemma 3.2 we can provide a first formula for the invariant ν_q .

Proposition 3.3 (The degree fromula). *Consider the involutive space* (\mathbb{S}^1 , ι) *along with a map* $q: \mathbb{S}^1 \to \mathbb{U}(2n)$ *such that*

$$\label{eq:definition} q\big(\iota(k)\big) \; = \; -q(k)^{\mathtt{t}} \; , \qquad \forall \; k \in \mathbb{S}^1 \; .$$

Then, the topological invariant ν_q of the map q described in Definition 2.7 can be computed with the formula

$$\nu_q \; = \; (-1)^{deg(p)} \; \in \; \mathbb{Z}_2 \; ,$$

independently of $p \in C(\mathbb{S}^1, \mathbb{U}(1))_q$.

Proof. First of all it is worth noting that all the conditions listed in Assumption 2.5 are satisfied and therefore the invariant v_q is well-defined. In view of Lemma 3.1 one gets

that

$$\nu_q \;\in\; H^1(\mathbb{S}^1,\mathbb{Z})/H^1_{\mathbb{Z}_2}\big(\mathbb{S}^1|(\mathbb{S}^1)^\iota,\mathbb{Z}(1)\big) \;\simeq\; \mathbb{Z}_2\;.$$

Since the isomorphism with \mathbb{Z}_2 is realized by the parity of the degree, we get the expression $\nu_q = (-1)^{deg\,p}$ in terms of $p \in C(\mathbb{S}^1,\mathbb{U}(1))_q$. Indeed, in view of Lemma 2.6 (1) we know that for every two $p,p' \in C(\mathbb{S}^1,\mathbb{U}(1))_q$ there is an equivariant map $r:\mathbb{S}^1 \to \mathbb{U}(1)$ such that p' = rp and $r(0) = r(\pi) = 1$. From Lemma 3.2 one infers that deg(r) is even and in turn deg(p) and deg(p') = deg(p) + deg(r) have the same parity.

The next result is preparatory for the proof of the Teo-Kane formula.

Lemma 3.4. Let $q:\mathbb{S}^1\to\mathbb{U}(2n)$ be a map defined on the involutive space (\mathbb{S}^1,ι) such that

$$q(\iota(k)) = -q(k)^{t}, \quad \forall k \in \mathbb{S}^{1}.$$

Then, there exists a map $\widetilde{q}:\mathbb{S}^1\times[0,1]\to\mathbb{U}(2n)$ such that

$$\widetilde{q}\big(\iota(k),t\big) \,=\, -\widetilde{q}(k,t)^{\mathtt{t}}\;, \qquad \widetilde{q}(k,0) \,=\, q(k)\;, \qquad \text{det}\big[\widetilde{q}(k,1)\big] \,=\, 1\;,$$

for all $k \in \mathbb{S}^1$ and $t \in [0, 1]$.

Proof. As anticipated in item ii) of Section 1, $det[q]: \mathbb{S}^1 \to \mathbb{U}(1)$ is an invariant map, i. e. $det[q(k)] = det[q(\iota(k))]$ for every $k \in \mathbb{S}^1$. Let $[\mathbb{S}^1, \mathbb{U}(1)]_{inv}$ be the set of classes of homotopy equivalent invariant functions. Then, one has that $[\mathbb{S}^1, \mathbb{U}(1)]_{inv} \simeq H^1_{\mathbb{Z}_2}(\mathbb{S}^1, \mathbb{Z}) = 0$ (cf. Appendix A). Therefore, there exists $f: \mathbb{S}^1 \times [0,1] \to \mathbb{U}(1)$ such that $f(k,t) = f(\iota(k),t)$, f(k,0) = 1 and $f(k,1) = det[q(k)]^{-1}$ for every $k \in \mathbb{S}^1$ and $t \in [0,1]$. The degree of the map $k \mapsto f(k,t)$ is constant in t and therefore is trivial for every $t \in [0,1]$ (since it is trivial for t = 0). This allows to define a square root

$$g(k,t) := f(k,t)^{\frac{1}{2}}$$

consistently. This provides a continuous map $g:\mathbb{S}^1\times[0,1]\to\mathbb{U}(1)$ such that $g(k,t)=g(\iota(k),t),\,g(k,0)=1$ and $g(k,1)^2=\det[q(k)]^{-1}$ for every $k\in\mathbb{S}^1$ and $t\in[0,1]$. Let us introduce the map $q:\mathbb{S}^1\times[0,1]\to\mathbb{U}(2n)$ defined by

$$\widetilde{q}(k,t) := \begin{pmatrix} g(k,t) & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \ddots \end{pmatrix} q(k) \begin{pmatrix} g(k,t) & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \ddots \end{pmatrix}. \tag{3.2}$$

By construction the map \tilde{q} meets the properties stated in the claim.

Theorem 3.5 (The Teo-Kane formula). Let $q: \mathbb{S}^1 \to \mathbb{U}(2n)$ be a map defined on the involutive space (\mathbb{S}^1, ι) such that

$$q\big(\iota(k)\big) \; = \; -q(k)^{\mathtt{t}} \; , \qquad \forall \; k \in \mathbb{S}^1 \; .$$

Then, the invariant $\nu_q \in \mathbb{Z}_2$ admits the expression

$$\nu_{q} = \frac{Pf[q(\pi)]}{Pf[q(0)]} \frac{det[q(0)]^{\frac{1}{2}}}{det[q(\pi)]^{\frac{1}{2}}}$$

where the branch $k \mapsto det[q(k)]^{\frac{1}{2}}$ must be chosen continuously between the two fixed points $k = 0, \pi$.

Proof. Theorem 2.8 (1) ensures that the invariant ν_q only depends on the homotopy class of q. By combining this fact with Lemma 3.4 one gets that $\nu_q = \nu_{q'}$ where $q'(k) := \widetilde{q}(k,1)$ according to the notation of Lemma 3.4. The crucial property of q' is that det[q'(k)] = 1 for every $k \in \mathbb{S}^1$. From Proposition 3.3 we know that $\nu_{q'} = (-1)^{deg(p')}$, where $p' : \mathbb{S}^1 \to \mathbb{U}(1)$ is such that $1 = p'(\iota(k))p'(k)$ for all $k \in \mathbb{S}^1$ and p'(k) = Pf[q'(k)] for $k = 0, \pi$. Then, one has $p'(0)^2 = 1 = p'(\pi)^2$ so that $p'(\pi)/p'(0) \in \mathbb{Z}_2$. In view of Lemma 3.2, one has that

$$\nu_{q'} = (-1)^{\deg(p')} = \frac{p'(\pi)}{p'(0)} = \frac{Pf[q'(\pi)]}{Pf[q'(0)]}.$$

From equation (3.2) which provides the link between q and q' and the well-known formula $Pf[BAB^t] = det[B]Pf[A]$ for generic matrices A, B with A skew-symmetric, one gets

$$Pf \big[q'(k) \big] \; = \; g(k,1) \; Pf \big[q(k) \big] \; = \; \frac{Pf \big[q(k) \big]}{det \big[q(k) \big]^{\frac{1}{2}}} \; , \qquad k = 0, \pi \; .$$

This concludes the proof.

For a concrete physical model of one-dimensional topological insulator of class DIII whose (non-trivial) topology is described be the Teo-Kane formula we refer to [TK, Section IV.C] or [CTSR, Section B.6.a]. The connection with the *Kramers polarization* is discussed in [BA].

3.2. The role as topological obstruction. In this section we will show that in dimension one the invariant ν_q can be interpreted as a *topological obstruction*. For this aim, we need to introduce the involutive space $(\mathbb{U}(2n),\theta)$ given by the unitary group on \mathbb{C}^{2n} endowed with the involution $\theta(U):=-U^t$. The fixed point set $\mathbb{U}(2n)^\theta$ coincides with the set of skew-symmetric matrices on \mathbb{C}^{2n} . Let us introduce the standard symplectic matrix

$$Q := \begin{pmatrix} 0 & -\mathbf{1}_n \\ +\mathbf{1}_n & 0 \end{pmatrix} \in \mathbb{U}(2n)^{\theta}$$
 (3.3)

and the *compact symplectic group* $Sp(n) = \{S \in \mathbb{U}(2n) \mid S^tQS = Q\}$. One has that $Pf[Q] = (-1)^{\frac{n(n+1)}{2}}$. By considering the left action of Sp(n) on $\mathbb{U}(2n)$ one can define the quotient space $\mathbb{U}(2n)/Sp(n)$. In view of the normal decomposition for skew-symmetric matrices, one gets the well-defined homeomorphism

$$\mathbb{U}(2n)/Sp(n) \ \ni \ [U] \ \longmapsto \ U^tQU \ \in \ \mathbb{U}(2n)^\theta \ .$$

In particular, $\mathbb{U}(2n)^{\theta}$ can be seen as the base space of a principal Sp(n)-bundle.

Let us start with two preliminary results.

Lemma 3.6. Let (X, τ) be an involutive space with a fixed point $* \in X^{\tau}$. Let $q: X \to \mathbb{U}(2n)$ be a map satisfying

$$q(\tau(x)) = -q(x)^{t}, \quad \forall x \in X.$$

Then there exists a map $\widetilde{q}: X \times [0,1] \to \mathbb{U}(2n)$ such that $\widetilde{q}(\tau(x),t) = -\widetilde{q}(x,t)^t$ and $\widetilde{q}(x,0) = q(x)$ for all $(x,t) \in X \times [0,1]$ and $\widetilde{q}(*,1) = Q$.

Proof. At the fixed point $q(*) \in \mathbb{U}(2n)^{\theta}$ is a skew-symmetric unitary matrix. Let $U \in \mathbb{U}(2n)$ be a unitary matrix such that $U^{t}QU = q(*)$. Since $\mathbb{U}(2n)$ is path connected, there is a path $u:[0,1] \to \mathbb{U}(2n)$ such that $u(0) = \mathbf{1}_{2n}$ and $u(1) = U^{-1}$. Now, let us define $\widetilde{q}: X \times [0,1] \to \mathbb{U}(2n)$ by

$$\tilde{q}(x,t) := u(t)^t q(x) u(t)$$
.

A straightforward check shows that this map has the prescribed properties.

Lemma 3.7. The space

$$SU(2n)^{\theta} := \{U \in U(2n)^{\theta} \mid det[U] = 1\}$$

has two connected components distinguished by their Pfaffians.

Proof. Under the homeomorphism $\mathbb{U}(2n)/Sp(n) \simeq \mathbb{U}(2n)^{\theta}$, one can identify the subspace $S\mathbb{U}(2n)^{\theta} \subset \mathbb{U}(2n)^{\theta}$ with

$$det^{-1}(+1)/Sp(n) \; \sqcup \; det^{-1}(-1)/Sp(n) \; \subset \; \mathbb{U}(2n)/Sp(n)$$

where $det^{-1}(\pm 1):=\{U\in \mathbb{U}(2n)\mid det(U)=\pm 1\}.$ In particular $det^{-1}(+1)=S\mathbb{U}(2n).$ Let

$$I := \left(egin{array}{ccc} i & 0 & 0 \ 0 & 1 & 0 \ 0 & 0 & \ddots \end{array}
ight) \in \, \mathbb{U}(2n) \; .$$

The map $U\mapsto IUI$ provides a homeomorphism $det^{-1}(-1)\simeq det^{-1}(+1)$. Thus, $S\mathbb{U}(2n)^{\theta}$ is homeomorphic to the disjoint union of two copies of $S\mathbb{U}(2n)/Sp(n)$, and in turn it has two connected components. From the form of the homeomorphism and the property of the Pfaffian one gets that $Pf:[U]\mapsto Pf[U^tQU]=\pm Pf[Q]$ for every $[U]\in det^{-1}(\pm 1)/Sp(n)$. Therefore, the Pfaffian distinguishes the two connected components of $S\mathbb{U}(2n)^{\theta}$.

The next result will shed light on the nature of the invariant ν_q as the topological obstruction for the existence of a homotopy transformation between two given sewing matrices.

Theorem 3.8 (Topological obstruction). Consider the involutive space (\mathbb{S}^1, ι) and two maps $q_j : \mathbb{S}^1 \to \mathbb{U}(n)$, j = 0, 1, satisfying the sewing matrix condition

$$q_{\mathfrak{j}}\big(\iota(k)\big) \; = \; -q_{\mathfrak{j}}(k)^{\mathtt{t}} \; , \qquad \forall \; k \in \mathbb{S}^1 \; , \quad \mathfrak{j} = 0,1 \; .$$

The maps q_0, q_1 are homotopy equivalent within the space of sewing matrices if and only if $\nu_{q_0} = \nu_{q_1}$.

Proof. Theorem 2.8 (1) provides the "only if" part of the claim. Therefore, we only need to show the "if" part. By Lemma 3.4 and Lemma 3.6, we can assume that $det[q_0(k)] = det[q_1(k)] = 1$ for all $k \in \mathbb{S}^1$ and $q_0(0) = q_1(0) = Q$. Under these assumptions, the coincidence of the invariants $\nu_{q_0} = \nu_{q_1}$ is equivalent to $Pf[q_0(\pi)] = Pf[q_1(\pi)]$ in view of Theorem 3.5. We will proceed to build a homotopy of sewing matrices $\widetilde{q}: \mathbb{S}^1 \times [0,1] \to \mathbb{U}(2n)$ such that $\widetilde{q}(k,j) = q_j(k)$ for j = 0,1 and $k \in \mathbb{S}^1$. We will do this in several steps: (1) As a first step let us choose a continuous map

$$\widetilde{Q}:[0,1]\longrightarrow S\mathbb{U}(2n)^{\theta}\subset \mathbb{U}(2n)\,,\qquad \widetilde{Q}(j):=q_{j}(\pi)\,,\;\;j=0,1\,.$$

This is possible, because Lemma 3.7 establishes that $SU(2n)^{\theta}$ has two connected components distinguished by the Pfaffian, and $Pf[q_0(\pi)] = Pf[q_1(\pi)]$ by hypothesis.

(2) As a second step, let us build the continuous map $\tilde{q}: \mathfrak{d}([0,\pi]\times[0,1])\to \mathbb{U}(2n)$ defined by

$$\widetilde{q}(k,t) \, := \, \begin{cases} q_0(0) \, = \, q_1(0) \, = \, Q & \text{ if } \quad k = 0 \, , \, \, 0 \leqslant t \leqslant 1 \, , \\ q_0(k) & \text{ if } \quad 0 \leqslant k \leqslant \pi \, , \, \, t = 0 \, , \\ q_1(k) & \text{ if } \quad 0 \leqslant k \leqslant \pi \, , \, \, t = 1 \, , \\ \widetilde{Q}(t) & \text{ if } \quad k = \pi \, , \, \, 0 \leqslant t \leqslant 1 \, . \end{cases}$$

The map above can be extended, without obstructions, to a continuous map $\widetilde{q}:[0,\pi]\times [0,1]\to \mathbb{U}(2n)$ (still denoted with the same symbol). Indeed, by identifying $\mathfrak{d}([0,\pi]\times [0,1])$ with \mathbb{S}^1 , one can think of \widetilde{q} as a loop in $S\mathbb{U}(2n)$. Since the fundamental group of $S\mathbb{U}(2n)$ is trivial, it is possible to get the invoked continuous extension.

(3) The third step consists in a further extension $\widetilde{q}:[-\pi,\pi]\times[0,1]\to\mathbb{U}(2n)$ defined by

$$\widetilde{q}(k,t) \; := \; \begin{cases} \widetilde{q}(k,t) & \text{if} \quad (k,t) \in [0,\pi] \times [0,1] \;, \\ -\widetilde{q}(-k,t)^t & \text{if} \quad (k,t) \in [-\pi,0] \times [0,1] \;. \end{cases}$$

Since $q_0(0) = q_1(0) = Q$ is skew-symmetric, \widetilde{q} is a well-defined continuous map on $[-\pi,\pi]\times[0,1]$. Moreover, since $\widetilde{q}(\pi,t)=\widetilde{Q}(t)$ is chosen to be skew-symmetric for all $t\in[0,1]$, one gets that $\widetilde{q}(\pi,t)=\widetilde{q}(-\pi,t)$ for all $t\in[0,1]$. This shows that \widetilde{q} descends to a continuous map $\widetilde{q}:\mathbb{S}^1\times[0,1]\to\mathbb{U}(2n)$.

(4) The fourth and final step consists in the direct check of the fact that the resulting map \tilde{q} provides the required homotopy of sewing matrices.

Remark 3.9 (A small generalization). It is likely that Theorem 3.8 could be generalized in the case of a 1-dimensional \mathbb{Z}_2 -CW complex (X,τ) such that the fixed point set X^τ is a finite collection of points. In this case it is expected that two sewing matrices $q_0, q_1: X \to \mathbb{U}(2n)$ are linked by a homotopy within the space of sewing matrices if and only if the following two conditions are satisfied:

- (1) $[det[q_0]] = [det[q_1]]$ in $H^1_{\mathbb{Z}_2}(X, \mathbb{Z}) \simeq [X, \mathbb{U}(1)]_{inv}$;
- (2) $\nu_{q_0} = \nu_{q_1}$ in $H^1(X, Z)/H^1_{\mathbb{Z}_2}(X|X^{\tau}, \mathbb{Z}(1))$.

Note that in this case the invariants ν_{q_j} are well-defined since $H^2_{\mathbb{Z}_2}(X|X^{\tau},\mathbb{Z}(1))=0$ as proved in [DG4, Proposition 4.1].

3.3. **The role as an index.** As an application of Theorem 3.8, we can establish an *index-type* theorem for the invariant for topological insulators of class DIII described in Definition 2.7. By this we mean that the quantity v_q , which is purely *topological* by definition, can be computed by an *analytic* expression. The latter is nothing more than a suitable index of an appropriate Fredholm operator.

In order to prove the main result of this section we need first to introduce some notation. Let $Mat_{2n}(\mathbb{C})$ be the C^* -algebra of $2n \times 2n$ matrices. Every continuous function $f: \mathbb{S}^1 \to Mat_{2n}(\mathbb{C})$ defines a bounded multiplication operator M_f acting on the Hilbert space $L^2(\mathbb{S}^1) \otimes \mathbb{C}^{2n}$ according to the prescription

$$(M_f\psi)(k) := f(k)\psi(k), \qquad \psi \in L^2(\mathbb{S}^1) \otimes \mathbb{C}^{2n}.$$

By construction the map $f\mapsto M_f$ is a homomorphism of C^* -algebras. In particular one has $M_f^*=M_{f^*}$. Using the standard complex conjugation on \mathbb{C}^{2n} , one can endow $L^2(\mathbb{S}^1)\otimes\mathbb{C}^{2n}$ with the real structure (or antiunitary involution) K defined by $(K\psi)(k):=\overline{\psi(-k)}$. Given a (bounded) operator A on $L^2(\mathbb{S}^1)\otimes\mathbb{C}^{2n}$, we define the complex conjugate of A as $\overline{A}:=KAK$, and the transpose of A as $A^t:=(\overline{A})^*=KA^*K$. The operator A will be called *real* if $\overline{A}=A$, *self-adjoint* if $A=A^*$, *complex symmetric* if $A=A^t$, *skew-adjoint* if $A=A^*$ and *skew-complex symmetric* if $A=A^t$ [LZ, KKL].

The next result shows that the multiplication operators associated to sewing matrices are automatically skew-complex symmetric.

Lemma 3.10. Let $q: \mathbb{S}^1 \to \mathbb{U}(2n)$ be a map defined on the involutive space (\mathbb{S}^1, ι) which satisfies the sewing matrix condition

$$q(\iota(k)) = -q(k)^{t}, \quad \forall k \in \mathbb{S}^{1}.$$

Then, it holds true that

$$(M_q)^t = -M_q$$
.

Proof. One has that

$$\begin{array}{ll} (KM_{\mathfrak{q}}^*K\psi)(k) \; = \; \overline{(M_{\mathfrak{q}^*}K\psi)(-k)} \; = \; \overline{\mathfrak{q}(\iota(k))^*} \, \overline{(K\psi)(-k)} \\ \\ & = \; -\mathfrak{q}(k)\psi(k) = -(M_{\mathfrak{q}}\psi)(k) \end{array}$$

for every $\psi \in L^2(\mathbb{S}^1) \otimes \mathbb{C}^{2n}$. This completes the proof.

The Fourier modes $\varphi_m(k) := e^{i\,mk}$, with $m \in \mathbb{Z}$, provides a basis for $L^2(\mathbb{S}^1)$. Let $\mathcal{H}_m := \mathbb{C}[\varphi_m] \otimes \mathbb{C}^{2n}$ be the subspace of $L^2(\mathbb{S}^1) \otimes \mathbb{C}^{2n}$ generated by $\varphi_m \otimes v$ with $v \in \mathbb{C}^{2n}$ and define the subspace of positive frequencies $\mathcal{H}_+ := \bigoplus_{m \in \mathbb{N}} \mathcal{H}_m$. The associated orthogonal projection $\Pi : L^2(\mathbb{S}^1) \otimes \mathbb{C}^{2n} \to \mathcal{H}_+$ is known as the *Hardy projection*. Its adjoint is given by the inclusion $\Pi^* : \mathcal{H}_+ \to L^2(\mathbb{S}^1) \otimes \mathbb{C}^{2n}$. Since $K\varphi_m = \varphi_m$ for every $m \in \mathbb{Z}$, it follows that \mathcal{H}_+ is preserved by the real structure K. Therefore (with a little abuse of notation) one has that $K\Pi = \Pi K$ and $K\Pi^* = \Pi^*K$. The *Toeplitz operator* associated to the continuous function $f : \mathbb{S}^1 \to \mathrm{Mat}_{2n}(\mathbb{C})$ is the bounded operator T_f on \mathcal{H}_+ defined by the compression $T_f := \Pi M_f \Pi^*$. As a matter of fact, if f vanishes nowhere, then T_f is a Fredholm operator [Mur, Corollary 3.5.12] with the associated Noether-Fredholm index [Mur, Section 1.4]

$$ind(T_f) := dim Ker(T_f) - dim Ker(T_f^*) \in \mathbb{Z}$$
.

However, when T_f is skew-complex symmetric one automatically gets $\operatorname{ind}(T_f) = 0$. In fact from $(T_f)^{\operatorname{t}} = -T_f$ one immediately infers the equality of the dimension of $\operatorname{Ker}(T_f)$ and $\operatorname{Ker}(T_f^*)$. Therefore, to capture the topology of skew-complex symmetric Toeplitz operators one needs a secondary index. Following [AS, DS, SB], we will consider the \mathbb{Z}_2 -(Fredholm) index defined by

$$ind_{\mathbb{Z}_2}(T_f) := (-1)^{dim \, Ker(T_f)} \in \mathbb{Z}_2 .$$
 (3.4)

This index turns out to be a deformation invariant for a skew-complex symmetric Fredholm operator. To justify the latter claim, let us consider the unitary operator on $L^2(\mathbb{S}^1) \otimes \mathbb{C}^{2n}$ defined by

$$\label{eq:definition} \mathfrak{I} \; := \; \boldsymbol{1}_{L^2} \; \otimes \; \left(\begin{array}{cc} 0 & -\boldsymbol{1}_n \\ +\boldsymbol{1}_n & 0 \end{array} \right) \; .$$

One can check that $\mathfrak{I}^{-1}=\mathfrak{I}^*=-\mathfrak{I}$, and $\mathsf{K}\mathfrak{I}\mathsf{K}=\mathfrak{I}$, namely \mathfrak{I} is *real skew-symmetric*. Moreover, \mathfrak{I} commutes with the Hardy projection \mathfrak{I} , and therefore it provides a real skew-symmetric operator also on the subspace \mathfrak{H}_+ . If A is skew-complex symmetric Fredholm operator then $A':=\mathfrak{I}A$ satisfies the relation $\mathfrak{I}^*(A')^{\mathsf{t}}\mathfrak{I}=A'$, namely A' is an *odd symmetric operator* in the parlance of [DS, SB]. In view of the bijection induced by the mapping $A\mapsto \mathfrak{I}A$, and the fact that the dimension of the kernels of A and $\mathfrak{I}A$ are the same, we can use [SB, Theorem 2] to state:

Lemma 3.11. The set of skew-complex symmetric Toeplitz operators on the Hilbert space \mathcal{H}_+ is the disjoint union of two open and connected components labelled by $\operatorname{ind}_{\mathbb{Z}_2}$.

Therefore, as a consequence of Lemma 3.10 and Lemma 3.11, it turns out that the \mathbb{Z}_2 -index is the correct invariant to detect the topology of the Toeplitz operator $T_q = \Pi M_q \Pi^*$ associated to a sewing matrix $q: \mathbb{S}^1 \to \mathbb{U}(2n)$. Since a homotopy of sewing matrices induces a homotopy of the associated Toeplitz operators in the space of skew-complex

symmetric Fredholm operators it follows that $ind_{\mathbb{Z}_2}(T_q)$ provides a homotopy invariant of the sewing matrix q. The next result shows that this invariant coincides with ν_q .

Theorem 3.12 (Index theorem). Let $q: \mathbb{S}^1 \to \mathbb{U}(2n)$ be a map defined on the involutive space (\mathbb{S}^1, ι) which satisfies the sewing matrix condition

$$q(\iota(k)) = -q(k)^{t}, \quad \forall k \in \mathbb{S}^{1}.$$

Then, it holds true that

$$\nu_q = ind_{\mathbb{Z}_2}(T_q)$$
.

In particular, this implies that

$$\frac{\operatorname{Pf}[q(\pi)]}{\operatorname{Pf}[q(0)]} \, \frac{\det[q(0)]^{\frac{1}{2}}}{\det[q(\pi)]^{\frac{1}{2}}} \, = \, (-1)^{\dim \operatorname{Ker}(T_q)} \, .$$

where the branch $k \mapsto det[q(k)]^{\frac{1}{2}}$ must be chosen continuously between the two fixed points $k = 0, \pi$.

Proof. It is enough to check the coincidence of the invariants for a representative of each of the *two* homotopy classes of sewing matrices (*cf.* Theorem 3.8). One homotopy class is represented by the constant map $q_+(k) = Q$ for all $k \in \mathbb{S}^1$, with Q defined by (3.3). An immediate application of Theorem 3.5 provides $\nu_{q_+} = +1$. On the other hand, since M_{q_0} is the multiplication operator by a constant invertible matrix, it follows that $T_{q_+} := \Pi M_{q_+} \Pi^*$ is invertible in \mathcal{H}_+ . As a consequence the dimension of $Ker(T_{q_+})$ is zero, and in turn $Ind_{\mathbb{Z}_2}(T_{q_+}) = +1$. This shows that $\nu_{q_+} = Ind_{\mathbb{Z}_2}(T_{q_+})$. The second homotopy class can be represented by

$$q_-(k) \, := \, \left(\begin{array}{c|ccc} 0 & + \, e^{+\, i \, k} & 0 & 0 \\ - \, e^{-\, i \, k} & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & -\mathbf{1}_{n-1} \\ 0 & 0 & +\mathbf{1}_{n-1} & 0 \end{array} \right) \; , \qquad k \in \mathbb{S}^1 \; .$$

Indeed, from

$$\begin{split} Pf\big[q_-(k)\big] &= Pf\left[\left(\begin{array}{cc} 0 & +e^{+\operatorname{i}\,k} \\ -e^{-\operatorname{i}\,k} & 0 \end{array} \right) \right] Pf\left[\left(\begin{array}{cc} 0 & -\mathbf{1}_{n-1} \\ +\mathbf{1}_{n-1} & 0 \end{array} \right) \right] \\ &= (-1)^{\frac{n(n-1)}{2}} Pf\left[\left(\begin{array}{cc} 0 & +e^{+\operatorname{i}\,k} \\ -e^{-\operatorname{i}\,k} & 0 \end{array} \right) \right] \end{split}, \qquad k=0,\pi \end{split}$$

one gets that

$$Pf[q_{-}(0)] = (-1)^{\frac{n(n-1)}{2}} = -Pf[q_{-}(\pi)],$$

and in turn one has $\nu_{q_-}=-1$ as a consequence of Theorem 3.5 and the constancy of the determinant

$$det\big[q_-(k)\big] \ = \ det\left[\left(\begin{array}{cc} 0 & -\textbf{1}_{n-1} \\ +\textbf{1}_{n-1} & 0 \end{array}\right)\right] \ = \ (-1)^{n-1} \ , \qquad \forall k \in \mathbb{S}^1 \ .$$

To compute the kernel of T_{q_-} let us consider the orthogonal decomposition $\mathcal{H}_+ = \mathcal{H}_{+,1} \oplus \mathcal{H}_{+,2}$ induced by the decomposition

$$\mathbb{C}^{2n} = \mathbb{C}[e_1, e_2] \oplus \mathbb{C}[e_3, \dots, e_{2n}] \simeq \mathbb{C}^2 \oplus \mathbb{C}^{2(n-1)}$$

where $\{e_1,\ldots,e_{2n}\}$ denotes the canonical basis of \mathbb{C}^{2n} . Evidently T_{q_-} is injective on $\mathcal{H}_{+,2}$. On the other hand, a generic element of $\psi\in\mathcal{H}_{+,1}$ can be represented in terms of the Fourier modes φ_m as

$$\psi \; = \; \left(\sum_{\mathfrak{m} \in \mathbb{N}} \alpha_{\mathfrak{m},1} \; \varphi_{\mathfrak{m}} \otimes e_1\right) + \left(\sum_{\mathfrak{m} \in \mathbb{N}} \alpha_{\mathfrak{m},2} \; \varphi_{\mathfrak{m}} \otimes e_2\right) \; , \qquad \alpha_{\mathfrak{m},\ell} \in \mathbb{C} \; , \quad \ell = 1,2$$

and a direct computation provides

$$M_{q_{-}}\Pi^{*}\psi \;=\; \left(\sum_{\mathfrak{m}\in\mathbb{N}}-\mathfrak{a}_{\mathfrak{m},1}\; \varphi_{\mathfrak{m}-1}\otimes e_{2}\right)+\left(\sum_{\mathfrak{m}\in\mathbb{N}}\mathfrak{a}_{\mathfrak{m},2}\; \varphi_{\mathfrak{m}+1}\otimes e_{1}\right)\;.$$

Therefore, one gets that the equation $T_{q_-}\psi$ has the non-trivial solution $\psi=\alpha(\varphi_1\otimes e_1)$, $\alpha\in\mathbb{C}$, and as a consequence the dimension of $Ker(T_{q_-})$ is one. This provides $ind_{\mathbb{Z}_2}(T_{q_-1})=-1$ and in turn $\nu_{q_-}=ind_{\mathbb{Z}_2}(T_{q_-})$. This concludes the proof.

Remark 3.13 (The bulk-edge correspondence). In the jargon of the theory of topological insulator an index formula of the type of Theorem 3.12 is referred as bulk-edge correspondence [PS]. In fact the topological invariant v_q is derived by the data concerning the system in the bulk, while the index $\operatorname{ind}_{\mathbb{Z}_2}(T_q)$ is computed by means of a Toeplitz operator which encodes the existence of an edge.

Remark 3.14 (The interpretation in K-theory). By combining the map $A\mapsto \mathbb{J}A$ introduced above with the argument used in the proof of [SB, Theorem 2] one obtain a homeomorphism between the set of skew-complex symmetric Toeplitz operators on the Hilbert space \mathcal{H}_+ and the classifying space $\mathcal{F}^2(\mathcal{H}_\mathbb{R})$ introduced in [AS]. As a consequence the K-group $KR^{-2}(*)$ can be identified with the set of homotopy class of skew-complex symmetric Fredholm operators, and the \mathbb{Z}_2 -index (3.4) realizes the isomorphism $KR^{-2}(*)\simeq \mathbb{Z}_2$. The construction of the skew-complex symmetric Fredholm operator T_q from q extends to a homomorphism of KR-theory

$$T \; : \; KR^{-3}(\mathbb{S}^1,\iota) \; \longrightarrow \; KR^{-2}(*) \; ,$$

which is bijective by the results obtained so far. Notice that the push-forward

$$\pi_* \; : \; KR^{-3}(\mathbb{S}^1,\iota) \; \longrightarrow \; KR^{-2}(*)$$

along the projection $\pi: (\mathbb{S}^1, \iota) \to *$ is also a bijection. Thus, we have $\pi_* = T$ which means that the construction with the Toeplitz operator realizes the push-forward.

3.4. **The relation with the KR-theory.** In this section we will prove that in the one-dimensional case the homomorphism (1.9) is indeed an isomorphism. In Lemma 3.1 it

has been proved that

$$H^1(\mathbb{S}^1,\mathbb{Z})/H^1_{\mathbb{Z}_2}\left(\mathbb{S}^1|(\mathbb{S}^1)^{\iota},\mathbb{Z}(1)\right) \simeq \mathbb{Z}_2$$
.

On the other hand, by using formula (B.4) and Table B.1 in [DG1] one gets

$$\begin{split} \mathsf{KR}^{-3}(\mathbb{S}^1,\iota) \; &\simeq \; \mathsf{KR}^{-3}(*) \; \oplus \; \mathsf{KR}^{-2}(*) \\ &\simeq \; \mathfrak{0} \; \oplus \; \mathbb{Z}_2 \; \simeq \; \mathbb{Z}_2 \; . \end{split}$$

Theorem 2.11 and Remark 2.12 establish the existence of the well-defined homomorphism

$$\nu : KR^{-3}(\mathbb{S}^1, \iota) \longrightarrow H^1(\mathbb{S}^1, \mathbb{Z})/H^1_{\mathbb{Z}_2}(\mathbb{S}^1 | (\mathbb{S}^1)^{\iota}, \mathbb{Z}(1)) . \tag{3.5}$$

Proposition 3.15. The homomorphism (3.5) is an isomorphism.

Proof. From the computations above one has that the homomorphism (3.5) has the form $\nu: \mathbb{Z}_2 \to \mathbb{Z}_2$. Moreover, in the proof of Theorem 3.12 we construct two sewing matrices q_{\pm} such that $\nu_{q_{\pm}} = \pm 1$. Let $[q_{\pm}] \in KR^{-3}(\mathbb{S}^1, \iota)$ be the class of the band insulators represented by the sewing matrix q_{\pm} . Since the homomorphism (3.5) corresponds to $\nu: [q_{\pm}] = \nu_{q_{\pm}}$ one gets the result.

3.5. The relation with the classification of "Real" gerbes. The aim of this section is to clarify the relation between the invariant ν and a *gerbe invariant* introduced in [GT].

Let us first introduce the gerbe invariant in a way slightly different from the original definition in [GT]. First of all, let us recall that a gerbe $\mathscr G$ over the space SU(2n) is classified by its Dixmier-Douady class

$$\mathfrak{D}\mathfrak{D}[\mathscr{G}] \;\in\; H^3\big(S\mathbb{U}(2\mathfrak{n}),\mathbb{Z}\big) \;\simeq\; \mathbb{Z} \;.$$

The basic gerbe \mathscr{G} over SU(2n) is a representative of the class $\mathcal{DD}[\mathscr{G}] \simeq 1$ which generates $H^3(SU(2n),\mathbb{Z})$. The basic gerbe \mathscr{G} admits only two "Real" structures (up to stable equivalences) over the involutive space $(SU(2n),\theta)$. We will denote with \mathscr{G}_+ and \mathscr{G}_- the basic gerbe equipped with the two different 'Real" structures. The two "Real" basic gerbes are classified by their "Real" Dixmier-Douady class

$$\mathfrak{D}\mathfrak{D}_R[\mathscr{G}_{\pm}] \;\in\; H^3_{\mathbb{Z}_2}\big(S\mathbb{U}(2n),\mathbb{Z}(1)\big)$$

and the two classes differ by an element of in $H^3_{\mathbb{Z}_2}(*,\mathbb{Z}(1))\subset H^3_{\mathbb{Z}_2}(S\mathbb{U}(2n),\mathbb{Z}(1))$, where $*\in S\mathbb{U}(2n)$ is any fixed point under the involution θ . As a result, we get a well-defined element

$$\widehat{\mathfrak{DD}}_R[\mathscr{G}_{\pm}] \;\in\; H^3_{\mathbb{Z}_2}\big(S\mathbb{U}(2\mathfrak{n}),\mathbb{Z}(1)\big)/H^3_{\mathbb{Z}_2}\big(*,\mathbb{Z}(1)\big)\;,$$

independently of the choice of the "Real" structures on \mathfrak{G} , as well as of the fixed point $* \in SU(2n)$.

Let (X, τ) be an involutive space which admits a fixed point $* \in X$, and $q: X \to SU(2n)$ an equivariant map. Such a map induces by pull-back the homomorphism

$$q^* : H^3_{\mathbb{Z}_2}\big(S\mathbb{U}(2n),\mathbb{Z}(1)\big)/H^3_{\mathbb{Z}_2}\big(*,\mathbb{Z}(1)\big) \ \longrightarrow \ H^3_{\mathbb{Z}_2}\big(X,\mathbb{Z}(1)\big)/H^3_{\mathbb{Z}_2}\big(*,\mathbb{Z}(1)\big) \ .$$

The *gerbe invariant* of the map $q: X \to SU(2n)$ is by definition the pull-back

$$dd_R[q] \; := \; q^* \widehat{\mathcal{D} \mathcal{D}}_R[\mathscr{G}_{\pm}] \; .$$

This is an invariant of the equivariant homotopy class of q. Therefore, dd_R provides a well-defined map (still denoted with the same symbol)

$$dd_R : [X, SU(2n)]_{\mathbb{Z}_2} \longrightarrow H^3_{\mathbb{Z}_2}(X, \mathbb{Z}(1))/H^3_{\mathbb{Z}_2}(*, \mathbb{Z}(1)).$$

The invariants $dd_R[q]$ and ν_q take values in different groups, and therefore we cannot generally compare these two invariants. However, in the case of the involutive space (\mathbb{S}^1, ι) , one has that

$$H^3_{\mathbb{Z}_2}(\mathbb{S}^1,\mathbb{Z}(1))/H^3_{\mathbb{Z}_2}(*,\mathbb{Z}(1)) \simeq \mathbb{Z}_2$$

in view of

$$H^3_{\mathbb{Z}_2}\big(\mathbb{S}^1,\mathbb{Z}(1)\big) \; \simeq \; \tilde{H}^3_{\mathbb{Z}_2}\big(\mathbb{S}^1,\mathbb{Z}(1)\big) \oplus H^3_{\mathbb{Z}_2}\big(*,\mathbb{Z}(1)\big) \; \simeq \; \mathbb{Z}_2 \oplus \mathbb{Z}_2$$

and $H^3_{\mathbb{Z}_2}(*,\mathbb{Z}(1))\simeq \mathbb{Z}_2$. This allows us to regard $dd_R[q]\in \mathbb{Z}_2$. Since also $\nu_q\in \mathbb{Z}_2$ we can compare the two \mathbb{Z}_2 -invariants $dd_R[q]$ and ν_q for equivariant maps $q:\mathbb{S}^1\to S\mathbb{U}(2n)$ over the involutive circle (\mathbb{S}^1,ι) .

Theorem 3.16 (Gerbe invariant). Let $q: \mathbb{S}^1 \to S\mathbb{U}(2n)$ be an equivariant map defined on the involutive space (\mathbb{S}^1, ι) satisfying the sewing matrix condition

$$q\big(\iota(k)\big)\,=\,-q(k)^{\mathtt{t}}\;,\qquad\forall\;k\in\mathbb{S}^1$$
 .

Then, it holds true that

$$\nu_q = dd_R[q]$$
.

Proof. The proof is similar to that of Theorem 3.12. By Lemma 3.6 and Theorem 3.8, the invariant ν induces a bijection $[\mathbb{S}^1, S\mathbb{U}(2n)]_{\mathbb{Z}_2} \simeq \mathbb{Z}_2$. In the proof of Theorem 3.12, we constructed two equivariant maps $q_{\pm}: \mathbb{S}^1 \to S\mathbb{U}(2n)$ such that $\nu_{q_{\pm}} = \pm 1$. Therefore, the claim is proved by showing that $dd_R[q_{\pm}] = \pm 1$. For the computation, we make use of an expression of $dd_R[q_{\pm}] \in \mathbb{Z}_2$ in terms of the *sign invariant* of "Real" gerbes [GT, Definition 4.4]. In general, for a "Real" gerbe \mathscr{G} on an involutive space (X, τ) and a path connected subspace $Y = Y^{\tau} \subset X^{\tau}$ in the fixed point set, there is a well-defined sign $\sigma(\mathscr{G}, Y) \in \mathbb{Z}_2$. Actually, this is the component in $H^0(Y, \mathbb{Z}_2) \simeq \mathbb{Z}_2$ of the "Real" Dixmier-Douady class $\mathcal{DD}_R(\mathscr{G}|_Y) \in H^3_{\mathbb{Z}_2}(Y, \mathbb{Z}(1))$ of the restricted gerbe $\mathscr{G}|_Y$ with respect to the decomposition

$$H^3_{\mathbb{Z}_2}\big(Y,\mathbb{Z}(1)\big) \; \simeq \; H^0(Y,\mathbb{Z}_2) \; \oplus \; H^2(Y,\mathbb{Z}_2)$$

as shown in [GT, Lemma B.1]. The fixed point set of the involutive space (\mathbb{S}^1, ι) consists of the points $k_+ = 0$ and $k_- = \pi$. Let \mathscr{G}_{\pm} be the two "Real" basic gerbes over $\mathbb{SU}(2n)$ and $q^*\mathscr{G}_{\pm}$ the pull-back gerbes over \mathbb{S}^1 . In view of [GT, Corolary 4.7] one has that

$$q^* \mathcal{D} \mathcal{D}_{R}(\mathscr{G}_{\pm}) = \mathcal{D} \mathcal{D}_{R}(q^* \mathscr{G}_{\pm}) = \sigma(q^* \mathscr{G}_{\pm}, \{k_{+}\}) \sigma(q^* \mathscr{G}_{\pm}, \{k_{-}\})$$
$$= \sigma(\mathscr{G}_{+}, \{q(k_{+})\}) \sigma(\mathscr{G}_{+}, \{q(k_{-})\})$$

is computed as a product of signs. Since the right-hand side of the equation above is not affected to the passage to the quotient which defines the map dd_R , one gets

$$dd_{R}[q] = \sigma(\mathscr{G}_{\sharp}, \{q(k_{+})\}) \ \sigma(\mathscr{G}_{\sharp}, \{q(k_{-})\}) \in \mathbb{Z}_{2}$$

where \mathscr{G}_{\sharp} stands irrelevantly for \mathscr{G}_{+} or \mathscr{G}_{-} . As seen in Lemma 3.7, the fixed point set of $(SU(2n),\theta)$ has two connected components. As it is pointed out in [GT, Remark 5.7], the sign invariant of \mathscr{G}_{\sharp} is constant on the connected components of $SU(2n)^{\theta}$ and differs on each of the two components. As a result, $dd_{R}[q]=+1$ if and only if $q(k_{+})$ and $q(k_{-})$ are in the same connected component and $dd_{R}[q]=-1$ if and only if $q(k_{+})$ and $q(k_{-})$ are in different connected components. This immediately proves that $dd_{R}[q_{\pm}]=\pm 1$ and the proof is over.

4. THE TWO-DIMENSIONAL CASE

The analysis of physical systems shows that there are (at least) two interesting definitions for two-dimensional band insulators of class DIII which differ for the involutive base space. By borrowing the discussion in [DG1, Section 2], we can distinguish between the *Dirac* (or *free*) case and the *Bloch* (or *periodic*) case. In the Dirac case the involutive space is (\mathbb{S}^2, ι) where

$$\mathbb{S}^2 \; := \; \left\{ (x_0, x_1, x_2) \in \mathbb{R}^3 \; \middle|\; x_0^2 + x_1^2 + x_2^2 = 1 \right\}$$

is the two-dimensional sphere with involution given by

$$\iota(x_0,x_1,x_2) \; := \; (x_0,-x_1,-x_2) \; , \qquad \forall \; (x_0,x_1,x_2) \in \mathbb{S}^2 \; .$$

In the Bloch case the involutive space is (\mathbb{T}^2, ι) where

$$\mathbb{T}^2 := \mathbb{S}^1 \times \mathbb{S}^1 = (\mathbb{R}/2\pi\mathbb{Z}) \times (\mathbb{R}/2\pi\mathbb{Z})$$

is the two-dimensional torus with involution given by

$$\iota(k_1, k_2) := (-k_1, -k_2), \quad \forall (k_1, k_2) \in \mathbb{T}^2.$$

In short, one has that $(\mathbb{T}^2, \iota) = (\mathbb{S}^1, \iota) \times (\mathbb{S}^1, \iota)$ is given by two copies of the involutive space for the one-dimensional case.

According to the terminology currently used in the theory of topological insulators, the topological phases detected in the Dirac case are called *strong*. Usually, the Bloch case present a richer family of topological phases. Some of these are inherited from the Dirac case via the pull-back of the equivariant map $\pi_0: (\mathbb{T}^2, \iota) \to (\mathbb{S}^2, \iota)$ described in [DG2, eq. (4.14)] and are still called strong. The remaining topological phases are known as *weak*.

It is worth noting that both spaces (\mathbb{S}^2, ι) and (\mathbb{T}^2, ι) meet the properties (a) and (b) listed in Assumption 2.5. However, the isomorphisms

$$H^2_{\mathbb{Z}_2}\left(\mathbb{S}^2|(\mathbb{S}^2)^{\iota},\mathbb{Z}(1)\right) \simeq \mathbb{Z}_2 \simeq H^2_{\mathbb{Z}_2}\left(\mathbb{T}^2|(\mathbb{T}^2)^{\iota},\mathbb{Z}(1)\right),$$

and the bijectivity of the FKMM invariant [DG2, Theorem 1.2], imply that the property (c) of Assumption 2.5 is not automatically guaranteed. This fact makes the analysis of the two-dimensional case slightly more complicated than the one-dimensional case.

4.1. **Strong invariant: the Dirac case.** Let us start by observing that for a two-dimensional sphere it holds true that $H^1(\mathbb{S}^2, \mathbb{Z}) = 0$ which immediately implies that

$$H^1(\mathbb{S}^2,\mathbb{Z})/H^1_{\mathbb{Z}_2}(\mathbb{S}^2|(\mathbb{S}^2)^\iota,\mathbb{Z}(1)) \; = \; 0 \; .$$

Therefore, in the Dirac case the map ν in (1.9) amounts to the trivial homomorphism, and for this reason it is unable to capture the topology of the related band insulator of class DIII.

On the other hand, from equations (B.3) and (B.9) and Tables B.1 and B.3 in [DG1] one gets

$$\mathsf{KR}^{-3}(\mathbb{S}^2, \iota) \simeq \widetilde{\mathsf{KR}}^{-3}(\mathbb{S}^2, \iota) \oplus \mathsf{KR}^{-3}(*)$$

 $\simeq \widetilde{\mathsf{KR}}^{0}(\mathbb{S}^6, \iota) \oplus 0 \simeq \mathbb{Z}_2.$

The latter equation shows that there is a non-trivial topology for band insulators of class DIII over (\mathbb{S}^2 , ι).

The topology of $KR^{-3}(\mathbb{S}^2, \iota)$ can be detected via the FKMM invariant by using the map κ in (2.13). First of all let us observe that

$$\text{Vec}_{\mathfrak{Q},0}^{2\mathfrak{n}}(\mathbb{S}^2,\iota) \ \simeq \ \text{Vec}_{\mathfrak{Q}}^2(\mathbb{S}^2,\iota) \ \simeq \ H^2_{\mathbb{Z}_2}\big(\mathbb{S}^2|(\mathbb{S}^2)^\iota,\mathbb{Z}(1)\big) \ \simeq \ \mathbb{Z}_2 \ .$$

The first isomorphism is a consequence of [DG2, Proposition 4.1] which provides the equality $\text{Vec}_{\mathfrak{Q},0}^{2n}(\mathbb{S}^2,\iota) = \text{Vec}_{\mathfrak{Q}}^{2n}(\mathbb{S}^2,\iota)$ (*i. e.* every "Quaternionic" vector bundle over (\mathbb{S}^2,ι) can be built over a product vector bundle) and [DG2, Corollary 2.1] which describes the stable range for low-dimensional "Quaternionic" vector bundles. The second isomorphism is provided by the FKMM invariant [DG2, Theorem 1.2]. With this information we can prove the following result:

Proposition 4.1 (Two-dimensional strong invariant). The homomorphism

$$\kappa \; : \; KR^{-3}(\mathbb{S}^2, \iota) \; \longrightarrow \; H^2_{\mathbb{Z}_2}\big(\mathbb{S}^2|(\mathbb{S}^2)^\iota, \mathbb{Z}(1)\big)$$

described in Theorem 2.11 is indeed an isomorphism.

Proof. In view of Theorem 2.11 and the results showed above we are considering a well-defined homomorphism of the form $\kappa: \mathbb{Z}_2 \to \mathbb{Z}_2$. Therefore, to prove the claim it is enough to show that κ is surjective. Let us use the additive representation of $\mathbb{Z}_2 \simeq \mathbb{Z}/2\mathbb{Z}$. The two distinct classes in $\text{Vec}_{\mathfrak{Q}}^2(\mathbb{S}^2, \mathfrak{t}) \simeq \mathbb{Z}_2$ can be represented by the "Quaternionic" vector bundles $\mathscr{E}_{\mathfrak{q}_0}$ and $\mathscr{E}_{\mathfrak{q}_1}$ obtained by endowing the product bundles $\mathbb{S}^2 \times \mathbb{C}^2$ with two inequivalent "Quaternionic" structures $\Theta_{\mathfrak{q}_0}$ and $\Theta_{\mathfrak{q}_1}$ associated to two sewing matrices $\mathfrak{q}_0, \mathfrak{q}_1: \mathbb{S}^2 \to \mathbb{U}(2)$. Let \mathfrak{q}_0 be the constant map defined as in Theorem 2.8 (2). Then, $\mathscr{E}_{\mathfrak{q}_0}$ turns out to be the trivial "Quaternionic" vector bundle with FKMM invariant $\kappa(\mathscr{E}_{\mathfrak{q}_0}) = 0$.

An explicit realization for the sewing matrix q_1 is described in [DG2, eq. (4.9)]. In this case \mathcal{E}_{q_1} provides a representative for the non trivial "Quaternionic" class and $\kappa(\mathcal{E}_{q_1}) = 1$. Now, consider the Hamiltonians $H_i : \mathbb{S}^2 \to \mathfrak{Herm}(\mathbb{C}^4)^{\times}$ given by

$$H_{\mathfrak{i}}(x) \;:=\; \left(\begin{array}{cc} 0 & q_{\mathfrak{i}}(x)^{-1} \\ q_{\mathfrak{i}}(x) & 0 \end{array}\right)\;, \qquad x \in \mathbb{S}^2\;,\;\; \mathfrak{i} = 0,1\;.$$

Then $[\mathbb{S}^2 \times \mathbb{C}^4, H_1, H_0]$ is a well-defined class in $KR^{-3}(\mathbb{S}^2, \iota)$ and by equation (2.14) one gets $\kappa([\mathbb{S}^2 \times \mathbb{C}^4, H_1, H_0]) = 1$. This concludes the proof.

Let $H: \mathbb{S}^2 \to \mathfrak{Herm}(\mathbb{C}^{4n})^{\times}$ be a band insulator of class DIII over (\mathbb{S}^2, ι) represented in a standard way. Denote with H_0 the trivial insulator constructed with the constant sewing matrix $q_0: \mathbb{S}^2 \to \mathbb{U}(2n)$ described in item (2) of Theorem 2.8. Then

$$\nu^s(H) \,:=\, \kappa\big([\mathbb{S}^2\times\mathbb{C}^{4n},H,H_0]\big)$$

will be called the *strong invariant* of H. This is nothing more than the FKMM invariant $\kappa(\mathscr{E}_{q_H})$ of the "Quaternionic" vector bundle associated to H with the sewing matrix q_H . In view of Proposition 4.1 this invariant totally captures the topology of two-dimensional band insulators of class DIII in the Dirac case.

4.2. **Weak-strong invariant: the Bloch case.** In order to study the classification of two-dimensional band insulator H of class DIII in the Bloch case we will specialize Theorem 2.11 for the involutive space (\mathbb{T}^2 , ι).

By using [DG1, eq. (B.7)] one gets

$$KR^{-3}(\mathbb{T}^{2}, \iota) \simeq KR^{-3}(*) \oplus (KR^{-2}(*))^{\oplus 2} \oplus KR^{-1}(*)$$

$$\simeq 0 \oplus (\mathbb{Z}_{2})^{\oplus 2} \oplus \mathbb{Z}_{2}$$

$$\simeq \mathbb{Z}_{2} \oplus \mathbb{Z}_{2} \oplus \mathbb{Z}_{2}.$$

$$(4.1)$$

On the other hand, the product structure $(\mathbb{T}^2, \iota) := (\mathbb{S}^1, \iota) \times (\mathbb{S}^1, \iota)$ and the use of the Künneth formula, provide

$$H^{1}(\mathbb{T}^{2},\mathbb{Z})/H^{1}_{\mathbb{Z}_{2}}(\mathbb{T}^{2}|(\mathbb{T}^{2})^{\iota},\mathbb{Z}(1)) \simeq \left(H^{1}(\mathbb{S}^{1},\mathbb{Z})/H^{1}_{\mathbb{Z}_{2}}(\mathbb{S}^{1}|(\mathbb{S}^{1})^{\iota},\mathbb{Z}(1))\right)^{\oplus 2}$$

$$\simeq (\mathbb{Z}_{2})^{\oplus 2} = \mathbb{Z}_{2} \oplus \mathbb{Z}_{2}.$$

$$(4.2)$$

Finally, from [DG2, Theorem 1.2] one knows that

$$\text{Vec}_{\mathfrak{D}}^{2\mathfrak{n}}(\mathbb{T}^2,\iota) \; \simeq \; \text{Vec}_{\mathfrak{D}}^2(\mathbb{T}^2,\iota) \; \simeq \; H^2_{\mathbb{Z}_2}(\mathbb{S}^1|(\mathbb{S}^1)^\iota,\mathbb{Z}(1)) \; \simeq \; \mathbb{Z}_2 \; .$$

Lemma 4.2. The homomorphism

$$\kappa \; : \; KR^{-3}(\mathbb{T}^2,\iota) \; \longrightarrow \; H^2_{\mathbb{Z}_2}\big(\mathbb{T}^2|(\mathbb{T}^2)^\iota,\mathbb{Z}(1)\big)$$

described in Theorem 2.11 is surjective.

Proof. The proof follows exactly as in Proposition 4.1 by constructing a class $[\mathbb{T}^2 \times \mathbb{C}^4, H_1', H_0']$ such that $\kappa([\mathbb{T}^2 \times \mathbb{C}^4, H_1', H_0']) = 1$. In this case the Hamiltonians H_1'

and H_0' can be obtained form the Hamiltonians H_1 and H_0 introduced in the proof of Proposition 4.1 via the pull-back under the equivariant map $\pi_0: (\mathbb{T}^2, \iota) \to (\mathbb{S}^2, \iota)$ described in [DG2, eq. (4.14)]. More precisely one has that $H_i'(k) := H_i(\pi_0(k))$ for i = 0, 1. This concludes the proof.

Let $H: \mathbb{T}^2 \to \mathfrak{Herm}(\mathbb{C}^{4n})^{\times}$ be a band insulator of class DIII over (\mathbb{T}^2, ι) represented in a standard way. As in the Dirac case we can define the *strong invariant*

$$v^{s}(H) := \kappa([\mathbb{T}^{2} \times \mathbb{C}^{4n}, H, H'_{0}]) \tag{4.3}$$

where H_0 is the trivial insulator constructed with the constant sewing matrix $q_0:\mathbb{T}^2\to \mathbb{U}(2n)$ described in item (2) of Theorem 2.8. Since this invariant coincides with the FKMM invariant $\kappa(\mathscr{E}_{q_H})$ of the "Quaternionic" vector bundle associated to H with the sewing matrix q_H , one can use the Fu–Kane–Mele index proved in [DG2, Theorem 4.2] to compute ν .

Proposition 4.3 (Fu–Kane–Mele formula for the strong invariant). The strong invariant v^s defined by (4.3) can be computed by the formula

$$\nu^{s}(H) = \prod_{k \in (\mathbb{T}^{2})^{\iota}} \frac{Pf[q_{H}(k)]}{\det[q_{H}(k)]^{\frac{1}{2}}} \in \{\pm 1\}$$

$$(4.4)$$

where the branch $k\mapsto det[q_H(k)]^{\frac{1}{2}}$ must be chosen continuously and that ν^s must be understood as a map with values into the multiplicative group $\mathbb{Z}_2\simeq\{\pm 1\}$.

Proof. According to [DG2, Theorem 4.2] one has that

$$v^{s}(H) = \prod_{k \in (\mathbb{T}^{2})^{t}} \frac{Pf[q_{H}(k)]}{p_{H}(k)}$$

$$(4.5)$$

where $p_H: \mathbb{T}^2 \to \mathbb{U}(1)$ meets $det[q_H(k)] = p_H(\iota(k))p_H(k)$ for all $k \in \mathbb{T}^2$. To obtain (4.4) form (4.5) we need to show that one can replace p_H with a map $p'_H: \mathbb{T}^2 \to \mathbb{U}(1)$ such that $det[q_H(k)] = p'_H(k)^2$ for all $k \in \mathbb{T}^2$. The following fact follows by observing that every continuous map $s: \mathbb{T}^2 \to \mathbb{U}(1)$ such that $s(k)^2 = det[q_H(k)]$ automatically satisfies $s(k) = s(\iota(k))$ for all $k \in \mathbb{T}^2$. The key to the proof is that every continuous map $f: \mathbb{T}^2 \to \mathbb{U}(1)$ can be uniquely expressed as

$$f(k_1,k_2) \; = \; e^{\,i\,\left(\alpha + \widetilde{f}(k_1,k_2) + n_1 k_1 + n_2 k_2\right)} \; , \qquad (k_1,k_2) \in \mathbb{T}^2$$

where $n_1, n_2 \in \mathbb{Z}$, $\alpha \in \mathbb{R}/2\pi\mathbb{Z}$ and $\widetilde{f}: \mathbb{T}^2 \to \mathbb{R}$ is a continuous function such that $\widetilde{f}(0,0)=0$. The integers n_1 and n_2 provide the winding numbers of f around the two standard homology cycles of \mathbb{T}^2 and α is determined by $f(0,0)=e^{i\,\alpha}$. By applying the same representation to $det[q_H]$ one gets

$$det[q_H(k_1,k_2)] \ = \ e^{i \left(\alpha_H + \widetilde{f}_H(k_1,k_2) + n_1 k_1 + n_2 k_2\right)} \ , \qquad (k_1,k_2) \in \mathbb{T}^2$$

By the invariance $det[q_H(k_1, k_2)] = det[q_H(-k_1, -k_2)]$, and the continuity of \widetilde{f}_H it follows that $n_1 = n_2 = 0$ and $\widetilde{f}_H(k_1, k_2) = \widetilde{f}_H(-k_1, -k_2)$. As a result, every continuous

function whose square coincides with det[qH] must be of the form

$$s(k_1, k_2) = \pm e^{i(\alpha_s + \frac{1}{2}\tilde{f}(k_1, k_2))},$$

with $\alpha_s=\frac{\alpha_H}{2}\in\mathbb{R}/2\pi\mathbb{Z}$. In particular one gets that $s(-k_1,-k_2)=s(k_1,k_2)$ as anticipated. \Box

It is worth noting equation (4.4) agrees with formula [CTSR, eq. (3.69)].

From Lemma 4.2 and (4.1) one infers that

$$Ker(\kappa) \simeq KR^{-3}(\mathbb{T}^2, \iota)/\mathbb{Z}_2 = \mathbb{Z}_2 \oplus \mathbb{Z}_2$$
.

Theorem 2.11 establishes the existence the of homomorphism

$$\nu : \operatorname{Ker}(\kappa) \longrightarrow H^{1}(\mathbb{T}^{2}, \mathbb{Z})/H^{1}_{\mathbb{Z}_{2}}(\mathbb{T}^{2}|(\mathbb{T}^{2})^{\iota}, \mathbb{Z}(1))$$

$$(4.6)$$

which is of the form $\nu : \mathbb{Z}_2 \oplus \mathbb{Z}_2 \to \mathbb{Z}_2 \oplus \mathbb{Z}_2$ in view of (4.2). This homomorphism turns out to be surjective, and therefore bijective.

Lemma 4.4. The homomorphism (4.6) described in Theorem 2.11 is an isomorphism.

Lemma 4.4 follows as a special case of Proposition 4.5 below. For that, let us introduce the equivariant inclusions

$$\phi_i : (\mathbb{S}^1, \iota) \hookrightarrow (\mathbb{T}^2, \iota), \qquad j = 1, 2$$

defined by $\phi_1(k)=(k,1)$ and $\phi_2(k)=(1,k)$. To every class $[\mathbb{T}^2\times\mathbb{C}^{2n},H_0,H_1]\in KR^{-3}(\mathbb{T}^2,\iota)$ we can associate via pull-back two classes

$$[\mathbb{T}^2 \times \mathbb{C}^{2n}, H_0, H_1]_{\mathfrak{f}} \; := \; [\mathbb{S}^1 \times \mathbb{C}^{2n}, \varphi_{\mathfrak{f}}^* H_0, \varphi_{\mathfrak{f}}^* H_1] \; \in \; KR^{-3}(\mathbb{S}^1, \iota) \; , \qquad \mathfrak{f} = 1, 2$$

where $\phi_j^*H_i(k):=H_i(\phi_j(k))$ for all $k\in\mathbb{S}^1$, j=1,2 and i=0,1. In view of Proposition 3.15 one can introduce the homomorphism

$$\nu^{\mathrm{w}} : \mathsf{KR}^{-3}(\mathbb{T}^2, \iota) \longrightarrow \left(\mathsf{H}^1(\mathbb{S}^1, \mathbb{Z}) / \mathsf{H}^1_{\mathbb{Z}_2} \big(\mathbb{S}^1 | (\mathbb{S}^1)^{\iota}, \mathbb{Z}(1) \big) \right)^{\oplus 2} \tag{4.7}$$

given by

$$\nu^{w} \,:\, [\mathbb{T}^{2} \times \mathbb{C}^{2n}, H_{0}, H_{1}] \,\mapsto\, \left(\nu([\mathbb{T}^{2} \times \mathbb{C}^{2n}, H_{0}, H_{1}]_{1}), \nu([\mathbb{T}^{2} \times \mathbb{C}^{2n}, H_{0}, H_{1}]_{2})\right).$$

Notice that ν^w is well-defined on the whole group $KR^{-3}(\mathbb{T}^2, \iota)$ and not only on $Ker(\kappa)$. We are now in position to provide a complete classification of two-dimensional band insulators of class DIII in the Bloch case.

Proposition 4.5 (Two-dimensional weak-strong invariant). The homomorphism ν^w extends the homomorphism ν . Moreover the composition $\underline{\nu} := (\nu^w, \kappa)$ of the homomorphisms ν^w and κ provides the isomorphism

$$\underline{\nu} \; : \; KR^{-3}(\mathbb{T}^2,\iota) \; \longrightarrow \; H^1(\mathbb{T}^2,\mathbb{Z})/H^1_{\mathbb{Z}_2}\big(\mathbb{T}^2|(\mathbb{T}^2)^\iota,\mathbb{Z}(1)\big) \; \oplus \; H^2_{\mathbb{Z}_2}\big(\mathbb{T}^2|(\mathbb{T}^2)^\iota,\mathbb{Z}(1)\big) \; \; .$$

Proof. The proof is based on the direct calculations. For that let us construct four sewing matrices q_{w_1} , q_{w_2} , q_s , q_o on \mathbb{T}^2 with values on $\mathbb{U}(2)$ as described below. Let π_1 and

 π_2 be the two canonical projections from $\mathbb{T}^2\simeq\mathbb{S}^1\times\mathbb{S}^1$ to \mathbb{S}^1 . Then, $q_{w_j}:=q_-\circ\pi_j$, for j=1,2, where $q_-:\mathbb{S}^1\to\mathbb{U}(2)$ is the sewing matrix introduced in the proof of Theorem 3.12. The sewing matrix $q_s:=q_1\circ\pi_0$ is the composition of the equivariant map $\pi_0:\mathbb{T}^2\to\mathbb{S}^2$ in the proof of Lemma 4.2 and the sewing matrix $q_1:\mathbb{S}^2\to\mathbb{U}(2)$ such that $\kappa(\mathscr{E}_{q_1})=1$ in Proposition 4.1. Finally, q_0 is the constant map as in Theorem 2.8 (2). These sewing matrices produce the following FKMM invariants:

$$\kappa(\mathscr{E}_{\mathfrak{q}_{w_1}}) \; = \; +1 \; , \quad \ \kappa(\mathscr{E}_{\mathfrak{q}_{w_2}}) \; = \; +1 \; , \quad \ \kappa(\mathscr{E}_{\mathfrak{q}_0}) \; = \; +1 \; , \quad \ \kappa(\mathscr{E}_{\mathfrak{q}_s}) \; = \; -1 \; .$$

In addition, one has that

$$\begin{array}{l} \nu_{q_{w_{1}} \circ \varphi_{1}} \; = \; -1 \; , \quad \nu_{q_{w_{2}} \circ \varphi_{1}} \; = \; +1 \; , \quad \nu_{q_{0} \circ \varphi_{1}} \; = \; +1 \; , \quad \nu_{q_{s} \circ \varphi_{1}} \; = \; +1 \; , \\ \nu_{q_{w_{1}} \circ \varphi_{2}} \; = \; +1 \; , \quad \nu_{q_{w_{2}} \circ \varphi_{2}} \; = \; -1 \; , \quad \nu_{q_{0} \circ \varphi_{2}} \; = \; +1 \; , \quad \nu_{q_{s} \circ \varphi_{2}} \; = \; +1 \; . \end{array}$$

In the above equations the computations for the cases $q_{w_j} \circ \varphi_i = q_- \circ (\pi_j \circ \varphi_i)$ follows just by observing that $\pi_j \circ \varphi_i$ is the identity map on \mathbb{S}^1 when i = j, and the restriction to a fixed point if $i \neq j$. The cases $q_0 \circ \varphi_i$ with i = 1, 2, are also trivial in view of the fact that q_0 is the constant map. For the last two cases $q_s \circ \varphi_i$, with i = 1, 2, we notice that the equivariant map $\pi_0 : \mathbb{T}^2 \to \mathbb{S}^2$ carries the three fixed points $(0,0), (0,\pi), (\pi,0)$ to (1,0,0) and the fixed point (π,π) to (-1,0,0). We also notice that q_1 takes values in $S\mathbb{U}(2) \subset \mathbb{U}(2)$. Since $q_s \circ \varphi_i : \mathbb{S}^1 \to S\mathbb{U}(2)$ carries the fixed points of \mathbb{S}^1 to the same point of $S\mathbb{U}(2)$, Theorem 3.5 leads to the results above. Now, let us denote with $H_{w_1}, H_{w_2}, H_{q_s}$ and H_0 the band insulators of class DIII associated to q_{w_1}, q_{w_2}, q_s and q_0 , respectively. One gets that

$$\begin{split} & \underline{\nu} \big([\mathbb{T}^2 \times \mathbb{C}^2, H_{w_1}, H_0] \big) \; = \; (-1, +1, +1) \; , \\ & \underline{\nu} \big([\mathbb{T}^2 \times \mathbb{C}^2, H_{w_2}, H_0] \big) \; = \; (+1, -1, +1) \; , \\ & \underline{\nu} \big([\mathbb{T}^2 \times \mathbb{C}^2, H_s, H_0] \big) \; = \; (+1, +1, -1), \end{split}$$

so that $\underline{\nu}$ is a surjection onto $\mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2$. Since $KR^{-3}(T^2) \simeq \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2$, the homomorphism $\underline{\nu}$ must be an isomorphism. As a result, one obtains that $Ker(\kappa)$ is generated by $[\mathbb{T}^2 \times \mathbb{C}^2, H_{w_i}, H_0]$, with i = 1, 2. This allows us to see that ν^w extends ν .

The novelty of Proposition 4.5 is that, in combination with Theorem 3.5, it provides a way to compute also the weak invariants

$$\boldsymbol{\nu}^{\boldsymbol{w}}(\boldsymbol{H}) \; := \; (\boldsymbol{\nu}_{\mathfrak{q}_{\varphi_1^*\boldsymbol{H}}}, \boldsymbol{\nu}_{\mathfrak{q}_{\varphi_2^*\boldsymbol{H}}})$$

of the two-dimensional band insulator H of class DIII over (\mathbb{T}^2, ι) .

APPENDIX A. EQUIVARIANT COHOMOLOGY AND HOMOTOPY

Let (X, τ) be an involutive space such that X has the structure of a (path connected) \mathbb{Z}_2 -CW complex. A map $f: X \to \mathbb{U}(1)$ is called *invariant* if $f(x) = f(\tau(x))$ for every $x \in X$. Similarly, it is called *equivariant* if $f(x) = \overline{f(\tau(x))}$ for every $x \in X$. Let $[X, \mathbb{U}(1)]$ be the set of classes of homotopy equivalent functions, $[X, \mathbb{U}(1)]_{inv}$ the set of

classes of homotopy equivalent invariant functions and $[X, \mathbb{U}(1)]_{\mathbb{Z}_2}$ the set of classes of homotopy equivalent equivariant functions. These three sets admit a representation in terms of cohomology groups. In fact, it holds true that

$$\begin{split} [X,\mathbb{U}(1)] &\simeq H^1(X,\mathbb{Z}) \\ [X,\mathbb{U}(1)]_{\text{inv}} &\simeq H^1_{\mathbb{Z}_2}(X,\mathbb{Z}) \\ [X,\mathbb{U}(1)]_{\mathbb{Z}_2} &\simeq H^1_{\mathbb{Z}_2}(X,\mathbb{Z}(1)) \end{split}$$
 (A.1)

where $H^{\bullet}(X,\mathbb{Z})$ is the usual cohomology ring of X with coefficients in \mathbb{Z} , and $H^{\bullet}_{\mathbb{Z}_2}(X,\mathbb{Z})$ and $H^{\bullet}_{\mathbb{Z}_2}(X,\mathbb{Z}(1))$ are the equivariant cohomology groups of X with system of coefficients in \mathbb{Z} and $\mathbb{Z}(1)$, respectively. For a review of the equivariant cohomology we refer to [DG1, Section 5] and references therein. The first isomorphism in (A.1) is a classical result in algebraic topology and follows by observing that $\mathbb{U}(1) = K(\mathbb{Z}, 1)$ is a Eilenberg-Maclane space and in turn $[X, K(\mathbb{Z}, 1)]$ is a representing space for the cohomology $H^1(X, \mathbb{Z})$ in view of [Hat, Theorem 4.57]. The proof of the second isomorphism is provided in [GT, Lemma A.3], while the third isomorphism is proved in [Gom, Proposition A.2].

APPENDIX B. STRONG EQUIVALENCE OF "QUATERNIONIC" STRUCTURES

Let $\operatorname{Vec}^n_{\mathfrak{Q}}(X,\tau)$ be the set of equivalence classes of "Quaternionic" vector bundles of rank n over the involutive space (X,τ) . The subset $\operatorname{Vec}^n_{\mathfrak{Q},0}(X,\tau) \subseteq \operatorname{Vec}^n_{\mathfrak{Q}}(X,\tau)$ consists of the equivalence classes of "Quaternionic" vector bundles whose underlying (complex) bundle structure is trivial. Let $\mathscr{E} := (X \times \mathbb{C}^n, \Theta)$ and $\mathscr{E}' := (X \times \mathbb{C}^n, \Theta')$ be two "Quaternionic" vector bundles defined on the trivial product bundle $X \times \mathbb{C}^n$. The two "Quaternionic" structures Θ and Θ' are defined by two sewing matrices \mathfrak{q} and \mathfrak{q}' according to (2.8). The isomorphism condition $\mathscr{E} \simeq \mathscr{E}'$ is equivalent to the existence of a continuous *intertwining* map $\mathfrak{q}: X \to \mathbb{U}(\mathfrak{n})$ such that

$$q'(x) = \phi(\tau(x))^{t} q(x) \phi(x), \quad \forall x \in X.$$
 (intertwining) (B.1)

In this case $\mathscr E$ and $\mathscr E'$ are representatives of the same equivalence class $[\mathscr E]\in Vec^n_{\mathfrak Q,0}(X,\tau).$

Definition B.1 (Strong isomorphism). We will say that the isomorphism between \mathcal{E} and \mathcal{E}' is *strong* if the intertwining map (B.1) meets the extra condition

$$\phi(x) = \phi(\tau(x)), \quad \forall x \in X. \quad (\tau\text{-invariance})$$
 (B.2)

In this case we will write $\mathscr{E} \stackrel{\mathtt{s}}{\simeq} \mathscr{E}'$.

The notion of strong isomorphism introduced with Definition B.1 provides a finer classification of $\text{Vec}_{\mathfrak{D},0}^n(X,\tau)$. The following example should clarify the difference between the two notions of isomorphism.

Example B.2. It is known that in the case of the involutive space (\mathbb{S}^1, ι) one has that $\operatorname{Vec}_{\mathfrak{Q},0}^2(\mathbb{S}^1, \iota) = \operatorname{Vec}_{\mathfrak{Q}}^2(\mathbb{S}^1, \iota) = 0$ [DG2, Proposition 2.6] meaning that every rank two

"Quaternionic" vector bundle over (\mathbb{S}^1, ι) is isomorphic to the trivial "Quaternionic" vector bundle $\mathscr{E}_0 := (\mathbb{S}^1 \times \mathbb{C}^2, \Theta_0)$ with *trivial* structure Θ_0 induced by the (constant) sewing matrix

$$q_0(k) \; \equiv \; q_0 \; := \; \left(\begin{array}{cc} 0 & -1 \\ 1 & 0 \end{array} \right) \; , \qquad \forall \; k \in \mathbb{S}^1 \; .$$

Consider the "Quaternionic" vector bundle $\mathscr{E}_0' := (\mathbb{S}^1 \times \mathbb{C}^2, \Theta_0')$ with "Quaternionic" structure Θ_0' induced by the (again constant) sewing matrix

$$q'_0(k) \equiv -q_0$$
, $\forall k \in \mathbb{S}^1$.

The (constant) intertwining map

$$\varphi_0(k) \; \equiv \; \varphi_0 \; := \; \left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array} \right) \; , \qquad \forall \; k \in \mathbb{S}^1 \; .$$

provides the isomorphism between \mathscr{E}_0 and \mathscr{E}_0' . Moreover, since the map φ_0 is constant, hence τ -invariant, one has that the "Quaternionic" vector bundles are *strongly* equivalent, $i.e. \mathscr{E}_0 \stackrel{\text{s}}{\simeq} \mathscr{E}_0'$. On the other hand, one can consider the "Quaternionic" vector bundle $\mathscr{E}_1 := (\mathbb{S}^1 \times \mathbb{C}^2, \Theta_1)$ with "Quaternionic" structure Θ_1 induced by the (non constant) sewing matrix

$$q_1(k) := \begin{pmatrix} \sin(k) & -\cos(k) \\ \cos(k) & \sin(k) \end{pmatrix}, \quad \forall \ k \in \mathbb{S}^1.$$

In this case the isomorphism $\mathcal{E}_1 \simeq \mathcal{E}_0$ is induced by the intertwining map

$$\varphi_1(k) \; := \; e^{i \frac{k}{2}} \left(\begin{array}{cc} sin\left(\frac{k}{2}\right) & -cos\left(\frac{k}{2}\right) \\ cos\left(\frac{k}{2}\right) & sin\left(\frac{k}{2}\right) \end{array} \right) \; , \qquad \forall \; k \in \mathbb{S}^1 \; .$$

Since $\phi_1(-k) \neq \phi_1(k)$ it follows that the "Quaternionic" structures Θ_0 and Θ_1 are only equivalent but *not* strongly equivalent, *i. e.* $\mathcal{E}_1 \not\simeq \mathcal{E}_0$.

In the next result we will use the definitions of unitary equivalence introduced in Definition 1.1. Let us observe that if V is a unitary equivalence between two band insulators of class DIII then the equation

$$\chi V(x) = V(x) \chi$$

holds true, independently of the nature of V.

Proposition B.3. Unitarily equivalent band insulators of class DIII define associated "Quaternionic" vector bundles which are isomorphic. When the unitary equivalence is strong then the associated "Quaternionic" vector bundles are strongly isomorphic.

Proof. Let H and H' be two unitarily equivalent band insulators of class DIII over (X, τ) . The functional calculus provides $Q_{H'}(x) = V(x)Q_H(x)V(x)^*$ where the operators $Q_{H'}(x)$

and Q_H are defined according to (2.5). According to Lemma 2.2, in the standard representation described in Preposition 2.1 the unitary V(x) takes the form

$$V(x) = \begin{pmatrix} \phi_V(x) & 0 \\ 0 & \overline{\phi_V(\tau(x))} \end{pmatrix}, \qquad (B.3)$$

for a suitable map $\phi_V: X \to \mathbb{U}(n)$. Then, the unitary equivalence between Q_H and $Q_{H'}$ translates into

$$q_{H'}(x) \; = \; \overline{\varphi_V \big(\tau(x)\big)} \; q_H(x) \; \varphi_V(x)^* \; = \; \big[\varphi_V \big(\tau(x)\big)^*\big]^\mathsf{t} \; q_H(x) \; \varphi_V(x)^*$$

where the function q_H and $q_{H'}$ are defined according to the off-diagonal decomposition (2.6). The latter equation shows that the "Quaternionic" vector bundles associated to H and H' are isomorphic. In the case of a strong unitary equivalence between H and H', the condition $V(\tau(x)) = V(x)$ translates into $\varphi_V(\tau(x)) = \varphi_V(x)$ and in turn in the strong isomorphism between the associated "Quaternionic" vector bundles.

REFERENCES

- [Ati] Atiyah, M. F.: K-theory and reality. Quart. J. Math. Oxford Ser. (2) 17, 367-386 (1966)
- [AS] Atiyah, M. F.; Singer, I. M.: *Index theory for skew-adjoint Fredholm operators*. Publ. Math. IHES **37**, 5-26 (1969)
- [AZ] Altland, A.; Zirnbauer, M.:: Non-standard symmetry classes in mesoscopic normal-superconducting hybrid structures. Phys. Rev.B **55**, 1142-1161 (1997)
- [BA] Budich, J. C.; Ardonne, E.: Topological invariant for generic one-dimensional time-reversal-symmetric superconductors in class DIII. Phys. Rev. B **88**, 134523 (2013)
- [CTSR] Chiu, C.-K.; Teo, J. C. Y.; Schnyder, A. P; Ryu, S.: Classification of topological quantum matter with symmetries. Rev. Mod. Phys. 88, 035005 (2016)
- [DG1] De Nittis, G.; Gomi, K.: Classification of "Real" Bloch-bundles: Topological Insulators of type AI. J. Geometry Phys. **86**, 303-338 (2014)
- [DG2] De Nittis, G.; Gomi, K.: Classification of "Quaternionic" Bloch-bundles: Topological Insulators of type AII. Commun. Math. Phys. **339**, 1-55 (2015)
- [DG3] De Nittis, G.; Gomi, K.: *The cohomological nature of the Fu-Kane–Mele invariant*. J. Geometry Phys. **124**, 124-164 (2018)
- [DG4] De Nittis, G.; Gomi, K.: *The FKMM-invariant in low dimension*. Lett. Math. Phys. **108**, 1225-1277 (2018)
- [DG5] De Nittis, G.; Gomi, K.: On the K-theoretic classification of dynamically stable systems. Rev. Math. Phys. **31**, 1950003 (2019)
- [DG6] De Nittis, G.; Gomi, K.: Chiral vector bundles. Math. Z. 290, 775-830 (2018)
- [DS] De Nittis, G.; Schulz-Baldes, H.: *Spectral Flows of Dilations of Fredholm Operators*. Canad. Math. Bull. **58**, 51-68 (2014)
- [DK] Donovan, D.; Karoubi, M.: *Graded Brauer groups and K-theory with local coefficients*. Inst. Hautes Études Sci. Publ. Math. **38**, 5-25 (1970)
- [Dup] Dupont, J. L.: Symplectic Bundles and KR-Theory. Math. Scand. 24, 27-30 (1969)
- [FHT] Freed, D. S.; Hopkins, M. J.; Teleman, C.: Loop groups and twisted K-theory. I. J. Topol. 4, 737-798 (2011)
- [FM] Freed, D. S.; Moore, G. W.: Twisted Equivariant Matter. Ann. Henri Poincaré 14, 1927-2023 (2013)

- [Gom] Gomi, K.: A Variant of K-Theory and Topological T-Duality for Real Circle Bundles. Commun. Math. Phys. **334**, 923-975 (2015)
- [Gom2] Gomi, K.: Freed-Moore K-theory. Accepted in Commun. Anal. Geom. E-print arXiv:1705.09134, (2017)
- [GT] Gomi, K.; Thiang, G. C.: 'Real'-gerbes and Dirac cones of topological insulators. E-print arXiv:2103.05350, (2021)
- [Hat] Hatcher, A.: Algebraic Topology. Cambridge University Press, Cambridge, 2002
- [Kah] Kahn, B.: Construction de classes de Chern équivariantes pour un fibré vectoriel Réel. Comm. Algebra. 15, 695-711 (1987)
- [Kit] Kitaev, A.: *Periodic table for topological insulators and superconductors.* AIP Conf. Proc. **1134**, 22-30 (2009)
- [KKL] Ko, E.; Ko, E.; Lee, J. E.: *Skew complex symmetric operator and Weyl type theorems.* Bull. Kor. Math. Soc. **52**, 1269-1283 (2015)
- [LZ] Li, C. G.; Zhu, S.: Skew symmetric normal operators. Proc. Amer. Math. Soc. 141, 2755-2762 (2013)
- [Mur] Murphy, G. J.: C*-Algebras and Operator Theory. Academic Press, Inc., 1990
- [PS] Prodan, E.; Schulz-Baldes, H.: *Bulk and Boundary Invariants for Complex Topological Insulators:* From K-Theory to Physics. Mathematical Physics Studies, Springer, 2016
- [Ros] Rosenberg, J.: Continuous-trace algebras from the bundle theoretic point of view. J. Austral. Math. Soc. Ser. A 47, 368-381 (1989)
- [RSFL] Ryu, S.; Schnyder, A. P.; Furusaki, A.; Ludwig, A. W. W.: *Topological insulators and superconductors: tenfold way and dimensional hierarchy.* New J. Phys. **12**, 065010 (2010)
- [SB] Schulz-Baldes, H.: Z₂-Indices and Factorization Properties of Odd Symmetric Fredholm Operators. Documenta Math. **20**, 1481-1500 (2015)
- [SRFL] Schnyder, A. P.; Ryu, S.; Furusaki, A.; Ludwig, A. W. W.: Classification of topological insulators and superconductors in three spatial dimensions. Phys. Rev. **B 78**, 195125 (2008)
- [TK] Teo, J. C. Y.; Kane, C. L.: *Topological defects and gapless modes in insulators and superconductors*. Phys. Rev. B **82**, 115120 (2010)
- (G. De Nittis) FACULTAD DE MATEMÁTICAS & INSTITUTO DE FÍSICA, PONTIFICIA UNIVERSIDAD CATÓLICA DE CHILE, SANTIAGO, CHILE.

Email address: gidenittis@mat.uc.cl

(K. Gomi) Department of Mathematics, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro-ku, Tokyo, 152-8551, Japan.

Email address: kgomi@math.titech.ac.jp