

Precoloring Extension. IV. General Bounds and List Colorings

Mihály Hujter; Zsolt Tuza

Abstract. A guaranteed upper bound is proved for the time complexity of the list-coloring problem on graphs.

For a positive integer n , the set $\{1, \dots, n\}$ is denoted by $[1, n]$. Throughout this note, n will denote the *order* of a graph, that is the number of vertices.

A *proper k -coloring* (or simply, a k -coloring) of a graph $G = (V, E)$ is a function $f : V \rightarrow [1, k]$ such that $f(v) \neq f(w)$ whenever $vw \in E$. The *vertex coloring problem*, whose input is a graph G and a positive integer k , consists of deciding whether G is properly k -colorable or not. In the *list-coloring problem*, every vertex v comes equipped with a list of permitted colors $L(v) \subseteq [1, \kappa]$ for some given positive integer κ , and we require the proper coloring to respect these lists, i.e., $f(v) \in L(v)$.

We say that a constant real number $t > 1$ *fits* G if for any m -element subset M of V , the *induced subgraph* $G[M]$ has at most t^m *inclusion-maximal independent sets*. It is well-known that 1.44225 fits any graph because $1.44225^3 > 3$; see [3] for a proof of the fact that an n -vertex graph contains at most $3^{n/3}$ inclusion-maximal independent sets. Similarly, 1.41422 fits any triangle-free graph because $1.41422^2 > 2$; see [8] for a proof of the fact that a triangle-free graph contains at most $2^{n/2}$ inclusion-maximal independent sets. Furthermore it was proved in [9] that, for any fixed natural number k , graphs not containing an induced matching with $k + 1$ edges have at most $O(n^{2k})$ inclusion-maximal independent sets. Hence for every k and every real $\epsilon > 0$ there exists an $n_0 = n_0(k, \epsilon)$ such that $t = 1 + \epsilon$ fits every such graph of order $n > n_0$.

A special case of the list-coloring problem is the *precoloring extension* with κ colors. Here each list is either $[1, \kappa]$ or a one-element subset of it. (See [5, 6, 7] for more details on precoloring extension.) Another special case is the *k -choosability* problem where each color list has size k for a fixed k . (See [4, 10] for more details on choosability.)

The main result in the present note, Theorem 1, claims that there exists a polynomial $p(n)$ such that if t fits a graph G with n vertices, then it can be

decided in $p(n) \cdot (1 + t)^n + O(\kappa n)$ time whether G is properly list-colorable or not. (Here the polynomial p is independent of t , and the $O(\kappa n)$ term is irrelevant except for reading an input with rather long vertex lists.) In case of a positive answer, we will construct a proper list-coloring needing no extra time. We will extend the methods and results of Lawler [1].

Given a graph $G = (V, E)$ on n vertices, we fix a permutation of the nonempty subsets of V for which $W, U \subseteq V$ and $|W| > |U|$ imply that W is before U in the fixed permutation. For any nonempty $W \subseteq V$ let $\langle W \rangle$ denote the position of W in the fixed permutation. As a special case, $\langle V \rangle = 1$, and $|W| = n - 1$ implies that $\langle W \rangle \in \{2, 3, \dots, n + 1\}$, furthermore $|W| = 1$ implies that $\langle W \rangle \in \{2^n - n, 2^n - n + 1, \dots, 2^n - 1\}$. We will consider a zero-one sequence (a_m) , $m = 1, 2, \dots, 2^n - 1$. We say that the sequence has the *partial increasing property* (PIP, for short) if $W \supset U$ implies $a_{\langle W \rangle} \leq a_{\langle U \rangle}$. In words, if W is before U in the permutation and if $a_{\langle W \rangle} = 1$, then $a_{\langle U \rangle} = 1$ holds by pip.

Initially, the sequence (a_m) , $m = 1, 2, \dots, 2^n - 1$ will be set as the *characteristic sequence of the independent sets* in G . In other words, $a_{\langle W \rangle} = 1$ holds if and only if the induced subgraph $G[W]$ is edgeless. Later some zero values in the sequence will be changed to one; however, the pip will be managed.

Clearly, list-colorability can be reformulated as follows: The vertex set V can be partitioned into independent set *color classes* V_1, \dots, V_κ such that for each $j \in [1, \kappa]$ and for each $v \in V_j$ the relation $j \in L(v)$ holds. Here we allow the empty set to occur among V_1, \dots, V_κ any number of times.

For each $j \in [1, \kappa]$ and for each $v \in V$ we define $L^j(v)$ as $L(v) \cap [1, j]$. Clearly $L^1(v) \subseteq \dots \subseteq L^\kappa(v) = L(v)$.

For each $j \in [1, \kappa]$ we will consider the j^{th} version of the list-coloring problem, denoted by \mathcal{LC}^j for all induced subgraphs $G[W]$ by considering the lists $L^j(v)$. Note that \mathcal{LC}^1 is trivial since $G[W]$ is list-colorable (with respect to the lists $L^j(v)$, $w \in W$) if and only if, on the one hand, W is an independent set in G , and on the other hand, $1 \in L(w)$ for all $w \in W$. Starting from the list \mathcal{I} of all inclusion-maximal independent sets of $G = (V, E)$, the sequence (a_m) , $m = 1, 2, \dots, 2^n - 1$ can be computed as follows:

1. Let all a_m get the value 0.
2. For each $I \in \mathcal{I}$, consider

$$I^{(1)} = \{v \in I : 1 \in L(v)\}$$

and set $a_{\langle I^{(1)} \rangle} = 1$.

3. Manage the above mentioned PIP of the sequence (a_m) .

For this purpose we go forward in the sequence and for any $a_{\langle W \rangle} = 1$ and for any $w \in W$ we set $a_{\langle W-w \rangle} = 1$. (Here $W - w$ simply denotes $W \setminus \{w\}$.)

Lemma 1. After performing the above three steps, the final values in the sequence $(a_{\langle W \rangle})$ give the true answer for the list-coloring problem for each $G[W]$ with respect to the lists $L^1(v)$. This means that $a_{\langle W \rangle} = 1$ holds if and only if $G[W]$ is list-colorable with respect to the lists $L^1(v)$, $v \in W$.

Proof. For any inclusion-maximal vertex set I , in step 2 the sequence element $a_{\langle I \rangle}$ gets value 1, and in step 3, for any other independent set W , the final value of $a_{\langle W \rangle}$ will also be 1. ■

Now for any fixed $j = 2, 3, \dots, \kappa$, we will recursively solve the list-colorability problem \mathcal{LC}^j for all $G[W]$ by using the lists consisting of the sets

$$I^{(j)} = \{v \in I : j \in L(v)\}$$

obtained from all inclusion-maximal independent sets I in G . For a fixed $j \in \{2, 3, \dots, \kappa\}$, we assume that the problem \mathcal{LC}^{j-1} is completely solved, and the result is properly recorded in the sequence (a_m) , $m = 1, 2, \dots, 2^n - 1$, that is for any $G[W]$ this induced subgraph is list-colorable with respect to the color lists $L^{j-1}(v)$, $v \in W$, if and only if $a_{\langle W \rangle} = 1$.

Lemma 2. The graph $G = (V, E)$ is list-colorable with respect to the color lists $L^j(v)$, $v \in V$, if and only if there is at least one independent set $I^{(j)}$ for which either $I^{(j)} = V$ or $a_{\langle V \setminus I^{(j)} \rangle} = 1$ holds as a result of the \mathcal{LC}^{j-1} problem.

Proof. If $I^{(j)} = V$, then G is clearly list-colorable since $f(v) = j$ suffices for all $v \in V$. If there is an $I^{(j)} \neq V$ for which $a_{\langle V \setminus I^{(j)} \rangle} = 1$ holds, then any list-coloring f of $G[V \setminus I^{(j)}]$ with respect to the lists $L^{j-1}(v)$, $v \in V \setminus I^{(j)}$, is clearly extendible to obtain a list-coloring of G by defining $f(v) = j$ for all $v \in I^{(j)}$. On the other hand, in any list-coloring f of G with respect to the lists $L^j(v)$, $v \in V$, the vertices v with $f(v) = j$, must form an independent set J , and this J must be contained in at least one $I^{(j)}$, and now the restriction of f onto $V \setminus I^{(j)}$ gives us a list-coloring of $G[V \setminus I^{(j)}]$ with respect to the lists $L^{j-1}(v)$, $v \in V \setminus I^{(j)}$. ■

Now Lemma 2 shows how we can solve the \mathcal{LC}^j problem for G . Similarly we can solve the \mathcal{LC}^j problem for any induced subgraph $G[W]$. Here we can use the

same sequence (a_m) , $m = 1, 2, \dots, 2^n - 1$. For each W we scan the sets $I^{(j)}$ for all inclusion-maximal independent sets I in $G[W]$, and update $a_{\langle W \rangle}$ from 0 to 1 if $a_{\langle W \setminus I^{(j)} \rangle} = 1$ holds in the sequence after round $j - 1$. In this procedure PIP ensures that the elements $a_{\langle U \rangle}$ may be updated only when they are not needed anymore for updates of any other elements.

If t fits G , then the total number of steps is not larger than $p(n)$ times the following number, where $p(n)$ is a polynomial independent of G and independent of t , too:

$$\sum_{W \neq \emptyset} \binom{n}{|W|} t^{|W|}$$

By the binomial theorem, the above sum is $(1 + t)^n - 1$. In summary, we obtain the following theorem.

Theorem 1. There exists a polynomial p for which if t fits G , an n -vertex graph, then all the problems \mathcal{LC}^j , $j = 1, 2, \dots, \kappa$ can be solved in

$$p(n) \cdot (1 + t)^n + O(\kappa n)$$

time.

Proof. While reading the input, we put aside all vertices whose lists have length at least n . Once the rest of the graph is properly list-colored, the vertices with long lists can be colored in an obvious greedy way. For this reason we can assume from now on that $\kappa < n$ holds.

It is proved in [2] that if a graph on n vertices has N inclusion-maximal independent sets, then those N sets can be listed in $O(n^3 N)$ time. We apply this method κ times for all induced subgraphs $G[W]$. The input graph G is list colorable if and only if $a_{\langle V \rangle} = 1$ holds after round κ . Thus there exists a suitable polynomial $p(n) = O(n^4)$. ■

If, in case of list-colorability, we want to output a proper list-coloring $f : V \rightarrow [1, \kappa]$, we simply have to memorize an independent set $I^{(\kappa)}$ for which $G[V \setminus I^{(\kappa)}]$ is list-colorable with respect to the lists $L^{\kappa-1}(v)$, $v \in V \setminus I^{(\kappa)}$, and for any $w \in I^{(\kappa)}$ we can set $f(w) = \kappa$. Then we decrease the value of κ by one, and we repeat the procedure for the decreased graph instead of the old one, and for the new \mathcal{LC}^κ problem instead of the old \mathcal{LC}^κ problem. And so on.

References

- [1] Lawler, E. L.: *A note on the complexity of the chromatic number problem*. Inf. Process. Lett. **5** (1976) 66–67.
- [2] Tsukiyama, S.; Ide, M.; Ariyoshi, H.; Shirakawa, I.: *A new algorithm for generating all the maximal independent sets*. SIAM J. Comput. **6** (1977) 505–517.
- [3] Moon, J.; Moser, L.: *On cliques in graphs*, Israel J. Math. **3** (1965) 23–28.
- [4] Erdős, P.; Rubin, A.L.; Taylor, H.: *Choosability in graphs*. Combinatorics, graph theory and computing, Proc. West Coast Conf., Arcata/Calif. 1979, (1980) 125–157.
- [5] Biró, M.; Hujter, M.; Tuza, Zs.: *Precoloring extension. I. Interval graphs*. Discrete Math. **100** (1992) 267–279.
- [6] Hujter, M.; Tuza, Zs.: *Precoloring extension. II. Graph classes related to bipartite graphs*. Acta Math. Univ. Comenianae **62** (1993) 1–11.
- [7] Hujter, M., and Tuza, Zs.: *Precoloring extension. III. Classes of perfect graphs*. Combin. Probab. Comput. **5** (1996) 35–56.
- [8] Hujter, M.; Tuza, Zs.: *The number of maximal independent sets in triangle-free graphs*. SIAM J. on Discrete Math. **6** (1993) 284–288.
- [9] Farber, M.; Hujter, M.; Tuza, Zs.: *An upper bound on the number of cliques in a graph*. Networks **23** (1993) 207–210.
- [10] Tuza, Zs.: *Graph colorings with local constraints – A survey*. Discuss. Math. Graph Theory **17** (1993) 161–228.