

Convergence of limit shapes for 2D near-critical first-passage percolation

Chang-Long Yao

Academy of Mathematics and Systems Science, CAS, Beijing, China. E-mail: deducemath@126.com

Abstract. We consider Bernoulli first-passage percolation on the triangular lattice in which sites have 0 and 1 passage times with probability p and $1 - p$, respectively. For each $p \in (0, p_c)$, let $\mathcal{B}(p)$ be the limit shape in the classical “shape theorem”, and let $L(p)$ be the correlation length. We show that as $p \uparrow p_c$, the rescaled limit shape $L(p)^{-1}\mathcal{B}(p)$ converges to a Euclidean disk. This improves a result of Chayes et al. [*J. Stat. Phys.* **45** (1986) 933–951]. The proof relies on the scaling limit of near-critical percolation established by Garban et al. [*J. Eur. Math. Soc.* **20** (2018) 1195–1268], and uses the construction of the collection of continuum clusters in the scaling limit introduced by Camia et al. [*Springer Proceedings in Mathematics & Statistics*, **299** (2019) 44–89].

Résumé. ???.

MSC2020 subject classifications: Primary 60K35; secondary 82B43

Keywords: first-passage percolation, near-critical percolation, scaling limit, correlation length, shape theorem

1. Introduction

1.1. The model and main result

First-passage percolation (FPP) is a stochastic growth model which was first introduced by Hammersley and Welsh [21] in 1965. For general background on FPP, we refer the reader to the recent survey [2]. In this paper, we will focus on FPP defined on the triangular lattice \mathbb{T} , since the proof of our main result relies on the scaling limit of near-critical site percolation on \mathbb{T} established by Garban, Pete and Schramm [16], while such result has not been proved for other planar lattices.

The model is defined as follows. Let \mathbb{T} be the triangular lattice embedded in \mathbb{C} , with site (vertex) set

$$V(\mathbb{T}) := \{x + ye^{\pi i/3} \in \mathbb{C} : x, y \in \mathbb{Z}\},$$

and bond (edge) set $E(\mathbb{T})$ obtained by connecting all pairs $u, v \in V(\mathbb{T})$ for which $|u - v| = 1$, where $|\cdot|$ denotes the Euclidean norm. We say that u and v are neighbors if $(u, v) \in E(\mathbb{T})$. A **path** is a sequence (v_0, \dots, v_n) of distinct sites such that v_{j-1} and v_j are neighbors for all $j = 1, \dots, n$. Let $\{t(v) : v \in V(\mathbb{T})\}$ be an i.i.d. family of nonnegative random variables with common distribution function F . For a path γ , we define its passage time by $T(\gamma) := \sum_{v \in \gamma} t(v)$. For $A, B \subset V(\mathbb{T})$, the **first-passage time** from A to B is defined by

$$T(A, B) := \inf\{T(\gamma) : \gamma \text{ is a path from a site in } A \text{ to a site in } B\}.$$

A **geodesic** from A to B is a path γ from A to B such that $T(\gamma) = T(A, B)$. If $x, y \in \mathbb{C}$, we define $T(x, y) := T(\{x'\}, \{y'\})$, where x' (resp. y') is the site in $V(\mathbb{T})$ closest to x (resp. y). Any possible ambiguity can be avoided by ordering $V(\mathbb{T})$ and taking the site in $V(\mathbb{T})$ smallest for this order.

In this paper, we concentrate on **Bernoulli FPP** on \mathbb{T} . More precisely, for each $p \in [0, 1]$, we consider the i.i.d. family $\{t(v) : v \in V(\mathbb{T})\}$ of Bernoulli random variables with parameter p , that is, $t(v) = 0$ with probability p and $t(v) = 1$ with probability $1 - p$. This gives rise to a product probability measure on the set of configurations $\{0, 1\}^{V(\mathbb{T})}$, which is denoted by \mathbf{P}_p , the corresponding expectation being \mathbf{E}_p . We refer to a site v with $t(v) = 0$ simply as **open**; otherwise, **closed**. So




FIG 1. Bernoulli FPP on \mathbb{T} . Each hexagon of the hexagonal lattice \mathbb{H} represents a site of \mathbb{T} , and is colored blue ($t(v) = 0$) or yellow ($t(v) = 1$). Here, the first-passage time $T(0, 9) = 2$.

one can view Bernoulli FPP as Bernoulli site percolation on \mathbb{T} (see, e.g., [7, 19] for background on percolation; note that for Bernoulli FPP, a site is open when it takes the value 0, while in the percolation literature, a site is open usually means that the site takes the value 1). We usually represent it as a random coloring of the faces of the dual regular hexagonal lattice \mathbb{H} , each face centered at $v \in V(\mathbb{T})$ being blue ($t(v) = 0$) or yellow ($t(v) = 1$); see Figure 1. Sometimes we view the site v as the hexagon H_v in \mathbb{H} centered at v .

Suppose $p \in [0, 1]$. It follows from the subadditive ergodic theorem that, for any $z \in \mathbb{C}$, there is a constant $\mu(p, z)$, such that

$$\lim_{n \rightarrow \infty} \frac{T(0, nz)}{n} = \mu(p, z) \quad \mathbf{P}_p\text{-a.s. and in } L^1. \quad (1)$$

We call $\mu(p) := \mu(p, 1)$ the **time constant**. As usual, we write $a_{0,n} := T(0, n)$.

It is well known (see, e.g., Kesten's Theorem 6.1 in [24] for general FPP) that

$$\mu(p) = 0 \quad \text{if and only if} \quad p \geq p_c, \quad (2)$$

where $p_c = p_c(\mathbb{T}) = 1/2$ is the critical point for Bernoulli site percolation on \mathbb{T} . In this paper, we will focus on the subcritical case, where $p < p_c$.

The fundamental object of study is

$$B(t) := \{z \in \mathbb{C} : T(0, z) \leq t\},$$

the set of points reached from the origin 0 within a time $t \geq 0$. Using (1) and (2), it is easy to deduce that $\mu(p, z)$ is a norm on \mathbb{C} for each fixed $p < p_c$. The unit ball in μ -norm is called the **limit shape** and will be denoted by

$$\mathcal{B}(p) := \{z \in \mathbb{C} : \mu(p, z) \leq 1\}.$$

It is the limit of $B(t)$ in the following sense: By the famous Cox-Durrett shape theorem (see, e.g., Theorem 2.17 in [2]), for each $p < p_c$ and each $\epsilon > 0$,

$$\mathbf{P}_p \left[(1 - \epsilon)\mathcal{B}(p) \subset \frac{B(t)}{t} \subset (1 + \epsilon)\mathcal{B}(p) \text{ for all large } t \right] = 1. \quad (3)$$

Moreover, $\mathcal{B}(p)$ is a convex, compact set with non-empty interior, and has the symmetries of \mathbb{T} that fix the origin. Apart from this, little is known about the geometry of $\mathcal{B}(p)$.

We want to study the asymptotics of $\mu(p)$ and $\mathcal{B}(p)$, as $p \uparrow p_c$. For this purpose, we will use a very useful concept from near-critical percolation: **correlation length**. Roughly speaking, the system “looks like” critical percolation on scales smaller than correlation length, while “notable” super/sub-critical behavior emerges above this length. There are several natural definitions of correlation length, and the corresponding lengths turn out to be of the same order of magnitude. See

Section 2.1 for two different definitions of it, denoted by $L(p)$ and $L_\epsilon(p)$, respectively. The former is defined in terms of the alternating 4-arm events, while the latter in terms of the box-crossing events.

Chayes, Chayes and Durrett [8] proved that,

$$\mu(p) \asymp L(p)^{-1} \quad \text{as } p \uparrow p_c. \quad (4)$$

This result together with (15) implies that $\mu(p) = (p_c - p)^{4/3+o(1)}$ as $p \uparrow p_c$. We note that (4) was proved in [8] for bond version of subcritical Bernoulli FPP on \mathbb{Z}^2 ; the same proof applies to our site version on \mathbb{T} .

Let $\mathbb{U} := \{z \in \mathbb{C} : |z| = 1\}$ denote the unit circle centered at 0. For $r > 0$ and $z \in \mathbb{C}$, let $\mathbb{D}_r(z) := \{x \in \mathbb{C} : |x - z| < r\}$ denote the open Euclidean disk of radius r centered at z , and let $\overline{\mathbb{D}}_r(z)$ denote its closure. Write $\mathbb{D}_r = \mathbb{D}_r(0)$ and $\mathbb{D} = \mathbb{D}_1$. Our main result below improves (4), and states that $\mathcal{B}(p)$ is asymptotically circular as $p \uparrow p_c$.

Theorem 1.1. *There exists a constant $\nu > 0$, such that*

$$\lim_{p \uparrow p_c} L(p)\mu(p, u) = \nu \quad \text{uniformly in } u \in \mathbb{U}.$$

In particular, when $p \uparrow p_c$, the normalized limit shape $L(p)^{-1}\mathcal{B}(p)$ tends to the Euclidean disk $\overline{\mathbb{D}}_{1/\nu}$ in the Hausdorff metric (22).

From the above theorem, it is easy to extract the following corollary (see Section 2.5 for its proof):

Corollary 1.2. *Suppose $\epsilon \in (0, 1/2)$. There exists a constant $\nu_\epsilon > 0$ depending on ϵ , such that*

$$\lim_{p \uparrow p_c} L_\epsilon(p)\mu(p, u) = \nu_\epsilon \quad \text{uniformly in } u \in \mathbb{U}.$$

In particular, when $p \uparrow p_c$, the normalized limit shape $L_\epsilon(p)^{-1}\mathcal{B}(p)$ tends to the Euclidean disk $\overline{\mathbb{D}}_{1/\nu_\epsilon}$ in the Hausdorff metric (22).

Remark 1.3. We cannot give the explicit values of the limits in Theorem 1.1 and Corollary 1.2. In our proof, the subadditive ergodic theorem plays a crucial role in showing the existence of the limit but gives no insight for the exact value of the limit. See Section 1.2 for a sketch of the proof.

Remark 1.4. It is believed that for each fixed $p < p_c$, the limit shape $\mathcal{B}(p)$ is not a Euclidean disk, since the anisotropy of \mathbb{T} may persist in the limit. Although we cannot prove such a statement for all $p < p_c$, we provide a short argument to show that this is indeed the case so long as p is sufficiently small. For general FPP, a theorem of Cox and Kesten (see [11] or Theorem 2.7 in [2]) states that, the time constant is continuous under weak convergence of the site-weight distributions of $t(v)$. This implies that $\mu(p)$ is continuous in p , since for each $p_0 \in [0, 1]$, the Bernoulli distribution $\text{Ber}(p)$ converges weakly to $\text{Ber}(p_0)$ as p tends to p_0 . Furthermore, by Remark 6.18 of [24], for each $u \in \mathbb{U}$, the function $\mu(p, u)$ is continuous in p , and this continuity is even uniform in u . Therefore, for any $\epsilon > 0$, we can choose $p \in (0, p_c)$ sufficiently small to make the Hausdorff distance between $\mathcal{B}(p)$ and $\mathcal{B}(0)$ smaller than ϵ . So for p small enough, $\mathcal{B}(p)$ is not a Euclidean disk since $\mathcal{B}(0)$ is a regular hexagon. We want to mention that in high dimensions, Kesten proved that the limit shape of the Eden model is not a Euclidean ball, and it is conjectured that this is true for all dimensions $d \geq 2$; see, e.g., Theorem 6.2 and Question 6.1.1 in [2].

1.2. Strategy of proof

Let us outline the proof of Theorem 1.1. Our method involves three steps:

Step 1. *Construction of the scaling limit of cluster ensemble.* Based on the scaling limit results in [15, 34] for critical percolation under the quad-crossing topology, Camia, Conijn and Kiss [9] constructed the scaling limit of (open) clusters for critical percolation, which is a collection of compact sets called the ‘‘continuum clusters’’. We extend this result to the near-critical case by using the approach from [9] and the scaling limit result for near-critical percolation from [16].

Step 2. *Study of FPP on the continuum cluster ensemble.* We need to define FPP for the scaling limit constructed in Step 1. For this purpose, we call a finite sequence Γ of distinct continuum clusters a (continuum) chain if any two consecutive clusters in Γ touch each other, and let $|\Gamma|$ denote the number of clusters in Γ . The first-passage time between two continuum clusters $\mathcal{C}, \mathcal{C}'$ is defined by

$$T(\mathcal{C}, \mathcal{C}') := \inf\{|\Gamma| - 1 : \Gamma \text{ is a chain from } \mathcal{C} \text{ to } \mathcal{C}'\}.$$

This enables us to define the ‘‘point-to-point’’ passage time $T_{m,n}$ (see Figure 4) as the first-passage time between two suitably chosen continuum clusters contained respectively in the disks $\mathbb{D}(m)$ and $\mathbb{D}(n)$. Similarly, for Bernoulli FPP on

the rescaled lattice $L(p)^{-1}\mathbb{T}$ we denote by $T_{m,n}^p$ the first-passage time between two suitably chosen discrete clusters contained respectively in the disks $\mathbb{D}(m)$ and $\mathbb{D}(n)$. Next, we show that under a coupling such that the percolation configuration on $L(p)^{-1}\mathbb{T}$ converges almost surely to the quad-crossing scaling limit as $p \uparrow p_c$, the first-passage time $T_{0,n}^p$ converges to $T_{0,n}$ in probability. Furthermore, we use the subadditive ergodic theorem to obtain a law of large numbers for $T_{0,n}$, that is, $T_{0,n}/n$ tends to a constant $\nu > 0$ almost surely as $n \rightarrow \infty$.

Step 3. Convergence of $L(p)\mu(p, u)$ to ν . In Step 2 we have showed that for any fixed n , the passage time $T_{0,n}$ is well-approximated by $T_{0,n}^p$ for all $p < p_c$ sufficiently close to p_c . This is a uniformly “local approximation” since n is fixed; we need a uniformly “global approximation”: For all $p < p_c$ sufficiently close to p_c , the passage time $T_{0,n}$ is well-approximated by $T_{0,n}^p$ for all large n . Indeed, we need to show that $L(p)\mu(p)$ tends to ν as $p \uparrow p_c$. We shall implement a standard renormalization argument to show that ν is an upper bound for the upper limit of $L(p)\mu(p)$ (see Figure 7): The results obtained in Step 2 allow us to get that, with high probability, the point-to-point passage times for paths constrained in suitably chosen blocks are well-approximated by the corresponding times for paths not constrained, leading to a K -dependent bond percolation. In order to show that ν is also a lower bound for the lower limit of $L(p)\mu(p)$, we use a “block approach”, which was introduced by Grimmett and Kesten in [20] to derive exponential large deviation bounds for passage times. Finally, due to the fact that the near-critical scaling limit is invariant under rotations, it is easy to show that $L(p)\mu(p, u) \rightarrow \nu$ uniformly in $u \in \mathbb{U}$ as $p \uparrow p_c$, which completes the proof.

1.3. Relations to previous works

In this subsection we wish to review some related works in the literature. For FPP, most works have focused on the subcritical regime, where $F(0) < p_c$. In this case $a_{0,n}$ grows linearly in n , and many results have been proved, such as shape theorems, fluctuations of first-passage times, geometry of geodesics. On the other hand, a number of open problems have been proposed. For example, although we know the existence of the time constant, it is an old question to find a non-trivial explicit distribution for which we can determine the time constant. See [2] for a recent survey.

In the following, we list a few works on Bernoulli FPP (on \mathbb{T}); exact asymptotics for this special model are closely related to the scaling limits of critical and near-critical percolation.

- *Critical Bernoulli FPP.* Chayes et al. [8] proved that $\mathbb{E}_{p_c} a_{0,n} \asymp \log n$. Subsequently, Kesten and Zhang [27] showed that $\text{Var} a_{0,n} \asymp \log n$, and derived a central limit theorem for $a_{0,n}$. Using the full scaling limit of critical percolation (i.e., the conformal loop ensemble CLE_6 ; for the general CLE_κ , $8/3 \leq \kappa \leq 8$, see [32, 33]) established by Camia and Newman [10], we obtained a law of large numbers for $a_{0,n}$ in [38]. The idea is to define FPP on CLE_6 by using “loop chains”, similarly to the cluster chains we used in the present paper; the application of the subadditive ergodic theorem relies on the scaling invariance of full-plane CLE_6 , similarly to the case in the present paper that the application of this theorem relies on the translation invariance of the continuum cluster ensemble. Except for these similarities, the proof here is much more complicated than that in [38]. In [39] we improved the result in [38], giving explicit limit theorem by identifying the exact values of the constants appearing in the first-order asymptotics of $a_{0,n}$, $\mathbb{E}_{p_c} a_{0,n}$ and $\text{Var}_{p_c} a_{0,n}$. (Recently, Damron, Hanson and Lam [12] have extended this result to general critical FPP.) Analogous limit theorem for critical Bernoulli FPP starting on the boundary was established in [23] by using SLE_6 . In [40], we constructed different subsequential limits for $a_{0,n}$, relying on the large deviation estimates on the nesting of CLE_6 loops. Moreover, as discussed in Section 4 in [39], it is expected that as $t \rightarrow \infty$, the outer boundary of $B(t)$, properly scaled, converges in distribution to a “typical” loop of full-plane CLE_6 .
- *Near-critical Bernoulli FPP: supercritical regime.* In [40], relying on the limit theorem for the critical case, we derived exact first-order asymptotics for $T(0, \mathcal{C}_\infty)$ together with its expectation and variance, as $p \downarrow p_c$, where \mathcal{C}_∞ is the infinite cluster with 0-time sites.
- *Near-critical Bernoulli FPP: subcritical regime.* As mentioned earlier, Chayes et al. [8] proved that $\mu(p) \asymp L(p)^{-1}$ as $p \uparrow p_c$. This motivates our present work.

Besides the works on Bernoulli FPP mentioned above, we now describe several works which are related to the present paper in the sense that the scaling limits are used to extract geometric information about the original discrete models.

- Duminil-Copin [13] used the scaling limit of near-critical percolation established in [16] to show that the Wulff crystal for subcritical percolation on \mathbb{T} converges to a Euclidean disk as $p \uparrow p_c$. Roughly speaking, it was showed that the typical shape of a cluster conditioned to be large becomes round. Our strategy of the proof of Theorem 1.1 is rather different from [13].
- For percolation on \mathbb{T} , we contract each blue cluster into a single vertex and define a new edge between any pair of blue clusters if there is a yellow hexagon touching both of them. Then we obtain a random graph called “cluster graph”. In [40], we applied the limit theorem for critical Bernoulli FPP to study the graph distance in the cluster

graph at criticality. The discrete cluster chains we used in the present paper are (self-avoiding) paths in the cluster graph. In the near-critical regime, we will prove that with high probability the first-passage time between two large blue clusters is equal to their graph distance in the cluster graph; the proof uses some techniques we developed for the cluster graph at criticality in [40].

- Based on the scaling limit of near-critical percolation in [16], Garban, Pete and Schramm [17] constructed the scaling limits of the minimal spanning tree and the invasion percolation tree. In the proof they also used the cluster graph, similarly to that we described above but with some differences; see Fig. 1 in [17] and the paragraph above it. Except for this similarity, our proof is quite different from that in [17].

1.4. Organization of the paper

The rest of the paper is organized as follows. In Section 2, we introduce the notation and basic definitions, including correlation length, quad-crossing topology and near-critical scaling limit, and collect some results about critical and near-critical percolation that will be used. Section 3 is devoted to the construction of the continuum cluster ensemble, which is the scaling limit of the collection of blue clusters for near-critical percolation. In Section 4 we define FPP on continuum cluster ensembles, and obtain a law of large numbers for the point-to-point passage times of this continuum FPP. We also show that some quantities of the discrete FPP approximate their corresponding quantities of the continuum FPP as $p \uparrow p_c$. In Section 5, based on the results for the continuum FPP together with the asymptotics of the corresponding discrete quantities, we use the renormalization method to prove our main result.

2. Notation and preliminaries

Throughout this paper, C, K, C_j or K_j stands for a positive constant that may change from line to line according to the context. $\mathbb{N} = \{1, 2, \dots\}$ denotes the set of natural integers and $\mathbb{Z}_+ = \{0\} \cup \mathbb{N}$. We identify the plane \mathbb{R}^2 with the set \mathbb{C} of complex numbers in the usual way.

For two positive functions f and g from a set \mathcal{X} to $(0, \infty)$, we write $f(x) \asymp g(x)$ to indicate that $f(x)/g(x)$ is bounded away from 0 and ∞ , uniformly in $x \in \mathcal{X}$.

A **circuit** is a path (v_1, \dots, v_n) with $n \geq 3$, such that v_1 and v_n are neighbors. Note that the bonds $(v_1, v_2), \dots, (v_n, v_1)$ of the circuit form a Jordan curve, and sometimes the circuit is viewed as this curve.

For a rectangle of the form $R = [x_1, x_2] \times [y_1, y_2]$, we call a path (v_0, v_1, \dots, v_k) of \mathbb{T} a **left-right (resp. top-bottom) crossing** of R , if $v_1, \dots, v_{k-1} \in R$, and the line segments $\overline{v_0 v_1}$ and $\overline{v_{k-1} v_k}$ intersect the left and right (resp. top and bottom) sides of R , respectively. For a left-right crossing of R , if all the sites in it are blue (resp. yellow), we call it a **blue (resp. yellow) left-right crossing** of R .

For $r > 0$, define the box $\Lambda_r := [-r, r]^2$. For $0 < r < R$, define the annulus $A(r, R) := \Lambda_R \setminus \Lambda_r$. For $z \in \mathbb{C}$ and $\theta \in [0, 2\pi]$, we set $\Lambda_r^\theta(z) := z + e^{i\theta} \cdot \Lambda_r$ and $A^\theta(z; r, R) := z + e^{i\theta} \cdot A(r, R)$. Write $\Lambda_r(z) := \Lambda_r^0(z)$ and $A(z; r, R) := A^0(z; r, R)$.

The so-called arm events play a central role in studying near-critical percolation. We write 0 and 1 for “blue” and “yellow”, respectively. A **color sequence** $\sigma = (\sigma_1, \sigma_2, \dots, \sigma_k)$ is an element of $\{0, 1\}^k$ with $k \geq 1$, and its length $|\sigma|$ is k . For ease of notation, we write $(\sigma_1 \sigma_2 \dots \sigma_k) := (\sigma_1, \sigma_2, \dots, \sigma_k)$. We say that a color sequence is **polychromatic** if the sequence contains at least one 0 and one 1. We identify two sequences if they are the same up to a cyclic permutation. Suppose $\theta \in [0, 2\pi]$. For an annulus $A^\theta(z; r, R)$, we denote by $\mathcal{A}_\sigma^\theta(z; r, R)$ the event that there exist $|\sigma|$ disjoint monochromatic paths called **arms** in $A^\theta(z; r, R)$ connecting the two boundary pieces of $A^\theta(z; r, R)$, whose colors are those prescribed by σ , when taken in counterclockwise order. The half-plane arm events are defined similarly: We denote by $\mathcal{A}_\sigma^{\theta,+}(z; r, R)$ the event that $\mathcal{A}_\sigma^\theta(z; r, R)$ occurs with the arms contained in the half-plane $z + e^{i\theta} \cdot \{u \in \mathbb{C} : \Im u \geq 0\}$. More generally, given $\vartheta \in (0, 2\pi)$, we denote by $\mathcal{A}_\sigma^{\theta,\vartheta}(z; r, R)$ the wedge arm event that $\mathcal{A}_\sigma^\theta(z; r, R)$ occurs with the arms contained in the wedge $W(z; \theta, \vartheta) := z + e^{i\theta} \cdot \{u \in \mathbb{C} : \arg(u) \in [-\vartheta/2, \vartheta/2]\}$.

For ease of notation, we write $\mathcal{A}_\sigma(z; r, R) := \mathcal{A}_\sigma^0(z; r, R)$ when $\theta = 0$, and write $\mathcal{A}_\sigma(r, R) := \mathcal{A}_\sigma(0; r, R)$ when $z = 0$. Analogous abbreviations apply to $\mathcal{A}_\sigma^{\theta,+}(z; r, R)$ and $\mathcal{A}_\sigma^{\theta,\vartheta}(z; r, R)$. For any $r \geq 1$, we let $\mathcal{A}_\sigma(z; r, r)$ be the entire sample space. Write $\mathcal{A}_4 := \mathcal{A}_{(0101)}$ and $\mathcal{A}_6 := \mathcal{A}_{(011011)}$.

2.1. Correlation length

First, let us define the **correlation length** $L(p)$ that we mainly work with in this paper: For any $p < p_c$, we set

$$L(p) := \inf \left\{ R \geq 1 : R^2 \mathbf{P}_{p_c}[\mathcal{A}_4(1, R)] \geq \frac{1}{p_c - p} \right\},$$

where we let $L(p) = 1/2$ if the above set is empty. We chose to work with $L(p)$ because our proof is based on Corollary 1.7 of [16] (see Theorem 2.2 below) which gives the scaling limit of near-critical percolation on $L(p)^{-1}\mathbb{T}$ as $p \uparrow p_c$.

Now we introduce another definition of correlation length: For each $\epsilon \in (0, 1/2)$ and $p < p_c$, let

$$L_\epsilon(p) := \inf\{R \geq 1 : \mathbf{P}_p[\text{there is a blue left-right crossing of } [0, R]^2] \leq \epsilon\}.$$

The particular choice of ϵ is not important in the above definition. Indeed, for any $\epsilon, \epsilon' \in (0, 1/2)$, we have $L_\epsilon(p) \asymp L_{\epsilon'}(p)$; see, e.g., Corollary 37 of [30]. We refer the interested reader to Section 7 of [30] for three natural definitions of correlation length (called ‘‘characteristic length’’ there), including $L_\epsilon(p)$.

Kesten (see Proposition 34 of [30]) proved that

$$(p_c - p)L_\epsilon(p)^2 \mathbf{P}_{p_c}[\mathcal{A}_4(1, L_\epsilon(p))] \asymp 1 \quad \text{as } p \uparrow p_c.$$

This together with the quasi-multiplicativity (6) and the inequality (9) at p_c implies that for any $\epsilon \in (0, 1/2)$,

$$L(p) \asymp L_\epsilon(p) \quad \text{as } p \uparrow p_c.$$

It is well known that $L(p) \rightarrow +\infty$ when $p \uparrow p_c$. For convenience, we will take the convention that $L(p_c) = L_\epsilon(p_c) = +\infty$. In the following, the expression ‘‘for any $1 \leq R \leq L(p)$ ’’ must be interpreted as ‘‘for any $R \geq 1$ ’’ when $p = p_c$. For $R \geq 1$, define

$$p(R) := \inf\{p : L(p') \geq R \text{ for all } p' \in [p, p_c]\}.$$

Note that $p(R) \in (0, p_c)$ for all $R \geq 1$, and $p(R) \uparrow p_c$ as $R \rightarrow \infty$.

2.2. Classical results for planar percolation

We assume that the reader is familiar with the FKG inequality, the BK (van den Berg-Kesten) inequality, Reimer’s inequality [31], and the RSW (Russo-Seymour-Welsh) technology. Here we collect some classical results in critical and near-critical percolation which will be used. See, e.g., Section 2.2 in [5], Section 2.2 in [6], and [25, 30, 35, 37].

- (i) *RSW bounds.* For any $k, K \geq 1$, there exists a constant $\delta = \delta(k, K) > 0$, such that for all $p \in (p(1), p_c]$ and $1 \leq n \leq KL(p)$,

$$\mathbf{P}_p[\text{there is a blue left-right crossing of } [0, kn] \times [0, n]] \geq \delta.$$

- (ii) *A-priori bounds on 1-arm events.* There exist constants $\lambda_1, \lambda'_1 > 0$, such that for any $K \geq 1$, there are constants $C_1(K), C_2(K) > 0$, such that for all $p \in (p(1), p_c]$ and $1 \leq r < R \leq KL(p)$,

$$C_1 \left(\frac{r}{R}\right)^{\lambda'_1} \leq \mathbf{P}_p[\mathcal{A}_{(0)}(r, R)] \leq \mathbf{P}_p[\mathcal{A}_{(1)}(r, R)] \leq C_2 \left(\frac{r}{R}\right)^{\lambda_1}. \quad (5)$$

- (iii) *Quasi-multiplicativity.* For any color sequence σ and $K \geq 1$, there exist constants $C_1(|\sigma|, K), C_2(|\sigma|, K) > 0$, such that for all $p \in (p(1), p_c]$ and $1 \leq r_1 < r_2 < r_3 \leq KL(p)$,

$$C_1 \mathbf{P}_p[\mathcal{A}_\sigma(r_1, r_2)] \mathbf{P}_p[\mathcal{A}_\sigma(r_2, r_3)] \leq \mathbf{P}_p[\mathcal{A}_\sigma(r_1, r_3)] \leq C_2 \mathbf{P}_p[\mathcal{A}_\sigma(r_1, r_2)] \mathbf{P}_p[\mathcal{A}_\sigma(r_2, r_3)]. \quad (6)$$

- (iv) *Stability for arm events near criticality.* For any color sequence σ and $K \geq 1$, there exist constants $C_1(|\sigma|, K), C_2(|\sigma|, K) > 0$, such that for all $p \in (p(1), p_c)$ and $1 \leq r < R \leq KL(p)$,

$$C_1 \mathbf{P}_{p_c}[\mathcal{A}_\sigma(r, R)] \leq \mathbf{P}_p[\mathcal{A}_\sigma(r, R)] \leq C_2 \mathbf{P}_{p_c}[\mathcal{A}_\sigma(r, R)]. \quad (7)$$

- (v) *Exponential decay with respect to $L(p)$.* There exist constants $C_1, C_2 > 0$, such that for all $p \in (p(1), p_c)$ and $R \geq L(p)$,

$$\mathbf{P}_p[\exists \text{ a yellow circuit surrounding } 0 \text{ in } A(R, 2R)] \geq 1 - C_1 \exp\left(-C_2 \frac{R}{L(p)}\right). \quad (8)$$

This follows from FKG and (7.23) in [30]. (See also (17) of [40] for the corresponding inequality at $p > p_c$.)

- (vi) *Lower bound on the 4-arm exponent.* There exist constants $\lambda_4, C > 0$, such that for all $p \in (p(1), p_c]$ and $1 \leq r < R \leq L(p)$,

$$\mathbf{P}_p[\mathcal{A}_4(r, R)] \geq C \left(\frac{r}{R}\right)^{2-\lambda_4}. \quad (9)$$

- (vii) *Upper bound on the 6-arm exponent.* There is a $\lambda_6 > 0$, such that for any $K \geq 1$ and polychromatic color sequence σ with $|\sigma| = 6$, there exists a constant $C(K) > 0$, such that for all $p \in (p(1), p_c]$ and $1 \leq r < R \leq KL(p)$,

$$\mathbf{P}_p[\mathcal{A}_\sigma(r, R)] \leq C \left(\frac{r}{R}\right)^{2+\lambda_6}. \quad (10)$$

- (viii) *Upper bounds on half-plane 2-arm and wedge 3-arm events.* For any $K \geq 1$ and color sequences σ_2, σ_3 with $|\sigma_2| = 2$ and $|\sigma_3| = 3$, there exists a constant $C(K) > 0$, such that for all $p \in (p(1), p_c]$, $1 \leq r < R \leq KL(p)$ and $\theta \in [0, 2\pi]$,

$$\mathbf{P}_p[\mathcal{A}_{\sigma_2}^{\theta,+}(r, R)] \leq C \left(\frac{r}{R}\right), \quad (11)$$

$$\mathbf{P}_p[\mathcal{A}_{\sigma_3}^{\theta,+}(r, R)] \leq C \left(\frac{r}{R}\right)^2. \quad (12)$$

See, e.g., Lemma 6.8 in [36] and Appendix A in [28] for the critical case. This combined with the stability for arm events near criticality gives the above inequalities for the near-critical case. Moreover, given any fixed $\vartheta \in (0, 2\pi)$ and $\epsilon \in (0, 1)$, there exists a constant $C(\vartheta, \epsilon) > 0$ such that for all $p \in (p(1), p_c]$, $1 \leq r < R \leq L(p)$ and $\theta \in [0, 2\pi]$,

$$\mathbf{P}_p[\mathcal{A}_{\sigma_3}^{\theta,\vartheta}(r, R)] \leq C \left(\frac{r}{R}\right)^{2\pi/\vartheta-\epsilon}. \quad (13)$$

The conformal invariance of the scaling limit gives $\mathbf{P}_{p_c}[\mathcal{A}_{\sigma_3}^{\theta,\vartheta}(r, R)] = (\mathbf{P}_{p_c}[\mathcal{A}_{\sigma_3}^{\theta,\pi}(r, R)])^{\pi/\vartheta+o(1)}$, which combined with (12) implies the above inequality in the critical case. Then the stability for (wedge) arm events near criticality yields (13).

- (ix) *Upper bounds on arm events with monochromatic color sequence.* For any polychromatic color sequence σ , there exist $\epsilon, C > 0$ (depending on $|\sigma|$), such that for all $p \in (p(1), p_c]$ and $1 \leq r < R \leq L(p)$,

$$\mathbf{P}_p[\mathcal{A}_{\underbrace{1 \dots 1}_{|\sigma|}}(r, R)] \leq C \left(\frac{r}{R}\right)^\epsilon \mathbf{P}_p[\mathcal{A}_\sigma(r, R)]. \quad (14)$$

This inequality at p_c (which was used in [40] as Lemma 1) follows from the proof of Theorem 5 in [3]; see in particular Step 1 of the proof. Combining the inequality at p_c and (7) yields (14).

- (x) When $p \uparrow p_c$,

$$L(p) = (p_c - p)^{-4/3+o(1)}. \quad (15)$$

2.3. Space of quad-crossings

To describe the scaling limits of planar percolation, Schramm and Smirnov introduced the quad-crossing topology in [34]. Let us briefly recall this topology in this subsection. We shall use the notation and definitions from [16].

When taking the scaling limit of percolation on the whole plane, it is convenient to compactify \mathbb{C} into $\hat{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$ (i.e., the Riemann sphere) as follows. First, we replace the Euclidean metric with a distance function $\Delta(\cdot, \cdot)$ defined on $\mathbb{C} \times \mathbb{C}$ by

$$\Delta(x, y) := \inf_{\varphi} \int (1 + |\varphi(s)|^2)^{-1} ds,$$

where the infimum is over all smooth curves $\varphi(s)$ joining x with y , parameterized by arc length s . This metric is equivalent to the Euclidean metric in bounded regions. Then, we add a single point ∞ at infinity to get the compact space $\hat{\mathbb{C}}$ which is isometric, via stereographic projection, to the two-dimensional sphere.

Let $D \subset \hat{\mathbb{C}}$ be open. A **quad** in the domain D can be considered as a homeomorphism Q from $[0, 1]^2$ into D . The space of all quads in D , denoted by \mathcal{Q}_D , can be equipped with the following metric:

$$d(Q_1, Q_2) := \inf_{\phi} \sup_{z \in \partial[0,1]^2} |Q_1(z) - Q_2(\phi(z))|,$$

where the infimum is over all homeomorphisms $\phi : [0, 1]^2 \rightarrow [0, 1]^2$ which preserve the four corners of the square. A **crossing** of a quad Q is a connected closed subset of $Q \setminus ([0, 1]^2)$ that intersects both $Q(\{0\} \times [0, 1])$ and $Q(\{1\} \times [0, 1])$. From the point of view of crossings, there is a natural partial order on \mathcal{Q}_D : We write $Q_1 \leq Q_2$ if any crossing of Q_2 contains a crossing of Q_1 . Furthermore, we write $Q_1 < Q_2$ if there are open neighborhoods \mathcal{N}_i of Q_i for $i \in \{1, 2\}$, such that $N_1 \leq N_2$ holds for any $N_i \in \mathcal{N}_i$. We say that a subset $\mathcal{S} \subset \mathcal{Q}_D$ is **hereditary** if, whenever $Q \in \mathcal{S}$ and $Q' \in \mathcal{Q}_D$ satisfies $Q' < Q$, we have $Q' \in \mathcal{S}$. The collection of all closed hereditary subsets of \mathcal{Q}_D will be denoted by \mathcal{H}_D .

By introducing a natural topology, \mathcal{H}_D can be made into a compact metric space. Indeed, let

$$\begin{aligned} \sqsupset_Q &:= \{\mathcal{S} \in \mathcal{H}_D : Q \in \mathcal{S}\} && \text{for any } Q \in \mathcal{Q}_D, \\ \sqsupset_U &:= \{\mathcal{S} \in \mathcal{H}_D : U \cap \mathcal{S} = \emptyset\} && \text{for any open subset } U \subset \mathcal{Q}_D. \end{aligned}$$

Then we endow \mathcal{H}_D with the topology \mathcal{T}_D which is the minimal topology that contains every \sqsupset_Q^c and \sqsupset_U^c as open sets. It was proved in [34], Theorem 1.13, that for any nonempty open subset $D \subset \hat{\mathbb{C}}$, the topological space $(\mathcal{H}_D, \mathcal{T}_D)$ is a compact metrizable Hausdorff space. In particular, $(\mathcal{H}_D, \mathcal{T}_D)$ is a Polish space. Furthermore, for any dense $\mathcal{Q}_0 \subset \mathcal{Q}_D$, the events $\{\sqsupset_Q : Q \in \mathcal{Q}_0\}$ generate the Borel σ -field of \mathcal{H}_D . An arbitrary metric generating the topology \mathcal{T}_D will be denoted by $d_{\mathcal{H}}$. The above compactness property implies that (see Corollary 1.15 of [34]), the space of Borel probability measures of $(\mathcal{H}_D, \mathcal{T}_D)$, equipped with the weak* topology is a compact metrizable Hausdorff space.

When $D = \hat{\mathbb{C}}$, we write $\mathcal{H} := \mathcal{H}_{\hat{\mathbb{C}}}$ and $\mathcal{T} := \mathcal{T}_{\hat{\mathbb{C}}}$. With a slight abuse of notation, when we refer to Q as a subset of $\hat{\mathbb{C}}$ in the following, we consider its range $Q \setminus ([0, 1]^2) \subset \hat{\mathbb{C}}$.

Note that any discrete percolation configuration $\omega_p^\eta := \eta\omega_p$ on $\eta\mathbb{T}$, considered as a union of blue hexagons in the plane, naturally induce an element in \mathcal{H} : the set of all quads for which ω_p^η contains a crossing. By a slight abuse of notation, we will still denote by ω_p^η the point in \mathcal{H} corresponding to the percolation configuration ω_p^η . It follows that ω_p^η induces a probability measure on \mathcal{H} , denoted by \mathbf{P}_p^η .

2.4. Near-critical scaling limit

As in [16], we define the following near-critical parameter scale: For $\eta > 0$ and $\lambda \in \mathbb{R}$, we set

$$p_\lambda(\eta) := p_c + \lambda \frac{\eta^2}{\alpha_4^\eta(\eta, 1)},$$

where $\alpha_4^\eta(r, R)$ stands for the probability of the alternating 4-arm event in $A(r, R)$ for critical site percolation on $\eta\mathbb{T}$.

Recall that for each $p \in [0, 1]$, ω_p stands for Bernoulli site percolation on \mathbb{T} with intensity p , and \mathbf{P}_p stands for the law of ω_p . The following are two natural ways to define near-critical percolation on rescaled lattices:

- For $\eta > 0$ and $\lambda \in \mathbb{R}$, let $\omega_{p_\lambda(\eta)}^\eta$ denote the percolation configuration on $\eta\mathbb{T}$ with intensity $p_\lambda(\eta)$, and let $\mathbf{P}_{p_\lambda(\eta)}^\eta$ denote the law of $\omega_{p_\lambda(\eta)}^\eta$.
- For $p < p_c$, let $\omega_p^{L(p)^{-1}} = L(p)^{-1}\omega_p$ denote the percolation configuration on $L(p)^{-1}\mathbb{T}$ with intensity p , and let $\mathbf{P}_p^{L(p)^{-1}}$ denote the law of $\omega_p^{L(p)^{-1}}$. (Such near-critical percolation for $p > p_c$ can be defined analogously.)

Note that for $\mathbf{P}_{p_\lambda(\eta)}^\eta$, the intensity $p_\lambda(\eta)$ is a function of the mesh size η , while for $\mathbf{P}_p^{L(p)^{-1}}$, the mesh size $L(p)^{-1}$ is a function of the intensity p . As discussed in Section 2.3, we also view $\omega_{p_\lambda(\eta)}^\eta$ (resp. $\omega_p^{L(p)^{-1}}$) as an element in \mathcal{H} , and view $\mathbf{P}_{p_\lambda(\eta)}^\eta$ (resp. $\mathbf{P}_p^{L(p)^{-1}}$) as the probability measure on \mathcal{H} induced by $\omega_{p_\lambda(\eta)}^\eta$ (resp. $\omega_p^{L(p)^{-1}}$).

The following theorem states that for fixed λ , the near-critical percolation $\omega_{p_\lambda(\eta)}^\eta$ has a scaling limit as $\eta \rightarrow 0$.

Theorem 2.1 (Theorem 1.4 and Corollary 10.5 in [16]). *Fix $\lambda \in \mathbb{R}$. As $\eta \rightarrow 0$, the near-critical percolation $\omega_{p_\lambda(\eta)}^\eta$ converges in law in $(\mathcal{H}, d_{\mathcal{H}})$ to a limiting random percolation configuration, denoted by $\omega^\infty(\lambda)$. Moreover, the law of $\omega^\infty(\lambda)$, denoted by $\mathbf{P}^{\infty, \lambda}$, is invariant under translations and rotations.*

Note that our definition of $\omega_{p_\lambda(\eta)}^\eta$ is slightly different from $\omega_\eta^{mc}(\lambda)$ in Theorem 1.4 of [16] (see Section 1.2 in [16] for the definition), but this makes no essential difference; the scaling limit $\omega^\infty(\lambda)$ in Theorem 2.1 is denoted by $\omega_\infty^{mc}(2\lambda)$ in [16]. We want to mention that in [1, 5, 13], the authors also used the scaling limit result for $\omega_{p_\lambda(\eta)}^\eta$.

Similarly to Theorem 2.1 on $\omega_{p_\lambda(\eta)}^\eta$, the following theorem states that the near-critical percolation $\omega_p^{L(p)^{-1}}$ also has a scaling limit as $p \uparrow p_c$, which is a key input for the proof of our main result.

Theorem 2.2 (Corollary 1.7 of [16]). *As $p \uparrow p_c$, the near-critical percolation $\omega_p^{L(p)^{-1}}$ converges in law in $(\mathcal{H}, d_{\mathcal{H}})$ to $\omega^\infty(-1)$.*

Theorems 2.1 and 2.2 state that the near-critical percolation measures converge. Moreover, the convergence of quad-crossing probabilities also holds:

Lemma 2.3. *Fix $\lambda \in \mathbb{R}$. Let $D \subset \mathbb{C}$ be a bounded domain. Let $Q \in \mathcal{Q}_D$. We have*

$$\lim_{\eta \rightarrow 0} \mathbf{P}_{p_\lambda(\eta)}^\eta[\Box_Q] = \mathbf{P}^{\infty, \lambda}[\Box_Q]. \quad ((9.2) \text{ in } [16]) \quad (16)$$

Moreover,

$$\lim_{p \uparrow p_c} \mathbf{P}_p^{L(p)^{-1}}[\Box_Q] = \mathbf{P}^{\infty, -1}[\Box_Q]. \quad (17)$$

The proof of (16) in [16] uses Theorem 2.1 and the proof of Corollary 5.2 in [34], relying on the RSW estimates for near-critical percolation; the same proof also works for (17) by using Theorem 2.2 and the proof of Corollary 5.2 in [34].

2.5. Proof of Corollary 1.2

In this subsection, we describe how to derive Corollary 1.2 from Theorem 1.1.

For $R > 0$, denote by \Box_R the left-right crossing event in the quad $[0, R]^2$. As in the discrete model, we define a notion of correlation length for $\omega^\infty(\lambda)$: Given $\epsilon \in (0, 1/2)$ and $\lambda < 0$, define

$$L_\epsilon^\infty(\lambda) := \inf\{R > 0 : \mathbf{P}^{\infty, \lambda}[\Box_R] \leq \epsilon\}.$$

For $\lambda \in \mathbb{R}$ and $R > 0$, by using (16) we can define

$$f(\lambda, R) := \lim_{\eta \rightarrow 0} \mathbf{P}_{p_\lambda(\eta)}^\eta[\Box_R] = \mathbf{P}^{\infty, \lambda}[\Box_R].$$

By (25) in [1] and the argument below it, we know that for fixed $R > 0$, $f(\lambda, R)$ is absolutely continuous and strictly increasing in λ , and it satisfies $f(\lambda, R) \in (0, 1)$ and

$$\lim_{\lambda \rightarrow -\infty} f(\lambda, R) = 0 \quad \text{and} \quad \lim_{\lambda \rightarrow \infty} f(\lambda, R) = 1.$$

Furthermore, it is clear that $f(0, R) = 1/2$ for all $R > 0$, by the well-known duality $f(-\lambda, R) = 1 - f(\lambda, R)$. By Corollary 10.5 of [16], for any scaling parameter $\rho > 0$, $\rho \cdot \omega^\infty(\lambda)$ has the same law as $\omega^\infty(\rho^{-3/4}\lambda)$. Therefore,

$$f(\rho\lambda, R) = f(\lambda, \rho^{4/3}R).$$

(See (26) in [1] for general quad.) The above argument implies that $f(\lambda, R)$ is absolutely continuous and strictly decreasing in R , and for any fixed $\lambda < 0$,

$$\lim_{R \rightarrow \infty} f(\lambda, R) = 0 \quad \text{and} \quad \lim_{R \rightarrow 0} f(\lambda, R) = 1/2.$$

This implies that for any fixed $\lambda < 0$ and $\epsilon \in (0, 1/2)$,

$$L_\epsilon^\infty(\lambda) = \text{the unique } R \text{ such that } f(\lambda, R) = \epsilon; \quad (18)$$

furthermore, $f(\lambda, R) > \epsilon$ when $R < L_\epsilon^\infty(\lambda)$ and $f(\lambda, R) < \epsilon$ when $R > L_\epsilon^\infty(\lambda)$.

It is well known that $L(p) \asymp L_\epsilon(p)$ as $p \uparrow p_c$ (see Section 2.1). The following is a refinement of this result.

Lemma 2.4. *For any fixed $\epsilon \in (0, 1/2)$, we have*

$$\lim_{p \uparrow p_c} \frac{L_\epsilon(p)}{L(p)} = L_\epsilon^\infty(-1).$$

Proof. By (17), for any $R > 0$, we have

$$\lim_{p \uparrow p_c} \mathbf{P}_p^{L(p)^{-1}}[\Box_R] = f(-1, R).$$

Then using (18) and the statement below it, we have

$$\lim_{p \uparrow p_c} \mathbf{P}_p^{L(p)^{-1}} [\Xi_{L_\epsilon^\infty(-1)}] = f(-1, L_\epsilon^\infty(-1)) = \epsilon,$$

and furthermore,

$$\lim_{p \uparrow p_c} \mathbf{P}_p^{L(p)^{-1}} [\Xi_R] = f(-1, R) > \epsilon \text{ when } R < L_\epsilon^\infty(-1),$$

$$\lim_{p \uparrow p_c} \mathbf{P}_p^{L(p)^{-1}} [\Xi_R] = f(-1, R) < \epsilon \text{ when } R > L_\epsilon^\infty(-1).$$

By the definition of $L_\epsilon(p)$ we have

$$\frac{L_\epsilon(p)}{L(p)} = \inf \left\{ R \geq L(p)^{-1} : \mathbf{P}_p^{L(p)^{-1}} [\Xi_R] \leq \epsilon \right\}.$$

Since $L(p) \rightarrow \infty$ as $p \uparrow p_c$, we conclude by the above argument that $\lim_{p \uparrow p_c} L_\epsilon(p)/L(p) = L_\epsilon^\infty(-1)$. \square

Proof of Corollary 1.2. Theorem 1.1 combined with Lemma 2.4 yields Corollary 1.2. \square

2.6. Basic properties of Bernoulli FPP

We say that a finite set $D \subset V(\mathbb{T})$ is simply connected if the union of the hexagons $H_v, v \in D$, is simply connected. For a simply connected set D of sites, we denote by $\partial^- D$ its inner site boundary, that is, the set of sites of D that are adjacent to some site of $V(\mathbb{T}) \setminus D$. We call a simply connected subset D of \mathbb{T} a discrete Jordan set if $\partial^- D$ is a circuit. A **discrete quad** is a discrete Jordan set D together with four distinct sites v_1, v_2, v_3, v_4 of $\partial^- D$, appearing in this order as $\partial^- D$ is traversed counterclockwise. Given a discrete quad $(D; v_1, v_2, v_3, v_4)$, we define the **arc** $(v_k v_{k+1})$ to be the path from v_k to v_{k+1} (with $v_5 = v_1$) in $\partial^- D$ as $\partial^- D$ is traversed counterclockwise.

For two disjoint finite sets $S, S' \subset V(\mathbb{T})$, we say that a path γ in $V(\mathbb{T}) \setminus (S \cup S')$ separates S from S' if any path from S to S' must intersect γ . For $V_0 \subset V(\mathbb{T})$ and $S, S' \subset V_0$, define

$$T(S, S')(V_0) := \inf \{ T(\gamma) : \gamma \text{ is a path from a site in } S \text{ to a site in } S' \text{ and } \gamma \subset V_0 \}.$$

The following topological or combinatorial properties of passage times are very useful in studying Bernoulli FPP on \mathbb{T} .

Proposition 2.5. *Consider Bernoulli FPP on \mathbb{T} with parameter $p \in (0, 1)$. The following statements hold.*

(i) *Let $(D; v_1, v_2, v_3, v_4)$ be a discrete quad. We have*

$$T((v_1 v_2), (v_3 v_4))(D) = \text{the maximal number of disjoint yellow paths from } (v_2 v_3) \text{ to } (v_4 v_1) \text{ in } D.$$

(ii) *Let \mathcal{C} and \mathcal{C}' be two fixed distinct finite blue clusters. Then almost surely*

$$T(\mathcal{C}, \mathcal{C}') = \text{the maximal number of disjoint yellow circuits separating } \mathcal{C} \text{ from } \mathcal{C}'.$$

This implies that, almost surely, there exist $T(\mathcal{C}, \mathcal{C}')$ disjoint yellow circuits separating \mathcal{C} from \mathcal{C}' , such that any geodesic from \mathcal{C} to \mathcal{C}' must intersect each of these circuits in exactly one site.

Proof. (i) is a quad version of property (i) for the annulus passage times in Proposition 2 of [40], and its proof is essentially the same as the proof of that property (see, e.g., the first part of the proof of Proposition 2.4 in [39]), which is omitted here.

(ii) is a slightly general version of (i) and (ii) in Proposition 2 of [40]; their proofs are the same when \mathcal{C} surrounds \mathcal{C}' or vice versa. We now prove the case where \mathcal{C} does not surround \mathcal{C}' and vice versa. Let N denote the maximal number of disjoint yellow circuits separating \mathcal{C} from \mathcal{C}' . The inequality $T(\mathcal{C}, \mathcal{C}') \geq N$ is trivial, since if there are N disjoint yellow circuits separating \mathcal{C} from \mathcal{C}' then any path connecting \mathcal{C} and \mathcal{C}' must intersect each of these circuits.

It remains to show that $T(\mathcal{C}, \mathcal{C}') \leq N$ almost surely. Suppose that there exist almost surely N disjoint yellow circuits separating \mathcal{C} from \mathcal{C}' , with N_1 of them surrounding \mathcal{C} and $N_2 = N - N_1$ of them surrounding \mathcal{C}' . Write $\mathcal{C}_0 := \mathcal{C}$ and

$\mathcal{C}'_0 := \mathcal{C}'$. For $1 \leq k \leq N_1$, let \mathcal{C}_k denote the innermost yellow circuit surrounding \mathcal{C}_{k-1} ; for $1 \leq k \leq N_2$, let \mathcal{C}'_k denote the innermost yellow circuit surrounding \mathcal{C}'_{k-1} . It is clear that \mathcal{C}_k , $1 \leq k \leq N_1$ and \mathcal{C}'_k , $1 \leq k \leq N_2$ are N disjoint yellow circuits separating \mathcal{C} from \mathcal{C}' . We claim that almost surely either there is a blue path γ such that its starting site has a neighbor $v_{N_1} \in \mathcal{C}_{N_1}$ and its ending site has a neighbor $v'_{N_2} \in \mathcal{C}'_{N_2}$, or there is a $v_{N_1} \in \mathcal{C}_{N_1}$ and a $v'_{N_2} \in \mathcal{C}'_{N_2}$ such that they are neighbors; in the latter case we let $\gamma = \emptyset$. Suppose the claim does not hold. Then it is easy to see that there is a.s. a yellow circuit separating \mathcal{C}_{N_1} from \mathcal{C}'_{N_2} , and thus there are a.s. $N + 1$ disjoint yellow circuits separating \mathcal{C} from \mathcal{C}' , a contradiction. Next, (since \mathcal{C}_{N_1} is the innermost circuit surrounding \mathcal{C}_{N_1-1}) we take a path γ_{N_1} from v_{N_1} to a site which has a neighbor $v_{N_1-1} \in \mathcal{C}_{N_1-1}$, with all sites of $\gamma_{N_1} \setminus \{v_{N_1}\}$ being blue, and then take a path γ_{N_1-1} from v_{N_1-1} to a site which has a neighbor $v_{N_1-2} \in \mathcal{C}_{N_1-2}$, with all sites of $\gamma_{N_1-1} \setminus \{v_{N_1-1}\}$ being blue, and so on. The process stops after N_1 steps, and γ_1 is a path from $v_1 \in \mathcal{C}_1$ to a site in \mathcal{C} , with all sites of $\gamma_1 \setminus \{v_1\}$ being blue. Similarly we take paths $\gamma'_{N_2}, \gamma'_{N_2-1}, \dots, \gamma'_1$. Then we concatenate the paths $\gamma_1, \dots, \gamma_{N_1}, \gamma, \gamma'_{N_2}, \dots, \gamma'_1$ to obtain a path Γ from \mathcal{C} to \mathcal{C}' , such that $T(\Gamma) = N$, which implies $T(\mathcal{C}, \mathcal{C}') \leq N$ almost surely. \square

In the rest of this paper, we mainly work on $\mathbf{P}_p^{L(p)^{-1}}$. For notational convenience, we write $\eta = \eta(p) := L(p)^{-1}$. Suppose that $w, h > 0$. For the rectangle $[0, w] \times [0, h]$, define the **line-to-line passage time** $l_{w,h}^p$ by

$$l_{w,h}^p = l_{w,h}^p(\omega_p^\eta) := \inf\{T(\gamma) : \gamma \text{ is a left-right crossing of } [0, w] \times [0, h] \text{ in } \eta\mathbb{T}\}.$$

More generally, the line-to-line passage time $l_{w,h}^{p,\theta}(z)$ corresponding to the rectangle $z + e^{i\theta}([0, w] \times [0, h])$, with $z \in \mathbb{C}$ and $\theta \in [0, 2\pi]$, is defined similarly as above. Note that $l_{w,h}^{p,0}(0) = l_{w,h}^p$.

Lemma 2.6. *There exist constants $C_1, C_2 > 0$ and $K \geq 2$, such that for all $p \in (p(10), p_c)$, $2\eta \leq h \leq 1$, $w \geq h$, $x \geq Kw/h$, $z \in \mathbb{C}$ and $\theta \in [0, 2\pi]$,*

$$\mathbf{P}_p^\eta \left[l_{w,h}^{p,\theta}(z) \geq x \right] \leq C_1 \exp(-C_2 x).$$

Proof. For simplicity, we shall show the lemma in the case $\theta = 0, z = 0$ and $h = 1$; the proof extends immediately to the general case.

Suppose $w \geq 1$ and $p \in (p(10), p_c)$. Let D_w^η be the largest discrete Jordan set of $\eta\mathbb{T}$ in $[0, w] \times [0, 1]$. Let $v_1, v_2, v_3, v_4 \in \partial^- D_w^\eta$ be four sites closest to the four points $(0, 1), (0, 0), (w, 0), (w, 1)$, respectively. Then we get a discrete quad $(D_w^\eta; v_1, v_2, v_3, v_4)$. It is easy to see that

$$l_{w,1}^p \leq T((v_1 v_2), (v_3 v_4))(D_w^\eta) + 4.$$

Using (i) of Proposition 2.5 and the above inequality, we have

$$\begin{aligned} l_{w,1}^p - 4 &\leq \text{the maximal number of disjoint yellow paths from } (v_2 v_3) \text{ to } (v_4 v_1) \text{ in } D_w^\eta \\ &\leq \text{the maximal number of disjoint yellow top-bottom crossings of } [0, w] \times [1/4, 3/4] \\ &:= \widetilde{l_{w,1}^p}. \end{aligned}$$

The argument in the following is analogous to Step 3 of the proof of Theorem 5 in [3]. Observe that any yellow top-bottom crossing of $[0, w] \times [1/4, 3/4]$ must either cross a rectangle in $\{[j/2, j/2 + 1] \times [1/4, 3/4] : j = 0, 1, \dots, \lfloor 2w \rfloor - 1\}$ from top to bottom, or a square in $\{[j/2, j/2 + 1/2] \times [1/4, 3/4] : j = 0, 1, \dots, \lfloor 2w \rfloor - 1\}$ from left to right. Therefore, if $\widetilde{l_{w,1}^p} \geq x$, then there are $\lfloor x \rfloor$ disjoint yellow top-bottom crossings of $[0, w] \times [1/4, 3/4]$, each one crossing a rectangle from the two families of rectangles above. By RSW, there exists a universal constant $\delta \in (0, 1)$, such that for any rectangle from the two families of rectangles above, the probability of the event that this rectangle has a yellow top-bottom or left-right crossing is bounded above by δ . This combined with the BK inequality implies that

$$\mathbf{P}_p^\eta \left[\widetilde{l_{w,1}^p} \geq x \right] \leq \sum_{n_1 + n_2 + \dots + n_{\lfloor 2w \rfloor} = \lfloor x \rfloor} \delta^{\lfloor x \rfloor} = \binom{\lfloor x \rfloor + 2\lfloor 2w \rfloor}{2\lfloor 2w \rfloor} \delta^{\lfloor x \rfloor},$$

where $n_1, n_2, \dots, n_{\lfloor 2w \rfloor} \in \mathbb{Z}_+$. Then using Stirling's formula we obtain that, there exist constants $C_3, C_4 > 0$ and $K_1 \geq 2$, such that for all $w \geq 1, x \geq K_1 w$ and $p \in (p(10), p_c)$,

$$\mathbf{P}_p^\eta \left[\widetilde{l_{w,1}^p} \geq x \right] \leq C_3 \exp(-C_4 x),$$

which implies the desired result for $l_{w,1}^p$ since $l_{w,1}^p \leq \widetilde{l_{w,1}^p} + 4$. \square

We will also consider annulus-crossing times. A path (v_0, v_1, \dots, v_k) of $\eta\mathbb{T}$ is called a crossing of $A(z; r_1, r_2)$ if $v_1, \dots, v_{k-1} \in A(z; r_1, r_2)$ and the line segments $\overline{v_0 v_1}$ and $\overline{v_{k-1} v_k}$ intersect the two boundary pieces of $A(z; r_1, r_2)$, respectively. Let

$$X^p(z; r_1, r_2) := \inf\{T(\gamma) : \gamma \text{ is a crossing of } A(z; r_1, r_2) \text{ in } \eta\mathbb{T}\}.$$

The next lemma is an annulus analog of Lemma 2.6, and is a near-critical analog of Corollary 2.3 in [38] for the critical case.

Lemma 2.7. *There exist constants $K_1, K_2 > 0$ and $K \geq 2$, such that for all $p \in (p(10), p_c)$, $\eta \leq r_1 \leq r_2/2 \leq 2$, $x \geq K \log_2(r_2/r_1)$ and $z \in \mathbb{C}$,*

$$\mathbf{P}_p^\eta[X^p(z; r_1, r_2) \geq x] \leq K_1 \exp(-K_2 x).$$

Proof. The proof is very similar to the proof of Lemma 2.6, so we only give a sketch. By using (i) of Proposition 2 in [40], proving the lemma boils down to estimate the maximal number of disjoint yellow circuits surrounding z in $A(z; r_1, r_2)$. Then one can use the argument in Step 3 of the proof of Theorem 5 in [3] to obtain it, based on BK inequality and near-critical RSW. \square

The square lattice has site set \mathbb{Z}^2 and bond set $E(\mathbb{Z}^2)$ obtained by connecting all pairs $u, v \in \mathbb{Z}^2$ for which $|u - v| = 1$. In a standard abuse of notation, we write \mathbb{Z}^2 to denote this graph. Let $\mathbf{P}_{\mathbb{Z}^2, p}^{\text{site}}$ (resp. $\mathbf{P}_{\mathbb{Z}^2, p}^{\text{bond}}$) denote the Bernoulli site (resp. bond) percolation measure on \mathbb{Z}^2 with parameter p , defined similarly as the measure \mathbf{P}_p on \mathbb{T} . Here we adapt the usual setting for Bernoulli FPP: Let each site (resp. bond) of \mathbb{Z}^2 take the value 0 (open) with probability p , and take the value 1 (closed) with probability $1 - p$.

In the following theorem, we will compare locally dependent fields with Bernoulli percolation measures. A family $Y = \{Y_v : v \in \mathbb{Z}^2\}$ of random variables is called k -dependent if any two sub-families $\{Y_v : v \in A\}$ and $\{Y_v : v \in A'\}$ are independent whenever the graph distance between v and v' is larger than k for all $v \in A$ and $v' \in A'$. We denote by $Z^p = \{Z_v^p : v \in \mathbb{Z}^2\}$ an i.i.d. family of Bernoulli random variables which has the law $\mathbf{P}_{\mathbb{Z}^2, p}^{\text{site}}$.

Theorem 2.8 (Theorem 7.65 of [19]). *Let $k \in \mathbb{N}$. There exists a nondecreasing function $\pi : [0, 1] \rightarrow [0, 1]$ satisfying $\pi(\delta) \rightarrow 1$ as $\delta \rightarrow 1$ such that the following assertion holds. If $Y = \{Y_v : v \in \mathbb{Z}^2\}$ is a k -dependent family of random variables satisfying*

$$\mathbf{P}[Y_v = 1] \geq \delta \quad \text{for all } v \in \mathbb{Z}^2,$$

then we have the stochastic domination: $Y \geq_{st} Z^{1-\pi(\delta)}$.

The following proposition is a site version of Theorem 2.3 in [26] in the case $d = 2$. (Note that Theorem 2.3 in [26] is a special case of Proposition 5.8 in [24].)

Proposition 2.9 (Theorem 2.3 of [26]). *If $p < p_c^{\text{site}}(\mathbb{Z}^2)$, then there are constants $\epsilon, C_1, C_2 > 0$ depending on p , such that for all $n \in \mathbb{N}$,*

$$\mathbf{P}_{\mathbb{Z}^2, p}^{\text{site}} \left[\begin{array}{l} \text{there exists a path starting from } 0 \text{ with at least} \\ n \text{ sites and fewer than } \epsilon n \text{ closed sites} \end{array} \right] \leq C_1 \exp(-C_2 n).$$

Let $\omega_{\mathbb{Z}^2}^{\text{bond}}$ denote the bond percolation configuration on \mathbb{Z}^2 . The **chemical distance** $D(u, v)(\omega_{\mathbb{Z}^2}^{\text{bond}})$ between two sites u and v in \mathbb{Z}^2 is defined by

$$D(u, v)(\omega_{\mathbb{Z}^2}^{\text{bond}}) := \inf\{\text{the number of bonds of } \gamma : \gamma \text{ is a closed path connecting } u \text{ and } v\}.$$

If u and v are not in the same closed cluster, we set $D(u, v) = \infty$. The following proposition is a corollary of Theorem 1.4 in [18].

Proposition 2.10. *For any $\epsilon > 0$, there exists $p_0 = p_0(\epsilon) \in (0, 1/2)$, such that for all $p \in [0, p_0]$ and all large n (depending on ϵ),*

$$\mathbf{P}_{\mathbb{Z}^2, p}^{\text{bond}}[D(0, n) \leq (1 + \epsilon)n] \geq 1 - \epsilon.$$

Proof. Given two sites $u, v \in \mathbb{Z}^2$, we denote by $u \leftrightarrow_c v$ the event that there is a closed path connecting u and v , and by $v \leftrightarrow_c \infty$ the event that v is in an infinite closed cluster. Let $\theta(p) := \mathbf{P}_{\mathbb{Z}^2, p}^{\text{bond}}[0 \leftrightarrow_c \infty]$.

It is well known that there is a.s. a unique infinite closed cluster when $0 \leq p < 1/2 = p_c^{\text{bond}}(\mathbb{Z}^2)$ and there is a.s. no infinite closed cluster when $1/2 \leq p \leq 1$; see e.g. [19]. From this and the FKG inequality, we obtain that for all $n \in \mathbb{N}$,

$$\mathbf{P}_{\mathbb{Z}^2, p}^{\text{bond}}[0 \leftrightarrow_c n] \geq \mathbf{P}_{\mathbb{Z}^2, p}^{\text{bond}}[0 \leftrightarrow_c \infty, n \leftrightarrow_c \infty] \geq \theta(p)^2. \quad (19)$$

It is also well known that $\theta(p)$ is a continuous function of p on the interval $[0, 1]$ (for bond percolation on \mathbb{Z}^2); see e.g. [19]. Moreover, it is clear that $\theta(0) = 1$. These facts and (19) imply that

$$\mathbf{P}_{\mathbb{Z}^2, p}^{\text{bond}}[0 \leftrightarrow_c n] \rightarrow 1 \quad \text{uniformly in } n \text{ as } p \rightarrow 0. \quad (20)$$

By Theorem 1.4 of [18], for each $\epsilon > 0$, there exists $p_1(\epsilon) \in (0, 1/2)$, such that for every $p \in [0, p_1(\epsilon))$,

$$\limsup_{n \rightarrow \infty} \frac{\log \mathbf{P}_{\mathbb{Z}^2, p}^{\text{bond}}[0 \leftrightarrow_c n, D(0, n) \geq (1 + \epsilon)n]}{n} < 0. \quad (21)$$

Combining (20) and (21), we obtain the desired result. \square

3. Scaling limit of near-critical percolation clusters

The proof of our main result relies on the scaling limit of near-critical percolation clusters. We start by setting notation in Section 3.1. By using the scaling limit results in [16] and the approach from [9], we construct the scaling limit of the collection of blue clusters for near-critical percolation in Section 3.2.

3.1. Setting and notation

For ease of notation, we write $\omega^\infty := \omega^\infty(-1)$ and $\mathbf{P}^\infty := \mathbf{P}^{\infty, -1}$, where $\omega^\infty(-1)$ and $\mathbf{P}^{\infty, -1}$ are defined in Theorem 2.1. Recall that $\eta = \eta(p) := L(p)^{-1}$.

Consider Bernoulli percolation on $\eta\mathbb{T}$ with $p \in (p(1), p_c)$. We shall view each site in $\eta\mathbb{T}$ as its corresponding (topologically closed) regular hexagons in $\eta\mathbb{H}$. **Clusters** are connected components of blue or yellow hexagons. In Section 3 we consider ω_p^η as a union of blue clusters, and construct the limit of large clusters in ω_p^η as $p \uparrow p_c$. For this purpose, we need to introduce some notation borrowed from [9], with small modifications adapted to our setup.

For a set $A \subset \mathbb{C}$, let $\text{diam}_\infty(A)$ denote the L^∞ -diameter of A . If D is a simply connected region with piecewise smooth boundary, we let $\mathcal{C}_D^p(\delta)$ denote the collection of (blue) clusters of ω_p^η , which are contained in D and have L^∞ -diameters at least δ . That is,

$$\mathcal{C}_D^p(\delta) := \{C \text{ is a cluster in } \omega_p^\eta : C \subset D \text{ and } \text{diam}_\infty(C) \geq \delta\}.$$

Let \mathcal{C}_D^p denote the collection of all clusters of ω_p^η which are contained in D , and let $\mathcal{C}^p(\delta)$ denote the collection of all clusters of ω_p^η with L^∞ -diameter at least δ . We write $\mathcal{C}_k^p(\delta) := \mathcal{C}_{\Lambda_k}^p(\delta)$ for short.

Let A, B be two subsets of \mathbb{C} . The Hausdorff distance between A, B is defined by

$$d_H(A, B) := \inf\{\epsilon > 0 : A + \Lambda_\epsilon \supset B \text{ and } B + \Lambda_\epsilon \supset A\}, \quad (22)$$

where $A + \Lambda_\epsilon := \{x + y \in \mathbb{C} : x \in A, y \in \Lambda_\epsilon\}$. Denote by \mathcal{S}_R the complete separable metric space of closed connected subsets of Λ_R with the metric (22).

Recall the definition of the distance function $\Delta(\cdot, \cdot)$ in Section 2.3. The distance D_H between subsets of $\hat{\mathbb{C}}$ is defined by

$$D_H(A, B) := \inf\{\epsilon > 0 : \forall x \in A, \exists y \in B \text{ such that } \Delta(x, y) \leq \epsilon \text{ and vice versa}\}. \quad (23)$$

Denote by \mathcal{S}_∞ the complete separable metric space of closed connected subsets of $\hat{\mathbb{C}}$ with the metric (23).

The distance $\widehat{\text{dist}}$ between finite collections i.e., sets of subsets of \mathbb{C} , denoted by $\mathcal{S}, \mathcal{S}'$, is defined by

$$\widehat{\text{dist}}(\mathcal{S}, \mathcal{S}') := \min_{\phi} \max_{S \in \mathcal{S}} d_H(S, \phi(S)), \quad (24)$$

where the infimum is taken over all bijections $\phi : \mathcal{S} \rightarrow \mathcal{S}'$. In case $|\mathcal{S}| \neq |\mathcal{S}'|$ we define the distance to be infinite. To account for possibly infinite collections, \mathcal{S} and \mathcal{S}' , of subsets of \mathbb{C} , we define

$$\text{dist}(\mathcal{S}, \mathcal{S}') := \inf\{\epsilon : \forall A \in \mathcal{S}, \exists B \in \mathcal{S}' \text{ such that } d_H(A, B) \leq \epsilon \text{ and vice versa}\}. \quad (25)$$

Similarly, for collections \mathcal{S} and \mathcal{S}' of subsets of $\hat{\mathbb{C}}$, we write

$$\text{Dist}(\mathcal{S}, \mathcal{S}') := \inf\{\epsilon : \forall A \in \mathcal{S}, \exists B \in \mathcal{S}' \text{ such that } D_H(A, B) \leq \epsilon \text{ and vice versa}\}. \quad (26)$$

Note that the metrics dist and Dist are equivalent on bounded regions, and convergence in $\widehat{\text{dist}}$ implies convergence in dist and Dist . Moreover, the space Ω_R (resp. Ω_∞) of closed subsets of \mathcal{S}_R (resp. \mathcal{S}_∞) with the metric dist (resp. Dist) is also a complete separable metric space. We denote by \mathcal{B}_R (resp. \mathcal{B}_∞) its Borel σ -algebra.

3.2. Scaling limit of the collection of clusters

In this subsection we show that the collection of large clusters of ω_p^η has a scaling limit as $p \uparrow p_c$.

Note that Theorem 2.2, combined with the fact that $(\mathcal{H}, d_{\mathcal{H}})$ is a Polish metric space (see Section 2.3), implies that there is a coupling of the measures (\mathbf{P}_p^η) and \mathbf{P}^∞ on $(\mathcal{H}, d_{\mathcal{H}})$ in which $\omega_p^\eta \rightarrow \omega^\infty$ a.s. as $p \uparrow p_c$.

The following theorem states that in a bounded region, the collection of clusters converges to a collection of closed connected sets (the ‘‘continuum clusters’’) as $p \uparrow p_c$. It is an analog of Theorems 1 and 11 in [9].

Theorem 3.1. *Let $k > \delta > 0$, and let \mathbf{P} be a coupling such that $\omega_p^\eta \rightarrow \omega^\infty$ in $(\mathcal{H}, d_{\mathcal{H}})$ a.s. as $p \uparrow p_c$. Then, as $p \uparrow p_c$, $\mathcal{C}_k^p(\delta)$ converges in \mathbf{P} -probability, in the metric $\widehat{\text{dist}}$, to a collection of closed connected sets in the interior of Λ_k , which we denote by $\mathcal{C}_k(\delta)$. Moreover, as $p \uparrow p_c$, \mathcal{C}_k^p converges in \mathbf{P} -probability, in the metric dist , to a collection of closed connected sets which we denote by \mathcal{C}_k . Furthermore, $\mathcal{C}_k(\delta)$ and \mathcal{C}_k are measurable functions of ω^∞ .*

The following theorem extends the above theorem to the case of the full plane and, moreover, states that the collection of full-plane continuum clusters is invariant under rotations and translations. It is an analog of Theorems 3 and 4 in [9].

Theorem 3.2. *Let \mathbb{P}_k denote the distribution of \mathcal{C}_k . There exists a unique probability measure \mathbb{P} on $(\Omega_\infty, \mathcal{B}_\infty)$ that is supported on collections of bounded (in the Euclidean metric), closed, connected subsets of $\hat{\mathbb{C}}$, which is the full plane limit of \mathbb{P}_k in the sense that, $\mathbb{P}|_{\Lambda_k} = \mathbb{P}_k$ for each $k \in \mathbb{N}$. Moreover, the following statements hold:*

- *Let \mathbf{P} be a coupling such that $\omega_p^\eta \rightarrow \omega^\infty$ in $(\mathcal{H}, d_{\mathcal{H}})$ a.s. as $p \uparrow p_c$. Then, as $p \uparrow p_c$, \mathcal{C}^p converges in \mathbf{P} -probability, in the metric Dist , to a collection of bounded, closed, connected subsets of $\hat{\mathbb{C}}$ which we denote by \mathcal{C} . Moreover, \mathcal{C} is a measurable function of ω^∞ and the distribution of \mathcal{C} is \mathbb{P} . Furthermore, for each $k \in \mathbb{N}$ and $\delta > 0$, $\mathcal{C}(\delta)|_{\Lambda_k}$ is a.s. a finite set, and $\widehat{\text{dist}}(\mathcal{C}^p(\delta)|_{\Lambda_k}, \mathcal{C}(\delta)|_{\Lambda_k})$, $\text{dist}(\mathcal{C}^p|_{\Lambda_k}, \mathcal{C}|_{\Lambda_k}) \rightarrow 0$ in probability as $p \uparrow p_c$.*
- *For any $k, n \in \mathbb{N}$ with $n \geq 2k + 1$, the configurations $\mathcal{C}|_{\Lambda_k}$ and $\mathcal{C}|_{\Lambda_{k(n)}}$ are independent.*
- *For $\theta \in [0, 2\pi]$ and $x \in \mathbb{C}$, let $f(z) := e^{i\theta}z + x$ be a map from \mathbb{C} to \mathbb{C} . Set $f(\mathcal{C}) := \{f(C) : C \in \mathcal{C}\}$. Then $f(\mathcal{C})$ and \mathcal{C} have the same distribution. That is, \mathbb{P} is invariant under rotations and translations.*

The proofs of Theorems 3.1 and 3.2 are analogous to those of the corresponding results in the critical case in [9]. Before moving to the proofs, we need to define the arm events that are measurable in the Borel σ -field of the quad-crossing topology. These events were first introduced in [15]. We shall borrow the notation and definitions from [9], with a slight modification adapted to our setting. (Note that for our FPP model a site is open when it takes the value 0, while in the standard percolation model a site is open when it takes the value 1, so the meaning of the color sequence in the present paper is opposite to that in [9].)

For $S \subset \hat{\mathbb{C}}$, let ∂S and $\text{int}(S)$ denote the boundary and interior of S , respectively. The arm events are defined as follows; see Figure 2.

Definition 3.3. Let $l \in \mathbb{Z}_+$ and $\kappa \in \{0, 1\}^l$. Let S be $\hat{\mathbb{C}}$, or a simply connected subset of $\hat{\mathbb{C}}$ with piecewise smooth boundary. Let D, E be two disjoint simply connected subsets of $\hat{\mathbb{C}}$ with piecewise smooth boundaries. Let $D \xleftrightarrow{\kappa, S} E$ denote the event that there are $\delta > 0$ and quads $Q_i \in \mathcal{Q}_{\text{int}(S)}$ for $i = 1, 2, \dots, l$ which satisfy the following conditions:

1. $\omega \in \Xi_{Q_i}$ for $i \in \{1, 2, \dots, l\}$ with $\kappa_i = 0$ and $\omega \in \Xi_{Q_i}^c$ for $i \in \{1, 2, \dots, l\}$ with $\kappa_i = 1$.
2. For all $i, j \in \{1, 2, \dots, l\}$ with $i \neq j$ and $\kappa_i = \kappa_j$, the quads Q_i and Q_j , viewed as subsets of $\hat{\mathbb{C}}$, are disjoint, and are at distance at least δ from each other and from the boundary of S .
3. $\Lambda_\delta + Q_i(\{0\} \times [0, 1]) \subset D$ and $\Lambda_\delta + Q_i(\{1\} \times [0, 1]) \subset E$ for $i \in \{1, 2, \dots, l\}$ with $\kappa_i = 0$.
4. $\Lambda_\delta + Q_i([0, 1] \times \{0\}) \subset D$ and $\Lambda_\delta + Q_i([0, 1] \times \{1\}) \subset E$ for $i \in \{1, 2, \dots, l\}$ with $\kappa_i = 1$.




FIG 2. Defining the arm event $D \xleftrightarrow{\kappa, S} E$ (in Definition 3.3) by using quad-crossings. Here, $\kappa = (1010)$, D is the gray square, S is the upper half-plane, E is the gray region outside the white region in the upper half-plane; the quads Q_2, Q_4 are crossed ($\kappa_2 = \kappa_4 = 0$), shown using blue curves, while the quads Q_1, Q_3 are not crossed ($\kappa_1 = \kappa_3 = 1$), shown using yellow curves (i.e., dual crossed).




FIG 3. Illustration of the event $\mathcal{A}_{(0),(101)}^3(a, b)$.

5. The intersections $Q_i \cap D$, for $i = 1, 2, \dots, l$, are at distance at least δ from each other; the same holds for $Q_i \cap E$.
6. A counterclockwise order of the quads Q_i , for $i = 1, 2, \dots, l$, is given by ordering counterclockwise the connected components of $Q_i \cap D$ containing $Q_i(0, 0)$.

We write $D \xleftrightarrow{\kappa} E$ for $D \xleftrightarrow{\kappa, \hat{\mathbb{C}}} E$.

In the following we consider some special arm events. For $z \in \mathbb{C}$ and $a > 0$, let $H_1(z, a), H_2(z, a), H_3(z, a), H_4(z, a)$ denote the left, lower, right, and upper half-planes which have the right, top, left and bottom sides of $\Lambda_a(z)$ on their boundaries, respectively. For $j = 1, 2, 3, 4$, $\kappa \in \{0, 1\}^l$ and $\kappa' \in \{0, 1\}^{l'}$ with $l, l' \geq 0$, we define the event $\mathcal{A}_{\kappa, \kappa'}^j(z; a, b)$ where there are $l + l'$ disjoint arms with color sequence $\kappa \vee \kappa' := (\kappa_1, \dots, \kappa_l, \kappa'_1, \dots, \kappa'_{l'})$ in $A(z; a, b)$ so that the l' arms, with color sequence κ' , are in the half-plane $H_j(z, a)$. That is,

$$\mathcal{A}_{\kappa, \kappa'}^j(z; a, b) := \left\{ \Lambda_a(z) \xleftrightarrow{\kappa \vee \kappa'} \left(\hat{\mathbb{C}} \setminus \Lambda_b(z) \right) \right\} \cap \left\{ \Lambda_a(z) \xleftrightarrow{\kappa', H_j(z, a)} \left(\hat{\mathbb{C}} \setminus \Lambda_b(z) \right) \right\}.$$

In the notation above, when z is omitted, it is assumed to be 0. When $\kappa' = \emptyset$, both the subscript κ' and the superscript j will be omitted. See Figure 3 for an illustration of an arm event.

Lemma 3.4 below is a near-critical analog of Lemma 10 of [9] in the critical case. As noted in [9], although Lemma 2.9 in [15] is a slightly weaker version of Lemma 10 in [9], the proof of the former extends immediately to that of the

latter. Similarly, Lemma 2.5 in [17] (see also Theorem 9.5 in [16]) is a slightly weaker version of Lemma 3.4, and its proof extends to this more general case. (The proof of Lemma 2.9 in [15] works, using the stability for arm events near criticality (7), together with the existence of the near-critical scaling limit, Theorem 2.2.)

Lemma 3.4. *Under a coupling \mathbf{P} of (\mathbf{P}_p^η) and \mathbf{P}^∞ on $(\mathcal{H}, d_{\mathcal{H}})$ such that $\omega_p^\eta \rightarrow \omega^\infty$ almost surely as $p \uparrow p_c$, we have for events $\mathcal{D} \in \{\{A \xleftarrow{(0),S} B\}, \{A \xleftarrow{(101),S} B\}, \mathcal{A}_{\kappa, \kappa'}^j(z; a, b)\}$,*

$$\mathbf{1}_{\mathcal{D}}(\omega_p^\eta) \rightarrow \mathbf{1}_{\mathcal{D}}(\omega^\infty) \quad \text{in } \mathbf{P}\text{-probability as } p \uparrow p_c,$$

for $(\kappa, \kappa') \in \{((0), \emptyset), ((0101), \emptyset), ((101010), \emptyset), (\emptyset, (101)), ((0), (101))\}$, rectangle $S \subset \mathbb{C}$, $j \in \{1, 2, 3, 4\}$, $0 < a < b$ and A, B disjoint simply connected subsets of \mathbb{C} with piecewise smooth boundaries.

The following bounds on the probabilities of some special arm events will be used later. Note that inequalities (10) and (12) imply (27) immediately; a combination of the right-hand inequality in (27), Reimer's inequality and (5) gives (28).

Lemma 3.5. *There exist constants $\lambda_6, \lambda_{1,3} > 0$, such that for any fixed $K \geq 1$, there exists a constant $C(K) > 0$, such that for all $p \in (p(1), p_c)$, $\eta \leq a < b \leq K$ and $j \in \{1, 2, 3, 4\}$,*

$$\mathbf{P}_p^\eta[\mathcal{A}_{(101010)}(a, b)] \leq C \left(\frac{a}{b}\right)^{2+\lambda_6}, \quad \mathbf{P}_p^\eta[\mathcal{A}_{\emptyset, (101)}^j(a, b)] \leq C \left(\frac{a}{b}\right)^2, \quad (27)$$

$$\mathbf{P}_p^\eta[\mathcal{A}_{(0), (101)}^j(a, b)] \leq C \left(\frac{a}{b}\right)^{2+\lambda_{1,3}}. \quad (28)$$

Now we are ready to prove Theorems 3.1 and 3.2. Since the proofs are analogous to those of the corresponding results in the critical case in [9], we only give a sketch.

Sketch of proof of Theorem 3.1. The proof is essentially the same as the proof of Theorem 11 (see also Theorem 1 in [9] for its weaker version) in [9], by using estimates on near-critical arm events and the existence of the near-critical scaling limit. We shall give a sketch of the proof below, and refer the reader to Sections 5 and 6 of [9] for the details.

Step 1. As in Section 5 of [9], we construct two approximations of blue clusters with L^∞ -diameters at least $\delta > 0$, which are completely contained in Λ_k . The first one relies solely on the arm events (see Definition 5 in [9]), while the other is simply the union of ϵ -boxes which intersect the cluster. It was showed in the critical case that when the mesh size is small, with high probability these two approximations coincide; see Propositions 1 and 2 in Section 5 of [9]). The results in Section 5 of [9] just use estimates on critical arm events given by Lemmas 4 and 5 in that paper. Lemma 3.5 is the corresponding lemma needed for our near-critical case. Then we can obtain the ‘‘clusters-approximations result’’ (i.e., analog of Propositions 1 and 2 in [9]) in the near-critical case by using Lemma 3.5 and the proof of Propositions 1 and 2 in [9].

Step 2. Following the proof of Theorem 11 given in Section 6 of [9], we can derive our Theorem 3.1 by using the clusters-approximations result in Step 1, Theorem 2.1 and Lemma 3.4. \square

Proof of Theorem 3.2. The proof of the existence and uniqueness of the measure \mathbb{P} is analogous to the proofs of Theorem 3 in [9] and Theorem 6 in [10]. Let $1 \leq k \leq k_1 < k_2$. First, we claim that the marginal distributions $\mathbb{P}_{k_1}|_{\Lambda_k}$ and $\mathbb{P}_{k_2}|_{\Lambda_k}$ are the same. For this, it suffices to show that under the coupling \mathbf{P} , we have $\text{dist}(\mathcal{C}_{k_2}|_{\Lambda_{k_1}}, \mathcal{C}_{k_1}) = 0$ with probability 1 for any integers $1 \leq k_1 < k_2$. By Theorem 3.1, for fixed $k \in \mathbb{N}$, $\text{dist}(\mathcal{C}_k^p, \mathcal{C}_k) \rightarrow 0$ in \mathbf{P} -probability as $p \uparrow p_c$. This combined with the fact (due to the half-plane 3-arm event having exponent larger than 1; see (12)) that for any fixed $\delta > 0$ and $k \in \mathbb{N}$, $\mathbf{P}[\mathcal{C}_{k+\epsilon}^p(\delta) \setminus \mathcal{C}_k^p(\delta) \neq \emptyset]$ tends to 0 uniformly for all $p \in (p(1), p_c)$ as $\epsilon \rightarrow 0$, implies our claim. Hence, the consistency relations needed to apply Kolmogorov's extension theorem (see, e.g., [14]) are satisfied. Since Ω_k and Ω_∞ are complete separable metric spaces, the measurable spaces $(\Omega_k, \mathcal{B}_k)$ and $(\Omega_\infty, \mathcal{B}_\infty)$ are standard Borel spaces, and \mathbb{P}_k is a probability measure on $(\Omega_k, \mathcal{B}_k)$, we can apply Kolmogorov's extension theorem and conclude that there is a unique probability measure \mathbb{P} on $(\Omega_\infty, \mathcal{B}_\infty)$ such that $\mathbb{P}|_{\Lambda_k} = \mathbb{P}_k$ for each $k \in \mathbb{N}$ and \mathbb{P} is supported on collections of bounded, closed and connected subsets of $\hat{\mathbb{C}}$.

Write $\mathcal{C} := \bigcup_k \mathcal{C}_k$. Then the above argument gives that the distribution of \mathcal{C} is \mathbb{P} . Using (8), it is easy to see that for each $\epsilon > 0$, there exists $K = K(\epsilon) \geq 1$ such that for all $k \geq K$ and $p \in (p(1), p_c)$,

$$\mathbf{P}[\text{Dist}(\mathcal{C}_k^p, \mathcal{C}^p) \leq \epsilon] \geq 1 - \epsilon. \quad (29)$$

Theorem 3.1 and (29) imply that $\text{Dist}(\mathcal{C}^p, \mathcal{C}) \rightarrow 0$ in probability as $p \uparrow p_c$. Moreover, the fact that $\mathcal{C}(\delta)|_{\Lambda_k} = \mathcal{C}_k(\delta)$ a.s. and $\text{dist}(\mathcal{C}|_{\Lambda_k}, \mathcal{C}_k) = 0$ a.s. (by the above argument) together with Theorem 3.1 implies the following statements immediately: \mathcal{C} is a measurable function of ω^∞ ; for each $k \in \mathbb{N}$ and $\delta > 0$, $\mathcal{C}(\delta)|_{\Lambda_k}$ is a.s. a finite set, and $\widehat{\text{dist}}(\mathcal{C}^p(\delta)|_{\Lambda_k}, \mathcal{C}(\delta)|_{\Lambda_k}), \text{dist}(\mathcal{C}^p|_{\Lambda_k}, \mathcal{C}|_{\Lambda_k}) \rightarrow 0$ in probability as $p \uparrow p_c$.

It follows easily from the above statement that for any $k, n \in \mathbb{N}$, $\text{dist}(\mathcal{C}^p|_{\Lambda_k(n)}, \mathcal{C}|_{\Lambda_k(n)}) \rightarrow 0$ in probability as $p \uparrow p_c$. It is clear that for any $k \in \mathbb{N}$, $n \geq 2k + 1$ and $p \in (p(1), p_c)$, the configurations $\mathcal{C}^p|_{\Lambda_k}$ and $\mathcal{C}^p|_{\Lambda_k(n)}$ are independent, so $\mathcal{C}|_{\Lambda_k}$ and $\mathcal{C}|_{\Lambda_k(n)}$ are independent.

We give the proof of the rotational invariance of \mathbb{P} below; similar proof works for the translation invariance, and we omit it. For $\theta \in [0, 2\pi]$, let $f(z) = e^{i\theta}z$ be a rotation of \mathbb{C} . By Theorem 2.1, $f(\omega^\infty)$ and ω^∞ have the same distribution. Therefore, similarly as the proof of Theorem 3.1, we can use a coupling such that $\omega_p^\eta \rightarrow f(\omega^\infty)$ in $(\mathcal{H}, d_{\mathcal{H}})$ a.s., to show that $\mathcal{C}_{f(\Lambda_k)}^p$ converges in distribution to $f(\mathcal{C}_k)$ with respect to the metric Dist (note that $f(\mathcal{C}_k)$ is constructed by using $f(\omega^\infty)$ and rotated boxes $f(\Lambda_{\epsilon/2}(z))$). We deduce from this, by letting $k \rightarrow \infty$, that \mathcal{C}^p converges in distribution to $f(\mathcal{C})$ with respect to Dist , as $p \uparrow p_c$. Since we have proved that \mathcal{C}^p also converges in distribution to \mathcal{C} with respect to Dist as $p \uparrow p_c$, it follows that $f(\mathcal{C})$ and \mathcal{C} have the same distribution. \square

Remark 3.6. Consider portions of clusters in a region, that is, the connected components in the region which come from the clusters intersecting the region but not completely contained in it. In Section 3.3 of the preprint version [41] of the present paper, we constructed the scaling limit of the collection of portions of clusters in a strip as $p \uparrow p_c$, which was used to prove Theorem 1.1 in that preprint.

For critical site percolation, the scaling limits in the loop-ensemble and quad-crossing spaces are equivalent in the sense that the associated σ -algebras are the same; see Section 2.3 of [15] for a proof that the loops determine the quad-crossing information, and see Theorem 6.10 in [22] for the converse result. To our knowledge, similar equivalence result has not been proved for near-critical percolation. Moreover, this equivalence result for critical percolation implies that the scaling limit of the collection of portions of clusters in Theorem 12 of [9] is not only a measurable function of the pair of quad-crossing and loop-ensemble scaling limits, but also measurable with respect to the single quad-crossing scaling limit.

4. FPP based on the scaling limit of near-critical percolation

In this section, we define a ‘‘continuum’’ FPP on the continuum cluster ensemble \mathcal{C} . We also define its discrete analog for \mathcal{C}^p on the lattice $\eta\mathbb{T}$. Then we show that as $p \uparrow p_c$, some quantities of the discrete FPP approach their corresponding quantities of the continuum FPP. Finally, we give a law of large numbers for the ‘‘point-to-point’’ passage times of the continuum FPP.

4.1. Definition of first-passage times

We first define first-passage times for \mathcal{C} . Then we define analogous quantities for \mathcal{C}^p with $p \in (p(1), p_c)$.

A **(continuum) chain** is a finite sequence $\Gamma = (\mathcal{C}_0, \mathcal{C}_1, \dots, \mathcal{C}_k)$ of distinct clusters in \mathcal{C} , such that $\mathcal{C}_{j-1} \cap \mathcal{C}_j \neq \emptyset$ for all $j \in \{1, 2, \dots, k\}$. Let $|\Gamma|$ denote the length, or, more precisely, the number of clusters in Γ . The **passage time** of Γ is defined by $T(\Gamma) := |\Gamma| - 1$. Similarly to the third item in Theorem 2 of [10] (which was called ‘‘finite chaining’’ in Proposition 2.7 in [15]) for the full-plane CLE_6 , \mathcal{C} also has the finite chaining property: Almost surely (under \mathbf{P}^∞), for any pair of clusters $\mathcal{C}, \mathcal{C}' \in \mathcal{C}$, there is a chain $(\mathcal{C}, \mathcal{C}_1, \dots, \mathcal{C}_k, \mathcal{C}')$ in \mathcal{C} connecting \mathcal{C} and \mathcal{C}' . It is not hard to show this property, but we will not give its proof here since we will not use it in our proofs.

The **first-passage time** between any pair of clusters $\mathcal{C}, \mathcal{C}' \in \mathcal{C}$ is defined by

$$T(\mathcal{C}, \mathcal{C}') := \inf\{T(\Gamma) : \Gamma \text{ is a chain in } \mathcal{C} \text{ that starts at } \mathcal{C} \text{ and ends at } \mathcal{C}'\},$$

where we use the convention that the infimum of the empty set is ∞ . It follows from the finite chaining property that the first-passage times between all pairs of clusters in \mathcal{C} are almost surely finite.

Remark 4.1. If we view each cluster of \mathcal{C} as a single vertex, and define a new edge between any pair of vertices if their corresponding clusters touch each other, then we get a graph called ‘‘(continuum) cluster graph’’. Observe that a chain corresponds to a (self-avoiding) path of the cluster graph, and the first-passage time between two clusters is just the graph distance between their corresponding vertices in the cluster graph. The finite chaining property implies that the cluster graph is almost surely connected.

Lemma 4.3 below states that, for any fixed point $z \in \mathbb{C}$, there exists almost surely a unique cluster whose Euclidean diameter is maximal among the clusters of \mathcal{C} in $\mathbb{D}_{1/2}(z)$, which is called the **largest (continuum) cluster in $\mathbb{D}_{1/2}(z)$** and denoted by $\mathcal{C}(z)$; if there is no such cluster, then let $\mathcal{C}(z) = \emptyset$. Now we can define the point-to-point passage times for \mathcal{C} : For $m, n \in \mathbb{Z}$ and $\theta \in [0, 2\pi]$, let

$$T_{m,n}^\theta = T_{m,n}^\theta(\mathcal{C}) := T(\mathcal{C}(me^{i\theta}), \mathcal{C}(ne^{i\theta})),$$




FIG 4. A chain Γ from $\mathcal{C}(m)$ to $\mathcal{C}(n)$ such that $T_{m,n}^{\theta=0} = T(\Gamma) = 5$.

where we let $T_{m,n}^{\theta} = \infty$ if $\mathcal{C}(me^{i\theta})$ or $\mathcal{C}(ne^{i\theta})$ is empty. See Figure 4. When $\theta = 0$, we write $T_{m,n} := T_{m,n}^0$. We remark that there are other ways to define point-to-point passage times for \mathcal{C} . The key point is that for each $z \in \mathbb{C}$ one should choose an appropriate cluster close to z . Different choice may lead to different technical arguments, but the strategy for the main result is essentially the same.

To study $T_{m,n}^{\theta}$, we need to define its discrete version. Suppose that $z \in \mathbb{C}$ and $p \in (p_c)$. In the following the sites in $\eta\mathbb{T}$ are viewed as hexagons. We let $\mathcal{C}^p(z)$ denote the cluster whose Euclidean diameter is maximal among the clusters of \mathcal{C}^p in $\mathbb{D}_{1/2}(z)$, and call it the **largest (discrete) cluster in $\mathbb{D}_{1/2}(z)$** . When there are more than one candidates for $\mathcal{C}^p(z)$, then we choose one of them in some deterministic way; when there are no clusters of \mathcal{C}^p in $\mathbb{D}_{1/2}(z)$, then we let $\mathcal{C}^p(z)$ denote the η -hexagon containing z (if z is on the boundary of two or three hexagons, then we choose one of them in some deterministic way). In fact, the arguments in Section 4.2 imply that with high probability the above two events do not occur for p close to p_c . For $m, n \in \mathbb{Z}$ and $\theta \in [0, 2\pi]$, let

$$T_{m,n}^{p,\theta} = T_{m,n}^{p,\theta}(\mathcal{C}^p) := T(\mathcal{C}^p(me^{i\theta}), \mathcal{C}^p(ne^{i\theta})).$$

When $\theta = 0$, we simply drop it from the notation, writing $T_{m,n}^p := T_{m,n}^{p,0}$.

Similarly to its continuum version above, a **(discrete) chain** is a finite sequence $(\mathcal{C}_0, \mathcal{C}_1, \dots, \mathcal{C}_k)$ of distinct clusters in \mathcal{C}^p , such that for each $j \in \{1, 2, \dots, k\}$, there is a yellow η -hexagon touching both of \mathcal{C}_{j-1} and \mathcal{C}_j .

4.2. Properties of large clusters

This subsection gives some properties of clusters with large diameters in region. In particular, Lemma 4.3 allows us to define $\mathcal{C}(z)$, and Lemma 4.4 says that with high probability $\mathcal{C}(z)$ is well-approximated by $\mathcal{C}^p(z)$ for p close to p_c , under a coupling. As mentioned in Section 4.1, one could use a different large cluster “close” to z instead of $\mathcal{C}(z)$ used here to study the point-to-point passage times for \mathcal{C} . For example, one may choose the outermost cluster surrounding z in $\mathbb{D}_{1/2}(z)$, as we did in [41], or the cluster having the largest “volume” in $\mathbb{D}_{1/2}(z)$. However, the former choice leads to a lot of technical issues, while the latter involves counting measures associated to continuum clusters (see [9]), which are rather complicated. The present definition of $\mathcal{C}(z)$ is more natural and makes the proofs cleaner.

For a set $S \subset \mathbb{C}$, let $\text{diam}(S)$ denote the Euclidean diameter of S . For convenience, we will just say diameter to mean Euclidean diameter in the rest of the paper. Recall that the L^∞ -diameter $\text{diam}_\infty(\cdot)$ is used in Section 3.

The following proposition will be used in the proofs of the next two lemmas. Roughly speaking, it says that, for p close p_c , with high probability the gaps between the diameters of clusters with large diameters are not too small. Note that Proposition 3.2 in [4] says that with high probability, the gaps between the volumes of critical clusters with large diameters are also not too small; similar result also holds for the near-critical case.

Proposition 4.2. *For all $\delta, \epsilon_1 \in (0, 1)$ there exists $\epsilon_2 = \epsilon_2(\delta, \epsilon_1) > 0$ such that for all $p \in (p(5/\epsilon_2), p_c)$,*

$$\mathbf{P}_p^\eta[\text{there exist distinct } \mathcal{C}, \mathcal{C}' \in \mathcal{C}^p(\delta)_{\Lambda_1} \text{ such that } |\text{diam}(\mathcal{C}) - \text{diam}(\mathcal{C}')| \leq \epsilon_2] \leq \epsilon_1.$$

Proof. The proof is based on upper bounds for arm probabilities and a counting argument. Let $\mathcal{A}_\sigma^{\theta,\vartheta}(z; r, R)$ denote the wedge arm event for ω_p^η defined at the beginning of Section 2. Write $\mathcal{A}_{(0),(101)}^{\theta,\vartheta}(z; r, R) := \mathcal{A}_{(0)}^{\theta,\vartheta}(z; r, R) \circ \mathcal{A}_{(101)}^{\theta,\vartheta}(z; r, R)$,




FIG 5. Illustration of arm events when $|u - u'| \in [10\epsilon, \delta/5]$ and $|v - v'| > \delta/5$. The clusters $\mathcal{C}, \mathcal{C}'$ and their site boundaries provide five independent arm events $\mathcal{A}_{(101)}^{\arg(v-u), \vartheta_0}(\tilde{u}; 2\epsilon, |u - u'|/4)$, $\mathcal{A}_{(101)}^{\arg(v'-u'), \vartheta_0}(\tilde{u}'; 2\epsilon, |u - u'|/4)$, $\mathcal{A}_{(0), (101)}^{\arg(v-u), \vartheta_0}(\tilde{u}; 3|u - u'|/2, 2\delta/5)$, $\mathcal{A}_{(101)}^{\arg(u-v), \vartheta_0}(\tilde{v}; 2\epsilon, \delta/20)$ and $\mathcal{A}_{(101)}^{\arg(u'-v'), \vartheta_0}(\tilde{v}'; 2\epsilon, \delta/20)$.

where \circ means the disjoint occurrence (see e.g. Chapter 2 of [19]). By (5) and (13), we know that there exist constants $C_0 > 0$, $\vartheta_0 \in (\pi, 3\pi/2)$ and $0 < 3\beta < \lambda_1 < 1/2$, such that for all $p \in (p(1), p_c)$, $\eta \leq r < R \leq 1$ and $\theta \in [0, 2\pi]$,

$$\mathbf{P}_p^\eta[\mathcal{A}_{(0)}(r, R)] \leq C_0 \left(\frac{r}{R}\right)^{\lambda_1} \quad \text{and} \quad \mathbf{P}_p^\eta[\mathcal{A}_{(101)}^{\theta, \vartheta_0}(r, R)] \leq C_0 \left(\frac{r}{R}\right)^{2-\beta}. \quad (30)$$

Letting $\lambda := \lambda_1 - \beta$, the above two inequalities combined with Reimer's inequality implies that

$$\mathbf{P}_p^\eta[\mathcal{A}_{(0), (101)}^{\theta, \vartheta_0}(z; r, R)] \leq C_0^2 \left(\frac{r}{R}\right)^{2+\lambda}. \quad (31)$$

Let $\delta \in (0, 1)$ and $\epsilon \in (0, \delta/50)$. Given a closed subset S of Λ_1 with $\text{diam}(S) \geq \delta$, we choose a pair $\{u(S), v(S)\}$ of vertices in $\epsilon\mathbb{Z}^2$ such that $\Lambda_{\epsilon/2}(u(S))$ and $\Lambda_{\epsilon/2}(v(S))$ contain two points of S whose distance is equal to the diameter of S . If there are more than one candidates for this pair, we choose one of them in some deterministic way. Write $\tilde{u}(u, v) := u + \epsilon \cdot (u - v)/|u - v|$ and $\tilde{v}(u, v) := v + \epsilon \cdot (v - u)/|v - u|$. It is easy to check that there exists $\epsilon_0 \in (0, \delta/50)$, such that for all $\epsilon \in (0, \epsilon_0)$ and all closed subset S of Λ_1 with $\text{diam}(S) \geq \delta$, the set $S + \mathbb{D}_{\epsilon/5}$ is contained in the intersection of the wedges $W(\tilde{u}; \arg(v - u), \vartheta_0)$ and $W(\tilde{v}; \arg(u - v), \vartheta_0)$.

In the rest of the proof, we suppose that $\delta \in (0, 1)$, $\epsilon \in (0, \epsilon_0)$ and $p \in (p(5/\epsilon), p_c)$. Assume that $\mathcal{C}, \mathcal{C}'$ are distinct clusters of $\mathcal{C}^p(\delta)$ in Λ_1 with $|\text{diam}(\mathcal{C}) - \text{diam}(\mathcal{C}')| \leq \epsilon$. For simplicity, we let u, u' denote $u(\mathcal{C}), u(\mathcal{C}')$, respectively, and let $\tilde{u} := \tilde{u}(u, v)$, $\tilde{u}' := \tilde{u}(u', v')$. We assume without loss of generality that $|u - u'| \leq \min\{|u - v'|, |v - v'|, |v - u'|\}$. By considering the distance between u and u' , we distinguish the following three cases; in each case the clusters $\mathcal{C}, \mathcal{C}'$ together with their yellow site boundaries produces the corresponding independent arm events. See Figure 5 for an illustration.

- If $|u - u'| \in [10\epsilon, \delta/5]$, then the events $\mathcal{A}_{(101)}^{\arg(v-u), \vartheta_0}(\tilde{u}; 2\epsilon, |u - u'|/4)$, $\mathcal{A}_{(101)}^{\arg(v'-u'), \vartheta_0}(\tilde{u}'; 2\epsilon, |u - u'|/4)$ and $\mathcal{A}_{(0), (101)}^{\arg(v-u), \vartheta_0}(\tilde{u}; 3|u - u'|/2, 2\delta/5)$ occur.
- If $|u - u'| > \delta/5$, then $\mathcal{A}_{(101)}^{\arg(v-u), \vartheta_0}(\tilde{u}; 2\epsilon, \delta/20)$ and $\mathcal{A}_{(101)}^{\arg(v'-u'), \vartheta_0}(\tilde{u}'; 2\epsilon, \delta/20)$ occur.
- If $|u - u'| \in [0, 10\epsilon)$, then $\mathcal{A}_{(0), (101)}^{\arg(v-u), \vartheta_0}(\tilde{u}; 12\epsilon, 2\delta/5)$ occurs.

Similar statements also hold for v and v' . It is easy to check that the arm events introduced in each case are independent; see, for example, the five arm events shown in Figure 5 are independent.

Note that $\text{diam}(\mathcal{C}) \in [\delta, 2\sqrt{2}]$. Now, assume that $\text{diam}(\mathcal{C}) \in [\delta_1, \delta_1 + \epsilon]$ for any fixed $\delta_1 \in [\delta, 2\sqrt{2}]$. Then, there are $O(\epsilon^{-3})$ choices for the pair $\{u, v\}$. Fix u and v . Then there are $O(1)$ choices for u' satisfying $|u - u'| \in [0, 10\epsilon)$,

$O(\epsilon^{-2})$ choices for u' satisfying $|u - u'| > \delta/5$, and $O(4^j)$ choices for u' satisfying $|u - u'| \in [10\epsilon, \delta/5]$ and $u' \in A(u; 2^j\epsilon, 2^{j+1}\epsilon)$ with $j \geq 3$. Now fix u' also. Then there are $O(1)$ choices for v' satisfying $|v - v'| \in [0, 10\epsilon]$, $O(\epsilon^{-1})$ choices for v' satisfying $|v - v'| > \delta/5$, and $O(2^k)$ choices for v' satisfying $|v - v'| \in [10\epsilon, \delta/5]$ and $v' \in A(v; 2^k\epsilon, 2^{k+1}\epsilon)$ with $k \geq 3$. The above arguments imply that there exist $C_1, C_2 > 0$ such that for all $\delta \in (0, 1)$, $\delta_1 \in [\delta, 2\sqrt{2}]$, $\epsilon \in (0, \epsilon_0)$ and $p \in (p(5/\epsilon), p_c)$,

$$\begin{aligned} & \mathbf{P}_p^\eta[\text{there exist distinct } \mathcal{C}, \mathcal{C}' \in \mathcal{C}^p(\delta)|_{\Lambda_1} \text{ such that } \text{diam}(\mathcal{C}) \in [\delta_1, \delta_1 + \epsilon] \text{ and } |\text{diam}(\mathcal{C}) - \text{diam}(\mathcal{C}')| \leq \epsilon] \\ & \leq \frac{C_1}{\epsilon^3} \left(\underbrace{\left(\left(\frac{\epsilon}{\delta} \right)^{2+\lambda} \right)}_{(1)} + \underbrace{\sum_{j=3}^{\lceil \log_2(\delta/\epsilon) \rceil} 4^j \left(\frac{\epsilon}{2^j\epsilon} \right)^{4-2\beta} \left(\frac{2^j\epsilon}{\delta} \right)^{2+\lambda}}_{(2)} \right) \left(\underbrace{\frac{1}{\epsilon} \left(\frac{\epsilon}{\delta} \right)^{4-2\beta}}_{(3)} + \underbrace{\sum_{k=3}^{\lceil \log_2(\delta/\epsilon) \rceil} 2^k \left(\frac{\epsilon}{2^k\epsilon} \right)^{4-2\beta} \left(\frac{2^k\epsilon}{\delta} \right)^{2+\lambda}}_{(4)} \right) \\ & \quad + \underbrace{\left(\frac{\epsilon}{\delta} \right)^{4+2\lambda}}_{(5)} + \underbrace{\frac{1}{\epsilon^3} \left(\frac{\epsilon}{\delta} \right)^{8-4\beta}}_{(6)} \quad \text{by (30) and (31)} \\ & \leq C_2 \epsilon^{1+\lambda-2\beta} / \delta^8, \end{aligned} \tag{32}$$

where the six terms (1)–(6) correspond respectively to the following six cases: (1) $|u - u'| \in [0, 10\epsilon)$, (2) $|u - u'| \in [10\epsilon, \delta/5]$, (3) $|v - v'| > \delta/5$, (4) $|v - v'| \in [10\epsilon, \delta/5]$, (5) $|u - u'|, |v - v'| \in [0, 10\epsilon)$, (6) $|u - u'|, |v - v'| > \delta/5$. Hence,

$$\begin{aligned} & \mathbf{P}_p^\eta[\text{there exist distinct } \mathcal{C}, \mathcal{C}' \in \mathcal{C}^p(\delta)|_{\Lambda_1} \text{ such that } |\text{diam}(\mathcal{C}) - \text{diam}(\mathcal{C}')| \leq \epsilon] \\ & \leq \sum_{j=0}^{\lfloor (2\sqrt{2}-\delta)/\epsilon \rfloor} \mathbf{P}_p^\eta[\exists \text{ distinct } \mathcal{C}, \mathcal{C}' \in \mathcal{C}^p(\delta)|_{\Lambda_1} \text{ s.t. } \text{diam}(\mathcal{C}) \in [\delta + j\epsilon, \delta + (j+1)\epsilon] \text{ and } |\text{diam}(\mathcal{C}) - \text{diam}(\mathcal{C}')| \leq \epsilon] \\ & \leq 4C_2 \epsilon^{\lambda-2\beta} / \delta^8 \quad \text{by (32)}. \end{aligned}$$

The lemma follows easily from this since $\lambda > 2\beta$. \square

Lemma 4.3. *Let $z \in \mathbb{C}$. Almost surely, there is a unique cluster whose diameter is maximal among the clusters of \mathcal{C} in $\mathbb{D}_{1/2}(z)$, which is called the **largest (continuum) cluster in $\mathbb{D}_{1/2}(z)$** and denoted by $\mathcal{C}(z)$. Moreover, for each $\epsilon > 0$, there exists $\delta = \delta(\epsilon) > 0$ such that*

$$\mathbf{P}^\infty[\text{diam}(\mathcal{C}(z)) \geq \delta] \geq 1 - \epsilon.$$

Proof. We shall prove the lemma when $z = 0$; the general case then follows since the law of \mathcal{C} is invariant under translations by Theorem 3.2. Let \mathbf{P} be a coupling such that $\omega_p^\eta \rightarrow \omega^\infty$ in $(\mathcal{H}, d_{\mathcal{H}})$ a.s. as $p \uparrow p_c$. Fix any $\delta \in (0, 1)$. For each $\epsilon_1 \in (0, 1)$, there exist $\epsilon_2 > 0, p_1 \in (0, p_c)$ such that

$$\begin{aligned} & \mathbf{P}[\text{there exist distinct } \mathcal{C}, \mathcal{C}' \in \mathcal{C}(\delta)|_{\Lambda_1} \text{ such that } |\text{diam}(\mathcal{C}) - \text{diam}(\mathcal{C}')| \leq \epsilon_2] \\ & \leq \mathbf{P}[\exists \text{ distinct } \mathcal{C}, \mathcal{C}' \in \mathcal{C}^{p_1}(\delta)|_{\Lambda_1} \text{ s.t. } |\text{diam}(\mathcal{C}) - \text{diam}(\mathcal{C}')| \leq 3\epsilon_2] + \mathbf{P}[\widehat{\text{dist}}(\mathcal{C}(\delta)|_{\Lambda_1}, \mathcal{C}^{p_1}(\delta)|_{\Lambda_1}) \geq \epsilon_2/4] \\ & \leq \epsilon_1 \quad \text{by Proposition 4.2 and Theorem 3.2.} \end{aligned} \tag{33}$$

Letting $\epsilon_1 \rightarrow 0$ gives that

$$\mathbf{P}[\text{there exist distinct } \mathcal{C}, \mathcal{C}' \in \mathcal{C}(\delta)|_{\Lambda_1} \text{ such that } |\text{diam}(\mathcal{C}) - \text{diam}(\mathcal{C}')| = 0] = 0. \tag{34}$$

Therefore, for each $\epsilon > 0$ there exist $\delta \in (0, 1/8)$ and $p_2 \in (0, p_c)$ such that

$$\begin{aligned} & \mathbf{P}[\mathcal{C}(\delta)|_{\mathbb{D}_{1/2}} \neq \emptyset \text{ and } \exists \text{ a unique cluster whose diameter is maximal among the clusters of } \mathcal{C}(\delta)|_{\mathbb{D}_{1/2}}] \\ & = \mathbf{P}[\mathcal{C}(\delta)|_{\mathbb{D}_{1/2}} \neq \emptyset] \quad \text{by (34)} \\ & \geq \mathbf{P}[\mathcal{C}^{p_2}(2\delta)|_{\mathbb{D}_{1/4}} \neq \emptyset] - \mathbf{P}[\text{dist}(\mathcal{C}|_{\Lambda_1}, \mathcal{C}^{p_2}|_{\Lambda_1}) \geq \delta/2] \\ & \geq 1 - \epsilon \quad \text{by RSW, FKG and Theorem 3.2.} \end{aligned}$$

The lemma follows from this immediately. \square

Lemma 4.4. *Let \mathbf{P} be a coupling such that $\omega_p^\eta \rightarrow \omega^\infty$ in $(\mathcal{H}, d_{\mathcal{H}})$ a.s. as $p \uparrow p_c$. For each $k \in \mathbb{N}$ and $\epsilon, \delta > 0$, there exists $p_0 \in (0, p_c)$, such that for all $p \in (p_0, p_c)$ and $z \in \Lambda_k$,*

$$\mathbf{P}[d_H(\mathcal{C}^p(z), \mathcal{C}(z)) \geq \delta] \leq \epsilon.$$

Proof. Let $0 < \delta_1 < \delta/8 < 1/10$, $p \in (p(5/\delta_1), p_c)$, $k \in \mathbb{N}$ and $z \in \Lambda_k$. Define the events

$$\mathcal{E}_1 = \mathcal{E}_1(\delta_1, p, k) := \{\text{dist}(\mathcal{C}|_{\Lambda_{k+1}}, \mathcal{C}^p|_{\Lambda_{k+1}}) < \delta_1\},$$

$$\mathcal{E}_2 = \mathcal{E}_2(z; \delta, \delta_1, p) := \{\text{diam}(\mathcal{C}^p(z)) \geq \delta \text{ and } \mathcal{C}^p(\delta/2)|_{\mathbb{D}_{1/2+2\delta_1}(z)} = \mathcal{C}^p(\delta/2)|_{\mathbb{D}_{1/2-2\delta_1}(z)}\},$$

$$\mathcal{E}_3 = \mathcal{E}_3(z; \delta, \delta_1) := \{\text{for any distinct } \mathcal{C}, \mathcal{C}' \in \mathcal{C}(\delta/2)|_{\Lambda_1(z)}, |\text{diam}(\mathcal{C}) - \text{diam}(\mathcal{C}')| \geq 9\delta_1\}.$$

First, we prove that $\mathcal{E}_1 \cap \mathcal{E}_2 \cap \mathcal{E}_3 \subset \{d_H(\mathcal{C}^p(z), \mathcal{C}(z)) < \delta_1\}$. Assume that the event $\mathcal{E}_1 \cap \mathcal{E}_2 \cap \mathcal{E}_3$ holds. Since $\mathcal{E}_1 \cap \mathcal{E}_2$ occurs, there is a continuum cluster \mathcal{C}_0 such that $d_H(\mathcal{C}^p(z), \mathcal{C}_0) < \delta_1$, and furthermore $\mathcal{C}^p(z) \subset \mathbb{D}_{1/2-2\delta_1}(z)$, $\mathcal{C}_0 \subset \mathbb{D}_{1/2}(z)$ and $\text{diam}(\mathcal{C}_0) > \text{diam}(\mathcal{C}^p(z)) - 4\delta_1 > \delta/2$. It remains to check that $\mathcal{C}_0 = \mathcal{C}(z)$. Assume for a contradiction that $\mathcal{C}_0 \neq \mathcal{C}(z)$. Since \mathcal{E}_3 occurs, we have $\text{diam}(\mathcal{C}(z)) \geq \text{diam}(\mathcal{C}_0) + 9\delta_1 > \text{diam}(\mathcal{C}^p(z)) + 5\delta_1$. Since \mathcal{E}_1 holds, there is a cluster \mathcal{C}_0^p of \mathcal{C}^p such that $d_H(\mathcal{C}_0^p, \mathcal{C}(z)) < \delta_1$, $\mathcal{C}_0^p \subset \mathbb{D}_{1/2+2\delta_1}(z)$ and $\text{diam}(\mathcal{C}_0^p) > \text{diam}(\mathcal{C}(z)) - 4\delta_1 > \text{diam}(\mathcal{C}^p(z)) + \delta_1$. Furthermore, the fact that \mathcal{E}_2 holds implies that $\mathcal{C}_0^p \subset \mathbb{D}_{1/2}(z)$. This together with $\mathcal{C}_0^p > \text{diam}(\mathcal{C}^p(z))$ contradicts the definition of $\mathcal{C}^p(z)$.

Therefore, for all $k \in \mathbb{N}$, $\epsilon > 0$ and $\delta \in (0, 1/2)$, there exist $\delta_1 \in (0, \delta/8)$ and $p_0 \in (0, p_c)$ such that for all $p \in (p_0, p_c)$ and $z \in \Lambda_k$,

$$\begin{aligned} \mathbf{P}[d_H(\mathcal{C}^p(z), \mathcal{C}(z)) < \delta] &\geq \mathbf{P}[d_H(\mathcal{C}^p(z), \mathcal{C}(z)) < \delta_1] \\ &\geq 1 - \mathbf{P}[\neg \mathcal{E}_1] - \mathbf{P}[\neg \mathcal{E}_2] - \mathbf{P}[\neg \mathcal{E}_3] \quad \text{by } \mathcal{E}_1 \cap \mathcal{E}_2 \cap \mathcal{E}_3 \subset \{d_H(\mathcal{C}^p(z), \mathcal{C}(z)) < \delta_1\} \\ &\geq 1 - \epsilon/2 - \mathbf{P}[\neg \mathcal{E}_2] \quad \text{by Theorem 3.2 and (33)} \\ &\geq 1 - \epsilon \quad \text{by RSW, FKG and (12), via a standard argument.} \end{aligned}$$

This completes the proof of the lemma. \square

4.3. Properties of first-passage times

Lemma 4.5. *There are constants $C_1, C_2 > 0$, such that for each $n \in \mathbb{Z}$, $k \in \mathbb{N}$, $\theta \in [0, 2\pi]$ and $p \in (p(10), p_c)$,*

$$\mathbf{P}_p^\eta[T_{n, n+1}^{p, \theta} \geq k] \leq C_1 \exp(-C_2 k). \quad (35)$$

Moreover, there is $C_0 > 0$ such that for each $\theta \in [0, 2\pi]$, $p \in (p(10), p_c)$ and $m, n \in \mathbb{Z}$ with $m < n$,

$$\mathbf{E}_p^\eta[T_{m, n}^{p, \theta}] \leq C_0(n - m). \quad (36)$$

Proof. For simplicity, we prove Lemma 4.5 in the case $\theta = 0$. The proof extends immediately to the general case.

Suppose that $p \in (p(10), p_c)$. Assume that $\mathcal{C}^p(n)$ and $\mathcal{C}^p(n+1)$ intersect boxes $\Lambda_r(u)$ and $\Lambda_r(v)$ for some $u, v \in 2r\mathbb{Z}^2$, respectively. If $r \in [\eta, 1/6]$ and $\text{diam}(\mathcal{C}^p(n)), \text{diam}(\mathcal{C}^p(n+1)) \geq 6r$, then $\mathcal{C}^p(n)$ and $\mathcal{C}^p(n+1)$ produce 1-arm events $\mathcal{A}_{(0)}(u; r, 3r)$ and $\mathcal{A}_{(0)}(v; r, 3r)$, respectively, and furthermore,

$$T_{n, n+1}^p \leq X^p(u; r, 4) + X^p(v; r, 4) + Y^p(n, u, v; r), \quad (37)$$

where $X^p(z; r_1, r_2)$ is the annulus-crossing time defined above Lemma 2.7, $Y^p(n, u, v; r) := Y^p(u; r) + Y^p(v; r) + Y^p(n)$ and

$$Y^p(z; r) := \inf\{T(\gamma)(\omega_p^\eta) : \gamma \text{ is a circuit surrounding } z \text{ in } A(z; r, 3r)\},$$

$$Y^p(n) := \inf\{T(\gamma)(\omega_p^\eta) : \gamma \text{ is a circuit surrounding } n + 1/2 \text{ in } A(n + 1/2; 1, 2)\}.$$

When $r = \eta$, it is easy to check that the following inequality always holds, although it could happen that $\text{diam}(\mathcal{C}^p(n)) < 6\eta$, or $\mathcal{C}^p(n)$ is even a yellow η -hexagon.

$$T_{n, n+1}^p \leq X^p(u; \eta, 4) + X^p(v; \eta, 4) + Y^p(n, u, v; \eta) + 6. \quad (38)$$

It follows easily from Lemma 2.6 that there exist $C_3, C_4 > 0$, such that for all $p \in (p(10), p_c)$, $n \in \mathbb{Z}$, $k \in \mathbb{N}$, $r \in [\eta, 1/6]$ and $u, v \in 2r\mathbb{Z}^2$,

$$\mathbf{P}_p^\eta[Y^p(n, u, v; r) \geq k] \leq C_3 \exp(-C_4 k). \quad (39)$$

For $x \geq 1$ and $n \in \mathbb{Z}$, define a set of pairs of points by

$$S(n; x) := \{(u, v) : u, v \in 2^{-x+1}\mathbb{Z}^2, \Lambda_{2^{-x}}(u) \text{ intersects } \mathbb{D}_{1/2}(n) \text{ and } \Lambda_{2^{-x}}(v) \text{ intersects } \mathbb{D}_{1/2}(n+1)\}.$$

It is clear that there is an absolute constant $C_5 > 0$ such that

$$\#S(n; x) \leq \exp(C_5 x). \quad (40)$$

Let K, K_2 be the constants from Lemma 2.7. Write

$$C_6 := \min\{C_4/3, K_2/3\} \quad \text{and} \quad C := \min\{1/(4K), C_6/(2C_5)\}.$$

Then, there is $C_7 > 0$ such that for all $p \in (p(10), p_c)$, $n \in \mathbb{Z}$, $k \in \mathbb{N}$, $u, v \in 2^{-Ck+1}\mathbb{Z}^2$ and $r^* = r^*(p, k) := \max\{2^{-Ck}, \eta\}$,

$$\begin{aligned} & \mathbf{P}_p^\eta[X^p(u; r^*, 4) + X^p(v; r^*, 4) + Y^p(n, u, v; r^*) + 6 \geq k] \\ & \leq \mathbf{P}_p^\eta[X^p(u; r^*, 4) \geq k/3 - 2] + \mathbf{P}_p^\eta[X^p(v; r^*, 4) \geq k/3 - 2] + \mathbf{P}_p^\eta[Y^p(n, u, v; r^*) \geq k/3 - 2] \\ & \leq C_7 \exp(-C_6 k) \quad \text{by Lemma 2.7 and (39)}. \end{aligned} \quad (41)$$

By a standard RSW and FKG argument, there exist $C_8, C_9 > 0$, such that for all $n \in \mathbb{Z}$, $p \in (p(10), p_c)$, $x \geq 1$ with $2^{-x} \geq \eta$,

$$\mathbf{P}_p^\eta[\text{diam}(\mathcal{C}^p(n)) < 6 \cdot 2^{-x} \text{ or } \text{diam}(\mathcal{C}^p(n+1)) < 6 \cdot 2^{-x}] \leq C_8 \exp(-C_9 x). \quad (42)$$

The above arguments imply that, for k satisfying $k \geq 3/C$ and $2^{-Ck} \geq \eta$,

$$\begin{aligned} & \mathbf{P}_p^\eta[T_{n, n+1}^p \geq k] \\ & \leq \mathbf{P}_p^\eta[\text{diam}(\mathcal{C}^p(n)) < 6 \cdot 2^{-Ck} \text{ or } \text{diam}(\mathcal{C}^p(n+1)) < 6 \cdot 2^{-Ck}] \\ & \quad + \mathbf{P}_p^\eta[\text{diam}(\mathcal{C}^p(n)), \text{diam}(\mathcal{C}^p(n+1)) \geq 6 \cdot 2^{-Ck} \text{ and } T_{n, n+1}^p \geq k] \\ & \leq C_8 \exp(-C_9 Ck) + \sum_{(u, v) \in S(n; Ck)} \mathbf{P}_p^\eta[X^p(u; r^*, 4) + X^p(v; r^*, 4) + Y^p(n, u, v; r^*) \geq k] \quad \text{by (42) and (37)} \\ & \leq C_8 \exp(-C_9 Ck) + \exp(C_5 Ck) \cdot C_7 \exp(-C_6 k) \quad \text{by (40) and (41)}, \end{aligned} \quad (43)$$

and for k satisfying $2^{-Ck} \leq \eta$,

$$\begin{aligned} \mathbf{P}_p^\eta[T_{n, n+1}^p \geq k] & \leq \sum_{(u, v) \in S(n; \log_2(1/\eta))} \mathbf{P}_p^\eta[X^p(u; \eta, 4) + X^p(v; \eta, 4) + Y^p(n, u, v; \eta) + 6 \geq k] \quad \text{by (38)} \\ & \leq \exp(C_5 \log_2(1/\eta)) \cdot C_7 \exp(-C_6 k) \quad \text{by (40) and (41)} \\ & \leq \exp(C_5 Ck) \cdot C_7 \exp(-C_6 k). \end{aligned} \quad (44)$$

Then (35) follows from (43) and (44) immediately since $C_5 C < C_6$. We get (36) from (35) since $T_{m, n}^p \leq \sum_{j=m}^{n-1} T_{j, j+1}^p$. \square

To study geometric properties of geodesics between clusters of \mathcal{C}^p , for $z \in \mathbb{C}$, $0 < 4r \leq R \leq 1$ and $p \in (p(4/r), p_c)$ we define the event

$$\tilde{\mathcal{A}}_6(z; r, R) := \mathcal{A}_6(z; r, R - 2\eta) \cup \mathcal{A}_{(111111)}(z; r + 2\eta, R) \cup \left\{ \begin{array}{l} \text{there exists } x \in [r + 2\eta, R - \eta] \text{ such that} \\ \mathcal{A}_6(z; r, x - \eta) \cap \mathcal{A}_{(111111)}(z; x, R) \text{ occurs} \end{array} \right\}. \quad (45)$$

Roughly speaking, the above event occurs when there is a geodesic γ between two large blue clusters with two yellow sites of γ “very” close to each other. We shall use the event in the proof of Lemma 4.7 below; see Figure 6 for an illustration.




FIG 6. Let $0 < 20\delta < \delta_1 < 1/4$ and $p \in (p(10/\delta), p_c)$. Assume that $\text{diam}(\mathcal{C}^p(0)), \text{diam}(\mathcal{C}^p(n)) \geq 4\delta_1$ and there is a geodesic γ from $\mathcal{C}^p(0)$ to $\mathcal{C}^p(n)$ with two yellow sites of γ contained in $\Lambda_{3\delta}(z)$. Then the event $\tilde{\mathcal{A}}_6(z; 4\delta, \delta_1)$ occurs.

Lemma 4.6. *Let $\lambda_6 > 0$ be the universal constant from (10). There is a constant $C > 0$ such that for all $z \in \mathbb{C}$, $0 < 4r \leq R \leq 1$ and $p \in (p(4/r), p_c)$, we have*

$$\mathbf{P}_p^\eta \left[\tilde{\mathcal{A}}_6(z; r, R) \right] \leq C \left(\frac{r}{R} \right)^{2+\lambda_6}.$$

Proof. Write $J := \lfloor \log_2(R/r) \rfloor$. Assume that there exists $x \in [r+2\eta, R-\eta]$ such that $\mathcal{A}_6(z; r, x-\eta) \cap \mathcal{A}_{(111111)}(z; x, R)$ holds, and x is the smallest one satisfying this event. Observe that if $x \leq 2r$, then $\mathcal{A}_{(111111)}(z; 2r, R)$ occurs; if $x \geq 2^{J-1}r$, then $\mathcal{A}_6(z; r, 2^{J-1}r - \eta)$ occurs; if $2^{j-1}r < x \leq 2^j r$ for $j \in \{2, \dots, J-1\}$, then $\mathcal{A}_6(z; r, 2^{j-1}r - \eta) \cap \mathcal{A}_{(111111)}(z; 2^j r, R)$ occurs. Therefore, there are constants $\epsilon, C, C_1 > 0$ such that for all $z \in \mathbb{C}$, $0 < 4r \leq R \leq 1$ and $p \in (p(4/r), p_c)$,

$$\begin{aligned} & \mathbf{P}_p^\eta \left[\tilde{\mathcal{A}}_6(z; r, R) \right] \\ & \leq \mathbf{P}_p^\eta \left[\mathcal{A}_{(111111)}(z; 2r, R) \right] + \mathbf{P}_p^\eta \left[\mathcal{A}_6(z; r, 2^{J-1}r - \eta) \right] + \sum_{j=2}^{J-1} \mathbf{P}_p^\eta \left[\mathcal{A}_6(z; r, 2^{j-1}r - \eta) \right] \mathbf{P}_p^\eta \left[\mathcal{A}_{(111111)}(z; 2^j r, R) \right] \\ & \leq C_1 \left(\frac{r}{R} \right)^{2+\lambda_6} + \sum_{j=2}^{J-1} C_1 \left(\frac{r}{R} \right)^{2+\lambda_6} \left(\frac{2^j r}{R} \right)^\epsilon \quad \text{by (10) and (14)} \\ & \leq C \left(\frac{r}{R} \right)^{2+\lambda_6}. \end{aligned}$$

□

For $n \in \mathbb{N}$, $p \in (p(1), p_c)$, $\delta \in (0, 1/2)$, $\theta \in [0, 2\pi]$ and $K \geq 2$, we define the event

$$\mathcal{F}_n^{p,\theta}(\delta, K) := \left\{ \begin{array}{l} \exists \text{ a chain } \Gamma \text{ from } \mathcal{C}^p(0) \text{ to } \mathcal{C}^p(ne^{i\theta}) \text{ such that } T_{0,n}^{p,\theta} = |\Gamma| - 1, \\ \text{diam}(\mathcal{C}^p) \geq \delta \text{ for all } \mathcal{C}^p \in \Gamma, \text{ and } \Gamma \subset \Lambda_{Kn} \end{array} \right\}.$$

Write $\mathcal{F}_n^p(\delta, K) := \mathcal{F}_n^{p,0}(\delta, K)$. Observe that for our FPP on $\eta\mathbb{T}$, if there is a geodesic between two large blue clusters with two yellow sites in this geodesic close to each other, then the blue cluster between these two yellow sites may be small and not be seen in the scaling limit. This creates a difference between the first-passage times in the discrete and continuum cases, since the continuum FPP is defined by chains of clusters. Nevertheless, thanks to the following lemma, for p close to p_c with high probability the first-passage time between two large blue clusters is realized by a discrete chain of large clusters. The proof relies on the estimate of the “6-arm” event in the previous lemma.

Lemma 4.7. *For each $R \geq 1$ and $\epsilon > 0$, there exist $K = K(\epsilon) \geq 3$ (independent of R), $\delta = \delta(\epsilon, R) \in (0, 1/4)$, such that for all $\theta \in [0, 2\pi]$, $p \in (p(10/\delta), p_c)$ and $n \in \mathbb{N}$ with $n \leq R$,*

$$\mathbf{P}_p^\eta[\mathcal{F}_n^{p,\theta}(\delta, K)] \geq 1 - \epsilon.$$

Proof. For simplicity, we prove Lemma 4.7 in the case $\theta = 0$. The proof extends immediately to the general case.

Let C_0 be the constant from (36). We use (36) and Markov's inequality to obtain that, for all $x > 0$, $n \in \mathbb{N}$ and $p \in (p(10), p_c)$,

$$\mathbf{P}_p^\eta[T_{0,n}^p \geq xC_0n] \leq \frac{1}{x}. \quad (46)$$

We claim that there are constants $\epsilon_0, C_1, C_2 > 0$ such that for each $p \in (p(10), p_c)$, $r \geq 1$ and the event

$$\mathcal{G}(r) := \{\exists \text{ a path } \gamma \text{ of } \omega_p^\eta \text{ starting at a hexagon in } \mathbb{D}_{1/2}, \text{ such that } T(\gamma) \leq \epsilon_0 r \text{ and } \gamma \not\subset \Lambda_r\},$$

we have

$$\mathbf{P}_p^\eta[\mathcal{G}(r)] \leq C_1 \exp(-C_2 r). \quad (47)$$

We use a standard renormalization argument to show (47). First, we define a family of random variables $\{X_z^p : z \in \mathbb{Z}^2\}$ as follows. We declare a vertex $z \in \mathbb{Z}^2$ to be **good** if there is a yellow circuit (viewed as a union of yellow hexagons) of ω_p^η surrounding the point Mz in $A(Mz; M/2, M)$, where M is a large constant that will be fixed later. The family $X^p = \{X_z^p : z \in \mathbb{Z}^2\}$ is defined by

$$X_z^p := \begin{cases} 1 & \text{if } z \text{ is good,} \\ 0 & \text{otherwise.} \end{cases}$$

Note that X^p is 2-dependent. It follows by (8) that

$$\lim_{M \rightarrow \infty} \mathbf{P}_p^\eta[z \text{ is good}] = 1 \quad \text{uniformly in } p \in (p(10), p_c) \text{ and } z \in \mathbb{Z}^2. \quad (48)$$

Then, by Theorem 2.8, there is a $p_0 \in (0, p_c^{\text{site}}(\mathbb{Z}^2))$ and a fixed constant $M \geq 1$, such that for each $p \in (p(10), p_c)$, X^p stochastically dominates the subcritical Bernoulli site FPP on \mathbb{Z}^2 with measure $\mathbf{P}_{\mathbb{Z}^2, p_0}^{\text{site}}$. (Recall that $\mathbf{P}_{\mathbb{Z}^2, p_0}^{\text{site}}[t(z) = 0] = p_0 = 1 - \mathbf{P}_{\mathbb{Z}^2, p_0}^{\text{site}}[t(z) = 1]$ for $z \in \mathbb{Z}^2$.)

Assume that the event $\mathcal{G}(r)$ holds. Then it is easy to see that there exists a path $\tilde{\gamma}$ of \mathbb{Z}^2 starting at the origin, such that $|\tilde{\gamma}| \geq r/M$ and $\Lambda_{M/2}(Mz) \cap \tilde{\gamma} \neq \emptyset$ for all $z \in \tilde{\gamma}$. Note that when a vertex $z \in \tilde{\gamma}$ is good, then γ passes through a yellow hexagon of ω_p^η in $A(Mz; M/2, M)$. Therefore,

$$T(\gamma) \geq (\text{the number of good vertices in } \tilde{\gamma})/9.$$

The above argument implies that there are constants $\epsilon_0, C_1, C_2 > 0$ such that for each $p \in (p(10), p_c)$ and $r \geq 1$,

$$\begin{aligned} \mathbf{P}_p^\eta[\mathcal{G}(r)] &\leq \mathbf{P}_p^\eta \left[\begin{array}{l} \exists \text{ a path } \tilde{\gamma} \text{ of } \mathbb{Z}^2 \text{ starting at the origin, such that } |\tilde{\gamma}| \geq r/M \\ \text{and the number of good vertices in } \tilde{\gamma} \text{ is smaller than } 10\epsilon_0 r \end{array} \right] \\ &\leq \mathbf{P}_{\mathbb{Z}^2, p_0}^{\text{site}} \left[\begin{array}{l} \exists \text{ a path } \tilde{\gamma} \text{ of } \mathbb{Z}^2 \text{ starting at the origin, such that } |\tilde{\gamma}| \geq r/M \\ \text{and the number of 1-valued vertices in } \tilde{\gamma} \text{ is smaller than } 10\epsilon_0 r \end{array} \right] \\ &\leq C_1 \exp(-C_2 r) \quad \text{by Proposition 2.9,} \end{aligned}$$

which gives the claim (47).

Let $\mathcal{O}(r)$ denote the event that there exists a yellow circuit of ω_p^η surrounding the origin in $A(r/2, r)$. Then for each $\epsilon \in (0, 1)$ there is $K \geq 3$ such that for all $n \in \mathbb{N}$, $p \in (p(10), p_c)$ and the event

$$\mathcal{H}_n(K) := \left\{ \begin{array}{l} \text{for each geodesic } \gamma \text{ from } \mathcal{C}^p(0) \text{ to } \mathcal{C}^p(n), \gamma \subset \Lambda_{Kn/2} \text{ and} \\ \text{each blue cluster which has a site of } \gamma \text{ is contained in } \Lambda_{Kn} \end{array} \right\},$$

we have

$$\begin{aligned} \mathbf{P}_p^\eta[\mathcal{H}_n(K)] &\geq \mathbf{P}_p^\eta[\{T_{0,n}^p \leq \epsilon_0 Kn/2\} \cap \mathcal{G}(Kn/2) \cap \mathcal{O}(Kn)] \\ &\geq 1 - \epsilon/3 \quad \text{by (46), (47) and (8).} \end{aligned} \quad (49)$$

It follows from a RSW and FKG argument that, for each $\epsilon \in (0, 1)$ there is $\delta_1 \in (0, 1/4)$ such that for each $p \in (p(10/\delta_1), p_c)$, $n \in \mathbb{N}$ and the event

$$\mathcal{E}_n(\delta_1) := \{\text{diam}(\mathcal{C}^p(0)), \text{diam}(\mathcal{C}^p(n)) \geq 4\delta_1\},$$

we have

$$\mathbf{P}_p^\eta[\mathcal{E}_n(\delta_1)] \geq 1 - \epsilon/3. \quad (50)$$

For $\delta \in (0, \delta_1/20)$ and $p \in (p(10/\delta), p_c)$, write

$$\mathcal{S}_n(\delta) := \left\{ \begin{array}{l} \text{for each geodesic } \gamma \text{ from } \mathcal{C}^p(0) \text{ to } \mathcal{C}^p(n), \text{ the } L^\infty \text{ distance} \\ \text{between any pair of yellow sites in } \gamma \text{ is larger than } 2\delta \end{array} \right\}.$$

Note that $\mathcal{F}_n^p(\delta, K) \supset \mathcal{H}_n(K) \cap \mathcal{E}_n(\delta_1) \cap \mathcal{S}_n(\delta)$. It remains to bound the probability of the event $\mathcal{H}_n(K) \cap \mathcal{E}_n(\delta_1) \cap \neg \mathcal{S}_n(\delta)$. Assume that this event holds. Then there exist two yellow sites v_1, v_2 in a geodesic γ from $\mathcal{C}^p(0)$ to $\mathcal{C}^p(n)$, such that $\|v_1 - v_2\|_\infty \leq 2\delta$ and $v_1, v_2 \in \Lambda_{Kn/2}$. It is clear that there is $z \in 2\delta\mathbb{Z}^2 \cap \Lambda_{Kn/2}$ such that $v_1, v_2 \in \Lambda_{3\delta}(z)$. By (ii) of Proposition 2.5, the following statements hold: If there exist no yellow sites of γ in $\Lambda_{\delta_1}(z)$ except v_1, v_2 , then the event $\mathcal{A}_6(z; 4\delta, \delta_1 - \eta)$ occurs. Otherwise, there exists a yellow site $v \in \gamma \setminus \{v_1, v_2\}$ which is contained in $\Lambda_{\delta_1}(z)$ with minimal L^∞ distance to z . Write $L := \|z - v\|_\infty$. If $L < 4\delta + 2\eta$, then $\mathcal{A}_{(111111)}(z; 4\delta + 2\eta, \delta_1)$ occurs; if $L > \delta_1 - \eta$, then $\mathcal{A}_6(z; 4\delta, \delta_1 - 2\eta)$ occurs; if $L \in [4\delta + 2\eta, \delta_1 - \eta]$, then $\mathcal{A}_6(z; 4\delta, L - \eta) \cap \mathcal{A}_{(111111)}(z; L, \delta_1)$ occurs. The above argument implies that the event $\tilde{\mathcal{A}}_6(z; 4\delta, \delta_1)$ defined by (45) occurs; see Figure 6. Therefore, there exist universal constants $\lambda_6, C_3 > 0$, such that for each $R \geq 1$ and $\epsilon > 0$, there exist $\delta_1 = \delta_1(\epsilon) \in (0, 1/4)$ (independent of R), $K = K(\epsilon) \geq 3$ (independent of R) and $\delta = \delta(\epsilon, R) \in (0, \delta_1/20)$, such that for all $p \in (p(10/\delta), p_c)$ and $n \in \mathbb{N}$ with $n \leq R$,

$$\begin{aligned} \mathbf{P}_p^\eta[\mathcal{H}_n(K) \cap \mathcal{E}_n(\delta_1) \cap \neg \mathcal{S}_n(\delta)] &\leq \sum_{z \in 2\delta\mathbb{Z}^2 \cap \Lambda_{Kn/2}} \mathbf{P}_p^\eta[\tilde{\mathcal{A}}_6(z; 4\delta, \delta_1)] \\ &\leq C_3 \left(\frac{KR}{\delta}\right)^2 \left(\frac{\delta}{\delta_1}\right)^{2+\lambda_6} \quad \text{by Lemma 4.6} \\ &\leq \epsilon/3. \end{aligned}$$

Combining this with (49) and (50), we obtain

$$\mathbf{P}_p^\eta[\mathcal{F}_n^p(\delta, K)] \geq \mathbf{P}_p^\eta[\mathcal{H}_n(K) \cap \mathcal{E}_n(\delta_1) \cap \mathcal{S}_n(\delta)] \geq 1 - \epsilon.$$

□

The next lemma says that for fixed n, θ and each $p < p_c$ sufficiently close to p_c , with high probability $T_{0,n}^\theta$ equals $T_{0,n}^{p,\theta}$.

Lemma 4.8. *Let \mathbf{P} be a coupling such that $\omega_p^\eta \rightarrow \omega^\infty$ in $(\mathcal{H}, d_{\mathcal{H}})$ a.s. as $p \uparrow p_c$. For each $R \geq 1$ and $\epsilon > 0$, there exists $p_0 = p_0(\epsilon, R) \in (0, p_c)$ such that for all $p \in (p_0, p_c)$, $\theta \in [0, 2\pi]$ and $n \in \mathbb{N}$ with $n \leq R$, we have*

$$\mathbf{P}[T_{0,n}^\theta = T_{0,n}^{p,\theta}] \geq 1 - \epsilon.$$

Proof. We start with an upper bound on the probability of the event that there exist two large blue clusters in a box such that they are close to each other but do not form a chain. For $S, S' \subset \mathbb{C}$, let $d_\infty(S, S') := \inf\{\|z - z'\|_\infty : z \in S, z' \in S'\}$ denote the L^∞ -distance between S and S' . There exist $C_1, \lambda_6 > 0$ such that for all $0 < 5r_1 \leq r_2 \leq 1$, $k \in \mathbb{N}$ and $p \in (p(5/r_1), p_c)$,

$$\begin{aligned} &\mathbf{P}[\exists \mathcal{C}_1^p, \mathcal{C}_2^p \in \mathcal{C}^p(r_2) \text{ in } \Lambda_k \text{ such that } 0 < d_\infty(\mathcal{C}_1^p, \mathcal{C}_2^p) \leq r_1 \text{ and } (\mathcal{C}_1^p, \mathcal{C}_2^p) \text{ is not a chain}] \\ &\leq \mathbf{P}_p^\eta[\mathcal{A}_6(z; r_1, r_2/4) \text{ occurs for some point } z \in \Lambda_k] \\ &\leq C_1 \left(\frac{k}{r_1}\right)^2 \left(\frac{r_1}{r_2}\right)^{2+\lambda_6} \quad \text{by (10)}. \end{aligned} \quad (51)$$

For the continuum configuration the probability of the corresponding event is bounded as follows: Since for any fixed $r_2 > 0$ and $k \in \mathbb{N}$, there are a.s. finitely many clusters of $\mathcal{C}(r_2)$ in Λ_k (by Theorem 3.2), we know that for any fixed $r_2 > 0$ and $k \in \mathbb{N}$,

$$\mathbf{P}[\exists \mathcal{C}_1, \mathcal{C}_2 \in \mathcal{C}(r_2) \text{ in } \Lambda_k \text{ such that } 0 < d_\infty(\mathcal{C}_1, \mathcal{C}_2) \leq r_1] \rightarrow 0 \quad \text{as } r_1 \rightarrow 0. \quad (52)$$

Let $\mathcal{F}_n^{p,\theta}(\delta, K)$ be the event defined above Lemma 4.7. Write

$$\mathcal{E}_n^{p,\theta}(\delta, K) := \left\{ \begin{array}{l} \exists \text{ a chain } \Gamma^p \text{ from } \mathcal{C}^p(0) \text{ to } \mathcal{C}^p(ne^{i\theta}), \text{ such that } T_{0,n}^{p,\theta} = |\Gamma^p| - 1, \\ \text{diam}(\mathcal{C}^p) \geq \delta \text{ for all } \mathcal{C}^p \in \Gamma^p, \text{ and } \Gamma^p \subset \Lambda_{Kn}; \\ \exists \text{ a chain } \Gamma \text{ from } \mathcal{C}(0) \text{ to } \mathcal{C}(ne^{i\theta}), \text{ such that } |\Gamma| = |\Gamma^p|, \\ \text{diam}(\mathcal{C}) \geq \delta/2 \text{ for all } \mathcal{C} \in \Gamma, \text{ and } \Gamma \subset \Lambda_{Kn} \end{array} \right\}$$

It is clear that $\mathcal{E}_n^{p,\theta}(\delta, K) \subset \{T_{0,n}^\theta \leq T_{0,n}^{p,\theta}\}$. Therefore, for each $R \geq 1$ and $\epsilon > 0$, there exist $K = K(\epsilon) \geq 3$ with $K \in \mathbb{N}$, $\delta = \delta(\epsilon, R) \in (0, 1/4)$, $\delta_1 = \delta_1(\epsilon, R) \in (0, \delta/100)$ and $p_1 = p_1(\epsilon, R) \in (0, p_c)$, such that for all $p \in (p_1, p_c)$, $\theta \in [0, 2\pi]$ and $n \in \mathbb{N}$ with $n \leq R$, we have

$$\begin{aligned} \mathbf{P}[T_{0,n}^\theta \leq T_{0,n}^{p,\theta}] &\geq \mathbf{P}[\mathcal{E}_n^{p,\theta}(\delta, K)] \\ &\geq \mathbf{P}[\mathcal{F}_n^{p,\theta}(\delta, K)] - \mathbf{P}[d_H(\mathcal{C}^p(0), \mathcal{C}(0)) \geq \delta_1] - \mathbf{P}[d_H(\mathcal{C}^p(ne^{i\theta}), \mathcal{C}(ne^{i\theta})) \geq \delta_1] \\ &\quad - \mathbf{P}[\text{dist}(\mathcal{C}^p|_{\Lambda_{Kn}}, \mathcal{C}|_{\Lambda_{Kn}}) \geq \delta_1] \\ &\quad - \mathbf{P}[\exists \mathcal{C}_1, \mathcal{C}_2 \in \mathcal{C}(\delta/4) \text{ in } \Lambda_{KR} \text{ such that } 0 < d_\infty(\mathcal{C}_1, \mathcal{C}_2) \leq 3\delta_1] \\ &\geq 1 - \epsilon/2 \quad \text{by (52), Theorem 3.2, Lemmas 4.7 and 4.4,} \end{aligned} \quad (53)$$

where the second inequality is due to the observation:

$$\begin{aligned} \mathcal{E}_n^{p,\theta}(\delta, K) &\supset \mathcal{F}_n^{p,\theta}(\delta, K) \cap \{d_H(\mathcal{C}^p(0), \mathcal{C}(0)) \leq \delta_1\} \cap \{d_H(\mathcal{C}^p(ne^{i\theta}), \mathcal{C}(ne^{i\theta})) \leq \delta_1\} \\ &\quad \cap \{\text{dist}(\mathcal{C}^p|_{\Lambda_{Kn}}, \mathcal{C}|_{\Lambda_{Kn}}) \leq \delta_1\} \cap \{\nexists \mathcal{C}_1, \mathcal{C}_2 \in \mathcal{C}(\delta/4) \text{ in } \Lambda_{KR} \text{ such that } 0 < d_\infty(\mathcal{C}_1, \mathcal{C}_2) \leq 3\delta_1\}. \end{aligned}$$

It remains to bound $\mathbf{P}[T_{0,n}^{p,\theta} \leq T_{0,n}^\theta]$ from below. Combining (53) and Lemma 4.5, we get that for each fixed $\theta \in [0, 2\pi]$ and $n \in \mathbb{N}$, $T_{0,n}^\theta$ is almost surely finite. Moreover, by Theorem 3.2 we know that all the continuum clusters of \mathcal{C} are almost surely bounded and the distribution of \mathcal{C} is invariant under rotations. Therefore, for each $R \geq 1$ and $\epsilon > 0$, there is an integer $N = N(\epsilon, R) \geq 2R$ and a $\delta_0 = \delta_0(\epsilon, R) \in (0, 1/4)$ such that for all $\theta \in [0, 2\pi]$, $n \in \mathbb{N}$ with $n \leq R$, and the event

$$\mathcal{G}_n^\theta(\delta_0, N) := \left\{ \begin{array}{l} \exists \text{ a chain } \Gamma \text{ from } \mathcal{C}(0) \text{ to } \mathcal{C}(ne^{i\theta}), \text{ such that } T_{0,n}^\theta = |\Gamma| - 1, \\ \text{diam}(\mathcal{C}) \geq \delta_0 \text{ for all } \mathcal{C} \in \Gamma, \text{ and } \Gamma \subset \Lambda_N \end{array} \right\},$$

we have

$$\mathbf{P}[\mathcal{G}_n^\theta(\delta_0, N)] \geq 1 - \epsilon/4. \quad (54)$$

Then, similarly to the proof of (53), for each $R \geq 1$ and $\epsilon > 0$ there exist $N = N(\epsilon, R) \geq 2R$ with $N \in \mathbb{N}$, $\delta_0 = \delta_0(\epsilon, R) \in (0, 1/4)$, $\delta_2 = \delta_2(\epsilon, R) \in (0, \delta_0/100)$ and $p_2 = p_2(\epsilon, R) \in (0, p_c)$, such that for all $p \in (p_2, p_c)$, $\theta \in [0, 2\pi]$ and $n \in \mathbb{N}$ with $n \leq R$,

$$\begin{aligned} &\mathbf{P}[T_{0,n}^{p,\theta} \leq T_{0,n}^\theta] \\ &\geq \mathbf{P} \left[\begin{array}{l} \mathcal{G}_n^\theta(\delta_0, N) \text{ occurs, and there is a chain } \Gamma^p \text{ from } \mathcal{C}^p(0) \text{ to } \mathcal{C}^p(ne^{i\theta}), \\ \text{such that } |\Gamma^p| = |\Gamma|, \text{diam}(\mathcal{C}^p) \geq \delta_0/2 \text{ for all } \mathcal{C}^p \in \Gamma^p, \text{ and } \Gamma^p \subset \Lambda_N^\theta \end{array} \right] \\ &\geq \mathbf{P}[\mathcal{G}_n^\theta(\delta_0, N)] - \mathbf{P}[d_H(\mathcal{C}^p(0), \mathcal{C}(0)) \geq \delta_2] - \mathbf{P}[d_H(\mathcal{C}^p(ne^{i\theta}), \mathcal{C}(ne^{i\theta})) \geq \delta_2] - \mathbf{P}[\text{dist}(\mathcal{C}^p|_{\Lambda_N}, \mathcal{C}|_{\Lambda_N}) \geq \delta_2] \\ &\quad - \mathbf{P}[\exists \mathcal{C}_1^p, \mathcal{C}_2^p \in \mathcal{C}^p(\delta_0/4) \text{ in } \Lambda_N \text{ such that } 0 < d_\infty(\mathcal{C}_1^p, \mathcal{C}_2^p) \leq 3\delta_2 \text{ and } (\mathcal{C}_1^p, \mathcal{C}_2^p) \text{ is not a chain}] \\ &\geq 1 - \epsilon/2 \quad \text{by (54), Lemma 4.4, Theorem 3.2 and (51).} \end{aligned} \quad (55)$$

Then Lemma 4.8 follows from a combination of inequalities (55) and (53). \square

Lemma 4.9. *Let $C_0 > 0$ be the absolute constant in Lemma 4.5. We have*

$$\mathbf{E}^\infty[T_{0,1}] \leq C_0. \quad (56)$$

Furthermore, there is a constant $C_1 > 0$ such that for each $\epsilon > 0$, there exists $N = N(\epsilon) > 0$ such that for all $n \in \mathbb{N}$ with $n > N$,

$$\mathbf{P}^\infty[T_{0,n} \geq C_1 n] \geq 1 - \epsilon. \quad (57)$$

Proof. Combining Lemmas 4.5 and 4.8 gives (56).

Inequality (47) implies that there exists a constant $C_1 > 0$ such that for each $\epsilon > 0$, there exists $N = N(\epsilon) > 0$ such that for all $p \in (p(10), p_c)$ and $n \in \mathbb{N}$ with $n > N$,

$$\mathbf{P}_p^\eta[T_{0,n}^p \geq C_1 n] \geq 1 - \epsilon/2.$$

Then (57) follows from this and Lemma 4.8. \square

The following is a law of large numbers for the point-to-point passage times of our continuum FPP.

Proposition 4.10. *Suppose $n \in \mathbb{N}$. There exists a constant $\nu > 0$ such that for any fixed $\theta \in [0, 2\pi]$,*

$$\lim_{n \rightarrow \infty} \frac{T_{0,n}^\theta}{n} = \nu \quad \mathbf{P}^\infty\text{-a.s. and} \quad \lim_{n \rightarrow \infty} \frac{\mathbf{E}^\infty T_{0,n}^\theta}{n} = \inf_n \frac{\mathbf{E}^\infty T_{0,n}^\theta}{n} = \nu.$$

Proof. Since the law of \mathcal{C} is invariant under rotations by Theorem 3.2, all the families $(T_{0,n}^\theta)_{n \in \mathbb{N}}$ for $\theta \in [0, 2\pi]$ have the same distribution. Thus, to prove Proposition 4.10, it suffices to prove it in the case $\theta = 0$. Suppose that $m, n \in \mathbb{Z}_+$ with $m < n$. We verify that the family of random variables $(T_{m,n})_{0 \leq m < n}$ satisfies the conditions of the subadditive ergodic theorem (see, e.g., [29] or Theorem 2.2 in [2]):

- $T_{0,n} \leq T_{0,m} + T_{m,n}$ for all $0 < m < n$.
Note that a concatenation of chains from $\mathcal{C}(0)$ to $\mathcal{C}(m)$ and from $\mathcal{C}(m)$ to $\mathcal{C}(n)$ yields a chain from $\mathcal{C}(0)$ to $\mathcal{C}(n)$. Combining this observation and the definition of $T_{m,n}$ gives the above triangle inequality.
- The distributions of the sequences $(T_{m,m+j})_{j \geq 1}$ and $(T_{m+1,m+j+1})_{j \geq 1}$ are the same for all $m \geq 0$.
The law of \mathcal{C} is translation invariant by Theorem 3.2, which implies this immediately.
- $(T_{nj,(n+1)j})_{n \geq 1}$ is a stationary ergodic sequence for each $j \geq 1$.
Define the horizontal shifts of the plane $\tau_j: \mathbb{C} \rightarrow \mathbb{C}, z \mapsto z - j$. Then $T_{nj,(n+1)j}(\mathcal{C}) = T_{0,j}(\tau_{nj}\mathcal{C})$. By Theorem 3.2, the law of \mathcal{C} is invariant under translations, so τ_j is measure preserving and $(T_{nj,(n+1)j})_{n \geq 1}$ is stationary. Next we show that τ_j is also mixing, which implies that $(T_{nj,(n+1)j})_{n \geq 1}$ is ergodic. When \mathcal{A}, \mathcal{B} are events which depend only on the realization of \mathcal{C} inside some box Λ_k , then $\lim_{n \rightarrow \infty} \mathbb{P}[\mathcal{A} \cap \tau_j^{-n}\mathcal{B}] = \mathbb{P}[\mathcal{A}]\mathbb{P}[\mathcal{B}]$ follows immediately since the events \mathcal{A} and $\tau_j^{-n}\mathcal{B}$ are independent for large n by Theorem 3.2. For arbitrary events \mathcal{A} and \mathcal{B} depending on \mathcal{C} , one approximates \mathcal{A} and \mathcal{B} by events which depend only on the realization of \mathcal{C} inside Λ_k , with $k \rightarrow \infty$. Then the result follows easily.
- $\mathbb{E}[T_{0,1}] < \infty$ and there is a constant $C > 0$ such that for each n , $\mathbb{E}[T_{0,n}] > -Cn$.
 $\mathbb{E}[T_{0,1}] < \infty$ follows from (56) and the rest of this item is obvious since $T_{0,n}$ is non-negative by definition.

Then by the subadditive ergodic theorem and (57) we obtain Proposition 4.10, where we use (57) to show that the limit ν is positive. \square

5. Convergence of normalized time constants

Let ν be as in Proposition 4.10, which is the ‘‘time constant’’ for the continuum FPP. First, in Section 5.1, we show that for each fixed $u \in \mathbb{U}$, ν is an upper bound for the upper limit of $L(p)\mu(p, u)$. Next, in Section 5.2, we show that ν is also a lower bound for the lower limit of $L(p)\mu(p, u)$. Finally, in Section 5.3 we prove our main result, Theorem 1.1, which is an immediate consequence of the above two results.

The basic idea is as follows. The results in Section 4 give that the discrete FPP on $\eta\mathbb{T}$ in any fixed box is ‘‘well-approximated’’ by the corresponding continuum FPP, as $p \uparrow p_c$. This, combined with appropriate renormalization arguments, implies that the approximation also holds in the whole plane.

5.1. Upper bound

The goal of this subsection is to prove:

Lemma 5.1. *For each $u \in \mathbb{U}$, we have $\limsup_{p \uparrow p_c} L(p)\mu(p, u) \leq \nu$.*

To show this, we need the following result on passage times of paths constrained in a box. For $x, y \in \mathbb{C}$ and $r > 0$, define

$$\text{Box}(x, y; r) := \{z \in \mathbb{C} : z \text{ is within } L^\infty\text{-distance } r \text{ of the straight line segment joining } x \text{ to } y\}.$$

Lemma 5.2. *There exists $K \in \mathbb{N}$ such that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for all $n \geq N$, there exists $p_0 \in (0, p_c)$ such that for all $p \in (p_0, p_c)$, $\theta \in [0, 2\pi]$ and $z \in \mathbb{C}$,*

$$\mathbf{P}_p^\eta \left[\text{there is a path } \gamma \text{ from } \mathcal{C}^p(z) \text{ to } \mathcal{C}^p(z + ne^{i\theta}) \text{ in } \text{Box}(z, z + ne^{i\theta}; Kn) \text{ s.t. } T(\gamma) \leq (\nu + \epsilon)n \right] \geq 1 - \epsilon.$$

Proof. Since the law of \mathcal{C} is invariant under rotations by Theorem 3.2, for any fixed n the passage times $T_{0,n}^\theta$ for $\theta \in [0, 2\pi]$ have the same distribution. This and Proposition 4.10 imply that for each $\epsilon > 0$, there exists $N_1 \in \mathbb{N}$ such that for all $n \geq N_1$ and $\theta \in [0, 2\pi]$,

$$\mathbf{P}^\infty [T_{0,n}^\theta \leq (\nu + \epsilon)n] \geq 1 - \epsilon/5.$$

Combining the above inequality and Lemma 4.8, we obtain that, for each $\epsilon > 0$, there exists $N_1 \in \mathbb{N}$ such that for all $n \geq N_1$, there exists $p_1 \in (p(10), p_c)$ such that for all $p \in (p_1, p_c)$ and $\theta \in [0, 2\pi]$,

$$\mathbf{P}_p^\eta [T_{0,n}^{p,\theta} \leq (\nu + \epsilon)n] \geq 1 - \epsilon/4. \quad (58)$$

By (47), there exists $K_1 \in \mathbb{N}$ such that for each $\epsilon \in (0, 1)$, there exists $N_2 \in \mathbb{N}$ such that for all $n \geq N_2$, $p \in (p(10), p_c)$ and $\theta \in [0, 2\pi]$,

$$\mathbf{P}_p^\eta [\exists \text{ a path } \gamma \text{ starting at a site in } \mathbb{D}_{1/2}, \text{ such that } T(\gamma) \leq (\nu + 1)n \text{ and } \gamma \not\subset \text{Box}(0, ne^{i\theta}; K_1 n)] \leq \epsilon/4. \quad (59)$$

Then (58) and (59) imply that there is $K_1 \in \mathbb{N}$ such that for each $\epsilon \in (0, 1)$, there exists $N = \max\{N_1, N_2\}$ such that for all $n \geq N$, there exists $p_1 \in (p(10), p_c)$ such that for all $p \in (p_1, p_c)$ and $\theta \in [0, 2\pi]$,

$$\mathbf{P}_p^\eta [\exists \text{ a path } \gamma \text{ from } \mathcal{C}^p(0) \text{ to } \mathcal{C}^p(ne^{i\theta}) \text{ in } \text{Box}(0, ne^{i\theta}; K_1 n) \text{ such that } T(\gamma) \leq (\nu + \epsilon)n] \geq 1 - \epsilon/2. \quad (60)$$

We need to generalize this result to pairs of points $z, z + ne^{i\theta}$ for $z \in \mathbb{C}$ and $\theta \in [0, 2\pi]$. It is easy to see that for each $\epsilon > 0$, there exist $\delta \in (0, 1/2)$ and $p_2 \in (p(10/\delta), p_c)$ such that for all $p \in (p_2, p_c)$ and $x \in \mathbb{C}$,

$$\begin{aligned} & \mathbf{P}_p^\eta [\mathcal{C}^p(y) = \mathcal{C}^p(x) \text{ for all } y \in \mathbb{D}_\eta(x)] \\ & \geq \mathbf{P}_p^\eta \left[\begin{array}{l} \text{diam}(\mathcal{C}^p(x)) \geq \delta \text{ and there is a unique open cluster having diameter } \text{diam}(\mathcal{C}^p(x)) \text{ in } \mathbb{D}_{1/2}(x); \\ \text{all the open clusters which have diameters at least } \delta \text{ in } \mathbb{D}_{1/2+\eta}(x) \text{ lie in } \mathbb{D}_{1/2-\eta}(x) \end{array} \right] \\ & \geq 1 - \epsilon/4 \quad \text{by FKG, RSW, Proposition 4.2 and (12)}. \end{aligned} \quad (61)$$

For $z \in \mathbb{C}$, let \tilde{z} denote the site of $\eta\mathbb{T}$ closest to z . Then, there exists $K = K_1 + 1 \in \mathbb{N}$ such that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for all $n \geq N$, there exists $p_0 = \max\{p_1, p_2\} \in (0, p_c)$ such that for all $p \in (p_0, p_c)$, $\theta \in [0, 2\pi]$ and $z \in \mathbb{C}$,

$$\begin{aligned} & \mathbf{P}_p^\eta \left[\text{there is a path } \gamma \text{ from } \mathcal{C}^p(z) \text{ to } \mathcal{C}^p(z + ne^{i\theta}) \text{ in } \text{Box}(z, z + ne^{i\theta}; Kn) \text{ such that } T(\gamma) \leq (\nu + \epsilon)n \right] \\ & \geq \mathbf{P}_p^\eta \left[\begin{array}{l} \mathcal{C}^p(\tilde{z}) = \mathcal{C}^p(z), \mathcal{C}^p(\tilde{z} + ne^{i\theta}) = \mathcal{C}^p(z + ne^{i\theta}), \text{ and there is a path } \gamma \text{ from} \\ \mathcal{C}^p(\tilde{z}) \text{ to } \mathcal{C}^p(\tilde{z} + ne^{i\theta}) \text{ in } \text{Box}(\tilde{z}, \tilde{z} + ne^{i\theta}; K_1 n) \text{ such that } T(\gamma) \leq (\nu + \epsilon)n \end{array} \right] \\ & \geq 1 - \epsilon \quad \text{by (61), (60) and the symmetry of } \eta\mathbb{T}, \end{aligned}$$

which concludes the proof. \square




FIG 7. Bernoulli site percolation configuration ω_p^η on $\eta\mathbb{T}$ induces the $6K$ -dependent family $X^p(\epsilon)$, which is a $6K$ -dependent bond percolation on \mathbb{Z}^2 .

Proof of Lemma 5.1. We will use a renormalization argument. For simplicity, we prove this lemma in the case $u = 1$; the proof for a general $u \in \mathbb{U}$ is analogous. By Lemma 5.2, there exists $K \in \mathbb{N}$ such that for each $\epsilon > 0$, we can choose $N = N(\epsilon) \in \mathbb{N}$ and $p_0 = p_0(\epsilon) \in (0, p_c)$, such that for each $p \in (p_0, p_c)$ and each bond $e = (e_-, e_+) \in E(\mathbb{Z}^2)$,

$$\mathbf{P}_p^\eta[\exists \text{ a path } \gamma \text{ from } \mathcal{C}^p(Ne_-) \text{ to } \mathcal{C}^p(Ne_+) \text{ in } \text{Box}(Ne_-, Ne_+; KN) \text{ such that } T(\gamma) \leq (\nu + \epsilon)N] \geq 1 - \epsilon. \quad (62)$$

We declare a bond $e \in E(\mathbb{Z}^2)$ to be ϵ -good if the event in (62) occurs. We call a (bond) path of \mathbb{Z}^2 ϵ -good if all the bonds of this path are ϵ -good. The family $X^p(\epsilon) = \{X_e^p(\epsilon) : e \in E(\mathbb{Z}^2)\}$ is defined by

$$X_e^p(\epsilon)(\omega_p^\eta) := \begin{cases} 1 & \text{if } e \text{ is } \epsilon\text{-good,} \\ 0 & \text{otherwise.} \end{cases}$$

It is easy to check that $X^p(\epsilon)$ is a $6K$ -dependent family; see Figure 7.

We have from Proposition 2.10 that for each $\epsilon > 0$, there is $p_1 = p_1(\epsilon) \in (0, 1/2)$ such that for all large n (depending on ϵ),

$$\mathbf{P}_{\mathbb{Z}^2, p_1}^{\text{bond}}[D(0, n) \leq (1 + \epsilon)n] \geq 3/4. \quad (63)$$

Our proof also works if the value $3/4$ in inequality (63) is replaced with any other fixed value in $(0, 1)$. By (62) and Theorem 2.8, for each $\epsilon > 0$, we can choose $\epsilon_1 = \epsilon_1(\epsilon) \in (0, \epsilon)$, such that for all $p \in (p_0(\epsilon_1), p_c)$, $X^p(\epsilon_1)$ stochastically dominates the Bernoulli bond percolation on \mathbb{Z}^2 with measure $\mathbf{P}_{\mathbb{Z}^2, p_1}^{\text{bond}}$. (Recall that in the present paper, each bond of \mathbb{Z}^2 takes the value 0 with probability p_1 under $\mathbf{P}_{\mathbb{Z}^2, p_1}^{\text{bond}}$.) Note that for the configuration ω_p^η , if there is an ϵ_1 -good path of \mathbb{Z}^2 from 0 to n so that the number of bonds of this path is not larger than $(1 + \epsilon)n$, then

$$T(\mathcal{C}^p(0), \mathcal{C}^p(Nn)) \leq (1 + \epsilon)(\nu + \epsilon_1)Nn \leq (1 + \epsilon)(\nu + \epsilon)Nn.$$

Therefore, for all $p \in (p_0(\epsilon_1), p_c)$ and all large n ,

$$\mathbf{P}_p^\eta[T(\mathcal{C}^p(0), \mathcal{C}^p(Nn)) \leq (1 + \epsilon)(\nu + \epsilon)Nn] \geq \mathbf{P}_{\mathbb{Z}^2, p_1}^{\text{bond}}[D(0, n) \leq (1 + \epsilon)n] \geq 3/4.$$

Then we have

$$\mathbf{P}_p^\eta[T(0, Nn) \leq (1 + \epsilon)(\nu + \epsilon)Nn + 2/\eta] \geq \mathbf{P}_p^\eta[T(\mathcal{C}^p(0), \mathcal{C}^p(Nn)) \leq (1 + \epsilon)(\nu + \epsilon)Nn] \geq 3/4.$$

By using this and the fact that \mathbf{P}_p -almost surely $a_{0,m}/m$ tends to $\mu(p)$ as $m \rightarrow \infty$ (see (1)), we obtain that $L(p)\mu(p) \leq (1 + \epsilon)(\nu + \epsilon)$ for all $p \in (p_0(\epsilon_1), p_c)$. Letting $\epsilon \rightarrow 0$, we have $\limsup_{p \uparrow p_c} L(p)\mu(p) \leq \nu$. \square

Remark 5.3. Lemma 5.2 is a key ingredient for the renormalization argument above. The following stronger version of Lemma 5.2 for a fixed θ was proved in [41]:

Fix any $\theta \in [0, 2\pi]$. For each $\epsilon > 0$, there exist $K, N \in \mathbb{N}$, such that for all $n \geq N$, there exists $p_0 = p_0(\theta, \epsilon, n) \in (0, p_c)$ such that for all $p \in (p_0, p_c)$ and $z \in \mathbb{C}$,

$$\mathbf{P}_p^\eta \left[\text{there is a path } \gamma \text{ from } \mathcal{C}^p(z) \text{ to } \mathcal{C}^p(z + ne^{i\theta}) \text{ in } \text{Box}(z, z + ne^{i\theta}; K) \text{ such that } T(\gamma) \leq (\nu + \epsilon)n \right] \geq 1 - \epsilon.$$

An application of this result may yield an 1-dependent bond percolation in a similar renormalization argument as above. However, to show this version of Lemma 5.2 we need to construct the scaling limit of the collection of pieces of clusters contained in a strip (see Remark 3.6) and study FPP in strips, which lead to many technical complications. See [41] for more details.

5.2. Lower bound

The goal of this subsection is to prove:

Lemma 5.4. *For each $u \in \mathbb{U}$, we have $\liminf_{p \uparrow p_c} L(p)\mu(p, u) \geq \nu$.*

In [20], Grimmett and Kesten used a ‘‘block approach’’ to obtain exponential large deviation bounds for first-passage times for a single FPP model. We will use their method with some modifications to prove Lemma 5.4, concerning the family $\{\mathbf{P}_p^\eta\}_{p \in (0, p_c)}$ of Bernoulli FPP measures. Similarly to the proof of Lemma 5.1, the proof of Lemma 5.4 also employs a type of renormalization, but it is quite different and more complicated.

The proof of Lemma 5.4 is divided into two parts: estimates for line-to-line passage times and a renormalization argument, which are given in Sections 5.2.1 and 5.2.2, respectively.

5.2.1. Line-to-line passage time

Let $\mathbb{D}^\eta(z)$ denote the union of all hexagons of $\eta\mathbb{H}$ which intersect $\mathbb{D}(z)$. We require the following lemma in order to study the line-to-line passage time $l_{n,m}^{p,\theta}(z)$ defined above Lemma 2.6. The main difficulty of its proof comes from the endpoints of a geodesic between $\mathbb{D}^\eta(x)$ and $\mathbb{D}^\eta(y)$. More precisely, we need to bound the passage time from $\mathcal{C}^p(\tilde{x})$ (resp. $\mathcal{C}^p(\tilde{y})$) to this geodesic, where \tilde{x} is a well-chosen point in $\mathbb{D}(x)$.

Lemma 5.5. *For each $\epsilon \in (0, \nu)$ and $\delta > 0$, there exists an integer $N = N(\epsilon, \delta) \geq 10$, such that for any integers n and m with $N \leq n \leq m$, there exists $p_0 = p_0(\epsilon, \delta, m) \in (0, p_c)$ such that for all $p \in (p_0, p_c)$ and $x, y \in \mathbb{C}$ with $n \leq |x - y| \leq m$,*

$$\mathbf{P}_p^\eta[T(\mathbb{D}^\eta(x), \mathbb{D}^\eta(y)) \leq (\nu - \epsilon)n] \leq \delta.$$

Proof. It suffices to prove the lemma in the case $y = 0$. Since the law of \mathcal{C} is invariant under rotations by Theorem 3.2, for any fixed n the passage times $T_{0,n}^\theta$ for $\theta \in [0, 2\pi]$ have the same distribution. This and Proposition 4.10 imply that for each $\epsilon \in (0, \nu)$ and $\delta > 0$, there exists an integer $N_1 = N_1(\epsilon, \delta) \geq 10$ such that for all $\theta \in [0, 2\pi]$ and all integers $n \geq N_1$,

$$\mathbf{P}^\infty[T_{0,n}^\theta \leq (\nu - \epsilon/2)n] \leq \delta/4. \quad (64)$$

Combining (64) and Lemma 4.8, we get that for each $\epsilon \in (0, \nu)$ and $\delta > 0$, there exists $N_1 \geq 10$ such that for any integers $m \geq N_1$, there exists $p_0 = p_0(\epsilon, \delta, m) \in (0, p_c)$ such that for all $p \in (p_0, p_c)$, $\theta \in [0, 2\pi]$ and $n \in \mathbb{N}$ with $N_1 \leq n \leq m$,

$$\mathbf{P}_p^\eta[T_{0,n}^{p,\theta} \leq (\nu - \epsilon/2)n] \leq \delta/2. \quad (65)$$

For $x \in \mathbb{C}$ and $p \in (p(10), p_c)$, we write

$$X^p(x)(\omega_p^\eta) := \inf\{T(\gamma) : \gamma \text{ is a circuit surrounding } x \text{ in } A(x; 2, 3)\},$$

$$Y^p(x)(\omega_p^\eta) := \inf\{T(\gamma) : \gamma \text{ is a path from a site in } \mathcal{C}^p(x) \text{ to a site outside } \Lambda_3(x)\}.$$

It is clear that for each $x \in \mathbb{C}$ with $|x| \geq 10$, there is $\tilde{x} \in \mathbb{C}$ such that $\arg(\tilde{x}) = \arg(x)$, $|x - \tilde{x}| \leq 1/2$ and $|\tilde{x}|$ is an integer. Suppose $p \in (p(10), p_c)$. Then we have $\Lambda_2(\tilde{x}) \supset \mathbb{D}(x) \supset \mathbb{D}_{1/2}(\tilde{x}) \supset \mathcal{C}^p(\tilde{x})$. Observe that for all $x \in \mathbb{C}$ with $|x| \geq 10$,

$$T_{0,|\tilde{x}|}^{p,\arg(\tilde{x})} \leq X^p(0) + Y^p(0) + X^p(\tilde{x}) + Y^p(\tilde{x}) + T(\mathbb{D}^\eta(0), \mathbb{D}^\eta(x)). \quad (66)$$

Applying Lemma 2.6, it is easy to obtain that for each $\delta > 0$, there is $N_2 = N_2(\delta) \geq 1$ such that for all $x \in \mathbb{C}$ and $p \in (p(10), p_c)$,

$$\mathbf{P}_p^\eta[X^p(x) \geq N_2] \leq \delta/10. \quad (67)$$

$$Y_k^p := \inf\{T(\gamma)(\omega_p^\eta) : \gamma \text{ is a top-bottom crossing of } [n-1, n] \times [k\epsilon_1 n, (k+1)\epsilon_1 n]\},$$

$$Z_{j,k}^p := T(\mathbb{D}^\eta(z_{0,j}), \mathbb{D}^\eta(z_{n,k}))(\omega_p^\eta) \text{ for points } z_{0,j} := (0, j\epsilon_1 n) \text{ and } z_{n,k} := (n, k\epsilon_1 n).$$

For ω_p^η , we let γ^p be a left-right crossing of $[0, n] \times [0, m]$ with $T(\gamma^p) = l_{n,m}^p$, $\gamma_{0,j}^p$ be a top-bottom crossing of $[0, 1] \times [j\epsilon_1 n, (j+1)\epsilon_1 n]$ with $T(\gamma_{0,j}^p) = X_j^p$, and $\gamma_{n,k}^p$ be a top-bottom crossing of $[n-1, n] \times [k\epsilon_1 n, (k+1)\epsilon_1 n]$ with $T(\gamma_{n,k}^p) = Y_k^p$. It is clear that $\mathbb{D}^\eta(z_{0,j}) \cap \gamma_{0,j}^p \neq \emptyset$ and $\mathbb{D}^\eta(z_{n,k}) \cap \gamma_{n,k}^p \neq \emptyset$. Moreover, γ^p intersects $\mathbb{D}^\eta(z_{0,j^*})$ or γ_{0,j^*}^p for some integer $j^* \in [0, \lceil m/(\epsilon_1 n) \rceil]$, and intersects $\mathbb{D}^\eta(z_{n,k^*})$ or γ_{n,k^*}^p for some integer $k^* \in [0, \lceil m/(\epsilon_1 n) \rceil]$. See Figure 8 for an illustration. Assume that the event

$$\left(\bigcap_{j=0}^{\lceil m/(\epsilon_1 n) \rceil} \{X_j^p \leq \epsilon n/3\} \right) \cap \left(\bigcap_{k=0}^{\lceil m/(\epsilon_1 n) \rceil} \{Y_k^p \leq \epsilon n/3\} \right) \cap \{l_{n,m}^p \leq n(\nu - \epsilon)\}$$

holds. Then

$$Z_{j^*,k^*}^p \leq T(\gamma^p) + T(\gamma_{0,j^*}^p) + T(\gamma_{n,k^*}^p) \leq n(\nu - \epsilon/3).$$

Note that $m/(\epsilon_1 n) \leq 3CK/\epsilon$ since $n \leq m \leq Cn$ and $\epsilon_1 = \epsilon/(3K)$. Then from the above argument, we have

$$\begin{aligned} \mathbf{P}_p^\eta[l_{n,m}^p \leq n(\nu - \epsilon)] &\leq \sum_{0 \leq j \leq \lceil 3CK/\epsilon \rceil} \mathbf{P}_p^\eta[X_j^p \geq \epsilon n/3] + \sum_{0 \leq k \leq \lceil 3CK/\epsilon \rceil} \mathbf{P}_p^\eta[Y_k^p \geq \epsilon n/3] \\ &\quad + \sum_{j,k \in [0, \lceil 3CK/\epsilon \rceil]} \mathbf{P}_p^\eta[Z_{j,k}^p \leq (\nu - \epsilon/3)n]. \end{aligned} \quad (69)$$

Let us bound the three terms on the right side of (69). By Lemma 2.6, there exists $N_0 = N_0(\epsilon, \delta, C) \geq 2/\epsilon_1$ such that for all $p \in (p(10), p_c)$, $j, k \in \mathbb{Z}_+$ and $n \geq N_0$, we have

$$\mathbf{P}_p^\eta[X_j^p \geq \epsilon n/3] \leq \delta\epsilon/(12CK) \quad \text{and} \quad \mathbf{P}_p^\eta[Y_k^p \geq \epsilon n/3] \leq \delta\epsilon/(12CK). \quad (70)$$

By Lemma 5.5, there is an integer $N = N(\epsilon, \delta, C) \geq N_0$ such that for all $n \geq N$, there exists $p_0 = p_0(\epsilon, \delta, C, n) \in (0, p_c)$ such that for all $p \in (p_0, p_c)$ and $j, k \in [0, \lceil 3CK/\epsilon \rceil]$,

$$\mathbf{P}_p^\eta[Z_{j,k}^p \leq (\nu - \epsilon/3)n] \leq \delta\epsilon^2/(48C^2K^2). \quad (71)$$

Plugging (70) and (71) into (69), we obtain $\mathbf{P}_p^\eta[l_{n,m}^p \leq n(\nu - \epsilon)] \leq \delta$ for all $N \leq n \leq m \leq Cn$ and all $p \in (p_0, p_c)$. \square

5.2.2. Renormalization

We now follow the main lines of the proof of Theorem 3.1 in [20] to prove Lemma 5.4. For simplicity, we will focus on the case when $u = 1$; the proof extends easily to the general case. The idea is, roughly speaking, a path joining 0 to a point far away from 0 with a ‘‘very short’’ passage time should cross ‘‘many’’ mesoscopic boxes which have ‘‘very short’’ box-crossing times (i.e., line-to-line passage times), and by estimates of box-crossing times in Lemma 5.6 and a counting argument we show that it is unlikely that such a path exists. Our notation in this subsection is analogous to that in [20] but with a slight difference to fit our context.

Step 1. Decomposition of a path into segments. For each $k = (k_1, k_2) \in \mathbb{Z}^2$ and integers M, N satisfying $M > N > 1$, define the boxes

$$S(k) := \{z \in \mathbb{R}^2 : Nk \leq z < N(k + (1, 1))\},$$

$$\widehat{S}(k) := \{z \in \mathbb{R}^2 : Nk - (M, M) \leq z < N(k + (1, 1)) + (M, M)\}.$$

Note that $\widehat{S}(k)$ contains $S(k)$ at its center. Later, we shall choose M such that it is much larger than N , but much smaller than n . In the following we assume that $p \in (p(10), p_c)$ and view the sites of $\eta\mathbb{T}$ as points in \mathbb{R}^2 . The boxes $S(k)$ ’s induce a partition of the sites of $\eta\mathbb{T}$, and will play the role of ‘‘renormalized sites’’.

Let $\gamma = (v(0), v(1), \dots, v(\xi))$ be a path in $\eta\mathbb{T}$ from $v(0) = 0$ to some site $v(\xi)$ in $\{(z_1, z_2) \in \mathbb{R}^2 : z_1 \geq n\}$. We associate to γ the following two sequences. First, let $k(0) = 0$ and $a(0) = 0$. Then let $v(a(1))$ be the first site along γ to be outside $\widehat{S}(k(0))$, and let $k(1)$ be the unique k such that $v(a(1)) \in S(k)$. Continue recursively to find sequences $(a(0), a(1), \dots, a(\tau))$ and $\sigma := (k(0), k(1), \dots, k(\tau))$ such that




FIG 9. The decomposition of a path γ into segments in Step 1. The black dots represent $v(a(i))$'s; the small crosses represent $k(i)$'s. In this picture, we have $\tilde{\sigma} = (l(0), \dots, l(\rho)) = (k(0), k(1), k(2), k(3), k(4), k(7), k(8), k(9))$, where $\tilde{\sigma}$ is the subsequence extracted from $\sigma = (k(0), \dots, k(\tau))$ by the loop removal process.

1. $0 = a(0) < a(1) < \dots < a(\tau) \leq \xi$,
2. $v(a(i)) \in S(k(i))$,
3. $a(i+1)$ is the smallest integer a larger than $a(i)$ such that $v(a) \notin \widehat{S}(k(i))$.

The final terms $a(\tau)$ and $k(\tau)$ satisfy

$$v(j) \in \widehat{S}(k(\tau)) \quad \text{if } a(\tau) \leq j \leq \xi.$$

By the classical “loop removal” process described in [20], we obtain a subsequence $\tilde{\sigma}$ of σ which is free of double points:

$$\tilde{\sigma} := (l(0), \dots, l(\rho)),$$

where $l(i) = k(j_i)$ for $i = 0, 1, \dots, \rho$ and $0 = j_0 < j_1 < \dots < j_\rho \leq \tau$. See Figure 9 for the construction. Note that although j_ρ and τ may not be equal, it is always true that $k(j_\rho) = k(\tau)$. By construction,

$$\|k(j+1) - k(j)\|_\infty \leq \frac{M}{N} + 1 \quad \text{for } j = 0, 1, \dots, \tau - 1.$$

This property is preserved by the loop removal process, in that

$$\|l(j+1) - l(j)\|_\infty \leq \frac{M}{N} + 1 \quad \text{for } j = 0, 1, \dots, \rho - 1.$$

Consider the portion $\gamma(i) := (v(a(i-1)), \dots, v(a(i)))$ of γ which stretches between $S(k(i-1))$ and $S(k(i))$, and define

$$L(i) := \|v(a(i)) - v(a(i-1))\|_\infty.$$

By construction,

$$M \leq L(i) < M + N + 1 \quad \text{for } 1 \leq i \leq \tau. \tag{72}$$

(A stronger version of (72) holds: $M \leq L(i) \leq M + N + \eta$ for $1 \leq i \leq \tau$.) Note that (72) is slightly different from the corresponding equation (3.5) “ $M \leq L(i) \leq M + N$ ” in [20] for a bond FPP on \mathbb{Z}^2 .

Step 2. Coloring of points and properties of white points. Let $0 < \epsilon < \nu/2$. For $1 \leq i \leq \rho$, consider the point $l(i) = k(j_i) \in \mathbb{Z}^2$ and the portion $\gamma(j_i)$ stretching between the two boxes $S(k(j_i - 1))$ and $S(k(j_i))$. We color $l(i)$ **white** if

$$T(\gamma(j_i)) \leq (\nu - 2\epsilon)L(j_i);$$

otherwise we color $l(i)$ **black**. Denote by w the number of white points in the sequence $(l(1), \dots, l(\rho))$. The next lemma gives estimates for w and ρ in terms of ϵ, n, M, N , and corresponds to Lemma 3.5 of [20].

Lemma 5.7 (Number of white points associated to a fast path). *Suppose that $p \in (p(10), p_c)$ and γ is a path in $\eta\mathbb{T}$ from 0 to some site in $\{(z_1, z_2) \in \mathbb{R}^2 : z_1 \geq n\}$. Suppose that ϵ, n, M, N and γ satisfy the following:*

- (i) ϵ is small: $0 < \epsilon < \nu/5$,
- (ii) M/N is large: $M(\nu - 3\epsilon) \geq (M + N + 1)(\nu - 4\epsilon)$,
- (iii) n is large: $n\epsilon \geq (M + 2N)(\nu - 4\epsilon)$,
- (iv) the passage time of γ is small: $T(\gamma) \leq n(\nu - 5\epsilon)$.

Then we have

$$w \geq \frac{\epsilon\rho}{2\nu} \quad \text{and} \quad \rho \geq \frac{n}{M + N} - 1.$$

Proof. Note that (i), (iii) and (iv) above are the same as those of Lemma 3.5 in [20], and (ii) is slightly different from the corresponding item of that lemma since (72) is slightly different from equation (3.5) in [20]. The proof of Lemma 5.7 is essentially the same as that of Lemma 3.5 in [20]. The details are omitted. \square

The following lemma is an analog of Lemma 3.6 in [20]. It says that a point is white implies that an annulus-crossing time is ‘‘short’’, and gives an upper bound on the probability that some given points are white, using the local dependence of white points.

Lemma 5.8 (Local dependence of white points). *Suppose that $p \in (p(10), p_c)$ and γ is a path in $\eta\mathbb{T}$ from 0 to some site in $\{(z_1, z_2) \in \mathbb{R}^2 : z_1 \geq n\}$. For any $1 \leq i \leq \rho$, the event $\{l(i) \text{ is white}\}$ is contained in the event*

$$\mathcal{E}^p(i) := \left\{ \begin{array}{l} \text{a site in } S(k(j_i)) \text{ is joined to a site outside the square} \\ \{z \in \mathbb{R}^2 : Nk(j_i) - (M - N, M - N) \leq z < Nk(j_i) + (M, M)\} \text{ by} \\ \text{a path in } \eta\mathbb{T} \text{ with passage time less than } (M + N + 1)(\nu - 2\epsilon) \end{array} \right\}.$$

For any subset \mathcal{S} of $\{l(1), \dots, l(\rho)\}$ we have

$$\mathbf{P}_p^\eta[\text{all points in } \mathcal{S} \text{ are white}] \leq \mathbf{P}_p^\eta[\mathcal{E}^p(i) \text{ occurs for each } l(i) \in \mathcal{S}] \leq \phi^{|\mathcal{S}|},$$

where

$$\phi = \phi(M, N, \epsilon, p) := \sup_{k \in \mathbb{Z}^2} \mathbf{P}_p^\eta \left[\begin{array}{l} \text{a site in } S(k) \text{ is joined to a site outside the square} \\ \{z \in \mathbb{R}^2 : Nk - (M - N, M - N) \leq z < Nk + (M, M)\} \text{ by} \\ \text{a path in } \eta\mathbb{T} \text{ with passage time less than } (M + N + 1)(\nu - 2\epsilon) \end{array} \right],$$

$$\text{and } \alpha = \alpha(M, N) := \left(\frac{N}{8(M+N)} \right)^2.$$

Proof. The proof is basically the same as for Lemma 3.6 in [20]. The details are omitted. \square

The next lemma corresponds to Lemma 3.7 in [20], with a slight modification adapted to our setting. It says that, the probability that a point is white can be made arbitrarily small by means of a suitable choice of N and M , uniformly in p close to p_c . This lemma is a key ingredient for the counting argument in the final proof.

Lemma 5.9 (Control of the probability that a point is white). *For each $\epsilon \in (0, \nu/5)$, $\delta > 0$ and $C > 2\nu/\epsilon$, there is an integer $N = N(\epsilon, \delta, C) \geq \nu/\epsilon$ and a $p_0 = p_0(\epsilon, \delta, C, N) \in (p(10), p_c)$, such that for each $p \in (p_0, p_c)$ and each integer $M \in [2\nu N/\epsilon, CN]$,*

$$\phi = \phi(M, N, \epsilon, p) \leq \delta.$$

Proof. The proof is analogous to the proof of Lemma 3.7 in [20]. Let B_1, B_2, B_3, B_4 denote the following boxes:

$$\begin{aligned} B_1 &= [-M + N, M] \times [N, M], & B_2 &= [N, M] \times [-M + N, M], \\ B_3 &= [-M + N, M] \times [-M + N, 0], & B_4 &= [-M + N, 0] \times [-M + N, M]. \end{aligned}$$

For each $k \in \mathbb{Z}^2$, if a site in $S(k)$ is joined to a site outside the square $\{z \in \mathbb{R}^2 : Nk - (M - N, M - N) \leq z < Nk + (M, M)\}$ by a path in $\eta\mathbb{T}$ with passage time not exceeding $(M + N + 1)(\nu - 2\epsilon)$, then one of the boxes $Nk + B_j$, $j \in \{1, 2, 3, 4\}$, is crossed between its longer sides by a path with passage time not exceeding $(M + N + 1)(\nu - 2\epsilon)$. Thus

$$\phi(M, N, \epsilon, p) \leq 4 \sup_{k \in \mathbb{Z}^2, \theta \in \{0, \pi/2\}} \mathbf{P}_p^\eta \left[l_{M-N, 2M-N}^{p, \theta}(k) \leq (M + N + 1)(\nu - 2\epsilon) \right].$$

If $0 < \epsilon < \nu/5$, $N \geq \nu/\epsilon$ and $M \geq 2\nu N/\epsilon$, then

$$2M - N \leq 3(M - N) \quad \text{and} \quad (M + N + 1)(\nu - 2\epsilon) \leq (M - N)(\nu - \epsilon),$$

giving that

$$\phi(M, N, \epsilon, p) \leq 4 \sup_{k \in \mathbb{Z}^2, \theta \in \{0, \pi/2\}} \mathbf{P}_p^\eta \left[l_{M-N, 3(M-N)}^{p, \theta}(k) \leq (M - N)(\nu - \epsilon) \right].$$

Applying Lemma 5.6 we obtain the desired result immediately. \square

Step 3. *A counting argument, end of the proof.* Now we are ready to prove Lemma 5.4 by using the preparatory lemmas in the previous step.

Proof of Lemma 5.4. We shall prove the lemma in the case $u = 1$; the proof extends easily to the general case by using the rotated lattice $u \cdot \mathbb{Z}^2$. We will follow the lines of the proof of Theorem 3.1 in [20], which is based on a counting argument.

Recall that each path γ in $\eta\mathbb{T}$ from 0 to a site in $\{(z_1, z_2) \in \mathbb{R}^2 : z_1 \geq n\}$ has an associated sequence $l(0), \dots, l(\rho)$. It is easy to see that the number of possible choices for this sequence is at most $(8(\frac{M}{N} + 1))^\rho$. Given the sequence of l 's, there are $\binom{\rho}{w}$ ways of choosing a set of cardinality w as the white points. Suppose that $\epsilon \in (0, \nu/5)$, and let δ be a fixed constant satisfying $\delta \in (0, 1)$ and $48\nu\delta\epsilon^3/(1152\nu^3) < \epsilon$. By Lemma 5.9, we can choose fixed $M, N \in \mathbb{N}$ and $p_0 \in (p(10), p_c)$ such that

$$N \geq \nu/\epsilon, \quad 2\nu N \leq M\epsilon \leq (3\nu - \epsilon)N \quad \text{and} \quad \phi(M, N, \epsilon, p) \leq \delta \quad \text{for all } p \in (p_0, p_c). \quad (73)$$

Then (i) and (ii) of Lemma 5.7 hold. Suppose also that n is large enough for (iii) of Lemma 5.7 to hold; by (i) and (iii), we have $n \geq M + 2N$, giving by Lemma 5.7 that $\rho \geq Kn$ where

$$K = K(M, N) := \frac{N}{(M + N)(M + 2N)}.$$

The point-to-line passage time $b_{0,n}^p$ is defined by

$$b_{0,n}^p := \inf \{T(\gamma)(\omega_p^\eta) : \gamma \text{ is a path in } \eta\mathbb{T} \text{ from } 0 \text{ to a site in } \{(z_1, z_2) \in \mathbb{R}^2 : z_1 \geq n\}\}.$$

Then, by (73), Lemmas 5.7 and 5.8, for all $p \in (p_0, p_c)$ and all n satisfying (iii) of Lemma 5.7, we have

$$\mathbf{P}_p^\eta[b_{0,n}^p < n(\nu - 5\epsilon)] \leq \sum_{\rho \geq Kn} \left(8 \left(\frac{M}{N} + 1\right)\right)^\rho \sum_{w \geq \rho/(2\nu)} \binom{\rho}{w} \delta^{\alpha w}, \quad (74)$$

where $\alpha = \alpha(M, N) = \left(\frac{N}{8(M+N)}\right)^2$. By an easy calculation (see the details at the end of the proof of Theorem 3.1 in [20]), we obtain from (74) that there are constants $K_1 > 0$ and $\delta_1 \in (0, 1)$, independent of n and p , such that

$$\mathbf{P}_p^\eta[b_{0,n}^p < n(\nu - 5\epsilon)] \leq K_1 \delta_1^n.$$

Hence, for each $p \in (p_0, p_c)$, we have $L(p)\mu(p) \geq \nu - 5\epsilon$ since $a_{0,m}/m$ tends to $\mu(p)$ \mathbf{P}_p -almost surely as $m \rightarrow \infty$ by (1). Letting $\epsilon \rightarrow 0$ yields $\liminf_{p \uparrow p_c} L(p)\mu(p) \geq \nu$. \square

5.3. Proof of Theorem 1.1

Finally, it is easy to derive Theorem 1.1 from Lemmas 5.1 and 5.4:

Proof of Theorem 1.1. Combining Lemmas 5.1 and 5.4, we obtain that for each $u \in \mathbb{U}$,

$$\lim_{p \uparrow p_c} L(p)\mu(p, u) = \nu. \quad (75)$$

It remains to prove that the convergence in (75) is uniform in $u \in \mathbb{U}$. By (4) (or (75)) and the fact that $\mu(p, z)$ is a norm on \mathbb{C} for each fixed $p \in (0, p_c)$ (see Section 1.1), there is a constant $C \geq 1$ such that

$$L(p)\mu(p, u) \leq C \quad \text{for all } p \in (0, p_c) \text{ and all } u \in \mathbb{U}. \quad (76)$$

Fix $\epsilon \in (0, 1)$. Let $K = K(\epsilon) := \lceil 2C\pi/\epsilon \rceil$. For $k \in \{1, 2, \dots, K\}$, let $u_k = u_k(\epsilon) := e^{2\pi ki/K}$. It follows from (75) that there is $p_0 = p_0(\epsilon) \in (0, p_c)$ such that for all $p \in (p_0, p_c)$ and all $k \in \{1, 2, \dots, K\}$,

$$|L(p)\mu(p, u_k) - \nu| \leq \epsilon/2. \quad (77)$$

It is obvious that for each $u \in \mathbb{U}$, there is $\tilde{u} \in \{u_1, \dots, u_K\}$ such that $|u - \tilde{u}| < \pi/K$. This, combined with (77), (76) and the fact that $\mu(p, z)$ is a norm on \mathbb{C} when $p \in (0, p_c)$, implies that for any $u \in \mathbb{U}$ with $u \notin \{u_1, \dots, u_K\}$ and $p \in (p_0, p_c)$, we have

$$\begin{aligned} L(p)\mu(p, u) &\leq L(p)\mu(p, \tilde{u}) + L(p)\mu(p, u - \tilde{u}) \\ &\leq \nu + \frac{\epsilon}{2} + |u - \tilde{u}|L(p)\mu\left(p, \frac{u - \tilde{u}}{|u - \tilde{u}|}\right) \leq \nu + \epsilon, \\ L(p)\mu(p, u) &\geq L(p)\mu(p, \tilde{u}) - L(p)\mu(p, \tilde{u} - u) \\ &\geq \nu - \frac{\epsilon}{2} - |\tilde{u} - u|L(p)\mu\left(p, \frac{\tilde{u} - u}{|\tilde{u} - u|}\right) \geq \nu - \epsilon. \end{aligned}$$

These two inequalities combined with (77) implies that for each $\epsilon \in (0, 1)$, there is $p_0 \in (0, p_c)$ such that

$$|L(p)\mu(p, u) - \nu| \leq \epsilon \quad \text{for all } p \in (p_0, p_c) \text{ and all } u \in \mathbb{U}.$$

Theorem 1.1 follows from this immediately. \square

Acknowledgments

The author is especially grateful to an anonymous referee for 1) pointing out that a preliminary version of the proof of Theorem 1.1 can be simplified, without using the scaling limit of the collection of portions of clusters in a strip, and 2) suggesting us another definition of $\mathcal{C}(z)$ which makes the proof cleaner. The referee's suggestions make the paper less technical and much shorter than its preprint version [41]. The author thanks Geoffrey Grimmett for an invitation to the Statistical Laboratory in Cambridge University, and thanks the hospitality of the Laboratory, where this project was initiated. The author was supported by the National Key R&D Program of China (No. 2020YFA0712700), the National Natural Science Foundation of China (No. 12288201 and No. 11601505) and the Key Laboratory of Random Complex Structures and Data Science, CAS (No. 2008DP173182).

References

- [1] Ahlberg, D., Steif, J.E.: Scaling limits for the threshold window: When does a monotone Boolean function flip its outcome? *Ann. Inst. H. Poincaré Probab. Statist.* **53**, 2135–2161 (2017)
- [2] Auffinger, A., Damron, M., Hanson, J.: 50 years of first-passage percolation. *University Lecture Series Vol. 68*, American Mathematical Society, (2017)
- [3] Beffara, V., Nolin, P.: On monochromatic arm exponents for 2D critical percolation. *Ann. Probab.* **39**, 1286–1304 (2011)
- [4] van den Berg, J., Conijn, R.: The gaps between the sizes of large clusters in 2D critical percolation. *Electron. Commun. Probab.* **18**, no. 92, 1–9 (2013)
- [5] van den Berg, J., Kiss, D., Nolin, P.: Two-dimensional volume-frozen percolation: deconcentration and prevalence of mesoscopic clusters. *Annales Scientifiques de l'École Normale Supérieure* **51**, 1017–1084 (2018)

- [6] van den Berg, J., Nolin, P.: Near-critical percolation with heavy-tailed impurities, forest fires and frozen percolation. *Probab. Theory Relat. Fields.* **181**, 211–290 (2021)
- [7] Bollobás, B., Riordan, O. *Percolation*. Cambridge University Press, New York (2006)
- [8] Chayes, J.T., Chayes, L., Durrett, R.: Critical behavior of the two-dimensional first passage time. *Journal of Stat. Physics* **45**, 933–951 (1986)
- [9] Camia, F., Conijn, R., Kiss, D.: Conformal measure ensembles for percolation and the FK-Ising model. *Sojourns in Probability Theory and Statistical Physics-II Brownian Web and Percolation, A Festschrift for Charles M. Newman*. Springer Proceedings in Mathematics & Statistics, Volume **299**, 44–89 (2019)
- [10] Camia, F., Newman, C.M.: Critical percolation: the full scaling limit. *Commun. Math. Phys.* **268**, 1–38 (2006)
- [11] Cox, J.T., Kesten, H.: On the continuity of the time constant of first-passage percolation. *J. Appl. Probab.* **18**, 809–819 (1981)
- [12] Damron, M., Hanson, J., Lam, W.-K.: Universality of the time constant for 2D critical first-passage percolation. arXiv:1904.12009v1 (2019). To appear in *Ann. Appl. Probab.*
- [13] Duminil-Copin, H.: Limit of the Wulff Crystal when approaching criticality for site percolation on the triangular lattice. *Electron. Commun. Probab.* **18**, no. 93, 1–9 (2013)
- [14] Durrett, R.: *Probability: Theory and Examples*, 4th edition. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge (2010)
- [15] Garban, C., Pete, G., Schramm, O.: Pivotal, cluster and interface measures for critical planar percolation. *J. Amer. Math. Soc.* **26**, 939–1024 (2013)
- [16] Garban, C., Pete, G., Schramm, O.: The scaling limits of near-critical and dynamical percolation. *J. Eur. Math. Soc.* **20**, 1195–1268 (2018)
- [17] Garban, C., Pete, G., Schramm, O.: The scaling limits of the minimal spanning tree and invasion percolation in the plane. *Ann. Probab.* **46**, 3501–3557 (2018)
- [18] Garett, O., Marchand, R.: Large deviations for the chemical distance in supercritical Bernoulli percolation. *Ann. Probab.* **35**, 833–866 (2007)
- [19] Grimmett, G.: *Percolation*, 2nd ed. Springer-Verlag Berlin (1999)
- [20] Grimmett, G., Kesten, H.: First-passage percolation, network flows and electrical resistances. *Z. Wahrsch. Verw. Gebiete* **66**, 335–366 (1984)
- [21] Hammersley, J.M., Welsh, D.J.A.: First-passage percolation, subadditive processes, stochastic networks, and generalized renewal theory. In *Bernoulli, Bayes, Laplace Anniversary Volume* (J. Neyman and L. LeCam, eds.) 61–110. Springer, Berlin. (1965)
- [22] Holden, N., Sun, X.: Convergence of uniform triangulations under the Cardy embedding. arXiv:1905.13207v2 (2020) To appear in *Acta Math.*
- [23] Jiang, J., Yao, C.-L.: Critical first-passage percolation starting on the boundary. *Stoch. Proc. Appl.* **129**, 2049–2065 (2019)
- [24] Kesten, H.: Aspects of first passage percolation. In *Lecture Notes in Math.*, Vol 1180, pp. 125–264 Berlin: Springer. (1986)
- [25] Kesten, H.: Scaling relations for 2D-percolation. *Comm. Math. Phys.* **109**, 109–156 (1987)
- [26] Kesten, H.: First-passage percolation. From classical to modern probability, *Progr. Probab.*, vol. 54, pp. 93–143. Birkhäuser, Basel (2003)
- [27] Kesten, H., Zhang, Y.: A central limit theorem for “critical” first-passage percolation in two-dimensions. *Probab. Theory Relat. Fields.* **107**, 137–160 (1997)
- [28] Lawler, G.F., Schramm, O., Werner, W.: One-arm exponent for critical 2D percolation. *Electron. J. Probab.* **7**, no. 2, 1–13 (2002)
- [29] Liggett, T.M.: An improved subadditive ergodic theorem. *Ann. Probab.* **13**, 1279–1285 (1985)
- [30] Nolin, P.: Near critical percolation in two-dimensions. *Electron. J. Probab.* **13**, 1562–1623 (2008)
- [31] Reimer, D.: Proof of the van den Berg-Kesten conjecture. *Combin. Probab. Comput.* **9**, 27–32 (2000)
- [32] Sheffield, S.: Exploration trees and conformal loop ensembles. *Duke Math. J.* **147**, 79–129 (2009)
- [33] Sheffield S., Werner W.: Conformal loop ensembles: the Markovian characterization and the loop-soup construction. *Ann. Math.* **176**, 1827–1917 (2012)
- [34] Schramm, O., Smirnov, S.: On the scaling limits of planar percolation (with an appendix by C. Garban). *Ann. Probab.* **39**, 1768–1814 (2011)
- [35] Smirnov, S., Werner, W.: Critical exponents for two-dimensional percolation. *Math. Res. Lett.* **8**, 729–744 (2001)
- [36] Sun, N.: Conformally invariant scaling limits in planar critical percolation. *Probab. Surveys* **8**, 155–209 (2011)
- [37] Werner, W.: Lectures on two-dimensional critical percolation. In: *Statistical mechanics*. IAS/Park City Math. Ser., Vol.16. Amer. Math. Soc., Providence, RI, 297–360 (2009)
- [38] Yao, C.-L.: Law of large numbers for critical first-passage percolation on the triangular lattice. *Electron. Commun. Probab.* **19**, no. 18, 1–14 (2014)
- [39] Yao, C.-L.: Limit theorems for critical first-passage percolation on the triangular lattice. *Stoch. Proc. Appl.* **128**, 445–460 (2018)
- [40] Yao, C.-L.: Asymptotics for 2D critical and near-critical first-passage percolation. *Probab. Theory Relat. Fields.* **175**, 975–1019 (2019)
- [41] Yao, C.-L.: Convergence of limit shapes for 2D near-critical first-passage percolation (original preprint version of this paper). arXiv:2104.01211v2