

SCREENING FOR BREAKTHROUGHS: OMITTED PROOFS

Gregorio Curello Ludvig Sinander
University of Bonn University of Oxford

21 February 2024

Abstract

This document contains all proofs omitted from our working paper ‘Screening for breakthroughs’ (Curello & Sinander, 2024); specifically, the February 2024 version of the paper ([version 8 on arXiv](#)).

Section 7.1 (p. 25)

The two utility possibility frontiers may be written as

$$F^0(u) = u - \lambda\phi^{-1}(u)$$

and $F^1(u) = u + \lambda \max_{L \geq 0} \{wL - \phi^{-1}(u + \kappa(L))\}.$

Proof of Lemma 1. Clearly F^0, F^1 are well-defined and continuous on $[0, \infty)$. It remains to show that they are strictly concave with peaks u^0, u^1 satisfying $u^1 < u^0$, and that the gap $F^1 - F^0$ is strictly decreasing.

Claim. For each $u > 0$, the maximisation problem in the expression for $F^1(u)$ has a unique solution $L^*(u) \in (0, \infty)$. Furthermore, $\lim_{u \rightarrow \infty} L^*(u) = 0$.

Proof. Fix $u > 0$, and write $f : \mathbf{R}_+ \rightarrow \mathbf{R}$ for the objective. Its (right-hand) derivative is

$$f'(L) = w - \frac{\kappa'(L)}{\phi'(\phi^{-1}(u + \kappa(L)))}.$$

Clearly f' is strictly decreasing, so there is at most one maximiser. We have $f'(0) = w > 0$, whereas $f'(L) < 0$ for any large enough $L > 0$ since

$$\lim_{L \rightarrow \infty} f'(L) \leq w - \lim_{L \rightarrow \infty} \frac{\kappa'(L)}{\phi'(\phi^{-1}(\kappa(L)))} = -\infty,$$

where the equality holds since the numerator is bounded away from zero as $L \rightarrow \infty$ while the denominator vanishes. Thus the unique maximiser $L^*(u)$ of f is interior, and therefore satisfies the first-order condition $f'(L^*(u)) = 0$. As $u \rightarrow \infty$ in the first-order condition, the denominator in the fraction vanishes since $\lim_{C \rightarrow \infty} \phi'(C) = 0$, requiring the numerator also to vanish, which since $\kappa' > 0$ on $(0, \infty)$ demands that $L^*(u) \rightarrow 0$. \square

F^0 is strictly concave since ϕ is. Its (right-hand) derivative

$$F^{0'}(u) = 1 - \frac{\lambda}{\phi'(\phi^{-1}(u))}$$

is strictly positive at $u = 0$ since $\lim_{C \rightarrow 0} \phi'(C) = \infty$, and is strictly negative for u large enough since $\phi^{-1}(u) \rightarrow \infty$ as $u \rightarrow \infty$ and $\lim_{C \rightarrow \infty} \phi'(C) = 0$. Thus F^0 is uniquely maximised at $u^0 \in (0, \infty)$ satisfying the first-order condition

$$\phi'(\phi^{-1}(u^0)) = \lambda.$$

F^1 is also strictly concave: for $u \neq u'$ in $[0, \infty)$ and $\eta \in (0, 1)$, we have

$$\begin{aligned} F^1(\eta u + (1 - \eta)u') &= \eta u + (1 - \eta)u' + \lambda \max_{L \geq 0} \{wL - \phi^{-1}(\eta u + (1 - \eta)u' + \kappa(L))\} \\ &> \eta u + (1 - \eta)u' \\ &\quad + \lambda \max_{L \geq 0} \{wL - \eta \phi^{-1}(u + \kappa(L)) - (1 - \eta) \phi^{-1}(u' + \kappa(L))\} \\ &\geq \eta \left[u + \lambda \max_{L \geq 0} \{wL - \phi^{-1}(u + \kappa(L))\} \right] \\ &\quad + (1 - \eta) \left[u' + \lambda \max_{L \geq 0} \{wL - \phi^{-1}(u' + \kappa(L))\} \right] \\ &= \eta F^1(u) + (1 - \eta) F^1(u'), \end{aligned}$$

where the strict inequality holds since $-\phi^{-1}$ is strictly concave. By the envelope theorem, we have

$$F^{1'}(u) = 1 - \frac{\lambda}{\phi'(\phi^{-1}(u + \kappa(L^*(u))))}.$$

This expression is strictly negative for u large enough because the denominator in the fraction vanishes as $u \rightarrow \infty$ since $L^*(u) \rightarrow 0$ by the claim. Thus F^1 has a unique maximiser $u^1 \in [0, \infty)$, which satisfies the first-order condition

$$\phi'(\phi^{-1}(u^1 + \kappa(L^*(u^1)))) \leq \lambda, \quad \text{with equality if } u^1 > 0.$$

To show that $u^0 > u^1$, consider two cases. If $u^1 = 0$, then $u^0 > 0 = u^1$. If instead $u^1 > 0$, then the first-order conditions for u^0 and u^1 together yield

$$\phi'(\phi^{-1}(u^0)) = \lambda = \phi'(\phi^{-1}(u^1 + \kappa(L^*(u^1)))).$$

Thus

$$u^0 = u^1 + \kappa(L^*(u^1)) > u^1$$

since $L^*(u^1) > 0$ by the claim and $\kappa > 0$ on $(0, \infty)$.

It remains only to show that $F^1 - F^0$ is strictly decreasing, so that $u^* = 0$. It suffices to show that $F^{1'} < F^{0'}$ on $(0, \infty)$. So fix any $u > 0$. Since $L^*(u) > 0$ by the claim, and $\kappa > 0$ on $(0, \infty)$, we have

$$F^{0'}(u) = 1 - \frac{\lambda}{\phi'(\phi^{-1}(u))} > 1 - \frac{\lambda}{\phi'(\phi^{-1}(u + \kappa(L^*(u))))} = F^{1'}(u). \quad \blacksquare$$

Supplemental appendix I.1 (p. 53)

To formalise and prove the trichotomy asserted in footnote 81 (p. 53), consider frontiers F^0, F^1 satisfying all of our model assumptions, except for the requirement that u^* be a strict local maximum of $F^1 - F^0$.

Definition 1. $\bar{u} \in (0, \infty)$ is a *saddle point* of a continuous function ψ defined on $[0, \infty)$ iff both

- (a) for all $\varepsilon > 0$, there are $u, u' \in (0, \infty)$ with $\bar{u} - \varepsilon < u < \bar{u} < u' < \bar{u} + \varepsilon$ such that $|\psi(u') - \psi(u)|/(u' - u) < \varepsilon$,¹ and
- (b) \bar{u} is neither a local maximum nor a local minimum of ψ .

Claim. If $u^* > 0$, then either (i) u^* is a local maximum of $F^1 - F^0$, (ii) u^* is a saddle point of $F^1 - F^0$, or (iii) both F^0 and F^1 are kinked at u^* .

Assuming that u^* is a strict local maximum of $F^1 - F^0$ thus amounts to ruling out the saddle-point case (ii), the (pathological) mutual-kink case (iii), and the case in which $F^1 - F^0$ is constant on a left-neighbourhood of u^* .

¹In case ψ is differentiable at \bar{u} , condition (a) reduces to the requirement that $\psi'(\bar{u}) = 0$. Our definition thus extends the usual definition of a saddle point, which applies only to differentiable functions.

Proof. Assume that $u^* > 0$ and that neither (i) nor (ii) holds; we shall deduce (iii). By its definition, u^* cannot be a local minimum of $F^1 - F^0$. Then by definition of ‘saddle point’, there must be an $\varepsilon > 0$ such that

$$\frac{|(F^1 - F^0)(u') - (F^1 - F^0)(u)|}{u' - u} \geq \varepsilon$$

for all $u, u' \in (0, \infty)$ satisfying $u^* - \varepsilon < u < u^* < u' < u^* + \varepsilon$. Since $F^1 - F^0$ is strictly decreasing on $[u^*, u^0]$, the expression in the numerator must be strictly negative for u sufficiently close to u^* , and thus $F^{1+}(u^*) \leq F^{0+}(u^*) - \varepsilon < F^{0+}(u^*)$ and (similarly) $F^{1-}(u^*) < F^{0-}(u^*)$. By definition of u^* , the frontiers F^0, F^1 share a supergradient $\eta \in \mathbf{R}$ at u^* , so that $F^{j+}(u^*) \leq \eta \leq F^{j-}(u^*)$ for both $j \in \{0, 1\}$. Then

$$F^{1+}(u^*) < F^{0+}(u^*) \leq \eta \leq F^{1-}(u^*) < F^{0-}(u^*),$$

implying that both F^0 and F^1 are kinked at u^* . ■

Supplemental appendix I.5 (p. 56)

For $u \in [0, \infty)$, write M_u for the space of all maps $\widehat{X} : \mathcal{F} \rightarrow [0, \infty)$ that satisfy $\mathbf{E}(\widehat{X}(\mathbf{F})) = u$.

Proof of Lemma 8. For concavity, take $u, u^\dagger \in [0, \infty)$ and $\lambda \in (0, 1)$. Let $\widehat{X} \in M_u$ and $\widehat{X}^\dagger \in M_{u^\dagger}$ be maximisers:

$$\mathbf{E}(\mathbf{F}(\widehat{X}(\mathbf{F}))) = F^1(u) \quad \text{and} \quad \mathbf{E}(\mathbf{F}(\widehat{X}^\dagger(\mathbf{F}))) = F^1(u^\dagger).$$

Then $\lambda\widehat{X} + (1 - \lambda)\widehat{X}^\dagger$ belongs to $M_{\lambda u + (1 - \lambda)u^\dagger}$, so that

$$\begin{aligned} & F^1(\lambda u + (1 - \lambda)u^\dagger) \\ & \geq \mathbf{E}(\mathbf{F}(\lambda\widehat{X}(\mathbf{F}) + (1 - \lambda)\widehat{X}^\dagger(\mathbf{F}))) \\ & \geq \mathbf{E}(\lambda\mathbf{F}(\widehat{X}(\mathbf{F})) + (1 - \lambda)\mathbf{F}(\widehat{X}^\dagger(\mathbf{F}))) \quad \text{since } \mathbf{F} \text{ is concave a.s.} \\ & = \lambda F^1(u) + (1 - \lambda)F^1(u^\dagger). \end{aligned}$$

To see that F^1 attains a maximum at u^1 , observe that F^1 is bounded above by $\mathbf{E}(\mathbf{F}(U^1(\mathbf{F})))$, and that it attains this value at u^1 . The peak is

unique because for $u \neq u^1$, any $\widehat{X} \in M_u$ must have $\widehat{X} \neq U^1$ on a non-null set, and thus

$$\mathbf{F}(\widehat{X}(\mathbf{F})) \leq (<) \mathbf{F}(U^1(\mathbf{F})) \quad \text{a.s. (with positive probability).}$$

For upper semi-continuity, observe that since F^1 is concave, it is continuous on the interior of its effective domain, which is a convex set that contains u^1 . F^1 is trivially continuous off the closure of its effective domain, where it is constant and equal to $-\infty$. It remains only to establish that

$$\limsup_{u' \rightarrow u} F^1(u') \leq F^1(u)$$

for u on the boundary of the effective domain (there are at most two such u). It suffices to show for an arbitrary decreasing sequence $(u_n)_{n \in \mathbf{N}}$ in the interior of the effective domain converging to some $u \in [0, u^1]$ that

$$\lim_{n \rightarrow \infty} F^1(u_n) \leq F^1(u)$$

(where the limit exists since $(F^1(u_n))_{n \in \mathbf{N}}$ is eventually monotone), and similarly for increasing sequences converging to $u \geq u^1$. We show the former, omitting the analogous argument for the latter.

For each $n \in \mathbf{N}$, let $\widehat{X}_n \in M_{u_n}$ be a maximiser at u_n :

$$\mathbf{E}(\mathbf{F}(\widehat{X}_n(\mathbf{F}))) = F^1(u_n).$$

Since $\bigcup_{u' \in [0, u^1]} M_{u'}$ is compact (because bounded), the sequence $(\widehat{X}_n)_{n \in \mathbf{N}}$ admits a convergent subsequence $(\widehat{X}_{n_k})_{k \in \mathbf{N}}$, whose limit we denote by \widehat{X} . We have

$$\begin{aligned} F^1(u) &\geq \mathbf{E}(\mathbf{F}(\widehat{X}(\mathbf{F}))) && \text{since } \widehat{X} \in M_u \\ &\geq \mathbf{E}\left(\lim_{k \rightarrow \infty} \mathbf{F}(\widehat{X}_{n_k}(\mathbf{F}))\right) && \text{since } \mathbf{F} \text{ is upper semi-continuous a.s.} \\ &= \lim_{k \rightarrow \infty} \mathbf{E}(\mathbf{F}(\widehat{X}_{n_k}(\mathbf{F}))) && \text{by bounded convergence} \\ &= \lim_{k \rightarrow \infty} F^1(u_{n_k}) \\ &= \lim_{n \rightarrow \infty} F^1(u_n) && \text{since } (F^1(u_n))_{n \in \mathbf{N}} \text{ is convergent.} \quad \blacksquare \end{aligned}$$

Supplemental appendix M (p. 65)

To prove Lemma 14 and the Gateaux lemma, we shall use the following standard integration-by-parts result:²

IBP lemma. Let ν be a finite measure on \mathbf{R}_+ , and let L be a ν -integrable function $\mathbf{R}_+ \rightarrow \mathbf{R}$ satisfying $L(t) = L(0) + \int_0^t l$ for some (Lebesgue-)integrable $l : \mathbf{R}_+ \rightarrow \mathbf{R}$. Then

$$\int_{[0,T]} L d\nu = L(T)\nu([0, T]) - \int_0^T \nu([0, t])l(t)dt \quad \text{for every } T \in \mathbf{R}_+.$$

1 Proof of Lemma 14 (p. 66)

Define $\Phi : \mathbf{R}_+ \rightarrow [0, \infty]$ by

$$\Phi(t) := r \int_0^t e^{-rs} \phi^0(s) ds.$$

We first show that $\mathbf{E}_G(\Phi(\tau)) < \infty$, so that Φ is G -integrable. Note that

$$\begin{aligned} \mathbf{E}_G \left(r \int_0^\tau \frac{e^{-rt}}{1-G(t)} dt \right) &= \lim_{T \rightarrow \infty} \int_{[0,T]} \left(r \int_0^s \frac{e^{-rt}}{1-G(t)} dt \right) G(ds) \\ &= \lim_{T \rightarrow \infty} r \int_0^T e^{-rt} \frac{G(T) - G(t)}{1-G(t)} dt = r \int_0^\infty e^{-rt} dt = 1, \end{aligned}$$

where the first and third equalities hold by monotone convergence, and the second follows from the IBP lemma with ν the measure associated with G and

$$l(t) := \begin{cases} re^{-rt}/[1-G(t)] & \text{for } t < T \\ 0 & \text{for } t \geq T \end{cases} \quad \text{if } G(T) < 1$$

and $l(t) := re^{-rt}$ if $G(T) = 1$. Thus since (x, X) satisfies the Euler equation with ϕ^0, ϕ^1 , (E) yields

$$\begin{aligned} \mathbf{E}_G(\Phi(\tau)) &= \mathbf{E}_G \left(r \int_0^\tau e^{-rt} \frac{-\int_{[0,t]} \phi^1 dG}{1-G(t)} dt \right) \\ &\leq \mathbf{E}_G \left(r \int_0^\tau e^{-rt} \frac{\int_{\mathbf{R}_+} |\phi^1| dG}{1-G(t)} dt \right) = \mathbf{E}_G(|\phi^1(\tau)|) < \infty, \end{aligned}$$

where the final inequality holds since ϕ^1 is G -integrable.

²See e.g. Theorem 18.4 in Billingsley (1995, p. 236).

To show that x belongs to \mathcal{X}_G , fix a $u \in (0, u^0)$; we shall prove that the maps $\psi_{x,u}^0$ and $\psi_{X,u}^1$ (defined on p. 66) are G -integrable. For the former, let \mathcal{T} be the set of $t \in \mathbf{R}_+$ at which $x_t \geq u$. The map $t \mapsto F^{0'}(x_t, u)$ is bounded on \mathcal{T} (by zero below and by $F^{0+}(u)$ above) since $x \leq u^0$ and F^0 is increasing and concave, so it suffices to show that

$$\varphi(t) := r \int_0^t e^{-rs} F^{0'}(x_s, u) \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}}(s) ds$$

is G -integrable. Since (x, X) satisfies the Euler equation, $\phi^0(t)$ is a supergradient of F^0 at x_t for a.e. $t \in \mathbf{R}_+$ such that $G(t) < 1$. Thus since $x < u$ on $\mathbf{R}_+ \setminus \mathcal{T}$ and F^0 is concave, we have

$$\phi^0(t) \geq F^{0+}(x_t) = F^{0'}(x_t, u) \quad \text{for a.e. } t \in \mathbf{R}_+ \setminus \mathcal{T} \text{ with } G(t) < 1,$$

so that

$$\varphi(t) \leq r \int_0^t e^{-rs} \phi^0(s) \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}}(s) ds \leq \Phi(t) \quad \text{for all } t \in \mathbf{R}_+ \text{ with } G(t) < 1.$$

Since φ and Φ are continuous and the former is non-negative, it follows that $0 \leq \varphi \leq \Phi$ on the support of G , so that φ is G -integrable since Φ is.

It remains to show that $\psi_{X,u}^1$ is G -integrable. Choose $\varepsilon \in (0, u \wedge [u^0 - u])$ so that $\varepsilon < u^1 \wedge (u^0 - u^1)$ if $u^1 > 0$, and let

$$\mathcal{T}' := \begin{cases} \{t \in \mathbf{R}_+ : X_t < u + \varepsilon\} & \text{if } u^1 = 0 \\ \{t \in \mathbf{R}_+ : (u \wedge u^1) - \varepsilon < X_t < (u \vee u^1) + \varepsilon\} & \text{if } u^1 > 0. \end{cases}$$

We will show that $t \mapsto \psi_{X,u}^1(t) = e^{-rt} F^{1'}(X_t, u)$ is bounded on \mathcal{T}' and G -integrable on $\mathbf{R}_+ \setminus \mathcal{T}'$. The former follows from the fact that

$$F^1 \text{ is Lipschitz continuous on } \begin{cases} [0, u + \varepsilon] & \text{if } u^1 = 0 \\ [(u \wedge u^1) - \varepsilon, (u \vee u^1) + \varepsilon] & \text{if } u^1 > 0, \end{cases}$$

so that $t \mapsto F^{1'}(X_t, u)$ is bounded on \mathcal{T}' . For the latter, note that (by definition of \mathcal{T}' ,) every $t \in \mathbf{R}_+ \setminus \mathcal{T}'$ has either $X_t < u, u^1$ or $X_t > u, u^1$. Since (x, X) satisfies the Euler equation, $\phi^1(t)$ is a supergradient of F^1 at X_t for G -a.e. $t \in \mathbf{R}_+$. Thus for G -a.e. $t \in \mathbf{R}_+ \setminus \mathcal{T}'$, we have

$$\begin{aligned} \text{either } & F^{1'}(X_t, u) = F^{1+}(X_t) \quad \text{and} \quad 0 < F^{1+}(X_t) \leq \phi^1(t) \\ \text{or } & F^{1'}(X_t, u) = F^{1-}(X_t) \quad \text{and} \quad 0 > F^{1-}(X_t) \geq \phi^1(t) \end{aligned}$$

by the concavity of F^0 , so that

$$\left| \psi_{X,u}^1(t) \right| \leq \left| F^{1'}(X_t, u) \right| \leq \left| \phi^1(t) \right| \quad \text{for } G\text{-a.e. } t \in \mathbf{R}_+ \setminus \mathcal{T}',$$

which implies that $\psi_{X,u}^1$ is G -integrable on $\mathbf{R}_+ \setminus \mathcal{T}'$ since ϕ^1 is. ■

2 Proof of the claim in the proof of Lemma 15 (p. 68)

For $\alpha \in (0, 1)$, define $h_\alpha^0, h_\alpha^1 : \mathbf{R}_+ \rightarrow (-\infty, \infty]$ by

$$h_\alpha^0(t) := re^{-rt} \frac{F^0(x_t + \alpha[u - x_t]) - F^0(x_t)}{\alpha}$$

and $h_\alpha^1(t) := e^{-rt} \frac{F^1(X_t + \alpha[u - X_t]) - F^1(X_t)}{\alpha}.$

Clearly both are measurable for each $\alpha \in (0, 1)$. By definition, we have

$$D\pi_G(x, x^\dagger - x) = \lim_{\alpha \downarrow 0} \left[\mathbf{E}_G \left(\int_0^\tau h_\alpha^0 \right) + \mathbf{E}_G \left(h_\alpha^1(\tau) \right) \right];$$

we must show that the limit exists and satisfies the asserted inequality.

As $\alpha \downarrow 0$, h_α^0 and h_α^1 converge pointwise to $h^0 : \mathbf{R}_+ \rightarrow (-\infty, \infty]$ and $h^1 : \mathbf{R}_+ \rightarrow [-\infty, \infty]$, respectively, given by

$$h^0(t) := re^{-rt} F^{0'}(x_t, u)(u - x_t) \quad \text{and} \quad h^1(t) := e^{-rt} F^{1'}(X_t, u)(u - X_t).$$

We shall show that

- (a) $t \mapsto \int_0^t h^0 \mathbf{1}_\mathcal{T}$ and $h^1 \mathbf{1}_{\mathcal{T}'}$ are G -integrable,
- (b) $\mathbf{E}_G(\int_0^\tau h^0)$ and $\mathbf{E}_G(h^1(\tau))$ exist and are $> -\infty$, and
- (c) $\mathbf{E}_G(\int_0^\tau h_\alpha^0) \rightarrow \mathbf{E}_G(\int_0^\tau h^0)$ and $\mathbf{E}_G(h_\alpha^1(\tau)) \rightarrow \mathbf{E}_G(h^1(\tau))$ as $\alpha \downarrow 0$.

This suffices since then $C := \mathbf{E}_G(\int_0^\tau h^0 \mathbf{1}_\mathcal{T}) + \mathbf{E}_G(h^1(\tau) \mathbf{1}_{\mathcal{T}'})$ is finite,

$$\begin{aligned} D\pi_G(x, x^\dagger - x) &= \mathbf{E}_G \left(\int_0^\tau h^0 \right) + \mathbf{E}_G \left(h^1(\tau) \right) \\ &= \mathbf{E}_G \left(\int_0^\tau h^0 \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}} \right) + \mathbf{E}_G \left(h^1(\tau) \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}'} \right) + C, \end{aligned}$$

and moreover any $t \in \mathbf{R}_+ \setminus \mathcal{T}$ has (by definition of \mathcal{T})

$$u - x_t \geq \varepsilon \quad \text{and thus} \quad h^0(t) \geq re^{-rt} F^{0'}(x_t, u)\varepsilon = re^{-rt} |F^{0'}(x_t, u)|\varepsilon,$$

while any $t \in \mathbf{R}_+ \setminus \mathcal{T}'$ has

$$\begin{aligned} &\text{either} \quad u - X_t \geq \varepsilon' \quad \text{and} \quad X_t < u^1 \quad (\text{hence } F^{1'}(X_t, u) > 0) \\ &\text{or (if } u^1 > 0) \quad u - X_t \leq \varepsilon' \quad \text{and} \quad X_t > u^1 \quad (\text{hence } F^{1'}(X_t, u) < 0) \end{aligned}$$

by definition of \mathcal{T}' , so that $h^1(t) \geq e^{-rt} |F^{1'}(X_t, u)|\varepsilon'$.

To obtain (a)–(c) for h^0 , observe that F^0 is K -Lipschitz on $[u - \varepsilon, u^0]$ for some $K > 0$ since it is concave and increasing on $[0, u^0]$. Thus

$$|h_\alpha^0(t)| \leq re^{-rt}K|x_t - u| \leq re^{-rt}Ku^0 \quad \text{for all } \alpha \in (0, 1) \text{ and } t \in \mathcal{T},$$

which implies that

$$\left| \int_0^t h_\alpha^0 \mathbf{1}_{\mathcal{T}} \right| \leq \int_0^t |h_\alpha^0| \mathbf{1}_{\mathcal{T}} \leq Ku^0 r \int_0^t e^{-rt} \mathbf{1}_{\mathcal{T}} \leq Ku^0 \quad \text{for all } \alpha \in (0, 1).$$

Thus applying the dominated convergence theorem twice yields

$$\lim_{\alpha \downarrow 0} \mathbf{E}_G \left(\int_0^\tau h_\alpha^0 \mathbf{1}_{\mathcal{T}} \right) = \mathbf{E}_G \left(\lim_{\alpha \downarrow 0} \int_0^\tau h_\alpha^0 \mathbf{1}_{\mathcal{T}} \right) = \mathbf{E}_G \left(\int_0^\tau h^0 \mathbf{1}_{\mathcal{T}} \right) \in \mathbf{R},$$

giving us property (a).

Meanwhile, $h_\alpha^0 \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}}$ is (pointwise) decreasing in α since F^0 is concave, and is non-negative since F^0 is increasing on $[0, u^0]$ and $x \leq u < u^0$ on $\mathbf{R}_+ \setminus \mathcal{T}$. Clearly the same is true of $t \mapsto \int_0^t h_\alpha^0 \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}}$. Thus applying the monotone convergence theorem twice yields

$$0 \leq \lim_{\alpha \downarrow 0} \mathbf{E}_G \left(\int_0^\tau h_\alpha^0 \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}} \right) = \mathbf{E}_G \left(\lim_{\alpha \downarrow 0} \int_0^\tau h_\alpha^0 \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}} \right) = \mathbf{E}_G \left(\int_0^\tau h^0 \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}} \right),$$

where the rightmost term is well-defined. Thus h^0 satisfies (b) and (c).

To derive (a)–(c) for h^1 , observe that

$$F^1 \text{ is Lipschitz continuous on } \begin{cases} [0, u + \varepsilon'] & \text{if } u^1 = 0 \\ [(u \wedge u^1) - \varepsilon', (u \vee u^1) + \varepsilon'] & \text{if } u^1 > 0 \end{cases}$$

since it is concave and maximised at u^0 . Hence the family $(h_\alpha^1)_{\alpha \in (0, 1)}$ is uniformly bounded on \mathcal{T}' , so that

$$\lim_{\alpha \downarrow 0} \mathbf{E}_G \left(h_\alpha^1(\tau) \mathbf{1}_{\mathcal{T}'(\tau)} \right) = \mathbf{E}_G \left(h^1(\tau) \mathbf{1}_{\mathcal{T}'(\tau)} \right) \in \mathbf{R} \quad (1)$$

by bounded convergence. Thus h^1 satisfies property (a).

Define $\bar{\alpha} := \varepsilon' / (\varepsilon' + |u^1 - u|)$. For any $t \in \mathbf{R}_+ \setminus \mathcal{T}'$ we have either $X_t \leq (u \wedge u^1) - \varepsilon' < u$ or (if $u^1 > 0$) $X_t \geq (u \vee u^1) + \varepsilon' > u$. Thus for any $\alpha \in (0, \bar{\alpha})$, either

$$X_t < (1 - \alpha)X_t + \alpha u \leq (1 - \alpha)(u \wedge u^1) - (1 - \alpha)\varepsilon' + \alpha u < u^1$$

(where the last inequality holds by $\alpha < \bar{\alpha}$) or (similarly)

$$X_t > (1 - \alpha)X_t + \alpha u \geq (1 - \alpha)(u \vee u^1) + (1 - \alpha)\varepsilon' + \alpha u > u^1.$$

In short, $(1 - \alpha)X_t + \alpha u$ lies between X_t and u^1 for any $t \in \mathbf{R}_+ \setminus \mathcal{T}'$ and $\alpha \in (0, \bar{\alpha})$, which since F^1 is concave and maximised at u^1 implies that $(0, \bar{\alpha}) \ni \alpha \mapsto h_\alpha^1 \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}'}$ is (pointwise) decreasing and non-negative. Hence

$$0 \leq \lim_{\alpha \downarrow 0} \mathbf{E}_G \left(h_\alpha^1(\tau) \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}'}(\tau) \right) = \mathbf{E}_G \left(h^1(\tau) \mathbf{1}_{\mathbf{R}_+ \setminus \mathcal{T}'}(\tau) \right)$$

by the monotone convergence theorem, where the right-hand side is well-defined. This together with (1) implies that h^1 satisfies (b) and (c). \square

3 Proof of the Gateaux lemma (p. 66)

For $\alpha \in (0, 1)$, define $h_\alpha^0, h_\alpha^1 : \mathbf{R}_+ \rightarrow (-\infty, \infty]$:

$$h_\alpha^0(t) := re^{-rt} \frac{F^0(x_t + \alpha[x_t^\dagger - x_t]) - F^0(x_t)}{\alpha}$$

and
$$h_\alpha^1(t) := e^{-rt} \frac{F^1(X_t + \alpha[X_t^\dagger - X_t]) - F^1(X_t)}{\alpha}.$$

Clearly both are measurable for each $\alpha \in (0, 1)$. We have

$$D\pi_G(x, x^\dagger - x) = \lim_{\alpha \downarrow 0} \left[\mathbf{E}_G \left(\int_0^\tau h_\alpha^0 \right) + \mathbf{E}_G \left(h_\alpha^1(\tau) \right) \right]$$

by definition; we must show that the limit exists and has the asserted value.

As $\alpha \downarrow 0$, h_α^0 and h_α^1 converge pointwise to $h^0 : \mathbf{R}_+ \rightarrow (-\infty, \infty]$ and $h^1 : \mathbf{R}_+ \rightarrow [-\infty, \infty]$, respectively, given by

$$h^0(t) := re^{-rt} F^{0\prime}(x_t, x_t^\dagger) (x_t^\dagger - x_t)$$

and
$$h^1(t) := e^{-rt} F^{1\prime}(X_t, X_t^\dagger) (X_t^\dagger - X_t).$$

Claim A. $D\pi_G(x, x^\dagger - x)$ exists and is finite, equal to $\mathbf{E}_G(\int_0^\tau h^0) + \mathbf{E}_G(h^1(\tau))$.

By claim A, we have

$$\begin{aligned}
D\pi_G(x, x^\dagger - x) &= \mathbf{E}_G\left(\int_0^\tau h^0\right) + \mathbf{E}_G(h^1(\tau)) \\
&= \mathbf{E}_G\left(r \int_0^\tau e^{-rs} \phi^0(s) [x_s^\dagger - x_s] ds\right) + \mathbf{E}_G\left(e^{-r\tau} \phi^1(\tau) [X_\tau^\dagger - X_\tau]\right) \\
&\quad + \mathbf{E}_G\left(r \int_0^\tau e^{-rt} [F^{0t}(x_t, x_t^\dagger) - \phi^0(t)] [x_t - x_t^\dagger] dt\right) \\
&\quad + \mathbf{E}_G\left(e^{-r\tau} [F^{1\tau}(X_\tau, X_\tau^\dagger) - \phi^1(\tau)] [X_\tau^\dagger - X_\tau]\right),
\end{aligned}$$

where the last equality holds since $t \mapsto r \int_0^t e^{-rs} \phi^0(s) ds$ and ϕ^1 are G -integrable and $x, x^\dagger, X, X^\dagger$ are bounded. The proof is completed by invoking the following claims to rewrite the first two terms in the desired form:

Claim B. It holds that

$$\mathbf{E}_G\left(r \int_0^\tau e^{-rs} \phi^0(s) [x_s^\dagger - x_s] ds\right) = r \int_0^\infty e^{-rt} [1 - G(t)] \phi^0(t) (x_t^\dagger - x_t) dt.$$

Claim C. It holds that

$$\mathbf{E}_G\left(e^{-r\tau} \phi^1(\tau) [X_\tau^\dagger - X_\tau]\right) = r \int_0^\infty e^{-rt} \left(\int_{[0,t]} \phi^1 dG\right) (x_t^\dagger - x_t) dt.$$

Proof of claim A. We must show that $t \mapsto \int_0^t h^0$ and h^1 are G -integrable and

$$\mathbf{E}_G\left(\int_0^\tau h_\alpha^0\right) \rightarrow \mathbf{E}_G\left(\int_0^\tau h^0\right) \quad \text{and} \quad \mathbf{E}_G\left(h_\alpha^1(\tau)\right) \rightarrow \mathbf{E}_G\left(h^1(\tau)\right) \quad \text{as } \alpha \downarrow 0.$$

Fix a $u \in (0, u^0)$.

For h^0 , define $\varphi : \mathbf{R}_+ \rightarrow [0, \infty]$ by

$$\varphi(t) := r e^{-rt} [F^{0t}(x_t, u) + F^{0t}(x_t^\dagger, u)].$$

Recall the definition of $\psi_{x,u}^0, \psi_{x^\dagger,u}^0$ (p. 66). Since x and x^\dagger belong to \mathcal{X}_G , the map

$$t \mapsto \int_0^t \varphi = \psi_{x,u}^0(t) + \psi_{x^\dagger,u}^0(t)$$

is G -integrable, and thus φ (being non-negative) is Lebesgue-integrable on $(0, t)$ for any $t \in \mathbf{R}_+$ with $G(t) < 1$.³ Since F^0 is concave, we have

$$|h_\alpha^0(t)| \leq \varphi(t) |x_t - x_t^\dagger| \leq \varphi(t) u^0 \quad \text{for all } \alpha \in (0, 1) \text{ and } t \in \mathbf{R}_+,$$

³If φ were not integrable on $(0, t)$, then $\psi_{x,u}^0(s) + \psi_{x^\dagger,u}^0(s) = \int_0^s \varphi \geq \int_0^t \varphi = \infty$ for every $s \geq t$, which if $G(t) < 1$ implies $\mathbf{E}_G(\psi_{x,u}^0(\tau) + \psi_{x^\dagger,u}^0(\tau)) = \infty$, a contradiction.

and thus

$$\left| \int_0^t h_\alpha^0 \right| \leq \left[\psi_{x,u}^0(t) + \psi_{x^\dagger,u}^0(t) \right] u^0 \quad \text{for all } \alpha \in (0,1) \text{ and } t \in \mathbf{R}_+.$$

Thus applying the dominated convergence theorem twice yields

$$\lim_{\alpha \downarrow 0} \mathbf{E}_G \left(\int_0^\tau h_\alpha^0 \right) = \mathbf{E}_G \left(\lim_{\alpha \downarrow 0} \int_0^\tau h_\alpha^0 \right) = \mathbf{E}_G \left(\int_0^\tau h^0 \right) \in \mathbf{R}.$$

Similarly, since F^1 is concave and X, X^\dagger take values in $[0, u^0]$, we have

$$\begin{aligned} |h_\alpha^1(t)| &\leq \left(\left| \psi_{X,u}^1(t) \right| + \left| \psi_{X^\dagger,u}^1(t) \right| \right) |X_t^\dagger - X_t| \\ &\leq \left(\left| \psi_{X,u}^1(t) \right| + \left| \psi_{X^\dagger,u}^1(t) \right| \right) u^0 \quad \text{for all } \alpha \in (0,1) \text{ and } t \in \mathbf{R}_+. \end{aligned}$$

The right-hand side is G -integrable by since x and x^\dagger belong to \mathcal{X}_G , so

$$\lim_{\alpha \downarrow 0} \mathbf{E}_G \left(h_\alpha^1(\tau) \right) = \mathbf{E}_G \left(h^1(\tau) \right) \in \mathbf{R}$$

by dominated convergence. \square

Proof of claim B. Fix a $T \in \mathbf{R}_+$ such that $\lim_{t \uparrow T} G(t) < 1$, and note that $l_T : \mathbf{R}_+ \rightarrow \mathbf{R}$ given by

$$l_T(t) := r e^{-rt} \phi^0(t) \left[x_t^\dagger - x_t \right] \mathbf{1}_{[0,T)}(t)$$

is (Lebesgue-)integrable since $t \mapsto r \int_0^t e^{-rs} \phi^0(s) ds$ is G -integrable and x, x^\dagger are bounded. We may therefore apply IBP lemma (p. 6) to l_T and the measure associated with G to obtain

$$\begin{aligned} \int_{[0,T]} \left(r \int_0^t e^{-rs} \phi^0(s) \left[x_s^\dagger - x_s \right] ds \right) G(dt) \\ = r \int_0^T [G(T) - G(t)] e^{-rt} \phi^0(t) \left[x_t^\dagger - x_t \right] dt. \end{aligned}$$

Define $\bar{T} := \sup\{t \in \mathbf{R}_+ : G(t) < 1\}$. If $\bar{T} < \infty$ and G has an atom at \bar{T} , then we obtain the desired equation by setting $T := \bar{T}$. Suppose for the remainder that this is not the case. It suffices to show that $\chi : \mathbf{R}_+ \rightarrow [0, \infty]$ given by

$$\chi(t) := \begin{cases} [1 - G(t)] r e^{-rt} \phi^0(t) & \text{for } t < \bar{T} \\ 0 & \text{for } t \geq \bar{T} \end{cases}$$

satisfies $\int_0^\infty \chi < \infty$ (so that it is Lebesgue-integrable), for then letting $T \uparrow \bar{T}$ above yields the desired result by the dominated convergence theorem since $t \mapsto r \int_0^t e^{-rs} \phi^0(s) ds$ is G -integrable and x, x^\dagger are bounded.

To show that $\int_0^\infty \chi$ is finite, note that applying the IBP lemma as above to x and $\tilde{x}^\dagger := x + k$ (where $k \in \mathbf{R}$ is a constant) yields

$$\int_{[0, T]} \left(r \int_0^t e^{-rs} \phi^0(s) ds \right) G(dt) = r \int_0^T [G(T) - G(t)] e^{-rt} \phi^0(t) dt$$

for any $T \in \mathbf{R}_+$ with $G(T) < 1$. Letting $T \uparrow \bar{T}$ and using the dominated (monotone) convergence theorem on the left-hand (right-hand) side yields

$$\int_{[0, \bar{T}]} \left(r \int_0^t e^{-rs} \phi^0(s) ds \right) G(dt) = \int_0^{\bar{T}} \chi,$$

which since G has no atom at \bar{T} implies that

$$\mathbf{E}_G \left(r \int_0^{\bar{T}} e^{-rt} \phi^0(t) dt \right) = \int_0^{\bar{T}} \chi = \int_0^\infty \chi.$$

Thus $\int_0^\infty \chi < \infty$ since $t \mapsto r \int_0^t e^{-rs} \phi^0(s) ds$ is G -integrable. \square

Proof of claim C. Define a measure ν_+ on \mathbf{R}_+ by

$$\nu_+(A) := \int_A \max\{\phi^1, 0\} dG \quad \text{for any measurable } A \subseteq \mathbf{R}_+.$$

Since ϕ^1 is G -integrable, ν_+ is absolutely continuous with respect to (the measure associated with) G , with Radon–Nikodým derivative $\max\{\phi^1, 0\}$. Furthermore, the function $L : \mathbf{R}_+ \rightarrow \mathbf{R}$ defined by

$$L(t) := e^{-rt} (X_t^\dagger - X_t) \quad \text{for each } t \in \mathbf{R}_+$$

is ν_+ -integrable since X^\dagger and X are bounded. Thus for any $T \in \mathbf{R}_+$, we have

$$\int_{[0, T]} L \max\{\phi^1, 0\} dG = \int_{[0, T]} L d\nu_+.$$

Furthermore, L satisfies $L(t) = L(0) + \int_0^t l$, where $l(t) := -e^{-rt} (x_t^\dagger - x_t)$. Thus for every $T \in \mathbf{R}_+$, the IBP lemma yields

$$\begin{aligned} & \int_{[0, T]} e^{-rt} (X_t^\dagger - X_t) \nu_+(dt) \\ &= e^{-rT} (X_T^\dagger - X_T) \nu_+([0, T]) + r \int_0^T \nu_+([0, t]) e^{-rt} (x_t^\dagger - x_t) dt. \end{aligned}$$

Applying the same argument to $\nu_-(A) := -\int_A \min\{\phi^1, 0\}dG$ and subtracting yields, for each $T \in \mathbf{R}_+$,

$$\begin{aligned} & \int_{[0,T]} e^{-rt} (X_t^\dagger - X_t) \phi^1(t) G(dt) \\ &= e^{-rT} (X_T^\dagger - X_T) \int_{[0,T]} \phi^1 dG + r \int_0^T \left(\int_{[0,t]} \phi^1 dG \right) e^{-rt} (x_t^\dagger - x_t) dt. \end{aligned}$$

Letting $T \rightarrow \infty$ yields the desired equation by boundedness of $x, x^\dagger, X, X^\dagger$ and dominated convergence. \square

With all three claims established, the proof is complete. \blacksquare

Supplemental appendix N.3 (p. 72)

Here's an example of a sequence of technologies satisfying properties (a)–(d). Choose an $N \in \mathbf{N}$ such that $1/N < (u^0 - u^1)/3$. Given $n \geq N$, let $I_n := [1/n, u^0 - 2/n]$. Fix $\gamma^n \in (0, 1/u^0 n)$ and $\delta^n \in (0, 1/n)$. For $j \in \{0, 1\}$, define $f_n^j : I_n \rightarrow \mathbf{R}$ by $f_n^j(u) := \frac{1}{\delta^n} \int_u^{u+\delta^n} F^{j+} - \gamma^n u$. Given $\zeta^n \in (0, 2/n)$, define $F_n^0, F_n^1 : I_n \rightarrow \mathbf{R}$ by

$$\begin{aligned} F_n^0(u) &:= F^0\left(u^* + \frac{1}{n}\right) + \int_{u^*+1/n}^u f_n^0 \\ \text{and } F_n^1(u) &:= F^1\left(u^* + \frac{1}{n}\right) + \int_{u^*+1/n}^u f_n^1 + \zeta^n. \end{aligned}$$

F_n^0, F_n^1 are strictly concave and differentiable, with (continuous) derivatives f_n^0, f_n^1 such that

$$F^{j-}(u + 1/n) - 1 \leq f_n^j(u) \leq F^{j+}(u) \quad \text{for } j \in \{0, 1\} \text{ and } u \in I_n. \quad (\natural)$$

As $n \rightarrow \infty$, $f_n^j \rightarrow F^{j+}$ pointwise for both $j \in \{0, 1\}$ since F^{j+} is right-continuous, so that $F_n^0 \rightarrow F^0$ and $F_n^1 - \zeta^n \rightarrow F^1$ pointwise by the bounded convergence theorem. In fact, the convergence is uniform (Rockafellar, 1970, Theorem 10.8, p. 90). Thus for any $\varepsilon^n \in (0, 1/n)$, provided δ^n and γ^n are sufficiently small, we have that $F_n^0 - F^0$ and $F_n^1 - \zeta^n - F^1$ are bounded by ε^n , so that choosing $\zeta^n > 2\varepsilon^n$ ensures $F_n^1 > F_n^0$ (as $F^1 \geq F^0$).

Since $F_n^1(1/n) > F_n^0(1/n)$, we may extend F_n^0, F_n^1 to $[0, u^0 - 2/n]$ while preserving strict concavity, (continuous) differentiability and $F_n^1 \geq F_n^0$ and ensuring that $F_n^1 - F_n^0$ is strictly increasing on a neighbourhood of 0 (which guarantees that $u_n^* > 0$) and that $F_n^{0'}, F_n^{1'}$ are bounded above on $(0, 1/n]$

by $\max_{j \in \{0,1\}} F^{j+}(1/n) + 1$. (The latter is possible because $F_n^{j'}(1/n) \leq F^{j+}(1/n)$ by (‡).) Further extend F_n^1 to all of $[0, \infty)$ by making it differentiable and strictly concave on $[u^0 - 2/n, \infty)$, with derivative bounded below by $F_n^{1'}(u^0 - 2/n) - 1/n$. Since $F_n^1(u^0 - 2/n) > F_n^0(u^0 - 2/n)$, we may also extend F_n^0 to $[0, \infty)$ while preserving strict concavity, (continuous) differentiability and $F_n^1 \geq F_n^0$ and ensuring that $u_n^0 \leq u^0$ and that $F_n^{0'}$ is bounded below by $F^{1-}(u^0 - 1/n) - 2$. (The latter is possible since $F_n^{1'}$ is bounded below by $F_n^{1'}(u^0 - 2/n) - 1/n \geq F^{1-}(u^0 - 1/n) - 2$ by (‡).) If u_n^* is not a strict local maximum of $F_n^1 - F_n^0$, then we may make it so by decreasing F_n^1 pointwise on $[0, u_n^*)$ by a small amount.

Claim. The sequence $(F_n^0, F_n^1)_{n=N}^\infty$ satisfies the model assumptions and properties (a)–(d) provided $\varepsilon^n \in (0, 1/n)$ is chosen sufficiently small.

Proof. It is clear that $(F_n^0, F_n^1)_{n=N}^\infty$ satisfies property (a), and that for each $n \in \mathbf{N}$, F_n^0, F_n^1 satisfy the all of the model assumptions except for $u_n^1 < u_n^0$ (the conflict of interest).

For (b) and the conflict of interest, since u^0 , u^1 and u^* are strict local maxima of (respectively) F^0 , F^1 and $F^1 - F^0$, we have

$$u^0 - \frac{1}{n} \leq u_n^0 \leq u^0, \quad |u_n^1 - u^1| \leq \frac{1}{n} \quad \text{and} \quad |u_n^* - u^*| \leq \frac{1}{n}$$

for $\varepsilon^n > 0$ sufficiently small, as $F_n^0 - F^0$ and $F_n^1 - (F^1 + \zeta^n)$ are bounded by ε^n on I_n . In particular, $u_n^1 < u_n^0$ (the conflict of interest) since $n \geq N$. Letting $n \rightarrow \infty$ yields (b).

For the first part of (c), fix a $u \in (0, u^0]$ at which F^{1-} is finite. If $u = u^0$, then for each $n \geq N$, $F_n^{0'}, F_n^{1'}$ are bounded below by $F^{1-}(u^0 - 1/n) - 2$ (by construction for $F_n^{0'}$, and by (‡) for $F_n^{1'}$). Suppose instead that $u < u^0$. Choose a $u' \in (u, u^0)$, and let $N' \geq N$ satisfy $1/N' < u < u^0 - 2/N'$ and $1/N' < u' - u$. Then u belongs to I_n for every $n \geq N'$, so that $F_n^{j'}(u) \geq F^{j-}(u + 1/n) - 1 \geq F^{j-}(u') - 1$ by (‡) for both $j \in \{0, 1\}$. The set

$$\bigcup_{j \in \{0,1\}} \left\{ F_n^{j'}(u) : n \in \{1, \dots, N'\} \right\}$$

is bounded since it is finite. Thus $(F_n^{0'})_{n \in \mathbf{N}}$ and $(F_n^{1'})_{n \in \mathbf{N}}$ are uniformly bounded below on $[0, u]$.

The argument for the second part of (c) is analogous: fix a $u \in [0, u^0)$ at which F^{0+}, F^{1+} are finite. If $u = 0$, then $F_n^{0'}, F_n^{1'}$ are bounded above by $\max_{j \in \{0,1\}} F_j^{0+}(1/n) + 1$ for all $n \geq N$ by construction. If instead $u > 0$,

then $F_n^{j'}(u) \leq F^{j+}(u)$ for all $n \geq N$ such that $1/n < u < u^0 - 2/n$ by (‡), and the set

$$\bigcup_{j \in \{0,1\}} \left\{ F_n^{j'}(u) : n \in \left\{ 1, \dots, \lfloor 1/u \rfloor \vee \left\lfloor \frac{2}{u^0 - u} \right\rfloor \right\} \right\}$$

is finite and thus bounded; so $(F_n^{0'})_{n \in \mathbb{N}}$ and $(F_n^{1'})_{n \in \mathbb{N}}$ are uniformly bounded above on $[u, u^0]$.

For (d), fix $j \in \{0, 1\}$ and a convergent sequence $(u_n)_{n=N}^\infty$ with limit $u \in [0, \infty)$ such that $0 < u_n \leq u_n^0$ for each $n \geq N$ and $\lim_{n \rightarrow \infty} F_n^{j'}(u_n)$ exists. Since $u_n^0 \leq u^0$ for each $n \geq N$, we have $u \leq u^0$. From above,

$$F_n^{j'}(u_n) = f_n^j(u_n) \begin{cases} \leq F^{j+}(u_n) & \text{if } u_n \geq 1/n \\ \geq F^{j-}(u_n + 1/n) - \gamma^n u_n & \text{if } u_n \leq u^0 - 2/n. \end{cases} \quad (\dagger)$$

Letting $n \rightarrow \infty$ yields

$$\lim_{n \rightarrow \infty} F_n^{j'}(u_n) \begin{cases} \leq F^{j-}(u) & \text{if } u > 0 \\ \geq F^{j+}(u) & \text{if } u < u^0. \end{cases}$$

If $u \in (0, u^0)$, then both cases are satisfied, so

$$F^{j+}(u) \leq \lim_{n \rightarrow \infty} F_n^{j'}(u_n) \leq F^{j-}(u).$$

If $u = 0$, then the second case applies, so that $F^{j+}(0) \leq \lim_{n \rightarrow \infty} F_n^{j'}(u_n)$, which suffices. If $u = u^0$, then the first case holds, so it remains only to show that $F^{j+}(u^0) \leq \lim_{n \rightarrow \infty} F_n^{j'}(u_n)$. For $j = 0$, we have $F^{0+}(u^0) \leq 0 \leq \lim_{n \rightarrow \infty} F_n^{0'}(u_n)$ since $u_n \leq u_n^0$ for each $n \geq N$. For $j = 1$, (†) implies

$$\begin{aligned} & F^{1-} \left(\min \left\{ u_n + 1/n, u^0 - 1/n \right\} \right) - \gamma^n u_n - 1/n \\ & \leq F_n^{1'} \left(\min \left\{ u_n, u^0 - 2/n \right\} \right) - 1/n \leq F_n^{1'}(u_n) \quad \text{for each } n \geq N, \end{aligned}$$

where the second inequality holds since $F_n^{1'}$ is bounded below by $F_n^{1'}(u^0 - 2/n) - 1/n$. Letting $n \rightarrow \infty$ yields $F^{1+}(u^0) \leq \lim_{n \rightarrow \infty} F_n^{1'}(u_n)$. ■

Supplemental appendix O.2 (p. 78)

Proof of Observation 10. It suffices to show that π_G is strictly concave. For distinct $x, x^\dagger \in \mathcal{X}$ (i.e. $x \neq x^\dagger$ on a non-null set) and any $\lambda \in (0, 1)$, we have

$$\begin{aligned} & \pi_G(\lambda x + (1 - \lambda)x^\dagger) \\ &= \mathbf{E}_G \left(r \int_0^\tau e^{-rt} F^0(\lambda x_t + (1 - \lambda)x_t^\dagger) dt + e^{-r\tau} F^1(\lambda X_\tau + (1 - \lambda)X_\tau^\dagger) \right) \\ &> \mathbf{E}_G \left(r \int_0^\tau e^{-rt} [\lambda F^0(x_t) + (1 - \lambda)F^0(x_t^\dagger)] dt \right. \\ &\quad \left. + e^{-r\tau} [\lambda F^1(X_\tau) + (1 - \lambda)F^1(X_\tau^\dagger)] \right) \\ &= \lambda \pi_G(x) + (1 - \lambda) \pi_G(x^\dagger), \end{aligned}$$

where the inequality holds since F^1 is concave, F^0 is strictly concave, G has unbounded support and $x \neq x^\dagger$ on a non-null set. ■

References

- Billingsley, P. (1995). *Probability and measure* (3rd). Wiley.
- Curello, G., & Sinander, L. (2024). *Screening for breakthroughs* [working paper, Feb 2024]. <https://arXiv.org/abs/2104.02044v8>
- Rockafellar, R. T. (1970). *Convex analysis*. Princeton University Press.