

A DYNAMICAL ARGUMENT FOR A RAMSEY PROPERTY

ENHUI SHI AND HUI XU

ABSTRACT. We show by a dynamical argument that there is a positive integer valued function q defined on positive integer set \mathbb{N} such that $q(\lfloor \log n \rfloor + 1)$ is a super-polynomial with respect to positive n and

$$\limsup_{n \rightarrow \infty} r((2n+1)^2, q(n)) < \infty,$$

where $r(\cdot, \cdot)$ is the opposite-Ramsey number function.

1. INTRODUCTION AND PRELIMINARIES

For positive integers p and q , we define the **opposite-Ramsey number** $r(p, q)$ to be the maximal number k for which every edge-coloring of the complete graph K_q with p colors yields a monochromatic complete subgraph of order k (the **order** of a graph means the number of its vertices).

The following is implied by the well-known Ramsey's theorem.

Theorem 1.1. *Let p be a fixed positive integer. Then*

$$\liminf_{q \rightarrow \infty} r(p, q) = \infty.$$

One may expect that if $p = p(n)$ and $q = q(n)$ are positive integer valued functions defined on \mathbb{N} and the speed of $q(n)$ tending to infinity is much faster than that of $p(n)$ as n tends to infinity, then we still have

$$\liminf_{n \rightarrow \infty} r(p(n), q(n)) = \infty.$$

The purpose of the paper is to show by a dynamical argument that this is not true in general even if $p(n)$ is a polynomial and $q(\lfloor \log n \rfloor + 1)$ is a super-polynomial. By a **super-polynomial**, we mean a function $f : \mathbb{N} \rightarrow \mathbb{R}$ such that for any polynomial $g(n)$,

$$\liminf_{n \rightarrow \infty} \frac{|f(n)|}{|g(n)|} = \infty.$$

Let (X, d) be a compact metric space. For any $\varepsilon > 0$, let $N(\varepsilon)$ denote the minimal number of subsets of diameter at most ε needed to cover X . The **lower box dimension** of X is defined to be

$$(1.1) \quad \underline{\dim}_B(X, d) = \liminf_{\varepsilon \rightarrow 0} \frac{\log N(\varepsilon)}{\log 1/\varepsilon}.$$

For a subset E of X and $\varepsilon > 0$, we say E is ε -**separated** if for any distinct $x, y \in E$, $d(x, y) \geq \varepsilon$. Let $S(\varepsilon)$ denote the cardinality of a maximal ε -separated subset of X . It is easy to verify $N(\varepsilon) \leq S(\varepsilon) \leq N(\varepsilon/2)$. Thus

Date: April 9, 2021.

$$(1.2) \quad \underline{\dim}_B(X, d) = \liminf_{\varepsilon \rightarrow 0} \frac{\log S(\varepsilon)}{\log 1/\varepsilon}.$$

Furthermore, it is easy to see that

$$(1.3) \quad \underline{\dim}_B(X, d) = \liminf_{n \rightarrow \infty} \frac{\log S(1/n)}{\log n}.$$

We use $\dim(X)$ to denote the topological dimension of X . It is well known that the topological dimension of X is always no greater than its lower box dimension with respect to any compatible metric.

A continuous action $G \curvearrowright X$ of group G on X is said to be **expansive** if there exists $c > 0$ such that for any two distinct points $x, y \in X$, $\sup_{g \in G} d(gx, gy) > c$. For $v = (v_1, \dots, v_k) \in \mathbb{Z}^k$, let $|v|$ denote $\max\{|v_1|, \dots, |v_k|\}$.

The following lemma is due to T. Meyerovitch and M. Tsukamoto.

Lemma 1.2. [4, Lemma 4.4] *Let k be a positive integer and $T : \mathbb{Z}^k \times X \rightarrow X$ be a continuous action of \mathbb{Z}^k on a compact metric space (X, d) . If the action is expansive, then there exist $\alpha > 1$ and a compatible metric D on X such that for any positive integer n and any two distinct points $x, y \in X$ satisfying $D(x, y) \geq \alpha^{-n}$, we have*

$$\max_{v \in \mathbb{Z}^k, |v| \leq n} D(T^v x, T^v y) \geq \frac{1}{4\alpha}.$$

Lemma 1.3. *If (X, d) is a compact metric space of infinite dimension, then $S(1/n)$ is a super-polynomial with respect to variable n .*

Proof. Since $\dim(X) \leq \underline{\dim}_B(X, d)$, we have

$$(1.4) \quad \underline{\dim}_B(X, d) = \liminf_{n \rightarrow \infty} \frac{\log S(1/n)}{\log n} = \infty.$$

Thus, for any positive integer k , $\liminf_{n \rightarrow \infty} \frac{S(1/n)}{n^k} = \infty$. □

2. MAIN RESULTS

For a positive real number x , we use $[x]$ to denote its integer part.

Theorem 2.1. *There is a function $q : \mathbb{N} \rightarrow \mathbb{R}$ such that $q([\log n] + 1)$ is a super-polynomial and*

$$\limsup_{n \rightarrow \infty} r((2n+1)^2, q(n)) < \infty.$$

Proof. Let $T : \mathbb{Z}^2 \times X \rightarrow X$ be an expansive continuous action on a compact metric space (X, d) of infinite dimension (see [5] where an expansive \mathbb{Z}^2 -action on \mathbb{T}^∞ was constructed). By Lemma 1.2, there exist $\alpha > 1$ and a compatible metric D on X such that for any positive integer n and any two distinct points $x, y \in X$ with $D(x, y) \geq \alpha^{-n}$,

$$\max_{v \in \mathbb{Z}^2, |v| \leq n} D(T^v x, T^v y) \geq \frac{1}{4\alpha}.$$

For each $n \in \mathbb{N}$, let V_n be a maximal α^{-n} -separated set of (X, D) . Hence $|V_n| = S(\alpha^{-n})$. Let G_n be the complete graph $K_{S(\alpha^{-n})}$ whose vertex set is V_n . Now we use the color set

$C_n = \{v \in \mathbb{Z}^2 : |v| \leq n\}$ to color the edges of G_n . Since V_n is α^{-n} -separated, for any two distinct points $x, y \in V_n$, $D(x, y) \geq \alpha^{-n}$. By Lemma 1.2, there exists $v \in C_n$ such that $D(T^v x, T^v y) \geq \frac{1}{4\alpha}$. Then we color the edge $\{x, y\}$ by v . By the definition of opposite-Ramsey number, there is a monochromatic complete subgraph H_n of order $r((2n+1)^2, S(\alpha^{-n}))$.

By Lemma 1.3, $S(1/n)$ is a super-polynomial. Let $q(n) = S(\alpha^{-n})$. Thus $q([\log n] + 1)$ is a super-polynomial with respect to positive n . Assuming that the conclusion of the Theorem is false, we have

$$\limsup_{n \rightarrow \infty} r((2n+1)^2, q(n)) = \infty.$$

Therefore, there is an increasing subsequence (n_i) of positive integers such the the sequence of orders of H_{n_i} is unbounded. Since H_{n_i} is monochromatic, there exists $v_{n_i} \in C_{n_i}$ such that the image of vertex set of H_{n_i} under $T^{v_{n_i}}$ is $\frac{1}{4\alpha}$ -separated. These imply that there are arbitrarily large $\frac{1}{4\alpha}$ -separated subsets of X , which contradicts the compactness of X . Thus we complete the proof. \square

3. COMPARISON WITH CLASSICAL RAMSEY NUMBER

For any positive integers k and g , the **Ramsey number** $R_g(k)$ is defined to be the minimal number n for which every edge-coloring of the complete graph K_n with g colors yields a monochromatic complete subgraph of order k .

By Corollary 3 of Greenwood and Gleason in [1], $R_g(k)$ has an upper bound g^{gk} . In [2] Lefmann and Rödl obtained a lower bound $2^{\Omega(gk)}$ for $R_g(k)$. Thus

$$(3.1) \quad 2^{\Omega((2n+1)^2 k)} \leq R_{(2n+1)^2}(k) \leq ((2n+1)^2)^{(2n+1)^2 k}.$$

Suppose $r((2n+1)^2, q(n)) = r < \infty$. Then it implies that

- (1) every edge-coloring of complete graph $K_{q(n)}$ with $(2n+1)^2$ colors yields a monochromatic complete subgraph of order r , hence

$$(3.2) \quad q(n) \geq R_{(2n+1)^2}(r);$$

- (2) there exists an edge-coloring of $K_{q(n)}$ with $(2n+1)^2$ colors such that there is no monochromatic complete subgraph of order $r+1$, hence

$$(3.3) \quad q(n) \leq R_{(2n+1)^2}(r+1).$$

Thus $q(n)$ gives a lower bound of $R_{(2n+1)^2}(r+1)$ and an upper bound of $R_{(2n+1)^2}(r)$. By Theorem 2.1, every expansive \mathbb{Z}^2 -action on a compact metric space of infinite dimension gives rise to such a $q(n)$. In addition, there is a positive integer r and an increasing subsequence (n_i) of positive integers such that for any $i \in \mathbb{N}$, $r((2n_i+1)^2, q(n_i)) = r$. Therefore, we obtain a lower bound of $R_{(2n_i+1)^2}(r+1)$ and an upper bound of $R_{(2n_i+1)^2}(r)$ for each $i \in \mathbb{N}$.

If $q(\log n)$ is a super-polynomial, then we claim that for any $A \geq 0$,

$$(3.4) \quad \liminf_{n \rightarrow \infty} \frac{q(n)}{A^n} = \infty.$$

In fact, take a positive integer m such that $e^m \geq A$. Then

$$\liminf_{n \rightarrow \infty} \frac{q(n)}{A^n} \geq \liminf_{n \rightarrow \infty} \frac{q(n)}{e^{mn}} = \liminf_{n \rightarrow \infty} \frac{q(\log(e^n))}{(e^n)^m} = \infty.$$

The lower bound of $R_{(2n+1)^2}(r+1)$ obtained by (3.1) is $2^{\Omega((2n+1)^2(r+1))}$ which is also faster than any exponential growth. If there is an expansive \mathbb{Z} -action on a compact metric space of infinite dimension, then we can get a lower bound for $R_{2n+1}(r+1)$ which is faster than the classical bound $2^{\Omega((2n+1)(r+1))}$. Unfortunately, in [3] Mañé showed that such action does not exist. If we can construct an expansive \mathbb{Z}^2 -action on a compact metric space of infinite dimension such that the condition $D(x, y) \geq \alpha^{-n}$ in Lemma 1.2 can be replaced by $D(x, y) \geq \alpha^{-n^2}$, then we can show that $q(n)$ obtained in Theorem 2.1 satisfies that $q([\sqrt{\log n}] + 1)$ is a super-polynomial. Then it satisfies $\liminf_{n \rightarrow \infty} \frac{q(n)}{A^{n^2}} = \infty$. Hence $q(n)$ is faster than the classical lower bound $2^{\Omega((2n+1)^2(r+1))}$. Therefore, we leave the following question.

Question 3.1. *Is there an expansive \mathbb{Z}^2 -action on a compact metric space (X, d) of infinite dimension and $\alpha > 1$ such that for any positive integer n and any two distinct points $x, y \in X$ satisfying $d(x, y) \geq \alpha^{-n^2}$, we have*

$$\max_{v \in \mathbb{Z}^2, |v| \leq n} d(T^v x, T^v y) \geq \frac{1}{4\alpha}.$$

A positive answer to Question 3.1 can give a better estimate of the lower bound of $R_{(2n_i+1)^2}(r+1)$, where (n_i) and r come from the system. By (3.2), a negative answer also gives a better estimate of the upper bound of $R_{(2n_i+1)^2}(r)$.

Finally, we remark that the above comparison between $q(n)$ and the bound of Ramsey is only for a subsequence of positive integers. However, dealing with a concrete system we may obtain more information and can get a special edge-coloring of $K_{q(n)}$ as the proof of Theorem 2.1. Our method may give a new direction to estimate the bounds of Ramsey numbers and construct edge-colorings of big graphs.

Acknowledgements. We would like to thank Professor Bingbing Liang and Professor Xin Wang for their helpful comments.

REFERENCES

- [1] R. E. Greenwood and A. M. Gleason, Combinatorial Relations and Chromatic Graphs. *Canadian Journal of Mathematics* 7(1955):1-7.
- [2] H. Lefmann and V. Rödl, On canonical Ramsey numbers for complete graphs versus paths. *J. Combin. Theory Ser. B* 58 (1993), no. 1, 1-13.
- [3] R. Mañé, Expansive homeomorphisms and topological dimension, *Trans. Amer. Math. Soc.* 252(1979), 313-319.
- [4] T. Meyerovitch and M. Tsukamoto, Expansive multiparameter actions and mean dimension. *Trans. Amer. Math. Soc.* 371 (2019), no. 10, 7275-7299.
- [5] E. Shi and L. Zhou, The nonexistence of expansive \mathbb{Z}^d actions on graphs. *Acta Math. Sin. (Engl. Ser.)* (2005) 1509-1514.

SOOCHOW UNIVERSITY, SUZHOU, JIANGSU 215006, CHINA
 Email address: ehshi@suda.edu.cn

SOOCHOW UNIVERSITY, SUZHOU, JIANGSU 215006, CHINA
 Email address: 20184007001@stu.suda.edu.cn