

Representing Polynomial Splines over Infinite Hierarchical Meshes by Finite Automata

Dmitry Berdinsky ^{*} Prohrak Kruengthomya [†]

Abstract We propose a data structure based on finite automata for representing polynomial splines over infinite hierarchical meshes. It allows to store and operate such splines using only finite amount of memory. This naturally extends a classical framework of hierarchical tensor product B-splines for infinite meshes in a way suitable for computing.

Keywords polynomial splines, hierarchical meshes, finite automata

1 Introduction

Polynomial [splines](#) over meshes which admit local refinement have been applied in geometric modelling, computer graphics and finite element analysis for the last forty years. The introduction of [isogeometric analysis](#) integrating computer-aided design and finite element analysis in the 2000s inspired new interest in this kind of splines as well as local refinement methods [10]. Let us briefly recall four approaches to polynomial splines over meshes admitting local refinement which were actively developed for the last two decades: hierarchical tensor product B-splines, polynomial splines over hierarchical T-meshes (PHT-splines), analysis-suitable T-splines (AS T-splines) and polynomial splines over locally refined box-partitions (LR-splines). Due to the vast amount of literature discussing these four approaches we provide only few references. The interested reader can find more references to some papers published before 2015 in a survey article [14].

The approach based on hierarchical tensor product B-splines was introduced by Forsey and Bartels in the end of 1980s [7]. In the mid 1990s Kraft proposed a selection mechanism¹ for generating basis functions and a quasi-interpolation operator for the spline space spanned by this basis [12]. In the 2010s Mokriš, Jüttler and Giannelli showed completeness of the basis generated by Kraft's selection mechanism under a certain geometric assumption on the shape of nested domains [15].

PHT-splines which are bicubic splines with continuity C^1 over hierarchical T-meshes and a construction of its basis functions were introduced by Deng et al. in the mid 2000s [4]. T-splines over T-meshes were introduced in the early 2000s by Sederberg et al. [20]. In order to address the problem of linear independence of T-splines, Li et al. introduced AS T-splines [13] and developed a local refinement algorithm for AS T-splines [19] in the early 2010s. LR-splines and an algorithm for generating its basis functions were introduced by Dokken, Lyche and Pettersen in the early 2010s [5]. The

^{*}Mahidol University, Faculty of Science and Centre of Excellence in Mathematics, CHE, Bangkok 10400, Thailand; e-mail: berdinsky@gmail.com.

[†]Mahidol University, Faculty of Science, Bangkok 10400, Thailand; e-mail: prohrakju@gmail.com.

¹This selection mechanism is usually referred to in the literature as Kraft's selection mechanism or Kraft's selection procedure

algorithm involves the dimension formula for spline spaces over T-meshes obtained via homological techniques by Mourrain in the early 2010s [16].

For all the above mentioned four approaches a mesh, by default, partitions a bounded domain $\Omega \subset \mathbb{R}^d$. To the best of our knowledge polynomial splines over meshes admitting local refinement that partition an unbounded domain have not been studied before. Moreover, anyone who would like to store and operate splines over infinite meshes using a computer immediately faces a challenge – a computer can use only a finite amount of memory. To resolve this challenge one needs to require some sort of regularity for both a mesh and a spline.

In this paper we propose a data structure which allows to store and operate polynomial splines over meshes admitting local refinement that partition \mathbb{R}^d , $d \geq 1$, but uses only a finite amount of memory. As a data structure we use a finite directed labelled graph which is a standard way to present a deterministic finite automaton. As an approach to polynomial splines over meshes admitting local refinement we use hierarchical tensor product B-splines. Note that only a special class of polynomial splines can be represented using only a finite amount of memory. This situation is inevitable as, for example, there exist already uncountably many two-level dyadic meshes on \mathbb{R}^d while a collection of all possible finite descriptions, finite strings over a finite alphabet, is countable.

We note that the construction proposed in this paper can be applied to represent splines over finite meshes which are used in applications. This is because regular splines, to be introduced in Section 6, include splines with bounded support. However, the main focus of this paper is on splines with unbounded support over infinite hierarchical meshes that partition \mathbb{R}^d .

Another component used in this paper is the theory of finite automata presentable (FA-presentable) structures in the sense of Khoussainov and Nerode [11]. We use a fundamental fact valid for FA-presentable structures: for a first order definition of a relation over domain of a FA-presentable structure there is an algorithm deciding this relation. In principle all algorithms we obtain in the paper come from this fundamental fact. For more on FA-presentable structures the reader is referred to the most recent survey article [22]. This paper also contributes to theory of FA-presentable structures from an application perspective; see also [8] where some applications of FA-presentable structures are considered.

A cell of a hierarchical mesh we associate with the coordinate tuple of the cell barycentre. Each element of this tuple we present by the string obtained from the standard representation of the abelian group $\mathbb{Z}[1/b] = \{s/b^\ell \mid s, \ell \in \mathbb{Z}, \ell \geq 0\}$ for some positive even integer b ; for example, the case $b = 10$ corresponds to the decimal representation. A tuple is then represented by the convolution of strings. We use the same representation for spline coefficients assuming that they are all in $\mathbb{Z}[1/b]$. However, we note that in principle one can use other representations of abelian subgroups of \mathbb{Q}^d for which the addition operation is FA-recognizable (see Section 3 for definition). For example, one can use nonstandard FA-presentations of abelian groups, see, e.g., [17].

We assume then that a collection of convoluted strings representing a mesh or a spline is a regular language, so it can be encoded by a deterministic finite automaton accepting this language. Now let us be given a deterministic finite automaton, corresponding to either a hierarchical mesh \mathcal{T} or a spline function f over this mesh, as an input. In this paper we describe the following procedures and algorithms:

- a) A N -level hierarchical mesh \mathcal{T} must be defined by a nested sequence of domains $\Omega^0 = \mathbb{R}^d \supseteq \Omega^1 \supseteq \dots \supseteq \Omega^{N-1} \supseteq \Omega^N = \emptyset$, where $\Omega^{N-1} \neq \emptyset$. We provide a verification procedure for this nestedness condition.
- b) Kraft's selection mechanism generate a complete basis of a spline space over hierarchical mesh \mathcal{T} if a certain geometric condition (see Assumption 2) on the shapes of the domains $\Omega^1, \dots, \Omega^{N-1}$ is satisfied [15]. We provide a verification procedure for this geometric condition.

- c) For a hierarchical mesh \mathcal{T} each basis function generated by Kraft's selection mechanism can be naturally associated with a certain cell of this mesh (see Figure 6 for the association rule). We show that a collection of cells (encoded by strings representing their barycentres) forms a regular language and provide a procedure for constructing deterministic finite automaton accepting this language.
- d) We provide an algorithm for computing $f(\bar{x})$, the value of a spline function f at any point $\bar{x} \in \mathbb{Z}[1/b]^d \subset \mathbb{R}^d$ given as an input.
- e) Suppose that a hierarchical mesh \mathcal{T} is refined by selecting a nonempty subdomain $\Omega^N \subseteq \Omega^{N-1}$. The domain Ω^N is composed of cells which we again encode by strings representing their barycentres. We assume that the language of such strings is regular and we are given a deterministic finite automaton recognizing this language. For the refined hierarchical mesh \mathcal{T}' the coefficients of f must be updated. We provide a procedure for constructing deterministic finite automaton corresponding to f with respect to \mathcal{T}' . The procedure is based on Boehm's knot insertion formula.

For verification procedures in a) and b) we provide first order formulas checking the nestedness and the geometric condition in Assumption 2, respectively. Similarly, for the procedure in c) we provide the first order formula defining a regular language. A concrete implementation of an algorithm constructing a deterministic finite automaton accepting this language mostly follows from Theorem 3. For an algorithm computing the value of a spline function at a given point in d) we use the fact that for a FA-recognizable function there is an algorithm computing it in linear time (see Section 3). For e) we simply show how to construct a FA-recognizable relation accepting a spline function over the refined mesh; a concrete implementation again mostly follows from Theorem 3.

In addition, we show that there is a procedure which for a given deterministic finite automata corresponding to given functions f_1 and f_2 constructs a deterministic finite automaton corresponding to the sum $f_1 + f_2$. Similarly, there is a procedure that for a given deterministic finite automaton corresponding to f and a fixed constant $\mu \in \mathbb{Z}[1/b]$ constructs a deterministic finite automaton corresponding to μf (see Section 6).

In Section 2 we define a general class of d -dimensional hierarchical meshes \mathcal{T} to be considered in this paper and spaces of splines of multi-degree (m, \dots, m) with maximum order smoothness, i.e., splines of class C^{m-1} , over these meshes $\mathcal{S}_m(\mathcal{T})$. We recall that in [15] the authors showed that under a certain geometric condition on nested domains defining a hierarchical mesh \mathcal{T} Kraft's selection mechanism generates a complete basis of the spline space $\mathcal{S}_m(\mathcal{T})$. Though the extension of this result for infinite hierarchical meshes that partition \mathbb{R}^d , $d \geq 1$ is straightforward, we present it in Section 2 for the sake of rigour and completeness. Note that for an infinite hierarchical mesh \mathcal{T} the space $\mathcal{S}_m(\mathcal{T})$ is not a finite-dimensional vector space like for the case of finite meshes. So Kraft's selection mechanism does not generate a basis of a spline space in the usual sense. However, for every $f \in \mathcal{S}_m(\mathcal{T})$ the sum $\sum_{\delta \in \mathcal{K}} \lambda_\delta \delta(\bar{x})$ is correctly defined and $f(\bar{x}) = \sum_{\delta \in \mathcal{K}} \lambda_\delta \delta(\bar{x})$ holds for some uniquely defined coefficients λ_δ at every $\bar{x} \in \mathbb{R}^d$, where \mathcal{K} is the collection of tensor product B-splines generated by Kraft's selection mechanism.

In Sections 3 and 4 we recall necessary definitions and facts from automata theory and the theory of FA-presentable structures, respectively. In Section 5 we introduce regular hierarchical meshes. In Subsections 5.1, 5.2 and 5.3 we describe the above mentioned procedures a), b) and c). In Section 6 we introduce regular splines. In Subsections 6.1 and 6.2 we describe the above mentioned algorithm d) and the procedure e). In Subsection 6.3 we show simple examples of regular splines.

2 Splines with Maximum Order Smoothness

For a given integer $\ell \geq 0$, we denote by T^ℓ a bi-infinite knot vector:

$$T^\ell = (\dots, t_{i-1}^\ell, t_i^\ell, t_{i+1}^\ell, \dots),$$

where $t_i^\ell = \frac{i}{2^\ell}$ for $i \in \mathbb{Z}$. Note that T^ℓ is uniform with the same distance between consecutive knots equal to $\frac{1}{2^\ell}$. Let d be a positive integer. We denote by \mathcal{G}_d^ℓ a d -dimensional grid consisting of the hyperplanes $H_{j,i}^\ell = \{(x_1, \dots, x_d) \mid x_j = t_i^\ell\}$ for $j = 1, \dots, d$ and $i \in \mathbb{Z}$. For a given integer $m \geq 0$, a grid \mathcal{G}_d^ℓ defines the set of tensor product B-splines $B_{d,m}^\ell$ each of which is the product:

$$P_{i_1, \dots, i_d, m}^\ell(x_1, \dots, x_d) = N_{i_1, m}^\ell(x_1) \dots N_{i_d, m}^\ell(x_d), \quad (1)$$

where $i_1, \dots, i_d \in \mathbb{Z}$ and, for $i \in \mathbb{Z}$, $N_{i, m}^\ell(t)$ is the i th B-spline basis function of degree m associated to the knot vector T^ℓ which is recursively defined by Cox-de Boor's formula:

$$N_{i, 0}^\ell(t) = \begin{cases} 1, & t_i^\ell \leq t < t_{i+1}^\ell, \\ 0, & \text{otherwise.} \end{cases}, \quad (2)$$

$$N_{i, j}^\ell(t) = \frac{t - t_i^\ell}{t_{i+j}^\ell - t_i^\ell} N_{i, j-1}^\ell(t) + \frac{t_{i+j+1}^\ell - t}{t_{i+j+1}^\ell - t_{i+1}^\ell} N_{i+1, j-1}^\ell(t), \quad (3)$$

where $j = 1, \dots, m$. Each tensor product B-spline $P_{i_1, \dots, i_d, m}^\ell$ has local support:

$$\{(x_1, \dots, x_d) \mid P_{i_1, \dots, i_d, m}^\ell(x_1, \dots, x_d) \neq 0\} = (t_{i_1}^\ell, t_{i_1+m+1}^\ell) \times \dots \times (t_{i_d}^\ell, t_{i_d+m+1}^\ell)$$

on which it takes positive values. Tensor product B-splines from $B_{d,m}^\ell$ are locally linear independent: for every open bounded set $U \subseteq \mathbb{R}^d$ the tensor product B-splines from $B_{d,m}^\ell$ having nonempty intersections of its support with U are linearly independent on U . For introduction to B-splines we refer the reader to, e.g., [18].

For a given ℓ we denote by \mathcal{C}_d^ℓ the collection of all closed d -dimensional cubes $\prod_{j=1}^d [t_{i_j}^\ell, t_{i_j+1}^\ell]$. Following [15] we call each of the cubes from \mathcal{C}_d^ℓ a cell of the grid \mathcal{G}_d^ℓ (or simply a cell). Let us consider a nested sequence of domains

$$\Omega^0 = \mathbb{R}^d \supseteq \Omega^1 \supseteq \dots \supseteq \Omega^{N-1} \supseteq \Omega^N = \emptyset,$$

where $\Omega^{N-1} \neq \emptyset$.

Assumption 1. *We assume that each Ω^ℓ , $\ell = 1, \dots, N-1$ is composed of cells from $\mathcal{C}_d^{\ell-1}$. That is, for each $\ell = 1, \dots, N-1$ there is a subset $M \subseteq \mathcal{C}_d^{\ell-1}$ for which $\Omega^\ell = \bigcup_{c \in M} c$.*

Remark 1. *Note that, since $\Omega_0 = \mathbb{R}^d$, this assumption is exactly equivalent to [15, Assumption 3.1]: each set $\Omega^0 \setminus \Omega^{\ell+1}$ for $\ell = 0, \dots, N-1$ must be composed of cells from \mathcal{C}_d^ℓ . However, in this paper we do not assume that the domains $\Omega_1, \dots, \Omega_{N-1}$ are bounded. They can be bounded or unbounded.*

A hierarchy of domains $\Omega^0 = \mathbb{R}^d \supseteq \Omega^1 \supseteq \dots \supseteq \Omega^{N-1} \supseteq \Omega^N = \emptyset$ satisfying Assumption 1 creates a subdivision of \mathbb{R}^d into the collection of cells $R^\ell \subseteq \mathcal{C}_d^\ell$ such that $\Omega^\ell \setminus \Omega^{\ell+1} = \bigcup_{c \in R^\ell} c$ for $\ell = 0, \dots, N-1$. We denote the subdivision of \mathbb{R}^d into the cells

from R^ℓ , $\ell = 0, \dots, N-1$ by \mathcal{T} . \mathcal{T} is also referred to as a d -dimensional box-partition. If $d = 2$, then \mathcal{T} is also called a T-mesh. We will simply call \mathcal{T} a hierarchical mesh. See Fig. 1 for an example of a 2-dimensional hierarchical mesh generated by a nested sequence of domains $\Omega^0 = \mathbb{R}^2 \supseteq \Omega_1 \supseteq \Omega_2 \supseteq \Omega_3 = \emptyset$. We denote by \mathcal{T}_d the collection of all d -dimensional cells of \mathcal{T} : $\mathcal{T}_d = \bigcup_{\ell=0}^{N-1} R^\ell$.




Figure 1: The figure on the left shows a portion of infinite domains Ω_1 (bounded by blue line segments) and Ω_2 (bounded by red line segments) satisfying Assumption 1. The grid lines of \mathcal{G}_2^0 , \mathcal{G}_2^1 and \mathcal{G}_2^2 are depicted as solid, dashed and dotted lines, respectively. The figure on the right shows the corresponding portion of a hierarchical mesh generated by a nested sequence of domains $\Omega^0 = \mathbb{R}^2 \supseteq \Omega_1 \supseteq \Omega_2 \supseteq \Omega_3 = \emptyset$.

Definition 1. We denote by $\mathcal{S}_m(\mathcal{T})$ the space of functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$ of the class C^{m-1} which are polynomials of multi-degree (m, \dots, m) in every cell from \mathcal{T}_d . That is, for every $c \in \mathcal{T}_d$, $f|_c = \sum_{i_1, \dots, i_d=0}^m a_{i_1 \dots i_d} x_1^{i_1} \dots x_d^{i_d}$. A function from $\mathcal{S}_m(\mathcal{T})$ is called a spline with maximum order of smoothness over \mathcal{T} .

In Definition 1 one can require a weaker assumption for a function f : the derivatives $\frac{\partial^{m-1} f}{\partial x_i^{m-1}}$ exist and continuous everywhere in \mathbb{R}^d for $i = 1, \dots, d$. Proposition 1 below shows that this weaker assumption does not affect the space of functions $\mathcal{S}_m(\mathcal{T})$. Moreover, as the same proposition shows, a stronger assumption: all derivatives $\frac{\partial^{k_1 + \dots + k_d} f}{\partial x_1^{k_1} \dots \partial x_d^{k_d}}$ exist and continuous everywhere in \mathbb{R}^d for $k_i = 0, \dots, m-1$ and $i = 1, \dots, d$, does not affect the space $\mathcal{S}_m(\mathcal{T})$ as well. In order to show this we use Lemma 1 below, see, e.g., [1].

Let us consider two d -dimensional cubes $c_1 = \prod_{j=1}^d [y_j', y_j'']$ and $c_2 = \prod_{j=1}^d [z_j', z_j'']$ in \mathbb{R}^d , i.e., $y_j' < y_j''$ and $z_j' < z_j''$ for $j = 1, \dots, d$. Suppose that the cubes c_1 and c_2 are adjacent such that the intersection $c_1 \cap c_2$ is a $(d-1)$ -dimensional cube, see Fig. 2 for the case of two 2-dimensional cubes in \mathbb{R}^2 . This intersection $c_1 \cap c_2$ is contained in some $(d-1)$ -dimensional hyperplane $x_i = x_0$ for some integer $i \in [1, d]$ and a constant $x_0 \in \mathbb{R}$. Let $f : c_1 \cup c_2 \rightarrow \mathbb{R}$ be a function such that $f|_{c_1} = p_1$ and $f|_{c_2} = p_2$ for some polynomials $p_1(x_1, \dots, x_d)$ and $p_2(x_1, \dots, x_d)$ of multi-degree (m, \dots, m) .

Lemma 1 ([1]). The derivative $\frac{\partial^k f}{\partial x_i^k}$ exists and continuous everywhere in $\text{dom } f = c_1 \cup c_2$ for some integer $0 \leq k \leq m-1$ if and only if $p_1 - p_2 = \lambda(x_i - x_0)^{k+1}$, where λ is a polynomial.

Proposition 1. Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a function which is a polynomial of multi-degree (m, \dots, m) in every cell from \mathcal{T}_d . Suppose that the derivatives $\frac{\partial^{m-1} f}{\partial x_i^{m-1}}$ exist and continuous everywhere in \mathbb{R}^d for $i = 1, \dots, d$. Then the derivatives $\frac{\partial^{k_1 + \dots + k_d} f}{\partial x_1^{k_1} \dots \partial x_d^{k_d}}$ exist and continuous for $k_i = 0, \dots, m-1$ and $i = 1, \dots, d$.

Proof. Let c_1 and c_2 be two adjacent cells from \mathcal{T}_d for which the intersection is a $(d-1)$ -dimensional cube contained in some hyperplane $x_i = x_0$. Let p_1 and p_2 be the polynomials of multi-degree (m, \dots, m) which are the restrictions of f to the cells




Figure 2: Adjacent 2-dimensional cells c_1 and c_2 for which the intersection $c_1 \cap c_2$ is the closed 1-dimensional line segment with the endpoints (y''_1, z'_2) and (z'_1, y''_2) .

c_1 and c_2 , respectively. Since the derivative $\frac{\partial^{m-1} f}{\partial x_i^{m-1}}$ exists and continuous, by Lemma 1 we obtain that $p_1 - p_2 = \lambda(x_i - x_0)^m$, where λ is a polynomial which depends on the variables $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_d$. Therefore, for every d -tuple of nonnegative integers (k_1, \dots, k_d) for which $k_i \leq m - 1$ we have that $\frac{\partial^{k_1 + \dots + k_d} p_1}{\partial x_1^{k_1} \dots \partial x_d^{k_d}} - \frac{\partial^{k_1 + \dots + k_d} p_2}{\partial x_1^{k_1} \dots \partial x_d^{k_d}} = \mu(x_i - x_0)^{m-k_i}$ for some polynomial μ . So, by Lemma 1, for each of such d -tuples (k_1, \dots, k_d) the derivative $\frac{\partial^{k_1 + \dots + k_d} f}{\partial x_1^{k_1} \dots \partial x_d^{k_d}}$ exists and continuous on $c_1 \cup c_2$. Considering all pairs of adjacent cells from \mathcal{T}_d which intersect in $(d-1)$ -dimensional cubes we see that the derivative $\frac{\partial^{k_1 + \dots + k_d} f}{\partial x_1^{k_1} \dots \partial x_d^{k_d}}$ exists and continuous everywhere in \mathbb{R}^d for every d -tuple of nonnegative integers (k_1, \dots, k_d) for which $k_j \leq m - 1$ for all $j = 1, \dots, d$. \square

Corollary 1. For every function $f \in \mathcal{S}_m(\mathcal{T})$ the derivatives $\frac{\partial^{k_1 + \dots + k_d} f}{\partial x_1^{k_1} \dots \partial x_d^{k_d}}$ exist and continuous for $k_i = 0, \dots, m - 1$ and $i = 1, \dots, d$.

Proof. The statement of the corollary directly follows from Proposition 1. \square

Remark 2. We note that Proposition 1 and Corollary 1 hold valid for every subdivision of \mathbb{R}^d into d -dimensional axis-aligned cubes, not necessarily the ones obtained from the grids \mathcal{G}_d^ℓ , $\ell = 0, \dots, N - 1$ and a nested sequence of domains $\Omega^0 = \mathbb{R}^d \supseteq \Omega^1 \supseteq \dots \supseteq \Omega^{N-1} \supseteq \Omega^N = \emptyset$ satisfying Assumption 1.

For a given $\ell = 0, \dots, N - 1$ let $\mathcal{M}^\ell = \Omega^0 \setminus \Omega^{\ell+1} = \mathbb{R}^d \setminus \Omega^{\ell+1}$. Then we have a nested sequence of domains $\mathcal{M}^{-1} = \emptyset, \mathcal{M}^0 = \mathbb{R}^d \setminus \Omega^1, \mathcal{M}^1 = \mathbb{R}^d \setminus \Omega^2, \dots, \mathcal{M}^{N-2} = \mathbb{R}^d \setminus \Omega^{N-1}, \mathcal{M}^{N-1} = \mathbb{R}^d$:

$$\emptyset = \mathcal{M}^{-1} \subseteq \mathcal{M}^0 \subseteq \mathcal{M}^1 \subseteq \dots \subseteq \mathcal{M}^{N-2} \subseteq \mathcal{M}^{N-1} = \mathbb{R}^d.$$

By Assumption 1, each domain \mathcal{M}^ℓ is composed of the cells from \mathcal{C}_d^ℓ for $\ell = 0, \dots, N - 1$. Below we recall some necessary definitions and results (see Definitions 2.3, 2.7 and 2.8, Lemma 2.4 and Proposition 2.10 in [15]).

Definition 2 ([15]). Let c_1 and c_2 be two different cells from \mathcal{C}_d^ℓ for some $\ell \geq 0$ such that the intersection $c_1 \cap c_2$ is non-empty. Let $p_1(x_1, \dots, x_d)$ and $p_2(x_1, \dots, x_d)$ be two polynomials of multi-degree (m, \dots, m) . We say that $p_1|_{c_1} \sim p_2|_{c_2}$ if $\frac{\partial^{k_1 + \dots + k_d} p_1}{\partial x_1^{k_1} \dots \partial x_d^{k_d}}(x_1, \dots, x_d) = \frac{\partial^{k_1 + \dots + k_d} p_2}{\partial x_1^{k_1} \dots \partial x_d^{k_d}}(x_1, \dots, x_d)$ for all $(x_1, \dots, x_d) \in c_1 \cap c_2$ and $k_i = 0, \dots, m - 1$ for $i = 1, \dots, d$.

Let $p(x_1, \dots, x_m)$ be a polynomial of multi-degree (m, \dots, m) and $c \in \mathcal{C}_d^\ell$ be a cell. The restriction $p|_c$ always can be expressed as a linear combination $p|_c = \sum_{\beta \in B_{d,m}^\ell} \lambda_c^\beta(p|_c) \beta|_c$, where $\lambda_c^\beta(p|_c)$ denote the coefficients of this linear combination.

Clearly, $\lambda_c^\beta(p|_c) = 0$ if $c \not\subset \overline{\text{supp } \beta}$. That is, a coefficient $\lambda_c^\beta(p|_c)$ can be nontrivial only if $c \subset \overline{\text{supp } \beta}$, so in the formal sum $\sum_{\beta \in B_{d,m}^\ell} \lambda_c^\beta(p|_c) \beta|_c$ only at most $(m+1) \times (m+1)$

terms can be nontrivial. The following lemma characterizes the relation $p_1|_{c_1} \sim p_2|_{c_2}$ in terms of the coefficients $\lambda_{c_1}^\beta(p_1|_{c_1})$ and $\lambda_{c_2}^\beta(p_2|_{c_2})$.

Lemma 2 ([15]). *Let c_1, c_2 and p_1, p_2 be two cells and two polynomials from Definition 2. Then $p_1|_{c_1} \sim p_2|_{c_2}$ if and only if for every $\beta \in B_{d,m}^\ell$, for which $\beta|_{c_1 \cap c_2} \neq 0$, $\lambda_{c_1}^\beta(p_1|_{c_1}) = \lambda_{c_2}^\beta(p_2|_{c_2})$.*

Definition 3 ([15]). *Let M be a collection of cells $M \subseteq \mathcal{C}_d^\ell$ for some $\ell \geq 0$ and \mathcal{M} be the domain covered by the cells from M : $\mathcal{M} = \bigcup_{c \in M} c$. We denote by $\mathbb{S}_m(M)$ the space of functions $f : \mathcal{M} \rightarrow \mathbb{R}$ which are polynomials of multi-degree (m, \dots, m) in every cell from M and for every pair of cells $c_1, c_2 \in M$ having nonempty intersection $c_1 \cap c_2 \neq \emptyset$: $f|_{c_1} \sim f|_{c_2}$.*

Definition 4 ([15]). *Let $\beta \in B_{d,m}^\ell$ and $M \subseteq \mathcal{C}_d^\ell$ for some $\ell \geq 0$. The coefficient graph Γ_β is defined as follows. The vertices of Γ_β are the cells $c \in M$ for which $c \subset \overline{\text{supp } \beta}$. Two vertices c_1 and c_2 in Γ_β are connected by an edge if $\beta|_{c_1 \cap c_2} \neq 0$.*




Figure 3: The figure on the left shows the cells of M (shaded in gray), the closure of the support $\overline{\text{supp } \beta}$ (bounded by red line segments) and the cells of M which are subsets of $\overline{\text{supp } \beta}$ (these cells are labeled by c_1, \dots, c_8). The figure on the right shows the graph Γ_β which consists of three connected components with the sets of vertices $\{c_2\}$, $\{c_1, c_3\}$ and $\{c_4, c_5, c_6, c_7, c_8\}$.

See Fig. 3 for an example of a graph Γ_β for $d = 2$ and $m = 3$ (in this case the support of a tensor product B-spline β is composed of 4×4 cells).

Let $M \subseteq \mathcal{C}_d^\ell$ and $\mathcal{M} = \bigcup_{c \in M} c$ be the domain covered by the cells from M . The following proposition is proved in [15] assuming, by default, that M is finite. Below we recall the original proof [15, Proposition 2.10] to show that one does not need to assume that M is finite.

Proposition 2 ([15]). *Let $f : \mathcal{M} \rightarrow \mathbb{R}$ be a function which is a polynomial of multi-degree (m, \dots, m) in every cell of M . Then $f \in \mathbb{S}_m(M)$ if and only if $\lambda_{c_1}^\beta(f|_{c_1}) = \lambda_{c_2}^\beta(f|_{c_2})$ for all $\beta \in B_{d,m}^\ell$ and c_1, c_2 belonging to the same connected component of Γ_β .*

Proof. Assume that there exist $\beta \in B_{d,m}^\ell$ and two cells c_1, c_2 in the same connected component of Γ_β for which $\lambda_{c_1}^\beta(f|_{c_1}) \neq \lambda_{c_2}^\beta(f|_{c_2})$. Then there exist two vertices (cells)

k_1 and k_2 of the same connected component of Γ_β for which $\lambda_{k_1}^\beta(f|_{k_1}) \neq \lambda_{k_2}^\beta(f|_{k_2})$. Since $f|_{k_1} \sim f|_{k_2}$, we get a contradiction with Lemma 2.

Now assume that $\lambda_{c_1}^\beta(f|_{c_1}) = \lambda_{c_2}^\beta(f|_{c_2})$ for all $\beta \in B_{d,m}^\ell$ and c_1, c_2 belonging to the same connected component of Γ_β . Therefore, by Lemma 2, for every $c, c' \in M$ having nonempty intersection $c \cap c' \neq \emptyset$, $f|_c \sim f|_{c'}$ which implies that $f \in \mathbb{S}_m(M)$. \square

Let $M \subseteq \mathcal{C}_d^\ell$, $\mathcal{M} = \bigcup_{c \in M} c$ and $\beta \in B_{d,m}^\ell$. Following the notation in [15] we denote by $K(\Gamma_\beta)$ the set of connected components of a graph Γ_β . For a given connected component $\Phi \in K(\Gamma_\beta)$ we denote by β_Φ the function $\beta_\Phi : \mathbb{R}^d \rightarrow \mathbb{R}$ defined as follows:

$$\beta_\Phi(\bar{x}) = \sum_{c \in V(\Phi)} \beta(\bar{x}) \chi_c^*(\bar{x}), \quad (4)$$

where $V(\Phi)$ is the set of vertices (cells) of the graph Φ , $\bar{x} = (x_1, \dots, x_d) \in \mathbb{R}^d$ and $\chi_c^*(\bar{x})$ is a normalized characteristic function: $\chi_c^*(\bar{x}) = \frac{\chi_c(\bar{x})}{\sum_{k \in M} \chi_k(\bar{x})}$ if $\bar{x} \in c$ and $\chi_c^*(\bar{x}) = 0$, otherwise. Clearly, $\beta_\Phi(\bar{x}) = \beta(\bar{x})$ if $\bar{x} \in \mathcal{M}$ so $\beta_\Phi|_{\mathcal{M}} = \beta|_{\mathcal{M}}$; $\beta_\Phi(\bar{x}) = 0$ if $\bar{x} \notin \mathcal{M}$.

For given $\ell \geq 0, d > 0, m > 0$ and $M \subseteq \mathcal{C}_d^\ell$ we denote by Δ the collection of $\beta_\Phi|_{\mathcal{M}}$ for all $\beta \in B_{d,m}^\ell$ and $\Phi \in K(\Gamma_\beta)$:

$$\Delta = \{\beta_\Phi|_{\mathcal{M}} \mid \beta \in B_{d,m}^\ell, \Phi \in K(\Gamma_\beta)\}.$$

It follows from the local linear independence of $B_{d,m}^\ell$ that Δ is locally linear independent. Assuming that M is finite, [15, Theorem 2.12] shows that Δ is a basis of a vector space $\mathbb{S}_m(M)$. When M is finite, Δ is finite so $\mathbb{S}_m(M)$ is a finite dimensional vector space. If M is infinite, then Δ is infinite. However, Δ is not a basis of $\mathbb{S}_m(M)$ if M is infinite. Nevertheless, for any collection of coefficients $\lambda_\delta, \delta \in \Delta$, the formal sum $\sum_{\delta \in \Delta} \lambda_\delta \delta$ is correctly defined as for every $x \in \mathcal{M}$ there are only finitely many $\delta \in \Delta$ for which $\delta(x) \neq 0$. So $\sum_{\delta \in \Delta} \lambda_\delta \delta$ is a function from $\mathbb{S}_m(M)$. Theorem 1 below shows that every function $f \in \mathbb{S}_m(M)$ can be uniquely represented as a formal sum $\sum_{\delta \in \Delta} \lambda_\delta \delta$. The proof of this theorem repeats the argument of [15, Theorem 2.12].

Theorem 1 ([15]). *For every $f \in \mathbb{S}_m(M)$, $f = \sum_{\delta \in \Delta} \lambda_\delta \delta$ for some uniquely defined coefficients $\lambda_\delta, \delta \in \Delta$.*

Proof. Let us consider $f \in \mathbb{S}_m(M)$. For every cell $c \in M$ we have that $f|_c = \sum_{\beta \in B_{d,m}^\ell} \lambda_c^\beta(f|_c) \beta|_c$. Therefore, $f(\bar{x}) = \sum_{\beta \in B_{d,m}^\ell} \lambda_c^\beta(f|_c) \beta(\bar{x})$ for $\bar{x} = (x_1, \dots, x_d) \in c$. Also, we have that $f(\bar{x}) = \sum_{c \in M} f(\bar{x}) \chi_c^*(\bar{x})$ for $x \in \mathcal{M}$. Therefore,

$$\begin{aligned} f(\bar{x}) &= \sum_{c \in M} \sum_{\beta \in B_{d,m}^\ell} \lambda_c^\beta(f|_c) \beta(\bar{x}) \chi_c^*(\bar{x}) = \sum_{\beta \in B_{d,m}^\ell} \sum_{c \in \Gamma_\beta} \lambda_c^\beta(f|_c) \beta(\bar{x}) \chi_c^*(\bar{x}) = \\ &= \sum_{\beta \in B_{d,m}^\ell} \sum_{\Phi \in K(\Gamma_\beta)} \sum_{c \in V(\Phi)} \lambda_c^\beta(f|_c) \beta(\bar{x}) \chi_c^*(\bar{x}) \end{aligned}$$

for $\bar{x} \in \mathcal{M}$. By Proposition 2, for fixed β and $\Phi \in K(\Gamma_\beta)$ the coefficients $\lambda_c^\beta(f|_c)$ are the same for all $c \in V(\Phi)$. Let us denote it by $\lambda_\Phi^\beta(f)$. Therefore, by (4), we obtain $f(\bar{x}) = \sum_{\beta \in B_{d,m}^\ell} \sum_{\Phi \in K(\Gamma_\beta)} \lambda_\Phi^\beta(f) \beta_\Phi(\bar{x})$ for $\bar{x} \in \mathcal{M}$. Changing β_Φ to δ and λ_Φ^β to λ_δ , in

the latter identity we conclude that $f = \sum_{\delta \in \Delta} \lambda_\delta \delta$. The uniqueness of coefficients λ_δ immediately follows from the local linear independence of Δ . \square

Let us denote by Δ^* the collection of $\beta|_{\mathcal{M}}$ for all $\beta \in B_{d,m}^\ell$ for which $\text{supp } \beta \cap \mathcal{M} \neq \emptyset$:

$$\Delta^* = \{\beta|_{\mathcal{M}} \mid \beta \in B_{d,m}^\ell, \text{supp } \beta \cap \mathcal{M} \neq \emptyset\}.$$

The analogue of [15, Corollary 2.13] which includes the case of infinite domains is as follows.

Corollary 2 ([15]). *If for each $\beta \in B_{d,m}^\ell$, for which $\text{supp } \beta \cap \mathcal{M} \neq \emptyset$, the intersection $\overline{\text{supp } \beta} \cap \mathcal{M}$ is connected, then every $f \in \mathbb{S}_m(M)$ is equal to $f = \sum_{\delta \in \Delta^*} \lambda_\delta \delta$ for some uniquely defined coefficients $\lambda_\delta, \delta \in \Delta^*$.*

Proof. If the assumption of the corollary is satisfied, then $\Delta = \Delta^*$. So the corollary follows directly from Theorem 1. \square

Now we are ready to show the analogue of [15, Theorem 3.5] for an infinite hierarchical mesh \mathcal{T} . Similarly to [15] we make the following additional assumption.

Assumption 2. *For a nested sequence of domains:*

$$\emptyset = \mathcal{M}^{-1} \subseteq \mathcal{M}^0 \subseteq \mathcal{M}^1 \subseteq \dots \subseteq \mathcal{M}^{N-2} \subseteq \mathcal{M}^{N-1} = \mathbb{R}^d$$

we assume that for each $\ell = 0, \dots, N-2$ the domain \mathcal{M}^ℓ satisfies the condition of Corollary 2, that is, for each $\beta \in B_{d,m}^\ell$, $\ell = 0, \dots, N-2$, for which $\text{supp } \beta \cap \mathcal{M}^\ell \neq \emptyset$, the intersection $\overline{\text{supp } \beta} \cap \mathcal{M}^\ell$ is connected.

We say that a hierarchical mesh \mathcal{T} satisfies Assumption 2 if it is generated by a nested sequence of domains $\Omega^0 = \mathbb{R}^d \supseteq \Omega^1 \supseteq \dots \supseteq \Omega^{N-1} \supseteq \Omega^N = \emptyset$ for which the domains $\mathcal{M}^\ell = \Omega^0 \setminus \Omega^{\ell+1}$, $\ell = 0, \dots, N-2$ satisfy Assumption 2.

For $\ell = 0, \dots, N-1$, let:

$$\mathcal{K}^\ell = \{\beta \in B_{d,m}^\ell \mid \text{supp } \beta \cap \mathcal{M}^{\ell-1} = \emptyset \wedge \text{supp } \beta \cap \mathcal{M}^\ell \neq \emptyset\} \text{ and } \mathcal{K} = \bigcup_{\ell=0}^{N-1} \mathcal{K}^\ell. \quad (5)$$

Remark 3. *The equation (5) defines a procedure usually known as Kraft's selection mechanism for generating basis functions. Informally, it can be described as follows. At the first iteration this mechanism takes all tensor product B-splines from $B_{d,m}^0$ (they are all tensor product B-splines with respect to the grid \mathcal{G}_d^0 with the support overlapping with the domain $\Omega_0 = \mathbb{R}^d$). At the second iteration it removes all tensor product B-splines with the support in the domain Ω_1 obtained at the previous iteration and add tensor product B-splines from $B_{d,m}^1$ with the support in the domain Ω_1 . At the third iteration it removes all tensor product B-splines with the support in the domain Ω_2 obtained at the previous iteration and add tensor product B-splines from $B_{d,m}^2$ with the support in the domain Ω_2 and etc. The process stops after the N th iteration.*

Each formal sum $\sum_{\delta \in \mathcal{K}} \lambda_\delta \delta$ defines a function from $\mathcal{S}_m(\mathcal{T})$. Theorem 2 below shows that every function $f \in \mathcal{S}_m(\mathcal{T})$ can be uniquely represented as a formal sum $\sum_{\delta \in \Delta} \lambda_\delta \delta$. The main argument of this theorem repeats the standard argument for the case of bounded domains, the reader may look it up in [15, Theorem 3.5].

Theorem 2 ([15]). *Assume that a hierarchical mesh \mathcal{T} satisfies Assumption 2. Then for every $f \in \mathcal{S}_m(\mathcal{T})$, $f = \sum_{\delta \in \mathcal{K}} \lambda_\delta \delta$ for some uniquely defined coefficients λ_δ .*

Proof. For $\ell = 0, \dots, N-1$ we denote by \mathcal{M}^ℓ the collection of cells from \mathcal{C}_d^ℓ covering a domain \mathcal{M}^ℓ : $\mathcal{M}^\ell = \bigcup_{c \in \mathcal{M}^\ell} c$. Let us consider $f \in \mathcal{S}_m(\mathcal{T})$. By Corollary 1, $f|_{\mathcal{M}^0} \in$

$\mathbb{S}_m(M^0)$. Therefore, by Corollary 2, $f|_{\mathcal{M}^0} = \sum_{\delta \in \mathcal{K}^0} \lambda_\delta \delta|_{\mathcal{M}^0}$ for some coefficients $\lambda_\delta, \delta \in \mathcal{K}^0$. We denote by f_0 the function $f_0 = \sum_{\delta \in \mathcal{K}^0} \lambda_\delta \delta$. Let consider now the function $(f - f_0) \in \mathcal{S}_m(\mathcal{T})$. By Corollary 1, $(f - f_0)|_{\mathcal{M}^1} \in \mathbb{S}_m(M^1)$. Therefore, by Corollary 2 and the identity $(f - f_0)|_{\mathcal{M}^0} = 0$, we obtain that $(f - f_0)|_{\mathcal{M}^1} = \sum_{\delta \in \mathcal{K}^1} \lambda_\delta \delta|_{\mathcal{M}^1}$ for some coefficients $\lambda_\delta, \delta \in \mathcal{K}^1$. We denote by f_1 the function $f_1 = \sum_{\delta \in \mathcal{K}^1} \lambda_\delta \delta$. Clearly, the function $(f - f_0 - f_1) \in \mathcal{S}_m(\mathcal{T})$ and $(f - f_0 - f_1)|_{\mathcal{M}^1} = 0$, so $(f - f_0 - f_1)|_{\mathcal{M}^2} = \sum_{\delta \in \mathcal{K}^2} \lambda_\delta \delta|_{\mathcal{M}^2}$ for some coefficients $\lambda_\delta, \delta \in \mathcal{K}^2$. We define $f_2 = \sum_{\delta \in \mathcal{K}^2} \lambda_\delta \delta$.

We repeat the process until all functions $f_i = \sum_{\delta \in \mathcal{K}^i} \lambda_\delta \delta, i = 0, \dots, N - 1$ are constructed. Finally, we have the identity $(f - \sum_{i=0}^{N-1} f_i)|_{\mathcal{M}^{N-1}} = 0$ which implies that $f = \sum_{i=0}^{N-1} f_i$, so $f = \sum_{\delta \in \mathcal{K}} \lambda_\delta \delta$. The uniqueness of coefficients λ_δ follows from local linear independence of tensor product B-splines. \square

3 Multitape Synchronous Finite Automata

Let us first recall the notion of finite automata and regular languages, see, e.g., [9]. Let Σ be a finite alphabet. We say that w is a string over the alphabet Σ if w is a finite sequence of symbols $\sigma_1 \sigma_2 \dots \sigma_n$, where $\sigma_i \in \Sigma$ for $i = 1, \dots, n$ and n is a nonnegative integer. We denote by $|w|$ the length of the string w : $|w| = n$. If $n = 0$, w is the empty string which we denote by ε . A collection of all strings over the alphabet Σ is denoted by Σ^* .

A nondeterministic finite automaton \mathcal{M} over the alphabet Σ consists of a finite set of states S , a set of initial states $I \subseteq S$, a transition functions $T : S \times \Sigma \rightarrow \mathcal{P}(S)$ and a set of accepting states $F \subseteq S$. The automaton \mathcal{M} accepts a string $w = \sigma_1 \dots \sigma_n$ if there exists a sequence of states $s_1, \dots, s_{n+1} \in S$ for which $s_1 \in I, s_{i+1} \in T(s_i, \sigma_i)$ for all $i = 1, \dots, n$ and $s_{n+1} \in F$. We say that a language $L \subseteq \Sigma^*$ is recognized by \mathcal{M} if L consists of all strings accepted by \mathcal{M} . A language recognized by a nondeterministic finite automaton is called regular. \mathcal{M} is called a deterministic finite automaton if for each state $s \in S$ and a symbol $\sigma \in \Sigma$ the set $T(s, \sigma)$ has exactly one element. Both deterministic and nondeterministic finite automata have the same computational power - they recognize the class of regular languages.

We denote by $\Sigma_\diamond = \Sigma \cup \{\diamond\}$ the alphabet $\Sigma_\diamond = \Sigma \cup \{\diamond\}$; it is assumed that the padding symbol \diamond is not in the alphabet Σ . We denote by Σ_\diamond^k the Cartesian product of k copies of Σ_\diamond . Let $w_1, \dots, w_k \in \Sigma^*$ be some strings over the alphabet Σ . The convolution $w = w_1 \otimes \dots \otimes w_k$ of the strings w_1, \dots, w_k is the string w over the alphabet $\Sigma_\diamond^k \setminus \{(\diamond, \dots, \diamond)\}$ such that for the i th symbol $(\sigma_i^1, \dots, \sigma_i^k)$ of w the symbol σ_i^j is the i th symbol of w_j if $i \leq |w_j|$ and \diamond , otherwise, for $i = 1, \dots, |w|$ and $j = 1, \dots, k$, where $|w| = \max\{|w_j| \mid j = 1, \dots, k\}$.

For example, the convolution of three strings $w_1 = 0001101, w_2 = 10100101110$ and $w_3 = 100101$ is as follows:

$$w_1 \otimes w_2 \otimes w_3 = \begin{array}{cccccccccccc} 0 & 0 & 0 & 1 & 1 & 0 & 1 & \diamond & \diamond & \diamond & \diamond \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 & 1 & \diamond & \diamond & \diamond & \diamond & \diamond \end{array}.$$

For a given relation $R \subseteq \Sigma^{*k}$, we denote by $\otimes R$ the relation:

$$\otimes R = \{w_1 \otimes \dots \otimes w_k \mid (w_1, \dots, w_k) \in R\} \subseteq \left(\Sigma_\diamond^k \setminus \{(\diamond, \dots, \diamond)\} \right)^*.$$

We say that a relation R is FA-recognizable if $\otimes R$ is a regular language over the alphabet $\Sigma_\diamond^k \setminus \{(\diamond, \dots, \diamond)\}$. One can think of a finite automaton recognizing $\otimes R$ as a read-only k -tape Turing machine with the input w_i written on the i th tape for each $i = 1, \dots, k$ and k heads moving synchronously from the left to the right until the whole input is read; after that the input is either accepted or rejected. Such an automaton is also called a k -tape synchronous finite automaton.

Let $f : D \rightarrow \Sigma^{*m}$ be a function from $D \subseteq \Sigma^{*n}$ to Σ^{*m} for any integers $n, m \geq 1$. We denote by $\text{Graph}(f)$ the graph of f , i.e., $\text{Graph}(f) = \{(\bar{u}, \bar{v}) \in \Sigma^{*n} \times \Sigma^{*m} \mid f(\bar{u}) = \bar{v}\} \subseteq \Sigma^{*(n+m)}$. We say that f is FA-recognizable, if $\text{Graph}(f)$ is FA-recognizable. Clearly, if f is FA-recognizable, $D \subseteq \Sigma^{*n}$ must be FA-recognizable. Assuming that $n = m = 1$, for a FA-recognizable function $f : D \rightarrow \Sigma^*$, where $D \subseteq \Sigma^*$, there is a linear-time algorithm which for a given input \bar{u} returns the output $\bar{v} = f(\bar{u})$, see, e.g., the proof of [6, Theorem 2.3.10]. Moreover, there is a characterization of FA-recognizable functions $f : D \rightarrow \Sigma^*$ as functions computed by a deterministic position-faithful one-tape Turing machine in linear time, see below.

A position-faithful one-tape Turing machine is a Turing machine which uses a semi-infinite tape with the left-most position containing the special symbol \boxplus which only occurs at this position and cannot be modified. The initial configuration of the tape is $\boxplus u \square^\infty$, where \square is a special blank symbol, and $u \in \Sigma^*$ for some alphabet Σ with $\Sigma \cap \{\boxplus, \square\} = \emptyset$. During the computation the Turing machine operates as usual, reading and writing cells to the right of the \boxplus symbol.

A function $f : D \rightarrow \Sigma^*$ from a regular domain $D \subseteq \Sigma^*$ to Σ^* is said to be computed by a position-faithful one-tape Turing machine, if when started with tape content being $\boxplus u \square^\infty$, where $u \in D$, the head initially being at \boxplus , the Turing machine eventually reaches an accepting state (and halts), with the tape content starting with $\boxplus f(u) \square$. There is no restriction on the output beyond the first appearance of \square .

Case, Jain, Seah and Stephan showed that a function $f : D \rightarrow \Sigma^*$, $D \subseteq \Sigma^*$, is FA-recognizable if and only if it is computed by a deterministic position-faithful one-tape Turing machine in linear time [3]. This characterization of FA-recognizable functions $f : D \rightarrow \Sigma^*$, $D \subseteq \Sigma^*$, apparently, holds valid for FA-recognizable functions $f : D \rightarrow \Sigma^{*m}$, $D \subseteq \Sigma^{*n}$. Namely, a function $f : D \rightarrow \Sigma^{*m}$, for $D \subseteq \Sigma^{*n}$, is FA-recognizable if and only if it is computed by a deterministic position-faithful one-tape Turing machine in linear time, where the input is a string from $\otimes D$ and the output is a string from $\otimes \Sigma^{*m}$. Furthermore, their result clearly holds valid for multivalued FA-recognizable functions f if it is assumed that the number of values that f can take for each argument in D is bounded from above by some fixed constant. That is, a multivalued function $f : D \rightarrow \Sigma^{*m}$, for $D \subseteq \Sigma^{*n}$, satisfying this assumption, is FA-recognizable if and only if for a given input $u \in \otimes D$ all values $f(u)$ are computed by a deterministic position-faithful one-tape Turing machine in linear time.

4 FA-Presented Structures

Now we are ready to discuss a key ingredient to be used in the following sections 5 and 6. Let us first recall the notion of FA-presented² structures as it was introduced by Khoussainov and Nerode [11]. Let

$$\mathcal{A} = (A; R_1^{m_1}, \dots, R_s^{m_s}, f_1^{n_1}, \dots, f_r^{n_r}, c_1, \dots, c_t)$$

be a structure, where $A \subseteq \Sigma^*$ for some alphabet Σ^* , $R_i^{m_i} \subseteq A^{m_i}$ are m_i -ary relations for $i = 1, \dots, s$, $f_j^{n_j} : A^{n_j} \rightarrow A$ are n_j -ary functions for $j = 1, \dots, r$, and c_k are constants for $k = 1, \dots, t$. The structure \mathcal{A} is said to be FA-presented if A is regular, the relations $R_i^{m_i}$ and the functions $f_j^{n_j}$ are FA-recognizable for all $i = 1, \dots, s$ and $j = 1, \dots, r$. A structure is said to be FA-presentable if it is isomorphic to a FA-presented structure.

²FA is a short for finite automata.

FA-presented structures enjoy the following fundamental properties, see [11, Corollary 4.2]. There exists an effective procedure that for a given first order definition of a relation R of a FA-presented structure \mathcal{A} yields an algorithm deciding R . The first order theory of a FA-presented structure \mathcal{A} is decidable. The proof of these two properties follows from the standard facts in automata theory which can be summarized as follows.

Theorem 3. (see [11, Theorem 4.4]) (1) Let R_1, R_2 and R be FA-recognizable relations. Then the relations corresponding to the expressions $(R_1 \vee R_2)$, $(R_1 \wedge R_2)$, $(R_1 \rightarrow R_2)$, $(\neg R_1)$, $\exists vR$ and $\forall vR$ are also FA-recognizable, where for a k -ary relation $R(v_1, \dots, v_k)$, for $k > 1$, and a variable v_i , $i = 1, \dots, k$:

$$\exists v_i R = \{(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_k) \mid (v_1, \dots, v_{i-1}, v_i, v_{i+1}, \dots, v_k) \in R\},$$

$$\forall v_i R = \{(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_k) \mid \forall v_i \in A((v_1, \dots, v_{i-1}, v_i, v_{i+1}, \dots, v_k) \in R)\}.$$

(2) The emptiness problem for finite automaton is decidable. That is, for a unary FA-recognizable relation $R(v)$ there is an algorithm which for a given deterministic finite automaton accepting R decides whether $\exists vR$ is true or false. Similarly, there is an algorithm deciding whether $\forall vR$ is true or false.

(3) There exists a procedure which for deterministic multi-tape synchronous finite automata recognizing R_1, R_2 and a k -ary relation $R(v_1, \dots, v_k)$, for $k > 1$, constructs deterministic multi-tape synchronous finite automata for recognizing the relations corresponding to the expressions $(R_1 \vee R_2)$, $(R_1 \wedge R_2)$, $(R_1 \rightarrow R_2)$, $(\neg R_1)$, $\exists v_i R$ and $\forall v_i R$ for $i = 1, \dots, k$.

A brief sketch of the proof of Theorem 3 is as follows. Part (1) follows from part (3). Part (2) for $\exists vR$ is the standard fact from automata theory, see, e.g., [9, Theorem 3.7]. For $\forall vR$ it follows from the equivalency of \forall and the composition $\neg \circ \exists \circ \neg$ (see the same argument used in the proof of [6, Theorem 1.4.6]). As for part (3), it is enough to show it only for the expressions $(R_1 \wedge R_2)$, $(\neg R_1)$ and $\exists vR$. For the expression $(R_1 \wedge R_2)$ it follows from the product construction for a deterministic finite automaton accepting the intersection of two regular languages. For $(\neg R_1)$ it follows from a construction of a deterministic finite automaton accepting the complement of a given regular language by swapping accepting and non-accepting states. For $\exists v_i R$ it follows from the standard Rabin-Scott powerset construction for converting a nondeterministic finite automaton into deterministic finite automaton (see, e.g., [9, Theorem 2.1]).

5 Regular Hierarchical Meshes

Let b be a positive integer divisible by 2. We denote by $\mathbb{Z}[1/b]$ the abelian group of all rational numbers of the form $\frac{s}{b^\ell}$ for $s, \ell \in \mathbb{Z}$ and $\ell \geq 0$. Each positive $z \in \mathbb{Z}[1/b]$ can be uniquely represented as the sum of its integral and fractional parts:

$$z = [z]_i + [z]_f = \sum_{i=1}^k \alpha_i b^{i-1} + \sum_{i=1}^k \beta_i b^{-i}, \quad (6)$$

where $\alpha_i, \beta_j \in \{0, 1, \dots, b-1\}$ for all $i = 1, \dots, k$ for which either $\alpha_k \neq 0$ or $\beta_k \neq 0$.

Let Σ_b be the alphabet consisting of the symbols $\frac{\alpha}{\beta}$, where $\alpha, \beta \in \{0, 1, \dots, b-1\}$.

Now, for a given positive $z \in \mathbb{Z}[1/b]$ we represent it as a string:

$$\begin{array}{cccccc} 0 & \alpha_1 & \alpha_2 & \dots & \alpha_k & \\ 0 & \beta_1 & \beta_2 & \dots & \beta_k & \end{array} \quad (7)$$

over the alphabet Σ_b . The first symbol $\begin{smallmatrix} 0 \\ 0 \end{smallmatrix}$ indicates that z is positive. Let $z \in \mathbb{Z}[1/b]$ be negative and $-z = \sum_{i=1}^{k'} \alpha'_i b^{i-1} + \sum_i^{k'} \beta'_i b^i$ be the decomposition of the form (6) for $-z > 0$. We represent z as a string:

$$\begin{matrix} 1 & \alpha'_1 & \alpha'_2 & \dots & \alpha'_{k'} \\ 1 & \beta'_1 & \beta'_2 & \dots & \beta'_{k'} \end{matrix} \quad (8)$$

over the alphabet Σ_b . The first symbol $\begin{smallmatrix} 1 \\ 1 \end{smallmatrix}$ indicates that z is negative.

For $z \in \mathbb{Z}[1/b]$ we denote by $(z)_b \in \Sigma_b^*$ the string (7) if $z > 0$, the string (8) if $z < 0$ and the string $\begin{smallmatrix} 0 \\ 0 \end{smallmatrix}$, if $z = 0$. For example, if $z = -\frac{27}{8}$, then $(z)_2 = \begin{smallmatrix} 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 \end{smallmatrix}$.

The language $\mathcal{L}_b = \{(z)_b \mid z \in \mathbb{Z}[1/b]\}$ is regular. We denote by $\psi_b : \mathcal{L}_b \rightarrow \mathbb{Z}[1/b]$ the bijection which maps a string $(z)_b \in \mathcal{L}_b$ to $z \in \mathbb{Z}[1/b]$. For both cases, \mathcal{L}_b and ψ_b , the subscript indicates the base b . For $b = 2$, the representation ψ_2 , up to minor modification, coincides with the representation of $\mathbb{Z}[1/2]$ described in [17, § 2]. Let us denote by Add the graph of the addition operation in $\mathbb{Z}[1/b]$ with respect to ψ_b , namely, $Add = \{(u, v, w) \in \mathcal{L}_b \times \mathcal{L}_b \times \mathcal{L}_b \mid \psi_b(u) + \psi_b(v) = \psi_b(w)\}$. The relation Add is FA-recognizable [17, § 2]. We denote by add the addition operation in \mathcal{L}_b , that is, $add : \mathcal{L}_b \times \mathcal{L}_b \rightarrow \mathcal{L}_b$ is a two-place function for which $add(u, v) = w$ if $\psi_b(u) + \psi_b(v) = \psi_b(w)$.

For a given d -tuple $\bar{z} = (z_1, \dots, z_d) \in \mathbb{Z}[1/b]^d$ let us denote by $(\bar{z})_b$ the convolution $(z_1)_b \otimes \dots \otimes (z_d)_b$ of strings $(z_1)_b, \dots, (z_d)_b \in \mathcal{L}_b$. Clearly, the language $\mathcal{L}_b^d = \{w_1 \otimes \dots \otimes w_d \mid w_i \in \mathcal{L}_b, i = 1, \dots, d\}$ is regular. We denote by $\psi_b^d : \mathcal{L}_b^d \rightarrow \mathbb{Z}[1/b]^d$ the bijection which maps a string $(\bar{z})_b \in \mathcal{L}_b^d$ to $\bar{z} \in \mathbb{Z}[1/b]^d$. For both cases, \mathcal{L}_b^d and ψ_b^d , the superscript indicates the dimension d . Let us denote by Add_d the graph of the addition operation in $\mathbb{Z}[1/b]^d$ with respect to ψ_b^d , namely, $Add_d = \{(u, v, w) \in \mathcal{L}_b^d \times \mathcal{L}_b^d \times \mathcal{L}_b^d \mid \psi_b^d(u) + \psi_b^d(v) = \psi_b^d(w)\}$. The relation Add_d is FA-recognizable. We denote by add_d the addition operation in \mathcal{L}_b^d , that is, $add_d : \mathcal{L}_b^d \times \mathcal{L}_b^d \rightarrow \mathcal{L}_b^d$ is a two-place function for which $add_d(u, v) = w$ if $\psi_b^d(u) + \psi_b^d(v) = \psi_b^d(w)$. Clearly, if $d = 1$, then $\mathcal{L}_b^1 = \mathcal{L}_b$, $\psi_b^1 = \psi_b$, $Add_1 = Add$ and $add_1 = add$.

Let \mathcal{T} be a d -dimensional hierarchical mesh defined by a nested sequence of domains:

$$\Omega^0 = \mathbb{R}^d \supseteq \Omega^1 \supseteq \dots \supseteq \Omega^{N-1} \supseteq \Omega^N = \emptyset,$$

where $\Omega^{N-1} \neq \emptyset$ and each $\Omega^\ell, \ell = 1, \dots, N-1$ is composed of cells from $\mathcal{C}_d^{\ell-1}$.

For each d -dimensional cube $c = \prod_{j=1}^d [t_{i_j}^\ell, t_{i_j+1}^\ell]$ we associate it with its barycentre $\bar{z}_c = (z_1, \dots, z_d)$, where $z_j = \frac{1}{2}(t_{i_j}^\ell + t_{i_j+1}^\ell)$ for $j = 1, \dots, d$ (see Fig. 4).




Figure 4: The figure shows a 2-dimensional cell and its barycentre (a black dot in the centre of the cell).

For each $\ell = 1, \dots, N-1$ we denote by $L_\ell \subset \mathcal{L}_b^d$ the language:

$$L_\ell = \{(\bar{z}_c)_b \mid c \in \mathcal{C}_d^{\ell-1} \wedge c \subseteq \Omega^\ell\}.$$

Definition 5. We say that a hierarchical mesh \mathcal{T} is regular if the language L_ℓ is regular for each $\ell = 1, \dots, N-1$.

The languages L_ℓ , $\ell = 1, \dots, N-1$ are pairwise disjoint: $L_i \cap L_j = \emptyset$ for $i, j = 1, \dots, N-1$ and $i \neq j$. Let $L = L_1 \cup \dots \cup L_{N-1}$. The following proposition shows that the hierarchical mesh \mathcal{T} is regular if and only if the language L is regular.

Proposition 3. The language L is regular if and only if each language L_ℓ , $\ell = 1, \dots, N-1$ is regular.

Proof. If each language L_ℓ , $\ell = 1, \dots, N-1$ is regular, then L is regular as the union of regular languages. Now assume that L is regular. For the sake of simplicity, let us assume that $b = 2$. If $b > 2$, the proof is analogous up to minor modification. For a given $\ell = 1, \dots, N-1$, a string w is in the language L_ℓ if and only if $w \in L$ and w is the convolution of d strings of the form $u_i \otimes v_i \in \Sigma$, $i = 1, \dots, d$, for which v_i is the concatenation of three strings $v_i = r_i 1 s_i$: a string $r_i \in \{0, 1\}^*$ consisting of $\ell-1$ symbols, a string consisting of a single symbol 1 and a string $s_i \in \{0\}^*$. The latter condition can be easily verified by a finite automaton for all $i = 1, \dots, d$. Therefore, L_ℓ is regular. \square

We note that if Ω_1 , the first domain defining \mathcal{T} , is bounded, then \mathcal{T} is regular. If Ω_1 is not bounded, then, informally speaking, \mathcal{T} is regular if, probably outside of some bounded region, it looks like a regular pattern. See Figure 5 for illustration of regular hierarchical meshes.




Figure 5: The figures show portions of regular 3-level hierarchical meshes.

5.1 Verification of Nestedness

Let us be given regular languages L_ℓ , $\ell = 1, \dots, N-1$ representing domains $\Omega^1, \dots, \Omega^{N-1}$ composed of cells from $\mathcal{C}_d^0, \dots, \mathcal{C}_d^{N-2}$, respectively. How one can verify that the domains $\Omega^1, \dots, \Omega^{N-1}$ are nested: $\Omega^1 \supseteq \dots \supseteq \Omega^{N-1}$?

In order to verify nestedness one has to verify that for each $\ell = 2, \dots, N-1$: $\Omega^\ell \subseteq \Omega^{\ell-1}$. Let $c \in \mathcal{C}_d^{\ell-1}$ be a cell for which $c \subseteq \Omega^\ell$ for some ℓ , $2 \leq \ell \leq N-1$. Then $c \subseteq \Omega^{\ell-1}$ if and only if there exists a cell $c' \in \mathcal{C}_d^{\ell-2}$ for which $c \subseteq c'$ and $c' \subseteq \Omega^{\ell-1}$. The inclusion $c \subseteq c'$ holds if and only if there is a vector $\bar{s} = (\pm \frac{1}{2^\ell}, \dots, \pm \frac{1}{2^\ell}) \in \mathbb{Z}[1/b]^d$ for which $\bar{z}_c + \bar{s} = \bar{z}_{c'}$, that is, $((\bar{z}_c)_b, (\bar{s})_b, (\bar{z}_{c'})_b) \in \text{Add}_d$. There are exactly 2^d vectors of the form $(\pm \frac{1}{2^\ell}, \dots, \pm \frac{1}{2^\ell})$. We denote these vectors by $\bar{s}_1^\ell, \dots, \bar{s}_k^\ell$, where $k = 2^d$. Let $s_i^\ell = (\bar{s}_i^\ell)_b \in \mathcal{L}_b^d$ for $i = 1, \dots, k$.

Therefore, $c \subseteq c'$ if and only if for a first order formula:

$$\Phi_\ell = \text{add}_d(u, s_1^\ell) \in L_{\ell-1} \vee \dots \vee \text{add}_d(u, s_k^\ell) \in L_{\ell-1},$$

the evaluation of Φ_ℓ is true for $u = (\bar{z}_c)_b$ and the constants $s_1^\ell, \dots, s_k^\ell$. Therefore, $\Omega^\ell \subseteq \Omega^{\ell-1}$ if and only if the following first order sentence:

$$\Upsilon_\ell = \forall u (u \in L_\ell \rightarrow \Phi_\ell)$$

is true for the structure $(\mathcal{L}_b^d; \text{add}_d, L_\ell, L_{\ell-1}, s_1^\ell, \dots, s_k^\ell)$. Thus, the domains $\Omega^1, \dots, \Omega^{N-1}$ are nested if the first order sentence:

$$\Upsilon_2 \wedge \dots \wedge \Upsilon_{N-1}$$

is true for the structure $(\mathcal{L}_b^d; \text{add}_d, L_1, \dots, L_{N-1}, s_1^2, \dots, s_k^{N-1})$.

Let M_1, \dots, M_{N-1} be deterministic finite automata recognizing the languages L_1, \dots, L_{N-1} , respectively. We denote by m_1, \dots, m_{N-1} the number of states of the automata M_1, \dots, M_{N-1} , respectively. For given $1 \leq \ell \leq N-1$ and $1 \leq i \leq k$, using carrying which is a part of the standard addition algorithm in $\mathbb{Z}[1/b]$, from the automaton $M_{\ell-1}$ one can construct a deterministic finite automaton $M_{\ell-1,i}$ recognizing the unary relation $\text{add}_d(u, s_i^\ell) \in L_{\ell-1}$. It can be seen that the number of states of $M_{\ell-1,i}$ is $O(m_{\ell-1})$. Therefore, using the product construction, one can construct a deterministic finite automaton recognizing the unary relation Φ_ℓ for which the number of states is $O(m_{\ell-1}^k)$. Therefore, one can construct a deterministic finite automaton recognizing the unary relation $u \in L_\ell \rightarrow \Phi_\ell$ for which the number of states is $O(m_\ell \cdot m_{\ell-1}^k)$. Since $\forall = \neg \circ \exists \circ \neg$ and the emptiness problem for a deterministic finite automaton with n states can be solved in $O(n^2)$ time, there is an algorithm deciding whether or not Υ_ℓ is true in $O(m_\ell^2 \cdot m_{\ell-1}^{2k})$ time. Thus, there is a polynomial-time algorithm that for given deterministic finite automata M_1, \dots, M_{N-1} decides whether or not $\Upsilon_2 \wedge \dots \wedge \Upsilon_{N-1}$ is true.

5.2 Verification of Assumption 2

Now for given regular languages L_ℓ , $\ell = 1, \dots, N-1$ how one can verify the condition of Theorem 2 (that is, Assumption 2) which ensures that for the collection of tensor product B-splines \mathcal{K} , see (5), and every $f \in \mathcal{S}_m(\mathcal{T})$, $f = \sum_{\delta \in \mathcal{K}} \lambda_\delta \delta$ for some uniquely defined coefficients λ_δ ?

In order to verify Assumption 2 one has to verify that for each $\ell = 0, \dots, N-2$ the domain $\mathcal{M}^\ell = \mathbb{R}^d \setminus \Omega^{\ell+1}$ satisfies the following: for each $\beta \in B_{d,m}^\ell$, for which $\text{supp } \beta \cap \mathcal{M}^\ell \neq \emptyset$, the intersection $\overline{\text{supp } \beta} \cap \mathcal{M}^\ell$ is connected.

Each $\beta \in B_{d,m}^\ell$ we associate with one of the $(m+1)^d$ cells from \mathcal{C}_d^ℓ composing $\overline{\text{supp } \beta}$, depending on the parity of $m+1$: if $m+1$ is odd then we associate β with the central cell of $\overline{\text{supp } \beta}$, if $m+1$ is even then we associate β with the cell which has the central vertex of $\overline{\text{supp } \beta}$ as its lower left corner³; for explanation see Fig. 6. For a given $\beta \in B_{d,m}^\ell$ we denote by $c_\beta \in \mathcal{C}_d^\ell$ the associated cell.

For a given $(i_1, \dots, i_d) \in \mathbb{Z}^d$ let $\bar{t}_{i_1 \dots i_d}^\ell$ be the vector $\bar{t}_{i_1 \dots i_d}^\ell = (\frac{i_1}{2^\ell}, \dots, \frac{i_d}{2^\ell}) \in \mathbb{Z}[1/b]^d$. Let $t_{i_1 \dots i_d}^\ell = (\bar{t}_{i_1 \dots i_d}^\ell)_b \in \mathcal{L}_b^d$. For a given $m \geq 0$ we denote I_m the set $I_m = \{(i_1, \dots, i_d) \in \mathbb{Z}^d \mid -\frac{m}{2} \leq i_k \leq \frac{m}{2}, k = 1, \dots, d\}$ if $m+1$ is odd and $I_m = \{(i_1, \dots, i_d) \in \mathbb{Z}^d \mid -\frac{m+1}{2} \leq i_k \leq \frac{m-1}{2}, k = 1, \dots, d\}$ if $m+1$ is even.

Let $\bar{i} = (i_1, \dots, i_d) \in I_m$. We denote by $\Phi_{\ell, m, \bar{i}}$ the following first order formula:

$$\Phi_{\ell, m, \bar{i}} = \text{add}_d(u, t_{i_1 \dots i_d}^\ell) \in L_{\ell+1}.$$

The condition that for a given $\beta \in B_{d,m}^\ell$ the intersection $\text{supp } \beta \cap \mathcal{M}^\ell \neq \emptyset$ holds if and only if the evaluation of the following formula:

$$\Psi_\ell = \bigvee_{\bar{i} \in I_m} \neg \Phi_{\ell, m, \bar{i}} \quad (9)$$

³We use the term lower left corner in the context of the case $d = 2$. If $d \neq 2$, we use the term lower left corner of a d -dimensional cell $[0, 1]^d$ for the vertex $(0, \dots, 0) \in \mathbb{R}^d$.




Figure 6: The figure on the left shows the support of $\beta \in B_{2,4}^\ell$ with the associated cell c_β shaded in gray. The figure on the right shows the support of $\beta \in B_{2,3}^\ell$ with the associated cell c_β shaded in gray; this cell has the central vertex of $\text{supp } \beta$ (shown as a black dot) as its lower left corner.

is true for $u = (\bar{z}_{c_\beta})_b$ and the constants $t_{i_1 \dots i_d}^\ell$. Moreover, the condition that the intersection $\overline{\text{supp } \beta} \cap \mathcal{M}^\ell$ is connected can be encoded by a first order formula as follows. Every possible nonempty intersection $\overline{\text{supp } \beta} \cap \mathcal{M}^\ell$ corresponds to a nonempty subset $J \subseteq I_m$ (see Fig. 7 for illustration) for which the evaluation of the following first order formula:

$$\Psi_{\ell,J} = \left(\bigwedge_{\bar{j} \in J} \neg \Phi_{\ell,m,\bar{j}} \right) \wedge \left(\bigwedge_{\bar{j} \in I_m \setminus J} \Phi_{\ell,m,\bar{j}} \right)$$

is true for $u = (\bar{z}_{c_\beta})_b$, the constants $t_{i_1 \dots i_d}^\ell$ and the domain \mathcal{L}_b^d .

We denote by \mathcal{J}_m the collection of all nonempty subsets $J \subseteq I_m$ that correspond to connected intersections. For example, in Fig. 7 the intersection on the left corresponding to the set $J = \{(-2, -1), (-1, -1), (0, -1), (1, -1), (2, -1), (-1, 0), (1, 0), (0, 1)\}$ is connected, so $J \in \mathcal{J}_4$; the intersection on the right corresponding to the set $J' = \{(-2, -1), (-1, -1), (0, -1), (1, -1), (2, -1), (-1, 0), (0, 0), (1, 0), (0, 1), (0, -2), (2, 2)\}$ is not connected, so $J' \notin \mathcal{J}_4$.




Figure 7: The figure on the left shows the support of some tensor product B-spline from $B_{2,4}^\ell$ and its intersection with \mathcal{M}^ℓ shaded in gray which is connected. The figure on the right shows the support of some tensor product B-spline from $B_{2,4}^\ell$ and its intersection with \mathcal{M}^ℓ shaded in gray which is not connected.

For given $d > 0$ and $\ell \geq 0$, we denote by $\tilde{\mathcal{L}}_\ell^d \subset \mathcal{L}_b^d$ the language $\tilde{\mathcal{L}}_\ell^d = \{(\bar{z}_c)_b \mid c \in \mathcal{C}_d^\ell\}$. For example, if $b = 2$, the language $\tilde{\mathcal{L}}_\ell^d$ consists of all convolutions of d strings of the form $u_i \otimes v_i \in \mathcal{L}_2$, $i = 1, \dots, d$ for which $v_i = r_i 1 s_i$, where $r_i \in \{0, 1\}^*$, $|r_i| = \ell$ and $s_i \in \{0\}^*$. A language $\tilde{\mathcal{L}}_\ell^d \subset \mathcal{L}_b^d$ is regular, see also Proposition 3. Finally, the condition that for every $\beta \in B_{d,m}^\ell$ such that $\text{supp } \beta \cap \mathcal{M}^\ell \neq \emptyset$ the intersection $\text{supp } \beta \cap \mathcal{M}^\ell$ is

connected holds if and only if the following first order formula:

$$\mathcal{X}_\ell = \forall u((u \in \tilde{L}_\ell^d \wedge \Psi_\ell) \rightarrow \bigvee_{J \in \mathcal{J}_m} \Psi_{\ell,J})$$

is true for the structure $(\mathcal{L}_b^d; \text{add}_d, \tilde{L}_\ell^d, L_{\ell+1}, \{t_{\tilde{i}}^\ell \mid \tilde{i} \in I_m\})$. Therefore, Assumption 2 holds for the domains $\mathcal{M}^0, \dots, \mathcal{M}^{N-2}$ if the first order sentence:

$$\mathcal{X}_0 \wedge \dots \wedge \mathcal{X}_{N-2}$$

is true for the structure:

$$(\mathcal{L}_b^d; \text{add}_d, \tilde{L}_0^d, \dots, \tilde{L}_{N-2}^d, L_1, \dots, L_{N-1}, \{t_{\tilde{i}}^\ell \mid \tilde{i} \in I_m, \ell = 0, \dots, N-2\}).$$

Similarly to Subsection 5.1, there is a polynomial-time algorithm that for given deterministic finite automata M_1, \dots, M_{N-1} decides whether or not $\mathcal{X}_0 \wedge \dots \wedge \mathcal{X}_{N-2}$ is true.

5.3 Regularity for \mathcal{K}

At each level $\ell = 0, \dots, N-1$ the collection of tensor product B-splines functions \mathcal{K}^ℓ generated by Kraft's selection mechanism, see (5), corresponds to the collection of cells $\mathcal{K}_c^\ell = \{c_\beta \mid \beta \in \mathcal{K}^\ell\} \subseteq C_d^\ell$ according to the rule for associating a tensor product B-spline β with the corresponding cell c_β , described in Subsection 5.2. We denote by \hat{L}_ℓ the language $\hat{L}_\ell = \{(\bar{z}_c)_b \mid c \in \mathcal{K}_c^\ell\} \subseteq \tilde{L}_\ell^d$. Below we will show that the languages $\hat{L}_0, \dots, \hat{L}_{N-1}$ are regular.

First we note that the language \hat{L}_0 is defined by the formula:

$$\Theta_0 = u \in \tilde{L}_0^d \wedge \Psi_0,$$

where Ψ_0 is given by (9). That is, \hat{L}_0 is the language of strings u from \mathcal{L}_b^d for which the evaluation of the formula Θ_0 is true. The formula Ψ_0 verifies whether the intersection of $\text{supp } \beta$ for $\beta \in B_{d,m}^0$ with \mathcal{M}^0 is nonempty. If it is nonempty, then $\beta \in \mathcal{K}^0$.

For given $\ell > 0$, $(i_1, \dots, i_d) \in I_m$ and $j = 1, \dots, k$, where $k = 2^d$, we denote by $\bar{r}_{i_1 \dots i_{dj}}^\ell$ the constant vectors $\bar{r}_{i_1 \dots i_{dj}}^\ell = \bar{t}_{i_1 \dots i_d}^\ell + \bar{s}_j^{\ell+1}$. Let $r_{i_1 \dots i_{dj}}^\ell = (\bar{r}_{i_1 \dots i_{dj}}^\ell)_b \in \mathcal{L}_b^d$. For a given $\ell > 0$, let:

$$\Theta_\ell = u \in \tilde{L}_\ell^d \wedge \Psi_\ell \wedge \bigwedge_{\tilde{i} \in I_m} \bigvee_{j=1}^k \text{add}_d(u, r_{i_1 \dots i_{dj}}^\ell) \in L_\ell.$$

The formula Θ_ℓ defines the language \hat{L}_ℓ for $\ell = 1, \dots, N-2$. For $\beta \in B_{d,m}^\ell$ the formula Ψ_ℓ verifies whether the intersection of $\text{supp } \beta$ with \mathcal{M}^ℓ is nonempty. The formula $\bigwedge_{\tilde{i} \in I_m} \bigvee_{j=1}^k \text{add}_d(u, r_{i_1 \dots i_{dj}}^\ell) \in L_\ell$ verifies whether $\text{supp } \beta \subseteq \Omega^\ell$. If for $\beta \in B_{d,m}^\ell$, $\text{supp } \beta \cap \mathcal{M}^\ell \neq \emptyset$ and $\text{supp } \beta \subseteq \Omega^\ell$, then $\beta \in \mathcal{K}^\ell$. For a given $\ell > 0$, let:

$$\Gamma_\ell = u \in \tilde{L}_\ell^d \wedge \bigwedge_{\tilde{i} \in I_m} \bigvee_{j=1}^k \text{add}_d(u, r_{i_1 \dots i_{dj}}^\ell) \in L_\ell. \quad (10)$$

Clearly, the formula Γ_{N-1} defines the language \hat{L}_{N-1} . For $\beta \in B_{d,m}^{N-1}$ the formula $\bigwedge_{\tilde{i} \in I_m} \bigvee_{j=1}^k \text{add}_d(u, r_{i_1 \dots i_{dj}}^{N-1}) \in L_{N-1}$ verifies whether $\text{supp } \beta \subseteq \Omega^{N-1}$. If for $\beta \in B_{d,m}^{N-1}$, $\text{supp } \beta \subseteq \Omega^{N-1}$, then $\beta \in \mathcal{K}^{N-1}$.

Since the languages $\widehat{L}_0, \dots, \widehat{L}_{N-1}$ are defined by the first order formulae $\Theta_0, \Theta_1, \dots, \Theta_{N-2}, \Gamma_{N-1}$, they must be regular for regular hierarchical meshes. Moreover, similarly to the argument in Subsection 5.1, for given deterministic finite automata M_1, \dots, M_{N-1} one can construct deterministic finite automata $\widehat{M}_0, \dots, \widehat{M}_{N-1}$ recognizing the languages $\widehat{L}_0, \dots, \widehat{L}_{N-1}$, respectively. Furthermore, these automata $\widehat{M}_0, \dots, \widehat{M}_{N-1}$ are constructed from the automata M_1, \dots, M_{N-1} in polynomial time.

6 Regular Splines

Let \mathcal{T} be a regular hierarchical d -dimensional mesh defined by a nested sequence of domains $\Omega^0 = \mathbb{R}^d \supseteq \Omega^1 \supseteq \dots \supseteq \Omega^{N-1} \supseteq \Omega^N = \emptyset$, where $\Omega^{N-1} \neq \emptyset$. Let $f = \sum_{\beta \in \mathcal{K}} \lambda_\beta \beta$

be a spline in $\mathcal{S}_m(\mathcal{T})$ defined by some coefficients λ_β , $\beta \in \mathcal{K}$, where $\mathcal{K} = \bigcup_{\ell=0}^{N-1} \mathcal{K}^\ell$ is

obtained by Kraft's selection mechanism, see the equation (5) and Remark 3. Recall that by Theorem 2, if \mathcal{T} satisfies Assumption 2, then each spline in $\mathcal{S}_m(\mathcal{T})$ can be written as the infinite sum $\sum_{\beta \in \mathcal{K}} \lambda_\beta \beta$. Each $\beta \in \mathcal{K}^\ell$ is associated with the cell $c_\beta \in \mathcal{K}_c^\ell \subseteq \mathcal{C}_d^\ell$

which is then associated with $(\bar{z}_{c_\beta})_b \in \widehat{L}_\ell$, see the notation in Subsection 5.3.

Definition 6. We say that a spline $f \in \mathcal{S}_m(\mathcal{T})$ is regular if the coefficients $\lambda_\beta \in \mathbb{Z}[1/b]$ for all $\beta \in \mathcal{K}$ and the relation $S_f = \{((\bar{z}_{c_\beta})_b, (\lambda_\beta)_b) \mid \beta \in \mathcal{K}\} \subset \mathcal{L}_b^d \times \mathcal{L}_b$ is FA-recognizable.

For a given $\ell = 0, \dots, N-1$, we denote by S_f^ℓ the relation:

$$S_f^\ell = \{((\bar{z}_{c_\beta})_b, (\lambda_\beta)_b) \mid \beta \in \mathcal{K}^\ell\}.$$

Similarly to Proposition 3, a spline $f \in \mathcal{S}_m(\mathcal{T})$ is regular if and only if each of the relation S_f^ℓ is FA-recognizable for $\ell = 0, \dots, N-1$.

Since the relation Add_d is FA-recognizable, for given regular splines $f_1, f_2 \in \mathcal{S}_m(\mathcal{T})$ the sum $f_1 + f_2 \in \mathcal{S}_m(\mathcal{T})$ is a regular spline. Moreover, for a constant $\mu \in \mathbb{Z}[1/b]$ the relation:

$$R_\mu = \{((\lambda)_b, (\mu\lambda)_b) \in \mathcal{L}_b \times \mathcal{L}_b \mid \lambda \in \mathbb{Z}[1/b]\} \quad (11)$$

is FA-recognizable. Therefore, for a regular spline $f \in \mathcal{S}_m(\mathcal{T})$, the spline $\mu f \in \mathcal{S}_m(\mathcal{T})$ is regular. Thus, the collection of all regular splines in $\mathcal{S}_m(\mathcal{T})$ is a module over the ring $\mathbb{Z}[1/b]$.

Remark 4. For given $f_1, f_2 \in \mathcal{S}_m(\mathcal{T})$ and $\ell = 0, \dots, N-1$, let $M_{1,\ell}$ and $M_{2,\ell}$ be deterministic finite automata recognizing the relations $S_{f_1}^\ell$ and $S_{f_2}^\ell$, respectively. We denote by $m_{1,\ell}$ and $m_{2,\ell}$ the number of states of $M_{1,\ell}$ and $M_{2,\ell}$, respectively. Let $f = f_1 + f_2$. The relation $S_f^\ell \subset \mathcal{L}_b^d \times \mathcal{L}_b$ is defined by the formula:

$$\Xi_\ell = \exists v_1 \exists v_2 (S_{f_1}^\ell(u, v_1) \wedge S_{f_2}^\ell(u, v_2) \wedge Add(v_1, v_2, v)).$$

That is, the evaluation of Ξ_ℓ is true for $u \in \mathcal{L}_b^d$ and $v \in \mathcal{L}_b$ if and only if $(u, v) \in S_f^\ell$. By Theorem 3, from the automata $M_{1,\ell}, M_{2,\ell}$ and an automaton recognizing Add one can construct a deterministic finite automaton recognizing the relation S_f^ℓ . However, the existential quantifiers in Ξ_ℓ require us to use the Rabin-Scott powerset construction which may lead to exponential growth in the number of states. In order to avoid this we propose to present $f = f_1 + f_2$ by the relation:

$$\mathbf{S}_f^\ell = \{(u, v_1, v_2, v) \in \mathcal{L}_b^d \times \mathcal{L}_b \times \mathcal{L}_b \times \mathcal{L}_b \mid S_{f_1}^\ell(u, v_1) \wedge S_{f_2}^\ell(u, v_2) \wedge Add(v_1, v_2, v)\}.$$

One then can construct a deterministic finite automaton recognizing \mathbf{S}_f^ℓ for which the number of states is $O(m_{1,\ell} \cdot m_{2,\ell})$. The same approach works for multiplication by a constant μ . For example, if f is then multiplied by μ , we can present μf by the relation:

$$\mathbf{S}_{\mu f}^\ell = \{(u, v_1, v_2, v, w) \in \mathcal{L}_b^d \times \mathcal{L}_b \times \mathcal{L}_b \times \mathcal{L}_b \times \mathcal{L}_b \mid \mathbf{S}_f^\ell(u, v_1, v_2, v) \wedge R_\mu(v, w)\}.$$

6.1 Computing Values of a Regular Spline

Let $f \in \mathcal{S}_m(\mathcal{T})$ be a regular spline given by a FA-recognizable relation S_f . For each $\ell = 0, \dots, N-1$, let us be given a deterministic finite automaton M_ℓ recognizing the relation S_f^ℓ . For a given point $\bar{x} = (x_1, \dots, x_d) \in \mathbb{Z}[1/b]^d$, how one can compute the value $f(\bar{x})$?

Let $R_f^\ell \subset \mathcal{L}_b^d \times \mathcal{L}_b^d \times \mathcal{L}_b$ be the relation that contains all triples $((\bar{x})_b, (\bar{z}_{c_\beta})_b, (\lambda_\beta)_b)$ of strings $(\bar{x})_b \in \mathcal{L}_b^d$, $(\bar{z}_{c_\beta})_b \in \mathcal{L}_b^d$ and $(\lambda_\beta)_b \in \mathcal{L}_b$ for $\beta \in \mathcal{K}^\ell$ such that $\bar{x} \in \text{supp } \beta$:

$$R_f^\ell = \{((\bar{x})_b, (\bar{z}_{c_\beta})_b, (\lambda_\beta)_b) \mid \bar{x} \in \mathbb{Z}[1/b]^d, \beta \in \mathcal{K}^\ell \wedge \bar{x} \in \text{supp } \beta\}.$$

Let $\bar{y} = (y_1, \dots, y_d) = \bar{z}_{c_\beta}$. The condition $\bar{x} \in \text{supp } \beta$ for $\beta \in \mathcal{K}^\ell$ is true if and only if the inequalities:

$$-\frac{m+2}{2^{\ell+1}} < x_i - y_i < \frac{m}{2^{\ell+1}}, \quad -\frac{m+1}{2^{\ell+1}} < x_i - y_i < \frac{m+1}{2^{\ell+1}} \quad (12)$$

hold for all $i = 1, \dots, d$, if m is odd and even, respectively, see Fig. 8.




Figure 8: The figure on the left shows the support of a spline $\beta \in B_{2,3}^\ell$, the points $\bar{x} \in \text{supp } \beta$, $\bar{y} = \bar{z}_{c_\beta}$ and the lower left corner of $\overline{\text{supp } \beta}$ – the point \bar{q} . The figure on the right shows the support of a spline $\beta \in B_{2,4}^\ell$, the points $\bar{x} \in \text{supp } \beta$, $\bar{y} = \bar{z}_{c_\beta}$ and the lower left corner \bar{q} .

For $\bar{r} = (r_1, \dots, r_d) \in \mathbb{Z}[1/b]^d$ and $\bar{s} = (s_1, \dots, s_d) \in \mathbb{Z}[1/b]^d$ we say that $\bar{r} < \bar{s}$ if $r_i < s_i$ for all $i = 1, \dots, d$. Let $R_{<}^d$ be the relation $R_{<}^d = \{((\bar{r})_b, (\bar{s})_b) \mid \bar{r}, \bar{s} \in \mathbb{Z}[1/b]^d, \bar{r} < \bar{s}\}$. The relation $R_{<}^d$ is FA-recognizable. Since Add_d and $R_{<}^d$ are FA-recognizable, the relation given by the inequalities (12) is FA-recognizable. Therefore, since S_f^ℓ is FA-recognizable, R_f^ℓ is FA-recognizable.

We denote by \bar{q}_β the lower left corner $\bar{q} = (q_1, \dots, q_d)$ of $\overline{\text{supp } \beta}$, see Fig. 8. We have that $y_i - q_i = \frac{m+2}{2^{\ell+1}}$ and $y_i - q_i = \frac{m+1}{2^{\ell+1}}$ for all $i = 1, \dots, d$, if m is odd and even, respectively. Since Add_d is FA-recognizable, the relation $Q_d^\ell = \{(\bar{z}_{c_\beta}, \bar{q}_\beta) \mid \beta \in B_{d,m}^\ell\}$ is FA-recognizable.

Now let $\tilde{R}_f^\ell \subset \mathcal{L}_b^d \times \mathcal{L}_b^d \times \mathcal{L}_b \times \mathcal{L}_b^d$ be the following relation:

$$\tilde{R}_f^\ell = \{((\bar{x})_b, (\bar{z}_{c_\beta})_b, (\lambda_\beta)_b, (\bar{x} - \bar{q}_\beta)_b) \mid \bar{x} \in \mathbb{Z}[1/b]^d, \beta \in \mathcal{K}^\ell \wedge \bar{x} \in \text{supp } \beta\}.$$

Since the relations R_f^ℓ , Q_d^ℓ and Add_d are FA-recognizable, the relation \tilde{R}_f^ℓ is FA-recognizable. From automata recognizing the relations $R_{<}^d$, Q_d^ℓ , Add_d and the automaton M_ℓ one can construct a deterministic finite automaton recognizing the relation \tilde{R}_f^ℓ for which the number of states is $O(m_\ell)$, where m_ℓ is the number of states of M_ℓ .

Note that for a given $\bar{x} \in \mathbb{Z}[1/b]^d$, there exist at most $(m+1)^d$ tensor product B-splines $\beta \in \mathcal{K}^\ell$ for which $\bar{x} \in \text{supp } \beta$. So \tilde{R}_f^ℓ can be seen as a multivalued function that for a given input \bar{x} returns at most $(m+1)^d$ pairs $((\lambda_\beta)_b, (\bar{x} - \bar{q}_\beta)_b)$ as an output.

Since \widetilde{R}_f^ℓ is FA-recognizable, this multivalued function is computed in linear time by a deterministic one-tape Turing machine, see Section 3.

We denote by $\mathcal{K}_{\bar{x}}^\ell$ the set $\mathcal{K}_{\bar{x}}^\ell = \{\beta \in \mathcal{K}^\ell \mid \bar{x} \in \text{supp } \beta\}$ and by $\mathcal{K}_{\bar{x}}$ the set $\mathcal{K}_{\bar{x}} = \bigcup_{\ell=0}^{N-1} \mathcal{K}_{\bar{x}}^\ell$. After all pairs $((\lambda_\beta)_b, (\bar{x} - \bar{q}_\beta)_b)$ for which $\bar{x} \in \text{supp } \beta$, where $\beta \in \mathcal{K}$, are computed, the value of the spline $f(\bar{x}) = \sum_{\beta \in \mathcal{K}_{\bar{x}}} \lambda_\beta \beta(\bar{x})$ at the point is obtained from the formula for $N_{0,m}^\ell(t)$ by evaluating $N_{0,m}^\ell(\bar{x} - \bar{q}_\beta)$ for every $\beta \in \mathcal{K}_{\bar{x}}^\ell$. Note that $\mathcal{K}_{\bar{x}}$ is a finite set containing at most $N(m+1)^d$ elements, so there are at most $N(m+1)^d$ terms in the sum $\sum_{\beta \in \mathcal{K}_{\bar{x}}} \lambda_\beta \beta(\bar{x})$.

For illustration below we provide concrete formulae for evaluating $N_{0,m}^0(t)$ for $m = 1, 2, 3$. If $\ell > 0$, $N_{0,m}^\ell(t) = N_{0,m}^0(2^\ell t)$ for $t \in (0, \frac{m+1}{2^\ell})$. By (2)–(3) one can obtain that (see, e.g., [21]):

$$N_{0,1}^0(t) = \begin{cases} t, & 0 \leq t < 1, \\ 2 - t, & 1 \leq t < 2, \\ 0, & \text{otherwise,} \end{cases} \quad (13)$$

$$N_{0,2}^0(t) = \begin{cases} \frac{1}{2}t^2, & 0 \leq t < 1, \\ -(t-1)^2 + (t-1) + \frac{1}{2}, & 1 \leq t < 2, \\ \frac{1}{2}(3-t)^2, & 2 \leq t < 3, \\ 0, & \text{otherwise,} \end{cases} \quad (14)$$

$$N_{0,3}^0(t) = \begin{cases} \frac{1}{6}t^3, & 0 \leq t < 1, \\ \frac{1}{6}(-3(t-1)^3 + 3(t-1)^2 + 3(t-1) + 1), & 1 \leq t < 2, \\ \frac{1}{6}(3(t-2)^3 - 6(t-2)^2 + 4), & 2 \leq t < 3, \\ \frac{1}{6}(-(t-3)^2 + 3(t-3)^2 - 3(t-3) + 1), & 3 \leq t < 4, \\ 0, & \text{otherwise.} \end{cases} \quad (15)$$

It follows from the formulae (13) and (14) that for $m = 1, 2$ and b divisible by 2, if $\bar{x} \in \mathbb{Z}[1/b]$, then $f(\bar{x}) \in \mathbb{Z}[1/b]$. However, in order to guarantee the same for $m = 3$, it is required that b is divisible by 6. That is, it follows from (15) that for $m = 3$ and b divisible by 6, if $\bar{x} \in \mathbb{Z}[1/b]$, then $f(\bar{x}) \in \mathbb{Z}[1/b]$.

If one applies the standard long multiplication algorithm to evaluate $N_{0,m}^\ell(\bar{x} - \bar{q}_\beta)$ and then multiply it by λ_β for each $\beta \in \mathcal{K}_{\bar{x}}$, the total computational complexity for evaluating $f(\bar{x})$ for a given input \bar{x} is quadratic; though it can be reduced if one applies a faster multiplication algorithm.

6.2 Refining Regular Hierarchical Meshes and Splines

Now let us refine a regular hierarchical mesh \mathcal{T} by selecting a nonempty subdomain $\Omega^N \subseteq \Omega^{N-1}$ composed of cells from \mathcal{C}_d^{N-1} . We assume that the language L_N corresponding to the subdomain Ω^N , according to the rule described in Section 5, is regular. So a hierarchical d -dimensional mesh \mathcal{T}' defined by a nested sequence of domains $\Omega^0 = \mathbb{R}^d \supseteq \Omega^1 \supseteq \dots \supseteq \Omega^{N-1} \supseteq \Omega^N \supseteq \Omega^{N+1} = \emptyset$ is regular.

Let $\mathcal{K}' = \bigcup_{\ell=0}^N \mathcal{K}'^\ell$ be the collection of tensor product B-splines generated by Kraft's selection mechanism for the hierarchical mesh \mathcal{T}' . If the languages $\widehat{L}_0, \dots, \widehat{L}_{N-1}$, corresponding to the collections $\mathcal{K}^0, \dots, \mathcal{K}^{N-1}$, respectively, are given as an input, how one does obtain the languages $\widehat{L}'_0, \dots, \widehat{L}'_N$ corresponding to the collections $\mathcal{K}'^0, \dots, \mathcal{K}'^N$, respectively?

Since $\mathcal{K}^0 = \mathcal{K}'^0, \dots, \mathcal{K}^{N-2} = \mathcal{K}'^{N-2}$, we have that $\widehat{L}'_0 = \widehat{L}_0, \dots, \widehat{L}'_{N-2} = \widehat{L}_{N-2}$. A tensor product B-spline $\beta \in \mathcal{K}'^{N-1}$ if and only if $\beta \in \mathcal{K}^{N-1}$ and $\text{supp } \beta \cap \mathcal{M}^{N-1} \neq \emptyset$, where $\mathcal{M}^{N-1} = \mathbb{R}^d \setminus \Omega^N$. The condition $\text{supp } \beta \cap \mathcal{M}^{N-1} \neq \emptyset$ is verified by the formula

Ψ_{N-1} , see (9). So the language \widehat{L}_{N-1} is defined by the formula $u \in \widehat{L}_{N-1} \wedge \Psi_{N-1}$. The formula Γ_N , see (10), verifies the condition that a tensor product B-spline $\beta \in \mathcal{K}'^N$. So the language \widehat{L}'_N is defined by the formula Γ_N . Thus, all languages $\widehat{L}'_0, \dots, \widehat{L}'_N$ are regular. Let \widehat{M}_{N-1} and M_N be deterministic finite automata recognizing the languages \widehat{L}_{N-1} and L_N , respectively. One can construct deterministic finite automata recognizing \widehat{L}'_{N-1} and \widehat{L}'_N for which the number of states is $O(\widehat{m}_{N-1} \cdot m_N)$ and $O(m_N)$, respectively, where \widehat{m}_{N-1} and m_N are the number of states of \widehat{M}_{N-1} and M_N , respectively.

Let $f \in \mathcal{S}_m(\mathcal{T})$ be regular spline given by a FA-recognizable relation S_f . How one does obtain a relation S'_f for the spline function f over the hierarchical mesh \mathcal{T}' ?

$$\text{We have that } f = \sum_{\ell=0}^{N-1} \sum_{\beta \in \mathcal{K}^\ell} \lambda_\beta \beta = \sum_{\ell=0}^{N-2} \sum_{\beta \in \mathcal{K}^\ell} \lambda_\beta \beta + \sum_{\beta \in \mathcal{K}'^{N-1}} \lambda_\beta \beta + \sum_{\beta \in \mathcal{K}'^N} \lambda_\beta \beta.$$

Therefore, $S_f^0 = S_f^0, \dots, S_f^{N-2} = S_f^{N-2}$ and $S_f^{N-1} = \{(\bar{u}, v) \in S_f^{N-1} \mid \bar{u} \in \widehat{L}'_{N-1}\}$. Clearly, S_f^0, \dots, S_f^{N-1} are FA-recognizable. Let $M_{f,N-1}$ be a deterministic finite automaton recognizing S_f^{N-1} . One can construct a deterministic finite automaton recognizing S_f^{N-1} for which the number of states is $O(m_{f,N-1} \cdot m_N)$, where $m_{f,N-1}$ is the number of states of $M_{f,N-1}$. Below we will show that the coefficients $\lambda_\beta \in \mathbb{Z}[1/b]$ for all $\beta \in \mathcal{K}'^N$ and the relation $S_f^N = \{((\bar{z}_{c_\beta})_b, (\lambda_\beta)_b) \mid \beta \in \mathcal{K}'^N\}$ is FA-recognizable.

For any given $\delta \in B_{d,m}^{\ell-1}$ each $\beta \in B_{d,m}^\ell$ for which $\text{supp } \beta \subset \text{supp } \delta$ corresponds to a multi-index $\bar{j}_{\delta,\beta} = (j_1, \dots, j_d)$, where $0 \leq j_k \leq m+1$ for all $k = 1, \dots, d$, that determines the position of $\text{supp } \beta$ inside $\text{supp } \delta$, see Fig. 9 for explanation. For given $\delta \in B_{d,m}^{\ell-1}$ and a multi-index $\bar{j} = (j_1, \dots, j_d)$ we denote by $\beta_{\delta,\bar{j}}$ the tensor product B-spline $\beta \in B_{d,m}^\ell$ for which $\bar{j}_{\delta,\beta} = \bar{j}$. Note that for the barycentres \bar{z}_{c_δ} and \bar{z}_{c_β} the following holds:

$$\bar{z}_{c_\delta} - \bar{z}_{c_\beta} = \begin{cases} \left(\frac{m+1}{2^{\ell+1}} - \frac{j_1}{2^\ell}, \dots, \frac{m+1}{2^{\ell+1}} - \frac{j_d}{2^\ell} \right), & \text{if } m \text{ is even,} \\ \left(\frac{m+2}{2^{\ell+1}} - \frac{j_1}{2^\ell}, \dots, \frac{m+2}{2^{\ell+1}} - \frac{j_d}{2^\ell} \right), & \text{if } m \text{ is odd.} \end{cases} \quad (16)$$




Figure 9: The left figure shows the support of a spline $\delta_1 \in B_{2,4}^{\ell-1}$, the supports of $\beta_1, \gamma_1 \in B_{2,4}^\ell$ (two hatched rectangles) and the points $\bar{p}_1 = \bar{z}_{c_{\beta_1}}$, $\bar{r}_1 = \bar{z}_{c_{\gamma_1}}$ and $\bar{q}_1 = \bar{z}_{c_{\delta_1}}$. The indices $\bar{j}_{\delta_1,\beta_1}, \bar{j}_{\delta_1,\gamma_1}$ are $\bar{j}_{\delta_1,\beta_1} = (0, 1)$ and $\bar{j}_{\delta_1,\gamma_1} = (4, 3)$. The right figure shows the support of a spline $\delta_2 \in B_{2,3}^{\ell-1}$, the supports of $\beta_2, \gamma_2 \in B_{2,3}^\ell$ (two hatched rectangles) and the points $\bar{p}_2 = \bar{z}_{c_{\beta_2}}$, $\bar{r}_2 = \bar{z}_{c_{\gamma_2}}$ and $\bar{q}_2 = \bar{z}_{c_{\delta_2}}$. The indices $\bar{j}_{\delta_2,\beta_2}, \bar{j}_{\delta_2,\gamma_2}$ are $\bar{j}_{\delta_2,\beta_2} = (0, 1)$ and $\bar{j}_{\delta_2,\gamma_2} = (4, 3)$.

We denote by $\mathcal{I}_{d,m}$ the set of multi-indices $\mathcal{I}_{d,m} = \{(j_1, \dots, j_d) \mid 0 \leq j_k \leq m+1, k = 1, \dots, d\}$. For each $\delta \in B_{d,m}^{\ell-1}$, we have that $\delta = \sum_{\bar{j} \in \mathcal{I}_{d,m}} \lambda_{\bar{j}} \beta_{\delta,\bar{j}}$. It can be verified directly from Boehm's knot insertion formula for B-splines [2] that for $d = 1$: $\lambda_j = \frac{1}{2^m} \binom{j}{m+1}$,

$j = 0, \dots, m+1$, where $\binom{j}{m+1} = \frac{(m+1)!}{j!(m+1-j)!}$ are the binomial coefficients. So all coefficients λ_j , $j = 0, \dots, m+1$ belong to $\mathbb{Z}[1/2]$. By the definition of multivariate tensor product B-splines, see (1), for $d > 1$, we immediately obtain that $\lambda_{\vec{j}} \in \mathbb{Z}[1/2]$ for all $\vec{j} \in \mathcal{I}_{d,m}$ as well.

Let $\bar{\mathcal{K}}^{N-1} = \mathcal{K}^{N-1} \setminus \mathcal{K}'^{N-1}$. We have that $\sum_{\delta \in \bar{\mathcal{K}}^{N-1}} \lambda_\delta \delta = \sum_{\beta \in \mathcal{K}'^N} \lambda_\beta \beta$. For a given $\beta \in \mathcal{K}'^N$, let $\Delta_\beta = \{(\delta, \vec{j}) \mid \delta \in \bar{\mathcal{K}}^{N-1}, \vec{j} \in \mathcal{I}_{d,m}, \text{supp } \beta \subset \text{supp } \delta \wedge \beta = \beta_{\delta, \vec{j}}\}$. Then, for any $\beta \in \mathcal{K}'^N$, $\lambda_\beta = \sum_{(\delta, \vec{j}) \in \Delta_\beta} \lambda_\delta \lambda_{\vec{j}}$. Since $\lambda_{\vec{j}} \in \mathbb{Z}[1/2]$ for all $\vec{j} \in \mathcal{I}_{d,m}$ and $\lambda_\delta \in \mathbb{Z}[1/b]$ for all $\delta \in \mathcal{K}$, then $\lambda_\beta \in \mathbb{Z}[1/b]$ for all $\beta \in \mathcal{K}'^N$.

For a given $\vec{j} \in \mathcal{I}_{d,m}$, let $q_{\vec{j}} \in \mathbb{Z}[1/2]^d$ be a constant vector given by the right-hand side of the equation (16). For a given $\ell \geq 1$ and $\vec{j} \in \mathcal{I}_{d,m}$, let:

$$R_{\ell, \vec{j}} = \{((\bar{z}_{c_\beta})_b, (\bar{z}_{c_\delta})_b) \mid \beta \in B_{d,m}^\ell, \delta \in B_{d,m}^{\ell-1} \wedge \bar{z}_{c_\delta} - \bar{z}_{c_\beta} = q_{\vec{j}}\}.$$

Since Add_d is FA-recognizable, the relation $R_{\ell, \vec{j}}$ are FA-recognizable for all $\ell \geq 1$ and $\vec{j} \in \mathcal{I}_{d,m}$. Therefore, since the language $L_{N-1} \setminus L'_{N-1}$ is regular and S_f^{N-1} is FA-recognizable, the relation:

$$Q_{f, \vec{j}} = \{((\bar{z}_{c_\beta})_b, (\lambda_\delta)_b) \mid \beta \in \mathcal{K}'^N, \delta \in \bar{\mathcal{K}}^{N-1}, ((\bar{z}_{c_\beta})_b, (\bar{z}_{c_\delta})_b) \in R_{N, \vec{j}}, \\ ((\bar{z}_{c_\delta})_b, (\lambda_\delta)_b) \in S_f^{N-1}\}$$

is FA-recognizable. Since multiplication by a constant in $\mathbb{Z}[1/b]$ is FA-recognizable, see (11), we finally obtain that:

$$S_f'^N = \{((\bar{z}_{c_\beta})_b, (\lambda_\beta)_b) \mid ((\bar{z}_{c_\beta})_b, (\lambda_\delta)_b) \in Q_{f, \vec{j}}, \vec{j} \in \mathcal{I}_{d,m} \wedge \lambda_\beta = \sum_{(\delta, \vec{j}) \in \Delta_\beta} \lambda_\delta \lambda_{\vec{j}}\} \quad (17)$$

is FA-recognizable.

Remark 5. Similarly to Remark 4, the use of the identity $\lambda_\beta = \sum_{(\delta, \vec{j}) \in \Delta_\beta} \lambda_\delta \lambda_{\vec{j}}$ in (17) may lead to exponential growth in the number of states. In order to avoid it, instead of $S_f'^N$ one can use $\mathbf{S}_f'^N$ defined below. For a given $\beta \in \mathcal{K}'^N$ and $\vec{j} \in \mathcal{I}_{d,m}$, let $\lambda_{\beta, \vec{j}} = \lambda_\beta$ if $(\delta, \vec{j}) \in \Delta_\beta$ for some δ and 0, otherwise. Clearly, we have $\lambda_\beta = \sum_{\vec{j} \in \mathcal{I}_{d,m}} \lambda_{\beta, \vec{j}} \lambda_{\vec{j}}$. Now

let us define $\mathbf{S}_f'^N \subset \mathcal{L}_b^d \times (\prod_{\vec{j} \in \mathcal{I}_{d,m}} \mathcal{L}_b) \times \mathcal{L}_b$ as:

$$\mathbf{S}_f'^N = \{((\bar{z}_{c_\beta})_b, (\lambda_{\beta, (0, \dots, 0)})_b, \dots, (\lambda_{\beta, (m+1, \dots, m+1)})_b, (\lambda_\beta)_b) \mid \beta \in \mathcal{K}'^N, \\ \lambda_\beta = \sum_{\vec{j} \in \mathcal{I}_{d,m}} \lambda_{\beta, \vec{j}} \lambda_{\vec{j}}\}.$$

One can construct a deterministic finite automaton recognizing $\mathbf{S}_f'^N$ for which the number of states is polynomial in $m_{f, N-1}$ and m_N .

6.3 Examples

First we note that a spline $f \in \mathcal{S}_m(\mathcal{T})$ over a regular hierarchical mesh \mathcal{T} with bounded support $\text{supp } f$ is regular.

Let \mathcal{T}_d^0 be a mesh defined by the grid \mathcal{G}_d^0 . A constant function over \mathcal{T}_d^0 which takes the value $\lambda \in \mathbb{Z}[1/b]$ for every point $x \in \mathbb{R}^d$ is a regular spline in $\mathcal{S}_m(\mathcal{T}_d^0)$ for $m \geq 0$. This follows from the partition of unity property for B-splines: $\sum_{i=-\infty}^{\infty} N_{i,m}^0 = 1$ for

$m \geq 0$. Moreover, a linear function $\sum_{i=1}^d \alpha_i x_i$, for $\alpha_i \in \mathbb{Z}[1/b]$, $i = 1, \dots, d$ is a regular spline in $\mathcal{S}_m(\mathcal{T})$ for $m \geq 1$. Since the collection of regular splines is closed under taking the sum and multiplication by a constant $\lambda \in \mathbb{Z}[1/b]$, it is enough to prove it for the functions $f_{i,d}(\bar{x}) = x_i$ for $\bar{x} = (x_1, \dots, x_d) \in \mathbb{R}^d$ and $i = 1, \dots, d$. In order to prove the latter, it is enough only to show that the linear function $f(t) = t$, $t \in \mathbb{R}$ is a regular spline in $\mathcal{S}_m(\mathcal{T}_1^0)$. This follows from the identity $\sum_{i=-\infty}^{+\infty} c_{i,m} N_{i,m}^0(t) = t$ for $m \geq 1$, where $c_{i,m} = i + \frac{m+1}{2}$. This identity is proved by induction. For $m = 1$, we recall that (see (13)):

$$N_{i,1}^0(t) = \begin{cases} t - i, & i \leq t < i + 1, \\ i + 2 - t, & i + 1 \leq t < i + 2, \\ 0, & \text{otherwise.} \end{cases}$$

Therefore, for $t \in [i, i + 1]$, we have $\sum_{i=-\infty}^{+\infty} c_{i,1} N_{i,1}^0(t) = (i + 1)N_{i,1}^0(t) + ((i - 1) + 1)N_{i-1,1}^0(t) = (i + 1)(t - i) + i(i + 1 - t) = t$. The inductive step follows from the Cox-de Boor's formula (3) as follows. Assume that $\sum_{i=-\infty}^{+\infty} c_{i,m} N_{i,m}^0(t) = t$ holds for some $m \geq 1$. By (3), $\sum_{i=-\infty}^{+\infty} c_{i,m+1} N_{i,m+1}^0(t) = \sum_{i=-\infty}^{+\infty} c_{i,m+1} \left(\frac{t-i}{m+1} N_{i,m}^0(t) + \frac{i+m+2-t}{m+1} N_{i+1,m}^0(t) \right) = \sum_{i=-\infty}^{+\infty} \left(c_{i,m+1} \frac{t-i}{m+1} + c_{i-1,m+1} \frac{i+m+1-t}{m+1} \right) N_{i,m}^0(t) = \sum_{i=-\infty}^{+\infty} \left(\frac{t}{m+1} + \frac{m}{m+1} c_{i,m} \right) N_{i,m}^0(t) = t$. From the formula $c_{i,m} = i + \frac{m+1}{2}$ it is clear that $f(t) = t$ is a regular spline $\mathcal{S}_m(\mathcal{T}_1^0)$. It follows from Section 6.2 that constant and linear functions are regular splines in $\mathcal{S}_m(\mathcal{T})$, for $m \geq 0$ and $m \geq 1$, respectively, for every regular hierarchical mesh \mathcal{T} .




Figure 10: The left figure shows the spline $g(t) = \sum_{j=-\infty}^{\infty} c_j N_{4j,3}^0(t)$, where $c_j = 1$ if j is even and $c_j = -1$ if j is odd. The right figure shows the spline $h(t) = \sum_{j=0}^{\infty} c'_j N_{8j,3}^2(t) + \sum_{j=-\infty}^{-1} c'_j N_{8j+4,3}^2(t)$, where $c'_j = j + 1$ for $j \geq 0$ and $c'_j = -j$ for $j \leq -1$.

We show other two simple examples below. Let us consider a spline $g(t) = \sum_{j=-\infty}^{\infty} c_j N_{4j,3}^0(t)$, where $c_j = 1$ if j is even and $c_j = -1$ if j is odd, see Fig.10 (left). Clearly, g is a regular spline in $\mathcal{S}_3(\mathcal{T}_1^0)$ as well as in $\mathcal{S}_3(\mathcal{T})$ for every one-dimensional regular hierarchical mesh \mathcal{T} . Now let \mathcal{T}' be a one-dimensional hierarchical mesh generated by the domains $\Omega_1 = \Omega_2 = \bigcup_{i=0}^{\infty} ([2i, 2i + 1] \cup [-2i - 1, -2i])$. A spline

function $h(t) = \sum_{j=0}^{\infty} c'_j N_{8j,3}^2(t) + \sum_{j=-\infty}^{-1} c'_j N_{8j+4,3}^2(t)$, where $c'_j = j + 1$ for $j \geq 0$ and $c'_j = -j$ for $j \leq -1$, see Fig. 10 (right), is a regular spline in $\mathcal{S}_3(\mathcal{T}')$ as well as in $\mathcal{S}_3(\mathcal{T}'')$ for every one-dimensional regular hierarchical mesh \mathcal{T}'' generated by domains $\Omega^0 = \mathbb{R}^1 \supseteq \Omega^1 \supseteq \Omega^2 \supseteq \dots \supseteq \Omega^{N-1} \supseteq \Omega^N = \emptyset$ for $N > 4$, where Ω_1 and Ω_2 are as above.

Acknowledgements

The authors thank Cesare Bracco and Andre Nies for useful discussions.

References

- [1] Billera, L.: Homology theory of smooth splines: generic triangulations and a conjecture of strang. *Trans. Amer. Math. Soc.* 310, 325–340 (1988)
- [2] Boehm, W.: Inserting new knots into B-spline curves. *Computer-Aided Design* 12:4, 199–201 (1980)
- [3] Case, J., Jain, S., Seah, S., Stephan, F.: Automatic functions, linear time and learning. *Logical Methods in Computer Science* 9(3:19) (2013)
- [4] Deng, J., Chen, F., Li, X., Hu, C., Tong, W., Yang, Z., Feng, Y.: Polynomial splines over hierarchical T-meshes. *Graphical Models* 70, 76–86 (2008)
- [5] Dokken, T., Lyche, T., Pettersen, K.: Polynomial splines over locally refined box-partitions. *Computer Aided Geometric Design* 30:3, 331–356 (2013)
- [6] Epstein, D.B.A., Cannon, J.W., Holt, D.F., Levy, S.V.F., Paterson, M.S., Thurston, W.P.: *Word Processing in Groups*. Jones and Barlett Publishers. Boston, MA (1992)
- [7] Forsey, D., Bartels, R.: Hierarchical B-spline refinement. *Computer Graphics* 22, 205–212 (1988)
- [8] Gao, Z., Jain, S., Qi, J., Schlicht, P., Stephan, F., Tarr, J.: Ordered semiautomatic rings with applications to geometry. *Lecture Notes in Computer Science*, vol. 12038, pp. 141–153. Springer (2020)
- [9] Hopcroft, J., Motwani, R., Ullman, J.D.: *Introduction to Automata Theory, Languages, and Computation*. Addison-Wesley (2001)
- [10] Hughes, T., Cottrell, J., Bazilevs, Y.: Isogeometric analysis: CAD, finite elements, NURBS, exact geometry and mesh refinement. *Computer Methods in Applied Mechanics and Engineering* 194, 4135–4195 (2005)
- [11] Khoussainov, B., Nerode, A.: Automatic presentations of structures. In: Leivant, D. (ed.) *Logic and Computational Complexity*, *Lecture Notes in Computer Science*, vol. 960, pp. 367–392. Springer Berlin Heidelberg (1995)
- [12] Kraft, R.: Adaptive and linearly independent multilevel B-splines. In: Méhauté, A.L., Rabut, C., Schumaker, L.L. (eds.) *Surface Fitting and Multiresolution Methods*, pp. 209–218. Vanderbilt University Press, Nashville (1997)
- [13] Li, X., Zheng, J., Sederberg, T., Hughes, T., Scott, M.: On linear independence of T-spline blending functions. *Computer Aided Geometric Design* 29, 63–76 (2012)
- [14] Li, X., Chen, F., Kang, H., Deng, J.: A survey on the local refinable splines. *Sci China Math* 59:4, 617–644
- [15] Mokriš, D., Jüttler, B., Giannelli, C.: On the completeness of hierarchical tensor-product B-splines. *Journal of Computational and Applied Mathematics* 271, 53–70 (2014)

- [16] Mourrain, B.: On the dimension of spline spaces on planar T-meshes. *Math. Comp.* 83, 847–871 (2014)
- [17] Nies, A., Semukhin, P.: Finite automata presentable abelian groups. *Annals of Pure and Applied Logic* 161(3), 458–467 (2009)
- [18] Prautzsch, H., Boehm, W., Paluszny, M.: *Bézier and B-Spline Techniques*. Springer (2002)
- [19] Scott, M., Li, X., Sederberg, T., Hughes, T.: Local refinement of analysis-suitable T-splines. *Computer Methods in Applied Mechanics and Engineering* 213–216, 206–222 (2012)
- [20] Sederberg, T., Zheng, J., Bakenov, A., Nasri, A.: T-splines and T-NURCCS. *ACM Trans. Graph.* 22, 477–484 (2003)
- [21] Shirley, P., Ashikhmin, M., Gleicher, M., Marschner, S., Reinhard, E., Sung, K., Thompson, W., Willemssen, P.: *Fundamentals of computer graphics*. AK Peters (2005)
- [22] Stephan, F.: Automatic structures — recent results and open questions. *Journal of Physics: Conference Series* 622, 012013 (jun 2015)