

Leap eccentric connectivity index of some graph operations with subdivided edges ^{*}

Ling Song, Zikai Tang[†]

School of Mathematics and Statistics, Hunan Normal University,
Changsha, Hunan 410081, P. R. China.

Abstract: The leap eccentric connectivity index of G is defined as

$$L\xi^C(G) = \sum_{v \in V(G)} d_2(v|G)e(v|G)$$

where $d_2(v|G)$ be the second degree of the vertex v and $e(v|G)$ be the eccentricity of the vertex v in G . In this paper, we first give a counterexample for that if G be a graph and $S(G)$ be its the subdivision graph, then each vertex $v \in V(G)$, $e(v|S(G)) = 2e(v|G)$ by Yarahmadi in [15] in Theorem 3.1. And we describe the upper and lower bounds of the leap eccentric connectivity index of four graphs based on subdivision edges, and then give the expressions of the leap eccentric connectivity index of join graph based on subdivision, finally, give the bounds of the leap eccentric connectivity index of four variants of the corona graph.

Keywords: Eccentricity; Second degrees; Leap eccentric connectivity index; Subdivision edges.

AMS Subject classification: 05C07

1 Introduction

In this paper, we only consider simple, undirected, finite graphs. Let G denote a graph with n vertices and m edge sets. Denoted by $d_G(u, v)$ the shortest path of length connecting u and v in G , for vertices $u, v \in V(G)$. For a vertex v and a positive integer k , we let $N_k(v|G)$ denote the open k -neighborhood of vertex v in G and defined as $N_k(v|G) = \{u \in V(G) | d_G(u, v) = k\}$. Let $d_k(v|G)$ denote the k degree of the vertex in G , expressed as the number of vertices in the open k -neighborhood of vertex v in G , that is, $d_k(v|G) = |N_k(v|G)|$. We can see that for any vertex v in G there are $d_1(v|G) = |N_1(v|G)|$

^{*}This work is supported by the Department of Education of Hunan Province(19A318).

[†]Corresponding author: zikaitang@163.com

and $d_2(v|G) = |N_2(v|G)|$. The eccentricity is defined as $e(v|G)$, for a vertex v in G , which represents the maximum distance from vertex v to other vertices in the graph, that is, $e(v|G) = \max\{d_G(u, v) | u \in V(G)\}$. For any vertex in the graph, we define the maximum eccentricity value as the diameter $\text{diam}(G)$. We let $V_e^\alpha(G) \subseteq V(G)$ denote the set of vertices in G where the eccentricity is equal to α , where $\alpha = 1, 2, \dots, \text{diam}(G)$, obviously $V_e^1(G)$ represents the set of vertices in G that have a eccentricity of 1 to other vertices, the degree of these vertices is $n - 1$, we call these vertices full vertices. Let $H \subseteq V(G)$ denote any subset of vertices of G , then the induced subgraph $\langle H \rangle$ of G is the graph that the vertex set is H , and the edge set is the edge in graph G with the vertex in H as the endpoint. If there are no graphs isomorphic to graph F in all induced subgraphs of graph G , we call graph G the F -free graph. For other terms and symbols that are not defined here, please refer to the reference [2].

Structure descriptors based on molecular graphs are often called topological indices and have very important meanings. In 1972, Gutman and Trinajestic [7] introduced the first, and elaborated it in [6]. The definition is as follows:

$$M_1(G) = \sum_{v \in V(G)} d_1^2(v|G).$$

Recently, some scholars replaced the vertex degree in the first Zagreb index with the second degree, and proposed a new index about Zagreb index, defined as the first leap Zagreb index [5]. The first leap Zagreb index is defined as:

$$LM_1(G) = \sum_{v \in V(G)} d_2^2(v|G)$$

In addition to the above mentioned degree-based topological indexes, some distance-based topological indexes have also caused extensive research. In 2004, Dankelmann introduced the eccentricity sum index [4], defined as:

$$\theta(G) = \sum_{v \in V(G)} e(v|G)$$

Sharma proposed the eccentric connectivity index [11], defined as:

$$\xi^C(G) = \sum_{v \in V(G)} d_1(v|G)e(v|G)$$

Recently, Naji proposed the leap eccentric connectivity index [10], defined as:

$$L\xi^C(G) = \sum_{v \in V(G)} d_2(v|G)e(v|G)$$

Regarding the study of the leap eccentric connectivity index, currently Manjunathe et al. [9] gave the values of the leap eccentric connectivity index for some graph operations contains cartesian product, composition, disjunctions, symmetric difference and corona product. The leap eccentric connectivity index of some other graphs has not been studied. This paper mainly studies the upper and lower bounds of the leap eccentric connectivity index of some graph operations with subdivided edges. Next, first introduce some basic graphs.

For a given graph G , the line graph is denoted by $L(G)$, and its vertex set is the edge set of G , where two vertices are adjacent if and only if they are adjacent edges in G . Cvetkovic [3] proposed four graphs based on subdivision edges, the subdivision graph $S(G)$, the $Q(G)$ graph, the $R(G)$ graph and the total graph $T(G)$. The subdivision graph of graph G is represented as $S(G)$, which represents the graph obtained by adding a vertex to each edge of G . The vertex set of $S(G)$ can be divided into two parts: one is the vertex set $V(G)$ in the original G , and the other is the vertex inserted on each edge, which is the original edge set $E(G)$ of G . The $Q(G)$ graph of G is a graph obtained by inserting new vertices on each edge of G , and then connecting these new vertex pairs located on adjacent edges of G . The $R(G)$ graph of G is defined as a graph obtained by adding new vertices corresponding to each edge to G and connecting each new vertex with the two endpoints of its corresponding edge, which can also be understood as the graph obtained by replacing each edge with a triangle in G . The total graph $T(G)$ of G is a graph whose vertex set is the union of the vertex set of G and the edge set of G , and the two vertices of $T(G)$ are adjacent if and only if the corresponding elements in G are adjacent or incident. In recent years, many researchers have paid attention to the subdivision join of several graphs. Indulal [8] proposed two new joins, subdivision vertex join $G \dot{\vee} H$, subdivision edge join $G \dot{\wedge} H$. F. Wen [13] proposed a new graph operation based on subdivision and join on three graphs (G and H_1, H_2), called subdivision vertex edge join $G^S \Delta (H_1^V \vee H_2^S)$, where G and H_1, H_2 vertex sets disjoint. Barik [1] proposed other variants of the corona graph, such as: subdivision double corona, $Q(G)$ graph double corona, $R(G)$ graph double

corona, total graph double corona.

In this paper, we first give a counterexample for that if G be a graph and $S(G)$ be its the subdivision graph, then each vertex $v \in V(G)$, $e(v|S(G)) = 2e(v|G)$ by Yarahmadi in [15] in Theorem 3.1. And we describe the upper and lower bounds of the leap eccentric connectivity index of four graphs based on subdivision edges, the subdivision graph $S(G)$, the $Q(G)$ graph, the $R(G)$ graph and the total graph $T(G)$, and then give the expressions of the leap eccentric connectivity index of join graph based on subdivision, finally, give the bounds of the leap eccentric connectivity index of four variants of the corona graph.

2 Leap eccentric connectivity index of four graphs based on subdivision edges

This section mainly introduces the expressions of the leap eccentric connectivity index of subdivision graphs $S(G)$, the $Q(G)$ graph, the $R(G)$ graph and the total graph $T(G)$ based on other invariants. If $|V(G)| = n$, $|E(G)| = m$, we find that $S(G)$, $Q(G)$, $R(G)$, and $T(G)$ all have $n + m$ vertices. In these four graphs, we record the original n vertices from graph G as $v \in V(G)$, and insert the m vertices corresponding to each edge of the graph G are denoted as $e \in E(G)$. Taking the graph $G = C_4$ as an example. In Figure 1, we describe the line graph $L(G)$, $S(G)$, $Q(G)$, $R(G)$, and $T(G)$, where black coloured vertices represent vertices from the vertex set in the original graph G , and blue coloured vertices represent newly inserted the vertex corresponding to each edge in G .




Figure 1: **Graphs** C_4 , $L(C_4)$, $S(C_4)$, $Q(C_4)$, $R(C_4)$, $T(C_4)$

In reference [15], Yarahmadi gave the inequality of the eccentricity of these four graphs, but we found that (i) in Theorem 3.1 for each vertex $v \in V(G)$, $e(v|S(G)) = 2e(v|G)$, the conclusion about the vertex eccentricity in the subdivision graph is incorrect. We can see in Figure 2 that for vertex v , $e(v|S(G)) = 2e(v|G) + 1$. In a graph that does not contain odd circles, $e(v|S(G)) = 2e(v|G)$ holds for vertex v .




Figure 2: **A graph satisfying $e(v|S(G)) = 2e(v|G) + 1$**

So we correct the conclusion of Theorem 3.1 in reference [15] to get the following lemma.

Lemma 2.1. *Let G be a connected graph, then*

(i) *For any vertex $v \in V(G)$, $2e(v|G) \leq e(v|S(G)) \leq 2e(v|G) + 1$;*

(ii) *For any vertex $e \in E(G)$, $2e(e|L(G)) \leq e(e|S(G)) \leq 2e(e|L(G)) + 1$.*

Then we introduce some conclusions related to distance in these four graphs.

Lemma 2.2. [14] *For any two vertices $v, v' \in V(G)$, $e, e' \in E(G)$, we have*

$$\frac{1}{2}d_{S(G)}(v, v') = d_{Q(G)}(v, v') - 1 = d_{R(G)}(v, v') = d_{T(G)}(v, v') = d_G(v, v')$$

$$\frac{1}{2}d_{S(G)}(e, e') = d_{Q(G)}(e, e') = d_{R(G)}(e, e') - 1 = d_{T(G)}(e, e') = d_{L(G)}(e, e')$$

Next, we introduce the conclusions about the second degree of vertices.

Lemma 2.3. [12] *Let G be a connected graph with n vertices and m edges, then*

$$\sum_{v \in V(G)} d_2(v|G) \leq M_1(G) - 2m$$

with equality if and only if G is a $\{C_3, C_4\}$ -free graph.

2.1 The subdivision graph $S(G)$

Observing graph $S(G)$ we can get the following conclusion.

Lemma 2.4. *Let G be a connected graph, then*

(i) *For any vertex $v \in V(G)$, $d_2(v|S(G)) = d_1(v|G)$;*

(ii) *For any vertex $e \in E(G)$, $d_2(e|S(G)) = d_1(e|L(G))$.*

Next we will give the bound of the leap eccentric connectivity index of the subdivision graph.

Theorem 2.5. *Let G be a connected graph, then*

$$2\xi^C(G) + 2\xi^C(L(G)) \leq L\xi^C(S(G)) \leq 2\xi^C(G) + 2|E(G)| + 2\xi^C(L(G)) + 2|E(L(G))|$$

Proof. From Lemma 2.1 and 2.4, we can get:

$$\begin{aligned} L\xi^C(S(G)) &= \sum_{v \in V(G)} d_2(v|S(G))e(v|S(G)) + \sum_{e \in E(G)} d_2(e|S(G))e(e|S(G)) \\ &\geq 2 \sum_{v \in V(G)} d_1(v|G)e(v|G) + 2 \sum_{e \in E(G)} d_1(e|L(G))e(e|L(G)) \\ &= 2\xi^C(G) + 2\xi^C(L(G)) \end{aligned}$$

and

$$\begin{aligned} L\xi^C(S(G)) &= \sum_{v \in V(G)} d_2(v|S(G))e(v|S(G)) + \sum_{e \in E(G)} d_2(e|S(G))e(e|S(G)) \\ &\leq \sum_{v \in V(G)} d_1(v|G)(2e(v|G) + 1) + \sum_{e \in E(G)} d_1(e|L(G))(2e(e|L(G)) + 1) \\ &= 2\xi^C(G) + 2|E(G)| + 2\xi^C(L(G)) + 2|E(L(G))| \end{aligned}$$

The proof is completed. \square

We find that when graph G does not contain odd circles, $e(v|S(G)) = 2e(v|G)$ holds. So for a connected bipartite graph, $e(v|S(G)) = 2e(v|G)$ is always true, so we can get the following conclusion.

Corollary 2.6. *Let G be a connected bipartite graph, then*

$$2\xi^C(G) + 2\xi^C(L(G)) \leq L\xi^C(S(G)) \leq 2\xi^C(G) + 2\xi^C(L(G)) + 2|E(L(G))|.$$

When the eccentricity of all vertices in graph G is reached at the pendant vertex, $e(e|S(G)) = 2e(e|L(G)) + 1$ holds for all vertices. Then we can get a connected bipartite graph where the eccentricity of all vertices is reached at the pendant vertex, the following equation holds.

Corollary 2.7. *Let G be a connected bipartite graph and the eccentricity of all vertices are reached at the pendant vertex, then*

$$L\xi^C(S(G)) = 2\xi^C(G) + 2\xi^C(L(G)) + 2|E(L(G))|.$$

2.2 The $Q(G)$ graph

Lemma 2.8. [15] *Let G be a connected graph, then*

(i) *For any vertex $v \in V(G)$, $e(v|Q(G)) = e(v|G) + 1$;*

(ii) *For any vertex $e \in E(G)$, $e(e|L(G)) \leq e(e|Q(G)) \leq e(e|L(G)) + 1$.*

Observing graph $Q(G)$ we can get the following conclusion.

Lemma 2.9. *Let G be a connected graph, then*

(i) *For any vertex $v \in V(G)$, $d_2(v|Q(G)) \geq d_1(v|G) + d_2(v|G)$, with equality if and only if G is a $\{C_3, C_4\}$ -free graph;*

(ii) *For any vertex $e \in E(G)$, $d_2(e|Q(G)) \leq d_1(e|L(G)) + d_2(e|L(G))$, with equality if and only if G is a $\{C_3\}$ -free graph.*

Theorem 2.10. *Let G be a $\{C_3, C_4\}$ -free connected graph, then the following inequality is established:*

(i) $L\xi^C(Q(G)) \geq \xi^C(G) + L\xi^C(G) + M_1(G) + \xi^C(L(G)) + L\xi^C(L(G));$

(ii) $L\xi^C(Q(G)) \leq \xi^C(G) + L\xi^C(G) + M_1(G) + \xi^C(L(G)) + L\xi^C(L(G)) + M_1(L(G)).$

Proof. G is a $\{C_3, C_4\}$ -free connected graph. According to Lemma 2.9, for any vertex $v \in V(G)$, $d_2(v|Q(G)) = d_1(v|G) + d_2(v|G)$ holds, for any vertex $e \in E(G)$, $d_2(e|Q(G)) = d_1(e|L(G)) + d_2(e|L(G))$ holds. And according to Lemma 2.8, we can get

$$\begin{aligned}
L\xi^C(Q(G)) &= \sum_{v \in V(G)} d_2(v|Q(G))e(v|Q(G)) + \sum_{e \in E(G)} d_2(e|Q(G))e(e|Q(G)) \\
&\geq \sum_{v \in V(G)} [d_1(v|G) + d_2(v|G)](e(v|G) + 1) \\
&\quad + \sum_{e \in E(G)} [d_1(e|L(G)) + d_2(e|L(G))]e(e|L(G)) \\
&= \xi^C(G) + 2|E(G)| + L\xi^C(G) + \sum_{v \in V(G)} d_2(v|G) + \xi^C(L(G)) + L\xi^C(L(G))
\end{aligned}$$

In Lemma 2.3, we know that when G is a $\{C_3, C_4\}$ -free connected graph, there is

$$\sum_{v \in V(G)} d_2(v|G) = M_1(G) - 2|E(G)|.$$

Taking this equation into the above formula, we can get

$$\begin{aligned}
L\xi^C(Q(G)) &\geq \xi^C(G) + 2|E(G)| + L\xi^C(G) + \sum_{v \in V(G)} d_2(v|G) + \xi^C(L(G)) + L\xi^C(L(G)) \\
&= \xi^C(G) + 2|E(G)| + L\xi^C(G) + M_1(G) - 2|E(G)| + \xi^C(L(G)) + L\xi^C(L(G)) \\
&= \xi^C(G) + L\xi^C(G) + M_1(G) + \xi^C(L(G)) + L\xi^C(L(G))
\end{aligned}$$

Inequality (i) is proved.

Similarly, we prove the second inequality (ii).

$$\begin{aligned}
L\xi^C(Q(G)) &= \sum_{v \in V(G)} d_2(v|Q(G))e(v|Q(G)) + \sum_{e \in E(G)} d_2(e|Q(G))e(e|Q(G)) \\
&\leq \sum_{v \in V(G)} [d_1(v|G) + d_2(v|G)](e(v|G) + 1) \\
&\quad + \sum_{e \in E(G)} [d_1(e|L(G)) + d_2(e|L(G))](e(e|L(G)) + 1) \\
&= \xi^C(G) + 2|E(G)| + L\xi^C(G) + \sum_{v \in V(G)} d_2(v|G) + \xi^C(L(G)) + L\xi^C(L(G)) \\
&\quad + 2|E(L(G))| + \sum_{e \in E(G)} d_2(e|L(G)) \\
&= \xi^C(G) + 2|E(G)| + L\xi^C(G) + M_1(G) - 2|E(G)| + \xi^C(L(G)) + L\xi^C(L(G)) \\
&\quad + 2|E(L(G))| + M_1(L(G)) - 2|E(L(G))| \\
&= \xi^C(G) + L\xi^C(G) + M_1(G) + \xi^C(L(G)) + L\xi^C(L(G)) + M_1(L(G))
\end{aligned}$$

The proof is completed. \square

When G is a $\{C_3, C_4\}$ -free connected bipartite graph and the eccentricity of all vertices in graph G is reached at the pendant vertex, the equal sign of the second inequality in Theorem 2.10 can be equal, that is, for any vertex $e \in E(G)$, $e(e|Q(G)) = e(e|L(G)) + 1$ established. So we can get the following corollary.

Corollary 2.11. *Let G is a $\{C_3, C_4\}$ -free connected bipartite graph and the eccentricity of all vertices in graph G is reached at the pendant vertex, then*

$$L\xi^C(Q(G)) = \xi^C(G) + L\xi^C(G) + M_1(G) + \xi^C(L(G)) + L\xi^C(L(G)) + M_1(L(G))$$

2.3 The $R(G)$ graph

Lemma 2.12. [15] *Let G be a connected graph, then*

(i) *For any vertex $v \in V(G)$, $e(v|G) \leq e(v|R(G)) \leq e(v|G) + 1$;*

(ii) For any vertex $e \in E(G)$, $e(e|R(G)) = e(e|L(G)) + 1$.

Observing graph $R(G)$ we can get the following conclusion.

Lemma 2.13. *Let G be a connected graph, then*

(i) For any vertex $v \in V(G)$, $d_2(v|R(G)) \geq 2d_2(v|G)$, with equality if and only if G is a $\{C_3, C_4\}$ -free graph;

(ii) For any vertex $e \in E(G)$, $d_2(e|R(G)) \leq 2d_1(e|L(G))$, with equality if and only if G is a $\{C_3\}$ -free graph.

Theorem 2.14. *Let G be a $\{C_3, C_4\}$ -free connected graph, then*

(i) $L\xi^C(R(G)) \geq 2L\xi^C(G) + 2\xi^C(L(G)) + 4|E(L(G))|$;

(ii) $L\xi^C(R(G)) \leq 2L\xi^C(G) + 2M_1(G) - 4|E(G)| + 2\xi^C(L(G)) + 4|E(L(G))|$.

Proof. G is a $\{C_3, C_4\}$ -free connected graph. According to Lemma 2.13, for any vertex $v \in V(G)$, $d_2(v|R(G)) = 2d_2(v|G)$ holds, for any vertex $e \in E(G)$, $d_2(e|R(G)) = 2d_1(e|L(G))$ holds. And according to Lemma 2.12, we can get

$$\begin{aligned} L\xi^C(R(G)) &= \sum_{v \in V(G)} d_2(v|R(G))e(v|R(G)) + \sum_{e \in E(G)} d_2(e|R(G))e(e|R(G)) \\ &\geq 2 \sum_{v \in V(G)} d_2(v|G)e(v|G) + 2 \sum_{e \in E(G)} d_1(e|L(G))(e(e|L(G)) + 1) \\ &= 2L\xi^C(G) + 2\xi^C(L(G)) + 4|E(L(G))| \end{aligned}$$

and

$$\begin{aligned} L\xi^C(R(G)) &= \sum_{v \in V(G)} d_2(v|R(G))e(v|R(G)) + \sum_{e \in E(G)} d_2(e|R(G))e(e|R(G)) \\ &\leq 2 \sum_{v \in V(G)} d_2(v|G)(e(v|G) + 1) + 2 \sum_{e \in E(G)} d_1(e|L(G))(e(e|L(G)) + 1) \\ &= 2L\xi^C(G) + 2 \sum_{v \in V(G)} d_2(v|G) + 2\xi^C(L(G)) + 4|E(L(G))| \end{aligned}$$

By Lemma 2.3, we have

$$\begin{aligned} L\xi^C(R(G)) &\leq 2L\xi^C(G) + 2(M_1(G) - 2|E(G)|) + 2\xi^C(L(G)) + 4|E(L(G))| \\ &= 2L\xi^C(G) + 2M_1(G) - 4|E(G)| + 2\xi^C(L(G)) + 4|E(L(G))| \end{aligned}$$

The proof is completed. \square

When G is a $\{C_3, C_4\}$ -free connected graph with an odd circle, the equal sign on the right side of Theorem 2.14 is equal, which means that for any vertex $v \in V(G)$,

$e(v|R(G)) = e(v|G) + 1$ holds. When G is a $\{C_3, C_4\}$ -free connected bipartite graph, then for all vertices $v \in V(G)$, $e(v|R(G)) = e(v|G)$ holds. So we can get the following corollary.

Corollary 2.15. *Let G is a $\{C_3, C_4\}$ -free connected bipartite graph, then*

$$L\xi^C(R(G)) = 2L\xi^C(G) + 2\xi^C(L(G)) + 4|E(L(G))|.$$

2.4 The $T(G)$ graph

Lemma 2.16. [15] *Let G be a connected graph, then*

(i) *For any vertex $v \in V(G)$, $e(v|G) \leq e(v|T(G)) \leq e(v|G) + 1$;*

(ii) *For any vertex $e \in E(G)$, $e(e|L(G)) \leq e(e|T(G)) \leq e(e|L(G)) + 1$.*

Observing graph $T(G)$ we can get the following conclusion.

Lemma 2.17. *Let G be a connected graph, then*

(i) *For any vertex $v \in V(G)$, $d_2(v|T(G)) \geq 2d_2(v|G)$, with equality if and only if G is a $\{C_3, C_4\}$ -free graph;*

(ii) *For any vertex $e \in E(G)$, $d_2(e|T(G)) \leq d_1(e|L(G)) + d_2(e|L(G))$, with equality if and only if G is a $\{C_3\}$ -free graph.*

Theorem 2.18. *Let G be a $\{C_3, C_4\}$ -free connected graph, then the following inequality is established:*

(i) $L\xi^C(T(G)) \geq 2L\xi^C(G) + \xi^C(L(G)) + L\xi^C(L(G))$;

(ii) $L\xi^C(T(G)) \leq 2L\xi^C(G) + 2M_1(G) - 4|E(G)| + \xi^C(L(G)) + L\xi^C(L(G)) + M_1(L(G))$.

Proof. G is a $\{C_3, C_4\}$ -free connected graph. According to Lemma 2.17, for any vertex $v \in V(G)$, $d_2(v|T(G)) = 2d_2(v|G)$ holds, for any vertex $e \in E(G)$, $d_2(e|T(G)) = d_1(e|L(G)) + d_2(e|L(G))$ holds. And according to Lemma 2.16, we can get

$$\begin{aligned} L\xi^C(T(G)) &= \sum_{v \in V(G)} d_2(v|T(G))e(v|T(G)) + \sum_{e \in E(G)} d_2(e|T(G))e(e|T(G)) \\ &\geq 2 \sum_{v \in V(G)} d_2(v|G)e(v|G) + \sum_{e \in E(G)} [d_1(e|L(G)) + d_2(e|L(G))]e(e|L(G)) \\ &= 2L\xi^C(G) + \xi^C(L(G)) + L\xi^C(L(G)) \end{aligned}$$

and

$$\begin{aligned}
L\xi^C(T(G)) &= \sum_{v \in V(G)} d_2(v|T(G))e(v|T(G)) + \sum_{e \in E(G)} d_2(e|T(G))e(e|T(G)) \\
&\leq 2 \sum_{v \in V(G)} d_2(v|G)(e(v|G) + 1) \\
&\quad + \sum_{e \in E(G)} [d_1(e|L(G)) + d_2(e|L(G))](e(e|L(G)) + 1) \\
&= 2L\xi^C(G) + 2 \sum_{v \in V(G)} d_2(v|G) + \xi^C(L(G)) + L\xi^C(L(G)) + 2|E(L(G))| \\
&\quad + \sum_{e \in E(G)} d_2(e|L(G))
\end{aligned}$$

By Lemma 2.3, we have

$$\begin{aligned}
L\xi^C(T(G)) &\leq 2L\xi^C(G) + 2(M_1(G) - 2|E(G)|) + \xi^C(L(G)) + L\xi^C(L(G)) + 2|E(L(G))| \\
&\quad + M_1(L(G)) - 2|E(L(G))| \\
&= 2L\xi^C(G) + 2M_1(G) - 4|E(G)| + \xi^C(L(G)) + L\xi^C(L(G)) + M_1(L(G))
\end{aligned}$$

The proof is completed. \square

We find that when the graph G is a $\{C_3, C_4\}$ -free connected graph that does not contain odd cycles, $e(v|T(G)) = e(v|G)$ holds for any vertex v in G . So for a $\{C_3, C_4\}$ -free connected bipartite graph, $e(v|T(G)) = e(v|G)$ is always true, so we can get the following conclusion.

Corollary 2.19. *Let G is a $\{C_3, C_4\}$ -free connected bipartite graph, then*

- (i) $L\xi^C(T(G)) \geq 2L\xi^C(G) + \xi^C(L(G)) + L\xi^C(L(G))$;
- (ii) $L\xi^C(T(G)) \leq 2L\xi^C(G) + \xi^C(L(G)) + L\xi^C(L(G)) + M_1(L(G))$.

When graph G is a $\{C_3, C_4\}$ -free connected graph where the eccentricity of all vertices is reached at the pendant vertex, then $e(e|T(G)) = e(e|L(G)) + 1$ holds for all vertices $e \in E(G)$. So we can get the following equation hold.

Corollary 2.20. *Let G is a $\{C_3, C_4\}$ -free connected bipartite graph and the eccentricity of all vertices is reached at the pendant vertex, then*

$$L\xi^C(T(G)) = 2L\xi^C(G) + \xi^C(L(G)) + L\xi^C(L(G)) + M_1(L(G)).$$

3 The leap eccentric connectivity index of join graph based on subdivision

In this section, we give the expressions of the leap eccentric connectivity index of several join graph based on subdivision.

3.1 Subdivision vertex join

Definition 3.1. Let G and H be two disjoint graphs with vertices n and n_1 , and edges m and m_1 , respectively. The subdivision vertex join of graphs G and H is denoted as $G\dot{\vee}H$, and is a graph obtained by connecting the i -th ($i = 1, 2, \dots, n$) vertex in original G in $S(G)$ and all vertices of H .

Lemma 3.2. Let G and H be two disjoint connected graphs with vertex numbers $n \geq 3$ and n_1 , and edge numbers m and m_1 respectively, then for any vertex $v \in V(G\dot{\vee}H)$, we have

(i) If graph $G \cong S_n$, then

$$e(v|G\dot{\vee}H) = \begin{cases} 2 \text{ or } 3, & \text{if } v \in V(G) \\ 3, & \text{if } e \in E(G) \\ 2, & \text{if } v \in V(H) \end{cases}$$

(ii) If graph $G \not\cong S_n$, then

$$e(v|G\dot{\vee}H) = \begin{cases} 3, & \text{if } v \in V(G) \\ 3 \leq e(e|G\dot{\vee}H) \leq 4, & \text{if } e \in E(G) \\ 2, & \text{if } v \in V(H) \end{cases}$$

(iii)

$$d_2(v|G\dot{\vee}H) = \begin{cases} n - 1, & \text{if } v \in V(G) \\ d_1(e|L(G)) + n_1, & \text{if } e \in E(G) \\ n_1 - 1 - d_1(v|H) + m, & \text{if } v \in V(H) \end{cases}$$

Theorem 3.3. Let G and H be two disjoint connected graphs with vertex numbers $n \geq 3$ and n_1 , and edge numbers m and m_1 , and $G \cong S_n$, then

$$L\xi^C(G\dot{\vee}H) = (3n - 1)(n - 1) + (3n + 2n_1 - 3)(n + n_1 - 2) - 4m_1$$

Proof. When the graph $G \cong S_n = K_{1,n-1}$, we know that there is $e(v|G\dot{\vee}H) = 2$ for the original center vertex in G , and $e(v|G\dot{\vee}H) = 3$ for the other vertices in G , and at this

time $m = n - 1$, the line graph of the star graph is complete graph K_{n-1} . Then,

$$\begin{aligned}
L\xi^C(G\dot{\vee}H) &= \sum_{v \in V(G\dot{\vee}H)} d_2(v|G\dot{\vee}H)e(v|G\dot{\vee}H) \\
&= 2(n-1) + 3(n-1)^2 + 3 \sum_{e \in E(G)} (n-2+n_1) \\
&\quad + 2 \sum_{v \in V(H)} (n_1-1-d_1(v|H)+n-1) \\
&= (n-1)(2+3n-3) + 3(n+n_1-2)(n-1) + 2(n+n_1-2)n_1 - 4m_1 \\
&= (3n-1)(n-1) + (3n+2n_1-3)(n+n_1-2) - 4m_1
\end{aligned}$$

The proof is completed. \square

Theorem 3.4. *Let G and H be two disjoint connected graphs with vertex numbers $n \geq 3$ and n_1 , and edge numbers m and m_1 , and $G \not\cong S_n$, then*

$$(i) L\xi^C(G\dot{\vee}H) \geq 3(n-1)n + 6|E(L(G))| + (2n_1 + 5m - 2)n_1 - 4m_1;$$

$$(ii) L\xi^C(G\dot{\vee}H) \leq 3(n-1)n + 8|E(L(G))| + 2(n_1 + 3m - 1)n_1 - 4m_1.$$

Proof.

$$\begin{aligned}
L\xi^C(G\dot{\vee}H) &= \sum_{v \in V(G\dot{\vee}H)} d_2(v|G\dot{\vee}H)e(v|G\dot{\vee}H) \\
&= (n-1) \sum_{v \in V(G)} 3 + \sum_{e \in E(G)} (d_1(e|L(G)) + n_1)e(e|G\dot{\vee}H) \\
&\quad + 2 \sum_{v \in V(H)} (n_1-1-d_1(v|H)+m)
\end{aligned}$$

For any vertex of $e \in E(G)$, we have $3 \leq e(e|G\dot{\vee}H) \leq 4$, from which we can calculate the upper and lower bounds of $L\xi^C(G\dot{\vee}H)$.

$$\begin{aligned}
L\xi^C(G\dot{\vee}H) &\geq 3(n-1)n + 3 \sum_{e \in E(G)} (d_1(e|L(G)) + n_1) + 2(n_1 + m - 1)n_1 - 4m_1 \\
&= 3(n-1)n + 6|E(L(G))| + 3n_1m + 2(n_1 + m - 1)n_1 - 4m_1 \\
&= 3(n-1)n + 6|E(L(G))| + (2n_1 + 5m - 2)n_1 - 4m_1
\end{aligned}$$

Similarly,

$$\begin{aligned}
L\xi^C(G\dot{\vee}H) &\leq 3(n-1)n + 4 \sum_{e \in E(G)} (d_1(e|L(G)) + n_1) + 2(n_1 + m - 1)n_1 - 4m_1 \\
&= 3(n-1)n + 8|E(L(G))| + 4n_1m + 2(n_1 + m - 1)n_1 - 4m_1 \\
&= 3(n-1)n + 8|E(L(G))| + 2(n_1 + 3m - 1)n_1 - 4m_1
\end{aligned}$$

The proof is completed. \square

When the eccentricity of all vertices in graph G is reached at the pendant vertex, $e(e|G\dot{\vee}H) = 3$ holds for all vertices $e \in E(G)$. So we can get the following corollary.

Corollary 3.5. *Let G and H be two disjoint connected graphs with vertex numbers $n \geq 3$ and n_1 , and edge numbers m and m_1 , and $G \not\cong S_n$ where the eccentricity of all vertices is reached at the pendant vertex, then*

$$L\xi^C(G\dot{\vee}H) = 3(n-1)n + 6|E(L(G))| + (2n_1 + 5m - 2)n_1 - 4m_1.$$

3.2 Subdivision edge join

Definition 3.6. *Let G and H be two disjoint graphs with vertices n and n_1 , and edges m and m_1 , respectively. The subdivision edge join of graphs G and H is denoted as $G\underline{\vee}H$, and is a graph obtained by connecting the newly inserted j -th ($j = 1, 2, \dots, m$) vertex corresponding to each edge of G in $S(G)$ and all vertices of H .*

Lemma 3.7. *Let G and H be two disjoint connected graphs with vertex numbers $n \geq 3$ and n_1 , and edge numbers m and m_1 respectively, then for any vertex $v \in V(G\underline{\vee}H)$, we have*

(i) *If graph $G \cong S_n$, then*

$$e(v|G\underline{\vee}H) = \begin{cases} 2 \text{ or } 4, & \text{if } v \in V(G) \\ 3, & \text{if } e \in E(G) \\ 2, & \text{if } v \in V(H) \end{cases}$$

(ii) *If graph $G \not\cong S_n$, then*

$$e(v|G\underline{\vee}H) = \begin{cases} 3, & \text{if } v \in V(G), e(v|G) = 1 \\ 4, & \text{if } v \in V(G), e(v|G) > 1 \\ 3, & \text{if } e \in E(G) \\ 2, & \text{if } v \in V(H) \end{cases}$$

(iii)

$$d_2(v|G\underline{\vee}H) = \begin{cases} d_1(v|G) + n_1, & \text{if } v \in V(G) \\ m - 1, & \text{if } e \in E(G) \\ n_1 - 1 - d_1(v|H) + n, & \text{if } v \in V(H) \end{cases}$$

Theorem 3.8. *Let G and H be two disjoint connected graphs with vertex numbers $n \geq 3$ and n_1 , and edge numbers m and m_1 , and $G \cong S_n$, then*

$$L\xi^C(G\underline{\vee}H) = (3n + 4n_1 - 2)(n - 1) + 2(n + n_1 - 1)(n_1 + 1) - 4m_1$$

Proof. When the graph $G \cong S_n = K_{1,n-1}$, we know that there is $e(v|G \vee H) = 2$ for the original center vertex in G , and $e(v|G \vee H) = 4$ for the other vertices in G , and at this time $m = n - 1$, the line graph of the star graph is complete graph K_{n-1} . Then,

$$\begin{aligned}
L\xi^C(G \vee H) &= \sum_{v \in V(G \vee H)} d_2(v|G \vee H)e(v|G \vee H) \\
&= 2(n + n_1 - 1) + 4(n_1 + 1)(n - 1) + 3(n - 2)(n - 1) \\
&\quad + 2 \sum_{v \in V(H)} (n_1 - 1 - d_1(v|H) + n) \\
&= 2(n + n_1 - 1) + (3n + 4n_1 - 2)(n - 1) + 2(n + n_1 - 1)n_1 - 4m_1 \\
&= (3n + 4n_1 - 2)(n - 1) + 2(n + n_1 - 1)(n_1 + 1) - 4m_1
\end{aligned}$$

The proof is completed. \square

Theorem 3.9. *Let G and H be two disjoint connected graphs with vertex numbers $n \geq 3$ and n_1 , and edge numbers m and m_1 , and $G \not\cong S_n$, then*

$$L\xi^C(G \vee H) = m(3m + 5) + 2n_1(n_1 + 3n - 1) - 4m_1 - (n + n_1 - 1)|V_e^1(G)|.$$

Proof. According to Lemma 3.7, for any vertex $v \in V(G)$, when $e(v|G) = 1$, $e(v|G \vee H) = 3$ holds, that is, when $v \in V_e^1(G)$, $e(v|G \vee H) = 3$, and $e(v|G \vee H) = 4$ holds for other original non-full vertices in G . Then,

$$\begin{aligned}
L\xi^C(G \vee H) &= \sum_{v \in V(G \vee H)} d_2(v|G \vee H)e(v|G \vee H) \\
&= 3 \sum_{v \in V_e^1(G)} (d_1(v|G) + n_1) + 4 \sum_{v \notin V_e^1(G)} (d_1(v|G) + n_1) + 3 \sum_{e \in E(G)} (m - 1) \\
&\quad + 2 \sum_{v \in V(H)} (n_1 - 1 - d_1(v|H) + n) \\
&= 4 \sum_{v \in V(G)} (d_1(v|G) + n_1) - \sum_{v \in V_e^1(G)} (d_1(v|G) + n_1) + 3m(m - 1) \\
&\quad + 2(n_1 - 1 + n)n_1 - 4m_1 \\
&= m(3m + 5) + 2n_1(n_1 + 3n - 1) - 4m_1 - (n + n_1 - 1)|V_e^1(G)|
\end{aligned}$$

The proof is completed. \square

3.3 Subdivision vertex edge join

Definition 3.10. Let G , H_1 and H_2 be three disjoint graphs with vertices n , n_1 and n_2 , and edges m , m_1 and m_2 respectively. The subdivision vertex edge join of graphs G , H_1 and H_2 is denoted as $G^S \Delta (H_1^V \vee H_2^S)$, and is the graph obtained by connecting the i -th ($i = 1, 2, \dots, n$) vertex in original G in $S(G)$ and all vertices of H_1 and the newly inserted j -th ($j = 1, 2, \dots, m$) vertex corresponding to each edge of G in $S(G)$ and all vertices of H_2 .

We can see that the number of vertices of $G^S \Delta (H_1^V \vee H_2^S)$ is $n + m + n_1 + n_2$ and the number of edges is $2m + nn_1 + mn_2 + m_1 + m_2$.

Lemma 3.11. Let G , H_1 and H_2 be the disjoint three connected graphs with vertex numbers $n \geq 3$, n_1 and n_2 , and edge numbers m , m_1 and m_2 , then for any vertex $v \in V(G^S \Delta (H_1^V \vee H_2^S))$, we have

(i) If graph $G \cong S_n$, then

$$e(v|G^S \Delta (H_1^V \vee H_2^S)) = \begin{cases} 2 \text{ or } 3, & \text{if } v \in V(G) \\ 3, & \text{if } e \in E(G) \text{ or } v \in V(H_i) \end{cases}$$

(ii) If graph $G \not\cong S_n$, then

$$e(v|G^S \Delta (H_1^V \vee H_2^S)) = 3, v \in V(G^S \Delta (H_1^V \vee H_2^S)).$$

(iii)

$$d_2(v|G^S \Delta (H_1^V \vee H_2^S)) = \begin{cases} n - 1 + n_2, & \text{if } v \in V(G) \\ m - 1 + n_1, & \text{if } e \in E(G) \\ n_1 - 1 - d_1(v|H_1) + m, & \text{if } v \in V(H_1) \\ n_2 - 1 - d_1(v|H_2) + n, & \text{if } v \in V(H_2) \end{cases}$$

Theorem 3.12. Let G , H_1 and H_2 be the disjoint three connected graphs with vertex numbers $n \geq 3$, n_1 and n_2 , and edge numbers m , m_1 and m_2 , and $G \cong S_n$, then

$$L\xi^C(G^S \Delta (H_1^V \vee H_2^S)) = 3(n + n_1 - 2)(n + n_1 - 1) + (n + n_2 - 1)(3n + 3n_2 - 1) - 6(m_1 + m_2)$$

Proof. When the graph $G \cong S_n = K_{1, n-1}$, we know that there is $e(v|G^S \Delta (H_1^V \vee H_2^S)) = 2$ for the original center vertex in G , and $e(v|G^S \Delta (H_1^V \vee H_2^S)) = 3$ for the other vertices in G , and at this time $m = n - 1$, the line graph of the star graph is complete graph K_{n-1} .

Then,

$$\begin{aligned}
L\xi^C(G^S \Delta (H_1^V \vee H_2^S)) &= \sum_{v \in V(G^S \Delta (H_1^V \vee H_2^S))} d_2(v|G^S \Delta (H_1^V \vee H_2^S))e(v|G^S \Delta (H_1^V \vee H_2^S)) \\
&= 2(n-1+n_2) + 3(n-1+n_2)(n-1) + 3 \sum_{e \in E(G)} (m-1+n_1) \\
&\quad + 3 \sum_{v \in V(H_1)} (n_1-1-d_1(v|H_1)+m) \\
&\quad + 3 \sum_{v \in V(H_2)} (n_2-1-d_1(v|H_2)+n) \\
&= 3(n-1+n_2)n - (n-1+n_2) + 3(n-2+n_1)(n-1) \\
&\quad + 3(n_1-2+n)n_1 - 6m_1 + 3(n_2-1+n)n_2 - 6m_2 \\
&= 3(n+n_1-2)(n+n_1-1) + (n+n_2-1)(3n+3n_2-1) \\
&\quad - 6(m_1+m_2)
\end{aligned}$$

The proof is completed. \square

Theorem 3.13. *Let G , H_1 and H_2 be the disjoint three connected graphs with vertex numbers $n \geq 3$, n_1 and n_2 , and edge numbers m , m_1 and m_2 , and $G \not\cong S_n$, then*

$$L\xi^C(G^S \Delta (H_1^V \vee H_2^S)) = 3(n+n_2-1)(n+n_2) + 3(m+n_1-1)(m+n_1) - 6(m_1+m_2)$$

Proof. According to Lemma 3.11, we have

$$\begin{aligned}
L\xi^C(G^S \Delta (H_1^V \vee H_2^S)) &= \sum_{v \in V(G^S \Delta (H_1^V \vee H_2^S))} d_2(v|G^S \Delta (H_1^V \vee H_2^S))e(v|G^S \Delta (H_1^V \vee H_2^S)) \\
&= 3 \sum_{v \in V(G)} (n-1+n_2) + 3 \sum_{e \in E(G)} (m-1+n_1) \\
&\quad + 3 \sum_{v \in V(H_1)} (n_1-1-d_1(v|H_1)+m) \\
&\quad + 3 \sum_{v \in V(H_2)} (n_2-1-d_1(v|H_2)+n) \\
&= 3(n-1+n_2)n + 3(m-1+n_1)m + 3(n_1-1+m)n_1 \\
&\quad - 6m_1 + 3(n_2-1+n)n_2 - 6m_2 \\
&= 3(n+n_2-1)(n+n_2) + 3(m+n_1-1)(m+n_1) - 6(m_1+m_2)
\end{aligned}$$

The proof is completed. \square

4 Leap eccentric connectivity index of four double corona graphs

In this section, we give the bounds of the leap eccentric connectivity index of four variants of the corona graph. First, an obvious conclusion is introduced.

Lemma 4.1. *Let G be a connected graph, then*

(i)

$$d_1(v|S(G)) = \begin{cases} d_1(v|G), & \text{if } v \in V(G) \\ 2, & \text{if } e \in E(G) \end{cases}$$

(ii)

$$d_1(v|Q(G)) = \begin{cases} d_1(v|G), & \text{if } v \in V(G) \\ 2 + d_1(e|L(G)), & \text{if } e \in E(G) \end{cases}$$

(iii)

$$d_1(v|R(G)) = \begin{cases} 2d_1(v|G), & \text{if } v \in V(G) \\ 2, & \text{if } e \in E(G) \end{cases}$$

(iv)

$$d_1(v|T(G)) = \begin{cases} 2d_1(v|G), & \text{if } v \in V(G) \\ 2 + d_1(e|L(G)), & \text{if } e \in E(G) \end{cases}$$

4.1 Subdivision double corona

Definition 4.2. *Let G be a connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with n_1 and n_2 vertices, respectively. The subdivision double corona graph composed of G , H_1 and H_2 is denoted as $G^{(S)} \circ \{H_1, H_2\}$, and is a graph obtained by taking one copy of $S(G)$, n_1 copies of H_1 , m copies of H_2 , then connecting the i^{th} vertex in original G in $S(G)$ to every vertex in the i^{th} copy of H_1 and the newly inserted j^{th} -th vertex corresponding to each edge of G in $S(G)$ to every vertex in the j^{th} copy of H_2 .*

We use H_1^i to represent the copied i -th H_1 , where $i = 1, 2, \dots, n$, H_2^j to represent the copied j -th H_2 , where $j = 1, 2, \dots, m$. Obviously, the number of vertices of the subdivision double corona graph $G^{(S)} \circ \{H_1, H_2\}$ is $n(n_1 + 1) + m(n_2 + 1)$.

Lemma 4.3. *Let G be a connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , respectively. Then*

(i)

$$e(v|G^{(S)} \circ \{H_1, H_2\}) = \begin{cases} e(v|S(G)) + 1, & \text{if } v \in V(G) \\ e(v|S(G)) + 1, & \text{if } e \in E(G) \\ e(v_i|S(G)) + 2, & \text{if } v \in V(H_1^i) \\ e(e_j|S(G)) + 2, & \text{if } v \in V(H_2^j) \end{cases}$$

(ii)

$$d_2(v|G^{(S)} \circ \{H_1, H_2\}) = \begin{cases} (n_2 + 1)d_1(v|G), & \text{if } v \in V(G) \\ 2n_1 + d_1(e|L(G)), & \text{if } e \in E(G) \\ (n_1 - 1) - d_1(v|H_1) + d_1(v_i|G), & \text{if } v \in V(H_1^i) \\ (n_2 - 1) - d_1(v|H_2) + 2, & \text{if } v \in V(H_2^j) \end{cases}$$

Theorem 4.4. *Let G be a connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , and edges m_1 and m_2 , respectively. Then*

(i)

$$\begin{aligned} L\xi^C(G^{(S)} \circ \{H_1, H_2\}) &\geq 2(n_1 + n_2 + 1)\xi^C(G) + 2(n_1^2 - n_1 - 2m_1)\theta(G) + 2\xi^C(L(G)) \\ &\quad + 2(2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 2|E(L(G))| \\ &\quad + 2m(3n_1 + n_2^2 + 2n_2 - 2m_2 + 1) + 2n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

(ii)

$$\begin{aligned} L\xi^C(G^{(S)} \circ \{H_1, H_2\}) &\leq 2(n_1 + n_2 + 1)\xi^C(G) + 2(n_1^2 - n_1 - 2m_1)\theta(G) + 2\xi^C(L(G)) \\ &\quad + 2(2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 4|E(L(G))| \\ &\quad + m(10n_1 + 3n_2^2 + 7n_2 - 6m_2 + 4) + 3n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

Proof. According to Lemma 4.3, we can get

$$\begin{aligned} L\xi^C(G^{(S)} \circ \{H_1, H_2\}) &= \sum_{v \in V(G^{(S)} \circ \{H_1, H_2\})} d_2(v|G^{(S)} \circ \{H_1, H_2\})e(v|G^{(S)} \circ \{H_1, H_2\}) \\ &= \sum_{v \in V(G)} (n_2 + 1)d_1(v|G)(e(v|S(G)) + 1) \\ &\quad + \sum_{e \in E(G)} (2n_1 + d_1(e|L(G)))(e(e|S(G)) + 1) \\ &\quad + \sum_{i=1}^n \sum_{v \in V(H_1^i)} (n_1 - 1 - d_1(v|H_1) + d_1(v_i|G))(e(v_i|S(G)) + 2) \\ &\quad + \sum_{j=1}^m \sum_{v \in V(H_2^j)} (n_2 - 1 - d_1(v|H_2) + 2)(e(e_j|S(G)) + 2) \end{aligned}$$

And according to Lemma 2.1, we know that for any vertex $v \in V(G)$, $2e(v|G) \leq e(v|S(G)) \leq 2e(v|G) + 1$; for any vertex $e \in E(G)$, $2e(e|L(G)) \leq e(e|S(G)) \leq 2e(e|L(G)) + 1$. So we can get the upper and lower bounds of the leap eccentric connectivity index of

the subdivision double corona graph $G^{(S)} \circ \{H_1, H_2\}$.

$$\begin{aligned}
L\xi^C(G^{(S)} \circ \{H_1, H_2\}) &\geq 2(n_2 + 1)\xi^C(G) + 2(n_2 + 1)m + 4n_1\theta(L(G)) + 2n_1m \\
&\quad + 2\xi^C(L(G)) + 2|E(L(G))| + 2n_1(n_1 - 1)\theta(G) + 2n_1(n_1 - 1)n \\
&\quad + 2n_1\xi^C(G) + 4n_1m - 4m_1\theta(G) - 4m_1n + 2n_2(n_2 + 1)\theta(L(G)) \\
&\quad + 2n_2(n_2 + 1)m - 4m_2\theta(L(G)) - 4m_2m \\
&= 2(n_1 + n_2 + 1)\xi^C(G) + 2(n_1^2 - n_1 - 2m_1)\theta(G) + 2\xi^C(L(G)) \\
&\quad + 2(2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 2|E(L(G))| \\
&\quad + 2m(3n_1 + n_2^2 + 2n_2 - 2m_2 + 1) + 2n(n_1^2 - n_1 - 2m_1)
\end{aligned}$$

Similarly, we can get the upper bound

$$\begin{aligned}
L\xi^C(G^{(S)} \circ \{H_1, H_2\}) &\leq 2(n_1 + n_2 + 1)\xi^C(G) + 2(n_1^2 - n_1 - 2m_1)\theta(G) + 2\xi^C(L(G)) \\
&\quad + 2(2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 4|E(L(G))| \\
&\quad + m(10n_1 + 3n_2^2 + 7n_2 - 6m_2 + 4) + 3n(n_1^2 - n_1 - 2m_1)
\end{aligned}$$

The proof is completed. \square

For a connected bipartite graph, $e(v|S(G)) = 2e(v|G)$ is always true, so we can get the following conclusion.

Corollary 4.5. *Let G be a connected bipartite graph with n vertices and m edges. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , and edges m_1 and m_2 , respectively. Then*

(i)

$$\begin{aligned}
L\xi^C(G^{(S)} \circ \{H_1, H_2\}) &\geq 2(n_1 + n_2 + 1)\xi^C(G) + 2(n_1^2 - n_1 - 2m_1)\theta(G) + 2\xi^C(L(G)) \\
&\quad + 2(2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 2|E(L(G))| \\
&\quad + 2m(3n_1 + n_2^2 + 2n_2 - 2m_2 + 1) + 2n(n_1^2 - n_1 - 2m_1)
\end{aligned}$$

(ii)

$$\begin{aligned}
L\xi^C(G^{(S)} \circ \{H_1, H_2\}) &\leq 2(n_1 + n_2 + 1)\xi^C(G) + 2(n_1^2 - n_1 - 2m_1)\theta(G) + 2\xi^C(L(G)) \\
&\quad + 2(2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 4|E(L(G))| \\
&\quad + m(8n_1 + 3n_2^2 + 5n_2 - 6m_2 + 2) + 2n(n_1^2 - n_1 - 2m_1)
\end{aligned}$$

When the eccentricity of all vertices in graph G is reached at the pendant vertex, $e(e|S(G)) = 2e(e|L(G)) + 1$ holds for all vertices. Then we can get the following equation for the connected bipartite graph and the eccentricity of all vertices is reached at the pendant vertex.

Corollary 4.6. *Let G be a connected bipartite graph and the eccentricity of all vertices are reached at the pendant vertex, where the number of vertices of G is n and the number of edges is m . Let H_1 and H_2 be graphs with vertices n_1 and n_2 , and edges m_1 and m_2 , respectively. Then*

$$\begin{aligned} L\xi^C(G^{(S)} \circ \{H_1, H_2\}) &= 2(n_1 + n_2 + 1)\xi^C(G) + 2(n_1^2 - n_1 - 2m_1)\theta(G) + 2\xi^C(L(G)) \\ &\quad + 2(2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 4|E(L(G))| \\ &\quad + m(8n_1 + 3n_2^2 + 5n_2 - 6m_2 + 2) + 2n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

4.2 $Q(G)$ graph double corona

Definition 4.7. *Let G be a connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with n_1 and n_2 vertices, respectively. The $Q(G)$ graph double corona composed of G , H_1 and H_2 is denoted as $G^{(Q)} \circ \{H_1, H_2\}$, and is a graph obtained by taking one copy of $Q(G)$, n_1 copies of H_1 , m copies of H_2 , then connecting the i^{th} vertex in original G in $Q(G)$ to every vertex in the i^{th} copy of H_1 and the newly inserted j^{th} -th vertex corresponding to each edge of G in $Q(G)$ to every vertex in the j^{th} copy of H_2 .*

We use H_1^i to represent the copied i -th H_1 , where $i = 1, 2, \dots, n$, H_2^j to represent the copied j -th H_2 , where $j = 1, 2, \dots, m$. Obviously, the number of vertices of the $Q(G)$ graph double corona $G^{(Q)} \circ \{H_1, H_2\}$ is $n(n_1 + 1) + m(n_2 + 1)$.

Lemma 4.8. *Let G be a connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , respectively. Then*

(i)

$$e(v|G^{(Q)} \circ \{H_1, H_2\}) = \begin{cases} e(v|Q(G)) + 1, & \text{if } v \in V(G) \\ e(v|Q(G)) + 1, & \text{if } e \in E(G) \\ e(v_i|Q(G)) + 2, & \text{if } v \in V(H_1^i) \\ e(e_j|Q(G)) + 2, & \text{if } v \in V(H_2^j) \end{cases}$$

(ii)

$$d_2(v|G^{(Q)} \circ \{H_1, H_2\}) = \begin{cases} d_2(v|Q(G)) + n_2d_1(v|G), & \text{if } v \in V(G) \\ d_2(e|Q(G)) + 2n_1 + n_2d_1(e|L(G)), & \text{if } e \in E(G) \\ (n_1 - 1) - d_1(v|H_1) + d_1(v_i|G), & \text{if } v \in V(H_1^i) \\ (n_2 - 1) - d_1(v|H_2) + 2 + d_1(e_j|L(G)), & \text{if } v \in V(H_2^j) \end{cases}$$

Theorem 4.9. *Let G be a $\{C_3, C_4\}$ -free connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , and edges m_1 and m_2 , respectively. Then*

(i)

$$\begin{aligned} L\xi^C(G^{(Q)} \circ \{H_1, H_2\}) &\geq L\xi^C(G) + (n_1 + n_2 + 1)\xi^C(G) + (n_1^2 - n_1 - 2m_1)\theta(G) \\ &\quad + 2M_1(G) + L\xi^C(L(G)) + (2n_2 + 1)\xi^C(L(G)) \\ &\quad + (2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + M_1(L(G)) + 6n_2|E(L(G))| \\ &\quad + 2m(4n_1 + n_2^2 + 3n_2 - 2m_2) + 3n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

(ii)

$$\begin{aligned} L\xi^C(G^{(Q)} \circ \{H_1, H_2\}) &\leq L\xi^C(G) + (n_1 + n_2 + 1)\xi^C(G) + (n_1^2 - n_1 - 2m_1)\theta(G) \\ &\quad + 2M_1(G) + L\xi^C(L(G)) + (2n_2 + 1)\xi^C(L(G)) \\ &\quad + (2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 2M_1(L(G)) + 10n_2|E(L(G))| \\ &\quad + m(10n_1 + 3n_2^2 + 7n_2 - 6m_2) + 3n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

Proof. When G is a $\{C_3, C_4\}$ -free connected graph, according to Lemma 2.9, we know that for any vertex $v \in V(G)$, $d_2(v|Q(G)) = d_1(v|G) + d_2(v|G)$, for any vertex $e \in E(G)$, $d_2(e|Q(G)) = d_1(e|L(G)) + d_2(e|L(G))$. And according to Lemma 2.3, we have when G is a $\{C_3, C_4\}$ -free connected graph,

$$\sum_{v \in V(G)} d_2(v|G) = M_1(G) - 2m.$$

In Lemma 2.8, we know that for any vertex $v \in V(G)$, $e(v|Q(G)) = e(v|G) + 1$, for any vertex $e \in E(G)$, $e(e|L(G)) \leq e(e|Q(G)) \leq e(e|L(G)) + 1$.

According to the above lemma and Lemma 4.8 and the definition of the leap eccentric connectivity index, we can calculate the upper and lower bounds of $L\xi^C(G^{(Q)} \circ \{H_1, H_2\})$.

The proof is completed. \square

When G is a $\{C_3, C_4\}$ -free connected bipartite graph and the eccentricity of all vertices in graph G is reached at the pendant vertex, the equal sign of the second inequality in Theorem 4.9 can be equal, that is, for any vertex $e \in E(G)$, $e(e|Q(G)) = e(e|L(G)) + 1$ established. So we can get the following corollary.

Corollary 4.10. *Let G is a $\{C_3, C_4\}$ -free connected bipartite graph where the eccentricity of all vertices in graph G is reached at the pendant vertex with n vertices and m edges. Let*

H_1 and H_2 be graphs with vertices n_1 and n_2 , and edges m_1 and m_2 , respectively. Then

$$\begin{aligned} L\xi^C(G^{(Q)} \circ \{H_1, H_2\}) &= L\xi^C(G) + (n_1 + n_2 + 1)\xi^C(G) + (n_1^2 - n_1 - 2m_1)\theta(G) \\ &\quad + 2M_1(G) + L\xi^C(L(G)) + (2n_2 + 1)\xi^C(L(G)) \\ &\quad + (2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 2M_1(L(G)) + 10n_2|E(L(G))| \\ &\quad + m(10n_1 + 3n_2^2 + 7n_2 - 6m_2) + 3n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

4.3 $R(G)$ graph double corona

Definition 4.11. Let G be a connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with n_1 and n_2 vertices, respectively. The $R(G)$ graph double corona composed of G , H_1 and H_2 is denoted as $G^{(R)} \circ \{H_1, H_2\}$, and is a graph obtained by taking one copy of $R(G)$, n_1 copies of H_1 , m copies of H_2 , then connecting the i^{th} vertex in original G in $R(G)$ to every vertex in the i^{th} copy of H_1 and the newly inserted j^{th} -th vertex corresponding to each edge of G in $R(G)$ to every vertex in the j^{th} copy of H_2 .

We use H_1^i to represent the copied i -th H_1 , where $i = 1, 2, \dots, n$, H_2^j to represent the copied j -th H_2 , where $j = 1, 2, \dots, m$. Obviously, the number of vertices of the $R(G)$ graph double corona $G^{(R)} \circ \{H_1, H_2\}$ is $n(n_1 + 1) + m(n_2 + 1)$.

Lemma 4.12. Let G be a connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , respectively. Then

(i)

$$e(v|G^{(R)} \circ \{H_1, H_2\}) = \begin{cases} e(v|R(G)) + 1, & \text{if } v \in V(G) \\ e(v|R(G)) + 1, & \text{if } e \in E(G) \\ e(v_i|R(G)) + 2, & \text{if } v \in V(H_1^i) \\ e(e_j|R(G)) + 2, & \text{if } v \in V(H_2^j) \end{cases}$$

(ii)

$$d_2(v|G^{(R)} \circ \{H_1, H_2\}) = \begin{cases} d_2(v|R(G)) + n_2d_1(v|G) + n_1d_1(v|G), & \text{if } v \in V(G) \\ d_2(e|R(G)) + 2n_1, & \text{if } e \in E(G) \\ (n_1 - 1) - d_1(v|H_1) + 2d_1(v_i|G), & \text{if } v \in V(H_1^i) \\ (n_2 - 1) - d_1(v|H_2) + 2, & \text{if } v \in V(H_2^j) \end{cases}$$

Theorem 4.13. Let G be a $\{C_3, C_4\}$ -free connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , and edges m_1 and m_2 , respectively. Then

(i)

$$\begin{aligned} L\xi^C(G^{(R)} \circ \{H_1, H_2\}) &\geq 2L\xi^C(G) + (3n_1 + n_2)\xi^C(G) + (n_1^2 - n_1 - 2m_1)\theta(G) + 2M_1(G) \\ &\quad + 2\xi^C(L(G)) + (2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 8|E(L(G))| \\ &\quad + m(14n_1 + 3n_2^2 + 5n_2 - 6m_2 - 4) + 2n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

(ii)

$$\begin{aligned} L\xi^C(G^{(R)} \circ \{H_1, H_2\}) &\leq 2L\xi^C(G) + (3n_1 + n_2)\xi^C(G) + (n_1^2 - n_1 - 2m_1)\theta(G) + 4M_1(G) \\ &\quad + 2\xi^C(L(G)) + (2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 8|E(L(G))| \\ &\quad + m(20n_1 + 3n_2^2 + 7n_2 - 6m_2 - 8) + 3n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

Proof. When G is a $\{C_3, C_4\}$ -free connected graph, according to Lemma 2.13, we know that for any vertex $v \in V(G)$, $d_2(v|R(G)) = 2d_2(v|G)$, for any vertex $e \in E(G)$, $d_2(e|R(G)) = 2d_1(e|L(G))$. And according to Lemma 2.3, we have when G is a $\{C_3, C_4\}$ -free connected graph,

$$\sum_{v \in V(G)} d_2(v|G) = M_1(G) - 2m.$$

In Lemma 2.12, we know that for any vertex $v \in V(G)$, $e(v|G) \leq e(v|R(G)) \leq e(v|G) + 1$, for any vertex $e \in E(G)$, $e(e|R(G)) = e(e|L(G)) + 1$.

According to the above lemma and Lemma 4.12 and the definition of the leap eccentric connectivity index, we can calculate the upper and lower bounds of $L\xi^C(G^{(R)} \circ \{H_1, H_2\})$.

The proof is completed. \square

When G is a $\{C_3, C_4\}$ -free connected graph with an odd circle, the equal sign on the right side of Theorem 4.13 is equal, which means that for any vertex $v \in V(G)$, $e(v|R(G)) = e(v|G) + 1$ holds. When G is a $\{C_3, C_4\}$ -free connected bipartite graph, then for all vertices $v \in V(G)$, $e(v|R(G)) = e(v|G)$ holds. So we can get the following corollary.

Corollary 4.14. *Let G is a $\{C_3, C_4\}$ -free connected bipartite graph with n vertices and m edges. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , and edges m_1 and m_2 , respectively. Then*

$$\begin{aligned} L\xi^C(G^{(R)} \circ \{H_1, H_2\}) &= 2L\xi^C(G) + (3n_1 + n_2)\xi^C(G) + (n_1^2 - n_1 - 2m_1)\theta(G) + 2M_1(G) \\ &\quad + 2\xi^C(L(G)) + (2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 8|E(L(G))| \\ &\quad + m(14n_1 + 3n_2^2 + 5n_2 - 6m_2 - 4) + 2n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

4.4 Total graph double corona

Definition 4.15. Let G be a connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with n_1 and n_2 vertices, respectively. The $T(G)$ graph double corona composed of G , H_1 and H_2 is denoted as $G^{(T)} \circ \{H_1, H_2\}$, and is a graph obtained by taking one copy of $T(G)$, n_1 copies of H_1 , m copies of H_2 , then connecting the i^{th} vertex in original G in $T(G)$ to every vertex in the i^{th} copy of H_1 and the newly inserted j^{th} -th vertex corresponding to each edge of G in $T(G)$ to every vertex in the j^{th} copy of H_2 .

We use H_1^i to represent the copied i -th H_1 , where $i = 1, 2, \dots, n$, H_2^j to represent the copied j -th H_2 , where $j = 1, 2, \dots, m$. Obviously, the number of vertices of the $T(G)$ graph double corona $G^{(T)} \circ \{H_1, H_2\}$ is $n(n_1 + 1) + m(n_2 + 1)$.

Lemma 4.16. Let G be a connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , respectively. Then

(i)

$$e(v|G^{(T)} \circ \{H_1, H_2\}) = \begin{cases} e(v|T(G)) + 1, & \text{if } v \in V(G) \\ e(v|T(G)) + 1, & \text{if } e \in E(G) \\ e(v_i|T(G)) + 2, & \text{if } v \in V(H_1^i) \\ e(e_j|T(G)) + 2, & \text{if } v \in V(H_2^j) \end{cases}$$

(ii)

$$d_2(v|G^{(T)} \circ \{H_1, H_2\}) = \begin{cases} d_2(v|T(G)) + n_2 d_1(v|G) + n_1 d_1(v|G), & \text{if } v \in V(G) \\ d_2(e|T(G)) + 2n_1 + n_2 d_1(e|L(G)), & \text{if } e \in E(G) \\ (n_1 - 1) - d_1(v|H_1) + 2d_1(v_i|G), & \text{if } v \in V(H_1^i) \\ (n_2 - 1) - d_1(v|H_2) + 2 + d_1(e_j|L(G)), & \text{if } v \in V(H_2^j) \end{cases}$$

Theorem 4.17. Let G be a $\{C_3, C_4\}$ -free connected graph with n vertices and m edges. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , and edges m_1 and m_2 , respectively. Then

(i)

$$\begin{aligned} L\xi^C(G^{(T)} \circ \{H_1, H_2\}) &\geq 2L\xi^C(G) + (3n_1 + n_2)\xi^C(G) + 2M_1(G) + (n_1^2 - n_1 - 2m_1)\theta(G) \\ &\quad + L\xi^C(L(G)) + (2n_2 + 1)\xi^C(L(G)) + M_1(L(G)) \\ &\quad + (2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 6n_2|E(L(G))| \\ &\quad + 2m(6n_1 + n_2^2 + 2n_2 - 2m_2 - 2) + 2n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

(ii)

$$\begin{aligned}
L\xi^C(G^{(T)} \circ \{H_1, H_2\}) &\leq 2L\xi^C(G) + (3n_1 + n_2)\xi^C(G) + 4M_1(G) + (n_1^2 - n_1 - 2m_1)\theta(G) \\
&\quad + L\xi^C(L(G)) + (2n_2 + 1)\xi^C(L(G)) + 2M_1(L(G)) \\
&\quad + (2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 10n_2|E(L(G))| \\
&\quad + m(20n_1 + 3n_2^2 + 7n_2 - 6m_2 - 8) + 3n(n_1^2 - n_1 - 2m_1)
\end{aligned}$$

Proof. When G is a $\{C_3, C_4\}$ -free connected graph, according to Lemma 2.17, we know that for any vertex $v \in V(G)$, $d_2(v|T(G)) = 2d_2(v|G)$, for any vertex $e \in E(G)$, $d_2(e|T(G)) = d_1(e|L(G)) + d_2(e|L(G))$. And according to Lemma 2.3, we have when G is a $\{C_3, C_4\}$ -free connected graph,

$$\sum_{v \in V(G)} d_2(v|G) = M_1(G) - 2m.$$

In Lemma 2.16, we know that for any vertex $v \in V(G)$, $e(v|G) \leq e(v|T(G)) \leq e(v|G) + 1$, for any vertex $e \in E(G)$, $e(e|L(G)) \leq e(e|R(G)) \leq e(e|L(G)) + 1$.

According to the above lemma and Lemma 4.16 and the definition of the leap eccentric connectivity index, we can calculate the upper and lower bounds of $L\xi^C(G^{(T)} \circ \{H_1, H_2\})$.

The proof is completed. \square

We find that when the graph G is a $\{C_3, C_4\}$ -free connected graph that does not contain odd cycles, $e(v|T(G)) = e(v|G)$ holds for any vertex v in G . So for a $\{C_3, C_4\}$ -free connected bipartite graph, $e(v|T(G)) = e(v|G)$ is always true, so we can get the following conclusion.

Corollary 4.18. *Let G is a $\{C_3, C_4\}$ -free connected bipartite graph with n vertices and m edges. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , and edges m_1 and m_2 , respectively. Then*

(i)

$$\begin{aligned}
L\xi^C(G^{(T)} \circ \{H_1, H_2\}) &\geq 2L\xi^C(G) + (3n_1 + n_2)\xi^C(G) + 2M_1(G) + (n_1^2 - n_1 - 2m_1)\theta(G) \\
&\quad + L\xi^C(L(G)) + (2n_2 + 1)\xi^C(L(G)) + M_1(L(G)) \\
&\quad + (2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 6n_2|E(L(G))| \\
&\quad + 2m(6n_1 + n_2^2 + 2n_2 - 2m_2 - 2) + 2n(n_1^2 - n_1 - 2m_1)
\end{aligned}$$

(ii)

$$\begin{aligned} L\xi^C(G^{(T)} \circ \{H_1, H_2\}) &\leq 2L\xi^C(G) + (3n_1 + n_2)\xi^C(G) + 2M_1(G) + (n_1^2 - n_1 - 2m_1)\theta(G) \\ &\quad + L\xi^C(L(G)) + (2n_2 + 1)\xi^C(L(G)) + 2M_1(L(G)) \\ &\quad + (2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 10n_2|E(L(G))| \\ &\quad + m(14n_1 + 3n_2^2 + 5n_2 - 6m_2 - 4) + 2n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

When graph G is a $\{C_3, C_4\}$ -free connected graph where the eccentricity of all vertices is reached at the pendant vertex, then $e(e|T(G)) = e(e|L(G)) + 1$ holds for all vertices $e \in E(G)$. So we can get the following equation hold.

Corollary 4.19. *Let G is a $\{C_3, C_4\}$ -free connected bipartite graph with n vertices and m edges, and the eccentricity of all vertices is reached at the pendant vertex. Let H_1 and H_2 be graphs with vertices n_1 and n_2 , and edges m_1 and m_2 , respectively. Then*

$$\begin{aligned} L\xi^C(G^{(T)} \circ \{H_1, H_2\}) &= 2L\xi^C(G) + (3n_1 + n_2)\xi^C(G) + 2M_1(G) + (n_1^2 - n_1 - 2m_1)\theta(G) \\ &\quad + L\xi^C(L(G)) + (2n_2 + 1)\xi^C(L(G)) + 2M_1(L(G)) \\ &\quad + (2n_1 + n_2^2 + n_2 - 2m_2)\theta(L(G)) + 10n_2|E(L(G))| \\ &\quad + m(14n_1 + 3n_2^2 + 5n_2 - 6m_2 - 4) + 2n(n_1^2 - n_1 - 2m_1) \end{aligned}$$

References

- [1] Barik S, Sahoo G, On the Laplacian spectra of some variants of corona, Linear Algebra and its Applications, 512(2017), 32-47.
- [2] Bondy J A, Murty U S R, "Graph theory(Graduate Texts in Mathematices)", Belin, Springer, 2008.
- [3] Cvetkovic D M, Doob M, Sachs H, "Spectra of graphs", Academic Press, New York, 1980.
- [4] Dankelmann P, Goddard W, Swart C S, The average eccentricity of a graph and its subgraphs, Utilitas Mathematica, 65(2004), 41-52.
- [5] Gutman I, Naji A M, Soner N D, On leap Zagreb indices of graphs, Communication in Combinatorics and Optimization, 2(2017), 99-117.
- [6] Gutman I, Rušćić B, Trinajstić N, et al, Graph theory and molecular orbitals. XII. Acyclic polyenes, The Journal of Chemical Physics, 62(1975), 3399-3405.
- [7] Gutman I, Trinajstić N, Graph theory and molecular orbitals. Total φ -electron energy of alternant hydrocarbons, Chemical Physics Letters. 17(1972), 535-538.

- [8] Indulal G. Spectrum of two new joins of graphs and infinite families of integral graphs, *Kragujevac journal of mathematics*, 36(2012), 133-139.
- [9] Manjunathe H R, Naji A M, Shiladhar P, et al. Leap eccentric connectivity index of some graph operations. *Int. J. Res. Anal. Reviews*, 6(2019), 882-887.
- [10] Naji A. M., Shiladhar P. and Soner N. D., On leap eccentric connectivity index of graphs, *Iranian J. Math. Sci. Inf. commun.*, in communication.
- [11] Sharma V, Goswami R, Madan A K, Eccentric connectivity index: a novel highly discriminating topological descriptor for structure- property and structure- activity studies, *Journal of chemical information and computer sciences*, 37(1997), 273-282.
- [12] Soner N D, Naji A M, The k-distance neighborhood polynomial of a graph//*Int. J. Math. Comput. Sci. WASET Conference Proceedings*, San Francico, USA, Sep. 2016, 26-27.
- [13] Wen F, Zhang Y, Li M. Spectra of subdivision vertex-edge join of three graphs, *Mathematics*, 7(2019), 171.
- [14] Yan W, Yang B Y, Yeh Y N, The behavior of Wiener indices and polynomials of graphs under five graph decorations, *Applied mathematics letters*, 20(2007), 290-295.
- [15] Yarahmadi Z, Moradi S, Doslic T, Eccentric connectivity index of graphs with subdivided edges, *Electronic Notes in Discrete Mathematics*, 45(2014), 167-176.