

On the first Banhatti-Sombor index ^{*}

Zhen Lin^{a†}, Ting Zhou^b, V.R. Kulli^c, Lianying Miao^b

^aSchool of Mathematics and Statistics, Beijing Key Laboratory on MCAACL,

Beijing Institute of Technology,

Beijing, 102488, P.R. China

^bSchool of Mathematics, China University of Mining and Technology,

Xuzhou, 221116, Jiangsu, P.R. China

^cDepartment of Mathematics, Gulbarga University,

Kalaburgi (Gulbarga)-585106, India

Abstract

Let d_v be the degree of the vertex v in a connected graph G . The first Banhatti-Sombor index of G is defined as $BSO(G) = \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}}$, which is a new vertex-degree-based topological index introduced by Kulli. In this paper, the mathematical relations between the first Banhatti-Sombor index and some other well-known vertex-degree-based topological indices are established. In addition, the trees extremal with respect to the first Banhatti-Sombor index on trees and chemical trees are characterized, respectively.

MSC Classification: 05C05, 05C07, 05C09, 05C92

Keywords: The first Banhatti-Sombor index; Degree; Tree

1 Introduction

Let G be a simple undirected connected graph with vertex set $V(G)$ and edge set $E(G)$. The number of vertices and edges of G is called order and size, respectively. Denote by \overline{G} the complement of G . For $v \in V(G)$, d_v denotes the degree of vertex v in G . The minimum and the maximum degree of G are denoted by $\delta(G)$ and $\Delta(G)$, or simply δ and Δ , respectively. A pendant vertex of G is a vertex of degree 1. A graph G is called

^{*}Supported by the National Natural Science Foundation of China (Nos. 12071411 and 11771443).

[†]Corresponding author. E-mail addresses: lnlzhen@163.com (Z. Lin), tb19080009b1@cumt.edu.cn (T. Zhou), vrkulli@gmail.com (V.R. Kulli), miaolianying@cumt.edu.cn (L. Miao).

(Δ, δ) -semiregular if $\{d_u, d_v\} = \{\Delta, \delta\}$ holds for all edges $uv \in E(G)$. Denote by K_n , C_n , P_n and $K_{1,n-1}$ the complete graph, cycle, path and star with n vertices, respectively.

The study of topological indices of various graph structures has been of interest to chemists, mathematicians, and scientists from related fields due to the fact that the topological indices play a significant role in mathematical chemistry especially in the QSPR/QSAR modeling. In 1975, the Randić index of a graph G introduced by Randić [16] is the most important and widely applied. It is defined as

$$R(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_u d_v}}.$$

The modified second Zagreb index of a graph G , introduced by Nikolić et al. [15], is defined as

$$M_2^*(G) = \sum_{uv \in E(G)} \frac{1}{d_u d_v}$$

The harmonic index and the inverse degree index of a graph G proposed by Fajtlowicz [6] are two the older vertex-degree-based topological indices. They are respectively defined as

$$H(G) = \sum_{uv \in E(G)} \frac{2}{d_u + d_v}, \quad ID(G) = \sum_{uv \in E(G)} \left(\frac{1}{d_u^2} + \frac{1}{d_v^2} \right).$$

The symmetric division deg index, the inverse sum indeg index and the geometric-arithmetic index of a graph G , introduced by Vukičević [20–22], Gašperov [22] and Furtula [21], are respectively defined as

$$SDD(G) = \sum_{uv \in E(G)} \frac{d_u^2 + d_v^2}{2d_u d_v}, \quad ISI(G) = \sum_{uv \in E(G)} \frac{d_u d_v}{d_u + d_v}, \quad GA(G) = \sum_{uv \in E(G)} \frac{2\sqrt{d_u d_v}}{d_u + d_v}.$$

The forgotten topological index, introduced by Furtula and Gutman [7], is defined as

$$F(G) = \sum_{uv \in E(G)} (d_u^2 + d_v^2)$$

In 2021, the Sombor index of a graph G is defined as

$$SO(G) = \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2},$$

which is a novel vertex-degree-based molecular structure descriptor proposed by Gutman [8]. The investigation of the Sombor index of graphs has quickly received much attention. In particular, Redžepović [18] showed that the Sombor index may be used successfully on modeling thermodynamic properties of compounds due to the fact that the Sombor index has satisfactory prediction potential in modeling entropy and enthalpy of vaporization of alkanes. Das et al. [3] and Wang et al. [23] gave the mathematical relations between the Sombor index and some other well-known vertex-degree-based topological indices. For other related results, one may refer to [1, 5, 10, 13, 14, 19] and the references therein.

Inspired by work on Sombor index, the first Banhatti-Sombor index of a connected graph G was introduced by Kulli [9] very recently and is defined as

$$BSO(G) = \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}}.$$

We find that the new index has close contact with numerous well-known vertex-degree-based topological indices. Moreover, the trees with the maximum and minimum first Banhatti-Sombor index among the set of trees with n vertices are determined, respectively. In particular, the extremal values of the first Banhatti-Sombor index for chemical trees are characterized.

2 Preliminaries

Lemma 2.1 *For any edge $uv \in E(G)$, $d_u^2 + d_v^2$ or $\frac{1}{d_u^2} + \frac{1}{d_v^2}$ is a constant if and only if G is a regular graph (when G is non-bipartite) or G is a (Δ, δ) -semiregular bipartite graph (when G is bipartite).*

Lemma 2.2 *For any positive real number a and b , we have*

$$\frac{2\sqrt{2}(a^2 + b^2 + ab)}{3(a + b)} \leq \sqrt{a^2 + b^2} \leq \frac{\sqrt{2}(a^2 + b^2)}{a + b}$$

with equality if and only if $a = b$.

Lemma 2.3 ([17]) *If $a_i > 0$, $b_i > 0$, $p > 0$, $i = 1, 2, \dots, n$, then the following inequality*

holds:

$$\sum_{k=1}^n \frac{a_k^{p+1}}{b_k^p} \geq \frac{\left(\sum_{i=1}^n a_i\right)^{p+1}}{\left(\sum_{i=1}^n b_i\right)^p}$$

with equality if and only if $\frac{a_1}{b_1} = \frac{a_2}{b_2} = \dots = \frac{a_n}{b_n}$.

Lemma 2.4 ([4]) Let a_1, a_2, \dots, a_n and b_1, b_2, \dots, b_n be real numbers such that $q \leq \frac{a_i}{b_i} \leq Q$ and $a_i \neq 0$ for $i = 1, 2, \dots, n$. Then there holds

$$\sum_{i=1}^n b_i^2 + Qq \sum_{i=1}^n a_i^2 \leq (Q + q) \sum_{i=1}^n a_i b_i$$

with equality if and only if $b_i = qa_i$ or $b_i = Qa_i$ for at least one i , $i = 1, 2, \dots, n$.

Lemma 2.5 ([2]) If $a = (a_1, a_2, \dots, a_n)$, $b = (b_1, b_2, \dots, b_n)$ are sequences of real numbers and $c = (c_1, c_2, \dots, c_n)$, $d = (d_1, d_2, \dots, d_n)$ are nonnegative, then

$$\sum_{i=1}^n d_i \sum_{i=1}^n c_i a_i^2 + \sum_{i=1}^n c_i \sum_{i=1}^n d_i b_i^2 \geq 2 \sum_{i=1}^n c_i a_i \sum_{i=1}^n d_i b_i$$

with equality if and only if $a = b = (k, k, \dots, k)$ is a constant sequence for positive c_i and d_i , $i = 1, 2, \dots, n$.

Lemma 2.6 ([12]) Let a_1, a_2, \dots, a_n and b_1, b_2, \dots, b_n be real numbers such that $a \leq a_i \leq A$ and $b \leq b_i \leq B$ for $i = 1, 2, \dots, n$. Then there holds

$$\left| \frac{1}{n} \sum_{i=1}^n a_i b_i - \frac{1}{n} \sum_{i=1}^n a_i \frac{1}{n} \sum_{i=1}^n b_i \right| \leq \frac{1}{n} \left[\frac{n}{2} \right] \left(1 - \frac{1}{n} \left[\frac{n}{2} \right] \right) (A - a)(B - b),$$

where $[x]$ denotes the integer part of x .

3 On relations between the first Bhatti-Sombor index and other degree-based indices

3.1 Bounds in terms of order, size and degree

Theorem 3.1 Let G be a connected graph of order n and size m with the minimum degree δ . Then

$$\frac{n}{\sqrt{2}} \leq BSO(G) \leq \frac{\sqrt{2}m}{\delta}$$

with equality if and only if G is a regular graph.

Proof. Note that

$$BSO(G) = \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \leq \sum_{uv \in E(G)} \sqrt{\frac{1}{\delta^2} + \frac{1}{\delta^2}} = \frac{\sqrt{2}m}{\delta}$$

with equality if and only if $d_u = \delta$ for any vertex u , that is, G is a regular graph.

By the Cauchy-Schwarz inequality, we have

$$BSO(G) = \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \geq \sum_{uv \in E(G)} \frac{1}{\sqrt{2}} \left(\frac{1}{d_u} + \frac{1}{d_v} \right) = \frac{1}{\sqrt{2}}n$$

with equality if and only if $d_u = d_v$ for any edge uv , that is, G is a regular graph.

Corollary 3.2 *Let G be a regular connected graph with n vertices. Then*

$$BSO(G) = \frac{n}{\sqrt{2}}.$$

Remark 3.3 *This implies that $BSO(G)$ dose not increase with the increase of the number of edges of G . Clearly, $BSO(K_n) = BSO(C_n)$.*

Corollary 3.4 *Let U_n be a unicyclic graph with n vertices. Then*

$$BSO(U_n) \geq \frac{n}{\sqrt{2}}$$

with equality if and only if $G \cong C_n$.

Corollary 3.5 *Let G be a connected graph of order n and size m with the maximum degree Δ and the minimum degree δ . Then*

$$\sqrt{2}n \leq BSO(G) + BSO(\overline{G}) \leq \sqrt{2} \left(\frac{m}{\delta} + \frac{n(n-1) - 2m}{2(n-1-\Delta)} \right)$$

with equality if and only if G is a regular graph.

Theorem 3.6 *Let G be a connected graph of order n and size m with the maximum degree Δ . Then*

$$BSO(G) \leq n - m(2 - \sqrt{2}) \frac{1}{\Delta}$$

with equality if and only if G is a regular graph.

Proof. Without loss of generality, we suppose that $d_u \geq d_v$. Then we have

$$\begin{aligned} BSO(G) &= \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \leq \sum_{uv \in E(G)} \left(\frac{1}{d_v} + (\sqrt{2} - 1) \frac{1}{d_u} \right) \\ &\leq \sum_{uv \in E(G)} \left(\frac{1}{d_v} + \frac{1}{d_u} \right) + m(\sqrt{2} - 2) \frac{1}{\Delta} = n - m(2 - \sqrt{2}) \frac{1}{\Delta} \end{aligned}$$

with equality if and only if G is a regular graph.

Corollary 3.7 *Let G be a connected graph of order n and size m with the maximum degree Δ and the minimum degree δ . Then*

$$\sqrt{2}n \leq BSO(G) + BSO(\overline{G}) \leq 2n - (2 - \sqrt{2}) \left(\frac{m}{\Delta} + \frac{n(n-1) - 2m}{2(n-1-\delta)} \right)$$

with equality if and only if G is a regular graph.

3.2 Bounds in terms of the Randić index, the modified second Zagreb index and the inverse degree index

Theorem 3.8 *Let G be a connected graph with the maximum degree Δ . Then*

$$\sqrt{2}R(G) \leq BSO(G) \leq \sqrt{2}\Delta M_2^*(G)$$

with equality if and only if G is a regular graph.

Proof. By the arithmetic geometric inequality, we have

$$BSO(G) = \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \geq \sum_{uv \in E(G)} \sqrt{\frac{2}{d_u d_v}} = \sqrt{2}R(G)$$

with equality if and only if $d_u = d_v$ for any edges, that is, G is a regular graph. It easy to see that

$$BSO(G) = \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \leq \sum_{uv \in E(G)} \frac{\sqrt{2}\Delta^2}{d_u d_v} = \sqrt{2}\Delta M_2^*(G)$$

with equality if and only if $d_u = d_v = \Delta$ for any edges, that is, G is a regular graph.

Theorem 3.9 *Let G be a connected graph with the maximum degree Δ and the minimum degree δ . Then*

$$BSO(G) \leq \sqrt{mID(G)}$$

with equality if and only if G is a regular graph (when G is non-bipartite) or G is a (Δ, δ) -semiregular bipartite graph (when G is bipartite).

Proof. By the Cauchy-Schwarz inequality, we have

$$BSO(G) = \sum_{uv \in E(G)} 1 \cdot \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \leq \sqrt{\sum_{uv \in E(G)} 1^2 \sum_{uv \in E(G)} \left(\frac{1}{d_u^2} + \frac{1}{d_v^2} \right)} = \sqrt{mID(G)},$$

with equality if and only if $\frac{1}{d_u^2} + \frac{1}{d_v^2}$ is a constant for any edge uv in a connected graph G . By Lemma 2.1, G is a regular graph (when G is non-bipartite) or G is a (Δ, δ) -semiregular bipartite graph (when G is bipartite).

3.3 Bounds in terms of the harmonic index, the symmetric division deg index and the modified second Zagreb index

Theorem 3.10 *Let G be a connected graph with the maximum degree Δ and the minimum degree δ . Then*

$$\sqrt{2}H(G) \leq BSO(G) \leq \frac{1}{\sqrt{2}} \left(\frac{\Delta}{\delta} + \frac{\delta}{\Delta} \right) H(G)$$

with equality if and only if G is a regular graph.

Proof. By Lemma 2.2, we have

$$\begin{aligned} BSO(G) &= \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \leq \sum_{uv \in E(G)} \frac{\sqrt{2} \left(\frac{d_v}{d_u} + \frac{d_u}{d_v} \right)}{d_u + d_v} \\ &\leq \sum_{uv \in E(G)} \frac{1}{\sqrt{2}} \left(\frac{\Delta}{\delta} + \frac{\delta}{\Delta} \right) \frac{2}{d_u + d_v} = \frac{1}{\sqrt{2}} \left(\frac{\Delta}{\delta} + \frac{\delta}{\Delta} \right) H(G). \end{aligned}$$

with equality if and only if $d_u = d_v$ for any edges, that is, G is a regular graph.

By Lemma 2.2, we have

$$\begin{aligned} BSO(G) &= \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \geq \sum_{uv \in E(G)} \frac{2\sqrt{2} \left(\frac{d_v}{d_u} + \frac{d_u}{d_v} + 1 \right)}{3(d_u + d_v)} \\ &\geq \sum_{uv \in E(G)} \frac{2\sqrt{2}(2+1)}{3(d_u + d_v)} = \sqrt{2}H(G) \end{aligned}$$

with equality if and only if $d_u = d_v$ for any edges, that is, G is a regular graph.

Theorem 3.11 *Let G be a connected graph with the maximum degree Δ and the minimum degree δ . Then*

$$\frac{2\sqrt{2}}{3\Delta}SDD(G) + \frac{\sqrt{2}}{3}H(G) \leq BSO(G) \leq \frac{\sqrt{2}}{\delta}SDD(G)$$

with equality if and only if G is a regular graph.

Proof. By Lemma 2.2, we have

$$\begin{aligned} BSO(G) &= \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \leq \sum_{uv \in E(G)} \frac{\sqrt{2} \left(\frac{d_v}{d_u} + \frac{d_u}{d_v} \right)}{d_u + d_v} \\ &\leq \sum_{uv \in E(G)} \frac{\sqrt{2} \left(\frac{d_v}{\delta} + \frac{d_u}{\delta} \right)}{\delta + \delta} = \frac{\sqrt{2}}{\delta}SDD(G). \end{aligned}$$

with equality if and only if $d_u = d_v = \delta$ for any edges, that is, G is a regular graph.

By Lemma 2.2, we have

$$\begin{aligned} BSO(G) &= \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \geq \sum_{uv \in E(G)} \frac{2\sqrt{2} \left(\frac{d_v}{d_u} + \frac{d_u}{d_v} + 1 \right)}{3(d_u + d_v)} \\ &= \sum_{uv \in E(G)} \frac{2\sqrt{2} \left(\frac{d_v}{d_u} + \frac{d_u}{d_v} \right)}{3(d_u + d_v)} + \sum_{uv \in E(G)} \frac{2\sqrt{2}}{3(d_u + d_v)} = \frac{2\sqrt{2}}{3\Delta}SDD(G) + \frac{\sqrt{2}}{3}H(G). \end{aligned}$$

with equality if and only if $d_u = d_v = \Delta$ for any edges, that is, G is a regular graph.

Theorem 3.12 *Let G be a connected graph with n vertices. Then*

$$BSO(G) \leq \sqrt{2M_2^*(G)SDD(G)}$$

with equality if and only if G is a regular graph (when G is non-bipartite) or G is a (Δ, δ) -semiregular bipartite graph (when G is bipartite).

Proof. Let $r = 1$, $a_i = \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}}$ and $b_i = \frac{1}{d_u d_v}$ in Lemma 2.3. Then we have

$$\frac{\left(\sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \right)^2}{\sum_{uv \in E(G)} \frac{1}{d_u d_v}} \leq \sum_{uv \in E(G)} \frac{\frac{1}{d_u^2} + \frac{1}{d_v^2}}{\frac{1}{d_u d_v}} = \sum_{uv \in E(G)} \left(\frac{d_v}{d_u} + \frac{d_u}{d_v} \right),$$

that is,

$$BSO(G) \leq \sqrt{2M_2^*(G)SDD(G)}$$

with equality if and only if $\sqrt{d_u^2 + d_v^2}$ is a constant for any edge uv in G , by Lemma 2.1, G is a regular graph (when G is non-bipartite) or G is a (Δ, δ) -semiregular bipartite graph (when G is bipartite).

Corollary 3.13 *Let G be a connected graph of order n and size m with the maximum degree Δ and the minimum degree δ . Then*

$$BSO(G) \leq \sqrt{mM_2^*(G) \left(\frac{\Delta}{\delta} + \frac{\delta}{\Delta} \right)}$$

with equality if and only if G is a regular graph or a (Δ, δ) -semiregular bipartite graph.

Proof. Without loss of generality, we assume that $d_u \geq d_v$. By the proof of Theorem 3.12, we have

$$\frac{\left(\sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \right)^2}{\sum_{uv \in E(G)} \frac{1}{d_u d_v}} \leq \sum_{uv \in E(G)} \left(\frac{d_v}{d_u} + \frac{d_u}{d_v} \right) \leq \left(\frac{\Delta}{\delta} + \frac{\delta}{\Delta} \right) m$$

with equality if and only if $d_u = \Delta$ and $d_v = \delta$ for any edge uv . This implies that G is a regular graph or a (Δ, δ) -semiregular bipartite graph. Conversely, it is easy to check that equality holds in Corollary 3.7 when G is a regular graph or a (Δ, δ) -semiregular bipartite graph.

3.4 Bounds in terms of the forgotten index

Theorem 3.14 *Let G be a connected graph of order n and size m with the maximum degree Δ and the minimum degree δ . Then*

$$BSO(G) \geq \frac{\sqrt{2}}{\Delta^3 + \delta^3} \left(\frac{m\delta^3}{\Delta} + \frac{F}{2} \right)$$

with equality if and only if G is a regular graph.

Proof. Let $a_i = \sqrt{d_u^2 + d_v^2}$ and $b_i = \frac{1}{d_u d_v}$ in Lemma 2.4. Then $q = \frac{1}{\sqrt{2}\Delta^3}$ and $Q = \frac{1}{\sqrt{2}\delta^3}$. By Lemma 2.4, we have

$$\sum_{uv \in E(G)} \frac{1}{d_u^2 d_v^2} + \frac{1}{2\Delta^3 \delta^3} \sum_{uv \in E(G)} (d_u^2 + d_v^2) \leq \frac{1}{\sqrt{2}} \left(\frac{1}{\Delta^3} + \frac{1}{\delta^3} \right) \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}},$$

that is,

$$\frac{m}{\Delta^4} + \frac{1}{2\Delta^3 \delta^3} F(G) \leq \frac{1}{\sqrt{2}} \left(\frac{1}{\Delta^3} + \frac{1}{\delta^3} \right) BSO(G),$$

that is,

$$BSO(G) \geq \frac{\sqrt{2}}{\Delta^3 + \delta^3} \left(\frac{m\delta^3}{\Delta} + \frac{F}{2} \right)$$

with equality if and only if $d_u = d_v = \Delta$ for any edge uv , that is, G is a regular graph.

Theorem 3.15 *Let G be a connected graph of order n and size m with the maximum degree Δ and the minimum degree δ . Then*

$$BSO(G) \leq \frac{2mSDD(G) + M_2^*(G)F(G)}{2SO(G)}$$

with equality if and only if G is a regular graph (when G is non-bipartite) or G is a (Δ, δ) -semiregular bipartite graph (when G is bipartite).

Proof. Let $a_i = b_i = \sqrt{d_u^2 + d_v^2}$, $c_i = \frac{1}{d_u d_v}$ and $d_i = 1$ in Lemma 2.5. Then we have

$$m \sum_{uv \in E(G)} \frac{d_u^2 + d_v^2}{d_u d_v} + \sum_{uv \in E(G)} \frac{1}{d_u d_v} \sum_{uv \in E(G)} (d_u^2 + d_v^2) \geq 2 \sum_{uv \in E(G)} \frac{\sqrt{d_u^2 + d_v^2}}{d_u d_v} \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2},$$

that is,

$$2mSDD(G) + M_2^*(G)F(G) \geq 2BSO(G)SO(G),$$

that is,

$$BSO(G) \leq \frac{2mSDD(G) + M_2^*(G)F(G)}{2SO(G)}$$

with equality if and only if $a_i = b_i = \sqrt{d_u^2 + d_v^2}$ for any edge uv in G , that is, $d_u^2 + d_v^2$ is a constant for any edge uv in G , by Lemma 2.1, G is a regular graph (when G is non-bipartite) or G is a (Δ, δ) -semiregular bipartite graph (when G is bipartite).

Corollary 3.16 *Let G be a connected graph of size m with the maximum degree Δ and the minimum degree δ . Then*

$$BSO(G) \leq \frac{m(\Delta^2\delta + \delta^2) + \Delta F(G)}{2\sqrt{2}\Delta\delta^3}$$

with equality if and only if G is a regular graph.

Corollary 3.17 *Let G be a connected graph of size m with the maximum degree Δ and the minimum degree δ . Then*

$$BSO(G) \leq \frac{m^2(2\Delta^3 + \Delta^2\delta + \delta^3)}{2\Delta\delta^2SO(G)}$$

with equality if and only if G is a regular graph.

3.5 Bounds in terms of the inverse sum indeg index and the geometric-arithmetic index

Theorem 3.18 *Let G be a connected graph of order n and size m with the maximum degree Δ and the minimum degree δ . Then*

$$BSO(G) \leq \frac{H(G)SDD(G) + 2M_2^*(G)ISI(G)}{\sqrt{2}GA(G)}$$

with equality if and only if G is a regular graph.

Proof. Let $a_i = \sqrt{d_u^2 + d_v^2}$, $b_i = \sqrt{2d_u d_v}$, $c_i = \frac{1}{d_u d_v}$ and $d_i = \frac{1}{d_u + d_v}$ in Lemma 2.5. Then we have

$$\begin{aligned} & \sum_{uv \in E(G)} \frac{1}{d_u + d_v} \sum_{uv \in E(G)} \frac{d_u^2 + d_v^2}{d_u d_v} + \sum_{uv \in E(G)} \frac{1}{d_u d_v} \sum_{uv \in E(G)} \frac{2d_u d_v}{d_u + d_v} \\ & \geq 2 \sum_{uv \in E(G)} \frac{\sqrt{d_u^2 + d_v^2}}{d_u d_v} \sum_{uv \in E(G)} \frac{\sqrt{2d_u d_v}}{d_u + d_v}, \end{aligned}$$

that is,

$$H(G)SDD(G) + 2M_2^*(G)ISI(G) \geq \sqrt{2}BSO(G)GA(G),$$

that is,

$$BSO(G) \leq \frac{H(G)SDD(G) + 2M_2^*(G)ISI(G)}{\sqrt{2}GA(G)}$$

with equality if and only if $\sqrt{d_u^2 + d_v^2} = \sqrt{2d_u d_v}$ for any edge uv , that is, G is a regular graph.

Corollary 3.19 *Let G be a connected graph of size m with the maximum degree Δ and the minimum degree δ . Then*

$$BSO(G) \leq \frac{m^2\Delta^2 + m^2\delta^2 + 4m\Delta ISI(G)}{2\sqrt{2}\Delta\delta^2 GA(G)}$$

with equality if and only if G is a regular graph.

3.6 Bounds in terms of the Sombor index and the modified second Zagreb index

Theorem 3.20 *Let G be a connected graph of size m with the maximum degree Δ and the minimum degree δ . Then*

$$\frac{2m^2}{SO(G)} \leq BSO(G) \leq \frac{1}{\delta^2}SO(G)$$

with equality if and only if G is a regular graph.

Proof. It is easy to see that

$$BSO(G) = \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \leq \frac{1}{\delta^2} \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2} \leq \frac{1}{\delta^2}SO(G),$$

with equality if and only if $d_u = d_v = \Delta$ for any edges, that is, G is a regular graph.

Let $a_i = b_i = \frac{1}{\sqrt{d_u d_v}}$ and $c_i = d_i = \sqrt{d_u^2 + d_v^2}$ in Lemma 2.5. Then

$$2 \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2} \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} \geq 2 \left(\sum_{uv \in E(G)} \sqrt{\frac{d_u^2 + d_v^2}{d_u d_v}} \right)^2 \geq 4m^2,$$

that is,

$$2SO(G)BSO(G) \geq 4m^2,$$

that is,

$$BSO(G) \geq \frac{2m^2}{SO(G)}$$

with equality if and only if G is a regular graph.

Theorem 3.21 *Let G be a connected graph of order n and size m with the maximum degree Δ and the minimum degree δ . Then*

$$\left| \frac{1}{m}BSO(G) - \frac{1}{m^2}SO(G)M_2^*(G) \right| \leq \xi(m) \frac{\sqrt{2}(\Delta + \delta)(\Delta - \delta)^2}{\Delta^2\delta^2},$$

where

$$\xi(m) = \frac{1}{4} \left(1 - \frac{1 + (-1)^{m+1}}{2m^2} \right).$$

Proof. Let $a_i = \sqrt{d_u^2 + d_v^2}$ and $b_i = \frac{1}{d_u d_v}$ in Lemma 2.6. Then $a = \sqrt{2}\delta$, $A = \sqrt{2}\Delta$, $b = \frac{1}{\Delta^2}$ and $B = \frac{1}{\delta^2}$. By Lemma 2.6, we have

$$\begin{aligned} & \left| \frac{1}{m} \sum_{uv \in E(G)} \sqrt{\frac{1}{d_u^2} + \frac{1}{d_v^2}} - \frac{1}{m^2} \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2} \sum_{uv \in E(G)} \frac{1}{d_u d_v} \right| \\ & \leq \frac{1}{m} \left\lfloor \frac{m}{2} \right\rfloor \left(1 - \frac{1}{m} \left\lfloor \frac{m}{2} \right\rfloor \right) \sqrt{2}(\Delta - \delta) \left(\frac{1}{\delta^2} - \frac{1}{\Delta^2} \right), \end{aligned}$$

that is,

$$\left| \frac{1}{m}BSO(G) - \frac{1}{m^2}SO(G)M_2^*(G) \right| \leq \xi(m) \frac{\sqrt{2}(\Delta + \delta)(\Delta - \delta)^2}{\Delta^2\delta^2},$$

where

$$\xi(m) = \frac{1}{m} \left\lfloor \frac{m}{2} \right\rfloor \left(1 - \frac{1}{m} \left\lfloor \frac{m}{2} \right\rfloor \right) = \frac{1}{4} \left(1 - \frac{1 + (-1)^{m+1}}{2m^2} \right).$$

4 The first Banhatti-Sombor index of trees

In this section, we determine the trees with the maximum and minimum first Banhatti-Sombor index among the set of trees of order n , respectively. For a tree T_n of order n

with maximum degree Δ , denote by n_i the number of vertices with degree i in T_n for $1 \leq i \leq \Delta$, and $m_{i,j}$ the number of edges of T_n connecting vertices of degree i and j , where $1 \leq i \leq j \leq \Delta$. Note that T_n is connected, so $m_{1,1} = 0$ for $n \geq 3$. Let $N = \{(i, j) \in \mathbb{N} \times \mathbb{N} : 1 \leq i \leq j \leq \Delta\}$. Then clearly the following relations hold:

$$|V(T_n)| = n = \sum_{i=1}^{\Delta} n_i, \quad (1)$$

$$|E(T_n)| = n - 1 = \sum_{(i,j) \in N} m_{i,j}, \quad (2)$$

and

$$\begin{cases} 2m_{1,1} + m_{1,2} + \dots + m_{1,\Delta} = n_1, \\ m_{1,2} + 2m_{2,2} + \dots + m_{2,\Delta} = 2n_2, \\ \dots \\ m_{1,\Delta} + m_{2,\Delta} + \dots + 2m_{\Delta,\Delta} = \Delta n_{\Delta}. \end{cases} \quad (3)$$

It follows easily from (1) and (3) that

$$n = \sum_{(i,j) \in N} \frac{i+j}{ij} m_{i,j}. \quad (4)$$

And the definition of the first Banhatti-Sombor index is equivalent to

$$SO(G) = \sum_{(i,j) \in P} \sqrt{\frac{1}{i^2} + \frac{1}{j^2}} m_{i,j}. \quad (5)$$

Theorem 4.1 *Let T_n be a tree with n -vertex. Then*

$$\frac{\sqrt{2}(n-3)}{2} + \sqrt{5} \leq BSO(T_n) \leq \sqrt{1 + (n-1)^2}.$$

The equality in the left hand side holds if and only if $T_n \cong P_n$, and the equality in the right hand side holds if and only if $T_n \cong K_{1,n-1}$.

Proof. First, we consider the equality in the left side. Let $N_1 = \{(i, j) \in N : (i, j) \neq (1, 1), (i, j) \neq (1, 2), (i, j) \neq (2, 2)\}$. By equation (4), we have

$$3m_{1,2} + 2m_{2,2} = 2n - \sum_{(i,j) \in N_1} \frac{2(i+j)}{ij} m_{i,j},$$

and by equation (2), we have

$$m_{1,2} + m_{2,2} = n - 1 - \sum_{(i,j) \in N_1} m_{i,j}.$$

Then we obtain the following expression for $m_{1,2}$ and $m_{2,2}$:

$$m_{1,2} = 2 + \sum_{(i,j) \in N_1} \left[2 - \frac{2(i+j)}{ij} \right] m_{i,j},$$

$$m_{2,2} = n - 3 + \sum_{(i,j) \in N_1} \left[\frac{2(i+j)}{ij} - 3 \right] m_{i,j}.$$

According to the expression (5), we have

$$\begin{aligned} BSO(T_n) &= m_{1,2} \sqrt{\frac{1}{4} + 1} + m_{2,2} \sqrt{\frac{1}{4} + \frac{1}{4}} + \sum_{(i,j) \in N_1} \sqrt{\frac{1}{i^2} + \frac{1}{j^2}} m_{i,j} \\ &= \sqrt{5} \left[1 + \sum_{(i,j) \in N_1} \left(1 - \frac{i+j}{ij} \right) m_{i,j} \right] + \frac{\sqrt{2}}{2} \left\{ n - 3 + \sum_{(i,j) \in N_1} \left[\frac{2(i+j)}{ij} - 3 \right] m_{i,j} \right\} \\ &\quad + \sum_{(i,j) \in N_1} \sqrt{\frac{1}{i^2} + \frac{1}{j^2}} m_{i,j} \\ &= \frac{\sqrt{2}}{2} (n - 3) + \sqrt{5} + \sum_{(i,j) \in N_1} \left[\sqrt{\frac{1}{i^2} + \frac{1}{j^2}} + (\sqrt{2} - \sqrt{5}) \frac{i+j}{ij} + \sqrt{5} - \frac{3\sqrt{2}}{2} \right] m_{i,j}. \end{aligned}$$

Let $f(x, y) = \sqrt{\frac{1}{x^2} + \frac{1}{y^2}} + (\sqrt{2} - \sqrt{5}) \frac{x+y}{xy} + \sqrt{5} - \frac{3\sqrt{2}}{2}$, where $(x, y) \in N$, it is easy to see that $f(1, 2) = 0$, $f(2, 2) = 0$ and $f(x, y) > 0$ for $(x, y) \in N_1$. Therefore, $BSO(T_n) = \frac{\sqrt{2}}{2}(n - 3) + \sqrt{5}$ if and only if $m_{i,j} = 0$ for all $(i, j) \in N_1$. And this occurs if and only if $T_n \cong P_n$. Conversely, if $T_n \cong P_n$, by (5), we obtain

$$BSO(P_n) = 2\sqrt{\frac{1}{4} + 1} + (n - 3)\sqrt{\frac{1}{4} + \frac{1}{4}} = \frac{\sqrt{2}}{2}(n - 3) + \sqrt{5}.$$

Thus, we have $BSO(T_n) \geq BSO(P_n)$ with equality if and only if $T_n \cong P_n$.

Now, we consider the equality on the right side. Let $N_2 = \left\{ (i, j) \in N : (i, j) \neq (1, 1), (i, j) \neq (1, \Delta), (i, j) \neq (\Delta, \Delta) \right\}$. Similar to the proof of the above, by equation (4), we have

$$(\Delta + 1)m_{1,\Delta} + 2m_{\Delta,\Delta} = \Delta n - \sum_{(i,j) \in N_2} \Delta \frac{i+j}{ij} m_{i,j},$$

and by equation (2), we have

$$m_{1,\Delta} + m_{\Delta,\Delta} = n - 1 - \sum_{(i,j) \in N_2} m_{i,j}.$$

Then we obtain the following expression for $m_{1,\Delta}$ and $m_{\Delta,\Delta}$:

$$(\Delta - 1)m_{1,\Delta} = (\Delta - 2)n + 2 - \sum_{(i,j) \in N_2} \left(\Delta \frac{i+j}{ij} - 2 \right) m_{i,j},$$

$$(\Delta - 1)m_{\Delta,\Delta} = n - (\Delta + 1) + \sum_{(i,j) \in N_2} \left(\Delta \frac{i+j}{ij} - (\Delta + 1) \right) m_{i,j}.$$

According to the expression (4), we have

$$\begin{aligned} BSO(T_n) &= m_{1,\Delta} \sqrt{\frac{1}{\Delta^2} + 1} + m_{\Delta,\Delta} \sqrt{\frac{1}{\Delta^2} + \frac{1}{\Delta^2}} + \sum_{(i,j) \in N_2} \sqrt{\frac{1}{i^2} + \frac{1}{j^2}} m_{i,j} \\ &= \frac{\sqrt{\Delta^2 + 1}}{\Delta(\Delta - 1)} \left[(\Delta - 2)n + 2 - \sum_{(i,j) \in N_2} \left(\Delta \frac{i+j}{ij} - 2 \right) m_{i,j} \right] \\ &\quad + \frac{\sqrt{1 + \Delta^2}}{\Delta(\Delta - 1)} \left[n - (\Delta + 1) + \sum_{(i,j) \in N_2} \left(\Delta \frac{i+j}{ij} - (\Delta + 1) \right) m_{i,j} \right] \\ &\quad + \sum_{(i,j) \in N_2} \sqrt{\frac{1}{i^2} + \frac{1}{j^2}} m_{i,j} \\ &= \frac{(\Delta - 2)n\sqrt{\Delta^2 + 1} + \sqrt{2}(n - \Delta - 1) + 2\sqrt{\Delta^2 + 1}}{\Delta(\Delta - 1)} \\ &\quad + \sum_{(i,j) \in N_2} \left[\sqrt{\frac{1}{i^2} + \frac{1}{j^2}} + \frac{\sqrt{2} - \sqrt{\Delta^2 + 1}}{\Delta - 1} \frac{i+j}{ij} + \frac{2\sqrt{\Delta^2 + 1} - \sqrt{2}(\Delta + 1)}{\Delta(\Delta - 1)} \right] m_{i,j}. \end{aligned}$$

Let $g(x, y) = \sqrt{\frac{1}{x^2} + \frac{1}{y^2}} + \frac{\sqrt{2} - \sqrt{\Delta^2 + 1}}{\Delta - 1} \frac{x+y}{xy} + \frac{2\sqrt{\Delta^2 + 1} - \sqrt{2}(\Delta + 1)}{\Delta(\Delta - 1)}$, where $(x, y) \in N$, it is easy to see that $f(1, \Delta) = 0$, $f(\Delta, \Delta) = 0$ and $f(x, y) < 0$ for $(x, y) \in N_2$. Therefore, $BSO(T_n) = \frac{(\Delta - 2)n\sqrt{\Delta^2 + 1} + \sqrt{2}(n - \Delta - 1) + 2\sqrt{\Delta^2 + 1}}{\Delta(\Delta - 1)}$ if and only if $m_{i,j} = 0$ for all $(i, j) \in N_2$. And this occurs if and only if $n_2 = n_3 = \dots = n_{\Delta-1} = 0$.

Let $h(x) = \frac{(x-2)n\sqrt{x^2+1} + \sqrt{2}(n-x-1) + 2\sqrt{x^2+1}}{x(x-1)}$. By derivative, we know that $h(x)$ is an increasing function for $[2, +\infty)$. Thus

$$h(\Delta) \leq h(n - 1) = \sqrt{1 + (n - 1)^2}.$$

Conversely, $BSO(K_{1,n-1}) = \sqrt{1 + (n-1)^2}$. Thus, we have $BSO(T_n) \leq BSO(K_{1,n-1})$ with equality if and only if $T_n \cong K_{1,n-1}$.

Similar to the method used in Theorem 4.1, we now give an upper bound on chemical trees without its proof.

Theorem 4.2 *Let T_n is a chemical tree with n vertices. If $n - 2 = 0 \pmod{3}$, then*

$$BSO(T_n) \leq \frac{2\sqrt{17}(n+1) + \sqrt{2}(n-5)}{12}$$

with equality if and only if $n_2 = n_3 = 0$.

References

- [1] R. Cruz, I. Gutman, J. Rada, Sombor index of chemical graphs, Appl. Math. Comput. 399 (2021) 126018.
- [2] S.S. Dragomir, On some inequalities (Romanian), "Caiete Metodico Ştiinţifice", No. 13, 1984, pp. 20. Faculty of Mathematics, Timişoara University, Romania.
- [3] K.Ch. Das, A.S. Çevik, I.N. Cangul, Y. Shang, On Sombor Index, Symmetry 13 (2021) 140.
- [4] J.B. Diaz, F.T. Metcalf, Stronger forms of a class of inequalities of G. Polja-G. Szego and L.V. Kantorovich, Bull. Amer. Math. Soc. 69 (1963) 415-418.
- [5] H. Deng, Z. Tang, R. Wu, Molecular trees with extremal values of Sombor indices, Int J Quantum Chem. DOI: 10.1002/qua.26622.
- [6] S. Fajtlowicz, On conjectures of Graffiti-II, Congr. Numer. 60 (1987) 187-197.
- [7] B. Furtula, I. Gutman, A forgotten topological index, J. Math. Chem. 53 (2015) 1184-1190.
- [8] I. Gutman, Geometric approach to degree-based topological indices: Sombor indices, MATCH Commun. Math. Comput. Chem. 86 (2021) 11-16.
- [9] V.R.Kulli, On Banhatti-Sombor indices, International Journal of Applied Chemistry, 8 (2021) 21-25.
- [10] V.R. Kulli, I. Gutman, Computation of Sombor Indices of Certain Networks, International Journal of Applied Chemistry, 8 (2021) 1-5.
- [11] J. Karamata, Sur une inégalité relative aux fonctions convexes, Publ. Math. Univ. Belgrade 1 (1932) 145-148.
- [12] X. Li, R.N. Mohapatra, R.S. Rodriguez, Grüss-type inequalities, J. Math. Anal. Appl. 267 (2002) 434-443.

- [13] Z. Lin, L. Miao, T. Zhou, On the spectral radius, energy and Estrada index of the Sombor matrix of graphs, arXiv:2102.03960.
- [14] I. Milovanović, E. Milovanović, M. Matejić, On some mathematical properties of sombor indices, Bull. Int. Math. Virtual Inst. 11 (2021) 341-353.
- [15] S. Nikolić, G. Kovačević, A. Miličević, N. Trinajstić, The Zagreb indices 30 years after, Croat. Chem. Acta 76 (2003) 113-124.
- [16] M. Randić, On characterization of molecular branching, J. Am. Chem. Soc. 97 (1975) 6609-6615.
- [17] J. Radon, Über die absolut additiven Mengenfunktionen, Wiener-Sitzungsber. (IIa) 122 (1913) 1295-1438.
- [18] I. Redžepović, Chemical applicability of Sombor indices, J. Serb. Chem. Soc. <https://doi.org/10.2298/JSC201215006R>.
- [19] T. Réti, T. Došlić, A. Ali, On the Sombor index of graphs, Contrib. Math. 3 (2021) 11-18.
- [20] D. Vukičević, Bond additive modelling 2. Mathematical properties of max-min rodeg index, Croat. Chem. Acta 83 (2010) 261-273.
- [21] D. Vukičević, B. Furtula, Topological index based on the ratios of geometrical and arithmetical means of end-vertex degrees of edges, J. Math. Chem. 46 (2000) 1369-1376.
- [22] D. Vukičević, M. Gašperov, Bond additive modelling 1. Adriatic indices, Croat. Chem. Acta 83 (2010) 261-273.
- [23] Z. Wang, Y. Mao, Y. Li, B. Furtula, On relations between Sombor and other degree-based indices, J. Appl. Math. Comput. <https://doi.org/10.1007/s12190-021-01516-x>.