

On the Gap-sum and Gap-product Sequences of Integer Sequences

Paul Barry
 School of Science
 Waterford Institute of Technology
 Ireland
 pbarry@wit.ie

Abstract

In this note, we explore two families of sequences associated to a suitable integer sequence: the gap-sum sequence and the gap-product sequence. These are the sums and the products of consecutive numbers not in the original sequence. We give closed forms for both, in terms of the original sequence, and in the case of Horadam sequences, we find the generating function of the gap-sum sequence. For some elementary sequences, we indicate that the gap-product sequences are given by the Fuss-Catalan-Raney numbers.

1 Introduction

Let A be the integer sequence $(a_n)_{n \geq 0}$. The consecutive numbers between, but not including, a_n to a_{n+1} will be called the n th A -gap, or just the n th gap, when the sequence in question is clear. Note that for sequences with $a_{n+1} > a_n + 1$, each gap is an arithmetic sequence with initial term $a_n + 1$, with $a_{n+1} - a_n - 1$ terms, and increment $+1$. If $a_{n+1} = a_n + 1$, then we say that the gap is 0.

Example 1. We consider the sequence $a_n = J_{n+2}$ which begins

$$1, 3, 5, 11, 21, 43, 85, 171, \dots$$

Here, J_n is the n -th Jacobsthal number, and hence

$$a_n = 4 \frac{2^n}{3} - \frac{(-1)^n}{3}.$$

The first gaps of this sequence are then

$$1, \underbrace{2}, 3, \underbrace{4}, 5, \underbrace{6, 7, 8, 9, 10}, 11, \underbrace{12, 13, \dots, 20}, 21, \dots$$

We define the *gap-sum sequence* of a_n to be the sequence whose n -th term is the sum of the elements of the n th-gap. Thus for the sequence above, the gap-sum sequence begins

$$2, 4, 40, 144, \dots$$

This sequence is often called the sequence of sums of consecutive non- A numbers. Similarly, we define the *gap-product sequence* of a_n to be the sequence whose n -th term is the product of the elements of the n -th gap. For the sequence of the above example, we have that the gap-product sequence begins

$$2, 4, 30240, 60949324800, \dots$$

A first result is the following.

Proposition 2. *Let S_n be the gap-sum sequence of a_n . Then we have*

$$S_n = \sum_{j=1}^{a_{n+1}-a_n-1} (a_n + j).$$

Let P_n be the gap-product sequence of a_n . Then we have

$$P_n = \prod_{j=1}^{a_{n+1}-a_n-1} (a_n + j).$$

Proof. This is a reformulation of the definitions of the gap-sum and the gap-product sequences. □

Example 3. We consider the Fibonacci numbers [A000045](#)

$$F_n = \frac{1}{\sqrt{5}} \left(\frac{1}{2} + \frac{\sqrt{5}}{2} \right)^n - \frac{1}{\sqrt{5}} \left(\frac{1}{2} - \frac{\sqrt{5}}{2} \right)^n.$$

We find that the formula for the gap-sum sequence $\sum_{j=1}^{F_{n+1}-F_n-1} F_n + j$ above in this case gives us

$$0, 0, 0, 0, 4, 13, 42, 119, 330, 890, 2376, \dots$$

This is [A109454](#) in the On-Line Encyclopedia of Integer Sequences (OEIS) [4, 5].

Example 4. We let A denote the sequence [A000040](#) of prime numbers

$$2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47, \dots$$

The prime gap-sum sequence then begins

$$0, 4, 6, 27, 12, 45, 18, 63, 130, 30, 170, \dots$$

This is the sequence [A054265](#) in the OEIS, where it is described as the sequence of sums of composite numbers between successive primes.

Using the formula for the sum of an arithmetic series, we obtain the following result for the gap-sum sequence.

Proposition 5. Let a_n be a sequence such that $a_{n+1} > a_n$. Then

$$S_n = \frac{a_{n+1} - a_n - 1}{2} (a_n + a_{n+1}).$$

Proof. The n -th gap is an arithmetic sequence beginning with $a_n + 1$ and having step $d = 1$, with $a_{n+1} - a_n - 1$ terms. Thus we have

$$S_n = \frac{a_{n+1} - a_n - 1}{2} (2(a_n + 1) + (a_{n+1} - a_n - 1) - 1) = \frac{a_{n+1} - a_n - 1}{2} (a_n + a_{n+1}).$$

□

For instance, if p_n denotes the n -th prime, then the prime gap-sum sequence has general term

$$S_n = \frac{p_{n+1} - p_n - 1}{2} (p_n + p_{n+1}).$$

We note that the gap-sum sequence can be the zero sequence. For example, if $a_n = n$, then the gap-sum sequence is the zero sequence

$$0, 0, 0, 0, \dots$$

Corollary 6. Let $a_n = rn$, $r \in \mathbb{N}$. Then the gap-sum sequence of a_n is given by

$$S_n = \frac{1}{2}(2n + 1)r(r - 1) = (2n + 1) \binom{r}{2}.$$

Corollary 7. Let $a_n = k^n$, then the gap-sum sequence for a_n is given by

$$S_n = \frac{1}{2} ((k^2 - 1)k^{2n} - (k + 1)k^n),$$

with generating function

$$S(x) = \frac{1}{2} \frac{(k + 1)(kx + k - 2)}{(1 - kx)(1 - k^2x)}.$$

For $k = 2$, this is the gap-sum sequence for 2^n . This is [A103897](#), which begins

$$0, 3, 18, 84, 360, 1488, 6048, 24384, \dots$$

This sequence has generating function

$$\frac{3x}{(1 - 2x)(1 - 4x)}.$$

We note that if $a_n = 2^n - 1$, then S_n is the sequence that begins

$$0, 2, 15, 77, 345, 1457, 5985, \dots$$

This sequence has the generating function

$$\frac{x(2 + x)}{(1 - x)(1 - 2x)(1 - 4x)}.$$

Example 8. We consider the case of $a_n = 2n^2$. We find that

$$S_n = 8n^3 + 10n^2 + 6n + 1,$$

with generating function

$$\frac{1 + 21x + 33x^2 + 3x^3}{(1-x)^4}.$$

This sequence begins

$$1, 25, 117, 325, 697, 1281, 2125, 3277, 4785, \dots$$

The gap-sum sequence for $a_n = n^2$ is [A048395](#). For this sequence we have

$$S_n = 2n^3 + 2n^2 + n,$$

with generating function

$$\frac{x(1+x)(5+x)}{(1-x)^4}.$$

We give below a short table of figurate numbers and their gap-sum sequences.

G.f. of a_n	a_n	S_n	G.f. of S_n
$\frac{x(1+x)}{(1-x)^3}$	n^2	$2n^3 + 2n^2 + n$	$\frac{x(1+x)(5+x)}{(1-x)^4}$
$\frac{x}{(1-x)^3}$	$\frac{n(n+1)}{2}$	$\frac{n(n+1)^2}{2}$	$\frac{x(2+x)}{(1-x)^4}$
$\frac{2x}{(1-x)^3}$	$n(n+1)$	$(2n+1)(n+1)^2$	$\frac{1+8x+3x^2}{(1-x)^4}$
$\frac{x(2+x)}{(1-x)^3}$	$\frac{n(3n+1)}{2}$	$\frac{(3n+1)(3n^2+4n+2)}{2}$	$\frac{1+14x+11x^2+x^3}{(1-x)^4}$
$\frac{x}{(1-x)^4}$	$\frac{n}{6}(n+1)(n+2)$	$\frac{n(n+1)(n+2)(n+3)(2n+3)}{4!}$	$\frac{5x(1+x)}{(1-x)^6}$
$\frac{x}{(1-x)^5}$	$\binom{n+3}{4}$	$\frac{n(n+1)(n+2)^2(n+3)(n^2+6n+11)}{144}$	$\frac{x(9+18x+7x^2+x^3)}{(1-x)^8}$
$\frac{x(1+2x)}{(1-x)^3}$	$\frac{n(3n-1)}{2}$	$\frac{3n(3n^2+2n+1)}{3}$	$\frac{3x(x^2+5x+3)}{(1-x)^4}$

The gap-sum sequence of the triangular numbers $\frac{n(n+1)}{2} = \binom{n+1}{2}$ is [A006002](#).

The gap-sum sequence of the tetrahedral numbers $\frac{n}{6}(n+1)(n+2) = \binom{n+2}{3}$ can be expressed as

$$5 \binom{n+3}{4} + 10 \binom{n+3}{5}.$$

This is 5 times [A005585](#), the 5-dimensional pyramidal numbers $\frac{n(n+1)(n+2)(n+3)(2n+3)}{5!}$.

The gap-sum sequence of $\binom{n+3}{4}$ can be expressed as

$$9 \binom{n+4}{5} + 36 \binom{n+4}{6} + 34 \binom{n+4}{7} + \binom{n+3}{7}.$$

2 The gap-product sequence

Proposition 9. Let a_n be a sequence with $a_{n+1} \geq a_n$. Then the gap-product sequence P_n is given by

$$P_n = \frac{(a_{n+1} - 1)!}{a_n!}.$$

Proof. The product of the terms of an arithmetic sequence with initial term t_1 , with r terms and increment d , is given by

$$P = \frac{\Gamma\left(\frac{t_1}{d} + r\right)}{\Gamma(t_1)}.$$

For P_n , we have $t_1 = a_n + 1$, $r = a_{n+1} - a_n - 1$, and $d = 1$. The result follows from this. \square

Example 10. We consider the Fibonacci numbers (whose terms satisfy $a_{n+1} > a_n$ after $n = 2$). We obtain that the gap-product sequence

$$P_n = \frac{(F_{n+1} - 1)!}{F_n!}$$

begins

$$1, 1, 1, 1, 4, 42, 11880, 390700800, 169958063987712000, \dots$$

Since $F_{n+1} = F_n + F_{n-1}$, we have that

$$P_n = \frac{\prod_{j=1}^{F_{n-1}-1} (F_{n+1} - j) F_n!}{F_n!} = \prod_{j=1}^{F_{n-1}-1} (F_{n+1} - j).$$

Example 11. For the sequence $a_n = 4\frac{2^n}{3} - \frac{(-1)^n}{3}$, we have

$$P_n = \frac{\left(8\frac{2^n}{3} + \frac{(-1)^n}{3} - 1\right)!}{\left(4\frac{2^n}{3} - \frac{(-1)^n}{3}\right)!} = \frac{(J_{n+3} - 1)!}{J_{n+2!}},$$

where $J_n = \frac{2^n}{3} - \frac{(-1)^n}{3}$ is the n -th Jacobsthal number ([A001045](#)).

3 The gap-sum sequences of Horadam sequences

A Horadam sequence is a second-order recurrence sequence $a_n = a_n(\alpha, \beta, r, s)$ that is defined by

$$a_n = ra_{n-1} + sa_{n-2},$$

with $a_0 = \alpha$, $a_1 = \beta$. For instance, $F_n = F_n(0, 1, 1, 1)$ and $J_n = J_n(0, 1, 1, 2)$. The generating function of $a_n a_n(\alpha, \beta, r, s)$ is given by

$$\frac{\alpha - (\alpha r - \beta)x}{1 - rx - sx^2}.$$

In this section we wish to find the generating function of the gap-sum sequence for a general Horadam sequence. We know that the gap-sum sequence for a general sequence a_n is given by

$$S_n = \frac{1}{2} (a_n + a_{n+1}) (a_{n+1} - a_n - 1) = \frac{1}{2} (a_{n+1}^2 - a_n^2 - a_n - a_{n+1}).$$

Thus the generating function of S_n is determined by those of a_n , a_{n+1} , a_n^2 , and a_{n+1}^2 . In the case of Horadam sequences, the sequence a_{n+1} is again a Horadam sequence, with generating function

$$\frac{\beta + \alpha s x}{1 - r x - s x^2},$$

with initial terms $\beta, r\alpha + s\beta$. Thus

$$a_{n+1}(\alpha, \beta, r, s) = a_n(\beta, r\alpha + s\beta, r, s).$$

Thus we are left with the problem of finding the generating function of the square of a Horadam sequence. This question has been addressed in [3].

Lemma 12. *The generating function of $a_n^2(\alpha, \beta, r, s)$ is given by*

$$g_2(\alpha, \beta, r, s) = \frac{\alpha^2 - (\alpha^2(r^2 + s) - \beta^2)x - s(\alpha^2 r^2 - 2\alpha\beta r + \beta^2)x^2}{1 - (r^2 + s)x - s(r^2 + s)x^2 + s^3 x^3},$$

and the generating function of $a_{n+1}^2(\alpha, \beta, r, s)$ is given by

$$\frac{\beta^2 + s(\alpha^2 s + 2\alpha\beta r - \beta^2)x - \alpha^2 s^3 x^2}{1 - (r^2 + s)x - s(r^2 + s)x^2 + s^3 x^3}.$$

Example 13. The generating function of the square of J_{n+2} is given by

$$g_2(1, 3, 1, 2) = \frac{1 + 6x - 8x^2}{1 - 3x - 6x^2 + 8x^3}.$$

The lemma then leads to the following result.

Proposition 14. *The generating sequence of the gap-sum sequence of the Horadam sequence $a_n = a_n(\alpha, \beta, r, s)$ is given as follows. Let*

$$W = \frac{\beta^2 + s(\alpha^2 s + 2\alpha\beta r - \beta^2)x - \alpha^2 s^3 x^2}{1 - (r^2 + s)x - s(r^2 + s)x^2 + s^3 x^3}.$$

Let

$$X = \frac{\alpha^2 - (\alpha^2(r^2 + s) - \beta^2)x - s(\alpha^2 r^2 - 2\alpha\beta r + \beta^2)x^2}{1 - (r^2 + s)x - s(r^2 + s)x^2 + s^3 x^3}.$$

Let

$$Y = \frac{\beta + \alpha s x}{1 - r x - s x^2},$$

and let

$$Z = \frac{\alpha - (\alpha r - \beta)x}{1 - r x - s x^2}.$$

Then the generating function of the gap-sum sequence of the Horadam sequence $a_n = a_n(\alpha, \beta, r, s)$ [A002605](#) which begins

$$1, 2, 6, 16, 44, 120, 328, \dots,$$

is given by

$$W - X - Y - Z.$$

Example 15. For the Horadam sequence $a_n(1, 2, 2, 2)$ we find that the gap-sum sequence has generating function

$$\frac{3x(4 + x - 2x^2)}{(1 - 2x - 2x^2)(1 - 6x - 12x^2 + 8x^3)}.$$

This expands to give the sequence S_n that begins

$$0, 12, 99, 810, 6150, 46368, 347004, \dots$$

Example 16. The table below summarizes the cases of the Fibonacci numbers $F_{n+1} = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k}$, the Jacobsthal numbers $J_{n+1} = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k} 2^k$, and the Pell numbers $P_{n+1} = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k} 2^{n-2k}$ [A000129](#). Note that the sequence from the previous example has $a_n(1, 2, 2, 2) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k} 2^{n-k}$.

Sequence	G.f.	G.f. of gap-sum	First terms of gap-sum
F_{n+1}	$\frac{1}{1-x-x^2}$	$-\frac{1-3x-x^2+x^3}{(1-x-x^2)(1-2x-2x^2+x^3)}$	$-1, 0, 0, 4, 13, 42, 119, 330, \dots$
J_{n+1}	$\frac{1}{1-x-2x^2}$	$-\frac{1-6x}{(1-2x)(1-2x-8x^2)}$	$-1, 2, 4, 40, 144, 672, 2624, 10880$
P_{n+1}	$\frac{1}{1-2x-x^2}$	$\frac{x(7+2x-x^2)}{(1-2x-x^2)(1-5x-5x^2+x^3)}$	$0, 7, 51, 328, 1980, 11711, 68663, 401184, \dots$

We note that the -1 's are artifacts arising from consecutive elements of the sequences in question being equal.

4 A special gap-product sequence related to the Fuss-Catalan numbers

We have seen that the gap-product sequence of the sequence a_n is given by

$$P_n = \frac{(a_{n+1} - 1)!}{a_n!}.$$

We take the special case of the one-parameter sequence

$$a_n = kn + 1.$$

We then have that P_n (which is again a one-parameter sequence) is given by

$$P_n = \frac{(k(n+1) + 1 - 1)!}{(kn + 1)!} = \frac{(k(n+1))!}{(kn + 1)!}.$$

Proposition 17. *The gap-product sequence of the sequence $a_n = kn + 1$ is given by*

$$P_n = k!FC(k, n)$$

where $F(n, k) = \frac{1}{kn+1} \binom{(k+1)n}{n}$ is the n, k Fuss Catalan number.

Proof. This follows since

$$P_n = \frac{(k(n+1) + 1 - 1)!}{(kn+1)!} = \frac{(k(n+1))!}{(kn+1)!} = \frac{k!}{k+1} \binom{(n+1)k}{k}.$$

□

In the following table, we show the gap-product sequences corresponding to the sequences $kn + 1$ on the left.

1	1	1	1	1	1	1	...
$n + 1$	1	1	1	1	1	1	...
$2n + 1$	2	4	6	8	10	12	...
$3n + 1$	6	30	72	132	210	306	...
$4n + 1$	24	336	1320	6840	12144	19656	...
$5n + 1$	120	5040	32760	116280	303600	657720	...

The following is a table of the corresponding Fuss-Catalan numbers.

	$\binom{n}{n}$	$\frac{1}{n+1} \binom{2n}{n}$	$\frac{1}{2n+1} \binom{3n}{n}$	$\frac{1}{3n+1} \binom{4n}{n}$	$\frac{1}{4n+1} \binom{5n}{n}$	$\frac{1}{5n+1} \binom{6n}{n}$...
1	1	1	1	1	1	1	...
$n + 1$	1	1	1	1	1	1	...
$2n + 1$	1	2	3	4	5	6	...
$3n + 1$	1	5	12	22	35	51	...
$4n + 1$	1	14	55	140	285	506	...
$5n + 1$	1	42	273	969	2530	5481	...

Reading left to right, these sequences are [A000012](#), [A000108](#), [A001764](#), [A002293](#), [A002294](#) and [A002295](#).

Example 18. We take the one-parameter sequence $a_n = kn + 2$. We find that

$$P_n = \frac{k! \binom{(n+1)k+1}{k}}{kn+2}.$$

For $k \geq 0$, this gives us the array that begins

2	1/2	1/2	1/2	1/2	1/2	1/2	...
$n + 2$	1	1	1	1	1	1	...
$2n + 2$	3	5	7	9	11	13	...
$3n + 2$	12	42	90	156	240	342	...
$4n + 2$	60	504	1716	4080	7980	13800	...
$5n + 1$	360	7920	43680	143640	358800	755160	...

The first two sequences here are [A001710](#) and [A102693](#). Dividing P_n by $\frac{k!}{2}$ now gives us the array with general element $\frac{2\binom{(n+1)k+1}{k}}{kn+2}$ that begins

	$\frac{2\binom{n+1}{n}}{2}$	$\frac{2\binom{2n+1}{n}}{n+2}$	$\frac{2\binom{3n+1}{2}}{2n+2}$	$\frac{2\binom{4n+1}{n}}{3n+2}$	$\frac{2\binom{5n+1}{n}}{4n+2}$	$\frac{2\binom{6n+1}{n}}{5n+2}$...
$\frac{2\binom{1}{0}}{2}$	1	1	1	1	1	1	...
$\frac{2\binom{n+2}{1}}{n+2}$	2	2	2	2	2	2	...
$\frac{2\binom{2n+3}{2}}{2n+2}$	3	5	7	9	11	13	...
$\frac{2\binom{3n+4}{3}}{3n+2}$	4	14	30	52	80	114	...
$\frac{2\binom{4n+5}{4}}{4n+2}$	5	42	143	340	665	1150	...
$\frac{2\binom{5n+6}{5}}{5n+2}$	6	136	728	2394	5980	12586	...

The Fuss-Catalan-Raney numbers [2] are the numbers

$$R_{p,r}(n) = \frac{r}{pn+r} \binom{pn+r}{n}.$$

We have

$$R_{p,2}(n) = \frac{2}{pn+2} \binom{pn+2}{n}.$$

Now we have

$$R_{p,2}(n) = \frac{2}{pn+2} \binom{pn+2}{n} = \frac{2}{(p-1)n+2} \binom{pn+1}{n}.$$

In general, we have the following result.

Proposition 19. *For $a_n = kn + r$, we have*

$$P_n = \frac{k!}{r} R_{k+1,r}(n).$$

We can reverse this identity, to obtain a characterization of the Fuss-Catalan-Raney numbers. This says that $R_{k,r}(n)$ is $\frac{r}{k!}$ times the gap-product sequence of the sequence $a_n = kn + r$.

5 Final comments

The formula

$$S_n = \sum_{j=1}^{a_{n+1}-a_n-1} (a_n + j) = \frac{a_{n+1} - a_n - 1}{2} (a_n + a_{n+1})$$

may still be used when the sequence a_n is not monotonic increasing, though the interpretation of the resulting sequence S_n is not immediate.

Example 20. We look at the sequence [A088748](#), which begins

$$1, 2, 3, 2, 3, 4, 3, 2, 3, 4, 5, 4, 3, 4, 3, 2, 3, 4, 5, 4, 5, 6, 5, \dots$$

We find that as with the natural numbers n , we have $S_n = 0$ (this assumes the convention that a summation from $j = 1$ to a negative number is empty). In the case of this sequence, the quantity $a_{n+1} - a_n - 1$ is only non-zero at those indices where the paper-folding sequence [A014707](#) [1] is non-zero (and equal to 1). At those indices, $a_{n+1} - a_n - 1$ is equal to -2 . We now look at the sequence $\tilde{S}_n = \sum_{j=1}^{|a_{n+1} - a_n - 1|} (a_n + j)$, which begins

$$0, 0, 9, 0, 0, 11, 9, 0, 0, 13, 11, 0, 11, 9, 0, 0, 0, 13, 0, 0, \dots$$

The sequence whose only non-zero elements of value $2a_n - 1$ occur when $a_{n+1} - a_n = -1$ begins

$$0, 0, 5, 0, 0, 7, 5, 0, 0, 0, 9, 7, 0, 7, 5, 0, 0, 0, 9, 0, 0, \dots$$

Subtracting these sequences, we get

$$0, 0, 4, 0, 0, 4, 4, 0, 0, 0, 4, 4, 0, 4, 4, 0, 0, 0, 4, 0, 0, \dots,$$

or 4 times the paper-folding sequence.

This happens since the quantity $|a_{n+1} - a_n - 1|$ is equal only to 2 when it is non-zero, the summation at indices where it is non-zero becomes $2a_n + 3$, thus giving us $2a_n + 3 - (2a_n - 1) = 4$ at those points where the paper-folding sequence is non-zero, and 0 otherwise.

References

- [1] J. P. Allouche and J. Shallit, *Automatic Sequences*, Cambridge University Press.
- [2] K. A. Penson and K. Życzkowski, Product of Ginibre matrices: Fuss-Catalan and Raney distributions, *Phys. Rev. E*, **83** 2011, 061118.
- [3] A formula for the generating functions of powers of Horadam's sequence, *Australasian Journal of Combinatorics* **30** (2004) 207–212.
- [4] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*. Published electronically at <http://oeis.org>, 2016.
- [5] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, *Notices Amer. Math. Soc.*, **50** (2003), 912–915.

2020 *Mathematics Subject Classification*: Primary 11B83; Secondary 11B25, 11B37, 11B39.
Keywords: Integer sequence, recurrence, generating function, gap-sum sequence, gap-product sequence, Horadam sequence, Fibonacci numbers, Jacobsthal numbers, Pell numbers, Fuss-Catalan numbers.

(Concerned with sequences [A000045](#), [A000012](#), [A000040](#), [A000108](#), [A000129](#), [A001045](#), [A001710](#), [A001764](#), [A002293](#), [A002294](#), [A002295](#), [A002605](#), [A005585](#), [A014707](#), [A006002](#), [A048395](#), [A054265](#), [A088748](#), [A102693](#), [A103897](#) and [A109454](#).)