TENSOR PRODUCT WEIGHT MODULES FOR THE MIRROR-TWISTED HEISENBERG-VIRASORO ALGEBRA

DONGFANG GAO AND KAIMING ZHAO

ABSTRACT. In this paper, we study irreducible weight modules with infinite dimensional weight spaces over the mirror-twisted Heisenberg-Virasoro algebra \mathcal{D} . More precisely, the necessary and sufficient conditions for the tensor products of irreducible highest weight modules and irreducible modules of intermediates series over \mathcal{D} to be irreducible are determined by using "shifting technique". This leads to a family of new irreducible weight modules over \mathcal{D} . Then we obtain that any two such tensor products are isomorphic if and only if the corresponding highest weight modules and modules of intermediate series are isomorphic respectively. Also we discuss submodules of the tensor product module when it is not irreducible.

1. INTRODUCTION

It is well-known that the Virasoro algebra \mathcal{V} is one of the most important Lie algebras in mathematics and in mathematical physics, because of its widespread applications, for example, in quantum physics (see [12]), conformal field theory (see [7]), vertex operator algebras (see [8, 11]), and so on. Many other interesting and important algebras are closely related to the Virasoro algebra, such as the Schrödinger-Virasoro algebra (see [14, 15]), the Heisenberg-Virasoro algebra (see [1, 20]) and the mirror-twisted Heisenberg-Virasoro algebra \mathcal{D} (see [3, 17]) which is the even part of the mirror-twisted N = 2 superconformal algebra (see [3]). The mirror-twisted Heisenberg-Virasoro algebra \mathcal{D} has a nice structure (see Definition 2.1) which is similar to the Heisenberg-Virasoro algebra. This algebra is the main object that we concern in this paper.

Highest weight modules and Harish-Chandra modules over many Lie algebras with triangular decompositions have been the most popular modules in representation theory. These modules are well understood for many infinite dimensional Lie algebras, for example, the Virasoro algebra in [2, 9, 22], the Heisenberg-Virasoro algebra in [1, 20], the Schrödinger-Virasoro algebra in [16, 23], and the mirror-twisted Heisenberg-Virasoro algebra in [17].

For the last two decades more and more mathematicians are interested in irreducible weight modules with infinite dimensional weight spaces, see for example, [4, 6, 24]. An efficient way to obtain irreducible weight module with infinite dimensional weight spaces is the tensor product of irreducible highest weight modules and modules of intermediate series (or loop modules), see for example, [4, 13] for affine Kac-Moody algebras, [5, 24] for the Virasoro algebra, [21] for the Heisenberg-Virasoro algebra. But it is generally not easy to determine the conditions for the tensor product to be irreducible (see [5, 18, 21]). In the present paper, we will determine the

²⁰⁰⁰ Mathematics Subject Classification. 17B10, 17B20, 17B65, 17B66, 17B68.

Key words and phrases. mirror-twisted Heisenberg-Virasoro algebra, tensor product, highest weight module, module of intermediate series, irreducible module.

necessary and sufficient conditions for the tensor product of an irreducible highest weight module and an irreducible module of intermediate series over \mathcal{D} to be irreducible.

The paper is organized as follows. In Section 2, we recall notations related to the mirror-twisted Heisenberg-Virasoro algebra \mathcal{D} and collect some known results on highest weight modules and module of intermediates series over \mathcal{D} , including also two important lemmas on tensor product modules for later use. In Section 3, we determine the necessary and sufficient conditions for the tensor product of an irreducible highest weight module and an irreducible module of intermediate series over \mathcal{D} to be irreducible. Consequently we obtain a lot of new irreducible weight modules with infinite dimensional weight spaces over \mathcal{D} , see Theorem 3.11. In Section 4, we show that two such tensor product modules to be isomorphic if and only if the corresponding highest weight modules and modules of intermediate series are isomorphic respectively, see Theorem 4.4. In Section 5, we apply the results established in Section 4 to construct irreducible weight \mathcal{D} -modules with infinite dimensional weight spaces. In these examples, when the tensor product module is not irreducible we also discuss its submodules.

Throughout this paper, we denote by \mathbb{Z} , \mathbb{N} , \mathbb{Z}_+ , \mathbb{C} and \mathbb{C}^* the sets of integers, positive integers, non-negative integers, complex numbers and nonzero complex numbers respectively. All vector spaces and Lie algebras are over \mathbb{C} . We denote by $\mathcal{U}(\mathcal{G})$ the universal enveloping algebra for a Lie algebra \mathcal{G} .

2. NOTATIONS AND PRELIMINARIES

In this section we recall some notations and collect some results for the future convenience.

Definition 2.1. *The* **mirror-twisted Heisenberg-Virasoro algebra** \mathcal{D} *is a Lie algebra with basis* $\{d_m, h_r, \mathbf{c}, \mathbf{l} | m \in \mathbb{Z}, r \in \frac{1}{2} + \mathbb{Z}\}$ defined by the following commutation relations

$$[d_m, d_n] = (m - n)d_{m+n} + \frac{m^3 - m}{12}\delta_{m+n,0}\mathbf{c},$$

$$[d_m, h_r] = -rh_{m+r},$$

$$[h_r, h_s] = r\delta_{r+s,0}\mathbf{l},$$

$$[\mathbf{c}, \mathcal{D}] = [\mathbf{l}, \mathcal{D}] = 0,$$

for $m, n \in \mathbb{Z}, r, s \in \frac{1}{2} + \mathbb{Z}$.

The Lie subalgebra spanned by $\{d_m, \mathbf{c} | m \in \mathbb{Z}\}$ is the Virasoro algebra \mathcal{V} , and the Lie subalgebra spanned by $\{h_r, \mathbf{l} | r \in \frac{1}{2} + \mathbb{Z}\}$ is the twisted Heisenberg algebra \mathcal{H} . Moreover, \mathcal{H} is an ideal of \mathcal{D} and \mathcal{D} is the semi-direct product of \mathcal{V} and \mathcal{H} . It is clear that $\mathcal{D}, \mathcal{V}, \mathcal{H}$ are all \mathbb{Z} -graded Lie algebras, and their universal enveloping algebras are also \mathbb{Z} -graded. The homogeneous spaces of $\mathcal{U}(\mathcal{D})$ are actually the eigenspaces with respect to $-d_0$. We call the eigenvalue of a homogeneous element the **degree**, say $\deg(x) = k$ if $x \in \mathcal{U}(\mathcal{D})_k$ for $k \in \frac{\mathbb{Z}}{2}$.

Note that \mathcal{D} has the triangular decomposition:

$$\mathcal{D} = \mathcal{D}^+ \oplus \mathcal{D}^0 \oplus \mathcal{D}^-,$$

where

$$\mathcal{D}^{\pm} = \bigoplus_{n \in \mathbb{N}} \mathbb{C}d_{\pm n} \oplus \bigoplus_{r \in \frac{1}{2} + \mathbb{Z}_{+}} \mathbb{C}h_{\pm r}, \quad \mathcal{D}^{0} = \mathbb{C}d_{0} \oplus \mathbb{C}\mathbf{c} \oplus \mathbb{C}\mathbf{l}.$$

Similar, \mathcal{V} and \mathcal{H} also have the triangular decompositions. For convenience, we define the following notations for $r, s \in \mathbb{Z}_+, k \in \frac{\mathbb{Z}}{2}, z \in \mathbb{C}$,

$$\mathcal{V}^{(r)} = \bigoplus_{i=r+1}^{\infty} \mathbb{C}d_i + \mathbb{C}\mathbf{c}, \quad \mathcal{D}^{(r,s)} = \bigoplus_{i=r+1}^{\infty} \mathbb{C}d_i \oplus \bigoplus_{j=s}^{\infty} \mathbb{C}h_{\frac{1}{2}+j} + \mathbb{C}\mathbf{c} + \mathbb{C}\mathbf{l},$$
$$\delta_{z \in \mathbb{Z}} = \begin{cases} 1 & \text{if } z \in \mathbb{Z} \\ 0 & \text{if } z \notin \mathbb{Z}, \end{cases} \quad \delta_{z \notin \mathbb{Z}} = \begin{cases} 0 & \text{if } z \in \mathbb{Z} \\ 1 & \text{if } z \notin \mathbb{Z}. \end{cases}$$

Definition 2.2. A \mathcal{D} -module V is called a weight module if $V = \bigoplus_{\lambda \in \mathbb{C}} V_{\lambda}$, where $V_{\lambda} = \{v \in V | d_0 v = \lambda v\}$. The subspace V_{λ} is called the weight space of V corresponding to the weight λ .

Definition 2.3. Let $\mathcal{G} = \bigoplus_{i \in \mathbb{Z}} \mathcal{G}_i$ be a \mathbb{Z} -graded Lie algebra. A \mathcal{G} -module V is called the **restricted** module if for any $v \in V$ there exists $n \in \mathbb{N}$ such that $\mathcal{G}_i v = 0$, for i > n. The category of restricted modules over \mathcal{G} will be denoted as $\mathcal{R}_{\mathcal{G}}$.

Note that if *V* is a \mathcal{V} -module, then *V* can be easily viewed as a \mathcal{D} -module by defining $\mathcal{H}V = 0$, the resulting module is denoted by $V^{\mathcal{D}}$. Thanks to [10], for any $H \in \mathcal{R}_{\mathcal{H}}$ with the action of **l** as a nonzero scalar *l*, we can give *H* a \mathcal{D} -module structure denoted by $H^{\mathcal{D}}$ via the following map

$$d_n \mapsto \frac{1}{2l} \sum_{k \in \mathbb{Z} + \frac{1}{2}} h_{n-k} h_k, \quad \forall n \in \mathbb{Z}, n \neq 0,$$
(2.1)

$$d_0 \mapsto \frac{1}{2l} \sum_{k \in \mathbb{Z} + \frac{1}{2}} h_{-|k|} h_{|k|} + \frac{1}{16},$$
(2.2)

$$h_r \mapsto h_r, \quad \forall r \in \frac{1}{2} + \mathbb{Z}, \quad \mathbf{c} \mapsto 1, \quad \mathbf{l} \mapsto l.$$
 (2.3)

2.1. Modules of intermediate series.

Definition 2.4. A weight D-module V is called a module of **intermediate series** if all its nonzero weight spaces are one-dimensional.

Recall that for all $\alpha, \beta \in \mathbb{C}$ and $\gamma \in \mathbb{C}^*$, $A(\alpha, \beta, \gamma) = \bigoplus_{k \in \frac{1}{2}\mathbb{Z}} \mathbb{C}v_k$ is an irreducible \mathcal{D} -module with the actions defined by

$$d_{m}v_{k} = (\alpha + \beta m - k)v_{m+k},$$

$$h_{r}v_{n} = v_{n+r},$$

$$h_{r}v_{s} = \gamma v_{r+s},$$

$$\mathbf{c}v_{k} = \mathbf{l}v_{k} = \mathbf{0},$$

(2.4)

for $m, n \in \mathbb{Z}$, $r, s \in \frac{1}{2} + \mathbb{Z}$, $k \in \frac{1}{2}\mathbb{Z}$.

Let us recall the \mathcal{V} -module of the intermediate series $A(\alpha, \beta) = \bigoplus_{k \in \mathbb{Z}} \mathbb{C}v_k$:

$$d_m v_k = (\alpha + \beta m - k) v_{m+k}, \quad \mathbf{c} v_k = 0, \tag{2.5}$$

for $m, k \in \mathbb{Z}$. We denote by $A'(\alpha, \beta)$ the unique nontrivial irreducible subquotient module of $A(\alpha, \beta)$. It is clear that $A(\alpha, \beta)^{\mathcal{D}}$ is an irreducible \mathcal{D} -module if and only if $A(\alpha, \beta)$ is an irreducible \mathcal{V} -module. Moreover, we have the following well-known results:

Lemma 2.5. (1) The \mathcal{V} -module $A(\alpha, \beta)$ is isomorphic to \mathcal{V} -module $A(\alpha + m, \beta)$ for $m \in \mathbb{Z}$.

- (2) The \mathcal{V} -module $A(\alpha, \beta)$ is irreducible if and only if $\alpha \notin \mathbb{Z}$ or $\beta \neq 0, -1$.
- (3) If $\alpha \in \mathbb{Z}$, $\beta = 0$, then $A(\alpha, \beta)$ has a unique nontrivial irreducible quotient module.
- (4) If $\alpha \in \mathbb{Z}$, $\beta = -1$, then $A(\alpha, \beta)$ has a unique nontrivial irreducible submodule.
- (5) $A'(\alpha, 0) \cong A'(\alpha, -1)$ for $\alpha \in \mathbb{C}$.

We have the classification for all irreducible modules of intermediate series over \mathcal{D} .

Lemma 2.6. (cf. [17, Theorem 3.9]) If V is an irreducible module of intermediate series over \mathcal{D} , then $V \cong A(\alpha, \beta, \gamma)$ or $V \cong A'(\alpha, \beta)^{\mathcal{D}}$ for some $\alpha, \beta \in \mathbb{C}$ and $\gamma \in \mathbb{C}^*$.

2.2. Verma modules.

Definition 2.7. A weight \mathcal{D} -module V is called a **highest weight module** with highest weight $\lambda \in \mathbb{C}$, if there exists a nonzero weight vector $v \in V_{\lambda}$ such that V is generated by v as \mathcal{D} -module and $\mathcal{D}^+v = 0$.

Definition 2.8. For $(c, h, l) \in \mathbb{C}^3$, let $\mathbb{C}\mathbf{1}$ be a $(\mathcal{D}^0 \oplus \mathcal{D}^+)$ -module with

$$d_0 \mathbf{1} = h \mathbf{1}, \quad \mathbf{c} \mathbf{1} = c \mathbf{1}, \quad \mathbf{l} \cdot \mathbf{1} = l \mathbf{1}, \quad \mathcal{D}^+ \mathbf{1} = 0.$$

The **Verma module** *over* \mathcal{D} *is defined by*

$$M(c, h, l) = U(\mathcal{D}) \otimes_{U(\mathcal{D}^0 \oplus \mathcal{D}^+)} \mathbb{C}\mathbf{1}$$

Let J(c, h, l) be the unique maximal submodule of M(c, h, l). Then we have the irreducible highest weight module L(c, h, l) = M(c, h, l)/J(c, h, l). We will use the **1** and **1** to denote the highest weight vectors of M(c, h, l) and L(c, h, l) respectively. Similarly, we have $M_{\mathcal{V}}(c, h), L_{\mathcal{V}}(c, h) = M_{\mathcal{V}}(c, h)/J_{\mathcal{V}}(c, h)$ for the Virasoro algebra \mathcal{V} , and $M_{\mathcal{H}}(l), L_{\mathcal{H}}(l) = M_{\mathcal{H}}(l)/J_{\mathcal{H}}(l)$ for the twisted Heisenberg algebra \mathcal{H} . It is clear that they are all restricted modules. Moreover, we have the following results about highest weight modules over \mathcal{D} .

Lemma 2.9. (cf. [17, Proposition 5.1]) Let $c, h \in \mathbb{C}$ and $l \in \mathbb{C}^*$. Then we have \mathcal{D} -module isomorphisms

$$\pi_1: M(c,h,l) \to M_{\mathcal{V}}(c-1,h-\frac{1}{16})^{\mathbb{D}} \otimes M_{\mathcal{H}}(l)^{\mathbb{D}},$$

$$\pi_2: L(c,h,l) \to L_{\mathcal{V}}(c-1,h-\frac{1}{16})^{\mathbb{D}} \otimes M_{\mathcal{H}}(l)^{\mathbb{D}}.$$

Lemma 2.10. (cf. [17, Corollary 5.2]) Let $c, h \in \mathbb{C}$ and $l \in \mathbb{C}^*$. Then the Verma module M(c, h, l) is irreducible if and only if $\phi_{r,s}(c-1, h-\frac{1}{16}) \neq 0$ for all $r, s \in \mathbb{Z}_+$, where

$$\begin{split} \phi_{r,s}(c,h) &= h - \frac{1}{48}(13-c)(r^2+s^2) \\ &- \frac{1}{48} \Big(\sqrt{(c-1)(c-25)}(r^2-s^2) - 24rs - 2 + 2c \Big). \end{split}$$

We conclude this section by recalling two familiar results for later use.

Lemma 2.11. (cf. [18, Lemma 6]) Let \mathcal{G} be a Lie algebra over \mathbb{C} with a countable basis, and V be an irreducible \mathcal{G} -module. For any $n \in \mathbb{N}$ and any linearly independent subset $\{v_1, v_2, \dots, v_n\} \subset V$, and any subset $\{v'_1, \dots, v'_n\} \subset V$, there exists some $u \in \mathcal{U}(\mathcal{G})$, such that

$$uv_i = v'_i, \forall i = 1, 2, \cdots, n.$$

Lemma 2.12. (cf. [18, Theorem 7]) Let \mathcal{G} be a Lie algebra over \mathbb{C} with a countable basis, and V_1, V_2 be \mathcal{G} -modules. Suppose that one of the following conditions holds:

- (1) The module V_1 is irreducible and $Ann_{\mathcal{G}}(v) + Ann_{\mathcal{G}}(S) = \mathcal{G}$ for all $v \in V_1$ and all finite subsets $S \subseteq V_2$;
- (2) For any finite subset $S \subseteq V_2$, V_1 is an irreducible $Ann_{\mathcal{G}}(S)$ -module.

Then the following statements hold.

- (a) Any submodule of $V_1 \otimes V_2$ is of the form $V_1 \otimes V'_2$ for a submodule V'_2 of V_2 ;
- (b) If V_1, V_2 are irreducible, then $V_1 \otimes V_2$ is irreducible.

3. IRREDUCIBILITY OF TENSOR PRODUCT WEIGHT MODULES

From Section 2, we know that all irreducible modules of intermediate series over \mathcal{D} are $A'(\alpha,\beta)^{\mathcal{D}}$ or $A(\alpha,\beta,\gamma)$ where $\alpha,\beta \in \mathbb{C}, \gamma \in \mathbb{C}^*$. In this section, we will determine the necessary and sufficient conditions for the tensor product modules $L(c, h, l) \otimes A'(\alpha,\beta)^{\mathcal{D}}$, $L(c, h, l) \otimes A(\alpha,\beta,\gamma)$ to be irreducible for any $c, h, l, \alpha, \beta \in \mathbb{C}, \gamma \in \mathbb{C}^*$.

Lemma 3.1. Suppose that V, W are \mathcal{V} -modules, and $H, K \in \mathbb{R}_{\mathcal{H}}$ are irreducible with nonzero action of **l**. Then

- (1) any D-submodule of V^D ⊗ H^D is of the form (V')^D ⊗ H^D for some V-submodule V' of V. In particular, V^D ⊗ H^D is an irreducible D-module if and only if V is an irreducible V-module;
- (2) $V^{\mathbb{D}} \otimes H^{\mathbb{D}} \cong W^{\mathbb{D}} \otimes K^{\mathbb{D}}$ if and only if $V \cong W$ and $H \cong K$.

Proof. (1) Note that $\mathcal{H} \subseteq Ann_{\mathcal{D}}(V^{\mathcal{D}})$, and $H^{\mathcal{D}}$ is an irreducible \mathcal{H} -module. Thus the result follows from Lemma 2.12.

(2) It is clear for the sufficiency.

Now we show the necessity. Suppose that $\varphi : V^{\mathcal{D}} \otimes H^{\mathcal{D}} \to W^{\mathcal{D}} \otimes K^{\mathcal{D}}$ is a \mathcal{D} -module isomorphism. Let's fix a nonzero $x_0 \in H$. For any $0 \neq v \in \mathcal{V}$, we write $\varphi(v \otimes x_0) = \sum_{i=1}^{t} w_i \otimes y_i$ with minimal *t*. We know that y_i 's are linearly independent. Then there exists some $u_0 \in \mathcal{U}(\mathcal{H})$ such that $u_0y_i = \delta_{1,i}y_i$, $i = 1, 2, \cdots, t$ by Lemma 2.11. Therefore,

$$\varphi(v \otimes uu_0 x_0) = \varphi(uu_0(v \otimes x_0)) = w_1 \otimes uy_1, \forall u \in \mathcal{U}(\mathcal{H}).$$

Denote $x_1 = u_0 x_0$. Then we have $\varphi(v \otimes u x_1) = w_1 \otimes u y_1$, $\forall u \in \mathcal{U}(\mathcal{H})$. Notice $\mathcal{U}(\mathcal{H}) x_1 = H$, $\mathcal{U}(\mathcal{H}) y_1 = K$, since H, K are irreducible. Define $\varphi_v : H \to K$ by $\varphi_v(ux_1) = uy_1$, for all $u \in \mathcal{U}(\mathcal{H})$. It is easy to see that φ_v is a $\mathcal{U}(\mathcal{H})$ -module isomorphism. That shows $H \cong K$ as \mathcal{H} -modules.

At the moment, we can assume that H = K. Then φ is an isomorphism from $V^{\mathbb{D}} \otimes H^{\mathbb{D}}$ to $W^{\mathbb{D}} \otimes H^{\mathbb{D}}$, and φ_{v} is an automorphism of H. In fact, φ_{v} is a scalar denoted by c_{v} , since H is irreducible and $\mathcal{U}(\mathcal{H})$ is countably generated. Then for any $v \in V, x \in H$,

$$\varphi(v\otimes x)=c_vw_v\otimes x,$$

for some $w_v \in H$. Define the map $\varpi : V \to W$ by $v \to c_v w_v$. It's clear that ϖ is bijective. Furthermore, we have that $d_n \varpi(v) \otimes x = \varpi(d_n v) \otimes x$, since $d_n \varphi(v \otimes x) = \varphi(d_n(v \otimes x))$, for all $v \in V, x \in H$. Therefore, $d_n \varpi(v) = \varpi(d_n v)$, that means ϖ is a \mathcal{V} -module isomorphism. So, $V \cong W$ as \mathcal{V} -modules. We complete the proof. **Proposition 3.2.** Let $c, h, l, \alpha, \beta \in \mathbb{C}, \gamma \in \mathbb{C}^*$.

- (1) Suppose $l \neq 0$. Then $L(c, h, l) \otimes A'(\alpha, \beta)^{\mathbb{D}}$ is an irreducible \mathbb{D} -module if and only if $L_{\mathcal{V}}(c-1, h-\frac{1}{16}) \otimes A'(\alpha, \beta)$ is an irreducible \mathcal{V} -module.
- (2) Suppose l = 0. Then $L(c, h, l) \otimes A'(\alpha, \beta)^{\mathbb{D}}$ is an irreducible \mathbb{D} -module if and only if $L_{\mathcal{V}}(c, h) \otimes A'(\alpha, \beta)$ is an irreducible \mathcal{V} -module.
- (3) Suppose l = 0, then $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$ is irreducible.

Proof. (1) From Lemma 2.9 we know that

$$L(c,h,l) \cong L_{\mathcal{V}}(c-1,h-\frac{1}{16})^{\mathcal{D}} \otimes M_{\mathcal{H}}(l)^{\mathcal{D}}$$

So we have

$$L(c,h,l) \otimes A'(\alpha,\beta)^{\mathcal{D}} \cong M_{\mathcal{H}}(l)^{\mathcal{D}} \otimes (L_{\mathcal{V}}(c-1,h-\frac{1}{16}) \otimes A'(\alpha,\beta))^{\mathcal{D}}$$

It is clear that $M_{\mathcal{H}}(l) \in \mathcal{R}_{\mathcal{H}}$ is irreducible when $l \neq 0$. Therefore, the result follows from Lemma 3.1.

(2) Since l = 0, we see that $L(c, h, l) \cong L_{\mathcal{V}}(c, h)^{\mathcal{D}}$ as \mathcal{D} -modules. Thus

~

$$L(c,h,l) \otimes A'(\alpha,\beta)^{\mathcal{D}} \cong L_{\mathcal{V}}(c,h)^{\mathcal{D}} \otimes A'(\alpha,\beta)^{\mathcal{D}} \cong (L_{\mathcal{V}}(c,h) \otimes A'(\alpha,\beta))^{\mathcal{D}}.$$

Then the result is clear.

(3) Since l = 0, we have $L(c, h, l) \cong L_{\mathcal{V}}(c, h)^{\mathcal{D}}$ as \mathcal{D} -modules. Notice that for any finite subset $S \subseteq L_{\mathcal{V}}(c, h)^{\mathcal{D}}$, there exists some *r* such that $\mathcal{V}^{(r)} + \mathcal{H} \subseteq Ann_{\mathcal{D}}(S)$. Moreover, we see that the module $A(\alpha, \beta, \gamma)$ defined in (2.4) is irreducible as $(\mathcal{V}^{(r)} + \mathcal{H})$ -module by the following two observations:

- (a) Each v_k can generate $A(\alpha, \beta, \gamma)$ as an \mathcal{H} -submodule;
- (b) $h_{\frac{1}{2}}h_{-m-\frac{1}{2}}d_mv_k = \gamma(\alpha k + m\beta)v_k$ for any m > r.

So the result follows from Lemma 2.12.

Remark 3.3. In the above theorem, we reduce the irreducibility of \mathbb{D} -module $L(c, h, l) \otimes A'(\alpha, \beta)^{\mathbb{D}}$ to the irreducibility of the \mathcal{V} -module $L_{\mathcal{V}}(c, h) \otimes A'(\alpha, \beta)$ or $L_{\mathcal{V}}(c - 1, h - \frac{1}{16}) \otimes A'(\alpha, \beta)$. Thanks to [5], the irreducibility of the \mathcal{V} -module $L_{\mathcal{V}}(c, h) \otimes A'(\alpha, \beta)$ has been completely determined for $c, h, \alpha, \beta \in \mathbb{C}$.

Now we have only the most difficult case left: $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$ with $l \neq 0$. To determine the necessary and sufficient conditions for this tensor product module to be irreducible we will break the arguments into several lemmas. From now on we assume that $c, h, \alpha, \beta \in \mathbb{C}, l, \gamma \in \mathbb{C}^*$, and we will use the definition for $A(\alpha, \beta, \gamma)$ in (2.4).

Lemma 3.4. (a). For any $k \in \frac{\mathbb{Z}}{2}$,

$$\mathbf{1} \otimes v_k \notin U(\mathcal{D})(\mathbf{1} \otimes v_{k+\frac{1}{2}}) \subseteq M(c,h,l) \otimes A(\alpha,\beta,\gamma).$$

(b). The \mathbb{D} -module $M(c, h, l) \otimes A(\alpha, \beta, \gamma)$ is not irreducible.

Proof. (a). Note that for any $k \in \frac{\mathbb{Z}}{2}$,

$$U(\mathcal{D})(\mathbf{1}\otimes v_{k+\frac{1}{2}})=U(\mathcal{D}^{-})U(\mathcal{D}^{+})(\mathbf{1}\otimes v_{k+\frac{1}{2}})\subseteq \sum_{i\in\mathbb{Z}_{+}}U(\mathcal{D}^{-})(\mathbf{1}\otimes v_{k+\frac{1}{2}+\frac{i}{2}}).$$

So, it is sufficient to show that $\mathbf{1} \otimes v_k \notin \sum_{i \in \mathbb{Z}_+} U(\mathcal{D}^-)(\mathbf{1} \otimes v_{k+\frac{1}{2}+\frac{i}{2}}).$

Assume that $\mathbf{1} \otimes v_k \in \sum_{i \in \mathbb{Z}_+} U(\mathcal{D}^-)(\mathbf{1} \otimes v_{k+\frac{1}{2}+\frac{i}{2}})$. Then by PBW Theorem we know that

$$\mathbf{1} \otimes v_k = \sum_{i=1}^r u_{-\frac{i}{2}} (\mathbf{1} \otimes v_{k+\frac{i}{2}}),$$

where $u_{-\frac{i}{2}} \in U(\mathcal{D}^{-})_{-\frac{i}{2}}$ with $u_{-\frac{r}{2}} \neq 0$. Thus we have

$$\sum_{i=1}^{\prime} u_{-\frac{i}{2}}(\mathbf{1} \otimes v_{k+\frac{i}{2}}) - \mathbf{1} \otimes v_k = 0,$$

i.e.,

$$\sum_{i=1}^{r} (u_{-\frac{i}{2}} \mathbf{1} \otimes v_{k+\frac{i}{2}}) + \sum_{i=1}^{r} (\mathbf{1} \otimes (u_{-\frac{i}{2}} v_{k+\frac{i}{2}}) - \mathbf{1} \otimes v_{k} = 0.$$

This is impossible since the nonzero vector $u_{-\frac{r}{2}} \mathbf{1} \otimes v_{k+\frac{r}{2}}$ cannot be in the span of the other vectors in the above formula.

(b) follows directly from (a).

Lemma 3.5. Let M be a nonzero \mathcal{D} -submodule of $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$.

- (a). There exists $k \in \frac{\mathbb{Z}}{2}$ such that $\overline{\mathbf{1}} \otimes v_k \in M$.
- (b). There exists $k \in \frac{\mathbb{Z}}{2}$ such that $\overline{\mathbf{1}} \otimes v_i \in M$ for all $i \in \frac{1}{2}\mathbb{Z}$ with $i \ge k$.

Proof. (a). Since M is a weight \mathcal{D} -module, we can take a nonzero weight vector

$$w = \sum_{i=0}^{s} w_{-\frac{i}{2}} \otimes v_{k+\frac{i}{2}} \in M, \text{ with } w_{0} = \bar{\mathbf{1}}, k \in \frac{\mathbb{Z}}{2}, w_{-\frac{i}{2}} \in U(\mathcal{D}^{-})_{-\frac{i}{2}} \bar{\mathbf{1}}$$

There is an $n \in \mathbb{N}$ such that $d_j w_{-\frac{i}{2}} = h_{\frac{1}{2}+j} w_{-\frac{i}{2}} = 0$ for all $j \ge n, i = 0, 1, \dots, s$. Since $A(\alpha, \beta, \gamma)$ is an irreducible $(\mathcal{V}^{(n)} + \mathcal{H})$ -module, by Lemma 2.11 we may take some $u \in U(\mathcal{V}^{(n)} + \mathcal{H})$ such that

$$uv_{k+\frac{i}{2}} = \delta_{0,\frac{i}{2}}v_0, \forall i = 0, 1, \dots, s.$$

We can write $u = \sum_i u_i u'_i$ with $u_i \in U(\mathcal{H})$ and $u'_i \in U(\mathcal{V}^{(n)})$ by PBW Theorem. For any $v \in A(\alpha, \beta, \gamma)$, we have

$$h_{j}h_{i}v = h_{\frac{i+j}{2}}h_{\frac{i+j}{2}}v, \quad \text{if} \quad i+j \in 2\mathbb{Z}+1,$$

$$h_{j}h_{i}v = h_{\frac{i+j}{2}-\frac{1}{2}}h_{\frac{i+j}{2}+\frac{1}{2}}v, \quad \text{if} \quad i+j \in 2\mathbb{Z}.$$

Take a sufficiently large *r*. By repeatedly and properly replacing $h_j h_i$ with $h_{\frac{i+j}{2}} h_{\frac{i+j}{2}-\frac{1}{2}} h_{\frac{i+j}{2}+\frac{1}{2}}$ in the expression of $h_r u$ we can result in an element $u' \in U(\mathcal{D}^{(n,n)})$ such that

$$u'v_{k+\frac{i}{2}} = h_r uv_{k+\frac{i}{2}} = \delta_{0,\frac{i}{2}}v_r, \forall i = 0, 1, \dots, s.$$

Since $u'w_{-\frac{i}{2}} = 0$ for all i = 0, 1, ..., s, we deduce that $\overline{\mathbf{1}} \otimes v_r = u'w \in M$.

(b). This follows by applying h_i to $\overline{\mathbf{1}} \otimes v_k \in M$ in (a) for all $i \in \frac{\mathbb{N}}{2}$.

Now, we need to introduce the "shifting technique". For any highest weight \mathcal{D} -module V with highest weight vector **v**, it is easy to check that $V \otimes \mathbb{C}[y^{\pm \frac{1}{2}}]$ can be endowed a \mathcal{D} -module structure by following actions

$$d_{n}(P\mathbf{v} \otimes y^{i}) = (d_{n} + \alpha + \beta n + \deg(P) - i)P\mathbf{v} \otimes y^{i+n},$$

$$h_{r}(P\mathbf{v} \otimes y^{i}) = (h_{r} + \delta_{i-\deg(P)\in\mathbb{Z}} + \gamma\delta_{i-\deg(P)\notin\mathbb{Z}})P\mathbf{v} \otimes y^{i+r},$$

$$\mathbf{c}(P\mathbf{v} \otimes y^{i}) = c(P\mathbf{v} \otimes y^{i}), \quad \mathbf{l}(P\mathbf{v} \otimes y^{i}) = l(P\mathbf{v} \otimes y^{i}),$$

(3.1)

for $n \in \mathbb{Z}, r \in \frac{1}{2} + \mathbb{Z}, i \in \frac{\mathbb{Z}}{2}$ and homogeneous element $P \in \mathcal{U}(\mathcal{D}^-)$. Then it is not hard to see that

$$V \otimes A(\alpha, \beta, \gamma) \cong V \otimes \mathbb{C}[y^{\pm \frac{1}{2}}]$$

as \mathcal{D} -modules. Specially, we have

$$\begin{split} M(c,h,l) \otimes A(\alpha,\beta,\gamma) &\cong M(c,h,l) \otimes \mathbb{C}[y^{\pm \frac{1}{2}}] \\ L(c,h,l) \otimes A(\alpha,\beta,\gamma) &\cong L(c,h,l) \otimes \mathbb{C}[y^{\pm \frac{1}{2}}]. \end{split}$$

Henceforth we may view $M(c, h, l) \otimes A(\alpha, \beta, \gamma)$, $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$ as $M(c, h, l) \otimes \mathbb{C}[y^{\pm \frac{1}{2}}]$, $L(c, h, l) \otimes \mathbb{C}[y^{$

$$M(c,h,l) \otimes y^k = \{ v \in M(c,h,l) \otimes \mathbb{C}[y^{\pm \frac{1}{2}}] \mid d_0 v = (\alpha + h - k)v \},$$

thus $M(c, h, l) \otimes \mathbb{C}[y^{\pm \frac{1}{2}}] = \bigoplus_{i \in \frac{\mathbb{Z}}{2}} (M(c, h, l) \otimes y^i)$. Denote

$$\begin{split} W^{(k)} &= \sum_{i \in \frac{\mathbb{Z}_+}{2}} \mathfrak{U}(\mathcal{D})(\mathbf{1} \otimes y^{k+i}) \subseteq M(c,h,l) \otimes \mathbb{C}[y^{\pm \frac{1}{2}}], \\ W^{(k)}_n &= W^{(k)} \cap (M(c,h,l) \otimes y^n), \forall n \in \frac{\mathbb{Z}}{2}. \end{split}$$

We have following observations:

Lemma 3.6. (1) $W^{(k)} = \sum_{i \in \frac{\mathbb{Z}_+}{2}} U(\mathcal{D}^-)(\mathbf{1} \otimes y^{k+i}).$ (2) $W^{(k)} \supseteq \bigoplus_{k \le i \in \frac{\mathbb{Z}}{2}} M(c, h, l) \otimes y^i.$ (3) $M(c, h, l) \otimes y^{k-\frac{1}{2}} = W^{(k)}_{k-\frac{1}{2}} \oplus \mathbb{C}(\mathbf{1} \otimes y^{k-\frac{1}{2}}).$ (4) If $P \in U(\mathcal{D}^-)$ is homogeneous such that

$$P\mathbf{1} \otimes y^{k-\frac{1}{2}} \in W_{k-\frac{1}{2}}^{(k)},$$

then
$$(U(\mathcal{D}^{-})P\mathbf{1}) \otimes y^{k-\frac{1}{2}} \subseteq W_{k-\frac{1}{2}}^{(k)}$$

Proof. (1) This follows from the following computations:

$$W^{(k)} = \sum_{i \in \frac{\mathbb{Z}_+}{2}} \mathcal{U}(\mathcal{D})(\mathbf{1} \otimes y^{k+i}) = \sum_{i \in \frac{\mathbb{Z}_+}{2}} \mathcal{U}(\mathcal{D}^-)\mathcal{U}(\mathcal{D}^+)(\mathbf{1} \otimes y^{k+i})$$
$$= \sum_{i \in \frac{\mathbb{Z}_+}{2}} \mathcal{U}(\mathcal{D}^-)(\sum_{j \in \frac{\mathbb{Z}_+}{2}} (\mathbf{1} \otimes y^{k+i+j})) = \sum_{i \in \frac{\mathbb{Z}_+}{2}} \mathcal{U}(\mathcal{D}^-)(\mathbf{1} \otimes y^{k+i}).$$

(2) By PBW Theorem, each nonzero element of $M(c, h, l) \otimes y^i$ can be written as linear combinations of vectors

$$v_{s,t} = h_{-p_s} \dots h_{-p_1} d_{-q_t} \dots d_{-q_1} \mathbf{1} \otimes y^t,$$

where $q_j \in \mathbb{N}, p_{j'} \in \frac{1}{2} + \mathbb{Z}_+$ for $1 \le j \le t, 1 \le j' \le s$.

Now, we show the result by induction on s + t. If s + t = 0, then $v_{s,t} = \mathbf{1} \otimes y^i \in W^{(k)}, \forall k \le i \in \frac{\mathbb{Z}}{2}$ is trivial. Suppose that the result is right when s + t = m. For s + t = m + 1, we may assume that

$$v_{s,t} = h_{-r}Q\mathbf{1} \otimes y^t$$
 where $Q \in U(\mathcal{D}^-), r > 0$, or

$$v_{s,t} = d_{-n}Q\mathbf{1} \otimes y^t$$
, where $Q \in U(\mathcal{V}^-), n > 0$.

Then by equations 3.1 and inductive hypothesis on Q, we have

$$v_{s,t} = h_{-r}Q\mathbf{1} \otimes y^{i}$$

= $h_{-r}(Q\mathbf{1} \otimes y^{i+r}) - (\delta_{i+r-\deg(Q)\in\mathbb{Z}} + \gamma\delta_{i+r-\deg(Q)\notin\mathbb{Z}})Q\mathbf{1} \otimes y^{i} \in W^{(k)}, \text{ or }$
 $v_{s,t} = d_{-n}Q\mathbf{1} \otimes y^{i}$
= $d_{-n}(Q\mathbf{1} \otimes y^{i+n}) - (\alpha - \beta n + \deg(Q) - i - n)Q\mathbf{1} \otimes y^{i} \in W^{(k)}.$

Thus $M(c, h, l) \otimes y^i \in W^{(k)}, \forall k \le i \in \frac{\mathbb{Z}}{2}$.

(3) $W_{k-\frac{1}{2}}^{(k)} \oplus \mathbb{C}(\mathbf{1} \otimes y^{k-\frac{1}{2}}) \subseteq M(c, h, l) \otimes y^{k-\frac{1}{2}}$ is clear by definition. $W_{k-\frac{1}{2}}^{(k)} \cap \mathbb{C}(\mathbf{1} \otimes y^{k-\frac{1}{2}}) = 0$ follows from the Lemma 3.4.

Now we show that $M(c, h, l) \otimes y^{k-\frac{1}{2}} \subseteq W_{k-\frac{1}{2}}^{(k)} \oplus \mathbb{C}(\mathbf{1} \otimes y^{k-\frac{1}{2}})$. Note that every nonzero element of $M(c, h, l) \otimes y^{k-\frac{1}{2}}$ can be written as linear combinations of vectors

$$v_{s,t} = h_{-p_s} \dots h_{-p_1} d_{-q_t} \dots d_{-q_1} \mathbf{1} \otimes y^{k-\frac{1}{2}},$$

where $q_j \in \mathbb{N}$, $p_{j'} \in \frac{1}{2} + \mathbb{Z}_+$ for $1 \le j \le t$, $1 \le j' \le s$ by PBW Theorem. By induction on s+t which is similar to (2), we obtain $v_{s,t} \in W^{(k)} \oplus \mathbb{C}(\mathbf{1} \otimes y^{k-\frac{1}{2}})$. Thus $M(c, h, l) \otimes y^{k-\frac{1}{2}} \subseteq W^{(k)} \oplus \mathbb{C}(\mathbf{1} \otimes y^{k-\frac{1}{2}})$. Then $M(c, h, l) \otimes y^{k-\frac{1}{2}} \subseteq W^{(k)}_{k-\frac{1}{2}} \oplus \mathbb{C}(\mathbf{1} \otimes y^{k-\frac{1}{2}})$.

(4) By PBW Theorem, every nonzero element of $(U(\mathcal{D}^-)P\mathbf{1}) \otimes y^{k-\frac{1}{2}}$ can be written as linear combinations of vectors

$$v_{s,t} = h_{-p_s} \dots h_{-p_1} d_{-q_t} \dots d_{-q_1} P \mathbf{1} \otimes y^{k-\frac{1}{2}},$$

where $q_j \in \mathbb{N}, p_{j'} \in \frac{1}{2} + \mathbb{Z}_+$ for $1 \le j \le t, 1 \le j' \le s$. By induction on s + t which is similar to (2), we obtain $v_{s,t} \in W^{(k)}$. Thus $(U(\mathcal{D}^-)P\mathbf{1}) \otimes y^{k-\frac{1}{2}} \in W^{(k)}_{k-\frac{1}{2}}$.

Now, similar to the definition of φ_n in [5], we want to define the linear map $\rho_n : \mathcal{U}(\mathcal{D}^-) \to \mathbb{C}$, for any $n \in \frac{\mathbb{Z}}{2}$. Let $T(\mathcal{D}^-)$ denote the tensor algebra of \mathcal{D}^- , i.e.,

$$T(\mathcal{D}^{-}) = \mathbb{C} \oplus (\mathcal{D}^{-}) \oplus (\mathcal{D}^{-} \otimes \mathcal{D}^{-}) \oplus \cdots,$$

which is a free associative algebra. For any $k \in \mathbb{N}$, define $\deg(d_{-k}) = -k$, $\deg(h_{\frac{1}{2}-k}) = \frac{1}{2} - k$ and $\deg(1) = 0$, then $T(\mathcal{D}^-)$ is a $\frac{\mathbb{Z}}{2}$ -graded algebra. For any $n \in \frac{\mathbb{Z}}{2}, \alpha, \beta \in \mathbb{C}$ and $\gamma \in \mathbb{C}^*$, we may

inductively define the linear map $\overline{\rho_n}$: $T(\mathcal{D}^-) \to \mathbb{C}$ as follows:

$$\overline{\rho_n}(1) = 1,$$

$$\overline{\rho_n}(h_{-r}u) = -(\delta_{n+r+k\in\mathbb{Z}} + \gamma\delta_{n+r+k\notin\mathbb{Z}})\overline{\rho_n}(u),$$

$$\overline{\rho_n}(d_{-i}u) = -(\alpha - \beta i - k - i - n)\overline{\rho_n}(u),$$

(3.2)

where r, i > 0 and u is an arbitrary homogeneous element of degree -k in $T(\mathcal{D}^-)$. Note that $\mathcal{U}(\mathcal{D}^-)$ is the quotient algebra $T(\mathcal{D}^-)/I$ where I is the two-sided ideal generated by

$$d_{-i}d_{-j} - d_{-j}d_{-i} - (j-i)d_{-i-j}$$

$$d_{-i}h_{-r} - h_{-r}d_{-i} - rh_{-i-r},$$

$$h_{-r}h_{-r'} - h_{-r'}h_{-r},$$

for $i, j \in \mathbb{N}, r, r' \in \frac{1}{2} + \mathbb{Z}_+$. It is not hard to see that $I \subseteq ker(\overline{\rho_n})$. So $\overline{\rho_n}$ induces a linear map $\mathcal{U}(\mathcal{D}^-) \to \mathbb{C}$, denoted by ρ_n . Now we have the following lemma.

Lemma 3.7. Let $P \in \mathcal{U}(\mathcal{D}^-), n \in \frac{\mathbb{Z}}{2}$. Then

(1) $P\mathbf{1} \otimes y^n \equiv \rho_n(P)\mathbf{1} \otimes y^n \pmod{W^{(n+\frac{1}{2})}};$ (2) $P\mathbf{1} \otimes y^n \in W^{(n+\frac{1}{2})}$ if and only if $\rho_n(P) = 0.$

Proof. (1) It is enough to show it for homogeneous *P* by induction on deg(*P*). If deg(*P*) = 0, i.e. $P \in \mathbb{C}$, then $P\mathbf{1} \otimes y^n \equiv \rho_n(P)\mathbf{1} \otimes y^n \pmod{W^{(n+\frac{1}{2})}}$ is trivial. Suppose $Q \in \mathcal{U}(\mathcal{D}^-)$ is a homogeneous element and the result holds for this *Q*. For $k \in \mathbb{N}$, $r \in \frac{1}{2} + \mathbb{Z}_+$, using Lemma 3.6 we have

$$d_{-k}(Q\mathbf{1} \otimes y^{n+k}) = (d_{-k} + \alpha - \beta k + \deg(Q) - n - k)Q\mathbf{1} \otimes y^n \in W^{(n+\frac{1}{2})},$$

$$h_{-r}(Q\mathbf{1} \otimes y^{n+r}) = (h_{-r} + \delta_{n+r-\deg(Q)\in\mathbb{Z}} + \gamma \delta_{n+r-\deg(Q)\notin\mathbb{Z}})Q\mathbf{1} \otimes y^n \in W^{(n+\frac{1}{2})}$$

By equations (3.2), these indicate that

$$d_{-k}Q\mathbf{1} \otimes y^{n} \equiv -(\alpha - \beta k + \deg(Q) - n - k)Q\mathbf{1} \otimes y^{n}$$

$$\equiv -(\alpha - \beta k + \deg(Q) - n - k)\rho_{n}(Q)\mathbf{1} \otimes y^{n}$$

$$\equiv \rho_{n}(d_{-k}Q)\mathbf{1} \otimes y^{n} \pmod{W^{(n+\frac{1}{2})}};$$

$$h_{-r}Q\mathbf{1} \otimes y^{n} \equiv -(\delta_{n+r-\deg(Q)\in\mathbb{Z}} + \gamma\delta_{n+r-\deg(Q)\notin\mathbb{Z}})Q\mathbf{1} \otimes y^{n}$$

$$\equiv -(\delta_{n+r-\deg(Q)\in\mathbb{Z}} + \gamma\delta_{n+r-\deg(Q)\notin\mathbb{Z}})\rho_{n}(Q)\mathbf{1} \otimes y^{n}$$

$$\equiv \rho_{n}(h_{-r}Q)\mathbf{1} \otimes y^{n} \pmod{W^{(n+\frac{1}{2})}}.$$

Then the result follows from induction on deg(Q).

(2) It follows from (1) and Lemma 3.4.

Remark 3.8. For the Verma module $M_{\mathcal{V}}(c,h)$ over \mathcal{V} , it is well-known that there exist two homogeneous elements $P_1, P_2 \in U(\mathcal{V}^-)$ such that $U(\mathcal{V}^-)P_1w_1 + U(\mathcal{V}^-)P_2w_1$ is the unique maximal proper submodule of $M_{\mathcal{V}}(c,h)$, where P_1, P_2 are allowed to be zero and w_1 is the highest weight vector in $M_{\mathcal{V}}(c,h)$.

Lemma 3.9. Let $c, h \in \mathbb{C}, l \in \mathbb{C}^*$. There exist two homogeneous elements $Q_1, Q_2 \in \mathcal{U}(\mathbb{D}^-)$ such that the unique maximal proper submodule J(c, h, l) of M(c, h, l) is generated by $Q_1\mathbf{1}, Q_2\mathbf{1}$, where Q_1, Q_2 are allowed to be zero.

Proof. We consider the D-module isomorphism in Lemma 2.9

$$\pi_1: M(c,h,l) \to M_{\mathcal{V}}(c-1,h-\frac{1}{16})^{\mathcal{D}} \otimes M_{\mathcal{H}}(l)^{\mathcal{D}}.$$

We know that the unique maximal proper submodule $J_{\mathcal{V}}(c,h)$ of $M_{\mathcal{V}}(c-1,h-\frac{1}{16})$ is generated by two singular weight vectors P_1w_1, P_2w_1 from the above remark, where w_1 is the highest weight vector of $M_{\mathcal{V}}(c-1,h-\frac{1}{16})$ and $P_1, P_2 \in U(\mathcal{V}^-)$. Then using Lemma 3.1 we know that the unique maximal proper submodule of $M_{\mathcal{V}}(c-1,h-\frac{1}{16})^{\mathcal{D}} \otimes M_{\mathcal{H}}(l)^{\mathcal{D}}$ is $J_{\mathcal{V}}(c,h) \otimes M_{\mathcal{H}}(l)^{\mathcal{D}}$ which is generated by $P_1w_1 \otimes w_2, P_2w_1 \otimes w_2$ as a \mathcal{D} -submodule, where w_2 is the highest weight vector of $M_{\mathcal{H}}(l)$. Then $J(c,h,l) = \pi^{-1}(J_{\mathcal{V}}(c,h) \otimes M_{\mathcal{H}}(l)^{\mathcal{D}})$ which is generated by $\pi_1^{-1}(P_1w_1 \otimes w_2), \pi_1^{-1}(P_2w_1 \otimes w_2)$ as a \mathcal{D} -submodule. Thus there exist two homogeneous elements $Q_1, Q_2 \in \mathcal{U}(\mathcal{D}^-)$ such that $Q_1\mathbf{1} = \pi_1^{-1}(P_1w_1 \otimes w_2), Q_2\mathbf{1} = \pi_1^{-1}(P_2w_1 \otimes w_2)$. So, the J(c,h,l) is generated by $Q_1\mathbf{1}, Q_2\mathbf{1}$.

Proposition 3.10. Suppose that $c, h, \alpha, \beta \in \mathbb{C}, l, \gamma \in \mathbb{C}^*$. Then $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$ is irreducible if and only if $(\rho_n(Q_1), \rho_n(Q_2)) \neq (0, 0)$ for all $n \in \mathbb{Z}$, where $Q_1, Q_2 \in \mathcal{U}(\mathbb{D}^-)$ are in Lemma 3.9

Proof. Let $J = U(\mathcal{D}^-)Q_1\mathbf{1} + U(\mathcal{D}^-)Q_2\mathbf{1}$ be the unique maximal proper submodule of M(c, h, l), π be the canonical map from $M(c, h, l) \otimes A(\alpha, \beta, \gamma)$ to $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$. First, we prove the following statement.

Claim 1. $L(c,h,l) \otimes A(\alpha,\beta,\gamma)$ is irreducible if and only if $M(c,h,l) \otimes y^n = W_n^{(n+\frac{1}{2})} + J \otimes y^n$ for all $n \in \frac{\mathbb{Z}}{2}$.

 (\Rightarrow) Since $W^{(n+\frac{1}{2})}$, $J \otimes A(\alpha, \beta, \gamma)$ are submodules of $M(c, h, l) \otimes A(\alpha, \beta, \gamma)$, and $W^{(n+\frac{1}{2})} \not\subseteq J \otimes A(\alpha, \beta, \gamma)$, we see that

$$J \otimes A(\alpha, \beta, \gamma) \subsetneq W^{(n+\frac{1}{2})} + J \otimes A(\alpha, \beta, \gamma) \subseteq M(c, h, l) \otimes A(\alpha, \beta, \gamma).$$

But $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$ is irreducible, so $W^{(n+\frac{1}{2})} + J \otimes A(\alpha, \beta, \gamma) = M(c, h, l) \otimes A(\alpha, \beta, \gamma)$. Thus we have $M(c, h, l) \otimes y^n = W_n^{(n+\frac{1}{2})} + J \otimes y^n$, $\forall n \in \frac{\mathbb{Z}}{2}$.

(\Leftarrow) Assume that V_1 is a nonzero submodule of $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$, then there exists $k \in \frac{\mathbb{Z}}{2}$ such that $\overline{\mathbf{1}} \otimes y^i \in V_1$, $\forall k \leq i \in \frac{\mathbb{Z}}{2}$ by Lemma 3.5. It shows that $\pi(W^{(k)} + J \otimes A(\alpha, \beta, \gamma)) \subseteq V_1$. From Lemma 3.6 (3) we know that

$$W_{k-\frac{1}{2}}^{(k)} \oplus \mathbb{C}(\mathbf{1} \otimes y^{k-\frac{1}{2}}) = M(c,h,l) \otimes y^{k-\frac{1}{2}} = W_{k-\frac{1}{2}}^{(k)} + J \otimes y^{k-\frac{1}{2}} \subseteq \pi^{-1}(V_1).$$

So $\mathbf{\overline{1}} \otimes y^{k-\frac{1}{2}} \in V_1$. Inductively, $\mathbf{\overline{1}} \otimes y^i \in V_1$ for all $i \in \frac{\mathbb{Z}}{2}$. Then it is easy to see that $V_1 = L(c, h, l) \otimes A(\alpha, \beta, \gamma)$. Therefore, $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$ is irreducible. Claim 1 is proved.

Using Claim 1 and Lemmas 3.4, 3.6, 3.7 we see that

$$\begin{split} L(c,h,l) \otimes A(\alpha,\beta,\gamma) \text{ is irreducible} \\ & \longleftrightarrow W_n^{(n+\frac{1}{2})} \oplus \mathbb{C}(\mathbf{1} \otimes y^n) = M(c,h,l) \otimes y^n = W_n^{(n+\frac{1}{2})} + J \otimes y^n, \forall n \in \frac{\mathbb{Z}}{2} \\ & \longleftrightarrow (U(\mathcal{D}^-)Q_1\mathbf{1} + U(\mathcal{D}^-)Q_2\mathbf{1}) \otimes y^n \notin W_n^{(n+\frac{1}{2})}, \forall n \in \frac{\mathbb{Z}}{2} \\ & \longleftrightarrow \{Q_1\mathbf{1} \otimes y^n, Q_2\mathbf{1} \otimes y^n\} \notin W_n^{(n+\frac{1}{2})}, \forall n \in \frac{\mathbb{Z}}{2} \\ & \longleftrightarrow \{\rho_n(Q_1), \rho_n(Q_2)\} \neq \{0,0\}, \forall n \in \frac{\mathbb{Z}}{2}. \end{split}$$

Now, we complete the proof of the theorem.

Now we combine Proposition 3.2 and Proposition 3.10 into the following main theorem.

Theorem 3.11. Let $c, h, l, \alpha, \beta \in \mathbb{C}, \gamma \in \mathbb{C}^*$.

- (1) If l = 0, then $L(c, h, l) \otimes A'(\alpha, \beta)^{\mathcal{D}}$ is an irreducible \mathcal{D} -module if and only if $L_{\mathcal{V}}(c, h) \otimes A'(\alpha, \beta)$ is an irreducible \mathcal{V} -module.
- (2) If $l \neq 0$, then $L(c, h, l) \otimes A'(\alpha, \beta)^{\mathbb{D}}$ is an irreducible \mathbb{D} -module if and only if $L_{\mathcal{V}}(c-1, h-\frac{1}{16}) \otimes A'(\alpha, \beta)$ is an irreducible \mathcal{V} -module.
- (3) If l = 0, then $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$ is irreducible.
- (4) If $l \neq 0$, then $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$ is irreducible if and only if $(\rho_n(Q_1), \rho_n(Q_2)) \neq (0, 0)$ for all $n \in \frac{\mathbb{Z}}{2}$ where $Q_1, Q_2 \in \mathcal{U}(\mathcal{D}^-)$ are in Lemma 3.9.

Remark 3.12. Suppose $l \neq 0$ and the unique maximal submodule of M(c, h, l) is generated by $Q_1\mathbf{1}, Q_2\mathbf{1}$ for some $Q_1, Q_2 \in \mathcal{U}(\mathcal{D}^-)$ as in Lemma 3.9. For $k \in \frac{\mathbb{Z}}{2}$, the image of $W^{(k)}$ under the canonical map $\pi : M(c, h, l) \otimes A(\alpha, \beta, \gamma) \to L(c, h, l) \otimes A(\alpha, \beta, \gamma)$ is denoted by $\overline{W^{(k)}}$. Then we have a sequence of submodules of $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$:

$$\cdots \subseteq \overline{W^{(k+\frac{1}{2})}} \subseteq \overline{W^{(k)}} \subseteq \overline{W^{(k-\frac{1}{2})}} \subseteq \cdots$$
(3.3)

From the proof of Proposition 3.10, the inclusion $W^{(k+\frac{1}{2})} \subseteq \overline{W^{(k)}}$ is proper if and only if $(\rho_k(Q_1), \rho_k(Q_2)) = (0, 0)$. Denote by Λ the set of all $n \in \frac{\mathbb{Z}}{2}$ such that $(\rho_n(Q_1), \rho_n(Q_2)) = (0, 0)$.

If $(Q_1, Q_2) = (0, 0)$, then $\Lambda = \frac{\mathbb{Z}}{2}$ and all inclusions $\overline{W^{(k+\frac{1}{2})}} \subseteq \overline{W^{(k)}}$ are proper for $k \in \frac{\mathbb{Z}}{2}$. Moreover, $\overline{W^{(k)}}/\overline{W^{(k+\frac{1}{2})}}$ is a highest weight module with highest weight $\alpha + h - k$ for $k \in \Lambda$.

If $(Q_1, Q_2) \neq (0, 0)$, then Λ is a finite subset of $\frac{\mathbb{Z}}{2}$. In sequence 3.3, the inclusion $\overline{W^{(k+\frac{1}{2})}} \subseteq \overline{W^{(k)}}$ is proper if and only if $k \in \Lambda$. Without lose of generality, we can assume that $\Lambda = \{k_1, k_2, \dots, k_t\}$ with $k_1 < k_2 < \dots < k_t$, then sequence 3.3 can be simplified as following:

$$0 \subsetneq \overline{W^{(k_{l+1})}} \subsetneq \overline{W^{(k_l)}} \subsetneq \cdots \subsetneq \overline{W^{(k_1)}} = L(c, h, l) \otimes A(\alpha, \beta, \gamma)$$

for any $k_{t+1} > k_t$. It shows that $\overline{W^{(k_{i+1})}} = \overline{W^{(k_i + \frac{1}{2})}}$ for $1 \le i \le t$. However, we know that $W^{(k_i)}$ is generated by $W^{(k_i + \frac{1}{2})}$ and $1 \otimes y^{k_i}$, thus $\overline{W^{(k_i)}}$ is generated by $\overline{W^{(k_{i+1})}}$ and $\overline{1 \otimes y^{k_i}}$. This implies that $\overline{W^{(k_i)}}/\overline{W^{(k_{i+1})}}$ is a highest weight module with highest weight $\alpha + h - k_i$. In particular, we obtain that $\overline{W^{(k_{i+1})}} = \overline{W^{(k_i + \frac{1}{2})}}$ is an irreducible weight \mathcal{D} -module.

4. Isomorphism theorem

In the section, we determine the necessary and sufficient conditions for two of the tensor product modules studied in Section 3 to be isomorphic.

Lemma 4.1. Let $\alpha, \beta \in \mathbb{C}, \gamma \in \mathbb{C}^*$. Then $A(\alpha, \beta, \gamma) \cong A(\alpha + n, \beta, \gamma)$ for any $n \in \mathbb{Z}_2$.

Proof. For any $n \in \frac{\mathbb{Z}}{2}$, it is straightforward to verify that the linear map

$$F: A(\alpha, \beta, \gamma) \to A(\alpha + n, \beta, \gamma), \quad v_k \to (\delta_{n \in \mathbb{Z}} + \delta_{n \notin \mathbb{Z}}(\delta_{k \in \mathbb{Z}} + \gamma \delta_{k \notin \mathbb{Z}}))v_{k+n}$$

is an isomorphism of \mathcal{D} -modules.

Proposition 4.2. Let $c, h, l, \alpha, \beta, c_1, h_1, l_1, \alpha_1, \beta_1, \in \mathbb{C}, \gamma, \gamma_1 \in \mathbb{C}^*$. Then $L(c, h, l) \otimes A(\alpha, \beta, \gamma) \cong L(c_1, h_1, l_1) \otimes A(\alpha_1, \beta_1, \gamma_1)$ if and only if $(\beta, \gamma, c, h, l) = (\beta_1, \gamma_1, c_1, h_1, l_1)$ and $\alpha - \alpha_1 \in \frac{\mathbb{Z}}{2}$.

Proof. From Lemma 4.1, we may assume that $0 \le \Re e \alpha$, $\Re e \alpha_1 < \frac{1}{2}$. Since the "if part" is trivial, we only need to show the "only if part".

We identify the \mathcal{D} -modules $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$ with $V_1 = L(c, h, l) \otimes \mathbb{C}[y^{\pm \frac{1}{2}}]$, and $L(c_1, h_1, l_1) \otimes A(\alpha_1, \beta_1, \gamma_1)$ with $V_2 = L(c_1, h_1, l_1) \otimes \mathbb{C}[y^{\pm \frac{1}{2}}]$. Let $\overline{\mathbf{1}}$ and $\overline{\mathbf{1}'}$ be the highest weight vectors of L(c, h, l) and $L(c_1, h_1, l_1)$ respectively. It is clear that

$$c = c_1, l = l_1.$$

Suppose $\varphi : V_1 \to V_2$ is a \mathcal{D} -module isomorphism. For $k \in \frac{\mathbb{Z}}{2}$, let $\varphi(\overline{\mathbf{1}} \otimes y^k) = \sum_{i=0}^{s} P_i \overline{\mathbf{1}'} \otimes y^i$ for some $t \in \frac{\mathbb{Z}}{2}$ where $P_i \in U(\mathcal{D}^-)_{-i}$ is homogeneous. Note that $\varphi(\overline{\mathbf{1}} \otimes y^k)$ and $\overline{\mathbf{1}} \otimes y^k$ have the same weight. We deduce that

$$\alpha + h - k = \alpha_1 + h_1 - t. \tag{4.1}$$

For any m, n > s, we compute

$$(\delta_{k\in\mathbb{Z}} + \gamma\delta_{k\notin\mathbb{Z}})\varphi(\overline{\mathbf{1}} \otimes y^{k+m+n+\frac{1}{2}}) = \varphi(h_{\frac{1}{2}+m+n}(\overline{\mathbf{1}} \otimes y^{k}))$$

$$= h_{\frac{1}{2}+m+n}\varphi(\overline{\mathbf{1}} \otimes y^{k}) = \sum_{i=0}^{s} (\delta_{t+i\in\mathbb{Z}} + \gamma_{1}\delta_{t+i\notin\mathbb{Z}})P_{i}\overline{\mathbf{1}'} \otimes y^{t+m+n+\frac{1}{2}},$$

$$(\delta_{k\in\mathbb{Z}} + \gamma\delta_{k\notin\mathbb{Z}})(\alpha + \beta m - k - n - \frac{1}{2})\varphi(\overline{\mathbf{1}} \otimes y^{k+m+n+\frac{1}{2}})$$

$$= \varphi(d_{m}h_{\frac{1}{2}+n}(\overline{\mathbf{1}} \otimes y^{k})) = d_{m}h_{\frac{1}{2}+n}\varphi(\overline{\mathbf{1}} \otimes y^{k}) = \sum_{i=0}^{s} d_{m}h_{\frac{1}{2}+n}(P_{i}\overline{\mathbf{1}'} \otimes y^{t})$$

$$= \sum_{i=0}^{s} (\delta_{t+i\in\mathbb{Z}} + \gamma_{1}\delta_{t+i\notin\mathbb{Z}})(\alpha_{1} + \beta_{1}m - i - t - n - \frac{1}{2})P_{i}\overline{\mathbf{1}'} \otimes y^{t+m+n+\frac{1}{2}},$$

$$(4.2)$$

Comparing the above two equations we obtain that

$$\alpha + \beta m - k - n - \frac{1}{2} = \alpha_1 + \beta_1 m - i - t - n - \frac{1}{2}, \forall m, n > s, 0 \le i \le s$$

with $P_i \mathbf{1}' \neq 0$. Combining with (4.1) we deduce that

$$\beta = \beta_1, \alpha = \alpha_1, i = h - h_1, t = k - h + h_1.$$

Since $i = -\deg(P_i) \ge 0$, so $h - h_1 \ge 0$. By considering φ^{-1} in the above arguments we deduce that $h_1 - h \ge 0$. We must have

$$h = h_1, t = k, s = 0.$$

Therefore, for any $k \in \frac{\mathbb{Z}}{2}$, we may assume that $\varphi(\overline{\mathbf{1}} \otimes y^k) = a_k \overline{\mathbf{1}'} \otimes y^k$ where $a_k \in \mathbb{C}^*$. From

$$\begin{split} \gamma a_{k+m+n+1} \overline{\mathbf{1}'} \otimes y^{k+m+n+1} &= \gamma \varphi(\overline{\mathbf{1}} \otimes y^{k+m+n+1}) = \varphi(h_{\frac{1}{2}+m} h_{\frac{1}{2}+n}(\overline{\mathbf{1}} \otimes y^k)) \\ &= h_{\frac{1}{2}+m} h_{\frac{1}{2}+n} \varphi(\overline{\mathbf{1}} \otimes y^k) = h_{\frac{1}{2}+m} h_{\frac{1}{2}+n}(a_k \overline{\mathbf{1}'} \otimes y^k) = a_k \gamma_1 \overline{\mathbf{1}'} \otimes y^{k+m+n+1}, \forall m, n \in \mathbb{Z}_+ \end{split}$$

We deduce that

$$\gamma a_{k+m+n+1} = a_k \gamma_1, \forall m, n \in \mathbb{Z}_+$$

yielding that $\gamma = \gamma_1$. We complete the proof.

Remark 4.3. In Proposition 4.2 we can have similar results for any highest weight modules, not only for irreducible highest weight modules.

Theorem 4.4. Let $c, h, l, \alpha, \beta, c_1, h_1, l_1, \alpha_1, \beta_1, \in \mathbb{C}, \gamma, \gamma_1 \in \mathbb{C}^*$. Then

- (1) $L(c,h,l) \otimes A'(\alpha,\beta)^{\mathbb{D}} \cong L(c_1,h_1,l_1) \otimes A'(\alpha_1,\beta_1)^{\mathbb{D}}$ if and only if $(c,h,l) = (c_1,h_1,l_1), \alpha \alpha_1 \in \mathbb{Z}$, and $\beta = \beta_1$ or $\{\beta,\beta_1\} = \{0,-1\}$.
- (2) $L(c, h, l) \otimes A(\alpha, \beta, \gamma) \cong L(c_1, h_1, l_1) \otimes A(\alpha_1, \beta_1, \gamma_1)$ if and only if $(\beta, \gamma, c, h, l) = (\beta_1, \gamma_1, c_1, h_1, l_1)$ and $\alpha - \alpha_1 \in \frac{\mathbb{Z}}{2}$.
- $(3) \ L(c,h,l)\otimes A(\alpha,\beta,\gamma)\not\cong L(c_1,h_1,l_1)\otimes A'(\alpha_1,\beta_1)^{\mathcal{D}}.$

Proof. (1) The "if part" is trivial, we only need to show the "only if part". It is clear that $l = l_1$ from $L(c, h, l) \otimes A'(\alpha, \beta)^{\mathbb{D}} \cong L(c_1, h_1, l_1) \otimes A'(\alpha_1, \beta_1)^{\mathbb{D}}$. If $l = l_1 \neq 0$, using Lemmata 2.9 and 3.1 we know that

$$L_{\mathcal{V}}(c-1,h-\frac{1}{16}) \otimes A'(\alpha,\beta) \cong L_{\mathcal{V}}(c_1-1,h_1-\frac{1}{16}) \otimes A'(\alpha_1,\beta_1),$$

$$M_{\mathcal{H}}(l) \cong M_{\mathcal{H}}(l_1).$$

Thanks to [5, Theorem 2],

$$L_{\mathcal{V}}(c-1,h-\frac{1}{16}) \cong L_{\mathcal{V}}(c_1-1,h_1-\frac{1}{16}), \quad A'(\alpha,\beta) \cong A'(\alpha_1,\beta_1).$$

Thus $(c, h, l) = (c_1, h_1, l_1), \alpha - \alpha_1 \in \mathbb{Z}, \beta = \beta_1 \text{ or } \{\beta, \beta_1\} = \{0, -1\}$ by Lemma 2.5.

If
$$l = l_1 = 0$$
, then $L(c, h, l) \cong L_V(c, h)^{\mathcal{D}}$, $L(c_1, h_1, l_1) \cong L_V(c_1, h_1)^{\mathcal{D}}$. Thus
 $L_V(c, h) \otimes A'(\alpha, \beta) \cong L_V(c_1, h_1) \otimes A'(\alpha_1, \beta_1)$.

So, $L_{\mathcal{V}}(c,h) \cong L_{\mathcal{V}}(c_1,h_1), A'(\alpha,\beta) \cong A'(\alpha_1,\beta_1)$ by Theorem 2 in [5]. We also have $(\gamma, c, h, l) = (\gamma_1, c_1, h_1, l_1), \alpha - \alpha_1 \in \mathbb{Z}, \beta = \beta_1 \text{ or } \{\beta, \beta_1\} = \{0, -1\}.$

(2) This is Proposition 4.2.

(3) From (3.1) we know that $h_{\frac{1}{2}}$ does not annihilate any weight vector in $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$, while it annihilates $\overline{\mathbf{1}'} \otimes A'(\alpha_1, \beta_1)^{\mathfrak{D}}$. So $L(c, h, l) \otimes A(\alpha, \beta, \gamma) \not\cong L(c_1, h_1, l_1) \otimes A'(\alpha_1, \beta_1)^{\mathfrak{D}}$. \Box

5. Examples

In this section, we show some concrete examples to illustrate Theorem 3.11, and to give irreducible weight modules with infinite dimensional weight spaces over the mirror-twisted Heisenberg-Virasoro algebra \mathcal{D} .

Example 5.1. We can find many irreducible Verma modules $M_{\mathcal{V}}(c,h)$ over \mathcal{V} in [2]. For all these irreducible Verma modules, the tensor product $M_{\mathcal{V}}(c,h)^{\mathbb{D}} \otimes A(\alpha,\beta,\gamma)$ is an irreducible weight \mathbb{D} -module with infinite dimensional weight spaces by Theorem 3.11 (3). Thus we obtain irreducible weight modules with infinite dimensional weight spaces.

In the rest of this section, we present some examples for the tensor product \mathcal{D} -modules $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$ with $l \neq 0$. We will use w_1 to denote the highest weight vector of $M_{\mathcal{V}}(c, h)$. Let us start with an example where the maximal proper submodule J(c, h, l) of M(c, h, l) is generated by one singular vector.

Example 5.2. Let $l \neq 0$. Then $M(2, \frac{1}{16}, l) \cong M_{\mathcal{V}}(1, 0)^{\mathcal{D}} \otimes M_{\mathcal{H}}(l)^{\mathcal{D}}$ by Lemma 2.9. From [2], we know that the unique maximal proper submodule of $M_{\mathcal{V}}(1, 0)$ is generated by $d_{-1}w_1$. Using (2.1) we deduce that the maximal proper submodule of $M(2, \frac{1}{16}, l)$ is generated by $(d_{-1} - \frac{1}{2l}h_{-\frac{1}{2}}^2)\mathbf{1}$.

Therefore, $L(2, \frac{1}{16}, l) \otimes A(\alpha, \beta, \gamma)$ *is irreducible if and only if*

$$\rho_n(d_{-1} - \frac{1}{2l}h_{-\frac{1}{2}}^2) = \beta - \alpha + 1 + n - \frac{1}{2l}\gamma \neq 0, \forall n \in \frac{\mathbb{Z}}{2}.$$

If $n = \alpha - \beta - 1 + \frac{1}{2l}\gamma \in \frac{\mathbb{Z}}{2}$, then $\overline{W^{(n+\frac{1}{2})}}$ is the unique minimal submodule of $L(2, \frac{1}{16}, l) \otimes A(\alpha, \beta, \gamma)$. Of course, it is the only irreducible submodule. Moreover, $(L(2, \frac{1}{16}, l) \otimes A(\alpha, \beta, \gamma))/\overline{W^{(n+\frac{1}{2})}}$ is a highest weight module with highest weight vector $\mathbf{1} \otimes y^n$ of weight $(2, \beta - \frac{\gamma}{2l} + \frac{17}{16}, l)$. By Lemma 2.10, we know that $(L(2, \frac{1}{16}, l) \otimes A(\alpha, \beta, \gamma))/\overline{W^{(n+\frac{1}{2})}}$ is irreducible if $\beta - \frac{\gamma}{2l} + 1 \neq (\frac{k}{2})^2$ for any $k \in \mathbb{Z}$.

Example 5.3. By Lemma 2.9, we have $M(2, \frac{5}{16}, l) \cong M_{\mathcal{V}}(1, \frac{1}{4})^{\mathfrak{D}} \otimes M_{\mathcal{H}}(l)^{\mathfrak{D}}$ when $l \neq 0$. From [2], we know that the unique maximal proper submodule of $M_{\mathcal{V}}(1, \frac{1}{4})$ is generated by $(d_{-1}^2 - d_{-2})w_1$. Using (2.1) we deduce that the maximal proper submodule of $M(2, \frac{5}{16}, l)$ is generated by $Q\mathbf{1}$ where

$$Q = d_{-1}^2 - d_{-2} - \frac{1}{l}d_{-1}h_{-\frac{1}{2}}^2 + \frac{1}{4l^2}h_{-\frac{1}{2}}^4 + \frac{3}{2l}h_{-\frac{3}{2}}h_{-\frac{1}{2}}$$

Therefore, $L(2, \frac{5}{16}, l) \otimes A(\alpha, \beta, \gamma)$ is irreducible if and only if $\rho_n(Q) \neq 0, \forall n \in \frac{\mathbb{Z}}{2}$. By direct computation, we have following equations

$$\begin{split} \rho_n(d_{-1}^2) &= (\alpha - \beta - 2 - n)(\alpha - \beta - 1 - n)\\ \rho_n(d_{-2}) &= -(\alpha - 2\beta - 2 - n),\\ \rho_n(d_{-1}h_{-\frac{1}{2}}^2) &= -\gamma(\alpha - \beta - 2 - n),\\ \rho_n(h_{-\frac{1}{2}}^4) &= \gamma^2,\\ \rho_n(h_{-\frac{3}{2}}^4h_{-\frac{1}{2}}) &= \gamma. \end{split}$$

Thus we have

$$\rho_n(Q) = \rho_n(d_{-1}^2 - d_{-2} - \frac{1}{l}d_{-1}h_{-\frac{1}{2}}^2 + \frac{1}{4l^2}h_{-\frac{1}{2}}^4 + \frac{3}{2l}h_{-\frac{3}{2}}h_{-\frac{1}{2}})$$

= $(\alpha - \beta - 2 - n)(\alpha - \beta - 1 - n) + (\alpha - 2\beta - 2 - n) + \frac{\gamma}{l}(\alpha - \beta - 2 - n) + \frac{1}{4l^2}\gamma^2 + \frac{3}{2l}\gamma$

Solving the equation $\rho_n(Q) = 0$, we get

$$n = (\alpha - \beta - 1 + \frac{\gamma}{2l}) \pm \sqrt{\beta - \frac{\gamma}{2l}} + 1.$$

Let

$$n_1 = (\alpha - \beta - 1 + \frac{\gamma}{2l}) + \sqrt{\beta - \frac{\gamma}{2l} + 1}, \quad n_2 = (\alpha - \beta - 1 + \frac{\gamma}{2l}) - \sqrt{\beta - \frac{\gamma}{2l} + 1}.$$

(1) We know that $L(2, \frac{5}{16}, l) \otimes A(\alpha, \beta, \gamma)$ is irreducible if and only if $n_1, n_2 \notin \frac{\mathbb{Z}}{2}$.

(2) Suppose only one of n_1 and n_2 is in $\frac{\mathbb{Z}}{2}$, say $n_1 \in \frac{\mathbb{Z}}{2}$. Then $\overline{W^{(n_1+\frac{1}{2})}}$ is the unique irreducible submodule of $L(2, \frac{5}{16}, l) \otimes A(\alpha, \beta, \gamma)$. And the quotient module $(L(2, \frac{5}{16}, l) \otimes A(\alpha, \beta, \gamma))/\overline{W^{(n_1+\frac{1}{2})}}$ is a highest weight module with highest weight $(2, \beta + \frac{21}{16} - \frac{\gamma}{2l} - \sqrt{\beta - \frac{\gamma}{2l} + 1}, l)$. By Lemma 2.10, we

know that $(L(2, \frac{5}{16}, l) \otimes A(\alpha, \beta, \gamma))/\overline{W^{(n_1 + \frac{1}{2})}}$ is irreducible if further $\beta - \frac{\gamma}{2l} + \frac{5}{4} - \sqrt{\beta - \frac{\gamma}{2l} + 1} \neq (\frac{k}{2})^2$ for any $k \in \mathbb{Z}$.

(3) Suppose that $n_1, n_2 \in \frac{\mathbb{Z}}{2}$. Without lose of generality, we can assume that $n_1 \leq n_2$, then we have a sequence of submodules of $L(2, \frac{5}{16}, l) \otimes A(\alpha, \beta, \gamma)$:

$$0 \subsetneq \overline{W^{(n_2 + \frac{1}{2})}} \subsetneq \overline{W^{(n_2)}} \subseteq \overline{W^{(n_1)}} = L(2, \frac{5}{16}, l) \otimes A(\alpha, \beta, \gamma)$$

where the inclusion $\overline{W^{(n_2)}} \subseteq \overline{W^{(n_1)}}$ is proper if and only if $n_1 < n_2$. Here, $\overline{W^{(n_2+\frac{1}{2})}}$ is the unique irreducible submodule of $L(2, \frac{5}{16}, l) \otimes A(\alpha, \beta, \gamma)$ and $\overline{W^{(n_2)}} / \overline{W^{(n_2+\frac{1}{2})}}$ is a highest weight module with highest weight $(2, \beta + \frac{21}{16} - \frac{\gamma}{2l} + \sqrt{\beta - \frac{\gamma}{2l} + 1}, l)$. By Lemma 2.10, we know that $\overline{W^{(n_2)}} / \overline{W^{(n_2+\frac{1}{2})}}$ is irreducible if $\beta - \frac{\gamma}{2l} + \frac{5}{4} + \sqrt{\beta - \frac{\gamma}{2l} + 1} \neq (\frac{k}{2})^2$ for any $k \in \mathbb{Z}$. As for quotient module $(L(2, \frac{5}{16}, l) \otimes A(\alpha, \beta, \gamma)) / \overline{W^{(n_2)}}$, it is also a highest weight module with highest weight $(2, \beta + \frac{29}{16} - \frac{\gamma}{2l} + \sqrt{\beta - \frac{\gamma}{2l} + 1} \neq (\frac{5}{16}, l) \otimes A(\alpha, \beta, \gamma))$.

To conclude this section, we give an example such that the unique maximal submodule J(c, h, l) of M(c, h, l) is generated by two singular vectors.

Example 5.4. Let $l \neq 0$, c = h = 0. By Lemma 2.9 we know that $M(1, \frac{1}{16}, l) \cong M_{\mathcal{V}}(0, 0)^{\mathbb{D}} \otimes M_{\mathcal{H}}(l)^{\mathbb{D}}$. It is clear that the unique maximal submodule of $M_{\mathcal{V}}(0, 0)$ is generated by $d_{-1}w_1, d_{-2}w_1$. Using (2.1) we deduce that the maximal proper submodule of $M(1, \frac{1}{16}, l)$ is generated by Q_1w and $Q_2\mathbf{1}$ where $Q_1 = d_{-1} - \frac{1}{2l}h_{-\frac{1}{2}}^2$, $Q_2 = d_{-2} - \frac{1}{l}h_{-\frac{3}{2}}h_{-\frac{1}{2}}$. We see that

$$\rho_n(Q_1) = \beta - \alpha + 1 + n - \frac{1}{2l}\gamma, \quad \rho_n(Q_2) = 2\beta - \alpha + 2 + n - \frac{\gamma}{l}$$

Taking $\rho_n(Q_1) = \rho_n(Q_2) = 0$, we have that $n = \alpha$ and $\beta + 1 = \frac{\gamma}{2l}$. Thus, if $n = \alpha \in \frac{\mathbb{Z}}{2}, \beta + 1 - \frac{\gamma}{2l} = 0$, then $\overline{W^{(\alpha+\frac{1}{2})}}$ is the unique irreducible submodule of $L(1, \frac{1}{16}, l) \otimes A(\alpha, \beta, \gamma)$ and $\overline{W^{(\alpha)}}/\overline{W^{(\alpha+\frac{1}{2})}}$ is a highest weight module with highest weight $(1, \frac{1}{16}, l)$. Otherwise, $L(1, \frac{1}{16}, l) \otimes A(\alpha, \beta, \gamma)$ is irreducible.

Remark 5.5. For any $c, h, \alpha, \beta \in \mathbb{C}, l, \gamma \in \mathbb{C}^*$, it is generally hard to determine the structure of nontrivial quotient modules $W^{(k)}/W^{(k+1)}$ for the \mathcal{D} -module $L(c, h, l) \otimes A(\alpha, \beta, \gamma)$. We also have the similar questions for the Virasoro algebra and Heisenberg-Virasoro algebra, see [5, 21].

Acknowledgments: The research in this paper was carried out during the visit of the first author at Wilfrid Laurier University in 2019-2021. The first author is partially supported by CSC of China (Grant 201906340096), and NSF of China (Grants 11771410 and 11931009). The second author is partially supported by NSFC (11871190) and NSERC (311907-2015). The first author wants to thank Y. Ma for helpful suggestions when he was preparing the paper.

References

 E. Arbarello, C. De Concini, V. G. Kac and C. Procesi, Moduli Spaces of Curves and Representation Theory, Comm. Math. Phys. 117 (1988), 1-36.

- [2] A. Astashkevich, On the structure of Verma modules over Virasoro and Neveu- Schwarz algebras, Comm. Math. Phys. 186 (1997), 531–562.
- [3] K. Barron, On twisted modules for N=2 supersymmetric vertex operator superalgebras, Lie theory and its applications in physics, 411-420 (2013), Springer Proc. Math. Stat., 36, Springer, Tokyo.
- [4] V. Chari and A. Pressley, A new family of irreducible, integrable modules for affine Lie algebras, Math. Ann. 277, 3 (1987) 543-562.
- [5] H. Chen, X. Guo and K. Zhao, Tensor product weight modules over the Virasoro algebra, J. London. Math. Soc. 88(3) (2013), 829-844.
- [6] C. Conley and C. Martin, A family of irreducible representations of the Witt Lie algebra with infinitedimensional weight spaces, Compos. Math., 128 (2) (2001), 153-175.
- [7] P. Di Francesco, P. Mathieu and D. Sénéchal, Conformal field theory, Springer-Verlag, New York, 1997.
- [8] C. Dong, G. Mason and Y. Zhu, Discrete series of the Virasoro algebra and the moonshine module, Algebraic groups and their generalizations: quantum and infinite-dimensional methods (University Park, PA), Proc. Sympos. Pure Math. (1991), 295–316, Part 2, Amer. Math. Soc. Providence, RI, 1994.
- [9] B. Feigin and D. Fuchs, Representations of the Virasoro algebra, Representation of Lie groups and related topics, Adv. Stud. Contemp. Math. Gordon and Breach, New York, (1990), 465–554.
- [10] I. Frenkel, J. Lepowsky and A. Meurman, Vertex operator algebras and the monster, Pure and Appl. Math. 134 (1998).
- [11] I. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, Duke Math. J. 66(1) (1992), 123–168.
- [12] P. Goddard and D. Olive, Kac-Moody and Virasoro algebras in relation to quantum physics, Internat. J. Modern Phys. A 1(2) (1986), 303-414.
- [13] X. Guo and K. Zhao, Irreducible representations of non-twisted affine Kac-Moody algebras, arXiv:1305.4059.
- [14] M. Henkel, Schrödinger invariance and strongly anisotropic critical systems, J. Stat. Phys. 75 (1994), 1023-1029.
- [15] M. Henkel, Phenomenology of local scale invariance: from conformal invariance to dynamical scaling, Nucl. Phys. B. 641 (2002), 405-410.
- [16] J. Li and Y. Su, Representations of the Schrödinger-Virasoro algebras. J. Math. Phys. 49(5) (2008), 053512, 15pp.
- [17] D. Liu, Y. Pei, L. Xia and K. Zhao, Irreducible modules over the mirror-twisted Heisenberg-Virasoro algebra, preprint, 2019.
- [18] D. Liu, and X. Zhang, Tensor product weight modules of Schrödinger-Virasoro algebras, Front. Math. China. 14 (2019), 381–393.
- [19] R. Lü, X. Guo and K. Zhao, Irreducible modules over the Virasoro algebra, Doc. Math. 16 (2011) 709–721.
- [20] R. Lü and K. Zhao, Classification of irreducible weight modules over the twisted Heisenberg-Virasoro algebras, Comm. Contemp. Math. 12(2) (2010), 183-205.
- [21] R. Lü and K. Zhao, Generalized oscillator representations of the twisted Heisenberg-Virasoro algebra, Algebr. Represent. Theory. 23 (2020) 1417-1442.
- [22] O. Mathieu, Classification of Harish-Chandra modules over the Virasoro Lie algebra, Invent. Math. 107(2) (1992), 225-234.
- [23] S. Tan and X. Zhang, Automorphisms and Verma modules for generalized Schrödinger-Virasoro algebras, J. Algebra. 322 (2009), 1379-1394.
- [24] H. Zhang, A class of representations over the Virasoro algebra, J. Algebra. 190(1) (1997), 1–10.

D. Gao: School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui 230026, P. R. China

Email address: gaodfw@mail.ustc.edu.cn

K. ZHAO: DEPARTMENT OF MATHEMATICS, WILFRID LAURIER UNIVERSITY, WATERLOO, ON N2L 3C5, CANADA

Email address: kzhao@wlu.ca