

On the Φ -Stability and Related Conjectures*

Lei Yu^{1*}

^{1*}School of Statistics and Data Science, LPMC, KLMDASR, and LEBPS,
Nankai University, 94 Weijin Rd, Tianjin, 300071, Tianjin, China.

Corresponding author(s). E-mail(s): leiyu@nankai.edu.cn;

Abstract

Given a convex function $\Phi : [0, 1] \rightarrow \mathbb{R}$ and the mean $\mathbb{E}f(\mathbf{X}) = a \in [0, 1]$, which Boolean function f maximizes the Φ -stability $\mathbb{E}[\Phi(\mathcal{T}_\rho f(\mathbf{X}))]$ of f ? Here \mathbf{X} is a random vector uniformly distributed on the discrete cube $\{-1, 1\}^n$ and \mathcal{T}_ρ is the Bonami-Beckner operator. Special cases of this problem include the (symmetric and asymmetric) α -stability problems and the “Most Informative Boolean Function” problem. In this paper, we provide several upper bounds for the maximal Φ -stability. When specializing Φ to some particular forms, by these upper bounds, we partially resolve Mossel and O’Donnell’s conjecture on α -stability with $\alpha > 2$, Li and Médard’s conjecture on α -stability with $1 < \alpha < 2$, and Courtade and Kumar’s conjecture on the “Most Informative Boolean Function” which corresponds to a conjecture on α -stability with $\alpha = 1$. Our proofs are based on discrete Fourier analysis, optimization theory, and improvements of the Friedgut–Kalai–Naor (FKN) theorem. Our improvements of the FKN theorem are sharp or asymptotically sharp for certain cases.

Keywords: Mossel–O’Donnell Conjecture, Courtade–Kumar Conjecture, Li–Médard Conjecture, Most Informative Boolean Function, Noise Stability, FKN Theorem, Boolean Function

MSC Classification: 60E15 , 68Q87 , 60G10

1 Introduction

Let \mathbf{X} be a random vector uniformly distributed on the discrete cube $\{-1, 1\}^n$. Let $\mathbf{Y} \in \{-1, 1\}^n$ be the random vector obtained by independently changing the sign of

*This arXiv version contains more proof details than the version published on Probability Theory and Related Fields.

each component of \mathbf{X} with the same probability $\frac{1-\rho}{2}$. Here, $\rho \in [0, 1]$ corresponds to the correlation coefficient between each component of \mathbf{X} and the corresponding one of \mathbf{Y} . Let T_ρ be the noise operator (or the *Bonami-Beckner operator*) which acts on Boolean functions $f : \{-1, 1\}^n \rightarrow \{0, 1\}$ such that $T_\rho f(\mathbf{x}) = \mathbb{E}[f(\mathbf{Y})|\mathbf{X} = \mathbf{x}]$. Let $\Phi : [0, 1] \rightarrow \mathbb{R}$ be a continuous and strictly convex function.

Definition 1. For a Boolean function $f : \{-1, 1\}^n \rightarrow \{0, 1\}$, the Φ -stability of f with respect to (w.r.t.) correlation coefficient ρ is defined as

$$\mathbf{Stab}_\Phi[f] = \mathbb{E}[\Phi(T_\rho f(\mathbf{X}))].$$

The Φ -stability is closely related to the Φ -entropy and the Φ -mutual information. Define the Φ -entropy of $f(\mathbf{Y})$ as

$$\text{Ent}_\Phi(f(\mathbf{Y})) := \mathbb{E}[\Phi(f(\mathbf{Y}))] - \Phi(\mathbb{E}f(\mathbf{Y})),$$

the *conditional* Φ -entropy of $f(\mathbf{Y})$ given \mathbf{X} as

$$\begin{aligned} \text{Ent}_\Phi(f(\mathbf{Y})|\mathbf{X}) &:= \mathbb{E}[\Phi(f(\mathbf{Y}))] - \mathbb{E}_\mathbf{X}\Phi(\mathbb{E}[f(\mathbf{Y})|\mathbf{X}]) \\ &= \mathbb{E}[\Phi(f(\mathbf{Y}))] - \mathbf{Stab}_\Phi[f], \end{aligned}$$

and the Φ -mutual information from $f(\mathbf{Y})$ to \mathbf{X} as

$$\begin{aligned} I_\Phi(f(\mathbf{Y}); \mathbf{X}) &:= \text{Ent}_\Phi(f(\mathbf{Y})) - \text{Ent}_\Phi(f(\mathbf{Y})|\mathbf{X}) \\ &= \mathbf{Stab}_\Phi[f] - \Phi(\mathbb{E}f(\mathbf{Y})) = \text{Ent}_\Phi(T_\rho f(\mathbf{X})). \end{aligned} \quad (1)$$

Hence, given the expectation of f , $\mathbf{Stab}_\Phi[f]$ is equal to the Φ -mutual information from $f(\mathbf{Y})$ to \mathbf{X} , up to a term which only depends on the expectation of f . In addition, observe that given the expectation of f , when there is no noise corruption (i.e., $\rho = 1$), $\mathbf{Stab}_\Phi[f] = \mathbb{E}[\Phi(f(\mathbf{X}))]$ which is also fixed. Hence, $\mathbf{Stab}_\Phi[f]$ also quantifies the change of the Φ -entropy of f after taking the noise operator.

In fact, the Φ -entropy is more related to the relative entropy than the Shannon entropy. Any nonnegative function g such that $\mathbb{E}_{\mathbf{Z} \sim P}[g(\mathbf{Z})] = 1$ can be seen as the Radon–Nikodym derivative of a probability distribution Q w.r.t. another probability distribution P , $\text{Ent}_\Phi(g)$ corresponds to a generalized relative entropy (called the “ f -divergence”) from Q to P [1–3].

The noise stability problem, in a general sense, concerns which Boolean functions (or measurable sets) are the “most stable” under the action of the noise operator. In terms of Φ -stability, the noise stability problem is formulated as follows.

Definition 2. The *maximal* Φ -stability w.r.t. volume $a \in [0, 1]$ is defined as

$$\mathbf{MaxStab}_\Phi(a) = \max_{f: \{-1, 1\}^n \rightarrow \{0, 1\}, \mathbb{E}f=a} \mathbf{Stab}_\Phi[f].$$

By the relation in (1), determining the maximal Φ -stability w.r.t. volume $a \in [0, 1]$ is equivalent to determining the maximum $I_\Phi(f(\mathbf{Y}); \mathbf{X})$ over all Boolean functions with $\mathbb{E}f = a$.

We next consider two common instances of Φ . For $\alpha \geq 1$, define $\Phi_\alpha, \Phi_\alpha^{\text{sym}} : [0, 1] \rightarrow \mathbb{R}$ as

$$\Phi_\alpha(t) := t \ln_\alpha(t) \quad \text{and} \quad \Phi_\alpha^{\text{sym}}(t) := t \ln_\alpha(t) + (1-t) \ln_\alpha(1-t).$$

where $0 \ln_\alpha(0) := 0$, and the function $\ln_\alpha : (0, +\infty) \rightarrow \mathbb{R}$ for $\alpha \in \mathbb{R}$ is defined as

$$\ln_\alpha(t) := \begin{cases} \frac{t^{\alpha-1}-1}{\alpha-1}, & \alpha \neq 1 \\ \ln(t), & \alpha = 1 \end{cases}$$

and is known as the α -*logarithm* (or q -*logarithm*) introduced by Tsallis [4], but with a slight reparameterization. Note that $\Phi_\alpha(t)$ and $\Phi_\alpha^{\text{sym}}(t)$ are non-decreasing in α given $t \in [0, 1]$ (since so is $\ln_\alpha(t)$), and Φ_1 and Φ_1^{sym} (with $\alpha = 1$) is the continuous extension of the ones with $\alpha > 1$. For brevity, we term the maximal Φ_α -stability and the maximal Φ_α^{sym} -stability respectively as the *maximal asymmetric and symmetric α -stabilities*¹, denoted as **MaxStab** $_\alpha$ and **MaxStab** $_\alpha^{\text{sym}}$. Similarly, the Φ_α -entropy and Φ_α^{sym} -entropy are respectively termed as the *asymmetric and symmetric α -entropies*, which are respectively denoted as Ent_α and $\text{Ent}_\alpha^{\text{sym}}$. Similar conventions also apply to the Φ_α -mutual information and Φ_α^{sym} -mutual information, denoted as I_α and I_α^{sym} . For Boolean f and for $\alpha = 1$, by definition, $\text{Ent}_1^{\text{sym}}(f)$, $\text{Ent}_1^{\text{sym}}(f(\mathbf{Y})|\mathbf{X})$, and $I_1^{\text{sym}}(f(\mathbf{Y}); \mathbf{X})$ respectively reduce to the *Shannon entropy* of $f(\mathbf{Y})$, the *conditional Shannon entropy* of $f(\mathbf{Y})$ given \mathbf{X} , and the *Shannon mutual information* (denoted as $I(f(\mathbf{Y}); \mathbf{X})$) between $f(\mathbf{Y})$ and \mathbf{X} . Furthermore, given f , $\text{Ent}_\alpha(f)$ and $\text{Ent}_\alpha^{\text{sym}}(f)$ are nondecreasing in α ; see [8, Theorem 33]. Moreover, it is easily seen that if *dictator functions* maximize the asymmetric α -stability over all balanced Boolean functions, then they also maximize the symmetric α -stability. Here, dictator functions refer to the functions $f_d := 1\{x_k = 1\}$ or $1\{x_k = -1\}$ for some $1 \leq k \leq n$.

The study of the noise stability problem, or more precisely, a two-function (or two-set) version of the noise stability problem called the *non-interactive correlation distillation (NICD) problem*, dates back to Gács and Körner's and Witsenhausen's seminal papers [9, 10]. By utilizing the tensorization property of the maximal correlation, Witsenhausen [10] showed that for $\alpha = 2$, the asymmetric and symmetric 2-stability w.r.t. $a = 1/2$ are attained by dictator functions. The symmetric α -stability problem with $\alpha \in \{3, 4, 5, \dots\}$ was studied by Mossel and O'Donnell [7], but only the case $\alpha = 3$ was solved by them. For $\alpha = 3$, the maximal symmetric α -stability w.r.t. $a = 1/2$ is attained by dictator functions, and moreover, Mossel and O'Donnell observed that this is not true for $\alpha = 10$. Mossel and O'Donnell conjectured² that dictator functions maximize the symmetric α -stability over all balanced Boolean functions for all $\alpha \in \{4, 5, \dots, 9\}$. Since for $\alpha = 1$, the symmetric 1-mutual information

¹The concept " α -stability" for real α was introduced in [5] for Gaussian distributions and in [6] for binary distributions. The special case with α being a positive integer was introduced in [7]. More precisely, the " α -stability" defined there refers to the Φ -stability with $\Phi : t \mapsto t^\alpha$ for the asymmetric version and $\Phi : t \mapsto t^\alpha + (1-t)^\alpha$ for the symmetric version. Hence, our definition is a linear transform version of theirs. We abuse the term " α -stability" here, since when the mean a is given, computing the maximal Φ -stabilities defined in [6] is equivalent to computing the maximal Φ -stabilities defined here.

²In fact, this is a stronger version of the original conjecture posed by Mossel and O'Donnell. In their original version, the Boolean functions are additionally restricted to be *antisymmetric* (or *odd*).

is nothing but the Shannon mutual information, the maximal 1-stability problem for this case can be interpreted as the problem of maximizing the Shannon mutual information $I(f(\mathbf{Y}); \mathbf{X})$ over all Boolean functions f with a given mean. In fact, this case was already studied by Courtade and Kumar [11, 12], but this problem still remains widely open, except for the extreme cases. Courtade and Kumar conjectured³ that for the mean $a = 1/2$, dictator functions maximize the symmetric 1-stability over all balanced Boolean functions. This conjecture attracts lots of interest from different fields [6, 13–17], and it is regarded as one of the most fundamental conjectures at the interface of information theory and the analysis of Boolean functions. Along these lines, Li and Médard [6] conjectured that for $\alpha \in (1, 2)$, the maximal asymmetric α -stability is still attained by dictator functions. Here we unify and slightly generalize the Mossel–O’Donnell conjecture, Courtade–Kumar conjecture, and Li–Médard conjecture in the following two conjectures.

Conjecture 1 (Maximal Asymmetric α -Stability Conjecture). *For $0 \leq \rho \leq 1$ and $\alpha \in [1, 9]$, $\text{MaxStab}_\alpha(\frac{1}{2})$ is attained by dictator functions.*

Conjecture 2 (Maximal Symmetric α -Stability Conjecture). *For $0 \leq \rho \leq 1$ and $\alpha \in [1, 9]$, $\text{MaxStab}_\alpha^{\text{sym}}(\frac{1}{2})$ is attained by dictator functions.*

Obviously, Conjecture 1 implies Conjecture 2. Furthermore, as mentioned above, Conjectures 1 and 2 with $\alpha = 2$ were proven by Witsenhausen [10]; Conjecture 2 with $\alpha = 3$ was proven by Mossel and O’Donnell [7]; the Mossel–O’Donnell conjecture corresponds to Conjecture 2 with $\alpha \in \{4, 5, \dots, 9\}$; the Courtade–Kumar conjecture corresponds to Conjecture 2 with $\alpha = 1$; and the Li–Médard conjecture corresponds to Conjecture 1 with $\alpha \in (1, 2)$. Conjecture 1 is open for $\alpha \in [1, 2) \cup (2, 9]$, and Conjecture 2 is open for $\alpha \in [1, 2) \cup (2, 3) \cup (3, 9]$.

Recently, Barnes and Özgür [18] showed that Conjecture 1 with $\alpha = 1$ and the same one but with $\alpha \in (1, 2)$ are equivalent, and Conjecture 2 with $\alpha = 1$ and the same one but with $\alpha \in (1, 2)$ are also equivalent. That is, the asymmetric (resp. symmetric) version of Courtade–Kumar conjecture and the asymmetric (resp. symmetric) version of Li–Médard conjecture are equivalent. Following Barnes and Özgür’s proofs in [18, Subsections III.C and III.D], one can obtain the following lemma, which slightly generalizes the “only if” parts of Barnes and Özgür’s Theorems 1 and 2 in [18].

Lemma 1. *For $a = 1/2$, there are two thresholds α_{\min} and α_{\max} satisfying $1 \leq \alpha_{\min} \leq 2 \leq \alpha_{\max}$ such that dictator functions are optimal in attaining the asymmetric max α -stability with $\alpha \geq 1$ if and only if $\alpha \in [\alpha_{\min}, \alpha_{\max}]$. This statement also holds for the symmetric max α -stability but with possibly different thresholds $\check{\alpha}_{\min}$ and $\check{\alpha}_{\max}$ satisfying the same condition $1 \leq \check{\alpha}_{\min} \leq 2 \leq \check{\alpha}_{\max}$.*

Here are some consequences of this lemma. Firstly, in terms of these thresholds, Conjectures 1 and 2 can be restated as that $\alpha_{\min} = \check{\alpha}_{\min} = 1$ and $\alpha_{\max}, \check{\alpha}_{\max} \geq 9$. Secondly, as mentioned previously, it was shown by Mossel and O’Donnell [7] that $\max\{\alpha_{\max}, \check{\alpha}_{\max}\} < 10$. Lastly, Since Conjecture 2 holds for $\alpha = 3$ [7], we actually have $\check{\alpha}_{\max} \geq 3$, i.e., Conjecture 2 holds for all $\alpha \in [2, 3]$. In other words, Conjecture 2 is only open for $\alpha \in [1, 2) \cup (3, 9]$. Combining all these points yields that $2 \leq \alpha_{\max} < 10$

³In fact, this is a weaker version of the original conjecture posed by Courtade and Kumar. In their original version, the Boolean functions are not restricted to be balanced.

and $3 \leq \check{\alpha}_{\max} < 10$. If Conjectures 1 and 2 are true, then the estimates of α_{\max} and $\check{\alpha}_{\max}$ can be improved to $9 \leq \alpha_{\max}, \check{\alpha}_{\max} < 10$.

1.1 Our Contributions

Our main contributions in this paper are:

Theorem 1 (Formally Stated in Corollary 2). *For (ρ, α) such that*

$$0 \leq \rho \leq \begin{cases} \rho^* & \alpha \in [1, 2) \\ 1 & \alpha \in [2, 5] \end{cases},$$

$\text{MaxStab}_{\alpha}^{\text{sym}}(\frac{1}{2})$ is attained by dictator functions, where $\rho^* \approx 0.461491$ (the solution to (16)).

Theorem 2 (Formally Stated in Corollary 3). *For (ρ, α) such that*

$$0 \leq \rho \leq \begin{cases} \frac{1-\theta(\alpha)}{1+\theta(\alpha)} & \alpha \in (1, 2) \\ 1 & \alpha \in [2, 3] \end{cases},$$

$\text{MaxStab}_{\alpha}(\frac{1}{2})$ is attained by dictator functions, where $\theta(\alpha)$ is some function (the solution to (20)).

For symmetric α -stability, our bound in Theorem 1 improves the previously best known bound $\check{\alpha}_{\max} \geq 3$ to $\check{\alpha}_{\max} \geq 5$, and for the asymmetric case, our bound in Theorem 2 improves the previously best known bound $\alpha_{\max} \geq 2$ to $\alpha_{\max} \geq 3$. In other words, we have verified the Mossel–O’Donnell conjecture for all $2 \leq \alpha \leq 5$ in the symmetric setting, and for all $2 \leq \alpha \leq 3$ in the asymmetric setting. Our results for $3 < \alpha \leq 5$ in the symmetric setting and for $2 < \alpha \leq 3$ in the asymmetric setting are new. As for the Courtade–Kumar conjecture and Li–Médard conjecture, we improve Samorodnitsky’s result [16] in the sense that we provide an explicit dimension-independent threshold $\rho^* \approx 0.461491$ for which the symmetric versions of the Courtade–Kumar and Li–Médard conjectures hold for all $\rho \in [0, \rho^*]$.

Theorems 1 and 2 are proven by combining discrete Fourier analysis and optimization theory. In fact, they are consequences of a general bound on the maximal Φ -stability derived in this paper by using these techniques. Furthermore, we also improve our bound for symmetric α -stability by incorporating improvements of Friedgut–Kalai–Naor (FKN) theorem [19] into our method. Our improved bounds are presented in the optimization form, which seems difficult to simplify. Numerical evaluation of this improved bound implies that the value of the threshold ρ^* in Theorem 1 can be improved to 0.83. Our improvements of the FKN theorem are sharp or asymptotically sharp for certain cases.

1.2 Related Works

We next summarize the literature on the noise stability problem. Although the study of the noise stability problem originated in the seminal papers [9, 10, 20], the term *noise stability* was first introduced by Benamini, Kalai, and Schramm [21]; see a brief

survey on [22, p. 68]. In [21], only the case of $\alpha = 2$ was studied. As mentioned previously, this was subsequently generalized to the cases of $\alpha > 2$, $\alpha = 1$, and $1 < \alpha < 2$ in [6, 7, 12], and the conjectures mentioned above were posed along with these generalizations. In fact, a weaker version (the two-function version) of Courtade–Kumar conjecture was solved by Pichler, Piantanida, and Matz [17] by using Fourier analysis. Specifically, they showed that $I(f(\mathbf{Y}); g(\mathbf{X}))$ is maximized by a pair of identical dictator functions over all Boolean functions (f, g) , including but not limited to balanced Boolean functions. In fact, if (f, g) are additionally restricted to be balanced, then this result is just a consequence of Witsenhausen’s maximal correlation bound [10]. In other words, Pichler, Piantanida, and Matz’s contribution is addressing the unbalanced case. However, the situation is totally different for the single-function version of Courtade–Kumar conjecture, since the latter is open even for the balanced case. A classic bound on the mutual information $I(f(\mathbf{Y}); \mathbf{X})$ for the balanced case is ρ^2 , proven by Witsenhausen and Wyner [23] (also see [24]). Such a bound can be proved via the so-called *Mrs. Gerber’s lemma* [25] or the *hypercontractivity inequality* [22]. Ordentlich, Shayevitz, and Weinstein [15] improve this bound to a sharper one for small ρ . This new bound turns out to be asymptotically sharp in the limiting case $\rho \rightarrow 0$. In 2016, Samorodnitsky [16] made a significant breakthrough on the Courtade–Kumar conjecture. Specifically, he proved the existence of a *dimension-independent* threshold ρ_0 for which the Courtade–Kumar conjecture holds for all $\rho \in [0, \rho_0]$. However, the value of ρ_0 was not explicitly given in his paper, and required to be “sufficiently small”. A weaker version of this result in which the threshold ρ_0 is replaced by a sequence that vanishes as $n \rightarrow \infty$ was also proven in [15, 26]. In addition, by considering a variant of noise model, Eldan, Mikulincer, and Raghavendra recently prove a variant version of the Courtade–Kumar conjecture [27]. Their proof is based on the so-called renormalized Brownian motion.

The noise stability of *unbalanced* Boolean functions was also widely investigated in the literature. For the mean $a = 1/4$, by combining Fourier analysis with a coding-theoretic result, the present author and Tan [28] showed that the indicator functions of $(n - 2)$ -subcubes (in n -dimensional discrete cube) maximize the 2-stability with $a = 1/4$. When the mean is small, hypercontractivity inequalities are an effective tool to address this case. In particular, Kahn, Kalai, and Linial [29] first applied the single-function version of (forward) hypercontractivity inequalities to obtain bounds for the 2-stability problem, by substituting the nonnegative functions in the hypercontractivity inequalities with the Boolean functions. Mossel and O’Donnell [22, 30] applied the two-function version of hypercontractivity inequalities to obtain bounds for the two-function version of α -stability problem in a similar way. Kahn, Kalai, and Linial’s result as well as Mossel and O’Donnell’s are known as *small-set expansion theorems*. Kamath and Anantharam [31] slightly strengthened the small-set expansion theorems via utilizing hypercontractivity inequalities in a slightly different way. All these bounds derived by hypercontractivity inequalities are asymptotically sharp in certain sense when the mean approaches zero [22]. A variant of the α -stability problem with the means vanishing exponentially as $n \rightarrow \infty$ was studied in [32–35]. Various stronger versions of hypercontractivity inequalities were derived or used to obtain sharper bounds on the α -stability in these papers. The optimal exponent for the α -stability problem in

this variant setting was explicitly given in entropy optimization forms in [34, 35], not only for the binary random vectors (\mathbf{X}, \mathbf{Y}) , but also for the random vectors defined on very general spaces (Polish spaces).

The maximal Φ -stability problem in the Gaussian setting with Φ restricted to be convex and increasing was fully resolved by Borell [20] in 1985. In particular, he showed that the Φ -stability is maximized by the indicators of half-spaces over all measurable Boolean functions $f : \mathbb{R}^n \rightarrow \{0, 1\}$ of the same measure; see an explicit statement of this result in [14]. Such a result is known as *Borell's Isoperimetric Theorem*. The Gaussian analogues of the Courtade–Kumar conjecture and Li–Médard conjecture were consequences of Borell's Isoperimetric Theorem, which were also proved respectively by Kindler, O'Donnell, and Witmer [14] and by Eldan [5] using alternative approaches.

1.3 Organization

This paper is organized as follows. In Sections 2 and 3, we present our main results and the improvements in detail. The proofs of the main results and related lemmas, propositions, and corollaries are provided in Sections 4-9.

2 Main Results

In this paper, we aim at determining the maximal Φ -stability for given volume $a \in (0, 1)$. By using Fourier analysis, we first prove a general bound on the maximal Φ -stability.

Theorem 3. For $a, \rho \in (0, 1)$, $\text{MaxStab}_\Phi(a) \leq \Gamma_\rho(a)$, where

$$\Gamma_\rho(a) := \sup_{P_{Z|S}} \mathbb{E}[\Phi(a + \rho Z)] \quad (2)$$

$$\text{s.t. } 0 \leq a + \rho Z \leq 1 \quad \text{a.s.} \quad (3)$$

$$\mathbb{E}[Z] = 0 \quad (4)$$

$$\mathbb{E}[Z^2] \leq \mathbb{E}[SZ] \quad (5)$$

with⁴

$$P_S(s) = \begin{cases} 1 - a & s = -a \\ a & s = 1 - a \end{cases}. \quad (6)$$

The supremum in (2) is taken over all conditional probability mass functions (conditional pmf) $P_{Z|S}$, and the random variables (S, Z) in the objective function and constraints follow the joint distribution $P_{SZ} := P_S P_{Z|S}$.

Note that the constraint in (5) effectively dominates (i.e., upper bounds) the energy of Z , since obviously, this constraint implies $\mathbb{E}[Z^2] \leq \mathbb{E}[SZ] \leq \mathbb{E}[S^2] = a(1-a)$. On the other hand, Φ is convex, and hence, the objective function in (2) is dominated if the energy of Z is dominated. In other words, the constraint (5) dominates the objective function via dominating the energy of Z .

⁴Throughout this paper, we use P_X to denote the probability mass function of a random variable X , and use $P_{Y|X}$ to denote the conditional probability mass function of a random variable Y given X .

Proof of Theorem 3. Our proof relies on Boolean Fourier analysis [22]. Consider the Fourier basis⁵ $\{\chi_S\}_{S \subseteq [n]}$ with $\chi_S(\mathbf{x}) := \prod_{i \in S} x_i$ for $S \subseteq [n]$. Then for a function $f : \{-1, 1\}^n \rightarrow \mathbb{R}$, define its Fourier coefficients as

$$\hat{f}_S := \mathbb{E}_{\mathbf{X} \sim \text{Unif}\{-1, 1\}^n} [f(\mathbf{X}) \chi_S(\mathbf{X})], \quad S \subseteq [n].$$

Then the Fourier expansion of the function f (cf. [22, Equation (1.6)]) is

$$f(\mathbf{x}) = \sum_{S \subseteq [n]} \hat{f}_S \chi_S(\mathbf{x}).$$

The *degree- k Fourier weight* of f is defined as

$$\mathbf{W}_k[f] := \sum_{S: |S|=k} \hat{f}_S^2, \quad k \in [0 : n].$$

For brevity, we denote $\mathbf{W}_k[f]$ as \mathbf{W}_k .

By the definition of Fourier weights, it is easily seen that for a Boolean function with volume $\mathbb{E}f = a$,

$$\mathbf{W}_0 = a^2 \quad \text{and} \quad \sum_{k=0}^n \mathbf{W}_k = a.$$

Define the *degree- k part* of f as

$$f_k(\mathbf{x}) := \sum_{S \subseteq [n]: |S|=k} \hat{f}_S \chi_S(\mathbf{x}).$$

Then it is easy to check that

$$\begin{aligned} T_\rho f(\mathbf{x}) &= \sum_{k=0}^n \rho^k f_k(\mathbf{x}) \\ \sum_{k=0}^n f_k(\mathbf{x}) &= f(\mathbf{x}) \in [0, 1], \quad \forall \mathbf{x} \\ f_0(\mathbf{x}) &= a, \quad \forall \mathbf{x} \\ \sum_{k=0}^n \mathbb{E}[f_k(\mathbf{X})^2] &= \sum_{k=0}^n \mathbf{W}_k = a \\ \mathbb{E}[f_k(\mathbf{X}) f_{k'}(\mathbf{X})] &= 0, \quad 0 \leq k' < k \leq n. \end{aligned}$$

Denote

$$Z := \sum_{k=1}^n \rho^{k-1} f_k(\mathbf{X}) \tag{7}$$

⁵Throughout this paper, we denote $[m : n] := \{m, m+1, \dots, n\}$. When $m = 1$, for brevity, we denote $[n] := [1 : n]$.

$$S := f - a = \sum_{k=1}^n f_k(\mathbf{X}). \quad (8)$$

Then

$$T_\rho f(\mathbf{X}) = a + \rho Z, \quad \mathbb{E}Z = 0,$$

and S follows the distribution in (6). Observe that

$$\mathbb{E}[Z^2] = \sum_{k=1}^n \rho^{2(k-1)} \mathbf{W}_k \leq \sum_{k=1}^n \rho^{k-1} \mathbf{W}_k = \mathbb{E}[SZ].$$

Hence $\mathbf{MaxStab}_\Phi(a) \leq \Gamma_\rho(a)$. \square

We next provide several properties of the optimization problem in (2). Denote $\mathcal{S} := \{-a, 1-a\}$. The set $A := \{(\Phi(a + \rho z), z, z^2 - sz) : 0 \leq a + \rho z \leq 1\}$ lies in the 3-dimensional Euclidean space, and $\mathbb{E}[(\Phi(a + \rho Z), Z, Z^2 - sZ) | S = s]$ lies in the convex hull of A . Since A is connected, by Bunt's extension of Carathéodory's theorem [36] or a more general version [37, Theorem 18], each point in the convex hull of A is a convex combination of at most 3 points in A . This further implies that without loss of optimality, it suffices to restrict $P_{Z|S}$ in the optimization in (2) such that $|\text{supp}(P_{Z|S=s})| \leq 3$ for each $s \in \mathcal{S}$. As a consequence, the supremum in (2) is actually a maximum, since the feasible region is compact and the objective and constraint functions are continuous.

For the optimization problem in (2), we also claim that without loss of optimality, it suffices to restrict $|\text{supp}(P_Z)| \leq 4$. First, observe that the $\sup_{P_{Z|S}}$ in (2) can be rewritten as three-fold optimizations: $\sup_{\mathcal{Z}} \sup_{P_{S|Z}} \sup_{P_Z}$, where the first sup is taken over all countable subsets \mathcal{Z} of \mathbb{R} with each element z satisfying $0 \leq a + \rho z \leq 1$, the second sup is taken over all conditional pmfs $P_{S|Z}$ (specifically, $\{P_{S|Z=z}, z \in \mathcal{Z}\}$), and the last sup is taken over all pmfs P_Z on \mathcal{Z} such that $\sum_{z \in \mathcal{Z}} P_{S|Z}(-a|z) P_Z(z) = P_S(-a)$ (the analogous equality for $s = 1-a$ automatically holds if this equality holds) and constraints (4) and (5) hold. Here by convention, if there is no feasible solution for a supremization, its value is set to $-\infty$. Given \mathcal{Z} and $P_{S|Z}$, the third sup above, \sup_{P_Z} , is in fact a linear program. Following arguments similar to the one in the last paragraph, one can restrict $|\text{supp}(P_Z)| \leq 4$.

Let $m \geq 3$. For $s \in \mathcal{S}$, denote $(z_{s,i})_{i \in [m]} \in \mathbb{R}^m$ as a vector consisting of all (possibly repetitions of) the elements in the support of $P_{Z|S=s}$. Denote $(p_{s,i})_{(s,i) \in \mathcal{S} \times [m]}$ as a joint pmf on $\mathcal{S} \times [m]$ whose marginal on \mathcal{S} is P_S . Then, the optimization in (2) is equivalently taken over all the tuples $(z_{s,i}, p_{s,i})_{(s,i) \in \mathcal{S} \times [m]}$ such that $\sum_{i \in [m]} p_{s,i} = P_S(s), \forall s \in \mathcal{S}$. Without loss of optimality, we may additionally assume $p_{s,i} > 0, \forall s \in \mathcal{S}, i \in [m]$, since if $p_{s,i} = 0, p_{s,j} > 0$, then we can re-choose $p_{s,i} \leftarrow \frac{p_{s,j}}{2}, p_{s,j} \leftarrow \frac{p_{s,j}}{2}, z_{s,i} \leftarrow z_{s,j}$ which leads to the same distribution P_{SZ} . Then, the arguments above lead us to define

$$\Lambda_\rho^{(m)}(a) := \max_{z_{s,i} \in \mathbb{R}, p_{s,i} > 0, (s,i) \in \mathcal{S} \times [m]} \sum_{(s,i) \in \mathcal{S} \times [m]} p_{s,i} \Phi(a + \rho z_{s,i}) \quad (9)$$

$$\text{s.t. } \sum_{i \in [m]} p_{s,i} = P_S(s), \forall s \in \mathcal{S}, \quad (10)$$

$$0 \leq a + \rho z_{s,i} \leq 1, \forall (s,i) \in \mathcal{S} \times [m],$$

$$\sum_{(s,i) \in \mathcal{S} \times [m]} p_{s,i} z_{s,i} = 0, \quad (11)$$

$$\sum_{(s,i) \in \mathcal{S} \times [m]} p_{s,i} (z_{s,i}^2 - s z_{s,i}) \leq 0. \quad (12)$$

As discussed above, without changing the value of the optimization, we can add the additional constraint $|\{z_{s,i}\}_{(s,i) \in \mathcal{S} \times [m]}| \leq 4$ into the maximization problem in (9). Combining all the points above, we arrive at the following results.

Proposition 1. *For $a, \rho \in (0, 1)$ and $m \geq 3$, the following hold.*

1. *It holds that $\Gamma_\rho(a) = \Lambda_\rho^{(m)}(a)$. (Hence we denote $\Lambda_\rho(a) := \Lambda_\rho^{(m)}(a)$ for $m \geq 3$.)*
2. *Any optimal solution to the optimization problem in (2) (or the equivalent one in (9)) satisfies $\mathbb{E}[Z^2] = \mathbb{E}[SZ]$.*
3. *Any optimal solution to the maximization problem in (9) satisfies that $z_{1-a,j} \geq z_{-a,i} + \frac{1}{2}$ for all $i, j \in [m]$ such that $z_{-a,i} > \frac{-a}{\rho}$, $z_{1-a,j} < \frac{1-a}{\rho}$. Moreover, this condition is satisfied by some $i, j \in [m]$.*
4. *The linear independence constraint qualification (LICQ)⁶ is satisfied for the maximization problem in (9).*

Statement 1 of Proposition 1 has already been derived above, and Statements 2-4 are proven in Section 4.

Proposition 1 enables us to employ the Karush–Kuhn–Tucker (KKT) conditions to simplify the bound in Theorem 3. In fact, the bound in Theorem 3 (or the equivalent one in (9)) is general enough to apply to several common cases, e.g., the asymmetric and symmetric α -stabilities, as follows.

2.1 Symmetric Φ

We first consider symmetric Φ . Here Φ is said to be symmetric (w.r.t. $\frac{1}{2}$) if $\Phi(\frac{1}{2} - t) = \Phi(\frac{1}{2} + t)$ for $t \in [0, \frac{1}{2}]$. Define

$$\bar{\Gamma}_\rho(a) := \sup_{(z_1, z_2) \in \bar{\mathcal{Z}}} (1 - a - p)\Phi(0) + p\Phi(a + \rho z_1) + q\Phi(a + \rho z_2) + (a - q)\Phi(1),$$

where

$$p = \frac{(1-a)a(1-\rho)}{(1-\rho-\rho z_1+\rho z_2)(a+\rho z_1)}, \quad (13)$$

$$q = \frac{(1-a)a(1-\rho)}{(1-\rho-\rho z_1+\rho z_2)(1-a-\rho z_2)}, \quad (14)$$

⁶The LICQ for a maximizer point \mathbf{x}^* is the condition that the gradients of the active inequality constraints and the gradients of the equality constraints are linearly independent at \mathbf{x}^* . We refer readers to [38, 39] for more details on LICQ.

and

$$\bar{Z} := \left\{ (z_1, z_2) \in \left(\frac{-a}{\rho}, \frac{1-a}{\rho} \right)^2 : z_2 - z_1 \geq \frac{1}{2}, \right. \\ \left. z_2 - z_1 \geq \frac{(1-\rho)z_2}{1-a-\rho z_2}, z_2 - z_1 \geq -\frac{(1-\rho)z_1}{a+\rho z_1} \right\}.$$

Define

$$\hat{\Gamma}_\rho(a) := \sup_{\max\{-\frac{a}{\rho}, c\} \leq \tilde{z} \leq \frac{-a(1+b)}{2(1-a)}} (1-a-p)\Phi(a+\rho\tilde{z}) + p\Phi(a+\rho(b-\tilde{z})) + a\Phi(a+\rho\hat{z}), \quad (15)$$

where $b = \frac{1-2a}{\rho}$, $c = \frac{1}{2}(b - \sqrt{\frac{a+2ab+b^2}{1-a}})$, and

$$\hat{z} = \frac{1}{2} \left(\sqrt{\frac{\Delta}{a}} + b + 1 \right), \\ p = \frac{-\sqrt{a\Delta} - (a(b+1-2\hat{z}) + 2\hat{z})}{2(b-2\hat{z})}$$

with $\Delta = a(b^2 - 4b\hat{z} + 2b + 4\hat{z}^2 + 1) + 4\hat{z}(b - \hat{z})$. For symmetric Φ , we prove the following result, whose proof is provided in Section 5.

Theorem 4. *Assume that Φ is symmetric and continuous on $[0, 1]$, and differentiable on $(0, 1)$ whose derivative Φ' is increasing and continuous. Assume $\rho \in (0, 1)$.*

1) *If additionally, Φ' is strictly concave on $(0, \frac{1}{2}]$, then for $a \in (0, \frac{1}{2}]$, $\mathbf{MaxStab}_\Phi(a) \leq \Lambda_\rho(a) = \bar{\Gamma}_\rho(a)$. Moreover, for $a = \frac{1}{2}$, we can additionally assume $z_1 + z_2 = 0$ in the supremization in (43).*

2) *If additionally, Φ' is strictly convex on $(0, \frac{1}{2}]$, then for $a \in (0, \frac{1}{2}]$, $\mathbf{MaxStab}_\Phi(a) \leq \Lambda_\rho(a) = \max\{\hat{\Gamma}_\rho(a), \hat{\Gamma}_\rho(1-a), \mathbb{E}[\Phi(a+\rho S)]\}$. In particular, for $a = \frac{1}{2}$, we have $\mathbf{MaxStab}_\Phi(\frac{1}{2}) \leq \Lambda_\rho(\frac{1}{2}) = \Phi(\frac{1-\rho}{2})$.*

Note that Φ_α^{sym} with $\alpha \in (1, 2) \cup (3, \infty)$ satisfies the assumption in Statement 1 of Theorem 4, and Φ_α^{sym} with $\alpha \in (2, 3)$ satisfies the assumption in Statement 2 of Theorem 4. We now apply Theorem 4 to the symmetric α -stability. Let ρ^* be the solution in $(0, 1)$ to the equation

$$\psi(\rho) := (1+\rho^2) \log\left(\frac{1+\rho}{2}\right) - (1-\rho)^2 \log\left(\frac{1-\rho}{2}\right) = 0. \quad (16)$$

We have that $\rho^* \approx 0.461491$. Then we prove the Courtade–Kumar conjecture for $\rho \leq \rho^*$. The proof is provided in Section 6.

Corollary 1. $\mathbf{MaxStab}_1^{\text{sym}}(\frac{1}{2})$ is attained by dictator functions for $0 \leq \rho \leq \rho^*$.

Samorodnitsky [16] proved the Courtade–Kumar conjecture for $\rho \in [0, \rho_0]$, where $0 < \rho_0 < 1$ is some absolute constant. However, the value of ρ_0 was not explicitly given in [16], and also it was assumed to be “sufficiently small”. In contrast, the value of ρ^* in Corollary 1 is explicitly given. As for the proof ideas, Samorodnitsky’s proof

in [16] and our proof in this paper are both based on Fourier analysis (and also quantitative versions of Friedgut–Kalai–Naor theorems which are used to further improve the threshold in Section 3.2), but use it in different ways. For example, different from the linear programs in [16], the duality for *nonconvex* programs is used in this paper.

It is worth noting that in [16], the mean a is not fixed. For this case, we numerically evaluate $\sup_{a \in [0, 1/2]} \bar{\Gamma}_\rho(a) - \Phi_1^{\text{sym}}(a)$, and observe that this supremum is attained at $a = 1/2$ for $0 \leq \rho \leq \rho^*$. This indicates that Statement 1 of Theorem 4 seems to imply the original Courtade–Kumar conjecture for $0 \leq \rho \leq \rho^*$, i.e., the conjecture that dictator functions maximize the mutual information $I(f(\mathbf{Y}); \mathbf{X})$ over all Boolean functions f . It remains to find a proof for this observation in the future.

We also generalize the corollary above to the case $\alpha \in [1, 5]$. The proof of the following corollary is provided in Section 7.

Corollary 2. *For (ρ, α) such that $0 \leq \rho \leq \begin{cases} \rho^* & \alpha \in [1, 2) \\ 1 & \alpha \in [2, 5] \end{cases}$, $\text{MaxStab}_\alpha^{\text{sym}}(\frac{1}{2})$ is attained by dictator functions.*

Corollary 2 resolves the Mossel–O’Donnell conjecture for $\alpha \in [2, 5]$. However, our result for cases $\alpha = 2, 3$ is not new. The case $\alpha = 2$ was resolved by Witsenhausen [10] by using the tensorization property of maximal correlation, and the case $\alpha = 3$ was resolved by Mossel and O’Donnell [7] by reducing the case $\alpha = 3$ to the case $\alpha = 2$. Note that Mossel and O’Donnell’s method seems difficult to extend to the case $\alpha > 3$. Moreover, the case $\alpha \in [2, 3]$ can be also obtained by combining Mossel and O’Donnell’s result and Lemma 1. Our Corollary 2, proven by using Fourier analysis, is a generalization of Witsenhausen’s and Mossel–O’Donnell’s results. Moreover, combining Corollary 2 and the counterexample example for $\alpha = 10$ in [7] yields that the estimation of $\check{\alpha}_{\max}$ is improved to $5 \leq \check{\alpha}_{\max} \leq 10$.

2.2 Asymmetric Φ

We next consider asymmetric Φ . Define⁷

$$\tilde{\Gamma}_\rho(a) = \sup_{0 \leq z_2 \leq 1-a} (1-a-p)\Phi(0) + p\Phi(a + \rho z_1) + a\Phi(a + \rho z_2),$$

where

$$z_1 = \frac{z_2(\rho(1-z_2) - a)}{1-a-\rho z_2}, \quad (17)$$

$$p = \frac{a(1-a-\rho z_2)^2}{a + \rho^2 z_2 - (a + \rho z_2)^2}. \quad (18)$$

Then following proof steps similar to those of Theorem 4, we have the following result. The proof is omitted.

⁷Indeed, the range of z_2 can be further restricted to $\frac{1-a}{2-\rho} \leq z_2 \leq 1-a$. For simplicity, we only restrict $0 \leq z_2 \leq 1-a$, which is sufficient to show our result on the asymmetric α -stability in Corollary 3.




Fig. 1: The region of $0 \leq \rho \leq \frac{1-\theta(\alpha)}{1+\theta(\alpha)}$.

Theorem 5. *If Φ is continuous on $[0, 1]$ and differentiable on $(0, 1)$ whose derivative Φ' is increasing, continuous, and strictly concave on $(0, 1)$, then for $a, \rho \in (0, 1)$,*

$$\mathbf{MaxStab}_{\Phi}(a) \leq \Lambda_{\rho}(a) = \tilde{\Gamma}_{\rho}(a). \quad (19)$$

Remark 1. If Φ' is increasing, continuous, and strictly convex, then by redefining $\tilde{\Phi} : t \in [0, 1] \mapsto \Phi(1 - t)$ and substituting $a \leftarrow 1 - a$, (19) still holds.

We now consider the asymmetric α -stability. Denote $\theta(\alpha)$ as the solution to

$$\theta^{2-\alpha} + \frac{1}{\alpha}(1 - \theta) = 1 \quad (20)$$

with unknown θ for given $\alpha \in (1, 2)$. Then the following holds, whose proof is provided in Section 8.

Corollary 3. *For (ρ, α) such that $0 \leq \rho \leq \begin{cases} \frac{1-\theta(\alpha)}{1+\theta(\alpha)} & \alpha \in (1, 2) \\ 1 & \alpha \in [2, 3] \end{cases}$, $\mathbf{MaxStab}_{\alpha}(\frac{1}{2})$ is attained by dictator functions.*

This result partially resolves the Li-Médard conjecture for the case of $0 \leq \rho \leq \frac{1-\theta(\alpha)}{1+\theta(\alpha)}$, $\alpha \in (1, 2)$. This region is plotted in Fig. 1. The question whether $\mathbf{MaxStab}_{\alpha}(\frac{1}{2})$ is attained by dictator functions for the case $\alpha = 3$ was posed in [6]. Our Corollary 3 gives a positive answer to this question. Combining Corollary 3 and the counterexample for $\alpha = 10$ in [7] implies $3 \leq \alpha_{\max} \leq 10$.

3 Further Improvement of Our Results

In this section, we further improve our bounds on the maximal Φ -stability by applying an improved version of the Friedgut–Kalai–Naor (FKN) theorem [19]. Hence, to this end, we improve the FKN theorem first.

3.1 Improvements of FKN Theorem

The FKN theorem concerns about which Boolean functions f on the discrete cube have Fourier coefficients concentrated at the lowest two levels. It states that such Boolean functions are close to either a constant function (i.e., $f = 0$ or 1) or a dictator function ($f = 1\{x_i = 1\}$ or $1\{x_i = -1\}$). Here we aim at improving the FKN theorem by focusing on the class of Boolean functions with a given mean. For this case, the FKN theorem can be also formulated as a theorem about maximizing the first-order Fourier weight of a Boolean function given the maximum of its correlations to all dictator functions. We next provide the formulation.

For $a, \beta \in 2^{-n}[0 : 2^n]$, define

$$W^{(n)}(a, \beta) := \max_{f: \{-1, 1\}^n \rightarrow \{0, 1\}: \mathbb{E}f = a, \max_{i \in [n]} |\hat{f}_{\{i\}}| = \beta} \mathbf{W}_1[f].$$

The quantity $W^{(n)}(a, \beta)$ was implicitly studied by Friedgut, Kalai, and Naor [19] who showed that for $a = \frac{1}{2}$, $W(\frac{1}{2}, \beta) \rightarrow \frac{1}{4}$ if and only if $\beta \rightarrow \frac{1}{2}$.

Similarly, we define

$$W^{(n)}(a) := \max_{f: \{-1, 1\}^n \rightarrow \{0, 1\}: \mathbb{E}f = a} \mathbf{W}_1[f].$$

Here without ambiguity, we use the same notation $W^{(n)}$ but with different numbers of parameters to denote two different functions $W^{(n)}(a, \beta)$ and $W^{(n)}(a)$. We first use $W^{(n)}(a)$ to bound $W^{(n)}(a, \beta)$ in the following lemma. The proof of Lemma 2 is provided in Section 9.

Lemma 2. *For any Boolean function $f : \{-1, 1\}^n \rightarrow \{0, 1\}$, we have*

$$\mathbf{W}_1[f] \leq \beta^2 + \left(\sqrt{W^{(n)}(a) - a^2} + \sqrt{W^{(n-1)}(a - \beta)} \right)^2 \quad (21)$$

where $a := \mathbb{E}f$ and $\beta := |\hat{f}_{\{1\}}|$. In particular, for balanced Boolean functions (i.e., $a = \frac{1}{2}$), we have

$$\mathbf{W}_1[f] \leq \beta^2 + W^{(n-1)}\left(\frac{1}{2} - \beta\right). \quad (22)$$

Moreover, for any possible values of β such that there exists at least one Boolean function with $\hat{f}_{\{1\}} = \beta$, equality in (22) is attained by some balanced Boolean function.

Remark 2. (21) implies that $W^{(n)}(a, \beta)$ is no larger than the RHS of (21), but (21) is indeed stronger than this conclusion in the sense that given β , (21) holds for any Boolean functions f such that there is at least one i satisfying $|\hat{f}_{\{i\}}| = \beta$, without requiring other $j \in [n] \setminus \{i\}$ satisfying $|\hat{f}_{\{j\}}| \leq \beta$.

In fact, the quantity $W^{(n)}(a)$ was relatively well studied in the literature. Define

$$\phi(t) := \begin{cases} 2t^2 \ln \frac{1}{t} & 0 < t \leq \frac{1}{4} \\ 2t^2(\frac{1}{\sqrt{t}} - 1) & \frac{1}{4} < t \leq \frac{1}{2} \\ \frac{t}{2} & \frac{1}{2} < t \leq 1 \end{cases}.$$

Define for $0 \leq t \leq 1$,

$$\varphi(t) := \phi(\min\{t, 1-t\}).$$

It is known that [22, 40, 41] for any $f : \{-1, 1\}^n \rightarrow \{0, 1\}$ with $\mathbb{E}f = a \in [0, 1]$,

$$W^{(n)}(a) \leq \varphi(a). \quad (23)$$

The bounds $\frac{t}{2}$ and $2t^2(\frac{1}{\sqrt{t}} - 1)$ were respectively proven by Fu, Wei, and Yeung [40] and the present author and Tan [41], by using linear programming methods (together with MacWilliams–Delsarte identities). The bound $2t^2 \ln \frac{1}{t}$ is called *Chang’s bound* [42], which can be proven by several methods, e.g., by hypercontractivity inequalities [22].

Combining Lemma 2 and (23) yields the following result.

Proposition 2. For $0 \leq \beta \leq a \leq \frac{1}{2}$,

$$W^{(n)}(a, \beta) \leq \beta^2 + (\sqrt{\varphi(a) - a^2} + \sqrt{\varphi(a - \beta)})^2. \quad (24)$$

In particular, for $0 \leq \beta \leq a = \frac{1}{2}$,

$$W^{(n)}(\frac{1}{2}, \beta) \leq \beta^2 + \varphi(\frac{1}{2} - \beta).$$

This bound improves the existing bound proven in [43]. Moreover, our bound is sharp for $a = \frac{1}{2}, \beta = \frac{1}{4}$, and asymptotically sharp for $a = \frac{1}{2}, \beta \uparrow \frac{1}{2}$. The sharpness for $a = \frac{1}{2}, \beta = \frac{1}{4}$ can be seen from that for $\beta = \frac{1}{4}$, $W^{(n-1)}(\frac{1}{4}) = \varphi(\frac{1}{4}) = \frac{1}{8}$ for $n \geq 3$, which is attained by $1\{(x_2, \dots, x_n) : x_2 = x_3 = 1\}$. The asymptotic sharpness for $a = \frac{1}{2}, \beta \uparrow \frac{1}{2}$ can be seen from that if we define $\mathcal{A}_n = \{(x_2, \dots, x_n) : \frac{1}{\sqrt{n-1}} \sum_{i=2}^n x_i \geq r_n\}$ with $\lim_{n \rightarrow \infty} r_n = Q^{-1}(\varepsilon)$ as a sequence of Hamming balls with volumes asymptotically approaching ε , where Q is the Q-function, then we have that for $\beta = \frac{1}{2} - \varepsilon$, $\lim_{n \rightarrow \infty} \mathbf{W}_1[1_{\mathcal{A}_n}] = \psi(Q^{-1}(\varepsilon)) \sim 2\varepsilon^2 \ln \frac{1}{\varepsilon}$ as $\varepsilon \downarrow 0$ [22], where ψ is the probability density function of the standard Gaussian.

As mentioned in Remark 2, in Lemma 2 we have not used the information $|\hat{f}_{\{i\}}| \leq \beta$ for all $i \in [2 : n]$. Hence, Lemma 2 (or Proposition 2) cannot provide a good bound when β is small, since for this case $|\hat{f}_{\{i\}}|$ could be significantly larger than β for some $i \in [2 : n]$. In order to obtain a good bound for this case, we use another method to derive the following quantitative (non-asymptotic) version of FKN theorem, which is a consequence of the variant of Khintchine’s inequality proven by König, Schütt, and Tomczak-Jaegermann [44]. It is worth noting that the variant of Khintchine’s inequality proven in [44] was also applied by Friedgut, Kalai, and Naor to prove the original (asymptotic) version of the FKN theorem [19].

Proposition 3. For $0 \leq \beta \leq a = \frac{1}{2}$,

$$W^{(n)}\left(\frac{1}{2}, \beta\right) \leq \frac{1}{4} \left(\sqrt{4\left(\frac{1}{2} - \frac{1}{\sqrt{2\pi}}\right)\beta} + \frac{1}{2\pi} + \frac{1}{\sqrt{2\pi}} \right)^2. \quad (25)$$

Proof. We have that

$$\mathbf{W}_1[f] = \mathbb{E} \left[\sum_{i=1}^n \hat{f}_{\{i\}} X_i f(\mathbf{X}) \right] \quad (26)$$

$$\leq \mathbb{E} \left[\sum_{i=1}^n \hat{f}_{\{i\}} X_i \mathbf{1} \left\{ \sum_{i=1}^n \hat{f}_{\{i\}} X_i \geq 0 \right\} \right] \quad (27)$$

$$\begin{aligned} &= \frac{1}{2} \mathbb{E} \left[\left| \sum_{i=1}^n \hat{f}_{\{i\}} X_i \right| \right] \\ &\leq \frac{1}{2} \left(\sqrt{\frac{2}{\pi}} \sqrt{\mathbf{W}_1[f]} + \left(1 - \sqrt{\frac{2}{\pi}}\right)\beta \right), \end{aligned} \quad (28)$$

where (27) follows since if we relax $(\hat{f}_{\{i\}})_{i \in [n]}$ and f to be independent quantities, then given $\hat{f}_{\{i\}}$, the Boolean function $f : \mathbf{x} \mapsto \mathbf{1} \left\{ \sum_{i=1}^n \hat{f}_{\{i\}} x_i \geq 0 \right\}$ maximizes the expectation in (26), and (28) follows from the following variant of Khintchine's inequality

$$\left| \mathbb{E} \left[\left| \sum_{i=1}^n c_i X_i \right| \right] - \sqrt{\frac{2}{\pi}} \|\mathbf{c}\|_2 \right| \leq \left(1 - \sqrt{\frac{2}{\pi}}\right) \|\mathbf{c}\|_\infty$$

with $\mathbf{c} := (c_1, c_2, \dots, c_n)$ which was proven by König, Schütt, and Tomczak-Jaegermann [44]. Solving the inequality in (28), we obtain (25). \square

3.2 Further Improvement of Our Bounds on Noise Stability

Based on upper bounds on $W^{(n)}(a, \beta)$ (e.g., (24) and (25)), we improve Theorem 3 as follows.

Theorem 6. Let $\omega(a, \beta)$ be an upper bound on $W^{(n)}(a, \beta)$. Then for $a, \rho \in (0, 1)$, $\text{MaxStab}_\Phi(a) \leq \Upsilon_\rho(a)$, where

$$\Upsilon_\rho(a) := \sup_{\beta, P_{Z|SX}} \mathbb{E}[\Phi(a + \rho Z)] \quad (29)$$

$$\text{s.t. } 0 \leq \beta \leq a, 1 - a,$$

$$0 \leq a + \rho Z \leq 1 \quad \text{a.s.}$$

$$\mathbb{E}Z = 0,$$

$$\mathbb{E}[XZ] = \beta, \quad (30)$$

$$\mathbb{E}[Z^2] \leq (1 - \rho)\omega(a, \beta) + \rho\mathbb{E}[SZ], \quad (31)$$

with

$$P_{SX}(s, x) = \begin{cases} \frac{1-a+\beta}{2} & (s, x) = (-a, -1) \\ \frac{1-a-\beta}{2} & (s, x) = (-a, 1) \\ \frac{a-\beta}{2} & (s, x) = (1-a, -1) \\ \frac{a+\beta}{2} & (s, x) = (1-a, 1) \end{cases}. \quad (32)$$

Proof. Recall that $\{\hat{f}_S\}$ are the Fourier coefficients of f . WLOG, we assume $|\hat{f}_{\{1\}}| \geq |\hat{f}_{\{i\}}|$ for all $2 \leq i \leq n$. If $\hat{f}_{\{1\}} \geq 0$, denote $\beta := \hat{f}_{\{1\}}$ and $X := X_1$; otherwise, $\beta := -\hat{f}_{\{1\}}$ and $X := -X_1$. Hence $\beta \geq 0$. Denote

$$Y = \sum_{i=2}^n \hat{f}_{\{i\}} X_i, \quad \hat{S} = \sum_{k=2}^n f_k(\mathbf{X}), \quad \hat{Z} = \sum_{k=2}^n \rho^{k-2} f_k(\mathbf{X}).$$

Recall the definitions of S, Z in (7) and (8). Then,

$$\begin{aligned} S &= f - a = \beta X + Y + \hat{S} \\ Z &= \beta X + Y + \rho \hat{Z} \\ \mathbb{E}[XZ] &= \beta \\ \mathbb{E}[SZ] &= \mathbf{W}_1 + \rho \mathbb{E}[\hat{S}\hat{Z}], \end{aligned} \quad (33)$$

and (S, X) follows the distribution in (32) (since $\mathbb{E}[SX] = \beta$). Observe that

$$\mathbb{E}[\hat{Z}^2] = \sum_{k=2}^n \rho^{2(k-2)} \mathbf{W}_k \leq \sum_{k=2}^n \rho^{k-2} \mathbf{W}_k = \mathbb{E}[\hat{S}\hat{Z}]. \quad (34)$$

Combining (33), (34), and the fact that $\mathbf{W}_1 \leq \omega(a, \beta)$, yields that

$$\begin{aligned} \mathbb{E}[Z^2] &= \mathbf{W}_1 + \rho^2 \mathbb{E}[\hat{Z}^2] \\ &\leq (1-\rho) \mathbf{W}_1 + \rho(\mathbf{W}_1 + \rho \mathbb{E}[\hat{S}\hat{Z}]) \\ &\leq (1-\rho) \omega(a, \beta) + \rho \mathbb{E}[SZ]. \end{aligned}$$

Hence $\mathbf{MaxStab}_\Phi(a) \leq \Upsilon_\rho(a)$. \square

Here $\omega(a, \beta)$ can be chosen as the upper bound in (24) for any a or as the minimum of the upper bounds in (24) and (25) for $a = 1/2$. Compared to Theorem 3, the bound in Theorem 6 is based on a more elaborate analysis, which introduces the maximum of absolute values of first-order Fourier coefficients as a parameter, i.e., β , and then optimizes the bound over all possible β . To better understand the intuition of Theorem 6, we focus on the case $a = 1/2$. For this case, if β is close to $1/2$, then (30) forces Z to be almost linearly dependent of the Boolean r.v. X (note that the energy of Z is controlled under the constraint and (31)). This in turn forces optimal solutions to (29) close to dictator functions (for which $f(\mathbf{X}) = \frac{1+X}{2}$, and $Y = \hat{S} = \hat{Z} = 0, S = Z = X/2$ where $X = X_i$ or $-X_i$ for some $i \in [n]$). On the other hand, if β is far from $1/2$,

then $\omega(a, \beta)$ becomes relatively small (see (24) and (25)), which, combined with (31), forces the energy of Z to be small. This in turn decreases the objective function, and hence, this kind of β is excluded from being optimal to the supremization in (29).

Denote $\mathcal{X} := \{\pm 1\}$. For simplicity, we next focus on the case $a = 1/2$, and suppose that $\omega(\beta) := \omega(1/2, \beta)$ is continuous in β and satisfies the trivial inequality $\omega(\beta) \leq \frac{1}{4}$ for all β (since $\mathbf{W}_0 + \mathbf{W}_1 \leq \mathbb{E}[f^2] = a$ and $\mathbf{W}_0 = a^2$). Similarly to the definition of $\Lambda_\rho^{(m)}(a)$ in (9), for such $\omega(\beta)$ and $m \geq 4$, we define

$$\begin{aligned} \Psi_\rho^{(m)} := & \max_{\substack{\beta, z_{s,x,i} \in \mathbb{R}, \\ p_{s,x,i} > 0, \\ (s,x,i) \in \mathcal{S} \times \mathcal{X} \times [m]}} \sum_{(s,x,i) \in \mathcal{S} \times \mathcal{X} \times [m]} p_{s,x,i} \Phi\left(\frac{1}{2} + \rho z_{s,x,i}\right) & (35) \\ \text{s.t. } & 0 \leq \beta \leq \frac{1}{2}, \\ & 0 \leq \frac{1}{2} + \rho z_{s,x,i} \leq 1, \forall (s,x,i) \in \mathcal{S} \times \mathcal{X} \times [m], \\ & \sum_{i \in [m]} p_{s,x,i} = P_{SX}(s,x), \forall (s,x) \in \mathcal{S} \times \mathcal{X}, \\ & \sum_{(s,x,i) \in \mathcal{S} \times \mathcal{X} \times [m]} p_{s,x,i} z_{s,x,i} = 0, \\ & \sum_{(s,x,i) \in \mathcal{S} \times \mathcal{X} \times [m]} p_{s,x,i} x z_{s,x,i} = \beta, \\ & \sum_{(s,x,i) \in \mathcal{S} \times \mathcal{X} \times [m]} p_{s,x,i} (z_{s,x,i}^2 - \rho s z_{s,x,i}) \leq (1 - \rho)\omega(\beta). \quad (36) \end{aligned}$$

Following proof steps similar to those of Proposition 1, one can show that the optimization problem in (29) satisfies the following properties. The proof is provided in Section 10.

Proposition 4. *Suppose that⁸ Φ is symmetric and continuous on $[0, 1]$, and differentiable on $(0, 1)$ whose derivative Φ' is increasing and continuous $(0, 1)$, and strictly concave on $(0, \frac{1}{2}]$. Suppose that $\omega(\beta)$ is continuous in β and satisfies that $\omega(\beta) \leq \frac{1}{4}$ for all β . For $a = 1/2$, $\rho \in (0, 1)$, and $m \geq 4$, the following hold.*

1. *Without loss of optimality, it suffices to restrict that $|\text{supp}(P_Z)| \leq 7$ and $|\text{supp}(P_Z|_{S=s, X=x})| \leq 4$ for $(s, x) \in \mathcal{S} \times \mathcal{X}$, which implies that $\Upsilon_\rho(\frac{1}{2}) = \Psi_\rho^{(m)}$.*
2. *Any optimal solution to the optimization problem in (29) with $a = 1/2$ (or the equivalent one in (35)) satisfies $\mathbb{E}[Z^2] = (1 - \rho)\omega(\beta) + \rho\mathbb{E}[SZ]$.*
3. *Any optimal solution to the maximization problem in (35) satisfies that $z_{\frac{1}{2}, x, j} \geq z_{-\frac{1}{2}, x, i} + \frac{1}{2}$ for all $i, j \in [m]$, $x \in \mathcal{X}$ such that $z_{-\frac{1}{2}, x, i} > \frac{-1}{2\rho}$, $z_{\frac{1}{2}, x, j} < \frac{1}{2\rho}$.*
4. *For even $m \geq 8$, there is an optimal solution P_{SXZ} to the maximization problem in (35) such that $P_{SXZ}(-s, -x, -z) = P_{SXZ}(s, x, z)$ for all (s, x, z) , which either satisfies*

⁸This guarantees the existence of the optimal solutions to the optimization problem in (29) or the equivalent one in (35).

the LICQ or can be expressed as

$$P_{SXXZ}(s, x, z) = \begin{cases} \frac{1+2\beta}{4} & (s, x, z) = (-\frac{1}{2}, -1, z_1) \\ \frac{1-2\beta}{4} & (s, x, z) = (\frac{1}{2}, -1, z_2) \\ \frac{1-2\beta}{4} & (s, x, z) = (-\frac{1}{2}, 1, -z_2) \\ \frac{1+2\beta}{4} & (s, x, z) = (\frac{1}{2}, 1, -z_1) \end{cases} \quad (37)$$

for some z_1, z_2 such that $-\frac{1}{2\rho} \leq z_1 \leq z_2 \leq \frac{1}{2\rho}$.

Define

$$\bar{\Upsilon}_\rho := \max_{\beta \in [0, \frac{1}{2}]} \max_{(z_1, z_2) \in \check{\mathcal{Z}}} h(\beta, z_1, z_2), \quad (38)$$

where

$$\check{\mathcal{Z}} := \{(z_1, z_2) : -\frac{1}{2\rho} \leq z_1 \leq z_2 \leq \frac{1}{2\rho}, 0 \leq p \leq \frac{1}{4} + \frac{\beta}{2}, 0 \leq q \leq \frac{1}{4} - \frac{\beta}{2}\},$$

and

$$h(\beta, z_1, z_2) := (1 - 2p - 2q)\Phi(0) + 2p\Phi(\frac{1}{2} + \rho z_1) + 2q\Phi(\frac{1}{2} + \rho z_2),$$

with

$$p = \frac{(1 - \rho)(1 + \rho - 4\rho^2\omega(\beta) + 2\beta(1 + 2\rho z_2 - \rho^2))}{4(1 + 2\rho z_1)(1 + \rho z_2 - \rho z_1 - \rho^2)}, \quad (39)$$

$$q = \frac{(1 - \rho)(1 + \rho - 4\rho^2\omega(\beta) - 2\beta(1 - 2\rho z_1 - \rho^2))}{4(1 - 2\rho z_2)(1 + \rho z_2 - \rho z_1 - \rho^2)}. \quad (40)$$

Theorem 7. *If Φ is symmetric and continuous on $[0, 1]$ and differentiable on $(0, 1)$ whose derivative Φ' is increasing and continuous $(0, 1)$, and strictly concave on $(0, \frac{1}{2}]$, then $\text{MaxStab}_\Phi(\frac{1}{2}) \leq \bar{\Upsilon}_\rho$.*

The proof of Theorem 7 is provided in Section 11 which is similar to that of Theorem 4.

We now focus on the case $\Phi = \Phi_1^{\text{sym}}$. Numerical results show that for the case $\Phi = \Phi_1^{\text{sym}}$, $\bar{\Upsilon}_\rho$ in Theorem 7 (with $\omega(\beta)$ taken as the minimum of the upper bounds in (24) and (25)) is (almost) attained by dictator functions for $0 \leq \rho \leq 0.83$, which means that the threshold ρ^* in Corollary 1 can be further improved to a value around 0.83. It remains to find a proof for this observation in the future. We are also interested in introducing new techniques to attack the case with $\rho > 0.83$, since the optimality of dictator functions for this case seemingly cannot be proven by our present method.

In following sections, we provide proofs for the results stated above.

4 Proof of Proposition 1

4.1 Statement 2

We use a perturbation method. Suppose $P_{Z|S}$ is an optimal solutions to (2) such that $\mathbb{E}[Z^2] < \mathbb{E}[SZ]$. We first assume that $a + \rho z = 0$ or 1 for all z such that $P_Z(z) > 0$,

which is equivalent to $Z \in \{-a/\rho, (1-a)/\rho\}$ (with probability one). By (4), we know that $P_Z(z) = \begin{cases} 1-a & z = -a/\rho \\ a & z = (1-a)/\rho \end{cases}$. From this, we have $\mathbb{E}[SZ] \leq \sqrt{\mathbb{E}[Z^2]\mathbb{E}[S^2]} = \rho\mathbb{E}[Z^2]$ which contradicts with (5). Hence, our assumption is false, or equivalently, $0 < a + \rho z^* < 1$ for some z^* such that $P_Z(z^*) > 0$.

Let s^* be such that $P_{SZ}(s^*, z^*) > 0$. We next construct a new conditional distribution $Q_{Z|S}$ by setting $Q_{Z|S}(z|s) = P_{Z|S}(z|s)$ for all $(s, z) \neq (s^*, z^*)$, and

$$Q_{Z|S}(z^* - \delta|s^*) = Q_{Z|S}(z^* + \delta|s^*) = \frac{1}{2}P_{Z|S}(z^*|s^*),$$

where $\delta > 0$ is small enough such that $0 < a + \rho(z^* \pm \delta) < 1$. By the choice of δ , (3) holds. It is easy to see that (4) still holds. Furthermore, observe that the RHS of (5) remains unchanged, and the LHS of (5) is continuous in δ . Hence for sufficiently small but positive δ , (5) still holds. Since Φ is strictly convex, $\mathbb{E}[\Phi(a + \rho Z)]$ increases after replacing $P_{Z|S}$ with $Q_{Z|S}$. This contradicts with the optimality of $P_{Z|S}$. This completes the proof of Statement 2.

4.2 Statement 3

We continue to use a perturbation method to prove Statement 3. Let $(z_{s,i}, p_{s,i})_{(s,i) \in \mathcal{S} \times [m]}$ be an optimal solution to the maximization in (9). Let P_{SZ} be the joint distribution induced by $(z_{s,i}, p_{s,i})_{(s,i) \in \mathcal{S} \times [m]}$. By Statement 2, $\mathbb{E}_P[Z^2] = \mathbb{E}_P[SZ]$ under P_{SZ} .

Suppose that $z_{-a,i} > z_{1-a,j}$ for some $i, j \in [m]$. Then, for a sufficiently small $\epsilon > 0$, we define a new distribution Q_{SZ} by replacing $(z_{-a,i}, p_{-a,i}) \leftarrow (z_{-a,i}, p_{-a,i} - \epsilon)$, $(z_{1-a,j}, p_{1-a,j}) \leftarrow (z_{1-a,j}, p_{1-a,j} - \epsilon)$ and introducing new points $(z_{-a,m+1}, p_{-a,m+1}) \leftarrow (z_{1-a,j}, \epsilon)$, $(z_{1-a,m+1}, p_{1-a,m+1}) \leftarrow (z_{-a,i}, \epsilon)$. We do not change other parameters. For this new distribution, it is easily seen that the marginal distributions of Z and S are unchanged. So, constraints in (10)-(11) are still satisfied by Q_{SZ} , and the value of the objective function induced by Q_{SZ} remains unchanged. As for constraint in (12), $\mathbb{E}[Z^2]$ remains the same, and $\mathbb{E}_Q[SZ] - \mathbb{E}_P[SZ] = \epsilon(z_{-a,i} - z_{1-a,j}) > 0$. So, $\mathbb{E}_Q[SZ] > \mathbb{E}_P[SZ] = \mathbb{E}_P[Z^2] = \mathbb{E}_Q[Z^2]$. Hence, (12) is satisfied by Q_{SZ} , which further implies that Q_{SZ} is an optimal solution to (9). However, by Statement 2, $\mathbb{E}_Q[SZ] = \mathbb{E}_Q[Z^2]$ should hold, which leads to a contradiction. Therefore, $z_{-a,i} \leq z_{1-a,j}$ for all $i, j \in [m]$.

Suppose that $z_{-a,i} \leq z_{1-a,j} < z_{-a,i} + \frac{1}{2}$ for some $i, j \in [m]$ such that $z_{-a,i} > \frac{-a}{\rho}$, $z_{1-a,j} < \frac{1-a}{\rho}$. Then, we define a new distribution Q_{SZ} by replacing $z_{-a,i} \leftarrow z_{-a,i} - \epsilon$, $z_{1-a,j} \leftarrow z_{1-a,j} + \epsilon'$ for some $\epsilon, \epsilon' > 0$ such that $p_{-a,i}\epsilon = p_{1-a,j}\epsilon'$. By definition, Q_{SZ} does not change $\mathbb{E}[Z]$, but it enlarges $\mathbb{E}[SZ] - \mathbb{E}[Z^2]$ for small enough $\epsilon, \epsilon' > 0$ since

$$\begin{aligned} & (\mathbb{E}_Q[SZ] - \mathbb{E}_Q[Z^2]) - (\mathbb{E}_P[SZ] - \mathbb{E}_P[Z^2]) \\ &= (p_{-a,i}\epsilon a + p_{1-a,j}\epsilon' \bar{a}) - (-p_{-a,i}\epsilon(2z_{-a,i} - \epsilon) + p_{1-a,j}\epsilon'(2z_{1-a,j} + \epsilon')) \\ &= p_{-a,i}\epsilon(1 - 2(z_{1-a,j} - z_{-a,i})) - (\epsilon + \epsilon'), \end{aligned}$$

which is positive for small enough $\epsilon, \epsilon' > 0$. Hence, Q_{SZ} is feasible. On the other hand, Q_{SZ} also enlarges the objective function. This is because, by denoting $\hat{\Phi}(z) = \Phi(a + \rho z)$,

$$\begin{aligned} & \mathbb{E}_Q[\hat{\Phi}(Z)] - \mathbb{E}_P[\hat{\Phi}(Z)] \\ &= p_{-a,i} \hat{\Phi}(z_{-a,i} - \epsilon) + p_{1-a,j} \hat{\Phi}(z_{1-a,j} + \epsilon') - p_{-a,i} \hat{\Phi}(z_{-a,i}) - p_{1-a,j} \hat{\Phi}(z_{1-a,j}) \\ &= (p_{1-a,j} \epsilon') \frac{\hat{\Phi}(z_{1-a,j} + \epsilon') - \hat{\Phi}(z_{1-a,j})}{\epsilon'} - (p_{-a,i} \epsilon) \frac{\hat{\Phi}(z_{-a,i}) - \hat{\Phi}(z_{-a,i} - \epsilon)}{\epsilon} \end{aligned}$$

which is positive since $p_{1-a,j} \epsilon' = p_{-a,i} \epsilon$, and by the convexity of Φ , $\frac{\hat{\Phi}(z_{1-a,j} + \epsilon') - \hat{\Phi}(z_{1-a,j})}{\epsilon'} > \frac{\hat{\Phi}(z_{-a,i}) - \hat{\Phi}(z_{-a,i} - \epsilon)}{\epsilon}$ (note that $z_{1-a,j} \geq z_{-a,i}$). Hence, Q_{SZ} induces a larger value of the objective function than P_{SZ} , which contradicts with the optimality of P_{SZ} . Hence, $z_{1-a,j} \geq z_{-a,i} + \frac{1}{2}$ for all $i, j \in [m]$ such that $z_{-a,i} > \frac{-a}{\rho}$, $z_{1-a,j} < \frac{1-a}{\rho}$.

If $\{z_{1-a,j}\}_{j \in [m]} = \{\frac{1-a}{\rho}\}$, then by (36), $\{z_{-a,i}\}_{i \in [m]} = \{\frac{-a}{\rho}\}$. Hence, we have $Z = \frac{S}{\rho}$, and hence, $\mathbb{E}[Z^2] = \frac{1}{\rho} \mathbb{E}[SZ]$, which contradicts with $\mathbb{E}[Z^2] \leq \mathbb{E}[SZ]$. Therefore, $z_{1-a,j} < \frac{1-a}{\rho}$ for some $j \in [m]$. Similarly, one can show that $z_{-a,i} > \frac{-a}{\rho}$ for some $i \in [m]$.

4.3 Statement 4

Suppose that LICQ is not satisfied. Denote $(z_{s,i}, p_{s,i})_{(s,i) \in \mathcal{S} \times [m]}$ as an optimal solution to the maximization in (9). Here and subsequently, we assume the first m components are indexed by $\{-a\} \times [m]$ and the last m indexed by $\{1-a\} \times [m]$. Denote $\mathcal{I} := \{(s,i) : a + \rho z_{s,i} = 0\}$ and $\mathcal{J} := \{(s,i) : a + \rho z_{s,i} = 1\}$. By Statement 3, $\mathcal{I} \subseteq \{-a\} \times [m]$ and $\mathcal{J} \subseteq \{1-a\} \times [m]$.

The gradients of the active inequality constraints (including (5) by Statement 2) and the gradients of the equality constraints constitute the $(|\mathcal{I}| + |\mathcal{J}| + 4) \times 4m$ matrix

$$G := \begin{bmatrix} \rho \mathbf{I}_{\mathcal{I}} & \mathbf{0}_{1 \times 2m} \\ -\rho \mathbf{I}_{\mathcal{J}} & \mathbf{0}_{1 \times 2m} \\ \mathbf{0}_{1 \times 2m} & (\mathbf{I}_{1 \times m}, \mathbf{0}_{1 \times m}) \\ \mathbf{0}_{1 \times 2m} & (\mathbf{0}_{1 \times m}, \mathbf{I}_{1 \times m}) \\ (p_{s,i})_{s,i} & (z_{s,i})_{s,i} \\ (p_{s,i}(2z_{s,i} - s))_{s,i} & (z_{s,i}^2 - s z_{s,i})_{s,i} \end{bmatrix}.$$

Here, $\mathbf{I}_{\mathcal{I}}$ denotes the $\{0, 1\}$ -matrix of size $|\mathcal{I}| \times 2m$ with each row containing exactly one “1”, and the “1” at the r -th row is located at the column indexed by the r -th element in \mathcal{I} . Obviously, the last m columns of $\mathbf{I}_{\mathcal{I}}$ consist of zeros. The matrix $\mathbf{I}_{\mathcal{J}}$ is defined similarly, and hence, the first m columns of $\mathbf{I}_{\mathcal{J}}$ consist of zeros. The assumption that LICQ is not satisfied implies that

$$\hat{G} := \begin{bmatrix} (p_{s,i})_{(s,i) \in (\mathcal{I} \cup \mathcal{J})^c} & (z_{s,i} - z_{s,1})_{(s,i) \in \mathcal{S} \times [2:m]} \\ (p_{s,i}(2z_{s,i} - s))_{(s,i) \in (\mathcal{I} \cup \mathcal{J})^c} & ((z_{s,i} - z_{s,1})(z_{s,i} + z_{s,1} - s))_{(s,i) \in \mathcal{S} \times [2:m]} \end{bmatrix}$$

is not of full rank. We next prove this is impossible.

Since the submatrix $\begin{bmatrix} (p_{s,i})_{(s,i) \in (\mathcal{I} \cup \mathcal{J})^c} \\ (p_{s,i}(2z_{s,i} - s))_{(s,i) \in (\mathcal{I} \cup \mathcal{J})^c} \end{bmatrix}$ of \hat{G} is not of full rank, we know that $z_{-a,i} = z_1$ or $-a/\rho$ and $z_{1-a,i} = z_2$ or $(1-a)/\rho$ for all $i \in [m]$ and some z_1, z_2 such that $-a - 2z_1 = 1 - a - 2z_2$. Furthermore, since the submatrix $\begin{bmatrix} (p_{-a,i})_{(-a,i) \in (\mathcal{I} \cup \mathcal{J})^c} & (z_{-a,i} - z_{-a,1})_{i \in [2:m]} \\ (p_{-a,i}(2z_1 + a))_{(-a,i) \in (\mathcal{I} \cup \mathcal{J})^c} & ((z_{-a,i} - z_{-a,1})(z_{-a,i} + z_{-a,1} + a))_{i \in [2:m]} \end{bmatrix}$ of \hat{G} is also not of full rank, we know that $z_{-a,i}$'s are identical for all i , and combining this with Statement 3 implies that $z_{-a,i} = z_1$ for all $i \in [m]$. Similarly, $z_{1-a,i} = z_2$ for all $i \in [m]$. Additionally, z_1, z_2 should satisfy $\mathbb{E}[Z] = 0, \mathbb{E}[Z^2] = \mathbb{E}[SZ]$, and hence, we have $Z = 0$ or $Z = S$. This contradicts with $-a - 2z_1 = 1 - a - 2z_2$, which implies that LICQ is satisfied.

5 Proof of Theorem 4

The Lagrangian of the optimization problem in (9) is

$$\begin{aligned} \mathcal{L} := & \sum_{s,i} p_{s,i} \left\{ \Phi(a + \rho z_{s,i}) + \theta_{s,i}(a + \rho z_{s,i}) + \theta'_{s,i}(1 - (a + \rho z_{s,i})) + \lambda(sz_{s,i} - z_{s,i}^2) + \eta z_{s,i} \right\} \\ & + \sum_s \mu_s \left(\sum_i p_{s,i} - P_S(s) \right). \end{aligned}$$

Since LICQ is satisfied, the KKT theorem, e.g., [39, Theorem 5.3.1], is valid, and then we obtain the following first-order necessary conditions for (local) optimal solutions:

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial z_{s,i}} &= p_{s,i} [\rho(\Phi'(a + \rho z_{s,i}) + \theta_{s,i} - \theta'_{s,i}) + \lambda(s - 2z_{s,i}) + \eta] = 0 \\ \frac{\partial \mathcal{L}}{\partial p_{s,i}} &= \Phi(a + \rho z_{s,i}) + \theta_{s,i}(a + \rho z_{s,i}) + \theta'_{s,i}(1 - (a + \rho z_{s,i})) + \lambda(sz_{s,i} - z_{s,i}^2) + \eta z_{s,i} + \mu_s \\ &= \Phi(a + \rho z_{s,i}) + \lambda(sz_{s,i} - z_{s,i}^2) + \eta z_{s,i} + \mu_s = 0 \\ 0 &\leq a + \rho z_{s,i} \leq 1, \forall (s,i) \in \mathcal{S} \times [m] \\ \sum_{s,i} p_{s,i} z_{s,i} &= 0 \\ \sum_{s,i} p_{s,i} (sz_{s,i} - z_{s,i}^2) &= 0 \\ \sum_i p_{s,i} &= P_S(s), \forall s \in \mathcal{S} \\ p_{s,i} &> 0, \forall (s,i) \in \mathcal{S} \times [m] \\ \theta_{s,i}(a + \rho z_{s,i}) &= 0 \\ \theta'_{s,i}(1 - a - \rho z_{s,i}) &= 0 \\ \lambda \geq 0, \vec{\theta}, \vec{\theta}' &\geq \vec{0}, \end{aligned} \tag{41}$$




Fig. 2: Plots of $z \mapsto \rho\Phi'(a + \rho z)$, $z \mapsto \lambda(2z + a) - \eta$, and $z \mapsto \lambda(2z + a - 1) - \eta$.

where the equality in (41) follows from Statement 2 of Proposition 1. Here, $\vec{\theta}, \vec{\theta}' \geq \vec{0}$ denotes all the components of $\vec{\theta}, \vec{\theta}'$ are nonnegative.

For (s, i) such that $-a/\rho < z_{s,i} < (1-a)/\rho$, we have $\theta_{s,i} = \theta'_{s,i} = 0$. Denote

$$g(s, z) := \rho\Phi'(a + \rho z) + \lambda(s - 2z) + \eta, \quad (42)$$

$$G(s, z) := \int_0^z g(s, t) dt.$$

Hence, for (s, i) such that $-a/\rho < z_{s,i} < (1-a)/\rho$, it holds that

$$\frac{1}{p_{s,i}} \frac{\partial \mathcal{L}}{\partial z_{s,i}} = g(s, z_{s,i}) = 0, \quad (43)$$

and moreover, for all (s, i) , it holds that

$$\frac{\partial \mathcal{L}}{\partial p_{s,i}} = G(s, z_{s,i}) + \Phi(a) + \mu_s = 0. \quad (44)$$

Equations (43) and (44) imply that given s , for all $z \in \{z_{s,i}\}_{i \in [m]}$, $g(s, z)$ is always equal to 0 and $G(s, z)$ remains the same.

5.1 Statement 1

By assumption, Φ is symmetric w.r.t. $\frac{1}{2}$ and Φ' is increasing, continuous on $(0, 1)$, and strictly concave on $(0, \frac{1}{2}]$. Hence, for each s , $g(s, z) = 0$ with z unknown has at most three distinct solutions. For the case of three distinct solutions, denote the solutions as z_1, z_2, z_3 such that $-a/\rho =: z_0 < z_1 < z_2 < z_3 < z_4 := (1-a)/\rho$. Obviously, for these solutions, the values of $G(s, z_i)$ and $G(s, z_{i+1})$ for $i \in \{1, 2\}$ are different (note that G is an antiderivative of g). Hence, given s , z_i and z_{i+1} cannot be solutions of (44) at the same time. This is also true when $g(s, z) = 0$ (with z unknown) has two distinct solutions.

On the other hand, by Statement 3 of Proposition 1, for each s , $\{z_{s,i}\}_{i \in [m]}$ contains at least one solution of (43) and (44). Moreover, the solutions to $g(-a, z) = 0$ are the

intersection of the curve $z \mapsto \rho\Phi'(a + \rho z)$ and the line $z \mapsto \lambda(2z + a) - \eta$ (with a positive slope), and the solutions to $g(1 - a, z) = 0$ are the intersection points of the same curve $z \mapsto \rho\Phi'(a + \rho z)$ and another line $z \mapsto \lambda(2z + a - 1) - \eta$ which is parallel to the previous one. See Fig. 2a.

From the points above, we claim that for each s , $\{z_{s,i}\}_{i \in [m]}$ cannot contain two distinct solutions of (43) and (44). This is because, for example, for $s = -a$, if $\{z_{s,i}\}_{i \in [m]}$ contains two distinct solutions of (43) and (44), then (43) and (44) must have three distinct solutions, and the smallest and the largest among them, denoted by z_1, z_3 respectively, must be contained in $\{z_{s,i}\}_{i \in [m]}$. In this case, all the solutions of (43) and (44) with $s = 1 - a$ are smaller than z_3 (by the observation in the paragraph above), and hence cannot be contained in $\{z_{1-a,i}\}_{i \in [m]}$ (by the first part of Statement 3 of Proposition 1). This contradicts with (the second part of) Statement 3 of Proposition 1. Hence, the claim holds for $s = -a$. For $s = 1 - a$, the claim can be proven similarly.

By the claim above,

$$\{z_{-a,i}\}_{i \in [m]} \subseteq \{-a/\rho, \hat{z}_1\} \quad \text{and} \quad \{z_{1-a,i}\}_{i \in [m]} \subseteq \{(1-a)/\rho, \hat{z}_2\}, \quad (45)$$

where \hat{z}_1 is a solution of (43) and (44) with $s = -a$, and \hat{z}_2 is a solution of (43) and (44) with $s = 1 - a$. By Statement 3 of Proposition 1, $\hat{z}_2 - \hat{z}_1 \geq \frac{1}{2}$. This means, optimal solutions P_{SZ} to the optimization problem in (9) satisfy

$$P_{SZ}(s, z) = \begin{cases} 1 - a - p & s = -a, z = \frac{-a}{\rho} \\ p & s = -a, z = \hat{z}_1 \\ q & s = 1 - a, z = \hat{z}_2 \\ a - q & s = 1 - a, z = \frac{1-a}{\rho} \end{cases}$$

for some $(p, q, \hat{z}_1, \hat{z}_2)$ such that $0 \leq p \leq 1 - a, 0 \leq q \leq a, -\frac{a}{\rho} < \hat{z}_1 \leq \hat{z}_2 < \frac{1-a}{\rho}$, $\hat{z}_2 - \hat{z}_1 \geq \frac{1}{2}$, and

$$\mathbb{E}Z = 0, \quad \mathbb{E}[SZ] = \mathbb{E}[Z^2]. \quad (46)$$

Solving equations in (46) with respect to unknowns (z_1, z_2) , we obtain p, q given in (13) and (14). Since $\hat{z}_2 \geq \hat{z}_1, 0 \leq p \leq 1 - a$, and $0 \leq q \leq a$, we know that

$$\begin{aligned} \hat{z}_2 - \hat{z}_1 &\geq \frac{(1-\rho)\hat{z}_2}{1-a-\rho\hat{z}_2} \\ \hat{z}_2 - \hat{z}_1 &\geq -\frac{(1-\rho)\hat{z}_1}{a+\rho\hat{z}_1}. \end{aligned}$$

Hence, the first part of Statement 1 holds.

We next prove the second part of Statement 1, in which $a = 1/2$. Denote δ_x as the Dirac measure at x . For an optimal solution $z_{s,i}, p_{s,i}, (s, i) \in \mathcal{S} \times [m]$ to the optimization problem in (9) which must satisfy (45), we now construct a new distribution $Q_{SZ} = \sum_{s,i} \frac{1}{2} p_{s,i} \cdot (\delta_{s,z_{s,i}} + \delta_{-s,-z_{s,i}})$ which is still optimal for the optimization in (9) with

m replaced by $2m$. Obviously, Q_{SZ} should also satisfy (45) since the above arguments still work when m is replaced by $2m$. Hence, $\hat{z}_1 + \hat{z}_2 = 0$ holds.

5.2 Statement 2

We next consider the case in which Φ' is increasing, continuous on $(0, 1)$, and strictly convex on $(0, \frac{1}{2}]$. Let $z_{s,i}, p_{s,i}, (s, i) \in \mathcal{S} \times [m]$ be an optimal solution to the maximization problem in (9). We next derive necessary conditions for the optimality of this solution. For this case, the equation (43) still has at most three distinct solutions.

We first consider the case $g(-a, -a/\rho) > 0$ with g defined in (42). For this case, at $(s, z) = (-a, -a/\rho)$, $\frac{1}{p_{s,i}} \frac{\partial \mathcal{L}}{\partial z_{s,i}} \Big|_{(-a, -a/\rho)} = g(-a, -a/\rho) + \theta > 0$. Hence, $\{z_{-a,i}\}_{i \in [m]}$ does not contain $-a/\rho$, and does contain at most two distinct solutions of (43) and (44) with $s = -a$ (by arguments similar to those in the previous subsection).

We now claim that if $g(-a, -a/\rho) > 0$, and meanwhile, $\{z_{-a,i}\}_{i \in [m]}$ contains exactly two distinct solutions of (43) and (44) (with $s = -a$), then $\{z_{-a,i}\}_{i \in [m]} = \{\tilde{z}, b - \tilde{z}\}$ with $b = \frac{1-2a}{\rho}$ for some $\tilde{z} > -a/\rho$, and moreover, $\{z_{1-a,i}\}_{i \in [m]}$ is a singleton. We next prove this claim. If $\{z_{-a,i}\}_{i \in [m]}$ contains exactly two distinct solutions of (43) and (44) (with $s = -a$), then $g(-a, z) = 0$ has three distinct solutions, denoted by $-a/\rho < z_1 < z_2 < z_3 \leq (1-a)/\rho$, and the solutions contained in $\{z_{-a,i}\}_{i \in [m]}$ are z_1, z_3 . See Fig. 2b. Moreover, by the facts that G is an antiderivative of g and Φ' is symmetric w.r.t. $1/2$, it holds that $G(-a, z_1) = G(-a, z_3)$ only if $a + \rho z_1 + a + \rho z_3 = 1$ (since only in this case, the two areas enclosed by the curve $z \mapsto \rho \Phi'(a + \rho z)$ and the line $z \mapsto \lambda(2z + a) - \eta$ have the same size). In this case, by Statement 3 of Proposition 1, for $s = 1 - a$, $\{z_{1-a,i}\}_{i \in [m]}$ only contain $(1-a)/\rho$ and/or the largest solution \hat{z}_3 to (43) and (44) with $s = 1 - a$. If $\hat{z}_3 < (1-a)/\rho$, obviously, $G(-a, \hat{z}_3) > G(1-a, (1-a)/\rho)$, and hence, \hat{z}_3 and $(1-a)/\rho$ cannot be solutions to (44) at the same time. Hence, $\{z_{1-a,i}\}_{i \in [m]}$ is a singleton and $\{z_{-a,i}\}_{i \in [m]} = \{\tilde{z}, b - \tilde{z}\}$ for some $\tilde{z} > -a/\rho$, completing the proof of the claim.

We next consider the case $g(-a, -a/\rho) \leq 0$. We claim that for this case, $\{z_{-a,i}\}_{i \in [m]}$ is a singleton. We next prove it. For this case, the equation (43) has at most two distinct solutions in $(-a/\rho, (1-a)/\rho)$. Since adjacent solutions cannot be both contained in $\{z_{-a,i}\}_{i \in [m]}$, only one solution can be contained in $\{z_{-a,i}\}_{i \in [m]}$. In other words, $\{z_{-a,i}\}_{i \in [m]} \subseteq \{-a/\rho, \tilde{z}\}$ for some solution \tilde{z} . Suppose that $\{z_{-a,i}\}_{i \in [m]} = \{-a/\rho, \tilde{z}\}$ with $\tilde{z} > -a/\rho$. For this case, by the fact that G is an antiderivative of g , we have that $G(-a, -a/\rho) = G(-a, \tilde{z})$ only if $g(-a, z) = 0$ has two distinct solutions in $(-a/\rho, (1-a)/\rho)$, denoted by $-a/\rho < z_1 < z_2 \leq (1-a)/\rho$, and the solution contained in $\{z_{-a,i}\}_{i \in [m]}$ is z_2 . Moreover, $G(-a, -a/\rho) = G(-a, z_2)$ if and only if the two areas enclosed by the curve $z \mapsto \rho \Phi'(a + \rho z)$ and the lines $z \mapsto \lambda(2z + a) - \eta$ and $z = -a/\rho$ have the same size. However, since Φ' is symmetric w.r.t. $1/2$, the area at the left side of $z = z_1$ is strictly larger than the one at the right side (note that z_2 is strictly smaller than $(1-a)/\rho$ by Statement 3 of Proposition 1). This leads to a contradiction, and hence, $\{z_{-a,i}\}_{i \in [m]}$ is a singleton and moreover, $\{z_{-a,i}\}_{i \in [m]} = \{\tilde{z}\}$ with $\tilde{z} > -a/\rho$.

By symmetry, similar conclusions can be drawn for $s = 1 - a$. Summarizing these points, optimal solutions to the maximization problem in (9) satisfy that $\{z_{-a,i}\}_{i \in [m]} \subseteq \{\tilde{z}, b - \tilde{z}\}$, $\{z_{1-a,i}\}_{i \in [m]} \subseteq \{\hat{z}, b - \hat{z}\}$, and $\{z_{-a,i}\}_{i \in [m]}$ or $\{z_{1-a,i}\}_{i \in [m]}$

is a singleton, where $b = \frac{1-2a}{\rho}$, $-a/\rho < \tilde{z} \leq \frac{b}{2} \leq \hat{z} < (1-a)/\rho$. Moreover, $\hat{z} \geq \frac{b+1}{2}$ if $\{z_{-a,i}\}_{i \in [m]}$ has size 2, and $\tilde{z} \leq \frac{b-1}{2}$ if $\{z_{1-a,i}\}_{i \in [m]}$ has size 2. We next divide the rest of the proof to three cases.

Case 1: We first consider the case in which both $\{z_{-a,i}\}_{i \in [m]}$ and $\{z_{1-a,i}\}_{i \in [m]}$ are singletons. Solving $\mathbb{E}Z = 0, \mathbb{E}[SZ] = \mathbb{E}[Z^2]$ yields $Z = 0$ or $Z = S$ with probability one. By Statement 3 of Proposition 1, $Z = 0$ is not an optimal solution to the optimization problem in (9). On the other hand, $Z = S$ results in the following value of the program:

$$\mathbb{E}[\Phi(a + \rho S)] = (1-a)\Phi(a + \rho(-a)) + a\Phi(a + \rho(1-a)).$$

Case 2: We next consider the case in which $\{z_{-a,i}\}_{i \in [m]}$ is of size 2 and $\{z_{1-a,i}\}_{i \in [m]}$ is a singleton. In this case, the optimal solution to the optimization problem in (9) is

$$P_{SZ}(s, z) = \begin{cases} 1-a-p & s = -a, z = \tilde{z} \\ p & s = -a, z = b - \tilde{z} \\ a & s = 1-a, z = \hat{z} \end{cases}$$

for some (p, \tilde{z}, \hat{z}) such that $0 < p < 1-a, -a/\rho < \tilde{z} \leq \frac{b}{2} \leq \hat{z} < (1-a)/\rho$, and $\hat{z} \geq \frac{b+1}{2}$. Solving $\mathbb{E}Z = 0, \mathbb{E}[SZ] = \mathbb{E}[Z^2]$ yields the feasible solution satisfying $\hat{z} \geq \frac{b+1}{2}$ which is given by

$$\hat{z} = \frac{1}{2}(\sqrt{\frac{\Delta}{a}} + b + 1), \quad (47)$$

$$p = \frac{-\sqrt{a\Delta} - (a(b+1-2\tilde{z}) + 2\tilde{z})}{2(b-2\tilde{z})} \quad (48)$$

with $\Delta = a(b^2 - 4b\tilde{z} + 2b + 4\tilde{z}^2 + 1) + 4\tilde{z}(b - \tilde{z})$. To ensure $\Delta \geq 0, p \geq 0$, it is required that $\tilde{z} \leq \frac{-a(1+b)}{2(1-a)}$. Hence, the optimal value of the program for this case is $\hat{\Gamma}_\rho(a)$.

Case 3: We lastly consider the case in which $\{z_{1-a,i}\}_{i \in [m]}$ is of size 2 and $\{z_{-a,i}\}_{i \in [m]}$ is a singleton. In this case, the optimal solution to the optimization problem in (9) is

$$P_{SZ}(s, z) = \begin{cases} 1-a & s = -a, z = \tilde{z} \\ p & s = 1-a, z = b - \tilde{z} \\ a-p & s = 1-a, z = \hat{z} \end{cases}$$

for some (p, \tilde{z}, \hat{z}) such that $0 < p < a, -a/\rho < \tilde{z} \leq \frac{b}{2} \leq \hat{z} < (1-a)/\rho$, and $\tilde{z} \leq \frac{b-1}{2}$. By symmetry, substituting $a \leftarrow 1-a, \tilde{z} \leftarrow -\tilde{z}, \hat{z} \leftarrow -\hat{z}$ (which implies $b \leftarrow -b$) into (47) and (48), we obtain the optimal p, \tilde{z} for this case, which results in the value $\hat{\Gamma}_\rho(1-a)$ of the program. Hence, the first part of Statement 2 holds.

For $a = 1/2$, we have $b = 0$ and both $\frac{-a(1+b)}{2(1-a)}$ and c are equal to $-1/2$. Hence, $\tilde{z} = -1/2$ is the unique feasible solution to the supremization in (15), which results in the value $\Phi(\frac{1-\rho}{2})$. Furthermore, for $a = 1/2$, $\mathbb{E}[\Phi(a + \rho S)]$ is the same value. Hence, the second part of Statement 2 follows.

6 Proof of Corollary 1

Excluding the trivial cases, we assume $\rho \in (0, \rho^*]$. By definition, $\Phi = \Phi_\alpha^{\text{sym}}$ with $\alpha \in (1, 2)$ satisfies the assumption in Statement 1 of Theorem 4, which means that Statement 1 of Theorem 4 can be applied to this case. For $a = 1/2$, we have $z_1 + z_2 = 0$, which further implies $p = q$. From $(z_1, z_2) \in \bar{\mathcal{Z}}$, we know $-\frac{1}{2} \leq z_1 \leq -\frac{1}{4}$. Denote the bound in (2) for $\Phi = \Phi_\alpha^{\text{sym}}$ as $\bar{\Gamma}_\rho^{(\alpha)}(a)$ where $\alpha \in [1, 2)$. Then, by the monotonicity of Φ_α^{sym} in α , we have for $\alpha \in (1, 2)$,

$$\bar{\Gamma}_\rho^{(1)}\left(\frac{1}{2}\right) \leq \lim_{\alpha \downarrow 1} \bar{\Gamma}_\rho^{(\alpha)}\left(\frac{1}{2}\right) = \lim_{\alpha \downarrow 1} \max_{-\frac{1}{2} \leq z_1 \leq -\frac{1}{4}} h_\alpha(z_1),$$

where

$$\begin{aligned} h_\alpha(z_1) &:= 2p\Phi_\alpha^{\text{sym}}\left(\frac{1}{2} + \rho z_1\right) \\ &= \frac{1 - \rho}{(1 + 2\rho z_1)(1 - \rho - 2\rho z_1)} \Phi_\alpha^{\text{sym}}\left(\frac{1}{2} + \rho z_1\right). \end{aligned}$$

We now swap the limit and maximization. Denote $\{\alpha_k\}$ as a decreasing sequence with limit 1. Denote $z_{1,k}$ as an optimal solution to the optimization $\max_{-\frac{1}{2} \leq z_1 \leq -\frac{1}{4}} h_\alpha(z_1)$ with $\alpha = \alpha_k$. By passing to a subsequence, we assume that $\{z_{1,k}\}$ converges to some z_1^* . Then, by the continuity of $h_\alpha(z_1)$ in (α, z_1) , $\bar{\Gamma}_\rho^{(\alpha_k)}\left(\frac{1}{2}\right) = h_{\alpha_k}(z_{1,k}) \rightarrow h_1(z_1^*) \leq \max_{-\frac{1}{2} \leq z_1 \leq -\frac{1}{4}} h_1(z_1)$, which implies $\bar{\Gamma}_\rho^{(1)}\left(\frac{1}{2}\right) \leq \max_{-\frac{1}{2} \leq z_1 \leq -\frac{1}{4}} h(z_1)$, with $h(z_1) := h_1(z_1)$ for brevity. More explicitly,

$$h(z_1) = \frac{1 - \rho}{(1 + 2\rho z_1)(1 - \rho - 2\rho z_1)} \left(\left(\frac{1}{2} + \rho z_1\right) \log\left(\frac{1}{2} + \rho z_1\right) + \left(\frac{1}{2} - \rho z_1\right) \log\left(\frac{1}{2} - \rho z_1\right) \right).$$

To compute the maximum of $h(z_1)$ over $-\frac{1}{2} \leq z_1 \leq -\frac{1}{4}$, we take its derivative which is a standard technique. That is,

$$h'(z_1) = \frac{-\rho(1 - \rho)\varphi_\rho(z_1)}{(1 + 2\rho z_1)^2(1 - \rho - 2\rho z_1)^2},$$

where

$$\varphi_\rho(z_1) := ((1 - 2\rho z_1)^2 - 2\rho) \log\left(\frac{1}{2} - \rho z_1\right) - (1 + 2\rho z_1)^2 \log\left(\frac{1}{2} + \rho z_1\right).$$

We claim that h is non-increasing given $\rho \in (0, \rho^*]$. We next prove it. It is easy to verify that

$$\varphi_\rho''(z_1) = \frac{8\rho^2(\rho + (1 - 2\rho z_1)^2 \log(\frac{1 - 2\rho z_1}{1 + 2\rho z_1}))}{(1 - 2\rho z_1)^2} \geq 0.$$

Hence, φ' is increasing. Furthermore, $\varphi'_\rho(-\frac{1}{2}) = \frac{4\rho g(\rho)}{\rho+1}$, where

$$g(\rho) := -(1 - \rho^2) \log\left(\frac{1 - \rho}{2}\right) - (1 + \rho)^2 \log\left(\frac{1 + \rho}{2}\right) - 1.$$

It is easy to verify that $g''(\rho) \leq 0$, and $g(0), g(\frac{1}{2}) \geq 0$. Hence, $g(\rho) \geq 0$ (i.e., $\varphi'_\rho(-\frac{1}{2}) \geq 0$) for $\rho \in (0, \frac{1}{2}]$. Combining this with the fact that φ'_ρ is increasing gives $\varphi'_\rho(z_1) \geq 0$ for $z_1 \in [-\frac{1}{2}, 0]$ and $\rho \in (0, \frac{1}{2}]$. Hence, for $\rho \in (0, \frac{1}{2}]$, φ_ρ is increasing on $[-\frac{1}{2}, 0]$.

Recall $\psi(\rho)$ defined in (16), and observe that $\varphi_\rho(-\frac{1}{2}) = \psi(\rho)$. We now prove $\psi(\rho) \geq 0$ for $\rho \in [0, \rho^*]$. It is easy to verify $\psi'''(\rho) \geq 0$, and hence $\psi'(\rho)$ is convex. Also, $\psi'(0) \geq 0, \psi'(\frac{1}{2}) \leq 0$. Hence, ψ' is first-positive-then-negative⁹ on $[0, \frac{1}{2}]$, which means that ψ is first-increasing-then-decreasing on $[0, \frac{1}{2}]$. Observe that $\psi(0), \psi(\rho^*) = 0$ and $\rho^* \in [0, \frac{1}{2}]$. Hence $\psi(\rho) \geq 0$ for $\rho \in (0, \rho^*]$, i.e., $\varphi_\rho(-\frac{1}{2}) \geq 0$.

Combining the facts that φ_ρ is increasing and $\varphi_\rho(-\frac{1}{2}) \geq 0$ for $\rho \in (0, \rho^*]$ yields that $\varphi_\rho(z_1) \geq 0$ for $z_1 \in [-\frac{1}{2}, 0]$ and $\rho \in (0, \rho^*]$. It means that $h'(z_1) \leq 0$ and hence h is non-increasing given $\rho \in (0, \rho^*]$, completing the proof of the claim above.

Therefore, by the claim above,

$$\bar{\Gamma}_\rho^{(1)}\left(\frac{1}{2}\right) \leq h\left(-\frac{1}{2}\right) = \left(\frac{1 - \rho}{2}\right) \log\left(\frac{1 - \rho}{2}\right) + \left(\frac{1 + \rho}{2}\right) \log\left(\frac{1 + \rho}{2}\right).$$

Observe that the most RHS above is attained by dictator functions, which completes the proof.

7 Proof of Corollary 2

Excluding the trivial cases, we assume $\rho \in (0, 1)$. We next prove Corollary 2. The case $\alpha \in (1, 2)$ is implied immediately by Lemma 1 and Corollary 1. For $\alpha \in [2, 5]$, by Lemma 1, it suffices to only prove the case $\alpha = 5$.

For $a = 1/2$ and $\Phi(t) = t^\alpha + (1 - t)^\alpha$ with $\alpha = 5$, similarly as in Section 16, by applying Statement 1 of Theorem 4, we have $z_1 + z_2 = 0$, and hence $p = q$, $-\frac{1}{2} \leq z_1 \leq -\frac{1}{4}$, and

$$\bar{\Gamma}_\rho\left(\frac{1}{2}\right) = \sup_{-\frac{1}{2} \leq z_1 \leq -\frac{1}{4}} 2\left[\frac{1}{2} - p + p\Phi\left(\frac{1}{2} + \rho z_1\right)\right]$$

with $p = \frac{1 - \rho}{2(1 - \rho - 2\rho z_1)(1 + 2\rho z_1)}$. Denoting $z = -2\rho z_1 \in [\frac{\rho}{2}, \rho]$, we have $\bar{\Gamma}_\rho\left(\frac{1}{2}\right) = \sup_{\frac{\rho}{2} \leq z \leq \rho} h(z)$, where

$$h(z) := \frac{-\rho + 5(\rho - 1)z^3 + 5(\rho - 1)z^2 + 15\rho z + z + 1}{16(-\rho + z + 1)}.$$

⁹We say a function f is *first-positive-then-negative* if there exists some real number a such that $f(x) \geq 0$ for $x < a$ and $f(x) \leq 0$ for $x > a$. We can define a function to be *first-negative-then-positive*, *first-increasing-then-decreasing*, and *first-decreasing-then-increasing* in a similar way.

By taking derivatives and using the standard arguments, it can be observed that $h(z)$ is non-decreasing. Hence $\bar{\Gamma}_\rho(a) = h(\rho) = \Phi(\frac{1-\rho}{2})$, completing the proof. We next provide the details on how to take derivatives to show that $h(z)$ is non-decreasing.

Taking derivative for h , we have $h'(z) = \frac{5\varphi(z)}{16(-\rho+z+1)^2}$, where

$$\varphi(z) = -3(\rho-1)\rho + 2(\rho-1)z^3 + (-3\rho^2 + 7\rho - 4)z^2 - 2(\rho-1)^2z.$$

Taking derivative again, we further have $\varphi'(z) = -2(1-\rho)(3z+1)(1-\rho+z) \leq 0$. Hence, $\varphi(z)$ is non-increasing, which implies that $\varphi(z) \geq \varphi(\rho) = (1-\rho)^3\rho \geq 0$, i.e., $h'(z) \geq 0$. Therefore, $h(z)$ is non-decreasing.

8 Proof of Corollary 3

Excluding the trivial cases, we assume $\rho \in (0, 1)$.

8.1 Case of $\alpha \in (1, 2)$

Here, we denote $\Phi(t) = t^\alpha$ with $\alpha \in (1, 2)$. Observe that Φ' is concave. By Theorem 5, for $a = 1/2$,

$$\tilde{\Gamma}_\rho\left(\frac{1}{2}\right) = \sup_{0 \leq z_2 \leq \frac{1}{2}} p\left(\frac{1}{2} + \rho z_1\right)^\alpha + \frac{1}{2}\left(\frac{1}{2} + \rho z_2\right)^\alpha.$$

where p, z_2 are defined in (17) and (18). Now we consider p as a free variable, and solving (17) and (18) w.r.t. z_1, z_2 , we obtain two solutions:

$$\begin{cases} z_1^{(1)} = \frac{p(1-2p-\rho)-T(p)}{2(1+2p)p\rho}, \\ z_2^{(1)} = \frac{\frac{1}{2}-p(1-\rho)+T(p)}{(1+2p)\rho}, \end{cases}$$

and

$$\begin{cases} z_1^{(2)} = \frac{p(1-2p-\rho)+T(p)}{2(1+2p)p\rho}, \\ z_2^{(2)} = \frac{\frac{1}{2}-p(1-\rho)-T(p)}{(1+2p)\rho}, \end{cases}$$

where

$$T(p) = \sqrt{p(p(2-2\rho+\rho^2)+\rho-1)}.$$

To ensure the term in the square root above is nonnegative, it is required that

$$\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}.$$

Observe that given p , $p z_1 + \frac{1}{2} z_2$ remains the same for $(z_1^{(1)}, z_2^{(1)})$ and $(z_1^{(2)}, z_2^{(2)})$, and moreover, $z_1^{(1)} \leq z_1^{(2)} \leq z_2^{(2)} \leq z_2^{(1)}$. By the convexity of Φ , we have that $(z_1^{(1)}, z_2^{(1)})$

leads to a larger value of $p(\frac{1}{2} + \rho z_1)^\alpha + \frac{1}{2}(\frac{1}{2} + \rho z_2)^\alpha$ than $(z_1^{(2)}, z_2^{(2)})$. Hence,

$$\tilde{\Gamma}_\rho(\frac{1}{2}) \leq \sup_{\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}} h_\alpha(p)$$

where

$$\begin{aligned} h_\alpha(p) &:= p(\frac{1}{2} + \rho z_1^{(1)})^\alpha + \frac{1}{2}(\frac{1}{2} + \rho z_2^{(1)})^\alpha \\ &= p\left(\frac{(2-\rho)p - T(p)}{2(1+2p)p}\right)^\alpha + \frac{1}{2}\left(\frac{1+p\rho + T(p)}{1+2p}\right)^\alpha. \end{aligned}$$

Define

$$\varphi(\alpha, p) := -A(p)D(p)^{\alpha-1} - \frac{B(p)}{\alpha} + C(p)$$

where

$$\begin{aligned} A(p) &:= p[(-1+6p)(1-\rho) + 2p\rho^2 - 2(2-\rho)T(p)] \\ B(p) &:= [(2-\rho)p - T(p)]2(1+2p)T(p) \\ C(p) &:= p[(3-2p-8p^2)(1-\rho) - 4\rho^2p^2 + 4p(2-\rho)T(p)] \\ D(p) &:= \frac{(p\rho + 1 + T(p))2p}{(2-\rho)p - T(p)}. \end{aligned}$$

It is easy to verify that $A(p), B(p), D(p) \geq 0$ for $\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}$. Then,

$$\frac{d}{dp} h_\alpha(p) = \left(\frac{(2-\rho)p - T(p)}{2(1+2p)p}\right)^{\alpha-1} \frac{-\alpha\varphi(\alpha, p)}{4(1+2p)^2pT(p)}. \quad (49)$$

Recall that the function θ is defined in (20). Obviously, it holds that $\theta^{-1}(\frac{1-\rho}{1+\rho}) \geq 1$ for $0 \leq \rho \leq 1$.

Lemma 3. For $\theta^{-1}(\frac{1-\rho}{1+\rho}) \leq \alpha < 2$ and $\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}$, we have $\varphi(\alpha, p) \leq 0$.

By this lemma and (49), for $\theta^{-1}(\frac{1-\rho}{1+\rho}) \leq \alpha < 2$, h_α is non-decreasing for $\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}$, which further implies that $\tilde{\Gamma}_\rho(\frac{1}{2}) \leq h_\alpha(\frac{1}{2})$. Observe that $h_\alpha(\frac{1}{2}) = \frac{1}{2}\Phi(\frac{1-\rho}{2}) + \frac{1}{2}\Phi(\frac{1+\rho}{2})$ is the Φ -stability of dictator functions. Hence, $\tilde{\Gamma}_\rho(\frac{1}{2})$ is attained by dictator functions for $\theta^{-1}(\frac{1-\rho}{1+\rho}) \leq \alpha < 2$ (or equivalently, for $\rho \leq \frac{1-\theta(\alpha)}{1+\theta(\alpha)}$), which completes the proof of Corollary 3 for the case $\alpha \in (1, 2)$. Hence, the rest is to prove the lemma above.

Proof of Lemma 3. Since $A(p), B(p), D(p) \geq 0$, by definition, given p , $\varphi(\alpha, p)$ is concave in α . It is easy to verify that $\varphi(1, p) = 0$. Hence, given p , one of the following two statements is true: 1) $\varphi(\alpha, p) \leq 0$ for any $\alpha > 0$; 2) the equation $\varphi(\alpha, p) = 0$ with

unknown α has exactly two distinct solutions, 1 and $\alpha^*(p) > 0$. If the first statement is true, then Lemma 3 holds for such p . Hence, it suffices to consider the case in which the second statement is true.

If p satisfies that $\alpha^*(p) < 1$, by the concavity of $\varphi(\alpha, p)$ in α , we have that $\varphi(\alpha, p) \leq 0$ for any $\alpha \geq 1$. Hence, Lemma 3 holds for such p .

We next consider the case $\alpha^*(p) > 1$. Similarly, by the concavity of $\varphi(\alpha, p)$ in α , we have that $\varphi(\alpha, p) \leq 0$ for any $\alpha \geq \alpha^*(p)$. Hence, Lemma 3 holds for $\alpha \geq \sup_{\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}} \alpha^*(p)$. To determine the supremization here, we need the following technical lemma. Its proof is deferred to Section 8.1.1.

Lemma 4. *For any $\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}$ such that $1 < \alpha^*(p) < 2$, we have $\frac{d}{dp} \alpha^*(p) \geq 0$.*

By the lemma above, $\sup_{\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}} \alpha^*(p) = \alpha^*(\frac{1}{2})$. It is easy to check that $\alpha^*(\frac{1}{2}) = \theta^{-1}(\frac{1-\rho}{1+\rho})$. Hence Lemma 3 holds for $\alpha \geq \theta^{-1}(\frac{1-\rho}{1+\rho})$.

Combining all the cases above completes the proof of Lemma 3. \square

8.1.1 Proof of Lemma 4

Denote $\varphi_1(\alpha, p) := \frac{\partial}{\partial \alpha} \varphi(\alpha, p)$ and $\varphi_2(\alpha, p) := \frac{\partial}{\partial p} \varphi(\alpha, p)$. By the assumption $\alpha^*(p) > 1$, the definition of $\alpha^*(p)$, and the concavity of $\varphi(\alpha, p)$ in α , we have $\varphi_1(\alpha^*(p), p) < 0$. By the implicit function theorem,

$$\frac{d}{dp} \alpha^*(p) = \frac{-\varphi_2(\alpha^*(p), p)}{\varphi_1(\alpha^*(p), p)}.$$

Hence, it suffices to show $\varphi_2(\alpha^*(p), p) \geq 0$.

Observe that

$$\varphi_2(\alpha, p) = -A'(p)D(p)^{\alpha-1} - (\alpha-1)A(p)D(p)^{\alpha-2}D'(p) - \frac{B'(p)}{\alpha} + C'(p) \quad (50)$$

Since $\alpha^* = \alpha^*(p)$ is a solution, i.e., $\varphi(\alpha^*, p) = 0$, we obtain $A(p)D(p)^{\alpha^*-1} = C(p) - \frac{B(p)}{\alpha^*}$. Substituting it into (50) with α replaced by α^* , we obtain

$$\begin{aligned} \varphi_2(\alpha^*, p) &= -A'(p)D(p)^{\alpha^*-1} - (\alpha^*-1)A(p)D(p)^{\alpha^*-2}D'(p) - \frac{B'(p)}{\alpha^*} + C'(p) \\ &= -A(p)D(p)^{\alpha^*-1} \left(\frac{A'(p)}{A(p)} + (\alpha^*-1) \frac{D'(p)}{D(p)} \right) - \frac{B'(p)}{\alpha^*} + C'(p) \\ &= -\left(C(p) - \frac{B(p)}{\alpha^*} \right) \left(\frac{A'(p)}{A(p)} + (\alpha^*-1) \frac{D'(p)}{D(p)} \right) - \frac{B'(p)}{\alpha^*} + C'(p) \\ &= \phi(\alpha^*, p), \end{aligned} \quad (51)$$

where

$$\phi(\alpha, p) := -C(p) \left(\frac{A'(p)}{A(p)} + (\alpha-1) \frac{D'(p)}{D(p)} \right) + \frac{B(p)}{\alpha} \left(\frac{A'(p)}{A(p)} - \frac{D'(p)}{D(p)} - \frac{B'(p)}{B(p)} \right)$$

$$+ B(p) \frac{D'(p)}{D(p)} + C'(p).$$

By standard arguments, it can be shown that

$$\frac{A'(p)}{A(p)} - \frac{D'(p)}{D(p)} - \frac{B'(p)}{B(p)} \leq 0; \quad (52)$$

see details below. Hence, given $\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}$, $\phi(\alpha, p)$ is concave in $\alpha > 0$.

On the other hand, using Mathematica, it is easy to show that $\phi(1, p) = 0$, $\phi(2, p) \geq 0$ for $\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}$. Hence, $\phi(\alpha, p) \geq 0$ for $1 < \alpha < 2$ and $\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}$, which, combined with (51), implies $\varphi_2(\alpha^*(p), p) \geq 0$ since by assumption, $1 < \alpha^*(p) < 2$. This completes the proof of Lemma 4.

For completeness, we next provide proof details of (52). Observe that

$$\frac{A'(p)}{A(p)} - \frac{D'(p)}{D(p)} - \frac{B'(p)}{B(p)} = \frac{p(F(p) + E(p)T(p))}{2(2p+1)T(p)^2(\rho p + 1 + T(p))A(p)},$$

where

$$\begin{aligned} E(p) := & -16(2\rho^4 - 9\rho^3 + 17\rho^2 - 16\rho + 6)p^3 \\ & - 8(\rho^4 + \rho^3 - 10\rho^2 + 14\rho - 7)p^2 - 8(\rho^3 - \rho^2 - 2\rho + 2)p - 2(\rho - 1)^2, \end{aligned}$$

and

$$\begin{aligned} F(p) := & -16(2\rho^5 - 11\rho^4 + 27\rho^3 - 36\rho^2 + 26\rho - 8)p^4 \\ & - 4(2\rho^5 + 4\rho^4 - 39\rho^3 + 86\rho^2 - 83\rho + 32)p^3 \\ & - 4(3\rho^4 - 4\rho^3 - 5\rho^2 + 13\rho - 7)p^2 - (\rho - 1)^2(5\rho + 4)p - (\rho - 1)^2. \end{aligned}$$

Using Mathematica¹⁰, one can show that $F(p) \leq 0$ and $F(p)^2 \geq E(p)^2T(p)^2$ for $\frac{1-\rho}{2-2\rho+\rho^2} \leq p \leq \frac{1}{2}$ (and $0 < \rho < 1$). Hence, $F(p) + E(p)T(p) \leq 0$, which in turn implies (52).

8.2 Case of $\alpha \in [2, 3]$

For $\alpha \in [2, 3]$, by Lemma 1, to prove Corollary 3, it suffices to prove the case $\alpha = 3$. Here we let $\Phi(t) = t^3$. Observe that Φ' is convex. By Remark 1, we have $\tilde{\Gamma}_\rho(a) = \sup_{0 \leq z_2 \leq a} g(z_2)$, where

$$g(z_2) := a - p + p(a - \rho z_1)^3 + (1 - a)(a - \rho z_2)^3$$

with

$$z_1 = \frac{z_2(\rho(1 - z_2) - (1 - a))}{a - \rho z_2},$$

¹⁰The code can be found in the link <https://www.dropbox.com/s/kll3157wfuwaw7c/code.nb?dl=0>

$$p = \frac{(1-a)(a-\rho z_2)^2}{1-a+\rho^2 z_2 - (1-a+\rho z_2)^2}.$$

For $a = 1/2$, by defining $z = 2\rho z_2 \in [0, \rho]$, we have $\tilde{\Gamma}_\rho(a) = \sup_{0 \leq z \leq \rho} h(z)$, where

$$h(z) := g\left(\frac{z}{2\rho}\right) = \frac{1 - (1-4\rho)z - (2\rho^2 + 2\rho + 1)z^2 - (1-2\rho)z^3}{8(1-z)}.$$

By standard arguments, it can be shown that $h'(z) \geq 0$; see details below. Hence, $h(z)$ is increasing, which implies $\tilde{\Gamma}_\rho(a) = h(\rho) = \frac{1}{2}\Phi\left(\frac{1+\rho}{2}\right) + \frac{1}{2}\Phi\left(\frac{1-\rho}{2}\right)$. Note that the most RHS is attained by dictator functions, completing the proof.

For completeness, we next provide proof details of $h'(z) \geq 0$. Taking derivative, we have $h'(z) = \frac{\varphi(z)}{4(z-1)^2}$, where

$$\varphi(z) := (1-2\rho)z^3 + (\rho^2 + 4\rho - 1)z^2 - (2\rho^2 + 2\rho + 1)z + 2\rho.$$

Taking derivative again, we have

$$\varphi'(z) = (3-6\rho)z^2 + 2(\rho^2 + 4\rho - 1)z - 2\rho^2 - 2\rho - 1.$$

For $\rho \leq 1/2$, $\varphi'(z)$ is convex in z . Hence for this case, $\varphi'(z) \leq \max\{\varphi'(0), \varphi'(\rho)\} \leq 0$. For $\rho > 1/2$, $\varphi'(z)$ is increasing in z for $z \leq \rho \leq \frac{\rho^2 + 4\rho - 1}{6\rho - 3}$. Hence for this case, $\varphi'(z) \leq \varphi'(\rho) \leq 0$. Combining these two cases, $\varphi'(z) \leq 0$ and hence $\varphi(z)$ is decreasing for $\rho \in [0, 1]$. Hence $\varphi(z) \geq \varphi(\rho) = (1-\rho)^3 \rho \geq 0$, which implies $h'(z) \geq 0$.

9 Proof of Lemma 2

Let \mathcal{B} be the support of f . Hence $|\mathcal{B}| = a2^n$. Denote $\mathcal{A}_1 := \{\mathbf{x} : x_1 = 1\}$ and $\mathcal{A}_1^c := \{-1, 1\}^n \setminus \mathcal{A}_1 = \{\mathbf{x} : x_1 = -1\}$. Then by definition, $|\mathcal{B} \cap \mathcal{A}_1| = \frac{a+\beta}{2}2^n$. Let $\mathcal{A} \subseteq \mathcal{A}_1$ be an arbitrary subset such that $\mathcal{B} \cap \mathcal{A}_1 \subseteq \mathcal{A}$ and $|\mathcal{A}| = a2^n$. Denote $g := 1_{\mathcal{A}}$, $h := 1_{\{\mathbf{x} \in \{-1, 1\}^{n-1} : (1, \mathbf{x}) \in \mathcal{A}, (1, \mathbf{x}) \notin \mathcal{B}\}}$, and $l := 1_{\{\mathbf{x} \in \{-1, 1\}^{n-1} : (-1, \mathbf{x}) \in \mathcal{B}\}}$. Then $\mathbb{E}h = \mathbb{E}l = \frac{a-\beta}{2}2^n$. Moreover, $f = g - h \cdot 1_{\{x_1 = 1\}} + l \cdot 1_{\{x_1 = -1\}}$, and hence, $\hat{f}_{\mathcal{S}} = \hat{g}_{\mathcal{S}} - \frac{1}{2}\hat{h}_{\mathcal{S} \setminus \{1\}} + \frac{1}{2}\hat{l}_{\mathcal{S} \setminus \{1\}}, \forall \mathcal{S} \subseteq [n]$. Note that h and l are Boolean functions on the $(n-1)$ -dimensional space. Hence, their Fourier coefficients $\hat{h}_{\mathcal{S}}, \hat{l}_{\mathcal{S}}, \mathcal{S} \subseteq [2:n]$ are also defined on the $(n-1)$ -dimensional space. By the Minkowski inequality,

$$\begin{aligned} \mathbf{W}_1[f] - \beta^2 &= \sum_{i=2}^n \hat{f}_{\{i\}}^2 = \sum_{i=2}^n \left(\hat{g}_{\{i\}} - \frac{1}{2}\hat{h}_{\{i\}} + \frac{1}{2}\hat{l}_{\{i\}} \right)^2 \\ &\leq \left(\sqrt{\sum_{i=2}^n \hat{g}_{\{i\}}^2} + \sqrt{\frac{1}{4}\sum_{i=2}^n \hat{h}_{\{i\}}^2} + \sqrt{\frac{1}{4}\sum_{i=2}^n \hat{l}_{\{i\}}^2} \right)^2 \\ &= \left(\sqrt{\mathbf{W}_1[g] - a^2} + \frac{1}{2}\sqrt{\mathbf{W}_1[h]} + \frac{1}{2}\sqrt{\mathbf{W}_1[l]} \right)^2 \end{aligned} \quad (53)$$

$$\leq \left(\sqrt{W^{(n)}(a) - a^2} + \sqrt{W^{(n-1)}(a - \beta)} \right)^2$$

where (53) follows by the facts that $\hat{g}_{\{1\}} = a$ and $\mathbf{W}_1[h] = \sum_{i=2}^n \hat{h}_{\{i\}}^2$, $\mathbf{W}_1[l] = \sum_{i=2}^n \hat{l}_{\{i\}}^2$ since the Fourier coefficients of h, l are defined on the $(n-1)$ -dimensional space. Therefore, (21) holds.

By Parseval's theorem, $W^{(n)}(a) \leq a - a^2$. Substituting this into (21) and specializing (21) to the case $a = \frac{1}{2}$, we obtain (22). On the other hand, this upper bound is attained by the Boolean function

$$f(\mathbf{x}) = 1\{\mathbf{x} : x_1 = 1, (x_2, x_3, \dots, x_n) \in \mathcal{A}^c\} + 1\{-\mathbf{x} : x_1 = 1, (x_2, x_3, \dots, x_n) \in \mathcal{A}\}$$

for some $\mathcal{A} \subseteq \{-1, 1\}^{n-1}$ such that $1_{\mathcal{A}}$ attains $W^{(n-1)}(\frac{1}{2} - k)$. This can be seen from the fact that

$$f(\mathbf{x}) = \frac{1+x_1}{2} - \sum_{S \subseteq [2:n]: |S| \text{ odd}} \hat{g}_S \chi_S(x_2, \dots, x_n) - x_1 \sum_{S \subseteq [2:n]: |S| \text{ even}} \hat{g}_S \chi_S(x_2, \dots, x_n)$$

where $\{\hat{g}_S\}$ are Fourier coefficients of $g(x_2, \dots, x_n) = 1_{\mathcal{A}}(x_2, \dots, x_n)$.

10 Proof of Proposition 4

The proofs of Statements 1-3 are exactly the same as those of Statements 1-3 of Proposition 1. Here we only provide the proof idea for Statement 4, and omit the details since it is similar to that of Statement 4 of Proposition 1. For even $m \geq 8$, denote $(z_{s,x,i}, p_{s,x,i})_{(s,x,i) \in \mathcal{S} \times \mathcal{X} \times [m]}$ as an optimal solution to the maximization in (9) such that $z_{-s,-x,i} = -z_{s,x,i}$, $p_{-s,-x,i} = p_{s,x,i}$ for all (s, x, i) . Such an optimal solution always exists since for any optimal solution $(\hat{z}_{s,x,i}, \hat{p}_{s,x,i})_{(s,x,i) \in \mathcal{S} \times \mathcal{X} \times [m/2]}$ to the optimization problem in (9) with $m \leftarrow m/2$, we can construct a desired distribution $P_{SXZ} = \sum_{s,x,i} \frac{1}{2} \hat{p}_{s,x,i} (\delta_{s,x,\hat{z}_{s,x,i}} + \delta_{-s,-x,-\hat{z}_{s,x,i}})$, which is optimal for the optimization in (35) and meanwhile satisfies $P_{SXZ}(-s, -x, -z) = P_{SXZ}(s, x, z)$ for all (s, x, z) .

Suppose that LICQ is not satisfied by the solution $(z_{s,x,i}, p_{s,x,i})_{(s,x,i) \in \mathcal{S} \times \mathcal{X} \times [m]}$. Denote

$$\mathcal{I} := \{(s, x, i) : a + \rho z_{s,x,i} = 0\} \text{ and } \mathcal{J} := \{(s, x, i) : a + \rho z_{s,x,i} = 1\}.$$

By Statement 3, we can write $\mathcal{I} = \mathcal{I}^- \cup \mathcal{I}^+$ and $\mathcal{J} = \mathcal{J}^- \cup \mathcal{J}^+$ for some $\mathcal{I}^- \subseteq \{(-a, -1)\} \times [m]$, $\mathcal{I}^+ \subseteq \{(-a, 1)\} \times [m]$, $\mathcal{J}^- \subseteq \{(1-a, -1)\} \times [m]$, and $\mathcal{J}^+ \subseteq \{(1-a, 1)\} \times [m]$. The gradients of the active inequality constraints (including (36) by Statement 2) and the gradients of the equality constraints constitute the $(|\mathcal{I}| + |\mathcal{J}| +$

$5) \times (4m)$ matrix

$$G := \begin{bmatrix} \rho \mathbf{I}_{\mathcal{I}} & \mathbf{0}_{1 \times 2m} \\ -\rho \mathbf{I}_{\mathcal{J}} & \mathbf{0}_{1 \times 2m} \\ \mathbf{0}_{1 \times 2m} & (\mathbf{I}_{1 \times m}, \mathbf{0}_{1 \times m}) \\ \mathbf{0}_{1 \times 2m} & (\mathbf{0}_{1 \times m}, \mathbf{I}_{1 \times m}) \\ (p_{s,x,i})_{s,x,i} & (z_{s,x,i})_{s,x,i} \\ (xp_{s,x,i})_{s,x,i} & (xz_{s,x,i})_{s,x,i} \\ (p_{s,x,i}(2z_{s,x,i} - \rho s))_{s,x,i} & (z_{s,x,i}^2 - \rho s z_{s,x,i})_{s,x,i} \end{bmatrix}.$$

The assumption that LICQ is not satisfied implies that G is not of full rank, or equivalently,

$$\hat{G} := \begin{bmatrix} (p_{s,-1,i})_{(s,i) \in (\mathcal{I}^- \cup \mathcal{J}^-)^c} & \mathbf{0}_{1 \times (2m - |\mathcal{I}^+| - |\mathcal{J}^+|)} & A^- & \mathbf{0}_{1 \times 2(m-1)} \\ \mathbf{0}_{1 \times (2m - |\mathcal{I}^-| - |\mathcal{J}^-|)} & (p_{s,1,i})_{(s,i) \in (\mathcal{I}^+ \cup \mathcal{J}^+)^c} & \mathbf{0}_{1 \times 2(m-1)} & A^+ \\ (p_{s,-1,i}(2z_{s,-1,i} - \rho s))_{(s,i) \in (\mathcal{I}^- \cup \mathcal{J}^-)^c} & (p_{s,1,i}(2z_{s,1,i} - \rho s))_{(s,i) \in (\mathcal{I}^+ \cup \mathcal{J}^+)^c} & B^- & B^+ \end{bmatrix}$$

is not of full rank where

$$\begin{aligned} A^- &:= (z_{s,-1,i} - z_{s,-1,1})_{(s,i) \in \mathcal{S} \times [2:m]}, \\ A^+ &:= (z_{s,1,i} - z_{s,1,1})_{(s,i) \in \mathcal{S} \times [2:m]}, \\ B^- &:= ((z_{s,-1,i} - z_{s,-1,1})(z_{s,-1,i} + z_{s,-1,1} - \rho s))_{(s,i) \in \mathcal{S} \times [2:m]}, \\ B^+ &:= ((z_{s,1,i} - z_{s,1,1})(z_{s,1,i} + z_{s,1,1} - \rho s))_{(s,i) \in \mathcal{S} \times [2:m]}. \end{aligned}$$

We next prove that under this condition, there is an optimal solution of the form in (37).

We first prove that for each (s, x) , $z_{s,x,i}$'s are identical for all $i \in [m]$. Since the submatrix

$$\begin{bmatrix} (p_{-\frac{1}{2},-1,i})_{(-\frac{1}{2},i) \in (\mathcal{I}^- \cup \mathcal{J}^-)^c} & (z_{-\frac{1}{2},-1,i} - z_{-\frac{1}{2},-1,1})_{i \in [2:m]} \\ (p_{-\frac{1}{2},-1,i}(2z_{-\frac{1}{2},-1,i} + \rho \frac{1}{2}))_{(-\frac{1}{2},i) \in (\mathcal{I}^- \cup \mathcal{J}^-)^c} & ((z_{-\frac{1}{2},-1,i} - z_{-\frac{1}{2},-1,1})(z_{-\frac{1}{2},-1,i} + z_{-\frac{1}{2},-1,1} + \rho \frac{1}{2}))_{i \in [2:m]} \end{bmatrix}$$

of \hat{G} is not of full rank (due to that $p_{s,x,i} > 0$ for all s, x, i), we know that $z_{-\frac{1}{2},-1,i}$'s are identical for all $i \in [m]$. Similarly, the same conclusion holds for other pairs (s, x) .

Therefore, Z is a function of (S, X) , or equivalently, can be expressed in the form of (37).

11 Proof of Theorem 7

The Lagrangian of the optimization problem in (35) is

$$\mathcal{L} := \sum_{s,x,i} p_{s,x,i} \left\{ \Phi\left(\frac{1}{2} + \rho z_{s,x,i}\right) + \theta_{s,x,i} \left(\frac{1}{2} + \rho z_{s,x,i}\right) + \theta'_{s,x,i} \left(1 - \left(\frac{1}{2} + \rho z_{s,x,i}\right)\right) \right\}$$

$$\begin{aligned}
& + \lambda(\rho s z_{s,x,i} - z_{s,x,i}^2 + (1-\rho)\omega(\beta)) + \eta z_{s,x,i} + \eta'(x z_{s,x,i} - \beta) \} \\
& + \sum_{s,x} \mu_{s,x} (\sum_i p_{s,x,i} - P_{SX}(s,x)).
\end{aligned}$$

Note that the distribution in (37) (by choosing proper z_1, z_2 so that it is feasible to the program in (35)) is also feasible in the program (38) (specifically, which corresponds to $p = \frac{1}{4} + \frac{\beta}{2}, q = \frac{1}{4} - \frac{\beta}{2}$). Hence the objective value induced by this solution is no more than \bar{Y}_ρ . From this observation and Statement 4 of Proposition 4, it suffices to consider the case in which the LICQ is satisfied for some symmetric optimal solution such that $P_{SXZ}(-s, -x, -z) = P_{SXZ}(s, x, z)$ for all (s, x, z) . We next prove the desired result for this case.

By the KKT theorem, we have the following first-order necessary conditions for (local) optimal solutions: For even $m \geq 8$,

$$\begin{aligned}
\frac{\partial \mathcal{L}}{\partial z_{s,x,i}} &= p_{s,x,i} [\rho(\Phi'(\frac{1}{2} + \rho z_{s,x,i}) + \theta_{s,x,i} - \theta'_{s,x,i}) + \lambda(s - 2z_{s,x,i}) + \eta + \eta'x] = 0 \\
\frac{\partial \mathcal{L}}{\partial p_{s,x,i}} &= \Phi(\frac{1}{2} + \rho z_{s,x,i}) + \lambda(\rho s z_{s,x,i} - z_{s,x,i}^2 + (1-\rho)\omega(\beta)) \\
&\quad + \eta z_{s,x,i} + \eta'(x z_{s,x,i} - \beta) + \mu_{s,x} = 0 \\
0 &\leq \frac{1}{2} + \rho z_{s,x,i} \leq 1, \forall (s, x, i) \in \mathcal{S} \times \mathcal{X} \times [m] \\
\sum_{s,x,i} p_{s,x,i} z_{s,x,i} &= 0, \\
\sum_{s,x,i} p_{s,x,i} x z_{s,x,i} &= \beta, \\
\sum_{s,x,i} p_{s,x,i} (z_{s,x,i}^2 - \rho s z_{s,x,i}) &= (1-\rho)\omega(\beta). \\
\sum_i p_{s,x,i} &= P_{SX}(s,x), \forall (s,x) \in \mathcal{S} \times \mathcal{X}, \\
p_{s,x,i} &> 0, \forall (s, x, i) \in \mathcal{S} \times \mathcal{X} \times [m] \\
\theta_{s,x,i} (\frac{1}{2} + \rho z_{s,x,i}) &= 0 \\
\theta'_{s,x,i} (\frac{1}{2} - \rho z_{s,x,i}) &= 0 \\
\lambda \geq 0, \vec{\theta}, \vec{\theta}' &\geq \vec{0}.
\end{aligned}$$

For (s, x, i) such that $-\frac{1}{2\rho} < z_{s,x,i} < \frac{1}{2\rho}$, we have $\theta_{s,x,i} = \theta'_{s,x,i} = 0$, and hence,

$$\frac{1}{p_{s,x,i}} \frac{\partial \mathcal{L}}{\partial z_{s,x,i}} = g(s, z_{s,x,i}) = 0 \tag{54}$$

$$\frac{\partial \mathcal{L}}{\partial p_{s,x,i}} = G(s, z_{s,x,i}) + \Phi\left(\frac{1}{2}\right) + \lambda(1-\rho)\omega(\beta) - \eta'\beta + \mu_{s,x} = 0, \quad (55)$$

where

$$g(s, x, z) := \rho\Phi'\left(\frac{1}{2} + \rho z\right) + \lambda(s - 2z) + \eta + \eta'x,$$

$$G(s, x, z) := \int_0^z g(s, x, z) dz.$$

Equations (43) and (44) imply that given (s, x) , $g(s, x, z)$ is always equal to 0 and $G(s, x, z)$ remains the same for all $z \in \{z_{s,x,i}\}_{i \in [m]}$.

By assumption, Φ is symmetric w.r.t. $\frac{1}{2}$ and Φ' is increasing, continuous on $(0, 1)$, and strictly concave on $(0, \frac{1}{2}]$. Hence, for each (s, x) , $g(s, x, z) = 0$ with z unknown has at most three distinct solutions. By the same argument used in the proof of Theorem 4, we have that given (s, x) , z_i and z_{i+1} cannot be solutions of (55) at the same time. Moreover, if given (s, x) , the set $\{z_{s,x,i}\}_{i \in [m]}$ contains two distinct solutions of (54) and (55), then in this case, (54) and (55) must have three distinct solutions, and the smallest and the largest among them, denoted by z_1, z_3 respectively, are contained in $\{z_{s,x,i}\}_{i \in [m]}$. Since G is the integral of g and $G(s, x, z_1) = G(s, x, z_3)$ and Φ' is symmetric w.r.t. $1/2$, we have $z_1 + z_3 = 0$. Hence, one of the following three cases could occur for some \hat{z}_1, \hat{z}_2 .

Case 1: $\{z_{-\frac{1}{2}, -1, i}\}_{i \in [m]} \subseteq \{-\frac{1}{2\rho}, \hat{z}_1\}$ and $\{z_{\frac{1}{2}, -1, i}\}_{i \in [m]} \subseteq \{\hat{z}_2, \frac{1}{2\rho}\}$.

Case 2: $\{z_{-\frac{1}{2}, -1, i}\}_{i \in [m]} \subseteq \{-\hat{z}_1, \hat{z}_1\}$ and $\{z_{\frac{1}{2}, -1, i}\}_{i \in [m]} \subseteq \{\frac{1}{2\rho}\}$.

Case 3: $\{z_{-\frac{1}{2}, -1, i}\}_{i \in [m]} \subseteq \{-\frac{1}{2\rho}\}$ and $\{z_{\frac{1}{2}, -1, i}\}_{i \in [m]} \subseteq \{-\hat{z}_2, \hat{z}_2\}$.

Since by assumption, the optimal solution P_{SXZ} is symmetric in the sense that $P_{SXZ}(-s, -x, -z) = P_{SXZ}(s, x, z)$ for all (s, x, z) , the values of $\{z_{s,1,i}\}_{(s,i) \in \mathcal{S} \times [m]}$ are determined by $\{z_{s,-1,i}\}_{(s,i) \in \mathcal{S} \times [m]}$. Specifically, for Case 1, P_{SXZ} can be expressed as

$$P_{SXZ}(s, x, z) = \begin{cases} \frac{1+2\beta}{4} - p & (s, x, z) = \pm(-\frac{1}{2}, -1, -\frac{1}{2\rho}) \\ p & (s, x, z) = \pm(-\frac{1}{2}, -1, \hat{z}_1) \\ \frac{1-2\beta}{4} - q & (s, x, z) = \pm(\frac{1}{2}, -1, \frac{1}{2\rho}) \\ q & (s, x, z) = \pm(\frac{1}{2}, -1, \hat{z}_2) \end{cases}$$

Solving $\mathbb{E}Z = 0, \mathbb{E}[XZ] = \beta$ gives the solution p, q in (39) and (40) with $z_1 \leftarrow \hat{z}_1, z_2 \leftarrow \hat{z}_2$. This solution yields $\bar{\Upsilon}_\rho$.

For Case 2,

$$P_{SXZ}(s, x, z) = \begin{cases} \frac{1+2\beta}{4} - p & (s, x, z) = \pm(-\frac{1}{2}, -1, -\hat{z}_1) \\ p & (s, x, z) = \pm(-\frac{1}{2}, -1, \hat{z}_1) \\ \frac{1-2\beta}{4} & (s, x, z) = \pm(\frac{1}{2}, -1, \frac{1}{2\rho}) \end{cases}$$

Solving $\mathbb{E}Z = 0, \mathbb{E}[XZ] = \beta$ gives $\hat{z}_1 = -\frac{4\beta\rho - 2\beta + 1}{4\beta\rho - 16\rho\rho + 2\rho}$. However, for this case, $\mathbb{E}[Z^2] - \rho\mathbb{E}[SZ] \geq (1-\rho)/4$, which is strictly larger than $(1-\rho)G(\beta)$ for $\beta \in [0, 1/2)$. Moreover,

$\mathbb{E}[Z^2] - \rho\mathbb{E}[SZ] = (1 - \rho)/4$ holds only if $\beta = p = 1/2$. For the case $\beta = p = 1/2$, we have $z_0 = 1/2$, which corresponds to the case in which the Boolean functions are in fact the dictator functions. This feasible solution is also feasible in the program (38) (specifically, which corresponds to $\beta = p = 1/2, -z_1 = z_2 = 1/2$). Hence the objective value induced by this solution is no more than $\bar{\Upsilon}_\rho$.

Similarly to Case 2, it can be checked that the objective value for Case 3 is also no more than $\bar{\Upsilon}_\rho$. (In fact, Case 3 is the same to Case 2 if we replace $\beta \leftarrow -\beta, X \leftarrow -X$ in Case 2. Note that, in this equivalent setting, β is nonpositive.)

Acknowledgments. This research was supported by by the NSFC grant 62101286 and the Fundamental Research Funds for the Central Universities of China (Nankai University).

Data availability. Data sharing is not applicable to this article as no data were created or analyzed in this study.

References

- [1] Csiszár, I.: Eine informationstheoretische ungleichung und ihre anwendung auf beweis der ergodizitaet von markoffschen ketten. *Magyer Tud. Akad. Mat. Kutato Int. Koezl.* **8**, 85–108 (1964)
- [2] Csiszár, I.: Information-type measures of difference of probability distributions and indirect observation. *Studia Scientiarum Mathematicarum Hungarica* **2**, 229–318 (1967)
- [3] Ali, S.M., Silvey, S.D.: A general class of coefficients of divergence of one distribution from another. *Journal of the Royal Statistical Society: Series B (Methodological)* **28**(1), 131–142 (1966)
- [4] Tsallis, C.: What are the numbers that experiments provide. *Quimica Nova* **17**(6), 468–471 (1994)
- [5] Eldan, R.: A two-sided estimate for the Gaussian noise stability deficit. *Inventiones Mathematicae* **201**(2), 561–624 (2015)
- [6] Li, J., Médard, M.: Boolean functions: Noise stability, non-interactive correlation distillation, and mutual information. *IEEE Trans. Inf. Theory* (2019)
- [7] Mossel, E., O’Donnell, R.: Coin flipping from a cosmic source: On error correction of truly random bits. *Random Structures & Algorithms* **26**(4), 418–436 (2005)
- [8] Sason, I., Verdú, S.: f -divergence inequalities. *IEEE Trans. Inf. Theory* **62**(11), 5973–6006 (2016)
- [9] Gács, P., Körner, J.: Common information is far less than mutual information. *Problems of Control and Information Theory* **2**(2), 149–162 (1973)

- [10] Witsenhausen, H.S.: On sequences of pairs of dependent random variables. *SIAM Journal on Applied Mathematics* **28**(1), 100–113 (1975)
- [11] Kumar, G.R., Courtade, T.A.: Which Boolean functions are most informative? In: 2013 IEEE International Symposium on Information Theory, pp. 226–230 (2013). IEEE
- [12] Courtade, T.A., Kumar, G.R.: Which Boolean functions maximize mutual information on noisy inputs? *IEEE Trans. Inf. Theory* **60**(8), 4515–4525 (2014)
- [13] Anantharam, V., Gohari, A.A., Kamath, S., Nair, C.: On hypercontractivity and the mutual information between Boolean functions. In: Communication, Control, and Computing (Allerton), 2013 51th Annual Allerton Conference On, pp. 13–19 (2013). IEEE
- [14] Kindler, G., O’Donnell, R., Witmer, D.: Remarks on the most informative function conjecture at fixed mean. arXiv preprint arXiv:1506.03167 (2015)
- [15] Ordentlich, O., Shayevitz, O., Weinstein, O.: An improved upper bound for the most informative Boolean function conjecture. In: 2016 IEEE International Symposium on Information Theory (ISIT), pp. 500–504 (2016). IEEE
- [16] Samorodnitsky, A.: On the entropy of a noisy function. *IEEE Trans. Inf. Theory* **62**(10), 5446–5464 (2016)
- [17] Pichler, G., Piantanida, P., Matz, G.: Dictator functions maximize mutual information. *The Annals of Applied Probability* **28**(5), 3094–3101 (2018)
- [18] Barnes, L.P., Özgür, A.: The Courtade-Kumar most informative Boolean function conjecture and a symmetrized Li-Médard conjecture are equivalent. In: 2020 IEEE International Symposium on Information Theory (ISIT), pp. 2205–2209 (2020). IEEE
- [19] Friedgut, E., Kalai, G., Naor, A.: Boolean functions whose fourier transform is concentrated on the first two levels. *Advances in Applied Mathematics* **29**(3), 427–437 (2002)
- [20] Borell, C.: Geometric bounds on the Ornstein–Uhlenbeck velocity process. *Probability Theory and Related Fields* **70**(1), 1–13 (1985)
- [21] Benjamini, I., Kalai, G., Schramm, O.: Noise sensitivity of Boolean functions and applications to percolation. *Publications Mathématiques de l’Institut des Hautes Etudes Scientifiques* **90**(1), 5–43 (1999)
- [22] O’Donnell, R.: *Analysis of Boolean Functions*. Cambridge University Press, ??? (2014)
- [23] Witsenhausen, H., Wyner, A.: A conditional entropy bound for a pair of discrete

- random variables. *IEEE Trans. Inf. Theory* **21**(5), 493–501 (1975)
- [24] Erkip, E.: The efficiency of information in investment. PhD thesis, Ph.D. dissertation, Dept. Electr. Eng., Stanford Univ. Press, Stanford, CA, USA (1996)
- [25] Wyner, A.D., Ziv, J.: A theorem on the entropy of certain binary sequences and applications: Part I. *IEEE Trans. Inf. Theory* **19**(6), 769–772 (1973)
- [26] Yang, H., Wesel, R.D.: On the most informative boolean functions of the very noisy channel. In: 2019 IEEE International Symposium on Information Theory (ISIT), pp. 1202–1206 (2019). IEEE
- [27] Eldan, R., Mikulincer, D., Raghavendra, P.: Noise stability on the Boolean hypercube via a renormalized Brownian motion. arXiv preprint arXiv:2208.06508 (2022)
- [28] Yu, L., Tan, V.Y.F.: On non-interactive simulation of binary random variables. *IEEE Trans. Inf. Theory* **67**(4), 2528–2538 (2021)
- [29] Kahn, J., Kalai, G., Linial, N.: The influence of variables on Boolean functions. In: 29th Annual Symposium on Foundations of Computer Science, pp. 68–80 (1988). IEEE
- [30] Mossel, E., O’Donnell, R., Regev, O., Steif, J.E., Sudakov, B.: Non-interactive correlation distillation, inhomogeneous Markov chains, and the reverse Bonami-Beckner inequality. *Israel Journal of Mathematics* **154**(1), 299–336 (2006)
- [31] Kamath, S., Anantharam, V.: On non-interactive simulation of joint distributions. *IEEE Trans. Inf. Theory* **62**(6), 3419–3435 (2016)
- [32] Ordentlich, O., Polyanskiy, Y., Shayevitz, O.: A note on the probability of rectangles for correlated binary strings. *IEEE Trans. Inf. Theory* **66**(12), 7878–7886 (2020)
- [33] Kirshner, N., Samorodnitsky, A.: A moment ratio bound for polynomials and some extremal properties of Krawchouk polynomials and Hamming spheres. arXiv preprint arXiv:1909.11929 (2019)
- [34] Yu, L., Anantharam, V., Chen, J.: Graphs of joint types, noninteractive simulation, and stronger hypercontractivity. arXiv preprint arXiv:2102.00668 (2021)
- [35] Yu, L.: Strong Brascamp-Lieb inequalities. arXiv preprint arXiv:2102.06935 (2021)
- [36] Bunt, L.N.H.: *Bijdrage Tot de Theorie der Convexe Puntverzamelingen*. Rijksuniversiteit te Groningen, ??? (1934)
- [37] Eggleston, H.G.: *Convexity*. Oxford University Press (1966)

- [38] Peterson, D.W.: A review of constraint qualifications in finite-dimensional spaces. *Siam Review* **15**(3), 639–654 (1973)
- [39] Bazaraa, M.S., Sherali, H.D., Shetty, C.M.: *Nonlinear Programming: Theory and Algorithms*. John Wiley & Sons, ??? (2013)
- [40] Fu, F.-W., Wei, V.K., Yeung, R.W.: On the minimum average distance of binary codes: Linear programming approach. *Discrete Applied Mathematics* **111**(3), 263–281 (2001)
- [41] Yu, L., Tan, V.Y.F.: An improved linear programming bound on the average distance of a binary code. arXiv preprint arXiv:1910.09416 (2019)
- [42] Chang, M.-C.: A polynomial bound in Freiman’s theorem. *Duke mathematical journal* **113**(3), 399–419 (2002)
- [43] Jendrej, J., Oleszkiewicz, K., Wojtaszczyk, J.O.: On some extensions of the FKN theorem. *Theory of Computing* **11**(1), 445–469 (2015)
- [44] König, H., Schütt, C., Tomczak-Jaegermann, N.: Projection constants of symmetric spaces and variants of Khintchine’s inequality. *Journal für die reine und angewandte Mathematik* **1999**(511), 1–42 (1999)