

# Semiparametric Marginal Regression for Clustered Competing Risks Data with Missing Cause of Failure

Wenxian Zhou

Department of Biostatistics, Indiana University

Giorgos Bakoyannis

Department of Biostatistics, Indiana University

Ying Zhang

Department of Biostatistics, University of Nebraska Medical Center

Constantin T. Yiannoutsos

Department of Biostatistics, Indiana University

April 20, 2021

## Abstract

Clustered competing risks data are commonly encountered in multicenter studies. The analysis of such data is often complicated due to informative cluster size, a situation where the outcomes under study are associated with the size of the cluster. In addition, cause of failure is frequently incompletely observed in real-world settings. To the best of our knowledge, there is no methodology for population-averaged analysis with clustered competing risks data with informative cluster size and missing causes of failure. To address this problem, we consider the semiparametric marginal proportional cause-specific hazards model and propose a maximum partial pseudolikelihood estimator under a missing at random assumption. To make the latter assumption more plausible in practice, we allow for auxiliary variables that may be related to

the probability of missingness. The proposed method does not impose assumptions regarding the within-cluster dependence and allows for informative cluster size. The asymptotic properties of the proposed estimators for both regression coefficients and infinite-dimensional parameters, such as the marginal cumulative incidence functions, are rigorously established. Simulation studies show that the proposed method performs well and that methods that ignore the within-cluster dependence and the informative cluster size lead to invalid inferences. The proposed method is applied to competing risks data from a large multicenter HIV study in sub-Saharan Africa where a significant portion of causes of failure is missing.

*Keywords:* Clustered data; Competing risks; Informative cluster size; Missing cause of failure.

# 1 Introduction

Clustered competing risks data are commonly encountered in multicenter studies (Zhou et al., 2012; Diao and Zeng, 2013). An important feature of such data is that the outcomes of individuals from the same cluster are typically dependent and, thus, the standard assumption of independence is violated (Balan and Putter, 2019; Bakoyannis, 2020). Therefore, standard methods for competing risks data are not applicable in the presence of clustering. In addition, cluster size is often informative, in the sense that the outcomes under study are associated with cluster size (Pavlou et al., 2013; Seaman et al., 2014; Cong et al., 2007). An example of a setting with informative cluster size is a study about dental health outcomes. In such studies, dental health outcomes are related to the total number of teeth (cluster size) in a given person (Williamson et al., 2008). With informative cluster size, the standard methods for clustered data lead to bias, since larger clusters are over-represented in the sample and have a larger influence on the parameter estimates. In addition, cause of failure is frequently incompletely observed in real-world settings due to nonresponse/missingness or by the study design (Bakoyannis et al., 2020). A complete case analysis which discards observations with missing event types is expected to lead to bias and efficiency loss (Lu and Tsiatis, 2001; Bakoyannis et al., 2010).

This work is motivated by a large multicenter HIV cohort study conducted by the East Africa Regional Consortium of the International Epidemiology Databases to Evaluate AIDS (EA-IeDEA). A major goal of the study was to evaluate HIV healthcare clinics in East Africa and study two important outcomes in HIV care: (i) disengagement from care and (ii) death while in care (i.e. prior to disengagement). In this study, patients who received antiretroviral

treatment (ART) at the same clinic are expected to have correlated outcomes. In addition, the number of patients in the clinic is expected to be associated with the outcomes of interest, since clinics with more patients are typically better staffed and are expected to provide better care. Furthermore, there was a severe death under-reporting issue in sub-Saharan Africa, which implies that a patient who was lost to clinic was either deceased (and the death was not reported to the clinic) or had disengaged from care. To deal with this issue, EA-IeDEA implemented a double-sampling design where a small subset of patients who were lost to clinic were intensively outreached in the community and their vital status was actively ascertained. This double-sampling design transformed the misclassification problem (i.e. unreported deaths) into a missing data problem by the study design, where cause of failure (i.e. death while in care or disengagement) was unknown for the non-outreached lost patients.

There are two main classes of models for dealing with the within-cluster dependence issue for survival data. One is frailty models (Clayton and Cuzick, 1985; Hougaard, 1986; Liu et al., 2011), which specify explicitly the within-cluster dependence via random effects and provide cluster-specific inference. Such models typically impose assumptions about the structure of the within-cluster dependence and the distribution of the random effects, and tend to be computationally intensive. Under this class of models, Katsahian et al. (2006) proposed a frailty proportional hazards model for the subdistribution of a competing risk, while Scheike et al. (2010) proposed a semiparametric random effects model for competing risks data where the interest is on a particular cause of failure. The other class of models is marginal models (Wei et al., 1989; Liang et al., 1993; Cai and Prentice, 1997; Spiekerman and Lin, 1998; Cai et al., 2000). These models do not rely on assump-

tions regarding the dependence structure and have a population-averaged interpretation. Following this idea, Zhou et al. (2012) proposed a marginal version of the Fine-Gray model (Fine and Gray, 1999) for population-averaged analysis of clustered competing risks. The issue of informative cluster size with survival outcomes has been addressed via a within-cluster resampling method (Cong et al., 2007) and a weighted score function approach (Williamson et al., 2008), where the weights are equal to the inverse of the number of observations in the corresponding cluster.

The issue of missing cause of failure with independent competing risks data has received considerable attention in the literature (Goetghebeur and Ryan, 1995; Lu and Tsiatis, 2001; Craiu and Duchesne, 2004; Gao and Tsiatis, 2005; Lu and Liang, 2008; Bakoyannis et al., 2010; Hyun et al., 2012; Bordes et al., 2014; Nevo et al., 2018; Bakoyannis et al., 2020). Recently, Bakoyannis et al. (2020) proposed a unified framework for semiparametric regression and risk prediction for competing risks data with missing at random (MAR) cause of failure, under the proportional cause-specific hazards model. Unlike previous methods, the approach by Bakoyannis et al. (2020) provides inference for both regression and functional parameters such as the cumulative incidence function. The latter quantity is key for risk prediction in modern medicine. Moreover, simulation studies have shown that the approach by Bakoyannis et al. (2020) provides substantially more efficient regression parameter estimates compared to augmented inverse probability weighting estimators (Gao and Tsiatis, 2005; Hyun et al., 2012) and the multiple-imputation estimator (Lu and Tsiatis, 2001). However, all the aforementioned methods did not consider a potential within-cluster dependence and are thus expected to lead to invalid inferences with clustered data. To the best of our knowledge, only Lee et al. (2017) have addressed the issue of analyzing clustered competing

risks data with missing cause of failure. Lee et al. (2017) proposed a frailty proportional cause-specific hazards model along with a hierarchical likelihood approach for estimation. Nevertheless, this approach does not allow for informative cluster size, it imposes strong assumptions regarding the within-cluster dependence and the distribution of the frailty, which may be violated in practice, and does not provide inference for the infinite-dimensional parameters such as the cumulative incidence function. In addition, the method provides cluster-specific inference and not population-averaged inference which is more scientifically relevant in many applications, including our motivating multicenter HIV study.

To the best of our knowledge, there is no general method for population-averaged inference based on clustered competing risks data with informative cluster size and missing causes of failure. To address this problem, we consider the semiparametric marginal proportional cause-specific hazards model and propose a maximum partial pseudolikelihood estimator under a MAR assumption. The proposed method does not impose assumptions regarding the within-cluster dependence and allows for informative cluster size. Moreover, the method can be easily implemented using off-the-shelf software that allows for case weights, such as the function `coxph` in the R package `survival` (details regarding computation using R are provided in Web Appendix A). The proposed estimators are shown to be strongly consistent and asymptotically normal. Closed-form variance estimators are provided and rigorous methodology for the calculation of simultaneous confidence bands for the infinite-dimensional parameters is proposed. Simulation studies show that the method performs well and that the previously proposed method for missing causes of failure by Bakoyannis et al. (2020), which ignores the within-cluster dependence and the potential informative cluster size, leads to invalid inferences. Finally, the method is applied to the data from the EA-IeDEA study

for illustration.

## 2 Methodology

### 2.1 Notation and Assumptions

Let  $i = 1, 2, \dots, n$  index the  $n$  clusters in the study and  $j = 1, 2, \dots, M_i$  index the subjects in the  $i$ th cluster. Also, let  $T_{ij}$  and  $U_{ij}$  denote the failure and right censoring times for the  $j$ th subject in the  $i$ th cluster. The corresponding observed counterparts are the minimum of the event or censoring times  $X_{ij} = T_{ij} \wedge U_{ij}$  and the failure (from any cause) indicator  $\Delta_{ij} = I(T_{ij} \leq U_{ij})$ . Here we consider the finite observation interval  $[0, \tau]$ , for an arbitrary  $\tau < \infty$ . Suppose that there are  $k$  competing causes of failure, with  $k < \infty$ , and let  $C_{ij} \in \{1, 2, \dots, k\}$  denote the cause of failure for the  $j$ th subject in the  $i$ th cluster. For the sake of generality, cluster size  $M$  is assumed to be random and informative, in the sense that there is an association between the event time and/or cause of failure and  $M$ . However, our proposed methodology applies trivially to simpler situations with non-informative or fixed cluster size. To incorporate missingness in the cause of failure, we define the missing indicator  $R_{ij}$ , with  $R_{ij} = 1$  indicating that the cause of failure for the  $j$ th subject in the  $i$ th cluster is observed, and  $R_{ij} = 0$  otherwise. As in previous works on missing cause of failure (Bakoyannis et al., 2020), we consider the situation where right censoring status is always observed, that is if  $\Delta_{ij} = 0$  then  $R_{ij} = 1$ . The cause of failure  $C_{ij}$  is only observed when both  $\Delta_{ij} = 1$  and  $R_{ij} = 1$ . Let  $\epsilon_{ij} = \Delta_{ij}R_{ij}C_{ij}$  be the observed cause of failure, with  $\epsilon_{ij} = 0$  denoting the cause of failure is missing or censored. The vector of covariates of scientific interest is denoted by

$\mathbf{Z}_{ij} \in \mathbb{R}^p$ . In addition, let  $\mathbf{A}_{ij} \in \mathbb{R}^q$  denote a vector of auxiliary variables, which may not be of scientific interest, but may be related to the probability of missingness. It has been argued that such auxiliary covariates can be used to make the MAR assumption more plausible in practice (Lu and Tsiatis, 2001; Nevo et al., 2018; Bakoyannis et al., 2019, 2020). As usual,  $(T_{ij}, C_{ij})$  and  $U_{ij}$  are assumed independent given  $\mathbf{Z}_{ij}$ . In addition,  $(T_{ij}, C_{ij}, U_{ij})$  are assumed independent across clusters conditionally on  $\mathbf{Z}_{ij}$ . However, within cluster  $i$ ,  $(T_{ij}, C_{ij})$ ,  $j = 1, \dots, M_i$ , are allowed to be dependent given  $\mathbf{Z}_{ij}$ , with an arbitrary dependence structure. Similarly, the right censoring times  $U_{ij}$  may be dependent within cluster  $i$ . The observed data are  $n$  i.i.d. copies of  $\mathbf{D}_i = (\mathbf{X}_i, \Delta_i, \epsilon_i, \mathbf{Z}_i, \mathbf{A}_i, \mathbf{R}_i, M_i)$ ,  $i = 1, \dots, n$ , where  $\mathbf{X}_i = \{X_{ij}\}_{j=1}^{M_i}$ ,  $\Delta_i = \{\Delta_{ij}\}_{j=1}^{M_i}$ ,  $\epsilon_i = \{\epsilon_{ij}\}_{j=1}^{M_i}$ ,  $\mathbf{Z}_i = \{\mathbf{Z}_{ij}\}_{j=1}^{M_i}$ ,  $\mathbf{A}_i = \{\mathbf{A}_{ij}\}_{j=1}^{M_i}$ , and  $\mathbf{R}_i = \{R_{ij}\}_{j=1}^{M_i}$ . To facilitate the presentation of the proposed estimator and its properties, we define the counting process  $N_{ij}(t) = I(X_{ij} \leq t, \Delta_{ij} = 1)$  and at-risk process  $Y_{ij}(t) = I(X_{ij} \geq t)$ . Additionally, we define the cause-specific counting process as  $N_{ijl}(t) = I(X_{ij} \leq t, \Delta_{ijl} = 1) = \Delta_{ijl} N_{ij}(t)$ , where  $\Delta_{ijl} = I(C_{ij} = l, \Delta_{ij} = 1)$ , for  $l = 1, \dots, k$ .

Letting  $\mathbf{W}_{ij} = (X_{ij}, \mathbf{Z}_{ij}, \mathbf{A}_{ij})$ , we impose the MAR assumption  $P(R_{ij} = 1 | C_{ij}, \Delta_{ij} = 1, \mathbf{W}_{ij}) = P(R_{ij} = 1 | \Delta_{ij} = 1, \mathbf{W}_{ij})$ . This assumption is equivalent to

$$\begin{aligned} P(C_{ij} = l | R_{ij} = 1, \Delta_{ij} = 1, \mathbf{W}_{ij}) &= P(C_{ij} = l | R_{ij} = 0, \Delta_{ij} = 1, \mathbf{W}_{ij}) \\ &= P(C_{ij} = l | \Delta_{ij} = 1, \mathbf{W}_{ij}) \\ &\equiv \pi_l(\mathbf{W}_{ij}, \gamma_0), \quad l = 1, \dots, k, \end{aligned}$$

where  $\pi_l(\mathbf{W}_{ij}, \gamma_0)$  is the marginal probability of the failure cause  $l$  given  $\mathbf{W}_{ij}$ , for a non-right-censored observation, and  $\gamma_0$  is assumed to be a finite-dimensional parameter. In Section 2.3 we provide a goodness-of-fit approach for evaluating the appropriateness of this model



in practice.

## 2.2 Estimation Approach

In this work, we provide estimators and inference methodology for both marginal cause-specific hazards and cumulative incidence functions. The covariate-specific marginal cause-specific hazards are defined as

$$\lambda_l(t; \mathbf{z}) = \lim_{h \rightarrow 0} \frac{P(t \leq T_{ij} < t + h, C_{ij} = l \mid T_{ij} \geq t, \mathbf{Z}_{ij} = \mathbf{z})}{h}, \quad l = 1, \dots, k,$$

and the covariate-specific marginal cumulative incidence functions are defined as

$$\begin{aligned} F_l(t; \mathbf{z}) &= P(T_{ij} \leq t, C_{ij} = l \mid \mathbf{Z}_{ij} = \mathbf{z}) \\ &= \int_0^t \exp \left\{ - \sum_{l=1}^k \Lambda_l(s; \mathbf{z}) \right\} \lambda_l(s; \mathbf{z}) ds, \quad l = 1, \dots, k, \end{aligned} \quad (1)$$

where  $\Lambda_l(t; \mathbf{z}) = \int_0^t \lambda_l(s; \mathbf{z}) ds$ , which is the covariate-specific cumulative hazard for the  $l$ th cause of failure. Here, we adopt the marginal proportional cause-specific hazards model

$$\lambda_l(t; \mathbf{z}) = \lambda_{0,l}(t) \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{z}), \quad l = 1, \dots, k,$$

where  $\lambda_{0,l}(t)$  is the  $l$ th unspecified baseline cause-specific hazards function.

When there are no missing causes of failure (i.e.  $R_{ij} = 1$  for  $j = 1, \dots, M_i$  and  $i = 1, \dots, n$ ), estimation for clustered competing risks data can be performed, under the working independence assumption, using the logarithm of the weighted partial likelihood for  $\boldsymbol{\beta} = (\boldsymbol{\beta}_1, \dots, \boldsymbol{\beta}_k)$

$$p l_n(\boldsymbol{\beta}) = \sum_{l=1}^k \sum_{i=1}^n \frac{1}{M_i} \sum_{j=1}^{M_i} \int_0^\tau \left[ \boldsymbol{\beta}_l^T \mathbf{Z}_{ij} - \log \left\{ \sum_{p=1}^n \frac{1}{M_p} \sum_{q=1}^{M_p} Y_{pq}(t) \exp(\boldsymbol{\beta}_l^T \mathbf{Z}_{pq}) \right\} \right] dN_{ijl}(t). \quad (2)$$

This can be seen as the competing risks analogue of the weighted log-partial likelihood by Cong et al. (2007), where the contribution of each subject is weighted by the inverse of the corresponding cluster size to account for informative cluster size.

When cause of failure is missing for some individuals, the weighted log-partial likelihood (2) cannot be evaluated for the observations with a missing cause of failure. For such situations, we propose a weighted partial pseudolikelihood estimator for  $\boldsymbol{\beta}$  which replaces the unobserved cause-specific counting processes with their conditional expectation given the observed data  $\mathbf{D}_{ij}$ . These conditional expectations are equal to

$$\tilde{N}_{ijl}(t; \gamma_0) \equiv E\{N_{ijl}(t) | \mathbf{D}_{ij}\} = \{R_{ij}\Delta_{ijl} + (1 - R_{ij})\pi_l(\mathbf{W}_{ij}, \gamma_0)\} N_{ij}(t),$$

The resulting logarithm of the expected partial pseudolikelihood conditional on the observed data  $\{\mathbf{D}_{ij}\}_{i=1, \dots, n; j=1, \dots, M_i}$  is

$$Q_n(\boldsymbol{\beta}) = \sum_{l=1}^k \sum_{i=1}^n \frac{1}{M_i} \sum_{j=1}^{M_i} \int_0^\tau \left[ \boldsymbol{\beta}_l^T \mathbf{Z}_{ij} - \log \left\{ \sum_{p=1}^n \frac{1}{M_p} \sum_{q=1}^{M_p} Y_{pq}(t) \exp(\boldsymbol{\beta}_l^T \mathbf{Z}_{pq}) \right\} \right] d\tilde{N}_{ijl}(t; \gamma_0). \quad (3)$$

The unknown parameter  $\gamma_0$  in (3) needs to be replaced with an estimate  $\hat{\gamma}_n$ . Such an estimate can be obtained by fitting the marginal binary or multinomial logistic model on the complete cases using generalized estimating equations and under a working independence assumption. Then, estimation of  $\boldsymbol{\beta}_0$  can be performed using the partial pseudoscore function

$$\mathbf{G}_{n,l}(\boldsymbol{\beta}; \hat{\gamma}_n) = \frac{1}{n} \sum_{i=1}^n \frac{1}{M_i} \sum_{j=1}^{M_i} \int_0^\tau \{\mathbf{Z}_{ij} - \mathbf{E}_n(t, \boldsymbol{\beta}_l)\} d\tilde{N}_{ijl}(t; \hat{\gamma}_n), \quad l = 1, \dots, k,$$

where

$$\mathbf{E}_n(t, \boldsymbol{\beta}_l) = \frac{\sum_{p=1}^n \frac{1}{M_p} \sum_{q=1}^{M_p} Y_{pq}(t) \exp(\boldsymbol{\beta}_l^T \mathbf{Z}_{pq}) \mathbf{Z}_{pq}}{\sum_{p=1}^n \frac{1}{M_p} \sum_{q=1}^{M_p} Y_{pq}(t) \exp(\boldsymbol{\beta}_l^T \mathbf{Z}_{pq})}.$$

The estimators  $\hat{\beta}_{n,l}$  are the solutions to  $\mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \hat{\gamma}_n) = \mathbf{0}$ ,  $l = 1, \dots, k$ . This estimation procedure can be easily implemented using `coxph` function in the R package `survival` with some data management. An illustration of the use of the `coxph` function to obtain parameter estimates with the proposed approach is provided in Web Appendix A.

For  $l = 1, \dots, k$  and  $t \in [0, \tau]$ , the Breslow-type estimator for the marginal cause-specific baseline cumulative hazard function is

$$\hat{\Lambda}_{n,l}(t) = \sum_{i=1}^n \frac{1}{M_i} \sum_{j=1}^{M_i} \int_0^t \frac{d\tilde{N}_{ijl}(u; \hat{\gamma}_n)}{\sum_{p=1}^n \frac{1}{M_p} \sum_{q=1}^{M_p} Y_{pq}(u) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_{pq})}.$$

Based on this estimator, the marginal covariate-specific cumulative incidence function can be estimated by

$$\hat{F}_{n,l}(t; \mathbf{z}_0) = \int_0^t \exp \left\{ - \sum_{l=1}^k \hat{\Lambda}_{n,l}(u-; \mathbf{z}_0) \right\} d\hat{\Lambda}_{n,l}(u; \mathbf{z}_0),$$

where  $\hat{\Lambda}_{n,l}(t; \mathbf{z}_0) = \hat{\Lambda}_{n,l}(t) \exp(\hat{\beta}_{n,l}^T \mathbf{z}_0)$ .

## 2.3 Asymptotic Properties

Here, we state the main theorems for the asymptotic properties of the proposed estimators  $\hat{\beta}_{n,l}$ ,  $\hat{\Lambda}_{n,l}(t)$  and  $\hat{F}_{n,l}(t; \mathbf{z}_0)$ . The detailed proofs of these theorems are provided in Web Appendix B. Before stating the regularity conditions, we define the negative of the true Hessian matrix as

$$\mathbf{H}_l(\beta) = E \left\{ \frac{1}{M} \sum_{j=1}^M \int_0^\tau \mathbf{V}(t, \beta) d\tilde{N}_{jl}(t; \gamma_0) \right\},$$

where

$$\mathbf{V}(t, \beta) = \frac{E\{\frac{1}{M} \sum_{j=1}^M Y_j(t) \exp(\beta^T \mathbf{Z}_j) \mathbf{Z}_j^{\otimes 2}\}}{E\{\frac{1}{M} \sum_{j=1}^M Y_j(t) \exp(\beta^T \mathbf{Z}_j)\}} - \left[ \frac{E\{\frac{1}{M} \sum_{j=1}^M Y_j(t) \exp(\beta^T \mathbf{Z}_j) \mathbf{Z}_j\}}{E\{\frac{1}{M} \sum_{j=1}^M Y_j(t) \exp(\beta^T \mathbf{Z}_j)\}} \right]^{\otimes 2}.$$

The following regularity conditions are assumed throughout the remainder of this paper.

- C1.  $\Lambda_{0,l}(t)$  is a non-decreasing continuous function with  $\Lambda_{0,l}(\tau) < \infty$ , for  $l = 1, \dots, k$  and  $E\{Y(\tau) \mid \mathbf{Z}, M\} > 0$  almost surely.
- C2. The true regression coefficients  $\beta_{0,l} \in \mathcal{B}_l \subset \mathbb{R}^{p_l}$ , where  $\mathcal{B}_l$  is bounded and convex set for  $l = 1, \dots, k$  and  $\beta_{0,l}$  is in interior of  $\mathcal{B}_l$ .
- C3. The inverse of the link function  $g$  for the marginal probability model of the cause of failure  $\pi_l(\mathbf{W}_{ij}, \gamma_0)$ ,  $l = 1, \dots, k$ , has continuous derivative  $\dot{g}$  with respect to  $\gamma_0$  on compact sets. The parameter space  $\Gamma$  of  $\gamma_0$  is a bounded subset of  $\mathbb{R}^{p_\gamma}$ .
- C4. The estimating function for the model of the cause of failure is Lipschitz continuous in  $\gamma$ , and the estimator  $\hat{\gamma}_n$  is strongly consistent and asymptotically linear, i.e.  $\sqrt{n}(\hat{\gamma}_n - \gamma_0) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{1}{M_i} \sum_{j=1}^{M_i} \omega_{ij} + o_p(1)$ , where  $\omega_{ij}$  is the influence function of  $j$ th subject in  $i$ th cluster, satisfying  $E(\omega_{ij}) = 0$  and  $E \|\omega_{ij}\|^2 < \infty$ .
- C5. The covariates of interest  $\mathbf{Z}$ , the auxiliary covariates  $\mathbf{A}$ , and the cluster size  $M$  are bounded, in the sense that there exist constants  $K \in \mathbb{R}_+$  and  $m_0 \in \mathbb{N}_+$  such that  $P(\|\mathbf{Z}\| \vee \|\mathbf{A}\| \leq K) = 1$  and  $P(M \leq m_0) = 1$ .
- C6. The true Hessian matrix  $-\mathbf{H}_l(\beta)$  is negative definite for all  $l = 1, \dots, k$ .

Regularity conditions C3 and C4 are satisfied when the marginal model for  $\pi_l(\mathbf{W}_{ij}, \gamma_0)$  is correctly specified with parameters estimated through generalized estimating equations under a working independence assumption. The assumptions on the parametric models for  $\pi_l(\mathbf{W}_{ij}, \gamma_0)$  can be evaluated using the cumulative residual processes

$$E \left[ \frac{1}{M} \sum_{j=1}^M R_j \{N_{jl}(t) - \pi_l(\mathbf{W}_j, \gamma_0) N_j(t)\} \right], \quad l = 1, \dots, k-1, \quad t \in [0, \tau],$$

which can be estimated by

$$\frac{1}{n} \sum_{i=1}^n \frac{1}{M_i} \sum_{j=1}^{M_i} R_{ij} \{N_{ijl}(t) - \pi_l(\mathbf{W}_{ij}, \hat{\gamma}_n) N_{ij}(t)\}.$$

If the model is correctly specified, the cumulative residual process is equal to 0 for  $t \in [0, \tau]$ .

A formal goodness of fit test can be conducted using the simulation-based approach by Pan and Lin (2005). A graphical evaluation of goodness of fit can also be performed by plotting the observed residual process and the 95% simultaneous confidence band around the line  $f(t) = 0$ ,  $t \in [0, \tau]$  (Bakoyannis et al., 2019, 2020).

Theorem 1 states the consistency of the proposed estimators  $\hat{\beta}_{n,l}$  and  $\hat{\Lambda}_{n,l}(t)$ . The proof of Theorem 1 is given in Web Appendix B.1.

**Theorem 1.** *Under the assumptions in Section 2.1 and regularity conditions C1 - C6,*

$$\sum_{l=1}^k \left\{ \|\hat{\beta}_{n,l} - \beta_{0,l}\| + \|\hat{\Lambda}_{n,l}(t) - \Lambda_{0,l}(t)\|_{\infty} \right\} \rightarrow_{as*} 0,$$

as  $n \rightarrow \infty$ , where  $\|f(t)\|_{\infty} = \sup_{t \in [0, \tau]} |f(t)|$ .

A corollary of Theorem 1 is the strong uniform consistency of  $\hat{F}_{n,l}(t; \mathbf{z}_0)$ ,  $l = 1, \dots, k$ , that is  $\sum_{l=1}^k \|\hat{F}_{n,l}(t; \mathbf{z}_0) - F_{0,l}(t; \mathbf{z}_0)\|_{\infty} \rightarrow_{as*} 0$ .

Theorem 2 provides the asymptotic distribution for the finite-dimensional parameter  $\hat{\beta}_{n,l}$ , which provides the basis for statistical inference about the regression coefficients  $\beta_{0,l}$ , for  $l = 1, \dots, k$ . The proof of Theorem 2 is given in Web Appendix B.2. Before providing the theorem, we define the following quantities

$$\psi_{ijl} = \mathbf{H}_l^{-1}(\beta_{0,l}) \int_0^{\tau} \{\mathbf{Z}_{ij} - \mathbf{E}(t, \beta_{0,l})\} d\tilde{M}_{ijl}(t; \beta_{0,l}, \gamma_0),$$

where

$$\mathbf{E}(t, \beta_{0,l}) = \frac{E\{\frac{1}{M} \sum_{j=1}^M Y_j(t) \exp(\beta_{0,l}^T \mathbf{Z}_j) \mathbf{Z}_j\}}{E\{\frac{1}{M} \sum_{j=1}^M Y_j(t) \exp(\beta_{0,l}^T \mathbf{Z}_j)\}},$$

and  $\tilde{M}_{ijl}(t; \boldsymbol{\beta}_{0,l}, \boldsymbol{\gamma}_0) = \tilde{N}_{ijl}(t; \boldsymbol{\gamma}_0) - \int_0^t Y_{ij}(u) \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{Z}_{ij}) d\Lambda_{0,l}(u)$ , where

$$\Lambda_{0,l}(t) = \int_0^t \frac{E\{\frac{1}{M} \sum_{j=1}^M d\tilde{N}_{jl}(u; \boldsymbol{\gamma}_0)\}}{E\{\frac{1}{M} \sum_{j=1}^M Y_j(u) \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{Z}_j)\}}.$$

Finally, we define the non-random quantity

$$\mathbf{R}_l = \mathbf{H}_l^{-1}(\boldsymbol{\beta}_{0,l}) E \left[ \frac{1}{M} \sum_{j=1}^M (1 - R_j) \int_0^\tau \{\mathbf{Z}_j - \mathbf{E}(t, \boldsymbol{\beta}_{0,l})\} dN_j(t) \dot{\pi}_l(\mathbf{W}_j, \boldsymbol{\gamma}_0)^T \right],$$

where  $\dot{\pi}_l(\mathbf{W}_j, \boldsymbol{\gamma}_0) = \partial\{\pi_l(\mathbf{W}_j, \boldsymbol{\gamma})\}(\partial\boldsymbol{\gamma})^{-1}|_{\boldsymbol{\gamma}=\boldsymbol{\gamma}_0}$ .

**Theorem 2.** *Under the assumptions in Section 2.1 and regularity conditions C1 - C6, for*

$$l = 1, \dots, k, \quad \sqrt{n}(\hat{\boldsymbol{\beta}}_{n,l} - \boldsymbol{\beta}_{0,l}) = n^{-1/2} \sum_{i=1}^n \left\{ M_i^{-1} \sum_{j=1}^{M_i} (\boldsymbol{\psi}_{ijl} + \mathbf{R}_l \boldsymbol{\omega}_{ij}) \right\} + o_p(1).$$

By Theorem 2,  $\sqrt{n}(\hat{\boldsymbol{\beta}}_{n,l} - \boldsymbol{\beta}_{0,l}) \rightarrow_d N(\mathbf{0}, \boldsymbol{\Sigma}_l)$ , where  $\boldsymbol{\Sigma}_l = E\{M^{-1} \sum_{j=1}^M (\boldsymbol{\psi}_{jl} + \mathbf{R}_l \boldsymbol{\omega}_j)\}^{\otimes 2}$ .

The covariance matrix  $\boldsymbol{\Sigma}_l$  can be consistently estimated using the empirical versions of the influence functions by

$$\hat{\boldsymbol{\Sigma}}_l = \frac{1}{n} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} (\hat{\boldsymbol{\psi}}_{ijl} + \hat{\mathbf{R}}_l \hat{\boldsymbol{\omega}}_{ij}) \right\}^{\otimes 2}.$$

The empirical versions of the influence functions can be obtained by replacing expectations with sample averages over clusters and unknown parameters with their consistent estimates.

Explicit formulas for the empirical versions of the influence functions are provided in Web Appendix B.5.

Theorems 3 and 4 provide the weak convergence of  $\hat{\Lambda}_{n,l}(t)$  and  $\hat{F}_{n,l}(t; \mathbf{z}_0)$ , respectively. Before providing these theorems, we define some useful quantities that appear in the influence functions of the estimators of the infinite-dimensional parameters. For  $l = 1, \dots, k$  and  $t \in [0, \tau]$ , define

$$\phi_{ijl}(t) = \int_0^t \frac{d\tilde{M}_{ijl}(s; \boldsymbol{\beta}_{0,j}, \boldsymbol{\gamma}_0)}{E\{\frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{Z}_j)\}} - (\boldsymbol{\psi}_{ijl} + \mathbf{R}_l \boldsymbol{\omega}_{ij})^T \int_0^t \mathbf{E}(s, \boldsymbol{\beta}_{0,l}) d\Lambda_{0,l}(s),$$

and the non-random function

$$\mathbf{R}_j^*(t) = E \left[ \frac{1}{M} \sum_{j=1}^M (1 - R_j) \dot{\pi}_j(\mathbf{W}_j, \gamma_0) \int_0^t \frac{dN_j(s)}{E\{\frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{Z}_j)\}} \right]^T.$$

In addition, we define the influence function

$$\begin{aligned} \phi_{ijl}^F(t; \mathbf{z}_0) = & \int_0^t \exp \left\{ - \sum_{l=1}^k \Lambda_{0,l}(s-; \mathbf{z}_0) \right\} d\phi_{ijl}^\Lambda(s; \mathbf{z}_0) \\ & - \int_0^t \left\{ \sum_{l=1}^k \phi_{ijl}^\Lambda(s-; \mathbf{z}_0) \right\} \exp \left\{ - \sum_{l=1}^k \Lambda_{0,l}(s-; \mathbf{z}_0) \right\} d\Lambda_{0,l}(s; \mathbf{z}_0), \end{aligned}$$

where  $\phi_{ijl}^\Lambda(t; \mathbf{z}_0) = \{\mathbf{z}_0^T(\boldsymbol{\psi}_{ijl} + \mathbf{R}_l \boldsymbol{\omega}_{ij}) \Lambda_{0,l}(t) + \phi_{ijl}(t) + \mathbf{R}_l^*(t) \boldsymbol{\omega}_{ij}\} \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{z}_0)$ . Finally,  $D[0, \tau]$  denotes the space of right-continuous functions with left-hand limits defined on  $[0, \tau]$ , and  $\{\xi_i\}_{i=1}^n$  are standard normal variables independent of the data. The proofs of the following theorems are given in Web Appendices B.3 and B.4.

**Theorem 3.** *Under the assumptions in Section 2.1 and regularity conditions C1 - C6, for  $l = 1, \dots, k$ , and  $t \in [0, \tau]$ ,*

$$\sqrt{n} \left\{ \hat{\Lambda}_{n,l}(t) - \Lambda_{0,l}(t) \right\} = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left[ \frac{1}{M_i} \sum_{j=1}^{M_i} \{ \phi_{ijl}(t) + \mathbf{R}_l^*(t) \boldsymbol{\omega}_{ij} \} \right] + o_p(1),$$

*with the influence functions belonging to a Donsker class, and conditional on the observed data,  $\hat{W}_{n,l}(\cdot) = n^{-1/2} \sum_{i=1}^n [M_i^{-1} \sum_{j=1}^{M_i} \{ \hat{\phi}_{ijl}(\cdot) + \hat{\mathbf{R}}_l^*(\cdot) \hat{\boldsymbol{\omega}}_{ij} \}] \xi_i$  converges weakly to the same limiting process as  $W_{n,l}(\cdot) = \sqrt{n} \{ \hat{\Lambda}_{n,l}(\cdot) - \Lambda_{0,l}(\cdot) \}$ .*

By Theorem 3,  $\sqrt{n} \{ \hat{\Lambda}_{n,l}(\cdot) - \Lambda_{0,l}(\cdot) \} \rightsquigarrow \mathbb{G}_{\Lambda_l}$  in  $D[0, \tau]$ , where  $\mathbb{G}_{\Lambda_l}$  is a tight mean zero Gaussian process with covariance function  $E[M^{-1} \sum_{j=1}^M \{ \phi_{jl}(t) + \mathbf{R}_l^*(t) \boldsymbol{\omega}_j \}][M^{-1} \sum_{j=1}^M \{ \phi_{jl}(s) + \mathbf{R}_l^*(s) \boldsymbol{\omega}_j \}]$ ,  $t, s \in [0, \tau]$ . A consistent estimator of the covariance function is

$$\frac{1}{n} \sum_{i=1}^n \left[ \frac{1}{M_i} \sum_{j=1}^{M_i} \left\{ \hat{\phi}_{ijl}(t) + \hat{\mathbf{R}}_l^*(t) \hat{\boldsymbol{\omega}}_{ij} \right\} \right] \left[ \frac{1}{M_i} \sum_{j=1}^{M_i} \left\{ \hat{\phi}_{ijl}(s) + \hat{\mathbf{R}}_l^*(s) \hat{\boldsymbol{\omega}}_{ij} \right\} \right], \quad t, s \in [0, \tau].$$

Explicit formulas for the empirical versions of the influence functions are provided in Web Appendix B.5. Calculation of confidence intervals and bands can be performed using an appropriate continuously differentiable transformation to avoid negative limits (Lin et al., 1994). A standard choice is the transformation  $g(x) = \log(x)$ . According to the functional delta method,  $\sqrt{n}q_l^\Lambda(t)[g\{\hat{\Lambda}_{n,l}(t)\} - g\{\Lambda_{0,l}(t)\}]$  is asymptotically equivalent to  $B_{n,l}(t) = q_l^\Lambda(t)\dot{g}\{\hat{\Lambda}_{n,l}(t)\}W_{n,l}(t)$ . Also, by Theorem 3,  $B_{n,l}(t)$  is asymptotically equivalent to  $\hat{B}_{n,l}(t) = q_l^\Lambda(t)\dot{g}\{\hat{\Lambda}_{n,l}(t)\}\hat{W}_{n,l}(t)$ , where  $q_l^\Lambda(t)$  is a weight function, with  $t \in [t_1, t_2]$ ,  $0 \leq t_1 \leq t_2 < \tau$ . The choice  $q_l^\Lambda(t) = \hat{\Lambda}_{n,l}(t)\{\hat{\sigma}_{\Lambda_l}(t)\}^{-1}$ , where  $\hat{\sigma}_{\Lambda_l}(t)$  is the square root of the estimated variance of  $\hat{\Lambda}_{n,l}(t)$ , gives the equal precision band (Nair, 1984); the choice  $q_l^\Lambda(t) = \hat{\Lambda}_{n,l}(t)\{1 + \hat{\sigma}_{\Lambda_l}^2(t)\}^{-1}$ , provides a Hall-Wellner-type band (Hall and Wellner, 1980). Now, a  $1 - \alpha$  confidence band for  $\Lambda_{0,l}(t)$  can be computed as

$$g^{-1} \left[ g \left\{ \hat{\Lambda}_{n,l}(t) \right\} \pm \frac{c_\alpha}{\sqrt{n}q_l^\Lambda(t)} \right], \quad t \in [t_1, t_2],$$

where  $c_\alpha$  is the  $1 - \alpha$  quantile of the distribution of  $\sup_{t \in [t_1, t_2]} |\hat{B}_{n,l}(t)|$ . This can be estimated using a large number of simulation realizations from the process  $\hat{B}_{n,l}(\cdot)$ , generated by repeatedly simulating sets of standard normal variables  $\{\xi_i\}_{i=1}^n$  (Spiekerman and Lin, 1998). Since confidence bands tend to be unstable at earlier and later time points, where there are fewer observed events, we suggest the restriction of the confidence band domain  $[t_1, t_2]$  to the 10th and 90th percentile of the event times.

**Theorem 4.** *Under the assumptions in Section 2.1 and regularity conditions C1 - C6, for  $l = 1, \dots, k$ , and  $t \in [0, \tau]$ ,*

$$\sqrt{n} \left\{ \hat{F}_{n,l}(t; \mathbf{z}_0) - F_{0,l}(t; \mathbf{z}_0) \right\} = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} \phi_{ijl}^F(t; \mathbf{z}_0) \right\} + o_p(1),$$



with the influence functions belonging to a Donsker class, and conditionally on the observed data,  $\hat{W}_{n,l}^F(\cdot; \mathbf{z}_0) = n^{-1/2} \sum_{i=1}^n \{M_i^{-1} \sum_{j=1}^{M_i} \hat{\phi}_{ijl}^F(\cdot; \mathbf{z}_0)\} \xi_i$  converges weakly to the same limiting process as  $W_{n,l}^F(\cdot; \mathbf{z}_0) = \sqrt{n} \{\hat{F}_{n,l}(\cdot; \mathbf{z}_0) - F_{0,l}(\cdot; \mathbf{z}_0)\}$ .

By Theorem 4,  $\sqrt{n} \{\hat{F}_{n,l}(\cdot; \mathbf{z}_0) - F_{0,l}(\cdot; \mathbf{z}_0)\} \rightsquigarrow \mathbb{G}_{F_l}$  in  $D[0, \tau]$ , where  $\mathbb{G}_{F_l}$  is a tight mean zero Gaussian process with covariance function  $E\{M^{-1} \sum_{j=1}^M \phi_{jl}^F(t; \mathbf{z}_0)\} \{M^{-1} \sum_{j=1}^M \phi_{jl}^F(s; \mathbf{z}_0)\}$ .

A consistent estimator for the covariance function is

$$\frac{1}{n} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} \hat{\phi}_{ijl}^F(t; \mathbf{z}_0) \right\} \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} \hat{\phi}_{ijl}^F(s; \mathbf{z}_0) \right\}, \quad t, s \in [0, \tau].$$

Explicit formulas for the empirical versions of the influence functions are provided in Web Appendix B.5. Similarly to the case of the cumulative baseline hazards  $\Lambda_{0,l}$ , a  $1 - \alpha$  confidence band for  $F_{0,l}(\cdot; \mathbf{z}_0)$  can be constructed as

$$g^{-1} \left[ g \left\{ \hat{F}_{0,l}(t; \mathbf{z}_0) \right\} \pm \frac{c_\alpha}{\sqrt{n} q_l^F(t; \mathbf{z}_0)} \right], \quad t \in [t_1, t_2],$$

where  $c_\alpha$  is the  $1 - \alpha$  quantile of the distribution of  $\sup_{t \in [t_1, t_2]} |\hat{B}_{n,l}^F(t; \mathbf{z}_0)|$ , with  $\hat{B}_{n,l}^F(t; \mathbf{z}_0) = q_l^F(t; \mathbf{z}_0) \dot{g}\{\hat{F}_{n,l}(t; \mathbf{z}_0)\} \hat{W}_{n,l}^F(t; \mathbf{z}_0)$ . A standard transformation to ensure that the limits of the bands for the cumulative incidence functions reside in  $[0, 1]$  is  $g(x) = \log\{-\log(x)\}$  (Cheng et al., 1998). The weight function choice  $q_l^F(t; \mathbf{z}_0) = \hat{F}_{n,l}(t; \mathbf{z}_0) \log\{\hat{F}_{n,l}(t; \mathbf{z}_0)\} \{\hat{\sigma}_{F_l}(t; \mathbf{z}_0)\}^{-1}$ , with  $\hat{\sigma}_{F_l}(t; \mathbf{z}_0)$  being the square root of the estimated variance of  $\hat{F}_{0,l}(t; \mathbf{z}_0)$ , provides an equal-precision-type band (Nair, 1984); the choice  $q_l^F(t; \mathbf{z}_0) = \hat{F}_{n,l}(t; \mathbf{z}_0) \log\{\hat{F}_{n,l}(t; \mathbf{z}_0)\} \{1 + \hat{\sigma}_{F_l}^2(t; \mathbf{z}_0)\}^{-1}$ , provides a Hall-Wellner-type band (Hall and Wellner, 1980).

### 3 Simulation Studies

To evaluate the finite-sample performance of the proposed estimators, and compare them with the estimators by Bakoyannis et al. (2020) which ignore the within-cluster dependence, we conducted a series of simulation experiments. We considered simulation settings similar to those used in Bakoyannis et al. (2020), with two competing risks, two covariates  $\mathbf{Z} = (Z_1, Z_2)^T$ , where  $Z_1 \sim N(0, 2^2)$  and  $Z_2 \sim \text{Bernoulli}(0.5)$ , and with the observation time interval being  $[0, 2]$ . The right censoring times were independently generated from the  $\text{Exp}(0.4)$  distribution. For each cause, the failure times were generated from Cox proportional hazards shared frailty models with a positive stable frailty (Hougaard, 1986; Cong et al., 2007; Liu et al., 2011) to introduce within-cluster dependence. These models had the form

$$\lambda_l(t \mid Z_{ij,l}, w_{i,l}) = \lambda_{0,l}(t) w_{i,l} \exp(\beta_{0,l} Z_{ij,l}), \quad l = 1, 2,$$

where  $w_{i,l}$  followed a positive stable distribution with parameter  $\alpha = 0.5$ , which induced a moderate within-cluster dependence. This simulation setup led to the marginal cause-specific hazard functions

$$\lambda_l(t \mid Z_{ij,l}) = \alpha \lambda_{0,l}(t) \Lambda_{0,l}(t)^{\alpha-1} \exp(\alpha \beta_{0,l} Z_{ij,l}), \quad l = 1, 2,$$

which were still proportional, owing to the positive stable frailty, with true parameters  $\beta'_{0,l} = \alpha \beta_{0,l}$ , and  $\Lambda'_{0,l}(t) = \Lambda_{0,l}(t)^\alpha$ .

In this simulation study we considered two scenarios. In both scenarios, the event time for the cause 1 was generated assuming  $\lambda_{0,1}(t) = 1$  and  $\beta_{0,1} = -0.5$ . In Scenario 1, the event time for cause 2 was simulated from a Gompertz distribution with baseline hazard  $\lambda_{0,2}(t) = \exp(-0.5 + 0.2t)$ , and  $\beta_{0,2} = -0.5$ . In Scenario 2, the event time for cause 2

was simulated from a Weibull distribution with baseline hazard  $\lambda_{0,1}(t) = \{2\sqrt{2t}\}^{-1}$  and  $\beta_{0,2} = -0.5$ . The simulation setup under Scenario 1 led to on average in 13.5% right-censored observations, 50.4% failures from cause 1, and 36.1% failures from cause 2. The corresponding figures for Scenario 2 were 12.7%, 41.8%, and 45.5%. The implied model for  $\pi_1(\mathbf{W}_{ij}, \boldsymbol{\gamma}_0)$  had approximately linear time effect with the form  $\text{logit}\{\pi_1(\mathbf{W}_{ij}, \boldsymbol{\gamma}_0)\} \approx \gamma_0 + \gamma_1 T + \gamma_2 Z_1 + \gamma_3 Z_2$  under Scenario 1, where  $\boldsymbol{\gamma}_0 \approx (0.25, -0.15, -0.25, 0.25)^T$ , and had the form  $\text{logit}\{\pi_1(\mathbf{W}_{ij}, \boldsymbol{\gamma}_0)\} = \gamma_0 + \gamma_1 \log(T) + \gamma_2 Z_1 + \gamma_3 Z_2$  under Scenario 2, where  $\boldsymbol{\gamma}_0 = (5 \log(2)/4, 0.25, -0.25, 0.25)^T$ . The missingness indicators  $R_{ij}$  were generated under the logistic model

$$\text{logit}\{P(R_{ij} = 1 \mid \Delta_{ij} = 1, \mathbf{W}_{ij})\} = (1, \mathbf{W}_{ij}^T) \boldsymbol{\theta},$$

where  $\mathbf{W}^T = (T, Z_1, Z_2)$ . We considered the parameter values  $\boldsymbol{\theta} = (0.7, 1, -1, 1)^T, (-0.2, 1, -1, 1)^T$ , and  $(-0.8, 1, -1, 1)^T$ , which resulted in 24.5%, 35.1%, and 42.8% missing causes of failure in Scenario 1, and 25.5%, 36.3%, and 44.1% missingness in Scenario 2.

In this simulation study we considered  $n = 50, 100$ , or  $200$  which correspond to situations with small to moderate number of clusters. To introduce informative cluster size, the cluster sizes  $M_i$  were generated from a mixture of discrete uniform distributions depending on the frailty, with  $M_i \sim \text{Unif}(20, 30)$  if  $w_{i,1} < \text{median}(w_1)$  and  $w_{i,2} < \text{median}(w_2)$ ,  $M_i \sim \text{Unif}(50, 60)$  if  $w_{i,1} \geq \text{median}(w_1)$  and  $w_{i,2} \geq \text{median}(w_2)$ , and  $M_i \sim \text{Unif}(30, 50)$ , otherwise.

For each simulation setting, we simulated 1000 datasets, and analyzed each dataset using the proposed method and the method by Bakoyannis et al. (2020). All analysis assumed the parametric model  $\text{logit}\{\pi_1(\mathbf{W}_{ij}, \boldsymbol{\gamma}_0)\} = \gamma_0 + \gamma_1 T + \gamma_2 Z_1 + \gamma_3 Z_2$ , which was approximately

correctly specified under Scenario 1, and misspecified under Scenario 2. The standard errors were estimated using the closed-form formulas provided in Section 2. The 95% confidence bands for  $\Lambda_{0,1}(t)$  and  $F_{0,1}(t)$  were computed based on 1000 simulation realizations standard normal variables  $\{\xi_i\}_{i=1}^n$  as described in Section 2. The limits of the time domain  $[t_1, t_2]$  for the confidence bands were chosen to be 10% and 90% percentile of the observed failure times.

The simulation results for the regression coefficient  $\beta_1$  under Scenario 1 are summarized in Table 1. The proposed estimator was approximately unbiased and the average standard errors were close to the Monte Carlo standard deviations. This provides numerical evidence for the consistency of our estimator of the regression coefficient  $\hat{\beta}_{n,1}$  and its associated standard error. The 95% coverage probabilities were close to the nominal level in all cases. In contrast, the method by Bakoyannis et al. (2020) provided biased estimates. The bias for  $\beta_1$  was relatively small under Scenario 1, with an increasing trend as the number of clusters increased. The average standard errors were smaller than the Monte Carlo standard deviation, which implies that the standard errors were under-estimated. This resulted in poor coverage probabilities of the corresponding 95% confidence intervals.

The simulation results for the pointwise estimates of the infinite-dimensional parameters  $\Lambda_{0,1}(t)$  and  $F_{0,1}(t)$  under Scenario 1 are provided in Web Appendix C. Our proposed estimators had good performance with small bias, average standard errors close to the Monte Carlo standard deviation, and 95% confidence interval coverage probabilities close to the nominal level. As expected, the method by Bakoyannis et al. (2020), which ignores the within-cluster dependence and the informative cluster size, provided estimators with large bias, severely under-estimated standard errors, and poor coverage probabilities of the 95%

Table 1: Simulation results for the regression coefficient  $\beta_1$  under Scenario 1 for the proposed approach and the approach by Bakoyannis et al. (2020) (BZY20) which ignores the within-cluster dependence.

$n$	$p_m(\%)$	Proposed				BZY20			
		Bias	MCSD	ASE	CP	Bias	MCSD	ASE	CP
50	25	-0.006	0.033	0.032	0.937	0.003	0.033	0.021	0.782
	35	-0.006	0.034	0.033	0.938	0.003	0.035	0.023	0.793
	43	-0.006	0.036	0.034	0.939	0.004	0.035	0.025	0.827
100	25	-0.002	0.022	0.022	0.949	0.007	0.022	0.015	0.777
	35	-0.002	0.023	0.023	0.941	0.007	0.023	0.016	0.803
	43	-0.002	0.024	0.024	0.948	0.007	0.024	0.018	0.822
200	25	-0.001	0.016	0.016	0.954	0.008	0.016	0.010	0.735
	35	-0.001	0.017	0.017	0.953	0.008	0.017	0.011	0.762
	43	-0.001	0.017	0.017	0.953	0.009	0.017	0.013	0.785

Note:  $n$ : number of clusters with cluster size  $M \in [30, 60]$ ;  $p_m$ : percentage of missingness; MCSD: Monte Carlo standard deviation; ASE: average estimated standard error; CP: coverage probability of 95% confidence interval

confidence intervals.

Table 2 presents the coverage probabilities of 95% simultaneous confidence bands for the infinite-dimensional parameters  $\Lambda_{0,1}(t)$  and  $F_{0,1}(t)$  under Scenario 1. The proposed 95% simultaneous confidence bands had coverage probabilities close to the nominal level, while the 95% simultaneous confidence bands by Bakoyannis et al. (2020) had a very poor coverage rate.

The simulation results under Scenario 2, where the model for  $\pi_1(\mathbf{W}_{ij}, \gamma)$  was misspecified, are provided in Web Appendix C. The results for point estimates under this scenario were similar to those from Scenario 1 (Table 1). This provides numerical evidence for the robustness of the proposed approach under some degree of model misspecification in  $\pi_l(\mathbf{W}_{ij}, \gamma)$ . However, the confidence bands had lower coverage rate under Scenario 2. This illustrates the importance to evaluate the goodness of fit of the model assumption for  $\pi_l(\mathbf{W}_{ij}, \gamma)$  in practice, as described in Section 2.3.

## 4 HIV Data Application

The proposed method was applied to the electronic health record data from the EA-IeDEA study to identify factors affecting disengagement from HIV care and death while in care (i.e. prior to a disengagement). Disengagement from care and death while in care were the two competing risks of interest. Disengagement from care was defined by the clinical investigators of the study as being alive and without HIV care for two months. The covariates of interest were sex, age, CD4 count at ART initiation, and HIV status disclosure. The dataset included 24373 HIV-infected adult patients from 31 clinics who initiated ART on/after January 1,

Table 2: Simulation results for the coverage probabilities of 95% simultaneous confidence bands for the infinite-dimensional parameters  $\Lambda_{0,1}(t)$  and  $F_{0,1}(t)$  under Scenario 1. Results from the proposed approach and the approach by Bakoyannis et al. (2020) (BZY20) which ignores the within-cluster dependence.

$n$	$p_m(\%)$	$\Lambda_{0,1}(t)$				$F_{0,1}(t)$			
		Proposed		BZY20		Proposed		BZY20	
		EP	HW	EP	HW	EP	HW	EP	HW
50	25	0.900	0.936	0.077	0.153	0.906	0.931	0.120	0.203
	35	0.912	0.936	0.105	0.185	0.904	0.928	0.146	0.245
	43	0.914	0.939	0.130	0.213	0.911	0.929	0.167	0.268
100	25	0.931	0.945	0.049	0.092	0.931	0.952	0.096	0.150
	35	0.931	0.947	0.064	0.114	0.932	0.951	0.106	0.173
	43	0.937	0.948	0.082	0.132	0.939	0.953	0.141	0.210
200	25	0.942	0.947	0.016	0.034	0.938	0.955	0.073	0.108
	35	0.940	0.950	0.025	0.046	0.945	0.954	0.085	0.129
	43	0.942	0.951	0.041	0.062	0.945	0.955	0.105	0.154

Note:  $n$ : number of clusters with cluster size  $M \in [30, 60]$ ;  $p_m$ : percentage of missingness; EP: equal precision bands; HW: Hall-Wellner-type bands

2010. Among these patients, 8082 were still in care, 84 died while in care (reported death), and 16207 were lost to clinic for at least two months. Among those 16207 patients who were lost to clinic, 5107 (31.5%) were intensively outreached in the community and their vital status was actively ascertained by outreach workers. Among them, 1867 (36.6%) patients were found to be deceased within two months from the last clinic visit, which indicates a substantial death under-reporting issue. The remaining 11100 lost patients who were not outreached had a missing cause of failure. Descriptive characteristics of the patients included in this analysis are presented in Table 3.

We assumed a marginal logistic model for  $\pi_l(\mathbf{W}_{ij}, \boldsymbol{\gamma}_0)$ ,  $l = 1, 2$  with time since ART initiation, sex, age, CD4 count at ART initiation, and HIV status disclosure as covariates. The goodness of fit evaluation for the parametric model  $\pi_1(\mathbf{W}_{ij}, \boldsymbol{\gamma}_0)$ , is depicted in Figure 1. The estimated residual process for death while in care fell within the 95% simultaneous confidence band under the null hypothesis (p-value = 0.862). This indicates that there is no evidence for a violation of the parametric model assumption imposed for this dataset.

The data analysis results using the proposed method and the method by Bakoyannis et al. (2020) are summarized in Table 4. Younger patients with a higher CD4 count at ART initiation had a higher hazard of disengagement from HIV care, while males and patients with a lower CD4 count at ART initiation had a higher hazard of death while in care. In contrast, the method by Bakoyannis et al. (2020) which ignores the within-cluster dependence and the informative cluster size, provided significant sex and HIV status disclosure effects on the hazard of disengagement from HIV care, and significant age and HIV status disclosure effects on the hazard of death while in care. The dubiously significant effects from the naïve analysis may be attributed to the under-estimation of standard errors, in addition to the



Table 3: Descriptive characteristics for the EA-IeDEA study sample included in the analysis

Variable	Right censoring	Cause of failure		
	In care	Disengagement	Death	Missing
	( <i>N</i> =8082)	( <i>N</i> =3240)	( <i>N</i> =1951 <sup>1</sup> )	( <i>N</i> =11100)
	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)
Gender				
Female	5334 (66.0)	2002 (61.8)	974 (49.9)	7363 (66.3)
Male	2748 (34.0)	1238 (38.2)	977 (50.1)	3737 (33.7)
HIV status disclosed				
Yes	5116 (63.3)	2022 (62.4)	1417 (72.6)	6917 (62.3)
No	2966 (36.7)	1218 (37.6)	534 (27.4)	4183 (37.7)
	Median (IQR)	Median (IQR)	Median (IQR)	Median (IQR)
Age <sup>2</sup>	38.0 (31.5, 46.1)	36.1 (29.4, 43.3)	39.1 (32.3, 48.2)	36.0 (29.3, 43.9)
CD4 <sup>3</sup>	196 (95, 297)	173 (72, 281)	66 (22, 168)	183 (82, 291)

Note: <sup>1</sup>: Included 84 reported deaths and 1867 unreported deaths confirmed through outreach; <sup>2</sup>: Age at

ART initiation in years; <sup>3</sup>: CD4 count at ART initiation in cells/ $\mu$ l



Figure 1: Plot for the cumulative residual process of the parametric model  $\pi_l(\mathbf{W}_{ij}, \gamma_0)$ ,  $l = 1, 2$  for death while in care based on the HIV data along with the 95% goodness-of-fit band (grey area) and the corresponding p-value.




Figure 2: Plot for predicted cumulative incidence functions (solid lines) of (a) disengagement from HIV care and (b) death while in care with the 95% simultaneous confidence bands based on equal precision bands (dotted lines) and Hall-Wellner bands (dashed lines), for a 40-year-old man with CD4 count of 150 cells/ $\mu$ l at ART initiation and undisclosed HIV status

bias due to the potential informative cluster size.

Figure 2 depicts the predicted cumulative incidence functions of (a) disengagement from care and (b) death while in care for a 40-year-old man with CD4 count of 150 cells/ $\mu$ l at ART initiation and undisclosed HIV status, along with the equal-precision-type and the Hall-Wellner-type bands.

## 5 Discussion

In this paper, we proposed a general framework for marginal semiparametric regression analysis of clustered competing risks data with missing cause of failure. Our approach utilizes the

Table 4: Data analysis of the EA-IeDEA study sample. Results from the proposed approach and the approach by Bakoyannis et al. (2020) (BZY20) which ignores the within-cluster dependence.

Covariates	Proposed <sup>1</sup>			BZY20 <sup>2</sup>		
	$\exp(\hat{\beta}_n)$	95% CI <sup>3</sup>	$p$ -value	$\exp(\hat{\beta}_n)$	95% CI <sup>3</sup>	$p$ -value
Disengagement from HIV care						
Sex (male = 1, female = 0)	0.94	(0.80, 1.10)	0.411	1.07	(1.01, 1.12)	0.016
Age (10 years)	0.79	(0.74, 0.84)	<0.001	0.77	(0.75, 0.79)	<0.001
CD4 (100 cells/ $\mu$ l)	1.06	(1.00, 1.12)	0.050	1.05	(1.04, 1.06)	<0.001
HIV status <sup>4</sup>	0.96	(0.84, 1.11)	0.606	0.83	(0.79, 0.87)	<0.001
Death while in care						
Sex (male = 1, female = 0)	1.40	(1.20, 1.41)	<0.001	1.30	(1.20, 1.41)	<0.001
Age (10 years)	0.98	(0.89, 1.09)	0.758	1.07	(1.03, 1.11)	0.001
CD4 (100 cells/ $\mu$ l)	0.63	(0.57, 0.70)	<0.001	0.67	(0.64, 0.71)	<0.001
HIV status <sup>4</sup>	1.15	(0.88, 1.50)	0.310	1.27	(1.16, 1.39)	<0.001

Note: <sup>1</sup>: The standard errors were estimated with cluster bootstrap; <sup>2</sup>: The standard errors were estimated with bootstrap at individual level; <sup>3</sup>: 95% confidence interval for the cause-specific hazard ratio  $\exp(\beta_0)$ ; <sup>4</sup>:

HIV status disclosed with Yes = 1, No = 0

marginal proportional cause-specific hazards model, and uses a partial pseudolikelihood approach for estimation under a MAR assumption. We provide estimators for both regression coefficients and infinite-dimensional parameters, such as the marginal cumulative incidence function. The proposed method does not impose assumptions regarding the within-cluster dependence and also accounts for informative cluster size. The proposed estimators were shown to be strongly consistent and asymptotically normal. Closed-form variance estimators were provided and rigorous methodology for the calculation of simultaneous confidence bands for the infinite-dimensional parameters was proposed. Our simulation studies showed that the performance of the method was satisfactory even with a very small number of clusters and, also, under a misspecified parametric model for the cause of failure  $\pi_1(W, \gamma_0)$ . In contrast, the previously proposed method by Bakoyannis et al. (2020), which ignores the within-cluster dependence and the informative cluster size, provided biased estimates, underestimated standard errors, and poor coverage probabilities. The analysis of the EA-IeDEA data illustrated that ignoring the within-cluster dependence and the informative cluster size may lead to dubiously significant results in practice. Last but not least, our proposed method can be easily implemented using the `coxph` function of the R package `survival`, via a simple data manipulation procedure. This is illustrated in Web Appendix A.

To the best of our knowledge, the issue of clustered competing risks data with missing cause of failure has only been addressed using frailty models by Lee et al. (2017). However, this methodology imposes strong assumptions regarding the within-cluster dependence and the distribution of the random effects, which may be violated in practice. Moreover, this approach does not account for informative cluster size, and does not provide inference about the infinite-dimensional parameters such as the cumulative incidence function. Nevertheless,

the covariate-specific cumulative incidence functions are essential for personalized risk prediction in modern medicine. Finally, the method is computationally intensive and provides cluster-specific inference, even though population-averaged inference is more scientifically relevant in many applications including our motivating EA-IeDEA study. The methodology presented in this paper effectively addresses all these limitations and is the first, to the best of our knowledge, rigorous approach for marginal analysis of clustered competing risks data with informative cluster size and missing causes of failure.

Our method adopted a parametric model for the marginal cause of failure probability  $\pi_l(\mathbf{W}_{ij}, \boldsymbol{\gamma}_0)$ ,  $l = 1, \dots, k$ , under a MAR assumption. Our simulation studies provided numerical evidence that our regression parameter estimators are robust against some degree of model misspecification. However, the confidence bands had lower coverage rate under a misspecified model for the cause of failure probability. This issue can be alleviated in practice by using flexible parametric models including regression B-splines (with fixed number of internal knots). We also proposed a goodness of fit procedure based on a cumulative residual process to evaluate the model assumption for  $\pi_l(\mathbf{W}_{ij}, \boldsymbol{\gamma}_0)$ ,  $l = 1, \dots, k$ . In our HIV data application, the use of this approach revealed that there was no evidence for a violation of the parametric model assumption.

There are many possible extensions on this work. One may consider nonparametric or semiparametric models for the marginal probability  $\pi_l(\mathbf{W}_{ij}, \boldsymbol{\gamma}_0)$ , or machine learning methods to predict the missing causes of failure. Moreover, extending the methodology for more general clustered and incomplete multi-state data (Liquet et al., 2012; Lan et al., 2017; Bakoyannis, 2020) is of interest from both practical and theoretical standpoints.

# Acknowledgements

Research reported in this publication was supported by the National Institute Of Allergy And Infectious Diseases (NIAID), Eunice Kennedy Shriver National Institute Of Child Health & Human Development (NICHD), National Institute On Drug Abuse (NIDA), National Cancer Institute (NCI), and the National Institute of Mental Health (NIMH), in accordance with the regulatory requirements of the National Institutes of Health under Award Numbers U01AI069911 and R21AI145662. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. This research has also been supported by Lilly Endowment, Inc., through its support for the Indiana University Pervasive Technology Institute, and by the President’s Emergency Plan for AIDS Relief (PEPFAR) through USAID under the terms of Cooperative Agreement No. AID-623-A-12-0001 it is made possible through joint support of the United States Agency for International Development (USAID). The contents of this journal article are the sole responsibility of AMPATH and do not necessarily reflect the views of USAID or the United States Government.

# References

- Bakoyannis, G. (2020). Nonparametric analysis of nonhomogeneous multistate processes with clustered observations. *Biometrics*, 1–14.
- Bakoyannis, G., F. Siannis, and G. Touloumi (2010). Modelling competing risks data with missing cause of failure. *Statistics in Medicine* 29, 3172–3185.

- Bakoyannis, G., Y. Zhang, and C. T. Yiannoutsos (2019). Nonparametric inference for Markov processes with missing absorbing state. *Statistica Sinica* 29, 2083–2104.
- Bakoyannis, G., Y. Zhang, and C. T. Yiannoutsos (2020). Semiparametric regression and risk prediction with competing risks data under missing cause of failure. *Lifetime Data Analysis* 26(4), 659–684.
- Balan, T. A. and H. Putter (2019). Nonproportional hazards and unobserved heterogeneity in clustered survival data: When can we tell the difference? *Statistics in medicine* 38(18), 3405–3420.
- Bordes, L., J. Y. Dauxois, and P. Joly (2014). Semiparametric inference of competing risks data with additive hazards and missing cause of failure under mcar or mar assumptions. *Electronic Journal of Statistics* 8, 41–95.
- Cai, J. and R. L. Prentice (1997). Regression estimation using multivariate failure time data and a common baseline hazard function model. *Lifetime Data Analysis* 3(3), 197–213.
- Cai, T., L. Wei, and M. Wilcox (2000). Semiparametric regression analysis for clustered failure time data. *Biometrika* 87(4), 867–878.
- Cheng, S. C., J. P. Fine, and L. J. Wei (1998). Prediction of cumulative incidence function under the proportional hazards model. *Biometrics* 54, 219–228.
- Clayton, D. and J. Cuzick (1985). Multivariate generalizations of the proportional hazards model. *Journal of the Royal Statistical Society: Series A (General)* 148(2), 82–108.



- Cong, X. J., G. Yin, and Y. Shen (2007). Marginal analysis of correlated failure time data with informative cluster sizes. *Biometrics* 63(3), 663–672.
- Craiu, R. V. and T. Duchesne (2004). Inference based on the em algorithm for the competing risks model with masked causes of failure. *Biometrika* 91, 543–558.
- Diao, G. and D. Zeng (2013). Clustered competing risks. *Handbook of Survival Analysis*, 511.
- Fine, J. P. and R. J. Gray (1999). A proportional hazards model for the subdistribution of a competing risk. *Journal of the American Statistical Association* 94, 496–509.
- Gao, G. and A. A. Tsiatis (2005). Semiparametric estimators for the regression coefficients in the linear transformation competing risks model with missing cause of failure. *Biometrika* 92, 875–891.
- Goetghebeur, E. and L. Ryan (1995). Analysis of competing risks survival data when some failure types are missing. *Biometrika* 82, 821–833.
- Hall, W. J. and J. A. Wellner (1980). Confidence bands for a survival curve from censored data. *Biometrika* 67, 133–143.
- Hougaard, P. (1986). A class of multivariate failure time distributions. *Biometrika* 73(3), 671–678.
- Hyun, S., J. Lee, and Y. Sun (2012). Proportional hazards model for competing risks data with missing cause of failure. *Journal of Statistical Planning and Inference* 142, 1767–1779.

- Katsahian, S., M. Resche-Rigon, S. Chevret, and R. Porcher (2006). Analysing multicentre competing risks data with a mixed proportional hazards model for the subdistribution. *Statistics in medicine* 25(24), 4267–4278.
- Kosorok, M. R. (2008). *Introduction to Empirical Processes and Semiparametric Inference*. New York: Springer.
- Lan, L., D. Bandyopadhyay, and S. Datta (2017). Non-parametric regression in clustered multistate current status data with informative cluster size. *Statistica Neerlandica* 71(1), 31–57.
- Lee, M., I. D. Ha, and Y. Lee (2017). Frailty modeling for clustered competing risks data with missing cause of failure. *Statistical methods in medical research* 26(1), 356–373.
- Liang, K.-Y., S. G. Self, and Y.-C. Chang (1993). Modelling marginal hazards in multivariate failure time data. *Journal of the Royal Statistical Society: Series B (Methodological)* 55(2), 441–453.
- Lin, D. Y., T. R. Fleming, and L. J. Wei (1994). Confidence bands for survival curves under the proportional hazards model. *Biometrika* 81, 73–81.
- Liquet, B., J.-F. Timsit, and V. Rondeau (2012). Investigating hospital heterogeneity with a multi-state frailty model: application to nosocomial pneumonia disease in intensive care units. *BMC medical research methodology* 12(1), 79.
- Liu, D., J. D. Kalbfleisch, and D. E. Schaubel (2011). A positive stable frailty model for clustered failure time data with covariate-dependent frailty. *Biometrics* 67(1), 8–17.

- Lu, K. and A. A. Tsiatis (2001). Multiple imputation methods for estimating regression coefficients in the competing risks model with missing cause of failure. *Biometrics* 57, 1191–1197.
- Lu, W. and Y. Liang (2008). Analysis of competing risks data with missing cause of failure under additive hazards model. *Statistica Sinica* 18, 219–234.
- Nair, V. N. (1984). Confidence bands for survival functions with censored data: a comparative study. *Technometrics* 26(3), 265–275.
- Nevo, D., R. Nishihara, S. Ogino, and M. Wang (2018). The competing risks cox model with auxiliary case covariates under weaker missing-at-random cause of failure. *Lifetime Data Analysis* 24, 425–442.
- Pan, Z. and D. Y. Lin (2005). Goodness-of-fit methods for generalized linear mixed models. *Biometrics* 61, 1000–1009.
- Pavlou, M., S. R. Seaman, and A. J. Copas (2013). An examination of a method for marginal inference when the cluster size is informative. *Statistica Sinica*, 791–808.
- Scheike, T. H., Y. Sun, M.-J. Zhang, and T. K. Jensen (2010). A semiparametric random effects model for multivariate competing risks data. *Biometrika* 97(1), 133–145.
- Seaman, S. R., M. Pavlou, and A. J. Copas (2014). Methods for observed-cluster inference when cluster size is informative: a review and clarifications. *Biometrics* 70(2), 449–456.
- Spiekerman, C. F. and D. Y. Lin (1998). Marginal regression models for multivariate failure time data. *Journal of the American Statistical Association* 93, 1164–1175.

- van der Vaart, A. W. and J. A. Wellner (1996). *Weak Convergence and Empirical Processes with Applications to Statistics*. New York: Springer-Verlag.
- Wei, L.-J., D. Y. Lin, and L. Weissfeld (1989). Regression analysis of multivariate incomplete failure time data by modeling marginal distributions. *Journal of the American statistical association* 84(408), 1065–1073.
- Williamson, J. M., H.-Y. Kim, A. Manatunga, and D. G. Addiss (2008). Modeling survival data with informative cluster size. *Statistics in medicine* 27(4), 543–555.
- Zhou, B., J. Fine, A. Latouche, and M. Labopin (2012). Competing risks regression for clustered data. *Biostatistics* 13(3), 371–383.

## A R Code

Our methodology can be easily implemented using standard software that allows for weights. In this Appendix we illustrate the use of the `coxph` function in the R package `survival`. Let `data` be a data set with the variables: cluster id `clusterid`, observed time `x`, cause of failure `c`, missingness indicator `r`, a covariate of interest `z`, an auxiliary variable `a`, and cluster size `clustersize`.

For illustration purposes, we analyze the cause specific hazard for cause 1 here. The first stage of the analysis is to estimate  $\gamma_0$  through generalized estimating equations (GEE) for logistic regression using the observations with an observed cause of failure. This can be done using the following code.

```
cause <- 1
```

```

data$include <- 1*(data$r==1 & data$c>0)/data$clustersize

data$outcome <- 1*(data$c==cause)

model <- geeglm( outcome ~ x + z + a, family = "binomial",

  data = data, id = clusterid, weight = include, corstr = "independence")

```

Then, one needs to calculate the estimated probability of the cause of failure  $\pi_1(\mathbf{W}_{ij}, \hat{\gamma}_n)$ .

```

data$yhat <- predict(model, data, type = "response")

```

The second stage of the analysis is to maximize the weighted partial pseudolikelihood. This can be implemented with the `coxph` function using the weight option and some simple data management. The data management steps are used to “remove” the weights from the risk sets for the observations with a missing cause of failure.

```

data$d <- data$r*(data$c == cause) + (1-data$r)*(data$yhat > 0)

data$weight <- data$r + (1-data$r)*data$yhat

data$weight <- data$weight + (data$weight == 0)

dt0 <- data[data$r==0, ]

dt0$weight <- 1 - dt0$weight

dt0$d <- 0

data1 <- rbind(data, dt0)

data1$weight <- data1$weight/data1$clustersize

```

Then the point estimates of the regression coefficient can be calculated using the augmented dataset `data1` as follows.

```

mod <- coxph(Surv(x, d) ~ z, weight = weight, data = data1)

```

```
beta1 <- coef(mod)
```

Standard errors of the proposed estimators can be estimated using cluster bootstrap. We plan to develop an R package to implement the proposed close-form variance estimators. The cause-specific baseline cumulative hazard function can be calculated with the `basehaz` function as follows.

```
H1 <- basehaz(mod, centered = FALSE)
```

Finally, we can get the estimated baseline cumulative incidence function using equation (1) in the main manuscript, based on the estimated regression coefficients and baseline cumulative hazard functions for all causes of failure. For example, in the case of two causes of failure, the baseline cumulative incidence function can be estimated as follows.

```
Haz1 <- H1$hazard
Haz2 <- H2$hazard
S <- exp(- Haz1 - Haz2)
S.minus <- c(1, S[1: (length(S) - 1)])
Haz1.minus <- c(0, Haz1[1: (length(Haz1) - 1)])
CIF1 <- cumsum(S.minus * (Haz1 - Haz1.minus))
```

Standard errors of the proposed estimators can again be estimated via cluster bootstrap.

## B Proofs for Theorems

The asymptotic properties of the proposed estimators are justified based on empirical process theory (Kosorok, 2008; van der Vaart and Wellner, 1996). We use the following standard

empirical process notations throughout the Appendix B. For any measurable functions  $f :$

$\mathcal{D} \mapsto \mathbb{R}$ ,

$$\mathbb{P}_n f = \frac{1}{n} \sum_{i=1}^n f(\mathbf{D}_i), \text{ and } Pf = \int_{\mathcal{D}} f dP = Ef.$$

Also, let  $K_0$  denote a generic constant that may change from place to place.

## B.1 Proof of Theorem 1

In light of condition C5, the expected partial pseudoscore function is

$$\mathbf{G}_l(\boldsymbol{\beta}) = P \left[ \sum_{j=1}^{m_0} \frac{1}{M} \int_0^\tau I(M \geq j) \{ \mathbf{Z}_j - \mathbf{E}(t, \boldsymbol{\beta}) \} d\tilde{N}_{jl}(t; \boldsymbol{\gamma}_0) \right].$$

First, we show that  $\mathbf{G}_l(\boldsymbol{\beta}_{0,l}) = \mathbf{0}$ ,

$$\begin{aligned} \mathbf{G}_l(\boldsymbol{\beta}_{0,l}) &= P \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \{ \mathbf{Z}_j - \mathbf{E}(t, \boldsymbol{\beta}_{0,l}) \} d\tilde{N}_{jl}(t; \boldsymbol{\gamma}_0) \right] \\ &= E \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \{ \mathbf{Z}_j - \mathbf{E}(t, \boldsymbol{\beta}_{0,l}) \} d\tilde{N}_{jl}(t; \boldsymbol{\gamma}_0) \right] \\ &= \int_0^\tau \sum_{j=1}^{m_0} E \left[ \frac{1}{M} I(M \geq j) \{ \mathbf{Z}_j - \mathbf{E}(t, \boldsymbol{\beta}_{0,l}) \} d\tilde{N}_{jl}(t; \boldsymbol{\gamma}_0) \right] \\ &= \int_0^\tau E \left[ \sum_{j=1}^{m_0} \frac{1}{M} I(M \geq j) \{ \mathbf{Z}_j - \mathbf{E}(t, \boldsymbol{\beta}_{0,l}) \} dE \left\{ \tilde{N}_{jl}(t; \boldsymbol{\gamma}_0) \mid \mathbf{Z}, M \right\} \right] \\ &= \int_0^\tau \left[ E \left\{ \sum_{j=1}^{m_0} \frac{1}{M} I(M \geq j) Y_j(t) \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{Z}_j) \mathbf{Z}_j d\Lambda_{0,l}(t) \right\} \right] \\ &\quad - \int_0^\tau \left[ E \left\{ \sum_{j=1}^{m_0} \frac{1}{M} I(M \geq j) Y_j(t) \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{Z}_j) \mathbf{Z}_j d\Lambda_{0,l}(t) \right\} \right] \\ &= \mathbf{0} \end{aligned}$$

Using empirical process notation, the empirical version of the pseudoscore function defined in Section 2.2 can be written as

$$\mathbf{G}_{n,l}(\boldsymbol{\beta}, \hat{\boldsymbol{\gamma}}_n) = \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \{ \mathbf{Z}_j - \mathbf{E}_n(t, \boldsymbol{\beta}) \} d\tilde{N}_{jl}(t; \hat{\boldsymbol{\gamma}}_n) \right].$$

The difference between the empirical partial pseudoscore function and expected partial pseudoscore function can be decomposed as

$$\mathbf{G}_{n,l}(\boldsymbol{\beta}, \hat{\boldsymbol{\gamma}}_n) - \mathbf{G}_l(\boldsymbol{\beta}) = \mathbf{A}_{n,l} + \mathbf{B}_{n,l} - \mathbf{C}_{n,l}(\boldsymbol{\beta}) - \mathbf{D}_{n,l}(\boldsymbol{\beta}) - \mathbf{E}_{n,l}(\boldsymbol{\beta}), \quad l = 1, \dots, k,$$

where

$$\begin{aligned} \mathbf{A}_{n,l} &= \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{Z}_j \{d\tilde{N}_{jl}(t; \hat{\boldsymbol{\gamma}}_n) - d\tilde{N}_{jl}(t; \boldsymbol{\gamma}_0)\} \right], \\ \mathbf{B}_{n,l} &= (\mathbb{P}_n - P) \left\{ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{Z}_j d\tilde{N}_{jl}(t; \boldsymbol{\gamma}_0) \right\}, \\ \mathbf{C}_{n,l}(\boldsymbol{\beta}) &= \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{E}_n(t, \boldsymbol{\beta}) \{d\tilde{N}_{jl}(t; \hat{\boldsymbol{\gamma}}_n) - d\tilde{N}_{jl}(t; \boldsymbol{\gamma}_0)\} \right], \\ \mathbf{D}_{n,l}(\boldsymbol{\beta}) &= \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \{\mathbf{E}_n(t, \boldsymbol{\beta}) - \mathbf{E}(t, \boldsymbol{\beta})\} d\tilde{N}_{jl}(t; \boldsymbol{\gamma}_0) \right], \end{aligned}$$

and

$$\mathbf{E}_{n,l}(\boldsymbol{\beta}) = (\mathbb{P}_n - P) \left\{ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{E}(t, \boldsymbol{\beta}) d\tilde{N}_{jl}(t; \boldsymbol{\gamma}_0) \right\}.$$

By conditions C3, C4 and C5,

$$\begin{aligned} \mathbf{A}_{n,l} &= \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{Z}_j \{d\tilde{N}_{jl}(t; \hat{\boldsymbol{\gamma}}_n) - d\tilde{N}_{jl}(t; \boldsymbol{\gamma}_0)\} \right] \\ &= \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{Z}_j (1 - R_j) \{\pi_l(\mathbf{W}_j, \hat{\boldsymbol{\gamma}}_n) - \pi_l(\mathbf{W}_j, \boldsymbol{\gamma}_0)\} dN_j(t) \right] \\ &\leq K_0 \|\hat{\boldsymbol{\gamma}}_n - \boldsymbol{\gamma}_0\| \mathbb{P}_n \left\{ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{Z}_j (1 - R_j) dN_j(t) \right\} \\ &\xrightarrow{as} \mathbf{0}. \end{aligned}$$

By the strong law of large numbers and condition C5,

$$\mathbf{B}_{n,l} = (\mathbb{P}_n - P) \left\{ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{Z}_j d\tilde{N}_{jl}(t; \boldsymbol{\gamma}_0) \right\} \xrightarrow{as} \mathbf{0}.$$



By conditions C2, C3 and C4,

$$\begin{aligned}
\sup_{\beta \in \mathcal{B}_l} |\mathbf{C}_{n,l}(\beta)| &= \sup_{\beta \in \mathcal{B}_l} \left| \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{E}_n(t, \beta) \{d\tilde{N}_{jl}(t; \hat{\gamma}_n) - d\tilde{N}_{jl}(t; \gamma_0)\} \right] \right| \\
&\leq \sup_{\beta \in \mathcal{B}_l} \left| \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{E}_n(t, \beta) \{\pi_l(\mathbf{W}_j, \hat{\gamma}_n) - \pi_l(\mathbf{W}_j, \gamma_0)\} dN_j(t) \right] \right| \\
&\leq K_0 \|\hat{\gamma}_n - \gamma_0\| \mathbb{P}_n \left\{ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{E}_n(t, \beta) dN_j(t) \right\} \\
&\rightarrow_{as^*} \mathbf{0}.
\end{aligned}$$

For  $\mathbf{D}_{n,l}(\beta)$ , the classes of functions  $\{M^{-1} \sum_{j=1}^{m_0} I(M \geq j) Y_j(t) \exp(\beta^T \mathbf{Z}_j), t \in [0, \tau], \beta \in \mathcal{B}_l\}$  and  $\{M^{-1} \sum_{j=1}^{m_0} I(M \geq j) Y_j(t) \exp(\beta^T \mathbf{Z}_j) \mathbf{Z}_j, t \in [0, \tau], \beta \in \mathcal{B}_l\}$  are Donsker by condition C5 and, thus, also Glivenko-Cantelli. Therefore

$$\begin{aligned}
\sup_{\beta \in \mathcal{B}_l} |\mathbf{D}_{n,l}(\beta_l)| &= \sup_{\beta \in \mathcal{B}_l} \left| \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \{\mathbf{E}_n(t, \beta_l) - \mathbf{E}(t, \beta_l)\} d\tilde{N}_{jl}(t; \gamma_0) \right] \right| \\
&\leq \sup_{t \in [0, \tau], \beta_l \in \mathcal{B}_l} \|\mathbf{E}_n(t, \beta_l) - \mathbf{E}(t, \beta_l)\| \sup_{\beta \in \mathcal{B}_l} \left| \mathbb{P}_n \left\{ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) d\tilde{N}_{ji}(t; \gamma_0) \right\} \right| \\
&\leq K_0 \sup_{t \in [0, \tau], \beta_l \in \mathcal{B}_l} \|\mathbf{E}_n(t, \beta_l) - \mathbf{E}(t, \beta_l)\| \\
&\rightarrow_{as^*} 0.
\end{aligned}$$

For  $\mathbf{E}_{n,l}(\beta_l)$ , consider the class of functions

$$\begin{aligned}
\mathcal{L}_l^{(p)} &= \left\{ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \mathbf{E}^{(p)}(t, \beta) d\tilde{N}_{jl}(t; \gamma_0), \beta \in \mathcal{B}_l \right\} \\
&= \left\{ \frac{1}{M} \sum_{j=1}^{m_0} \{R_j \Delta_{jt} + (1 - R_j) \pi_l(\mathbf{W}_j, \gamma_0)\} \int_0^\tau I(M \geq j) E^{(p)}(t, \beta) dN_j(t), \beta \in \mathcal{B}_l \right\},
\end{aligned}$$

where

$$E^{(p)}(t, \beta) = \frac{E\{\frac{1}{M} \sum_{j=1}^{m_0} I(M \geq j) Y_j(t) \exp(\beta^T \mathbf{Z}_j) Z_j^{(p)}\}}{E\{\frac{1}{M} \sum_{j=1}^{m_0} I(M \geq j) Y_j(t) \exp(\beta^T \mathbf{Z}_j)\}},$$

with  $Z_j^{(p)}$  being the  $p$ th component of  $\mathbf{Z}_j$ . For an arbitrary probability measure  $Q$ , define the norm  $\|f\|_{Q,2} = (\int f^2 dQ)^{1/2}$ . Now, for any finitely discrete probability measure  $Q$  and

$\forall \beta_1, \beta_2 \in \mathcal{B}_l$  and  $f_{\beta_1}^{(p)}, f_{\beta_2}^{(p)} \in \mathcal{L}_l^{(p)}$

$$\begin{aligned}
& \left\| f_{\beta_1}^{(p)} - f_{\beta_2}^{(p)} \right\|_{Q,2} \\
& \leq \left\| \frac{1}{M} \sum_{j=1}^{m_0} \{R_j \Delta_{jl} + (1 - R_j) \pi_l(\mathbf{W}_j, \gamma_0)\} \int_0^\tau I(M \geq j) |E^{(p)}(t, \beta_1) - E^{(p)}(t, \beta_2)| dN_j(t) \right\|_{Q,2} \\
& \leq \left\| \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau |E^{(p)}(t, \beta_1) - E^{(p)}(t, \beta_2)| dN_j(t) \right\|_{Q,2} \\
& \leq K_0 \|\beta_1 - \beta_2\|,
\end{aligned}$$

by the Lipschitz continuity of  $E^{(p)}(t, \beta)$ . Then for  $\forall \beta \in \mathcal{B}_l$  there exists a  $\beta_i, i = 1, \dots, N(\epsilon, \mathcal{B}_l, \|\cdot\|)$  such that  $\|\beta_i - \beta\| < \epsilon$ . Therefore,  $\forall f_{\beta}^{(p)} \in \mathcal{L}_l^{(p)}$  there exists a  $f_{\beta_i}^{(p)}$  such that

$$\|f_{\beta_i}^{(p)} - f_{\beta}^{(p)}\|_{Q,2} \leq K_0 \|\beta_i - \beta\| \equiv \epsilon'.$$

Therefore,  $\mathcal{L}_l^{(p)}$  can be covered by  $N(\epsilon, \mathcal{B}_l, \|\cdot\|)$   $L_2(Q)$   $\epsilon'$ -balls centered at  $f_{\beta_i}^{(p)}$ . Thus, the covering number for  $\mathcal{L}_l^{(p)}$  is  $N(\epsilon', \mathcal{L}_l^{(p)}, L_2(Q)) \leq N(\epsilon, \mathcal{B}_l, \|\cdot\|)$ , where  $\mathcal{B}_l$  is a Donsker class as a consequence of condition C2. In addition, using similar arguments to those in page 142 in Kosorok (2008), the class  $\mathcal{L}_l^{(p)}$  is pointwise measurable. Consequently,  $\mathcal{L}_l^{(p)}$  is Donsker and, thus, also Glivenko-Cantelli. Therefore,  $\sup_{\beta_l \in \mathcal{B}_l} \|\mathbf{E}_{n,l}(\beta_l)\| \rightarrow_{as^*} 0$  and, thus,  $\sup_{\beta_l \in \mathcal{B}_l} \|\mathbf{G}_{n,l}(\beta_l, \hat{\gamma}_n) - \mathbf{G}_l(\beta_l)\| \rightarrow_{as^*} 0$ . This concludes the proof that  $\|\hat{\beta}_{n,l} - \beta_{0,l}\| \rightarrow_{as^*} 0$ .

Next, we prove the uniform consistency of  $\hat{\Lambda}_1(t)$ . After some algebra we have that

$$\hat{\Lambda}_{n,1}(t) - \Lambda_{0,1}(t) = A_{n,1}^*(t) + B_{n,1}^*(t), \quad l = 1, \dots, k, \quad (4)$$

where

$$\begin{aligned}
A_{n,l}^*(t) &= \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^M (1 - R_j) \{ \pi_l(\mathbf{W}_j, \hat{\gamma}_n) - \pi_l(\mathbf{W}_j, \gamma_0) \} \int_0^t \frac{dN_j(t)}{\mathbb{P}_n \{ \frac{1}{M} \sum_{j=1}^M Y_j(t) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_j) \}} \right], \\
B_{n,l}^*(t) &= \left[ \int_0^t \frac{\mathbb{P}_n \{ \frac{1}{M} \sum_{j=1}^M d\tilde{N}_{jl}(t; \gamma_0) \}}{\mathbb{P}_n \{ \frac{1}{M} \sum_{j=1}^M Y_j(t) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_j) \}} - \int_0^t \frac{P \{ \frac{1}{M} \sum_{j=1}^M d\tilde{N}_{jl}(t; \gamma_0) \}}{P \{ \frac{1}{M} \sum_{j=1}^M Y_j(t) \exp(\beta_{0,l}^T \mathbf{Z}_j) \}} \right].
\end{aligned}$$

By conditions C3, C4 and C5,  $\|A_{n,l}^*(t)\|_\infty \rightarrow_{as^*} 0$ . By a similar expansion and arguments in Kosorok (2008),  $\|B_{n,l}^*(t)\|_\infty \rightarrow_{as^*} 0$ . Therefore,  $\|\hat{\Lambda}_{n,l}(t) - \Lambda_{0,l}(t)\|_\infty \rightarrow_{as^*} 0$ .

## B.2 Proof of Theorem 2

The estimator  $\hat{\beta}_{n,l}$  satisfies

$$\begin{aligned} \mathbf{0} &= \sqrt{n} \mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \hat{\gamma}_n) \\ &= \sqrt{n} \{ \mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \hat{\gamma}_n) - \mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \gamma_0) \} + \sqrt{n} \mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \gamma_0). \end{aligned} \quad (5)$$

The first term in the right side of (5) can be expressed as

$$\sqrt{n} \{ \mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \hat{\gamma}_n) - \mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \gamma_0) \} = \mathbf{A}'_{n,l} - \mathbf{B}'_{n,l} - \mathbf{C}'_{n,l} + \mathbf{D}'_{n,l},$$

where

$$\begin{aligned} \mathbf{A}'_{n,l} &= \sqrt{n}(\mathbb{P}_n - P) \left[ \frac{1}{M} \sum_{j=1}^{m_0} I(M \geq j) \mathbf{Z}_j N_j(\tau) (1 - R_j) \{ \pi_l(\mathbf{W}_j, \hat{\gamma}_n) - \pi_l(\mathbf{W}_j, \gamma_0) \} \right], \\ \mathbf{B}'_{n,l} &= \sqrt{n}(\mathbb{P}_n - P) \left[ \frac{1}{M} \sum_{j=1}^{m_0} I(M \geq j) (1 - R_j) \{ \pi_l(\mathbf{W}_j, \hat{\gamma}_n) - \pi_l(\mathbf{W}_j, \gamma_0) \} \int_0^\tau \mathbf{E}(t, \beta_{0,l}) dN_j(t) \right], \\ \mathbf{C}'_{n,l} &= \sqrt{n}(\mathbb{P}_n - P) \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \{ \mathbf{E}_n(t, \beta_l) - \mathbf{E}(t, \beta_{0,l}) \} d\{ \tilde{N}_{jl}(t; \hat{\gamma}_n) - \tilde{N}_{jl}(t; \gamma_0) \} \right], \end{aligned}$$

and

$$\mathbf{D}'_{n,l} = P \left[ \frac{1}{M} \sum_{j=1}^{m_0} I(M \geq j) (1 - R_j) \int_0^\tau \{ \mathbf{Z}_j - \mathbf{E}_n(t, \beta_{0,l}) \} dN_j(t) \dot{\pi}_l(\mathbf{W}_j, \gamma_0)^T \right] \times \sqrt{n}(\hat{\gamma}_n - \gamma_0).$$

By conditions C3, C5 and the continuous mapping theorem it follows that  $\mathbf{A}'_{n,l} \rightarrow_p \mathbf{0}$  and

$\mathbf{B}'_{n,l} \rightarrow_p \mathbf{0}$ . Using Lemma 4.2 of Kosorok (2008), it follows that  $\mathbf{C}'_{n,l} \rightarrow_p \mathbf{0}$ . Therefore, the

first term in the right side of (5) is  $\sqrt{n} \{ \mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \hat{\gamma}_n) - \mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \gamma_0) \} = \mathbf{A}'_{n,l} - \mathbf{B}'_{n,l} - \mathbf{C}'_{n,l} +$

$\mathbf{D}'_{n,l} = \mathbf{D}'_{n,l} + o_p(1)$ . The second term in the right side of (5) can be expressed as

$$\sqrt{n}\mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \gamma_0) = \sqrt{n}\mathbf{G}_{n,l}(\beta_{0,l}, \gamma_0) - \mathbf{H}_l(\beta_{0,l})\sqrt{n}(\hat{\beta}_{n,l} - \beta_{0,l}) + o_p(1 + \sqrt{n}\|\hat{\beta}_{n,l} - \beta_{0,l}\|).$$

By condition C6,  $\mathbf{H}_l(\beta_{0,l})$  is invertible and, thus, there exists a constant  $K_0 > 0$  such that for any  $\beta_l \in \mathcal{B}_l$ ,  $\|\mathbf{H}_l(\beta_{0,l})(\beta_l - \beta_{0,l})\| \geq K_0\|\beta_l - \beta_{0,l}\|$ . Therefore, by Taylor expansion around  $\beta_{0,l}$ ,  $\|\mathbf{G}_{n,l}(\beta_l, \gamma_0) - \mathbf{G}_l(\beta_{0,l}, \gamma_0)\| \geq K_0\|\beta_l - \beta_{0,l}\| + o_p(\|\beta_l - \beta_{0,l}\|)$ . Now,

$$\begin{aligned} & \sqrt{n}\{\mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \gamma_0) - \mathbf{G}_l(\beta_{0,l}, \gamma_0)\} \\ &= \sqrt{n}(\mathbb{P}_n - P) \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \{\mathbf{Z}_j - E(t, \beta_{0,l})\} d\tilde{N}_{jl}(t; \gamma_0) \right] \\ & \quad + o_p(1 + \sqrt{n}\|\beta_{n,l} - \beta_{0,l}\|) + o_p(1) \\ &= O_p(1) + o_p(1 + \sqrt{n}\|\hat{\beta}_{n,l} - \beta_{0,l}\|) + o_p(1). \end{aligned}$$

Consequently,  $\{K_0 + o_p(1)\}\sqrt{n}\|\hat{\beta}_{n,l} - \beta_{0,l}\| \leq O_p(1) + o_p(1 + \sqrt{n}\|\hat{\beta}_{n,l} - \beta_{0,l}\|) + o_p(1)$ , and thus  $\sqrt{n}\|\hat{\beta}_{n,l} - \beta_{0,l}\| = O_p(1)$ . This leads to the conclusion that  $\sqrt{n}\mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \gamma_0) = \sqrt{n}\mathbf{G}_{n,l}(\beta_{0,l}, \gamma_0) - \mathbf{H}_l(\beta_{0,l})\sqrt{n}(\hat{\beta}_{n,l} - \beta_{0,l}) + o_p(1)$ . Recalling that  $\tilde{M}_{jl}(t; \beta_{0,l}, \gamma_0) = \tilde{N}_{jl}(t; \gamma_0) - \int_0^t I(X_j \geq u) \exp(\beta_{0,l}^T \mathbf{Z}_j) d\Lambda_{0,l}(u)$  we have

$$\sqrt{n}\mathbf{G}_{n,l}(\beta_{0,l}, \gamma_0) = \sqrt{n}\mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \{\mathbf{Z}_j - \mathbf{E}(t, \beta_{0,l})\} d\tilde{M}_{jl}(t; \beta_{0,l}, \gamma_0) \right] + o_p(1).$$

Taking all the pieces together we have that

$$\begin{aligned} \mathbf{0} &= \sqrt{n}\mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \hat{\gamma}_n) \\ &= \sqrt{n}\{\mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \hat{\gamma}_n) - \mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \gamma_0)\} + \sqrt{n}\mathbf{G}_{n,l}(\hat{\beta}_{n,l}, \gamma_0) \\ &= P \left\{ \frac{1}{M} \sum_{j=1}^{m_0} I(M \geq j)(1 - R_j) \int_0^\tau [\mathbf{Z}_j - \mathbf{E}_n(t, \beta_{0,l})] dN_j(t) \dot{\pi}_l(\mathbf{W}_j, \gamma_0)^T \right\} \times \sqrt{n}(\hat{\gamma}_n - \gamma_0) \\ & \quad + \sqrt{n}\mathbb{P}_n \left\{ \frac{1}{M} \sum_{j=1}^{m_0} \int_0^\tau I(M \geq j) \{\mathbf{Z}_j - \mathbf{E}(t, \beta_{0,l})\} d\tilde{M}_{jl}(t; \beta_{0,l}, \gamma_0) \right\} \\ & \quad - \mathbf{H}_l(\beta_{0,l})\sqrt{n}(\hat{\beta}_{n,l} - \beta_{0,l}) + o_p(1). \end{aligned}$$

Rearranging the terms and according to conditions C4 and C6 leads to

$$\sqrt{n}(\hat{\beta}_{n,l} - \beta_{0,l}) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} (\psi_{ijl} + \mathbf{R}_l \omega_{ij}) \right\} + o_p(1),$$

where

$$\psi_{ijl} = \mathbf{H}_l^{-1}(\beta_{0,l}) \int_0^\tau \{ \mathbf{Z}_{ij} - \mathbf{E}(t, \beta_{0,l}) \} d\tilde{M}_{ijl}(t; \beta_{0,l}, \gamma_0),$$

and

$$\mathbf{R}_l = \mathbf{H}_l^{-1}(\beta_{0,l}) E \left[ \frac{1}{M} \sum_{j=1}^M (1 - R_j) \int_0^\tau \{ \mathbf{Z}_j - \mathbf{E}(t, \beta_{0,l}) \} dN_j(t) \dot{\pi}_l(\mathbf{W}_j, \gamma_0)^T \right].$$

### B.3 Proof of Theorem 3

By Taylor expansion and the consistency of  $\hat{\beta}_{n,l}$  and  $\hat{\gamma}_n$ , the first term in the right side of expansion (4) can be written as

$$\begin{aligned} A_{n,l}^*(t) &= \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^M (1 - R_j) \{ \pi_l(\mathbf{W}_j, \hat{\gamma}_n) - \pi_l(\mathbf{W}_j, \gamma_0) \} \int_0^t \frac{dN_j(s)}{\mathbb{P}_n \{ \frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_j) \}} \right] \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} \mathbf{R}_l^*(t) \omega_{ij} \right\} + o_p(n^{-1/2}), \end{aligned}$$

where

$$\mathbf{R}_l^*(t) = E \left[ \frac{1}{M} \sum_{j=1}^M (1 - R_j) \dot{\pi}_j(\mathbf{W}_j, \gamma_0) \int_0^t \frac{dN_j(s)}{E \{ \frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\beta_{0,l}^T \mathbf{Z}_j) \}} \right]^T.$$

By similar analysis to that provided in Page 57 of Kosorok (2008), the second term in (4)

can be written as

$$\begin{aligned} B_{n,l}^*(t) &= \left[ \int_0^t \frac{\mathbb{P}_n \{ \frac{1}{M} \sum_{j=1}^M d\tilde{N}_{jl}(t; \gamma_0) \}}{\mathbb{P}_n \{ \frac{1}{M} \sum_{j=1}^M Y_j(t) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_j) \}} - \int_0^t \frac{P \{ \frac{1}{M} \sum_{j=1}^M d\tilde{N}_{jl}(t; \gamma_0) \}}{P \{ \frac{1}{M} \sum_{j=1}^M Y_j(t) \exp(\beta_{0,l}^T \mathbf{Z}_j) \}} \right] \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} \phi_{ijl}(t) \right\} + o_p(n^{-1/2}), \end{aligned}$$

where

$$\phi_{ijl}(t) = \int_0^t \frac{d\tilde{M}_{ijl}(s; \beta_{0,j}, \gamma_0)}{E\{\frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\beta_{0,l}^T \mathbf{Z}_j)\}} - (\boldsymbol{\psi}_{ijl} + \mathbf{R}_l \boldsymbol{\omega}_{ij})^T \int_0^t E(s, \beta_{0,l}) d\Lambda_{0,l}(s).$$

Therefore,

$$\sqrt{n} \left\{ \hat{\Lambda}_{n,l}(t) - \Lambda_{0,l}(t) \right\} = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left[ \frac{1}{M_i} \sum_{j=1}^{M_i} \{ \phi_{ijl}(t) + \mathbf{R}_l^*(t) \boldsymbol{\omega}_{ij} \} \right] + o_p(1).$$

By conditions C1, C4, and C5, and lemmas 1 and 2 in the supporting information of Bakoyannis (2020), the class of functions

$$\left[ \frac{1}{M} \sum_{j=1}^M \{ \phi_{jl}(t) + \mathbf{R}_l^*(t) \boldsymbol{\omega}_j \} : t \in [0, \tau] \right]$$

is Donsker. We now show that conditional on the data, the estimated multiplier process

$$\hat{W}_{n,l}(\cdot) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left[ \frac{1}{M_i} \sum_{j=1}^{M_i} \{ \hat{\phi}_{ijl}(\cdot) + \hat{\mathbf{R}}_l^*(\cdot) \hat{\boldsymbol{\omega}}_{ij} \} \right] \xi_i$$

converges weakly to the same limiting process as  $W_{n,l}(\cdot) = \sqrt{n} \{ \hat{\Lambda}_{n,l}(\cdot) - \Lambda_{0,l}(\cdot) \}$ . Define

$$\tilde{W}_{n,l}(\cdot) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left[ \frac{1}{M_i} \sum_{j=1}^{M_i} \{ \phi_{ijl}(\cdot) + \mathbf{R}_l^*(\cdot) \boldsymbol{\omega}_{ij} \} \right] \xi_i.$$

By the Donsker property of the class of influence functions and the conditional multiplier central limit theorem (van der Vaart and Wellner, 1996),  $\tilde{W}_{n,l}(\cdot)$  converges weakly, conditionally on the data, to the same limiting process as  $W_{n,l}(\cdot)$ . To complete the proof, we need to show

$$\|\hat{W}_{n,l}(t) - \tilde{W}_{n,l}(t)\|_\infty = o_p(1), \quad l = 1, \dots, k,$$

unconditionally. After some algebra

$$\|\hat{W}_{n,l}(t) - \tilde{W}_{n,l}(t)\|_\infty \leq A_{n,l}'' + B_{n,l}'' + C_{n,l}'',$$

where

$$A''_{n,l} = \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \left[ \frac{1}{M_i} \sum_{j=1}^{M_i} \left\{ \hat{\phi}_{ijl}(t) - \phi_{ijl}(t) \right\} \right] \xi_i \right\|_{\infty},$$

$$B''_{n,l} = \sup_{t \in [0, \tau]} \left\| \hat{\mathbf{R}}_l^*(t) - \mathbf{R}_l^*(t) \right\| \times \left( \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} (\hat{\omega}_{ij} - \omega_{ij}) \right\} \xi_i \right\| + \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \left( \frac{1}{M_i} \sum_{j=1}^{M_i} \omega_{ij} \right) \xi_i \right\| \right),$$

and

$$C''_{n,l} = \sup_{t \in [0, \tau]} \left\| \mathbf{R}_l^*(t) \right\| \times \left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} (\hat{\omega}_{ij} - \omega_{ij}) \right\} \xi_i \right\|.$$

Using the same arguments to those used in the proof of Theorem 4 in Spiekerman and Lin (1998) and regularity conditions C3 and C4,  $A''_{n,l} = o_p(1)$ .

For  $B''_{n,l}$ , using the same arguments to those used in the proof of Lemma A.3 in Spiekerman and Lin (1998) leads to the conclusion that

$$\left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} (\hat{\omega}_{ij} - \omega_{ij}) \right\} \xi_i \right\| = o_p(1).$$

Next, by condition C4 and the central limit theorem

$$\left\| \frac{1}{\sqrt{n}} \sum_{i=1}^n \left( \frac{1}{M_i} \sum_{j=1}^{M_i} \omega_{ij} \right) \xi_i \right\| = O_p(1).$$

Also, after some algebra, we have that

$$\sup_{t \in [0, \tau]} \left\| \hat{\mathbf{R}}_l^*(t) - \mathbf{R}_l^*(t) \right\| \leq A'''_{n,l} + B'''_{n,l} + C'''_{n,l},$$

where

$$A'''_{n,l} = \sup_{t \in [0, \tau]} \left\| \mathbb{P}_n \left[ \frac{1}{M} \sum_{j=1}^M \left\{ \hat{\pi}_l(\mathbf{W}_j, \hat{\gamma}_n) - \hat{\pi}_l(\mathbf{W}_j, \gamma_0) \right\} \int_0^t \frac{dN_j(s)}{\mathbb{P}_n \left\{ \frac{1}{M} \sum_{j=1}^M Y_j(s) \exp \left( \hat{\beta}_{n,l}^T \mathbf{Z}_j \right) \right\}} \right] \right\|,$$

$$B'''_{n,l} = \sup_{t \in [0, \tau]} \left\| \mathbb{P}_n \frac{1}{M} \sum_{j=1}^M \hat{\pi}_l(\mathbf{W}_j, \gamma_0) \int_0^t \left[ \frac{1}{\mathbb{P}_n \left\{ \frac{1}{M} \sum_{j=1}^M Y_j(s) e^{\hat{\beta}_{n,l}^T \mathbf{Z}_j} \right\}} - \frac{1}{P \left\{ \frac{1}{M} \sum_{j=1}^M Y_j(s) e^{\beta_{0,l}^T \mathbf{Z}_j} \right\}} \right] dN_j(s) \right\|,$$

and

$$C_{n,l}''' = \sup_{t \in [0, \tau]} \left\| (\mathbb{P}_n - P) \left[ \frac{1}{M} \sum_{j=1}^M \dot{\pi}_l(\mathbf{W}_j, \gamma_0) \int_0^t \frac{1}{P \left\{ \frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\beta_{0,l}^T \mathbf{Z}_j) \right\}} dN_j(s) \right] \right\|.$$

By conditions C3, C4, C5, and the continuous mapping theorem,

$$\max_{ij} \|\dot{\pi}_l(\mathbf{W}_{ij}, \hat{\gamma}_n) - \dot{\pi}_l(\mathbf{W}_{ij}, \gamma_0)\| = o_{as}(1).$$

By Theorem 2 and conditions C1, C2, and C5,

$$\left\| \frac{1}{\mathbb{P}_n \left\{ \frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_j) \right\}} \right\|_{\infty} = \left\| \frac{1}{P \left\{ \frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\beta_{0,l}^T \mathbf{Z}_j) \right\} + o_{as^*}(1)} \right\| = O_{as^*}(1).$$

Therefore,  $A_{n,l}''' = o_{as^*}(1)$ . Next, by conditions C3 and C5,  $\max_{ij} \|\dot{\pi}_l(\mathbf{W}_{ij}, \gamma_0)\| = O_{as}(1)$ .

Also, by conditions C2, C5, and the Donsker property of  $\{Y_j(t) : t \in [0, \tau]\}$ ,

$$\left\| \frac{1}{\mathbb{P}_n \left\{ \frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_j) \right\}} - \frac{1}{P \left\{ \frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\beta_{0,l}^T \mathbf{Z}_j) \right\}} \right\|_{\infty} = o_{as^*}(1),$$

and, thus,  $B_{n,l}''' = o_{as^*}(1)$ . For  $C_{n,l}'''$ , consider the classes of functions  $\mathcal{F} = \{N_j(s) : t \in [0, \tau]\}$

and

$$\mathcal{L}_{l,1} = \left\{ f_{t,l} = \frac{1}{M} \sum_{j=1}^{m_0} I(M \geq j) \dot{\pi}_l(\mathbf{W}_j, \gamma_0) \int_0^t \frac{1}{P \left\{ \frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\beta_{0,l}^T \mathbf{Z}_j) \right\}} dN_j(s), t \in [0, \tau] \right\}.$$

For any finitely discrete probability measure  $Q$  and any  $t_1, t_2 \in [0, \tau]$  we have that

$$\begin{aligned} \|f_{t_1,l} - f_{t_2,l}\|_{Q,2} &\leq \left\| \frac{1}{M} \sum_{j=1}^{m_0} I(M \geq j) \dot{\pi}_l(\mathbf{W}_j, \gamma_0) \int_{t_1}^{t_2} \frac{dN_j(s)}{P \left\{ \frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\beta_{0,l}^T \mathbf{Z}_j) \right\}} \right\|_{Q,2} \\ &\leq K_0 \|N_j(t_2) - N_j(t_1)\|_{Q,2}. \end{aligned}$$

Therefore,  $\mathcal{L}_{l,1}$  can be covered by  $N(\epsilon, \mathcal{F}, L_2(Q))$   $L_2(Q)$   $\epsilon'$ -balls centered at  $f_{t_i,l}$ , where  $\mathcal{F}$  is a Donsker class by lemma 4.1 in Kosorok (2008). In addition, using similar arguments to those in page 142 in Kosorok (2008), the class  $\mathcal{L}_{l,1}$  is pointwise measurable. Consequently,



the class  $\mathcal{L}_{l,1}$  is Donsker and, thus, also Glivenko-Cantelli, which leads to the conclusion that  $C'''_{n,l} = o_{as^*}(1)$ . Therefore,  $\sup_{t \in [0, \tau]} \|\hat{\mathbf{R}}_l^*(t) - \mathbf{R}_l^*(t)\| = o_p(1)$  and, thus,  $B''_{n,l} = o_p(1)$ . Similar arguments lead to the conclusion that  $C'''_{n,l} = o_p(1)$ . Thus,

$$\left\| \hat{W}_{n,l}(t) - \tilde{W}_{n,l}(t) \right\|_{\infty} = o_p(1),$$

which completes the proof of the last statement in Theorem 3.

## B.4 Proof of Theorem 4

It is easy to show that

$$\begin{aligned} \tilde{W}_{n,l}(t; \mathbf{z}_0) &= \sqrt{n} \{ \hat{\Lambda}_{n,l}(t; \mathbf{z}_0) - \Lambda_{0,l}(t; \mathbf{z}_0) \} \\ &= \sqrt{n} \{ \hat{\Lambda}_{n,l}(t) \exp(\hat{\boldsymbol{\beta}}_{n,l}^T \mathbf{z}_0) - \Lambda_{0,l}(t) \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{z}_0) \} \\ &= \sqrt{n} \{ \hat{\Lambda}_{n,l}(t) - \Lambda_{0,l}(t) + \mathbf{z}_0^T (\hat{\boldsymbol{\beta}}_{n,l} - \boldsymbol{\beta}_{0,l}) \Lambda_{0,l}(t) \} \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{z}_0) + o_p(1) \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{1}{M_i} \sum_{j=1}^{M_i} \{ \mathbf{z}_0^T (\boldsymbol{\psi}_{ijl} + \mathbf{R}_l \boldsymbol{\omega}_{ij}) \Lambda_{0,l}(t) + \phi_{ijl}(t) + \mathbf{R}_l^*(t) \boldsymbol{\omega}_{ij} \} \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{z}_0) + o_p(1) \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{1}{M_i} \sum_{j=1}^{M_i} \phi_{ijl}^{\Lambda}(t; \mathbf{z}_0) + o_p(1). \end{aligned}$$

Similarly to the decomposition in Cheng et al. (1998),

$$\sqrt{n} \left\{ \hat{F}_{n,l}(t; \mathbf{z}_0) - F_{0,l}(t; \mathbf{z}_0) \right\} = \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} \phi_{ijl}^F(t; \mathbf{z}_0) \right\} + o_p(1),$$

where

$$\begin{aligned} \phi_{ijl}^F(t; \mathbf{z}_0) &= \int_0^t \exp \left\{ - \sum_{l=1}^k \Lambda_{0,l}(s-; \mathbf{z}_0) \right\} d\phi_{ijl}^{\Lambda}(s; \mathbf{z}_0) \\ &\quad - \int_0^t \left\{ \sum_{l=1}^k \phi_{ijl}^{\Lambda}(s-; \mathbf{z}_0) \right\} \exp \left\{ - \sum_{l=1}^k \Lambda_{0,l}(s-; \mathbf{z}_0) \right\} d\Lambda_{0,l}(s; \mathbf{z}_0), \end{aligned}$$

and  $\phi_{ijl}^\Lambda(t; \mathbf{z}_0) = \{\mathbf{z}_0^T(\boldsymbol{\psi}_{ijl} + \mathbf{R}_l \boldsymbol{\omega}_{ij})\Lambda_{0,l}(t) + \phi_{ijl}(t) + \mathbf{R}_l^*(t)\boldsymbol{\omega}_{ij}\} \exp(\boldsymbol{\beta}_{0,l}^T \mathbf{z}_0)$ . The class of functions  $\{\phi_{jl}^F(t; \mathbf{z}_0 : t \in [0, \tau])\}$  is Donsker by conditions C1, C4, and C5, lemmas 1 and 2 in the supporting information of Bakoyannis (2020), and corollary 9.32 in Kosorok (2008).

To conclude the proof of Theorem 4, we show that, conditionally on the data,

$$\hat{W}_{n,l}^F(\cdot; \mathbf{z}_0) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{1}{M_i} \sum_{j=1}^{M_i} \hat{\phi}_{ijl}^F(\cdot; \mathbf{z}_0) \xi_i$$

converges weakly to the same limiting process as  $W_{n,l}^F(\cdot; \mathbf{z}_0) = \sqrt{n}\{\hat{F}_{n,l}(\cdot; \mathbf{z}_0) - F_{0,l}(\cdot; \mathbf{z}_0)\}$ .

Now, define

$$\tilde{W}_{n,l}^F(\cdot; \mathbf{z}_0) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{1}{M_i} \sum_{j=1}^{M_i} \phi_{ijl}^F(\cdot; \mathbf{z}_0) \xi_i.$$

By the Donsker property of the class of influence functions and the conditional multiplier central limit theorem (van der Vaart and Wellner, 1996),  $\tilde{W}_{n,l}^F(\cdot; \mathbf{z}_0)$  converges weakly, conditionally on the data, to the same limiting process as  $W_{n,l}^F(\cdot; \mathbf{z}_0)$ . To complete the proof, we need to show that

$$\|\hat{W}_{n,l}^F(t; \mathbf{z}_0) - \tilde{W}_{n,l}^F(t; \mathbf{z}_0)\|_\infty = o_p(1), \quad l = 1, \dots, k,$$

unconditionally. Some algebra leads to the following bound

$$\|\hat{W}_{n,l}^F(t; \mathbf{z}_0) - \tilde{W}_{n,l}^F(t; \mathbf{z}_0)\|_\infty \leq A_{n,l}'''' + B_{n,l}'''' + C_{n,l}'''' + D_{n,l}'''' + E_{n,l}'''' + F_{n,l}'''' ,$$

where

$$A_{n,l}'''' = \left\| \int_0^t \exp \left\{ - \sum_{l=1}^k \hat{\Lambda}_{n,l}(s-; \mathbf{z}_0) \right\} d \left[ \sqrt{n} \mathbb{P}_n \frac{1}{M} \sum_{j=1}^M \left\{ \hat{\phi}_{jl}^A(s; \mathbf{z}_0) - \phi_{jl}^A(s; \mathbf{z}_0) \right\} \xi \right] \right\|_\infty ,$$

$$B_{n,l}'''' = \left\| \int_0^t \exp \left\{ - \sum_{l=1}^k \hat{\Lambda}_{n,l}(s-; \mathbf{z}_0) \right\} - \exp \left\{ - \sum_{l=1}^k \Lambda_{0,l}(s-; \mathbf{z}_0) \right\} d \left\{ \sqrt{n} \mathbb{P}_n \frac{1}{M} \sum_{j=1}^M \phi_{jl}^A(s; \mathbf{z}_0) \xi \right\} \right\|_\infty ,$$

$$\begin{aligned}
C_{n,j}''' &= \left\| \sum_{l=1}^k \int_0^t \left[ \mathbb{P}_n \frac{1}{M} \sum_{j=1}^M \left\{ \hat{\phi}_{jl}^A(s-; \mathbf{z}_0) - \phi_{jl}^A(s-; \mathbf{z}_0) \right\} \xi \right] \exp \left\{ - \sum_{l=1}^k \hat{\Lambda}_{n,l}(s-; \mathbf{z}_0) \right\} \right. \\
&\quad \left. \times d \left[ \sqrt{n} \left\{ \hat{\Lambda}_{n,l}(s; \mathbf{z}_0) - \Lambda_{0,l}(s; \mathbf{z}_0) \right\} \right] \right\|_{\infty}, \\
D_{n,j}''' &= \left\| \sum_{l=1}^k \int_0^t \left\{ \mathbb{P}_n \frac{1}{M} \sum_{j=1}^M \phi_{jl}^A(s-; \mathbf{z}_0) \xi \right\} \left[ \exp \left\{ - \sum_{l=1}^k \hat{\Lambda}_{n,l}(s-; \mathbf{z}_0) \right\} - \exp \left\{ - \sum_{l=1}^k \Lambda_{0,l}(s-; \mathbf{z}_0) \right\} \right] \right. \\
&\quad \left. \times d \left[ \sqrt{n} \left\{ \hat{\Lambda}_{n,l}(s; \mathbf{z}_0) - \Lambda_{0,l}(s; \mathbf{z}_0) \right\} \right] \right\|_{\infty}, \\
E_{n,l}''' &= \left\| \sum_{l=1}^k \int_0^t \left[ \sqrt{n} \mathbb{P}_n \frac{1}{M} \sum_{j=1}^M \left\{ \hat{\phi}_{jl}^A(s-; \mathbf{z}_0) - \phi_{jl}^A(s-; \mathbf{z}_0) \right\} \xi \right] d\Lambda_{0,l}(s; \mathbf{z}_0) \right\|_{\infty}, \\
F_{n,l}''' &= O_p(1) \left| \sqrt{n} \mathbb{P}_n \xi \right| \left\| \int_0^t \left[ \exp \left\{ - \sum_{l=1}^k \hat{\Lambda}_{n,l}(s-; \mathbf{z}_0) \right\} - \exp \left\{ - \sum_{l=1}^k \Lambda_{0,l}(s-; \mathbf{z}_0) \right\} \right] d\Lambda_{0,l}(s; \mathbf{z}_0) \right\|_{\infty}.
\end{aligned}$$

By integration by parts, Theorem 3, Lemma A.3 in Spiekerman and Lin (1998) and the boundedness conditions,  $A_{n,l}''' = o_p(1)$ . By Theorem 1, the Donsker property of the class  $\{\phi_{jl}^{\Lambda}(t; \mathbf{z}_0) : t \in [0, \tau]\}$ , and arguments similar to those used in the proof of proposition 7.27 in Kosorok (2008),  $B_{n,l}''' = o_p(1)$ . For  $C_{n,l}'''$ , the integrand converges uniformly to 0 in probability and, thus, by Theorem 3 and arguments similar to those used in the proof of proposition 7.27 in Kosorok (2008), it follows that  $C_{n,l}''' = o_p(1)$ . Using similar arguments, it can be shown that  $D_{n,l}''' = o_p(1)$ . For  $E_{n,l}'''$ , the integrand converges uniformly to 0 in probability and, thus, by condition C1 it follows that  $E_{n,l}''' = o_p(1)$ . Finally, by Theorem 1, condition C1, and the central limit theorem, it follows that  $F_{n,l}''' = o_p(1)$ . Therefore,

$$\left\| \hat{W}_{n,l}^F(t; \mathbf{z}_0) - \tilde{W}_{n,l}^F(t; \mathbf{z}_0) \right\|_{\infty} = o_p(1),$$

for all  $l = 1, \dots, k$ , which completes the proof of the last statement in Theorem 4.

## B.5 Standard Error Estimators

The covariance matrix  $\Sigma_l$  can be consistently estimated using the empirical versions of the influence functions by

$$\hat{\Sigma}_l = \frac{1}{n} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} (\hat{\psi}_{ijl} + \hat{\mathbf{R}}_l \hat{\omega}_{ij}) \right\}^{\otimes 2}.$$

The first component is

$$\hat{\psi}_{ijl} = \mathbf{H}_{n,l}^{-1}(\hat{\beta}_{n,l}) \int_0^\tau \{ \mathbf{Z}_{ij} - \mathbf{E}_n(t, \hat{\beta}_{n,l}) \} d\hat{M}_{ijl}(t; \hat{\beta}_{n,l}, \hat{\gamma}_n),$$

where

$$\begin{aligned} \mathbf{H}_{n,l}(\hat{\beta}_{n,l}; \hat{\gamma}_n) &= \frac{1}{n} \sum_{i=1}^n \frac{1}{M_i} \sum_{j=1}^{M_i} \int_0^\tau \mathbf{V}_{n,l}(t, \hat{\beta}_{n,l}) d\tilde{N}_{ijl}(t; \hat{\gamma}_n), \\ \mathbf{V}_{n,l}(t, \hat{\beta}_{n,l}) &= \frac{\sum_{p=1}^n \frac{1}{M_p} \sum_{q=1}^{M_p} Y_{pq}(t) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_{pq}) \mathbf{Z}_{pq}^{\otimes 2}}{\sum_{p=1}^n \frac{1}{M_p} \sum_{q=1}^{M_p} Y_{pq}(t) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_{pq})} - \left\{ \frac{\sum_{p=1}^n \frac{1}{M_p} \sum_{q=1}^{M_p} Y_{pq}(t) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_{pq}) \mathbf{Z}_{pq}}{\sum_{p=1}^n \frac{1}{M_p} \sum_{q=1}^{M_p} Y_{pq}(t) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_{pq})} \right\}^{\otimes 2}, \\ \mathbf{E}_n(t, \hat{\beta}_{n,l}) &= \frac{\sum_{p=1}^n \frac{1}{M_p} \sum_{q=1}^{M_p} Y_{pq}(t) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_{pq}) \mathbf{Z}_{pq}}{\sum_{p=1}^n \frac{1}{M_p} \sum_{q=1}^{M_p} Y_{pq}(t) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_{pq})}, \end{aligned}$$

and

$$\hat{M}_{ijl}(t; \hat{\beta}_{n,l}, \hat{\gamma}_n) = \tilde{N}_{ijl}(t; \hat{\gamma}_n) - \int_0^t Y_{ij}(u) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_{ij}) d\hat{\Lambda}_{n,l}(u).$$

The second component is

$$\hat{\mathbf{R}}_l = \mathbf{H}_{n,l}^{-1}(\hat{\beta}_{n,l}) \frac{1}{n} \sum_{i=1}^n \left[ \frac{1}{M_i} \sum_{j=1}^{M_i} (1 - R_{ij}) \int_0^\tau \{ \mathbf{Z}_{ij} - \mathbf{E}_n(t, \hat{\beta}_{n,l}) \} dN_{ij}(t) \dot{\pi}_l(\mathbf{W}_{ij}, \hat{\gamma}_n)^T \right].$$

A consistent estimator for the covariance function for the baseline cumulative cause-specific hazard is

$$\frac{1}{n} \sum_{i=1}^n \left[ \frac{1}{M_i} \sum_{j=1}^{M_i} \left\{ \hat{\phi}_{ijl}(t) + \hat{\mathbf{R}}_l^*(t) \hat{\omega}_{ij} \right\} \right] \left[ \frac{1}{M_i} \sum_{j=1}^{M_i} \left\{ \hat{\phi}_{ijl}(s) + \hat{\mathbf{R}}_l^*(s) \hat{\omega}_{ij} \right\} \right], \quad t, s \in [0, \tau],$$

where

$$\hat{\phi}_{ijl}(t) = \int_0^t \frac{d\hat{M}_{ijl}(t; \hat{\beta}_{n,l}, \hat{\gamma}_n)}{\frac{1}{n} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} Y_{ij}(s) \exp(\hat{\beta}_{n,l}^T \mathbf{Z}_{ij}) \right\}} - (\hat{\psi}_{ijl} + \hat{\mathbf{R}}_l \hat{\omega}_{ij})^T \int_0^t \mathbf{E}_n(s, \hat{\beta}_{n,l}) d\hat{\Lambda}_{n,l}(s),$$

and

$$\hat{\mathbf{R}}_l^*(t) = E \left[ \frac{1}{M} \sum_{j=1}^M (1 - R_j) \dot{\pi}_j(\mathbf{W}_j, \gamma_0) \int_0^t \frac{dN_j(s)}{E \left\{ \frac{1}{M} \sum_{j=1}^M Y_j(s) \exp(\beta_{0,l}^T \mathbf{Z}_j) \right\}} \right]^T.$$

A consistent estimator for the covariance function for the covariate-specific cumulative incidence function is

$$\frac{1}{n} \sum_{i=1}^n \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} \hat{\phi}_{ijl}^F(t; \mathbf{z}_0) \right\} \left\{ \frac{1}{M_i} \sum_{j=1}^{M_i} \hat{\phi}_{ijl}^F(s; \mathbf{z}_0) \right\}, \quad t, s \in [0, \tau],$$

where

$$\begin{aligned} \hat{\phi}_{ijl}^F(t; \mathbf{z}_0) &= \int_0^t \exp \left\{ - \sum_{l=1}^k \hat{\Lambda}_{n,l}(s-; \mathbf{z}_0) \right\} d\hat{\phi}_{ijl}^\Lambda(s; \mathbf{z}_0) \\ &\quad - \int_0^t \left\{ \sum_{l=1}^k \hat{\phi}_{ijl}^\Lambda(s-; \mathbf{z}_0) \right\} \exp \left\{ - \sum_{l=1}^k \hat{\Lambda}_{n,l}(s-; \mathbf{z}_0) \right\} d\hat{\Lambda}_{n,l}(s; \mathbf{z}_0), \end{aligned}$$

and

$$\hat{\phi}_{ijl}^\Lambda(t; \mathbf{z}_0) = \{ \mathbf{z}_0^T (\hat{\psi}_{ijl} + \hat{\mathbf{R}}_l \hat{\omega}_{ij}) \hat{\Lambda}_{n,l}(t) + \hat{\phi}_{ijl}(t) + \hat{\mathbf{R}}_l^*(t) \hat{\omega}_{ij} \} \exp(\hat{\beta}_{n,l}^T \mathbf{z}_0).$$

## C Additional Simulation Results

The simulation results for the pointwise estimates of the infinite-dimensional parameters  $\Lambda_{0,1}(t)$  and  $F_{0,1}(t)$  under scenario 1 are provided in Table 5 and 6. Simulation results under scenario 2 are presented in Tables 7, 8, 9 and 10.

Table 5: Simulation results for the infinite-dimensional parameters  $\Lambda_{0,1}(t)$  at selected time points under Scenario 1. Results from the proposed approach and the approach by Bakoyannis et al. (2020) (BZY20) which ignores the within-cluster dependence.

$n$	$p_m(\%)$	$t$	Proposed				BZY20			
			Bias	MCSD	ASE	CP	Bias	MCSD	ASE	CP
50	25	0.1	0.001	0.084	0.080	0.915	0.090	0.104	0.025	0.252
		0.2	0.000	0.105	0.101	0.917	0.120	0.127	0.034	0.259
		0.4	0.004	0.135	0.130	0.934	0.160	0.156	0.049	0.284
		0.8	0.011	0.185	0.176	0.947	0.203	0.201	0.081	0.385
	43	0.1	0.000	0.086	0.081	0.915	0.091	0.106	0.031	0.306
		0.2	-0.001	0.108	0.104	0.926	0.121	0.130	0.042	0.318
		0.4	0.003	0.139	0.134	0.935	0.160	0.160	0.058	0.355
		0.8	0.011	0.188	0.181	0.942	0.203	0.203	0.093	0.441
200	25	0.1	0.001	0.040	0.040	0.954	0.090	0.049	0.013	0.074
		0.2	0.001	0.051	0.051	0.951	0.122	0.062	0.017	0.055
		0.4	0.003	0.066	0.066	0.955	0.159	0.077	0.024	0.055
		0.8	0.006	0.093	0.088	0.950	0.200	0.101	0.040	0.096
	43	0.1	0.001	0.040	0.041	0.952	0.091	0.051	0.015	0.096
		0.2	0.002	0.053	0.052	0.952	0.123	0.064	0.021	0.081
		0.4	0.003	0.069	0.068	0.952	0.161	0.079	0.029	0.077
		0.8	0.007	0.096	0.091	0.947	0.199	0.103	0.046	0.138

Note:  $n$ : number of clusters with cluster size  $M \in [30, 60]$ ;  $p_m$ : percentage of missingness;  $t$ : selected time points ; MCSD: Monte Carlo standard deviation; ASE: average estimated standard error; CP: coverage

Table 6: Simulation results for the infinite-dimensional parameters  $F_{0,1}(t)$  at selected time points under Scenario 1. Results from the proposed approach and the approach by Bakoyannis et al. (2020) (BZY20) which ignores the within-cluster dependence.

$n$	$p_m(\%)$	$t$	Proposed				BZY20			
			Bias	MCSD	ASE	CP	Bias	MCSD	ASE	CP
50	25	0.1	-0.002	0.049	0.047	0.924	0.036	0.055	0.014	0.300
		0.2	-0.003	0.051	0.049	0.926	0.033	0.057	0.015	0.343
		0.4	-0.002	0.053	0.051	0.933	0.025	0.058	0.017	0.388
		0.8	-0.002	0.053	0.051	0.938	0.015	0.058	0.018	0.429
	43	0.1	-0.002	0.049	0.047	0.924	0.036	0.056	0.018	0.381
		0.2	-0.003	0.053	0.051	0.924	0.033	0.059	0.020	0.427
		0.4	-0.002	0.055	0.053	0.934	0.026	0.060	0.023	0.493
		0.8	-0.002	0.056	0.054	0.936	0.015	0.060	0.024	0.555
200	25	0.1	0.000	0.023	0.024	0.954	0.038	0.027	0.007	0.177
		0.2	0.000	0.025	0.025	0.950	0.035	0.028	0.008	0.198
		0.4	0.000	0.026	0.026	0.952	0.028	0.029	0.008	0.287
		0.8	0.000	0.027	0.026	0.945	0.016	0.029	0.009	0.402
	43	0.1	0.000	0.024	0.024	0.952	0.038	0.028	0.009	0.204
		0.2	0.000	0.026	0.026	0.953	0.036	0.029	0.010	0.276
		0.4	0.000	0.028	0.027	0.949	0.028	0.030	0.011	0.368
		0.8	0.000	0.028	0.027	0.945	0.017	0.030	0.012	0.503

Note:  $n$ : number of clusters with cluster size  $M \in [30, 60]$ ;  $p_m$ : percentage of missingness;  $t$ : selected time points ; MCSD: Monte Carlo standard deviation; ASE: average estimated standard error; CP: coverage probability of 95% pointwise confidence interval

Table 7: Simulation results for the regression coefficient  $\beta_1$  under Scenario 2 for the proposed approach and the approach by Bakoyannis et al. (2020) (BZY20) which ignores the within-cluster dependence.

$n$	$p_m(\%)$	Proposed				BZY20			
		Bias	MCSD	ASE	CP	Bias	MCSD	ASE	CP
50	26	-0.002	0.035	0.033	0.938	0.008	0.035	0.023	0.781
	36	0.000	0.036	0.035	0.936	0.011	0.036	0.025	0.803
	44	0.002	0.037	0.036	0.936	0.013	0.037	0.027	0.832
100	26	0.002	0.024	0.024	0.950	0.012	0.023	0.016	0.740
	36	0.004	0.025	0.025	0.937	0.014	0.024	0.018	0.771
	44	0.005	0.026	0.026	0.937	0.016	0.025	0.019	0.782
200	26	0.003	0.017	0.017	0.945	0.013	0.016	0.011	0.681
	36	0.005	0.017	0.018	0.943	0.016	0.017	0.012	0.672
	44	0.007	0.018	0.018	0.940	0.018	0.018	0.014	0.663

Note:  $n$ : number of clusters with cluster size  $M \in [30, 60]$ ;  $p_m$ : percentage of missingness; MCSD: Monte Carlo standard deviation; ASE: average estimated standard error; CP: coverage probability of 95% confidence interval



Table 8: Simulation results for the infinite-dimensional parameters  $\Lambda_{0,1}(t)$  at selected time points under Scenario 2. Results from the proposed approach and the approach by Bakoyannis et al. (2020) (BZY20) which ignores the within-cluster dependence.

$n$	$p_m(\%)$	$t$	Proposed				BZY20			
			Bias	MCSD	ASE	CP	Bias	MCSD	ASE	CP
50	26	0.1	-0.004	0.088	0.083	0.911	0.084	0.106	0.028	0.308
		0.2	-0.013	0.109	0.104	0.907	0.104	0.129	0.037	0.317
		0.4	-0.014	0.141	0.134	0.922	0.139	0.161	0.052	0.339
		0.8	0.004	0.193	0.182	0.939	0.200	0.210	0.083	0.405
	44	0.1	-0.008	0.088	0.083	0.907	0.078	0.106	0.034	0.376
		0.2	-0.024	0.110	0.105	0.897	0.090	0.130	0.044	0.403
		0.4	-0.028	0.142	0.136	0.911	0.121	0.162	0.061	0.422
		0.8	-0.001	0.196	0.186	0.945	0.194	0.214	0.095	0.461
200	26	0.1	-0.004	0.041	0.042	0.955	0.085	0.050	0.014	0.101
		0.2	-0.012	0.053	0.052	0.931	0.106	0.062	0.018	0.109
		0.4	-0.015	0.068	0.067	0.934	0.139	0.078	0.026	0.111
		0.8	-0.001	0.096	0.091	0.948	0.196	0.104	0.041	0.111
	44	0.1	-0.007	0.041	0.042	0.942	0.079	0.050	0.017	0.163
		0.2	-0.021	0.053	0.053	0.916	0.092	0.063	0.022	0.201
		0.4	-0.029	0.069	0.068	0.916	0.121	0.079	0.030	0.205
		0.8	-0.006	0.098	0.093	0.940	0.188	0.106	0.047	0.168

Note:  $n$ : number of clusters with cluster size  $M \in [30, 60]$ ;  $p_m$ : percentage of missingness;  $t$ : selected time points ; MCSD: Monte Carlo standard deviation; ASE: average estimated standard error; CP: coverage probability of 95% pointwise confidence interval

Table 9: Simulation results for the infinite-dimensional parameters  $F_{0,1}(t)$  at selected time points under Scenario 2. Results from the proposed approach and the approach by Bakoyannis et al. (2020) (BZY20) which ignores the within-cluster dependence.

$n$	$p_m(\%)$	$t$	Proposed				BZY20			
			Bias	MCSD	ASE	CP	Bias	MCSD	ASE	CP
50	26	0.1	0.001	0.045	0.043	0.928	0.024	0.051	0.013	0.368
		0.2	-0.002	0.048	0.046	0.925	0.017	0.054	0.015	0.409
		0.4	-0.003	0.051	0.049	0.931	0.008	0.056	0.016	0.425
		0.8	-0.001	0.052	0.050	0.932	-0.002	0.057	0.017	0.448
	44	0.1	0.004	0.046	0.044	0.929	0.026	0.052	0.018	0.452
		0.2	-0.002	0.050	0.048	0.922	0.017	0.055	0.020	0.483
		0.4	-0.004	0.053	0.051	0.927	0.008	0.058	0.022	0.533
		0.8	-0.001	0.054	0.052	0.933	-0.001	0.058	0.023	0.567
200	26	0.1	0.003	0.022	0.022	0.957	0.025	0.025	0.007	0.270
		0.2	0.000	0.024	0.024	0.947	0.019	0.027	0.007	0.347
		0.4	-0.002	0.025	0.025	0.943	0.009	0.028	0.008	0.420
		0.8	-0.001	0.026	0.025	0.944	-0.001	0.029	0.009	0.442
	44	0.1	0.006	0.022	0.023	0.952	0.029	0.026	0.009	0.306
		0.2	0.000	0.025	0.024	0.942	0.019	0.028	0.010	0.434
		0.4	-0.002	0.026	0.026	0.945	0.009	0.029	0.011	0.517
		0.8	0.000	0.027	0.027	0.939	0.000	0.030	0.012	0.554

Note:  $n$ : number of clusters with cluster size  $M \in [30, 60]$ ;  $p_m$ : percentage of missingness;  $t$ : selected time points ; MCSD: Monte Carlo standard deviation; ASE: average estimated standard error; CP: coverage probability of 95% pointwise confidence interval

Table 10: Simulation results for the coverage probabilities of 95% simultaneous confidence bands for the infinite-dimensional parameters  $\Lambda_{0,1}(t)$  and  $F_{0,1}(t)$  under Scenario 2. Results from the proposed approach and the approach by Bakoyannis et al. (2020) (BZY20) which ignores the within-cluster dependence.

$n$	$p_m(\%)$	$\Lambda_{0,1}(t)$				$F_{0,1}(t)$			
		Proposed		BZY20		Proposed		BZY20	
		EP	HW	EP	HW	EP	HW	EP	HW
50	26	0.768	0.914	0.078	0.152	0.803	0.924	0.087	0.171
	36	0.684	0.908	0.073	0.166	0.731	0.923	0.081	0.169
	44	0.598	0.915	0.078	0.163	0.661	0.927	0.078	0.172
100	26	0.755	0.942	0.038	0.098	0.796	0.935	0.035	0.113
	36	0.568	0.938	0.026	0.089	0.609	0.931	0.024	0.089
	44	0.421	0.941	0.013	0.069	0.452	0.927	0.010	0.060
200	26	0.684	0.946	0.009	0.045	0.717	0.940	0.013	0.065
	36	0.348	0.942	0.002	0.035	0.358	0.929	0.001	0.034
	44	0.169	0.922	0.000	0.012	0.179	0.914	0.000	0.013

Note:  $n$ : number of clusters with cluster size  $M \in [30, 60]$ ;  $p_m$ : percentage of missingness; EP: equal precision bands; HW: Hall-Wellner-type bands