

Identification of Peer Effects with Miss-specified Peer Groups: Missing Data and Group Uncertainty

Christiern Rose
University of Queensland

Abstract

We consider identification of peer effects under peer group miss-specification. Our model of group miss-specification allows for missing data and peer group uncertainty. Missing data can take the form of some individuals being completely absent from the data, and the researcher need not have any information on these individuals and may not even know that they are missing. We show that peer effects are nevertheless identifiable if these individuals are missing completely at random, and propose a GMM estimator which jointly estimates the sampling probability and peer effects. In practice this means that the researcher need only have access to an individual/household level sample with group identifiers. The researcher may also be uncertain as to what is the relevant peer group for the outcome under study. We show that peer effects are nevertheless identifiable provided that the candidate peer groups are nested within one another (e.g. classroom, grade, school) and propose a non-linear least squares estimator. We conduct a Monte-Carlo experiment to demonstrate our identification results and the performance of the proposed estimators in a setting tailored to real data (the Dartmouth room-mate data).

Key words: Peer effects, social interactions, identification, missing data, measurement error, networks

1. INTRODUCTION

Empirical researchers studying peer effects face two challenges which this paper aims to address through providing identification results which lead to an estimator in a straightforward way. First, most existing methods require that the researcher has access to *group data*: a sample of groups such that within each group every member is observed (see [Bramoullé et al. \(2020\)](#) for a recent review). Such data are often proprietary and/or restricted-access due to anonymity concerns ([Breza et al., 2020](#)), and widely available individual/household survey data cannot be used because they are not designed to sample every group member (e.g. neighbourhoods or industries in national household surveys/census microsamples, grades in PISA data). The problem is particularly acute in models of peer effects because, if even one peer is missing, the peer average cannot be constructed and hence every individual which has at least one missing peer has to be discarded. In this setting it may be tempting to use the average of the observed peers. However, this introduces measurement error, which can be large if peer groups are small. Moreover, the researcher may not even know that an individual is missing, hence might not even be aware that there is measurement error in the first place. We refer to this as the *missing data* problem.

Second, the researcher often has to make a choice regarding which is the relevant peer group for the particular outcome under study. Often there will be multiple groups but the researcher will not know which is appropriate. For example, studies based on the Dartmouth room-mate data have noted that it isn't clear whether peer effects ought to

operate at the room level, the floor level or the dorm level (Sacerdote, 2001; Glaeser et al., 2003; Angrist, 2014), and the relevant group may be different for different outcomes such as fraternity membership (Sacerdote, 2001).¹ Standard empirical practice would then be to re-estimate the effects for each possible peer definition. However, this implicitly assumes that peer definition is the same for all individuals (i.e. it is deterministic) and there is no reason for this to be the case. In the context of the Dartmouth data, some dorms may simply be more sociable than others, and the researcher is unlikely to know which ones. We refer to this as the *group uncertainty* problem.

We first show that missing data and group uncertainty are both examples of a wider class of miss-specification in which the members of the specified group are a subset of the members of the (true) group. This is clear in the missing data case, and in the group uncertainty case, it is applicable in the common empirical setting in which the possible groups are nested. This is satisfied in the Dartmouth data because rooms are nested in floors which are nested in dorms. For peer effects in education, interactions could take place at the gender-grade level (Hoxby, 2000; Lavy and Schlosser, 2011) or the grade level. Similarly, neighbourhood effects might operate at the street, suburb or city levels. The members of the specified group can then be guaranteed to be members of the group by specifying the smallest group.

Next we show that, in the group interactions model we consider, if the specified groups are a subset of the groups and specified-group fixed effects are used, group miss-specification reduces to knowledge of the group size. If the group size is observed then peer effects are identifiable. The reason for this is that the specified-group fixed effect controls for part of the miss-specification. This result was first noted by Davezies et al. (2009) in the context of missing data,² but we extend it to also consider the case in which the group size is unobserved. This is important, because it allows for settings in which the researcher has no information on missing individuals and need not even know whether they exist. One such example is individual/household level survey data which are not designed to sample all group members. We show that peer effects can be identified through assumptions relating the group size to the size of the specified group. We consider two such assumptions: one for missing data and another for group uncertainty.

For missing data, we consider a Missing Completely at Random (MCAR) assumption. This allows the researcher to use individual/household level survey data with group indicators, or, if group data are available, for some individuals to be missing. The MCAR assumption implies that the distribution of the specified group size conditional on the group size follows a Binomial distribution,³ hence the specified group size provides information on the group size, which can be used to identify the peer effects. We show that the parameters are identified if (i) the distribution of group sizes has at least three points of support (i.e. there are at least three group sizes), (ii) the endogenous and contextual effects are not such that they exactly offset one another (yielding a zero reduced form peer effect) and (iii) the distribution of group sizes is bounded. Conditions (i) and (ii) are necessary even when there is no miss-specification (Lee, 2007; Davezies et al., 2009; Bramoullé et al., 2009), whereas condition (iii) is a mild requirement to allow for miss-specification. We then propose a GMM estimator which combines Maximum Likelihood estimation of the MCAR sampling probability and non-linear least squares estimation of the reduced form of the outcome equation.

For group uncertainty, we consider a setting in which the group is one of two types

¹These data are widely used because of conditionally random allocation of students to dorm rooms.

²Davezies et al. (2009) do not study group uncertainty.

³Our arguments also account for the fact that we do not observe any groups of size zero (i.e. with all members missing).

(e.g. dorm rooms and floors) at random. We can also allow the type probability to depend on the exogenous characteristics. An extension to three or more groups is also straightforward. We show that the parameters are identified under conditions (i) and (ii) from the previous paragraph and a condition on the support of the joint distribution of the sizes of the two group types. This support condition is similar in spirit to that concerning identifiability of bi-partite fixed effects models (e.g. the ‘connected set’ for the worker-firm fixed effect model of [Abowd et al. \(1999\)](#)). The basic idea is that the reduced form peer effect is a weighted average of the reduced form peer effects under each group type, each of which depends on the size of the group type. Hence if there is sufficient joint variation in the group sizes (e.g. room size and floor size), we can disentangle their reduced form peer effects, and hence identify the endogenous and contextual effects.

1.1. Related literature

We now relate our work to the literature. The literature on sampled networks, mismeasured networks and missing data, has recently attracted much interest. [Lewbel et al. \(2019\)](#) and [Hardy et al. \(2019\)](#) consider the implications of measurement error in the network through which peer effects operate. [Lewbel et al. \(2019\)](#) additionally consider the case in which only group identifiers are available and no information on the network is available beyond group identifiers. [Boucher and Houndetoungan \(2020\)](#) study peer effects when the network is unknown but that its distribution is known or can be consistently estimated. [Chandrasekhar and Lewis \(2011\)](#) studies the implications of using sampled network data in applied work, provide analytical corrections in some examples of interest and develop a more general graphical reconstruction approach. In the context of peer effects, the analytical correction can be applied if the researcher has data on the network, outcomes and exogenous characteristics for a subsample of individuals and knows the identities, outcomes and exogenous characteristics of all of their peers. This means that data are only missing for peers-of-peers. To apply graphical reconstruction, for every individual, the researcher must observe the outcomes, exogenous characteristics and variables which can be used to predict the propensity of individuals to form links in a network formation model. [Breza et al. \(2020\)](#) use a parametric network formation model to show that network structure can be identified and used to construct network statistics (e.g. centrality measures) if the researcher does not collect network data for all individuals, but instead collects relational data for a sub-sample of individuals and basic characteristics of all individuals.⁴ [Sojourner \(2013\)](#), [Wang and Lee \(2013\)](#) and [Liu et al. \(2017\)](#) consider the case in which the network and all individuals are observed without error but there are missing data in the outcomes and/or exogenous characteristics.

All of these papers suppose that the researcher observes at least some data on every individual in the group.⁵ In contrast, we allow for some individuals to be entirely missing and do not require knowledge on whether there even exist such individuals. These gains come from exploiting the structure of the group interactions model we consider. Hence, though we consider a less general network structure than other research, we also require weaker assumptions on data availability.⁶

Our work is also related to the literature on heterogeneous peer effects. [Goldsmith-Pinkham and Imbens \(2013\)](#), [Arduini et al. \(2020\)](#) and [Reza et al. \(2021\)](#)

⁴Relational data are collected by asking questions of the form ‘Consider all the individuals in your group with whom you do activity X. How many of these have trait Y?’ ([Breza et al., 2020](#))

⁵Some additionally require specification and estimation of one or more network formation model.

⁶We discuss this issue further in Section 7

study peer effects which operate through more than one network. These papers suppose that every individual is subject to the same peer effects, but that they operate through two or more networks. In contrast, we suppose that individuals are exposed to peer effects operating through only one network, but there is uncertainty as to which is the relevant network. Our model is thus not a model of heterogeneous peer effects, but one of peer effects with uncertain peer groups.

2. MODEL

We consider the group-interactions framework in which the data comprise N_0 individuals from an i.i.d. sample of G_0 groups. Though our focus is on identification, the asymptotic we have in mind is $G_0 \rightarrow \infty$, which is analogous to the large- N fixed- T asymptotic used for microeconomic panel data. This model and asymptotic has been previously studied by Lee (2007), Bramoullé et al. (2009) and Davezies et al. (2009), among others, and can be microfounded based on a game with quadratic utility in one's own outcome and the average outcome of the others in the group (see Blume et al. (2015)).

Individual $i \in \{1, 2, \dots, N_0\}$ is in exactly one group denoted $g_0(i) \in \{1, 2, \dots, G_0\}$. The size of i 's group is $n_{g_0(i)} \geq 2$. When it is not required, we omit the dependence on i and simply refer to g_0 and n_{g_0} . The continuous outcome y_i of individual i is determined by

$$y_i = \alpha_{g_0} + \left(\frac{1}{n_{g_0} - 1} \sum_{\substack{j: g_0(j)=g_0(i) \\ j \neq i}} y_j \right) \beta + \left(\frac{1}{n_{g_0} - 1} \sum_{\substack{j: g_0(j)=g_0(i) \\ j \neq i}} x_j \right) \delta + x_i \gamma + \epsilon_i, \quad (2.1)$$

where $\alpha_{g_0(i)}$ is group unobserved heterogeneity, x_i are exogenous characteristics and ϵ_i is an error term satisfying

$$\mathbb{E}[\epsilon_i | \alpha_{g_0}, n_{g_0}, \mathbf{x}_{g_0}] = 0 \quad \forall i \in \{1, 2, \dots, N_0\}, \quad (2.2)$$

where $\mathbf{x}_{g_0} = (x_j)_{j: g_0(j)=g_0(i)}$ is $n_{g_0} \times 1$. The moment condition supposes that group size and the characteristics of the group members are strictly exogenous with respect to ϵ_i . However, n_{g_0} and \mathbf{x}_{g_0} can be arbitrarily dependent on α_{g_0} . The model includes correlated effects (captured by α_{g_0}), endogenous peer effects (β) and contextual peer effects (δ). For simplicity of exposition, following Bramoullé et al. (2009) results are presented for the case where there is a single characteristic x_i . All results continue to apply in the more general case in which x_i , γ and δ are K -dimensional vectors. To ensure that the reduced form of (2.1) exists, we additionally assume that $|\beta| < 1$, in which case the reduced form is

$$y_i = \frac{\alpha_{g_0}}{1 - \beta} + \left(\sum_{\substack{j: g_0(j)=g_0(i) \\ j \neq i}} x_j \right) \pi_1(n_{g_0}) + x_i \pi_2(n_{g_0}) + u_i, \quad (2.3)$$

$$\pi_1(n) = \left(\frac{(\delta + \beta\gamma)(n-1)^{-1}}{1 - \beta(\beta + n - 2)(n-1)^{-1}} \right), \quad \pi_2(n) = \left(\frac{\gamma + \beta(\delta - \gamma(n-2))(n-1)^{-1}}{1 - \beta(\beta + n - 2)(n-1)^{-1}} \right),$$

where u_i is the reduced form error. Applying the within-group transformation

$$\tilde{y}_i = y_i - \frac{1}{n_{g_0}} \sum_{j: g_0(j)=g_0(i)} y_j$$

to the reduced form yields,

$$\tilde{y}_i = \tilde{x}_i \pi(n_{g_0}) + \tilde{u}_i, \quad \pi(n) = \left(\frac{\gamma - \delta(n-1)^{-1}}{1 + \beta(n-1)^{-1}} \right) \quad (2.4)$$

3. (POSSIBLY) MISS-SPECIFIED GROUPS

We suppose now that the researcher specifies the group $g(i)$ for individual i , with corresponding group size $n_{g(i)}$. These groups may be miss-specified, and may also cover only a subset of individuals of cardinality $N \leq N_0$. The latter could arise, for example, if the available data comprise a random sample drawn from the N_0 individuals.

Due to the group heterogeneity term (α_{g_0}), it is immediately clear that identification fails whenever the specified groups are not a subset of the groups (i.e. $g(i) = g(j)$ does not imply $g_0(i) = g_0(j)$). This is because the within-specified-group transformation

$$\bar{y}_i = y_i - \frac{1}{n_g} \sum_{j:g(j)=g(i)} y_j$$

does not eliminate α_{g_0} . Moreover, identification fails even in the absence of correlated effects (i.e. even if $\alpha_{g_0} = \alpha$) due to an omitted variables problem. To see this, consider an example in which the researcher has data on all individuals $N = N_0$ for some even number N_0 and specifies $G = 1$ group containing all individuals, when in fact there are $G_0 = 2$ groups of equal size. Solving for the reduced form and applying the within-specified-group transformation yields, for individual i in group $g_0(i) = 1$,

$$\bar{y}_i = \bar{x}_i \pi(N) + t_1 + t_2 + t_3 + \bar{u}_i \quad (3.1)$$

$$= \bar{x}_i \pi(N) + \bar{v}_i \quad (3.2)$$

where

$$t_1 = \frac{\alpha_1 - \alpha_2}{2(1 - \beta)}, \quad t_2 = \left(\frac{\gamma + \delta}{1 - \beta} \right) \frac{1}{N} \left(\sum_{j:g_0(j)=1} x_j - \sum_{j:g_0(j)=2} x_j \right),$$

$$t_3 = -\frac{\bar{x}_i N (\gamma \beta + \delta)}{2(N - 1 + \beta)(N/2 - 1 + \beta)}$$

From the above, it is immediately clear that $\mathbb{E}[\bar{v}_i | N, \mathbf{x}_g] \neq 0$. There are three reasons for this, each corresponding to a term in equation (3.1). First, the researcher specifies $\alpha_1 = \alpha_2$, which may be violated. This yields the term t_1 . Second, the researcher includes the exogenous characteristics and outcomes of those in group 2 on the right hand side of the structural equations for the group 1 outcomes. This yields the term t_2 . Finally, the researcher miss-specifies the group size to be N instead of $N/2$. This, leads the reduced form parameter to be incorrectly specified as $\pi(N)$ rather than $\pi(N/2)$, yielding the term t_3 . All of these terms are correlated with \bar{x}_i , hence any estimator (e.g. non-linear least squares) based on $\mathbb{E}[\bar{v}_i | N, \mathbf{x}_g] = 0$ is inconsistent for γ, δ, β .

The example above considered the case in which the specified group contained the group. In contrast, if the groups contain the specified groups, the first two sources of miss-specification are eliminated. This is because, if all individuals in specified group g are in group g_0 then they share the same correlated effect, and, the sum of the outcomes and exogenous characteristics over all those in group g_0 but not in group g is common to all those in group g . Hence, the terms t_1 and t_2 do not arise if the specified groups are

contained within the groups. Since it is unclear how to otherwise mitigate these sources of miss-specification, we hence proceed under the following assumption

Assumption 1 (Subset miss-specification)

$$g(i) = g(j) \Rightarrow g_0(i) = g_0(j) \text{ for all } i, j.$$

and the remainder of the paper focuses on miss-specification arising due to incorrect specification of the group size (i.e. due to t_3).

Assumption 1 covers at least two empirically important cases. First, it allows for the case in which the researcher observes only a sample of $N < N_0$ individuals drawn from all N_0 individuals. This means that the researcher may only have access to a random sample of individuals with outcomes, characteristics and group identifiers, but that the sample has not been designed to cover all members of each group. For example, in the context of education, the researcher might access a sample of students' test-scores, characteristics and classroom/school identifiers (Davezies et al., 2009) or have access to a large educational survey such as the student level PISA survey, which contains school and grade identifiers. In the context of risky behaviours, survey data on smoking, drinking and illicit drug use among students can be incomplete (Lundborg, 2006). In the context of neighbourhood effects, the neighbourhood average outcome may be measured using survey data which includes geographic identifiers (e.g. census tract). For example, Bertrand et al. (2000) use the 5% public use microsample of the 1990 Census to study welfare take-up. Glaeser et al. (2003) use the same data for wages.

In all of these cases the researcher knows the group structure but only observes the outcomes and exogenous characteristics of a subset of the relevant individuals. This implies that the group average outcome/characteristic is measured with error due to a missing data problem. As will be shown below, the form of the measurement error is such that the structural parameters are identifiable either when the group size is known or when additional assumptions are made on the missingness mechanism.⁷ For example, if data are known to result from a random sample (e.g. a survey of students or a household survey), it may be reasonable to assume that the missingness mechanism is completely at random (MCAR).

Second, Assumption 1 covers the case in which there is group uncertainty but the candidate peer groups are naturally nested. It is satisfied provided that the researcher specifies the smallest of the groups. For example, Glaeser et al. (2003) and Angrist (2014) estimate their model of peer effects for peer groups based on dorm rooms and floors. As will be shown below, the structural parameters are identifiable even if the relevant peer group is the room, the floor, or sometimes the room and sometimes the floor.

4. IDENTIFICATION

We now study identification under Assumption 1. Our identification analysis follows Bramoullé et al. (2009), hence we say that the structural parameters are (point) identified if and only if they can be uniquely recovered from the reduced form parameters presented below (i.e. there is an injective relationship). Our results are thus asymptotic in nature (see Manski (1995)), and hence characterise whether endogenous, contextual and correlated effects can be disentangled if there is no limit to the number of groups G_0 .

Under Assumption 1, the reduced form for individual i in specified group g is

$$\bar{y}_i = \bar{x}_i \pi(n_{g_0}) + \bar{u}_i \tag{4.1}$$

⁷Identification under known group sizes was first established by Davezies et al. (2009), whereas identification based on restricting the missingness mechanism is novel.

Taking conditional expectations yields

$$\mathbb{E}[\bar{y}_i | n_g, \mathbf{x}_g] = \bar{x}_i \mathbb{E}[\pi(n_{g_0}) | n_g, \mathbf{x}_g] + \mathbb{E}[\bar{u}_i | n_g, \mathbf{x}_g] \quad (4.2)$$

Equation (4.2) shows that identification prospects depend on whether there is within-specified-group variation in the exogenous characteristic, on whether $\mathbb{E}[\bar{u}_i | n_g, \mathbf{x}_g] = 0$, and if both of these conditions are satisfied, on whether the structural parameters can be uniquely recovered from the reduced form parameters $\mathbb{E}[\pi(n_{g_0}) | n_g = n, \mathbf{x}_g = \mathbf{x}]$ for all $(\mathbf{x}, n \geq 2)$ in the support of (\mathbf{x}_g, n_g) .⁸

In general, even under Assumption 1, the moment condition in (2.2) does not imply $\mathbb{E}[\bar{u}_i | n_g, \mathbf{x}_g] = 0$. Suppose however that we can make a restriction such that $\mathbb{E}[\bar{u}_i | n_g, \mathbf{x}_g] = 0$, and also that there is within-specified-group variation in $\mathbf{x}_g | n_g = n$ for all $n \geq 2$ in the support of n_g . Then equation (4.2) identifies the reduced form parameter $\mathbb{E}[\pi(n_{g_0}) | n_g = n, \mathbf{x}_g = \mathbf{x}]$, so the structural parameters are point identified if they can be uniquely recovered from

$$\mathbb{E}[\pi(n_{g_0}) | n_g = n, \mathbf{x}_g = \mathbf{x}] = \sum_{m=2}^{\infty} \mathbb{P}[n_{g_0} = m | n_g = n, \mathbf{x}_g = \mathbf{x}] \pi(n) \quad (4.3)$$

for all $(\mathbf{x}, n \geq 2)$ in the support of (\mathbf{x}_g, n_g) . The distribution $n_{g_0} | n_g, \mathbf{x}_g$ is not observed, hence the structural parameters are not point identified unless it can be restricted.⁹

We now consider three assumptions, each of which guarantees $\mathbb{E}[\bar{u}_i | n_g, \mathbf{x}_g] = 0$ and restricts the distribution of $n_{g_0} | n_g, \mathbf{x}_g$. Each assumption includes the standard model of peer effects with known peer groups as a special case, respectively when $n_g = n_{g_0}$ (Assumption 2), when $\rho = 1$ (Assumption 3) and when $\psi \in \{0, 1\}$ (Assumption 4).

Assumption 2 (Missing data: Known group sizes)

Assumption 1 holds and n_{g_0} is observed by the researcher or $n_{g_0} = f(n_g, \mathbf{x}_g)$ for a known function f .

Assumption 2 covers the simplest case in which the group size is known. Identification under Assumption 2 was first considered by Davezies et al. (2009), but we include a brief discussion for comparison with our new results below. If n_{g_0} is directly observed in the data, one simply replaces n_g with n_{g_0} in equations (4.2) and (4.3). If instead $n_{g_0} = f(n_g, \mathbf{x}_g)$, one uses (4.2) and (4.3) as stated. In either case, $\pi(m)$ is identified for all m in the support of n_{g_0} , such that the structural parameters are identified when they can be uniquely recovered given from these parameters.

Assumption 3 (Missing data: Missing Completely at Random)

For each individual $i \in \{1, 2, \dots, N_0\}$, the researcher independently observes $(y_i, x_i, g_0(i))$ with probability p_i and does not observe $(y_i, x_i, g_0(i))$ with probability $1 - p_i$, $\mathbb{E}[p_i | \mathbf{x}_{g_0}, n_{g_0}] = \rho$ and $\text{COW}[p_i, p_j | \mathbf{x}_{g_0}, n_{g_0}] = 0$ for $j \in g_0(i), j \neq i$, where $\rho \in (0, 1]$ is an unknown parameter. The researcher groups observed individuals by g_0 .

Assumption 3 covers the case in which the researcher observes a sample of size N drawn from the N_0 individuals and knows the correct group for each observed individual. It corresponds to the well known missing completely at random (MCAR) assumption which is widely used (often implicitly) in empirical work. In practice, MCAR means

⁸We require $n \geq 2$ because the term $\mathbb{E}[\pi(n_{g_0}) | n_g = 1, \mathbf{x}_g]$ is not identifiable since $n_g = 1$ implies $\bar{y}_i = \bar{x}_i = 0$.

⁹Partial identification is possible because the probabilities $\mathbb{P}[n_{g_0} = m | n_g, \mathbf{x}_g]$ for $n = 2, 3, \dots$ are non-negative and sum to 1. We do not pursue partial identification, since, as shown below, typical empirical settings faced by researchers can lead to point identification.

that the researcher has access to a sample with outcomes, characteristics and group membership indicators. However, though the researcher knows the number of individuals sampled from each group (n_g), they does not know the group sizes (n_{g_0}), nor whether there are any missing individuals in each group.¹⁰

In this setting, in an attempt to correct the measurement error induced by possibly missing individuals, it might be tempting to proceed by dropping any groups with missing members prior to estimation. There are four problems with this. First, the researcher typically would not know which groups having missing members and which do not (i.e. they may not even know if $n_g < n_{g_0}$), so it might not be possible. Second, though individuals are missing completely at random, dropping every group with at least one missing individual leads to an estimation sample of groups selected systematically on group size.¹¹ Third, it is not an efficient use of the data, because many of the groups that would be dropped still provide information on the parameters. Moreover, if ρ is small and the group sizes are large, the vast majority of observed groups will be missing at least one member. Fourth, as discussed above, with group fixed effects, the researcher need only worry about the group size, rather than missing exogenous characteristics and outcomes. As discussed below, this problem is not particularly difficult to solve, and we do not need to drop any observations to do so.

We now discuss the informational content of Assumption 3, and how this can be exploited for identification. Under Assumption 3 we have

$$n_g | n_{g_0}, \mathbf{x}_g \sim \text{Binomial}(n_{g_0}, \rho), \quad (4.4)$$

which, together with the observed distribution of $n_g | n_g \geq 1, \mathbf{x}_g$, can lead to identification of ρ and the other structural parameters.¹² As shown in the proof of Theorem 1 below, point identification of ρ comes from restrictions on the support of n_{g_0} . The result uses that $n_{g_0} \geq 2$, which is maintained throughout the literature (see e.g. Bramoullé et al. (2009), Lee (2007), Davezies et al. (2009)).¹³ Indeed, groups of size one cannot provide information on peer effects and are not consistent with our model. However, if individuals are sampled, then we may *observe* groups with only one member even though all groups have at least two members (i.e. $\mathbb{P}[n_g = 1] > 0$ if $\rho < 1$). Hence the distribution of n_g provides information on ρ , which combined with (4.4), leads to its identification. Equation (4.3) can then be used to identify the peer effects.

Assumption 4 (Group uncertainty)

For each individual $i \in \{1, 2, \dots, N_0\}$ the researcher observes $(y_i, x_i, g_1(i), g_2(i))$, $2 \leq n_{g_1}$ and $g_1(i) = g_1(j) \Rightarrow g_2(i) = g_2(j)$ for all $(i, j) \in \{1, 2, \dots, N_0\}^2$. The researcher groups individuals by g_1 , $\mathbb{E}[\epsilon_i | n_{g_1}, n_{g_2}, \mathbf{x}_{g_0}] = 0$ and $\mathbb{P}[n_{g_0} = n_{g_1} | n_{g_1}, n_{g_2}, \mathbf{x}_{g_0}] = \psi$, $\mathbb{P}[n_{g_0} = n_{g_2} | n_{g_1}, n_{g_2}, \mathbf{x}_{g_0}] = 1 - \psi$ for an unknown parameter $\psi \in [0, 1]$.

Assumption 4 covers the case in which there is group uncertainty but the candidate groups are naturally nested. For simplicity of exposition we present the case of two potential groups. Though an extension to three or more is straightforward, we do not pursue it formally. As is made clear by the proof of Theorem 1, we could also allow for ψ to depend on \mathbf{x}_g .

In the Dartmouth setting, Assumption 4 allows that peer effects operate at the room level on some floors and at the floor level on others. We also require a minor strength-

¹⁰The group sizes would not be known even if ρ were known.

¹¹This is because larger groups are more likely to have at least one missing member than smaller groups.

¹²Conditioning on $n_g \geq 1$ is important, because for some groups the researcher may not observe any individuals. Hence it is the distribution $n_g | n_g \geq 1, \mathbf{x}_g$ which is observed rather than $n_g | \mathbf{x}_g$.

¹³In practice, depending on the application one may wish to impose a lower bound larger than two, which can improve the performance of the estimator presented below.

ening of the moment condition, replacing (2.2) with $\mathbb{E}[\epsilon_i | n_{g_1}, n_{g_2}, \mathbf{x}_{g_0}] = 0$, and hence n_g with n_{g_1}, n_{g_2} in equations (4.2) and (4.3). The stronger moment condition requires that both potential group sizes be strictly exogenous with respect to the structural error. Under Assumption 4, identification hinges on the support of the joint distribution of n_{g_1} and n_{g_2} . Since the reduced form parameters are

$$\mathbb{E}[\pi(n_{g_0}) | n_{g_1} = n_1, n_{g_2} = n_2, \mathbf{x}_g = \mathbf{x}] = \psi\pi(n_1) + (1 - \psi)\pi(n_2) \quad (4.5)$$

the first part of the identification argument identifies $\psi\pi(n_1)$ and $(1 - \psi)\pi(n_2)$ for all (n_1, n_2) in a subset of the support of (n_{g_1}, n_{g_2}) using well known arguments for bi-partite fixed effects models (e.g. (Abowd et al., 1999)). In particular, we use the notion of the ‘connected-set’, which in our context corresponds to restrictions on the bi-partite graph representing the support of (n_{g_1}, n_{g_2}) . We thus use the following definition.

Definition For two discrete random variables X and Y with supports \mathcal{X} and \mathcal{Y} , the bi-partite graph $\mathcal{G}(X, Y)$ has $|\mathcal{X}| + |\mathcal{Y}|$ vertices, with each vertex corresponding to a support point of either X or Y . For two vertices, one corresponding to the support point $x \in \mathcal{X}$ and another to the support point $y \in \mathcal{Y}$, an undirected edge exists if $\mathbb{P}[X = x, Y = y] > 0$ and does not exist otherwise. There are no further edges in $\mathcal{G}(X, Y)$.

We now present the main identification result.

Theorem 1

(γ, β, δ) are point identified if $\gamma\beta + \delta \neq 0$ and any one of the following holds

1. Assumption 2 holds and the support of the distribution of n_{g_0} has at least three elements (Davezies et al., 2009).
2. Assumption 3 holds and the support of the distribution of n_{g_0} is bounded and has at least three elements.
3. Assumption 4 holds, the support of the distribution of n_{g_1} has at least three elements, the support of the distribution of n_{g_2} has at least three elements and either exists a connected component of $\mathcal{G}(n_{g_1}, n_{g_2})$ with at least 3 vertices corresponding to the support of n_{g_1} or there exists a connected component of $\mathcal{G}(n_{g_1}, n_{g_2})$ with at least 3 vertices corresponding to the support of n_{g_2} .

Theorem 1 shows that the structural parameters are identifiable if $\gamma\beta + \delta \neq 0$ and the distribution of group sizes has sufficient support. The former is a well known necessary identification condition in models with correctly specified groups and both endogenous and contextual effects (see for example Bramoullé et al. (2009) or Rose (2017)), which is generically satisfied over the parameter space. If it is violated then the endogenous and contextual effects exactly offset one another such that the reduced form effect is $\pi(n) = \gamma$, which does not vary with n , and hence no amount of variation in group sizes can identify β and δ . All three results in Theorem 1 also require that the support of the distribution of n_{g_0} has at least three elements. This is also necessary condition for identification when the groups are correctly specified and there are endogenous, contextual and correlated effects (Lee, 2007; Davezies et al., 2009; Bramoullé et al., 2009), hence it is also necessary if the groups are (possibly) miss-specified.

Result 2 also uses the mild assumption that the distribution of n_{g_0} be bounded, which is needed to identify ρ and the distribution of $n_{g_0} | \mathbf{x}_g$ from the distribution of $n_g | n_g \geq 1, \mathbf{x}_g$ as a solution of an eigensystem.¹⁴ This assumption is also used, for example, by

¹⁴The lower bound on the support of n_{g_0} is 2 by assumption, and the upper bound need not be known since it is identifiable from the distribution of $n_g | n_g \geq 1$.

Boucher and Houndetoungan (2020). Result 3 uses a mild condition on the joint support of the candidate group sizes (n_{g_1}, n_{g_2}) . It is satisfied, for example, whenever there exists n in the support of n_{g_1} such that the support of $n_{g_2}|n_{g_1} = n$ has at least three elements or the converse, reversing the roles of n_{g_1} and n_{g_2} , holds.

5. ESTIMATION

If the group sizes are known (Assumption 2), we estimate γ, δ, β by non-linear least squares based on

$$\mathbb{E}[\bar{y}_i | n_{g_0}, \mathbf{x}_g] = \bar{x}_i \pi(n_{g_0}).$$

Similarly, for group uncertainty (Assumption 4), we apply non-linear least squares to based on

$$\mathbb{E}[\bar{y}_i | n_{g_1}, n_{g_2}, \mathbf{x}_g] = \bar{x}_i [\psi \pi(n_{g_1}) + (1 - \psi) \pi(n_{g_2})].$$

If the researcher has a random sample of individuals (Assumption 3), we apply maximum likelihood for ρ and non-linear least squares for γ, δ, β . For maximum likelihood estimation of ρ , we use the sample of G observed group sizes n_g with known distribution

$$\mathbb{P}[n_g = n | n_g \geq 1] = \frac{\sum_{m=n}^{\bar{n}} \mathbb{P}[n_{g_0} = m] \binom{m}{n} \rho^n (1 - \rho)^{1-n}}{\sum_{l=1}^{\bar{n}} \sum_{m=l}^{\bar{n}} \mathbb{P}[n_{g_0} = m] \binom{m}{l} \rho^l (1 - \rho)^{1-l}} \quad (5.1)$$

for $n = 1, \dots, \bar{n}$. Thus, since $\mathbb{P}[n_{g_0} = 1] = 0$ by assumption and $\mathbb{P}[n_{g_0} = \bar{n}] = 1 - \sum_{m=2}^{\bar{n}-1} \mathbb{P}[n_{g_0} = m]$ we jointly estimate the $\bar{n} - 1$ parameters $\rho, \mathbb{P}[n_{g_0} = 2], \dots, \mathbb{P}[n_{g_0} = \bar{n} - 1]$. To estimate γ, δ, β , we apply non-linear least squares to

$$\mathbb{E}[\bar{y}_i | n_g, \mathbf{x}_g] = \bar{x}_i \sum_{n=2}^{\bar{n}} \mathbb{P}[n_{g_0} = n | n_g] \pi(n) \quad (5.2)$$

where $\mathbb{P}[n_{g_0} = n | n_g]$ is a known function of the parameters ρ and $\mathbb{P}[n_{g_0} = 2], \dots, \mathbb{P}[n_{g_0} = \bar{n} - 1]$, which we do not make explicit for simplicity of exposition.¹⁵ In practice we do not sequentially estimate the parameters by maximum likelihood and then by non-linear least squares. Instead, we estimate the parameters jointly by GMM based on the non-linear least squares moment condition and equality to zero of the expectation of the score of the log-likelihood function based on (5.1). This framework allows the researcher to include additional information when it is available. For example, if the researcher knows which observed groups have missing members (i.e. they observes an indicator for $n_g = n_{g_0}$), they can modify the likelihood to also use $\mathbb{P}[n_g = n_{g_0} | n_g] = \rho^{n_g}$. In the same way, it is also straightforward to make a parametric assumption on the distribution of n_{g_0} .

6. MONTE-CARLO

We tailor the design to the well known data on roommates at Dartmouth college studied by Sacerdote (2001), Glaeser et al. (2003) and Angrist (2014), among others. The original data comprise 1589 freshman at Dartmouth college who were randomly assigned to dorm-rooms. Fifty three percent of dorm-rooms were doubles, 44 percent were triples and the remaining 3 percent were quads. The first part of our Monte-Carlo experiment

¹⁵To control the number of parameters to be estimated we make the additional restriction $\mathbb{P}[n_{g_0} = n | n_g, \mathbf{x}_g] = \mathbb{P}[n_{g_0} = n | n_g]$ for $n = 1, \dots, \bar{n}$ to derive (5.2). This contrasts with our identification results, for which this additional restriction is not made.

considers what would have happened had only a random sample of these students been available. The second part considers the issue of the definition of the relevant peer group, which could be the room or the floor (Sacerdote, 2001; Glaeser et al., 2003; Angrist, 2014).

6.1. Missing data

We now suppose that only a random sample from the Dartmouth data were available. The original Dartmouth data comprise a complete sample of freshman (i.e. $\rho = 1$). Our design varies $\rho \in \{0.3, 0.5, 0.7, 0.9, 1\}$. For $\rho = 1$ we set the sampling probability $p_i = 1$ for all $i \in \{1, \dots, N_0\}$ and for $\rho < 1$ we set $p_i = \rho + q_i$, where $q_i \stackrel{i.i.d.}{\sim} \text{Uniform}(-0.1, 0.1)$. We set G_0 to be the nearest integer to $M/(\rho\mathbb{E}[n_{g_0}])$, where M is discussed below. For each dataset, we draw G_0 dorm-rooms from the distribution of dorm-room sizes, which we take to be $n_{g_0} - 2 \sim \text{Binomial}(2, 0.25)$ so as to match the sample mean and support of the Dartmouth data.¹⁶ We use a parametric distribution so as to evaluate the performance of the estimator both with and without a parametric restriction. We use the Binomial distribution because it is bounded, and hence satisfies the requirements of the identification in Theorem 1.

Since we draw a fixed number of dorm rooms, each with a random size, the number of students N_0 varies from one dataset to the next. We then draw an independent random sample of N students, which includes each of the N_0 students independently with probability ρ . Hence N also varies from one dataset to the next. Allowing G_0 to depend on ρ as above implies that the size of the observed sample is $N \approx M$ no matter the value of ρ . Setting $M = 1600$, our design matches the sample size of the Dartmouth data. We thus consider the performance of our method had the Dartmouth data been obtained from a random sample of freshman, rather than all freshman, holding the number of observations constant as we vary ρ . We also consider $M = 8000$ to study the performance of the method under different sample sizes.

Sacerdote (2001) exploits random assignment of freshman to dorms in the Dartmouth data to identify peer effects in educational attainment, measured by freshman GPA. It is argued that endogenous effects are difficult to identify due to the reflection problem, though in principle the variation in dorm-room sizes could be used to achieve this because there are three room-sizes (Lee, 2007; Davezies et al., 2009; Bramoullé et al., 2009). Sacerdote (2001) focuses instead on credibly identifying the reduced form effect of average room-mate high school attainment (Table 3, column 5) on freshman GPA. The estimated peer effect is positive and statistically significant, though smaller in magnitude than own high-school attainment. This reduced form regression has $R^2 = 0.19$.

To match this setting as well as possible, in our design we set the own effect to be $\gamma = 1$, and the endogenous effect to be $\beta = 0$, hence the reduced form effect of room-mate's high-school attainment is equal to the contextual effect, which we set to be $\delta = 0.5$. We set $x_i \stackrel{i.i.d.}{\sim} \mathcal{N}(0, 1)$, $\epsilon_i \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \sigma^2)$ and $\alpha_{g_0} \stackrel{i.i.d.}{\sim} \mathcal{N}(1, \sigma^2)$ and choose $\sigma^2 = 2(\gamma^2 + \delta^2/\mathbb{E}[n_{g_0} - 1])$. For the expected dorm-room size, this choice corresponds to fraction 0.8 of $\mathbb{V}[y_i]$ being due to $\alpha_{g_0(i)} + \epsilon_i$ and 0.2 being due to $x_i\gamma + (1 - n_{g_0})^{-1} \sum_{j:g_0(j)=g_0(i), j \neq i} x_j$, hence a population R^2 of 0.2 in a reduced form regression without dorm nor dorm-room fixed effects, as estimated by Sacerdote (2001). The specification that we estimate differs from that of Sacerdote (2001) because we include dorm-room fixed effects, which are necessary so as to allow for miss-specification.

¹⁶In this design, in expectation, 56.25% of rooms are doubles, 37.5% are triples and 6.25% are quads.

6.1.1 Results

Table 1 presents the results. We consider four estimators; one for which the observed groups are treated as if they are the groups ('Miss-specified'), one for which the group size is known ('Known'), one for which the group size is unknown but Assumption 3 holds ('MCAR') and a modification of MCAR under which the parametric assumption $n_{g_0} - 2 \sim \text{Binomial}(2, \omega)$ is additionally imposed ('MCAR-P'). The MCAR and MCAR-P specifications take the upper bound on the support of the group sizes to be known as $\bar{n} = 4$, which, in the context of the Dartmouth data implies that the researcher knows that there are no rooms of more than four students. The Miss-specified, MCAR and MCAR-P estimators only use the outcome, exogenous characteristic and group indicator for the N sampled individuals. The Known estimator additionally uses knowledge of the group size.

Beginning with the contextual effects only specifications ($\beta = 0$ is imposed) and $\rho = 1$, we see that the results are very similar (identical to three decimal places) for all four estimators. This suggests that there is little cost in applying the MCAR approach when there are no missing data. As ρ decreases, the distribution of the miss-specified estimator δ and γ both shift closer to zero. For $\rho \leq 0.5$, the bias becomes sizeable. In contrast, the other estimators remain approximately unbiased for all values of ρ , though their RMSE increases as ρ decreases. This is likely because, as ρ decreases the number of groups with only one observed member increases, and these groups provide no information on the parameters due to the within-observed-group transformation.

The estimator with known group sizes performs at least as well as the MCAR estimator in terms of RMSE. This is unsurprising since it also requires more information in order to be implemented. For $\rho = 0.9$ the difference in RMSE is small (0.087 vs 0.1 for δ when $N \approx M = 8000$), though the gap widens as ρ decreases. This is likely because, as ρ decreases, there is less variation in the distribution of observed group sizes, and hence, for a given observed group size, more uncertainty as to the group size. The additional parametric assumption on the group size distribution makes little difference when $\rho \geq 0.7$, but when $\rho \leq 0.5$ the gap in RMSE between the MCAR and MCAR-P estimators becomes non-negligible.

Moving on to the specifications with contextual and endogenous effects, it is clear that, though identified due to there being at least three group sizes, there is insufficient group size variation in this design to reliably separate the two. In particular, the RMSE on β is large and all estimators of β are biased upwards, whereas estimators of δ tend to be biased downwards. Nevertheless, the qualitative conclusions regarding the relative performance of the four estimators are the same as for the specifications with contextual effects only.

6.2. Group uncertainty

Sacerdote (2001), Glaeser et al. (2003) and Angrist (2014) all discuss the appropriate definition of the peer group, which may either be the room, the floor or the entire dorm. It is also possible that the appropriate peer group for academic attainment may be different to that of social outcomes such as fraternity membership (Sacerdote, 2001). For this reason, these papers repeat their empirical analysis for interactions at the room, floor and dorm levels.

We focus here on the distinction between the room and the floor. To match the Dartmouth data, we first draw the rooms as described above. We then draw $f_1 \in \{1, 2, \dots, 5\}$ uniformly, and take rooms $\{1, 2, \dots, f_1\}$ to comprise the first floor. We then draw another integer $f_2 \in \{1, 2, \dots, 5\}$ and take rooms $\{f_1 + 1, \dots, f_1 + f_2\}$ to comprise the second floor.

We proceed in this way until every room has been allocated to a floor. In the Dartmouth data, the sample mean number of students per floor is 8. This data generating process yields an expected number of students per floor of 7.5. All other aspects of the data generating process remain unchanged and we vary $\psi \in \{0.2, 0.4, 0.6, 0.8\}$.

6.2.1 Results

Table 2 presents the results. We consider four estimators; one in which the room is specified as the peer group ('Room'), one in which the floor is specified as the peer group ('Floor'), one in which the researcher knows whether the group is the room or the floor ('Known') and one in which the group is unknown but Assumption 4 is assumed to hold ('Uncertain').

For brevity, the discussion focuses on the contextual effects only specifications. Specifying the group to be the room leads to downwards bias of δ . The bias and RMSE grow as ψ decreases. This is because the proportion of incorrectly specified groups grows as ψ decreases. For the same reason the RMSE from specifying the group to be the dorm decreases as ρ decreases. The results suggest that the estimator based on specifying the floor is unbiased. This is coincidental. It just so happens that in this design the three sources of bias (the terms t_1, t_2 and t_3 discussed in Section 3) offset one another. To demonstrate this, Table 3 modifies the design by setting $\alpha_{g_0} \stackrel{i.i.d.}{\sim} \mathcal{N}(n_{g_0}^{-1} \sum_{j:j \in g_0} x_j, \sigma^2)$ instead of $\alpha_{g_0} \stackrel{i.i.d.}{\sim} \mathcal{N}(1, \sigma^2)$. This has no impact on the estimators Room, Known and Uncertain because all are based on room fixed effects.¹⁷ However it introduces substantial bias for the Floor estimator which is based on floor fixed effects.

Comparing the Known estimator to the Uncertain estimator, as expected the RMSE of the latter is larger since it requires less information to be implemented. Nevertheless, depending on the sample size and value of ψ , the estimator based on Assumption 4 has small bias and sufficiently small RMSE so as to be distinguishable from zero. Considering now the specifications with endogenous and contextual effects, though not well identified, the main qualitative conclusions regarding the relative performance of the four estimators are the same as for the specifications with contextual effects only. Notice also that once endogenous effects are introduced the estimator based on specifying the floor often has larger bias and RMSE than that based on Assumption 4. This is particularly true for δ .

7. DISCUSSION

Our identification and Monte-Carlo results demonstrate that it is possible to conduct empirical analysis of peer effects despite missing data and group uncertainty. Regarding missing data, we require only that the researcher has access to a sample of individuals with outcome, exogenous characteristic and group membership data. Under the MCAR assumption, we require no further information on group size nor do we require any information on the individuals which were not sampled, nor do we even need to know if such individuals exist or not. In principal, and subject to the limitations discussed below, this opens up the possibility of peer effects studies based on widely available individual level survey data (provided that it contains group identifiers). Two examples of such data are the PISA sample of students, for which school-grade identifiers are available, and household surveys with local area identifiers (e.g. Census tracts or suburb identifiers). We also show that peer effects can be identified under group uncertainty.

¹⁷For these estimators the results are numerically identical to results reported in Table 2 because identical seeds were used to draw the data for each replication.

Our results could also be extended to incorporate both missing individuals and group uncertainty simultaneously.

A limitation of this paper is that our results are specific to the group interactions model we consider. Under more general network structures (e.g. social networks), it is not the case the specified group fixed effects reduce the problem of miss-specification to the problem of inferring the group size. This is because the missing individuals do not all exert the same peer effect on the sampled individuals.

The specificity of our results to the group interactions model also has implications regarding applicability to survey data as discussed above. The reason for this is that, even if the groups are correctly specified, there is a weak identification problem when group sizes are large (Lee, 2007). Hence in order to use our approach with survey data, the researcher would need to be sure that groups are small. For example, neighbourhood effects based on census-tract identifiers would likely be weakly identified, though the prospects would be better for student survey data with classroom/grade identifiers such as PISA. Future work may seek to resolve these issues, either through appealing to additional data and/or through specification of a network formation model. Another concern using survey data, particularly for studies of neighbourhood effects, might be that there is selection of individuals into groups. Though our focus is not on network endogeneity, our specification of the correlated effect allows for arbitrary dependence of group members' characteristics and the number of individuals in the group with unobserved group heterogeneity, which considerably mitigates this issue.

REFERENCES

- ABOWD, J. M., F. KRAMARZ AND D. N. MARGOLIS, "High wage workers and high wage firms," *Econometrica* 67 (1999), 251–333.
- ANGRIST, J. D., "The perils of peer effects," *Labour Economics* 30 (2014), 98–108.
- ARDUINI, T., E. PATAcCHINI AND E. RAINONE, "Identification and estimation of network models with heterogeneous interactions," in *The Econometrics of Networks* (Emerald Publishing Limited, 2020).
- BERTRAND, M., E. F. LUTTMER AND S. MULLAINATHAN, "Network effects and welfare cultures," *The Quarterly Journal of Economics* 115 (2000), 1019–1055.
- BLUME, L. E., W. A. BROCK, S. N. DURLAUF AND R. JAYARAMAN, "Linear social interactions models," *Journal of Political Economy* 123 (2015), 444–496.
- BOUCHER, V. AND A. HOUNDETOUNGAN, "Estimating peer effects using partial network data," *Working paper* (2020).
- BRAMOULLÉ, Y., H. DJEBBARI AND B. FORTIN, "Identification of peer effects through social networks," *Journal of econometrics* 150 (2009), 41–55.
- , "Peer effects in networks: A survey," *Annual Review of Economics* 12 (2020), 603–629.
- BREZA, E., A. G. CHANDRASEKHAR, T. H. MCCORMICK AND M. PAN, "Using aggregated relational data to feasibly identify network structure without network data," *American Economic Review* 110 (2020), 2454–84.
- CHANDRASEKHAR, A. AND R. LEWIS, "Econometrics of sampled networks," *Unpublished manuscript, MIT.[422]* (2011).

- DAVEZIES, L., X. D'HAULTFOEUILLE AND D. FOUGÈRE, "Identification of peer effects using group size variation," *The Econometrics Journal* 12 (2009), 397–413.
- GLAESER, E. L., B. I. SACERDOTE AND J. A. SCHEINKMAN, "The social multiplier," *Journal of the European Economic Association* 1 (2003), 345–353.
- GOLDSMITH-PINKHAM, P. AND G. W. IMBENS, "Social networks and the identification of peer effects," *Journal of Business & Economic Statistics* 31 (2013), 253–264.
- HARDY, M., R. M. HEATH, W. LEE AND T. H. McCORMICK, "Estimating spillovers using imprecisely measured networks," *arXiv preprint arXiv:1904.00136* (2019).
- HOXBY, C., "Peer effects in the classroom: Learning from gender and race variation," Technical Report, National Bureau of Economic Research, 2000.
- LAVY, V. AND A. SCHLOSSER, "Mechanisms and impacts of gender peer effects at school," *American Economic Journal: Applied Economics* 3 (2011), 1–33.
- LEE, L.-F., "Identification and estimation of econometric models with group interactions, contextual factors and fixed effects," *Journal of Econometrics* 140 (2007), 333–374.
- LEWBEL, A., X. QU, X. TANG ET AL., "Social networks with misclassified or unobserved links," *Unpublished manuscript* (2019).
- LIU, X., E. PATACCINI AND E. RAINONE, "Peer effects in bedtime decisions among adolescents: a social network model with sampled data," *The econometrics journal* 20 (2017), S103–S125.
- LUNDBORG, P., "Having the wrong friends? Peer effects in adolescent substance use," *Journal of health economics* 25 (2006), 214–233.
- MANSKI, C. F., *Identification problems in the social sciences* (Harvard University Press, 1995).
- REZA, S., P. MANCHANDA AND J.-K. CHONG, "Identification and Estimation of Endogenous Peer Effects Using Partial Network Data from Multiple Reference Groups," *Management Science* (2021).
- ROSE, C. D., "Identification of peer effects through social networks using variance restrictions," *The Econometrics Journal* 20 (2017), S47–S60.
- SACERDOTE, B., "Peer effects with random assignment: Results for Dartmouth roommates," *The Quarterly journal of economics* 116 (2001), 681–704.
- SOJOURNER, A., "Identification of peer effects with missing peer data: Evidence from Project STAR," *The Economic Journal* 123 (2013), 574–605.
- WANG, W. AND L.-F. LEE, "Estimation of spatial autoregressive models with randomly missing data in the dependent variable," *The Econometrics Journal* 16 (2013), 73–102.

APPENDIX

Proof of Theorem 1

The first result was established by (Davezies et al., 2009) and so we do not repeat their proof here.

To establish the second result, first note that $\mathbb{E}[\epsilon_i | n_{g_0}, \mathbf{x}_{g_0}] = 0$ and Assumption 3 imply $\mathbb{E}[\epsilon_i | n_g, \mathbf{x}_g] = 0$, hence $\mathbb{E}[\bar{u}_i | n_g, \mathbf{x}_g] = 0$. Let us now fix $\mathbf{x}_g = \mathbf{x}$ for some \mathbf{x} in its support. We begin by identifying ρ and the distribution of $n_{g_0} | \mathbf{x}_g = \mathbf{x}$ using the observable distribution of $n_g | n_g \geq 1, \mathbf{x}_g = \mathbf{x}$ and the fact that Assumption 3 implies

$$n_g | n_{g_0}, \mathbf{x}_g = \mathbf{x} \sim \text{Binomial}(n_{g_0}, \rho)$$

By the boundedness assumption, there exists $\bar{n} < \infty : \mathbb{P}[n_{g_0} = \bar{n}] > 0, \mathbb{P}[n_{g_0} \leq \bar{n}] = 1$. Note that, because $\rho > 0$, \bar{n} is identifiable as the integer n such that $\mathbb{P}[n_g = n | n_g \geq 1] > 0$. This implies that there exists identifiable $\bar{n}(\mathbf{x}) < \infty : \mathbb{P}[n_{g_0} = \bar{n}(\mathbf{x}) | \mathbf{x}_g = \mathbf{x}] > 0, \mathbb{P}[n_{g_0}(\mathbf{x}) \leq \bar{n}(\mathbf{x}) | \mathbf{x}_g = \mathbf{x}] = 1$.

Let $\mathbf{p}_n(\mathbf{x}) = \mathbb{P}[n_g = n | n_g \geq 1, \mathbf{x}_g = \mathbf{x}]$, $\mathbf{q}_m(\mathbf{x}) = \mathbb{P}[n_{g_0} = m | \mathbf{x}_g = \mathbf{x}]$. Under Assumption 3, $\mathbf{p}(\mathbf{x}) = (\mathbf{p}_n(\mathbf{x}))_{n=1, \dots, \bar{n}(\mathbf{x})}$ and $\mathbf{q}(\mathbf{x}) = (\mathbf{q}_m(\mathbf{x}))_{m=1, \dots, \bar{n}(\mathbf{x})}$ verify

$$\mathbf{p}(\mathbf{x}) = \frac{\mathbf{A}(\mathbf{x}, \rho) \mathbf{q}(\mathbf{x})}{\iota'_{\bar{n}(\mathbf{x})} \mathbf{A}(\mathbf{x}, \rho) \mathbf{q}(\mathbf{x})} \quad (7.1)$$

where $\mathbf{A}(\mathbf{x}, \rho) = (\mathbf{A}_{ij}(\mathbf{x}, \rho))_{i=1, \dots, \bar{n}(\mathbf{x}), j=1, \dots, \bar{n}(\mathbf{x})}$,

$$\mathbf{A}_{ij}(\mathbf{x}, \rho) = \begin{cases} \binom{j}{i} \rho^i (1 - \rho)^{j-i} & i \leq j \\ 0 & \text{otherwise} \end{cases}$$

and $\iota'_{\bar{n}(\mathbf{x})}$ is the $\bar{n}(\mathbf{x})$ dimensional vector of ones. Rearranging (7.1) yields

$$(\mathbf{p}(\mathbf{x}) \iota'_{\bar{n}(\mathbf{x})} \mathbf{A}(\mathbf{x}, \rho) - \mathbf{A}(\mathbf{x}, \rho)) \mathbf{q}(\mathbf{x}) = \mathbf{0} \quad (7.2)$$

The inverse of $\mathbf{A}(\mathbf{x}, \rho)$ exists since $|\mathbf{A}(\mathbf{x}, \rho)| = \rho^{T(\bar{n}(\mathbf{x}))}$ where $T(\bar{n}(\mathbf{x}))$ is the $\bar{n}(\mathbf{x})^{\text{th}}$ triangular number and $\rho > 0$, hence

$$(\mathbf{A}(\mathbf{x}, \rho)^{-1} \mathbf{p}(\mathbf{x}) \iota'_{\bar{n}(\mathbf{x})} \mathbf{A}(\mathbf{x}, \rho) - \mathbf{I}_{\bar{n}(\mathbf{x})}) \mathbf{q}(\mathbf{x}) = \mathbf{0} \quad (7.3)$$

Denote $\mathbf{B}(\mathbf{x}, \rho) = \mathbf{A}(\mathbf{x}, \rho)^{-1} \mathbf{p}(\mathbf{x}) \iota'_{\bar{n}(\mathbf{x})} \mathbf{A}(\mathbf{x}, \rho)$. Then clearly the rank of $\mathbf{B}(\mathbf{x}, \rho)$ cannot exceed 1. Moreover, since $\rho > 0$ we have $\mathbf{B}(\mathbf{x}, \rho) \neq \mathbf{0}$, hence $\mathbf{B}(\mathbf{x}, \rho)$ has rank 1. From (7.3), we have that 1 is an eigenvalue of $\mathbf{B}(\mathbf{x}, \rho)$. Hence $\mathbf{B}(\mathbf{x}, \rho)$ has eigenvalues 1 of multiplicity 1 and 0 of multiplicity $\bar{n}(\mathbf{x}) - 1$ and $\mathbf{q}(\mathbf{x})$ is an eigenvector associated with the eigenvalue 1. Let $\mathbf{v}(\mathbf{x}, \rho)$ be an arbitrary eigenvector associated with the eigenvalue 1. Then, since $\mathbf{q}(\mathbf{x})$ is a vector of probabilities summing to one, we have $\mathbf{q}(\mathbf{x}) = \mathbf{v}(\mathbf{x}, \rho) / (\iota'_{\bar{n}(\mathbf{x})} \mathbf{v}(\mathbf{x}, \rho))$. Hence, given ρ we can identify $\mathbf{q}(\mathbf{x})$.

To identify ρ , we use the restriction $\mathbf{q}_1(\mathbf{x}) = 0$, which is known because $n_{g_0} \geq 2$. Since $\mathbf{B}(\mathbf{x}, \rho)$ is a smooth function of ρ for $\rho \in (0, 1]$ and 1 is always a simple eigenvalue of $\mathbf{B}(\mathbf{x}, \rho)$, the eigenvector $\mathbf{v}(\rho)$ is a smooth function of ρ for $\rho \in (0, 1]$, hence so is $\mathbf{q}_1(\mathbf{x})$. Moreover, it is clear that $\mathbf{q}_1(\mathbf{x})$ is strictly decreasing in ρ if $\mathbf{p}(\mathbf{x})$ is fixed. Hence the restriction $\mathbf{q}_1(\mathbf{x}) = 0$ identifies ρ . By these arguments, ρ and $\mathbf{q}(\mathbf{x})$ are identified.

Provided that there is variation in the elements of \mathbf{x} , the conditional moment

$$\mathbb{E}[\bar{y}_i | n_g = n, \mathbf{x}_g = \mathbf{x}] = \bar{x}_i \mathbb{E}[\pi(n_{g_0}) | n_g = n, \mathbf{x}_g = \mathbf{x}] \quad (7.4)$$

identifies $\mathbb{E}[\pi(n_{g_0}) | n_g = n, \mathbf{x}_g = \mathbf{x}]$ for all such $(\mathbf{x}, n \geq 2)$ in the support of (\mathbf{x}_g, n_g) (it is not identified when $n = 1$ nor when $\mathbf{x} = c \iota_n$ for some constant c because in these cases $\bar{y}_i = \bar{x}_i = 0$). The rest of the proof restricts attention to the case in which there is variation in the elements of \mathbf{x} .

Define $\mathbf{r}_n(\mathbf{x}) = \mathbb{P}[n_g = n | \mathbf{x}_g = \mathbf{x}]$ and $\mathbf{r}(\mathbf{x}) = (\mathbf{r}_n(\mathbf{x}))_{n=1, \dots, \bar{n}(\mathbf{x})}$. So

$$\mathbf{r}(\mathbf{x}) = (\ell'_{\bar{n}(\mathbf{x})} \mathbf{A}(\mathbf{x}, \rho) \mathbf{q}(\mathbf{x})) \mathbf{p}(\mathbf{x})$$

is also identified. Now denote $\mathcal{Q}(\mathbf{x}) = \{m \in \{2, \dots, \bar{n}(\mathbf{x})\} : \mathbf{q}_m(\mathbf{x}) > 0\}$, the largest element of which is $\bar{n}(\mathbf{x})$. Moreover, since $\rho > 0$, the support of $\mathbf{r}(\mathbf{x})$ is $\mathcal{R}(\mathbf{x}) = \{1, \dots, \bar{n}(\mathbf{x})\}$ if $\rho \in (0, 1)$ and $\mathcal{R}(\mathbf{x}) = \mathcal{Q}(\mathbf{x})$ if $\rho = 1$. Let $\hat{\boldsymbol{\pi}}(\mathbf{x}) = (\pi(m))_{m \in \mathcal{Q}(\mathbf{x})}$, $\hat{\mathbf{q}}(\mathbf{x}) = (\mathbf{q}_m(\mathbf{x}))_{m \in \mathcal{Q}(\mathbf{x})}$, $\hat{\mathbf{r}}(\mathbf{x}) = (\mathbf{r}_n(\mathbf{x}))_{n \in \mathcal{R}(\mathbf{x}) \setminus 1}$ and $\hat{\mathbf{A}}(\mathbf{x}, \rho)$ be the sub-matrix of $\mathbf{A}(\mathbf{x}, \rho)$ formed from rows $\mathcal{R}(\mathbf{x}) \setminus 1$ and columns $\mathcal{Q}(\mathbf{x})$. Then we have the equations

$$(\mathbb{E}[\pi(n_{g_0}) | n_g = n, \mathbf{x}_g = \mathbf{x}])_{n \in \mathcal{R}(\mathbf{x}) \setminus 1} = \text{diag}(\hat{\mathbf{r}}(\mathbf{x}))^{-1} \hat{\mathbf{A}}(\mathbf{x}, \rho) \text{diag}(\hat{\mathbf{q}}(\mathbf{x})) \hat{\boldsymbol{\pi}}(\mathbf{x}), \quad (7.5)$$

for which the left hand side is identified. Since $\mathbf{A}(\mathbf{x}, \rho)$ has full rank, $\hat{\mathbf{A}}(\mathbf{x}, \rho)$ has rank $|\mathcal{Q}(\mathbf{x})|$. This is because, when $\rho = 1$, we have $\mathcal{R}(\mathbf{x}) = \mathcal{Q}(\mathbf{x})$ and $\hat{\mathbf{A}}(\mathbf{x}, \rho) = \mathbf{I}_{|\mathcal{Q}(\mathbf{x})|}$. When $\rho \in (0, 1)$, $\mathcal{R} = \{1, \dots, \bar{n}(\mathbf{x})\}$, hence $\hat{\mathbf{A}}(\mathbf{x}, \rho)$ comprises rows $\{2, \dots, \bar{n}(\mathbf{x})\}$ and a subset of columns $\{2, \dots, \bar{n}(\mathbf{x})\}$ of $\mathbf{A}(\mathbf{x}, \rho)$. The sub-matrix of $\mathbf{A}(\mathbf{x}, \rho)$ with rows $\{2, \dots, \bar{n}(\mathbf{x})\}$ and columns $\{2, \dots, \bar{n}(\mathbf{x})\}$ has determinant $\rho^{T(\bar{n}(\mathbf{x})-1)} > 0$, which implies that $\hat{\mathbf{A}}(\mathbf{x}, \rho)$ has rank $|\mathcal{Q}(\mathbf{x})|$. Moreover, $\text{diag}(\hat{\mathbf{q}}(\mathbf{x}))$ is invertible by construction, hence (7.5) identifies $\hat{\boldsymbol{\pi}}(\mathbf{x})$ for all \mathbf{x} in the support of \mathbf{x}_g .

By assumption there are at least three different group sizes in the support of n_{g_0} , so $|\cup_{\mathbf{x}} \mathcal{Q}(\mathbf{x})| \geq 3$. This implies that $\pi(n_1), \pi(n_2), \pi(n_3)$ are identified for at least three different group sizes. We conclude by applying the well known result established by Lee (2007), Davezies et al. (2009) and Bramoullé et al. (2009) that (γ, δ, β) are identified when there are at least three different group sizes and $\gamma\beta + \delta \neq 0$.

To establish the third result, again fix $\mathbf{x}_g = \mathbf{x}$ where \mathbf{x} is some point in the support of \mathbf{x}_g . Provided that there is variation in the elements of \mathbf{x} , the conditional moment

$$\mathbb{E}[\bar{y}_i | n_{g_1} = n_1, n_{g_2} = n_2, \mathbf{x}_g = \mathbf{x}] = \bar{x}_i \mathbb{E}[\pi(n_{g_0}) | n_{g_1} = n_1, n_{g_2} = n_2, \mathbf{x}_g = \mathbf{x}] \quad (7.6)$$

identifies $\mathbb{E}[\pi(n_{g_0}) | n_{g_1} = n_1, n_{g_2} = n_2, \mathbf{x}_g = \mathbf{x}]$ for all such $(\mathbf{x}, n_1 \geq 2, n_2)$ in the support of $(\mathbf{x}_g, n_{g_1}, n_{g_2})$ (it is not identified when $n_1 = 1$ nor when $\mathbf{x} = c\mathbf{1}_{n_1}$ for some constant c because in these cases $\bar{y}_i = \bar{x}_i = 0$). We obtain

$$\mathbb{E}[\pi(n_{g_0}) | n_{g_1} = n_1, n_{g_2} = n_2, \mathbf{x}_g = \mathbf{x}] = \psi \pi(n_1) + (1 - \psi) \pi(n_2) \quad (7.7)$$

We now consider the cases $\psi = 0$, $\psi = 1$ and $\psi \in (0, 1)$ separately. If $\psi = 0$ then (γ, δ, β) are identified because $\gamma\beta + \delta \neq 0$ and n_{g_2} has at least three points in the support of its distribution. If $\psi = 1$ then (γ, δ, β) are identified because $\gamma\beta + \delta \neq 0$ and n_{g_1} has at least three points in the support of its distribution. The rest of the proof now considers $\psi \in (0, 1)$. By standard arguments for two-way bi-partite fixed effects models (e.g. Abowd et al. (1999)), equation (7.7) identifies $\mu(n_1) = \psi \pi(n_1) + c$ and $\mu(n_2) = (1 - \psi) \pi(n_2) - c$ for every pair (n_1, n_2) corresponding to two vertices in a connected component of $\mathcal{G}(n_{g_1}, n_{g_2})$, where c is an unknown constant.

Consider a connected component of $\mathcal{G}(n_{g_1}, n_{g_2})$ with at least three vertices corresponding to the support of n_{g_1} . If such a component does not exist we can instead consider a connected component of $\mathcal{G}(n_{g_1}, n_{g_2})$ with at least three vertices corresponding to the support of n_{g_2} . One of these two exists by assumption, so without loss of generality we present the first. Let the vertices in this connected component respectively correspond to \mathcal{N}_1 and \mathcal{N}_2 , where \mathcal{N}_1 is a subset of the support of n_{g_1} and \mathcal{N}_2 is a subset of the support of n_{g_2} . Then it must be that $|\mathcal{N}_1| \geq 3$ and $|\mathcal{N}_2| \geq 1$ (the

latter is because $\mathcal{G}(n_{g_1}, n_{g_2})$ is bi-partite). So there exist $2 \leq n_{1,1} < n_{1,2} < n_{1,3}$ and $2 \leq n_2$ such that $\{n_{1,1}, n_{1,2}, n_{1,3}\} \subseteq \mathcal{N}_1$ and $\{n_2\} \subseteq \mathcal{N}_2$. Moreover, $\mu(n)$ is identified for $n \in \{n_{1,1}, n_{1,2}, n_{1,3}, n_2\}$. For $i \in \{1, 2, 3\}$, denote

$$\Delta_i = \mu(n_{1,i}) + \mu(n_2) = \frac{\gamma(n_2 - 1) - \delta}{n_2 - 1 + \beta} + \frac{\psi(\gamma\beta + \delta)(n_{1,i} - n_2)}{(n_{1,i} - 1 + \beta)(n_2 - 1 + \beta)} \quad (7.8)$$

Then after some algebra, for $j \in \{1, 2, 3\}$ we have

$$\Delta_i - \Delta_j = \frac{\psi(\gamma\beta + \delta)(n_{1,i} - n_{1,j})}{(n_{1,i} - 1 + \beta)(n_{1,j} - 1 + \beta)} \quad (7.9)$$

Now since $\psi \neq 0$ and $\delta + \beta\gamma \neq 0$,

$$\Delta \equiv \frac{\Delta_1 - \Delta_2}{\Delta_1 - \Delta_3} = \frac{(n_{1,1} - n_{1,2})(n_{1,3} - 1 + \beta)}{(n_{1,1} - n_{1,3})(n_{1,2} - 1 + \beta)} \quad (7.10)$$

Since Δ is identified, β is identified by (7.10) if $\Delta(n_{1,1} - n_{1,3}) \neq n_{1,1} - n_{1,2}$. Suppose that $\Delta(n_{1,1} - n_{1,3}) = n_{1,1} - n_{1,2}$. Then (7.10) implies $n_{1,2} = n_{1,3}$, but here we have $n_{1,2} < n_{1,3}$, a contradiction. So β is identified. Since β is identified, $\psi(\gamma\beta + \delta)$ is identified from (7.9), hence γ and δ are identified from (7.8). Finally, ψ is identified because $\psi(\gamma\beta + \delta)$ and (γ, δ, β) are identified and $\gamma\beta + \delta \neq 0$. Notice also that one can replace ψ with $\psi(\mathbf{x})$ and $\mathcal{G}(n_{g_1}, n_{g_2})$ with $\mathcal{G}(n_{g_1}, n_{g_2} | \mathbf{x}_g = \mathbf{x})$ everywhere in the above proof, hence we can also allow for $\psi = \psi(\mathbf{x}_g)$.

Table 1: Monte Carlo Results: Missing data

Contextual effect only ($\beta = 0$ imposed), $N \approx M = 8000$									
ρ		Miss-specified		Known		MCAR		MCAR-P	
		Mean	RMSE	Mean	RMSE	Mean	RMSE	Mean	RMSE
1	δ	0.502	0.084	0.502	0.084	0.502	0.084	0.502	0.084
	γ	1.002	0.06	1.002	0.06	1.002	0.06	1.002	0.06
0.9	δ	0.428	0.112	0.5	0.087	0.501	0.1	0.501	0.1
	γ	0.971	0.071	1	0.063	1.001	0.071	1.001	0.071
0.7	δ	0.329	0.197	0.498	0.097	0.504	0.152	0.501	0.149
	γ	0.931	0.105	0.999	0.067	1.002	0.099	1.001	0.098
0.5	δ	0.257	0.271	0.501	0.112	0.507	0.286	0.491	0.234
	γ	0.9	0.146	1.001	0.078	1.001	0.168	0.994	0.149
0.3	δ	0.205	0.345	0.497	0.14	0.707	1.538	0.489	0.451
	γ	0.873	0.21	0.996	0.096	1.152	2.362	0.986	0.269
Contextual effect only ($\beta = 0$ imposed), $N \approx M = 1600$									
1	δ	0.498	0.184	0.498	0.184	0.498	0.184	0.498	0.184
	γ	0.999	0.132	0.999	0.132	0.999	0.132	0.999	0.132
0.9	δ	0.419	0.221	0.496	0.207	0.492	0.242	0.491	0.241
	γ	0.964	0.157	0.997	0.146	0.994	0.167	0.994	0.166
0.7	δ	0.328	0.28	0.502	0.213	0.517	0.365	0.506	0.345
	γ	0.929	0.191	1	0.146	1.006	0.233	1.001	0.225
0.5	δ	0.259	0.362	0.486	0.247	0.691	1.303	0.52	0.561
	γ	0.903	0.256	0.993	0.171	1.083	0.643	1.007	0.344
Contextual and endogenous effects, $N \approx M = 8000$									
1	β	0.095	0.772	0.095	0.772	0.095	0.772	0.095	0.772
	δ	0.471	0.245	0.471	0.245	0.471	0.245	0.471	0.245
	γ	1.02	0.182	1.02	0.182	1.02	0.182	1.02	0.182
0.9	β	0.146	0.845	0.087	0.784	0.096	0.831	0.096	0.832
	δ	0.379	0.333	0.476	0.243	0.473	0.267	0.472	0.267
	γ	1	0.178	1.019	0.189	1.021	0.199	1.021	0.199
0.7	β	0.116	0.891	0.117	0.799	0.044	0.875	0.045	0.874
	δ	0.277	0.474	0.463	0.261	0.482	0.319	0.48	0.319
	γ	0.948	0.17	1.023	0.19	1.006	0.218	1.005	0.217
0.5	β	0.06	0.93	0.035	0.811	-0.001	0.914	0.004	0.912
	δ	0.234	0.561	0.492	0.265	0.5	0.399	0.492	0.387
	γ	0.913	0.19	1.009	0.199	0.998	0.262	0.996	0.258
0.3	β	0	0.938	0.048	0.849	0.015	0.905	0.004	0.908
	δ	0.209	0.644	0.482	0.3	0.568	1.457	0.459	0.597
	γ	0.876	0.241	1.005	0.208	1.093	2.355	0.972	0.345
Contextual and endogenous effects, $N \approx M = 1600$									
1	β	0.066	0.888	0.066	0.888	0.066	0.888	0.066	0.888
	δ	0.481	0.343	0.481	0.343	0.481	0.343	0.481	0.343
	γ	1.015	0.245	1.015	0.245	1.015	0.245	1.015	0.245
0.9	β	0.073	0.932	0.062	0.909	0.053	0.926	0.054	0.926
	δ	0.404	0.426	0.482	0.371	0.484	0.402	0.483	0.402
	γ	0.985	0.253	1.013	0.261	1.01	0.28	1.01	0.279
0.7	β	0.04	0.947	0.025	0.91	0.01	0.939	0.014	0.941
	δ	0.321	0.53	0.506	0.362	0.517	0.511	0.507	0.503
	γ	0.944	0.26	1.013	0.269	1.013	0.341	1.01	0.335
0.5	β	0.029	0.958	0.014	0.913	0.011	0.944	0.018	0.944
	δ	0.256	0.634	0.489	0.413	0.605	1.083	0.51	0.711
	γ	0.919	0.317	1.001	0.274	1.052	0.585	1.013	0.453

Notes: Based on 1000 replications and reported to 3 decimal places. ‘Miss-specified’ refers to the NLS estimator obtained by treating the observed groups as if they were the groups (i.e. based on the moment $E[\bar{y}_i | n_g, \mathbf{x}_g] = \bar{x}_i \pi(n_g)$). ‘Known’ is the NLS estimator under Assumption 2 (i.e. based on the moment $E[\bar{y}_i | n_{g_0}, \mathbf{x}_g] = \bar{x}_i \pi(n_{g_0})$). ‘MCAR’ is the GMM estimator under Assumption 3. ‘MCAR-P’ is the GMM estimator under Assumption 3 imposing the parametric restriction $n_{g_0} - 2 \sim \text{Binomial}(2, \omega)$. MCAR and MCAR-P are based on the upper bound on the support of n_{g_0} being known to be $\bar{n} = 4$. Results are presented for (β, δ, γ) only. Their true values are $(0, 0.5, 1)$. Results for ρ and ω are available on request.

Table 2: Monte Carlo Results: Uncertain groups

Contextual effect only ($\beta = 0$ imposed), $N = 8000$									
ψ		Room		Floor		Known		Uncertain	
		Mean	RMSE	Mean	RMSE	Mean	RMSE	Mean	RMSE
0.8	δ	0.41	0.123	0.501	0.135	0.496	0.066	0.516	0.131
	γ	0.992	0.062	1.001	0.034	0.999	0.04	1.002	0.062
0.6	δ	0.321	0.198	0.498	0.135	0.5	0.061	0.504	0.142
	γ	0.983	0.064	1	0.032	1	0.033	1.001	0.063
0.4	δ	0.232	0.281	0.497	0.133	0.5	0.064	0.501	0.145
	γ	0.973	0.067	1	0.031	1	0.028	1.001	0.063
0.2	δ	0.144	0.366	0.499	0.129	0.497	0.08	0.503	0.145
	γ	0.964	0.071	1	0.028	1	0.026	1.004	0.058
Contextual effect only ($\beta = 0$ imposed), $N = 1600$									
0.8	δ	0.402	0.208	0.494	0.333	0.495	0.14	0.564	0.29
	γ	0.983	0.132	0.997	0.077	0.997	0.088	1	0.135
0.6	δ	0.313	0.263	0.503	0.327	0.493	0.137	0.538	0.31
	γ	0.975	0.134	1.001	0.075	0.996	0.073	0.999	0.135
0.4	δ	0.226	0.331	0.505	0.319	0.492	0.146	0.516	0.332
	γ	0.966	0.136	1	0.071	0.997	0.063	0.999	0.134
0.2	δ	0.134	0.409	0.507	0.308	0.499	0.174	0.503	0.357
	γ	0.955	0.139	1.001	0.065	0.999	0.058	1.006	0.13
Contextual and endogenous effects, $N = 8000$									
0.8	β	0.041	0.814	0.127	0.683	0.054	0.46	0.055	0.799
	δ	0.392	0.36	0.429	0.384	0.466	0.214	0.492	0.302
	γ	0.998	0.165	1.007	0.051	1.001	0.064	1.01	0.167
0.6	β	0.039	0.848	0.112	0.663	0.07	0.461	0.099	0.8
	δ	0.301	0.495	0.431	0.385	0.462	0.227	0.458	0.354
	γ	0.989	0.142	1.004	0.049	1.004	0.049	1.013	0.143
0.4	β	0.042	0.873	0.115	0.679	0.072	0.491	0.106	0.769
	δ	0.205	0.64	0.427	0.394	0.458	0.256	0.446	0.379
	γ	0.978	0.119	1.004	0.048	1.003	0.043	1.01	0.117
0.2	β	0.032	0.881	0.119	0.653	0.106	0.561	0.123	0.724
	δ	0.123	0.772	0.422	0.391	0.436	0.303	0.433	0.393
	γ	0.969	0.099	1.004	0.044	1.004	0.042	1.011	0.093
Contextual and endogenous effects, $N = 1600$									
0.8	β	0.038	0.913	0.067	0.841	0.097	0.709	0.019	0.896
	δ	0.393	0.449	0.465	0.61	0.444	0.343	0.559	0.428
	γ	0.995	0.228	1.001	0.098	1.002	0.123	1.005	0.239
0.6	β	0.024	0.927	0.088	0.833	0.065	0.697	0.051	0.9
	δ	0.311	0.568	0.458	0.604	0.455	0.358	0.522	0.485
	γ	0.986	0.209	1.006	0.096	0.998	0.096	1.009	0.22
0.4	β	0.019	0.94	0.096	0.833	0.107	0.716	0.045	0.888
	δ	0.223	0.7	0.457	0.579	0.435	0.386	0.493	0.55
	γ	0.974	0.192	1.006	0.091	1.001	0.081	1.005	0.198
0.2	β	-0.004	0.94	0.091	0.825	0.078	0.754	0.06	0.863
	δ	0.147	0.826	0.456	0.57	0.451	0.429	0.467	0.606
	γ	0.962	0.177	1.004	0.085	1.001	0.075	1.011	0.176

Notes: Based on 1000 replications and reported to 3 decimal places. ‘Room’ refers to the NLS estimator obtained by treating the observed group 1 as if they were the groups (i.e. based on the moment $\mathbb{E}[\bar{y}_i | n_{g_1}, n_{g_2}, \mathbf{x}_g] = \bar{x}_i \pi(n_{g_1})$). ‘Floor’ refers to the NLS estimator obtained by treating the observed group 2 as if they were the groups (i.e. based on the moment $\mathbb{E}[\bar{y}_i | n_{g_1}, n_{g_2}, \mathbf{x}_g] = \bar{x}_i \pi(n_{g_2})$). ‘Known’ is the NLS estimator under Assumption 2 (i.e. based on the moment $\mathbb{E}[\bar{y}_i | n_{g_0}, \mathbf{x}_g] = \bar{x}_i \pi(n_{g_0})$). ‘Uncertain’ is the NLS estimator based on $\mathbb{E}[\bar{y}_i | n_{g_1}, n_{g_2}, \mathbf{x}_g] = \bar{x}_i [\psi \pi(n_{g_1}) + (1 - \psi) \pi(n_{g_2})]$. Results are presented for (β, δ, γ) only. Their true values are $(0, 0.35, 1)$. Results for ψ are available on request.

Table 3: Monte Carlo Results: Uncertain groups (FE)

Contextual effect only ($\beta = 0$ imposed), $N = 8000$									
ψ		Room		Floor		Known		Uncertain	
		Mean	RMSE	Mean	RMSE	Mean	RMSE	Mean	RMSE
0.8	δ	0.41	0.123	0.847	0.375	0.496	0.066	0.516	0.131
	γ	0.992	0.062	1.3	0.303	0.999	0.04	1.002	0.062
0.6	δ	0.321	0.198	0.756	0.292	0.5	0.061	0.504	0.142
	γ	0.983	0.064	1.224	0.227	1	0.033	1.001	0.063
0.4	δ	0.232	0.281	0.668	0.218	0.5	0.064	0.501	0.145
	γ	0.973	0.067	1.149	0.153	1	0.028	1.001	0.063
0.2	δ	0.144	0.366	0.584	0.156	0.497	0.08	0.503	0.145
	γ	0.964	0.071	1.075	0.081	1	0.026	1.004	0.058
Contextual effect only ($\beta = 0$ imposed), $N = 1600$									
0.8	δ	0.402	0.208	0.841	0.488	0.495	0.14	0.564	0.29
	γ	0.983	0.132	1.297	0.309	0.997	0.088	1	0.135
0.6	δ	0.313	0.263	0.762	0.43	0.493	0.137	0.538	0.31
	γ	0.975	0.134	1.225	0.24	0.996	0.073	0.999	0.135
0.4	δ	0.226	0.331	0.679	0.375	0.492	0.146	0.516	0.332
	γ	0.966	0.136	1.151	0.169	0.997	0.063	0.999	0.134
0.2	δ	0.134	0.409	0.596	0.328	0.499	0.174	0.503	0.357
	γ	0.955	0.139	1.076	0.102	0.999	0.058	1.006	0.13
Contextual and endogenous effects, $N = 8000$									
0.8	β	0.041	0.814	0.224	0.558	0.054	0.46	0.055	0.799
	δ	0.392	0.36	0.725	0.375	0.466	0.214	0.492	0.302
	γ	0.998	0.165	1.318	0.324	1.001	0.064	1.01	0.167
0.6	β	0.039	0.848	0.198	0.565	0.07	0.461	0.099	0.8
	δ	0.301	0.495	0.645	0.347	0.462	0.227	0.458	0.354
	γ	0.989	0.142	1.237	0.244	1.004	0.049	1.013	0.143
0.4	β	0.042	0.873	0.17	0.602	0.072	0.491	0.106	0.769
	δ	0.205	0.64	0.571	0.347	0.458	0.256	0.446	0.379
	γ	0.978	0.119	1.159	0.168	1.003	0.043	1.01	0.117
0.2	β	0.032	0.881	0.152	0.612	0.106	0.561	0.123	0.724
	δ	0.123	0.772	0.492	0.36	0.436	0.303	0.433	0.393
	γ	0.969	0.099	1.082	0.094	1.004	0.042	1.011	0.093
Contextual and endogenous effects, $N = 1600$									
0.8	β	0.038	0.913	0.138	0.747	0.097	0.709	0.019	0.896
	δ	0.393	0.449	0.785	0.62	0.444	0.343	0.559	0.428
	γ	0.995	0.228	1.31	0.331	1.002	0.123	1.005	0.239
0.6	β	0.024	0.927	0.129	0.772	0.065	0.697	0.051	0.9
	δ	0.311	0.568	0.715	0.582	0.455	0.358	0.522	0.485
	γ	0.986	0.209	1.237	0.263	0.998	0.096	1.009	0.22
0.4	β	0.019	0.94	0.138	0.789	0.107	0.716	0.045	0.888
	δ	0.223	0.7	0.617	0.565	0.435	0.386	0.493	0.55
	γ	0.974	0.192	1.161	0.191	1.001	0.081	1.005	0.198
0.2	β	-0.004	0.94	0.136	0.804	0.078	0.754	0.06	0.863
	δ	0.147	0.826	0.521	0.563	0.451	0.429	0.467	0.606
	γ	0.962	0.177	1.083	0.123	1.001	0.075	1.011	0.176

Notes: Based on 1000 replications and reported to 3 decimal places. ‘Room’ refers to the NLS estimator obtained by treating the observed group 1 as if they were the groups (i.e. based on the moment $\mathbb{E}[\bar{y}_i | n_{g_1}, n_{g_2}, \mathbf{x}_g] = \bar{x}_i \pi(n_{g_1})$). ‘Floor’ refers to the NLS estimator obtained by treating the observed group 2 as if they were the groups (i.e. based on the moment $\mathbb{E}[\bar{y}_i | n_{g_1}, n_{g_2}, \mathbf{x}_g] = \bar{x}_i \pi(n_{g_2})$). ‘Known’ is the NLS estimator under Assumption 2 (i.e. based on the moment $\mathbb{E}[\bar{y}_i | n_{g_0}, \mathbf{x}_g] = \bar{x}_i \pi(n_{g_0})$). ‘Uncertain’ is the NLS estimator based on $\mathbb{E}[\bar{y}_i | n_{g_1}, n_{g_2}, \mathbf{x}_g] = \bar{x}_i [\psi \pi(n_{g_1}) + (1 - \psi) \pi(n_{g_2})]$. Results are presented for (β, δ, γ) only. Their true values are $(0, 0.35, 1)$. Results for ψ are available on request.