

Quadratic-stretch elasticity

E. Vitral and J. A. Hanna

*Department of Mechanical Engineering, University of Nevada,
1664 N. Virginia St. (0312), Reno, NV 89557-0312, U.S.A.*

(Dated: July 12, 2021)

A nonlinear small-strain elastic theory is constructed from a systematic expansion in Biot strains, truncated at quadratic order. The primary motivation is the desire for a clean separation between stretching and bending energies for shells, which appears to arise only from reduction of a bulk energy of this type. An approximation of isotropic invariants, bypassing the solution of a quartic equation or computation of tensor square roots, allows stretches, rotations, stresses, and balance laws to be written in terms of derivatives of position. Two-field formulations are also presented. Extensions to anisotropic theories are briefly discussed.

I. INTRODUCTION

In a finite deformation, described in terms of positions or displacements, information about strains is polluted with information about rotations. In statics, the strains are important but the rotations are irrelevant, and any hyperelastic energy and resulting constitutive relations must respect this.

A common solution is to symmetrize the deformation by “squaring” it with its transpose to obtain particular objective strain measures, either the right or left Cauchy-Green deformation tensors, the respective squares of the right and left stretch tensors. This has the additional advantages that these measures, and the closely related Green-Lagrange or Euler-Almansi strain tensors, may be written straightforwardly using derivatives of position, and interpreted quite intuitively in terms of metric tensor components in referential and present configurations of the body. Many other objective strain measures may be constructed as well, and a general nonlinear elastic energy can be written in terms of any of these. In the continuum mechanics literature, this is often the end of the discussion. One finds results derived for general stored energy functions, as well as a variety of phenomenological models of nonlinearly elastic materials, often expressed in terms of an incomplete collection of powers of principal stretches, for example Neo-Hookean, Mooney-Rivlin, or Ogden materials [1].

However, in the physics literature, there is a desire to construct field theories by a systematic expansion in a small quantity. In linear elasticity, this would be the displacement, but in a nonlinear elasticity theory the choice of appropriate field variables is not immediately clear, and constitutes a fundamental unanswered question. Further motivation and clues for this question may be found in the study of thin shells, where it is often the case that strains are small but deformations are far from linear. Due to its ease of representation and interpretation, the Green-Lagrange strain has been widely adopted as the small field for expansion of elastic energies by soft condensed matter physicists [2–6]. However, the squaring of the deformation has the unfortunate consequence that the leading-order terms in such energies are quartic in stretch. As has been rediscovered several times, surface elastic energies derived from dimensional reduction of such bulk elastic energies have undesirable features, including a mixing of stretching and bending contents [7–11]. This stands in unsatisfying contrast to simple direct theories [12–14] that employ particular natural kinematic measures of surface stretching and bending to obtain separate and distinct energies. Selecting a simple bulk theory that corresponds to such simple direct theories requires a careful construction of energy quadratic in stretch, but this process has only been performed for one-dimensional and axisymmetric bodies [7, 8, 10, 15–17], and indeed appropriate bending measures have only been defined for such special cases¹ of thin bodies [8, 10, 18, 19]. Finally, quadratic-stretch elastic energies arise naturally in bead-spring descriptions of soft matter meso-structures such as biological membranes [20, 21].

It is our intent in this paper to lay the groundwork for development of theories applicable to general shells, by first considering bulk elasticity from the perspective of a comprehensive expansion in stretches, truncated at quadratic order. More precisely, we make use of the Biot strain, the deviation of the right stretch from the identity, as our small quantity. As will become apparent below, this choice classifies our theory as “referential”, but there is no reason why a complementary left-handed theory making use of tensors in the present configuration could not be constructed. We also leave the analysis of reduced energies for thin bodies to subsequent work. A significant challenge in constructing a stretch-based theory is that stretch has an indirect dependence on position and its derivatives, requiring simultaneous consideration of an additional field, the rotation. Our synthesis leads us to connect several areas in theoretical and computational elasticity, including the kinematics of stretch and rotation, variational principles with auxiliary

¹ E. G. Virga has recently shared with us an unpublished note in which he derived more general bending measures.

fields, and relations between isotropic invariants of different strain measures. We employ convected coordinates and a mixture of referential and present bases, revealing interesting relationships between components of various tensors. Our Biot-quadratic energy is surprisingly rare in the nonlinear elasticity literature, although its roots go back to early work by Lurie [22] and John [23]. Inspired by the works of Atluri and Murakawa [24], Wiśniewski [25], and Merlini [26], we propose a variational principle in terms of position, Biot strain, and rotation fields, and show that constitutive stress-strain relations and balance equations can be written in terms of positions and stretches alone. A new result is that the neglect of terms cubic in Biot strain allows for an explicit representation of stretch without either taking tensor square roots or solving a quartic equation, leading to an approximate description of all fields in terms of positions alone.

Our kinematic treatment may have broader application to theories in which rotations must be tracked, such as models of nematic elastomers [27, 28] or tension-field theory [29]. And although it is not the primary motivation for this work, we anticipate that it may have value with regard to the systematic construction of general nonlinear small-strain field theories of elasticity. However, one disadvantage of the present method as it currently stands is that it is not immediately clear how to adapt it to incompatible elasticity, in which no stress-free reference configuration or corresponding set of basis vectors exists but nonetheless the idea of a reference metric is still useful and intuitive. We reserve this question for future work, and note that while the idea of an incompatible deformation gradient appears in previous works [30, 31] it is not clear how such an object would be constructed in practice.

An outline of the paper follows. Section II defines relevant tensorial objects and derives kinematic relations between them, their components, and their invariants. Small-strain approximations lead to algebraic expressions for invariants and explicit expressions for all the relevant tensors in terms of derivatives of position. Section III presents stresses, balance laws, and constitutive relations obtained from a mixed variational principle in terms of position, Biot strain, and rotation. In Section IV, a general isotropic energy quadratic in Biot strains is constructed. Further constitutive relations are derived, in particular one for Biot stress in terms of Biot strain and its invariants, and then specialized to this energy. Formulations are presented in terms of positions, positions and rotations, or positions and stretches. Extensions to anisotropic theories are sketched in Section V. Appendices A-D discuss alternate decompositions of the deformation gradient, the symmetry of a certain rotated stress, details of the energy variation, and the strain and stress of Bell that complement those of Biot.

II. KINEMATICS

A. Definitions and notation

We denote material coordinates as η^i and (noncovariant) material derivatives with respect to these as d_i . An elastic body \mathcal{B} with boundary $\partial\mathcal{B}$ has reference configuration $\mathbf{X}(\eta^i)$ and present configuration $\mathbf{x}(\eta^i)$, both in \mathbb{E}^3 . We define referential and present coordinate bases $\mathbf{G}_I = d_i\mathbf{X}$ and $\mathbf{g}_i = d_i\mathbf{x}$, reciprocal bases through the relations $\mathbf{G}^I \cdot \mathbf{G}_J = \delta_J^I$ and $\mathbf{g}^i \cdot \mathbf{g}_j = \delta_j^i$ using the Kronecker delta, and covariant and contravariant components of the corresponding metric tensors $G_{IJ} = \mathbf{G}_I \cdot \mathbf{G}_J$, $g_{ij} = \mathbf{g}_i \cdot \mathbf{g}_j$, $G^{IJ} = \mathbf{G}^I \cdot \mathbf{G}^J$, $g^{ij} = \mathbf{g}^i \cdot \mathbf{g}^j$. Both metric tensors correspond to the identity, $\mathbf{G} = \mathbf{g} = \mathbf{I}$. In all of these expressions, capitalization of some indices is just a reminder that these should be raised and lowered with components of the reference metric; summation ignores case. We denote covariant derivatives corresponding to the referential and present metrics as $\bar{\nabla}_I$ and ∇_i , respectively, and corresponding Christoffel symbols as $\bar{\Gamma}_{JK}^I$ and Γ_{jk}^i . We define referential and present gradients $\bar{\nabla}() = \bar{\nabla}_i() \mathbf{G}^I$ and $\nabla() = \nabla_i() \mathbf{g}^i$. The metric determinants $G = \det[G_{ij}]$ and $g = \det[g_{ij}]$, where $[\]$ denotes a matrix, are found in the referential and present volume forms $dV = \sqrt{G} \prod_i d\eta^i$ and $dv = \sqrt{g/G} dV$. Surface forms dA and da are defined analogously using two surface coordinates only.

B. Deformation measures, polar decomposition, and shifter

The deformation gradient $\bar{\nabla}\mathbf{x}$ takes the simple form $\mathbf{g}_i \mathbf{G}^I$ in material coordinates. While this is straightforwardly expressed purely in terms of derivatives of position, it is far from trivial to do the same for its rotationally-invariant stretching content.

Any non-singular second order tensor, such as the deformation gradient when $\sqrt{g/G} > 0$, admits unique right and left decompositions into an element of the special orthogonal group $\text{SO}(3)$ of rotations and an element of the set Sym^+ of positive-definite symmetric tensors. In terms of the rotation tensor $\mathbf{Q} \in \text{SO}(3)$ and the right \mathbf{U} or left \mathbf{V} stretch

tensors $\in \text{Sym}^+$,

$$\mathbf{g}_i \mathbf{G}^I = \mathbf{Q} \cdot \mathbf{U} = \mathbf{V} \cdot \mathbf{Q}. \quad (1)$$

This is the most commonly adopted decomposition in continuum mechanics, but in Appendix A we consider alternatives. The orthogonal rotation's transpose is also its inverse,

$$\mathbf{Q}^\top \cdot \mathbf{Q} = \mathbf{Q} \cdot \mathbf{Q}^\top = \mathbf{I}, \quad (2)$$

a fact used repeatedly in the sequel. The right and left Cauchy-Green deformation tensors are $\mathbf{C} = \mathbf{G}^I \mathbf{g}_i \cdot \mathbf{g}_j \mathbf{G}^J = g_{ij} \mathbf{G}^I \mathbf{G}^J$ and $\mathbf{B} = \mathbf{g}_i \mathbf{G}^I \cdot \mathbf{G}^J \mathbf{g}_j = G^{IJ} \mathbf{g}_i \mathbf{g}_j$, respectively. Further relationships include $\mathbf{C} = \mathbf{U}^2$, $\mathbf{B} = \mathbf{V}^2$, $\mathbf{U} = \mathbf{Q}^\top \cdot \mathbf{V} \cdot \mathbf{Q}$, $\mathbf{V} = \mathbf{Q} \cdot \mathbf{U} \cdot \mathbf{Q}^\top$, $\mathbf{C} = \mathbf{Q}^\top \cdot \mathbf{B} \cdot \mathbf{Q}$, and $\mathbf{B} = \mathbf{Q} \cdot \mathbf{C} \cdot \mathbf{Q}^\top$.

A natural way to represent the rotation and stretch tensors is

$$\mathbf{Q} = Q^i{}_J \mathbf{g}_i \mathbf{G}^J, \quad \mathbf{U} = U_{IJ} \mathbf{G}^I \mathbf{G}^J, \quad \mathbf{V} = V^{ij} \mathbf{g}_i \mathbf{g}_j, \quad (3)$$

such that \mathbf{Q} has the same mixed character as the deformation gradient, taking quantities in the reference configuration into the present configuration. From (1), we find that the components of rotation, stretches, and metrics are related,

$$Q^i{}_J G^{JK} U_{KL} = Q^i{}_J U^J{}_L = \delta_L^i = V^i{}_k Q^k{}_L = V^{ij} g_{jk} Q^k{}_L. \quad (4)$$

We further see that

$$Q^i{}_J g_{ik} Q^k{}_L = G_{JL}, \quad Q^i{}_J G^{JL} Q^k{}_L = g^{ik}, \quad (5)$$

which may be compared with

$$U_{IJ} G^{IL} U_{KL} = g_{jk} \quad \text{or} \quad U^I{}_J U^J{}_K = G^{IJ} g_{jk}. \quad (6)$$

Ericksen and Truesdell [32] describe this mixed-basis representation of rotation as ‘‘shifted’’. The components of the shifter [33] $\boldsymbol{\mu} = (\mathbf{g}^k \cdot \mathbf{G}_J) \mathbf{g}_k \mathbf{G}^J = \mu^k{}_J \mathbf{g}_k \mathbf{G}^J$ can be used to relate the mixed-basis components of rotation with their representation in either the reference or present configuration. The shifter can be thought of as the identity in a mixed-basis representation; the statement $\mathbf{Q} \cdot \boldsymbol{\mu} = \mathbf{Q} = \boldsymbol{\mu} \cdot \mathbf{Q}$ is, in components, $Q^i{}_k \mu^k{}_J = Q^i{}_J = \mu^i{}_L Q^L{}_J$. Invariants are computed along similar lines, for example $\text{Tr } \mathbf{Q} = \mathbf{Q} : \mathbf{I} = Q^i{}_J \mathbf{g}_i \mathbf{G}^J : \mu^k{}_L \mathbf{g}_k \mathbf{G}^L = Q^i{}_J \mu^i{}_J = Q^i{}_L \mu^k{}_L = Q^i{}_i = Q^L{}_L$.

C. Stretch and rotation

Due to the uniqueness of the square root of a symmetric positive-definite tensor [34], the right stretch \mathbf{U} can be obtained as the square root of the right Cauchy-Green deformation $\sqrt{\mathbf{C}}$, or similarly, the left stretch \mathbf{V} can be obtained as the square root of the left Cauchy-Green deformation $\sqrt{\mathbf{B}}$. However, this operation is inconvenient, and adds an additional layer of complexity, as these square roots, unlike \mathbf{C} and \mathbf{B} themselves, cannot be written in terms of derivatives of position. This has led many to explore other representations for \mathbf{U} and \mathbf{Q} that avoid the computation of tensor square roots, but instead make use of relationships between invariants of different strain tensors. While two-dimensional derivations such as that of Biot [35] are not too cumbersome, three-dimensional representations generally require another difficult step such as the selection of a root of a quartic equation [36–46].

Our starting point is Ting's relatively simple expression [38] for the right stretch \mathbf{U} in terms of \mathbf{C} and the invariants of \mathbf{U} . The three principal invariants $i_k^{\mathbf{U}}$ of \mathbf{U} are

$$\begin{aligned} i_1^{\mathbf{U}} &= \text{Tr } \mathbf{U} &= \lambda_1 + \lambda_2 + \lambda_3, \\ i_2^{\mathbf{U}} &= \frac{1}{2} \left[(\text{Tr } \mathbf{U})^2 - \text{Tr } (\mathbf{U}^2) \right] &= \lambda_1 \lambda_2 + \lambda_2 \lambda_3 + \lambda_1 \lambda_3, \\ i_3^{\mathbf{U}} &= \text{Det } \mathbf{U} &= \lambda_1 \lambda_2 \lambda_3, \end{aligned} \quad (7)$$

where the principal stretches λ_k are the eigenvalues of \mathbf{U} . Since \mathbf{U} is positive-definite, its eigenvalues and principal invariants are all positive. The stretch tensors \mathbf{U} and \mathbf{V} have the same eigenvalues and invariants. The Cauchy-Green strain tensors \mathbf{C} and \mathbf{B} also share eigenvalues, which are the squares λ_k^2 of those of \mathbf{U} , and invariants $i_k^{\mathbf{C}}$, constructed in an analogous manner to those of \mathbf{U} . Just like \mathbf{C} itself, all three $i_k^{\mathbf{C}}$ can be written in terms of derivatives of position.

In our notation, Ting's expression is

$$\mathbf{U} = (i_1^{\mathbf{U}} i_2^{\mathbf{U}} - i_3^{\mathbf{U}})^{-1} \left(i_1^{\mathbf{U}} i_3^{\mathbf{U}} G_{IJ} + \left[(i_1^{\mathbf{U}})^2 - i_2^{\mathbf{U}} \right] g_{ij} - g_{ik} G^{KL} g_{lj} \right) \mathbf{G}^I \mathbf{G}^J. \quad (8)$$

Similarly, the left stretch \mathbf{V} may be written

$$\mathbf{V} = (i_1^{\mathbf{U}} i_2^{\mathbf{U}} - i_3^{\mathbf{U}})^{-1} \left(i_1^{\mathbf{U}} i_3^{\mathbf{U}} g^{ij} + \left[(i_1^{\mathbf{U}})^2 - i_2^{\mathbf{U}} \right] G^{IJ} - G^{IK} g_{kl} G^{LJ} \right) \mathbf{g}_i \mathbf{g}_j. \quad (9)$$

The relations $\mathbf{Q} = \mathbf{Q}^{-\top} = (\mathbf{g}_i \mathbf{G}^I \cdot \mathbf{U})^{-\top} = \mathbf{g}^i \mathbf{G}_I \cdot \mathbf{U}$ yield the rotation

$$\mathbf{Q} = (i_1^{\mathbf{U}} i_2^{\mathbf{U}} - i_3^{\mathbf{U}})^{-1} \left(i_1^{\mathbf{U}} i_3^{\mathbf{U}} G_{KJ} g^{ki} + \left[(i_1^{\mathbf{U}})^2 - i_2^{\mathbf{U}} \right] \delta_j^i - G^{IK} g_{kj} \right) \mathbf{g}_i \mathbf{G}^J. \quad (10)$$

The issue now becomes how to write the invariants $i_k^{\mathbf{U}}$ of \mathbf{U} . The third invariant is simple, as $i_3^{\mathbf{U}} = \sqrt{i_3^{\mathbf{C}}} = \sqrt{g/G}$. However, to obtain the other two invariants requires finding the eigenvalues of \mathbf{C} , or solving a quartic equation for $i_1^{\mathbf{U}}$ obtained from the trace of (8) [36, 39].

In the following section, we show that neglect of terms higher order than quadratic in Biot strain allows for the computation of relevant strain invariants from a quadratic equation in which the single relevant root is easily identifiable. This leads to explicit approximate representations of stretch and rotation in terms of derivatives of position.

D. Quadratic-Biot theory: explicit approximate representation of stretch and rotation in terms of derivatives of position

The central quantity in our derivations will be the Biot strain

$$\mathbf{E}_B = \mathbf{U} - \mathbf{I}, \quad (11)$$

whose eigenvalues are $\Delta_k = \lambda_k - 1$ and whose principal invariants are related to those of the right stretch by

$$\begin{aligned} i_1^{\mathbf{E}_B} &= \Delta_1 + \Delta_2 + \Delta_3 &= i_1^{\mathbf{U}} - 3, \\ i_2^{\mathbf{E}_B} &= \Delta_1 \Delta_2 + \Delta_1 \Delta_3 + \Delta_2 \Delta_3 &= i_2^{\mathbf{U}} - 2i_1^{\mathbf{U}} + 3, \\ i_3^{\mathbf{E}_B} &= \Delta_1 \Delta_2 \Delta_3 &= i_3^{\mathbf{U}} - i_2^{\mathbf{U}} + i_1^{\mathbf{U}} - 1. \end{aligned} \quad (12)$$

The simplification we seek will come from neglect of the cubic invariant. Note that

$$i_3^{\mathbf{U}} = i_3^{\mathbf{E}_B} + i_2^{\mathbf{E}_B} + i_1^{\mathbf{E}_B} + 1, \quad (13)$$

and, as $\mathbf{E}_B^2 = \mathbf{C} - 2\mathbf{U} + \mathbf{I}$,

$$i_2^{\mathbf{E}_B} = \frac{1}{2} \left[(\text{Tr } \mathbf{E}_B)^2 - \text{Tr} (\mathbf{E}_B^2) \right] = \frac{1}{2} (i_1^{\mathbf{E}_B})^2 - \frac{1}{2} i_1^{\mathbf{C}} + i_1^{\mathbf{E}_B} + \frac{3}{2}. \quad (14)$$

These expressions may be combined into a quadratic equation for the first invariant of Biot strain,

$$(i_1^{\mathbf{E}_B})^2 + 4i_1^{\mathbf{E}_B} = i_1^{\mathbf{C}} + 2i_3^{\mathbf{U}} - 2i_3^{\mathbf{E}_B} - 5. \quad (15)$$

The only term on the right hand side of (15) that cannot be expressed in terms of derivatives of position is $i_3^{\mathbf{E}_B} = \text{Det}(\mathbf{U} - \mathbf{I})$, but this term is $O(\Delta^3)$ (and identically zero in the case of plane strain). Thus,

$$(i_1^{\mathbf{E}_B})^2 + 4i_1^{\mathbf{E}_B} = g_{ij} G^{IJ} + 2\sqrt{g/G} - 5 + O(i_3^{\mathbf{E}_B}), \quad (16)$$

and the only relevant root is that which connects with the solution for vanishing deformation, for which $i_1^{\mathbf{E}_B}$ and the right hand side of (16) are both zero, namely

$$i_1^{\mathbf{E}_B} = -2 + \sqrt{g_{ij} G^{IJ} + 2\sqrt{g/G} - 1} + O(i_3^{\mathbf{E}_B}). \quad (17)$$

This result allows us to approximately express the three invariants appearing in expressions for the stretches (8-9) and rotation (10) purely in terms of derivatives of position,

$$\begin{aligned} i_1^{\mathbf{U}} &= 1 + \sqrt{g_{ij}G^{IJ} + 2\sqrt{g/G} - 1} + O(i_3^{\mathbf{E}_B}), \\ i_2^{\mathbf{U}} &= \sqrt{g_{ij}G^{IJ} + 2\sqrt{g/G} - 1} + \sqrt{g/G} + O(i_3^{\mathbf{E}_B}), \\ i_3^{\mathbf{U}} &= \sqrt{g/G}. \end{aligned} \quad (18)$$

Note that neglect of $O(\Delta^3)$ terms makes these three quantities linearly dependent, as may have been gleaned from (12). The two quantities $g_{ij}G^{IJ}$ and g/G appearing in (18) are the first and third invariants of \mathbf{C} . Using (12) and (18) and collecting results,

$$\begin{aligned} i_1^{\mathbf{E}_B} &= -2 + \sqrt{g_{ij}G^{IJ} + 2\sqrt{g/G} - 1} + O(i_3^{\mathbf{E}_B}), \\ i_2^{\mathbf{E}_B} &= 1 - \sqrt{g_{ij}G^{IJ} + 2\sqrt{g/G} - 1} + \sqrt{g/G} + O(i_3^{\mathbf{E}_B}), \\ i_3^{\mathbf{E}_B} &= O(\Delta_1\Delta_2\Delta_3). \end{aligned} \quad (19)$$

III. CONSERVATION LAWS AND CONSTITUTIVE RELATIONS

A. Stresses

The force per unit area in the reference and present configurations is respectively given by the tractions \mathbf{T} and \mathbf{t} , which are related to the first Piola-Kirchhoff and Cauchy stresses \mathbf{P} and $\boldsymbol{\sigma}$,

$$\mathbf{t} da = \mathbf{n} \cdot \boldsymbol{\sigma} da = \boldsymbol{\sigma}^\top \cdot \mathbf{n} da = \sqrt{g/G} \boldsymbol{\sigma}^\top \cdot \mathbf{g}^i \mathbf{G}_I \cdot \mathbf{N} dA = \mathbf{P} \cdot \mathbf{N} dA = \mathbf{N} \cdot \mathbf{P}^\top dA = \mathbf{T} dA, \quad (20)$$

where \mathbf{N} and \mathbf{n} are the referential and present unit normals. The stresses are related by $\mathbf{P} = \sqrt{g/G} \boldsymbol{\sigma}^\top \cdot \mathbf{g}^i \mathbf{G}_I$. Because $\boldsymbol{\sigma}$ is often assumed symmetric, there is some inconsistency in the literature as to the definition of \mathbf{P} . For instance, both Lurie [22] and Atluri [47] define \mathbf{P} as the transpose of our definition. While \mathbf{P} is generally not symmetric, $\mathbf{P} \cdot \mathbf{G}^I \mathbf{g}_i$ is symmetric whenever $\boldsymbol{\sigma}$ is symmetric. In Appendix B, we further show that $\mathbf{P}^\top \cdot \mathbf{Q}$ is always symmetric for an isotropic material with symmetric $\boldsymbol{\sigma}$.

B. Variational framework

We adopt, with slight modifications, the variational principle of Atluri and Murakawa [24, 47, 48] to write a referential stored energy (density) \mathcal{W} in terms of Biot strain, with position derivatives, rotation, and right stretch linked by a multiplier so that auxiliary fields may be varied independently. The energy is

$$\mathcal{E}(\mathbf{x}, \mathbf{Q}, \mathbf{E}_B; \mathbf{P}) = \int_B [\mathcal{W}(\mathbf{E}_B) + \mathbf{P} : (\mathbf{g}_i \mathbf{G}^I - \mathbf{Q} \cdot \mathbf{U})] dV. \quad (21)$$

Our choice of symbol anticipates the identification of the multiplier \mathbf{P} with the first Piola-Kirchhoff stress. Three expressions are obtained from stationarity of (21) under variation of $\mathbf{x} \in \mathbb{E}^3$, $\mathbf{Q} \in \text{SO}(3)$, $\mathbf{U} \in \text{Sym}^+$, noting that $\delta \mathbf{U} = \delta \mathbf{E}_B$. Details of the variation, and the derivation of the equations and single boundary condition on stress, are reserved for Appendix C. We obtain the balances of linear and angular momentum (22-23) and the constitutive relation (24) for the Biot stress $\boldsymbol{\Sigma}_{\text{Biot}}$,

$$\bar{\nabla} \cdot \mathbf{P}^\top = \mathbf{0}, \quad (22)$$

$$\mathbf{g}_i \mathbf{G}^I \cdot \mathbf{P}^\top = \mathbf{P} \cdot \mathbf{G}^J \mathbf{g}_j, \quad (23)$$

$$\boldsymbol{\Sigma}_{\text{Biot}} \equiv \frac{\partial \mathcal{W}}{\partial \mathbf{E}_B} = \frac{1}{2} (\mathbf{P}^\top \cdot \mathbf{Q} + \mathbf{Q}^\top \cdot \mathbf{P}), \quad (24)$$

to be used alongside the compatibility constraint (1) and the restriction on rotation (2). An equivalent set of equations was presented by Wiśniewski [25]. One possible set of component forms is

$$\bar{\nabla}_J P^{iJ} = d_J P^{iJ} + \Gamma_{kj}^i P^{kJ} + \bar{\Gamma}_{JL}^J P^{iL} = 0, \quad (25)$$

$$P^{iJ} = P^{jI}, \quad (26)$$

$$(\boldsymbol{\Sigma}_{\text{Biot}})^{IJ} = \frac{1}{2} (P^{kI} Q_k^J + Q_k^I P^{kJ}), \quad (27)$$

to accompany (4) and (5), where of course $Q_i^J = g_{ik} Q_k^L G^{LJ}$ and so on.

We have defined the Biot stress $\boldsymbol{\Sigma}_{\text{Biot}}$ in a natural way, as the derivative of the stored energy \mathcal{W} with respect to the Biot strain \mathbf{E}_B , rather than the more traditional [1] derivative with respect to the right stretch \mathbf{U} . The Biot stress is the symmetric part of $\mathbf{P}^\top \cdot \mathbf{Q}$. The unsymmetrized quantity has a particular interpretation [1], which can be seen from the relation $\mathbf{N} \cdot \mathbf{P}^\top \cdot \mathbf{Q} = \mathbf{T} \cdot \mathbf{Q}$, indicating that its associated local load is a rotation of the referential traction \mathbf{T} . In Appendix D we discuss the Bell stress and strain, which would appear in a complementary formulation based on the left stretch \mathbf{V} .

When \mathcal{W} is specified, the constitutive relation (24) and, if necessary, the angular momentum balance (23), will provide the first Piola-Kirchhoff stress \mathbf{P} in terms of \mathbf{Q} and \mathbf{U} . Then the unknowns in the remaining equations are the position \mathbf{x} , rotation \mathbf{Q} , and stretch \mathbf{U} . In Section IV C, different formulations will be presented in terms of one or two out of three of these quantities.

IV. ISOTROPIC QUADRATIC-BIOT THEORY

In this section we consider an elastic energy quadratic in the isotropic invariants of right stretch or, equivalently, Biot strain, constitutive relations for this specific energy, and forms of the field equations in terms of positions alone or combined with either rotations or stretches.

A. Energy

A quadratic-Biot energy may be identified with that of John's two-dimensional "harmonic" materials [23] and Lurie's "semilinear" materials [22]. This energy was considered by Neff and co-workers in [49]. It also appears in Ozenda and Virga [11], who however lose its desirable qualities by subsequently expanding the right stretch in powers of the Green-Lagrange strain. Carroll [50] introduced more general classes of energies defined by functions of stretch invariants. Another setting in which stretch-based energies have been explored is that of theories with independent rotational degrees of freedom, introduced either for computational convenience or to describe micropolar continua [49, 51–59].

Consider small Biot strains around a stress-free reference configuration at which the three isotropic invariants $i_1^{\mathbf{E}_B} = i_2^{\mathbf{E}_B} = i_3^{\mathbf{E}_B} = 0$. As the third invariant is of cubic order, a general isotropic quadratic energy is of the form

$$\mathcal{W}(i_1^{\mathbf{E}_B}, i_2^{\mathbf{E}_B}) = c_1 (i_1^{\mathbf{E}_B})^2 + c_2 i_2^{\mathbf{E}_B}, \quad (28)$$

$$= c_1 \left(-2 + \sqrt{g_{ij} G^{IJ} + 2\sqrt{g/G} - 1} \right)^2 + c_2 \left(1 - \sqrt{g_{ij} G^{IJ} + 2\sqrt{g/G} - 1} + \sqrt{g/G} \right) + O(i_3^{\mathbf{E}_B}), \quad (29)$$

$$= \left(c_1 + \frac{c_2}{2} \right) (\text{Tr } \mathbf{E}_B)^2 - \frac{c_2}{2} \text{Tr} (\mathbf{E}_B^2), \quad (30)$$

where $c_1 \geq -c_2/2$ and $c_2 \leq 0$ are constant material parameters. In terms of an elastic tensor, with $\mathcal{W} = \frac{1}{2} \mathbf{E}_B : \mathbf{A} : \mathbf{E}_B$, we have $\frac{1}{2} A^{IJKL} = (c_1 + \frac{c_2}{2}) G^{IJ} G^{KL} - \frac{c_2}{4} (G^{IK} G^{JL} + G^{IL} G^{JK})$. The approximate form (29) of the energy suggests a definition in terms of the alternate invariants $i_1^{\mathbf{E}_B} = \sqrt{g_{ij} G^{IJ} + 2\sqrt{g/G} - 1} - 2$ and $i_2^{\mathbf{E}_B} + i_1^{\mathbf{E}_B} = \sqrt{g/G} - 1$. An incompressible ($\sqrt{g/G} = 1$) form of the quadratic-Biot energy would depend only on the first of these, in the simplified form $\sqrt{g_{ij} G^{IJ} + 1} - 2$. It is illustrative to compare this with the incompressible neo-Hookean energy

$$\mathcal{W}_{\text{NH}} = c_{\text{NH}} (-3 + g_{ij} G^{IJ}), \quad (31)$$

in which only some of the possible quadratic-stretch terms appear. Incompressibility needs to be implemented with an additional constraint (pressure) term, but the energetic parts of quadratic-Biot with $c_2 = -4c_1$ and neo-Hookean would agree to $O(i_3^{\mathbf{E}_B})$.

B. Constitutive relations

For an isotropic material with symmetric Cauchy stress $\boldsymbol{\sigma}$, the terms in the Biot stress are identically symmetric, so that $\boldsymbol{\Sigma}_{\text{Biot}} = \mathbf{P}^\top \cdot \mathbf{Q} = \mathbf{Q}^\top \cdot \mathbf{P}$. The first Piola-Kirchhoff stress \mathbf{P} can now be written as a function of \mathbf{Q} , \mathbf{U} , and derivatives of \mathcal{W} with respect to the invariants of \mathbf{U} [60–62]. The derivation, as carefully detailed by Wheeler [60], leads to the expression

$$\mathbf{P} = \left(\frac{\partial \mathcal{W}}{\partial i_1^{\mathbf{U}}} + i_1^{\mathbf{U}} \frac{\partial \mathcal{W}}{\partial i_2^{\mathbf{U}}} \right) \mathbf{Q} - \frac{\partial \mathcal{W}}{\partial i_2^{\mathbf{U}}} \mathbf{Q} \cdot \mathbf{U} + \frac{\partial \mathcal{W}}{\partial i_3^{\mathbf{U}}} i_3^{\mathbf{U}} (\mathbf{Q} \cdot \mathbf{U})^{-\top}. \quad (32)$$

Using the Cayley-Hamilton theorem [62], $\mathbf{U}^3 - i_1^{\mathbf{U}} \mathbf{U}^2 + i_2^{\mathbf{U}} \mathbf{U} - i_3^{\mathbf{U}} \mathbf{I} = \mathbf{0}$, which allows replacement of \mathbf{U}^{-1} thus:

$$\mathbf{P} = \left(\frac{\partial \mathcal{W}}{\partial i_1^{\mathbf{U}}} + i_1^{\mathbf{U}} \frac{\partial \mathcal{W}}{\partial i_2^{\mathbf{U}}} + i_2^{\mathbf{U}} \frac{\partial \mathcal{W}}{\partial i_3^{\mathbf{U}}} \right) \mathbf{Q} - \left(\frac{\partial \mathcal{W}}{\partial i_2^{\mathbf{U}}} + i_1^{\mathbf{U}} \frac{\partial \mathcal{W}}{\partial i_3^{\mathbf{U}}} \right) \mathbf{Q} \cdot \mathbf{U} + \frac{\partial \mathcal{W}}{\partial i_3^{\mathbf{U}}} \mathbf{Q} \cdot \mathbf{U}^2. \quad (33)$$

The relationships between invariants (12), those between stresses, and the definition of the Biot strain, convert this expression into one for the Biot stress in terms of the Biot strain and its invariants,

$$\boldsymbol{\Sigma}_{\text{Biot}} = \left(\frac{\partial \mathcal{W}}{\partial i_1^{\mathbf{E}_B}} + i_1^{\mathbf{E}_B} \frac{\partial \mathcal{W}}{\partial i_2^{\mathbf{E}_B}} + i_2^{\mathbf{E}_B} \frac{\partial \mathcal{W}}{\partial i_3^{\mathbf{E}_B}} \right) \mathbf{I} - \left(\frac{\partial \mathcal{W}}{\partial i_2^{\mathbf{E}_B}} + i_1^{\mathbf{E}_B} \frac{\partial \mathcal{W}}{\partial i_3^{\mathbf{E}_B}} \right) \mathbf{E}_B + \frac{\partial \mathcal{W}}{\partial i_3^{\mathbf{E}_B}} \mathbf{E}_B^2. \quad (34)$$

Specifically for the quadratic-Biot energy (28),

$$\begin{aligned} \boldsymbol{\Sigma}_{\text{Biot}} &= (2c_1 + c_2) i_1^{\mathbf{E}_B} \mathbf{I} - c_2 \mathbf{E}_B, \\ &= (2c_1 + c_2) \left(-2 + \sqrt{g_{ij} G^{IJ} + 2\sqrt{g/G} - 1} \right) \mathbf{I} - c_2 \mathbf{E}_B + O(i_3^{\mathbf{E}_B}), \end{aligned} \quad (35)$$

and $\mathbf{P} = \mathbf{Q} \cdot \boldsymbol{\Sigma}_{\text{Biot}}$.

C. Formulations in terms of rotations or stretches

Either the rotation or stretch field may be eliminated to form a set of equations for two fields, one of which must be obtained by solving a quadratic equation, or both rotation and stretch can be eliminated to obtain an approximate description in terms of position derivatives alone. Formulations involving independent rotations have been developed in [25, 26, 55]. While these were motivated by computational concerns, the equations presented below are relatively simple from an analytical point of view.

A description in terms of position derivatives and rotations is the linear momentum equation (22) with

$$\mathbf{P} = \left[(2c_1 + c_2) i_1^{\mathbf{E}_B} + c_2 \right] \mathbf{Q} - c_2 \mathbf{g}_i \mathbf{G}^I, \quad (36)$$

$$P^{iJ} = \left[(2c_1 + c_2) i_1^{\mathbf{E}_B} + c_2 \right] Q^i_K G^{KJ} - c_2 G^{IJ}, \quad (37)$$

with $i_1^{\mathbf{E}_B} = Q_k^K - 3 = g_{ki} Q^i_J G^{KJ} - 3 = -2 + \sqrt{g_{ij} G^{IJ} + 2\sqrt{g/G} - 1} + O(i_3^{\mathbf{E}_B})$. The rotations may be obtained either exactly by solving the constraint (2) or (5), or approximately by using the explicit form from (10) and (18).

A description in terms of position derivatives and stretches is the linear momentum equation (22) with

$$\mathbf{P} = \left[(2c_1 + c_2) i_1^{\mathbf{E}_B} + c_2 \right] \mathbf{g}^i \mathbf{G}_I \cdot \mathbf{U} - c_2 \mathbf{g}_i \mathbf{G}^I, \quad (38)$$

$$P^{iJ} = \left[(2c_1 + c_2) i_1^{\mathbf{E}_B} + c_2 \right] g^{ik} U_{KL} G^{LJ} - c_2 G^{IJ}, \quad (39)$$

with $i_1^{\mathbf{E}_B} = U^I_I - 3 = G^{IJ}U_{JI} - 3 = -2 + \sqrt{g_{ij}G^{IJ} + 2\sqrt{g/G} - 1} + O(i_3^{\mathbf{E}_B})$. We have used $\mathbf{Q} = \mathbf{Q}^{-\top}$ to rewrite \mathbf{P} in terms of \mathbf{U} . The stretches may be obtained either exactly by solving (6), or approximately by using the explicit form from (8) and (18).

V. TOWARDS ANISOTROPIC THEORIES

Many soft material structures of the type our theory is intended to address are anisotropic. In this section, we briefly indicate how to construct energies and balance equations for such materials, through the example of a transversely isotropic elastic solid such as a fiber-reinforced material.

Let the anisotropic material have a distinguished direction, in the reference configuration, identified with the unit vector \mathbf{D} . Following the pattern for constructing a transversely isotropic energy using this quantity and a strain tensor [63, 64], we obtain to quadratic order in stretch

$$\mathcal{W}(\mathbf{E}_B, \mathbf{D}) = \left(c_1 + \frac{c_2}{2}\right) (\text{Tr } \mathbf{E}_B)^2 - \frac{c_2}{2} \text{Tr} (\mathbf{E}_B^2) + c_3 (\text{Tr } \mathbf{E}_B) (\mathbf{E}_B : \mathbf{D}\mathbf{D}) + c_4 (\mathbf{E}_B : \mathbf{D}\mathbf{D})^2 + c_5 \mathbf{E}_B^2 : \mathbf{D}\mathbf{D}, \quad (40)$$

where the c_i are constant material parameters. In terms of an elastic tensor, with $\mathcal{W} = \frac{1}{2} \mathbf{E}_B : \mathbf{A} : \mathbf{E}_B$, we have $\frac{1}{2} A^{IJKL} = (c_1 + \frac{c_2}{2}) G^{IJ} G^{KL} - \frac{c_2}{4} (G^{IK} G^{JL} + G^{IL} G^{JK}) + c_3 G^{IJ} D^K D^L + c_4 D^I D^J D^K D^L + \frac{c_5}{4} (G^{IK} D^J D^L + G^{IL} D^J D^K + G^{JK} D^I D^L + G^{JL} D^I D^K)$. The Biot stress is still computed as the derivative of \mathcal{W} with respect to \mathbf{E}_B ,

$$\Sigma_{\text{Biot}} = (2c_1 + c_2) i_1^{\mathbf{E}_B} \mathbf{I} - c_2 \mathbf{E}_B + c_3 (\mathbf{I} \mathbf{E}_B : \mathbf{D}\mathbf{D} + i_1^{\mathbf{E}_B} \mathbf{D}\mathbf{D}) + 2c_4 \mathbf{E}_B : \mathbf{D}\mathbf{D}\mathbf{D}\mathbf{D} + c_5 (\mathbf{D}\mathbf{D} \cdot \mathbf{E}_B + \mathbf{E}_B \cdot \mathbf{D}\mathbf{D}), \quad (41)$$

but $\mathbf{P}^\top \cdot \mathbf{Q}$ is no longer symmetric. The first Piola-Kirchhoff stress can be written [25] in terms of the symmetric Biot stress plus an additional anisotropic contribution Σ_{an} ,

$$\mathbf{P} = \mathbf{Q} \cdot (\Sigma_{\text{Biot}} + \Sigma_{\text{an}}), \quad (42)$$

whose determination will require use of the angular momentum balance (23) alongside the linear momentum balance (22) and constitutive equation (24).

VI. CONCLUSIONS

This paper has established a basis for small-strain nonlinear-elastic theories with energies quadratic in stretch, reflecting a systematic expansion in Biot strains. Results on the kinematics of stretch and rotation, and variational principles for elasticity with auxiliary fields, have been further developed and combined within this framework. Neglect of higher order strains results in simple algebraic expressions for isotropic invariants that bypass the need for complex operations involving quartic equations or tensor square roots. Stresses, balance laws, and constitutive relations are expressed in terms of derivatives of position, with optional simultaneous consideration of a stretch or rotation field. The ideas are developed in the context of an isotropic material, with a brief sketch of anisotropic extensions.

ACKNOWLEDGMENTS

This work was supported by U.S. National Science Foundation grant CMMI-2001262. Helpful conversations with O. Oshri, D. J. Steigmann, E. G. Virga, and A. Yavari are acknowledged.

Appendix A: Other decompositions of the deformation gradient

The multiplicative polar decomposition of the deformation gradient (1) is a cornerstone of continuum mechanics, possessing several desirable features, including unique symmetric positive-definite right and left stretches. However, it is not the only option. In micropolar continuum theories [26, 65, 66] the multiplicative decomposition involves an independent micropolar rotation and a non-symmetric stretch. Chen [67] proposed an additive decomposition into a

symmetric strain tensor and an orthogonal rotation tensor,

$$\mathbf{g}_i \mathbf{G}^I = \mathbf{S} + \mathbf{R}. \quad (\text{A1})$$

The strain \mathbf{S} is unique. By contrast, there are two choices in the polar decomposition, corresponding to the order of application of strain and rotation, with corresponding natural notions of referential (Biot) and present (Bell) strains. The tensors in the decomposition (A1) can be computed directly from derivatives of position [67],

$$\mathbf{S} = -\mathbf{I} + \frac{1}{2} (\mathbf{g}_i \mathbf{G}^I + \mathbf{G}^I \mathbf{g}_i) - \frac{1}{1 + \cos\theta} \mathbf{W} \cdot \mathbf{W}, \quad (\text{A2})$$

$$\mathbf{R} = \mathbf{I} + \frac{1}{2} \mathbf{W} + \frac{1}{1 + \cos\theta} \mathbf{W} \cdot \mathbf{W}, \quad (\text{A3})$$

$$\mathbf{W} = \frac{1}{2} (\mathbf{g}_i \mathbf{G}^I - \mathbf{G}^I \mathbf{g}_i), \quad (\text{A4})$$

$$\cos\theta = (1 - \mathbf{W} : \mathbf{W})^{1/2}, \quad (\text{A5})$$

However, in contrast to the stretches from the polar decomposition, the strain \mathbf{S} is not a pure measure of material distortion but is still corrupted by irrelevant rotational information, and the functional form of an objective elastic energy must include some combination of both strain \mathbf{S} and rotation \mathbf{R} . This approach has, to our knowledge, not been explored. In micropolar theories [26, 65, 66], a quantity of the type $\mathbf{S} = \mathbf{g}_i \mathbf{G}^I - \mathbf{R}$ is one possible definition for the linear strain, to be accompanied by an angular strain.

Appendix B: Symmetry of $\mathbf{P}^\top \cdot \mathbf{Q}$ for isotropic materials

The following line of reasoning may be found in Lurie [22] and Ogden [1]. For isotropic materials with symmetric Cauchy stress $\boldsymbol{\sigma}$, there exists a representation $\boldsymbol{\sigma} = c_0 \mathbf{I} + c_1 \mathbf{V} + c_2 \mathbf{V}^2$, where the c_i are functions of the invariants of \mathbf{V} . This means that $\boldsymbol{\sigma}$ is coaxial to (shares eigenvectors with) \mathbf{V} and \mathbf{B} and, consequently, $\mathbf{Q}^\top \cdot \boldsymbol{\sigma} \cdot \mathbf{Q}$ is coaxial to \mathbf{U} and \mathbf{C} . Noting further that \mathbf{U} is coaxial with its inverse, and that $\mathbf{P}^\top \cdot \mathbf{Q} = \sqrt{g/G} \mathbf{U}^{-1} \cdot \mathbf{Q}^\top \cdot \boldsymbol{\sigma} \cdot \mathbf{Q}$ and $\mathbf{Q}^\top \cdot \mathbf{P} = \sqrt{g/G} \mathbf{Q}^\top \cdot \boldsymbol{\sigma} \cdot \mathbf{Q} \cdot \mathbf{U}^{-1}$, the coaxiality of $\mathbf{Q}^\top \cdot \boldsymbol{\sigma} \cdot \mathbf{Q}$ and \mathbf{U}^{-1} implies the symmetry of $\mathbf{P}^\top \cdot \mathbf{Q}$, which can thus be identified with the Biot stress *via* (24).

Appendix C: Variation of the energy

Here we detail the first variation of the energy (21) under independent shifts in $\mathbf{x} \in \mathbb{E}^3$, $\mathbf{Q} \in \text{SO}(3)$, $\mathbf{U} \in \text{Sym}^+$. The approach follows Atluri and Murakawa [24].

Noting that $\delta \mathbf{U} = \delta \mathbf{E}_B$ and using $\mathbf{P} : \delta \mathbf{Q} \cdot \mathbf{U} = \delta \mathbf{Q} : \mathbf{P} \cdot \mathbf{U}$ and $\mathbf{P} : \mathbf{Q} \cdot \delta \mathbf{E}_B = \mathbf{Q}^\top \cdot \mathbf{P} : \delta \mathbf{E}_B$, we obtain

$$\delta \mathcal{E} = \int_B \left[-\bar{\nabla} \cdot \mathbf{P}^\top \cdot \delta \mathbf{x} - \delta \mathbf{Q} : \mathbf{P} \cdot \mathbf{U} + \left(\frac{\partial \mathcal{W}}{\partial \mathbf{E}_B} - \mathbf{Q}^\top \cdot \mathbf{P} \right) : \delta \mathbf{E}_B \right] dV + \int_{\partial B} \mathbf{N} \cdot \mathbf{P}^\top \cdot \delta \mathbf{x} dA. \quad (\text{C1})$$

The terms conjugate to $\delta \mathbf{x}$ directly provide the linear momentum balance (22) and a boundary condition $\mathbf{N} \cdot \mathbf{P}^\top = \mathbf{0}$.

Using $\delta \mathbf{Q} \cdot \mathbf{Q}^\top = -\mathbf{Q} \cdot \delta \mathbf{Q}^\top$, we rewrite the quantity involving $\delta \mathbf{Q}$ as $-\delta \mathbf{Q} : \mathbf{P} \cdot \mathbf{U} = (\mathbf{Q} \cdot \delta \mathbf{Q}^\top \cdot \mathbf{Q}) : \mathbf{P} \cdot \mathbf{U} = \delta \mathbf{Q}^\top \cdot \mathbf{Q} : (\mathbf{Q}^\top \cdot \mathbf{P} \cdot \mathbf{U})$. Because $\delta \mathbf{Q}^\top \cdot \mathbf{Q}$ is antisymmetric, only the antisymmetric part of the conjugate quantity need vanish,

$$\mathbf{U} \cdot \mathbf{P}^\top \cdot \mathbf{Q} = \mathbf{Q}^\top \cdot \mathbf{P} \cdot \mathbf{U}. \quad (\text{C2})$$

Left dotting with \mathbf{Q} and right dotting with \mathbf{Q}^\top provides the form of the angular momentum balance shown in (23).

Because $\delta \mathbf{E}_B$ is symmetric, only the symmetric part of its conjugate quantity need vanish, leading to the constitutive equation (24).

Appendix D: Bell strain and stress

Our framework has employed the referential right stretch and associated Biot strain. Their counterparts are the present left stretch \mathbf{V} and the Bell strain, defined as

$$\mathbf{E}_{\text{Bell}} = \mathbf{V} - \mathbf{I}. \quad (\text{D1})$$

The Biot and Bell strains share eigenvalues and invariants. A variational principle based on the energy

$$\mathcal{E}(\mathbf{x}, \mathbf{Q}, \mathbf{E}_{\text{Bell}}; \mathbf{P}) = \int_{\mathcal{B}} [\mathcal{W}(\mathbf{E}_{\text{Bell}}) + \mathbf{P} : (\mathbf{g}_i \mathbf{G}^I - \mathbf{V} \cdot \mathbf{Q})] dV \quad (\text{D2})$$

would result in a constitutive relation for the symmetric Bell stress [62, 68, 69],

$$\boldsymbol{\Sigma}_{\text{Bell}} \equiv \frac{\partial \mathcal{W}}{\partial \mathbf{E}_{\text{Bell}}} = \frac{1}{2} (\mathbf{Q} \cdot \mathbf{P}^\top + \mathbf{P} \cdot \mathbf{Q}^\top). \quad (\text{D3})$$

For isotropic materials with symmetric Cauchy stress, $\boldsymbol{\Sigma}_{\text{Biot}} = \mathbf{P}^\top \cdot \mathbf{Q}$ and $\boldsymbol{\Sigma}_{\text{Bell}} = \mathbf{Q} \cdot \mathbf{P}^\top$, so the first Piola-Kirchhoff stress \mathbf{P} admits the decomposition

$$\mathbf{P} = \mathbf{Q} \cdot \boldsymbol{\Sigma}_{\text{Biot}} = \boldsymbol{\Sigma}_{\text{Bell}} \cdot \mathbf{Q}. \quad (\text{D4})$$

Despite its resemblance to the polar decomposition of the deformation gradient, this decomposition for \mathbf{P} is not unique, since $\boldsymbol{\Sigma}_{\text{Biot}}$ and $\boldsymbol{\Sigma}_{\text{Bell}}$ are generally neither positive- nor negative-definite.

-
- [1] R. W. Ogden. *Non-Linear Elastic Deformations*. Courier, 1997.
 - [2] E. Efrati, E. Sharon, and R. Kupferman. Elastic theory of unconstrained non-Euclidean plates. *Journal of the Mechanics and Physics of Solids*, 57:762–775, 2009.
 - [3] M. A. Dias, J. A. Hanna, and C. D. Santangelo. Programmed buckling by controlled lateral swelling in a thin elastic sheet. *Physical Review E*, 84:036603, 2011.
 - [4] M. Pezulla, N. Stoop, X. Jiang, and D. P. Holmes. Morphing of geometric composites *via* residual swelling. *Soft Matter*, 11:5812–5820, 2015.
 - [5] T.-S. Nguyen and J. V. Selinger. Theory of liquid crystal elastomers and polymer networks. *The European Physical Journal E*, 40:76, 2017.
 - [6] W. M. van Rees, E. Vouga, and L. Mahadevan. Growth patterns for shape-shifting elastic bilayers. *PNAS*, 114(44):11597–11602, 2017.
 - [7] H. Irschik and J. Gerstmayr. A continuum mechanics based derivation of Reissner’s large-displacement finite-strain beam theory: the case of plane deformations of originally straight Bernoulli-Euler beams. *Acta Mechanica*, 206:1–21, 2009.
 - [8] O. Oshri and H. Diamant. Strain tensor selection and the elastic theory of incompatible thin sheets. *Physical Review E*, 95:053003, 2017.
 - [9] J. A. Hanna. Some observations on variational elasticity and its application to plates and membranes. *Zeitschrift für angewandte Mathematik und Physik*, 70:76, 2019.
 - [10] H. G. Wood and J. A. Hanna. Contrasting bending energies from bulk elastic theories. *Soft Matter*, 15:2411–2417, 2019.
 - [11] O. Ozenda and E. G. Virga. On the Kirchhoff-Love hypothesis (revised and vindicated). *Journal of Elasticity*, 143:359–384, 2021.
 - [12] S. Antman. General solutions for plane extensible elasticae having nonlinear stress-strain laws. *Quarterly of Applied Mathematics*, 26(1):35–47, 1968.
 - [13] E. Reissner. On one-dimensional finite-strain beam theory: the plane problem. *Zeitschrift für angewandte Mathematik und Physik*, 23:795–804, 1972.
 - [14] A. B. Whitman and C. N. DeSilva. An exact solution in a nonlinear theory of rods. *Journal of Elasticity*, 4(4):265–280, 1974.
 - [15] T. Iwakuma and S. Kuranishi. How much contribution does the shear deformation have in a beam theory? *Structural Engineering/Earthquake Engineering, Proceedings of the Japan Society of Civil Engineers*, 344:141–151, 1984.
 - [16] T. Chaisomphob, F. Nishino, A. Hasegawa, and A. G. A. Abdel-Shafy. An elastic finite displacement analysis of plane beams with and without shear deformation. *Structural Engineering/Earthquake Engineering, Proceedings of the Japan Society of Civil Engineers*, 368:169–177, 1986.
 - [17] A. Magnusson, M. Ristinmaa, and C. Ljung. Behaviour of the extensible elastica solution. *International Journal of Solids and Structures*, 38:8441–8457, 2001.
 - [18] S. S. Antman. *Nonlinear Problems of Elasticity*. Springer, New York, second edition, 2005.

- [19] S. Knoche and J. Kierfeld. Buckling of spherical capsules. *Physical Review E*, 84:046608, 2011.
- [20] H. S. Seung and D. R. Nelson. Defects in flexible membranes with crystalline order. *Physical Review A*, 38(2):1005–1018, 1988.
- [21] M. Deserno. Fluid lipid membranes: From differential geometry to curvature stresses. *Chemistry and Physics of Lipids*, 185:11–45, 2015.
- [22] A. I. Lur'e. Theory of elasticity for a semilinear material. *Journal of Applied Mathematics and Mechanics*, 32(6):1068–1085, 1968.
- [23] F. John. Plane strain problems for a perfectly elastic material of harmonic type. *Communications on Pure and Applied Mathematics*, 13(2):239–296, 1960.
- [24] S. N. Atluri and H. Murakawa. On hybrid finite element models in nonlinear solid mechanics. In P.G. Bergan et al., editor, *Finite Elements in Nonlinear Mechanics*, pages 3–41. Tapir, 1977.
- [25] K. Wisniewski. A shell theory with independent rotations for relaxed Biot stress and right stretch strain. *Computational Mechanics*, 21(2):101–122, 1998.
- [26] T. Merlini. A variational formulation for finite elasticity with independent rotation and Biot-axial fields. *Computational Mechanics*, 19(3):153–168, 1997.
- [27] C. D. Modes, K. Bhattacharya, and M. Warner. Gaussian curvature from flat elastica sheets. *Proceedings of the Royal Society A*, 467:1121–1140, 2011.
- [28] P. Plucinsky, B. A. Kowalski, T. J. White, and K. Bhattacharya. Patterning nonisometric origami in nematic elastomer sheets. *Soft Matter*, 14:3127–3134, 2018.
- [29] D. J. Steigmann. Tension-field theory. *Proceedings of the Royal Society of London A*, 429:141–173, 1990.
- [30] S. Sadik and A. Yavari. On the origins of the idea of the multiplicative decomposition of the deformation gradient. *Mathematics and Mechanics of Solids*, 22:771–772, 2017.
- [31] P. Nardinocchi, L. Teresi, and V. Varano. The elastic metric: A review of elasticity with large distortions. *International Journal of Non-Linear Mechanics*, 56:34–42, 2013.
- [32] J. L. Ericksen and C. Truesdell. Exact theory of stress and strain in rods and shells. *Archive for Rational Mechanics and Analysis*, 1(1):295–323, 1957.
- [33] L. E. Malvern. *Introduction to Continuum Mechanics*. Prentice Hall, 1969.
- [34] R. A. Stephenson. On the uniqueness of the square-root of a symmetric, positive-definite tensor. *Journal of Elasticity*, 10(2), 1980.
- [35] M. A. Biot. *Mechanics of Incremental Deformations*. John Wiley & Sons, 1965.
- [36] A. Hoger and D. E. Carlson. Determination of the stretch and rotation in the polar decomposition of the deformation gradient. *Quarterly of Applied Mathematics*, 42(1):113–117, 1984.
- [37] K. Sawyers. Comments on the paper *Determination of the stretch and rotation in the polar decomposition of the deformation gradient* by A. Hoger and D. E. Carlson. *Quarterly of Applied Mathematics*, 44(2):309–311, 1986.
- [38] T. C. T. Ting. Determination of $C^{1/2}$, $C^{-1/2}$ and more general isotropic tensor functions of C . *Journal of Elasticity*, 15(3):319–323, 1985.
- [39] J. Stickforth. The square root of a three-dimensional positive tensor. *Acta Mechanica*, 67(1-4):233–235, 1987.
- [40] J. Lu and P. Papadopoulos. On the direct determination of the rotation tensor from the deformation gradient. *Mathematics and Mechanics of Solids*, 2(1):17–26, 1997.
- [41] D. Guan-Suo. Determination of the rotation tensor in the polar decomposition. *Journal of Elasticity*, 50(3):197–207, 1998.
- [42] C. S. Jog. On the explicit determination of the polar decomposition in n -dimensional vector spaces. *Journal of Elasticity*, 66(2):159–169, 2002.
- [43] C. Bouby, D. Fortuné, W. Pietraszkiewicz, and C. Vallée. Direct determination of the rotation in the polar decomposition of the deformation gradient by maximizing a Rayleigh quotient. *Zeitschrift für Angewandte Mathematik und Mechanik*, 85(3):155–162, 2005.
- [44] A. Norris. Invariants of $C^{1/2}$ in terms of the invariants of C . *Journal of Mechanics of Materials and Structures*, 2(9):1805–1812, 2007.
- [45] N. H. Scott. $U = C^{1/2}$ and its invariants in terms of C and its invariants. *Journal of Elasticity*, 141(2):363–379, 2020.
- [46] B.-H. Sun. Explicit representation for the $SO(3)$ rotation tensor of deformable bodies. *Applied Mathematics Letters*, 111:106606, 2021.
- [47] S. N. Atluri. Alternate stress and conjugate strain measures, and mixed variational formulations involving rigid rotations, for computational analyses of finitely deformed solids, with application to plates and shells—I: Theory. *Computers & Structures*, 18(1):93–116, 1984.
- [48] S. N. Atluri and A. Cazzani. Rotations in computational solid mechanics. *Archives of Computational Methods in Engineering*, 2(1):49–138, 1995.
- [49] P. Neff, A. Fischle, and I. Münch. Symmetric Cauchy stresses do not imply symmetric Biot strains in weak formulations of isotropic hyperelasticity with rotational degrees of freedom. *Acta Mechanica*, 197(1):19–30, 2008.
- [50] M. M. Carroll. Finite strain solutions in compressible isotropic elasticity. *Journal of Elasticity*, 20(1):65–92, 1988.
- [51] B. Fraeijns de Veubeke. A new variational principle for finite elastic displacements. *International Journal of Engineering Science*, 10(9):745–763, 1972.
- [52] H. Buefer. On the work theorems for finite and incremental elastic deformations with discontinuous fields: a unified treatment of different versions. *Computer Methods in Applied Mechanics and Engineering*, 36:95–124, 1983.

- [53] H. Buefler. The Biot stresses in nonlinear elasticity and the associated generalized variational principles. *Ingenieur-Archiv*, 55(6):450–462, 1985.
- [54] J. Makowski and H. Stumpf. Buckling equations for elastic shells with rotational degrees of freedom undergoing finite strain deformation. *International Journal of Solids and Structures*, 26(3):353–368, 1990.
- [55] J. C. Simo, D. D. Fox, and T. J. R. Hughes. Formulations of finite elasticity with independent rotations. *Computer Methods in Applied Mechanics and Engineering*, 95(2):277–288, 1992.
- [56] C. Sansour and H. Buefler. An exact finite rotation shell theory, its mixed variational formulation and its finite element implementation. *International Journal for Numerical Methods in Engineering*, 34(1):73–115, 1992.
- [57] H. Buefler. On drilling degrees of freedom in nonlinear elasticity and a hyperelastic material description in terms of the stretch tensor. Part 1: Theory. *Acta Mechanica*, 113(1):21–35, 1995.
- [58] C. Sansour, H. Buefler, and H. Müllerschön. On drilling degrees of freedom in nonlinear elasticity and a hyperelastic material description in terms of the stretch tensor. Part 2: Application to membranes. *Acta Mechanica*, 115(1):103–117, 1996.
- [59] I. Chao, U. Pinkall, P. Sanan, and P. Schröder. A simple geometric model for elastic deformations. *ACM Transactions on Graphics*, 29(4):Article 38, 2010.
- [60] L. T. Wheeler. On the derivatives of the stretch and rotation with respect to the deformation gradient. *Journal of Elasticity*, 24(1-3):129–133, 1990.
- [61] K. Wiśniewski and E. Turska. A note on the hyperelastic constitutive equation for rotated Biot stress. *Archives of Mechanics*, 48(5):947–953, 1996.
- [62] D. J. Steigmann. Invariants of the stretch tensors and their application to finite elasticity theory. *Mathematics and Mechanics of Solids*, 7(4):393–404, 2002.
- [63] A. J. M. Spencer. Constitutive theory for strongly anisotropic solids. In A. J. M. Spencer, editor, *Continuum Theory of the Mechanics of Fibre-Reinforced Composites*, pages 1–32. Springer, 1984.
- [64] J. G. Murphy. Transversely isotropic biological, soft tissue must be modelled using both anisotropic invariants. *European Journal of Mechanics A/Solids*, 42:90–96, 2013.
- [65] P. Steinmann and E. Stein. A unifying treatise of variational principles for two types of micropolar continua. *Acta Mechanica*, 121(1):215–232, 1997.
- [66] W. Pietraszkiewicz and V. A. Eremeyev. On natural strain measures of the non-linear micropolar continuum. *International Journal of Solids and Structures*, 46(3-4):774–787, 2009.
- [67] Z. Chen. On the representation of finite rotation in nonlinear field theory of continuum mechanics. *Applied Mathematics and Mechanics*, 7(11):1017–1026, 1986.
- [68] J. F. Bell. Continuum plasticity at finite strain for stress paths of arbitrary composition and direction. *Archive for Rational Mechanics and Analysis*, 84:139–170, 1983.
- [69] M. F. Beatty and M. A. Hayes. Deformations of an elastic, internally constrained material. Part 1: Homogeneous deformations. *Journal of Elasticity*, 29(1):1–84, 1992.