POLYTIME REDUCTIONS OF AF-ALGEBRAIC PROBLEMS

DANIELE MUNDICI

ABSTRACT. We assess the computational complexity of several decision problems concerning (Murray-von Neumann) equivalence classes of projections of AF-algebras whose Elliott classifier is lattice-ordered. We construct polytime reductions among many of these problems.

1. Introduction

Let $\mathfrak A$ be an AF-algebra, [4]. The partial addition + in Elliott's local semigroup [9] of $\mathfrak A$ is uniquely extendible to a total operation that preserves all algebraic and order properties of + iff the Murray-von Neumann order \leq of $\mathfrak A$ is a lattice, for short, $\mathfrak A$ is an $AF\ell$ -algebra. See Theorem 1.1. $AF\ell$ -algebras have a preeminent role in the literature on AF-algebras. The Elliott classifier $E(\mathfrak A)$ of any $AF\ell$ -algebra $\mathfrak A$ is an MV-algebra, i.e., a Lindenbaum algebra in Lukasiewicz logic L_{∞} , [6, 19]. All countable MV-algebras arise as $E(\mathfrak A)$ for some $AF\ell$ -algebra $\mathfrak A$. Since $\mathfrak A$ is a quotient of the universal $AF\ell$ -algebra $\mathfrak M$ of [16, §8], and $E(\mathfrak M)$ is the free MV-algebra over countably many generators, each L_{∞} -formula ϕ naturally codes an equivalence class $\phi^{\mathfrak A}$ of projections in $\mathfrak A$.

We equip the Elliott monoid $E(\mathfrak{A})$ of every $AF\ell$ -algebra \mathfrak{A} with a partial order $\phi^{\mathfrak{A}} \sqsubseteq \psi^{\mathfrak{A}}$ intuitively meaning that all projections in $\phi^{\mathfrak{A}}$ are "less eccentric" than those in $\psi^{\mathfrak{A}}$. We prove that p is central iff its (always Murray-von Neumann) equivalence class [p] is \sqsubseteq -minimal iff it is a (Freudenthal) characteristic element of $K_0(\mathfrak{A})$ iff $[p] \wedge [1-p] = 0$. Among others, we consider the following decision problems: (a) $\phi^{\mathfrak{A}} : \models ? \psi^{\mathfrak{A}}$, (b) $\phi^{\mathfrak{A}} : \not \vdash ? \psi^{\mathfrak{A}}$, (c) $\phi^{\mathfrak{A}} : \not \vdash ? \psi^{\mathfrak{A}}$, (d) $\phi^{\mathfrak{A}} : \not \vdash ? 0$, (e) Is $\phi^{\mathfrak{A}}$ central? We prove that problems (a)-(d) for \mathfrak{A} have the same computational complexity, up to a polytime reduction. If \mathfrak{A} is simple, or if \mathfrak{A} has no quotient isomorphic to \mathbb{C} , then also problem (e) for \mathfrak{A} has the same complexity as (a)-(d). The complexity of each problem (a)-(e) is polytime for many relevant AF-algebras in the literature, including the Behncke-Leptin algebras $\mathcal{A}_{m,n}$ and every Effros-Shen algebra \mathfrak{F}_{θ} , for $\theta \in [0,1] \setminus \mathbb{Q}$ a real algebraic integer, or for $\theta = 1/e$.

For every AF ℓ -algebra \mathfrak{A} we let $\mathsf{prim}(\mathfrak{A})$ denote the set of primitive (always closed and two-sided) ideals of \mathfrak{A} equipped with the Jacobson topology. By [4, 3.8], an ideal \mathfrak{P} of \mathfrak{A} is primitive iff it is prime , in the sense that whenever ideals $\mathfrak{P}_1, \mathfrak{P}_2$ of \mathfrak{A} satisfy $\mathfrak{P}_1 \cap \mathfrak{P}_2 = \mathfrak{P}$ then either $\mathfrak{P}_1 = \mathfrak{P}$ or $\mathfrak{P}_2 = \mathfrak{P}$.

For any MV-algebra A we let Spec(A) denote the space of prime ideals of A endowed with the Zariski (hull-kernel) topology, ([19, 4.14]). As shown in [6, 1.2.14],

$$\bigcap \mathsf{Spec}(A) = \{0\}. \tag{1}$$

The basic properties of AF ℓ -algebras are summarized by the following two results:

Date: April 30, 2021.

²⁰¹⁰ Mathematics Subject Classification. Primary: 46L35. Secondary: 06D35, 06F20, 19A49, 46L80, 47L40.

 $Key\ words\ and\ phrases.$ AF algebra, Elliott classification, Elliott local semigroup, Murray-von Neumann order, Grothendieck K_0 group, MV algebra, liminary C*-algebra, Effros-Shen algebra, Behnke-Leptin algebras, Farey AF algebra.

Theorem 1.1. Let \mathfrak{A} be an AF-algebra.

- (i) [20, Theorem 1] Elliott's partial addition + in $L(\mathfrak{A})$ has at most one extension to an associative, commutative, monotone operation \oplus : $L(\mathfrak{A})^2 \to L(\mathfrak{A})$ such that for each projection $p \in \mathfrak{A}$, $[1_{\mathfrak{A}}-p]$ is the smallest $[q] \in L(\mathfrak{A})$ satisfying $[p] \oplus [q] = [1_{\mathfrak{A}}]$. The semigroup $(S(\mathfrak{A}), \oplus)$ expanding $L(\mathfrak{A})$ exists iff \mathfrak{A} is an AFl-algebra.
- (ii) [9] Let \mathfrak{A}_1 and \mathfrak{A}_2 be AFl-algebras. For each j=1,2 let \oplus_j be the extension of Elliott's addition given by (i). Then the semigroups $(S(\mathfrak{A}_1), \oplus_1)$ and $(S(\mathfrak{A}_2), \oplus_2)$ are isomorphic iff so are \mathfrak{A}_1 and \mathfrak{A}_2 .
- (iii) [20, Theorem 2, Proposition 2.2] $(S(\mathfrak{A}), \oplus)$ has the richer structure of a monoid $(E(\mathfrak{A}), 0, *, \oplus)$ with an involution $[p]^* = [1_{\mathfrak{A}} p]$. The Murray-von Neumann lattice-order of equivalence classes of projections $[p], [q] \in E(\mathfrak{A})$ is definable by the involutive monoidal operations of $E(\mathfrak{A})$ upon setting $[p] \vee [q] = ([p]^* \oplus [q])^* \oplus [q]$ and $[p] \wedge [q] = ([p]^* \vee [q]^*)^*$. We say that $E(\mathfrak{A})$ is the Elliott monoid of \mathfrak{A} .
- (iv) [16, Theorem 3.9] The map $\mathfrak{A} \mapsto (E(\mathfrak{A}), 0, {}^*, \oplus)$ is a bijective correspondence between isomorphism classes of AFl-algebras and isomorphism classes of countable abelian monoids with a unary operation * satisfying the equations: $x^{**} = x$, $0^* \oplus x = 0^*$, and $(x^* \oplus y)^* \oplus y = (y^* \oplus x)^* \oplus x$. The involutive monoids defined by these equations are known as MV-algebras, [5, 6, 19]. Let Γ be the categorical equivalence between unital ℓ -groups and MV-algebras defined in [16, §3]. Then $(E(\mathfrak{A}), 0, {}^*, \oplus)$ is isomorphic to $\Gamma(K_0(\mathfrak{A}), K_0(\mathfrak{A})^+, [1_{\mathfrak{A}}])$.
 - (v) ([7, 8, 11]) For any $AF\ell$ -algebra \mathfrak{A} the dimension group

$$K_0(\mathfrak{A}) = (K_0(\mathfrak{A}), K_0(\mathfrak{A})^+, [1_{\mathfrak{A}}])$$

is a countable lattice-ordered abelian group with a distinguished strong order unit (for short, a unital ℓ -group). All countable unital ℓ -groups arise in this way. Let \mathfrak{A}_1 and \mathfrak{A}_2 be $AF\ell$ -algebras. Then $K_0(\mathfrak{A}_1)$ and $K_0(\mathfrak{A}_2)$ are isomorphic unital ℓ -groups iff \mathfrak{A}_1 and \mathfrak{A}_2 are isomorphic.

Theorem 1.2. In any $AF\ell$ -algebra \mathfrak{A} we have:

- (i) The map $\eta: \mathfrak{I} \mapsto K_0(\mathfrak{I}) \cap E(\mathfrak{A})$ is an isomorphism of the lattice of ideals of \mathfrak{A} onto the lattice of ideals of $E(\mathfrak{A})$. Primitive ideals of \mathfrak{A} correspond via η to prime ideals of $E(\mathfrak{A})$.
 - (ii) In particular, η is a homeomorphism of $prim(\mathfrak{A})$ onto $Spec(E(\mathfrak{A}))$.
- (iii) Suppose j is an ideal of the countable MV-algebra A. Let the AF ℓ -algebra \mathfrak{A} be defined by $E(\mathfrak{A}) = A$, as in Theorem 1.1(iv). Let the ideal \mathfrak{J} of \mathfrak{A} be defined by $\eta(\mathfrak{J}) = j$. Then A/j is isomorphic to $E(\mathfrak{A}/\mathfrak{J})$.
- (iv) For every ideal \mathfrak{I} of \mathfrak{A} , the map $[p/\mathfrak{I}] \mapsto [p]/\eta(\mathfrak{I})$, (with p ranging over all projections in \mathfrak{A}), is an isomorphism of $E(\mathfrak{A}/\mathfrak{I})$ onto $E(\mathfrak{A})/\eta(\mathfrak{I})$. In particular, for every $\mathfrak{P} \in \text{prim}(\mathfrak{A})$ the MV-algebra $E(\mathfrak{A}/\mathfrak{P})$ is totally ordered and $\mathfrak{A}/\mathfrak{P}$ has comparability of projections in the sense of Murray-von Neumann.
- (v) The map $\mathfrak{l} \mapsto \mathfrak{l} \cap \Gamma(K_0(\mathfrak{A}))$ is an isomorphism of the lattice of ideals of $K_0(\mathfrak{A})$ (i.e., kernels of unit preserving ℓ -homomorphisms of $K_0(\mathfrak{A})$) onto the lattice of ideals of $E(\mathfrak{A})$. Further,

$$\Gamma\left(\frac{K_0(\mathfrak{A})}{\mathfrak{l}}\right) \;\cong\; \frac{\Gamma(K_0(\mathfrak{A}))}{\mathfrak{l}\;\cap\; \Gamma(K_0(\mathfrak{A}))}\,.$$

Proof. (i) From [7, Proposition IV.5.1] and [11, p.196, 21H] one gets an isomorphism of the lattice of ideals of \mathfrak{A} onto the lattice of ideals of the unital ℓ -group $K_0(\mathfrak{A})$. The preservation properties of Γ , [6, Theorems 7.2.2, 7.2.4], then yield the

desired isomorphism. The second statement immediately follows from [4, Theorem 3.8] and the above mentioned characterization of prime ideals in MV-algebras, ([19, Proposition 4.13]).

- (ii) follows from (i), by definition of the topologies of $prim(\mathfrak{A})$ and $Spec(E(\mathfrak{A}))$.
- (iii) We have an exact sequence $0 \to \mathfrak{J} \to \mathfrak{A} \to \mathfrak{A}/\mathfrak{J} \to 0$. Correspondingly ([7, IV.15] or [8, Corollary 9.2]), we have an exact sequence $0 \to K_0(\mathfrak{J}) \to K_0(\mathfrak{A}) \to K_0(\mathfrak{A}/\mathfrak{J}) \to 0$, whence the unital ℓ -groups $K_0(\mathfrak{A}/\mathfrak{J})$ and $K_0(\mathfrak{A})/K_0(\mathfrak{J})$ are isomorphic. The preservation properties of Γ under quotients [6, Theorem 7.2.4], together with Theorem 1.1(iv)-(v) yield

$$E\left(\frac{\mathfrak{A}}{\mathfrak{J}}\right) \cong \Gamma\left(K_0\left(\frac{\mathfrak{A}}{\mathfrak{J}}\right)\right) \cong \frac{\Gamma(K_0(\mathfrak{A}))}{K_0(\mathfrak{J}) \cap \Gamma(K_0(\mathfrak{A}))} \cong \frac{E(\mathfrak{A})}{\eta(\mathfrak{J})} = \frac{A}{\mathfrak{f}}.$$

(iv) Combine (i) and (iii) with the preservation properties of K_0 for exact sequences and the preservation properties of Γ under quotients. The MV-algebra $E(\mathfrak{A}/\mathfrak{P})$ is totally ordered. As a matter of fact, by (ii), $\eta(\mathfrak{P})$ belongs to $\mathsf{Spec}(E(\mathfrak{A}))$ whenever \mathfrak{P} beongs to $\mathsf{prim}(\mathfrak{A})$. By Theorem 1.1(iv), $\mathfrak{A}/\mathfrak{P}$ has comparability of projections.

2. Characterizing central projections in AF\ell-algebras

Following [12, p.22], by a characteristic element in a partially ordered abelian group G with order unit u, we mean an element c with $0 \le c \le u$ such that the greatest lower bound of the set $\{c, u-c\}$ exists and equals 0. In the framework of real vector lattices, with u an arbitrary positive element, c is known as a "component of u", [1, p.13], [13, p.284]. This notion goes back to Freudenthal, [10, p. 643, (5.2)].

In every MV-algebra A one defines $a\odot b=(a^*\oplus b^*)^*$ and $\mathsf{dist}(a,b)=(a\odot b^*)\oplus (b\odot a^*)$. Chang [5] writes $x\cdot y$ instead of $x\odot y$, and d(x,y) instead of $\mathsf{dist}(x,y)$. The latter is known as *Chang's distance function*. See [6, p.8 and 1.2.4] and [5, p.477]. Further, let us use the notation $\mathsf{B}(A)$ for the set of *idempotent* elements of the MV-algebra A,

$$\mathsf{B}(A) = \{ a \in A \mid a \oplus a = a \}. \tag{2}$$

As observed by Chang [5, Theorems 1.16-1.17], $\mathsf{B}(A)$ is a subalgebra of A which turns out to be a boolean algebra. By an MV-chain we mean an MV-algebra D whose underlying order is total, [6, p.10].

The binary relation "p is closer than q to the center". Let \mathfrak{P} be a primitive ideal of an AF ℓ -algebra \mathfrak{A} . By Theorem 1.2 the quotient AF ℓ -algebra $\mathfrak{A}/\mathfrak{P}$ has comparability of projections. Thus for any projections $p, q \in \mathfrak{A}$ we have the following three mutually incompatible cases:

$$p/\mathfrak{P} \prec q/\mathfrak{P}, \ p/\mathfrak{P} \succ q/\mathfrak{P}, \ p/\mathfrak{P} \sim q/\mathfrak{P}.$$

In particular, we have incompatible cases:

$$p/\mathfrak{P} \prec p^*/\mathfrak{P}, \ p/\mathfrak{P} \succ p^*/\mathfrak{P}, \ p/\mathfrak{P} \sim p^*/\mathfrak{P}.$$

The proof of the following result will appear elsewhere:

Theorem 2.1 (Characterization of central projections). Suppose \mathfrak{A} is an AF ℓ -algebra. In view of Theorem 1.1(iv)-(v), let us identify $E(\mathfrak{A})$ with the unit interval $\Gamma(K_0(\mathfrak{A}))$ of the unital ℓ -group $K_0(\mathfrak{A})$.

- (I) For every projection p of \mathfrak{A} the following conditions are equivalent:
 - (i) $p/\mathfrak{P} \in \{0,1\}$, (the trivial elements of $\mathfrak{A}/\mathfrak{P}$) for all $\mathfrak{P} \in \mathsf{prim}(\mathfrak{A})$.

- (ii) $[p] \in \mathsf{B}(E(\mathfrak{A})).$
- (iii) p is central in \mathfrak{A} .
- (iv) [p] is a characteristic element of $K_0(\mathfrak{A})$.
- (II) For any $x, y \in E(\mathfrak{A})$ let us write $x \sqsubseteq y$ iff for every prime ideal \mathfrak{p} of $E(\mathfrak{A})$ $(y/\mathfrak{p} < y^*/\mathfrak{p} \text{ implies } x/\mathfrak{p} \le y/\mathfrak{p})$ and $(y/\mathfrak{p} > y^*/\mathfrak{p} \text{ implies } x/\mathfrak{p} \ge y/\mathfrak{p})$,

with \leq the underlying total order of $E(\mathfrak{A})/\mathfrak{p}$ defined in Theorem 1.1(iii). Then \sqsubseteq endows $E(\mathfrak{A})$ with a partial order relation. Moreover, for every projection p in \mathfrak{A} , the equivalent the conditions in (I) are also equivalent to [p] being \sqsubseteq -minimal in $E(\mathfrak{A})$.

3. The algorithmic theory of AF\ell-algebras

Our standard reference for computability theory is [14]. For L_{∞} we refer to [6, 19].

The syntax of TERM_n and TERM_{\omega}. The set $\{0,^*,\oplus\}$ of constant and operation symbols of involutive monoids is enriched by adding countably many variable symbols X_1, X_2, \ldots . Henceforth, the set $\mathcal{A} = \{0,^*, \oplus, X_1, X_2, \ldots,), (, \}$ is our alphabet. Parentheses are added to construct a non-ambiguous readable syntax. The set \mathcal{A}^* of strings over \mathcal{A} is defined by $\mathcal{A}^* = \{(s_1, s_2, ..., s_l) \in \mathcal{A}^l \mid l = 0, 1, 2, ...\}$. Let $n = 1, 2, \ldots$ By a term in the variables X_1, \ldots, X_n , we mean a string $\phi \in \mathcal{A}^*$ obtainable by the following inductive definition: (*) 0 and X_1, X_2, \ldots, X_n are terms; (**) if α and β are terms, then so are α^* and $(\alpha \oplus \beta)$. We let TERM_n be the set of terms in the variables X_1, X_2, \ldots, X_n . Elements of TERM_n are also known as n-variable Lukasiewicz formulas. We also let TERM_{\omega} = \int \mathbb{I}_n \text{TERM}_n.

Coding equivalence classes of projections by \mathbf{L}_{∞} -formulas. Fix $n = 1, 2, \ldots$ By McNaughton's theorem [15], the coordinate functions $\{\pi_1, \ldots, \pi_n\}$ are a distinguished free generating set of the free MV-algebra $\mathcal{M}([0,1]^n) \cong E(\mathfrak{M}_n)$. In view of Theorem 1.1(iv), the AF ℓ -algebra \mathfrak{M}_n is defined by $E(\mathfrak{M}_n) = \mathcal{M}([0,1]^n)$.

Arbitrarily pick representative projections $\mathsf{p}_1,\ldots,\mathsf{p}_n\in\mathfrak{M}_n$ such that the equivalence classes $[\mathsf{p}_1],\ldots,[\mathsf{p}_n]\in E(\mathfrak{M}_n)$ correspond to π_1,\ldots,π_n via Elliott's classification. We may naturally say that the variable symbol X_i codes both the coordinate function $\pi_i\in\mathcal{M}([0,1]^n)$ and the equivalence class $[\mathsf{p}_i]\in E(\mathfrak{M}_n)$. For definiteness, we will say that π_i is the interpretation of X_i in (the Elliott monoid of) \mathfrak{M}_n , in symbols, $X_i^{\mathfrak{M}_n}=\pi_i$. For every $\phi\in\mathsf{TERM}_n$, the interpretation $\phi^{\mathfrak{M}_n}$ of ϕ in (the Elliott monoid of) \mathfrak{M}_n is then defined by (*) $0^{\mathfrak{M}_n}=$ the constant zero function over $[0,1]^n$ and inductively, (**) $(\psi^*)^{\mathfrak{M}_n}=(\psi^{\mathfrak{M}_n})^*$, $(\alpha\oplus\beta)^{\mathfrak{M}_n}=(\alpha^{\mathfrak{M}_n}\oplus\beta^{\mathfrak{M}_n})$. We also say that ϕ codes $\phi^{\mathfrak{M}_n}$ in (the Elliott monoid of) \mathfrak{M}_n . By a traditional abuse of notation, the MV-algebraic operation symbols also denote their corresponding operations. More generally, let $\mathfrak{A}=\mathfrak{M}_n/\mathfrak{I}$ be an AF ℓ -algebra, for some ideal \mathfrak{I} of \mathfrak{M}_n . Let

$$i = K_0(\mathfrak{I}) \cap E(\mathfrak{M}_n). \tag{3}$$

be the ideal of $\mathcal{M}([0,1]^n)$ corresponding to \mathfrak{I} by Theorem 1.2(v). Via the identification $E(\mathfrak{M}_n/\mathfrak{I})=\mathcal{M}([0,1]^n)$ /i, the interpretation $\phi^{\mathfrak{A}}$ of ϕ in (the Elliott monoid of) \mathfrak{A} is defined by: $0^{\mathfrak{A}}=\{0\}\subseteq \mathfrak{A}, X_i^{\mathfrak{A}}=X_i^{\mathfrak{M}_n/\mathfrak{I}}=X_i^{\mathfrak{M}_n/\mathfrak{I}}=\pi_i/\mathfrak{i}$ and inductively, $(\psi^*)^{\mathfrak{A}}=(\psi^{\mathfrak{A}})^*, \quad (\alpha\oplus\beta)^{\mathfrak{A}}=(\alpha^{\mathfrak{A}}\oplus\beta^{\mathfrak{A}})$. The following identities are easily verified by induction on the syntactical complexity of terms (i.e., the number of symbols occurring in each term), using the unique readability property of the syntax: $(\psi^*)^{\mathfrak{A}}=\frac{(\psi^*)^{\mathfrak{M}_n}}{\mathfrak{i}}=\left(\frac{\psi^{\mathfrak{M}_n}}{\mathfrak{i}}\right)^*=\frac{(\psi^{\mathfrak{M}_n})^*}{\mathfrak{i}}$ and $(\alpha\oplus\beta)^{\mathfrak{A}}=\frac{(\alpha\oplus\beta)^{\mathfrak{M}_n}}{\mathfrak{i}}=\frac{(\alpha\oplus\beta)^{\mathfrak{M}_n}}{\mathfrak{i}}$

 $\frac{\alpha^{\mathfrak{M}_n}}{\mathfrak{i}} \oplus \frac{\beta^{\mathfrak{M}_n}}{\mathfrak{i}} = \frac{\alpha^{\mathfrak{M}_n} \oplus \beta^{\mathfrak{M}_n}}{\mathfrak{i}}$. We also say that ϕ codes $\phi^{\mathfrak{A}}$ in (the Elliott monoid of)

The interpretation $\phi^{\mathfrak{M}}$ of $\phi \in \mathsf{TERM}_{\omega}$ in (the Elliott monoid of) the universal $\mathsf{AF}\ell$ -algebra \mathfrak{M} and its quotients is similarly defined.

Decision problems for AFℓ-algebras. Fix a cardinal $\kappa = 1, 2, ..., \omega$. Let $\mathfrak{A} = \mathfrak{M}_{\kappa}/\mathfrak{I}$ for some ideal \mathfrak{I} of \mathfrak{M}_{κ} . (Here $\mathfrak{M}_{\omega} = \mathfrak{M}$.) The word problem P_1 of \mathfrak{A} is defined by $\mathsf{P}_1 = \{(\phi, \psi) \in \mathcal{A}^* \mid (\phi, \psi) \in \mathsf{TERM}_{\kappa}^2 \text{ and } \phi^{\mathfrak{A}} = \psi^{\mathfrak{A}}\}$. Intuitively, on input strings ϕ and ψ , P_1 checks if ϕ and ψ are strings in TERM_{κ} coding the same equivalence class of projections of \mathfrak{A} . Likewise, the order problem P_2 of \mathfrak{A} is the subset of \mathcal{A}^* given by $\{(\phi, \psi) \in \mathsf{TERM}_{\kappa}^2 \mid \phi^{\mathfrak{A}} \leq \psi^{\mathfrak{A}}\}$. Problem P_2 checks if ϕ codes an equivalence class of projections $\phi^{\mathfrak{A}}$ in \mathfrak{A} that precedes $\psi^{\mathfrak{A}}$ in the Murray-von Neumann order \preceq of projections in \mathfrak{A} . The eccentricity problem $\mathsf{P}_3 = \{(\phi, \psi) \in \mathsf{TERM}_{\kappa}^2 \mid \phi^{\mathfrak{A}} \sqsubseteq \psi^{\mathfrak{A}}\}$ of \mathfrak{A} checks whether ϕ codes an equivalence class of projections $\phi^{\mathfrak{A}}$ in \mathfrak{A} that precedes $\psi^{\mathfrak{A}}$ in the \sqsubseteq -order of \mathfrak{A} . Further, the zero problem $\mathsf{P}_4 = \{\phi \in \mathsf{TERM}_{\kappa} \mid \phi^{\mathfrak{A}} = 0\}$ of \mathfrak{A} checks if $\phi^{\mathfrak{A}} = 0$. The central projection problem P_5 of \mathfrak{A} checks if $\phi^{\mathfrak{A}}$ is an equivalence class of central projections in \mathfrak{A} . The nontrivial projection problem P_6 of \mathfrak{A} checks if $\phi^{\mathfrak{A}}$ different from 0 and 1. The central nontrivial projection problem P_7 of \mathfrak{A} checks if $\phi^{\mathfrak{A}}$ is an equivalence class of central projections of \mathfrak{A} other than 0 or 1.

Polytime problems and (Turing) reductions. For any formula $\phi \in \mathsf{TERM}_{\kappa}$ we let $||\phi|| =$ the number of occurrences of symbols in ϕ . Let $i = 1, \ldots, 7$. We say that P_i is decidable in polytime if there is a polynomial $q \colon \{0, 1, 2, \ldots\} \to \{0, 1, 2, \ldots\}$ and a Turing machine \mathcal{T} with the following property: Over any input string $\sigma \in \mathcal{A}^*$, machine \mathcal{T} decides if σ belongs to P_i within a number of steps $\leq q(||\sigma||)$. Given problems $\mathsf{P}', \mathsf{P}'' \subseteq \mathcal{A}^*$, a reduction ρ of P' to P'' is a map $\rho \colon \mathcal{A}^* \to \mathcal{A}^*$ such that for every string $\sigma \in \mathcal{A}^*$, $\sigma \in \mathsf{P}'$ iff $\rho(\sigma) \in \mathsf{P}''$. When ρ is computable in polynomial time we say it is a polytime reduction. Compare with [14, p.211].

Proposition 3.1. Let \mathfrak{A} be an AF ℓ -algebra whose Elliott monoid is finitely generated. Let \mathfrak{I} (resp., \mathfrak{J}) be an ideal of \mathfrak{M}_m (resp., of \mathfrak{M}_n) such that $\mathfrak{A} \cong \mathfrak{M}_m/\mathfrak{I} \cong \mathfrak{M}_m/\mathfrak{I}$. Let $P \subseteq \mathcal{A}^*$ be any problem among P_1, \ldots, P_7 . Then P for $\mathfrak{M}_m/\mathfrak{I}$ is polytime reducible to P for $\mathfrak{M}_n/\mathfrak{I}$. Thus in particular, problem P for $\mathfrak{M}_m/\mathfrak{I}$ is decidable in polytime iff so is P for $\mathfrak{M}_n/\mathfrak{I}$.

Proof. A rational polyhedron $P \subseteq [0,1]^n$ is the union of finitely many simplexes $T_i \subseteq [0,1]^n$ with rational vertices. Two rational polyhedra P and Q are \mathbb{Z} -homeomorphic if Q is the image of P under a piecewise linear homeomorphism η such that all linear pieces of η and η^{-1} have integer coefficients. Theorem 1.2 yields ideals i, j such that $E(\mathfrak{A}) \cong \mathcal{M}([0,1]^m)/\mathfrak{i} \cong \mathcal{M}([0,1]^n)/\mathfrak{j}$. Elliott's classification, and the \mathbb{Z} -homeomorphism in [19,8.7], (coded by an m-tuple ψ_{η} of n-variable terms) yields a polytime reduction ρ' : $\phi \in \mathsf{TERM}_m \mapsto \phi(\psi_{\eta}) \in \mathsf{TERM}_n$ of problem P_i for $\mathfrak{M}_m/\mathfrak{I}$ to P_i for $\mathfrak{M}_n/\mathfrak{I}$, and a polytime reduction ρ'' in the opposite direction. \square

Theorem 3.2. Fix an AF ℓ -algebra $\mathfrak A$ whose Elliott monoid is finitely generated. Then for all $i, j \in \{1, ..., 4\}$ problem P_i for $\mathfrak A$ is polytime reducible to problem P_j . Thus in particular, P_i for $\mathfrak A$ is polytime iff so is P_j for $\mathfrak A$.

Proof. Following [6, p.8], for every MV-algebra A we define the operation \ominus : $A^2 \to A$ by $x \ominus y = x \odot y^*$. By [6, Lemma 1.1.2],

$$x \le y \quad \text{iff} \quad x \ominus y = 0.$$
 (4)

For all input terms α, β , let us write for short $\alpha^{\mathfrak{A}} = a$ and $\beta^{\mathfrak{A}} = b$. By definition of the \sqsubseteq -order, $a \sqsubseteq b$ iff for every prime ideal \mathfrak{p} of the MV-algebra $E(\mathfrak{A})$ the following holds:

$$(b/\mathfrak{p} < b^*/\mathfrak{p} \text{ implies } a/\mathfrak{p} \le b/\mathfrak{p})$$
 and $(b/\mathfrak{p} > b^*/\mathfrak{p} \text{ implies } a/\mathfrak{p} \ge b/\mathfrak{p}).$ (5)

This is equivalent to $(b/\mathfrak{p} \leq b^*/\mathfrak{p})$ and $a/\mathfrak{p} \leq b/\mathfrak{p}$ or $(b/\mathfrak{p} \geq b^*/\mathfrak{p})$ and $a/\mathfrak{p} \geq b/\mathfrak{p}$, which by (4) can be reformulated as

$$(b/\mathfrak{p} \ominus b^*/\mathfrak{p} = 0 = a/\mathfrak{p} \ominus b/\mathfrak{p}) \text{ or } (b^*/\mathfrak{p} \ominus b/\mathfrak{p} = 0 = b/\mathfrak{p} \ominus a/\mathfrak{p}).$$
 (6)

By definition of the lattice operations in the MV-chain $E(\mathfrak{A})/\mathfrak{p}$, (Theorem 1.1(iii)), (5)-(6) equivalently state $((b/\mathfrak{p} \ominus b^*/\mathfrak{p}) \lor (a/\mathfrak{p} \ominus b/\mathfrak{p})) \land ((b^*/\mathfrak{p} \ominus b/\mathfrak{p}) \lor (b/\mathfrak{p} \ominus a/\mathfrak{p})) = 0$. By (1), the Elliott monoid $E(\mathfrak{A})$ satisfies: $a \sqsubseteq b$ iff $((b \ominus b^*) \lor (a \ominus b)) \land ((b^* \ominus b) \lor (b \ominus a)) = 0$. By definition of $a, b, a \sqsubseteq b$ iff $(((\beta \ominus \beta^*) \lor (\alpha \ominus \beta)) \land ((\beta^* \ominus \beta) \lor (\beta \ominus \alpha)))^{\mathfrak{A}} = 0$. We have just constructed a polytime reduction of the eccentricity problem P_3 for \mathfrak{A} to the zero problem P_4 for \mathfrak{A} . A converse polytime reduction is immediately obtained by noting that $\phi^{\mathfrak{A}} = [0]$ iff $\phi^{\mathfrak{A}} \sqsubseteq [0]$, whence P_4 is a subproblem of P_3 .

A polytime reduction of the order problem P_2 to the zero problem P_4 for $\mathfrak A$ trivially follows by noting that $\phi^{\mathfrak A} \preceq \psi^{\mathfrak A}$ iff $(\phi \ominus \psi)^{\mathfrak A} = 0$. Conversely, a polytime reduction of P_4 to P_2 for $\mathfrak A$ follows by noting that $\phi^{\mathfrak A} = 0$ iff $\phi^{\mathfrak A} \preceq 0$.

A polytime reduction of the word problem P_1 to the zero problem P_4 for \mathfrak{A} follows from $\phi^{\mathfrak{A}} = \psi^{\mathfrak{A}}$ iff $\operatorname{dist}(\phi^{\mathfrak{A}}, \psi^{\mathfrak{A}}) = 0$ iff $((\phi \ominus \psi) \oplus (\psi \ominus \phi))^{\mathfrak{A}} = 0$. Conversely, a polytime reduction of P_4 to P_1 for \mathfrak{A} trivially follows from the fact that the former problem is a special case of the latter.

Concerning the central projection problem P_5 we have:

Theorem 3.3. Arbitrarily fix an AF ℓ -algebra $\mathfrak A$ whose Elliott monoid $E(\mathfrak A)$ is finitely generated.

- (i) There is a polytime reduction of the central projection problem P_5 to the zero problem P_4 for $\mathfrak A$.
- (ii) The converse polytime reduction exists if $\mathfrak A$ has no quotient isomorphic to $\mathbb C$, or if $\mathfrak A$ is simple.

Proof. (i) As a matter of fact, with the notation of (2), we have:

 $\phi^{\mathfrak{A}}$ is the equivalence class of a central projection in ${\mathfrak{A}}$

iff $\phi^{\mathfrak{A}}$ belongs to $\mathsf{B}(E(\mathfrak{A}))$, by Theorem 2.1

iff
$$\phi^{\mathfrak{A}} \wedge (\phi^{\mathfrak{A}})^* = 0$$
, by [5, Theorem 1.16], (with \wedge as in Theorem 1.1(iii))

iff
$$(\phi \wedge \phi^*)^{\mathfrak{A}} = 0$$
.

The desired polytime reduction transforms ϕ into $\phi \wedge \phi^*$.

(ii) Let us first assume

$$\mathfrak{A}$$
 has no quotient isomorphic to \mathbb{C} . (7)

Pick an $n=1,2,\ldots$ and an ideal \Im of \mathfrak{M}_n such that $\mathfrak{A}\cong \mathfrak{M}_n/\Im$. By Proposition 3.1, our proof will not depend on the actual choice of n and \Im . Let $A=E(\mathfrak{A})$ and \mathfrak{i} be the ideal of $E(\mathfrak{M}_n)$ corresponding to \Im by Theorem 1.2(v), $\mathfrak{i}=K_0(\Im)\cap E(\mathfrak{M}_n)$. We have $E(\mathfrak{A})=E(\mathfrak{M}_n/\Im)=E(\mathfrak{M}_n)/\mathfrak{i}=\mathcal{M}([0,1]^n)/\mathfrak{i}=A$. Setting $X_i^{\mathfrak{A}}=\pi_i/\mathfrak{i}, \quad (i=1,\ldots,n),$ every $\phi\in\mathsf{TERM}_n$ is interpreted as $\phi^{\mathfrak{A}}=\phi^{\mathfrak{M}_n}/\mathfrak{i}$.

To obtain the desired polytime reduction, let us set

$$\rho \colon \phi \mapsto \phi \land \bigvee_{i=1}^{n} (X_i \land X_i^*), \quad \text{for each } \phi \in \mathsf{TERM}_n.$$
 (8)

We must prove

$$\phi^{\mathfrak{A}} = 0 \Leftrightarrow \left(\phi \wedge \bigvee_{i=1}^{n} (X_i \wedge X_i^*)\right)^{\mathfrak{A}} \in \{0, 1\}. \tag{9}$$

Indeed, by Theorem 2.1, equation (9) amounts to saying that $\phi^{\mathfrak{A}} = 0$ iff the term $\rho(\phi)$ given by $\phi \wedge \bigvee_{i=1}^{n} (X_i \wedge X_i^*)$ codes the equivalence class of a central projection of \mathfrak{A} .

The (\Rightarrow) direction of (9) is trivial. For the (\Leftarrow) direction, arguing by way of contradiction, let us assume

$$\left(\phi \wedge \bigvee_{i=1}^{n} (X_i \wedge X_i^*)\right)^{\mathfrak{A}} \subseteq \{0,1\} \quad \text{and} \quad \phi^{\mathfrak{A}} \neq 0. \tag{10}$$

By (1), for some prime ideal \mathfrak{p} of A we have

$$\phi^{\mathfrak{A}}/\mathfrak{p} \neq 0$$
 in the MV-chain A/\mathfrak{p} . (11)

On the other hand, every prime ideal \mathfrak{q} satisfies $(\phi \wedge \bigvee_{i=1}^{n} (X_i \wedge X_i^*))^{\mathfrak{A}}/\mathfrak{q} \in \{0,1\}$ in the MV-chain A/\mathfrak{q} . In particular,

$$\frac{\left(\phi \wedge \bigvee_{i=1}^{n} (X_i \wedge X_i^*)\right)^{\mathfrak{A}}}{\mathfrak{p}} \in \{0, 1\} \text{ in the MV-chain } A/\mathfrak{p} . \tag{12}$$

Another application of Theorem 1.2 yields a unique primitive ideal \mathfrak{P} of \mathfrak{A} such that $E(\mathfrak{A}/\mathfrak{P}) = E(\mathfrak{A})/\mathfrak{p}$

Claim: For each i = 1, ..., n the interpretation $X_i^{\mathfrak{A}}/\mathfrak{p}$ of the variable symbol X_i in $\mathfrak{A}/\mathfrak{P}$, as well as in its Elliott monoid $\in A/\mathfrak{p}$, is a trivial element 0 or 1.

For otherwise (absurdum hypothesis), say without loss of generality

$$X_1^{\mathfrak{A}}/\mathfrak{p} \notin \{0,1\}$$
, whence also $(X_1^*)^{\mathfrak{A}}/\mathfrak{p}$ does not belong to $\{0,1\}$. (13) By (12),

$$\{0,1\} \quad \ni \quad \frac{\left(\phi \wedge \bigvee_{i=1}^{n} (X_{i} \wedge X_{i}^{*})\right)^{\mathfrak{A}}}{\mathfrak{p}}$$

$$= \quad \frac{\phi^{\mathfrak{A}} \wedge \left(\bigvee_{i=1}^{n} (X_{i} \wedge X_{i}^{*})\right)^{\mathfrak{A}}}{\mathfrak{p}}$$

$$= \quad \frac{\phi^{\mathfrak{A}}}{\mathfrak{p}} \wedge \bigvee_{i=1}^{n} \frac{\left(X_{i} \wedge X_{i}^{*}\right)^{\mathfrak{A}}}{\mathfrak{p}}.$$

In the MV-chain A/\mathfrak{p} , from (13) we get

$$0<\frac{(X_1\wedge X_1^*)^{\mathfrak{A}}}{\mathfrak{p}}<1 \text{ and for each } j=2,\dots,n,\ \frac{(X_j\wedge X_j^*)^{\mathfrak{A}}}{\mathfrak{p}}<1.$$

As a consequence, $0 < \left(\bigvee_{i=1}^n \frac{(X_i \wedge X_i^*)^{\mathfrak{A}}}{\mathfrak{p}}\right) < 1$. From (11), we obtain $0 < \frac{\phi^{\mathfrak{A}}}{\mathfrak{p}} \wedge \bigvee_{i=1}^n \frac{(X_i \wedge X_i^*)^{\mathfrak{A}}}{\mathfrak{p}} < 1$, against (10).

Having thus settled our claim, for each $i=1,\ldots,n$ we have $X_i^{\mathfrak{A}}/\mathfrak{p}\in\{0,1\}$ in the MV-chain A/\mathfrak{p} . Let \mathfrak{n} be the only maximal ideal of A above the prime ideal \mathfrak{p} , as given by [6, 1.2.12]. Then a fortiori, $\left\{\frac{X_i^{\mathfrak{A}}}{\mathfrak{n}}, \frac{X_i^{*\mathfrak{A}}}{\mathfrak{n}}\right\} = \{0,1\}$ in $\frac{A}{\mathfrak{n}}$ for each $i=1,\ldots,n$. Since the set of elements $\{X_1^{\mathfrak{A}}/\mathfrak{n},\ldots,X_n^{\mathfrak{A}}/\mathfrak{n}\}$ generates the simple MV-chain A/\mathfrak{n} , we obtain $A/\mathfrak{n}=\{0,1\}$ = the set of trivial elements of A/\mathfrak{n} . In correspondence with the maximal ideal \mathfrak{n} of A, Theorem 1.2(v) yields a maximal ideal \mathfrak{N} of \mathfrak{A} such that $\mathfrak{A}/\mathfrak{N} \cong \mathbb{C}$. This contradicts our standing assumption (7), thus

proving (9). The map ρ defined in (8) is the desired polytime reduction of problem P_4 to P_5 for $\mathfrak A$ whenever $\mathfrak A$ has no quotient isomorphic to $\mathbb C$.

To conclude the proof, there remains to consider the case when $\mathfrak A$ is simple.

If $\mathfrak{A} \cong \mathbb{C}$ then both problems P_4 and P_5 are polytime, whence they are (trivially) polytime reducible each to the other. On the other hand, if $\mathfrak{A} \ncong \mathbb{C}$, the assumed simplicity of \mathfrak{A} ensures that \mathfrak{A} has no quotient isomorphic to \mathbb{C} . The desired polytime reduction of problem P_4 to P_5 for \mathfrak{A} is then obtained arguing as in the first part of the proof of (ii).

Corollary 3.4. Let \mathfrak{M} be the universal AF- algebra of [16, §8].

- (i) For each i = 1, ..., 5, problem P_i for \mathfrak{M} is coNP-complete.
- (ii) The nontriviality problem P₆ is NP-complete.
- (iii) If there is a polytime reduction of P_i to P_6 then NP = coNP.
- (iv) The central nontrivial projection problem P₇ is (trivially) polytime.

Corollary 3.5. Let \mathfrak{M}_1 be the "Farey" AF ℓ -algebra of [17, 3, 18].

Problems P_1, \ldots, P_7 for \mathfrak{M}_1 are decidable in polytime.

Corollary 3.6. Problems P_1, \ldots, P_7 are decidable in polynomial time for the following $AF\ell$ -algebras:

- (i) The Effros-Shen algebra \mathfrak{F}_{θ} for θ a quadratic irrational, or $\theta = 1/e$, or $\theta \in [0,1] \setminus \mathbb{Q}$ a real algebraic integer.
- (ii) The Effros-Shen algebra \mathfrak{F}_{θ} for any irrational $\theta \in [0,1]$ having the following property: There is a real $\kappa > 0$ such that for every $n = 0, 1, \ldots$ the sequence $[a_0, \ldots, a_n]$ of partial quotients of θ is computable (as a finite list of binary integers) in less than $2^{\kappa n}$ steps.

Corollary 3.7. Problems P_1, \ldots, P_7 are decidable in polynomial time for all Behncke-Leptin [2] algebras $A_{m,n}$.

References

- C. D. Aliprantis, O. Burkinshaw, Locally solid Riesz spaces, Pure and Applied Mathematics, Vol. 76, Academic Press, NY, 1978.
- [2] H. Behncke, H. Leptin, C*-algebras with a two-point dual, J. Functional Analysis, 10 (1972) 330–335.
- [3] F. Boca, An AF algebra associated with the Farey tessellation, Canad. J. Math, 60 (2008) 975–1000.
- [4] O. Bratteli, Inductive limits of finite-dimensional C*-algebras, Trans. Amer. Math. Soc., 171 (1972) 195-234.
- [5] C. C. Chang, Algebraic analysis of many valued logics, Trans. Amer. Math. Soc., 88 (1958) 467-490.
- [6] R. Cignoli, I. M. L. D'Ottaviano, D. Mundici, Algebraic Foundations of many-valued Reasoning, Trends in Logic, Vol.7, Kluwer, Dordrecht, 2000. Reprinted, Springer Science & Business Media, (2013).
- [7] K. R. Davidson, C*-Algebras by Example, Fields Institute Monographs, Vol. 6, 1996.
- [8] E. G. Effros, Dimensions and C*-algebras, CBMS Regional Conf. Series in Math., Vol. 46, American Mathematical Society, Providence, RI, 1981.
- [9] G. A. Elliott, On the classification of inductive limits of sequences of semisimple finitedimensional algebras, J. Algebra, 38 (1976) 29-44.
- [10] H. Freudenthal, Teilweise geordnete Moduln, Proc. Akad. Wetenschappen Amsterdam, 39 (1936) 641-651.
- [11] K. R. Goodearl, Notes on Real and Complex C*-Algebras, Shiva Math. Series, Vol. 5, Birkhäuser, Boston, 1982.
- [12] K. R. Goodearl, D. E. Handelman, J. W. Lawrence, Affine Representations of Grothendieck Groups and Applications to Rickart C*-Algebras and ℵ₀-Continuous Regular Rings, Memoirs of the American Mathematical Society, 1980.

- [13] W. A. J. Luxemburg, A. C. Zaanen, Riesz spaces, Vol. I., North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., New York, 1971.
- $\left[14\right]\,$ M. Machtey, P. Young, An introduction to the general theory of algorithms. The computer science library. Theory of computation series. North-Holland, New York, Oxford, and Shannon, 1978. Published online by Cambridge University Press: 12 March 2014.
- [15] R. McNaughton, A theorem about infinite-valued sentential logic, J. Symbolic Logic, 16 (1951) 1-13.
- [16] D. Mundici, Interpretation of AF C*-algebras in Lukasiewicz sentential calculus, J. Functional Analysis, 65 (1986) 15-63.
- [17] D. Mundici, Farey stellar subdivisions, ultrasimplicial groups, and K_0 of AF C*-algebras, Advances in Mathematics, 68 (1988) 23-39.
- [18] D. Mundici, Recognizing the Farey-Stern-Brocot AF algebra, in memoriam Renato Caccioppoli, Rendiconti Lincei Mat. Appl., 20 (2009) 327-338.
- [19] D. Mundici, Advanced Łukasiewicz calculus and MV-algebras, Trends in Logic, Vol. 35, Springer, Berlin, 2011.
- [20] D. Mundici, G. Panti, Extending addition in Elliott's local semigroup, Journal of Functional Analysis, 117 (1993) 461-471.
- (D. Mundici) Department of Mathematics and Computer Science "Ulisse Dini", Uni-VERSITY OF FLORENCE, VIALE MORGAGNI 67/A, I-50134 FLORENCE, ITALY

Email address: daniele.mundici@unifi.it