

# PRESCRIBED WEBSTER SCALAR CURVATURE FLOW ON CR MANIFOLD

PAK TUNG HO, QUỐC ANH NGÔ, AND HONG ZHANG

ABSTRACT. Using the prescribed Webster scalar curvature flow, we prove some existence results on the 3-dimensional compact CR manifold with nonnegative CR Yamabe constant.

## 1. INTRODUCTION

Let  $(M, \theta_0)$  be a compact strictly pseudoconvex CR manifold of real dimension  $2n + 1$  with a background contact form  $\theta_0$ . Similar to the Yamabe problem, the CR Yamabe problem is to find a contact form  $\theta$  conformal to  $\theta_0$  such that its Webster scalar curvature is constant. This was first introduced by Jerison and Lee in [10], and was solved in [4, 5, 9, 11, 12, 20].

The CR Yamabe flow is the geometric flow introduced to solve the CR Yamabe problem. It is defined as the evolution of  $\theta(t)$ :

$$\frac{\partial}{\partial t} \theta(t) = -(R_{\theta(t)} - \bar{R}_{\theta(t)})\theta(t), \quad (1.1)$$

where  $R_{\theta(t)}$  is the Webster scalar curvature of the contact form  $\theta(t)$ , and  $\bar{R}_{\theta(t)}$  is the average of  $R_{\theta(t)}$ :

$$\bar{R}_{\theta(t)} = \frac{\int_M R_{\theta(t)} \theta_0 \wedge (d\theta_0)^n}{\int_M \theta_0 \wedge (d\theta_0)^n}.$$

We refer the readers to [1–3, 13, 16, 21, 22, 29] for results related to the CR Yamabe flow.

Let  $f$  be a smooth function on  $M$ . As a generalization of the CR Yamabe problem, the prescribed Webster scalar curvature problem is to find a contact form  $\theta$  conformal to  $\theta_0$  such that its Webster scalar curvature  $R_\theta$  is the given function  $f$ . If we write  $\theta = u^{2/n}\theta_0$ , then the prescribed Webster scalar curvature problem is equivalent to solving the semi-linear equation

$$-(2 + 2/n)\Delta_{\theta_0} u + R_{\theta_0} u = f u^{1+2/n}, \quad (1.2)$$

where  $\Delta_{\theta_0}$  and  $R_{\theta_0}$  are, respectively, the sub-Laplacian and the Webster scalar curvature of the contact form  $\theta_0$ . See [6, 8, 19, 24, 26–28] and references therein for results related to the prescribed Webster scalar curvature problem.

Similar to the CR Yamabe flow, the prescribed Webster scalar curvature flow is introduced to study the prescribed Webster scalar curvature problem, and is defined as

$$\frac{\partial}{\partial t} \theta(t) = -(R_{\theta(t)} - \lambda(t)f)\theta(t).$$

---

*Date:* 4<sup>th</sup> May, 2021 at 01:17.

*2020 Mathematics Subject Classification.* 32V05, 32V20, 53E99.

*Key words and phrases.* Webster scalar curvature; CR Yamabe problem; flow.

Note that when  $f \equiv 1$ , the prescribed Webster scalar curvature flow reduces to the CR Yamabe flow. In [14, 17, 18], the first author studied the prescribed Webster scalar curvature flow on the sphere  $S^{2n+1}$ , when the prescribed function  $f$  is positive. On the other hand, in [15], the first author studied the prescribed Webster scalar curvature flow on the 3-dimensional CR manifold with negative CR Yamabe constant, when  $f$  is assumed to be negative.

In this paper, we study the prescribing Webster scalar problem on 3-dimensional CR manifold with nonnegative CR Yamabe constant. By using the prescribed Webster scalar flow, we prove some existence results. Unlike the results mentioned above, the prescribed function  $f$  here in our existence results is allowed to change sign. More precisely, our main results read as

**Theorem 1.1.** *Let  $(M, \theta_0)$  be a compact strictly pseudoconvex CR manifold of real dimension 3 with positive CR Yamabe constant. Suppose that  $M$  is not conformally equivalent to the CR sphere  $\mathbb{S}^3$  and the CR Paneitz operator of  $M$  is nonnegative. Also, let  $f$  be a smooth function on  $(M, \theta_0)$  with  $\sup_M f > 0$ . Then there exists a positive constant  $c_*$  depending only on  $(M, \theta_0)$  such that if*

$$\left\{ x \in M : f(x) = \sup_M f \right\} \subseteq \left\{ x \in M : f(x) > 0 \text{ and } \Delta_{\theta_0} f(x)/f(x) > -c_* \right\}, \quad (1.3)$$

then there exist a positive solution to the prescribed Webster scalar curvature equation (1.2).

Comparing Theorem 1.1 with the result of Escobar and Schoen in [7] where  $f$  is only assumed to be positive somewhere in the dimension 3, one can see that we need an extra condition (1.3), which shows the difference between the Riemannian case and the CR case. The reason why we need this extra condition (1.3) is as follows: In view of Lemma 5.15 below, the dominant term is  $O(\varepsilon^{2n})$ , and the expansion of the prescribed function  $f$  around a blow-up point  $a$  is something like  $f(a) + A\varepsilon + B\varepsilon^2 + O(\varepsilon^3)$ , where  $A$  and  $B$  involve the gradient and Laplacian of  $f$  at  $a$ . When  $n = 1$ , we can see that the dominant term is  $O(\varepsilon^2)$ , but  $\varepsilon^2$  also appears in the expansion of  $f$ . Since  $B$  is generally not small, we would lose control if we do not impose any extra condition on  $f$ . While in the Riemannian case, the dominant term is  $O(\varepsilon^{n-2})$ , and the expansion of the prescribed function  $f$  is the same as above. When  $n = 3$ , the dominant term becomes  $\varepsilon$ . Although  $\varepsilon$  appears in the expansion of  $f$ , the coefficient  $A$  is  $o(1)$  as time goes to infinity, since  $A$  involves only gradient of  $f$  at  $a$ . Therefore, in the Riemannian case, one has full control without any extra condition.

**Theorem 1.2.** *Let  $(M, \theta_0)$  be a compact strictly pseudoconvex CR manifold of real dimension 3 with vanishing CR Yamabe constant. Also, let  $f$  be a smooth function on  $(M, \theta_0)$  satisfying*

$$\sup_M f > 0 \text{ and } \int_M f \theta_0 \wedge (d\theta_0)^n < 0. \quad (1.4)$$

Then there exists a positive solution to the prescribed Webster scalar curvature equation (1.2).

We remark that Theorem 1.2 is sharp in the following sense: if (1.2) is solvable on  $(M, \theta_0)$  with vanishing CR Yamabe constant, then either  $f \equiv 0$  or  $f$  must satisfies (1.4). To see this, suppose that  $u$  is a positive solution to (1.2) with  $f \not\equiv 0$ .

Since  $(M, \theta_0)$  has vanishing CR Yamabe constant, we may assume, by a conformal change if necessary, that  $R_{\theta_0} \equiv 0$ . Then (1.2) reduces to

$$-(2 + 2/n)\Delta_{\theta_0}u = fu^{1+2/n}. \quad (1.5)$$

Integrating (1.5) over  $M$  yields  $\int_M fu^{1+2/n}\theta_0 \wedge (d\theta_0)^n = 0$ . Since  $u > 0$  and  $f \not\equiv 0$ , this implies that  $f$  changes sign, and in particular,  $\sup_M f > 0$ . On the other hand, multiplying  $u^{-1-2/n}$  to (1.5) and integrating it over  $M$ , we obtain

$$\int_M f\theta_0 \wedge (d\theta_0)^n = -(2 + 2/n)(1 + 2/n) \int_M u^{-2-2/n}|\nabla_{\theta_0}u|_{\theta_0}\theta_0 \wedge (d\theta_0)^n < 0.$$

We also remark that most of the arguments in the proof of Theorems 1.1 and 1.2 are still valid in dimension  $2n + 1$ . In fact, we wonder if Theorems 1.1 and 1.2 are still true when the dimension is  $2n + 1$ . We hope to come back to this in future. As a result, we will try to write down the proof in the general case when the dimension is  $2n + 1$ , and point out exactly the part when the assumption  $n = 1$  is needed.

The method in this paper is inspired by Mayer [25], in which he considered the prescribed scalar curvature problem on a closed  $n$ -dimensional Riemannian manifold of positive Yamabe invariant, where  $3 \leq n \leq 5$ .

#### ACKNOWLEDGEMENT

The first author was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (Grant No. 2020R1A6A1A03047877 and 2019R1F1A1041021). The third author would like to acknowledge the support from the Young Scientists Fund of the NSFC (Grant No. 11701544).

## 2. PRESCRIBED WEBSTER SCALAR CURVATURE FLOW

Let  $(M, \theta_0)$  be a compact strictly pseudoconvex CR manifold of real dimension  $2n + 1$  with a contact form  $\theta_0$ ,  $f$  a smooth function on  $M$ . As in Theorems 1.1 and 1.2, the CR Yamabe constant of  $(M, \theta_0)$  is assumed to be nonnegative. Therefore, by a conformal change if necessary, we may assume that the Webster scalar curvature of  $\theta_0$  is positive (is zero respectively) when the CR Yamabe constant is positive (is zero respectively).

We consider the prescribed Webster scalar curvature flow, which is the evolution equation of  $\theta(t)$  satisfying

$$\begin{cases} \frac{\partial}{\partial t}\theta(t) = -(R_{\theta(t)} - \lambda(t)f)\theta(t), \\ \theta(0) = \theta^0, \end{cases} \quad (2.1)$$

where  $R_{\theta(t)}$  is the Webster scalar curvature of the contact form  $\theta(t)$ ,  $\lambda(t)$  is to be determined and  $\theta^0$  is an initial conformal contact form. If we write  $\theta(t) = u(t)^{2/n}\theta_0$  and  $\theta^0 = u_0^{2/n}\theta_0$  for some smooth functions  $u(t)$  and  $u_0$ , then we can obtain an evolution equation for  $u(t)$  from (2.1).

$$\begin{cases} \frac{\partial}{\partial t}u(t) = -\frac{n}{2}(R_{\theta(t)} - \lambda(t)f)u(t), \\ u(0) = u_0. \end{cases} \quad (2.2)$$

Here, in terms of  $u$ ,  $R_{\theta(t)}$  can be written as

$$R_{\theta(t)} = u^{-1-\frac{2}{n}} \left( - (2 + 2/n) \Delta_{\theta_0} u(t) + R_{\theta_0} u(t) \right). \quad (2.3)$$

To deal with the case where  $f$  changes sign, it is convenient to consider the energy functional associated with the prescribed Webster scalar curvature problem

$$\mathcal{E}(u) = \int_M \left( (2 + 2/n) |\nabla_{\theta_0} u|_{\theta_0}^2 + R_{\theta_0} u^2 \right) \theta_0 \wedge (d\theta_0)^n \quad (2.4)$$

with the constraint condition

$$X_f := \left\{ u \in \mathbf{S}_1^2(M) : u \geq 0, \int_M f u^{2+2/n} \theta_0 \wedge (d\theta_0)^n = 1 \right\}. \quad (2.5)$$

The constraint condition (2.5) above reminds us that we should keep the quantity  $\int_M f u(t)^{2+2/n} \theta_0 \wedge (d\theta_0)^n$  unchanged during the evolution. That is

$$\begin{aligned} 0 &= \frac{d}{dt} \int_M f u(t)^{2+2/n} \theta_0 \wedge (d\theta_0)^n = (2 + 2/n) \int_M f u_t u(t)^{1+2/n} \theta_0 \wedge (d\theta_0)^n \\ &= -(n+1) \int_M \left( f R_{\theta(t)} - \lambda(t) f^2 \right) \theta \wedge (d\theta)^n. \end{aligned} \quad (2.6)$$

Solving (2.6), we find that

$$\lambda(t) = \frac{\int_M f R_{\theta(t)} \theta \wedge (d\theta)^n}{\int_M f^2 \theta \wedge (d\theta)^n}. \quad (2.7)$$

If we let  $u_0 \in X$ , from (2.6) we then have

$$\int_M f u(t)^{2+2/n} \theta_0 \wedge (d\theta_0)^n \equiv 1. \quad (2.8)$$

To end this section, we collect some well-known results for later reference.

- (i) Given a closed CR-manifold  $(M, \theta)$ , the CR-Yamabe constant, denoted by  $\mu(M)$ , is defined by (see [10] for example)

$$\mu(M) = \inf_{\{u \geq 0, 0 \neq u \in \mathbf{S}_1^2(M)\}} \frac{\int_M (2 + 2/n) |\nabla_{\theta} u|^2 + R_{\theta} u^2 \theta \wedge (d\theta)^n}{\left( \int_M u^{2+2/n} \theta \wedge (d\theta)^n \right)^{\frac{n}{n+1}}}. \quad (2.9)$$

Here,  $\mathbf{S}_1^2(M)$  is the Folland-Stein space of  $(M, \theta)$ .

- (ii) Given any  $\varepsilon > 0$ , there exist  $B > 0$  such that for  $u \in \mathbf{S}_1^2(M)$

$$\begin{aligned} \left( \int_M u^{2+2/n} \theta \wedge (d\theta)^n \right)^{\frac{n}{n+1}} &\leq (K_n + \varepsilon) \int_M |\nabla_{\theta} u|^2 \theta \wedge (d\theta)^n \\ &\quad + B \int_M u^2 \theta \wedge (d\theta)^n, \end{aligned} \quad (2.10)$$

where  $K_n = 1/(2\pi n^2)$  (see Theorem A.1 in [20]). In particular, there exists a positive constant  $\mathcal{C}$  such that

$$\left( \int_M u^{2+2/n} \theta \wedge (d\theta)^n \right)^{\frac{n}{n+1}} \leq \mathcal{C} \left( \int_M (|\nabla_{\theta} u|^2 + u^2) \theta \wedge (d\theta)^n \right). \quad (2.11)$$

## 3. GLOBAL EXISTENCE OF THE FLOW

**Lemma 3.1.** *For a smooth solution  $u(t)$  of the flow (2.2) there holds*

$$\frac{d}{dt} \mathcal{E}(u(t)) = -n \int_M (R_{\theta(t)} - \lambda(t)f)^2 \theta \wedge (d\theta)^n.$$

*In particular, the energy function  $\mathcal{E}$  is non-increasing along the flow.*

**Proof.** By (2.2) and (2.3), we can compute

$$\begin{aligned} \frac{d}{dt} \mathcal{E}(u(t)) &= 2 \int_M (2 + 2/n) \langle \nabla_{\theta_0} u, \nabla_{\theta_0} u_t \rangle + R_{\theta_0} u u_t \theta_0 \wedge (d\theta_0)^n \\ &= 2 \int_M \left( - (2 + 2/n) \Delta_{\theta_0} u + R_{\theta_0} \right) u_t \theta_0 \wedge (d\theta_0)^n \\ &= -n \int_M R_{\theta(t)} (R_{\theta(t)} - \lambda(t)f) \theta \wedge (d\theta)^n \\ &= -n \int_M (R_{\theta(t)} - \lambda(t)f)^2 \theta \wedge (d\theta)^n, \end{aligned}$$

where we have used (2.6) in the last equality. □

Lemma 3.1 implies a uniform priori bound

$$\mathcal{E}(u(t)) + n \int_0^t \int_M (R_{\theta(t)} - \lambda(t)f)^2 \theta \wedge (d\theta)^n dt \leq \mathcal{E}(u_0), \quad \text{for all } t > 0. \quad (3.1)$$

Now, we derive the evolution equation for the Webster scalar curvature  $R_{\theta(t)}$ .

**Lemma 3.2.** *Assume that  $u(t)$  is a smooth solution of the flow (2.2). Then one has*

$$\frac{\partial}{\partial t} (\lambda(t)f - R_{\theta(t)}) = (n+1) \Delta_{\theta(t)} (\lambda(t)f - R_{\theta(t)}) + (\lambda(t)f - R_{\theta(t)}) R_{\theta(t)} + \lambda'(t)f,$$

where  $\lambda'(t)$  is given by

$$\begin{aligned} \lambda'(t) &= - \left( \int_M f^2 \theta \wedge (d\theta)^n \right)^{-1} \left[ (n+1) \int_M f \Delta_{\theta(t)} (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \right. \\ &\quad \left. + n \int_M f (\lambda(t)f - R_{\theta(t)})^2 \theta \wedge (d\theta)^n + \lambda(t) \int_M f^2 (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \right]. \end{aligned}$$

**Proof.** Using (2.3) and (2.2) we have

$$\begin{aligned}
\partial_t R_{\theta(t)} &= \left( -1 - \frac{2}{n} \right) u^{-2-\frac{2}{n}} u_t \left( -(2+2/n)\Delta_{\theta_0} u + R_{\theta_0} u \right) \\
&\quad + u^{-1-\frac{2}{n}} \left( -(2+2/n)\Delta_{\theta_0} u_t + R_{\theta_0} u_t \right) \\
&= \left( -\frac{n}{2} - 1 \right) (\lambda(t)f - R_{\theta(t)}) R_{\theta(t)} - (n+1) u^{-1-\frac{2}{n}} \Delta_{\theta_0} \left( (\lambda(t)f - R_{\theta(t)}) u \right) \\
&\quad + \frac{n}{2} u^{-\frac{2}{n}} R_{\theta_0} (\lambda(t)f - R_{\theta(t)}) \\
&= \left( -\frac{n}{2} - 1 \right) (\lambda(t)f - R_{\theta(t)}) R_{\theta(t)} - (n+1) u^{-1-\frac{2}{n}} \left[ u \Delta_{\theta_0} (\lambda(t)f - R_{\theta(t)}) \right. \\
&\quad \left. + 2 \langle \nabla_{\theta_0} (\lambda(t)f - R_{\theta(t)}), \nabla_{\theta_0} u \rangle \right] + \left[ -(n+1) u^{-1-\frac{2}{n}} (\lambda(t)f - R_{\theta(t)}) \Delta_{\theta_0} u \right. \\
&\quad \left. + \frac{n}{2} u^{-\frac{2}{n}} R_{\theta_0} (\lambda(t)f - R_{\theta(t)}) \right] \\
&= \left( -\frac{n}{2} - 1 \right) (\lambda(t)f - R_{\theta(t)}) R_{\theta(t)} - (n+1) \Delta_{\theta} (\lambda(t)f - R_{\theta(t)}) \\
&\quad + \frac{n}{2} (\lambda(t)f - R_{\theta(t)}) R_{\theta(t)} \\
&= -(n+1) \Delta_{\theta} (\lambda(t)f - R_{\theta(t)}) - (\lambda(t)f - R_{\theta(t)}) R_{\theta(t)}.
\end{aligned}$$

Notice that in the second last equality above we have used the identity

$$\Delta_{\theta} \varphi = u^{-1-\frac{2}{n}} \left( u \Delta_{\theta_0} \varphi + 2 \langle \nabla_{\theta_0} \varphi, \nabla_{\theta_0} u \rangle \right) \text{ for any } \varphi.$$

As for  $\lambda'(t)$ , we use (2.7), the evolution equation for  $R_{\theta(t)}$  above and (2.2) to calculate it as follows.

$$\begin{aligned}
\lambda'(t) &= \left( \int_M f^2 \theta \wedge (d\theta)^n \right)^{-1} \left[ \int_M f \partial_t R_{\theta(t)} \theta \wedge (d\theta)^n + \int_M f R_{\theta(t)} \partial_t (u^{2+2/n}) \theta_0 \wedge (d\theta_0)^n \right] \\
&\quad - \left( \int_M f^2 \theta \wedge (d\theta)^n \right)^{-2} \int_M f R_{\theta(t)} \theta \wedge (d\theta)^n \int_M f^2 \partial_t (u^{2+2/n}) \theta_0 \wedge (d\theta_0)^n \\
&= \left( \int_M f^2 \theta \wedge (d\theta)^n \right)^{-1} \left[ -(n+1) \int_M f \Delta_{\theta(t)} (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \right. \\
&\quad \left. + n \int_M f R_{\theta(t)} (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n - (n+1) \lambda(t) \int_M f^2 (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \right] \\
&= - \left( \int_M f^2 \theta \wedge (d\theta)^n \right)^{-1} \left[ (n+1) \int_M f \Delta_{\theta(t)} (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \right. \\
&\quad \left. + n \int_M f (\lambda(t)f - R_{\theta(t)})^2 \theta \wedge (d\theta)^n + \lambda(t) \int_M f^2 (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \right].
\end{aligned}$$

□

**Lemma 3.3.** *There exist two positive constant  $c_1$  and  $c_2$  such that*

$$c_1 \leq \text{vol}(M, \theta(t)) \leq c_2,$$

where  $c_1$  depends only on  $f, M$ , and  $c_2$  depends only on  $n, f, M, u_0$ .

**Proof.** First, we show the lower bound. To see this, from (2.8) it follows that

$$1 = \int_M f \theta \wedge (d\theta)^n \leq \left( \sup_M f \right) \text{vol}(M, \theta(t)).$$

This implies that

$$\text{vol}(M, \theta(t)) \geq \frac{1}{\sup_M f} =: c_1.$$

Next, we prove the upper bound. We split the argument into two cases.

**Case 1.**  $R_{\theta_0} > 0$ , i.e., positive case. In this case, it follows from Folland-Stein embedding (2.11) and the energy bound (3.1) that

$$\text{vol}(M, \theta(t)) = \|u\|_{L^{2+2/n}}^{2+2/n} \leq c_0 \mathcal{E}(u)^{1+1/n} \leq c_0 \mathcal{E}(u_0)^{1+1/n},$$

for some positive constant  $c_0$ .

**Case 2.**  $R_{\theta_0} \equiv 0$ , i.e., null case. In this case, we first use Poincaré type inequality (see [23] for example) and (3.1) to obtain

$$\|u - \bar{u}\|_{L^{2+2/n}} \leq c_0 \|\nabla_{\theta_0} u\|_{L^2} = c_0 \sqrt{\frac{n \mathcal{E}(u)}{2n+2}} \leq c_0 \sqrt{\frac{n \mathcal{E}(u_0)}{2n+2}}. \quad (3.2)$$

It remains to bound  $\bar{u}$ . Noting that  $\int_M f \theta_0 \wedge (d\theta_0)^n < 0$ , we have by (2.8) that

$$\begin{aligned} \bar{u}^{2+2/n} \int_M -f \theta_0 \wedge (d\theta_0)^n &= -1 + \int_M f(u^{2+2/n} - \bar{u}^{2+2/n}) \theta_0 \wedge (d\theta_0)^n \\ &\leq \sup_M |f| \int_M |u^{2+2/n} - \bar{u}^{2+2/n}| \theta_0 \wedge (d\theta_0)^n \end{aligned} \quad (3.3)$$

Recall that for any  $a, b > 0$ , we have the elementary inequality

$$\begin{aligned} |a^{2+2/n} - b^{2+2/n}| &\leq (2+2/n)(a+b)^{1+2/n}|a-b| \\ &\leq (2+2/n)2^{2/n}(|a-b|^{1+2/n} + (2b)^{1+2/n})|a-b| \\ &\leq c(n)(|a-b|^{2+2/n} + b^{1+2/n}|a-b|), \end{aligned}$$

where  $c(n)$  is a positive constant depending only on  $n$ . Applying this inequality with  $a = u$  and  $b = \bar{u}$  we get

$$|u^{2+2/n} - \bar{u}^{2+2/n}| \leq c(n)(|u - \bar{u}|^{2+2/n} + \bar{u}^{1+2/n}|u - \bar{u}|). \quad (3.4)$$

Substituting (3.4) into (3.3) and using Hölder's inequality and Young's inequality, we obtain

$$\begin{aligned} \bar{u}^{2+2/n} &\leq \frac{c(n) \sup_M f}{\int_M -f \theta_0 \wedge (d\theta_0)^n} (\|u - \bar{u}\|_{L^{2+2/n}}^{2+2/n} + \bar{u}^{1+2/n} \text{vol}(M, \theta_0)^{\frac{n+2}{2(n+1)}} \|u - \bar{u}\|_{L^{2+2/n}}) \\ &\leq \frac{1}{2} \bar{u}^{2+2/n} + c(n, f, M) \|u - \bar{u}\|_{L^{2+2/n}}^{2+2/n}, \end{aligned}$$

for some positive constant  $c(n, f, M)$  depending only on  $n, f$  and  $M$ . This together with (3.2) yields

$$\bar{u} \leq c(n, f, M, u_0),$$

for some positive constant  $c(n, f, M, u_0)$  depending only on  $n, f, M$  and  $u_0$ . Then from Minkowski's inequality and (3.2), it follows that there exists a positive constant  $c_2$  depending only on  $n, f, M$  and  $u_0$  such that

$$\text{vol}(M, \theta(t)) \leq c_2,$$

which completes the proof.  $\square$

The proof of the lemma above also implies the  $S_1^2$ -bound.

**Corollary 3.4.** *There exists a positive constant  $c_3$  depending only on  $n, M, f, u_0$  such that*

$$\|u(t)\|_{\mathbf{S}_1^2(M)} \leq c_3.$$

Next lemma shows that the normalized factor  $\lambda$  keeps bounded as long as the flow exists.

**Lemma 3.5.** *There exists a positive constant  $\Lambda_0$  depending only on  $n, f, M, u_0$  such that*

$$|\lambda(t)| \leq \Lambda_0.$$

**Proof.** On one hand, by (2.3) and integration by parts we can write

$$\begin{aligned} \int_M f R_{\theta(t)} \theta \wedge (d\theta)^n &= \int_M f u \left( - (2 + 2/n) \Delta_{\theta_0} u + R_{\theta_0} u \right) \theta_0 \wedge (d\theta_0)^n \\ &= \int_M (2 + 2/n) \langle \nabla_{\theta_0} (f u), \nabla_{\theta_0} u \rangle + f R_{\theta_0} u^2 \theta_0 \wedge (d\theta_0)^n \\ &= \int_M f \left[ (2 + 2/n) |\nabla_{\theta_0} u|^2 + R_{\theta_0} u^2 \right] \theta_0 \wedge (d\theta_0)^n \\ &\quad + \int_M (2 + 2/n) u \langle \nabla_{\theta_0} f, \nabla_{\theta_0} u \rangle \theta_0 \wedge (d\theta_0)^n. \end{aligned}$$

Hence, from Corollary 3.4 it follows that

$$\begin{aligned} \left| \int_M f R_{\theta(t)} \theta \wedge (d\theta)^n \right| &\leq \sup_M (|f|(2 + 2/n + R_{\theta_0})) \|u\|_{\mathbf{S}_1^2(M)}^2 + (2 + 2/n) \sup_M |\nabla_{\theta_0} f| \|u\|_{\mathbf{S}_1^2(M)}^2 \\ &\leq c_3 \left[ \sup_M (|f|(2 + 2/n + R_{\theta_0})) + (2 + 2/n) \sup_M |\nabla_{\theta_0} f| \right]. \end{aligned}$$

On the other hand, using Hölder's inequality, (2.8) and Lemma 3.3 we can obtain

$$1 = \left( \int_M f \theta \wedge (d\theta)^n \right)^2 \leq \text{vol}(M, \theta(t)) \int_M f^2 \theta \wedge (d\theta)^n \leq c_2 \int_M f^2 \theta \wedge (d\theta)^n,$$

that is

$$\int_M f^2 \theta \wedge (d\theta)^n \geq c_2^{-1}. \quad (3.5)$$

Substituting the estimates above into (2.7) we thus find a positive constant  $\Lambda_0$  depending only on  $n, f, M, u_0$  such that  $|\lambda(t)| \leq \Lambda_0$ .  $\square$

To obtain the global existence of the flow (2.2), we need to bound the derivative of  $\lambda(t)$ . To this end, we let

$$F_p(\theta(t)) = \int_M |R_{\theta(t)} - \lambda(t)|^p \theta \wedge (d\theta)^n, \quad p > 0$$



and try to show that  $F_2(\theta(t))$  is bounded. We calculate the derivative of  $F_p(\theta(t))$ . By Lemma 3.2, we have

$$\begin{aligned}
 \frac{d}{dt}F_p(\theta(t)) &= \frac{d}{dt} \int_M |\lambda(t)f - R_{\theta(t)}|^p \theta \wedge (d\theta)^n \\
 &= p \int_M (\lambda(t)f - R_{\theta(t)})_t (\lambda(t)f - R_{\theta(t)}) |\lambda(t)f - R_{\theta(t)}|^{p-2} \theta \wedge (d\theta)^n \\
 &\quad + (n+1) \int_M |\lambda(t)f - R_{\theta(t)}|^p (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \\
 &= -p(n+1) \int_M \Delta_{\theta(t)} (\lambda(t)f - R_{\theta(t)}) (\lambda(t)f - R_{\theta(t)}) \\
 &\quad |\lambda(t) - R_{\theta(t)}|^{p-2} \theta \wedge (d\theta)^n + p \int_M R_{\theta(t)} |\lambda(t)f - R_{\theta(t)}|^p \theta \wedge (d\theta)^n \\
 &\quad + \lambda'(t)p \int_M f (\lambda(t)f - R_{\theta(t)}) |\lambda(t)f - R_{\theta(t)}|^{p-2} \theta \wedge (d\theta)^n \\
 &\quad + (n+1) \int_M |\lambda(t)f - R_{\theta(t)}|^p (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \\
 &= -\frac{4(p-1)(n+1)}{p} \int_M \left| \nabla_{\theta(t)} |\lambda(t)f - R_{\theta(t)}|^{p/2} \right|^2 \theta \wedge (d\theta)^n \\
 &\quad + p \int_M \lambda(t)f |\lambda(t)f - R_{\theta(t)}|^p \theta \wedge (d\theta)^n \\
 &\quad + \lambda'(t)p \int_M f (\lambda(t)f - R_{\theta(t)}) |\lambda(t)f - R_{\theta(t)}|^{p-2} \theta \wedge (d\theta)^n \\
 &\quad + (n+1-p) \int_M |\lambda(t)f - R_{\theta(t)}|^p (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n. \tag{3.6}
 \end{aligned}$$

**Lemma 3.6.** *There exists a positive constant  $c_4(n, f, M, u_0)$  such that  $F_2(\theta(t)) \leq c_4$ .*

**Proof.** Setting  $p = 2$  in (3.6) and using (2.6) and Lemma 3.5 we have

$$\begin{aligned}
 \frac{d}{dt}F_2(\theta(t)) &= -2(n+1) \int_M \left| \nabla_{\theta(t)} (\lambda(t)f - R_{\theta(t)}) \right|^2 \theta \wedge (d\theta)^n \\
 &\quad + 2 \int_M \lambda(t)f (\lambda(t)f - R_{\theta(t)})^2 \theta \wedge (d\theta)^n \\
 &\quad + 2\lambda'(t) \int_M f (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \\
 &\quad + (n-1) \int_M (\lambda(t)f - R_{\theta(t)})^3 \theta \wedge (d\theta)^n \\
 &= (n+1) \int_M \lambda(t)f (\lambda(t)f - R_{\theta(t)})^2 \theta \wedge (d\theta)^n \\
 &\quad - (2 + 2/n) \int_M \left| \nabla_{\theta(t)} (\lambda(t)f - R_{\theta(t)}) \right|^2 \theta \wedge (d\theta)^n \\
 &\quad - (n-1) \left[ (2 + 2/n) \int_M \left| \nabla_{\theta(t)} (\lambda(t)f - R_{\theta(t)}) \right|^2 \right. \\
 &\quad \left. + R_{\theta(t)} (\lambda(t)f - R_{\theta(t)})^2 \theta \wedge (d\theta)^n \right] \\
 &\leq (n+1)\Lambda_0 \left( \sup_M f \right) F_2(\theta(t)),
 \end{aligned}$$

where the term in the bracket in the second equality is non-negative due to the conformal invariant and the assumption of non-negative CR-Yamabe constant. Now, integrating both sides of the above differential inequality from 0 to  $t$  and using (3.1), we can obtain

$$\begin{aligned} F_2(\theta(t)) &\leq F_2(\theta^0) + (n+1)\Lambda_0 \left( \sup_M f \right) \int_0^t F_2(\theta(s)) \, ds \\ &\leq F_2(\theta^0) + (n+1)\Lambda_0 \left( \sup_M f \right) \mathcal{E}(u_0) =: c_4. \end{aligned}$$

It is clear that  $c_4$  depends only on  $n, f, M$ , and  $u_0$ .  $\square$

With the help of the boundedness of  $F_2(\theta(t))$ , we can control the derivative of  $\lambda(t)$ .

**Lemma 3.7.** *There exists a positive constant  $C$  depending only on  $n, f, M$ , and  $u_0$  such that  $|\lambda'(t)| \leq C\sqrt{F_2(\theta(t))}$ . In particular, there exists a positive constant  $\Lambda_1$  depending only on  $n, f, M$ , and  $u_0$  such that  $|\lambda'(t)| \leq \Lambda_1$ .*

**Proof.** It follows from (3.5) and Lemmas 3.2, 3.3 and 3.5 that

$$\begin{aligned} |\lambda'(t)| &\leq (n+1)c_2 \left| \int_M f \Delta_{\theta(t)}(\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \right| + nc_2 \left( \sup_M |f| \right) F_2(\theta(t)) \\ &\quad + c_2^{3/2} \Lambda_0 \left( \sup_M f^2 \right) \sqrt{F_2(\theta(t))}. \end{aligned} \quad (3.7)$$

It remains to estimate the first term on the right hand side of the inequality above. By integration by parts, we rewrite the integral as follows:

$$\begin{aligned} &\int_M f \Delta_{\theta(t)}(\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \\ &= \int_M (\lambda(t)f - R_{\theta(t)}) \Delta_{\theta(t)} f \theta \wedge (d\theta)^n \\ &= \int_M (u^2 \Delta_{\theta_0} f + 2u \langle \nabla_{\theta_0} f, \nabla_{\theta_0} u \rangle) (\lambda(t)f - R_{\theta(t)}) \theta_0 \wedge (d\theta_0)^n \\ &= \int_M u^2 \Delta_{\theta_0} f (\lambda(t)f - R_{\theta(t)}) \theta_0 \wedge (d\theta_0)^n \\ &\quad + \int_M 2u \langle \nabla_{\theta_0} f, \nabla_{\theta_0} u \rangle (\lambda(t)f - R_{\theta(t)}) \theta_0 \wedge (d\theta_0)^n \\ &=: \text{I} + \text{II}. \end{aligned} \quad (3.8)$$

**Estimate of I:** By Hölder's inequality and Lemma 3.3 we can estimate

$$\begin{aligned} \text{II} &\leq \sup_M |\Delta_{\theta_0} f| \int_M u^{1-\frac{1}{n}} |\lambda(t)f - R_{\theta(t)}| u^{1+\frac{1}{n}} \theta_0 \wedge (d\theta_0)^n \\ &\leq \sup_M |\Delta_{\theta_0} f| \left( \int_M u^{2-\frac{2}{n}} \theta_0 \wedge (d\theta_0)^n \right)^{\frac{1}{2}} \sqrt{F_2(\theta(t))} \\ &\leq \sup_M |\Delta_{\theta_0} f| \text{vol}(M, \theta(t))^{\frac{n-1}{2(n+1)}} \text{vol}(M, \theta_0)^{\frac{1}{n+1}} \sqrt{F_2(\theta(t))} \\ &\leq \sup_M |\Delta_{\theta_0} f| c_2^{\frac{n-1}{2(n+1)}} \text{vol}(M, \theta_0)^{\frac{1}{n+1}} \sqrt{F_2(\theta(t))}. \end{aligned} \quad (3.9)$$

**Estimate of II:** From Hölder's inequality we deduce that

$$\begin{aligned} \text{III} &\leq 2 \sup_M |\nabla_{\theta_0} f| \int_M u^{-1/n} |\nabla_{\theta_0} u| |\lambda(t)f - R_{\theta(t)}| u^{1+1/n} \theta_0 \wedge (d\theta_0)^n \\ &\leq 2 \sup_M |\nabla_{\theta_0} f| \left( \int_M u^{-2/n} |\nabla_{\theta_0} u|^2 \theta_0 \wedge (d\theta_0)^n \right)^{\frac{1}{2}} \sqrt{F_2(\theta(t))}. \end{aligned} \quad (3.10)$$

Now, we have to control the term  $\int_M u^{-2/n} |\nabla_{\theta_0} u|^2 \theta_0 \wedge (d\theta_0)^n$ . To see this, we split the argument into two cases.

**Case 1:**  $n \neq 2$ . In this case, by integrating by parts, (2.3) and the fact that  $R_{\theta_0} \geq 0$  we have

$$\begin{aligned} \int_M u^{-2/n} |\nabla_{\theta_0} u|^2 \theta_0 \wedge (d\theta_0)^n &= \frac{n}{n-2} \int_M \langle \nabla_{\theta_0} (u^{1-2/n}), \nabla_{\theta_0} u \rangle \theta_0 \wedge (d\theta_0)^n \\ &= \frac{n^2}{2(n+1)(n-2)} \int_M u^{1-2/n} \left[ -(2+2/n)\Delta_{\theta_0} u \right] \theta_0 \wedge (d\theta_0)^n \\ &= \frac{n^2}{2(n+1)(n-2)} \int_M R_{\theta(t)} u^2 - R_{\theta_0} u^{2-2/n} \theta_0 \wedge (d\theta_0)^n \\ &\leq \frac{n^2}{2(n+1)|n-2|} \int_M |R_{\theta(t)} - \lambda(t)f| u^2 + |\lambda(t)f| u^2 \\ &\quad + R_{\theta_0} u^{2-2/n} \theta_0 \wedge (d\theta_0)^n. \end{aligned} \quad (3.11)$$

In the same way as in the estimate of I, we obtain by Lemma 3.6 that

$$\int_M |R_{\theta(t)} - \lambda(t)f| u^2 \theta_0 \wedge (d\theta_0)^n \leq c_2^{\frac{n-1}{2(n+1)}} \text{vol}(M, \theta_0)^{\frac{1}{n+1}} \sqrt{c_4}.$$

and

$$\int_M R_{\theta_0} u^{2-2/n} \theta_0 \wedge (d\theta_0)^n \leq c_2^{\frac{n-1}{2(n+1)}} \text{vol}(M, \theta_0)^{\frac{2}{n+1}} \sup_M R_{\theta_0}.$$

Moreover, we use Corollary 3.4 to bound

$$\int_M |\lambda(t)f| u^2 \theta_0 \wedge (d\theta_0)^n \leq \Lambda_0 \left( \sup_M |f| \right) \|u\|_{S_1^2(M)}^2 \leq \Lambda_0 \left( \sup_M |f| \right) c_3^2.$$

Substituting these estimates into (3.11) gives

$$\int_M u^{-2/n} |\nabla_{\theta_0} u|^2 \theta_0 \wedge (d\theta_0)^n \leq C.$$

Hereafter, we use  $C$  to denote the constant depending only on  $n, f, M$  and  $u_0$  which may vary from line to line.

**Case 2:**  $n = 2$ . In this case, it follows from integration by parts and (2.3) that

$$\begin{aligned} \int_M u^{-1} |\nabla_{\theta_0} u|^2 \theta_0 \wedge (d\theta_0)^n &= \int_M \langle \nabla_{\theta_0} \log u, \nabla_{\theta_0} u \rangle \theta_0 \wedge (d\theta_0)^n \\ &= \frac{1}{3} \int_M \log u \left( -3\Delta_{\theta_0} u \right) \theta_0 \wedge (d\theta_0)^n \\ &= \frac{1}{3} \int_M \log u \left( R_{\theta(t)} u^2 - R_{\theta_0} u \right) \theta_0 \wedge (d\theta_0)^n \\ &\leq \frac{1}{3} \int_M |\log u| \left( |R_{\theta(t)} - \lambda(t)f| u^2 + |\lambda(t)f| u^2 + R_{\theta_0} u \right) \theta_0 \wedge (d\theta_0)^n. \end{aligned} \quad (3.12)$$

To estimate the terms on the right hand side of the inequality above, we consider the function

$$\sigma(s) = s^\alpha |\log s|^\beta, \quad s > 0, \alpha, \beta > 0.$$

It is easy to see that

$$0 \leq \sigma(s) \leq s^{\alpha+\beta}, \text{ if } s \geq 1 \quad \text{and} \quad 0 < \sigma(s) \leq ((\alpha e)^{-1} \beta)^\beta, \text{ if } s \in (0, 1). \quad (3.13)$$

Now, using Hölder's inequality, (3.13) with  $\alpha = 1, \beta = 2$ , (3.6) and Lemma 3.3, we can obtain

$$\begin{aligned} & \int_M |\log u| R_{\theta(t)} - \lambda(t) f |u|^2 \theta_0 \wedge (d\theta_0)^n \\ &= \int_M u^{\frac{1}{2}} |\log u| R_{\theta(t)} - \lambda(t) f |u|^{3/2} \theta_0 \wedge (d\theta_0)^n \\ &\leq \left( \int_M u |\log u|^2 \theta_0 \wedge (d\theta_0)^n \right)^{\frac{1}{2}} \sqrt{F_2(\theta(t))} \\ &\leq \left( \int_M u^3 + 4e^{-2} \theta_0 \wedge (d\theta_0)^n \right)^{\frac{1}{2}} \sqrt{c_4} \\ &= \left( \text{vol}(M, \theta(t)) + 4e^{-2} \text{vol}(M, \theta_0) \right)^{\frac{1}{2}} \sqrt{c_4} \\ &\leq \left( c_2 + 4e^{-2} \text{vol}(M, \theta_0) \right)^{\frac{1}{2}} \sqrt{c_4}. \end{aligned}$$

Applying (3.13) with  $\alpha = 2, \beta = 1$  and using Lemmas 3.5 and 3.3, we have

$$\begin{aligned} \int_M |\lambda(t) f |u|^2 | \log u| \theta_0 \wedge (d\theta_0)^n &\leq \Lambda_0 \sup_M |f| \int_M u^3 + (2e)^{-1} \theta_0 \wedge (d\theta_0)^n \\ &\leq \Lambda_0 \sup_M |f| \left( c_2 + (2e)^{-1} \text{vol}(M, \theta_0) \right). \end{aligned}$$

Finally, by (3.13) with  $\alpha = 1, \beta = 1$  and Corollary 3.4 we get

$$\begin{aligned} \int_M R_{\theta_0} u |\log u| \theta_0 \wedge (d\theta_0)^n &\leq \sup_M R_{\theta_0} \int_M u^2 + e^{-1} \theta_0 \wedge (d\theta_0)^n \\ &\leq \sup_M R_{\theta_0} \left( c_3^2 + e^{-1} \text{vol}(M, \theta_0) \right). \end{aligned}$$

Substituting the three estimates above into (3.12) yields

$$\int_M u^{-2/n} |\nabla_{\theta_0} u|^2 \theta_0 \wedge (d\theta_0)^n \leq C.$$

Plugging the estimates in Case 1 and Case 2 into (3.10), we get

$$|\text{II}| \leq C \sqrt{F_2(\theta(t))}.$$

This together with (3.7), (3.8) and (3.9) gives the assertion.  $\square$

Now, we are ready to show that the Webster scalar curvature is bounded from the below uniformly. Before doing so, we define

$$\gamma = \min \left\{ -\Lambda_0 \sup_M |f|, \inf_M (R_{\theta^0} - \lambda(0) f), -\sqrt{\frac{4}{3} (\Lambda_0^2 (\sup_M f)^2 + \Lambda_1 \sup_M |f|)} \right\} \quad (3.14)$$

**Lemma 3.8.** *The Webster scalar curvature  $R_{\theta(t)}$  of the contact form  $\theta(t)$  satisfies*

$$R_{\theta(t)} - \lambda(t) f \geq \gamma.$$

**Proof.** Define the differential operator

$$\mathcal{L} = \partial_t - (n+1)\Delta_{\theta(t)} + (\lambda(t)f - \gamma).$$

Then it follows from Lemmas 3.2, 3.5 and 3.7 and the choice of  $\gamma$  in (3.14) that

$$\begin{aligned} \mathcal{L}(\lambda(t)f - R_{\theta(t)} + \gamma) &= \partial_t(\lambda(t)f - R_{\theta(t)}) - (n+1)\Delta_{\theta(t)}(\lambda(t)f - R_{\theta(t)}) \\ &\quad + (\lambda(t)f - \gamma)(\lambda(t)f - R_{\theta(t)} + \gamma) \\ &= (\lambda(t)f)^2 - \gamma^2 + (\gamma - R_{\theta(t)})R_{\theta(t)} + \lambda'(t)f \\ &\leq \Lambda_0^2 \left( \sup_M f \right)^2 - \frac{3}{4}\gamma^2 + \Lambda_1 \sup_M |f| \leq 0. \end{aligned}$$

Since  $\lambda(t)f - \gamma \geq 0$  and  $\lambda(t)f - R_{\theta(t)} + \gamma|_{t=0} \leq 0$  by (3.14), we can apply maximum principle to the operator  $\mathcal{L}$  to get the result.  $\square$

With all the preparations above, we are in a position to prove the global existence of the flow (2.2). The key step is to bound the conformal factor  $u(\cdot, t)$ .

**Lemma 3.9.** *For any  $T > 0$ , there exists a positive constant  $C = C(T)$  such that*

$$C^{-1} \leq u(\cdot, t) \leq C, \quad \text{for all } t \in [0, T].$$

**Proof.** From (2.2) and Lemma 3.8, it follows that

$$\frac{\partial}{\partial t} \log u \leq -\frac{n}{2}\gamma.$$

So, for  $t \in [0, T]$ , we have

$$u(x, t) \leq \left( \sup_M u_0 \right) \exp\left(-\frac{n}{2}\gamma T\right).$$

Next, we get the lower bound. To do so, we let

$$P := R_{\theta_0} + \sup_{t \in [0, T]} \sup_M \left( -(\lambda(t)f + \gamma)u^{2/n} \right).$$

Then by (2.3) and Lemma 3.8 we can obtain

$$\begin{aligned} 0 &\leq (R_{\theta(t)} - \lambda(t)f - \gamma)u^{1+2/n} \\ &= R_{\theta(t)}u^{1+2/n} - (\lambda(t)f + \gamma)u^{1+2/n} \\ &= -(2 + 2/n)\Delta_{\theta_0}u + R_{\theta_0}u - (\lambda(t)f + \gamma)u^{1+2/n} \\ &\leq -(2 + 2/n)\Delta_{\theta_0}u + Pu. \end{aligned}$$

Thus, we can apply Proposition A.1 in [16] to conclude that

$$C \left( \inf_M u(t) \right) \left( \sup_M u(t) \right)^{1+2/n} \geq \int_M u(t)^{2+2/n} \theta_0 \wedge (d\theta_0)^n.$$

This together with Lemma 3.3 and the upper bound above implies that

$$\inf_M u(t) \geq C^{-1}.$$

This proves the assertion.  $\square$

**Lemma 3.10.** *For  $p > n + 1$ , there exists a positive constant  $C$  such that*

$$\frac{d}{dt} F_p(\theta(t)) \leq C F_p(\theta(t)) + C F_p(\theta(t))^{\frac{p-n}{p-n-1}}.$$

Here, if  $R_{\theta_0} > 0$ , then  $C$  is independent of time; while if  $R_{\theta_0} \equiv 0$ , then the differential inequality above holds on  $[0, T]$  for any given  $T > 0$  with  $C$  depending on  $T$ .

**Proof. Case 1.**  $R_{\theta_0} > 0$ . In this case, from (2.10), (3.6) and Lemmas 3.5 and 3.7, it follows that

$$\begin{aligned}
\frac{d}{dt}F_p(\theta(t)) &= -\frac{2n(p-1)}{p} \int_M (2+2/n) \left| \nabla_{\theta(t)} |\lambda(t)f - R_{\theta(t)}|^{p/2} \right|^2 \\
&\quad + R_{\theta(t)} |\lambda(t)f - R_{\theta(t)}|^p \theta \wedge (d\theta)^n \\
&\quad + \left( p + \frac{2n(p-1)}{p} \right) \int_M R_{\theta(t)} |\lambda(t)f - R_{\theta(t)}|^p \theta \wedge (d\theta)^n \\
&\quad + \lambda'(t)p \int_M f(\lambda(t)f - R_{\theta(t)}) |\lambda(t)f - R_{\theta(t)}|^{p-2} \theta \wedge (d\theta)^n \\
&\quad + (n+1) \int_M |\lambda(t)f - R_{\theta(t)}|^p (\lambda(t)f - R_{\theta(t)}) \theta \wedge (d\theta)^n \\
&\leq -\frac{2n(p-1)\mu(M)}{p} F_{\frac{p(n+1)}{n}}(\theta(t))^{\frac{n}{n+1}} + \left( p + \frac{2n(p-1)}{p} \right) \Lambda_0 \sup_M |f| F_p(\theta(t)) \\
&\quad + Cp \sqrt{F_2(\theta(t))} F_{p-1}(\theta(t)) + \left( n+1+p + \frac{2n(p-1)}{p} \right) F_{p+1}(\theta(t)).
\end{aligned} \tag{3.15}$$

By Hölder's inequality we can estimate

$$\sqrt{F_2(\theta(t))} F_{p-1}(\theta(t)) \leq C F_p(\theta(t))^{\frac{1}{p}} F_p(\theta)^{\frac{p-1}{p}} = C F_p(\theta(t)).$$

Furthermore, using the Hölder's inequality and Young's inequality we obtain

$$\begin{aligned}
F_{p+1}(\theta(t)) &\leq F_{\frac{p(n+1)}{n}}(\theta(t))^{\frac{n}{p}} F_p(\theta(t))^{\frac{p-n}{p}} \\
&\leq \varepsilon F_{\frac{p(n+1)}{n}}(\theta(t))^{\frac{n}{n+1}} + C(\varepsilon) F_p(\theta(t))^{\frac{p-n}{p-n-1}}.
\end{aligned}$$

Substituting the two estimates above with a suitable  $\varepsilon$  into (3.15), we thus complete the proof of case 1.

**Case 2.**  $R_{\theta_0} \equiv 0$ . In this case, the method in case 1 does not work since the CR-Yamabe constant  $\mu(M) = 0$ . However, we may make use of the time-dependent bound for the conformal factor  $u$  and Folland-Stein embedding theorem. Precisely speaking, we use Lemma 3.9 and (2.11) to obtain

$$\begin{aligned}
\int_M \left| \nabla_{\theta(t)} |\lambda(t)f - R_{\theta(t)}|^{p/2} \right|^2 \theta \wedge (d\theta)^n &\geq C(T) \int_M \left| \nabla_{\theta_0} |\lambda(t)f - R_{\theta(t)}|^{p/2} \right|^2 \theta_0 \wedge (d\theta_0)^n \\
&= C(T) \left\| |\lambda(t)f - R_{\theta(t)}|^{p/2} \right\|_{\mathbb{S}^2_1(M)}^2 + C(T) F_p(\theta(t)) \\
&\geq C(T) F_{\frac{p(n+1)}{n}}(\theta(t))^{\frac{n+1}{n}} + C(T) F_p(\theta(t)).
\end{aligned}$$

With help of the inequality above, we then follow the proof of case 1 to obtain the assertion.  $\square$

With all these, we can follow the exactly same scheme in [17] to prove the global existence of the flow (2.2).

**Proposition 3.11.** *Given an initial smooth function  $u_0 \in X_f$ , there exists a smooth solution  $u(\cdot, t)$  of the flow (2.2) on  $[0, +\infty)$ .*

4.  $L^p$  CONVERGENCE

In this section, we devote ourselves to proving the  $L^p$  convergence of the flow (2.2), that is  $F_p(\theta(t)) \rightarrow 0$ , as  $t \rightarrow +\infty$ . To realize this, we first show the following differential inequality.

**Lemma 4.1.** *There exists a positive constant  $C$ , independent of time  $t$ , such that*

$$\frac{d}{dt}F_p(\theta(t)) \leq CF_p(\theta(t)), \quad 2 \leq p \leq n+1. \quad (4.1)$$

**Proof.** Recalling (3.6) and using Lemmas 3.3, 3.5, 3.7, 3.8 and Hölder's inequality, we obtain

$$\begin{aligned} \frac{d}{dt}F_p(\theta(t)) &\leq p\Lambda_0\left(\sup_M f\right)F_p(\theta(t)) + p\left(\sup_M f\right)C\sqrt{F_2(\theta(t))}F_{p-1}(\theta(t)) \\ &\quad - \gamma(n+1-p)F_p(\theta(t)) \\ &\leq CF_p(\theta(t)). \end{aligned}$$

Here, we have used the fact that  $\gamma < 0$  and  $p \leq n+1$ . In addition, it is clear that  $C$  is independent of time  $t$ .  $\square$

**Lemma 4.2.** *If either of two conditions below holds*

- (i)  $R_{\theta_0} > 0$  and  $0 < p < \infty$ ;
- (ii)  $R_{\theta_0} \equiv 0$  and  $0 < p \leq n+1$ ,

then  $F_p(\theta(t)) \rightarrow 0$ , as  $t \rightarrow +\infty$ .

**Proof. Case 1.**  $R_{\theta_0} > 0$ . In this case, one may follow the proof of [17, Lemma 3.2]. Hence, we omit the details.

**Case 2.**  $R_{\theta_0} \equiv 0$ . Note that the volume of  $(M, \theta(t))$  is bounded. Hence, by the Hölder's inequality, it suffices to show the conclusion for any  $2 \leq p \leq n+1$ . And in view of (4.1), we only need to show that  $F_p$  is integrable on  $[0, +\infty)$ . Indeed, if this is the case, then there would be a time sequence  $(t_j)_j \subset (0, +\infty)$  such that  $F_p(\theta(t_j)) \rightarrow 0$  as  $j \rightarrow +\infty$ . Integrating (4.1) from  $t_j$  to  $t$  with  $t > t_j$  yields

$$F_p(\theta(t)) \leq F_p(\theta(t_j)) + \int_{t_j}^t F_p(\theta(t)) dt.$$

Then sending  $j$  to the infinity gives that  $F_p(\theta(t)) \rightarrow 0$  as  $t \rightarrow +\infty$ . To see the integrability of  $F_p(\theta(t))$ , we will perform mathematical induction for  $2 \leq p \leq n+1$  with  $n \geq 2$  (it is trivial when  $n = 1$ ). When  $p = 2$ , the integrability on  $[0, +\infty)$  follows from (3.1). Now, we assume that  $F_k(\theta(t))$  is integrable for some  $2 < k < n+1$  and claim that  $F_{k+1}(\theta(t))$  is integrable. Indeed, setting  $p = k$  in (3.6), we get the estimate

$$\int_M |\lambda(t)f - R_{\theta(t)}|^k (R_{\theta(t)} - \lambda(t)f) \theta \wedge (d\theta)^n \leq CF_k(\theta(t)) - \frac{d}{dt}F_k(\theta(t)).$$

This together the induction assumption and (4.1) with  $p = k$  implies that

$$\int_0^\infty \int_M |\lambda(t)f - R_{\theta(t)}|^k (R_{\theta(t)} - \lambda(t)f) \theta \wedge (d\theta)^n dt \leq C.$$

By Lemma 3.8, we have

$$\begin{aligned}
& \int_M |\lambda(t)f - R_{\theta(t)}|^{k+1} \theta \wedge (d\theta)^n \\
&= \int_M |\lambda(t)f - R_{\theta(t)}|^k (R_{\theta(t)} - \lambda(t)f) \theta \wedge (d\theta)^n \\
&\quad + 2 \int_M |\lambda(t)f - R_{\theta(t)}|^k (R_{\theta(t)} - \lambda(t)f)^- \theta \wedge (d\theta)^n \\
&\leq \int_M |\lambda(t)f - R_{\theta(t)}|^k (R_{\theta(t)} - \lambda(t)f) \theta \wedge (d\theta)^n + 2(-\gamma)F_k(\theta(t)),
\end{aligned}$$

where  $h^- = \max\{-h, 0\}$  for a function  $h$ . From this estimate, we conclude that  $F_{k+1}(\theta(t))$  is integrable on  $[0, +\infty)$ .  $\square$

## 5. BLOW-UP ANALYSIS

In this section, we perform the so-called blow-up analysis. To do so, let us start with the CR version of concentration-compactness theorem.

**5.1. Concentration-Compactness.** Consider the semi-linear equation

$$-\Delta_{\theta}u + hu = fu^{1+2/n} \quad (5.1)$$

and the associated functional

$$I_f[u] = \frac{1}{2} \int_M (|\nabla u|_{\theta}^2 + hu^2) \theta \wedge (d\theta)^n - \frac{n}{2(n+1)} \int_M fu^{2+2/n} \theta \wedge (d\theta)^n.$$

It is known (c.f. [9]) that any nonnegative and nontrivial solution of the equation on Heisenberg group

$$-\Delta_{\theta_{\mathbb{H}^n}}u = u^{1+2/n} \quad (5.2)$$

must be of the form

$$\mathcal{W}_{\zeta}^{\mu} = \mu^n \left| (s - s_0) + i(|z - z_0|^2 + (\mu/n)^2) \right|^{-n}, \quad (5.3)$$

where  $\zeta = (z_0, s_0)$  with  $z_0 \in \mathbb{C}^n$ ,  $s_0 \in \mathbb{R}$  and  $\mu > 0$ . Furthermore, the functions of the form (5.3) are, up to left translations and dilations  $\mathcal{T}_{\delta} : (z, s) \mapsto (\delta z, \delta^2 s)$ , the only extremals of the the sharp Sobolev inequality on Heisenberg group

$$\left( \int_{\mathbb{H}^n} |u|^{2+2/n} \theta \wedge (d\theta)^n \right)^{\frac{n}{n+1}} \leq K_n \int_{\mathbb{H}^n} |\nabla u|_{\theta}^2 \theta \wedge (d\theta)^n, \quad (5.4)$$

where  $K_n$  is the same constant as in (2.10). The energy on Heisenberg group is denoted by

$$E(u) = \frac{1}{2} \int_{\mathbb{H}^n} |\nabla u|_{\theta_{\mathbb{H}^n}}^2 \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n - \frac{n}{2(n+1)} \int_{\mathbb{H}^n} |u|^{2+2/n} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n.$$

If we choose  $\mu > 0$  such that

$$\int_{\mathbb{H}^n} |\mathcal{W}_{\zeta}^{\mu}|^{2+2/n} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n = K_n^{-(n+1)}, \quad (5.5)$$

then by (5.2) and (5.4) we get

$$E(\mathcal{W}_{\zeta}^{\mu}) = \frac{K_n^{-(n+1)}}{2(n+1)}.$$



**Theorem 5.1.** *Let  $(u_k)_{k \in \mathbb{N}}$  be an  $\mathbf{S}_1^2(M)$ -bounded Palais-Smale sequence of non-negative functions for  $I_f$ , where  $h, f \in C^\infty(M)$  with  $\sup_M f > 0$ . Then, there exist  $m \in \mathbb{N}$  and a nonnegative weak solution  $u_\infty \in \mathbf{S}_1^2(M)$  of (5.1), such that, up to a subsequence, the following conclusions hold for  $1 \leq j \leq m$ :*

- (i) *there exist  $x_k^j \rightarrow x^j \in M$  such that  $f(x^j) > 0$  as  $k \rightarrow +\infty$ ;*
- (ii) *there exist  $\varepsilon_k^j > 0$  such that  $\varepsilon_k^j \rightarrow 0$  and*

$$\frac{\varepsilon_k^i}{\varepsilon_k^j} + \frac{\varepsilon_k^j}{\varepsilon_k^i} + \frac{d(x_k^i, x_k^j)^2}{\varepsilon_k^i \varepsilon_k^j} \rightarrow +\infty, \text{ for all } i \neq j, \text{ as } k \rightarrow +\infty, \quad (5.6)$$

where  $d(\cdot, \cdot)$  is the Carnot-Carathéodory distance on  $M$  with respect to the contact form  $\theta_0$ ;

- (iii) *with*

$$\mathcal{U}_k^j(x) = (\varepsilon_k^j)^{-n} f(x^j)^{-n/2} \mathcal{U}^j \left( \mathcal{T}_{(\varepsilon_k^j)^{-1}} \exp_{x_k^j}^{-1}(x) \right)$$

for  $x \in \exp_{x_k^j}(B_{2r_0}^+(0))$ , there holds

$$\left\| u_k - u_\infty - \sum_{j=1}^m \eta_k^j \mathcal{U}_k^j \right\|_{\mathbf{S}_1^2(M)} \rightarrow 0, \text{ as } k \rightarrow +\infty, \quad (5.7)$$

where  $\mathcal{U}^j = \mathcal{U}_{\zeta_j}^{\mu_j}$  satisfies (5.5),  $r_0 > 0$  is small,  $\exp_{x_k^j} : B_{2r_0}(0) \subset \mathbb{H}^n \mapsto M$  are the local charts centered at  $x_k^j \in M$ ,  $\mathcal{T}$  is the dilation as before, and  $\eta_k^j$  are smooth cut-off functions such that  $\eta_k^j \equiv 1$  in  $\exp_{x_k^j}(B_{r_0}(0))$  and  $\eta_k^j \equiv 0$  in  $M \setminus \exp_{x_k^j}(B_{2r_0})$ . Moreover,

$$I_f[u_k] = I_f[u_\infty] + \frac{K_n^{-(n+1)}}{2(n+1)} \sum_{j=1}^m f(x^j)^{-n} + o(1), \quad (5.8)$$

where  $o(1) \rightarrow 0$  as  $k \rightarrow +\infty$ .

**Proof.** To prove Theorem 5.1, one can just follow the proof of Proposition 5.1 in [21] which is contained in Appendix B of [21]; the only difference between Theorem 5.1 here and Proposition 5.1 in [21] is that we are considering (5.1) in which  $f$  is a function, while in Proposition 5.1 in [21] they are considering (5.1) in which  $f$  is a positive constant. Note that the assertion  $f(x_j) > 0$  in Theorem 5.1 (i) follows from the fact that if

$$-\Delta_{\theta_{\mathbb{H}^n}} u = cu^{1+2/n}$$

has a positive solution for some constant  $c$ , then  $c$  must be positive.  $\square$

**5.2. Weak solutions.** A simple characterization of the asymptotic behavior of energy  $\mathcal{E}(u)$  is given as below

**Lemma 5.2.** *Let  $u(\cdot, t)$  be a smooth solution of the flow (2.2). Then there hold*

- (i)  $\mathcal{E}(u(t)) = \lambda(t) + o(1)$ , where  $o(1) \rightarrow 0$  as  $t \rightarrow \infty$ ;
- (ii) *there exists a constant  $\lambda_\infty$  such that  $\lambda(t) \rightarrow \lambda_\infty$  as  $t \rightarrow \infty$ .*

**Proof.** (i) From (2.3), (2.8) and Lemma 4.2 it follows that

$$\mathcal{E}(u(t)) = \int_M R_{\theta(t)} \theta \wedge (d\theta)^n = \int_M (R_{\theta(t)} - \lambda(t)f) \theta \wedge (d\theta)^n + \lambda(t) = \lambda(t) + o(1),$$

where  $o(1) \rightarrow 0$ , as  $t \rightarrow \infty$ .

(ii) By Lemma 3.1, we know that  $\mathcal{E}(u(t))$  is monotone non-increasing. In addition, (2.10) implies that  $\mathcal{E}(u(t))$  has a lower bound. Hence  $\mathcal{E}(u(t))$  converges as  $t \rightarrow \infty$ . Then the assertion in (i) concludes that  $\lambda(t) \rightarrow \lambda_\infty$  for some constant  $\lambda_\infty$  as  $t \rightarrow \infty$ .  $\square$

For a nonnegative function  $w \in \mathbf{S}_1^2(M)$ , we define the energy functional

$$J_f[w] = \frac{1}{2} \int_M \left( (2 + 2/n) |\nabla w|_{\theta_0}^2 + R_{\theta_0} w^2 - \frac{n}{n+1} \lambda_\infty f w^{2+2/n} \right) \theta_0 \wedge (d\theta_0)^n.$$

Then, it is easy to compute

$$dJ_f[w](\phi) = \int_M (2 + 2/n) \langle \nabla w, \nabla \phi \rangle_{\theta_0} + R_{\theta_0} w \phi - \lambda_\infty f w^{1+2/n} \phi \theta_0 \wedge (d\theta_0)^n. \quad (5.9)$$

Moreover, we have

**Lemma 5.3.** *For any smooth solution  $u(\cdot, t)$  of the flow (2.2), there hold*

$$J_f[u(t)] = \frac{1}{2} \mathcal{E}(u(t)) - \frac{n}{2(n+1)} \lambda_\infty \quad \text{and} \quad J_f[u(t)] = \frac{1}{2(n+1)} \lambda_\infty + o(1),$$

where  $o(1) \rightarrow 0$  as  $t \rightarrow +\infty$ .

**Proof.** From (2.8) and definition of  $J_f$ , it follows that

$$\begin{aligned} J_f[u(t)] &= \int_M (1 + 1/n) |\nabla u|_{\theta_0}^2 + \frac{1}{2} R_{\theta_0} u^2 - \frac{n}{2(n+1)} \lambda_\infty f u^{2+2/n} \theta_0 \wedge (d\theta_0)^n \\ &= \frac{1}{2} \mathcal{E}(u(t)) - \frac{n}{2(n+1)} \lambda_\infty. \end{aligned}$$

Rewriting the above equality as

$$J_f[u(t)] = \frac{1}{2} (\mathcal{E}(u(t)) - \lambda_\infty) + \frac{1}{2(n+1)} \lambda_\infty.$$

Thanks to Lemma 5.2(i) that  $\mathcal{E}(u(t)) \rightarrow \lambda_\infty$  as  $t \rightarrow +\infty$ , we thus finished the proof.  $\square$

Let  $\{t_k : k \in \mathbb{N}\}$  be a time sequence such that  $t_k \rightarrow +\infty$  as  $k \rightarrow +\infty$ . Also, we set  $u_k := u(t_k)$ ,  $\theta_k := \theta(t_k)$ ,  $\lambda_k := \lambda(t_k)$  and  $R_k := R_{\theta_k}$ . Then we will show the following:

**Lemma 5.4.**  *$(u_k)_k$  is a Palais-Smale sequence for the energy functional  $J_f$ .*

**Proof.** We need to show that

- The sequence  $(J_f[u_k])_k$  is bounded;
- $dJ_f[u_k] \rightarrow 0$  strongly in  $\mathbf{S}_1^2(M)^{-1}$  as  $k \rightarrow +\infty$ .

The boundedness of  $(J_f[u_k])_k$  is straightforward by Lemma 5.3 and (3.1). Hence, we are left to check the strong convergence  $dJ_f[u_k] \rightarrow 0$  in  $\mathbf{S}_1^2(M)^{-1}$  as  $k \rightarrow +\infty$ . To see this, we fix any  $\phi \in \mathbf{S}_1^2(M)$ . Then (2.3), (2.11) and Hölder's inequality yield

$$\begin{aligned} dJ_f[u_k](\phi) &= \int_M \left[ -(2 + 2/n) \Delta_{\theta_0} u_k + R_{\theta_0} u_k - \lambda_\infty f u_k^{1+2/n} \right] \phi \theta_0 \wedge (d\theta_0)^n \\ &= \int_M (R_k - \lambda_\infty f) u_k^{1+2/n} \phi \theta_0 \wedge (d\theta_0)^n \\ &\leq \|R_k - \lambda_k f + (\lambda_k - \lambda_\infty) f\|_{L^{\frac{2(n+1)}{n+2}}(M, \theta_k)} \|\phi\|_{L^{\frac{2(n+1)}{n}}(M, \theta_0)} \\ &\leq C \left( F_{\frac{2(n+1)}{n+2}}(\theta_k)^{\frac{n+2}{2(n+1)}} + |\lambda_k - \lambda_\infty| \sup_M f c_2^{\frac{n+2}{2(n+1)}} \right) \|\phi\|_{\mathbf{S}_1^2(M)}. \end{aligned}$$

Notice that  $2(n+1)/(n+2) < 2$ , it follows from Lemma 4.2 that  $F_{\frac{2n}{n+1}}(g_k) = o(1)$  as  $k \rightarrow +\infty$ . Moreover, by Lemma 5.2, we have  $\lambda_k - \lambda_\infty = o(1)$ , as  $k \rightarrow +\infty$ . Hence, the assertion follows.  $\square$

**Lemma 5.5.** *There exists  $u_\infty \in \mathbf{S}_1^2(M)$  such that  $u_k$  converges to  $u_\infty$  weakly in  $\mathbf{S}_1^2(M)$  as  $k \rightarrow +\infty$ . Moreover, if  $u_\infty \not\equiv 0$ , then (1.2) possess a positive solution.*

**Proof.** Since  $(u_k)_k$  is bounded in  $\mathbf{S}_1^2(M)$  by Corollary 3.4, there exists  $u_\infty \in \mathbf{S}_1^2(M)$  such that, up to a subsequence,  $u_k \rightharpoonup u_\infty$  weakly in  $\mathbf{S}_1^2(M)$  as  $k \rightarrow +\infty$ . By Lemma 5.4, we have  $dJ_f[u_k] = o(1)$  in  $\mathbf{S}_1^2(M)^{-1}$ . Then, it follows from (5.9) that

$$\int_M (2+2/n)\langle \nabla u_k, \nabla \phi \rangle_{\theta_0} + R_{\theta_0} u_k \phi \theta_0 \wedge (d\theta_0)^n = \int_M \lambda_\infty f u_k^{1+2/n} \phi \theta_0 \wedge (d\theta_0)^n + o(1).$$

Letting  $k \rightarrow +\infty$ , we obtain that  $u_\infty$  solves the Eq.(1.2) in the weak sense

$$\int_M (2+2/n)\langle \nabla u_\infty, \nabla \phi \rangle_{\theta_0} + R_{\theta_0} u_\infty \phi \theta_0 \wedge (d\theta_0)^n = \int_M \lambda_\infty f u_\infty^{1+2/n} \phi \theta_0 \wedge (d\theta_0)^n.$$

Moreover, standard regularity theory (c.f. Theorem 5.15 in [10]) implies that  $u_\infty$  is smooth if  $f$  is smooth. Hence,  $u_\infty$  satisfies

$$-(2+2/n)\Delta_{\theta_0} u_\infty + R_{\theta_0} u_\infty = \lambda_\infty f u_\infty^{1+2/n}. \quad (5.10)$$

Now, if  $u_\infty \not\equiv 0$ , we can apply Proposition A.1 in [16] to conclude that  $u_\infty > 0$ .

Next, we claim that  $\lambda_\infty > 0$ . If  $R_{\theta_0} > 0$ , it follows from (2.8), (2.10) and Lemma 3.3 that

$$\mathcal{E}(u(t)) \geq \mu(M) \text{vol}(M, \theta(t))^{\frac{n}{n+1}} \geq \mu(M) \left( \sup_M f \right)^{-\frac{n}{n+1}}.$$

Then by Lemma 5.2 we get that

$$\lambda_\infty = \lim_{t \rightarrow \infty} \mathcal{E}(u(t)) \geq \mu(M) \left( \sup_M f \right)^{-\frac{n}{n+1}} > 0.$$

If  $R_{\theta_0} \equiv 0$ , we first note that there exists a subsequence, still denoted by  $\{t_k\}$ , such that  $u_k \rightharpoonup u_\infty$  weakly in  $\mathbf{S}_1^2(M)$  and strongly in  $L^2(M)$ . Now, our argument is split into two cases.

**Case 1.**  $u_\infty \equiv 0$ . In this case, using (2.8), (2.11), Lemma 3.3 and the fact that  $\|u_k\|_{L^2(M)} \rightarrow 0$  as  $k \rightarrow \infty$ , we obtain

$$\begin{aligned} \int_M |\nabla_{\theta_0} u_k|^2 \theta_0 \wedge (d\theta_0)^n &\geq \mathcal{C}^{-1} \text{vol}(M, \theta_k)^{\frac{n}{n+1}} - \int_M u_k^2 \theta_0 \wedge (d\theta_0)^n \\ &\geq \mathcal{C}^{-1} \left( \sup_M f \right)^{-\frac{n}{n+1}} + o(1). \end{aligned}$$

This implies that

$$\begin{aligned} \lambda_\infty &= \lim_{v \rightarrow \infty} \mathcal{E}(u_k) = \lim_{k \rightarrow \infty} \int_M (2+2/n) |\nabla_{\theta_0} u_k|^2 \theta_0 \wedge (d\theta_0)^n \\ &\geq (2+2/n) \mathcal{C}^{-1} \left( \sup_M f \right)^{-\frac{n}{n+1}} > 0. \end{aligned}$$

**Case 2.**  $u_\infty \not\equiv 0$ . In this case, it follows from (5.10) that  $u_\infty > 0$  solves

$$-(2+2/n)\Delta_{\theta_0} u_\infty = \lambda_\infty f u_\infty^{1+2/n}.$$

To prove that  $\lambda_\infty > 0$ , we suppose on contrary  $\lambda_\infty = 0$ , which gives  $\Delta_{\theta_0} u_\infty = 0$ . Hence,  $u_\infty$  must be a positive constant, say  $c > 0$ . Since  $\lambda_\infty = 0$ , it follows from Lemma 5.2(i) that

$$\int_M |\nabla_{\theta_0}(u_k - u_\infty)|^2 \theta_0 \wedge (d\theta_0)^n = \int_M |\nabla_{\theta_0} u_k|^2 \theta_0 \wedge (d\theta_0)^n = \frac{n \mathcal{E}(u_k)}{2n+2} \rightarrow 0$$

as  $k \rightarrow \infty$ . This together with (2.11) implies that

$$\begin{aligned} & \left( \int_M |u_k - u_\infty|^{2+2/n} \theta_0 \wedge (d\theta_0)^n \right)^{\frac{n}{n+1}} \\ & \leq \mathcal{C}^{-1} \left( \int_M |\nabla_{\theta_0}(u_k - u_\infty)|^2 \theta_0 \wedge (d\theta_0)^n + \int_M |u_k - u_\infty|^2 \theta_0 \wedge (d\theta_0)^n \right) \rightarrow 0 \end{aligned}$$

as  $k \rightarrow \infty$ . This together with (2.8) implies that  $\int_M f u_\infty^{2+2/n} \theta_0 \wedge (d\theta_0)^n = 1$ . Since  $u_\infty \equiv c > 0$ , we have  $\int_M f \theta_0 \wedge (d\theta_0)^n = c^{-2-2/n} > 0$ , contradicting the assumption  $\int_M f \theta_0 \wedge (d\theta_0)^n < 0$ . This proves the claim, i.e.  $\lambda_\infty > 0$ .

Therefore, if  $u_\infty \neq 0$ , then  $\lambda_\infty^{n/2} u_\infty$  produces a positive solution to (1.2).  $\square$

Lemma 5.5 shows that Theorems 1.1 and 1.2 will immediately follow provided that we can show that there exists a time sequence  $(t_k)_k$  such that  $u_k \rightarrow u_\infty \neq 0$  weakly in  $\mathbf{S}_1^2(M)$ . However, this is generally hard to be proved via direct estimates. Hence, we shall adopt a contradiction argument. To do so, we assume, from now on, that  $u(t) \rightarrow 0$  weakly in  $\mathbf{S}_1^2(M)$  as  $t \rightarrow +\infty$ . Under this assumption, it is easy to see from (2.11) and Lemma 3.3 that  $u_k$  cannot converge to 0 strongly in  $\mathbf{S}_1^2(M)$ , and the concentration phenomenon will occur.

**5.3. Single concentration point.** Set

$$\lambda_* = \frac{2(n+1)}{n} \left( \sup_M f \right)^{-\frac{n}{n+1}} K_n^{-1},$$

where  $K_n = 1/(2\pi n^2)$  is the optimal constant appeared in (2.10). Then we have the following:

**Lemma 5.6.**  $\lambda_\infty \geq \lambda_*$ .

**Proof.** Since  $u \rightarrow 0$  weakly in  $\mathbf{S}_1^2(M)$ ,  $u \rightarrow 0$  strongly in  $L^2(M)$  as  $t \rightarrow 0$ . This fact together with Lemma 5.2 implies that

$$\begin{aligned} \lambda_\infty &= \lim_{t \rightarrow +\infty} \mathcal{E}(u(t)) = \lim_{t \rightarrow +\infty} \int_M (2 + 2/n) |\nabla u|_{\theta_0}^2 + R_{\theta_0} u^2 \theta_0 \wedge (d\theta_0)^n \\ &= \lim_{t \rightarrow +\infty} \int_M (2 + 2/n) |\nabla u|_{\theta_0}^2 \theta_0 \wedge (d\theta_0)^n \end{aligned} \quad (5.11)$$

On the other hand, given  $\varepsilon > 0$ , by (2.8), (2.10), Lemma 3.3 and the fact that  $\|u\|_{L^2} = o(1)$  we get

$$\begin{aligned} \int_M (2 + 2/n) |\nabla u|_{\theta_0}^2 \theta_0 \wedge (d\theta_0)^n &\geq \frac{2(n+1)}{n(K_n + \varepsilon)} \left( \text{vol}(M, \theta(t)) \right)^{\frac{n}{n+1}} + o(1) \\ &\geq \frac{2(n+1)}{n(K_n + \varepsilon)} \left( \sup_M f \right)^{-\frac{n}{n+1}} + o(1) \end{aligned} \quad (5.12)$$

Combining (5.11) and (5.12) yields

$$\lambda_\infty \geq \frac{2(n+1)}{n(K_n + \varepsilon)} \left( \sup_M f \right)^{-\frac{n}{n+1}}.$$

Letting  $\varepsilon$  to zero, we thus complete the proof.  $\square$

With a suitable choice of initial data, we can guarantee the single concentration point of the flow.

**Lemma 5.7.** *If the initial data  $u_0$  satisfies  $\mathcal{E}(u_0) < 2^{\frac{1}{n+1}} \lambda_*$ , then  $m = 1$ , where  $m$  is the positive integer in Theorem 5.1.*

**Proof.** First, note that  $J_f[w] = (2 + 2/n)I_{\frac{n}{2(n+1)}\lambda_\infty f}[w]$ . Thus, we can apply Theorem 5.1 to the functional  $J_f[u(t)]$  to obtain

$$J_f[u(t)] = \frac{K_n^{-(n+1)}}{2(n+1)} \sum_{j=1}^m \left( \frac{2(n+1)}{n} \right)^{n+1} (\lambda_\infty f(x_j))^{-n} + o(1).$$

This together with Lemma 5.3 implies that

$$\begin{aligned} \frac{\lambda_\infty}{2(n+1)} &= \lim_{t \rightarrow +\infty} J_f[u(t)] = \frac{K_n^{-(n+1)}}{2(n+1)} \sum_{j=1}^m \left( \frac{2(n+1)}{n} \right)^{n+1} (\lambda_\infty f(x_j))^{-n} \quad (5.13) \\ &\geq \frac{mK_n^{-(n+1)}}{2(n+1)} \left( \frac{2(n+1)}{n} \right)^{n+1} \lambda_\infty^{-n} (\sup_M f)^{-n}. \end{aligned}$$

Hence

$$\lambda_\infty^{\frac{1}{n+1}} \geq m^{\frac{1}{n+1}} \frac{2(n+1)}{n} (\sup_M f)^{-\frac{n}{n+1}} K_n^{-1} = m^{\frac{1}{n+1}} \lambda_*.$$

On the other hand, it follows from Lemma 5.2 and (3.1) that

$$\lambda_\infty = \lim_{t \rightarrow +\infty} \mathcal{E}(u(t)) \leq \mathcal{E}(u_0) < 2^{\frac{1}{n+1}} \lambda_*.$$

Combining the estimates above yields  $m < 2$ , i.e.  $m = 1$ .  $\square$

We have the following convention: For the rest of the paper, our initial data is chosen to satisfy

$$\int_M f u_0^{1+2/n} \theta_0 \wedge (d\theta_0)^n = 1 \text{ and } \mathcal{E}(u_0) < 2^{\frac{1}{n+1}} \lambda_*. \quad (5.14)$$

With such a choice of initial data, we know by Lemma 5.7 that there would be only one concentration point denoted by  $a_\infty$ . Then (5.13) can be reduced as

$$\frac{\lambda_\infty}{2(n+1)} = \frac{K_n^{-(n+1)}}{2(n+1)} \left( \frac{2(n+1)}{n} \right)^{n+1} (\lambda_\infty f(a_\infty))^{-n}.$$

Solving this gives

$$\lambda_\infty = \frac{2(n+1)}{n} K_n^{-1} f(a_\infty)^{-\frac{n}{n+1}}. \quad (5.15)$$

Moreover, we may choose suitable cut-off function  $\eta(t)$ ,  $\varepsilon^*(t) > 0$  with  $\varepsilon^*(t) \rightarrow 0$  and  $a^*(t) \in M$  with  $a^*(t) \rightarrow a_\infty$  such that as  $t \rightarrow \infty$

$$\|u(t) - \varphi^*(t)\|_{S^2_1(M)} \rightarrow 0, \quad (5.16)$$

where

$$\varphi^*(t, x) = \left( \frac{n\lambda_\infty f(a_\infty)}{2(n+1)} \right)^{-n/2} \eta(t) (\varepsilon^*(t))^{-n} \mathcal{U}(\mathcal{T}_{\varepsilon^*(t)^{-1}} \exp_{a^*(t)}^{-1}(x))$$

and

$$\mathcal{U} = |s + i(|z|^2 + (1/n)^2)|^{-n}.$$

#### 5.4. Bubbles.

**Case 1**  $\mu(M) > 0$ . Given any  $a \in M$ , we can find a contact form  $\theta_a = \omega_a^{*2/n} \theta_0$  conformal to  $\theta_0$  such that it induces CR normal coordinates  $(z, s)$  around  $a \in M$ . Let  $G_a^*$  be the Green's function with pole at  $a$ . Then we have

$$-(2 + 2/n)\Delta_{\theta_a} G_a^*(x) + R_{\theta_a}^* G_a^*(x) = \delta_a. \quad (5.17)$$

where  $R_{\theta_a}^*$  is the Webster scalar curvature of  $\theta_a$ . Moreover, the Green's function satisfies the estimates (see Appendix A in [21])

$$|G_a^*(x) - \rho_a(x)^{-2n} - A_a^*| \leq C\rho_a(x) \quad (5.18)$$

and

$$|\nabla(G_a^*(x) - \rho_a(x)^{-2n})|_{\theta_a} \leq C, \quad (5.19)$$

where  $A_a^*$  is the CR mass satisfying  $\inf_{a \in M} A_a^* > 0$  under the assumptions in Theorem 1.1, and  $\rho_a(x)$  the distance in the CR normal coordinates at  $a$  defined as

$$\rho_a(x) = (s^2 + |z|^4)^{1/4}.$$

We define  $\chi : \mathbb{R} \mapsto [0, 1]$  to be a smooth cut-off function satisfying  $\chi(\tau) = 1$  for  $\tau \leq 1$  and  $\chi(\tau) = 0$  for  $\tau \geq 2$ . For  $\delta > 0$ , we then define  $\chi_\delta(\tau) = \chi(\tau/\delta)$ . And, for  $\varepsilon \ll \delta$  we let

$$\varphi_{a,\varepsilon} = \omega_a^*(x) \psi_{a,\varepsilon}, \quad (5.20)$$

where

$$\psi_{a,\varepsilon} = \varepsilon^n \left[ \frac{\chi_\delta(\rho_a)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2})} + (1 - \chi_\delta(\rho_a))G_a^* \right]. \quad (5.21)$$

**Case 2**  $\mu(M) = 0$ . Given any  $a \in M$ , we consider the local charts  $(z, s)$  around  $a$ . Motivated by [7], we let

$$\varphi_{a,\varepsilon} = \varepsilon^n \left[ \frac{\chi_\delta(\rho_a)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2})} + (1 - \chi_\delta(\rho_a))(\rho_a^{-2n} + \Lambda) \right],$$

where  $\Lambda$  is a positive constant to be determined,  $\chi_\delta$  and  $\rho_a$  are the same notations as in the Case 1.

To provide a unified proof for both cases, we hereafter let

$$\omega_a = \begin{cases} \omega_a^*, & \text{if } \mu(M) > 0, \\ 1, & \text{if } \mu(M) = 0, \end{cases} \quad R_{\theta_a} = \begin{cases} R_{\theta_a}^*, & \text{if } \mu(M) > 0, \\ 0, & \text{if } \mu(M) = 0, \end{cases}$$

and

$$G_a = \begin{cases} G_a^*, & \text{if } \mu(M) > 0, \\ \rho_a^{-2n} + \Lambda, & \text{if } \mu(M) = 0, \end{cases} \quad A_a = \begin{cases} A_a^*, & \text{if } \mu(M) > 0, \\ \Lambda, & \text{if } \mu(M) = 0. \end{cases}$$

We have the following:

**Definition 5.8.** For  $u \in \mathbf{S}_1^2(M)$  and sufficiently small  $\mu > 0$ , we define

$$D_u(\mu) = \left\{ (\alpha, \varepsilon, a) \in (\mathbb{R}_+, \mathbb{R}_+, M) : \varepsilon, \left| \frac{n\lambda\alpha^{2/n}f(a)}{2(n+1)} - 1 \right|, \|u - \alpha\varphi_{a,\varepsilon}\| < \mu \right\}$$

From (5.16), it follows that

$$D_{u(t)}(\mu) \neq \emptyset$$

for all sufficiently small  $\mu > 0$  and all sufficiently large  $t$ .

**Lemma 5.9.** *There holds*

$$\begin{aligned} \mathcal{L}_{\theta_0} \varphi_{a,\varepsilon} &= (2 + 2/n)\varphi_{a,\varepsilon}^{1+2/n} + (2 + 2/n)\varepsilon^n A_a \omega_a^{1+2/n} \Delta_{\theta_a} \chi_\delta(\rho_a) \\ &\quad + \omega_a^{1+2/n} \varepsilon^n (I_1 + I_2 + I_3 + I_4 + I_5 + I_6 + I_7 + I_8), \end{aligned}$$

where

$$\begin{aligned} I_1 &= -(2 + 2/n)\chi_\delta(\rho_a) \left[ \Delta_{\theta_a} \left( \frac{1}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2}} \right) + \frac{\varepsilon^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right], \\ I_2 &= \frac{\chi_\delta(\rho_a) R_{\theta_a}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2}}, \\ I_3 &= (2 + 2/n) \Delta_{\theta_a} \chi_\delta(\rho_a) (G_a - \rho_a^{-2n} - A_a), \\ I_4 &= -(2 + 2/n) \Delta_{\theta_a} \chi_\delta(\rho_a) \left( \frac{1}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2}} - \rho_a^{-2n} \right), \\ I_5 &= 2(2 + 2/n) \left\langle \nabla \chi_\delta(\rho_a), \nabla (G_a - \rho_a^{-2n}) \right\rangle_{\theta_a}, \\ I_6 &= -2(2 + 2/n) \left\langle \nabla \chi_\delta(\rho_a), \nabla \left( \frac{1}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2}} - \rho_a^{-2n} \right) \right\rangle_{\theta_a}, \\ I_7 &= -(2 + 2/n) \varepsilon^2 \left[ \left( \frac{\chi_\delta(\rho_a)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2}} + (1 - \chi_\delta(\rho_a)) G_a \right)^{1+2/n} \right. \\ &\quad \left. - \frac{\chi_\delta(\rho_a)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right], \\ I_8 &= (1 - \chi_\delta(\rho_a)) \left[ (2 + 2/n) \Delta_{\theta_a} G_a - R_{\theta_a} G_a \right]. \end{aligned}$$

Hereafter,  $\mathcal{L}_{\theta_0}$  is the conformal sub-Laplacian with respect to  $\theta_0$ .

**Proof.** See [21, Proposition A.1].  $\square$

Now, we will choose some suitable parameters  $\alpha$ ,  $\varepsilon$  and  $a$  to simplify the proceeding arguments.

**Proposition 5.10.** *For each  $t > 0$ , there exists  $\mu_0 = \mu_0(t)$  with  $\mu_0(t) \rightarrow 0$  as  $t \rightarrow +\infty$  such that*

$$\inf_{(\tilde{\alpha}, \tilde{\varepsilon}, \tilde{a}) \in D_{u(t)}(2\mu_0)} \int_M u(t)^{2/n} |u(t) - \tilde{\alpha} \varphi_{\tilde{a}, \tilde{\varepsilon}}|^2 \theta_0 \wedge (d\theta_0)^n$$

admits a unique minimizer  $(\alpha(t), \varepsilon(t), a(t)) \in D_{u(t)}(\mu_0)$ .

**Proof.** Since the proof is almost identical with the one in [25, Proposition 3.10], we omit the proof and refer the readers to [25].  $\square$

For simplicity, we set

$$\varphi = \varphi_{a,\varepsilon}, \quad v = u - \alpha\varphi.$$

**Definition 5.11.** For any  $\varepsilon > 0$  and  $a \in M$ , we define

- (i)  $(d_1, d_2, d_3) = (1, \varepsilon \partial_\varepsilon, \varepsilon \nabla_a)$ ;
- (ii)  $\phi_k = d_k \varphi$  for  $k = 1, 2, 3$ .

Here, we regard  $a$  as a point in  $\mathbb{H}^n$  by using CR normal coordinates  $(z, s)$ , and  $\nabla_a$  is the usual sub-gradient in  $\mathbb{H}^n$ .

The next lemma provides us estimates on  $\phi_k$ , which is vital for proving the main theorems.

**Lemma 5.12.** *One has*

- (i)  $|d_k \phi_l| \leq C\varphi$  for  $k, l = 1, 2, 3$ ;

- (ii)  $\int_M \varphi^{2/n} \phi_k^2 \theta_0 \wedge (d\theta_0)^n = c_k + O(\varepsilon^{2n+2} + \varepsilon^4)$  for  $k = 1, 2, 3$ ;
- (iii)  $\int_M \varphi^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n = O(\varepsilon^{2n+2})$  for  $k = 2, 3$ ;
- (iv)  $\int_M \varphi^{2/n} \phi_k \phi_l \theta_0 \wedge (d\theta_0)^n = O(\varepsilon^{2n+2} + \varepsilon^4)$  for all  $k \neq l$ .

Here  $c_k$  are some positive constants.

Due to the length of the proof, Lemma 5.12 will be proved in Appendix A.

**Lemma 5.13.** *Suppose that  $n = 1$ . For  $t$  large, one has*

- (i)  $\int_M \mathcal{L}_{\theta_0} \phi_k v \theta_0 \wedge (d\theta_0)^n = O(\delta^{6-2n} \varepsilon^{2n} + \varepsilon^6) + O(\|v\|^2)$ ,
- (ii)  $\int_M f u^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n = \int_M f(\alpha\varphi)^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n + O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \|v\|^2)$ .

**Proof.** (i) In view of Lemma 5.9, we have

$$\begin{aligned} \int_M \mathcal{L}_{\theta_0} \phi_k v \theta_0 \wedge (d\theta_0)^n &= \int_M (d_k \mathcal{L}_{\theta_0} \varphi) v \theta_0 \wedge (d\theta_0)^n \\ &= \int_M \left[ (2 + 2/n) d_k \varphi^{1+2/n} + (2 + 2/n) d_k (\varepsilon^n A_a \omega_a^{1+2/n} \Delta_{\theta_a} \chi_\delta(\rho_a)) \right. \\ &\quad \left. + \sum_{i=1}^8 d_k (\omega_a^{1+2/n} \varepsilon^n I_i) \right] v \theta_0 \wedge (d\theta_0)^n. \end{aligned}$$

**Case  $k = 1$ .** First, notice that  $R_{\theta_a} = 0$  and

$$\Delta_{\theta_a} G_a = \left[ \Delta_{\theta_a} + O(\rho_a^2) \right] \rho_a^{-2n} = O(1),$$

if  $\mu(M) = 0$  and  $n = 1$ . Thus,  $I_8 = O(1) 1_{\delta \leq \rho_a \leq 2\delta}$  if  $\mu(M) = 0$  and  $n = 1$ . On the other hand, by (5.17),  $I_8 = 0$  if  $\mu(M) > 0$ . Recalling that  $d_1 = 1$ , one may follow the proof in [21, Proposition A.1] to obtain

$$\begin{aligned} \int_M \mathcal{L}_{\theta_0} \phi_1 v \theta_0 \wedge (d\theta_0)^n &= \int_M (2 + 2/n) \varphi^{1+2/n} v \theta_0 \wedge (d\theta_0)^n \\ &\quad + \int_M (2 + 2/n) \varepsilon^n A_a \omega_a^{1+2/n} \Delta_{\theta_a} \chi_\delta(\rho_a) v \theta_0 \wedge (d\theta_0)^n \\ &\quad + O\left( \int_M \left[ \left( \frac{\varepsilon^2}{s^2 + ((\varepsilon/n)^2 + |z|^2)^2} \right)^{n/2} \rho_a^2 1_{\{\rho_a \leq 2\delta\}} + \delta^{-1} \varepsilon^n 1_{\{\delta \leq \rho_a \leq 2\delta\}} \right. \right. \\ &\quad \left. \left. + \left( \frac{\varepsilon^2}{s^2 + ((\varepsilon/n)^2 + |z|^2)^2} \right)^{1+n/2} 1_{\{\rho_a \geq \delta\}} \right] v \theta_a \wedge (d\theta_a)^n \right). \end{aligned} \quad (5.22)$$

Using Hölder's inequality, we can estimate

$$\begin{aligned} &\int_M \left( \frac{\varepsilon^2}{s^2 + ((\varepsilon/n)^2 + |z|^2)^2} \right)^{n/2} \rho_a^2 1_{\{\rho_a \leq 2\delta\}} v \theta_a \wedge (d\theta_a)^n \\ &= O\left( \varepsilon^{2n} \int_{\rho_a \leq 2\delta} \frac{\rho_a^4}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^n} \theta_a \wedge (d\theta_a)^n + \|v\|^2 \right) \\ &= O(\delta^{6-2n} \varepsilon^{2n} + \|v\|^2), \end{aligned} \quad (5.23)$$



whenever  $n \leq 3$ . Similarly, we can obtain

$$\begin{aligned}
 & \int_M \delta^{-1} \varepsilon^n 1_{\{\delta \leq \rho_a \leq 2\delta\}} v \theta_a \wedge (d\theta_a)^n \\
 &= O\left(\delta^{-2} \varepsilon^{2n} \int_{\delta \leq \rho_a \leq 2\delta} \theta_a \wedge (d\theta_a)^n + \|v\|^2\right) \\
 &= O\left(\delta^{2n} \varepsilon^{2n} + \|v\|^2\right)
 \end{aligned} \tag{5.24}$$

and

$$\begin{aligned}
 & \int_M \left( \frac{\varepsilon^2}{s^2 + ((\varepsilon/n)^2 + |z|^2)^2} \right)^{1+n/2} 1_{\{\rho_a \geq \delta\}} v \theta_a \wedge (d\theta_a)^n \\
 &= O\left(\varepsilon^{2n+4} \int_{\rho_a \geq \delta} \frac{1}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_a \wedge (d\theta_a)^n + \|v\|^2\right) \\
 &= O\left(\varepsilon^{2n+4} + \|v\|^2\right).
 \end{aligned} \tag{5.25}$$

Now, by integrating by parts and Hölder's inequality, we have

$$\begin{aligned}
 & \int_M \varepsilon^n A_a \omega_a^{1+2/n} \Delta_{\theta_a} \chi_\delta(\rho_a) v \theta_0 \wedge (d\theta_0)^n \\
 &= \int_M \varepsilon^n A_a \omega_a^{-1} \Delta_{\theta_a} \chi_\delta(\rho_a) v \theta_a \wedge (d\theta_a)^n \\
 &= \int_M \varepsilon^n A_a \left[ \langle \nabla \omega_a^{-1}, \nabla \chi_\delta(\rho_a) \rangle_{\theta_a} v + \omega_a^{-1} \langle \nabla v, \nabla \chi_\delta(\rho_a) \rangle_{\theta_a} \right] \theta_a \wedge (d\theta_a)^n \\
 &= O\left( \int_{\delta \leq \rho_a \leq 2\delta} \varepsilon^n |\nabla \chi_\delta(\rho_a)| (|v| + |\nabla v|_{\theta_a}) \theta_a \wedge (d\theta_a)^n \right) \\
 &= O\left( \int_{\delta \leq \rho_a \leq 2\delta} \varepsilon^{2n} \theta_a \wedge (d\theta_a)^n + \|v\|^2 \right) \\
 &= O\left(\delta^{2n+2} \varepsilon^{2n} + \|v\|^2\right).
 \end{aligned} \tag{5.26}$$

Substituting (5.23)-(5.26) into (5.22) gives

$$\int_M \mathcal{L}_{\theta_0} \phi_1 v \theta_0 \wedge (d\theta_0)^n = \int_M (2 + 2/n) \varphi^{1+2/n} v \theta_0 \wedge (d\theta_0)^n + O(\delta^{6-2n} \varepsilon^{2n}) + O(\|v\|^2).$$

**Case  $k = 2$ .** Recalling that  $d_2 = \varepsilon \partial_\varepsilon$  we have

$$\begin{aligned}
 & \int_M \mathcal{L}_{\theta_0} \phi_2 v \theta_0 \wedge (d\theta_0)^n = \int_M 2(1 + 1/n)(1 + 2/n) \varphi^{2/n} \phi_2 v \theta_0 \wedge (d\theta_0)^n \\
 &+ \int_M 2(n + 1) \varepsilon^n A_a \omega_a^{1+2/n} \Delta_{\theta_a} \chi_\delta(\rho_a) v \theta_0 \wedge (d\theta_0)^n \\
 &+ \int_M n \omega_a^{1+2/n} \varepsilon^n \left( \sum_{i=1}^8 I_i \right) v \theta_0 \wedge (d\theta_0)^n + E
 \end{aligned} \tag{5.27}$$

where

$$E = \int_M \omega_a^{1+2/n} \varepsilon^n \left( \sum_{i=1}^7 d_2 I_i \right) v \theta_0 \wedge (d\theta_0)^n,$$

thanks to the facts that  $I_8 = 0$ , if  $\mu(M) > 0$  and  $d_2 I_8 = 0$ , if  $\mu(M) = 0$ . In view of the proof of **Case  $k = 1$** , we are left to estimate the term  $E$ . From Lemma 5.9, it

follows that

$$\begin{aligned}
E &= \int_M \omega_a^{-1} \varepsilon^n d_2 \left\{ - (2 + 2/n) \chi_\delta(\rho_a) \left( \Delta_{\theta_a} \left( \frac{1}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2}} \right) \right. \right. \\
&\quad \left. \left. + \frac{\varepsilon^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right) + \frac{\chi_\delta(\rho_a) R_{\theta_a} - (2 + 2/n) \Delta_{\theta_a} \chi_\delta(\rho_a)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2}} \right. \\
&\quad \left. - 2(2 + 2/n) \left\langle \nabla \chi_\delta(\rho_a), \nabla \left( \frac{1}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2}} \right) \right\rangle_{\theta_a} \right. \\
&\quad \left. - (2 + 2/n) n \varepsilon^2 \left[ \left( \frac{\chi_\delta(\rho_a)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2}} + (1 - \chi_\delta(\rho_a)) G_a \right)^{1+2/n} \right. \right. \\
&\quad \quad \left. \left. - \frac{\chi_\delta(\rho_a)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right] \right\} v \theta_a \wedge (d\theta_a)^n \\
&= \int_M \omega_a^{-1} \varepsilon^{n+2} \left\{ \frac{4(n+1)}{n^2} \chi_\delta(\rho_a) \left[ \Delta_{\theta_a} \left( \frac{(\varepsilon/n)^2 + |z|^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right) \right. \right. \\
&\quad \left. \left. - \frac{n(s^2 + ((\varepsilon/n)^2 + |z|^2)^2) - (1 + 2/n) \varepsilon^2 ((\varepsilon/n)^2 + |z|^2)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+2}} \right] \right. \\
&\quad \left. - \frac{(\chi_\delta(\rho_a) R_{\theta_a} - (2 + 2/n) \Delta_{\theta_a} \chi_\delta(\rho_a)) ((\varepsilon/n)^2 + |z|^2)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right\} v \\
&\quad - \frac{8(n+1)}{n^2} \varepsilon^{n+2} \left( \left\langle \nabla \chi_\delta(\rho_a), \nabla(\omega_a^{-1} v) \right\rangle_{\theta_a} + \Delta_{\theta_a}(\chi_\delta(\rho_a)) \omega_a^{-1} v \right) \\
&\quad \times \left( \frac{(\varepsilon/n)^2 + |z|^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right) \\
&\quad - (2 + 2/n) \omega_a^{-1} \varepsilon^n d_2 \left\{ \varepsilon^2 \left[ \left( \frac{\chi_\delta(\rho_a)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2}} + (1 - \chi_\delta(\rho_a)) G_a \right)^{1+2/n} \right. \right. \\
&\quad \quad \left. \left. - \frac{\chi_\delta(\rho_a)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right] \right\} v \theta_a \wedge (d\theta_a)^n \\
&= \int_{\{\rho_a \leq \delta\}} \frac{4(n+1)}{n^2} \omega_a^{-1} \varepsilon^{n+2} \left[ \Delta_{\theta_a} \left( \frac{(\varepsilon/n)^2 + |z|^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right) \right. \\
&\quad \left. - \frac{n(s^2 + ((\varepsilon/n)^2 + |z|^2)^2) - (1 + 2/n) \varepsilon^2 ((\varepsilon/n)^2 + |z|^2)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+2}} \right] v \theta_a \wedge (d\theta_a)^n \\
&\quad + \int_{\{\rho_a \leq \delta\}} - \frac{\omega_a^{-1} \varepsilon^{n+2} R_{\theta_a} ((\varepsilon/n)^2 + |z|^2)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} v \theta_a \wedge (d\theta_a)^n + O(\varepsilon^{2n+4} + \|v\|^2) \\
&:= E_1 + E_2 + O(\varepsilon^{2n+4} + \|v\|^2).
\end{aligned}$$

Observing that

$$\begin{aligned}
\Delta_{\theta_a} \left( \frac{(\varepsilon/n)^2 + |z|^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right) &= (\Delta_{\mathbb{H}^n} + O(\rho_a^2)) \left( \frac{(\varepsilon/n)^2 + |z|^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right) \\
&= \frac{n(s^2 + ((\varepsilon/n)^2 + |z|^2)^2) - (1 + 2/n) \varepsilon^2 ((\varepsilon/n)^2 + |z|^2)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+2}} + \frac{O(\rho_a^2) ((\varepsilon/n)^2 + |z|^2)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}},
\end{aligned}$$

we have the estimate

$$E_1 = O \left( \varepsilon^{2n+4} \int_{\{\rho_a \leq \delta\}} \frac{\rho_a^4 ((\varepsilon/n)^2 + |z|^2)^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_a \wedge (d\theta_a)^n + \|v\|^2 \right)$$

$$= O(\varepsilon^6 + \|v\|^2).$$

Using the fact that  $R_{\theta_a} = O(\rho_a^2)$  we obtain

$$\begin{aligned} E_2 &= O\left(\varepsilon^{2n+4} \int_{\{\rho_a \leq \delta\}} \frac{\rho_a^4((\varepsilon/n)^2 + |z|^2)^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_a \wedge (d\theta_a)^n + \|v\|^2\right) \\ &= O(\varepsilon^6 + \|v\|^2). \end{aligned}$$

Substituting all the estimates above into (5.27) gives

$$\begin{aligned} \int_M \mathcal{L}_{\theta_0} \phi_2 v \theta_0 \wedge (d\theta_0)^n &= \int_M 2(1 + 1/n)(1 + 2/n) \varphi^{2/n} \phi_2 v \theta_0 \wedge (d\theta_0)^n \\ &\quad + O(\delta^{6-2n} \varepsilon^{2n}) + O(\varepsilon^6) + O(\|v\|^2). \end{aligned}$$

**Case  $k = 3$ .** Recalling that  $d_3 = \varepsilon \nabla_a$  where  $a = (z_0, s_0)$ , we have

$$\begin{aligned} \int_M \mathcal{L}_{\theta_0} \phi_3 v \theta_0 \wedge (d\theta_0)^n &= \int_M 2(1 + 1/n)(1 + 2/n) \varphi^{2/n} \phi_3 v \theta_0 \wedge (d\theta_0)^n \\ &\quad + \int_M (2 + 2/n) \varepsilon^{n+1} \nabla_a (A_a \omega_a^{1+2/n} \Delta_{\theta_a} \chi_\delta(\rho_a)) v \theta_0 \wedge (d\theta_0)^n \\ &\quad + \int_M (1 + 2/n) \omega_a^{2/n} \nabla_a (\omega_a) \varepsilon^{n+1} \left( \sum_{i=1}^8 I_i \right) v \theta_0 \wedge (d\theta_0)^n + \bar{E} \end{aligned} \tag{5.28}$$

where

$$\bar{E} = \int_M \omega_a^{1+2/n} \varepsilon^n \left( \sum_{i=1}^8 d_3 I_i \right) v \theta_0 \wedge (d\theta_0)^n.$$

Following the proof of **Case  $k = 1$**  we can get

$$\int_M (2 + 2/n) \varepsilon^{n+1} \nabla_a (A_a \omega_a^{1+2/n} \Delta_{\theta_a} \chi_\delta(\rho_a)) v \theta_0 \wedge (d\theta_0)^n = O(\varepsilon^{2n+2} + \|v\|^2)$$

and

$$\int_M (1 + 2/n) \omega_a^{2/n} \nabla_a (\omega_a) \varepsilon^{n+1} \left( \sum_{i=1}^8 I_i \right) v \theta_0 \wedge (d\theta_0)^n = O(\varepsilon^{2n+2} + \|v\|^2).$$

Now, if  $\mu(M) > 0$ , it follows from the fact  $I_8 = 0$  that

$$\begin{aligned} \bar{E} &= \int_{\{\rho_a \leq \delta\}} \omega_a^{-1} \varepsilon^n \left( \sum_{i=1}^7 d_3 I_i \right) v \theta_0 \wedge (d\theta_0)^n + O(\varepsilon^{2n+2} + \|v\|^2) \\ &= \int_{\{\rho_a \leq \delta\}} \omega_a^{-1} \varepsilon^n d_3 \left\{ - (2 + 2/n) \left[ \Delta_{\theta_a} \left( \frac{1}{((s - s_0)^2 + ((\varepsilon/n)^2 + |z - z_0|^2)^2)^{n/2}} \right) \right. \right. \\ &\quad \left. \left. + \frac{\varepsilon^2}{((s - s_0)^2 + ((\varepsilon/n)^2 + |z - z_0|^2)^2)^{n/2+1}} \right] + \frac{R_{\theta_a}}{((s - s_0)^2 + ((\varepsilon/n)^2 + |z - z_0|^2)^2)^{n/2}} \right\} \\ &\quad v \theta_a \wedge (d\theta_a)^n \Big|_{a=0} + O(\varepsilon^{2n+2} + \|v\|^2). \\ &= \int_{\{\rho_a \leq \delta\}} \omega_a^{-1} \varepsilon^{n+1} \left\{ - (2n + 2) \left[ \Delta_{\theta_a} \left( \frac{\bar{z}((\varepsilon/n)^2 + |z|^2)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right) \right. \right. \\ &\quad \left. \left. + \frac{(1 + 2/n) \varepsilon^2 ((\varepsilon/n)^2 + |z|^2) \bar{z}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+2}} \right] + \frac{\nabla_a R_{\theta_a} (s^2 + ((\varepsilon/n)^2 + |z|^2)^2) - n R_{\theta_a} ((\varepsilon/n)^2 + |z|^2) \bar{z}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right\} \end{aligned}$$

$$\begin{aligned}
& v\theta_a \wedge (d\theta_a)^n + O(\varepsilon^{2n+2} + \|v\|^2) \\
= & O\left(\varepsilon^{2n+2} \int_{\{\rho_a \leq \delta\}} \frac{\rho_a^4((\varepsilon/n)^2 + |z|^2)^2 |z|^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_a \wedge (d\theta_a)^n\right) \\
& + O\left(\varepsilon^{2n+2} \int_{\{\rho_a \leq \delta\}} \frac{\rho_a^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^n} \theta_a \wedge (d\theta_a)^n\right) + O(\varepsilon^{2n+2} + \|v\|^2) \\
= & O(\varepsilon^6 + \varepsilon^{2n+2} + \|v\|^2).
\end{aligned}$$

While  $\mu(M) = 0$ , we notice that

$$\begin{aligned}
d_3 I_8 &= -\varepsilon \left[ \left( \nabla_a (\chi_\delta(\rho_a)) + \frac{n(1 - \chi_\delta(\rho_a))}{2} \right) (\Delta_{\theta_{\mathbb{H}^n}} + O(\rho_a^2)) (\rho_a^{-2n} + \rho_a^{-2n-4} \bar{z}|z|^2) \right] \Big|_{a=0} \\
&= O(\delta^{1-2n} \varepsilon),
\end{aligned}$$

and hence  $\bar{E}$  can be estimated as before. Substituting all the estimates above into (5.28) yields

$$\begin{aligned}
\int_M \mathcal{L}_{\theta_0} \phi_3 v \theta_0 \wedge (d\theta_0)^n &= \int_M 2(1 + 1/n)(1 + 2/n) \varphi^{2/n} \phi_3 v \theta_0 \wedge (d\theta_0)^n \\
&\quad + O(\varepsilon^{2n+2}) + O(\varepsilon^6) + O(\|v\|^2).
\end{aligned}$$

Finally, we are going to estimate the first term on the right hand side of (5.22), (5.27) and (5.28). To this end, we use Proposition (5.10) and expand  $u^{2/n} = (\alpha\varphi + v)^{2/n}$  to obtain

$$0 = \int_M u^{2/n} \phi_k v \theta_0 \wedge (d\theta_0)^n = \int_M (\alpha\varphi)^{2/n} \phi_k v \theta_0 \wedge (d\theta_0)^n + O(\|v\|^2).$$

This implies that

$$\int_M \varphi^{2/n} \phi_k v \theta_0 \wedge (d\theta_0)^n = O(\|v\|^2). \quad (5.29)$$

By inserting (5.29) into (5.22), (5.27) and (5.28), we thus complete the proof of (i).

(ii) By expansion as above, we have

$$\begin{aligned}
\int_M f u^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n &= \int_M f(\alpha\varphi)^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n \\
&\quad + (1 + 2/n) \int_M f(\alpha\varphi)^{2/n} \phi_k v \theta_0 \wedge (d\theta_0)^n + O(\|v\|^2).
\end{aligned} \quad (5.30)$$

It remains to estimate the second term on the right hand side of (5.30). From (5.29) and Lemma 5.12, it follows that

$$\begin{aligned}
& \int_M f \varphi^{2/n} \phi_k v \theta_0 \wedge (d\theta_0)^n \\
&= \int_M (f - f(a)) \varphi^{2/n} \phi_k v \theta_0 \wedge (d\theta_0)^n + \int_M f(a) \varphi^{2/n} \phi_k v \theta_0 \wedge (d\theta_0)^n \\
&= \int_{B_\delta(a)} (f - f(a)) \varphi^{2/n} \phi_k v \theta_0 \wedge (d\theta_0)^n + O(\varepsilon^{2n+4} + \|v\|^2) \\
&= O\left( \int_{B_\delta(a)} |f - f(a)| \varphi^{1+2/n} v \theta_0 \wedge (d\theta_0)^n \right) + O(\varepsilon^{2n+4} + \|v\|^2).
\end{aligned}$$

By Hölder's inequality, we can estimate

$$\begin{aligned}
 & \int_{B_\delta(a)} |f - f(a)| \varphi^{1+2/n} v \theta_0 \wedge (d\theta_0)^n \\
 & \leq \left[ \int_{B_\delta(a)} |f - f(a)|^{\frac{2(n+1)}{n+2}} \varphi^{2+2/n} \theta_0 \wedge (d\theta_0)^n \right]^{\frac{n+2}{2(n+1)}} \left[ \int_M |v|^{2+2/n} \theta_0 \wedge (d\theta_0)^n \right]^{\frac{n}{2(n+1)}} \\
 & = O\left( \left[ \int_{B_\delta(a)} |f - f(a)|^{\frac{2(n+1)}{n+2}} \frac{\varepsilon^{2n+2}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^{n+1})} \theta_a \wedge (d\theta_a)^n \right]^{\frac{n+2}{(n+1)}} + \|v\|^2 \right) \\
 & = O\left( \left[ \int_{B_{\frac{\delta}{\varepsilon}}(a)} \frac{|f(\varepsilon z, \varepsilon^2 s) - f(a)|^{\frac{2(n+1)}{n+2}}}{(s^2 + ((1/n)^2 + |z|^2)^{n+1})} \theta_a \wedge (d\theta_a)^n \right]^{\frac{n+2}{(n+1)}} + \|v\|^2 \right) \\
 & = O\left( \left[ \int_{B_{\frac{\delta}{\varepsilon}}(a)} \frac{(\varepsilon |\nabla f(a)|_{\theta_0}|(z, \varepsilon s)|)^{\frac{2(n+1)}{n+2}} + O((\varepsilon|(z, \varepsilon s)|)^{\frac{4(n+1)}{n+2}})}{(s^2 + ((1/n)^2 + |z|^2)^{n+1})} \theta_a \wedge (d\theta_a)^n \right]^{\frac{n+2}{(n+1)}} + \|v\|^2 \right) \\
 & = O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \varepsilon^4 + \|v\|^2) = O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \|v\|^2).
 \end{aligned}$$

Plugging the two estimates above into (5.30) gives

$$\int_M f u^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n = \int_M f(\alpha\varphi)^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n + O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \|v\|^2).$$

This completes the proof of (ii).  $\square$

To proceed further, we consider the so-called shadow flow, which will build the relation between the parameters such as  $a, \varepsilon, \alpha$  and the prescribed function  $f$ .

**Lemma 5.14.** *Set*

$$\sigma_k = - \int_M (\mathcal{L}_{\theta_0} u - \lambda f u^{1+2/n}) \phi_k \theta_0 \wedge (d\theta_0)^n, \quad k = 1, 2, 3.$$

For  $t$  sufficiently large, there hold

- (i)  $\dot{\alpha}/\alpha = c_1^{-1} \alpha^{-(1+2/n)} \sigma_1 (1 + O(\varepsilon)) + R_1$ ,
- (ii)  $\dot{\varepsilon}/\varepsilon = c_2^{-1} \alpha^{-(1+2/n)} \sigma_2 (1 + O(\varepsilon)) + R_2$ ,
- (iii)  $\dot{a}/\varepsilon = c_3^{-1} \alpha^{-(1+2/n)} \sigma_3 (1 + O(\varepsilon)) + R_3$ ,

where  $R_k = O(\|v\|^2 + F_2(\theta(t)))_k$  and  $c_k$  are constants appeared in Lemma 5.12 (ii).

**Proof.** Let  $(\zeta^1, \zeta^2, \zeta^3) = (\dot{\alpha}, \alpha \dot{\varepsilon} \varepsilon^{-1}, \alpha \dot{a} \varepsilon^{-1})$ . Using the flow  $u_t = -(R_{\theta(t)} - \lambda f)u$  and the fact that  $\int_M u^{2/n} \phi_k v \theta_0 \wedge (d\theta_0)^n = 0$  and recalling  $(\phi_1, \phi_2, \phi_3) = (\varphi, \varepsilon \partial_\varepsilon \varphi, \varepsilon \nabla_a \varphi)$ , we obtain

$$\begin{aligned}
 \sigma_k & = \int_M u_t u^{2/n} \phi_k \theta_0 \wedge (d\theta_0)^n = \int_M \partial_t (\alpha\varphi + v) u^{2/n} \phi_k \theta_0 \wedge (d\theta_0)^n \\
 & = \zeta^l \int_M u^{2/n} \phi_k \phi_l \theta_0 \wedge (d\theta_0)^n - \int_M (\partial_t u^{2/n}) \phi_k v \theta_0 \wedge (d\theta_0)^n \\
 & \quad - \int_M u^{2/n} (\partial_t \phi_k) v \theta_0 \wedge (d\theta_0)^n \\
 & := \zeta^l I - II - III.
 \end{aligned} \tag{5.31}$$

**Estimate of I.** It follows from expansion and Lemma 5.12 (i) and (iv) that

$$I = \int_M (\alpha\varphi)^{2/n} \phi_k \phi_l \theta_0 \wedge (d\theta_0)^n + O(\|v\|)_{k,l}$$

$$\begin{aligned}
&= \alpha^{2/n} \delta_{lk} \int_M \varphi^{2/n} \phi_k^2 \theta_0 \wedge (d\theta_0)^n + (1 - \delta_{lk}) \alpha^{2/n} \int_M \varphi^{2/n} \phi_k \phi_l \theta_0 \wedge (d\theta_0)^n + O(\|v\|)_{k,l} \\
&= \alpha^{2/n} c_k \delta_{lk} + O(\varepsilon + \|v\|)_{k,l}.
\end{aligned}$$

**Estimate of II.** By (3.3), Hölder's inequality and Lemma 5.12 (i), we have

$$\begin{aligned}
|II| &= \left| \frac{2}{n} \int_M (R_{\theta(t)} - \lambda f) u^{2/n} \phi_k v \theta_0 \wedge (d\theta_0)^n \right| \\
&\leq C \int_M |R_{\theta(t)} - \lambda f| u^{2/n} \varphi |v| \theta_0 \wedge (d\theta_0)^n \\
&\leq C \int_M |R_{\theta(t)} - \lambda f| u^{2/n} |u - v| |v| \theta_0 \wedge (d\theta_0)^n \\
&\leq C \int_M |R_{\theta(t)} - \lambda f| u^3 |v| \theta_0 \wedge (d\theta_0)^n + C \int_M |R - \lambda f| u^2 |v|^2 \theta_0 \wedge (d\theta_0)^n \\
&\leq C \|v\| \left( \int_M (R_{\theta(t)} - \lambda f)^2 u^4 \theta_0 \wedge (d\theta_0)^n \right)^{1/2} \left( \int_M u^4 \theta_0 \wedge (d\theta_0)^n \right)^{1/4} \\
&\quad + C \|v\|^2 \left( \int_M (R_{\theta(t)} - \lambda f)^2 u^4 \theta_0 \wedge (d\theta_0)^n \right)^{1/2} \\
&\leq C (F_2(\theta(t)) + \|v\|^2),
\end{aligned}$$

since  $n = 1$ .

**Estimate of III.** Using Lemma 5.12 (i) again, we get

$$III = \zeta^l \int_M u^{2/n} d_l \phi_k v \theta_0 \wedge (d\theta_0)^n = O\left( \int_M u^{1+2/n} v \theta_0 \wedge (d\theta_0)^n \right)_{k,l} \zeta^l = O(\|v\|)_{k,l} \zeta^l.$$

Now, substituting the estimates for  $I, II, III$  into (5.31) gives

$$\sigma_k = \left[ \alpha^{2/n} c_k \delta_{kl} + O(\varepsilon + \|v\|)_{k,l} \right] \zeta^l + O(F_2(\theta(t)) + \|v\|^2)_k.$$

Setting  $E_{k,l} = \alpha^{2/n} c_k \delta_{kl} + O(\varepsilon + \|v\|)_{k,l}$ , one then has

$$E_{k,l} \zeta^l = \sigma_k + O(F_2(\theta(t)) + \|v\|^2)_k$$

Using the fact that

$$E_{k,l}^{-1} = \alpha^{-2/n} c_k^{-1} \delta_{kl} + O(\varepsilon + \|v\|)_{k,l}$$

and using the definition of  $\sigma_k$  that

$$\sigma_k = O(F_2(\theta(t))^{1/2}),$$

we can obtain

$$\begin{aligned}
\zeta^l &= \sigma_k \left[ \alpha^{-2/n} c_k^{-1} \delta_{kl} + O(\varepsilon + \|v\|)_{k,l} \right] + O(F_2(\theta(t)) + \|v\|^2)_l \\
&= \alpha^{-2/n} c_l^{-1} \sigma_l (1 + O(\varepsilon)) + O(F_2(\theta(t)) + \|v\|^2)_l.
\end{aligned}$$

In view of the definition of  $\zeta^l$ , the proof is thus complete.  $\square$

To relate the parameters  $\alpha, \varepsilon$  and  $a$  with the prescribed function  $f$ , it remains to refine the estimate of the shadow flows  $\sigma_k$ .

**Lemma 5.15.** *There exist positive constants  $d_1, d_2, \dots, e_4$  such that, for  $t$  large, there hold*

$$\begin{aligned}
 \text{(i)} \quad \sigma_1 &= (2 + 2/n)\alpha \left( \frac{n\lambda\alpha^{2/n}f(a)}{2(n+1)} - 1 \right) \int_M \varphi^{2+2/n} \theta_0 \wedge (d\theta_0)^n + \alpha d_1 A_a \varepsilon^{2n} \\
 &\quad + \alpha^{1+2/n} \lambda e_1 \varepsilon^2 \Delta_{\theta_0} f(a) + R_1; \\
 \text{(ii)} \quad \sigma_2 &= (2 + 2/n)\alpha \left( \frac{n\lambda\alpha^{2/n}f(a)}{2(n+1)} - 1 \right) \int_M \varphi^{1+2/n} \phi_2 \theta_0 \wedge (d\theta_0)^n + \alpha d_2 A_a \varepsilon^{2n} \\
 &\quad + \alpha^{1+2/n} \lambda e_2 \varepsilon^2 \Delta_{\theta_0} f(a) + R_2; \\
 \text{(iii)} \quad \sigma_3 &= (2 + 2/n)\alpha \left( \frac{n\lambda\alpha^{2/n}f(a)}{2(n+1)} - 1 \right) \int_M \varphi^{1+2/n} \phi_3 \theta_0 \wedge (d\theta_0)^n \\
 &\quad + \alpha^{1+2/n} \lambda (e_3 \varepsilon \nabla_{\theta_0} f(a) + e_4 \varepsilon^3 \nabla_{\theta_0} \Delta_{\theta_0} f(a)) + R_3,
 \end{aligned}$$

where

$$R_k = O\left(\delta^{4-2n} \varepsilon^{2n} + \delta \varepsilon^{2n} + \varepsilon^4 + |\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \|v\|^2\right)_k \text{ for } k = 1, 2, 3.$$

**Proof.** Recall that

$$\sigma_k = - \int_M (\mathcal{L}_{\theta_0} u - \lambda f u^{1+2/n}) \phi_k \theta_0 \wedge (d\theta_0)^n.$$

**Estimate of**  $\int_M \mathcal{L}_{\theta_0} u \phi_k \theta_0 \wedge (d\theta_0)^n$ . By integrating by parts, we may write

$$\begin{aligned}
 \int_M \mathcal{L}_{\theta_0} u \phi_k \theta_0 \wedge (d\theta_0)^n &= \int_M \mathcal{L}_{\theta_0} (\alpha \varphi + v) \phi_k \theta_0 \wedge (d\theta_0)^n \\
 &= \alpha \int_M (\mathcal{L}_{\theta_0} \varphi) \phi_k \theta_0 \wedge (d\theta_0)^n + \int_M (\mathcal{L}_{\theta_0} \phi_k) v \theta_0 \wedge (d\theta_0)^n. \quad (5.32)
 \end{aligned}$$

The first term on the right hand side of (5.32) can be estimated as follows. Using Lemmas 5.9 and 5.12 and following the proof of (5.23)-(5.25), we have

$$\begin{aligned}
 &\alpha \int_M (\mathcal{L}_{\theta_0} \varphi) \phi_k \theta_0 \wedge (d\theta_0)^n \\
 &= (2 + 2/n)\alpha \int_M \varphi^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n \\
 &\quad + (2 + 2/n)\alpha \varepsilon^n \int_M \omega_a^{1+2/n} A_a \Delta_{\theta_a} \chi_\delta(\rho_a) \phi_k \theta_0 \wedge (d\theta_0)^n \\
 &\quad + O\left( \int_M \left[ \left( \frac{\varepsilon^2}{s^2 + ((\varepsilon/n)^2 + |z|^2)^2} \right)^{n/2} \rho_a^2 1_{\{\rho_a \leq 2\delta\}} + \delta^{-1} \varepsilon^n 1_{\{\delta \leq \rho_a \leq 2\delta\}} \right. \right. \\
 &\quad \left. \left. + \left( \frac{\varepsilon^2}{s^2 + ((\varepsilon/n)^2 + |z|^2)^2} \right)^{1+n/2} 1_{\{\rho_a \geq \delta\}} \right] \phi_k \theta_a \wedge (d\theta_a)^n \right) \\
 &:= (2 + 2/n)\alpha \int_M \varphi^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n + I + O\left(\delta^{4-2n} \varepsilon^{2n} + \delta \varepsilon^{2n} + \varepsilon^{2n+2}\right).
 \end{aligned}$$

It remains to estimate the term  $I$  above. As always, we split the argument into three cases.

**Case  $k = 1$ .** By denoting that  $\psi = \psi_{a,\varepsilon}$  which was defined in (5.21), we may write  $I$  as

$$\begin{aligned}
 I &= (2 + 2/n)\alpha \varepsilon^n A_a \int_M \Delta_{\theta_a} \chi_\delta(\rho_a) \psi \theta_a \wedge (d\theta_a)^n \\
 &= (2 + 2/n)\alpha \varepsilon^n A_a \left[ \int_M \Delta_{\theta_a} \chi_\delta(\rho_a) \varepsilon^n \rho_a^{-2n} \theta_a \wedge (d\theta_a)^n \right. \\
 &\quad \left. + \int_M \Delta_{\theta_a} \chi_\delta(\rho_a) (\psi - \varepsilon^n \rho_a^{-2n}) \theta_a \wedge (d\theta_a)^n \right]
 \end{aligned}$$

$$:= (2 + 2/n)\alpha\varepsilon^n A_a(I_1 + I_2).$$

To estimate  $I_1$ , we use integration by parts to get

$$\begin{aligned} I_1 &= -\varepsilon^n \int_M \langle \nabla \rho_a^{-2n}, \nabla \chi_\delta(\rho_a) \rangle_{\theta_a} \theta_a \wedge (d\theta_a)^n \\ &= 2n \int_{\{\delta \leq \rho_a \leq 2\delta\}} \varepsilon^n \partial_{\rho_a}(\chi_\delta(\rho_a)) \rho_a^{-2n-1} |\nabla \rho_a|_{\theta_a}^2 \theta_a \wedge (d\theta_a)^n \\ &= 2n\varepsilon^n |B^n| \int_\delta^{2\delta} \partial_\rho(\chi_\delta(\rho)) d\rho = -2n|\mathbb{S}^{2n+1}| \varepsilon^n, \end{aligned}$$

where  $|B^n|$  is the volume of the unit ball in  $\mathbb{H}^n$ . For  $I_2$ , in view of the definition of  $\psi$ , we can estimate

$$\begin{aligned} |I_2| &\leq \frac{C\varepsilon^n}{\delta^2} \left[ \int_{\{\delta \leq \rho_a \leq 2\delta\}} \left| \frac{1}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2})} - \frac{1}{(s^2 + |z|^4)^{n/2}} \right| \theta_a \wedge (d\theta_a)^n \right. \\ &\quad \left. + \int_{\{\delta \leq \rho_a \leq 2\delta\}} \left| G_a - \rho_a^{-2n} \right| \theta_a \wedge (d\theta_a)^n \right] \\ &\leq \frac{C\varepsilon^n}{\delta^2} \left[ \int_{\{\delta \leq \rho_a \leq 2\delta\}} \left| \frac{\varepsilon^2}{(s^2 + |z|^4)^{n/2+1}} \right| \theta_a \wedge (d\theta_a)^n \right. \\ &\quad \left. + \int_{\{\delta \leq \rho_a \leq 2\delta\}} A_a + O(\rho_a) \theta_a \wedge (d\theta_a)^n \right] \\ &= O(\varepsilon^{n+2} + \delta^{2n} \varepsilon^n). \end{aligned}$$

Combining all the estimates yields

$$I = -4(n+1)\alpha|B^n|A_a\varepsilon^{2n} + O(\delta^{2n}\varepsilon^{2n} + \varepsilon^{2n+2}).$$

**Case  $k = 2$ .** From (A.1) and the conclusion in Case  $k = 1$ , it follows that

$$\begin{aligned} I &= (2 + 2/n)\alpha A_a \varepsilon^n \int_M \omega_a^{1+2/n} \Delta_{\theta_a} \chi_\delta(\rho_a) \left[ n\varphi - \frac{2n\omega_a \varepsilon^{n+2} \chi_\delta(\rho_a) ((\varepsilon/n)^2 + |z|^2)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2+1}} \right] \theta_0 \wedge (d\theta_0)^n \\ &= 2(n+1)\alpha A_a \varepsilon^n \int_M \Delta_{\theta_a} \chi_\delta(\rho_a) \psi \theta_a \wedge (d\theta_a)^n + O(\varepsilon^{2n+2}) \\ &= -4n(n+1)\alpha|B^n|A_a\varepsilon^{2n} + O(\delta^{2n}\varepsilon^{2n} + \varepsilon^{2n+2}). \end{aligned}$$

**Case  $k = 3$ .** From (A.2), it is easy to see that

$$I = (2 + 2/n)\alpha A_a \varepsilon^n \int_M \omega_a^{1+2/n} \Delta_{\theta_a} \chi_\delta(\rho_a) \phi_3 \theta_0 \wedge (d\theta_0)^n = O(\varepsilon^{2n+1}).$$

Plugging all the estimates of three cases and the estimate in Lemma 5.13 (i) into (5.32) yields

$$\begin{aligned} - \int_M \mathcal{L}_{\theta_0} u \phi_k \theta_0 \wedge (d\theta_0)^n &= -(2 + 2/n)\alpha \int_M \varphi^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n + \alpha d_k A_a \varepsilon^{2n} \\ &\quad + O(\delta^{4-2n} \varepsilon^{2n} + \delta \varepsilon^{2n} + \varepsilon^{2n+1} + \varepsilon^6 + \|v\|^2), \end{aligned} \tag{5.33}$$

where  $d_1 = 4(n+1)|B^n|$ ,  $d_2 = 4n(n+1)|B^n|$  and  $d_3 = 0$ .



**Estimate of  $\int_M fu^{1+2/n}\phi_k\theta_0 \wedge (d\theta_0)^n$ .** By Lemma 5.13, we can get

$$\begin{aligned}
 & \int_M fu^{1+2/n}\phi_k\theta_0 \wedge (d\theta_0)^n \\
 &= \int_M f(\alpha\varphi)^{1+2/n}\phi_k\theta_0 \wedge (d\theta_0)^n + O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \|v\|^2) \\
 &= \alpha^{1+2/n} \int_M (f - f(a))\varphi^{1+2/n}\phi_k\theta_0 \wedge (d\theta_0)^n \\
 &\quad + \alpha^{1+2/n} f(a) \int_M \varphi^{1+2/n}\phi_k\theta_0 \wedge (d\theta_0)^n + O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \|v\|^2) \\
 &:= \alpha^{1+2/n} I + \alpha^{1+2/n} f(a) \int_M \varphi^{1+2/n}\phi_k\theta_0 \wedge (d\theta_0)^n + O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \|v\|^2).
 \end{aligned} \tag{5.34}$$

To estimate  $I$  in (5.34), we expand  $f(\varepsilon z, \varepsilon^2 s) - f(a)$  in  $B_{\delta/\varepsilon}(a)$  to obtain

$$\begin{aligned}
 f(\varepsilon z, \varepsilon^2 s) - f(a) &= \nabla_{\theta_0} f(a) \cdot (\varepsilon z, \varepsilon^2 s) + \frac{1}{2} (\nabla d)_{\theta_0} f(a) \cdot (\varepsilon z, \varepsilon^2 s)^2 \\
 &\quad + \frac{1}{6} (\nabla^2 d)_{\theta_0} f(a) \cdot (\varepsilon z, \varepsilon^2 s)^3 + O(|(\varepsilon z, \varepsilon^2 s)|^4).
 \end{aligned}$$

**Case  $k = 1$ .** Using the expansion above and symmetry, we have

$$\begin{aligned}
 I &= \int_M (f - f(a))\varphi^{2+2/n}\theta_0 \wedge (d\theta_0)^n \\
 &= \int_{B_{\delta}(a)} (f - f(a)) \frac{\varepsilon^{2n+2}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+1}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+2}) \\
 &= \int_{B_{\frac{\delta}{\varepsilon}}(a)} \frac{(f(\varepsilon x, \varepsilon^2 s) - f(a))}{(s^2 + ((1/n)^2 + |z|^2)^2)^{n+1}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+2}) \\
 &= e_1 \Delta_{\theta_0} f(a) \varepsilon^2 + O(\varepsilon^4 + \varepsilon^{2n+2}),
 \end{aligned}$$

where

$$e_1 = \frac{1}{2n} \int_{\mathbb{H}^n} \frac{|z|^2}{(s^2 + ((1/n)^2 + |z|^2)^2)^{n+1}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n.$$

**Case  $k = 2$ .** By (A.1), the expansion above and symmetry, we obtain

$$\begin{aligned}
 I &= \int_M (f - f(a))\varphi^{1+2/n}\phi_2\theta_0 \wedge (d\theta_0)^n \\
 &= \int_M (f - f(a))\varphi^{1+2/n} \left( n\varphi - \frac{2\omega_a \varepsilon^{n+2} \chi_{\delta}(\rho_a) ((\varepsilon/n)^2 + |z|^2)}{n(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right) \theta_0 \wedge (d\theta_0)^n \\
 &= n \int_M (f - f(a))\varphi^{2+2/n}\theta_0 \wedge (d\theta_0)^n \\
 &\quad - \int_M (f - f(a))\varphi^{1+2/n} \frac{2\omega_a \varepsilon^{n+2} \chi_{\delta}(\rho_a) ((\varepsilon/n)^2 + |z|^2)}{n(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \theta_0 \wedge (d\theta_0)^n \\
 &= n \int_{B_{\delta}(a)} (f - f(a)) \frac{\varepsilon^{2n+2}(s^2 + |z|^4 - (\varepsilon/n)^4)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_a \wedge (d\theta_a)^n + O(\varepsilon^{2n+2}) \\
 &= n \int_{B_{\frac{\delta}{\varepsilon}}(a)} \frac{(f(\varepsilon z, \varepsilon^2 s) - f(a))(s^2 + |z|^4 - (1/n)^4)}{(s^2 + ((1/n)^2 + |z|^2)^2)^{n+2}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+2})
 \end{aligned}$$

$$= e_2 \Delta_{\theta_0} f(a) \varepsilon^2 + O(\varepsilon^4 + \varepsilon^{2n+2}),$$

where

$$e_2 = \frac{1}{2n} \int_{\mathbb{H}^n} \frac{|z|^2 (s^2 + |z|^4 - (1/n)^4)}{(s^2 + ((1/n)^2 + |z|^2)^2)^{n+2}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n.$$

**Case  $k = 3$ .** Recalling that  $\phi_3 = d_3 \varphi = \varepsilon \nabla_a \varphi|_{a=0}$  and applying the conclusion in **Case  $k = 1$** , we have

$$\begin{aligned} I &= \int_M (f - f(a)) \varphi^{1+2/n} \phi_3 \theta_0 \wedge (d\theta_0)^n = \int_M (f - f(a)) \varphi^{1+2/n} d_3 \varphi \theta_0 \wedge (d\theta_0)^n \\ &= \frac{n}{2(n+1)} \left( \int_M d_3(f(a)) \varphi^{2+2/n} \theta_0 \wedge (d\theta_0)^n + d_3 \int_M (f - f(a)) \varphi^{2+2/n} \theta_0 \wedge (d\theta_0)^n \right) \\ &= e_3 \nabla_{\theta_0} f(a) \varepsilon + e_4 \nabla_{\theta_0} \Delta_{\theta_0} f(a) \varepsilon^3 + O(\varepsilon^4 + \varepsilon^{2n+2}), \end{aligned}$$

where

$$e_3 = \frac{n}{2(n+1)} \int_{\mathbb{H}^n} \frac{1}{(s^2 + ((1/n)^2 + |z|^2)^2)^{n+1}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n$$

and

$$e_4 = \frac{1}{4(n+1)} \int_{\mathbb{H}^n} \frac{|z|^2}{(s^2 + ((1/n)^2 + |z|^2)^2)^{n+1}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n.$$

Substituting all the estimates above into (5.34) we obtain

$$\begin{aligned} \lambda \int_M f u^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n &= \lambda \alpha^{1+2/n} D_k + \lambda \alpha^{1+2/n} f(a) \int_M \varphi^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n \\ &\quad + O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \varepsilon^4 + \varepsilon^{2n+2} + \|v\|^2) \end{aligned} \quad (5.35)$$

where

$$D_k = \begin{cases} e_1 \Delta_{\theta_0} f(a) \varepsilon^2, & \text{for } k = 1, \\ e_2 \Delta_{\theta_0} f(a) \varepsilon^2, & \text{for } k = 2, \\ e_3 \nabla_{\theta_0} f(a) \varepsilon + e_4 \nabla_{\theta_0} \Delta_{\theta_0} f(a) \varepsilon^3, & \text{for } k = 3. \end{cases}$$

Now, plugging (5.33) and (5.35) into the expression of  $\sigma_k$ , we thus complete the proof.  $\square$

Since  $\sigma_1$  can be controlled by  $F_2(\theta(t))$ , we may simplify the estimates of  $\sigma_k$  as follows.

**Corollary 5.16.** *For  $t$  large, there hold*

$$(i) \quad \sigma_2 = d_2 \alpha A_a \varepsilon^{2n} + e_2 \alpha^{1+2/n} \lambda \Delta_{\theta_0} f(a) \varepsilon^2 + R_2$$

$$(ii) \quad \sigma_3 = \alpha^{1+2/n} \lambda (e_3 \nabla_{\theta_0} f(a) \varepsilon + e_4 \nabla_{\theta_0} \Delta_{\theta_0} f(a) \varepsilon^3) + R_3,$$

where

$$R_k = (\delta^{4-2n} \varepsilon^{2n} + \delta \varepsilon^{2n} + \varepsilon^4 + F_2(\theta(t)) + \|v\|^2)_k, \quad \text{for } k = 2, 3.$$

**Proof.** Notice by (3.3) and Hölder's inequality that

$$\begin{aligned} |\sigma_1| &= \left| \int_M (R_{\theta(t)} - \lambda f) u^3 \varphi \theta_0 \wedge (d\theta_0)^n \right| \\ &\leq C \left( \int_M (R_{\theta(t)} - \lambda f)^2 u^4 \theta_0 \wedge (d\theta_0)^n \right)^{1/2} \left( \int_M u^4 \theta_0 \wedge (d\theta_0)^n \right)^{1/4} \leq C F_2(\theta(t))^{1/2}, \end{aligned}$$

since  $n = 1$ . Hence, it follows from Lemma 5.15 (i) that

$$\frac{n\lambda\alpha^{2/n}f(a)}{2(n+1)} = 1 + O(\varepsilon^{2n} + [|\nabla f(a)|_{\theta_0}^2 + \Delta_{\theta_0}f(a)]\varepsilon^2 + F_2(\theta(t))^{1/2} + \|v\|^2). \quad (5.36)$$

Moreover, by Lemma 5.12 (iii), we have

$$\int_M \varphi^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n = O(\varepsilon^{2n+2}), \quad \text{for } k = 2, 3.$$

Plugging this and (5.36) into Lemma 5.15 (ii) and (iii) yields the conclusion.  $\square$

To proceed further, we consider the functional

$$\mathcal{E}_f(u) = \frac{\mathcal{E}(u)}{\left(\int_M f u^{2+2/n} \theta_0 \wedge (d\theta_0)^n\right)^{\frac{n}{n+1}}} := \frac{\mathcal{E}(u)}{f_u^{\frac{n}{n+1}}},$$

where  $\mathcal{E}(u)$  is defined in (2.4). Also, for each  $t$ , we set

$$H_u(\mu_0) = \left\{ \psi : \int_M \psi \phi_k u^{2/n} \theta_0 \wedge (d\theta_0)^n = 0 \text{ for all } k = 1, 2, 3 \right\},$$

where  $\mu_0$  is defined in Proposition 5.10. Recalling that  $u = \alpha\varphi + v$ , we then have

**Lemma 5.17.** *For  $t$  large and for  $h_1, h_2 \in H_u(\mu_0)$ , one has*

$$\begin{aligned} \text{(i)} \quad & \|\partial \mathcal{E}_f(\alpha\varphi)|_{H_u(\mu)}\| = O(\delta^{4-3n}\varepsilon^n + |\nabla f(a)|_{\theta_0}\varepsilon + |\Delta_{\theta_0}f(a)|\varepsilon^2 + \varepsilon^3 + \varepsilon^{n+2} \\ & \quad + F_2(\theta(t))^{1/2} + \|v\|^2); \\ \text{(ii)} \quad & \frac{1}{2}\partial^2 \mathcal{E}_f(\alpha\varphi)h_1h_2 = f_{\alpha\varphi}^{-\frac{n}{n+1}} \left[ \int_M (\mathcal{L}_{\theta_0} h_1)h_2 \theta_0 \wedge (d\theta_0)^n \right. \\ & \quad \left. - \frac{2(n+1)(n+2)}{n^2} \int_M \varphi^{2/n} h_1 h_2 \theta_0 \wedge (d\theta_0)^n \right] + O(\mu_0)\|h_1\|\|h_2\|. \end{aligned}$$

**Proof.** Given  $h \in H_u(\mu_0)$  with  $\|h\| = 1$ , a straightforward calculation then shows that

$$\frac{1}{2}\partial \mathcal{E}_f(\alpha\varphi)h = f_{\alpha\varphi}^{-\frac{n}{n+1}} \left[ \int_M \mathcal{L}_{\theta_0}(\alpha\varphi)h \theta_0 \wedge (d\theta_0)^n - \frac{\mathcal{E}(\alpha\varphi)}{f_{\alpha\varphi}} \int_M f(\alpha\varphi)^{1+2/n} h \theta_0 \wedge (d\theta_0)^n \right] \quad (5.37)$$

and

$$\begin{aligned} & \frac{1}{2}\partial^2 \mathcal{E}_f(\alpha\varphi)h_1h_2 \\ & = f_{\alpha\varphi}^{-\frac{n}{n+1}} \left[ \int_M (\mathcal{L}_{\theta_0} h_1)h_2 \theta_0 \wedge (d\theta_0)^n - (1+2/n)\frac{\mathcal{E}(\alpha\varphi)}{f_{\alpha\varphi}} \int_M f(\alpha\varphi)^{2/n} h_1 h_2 \theta_0 \wedge (d\theta_0)^n \right] \\ & \quad - 2f_{\alpha\varphi}^{-\frac{2n+1}{n+1}} \left[ \int_M \mathcal{L}_{\theta_0}(\alpha\varphi)h_1 \theta_0 \wedge (d\theta_0)^n \int_M f(\alpha\varphi)^{1+2/n} h_2 \theta_0 \wedge (d\theta_0)^n \right. \\ & \quad \left. + \int_M \mathcal{L}_{\theta_0}(\alpha\varphi)h_2 \theta_0 \wedge (d\theta_0)^n \int_M f(\alpha\varphi)^{1+2/n} h_1 \theta_0 \wedge (d\theta_0)^n \right] \\ & \quad + (4+2/n)\frac{\mathcal{E}(\alpha\varphi)}{f_{\alpha\varphi}^{\frac{3n+2}{n+1}}} \int_M f(\alpha\varphi)^{1+2/n} h_1 \theta_0 \wedge (d\theta_0)^n \int_M f(\alpha\varphi)^{1+2/n} h_2 \theta_0 \wedge (d\theta_0)^n. \end{aligned} \quad (5.38)$$

By (2.8), we may write

$$\begin{aligned}\mathcal{E}(u) &= \int_M R_{\theta(t)} u^{2+2/n} \theta_0 \wedge (d\theta_0)^n \\ &= \int_M (R_{\theta(t)} - \lambda f) u^{2+2/n} \theta_0 \wedge (d\theta_0)^n + \lambda = \lambda + O(F_2(\theta(t))^{1/2}).\end{aligned}\quad (5.39)$$

This together with Lemma 5.13 (i) implies that

$$\begin{aligned}\mathcal{E}(\alpha\varphi) &= \int_M \mathcal{L}_{\theta_0}(\alpha\varphi)(\alpha\varphi) \theta_0 \wedge (d\theta_0)^n = \int_M \mathcal{L}_{\theta_0}(u-v)(u-v) \theta_0 \wedge (d\theta_0)^n \\ &= \int_M u \mathcal{L}_{\theta_0} u \theta_0 \wedge (d\theta_0)^n - 2 \int_M v \mathcal{L}_{\theta_0} u \theta_0 \wedge (d\theta_0)^n + \int_M v \mathcal{L}_{\theta_0} v \theta_0 \wedge (d\theta_0)^n \\ &= \mathcal{E}(u) - 2 \int_M v \mathcal{L}_{\theta_0}(\alpha\varphi) \theta_0 \wedge (d\theta_0)^n - \int_M v \mathcal{L}_{\theta_0} v \theta_0 \wedge (d\theta_0)^n \\ &= \lambda + O(\delta^{6-2n} \varepsilon^{2n} + \varepsilon^6 + F_2(\theta(t))^{1/2} + \|v\|^2).\end{aligned}$$

By Lemma 5.13 (ii) and (2.8), we can get

$$\begin{aligned}f_{\alpha\varphi} &= \int_M f(\alpha\varphi)^{2+2/n} \theta_0 \wedge (d\theta_0)^n = \int_M f(\alpha\varphi)^{1+2/n} \alpha\varphi_1 \theta_0 \wedge (d\theta_0)^n \\ &= \int_M f u^{1+2/n} \alpha\varphi \theta_0 \wedge (d\theta_0)^n + O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \|v\|^2) \\ &= \int_M f u^{1+2/n} (u-v) \theta_0 \wedge (d\theta_0)^n + O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \|v\|^2) \\ &= 1 - \int_M f u^{1+2/n} v \theta_0 \wedge (d\theta_0)^n + O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \|v\|^2) \\ &= 1 + O(|\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + \|v\|^2).\end{aligned}$$

Combining the two estimates above yields

$$\frac{\mathcal{E}(\alpha\varphi)}{f_{\alpha\varphi}} = \lambda + O(\delta^{6-2n} \varepsilon^{2n} + \varepsilon^6 + |\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + F_2(\theta(t))^{1/2} + \|v\|^2). \quad (5.40)$$

Substituting this into (5.37) gives

$$\begin{aligned}\frac{1}{2} \partial \mathcal{E}_f(\alpha\varphi) h &= f_{\alpha\varphi}^{-\frac{n}{n+1}} \left[ \int_M \mathcal{L}_{\theta_0}(\alpha\varphi) h \theta_0 \wedge (d\theta_0)^n - \lambda \int_M f(\alpha\varphi)^{1+2/n} h \theta_0 \wedge (d\theta_0)^n \right] \\ &\quad + O(\delta^{6-2n} \varepsilon^{2n} + \varepsilon^6 + |\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + F_2(\theta(t))^{1/2} + \|v\|^2).\end{aligned}\quad (5.41)$$

By Lemma 5.9 and following the proof of Lemma 5.13 (i), we have

$$\int_M \mathcal{L}_{\theta_0}(\alpha\varphi) h \theta_0 \wedge (d\theta_0)^n = (2 + 2/n) \alpha \int_M \varphi^{1+2/n} h \theta_0 \wedge (d\theta_0)^n + O(\delta^{4-3n} \varepsilon^n). \quad (5.42)$$

By the expansion of  $f$  around  $a$  and the proof of Lemma 5.15 (i), we can obtain

$$\begin{aligned}\int_M \lambda f(\alpha\varphi)^{1+2/n} h \theta_0 \wedge (d\theta_0)^n \\ &= \lambda \alpha^{1+2/n} f(a) \int_M \varphi^{1+2/n} h \theta_0 \wedge (d\theta_0)^n + \int_M \lambda (f - f(a)) (\alpha\varphi)^{1+2/n} h \theta_0 \wedge (d\theta_0)^n \\ &= \lambda \alpha^{1+2/n} f(a) \int_M \varphi^{1+2/n} h \theta_0 \wedge (d\theta_0)^n + O(|\nabla f(a)|_{\theta_0} \varepsilon + |\Delta_{\theta_0} f(a)| \varepsilon^2 + \varepsilon^3 + \varepsilon^{n+2}).\end{aligned}\quad (5.43)$$

Plugging the estimates (5.42) and (5.43) into (5.41) gives

$$\begin{aligned} \frac{1}{2} \partial \mathcal{E}_f(\alpha\varphi)h &= f_{\alpha\varphi}^{-\frac{n}{n+1}}(2 + 2/n)\alpha \left(1 - \frac{n\lambda\alpha^{2/n}f(a)}{2(n+1)}\right) \int_M \varphi^{1+2/n}h\theta_0 \wedge (d\theta_0)^n \\ &\quad + O(\delta^{4-3n}\varepsilon^n + |\nabla f(a)|_{\theta_0}\varepsilon + |\Delta_{\theta_0}f(a)|\varepsilon^2 + \varepsilon^3 + \varepsilon^{n+2} + F_2(\theta(t))^{1/2} + \|v\|^2). \end{aligned}$$

This together with (5.36) proves the claim (i).

To prove claim (ii), we notice that for  $h \in H_u(\mu_0)$

$$\begin{aligned} &\int_M \mathcal{L}_{\theta_0}(\alpha\varphi)h\theta_0 \wedge (d\theta_0)^n \\ &= \int_M (\mathcal{L}_{\theta_0}u)h\theta_0 \wedge (d\theta_0)^n - \int_M (\mathcal{L}_{\theta_0}v)h\theta_0 \wedge (d\theta_0)^n \\ &= \int_M \lambda f u^{1+2/n}h\theta_0 \wedge (d\theta_0)^n + O(F_2(\theta(t))^{1/2} + \|v\|)\|h\| \\ &= \int_M \lambda(f - f(a))u^{1+2/n}h\theta_0 \wedge (d\theta_0)^n + \lambda f(a) \int_M u^{1+2/n}h\theta_0 \wedge (d\theta_0)^n \\ &\quad + O(F_2(\theta(t))^{1/2} + \|v\|)\|h\| \\ &= \int_M \lambda(f - f(a))u^{1+2/n}h\theta_0 \wedge (d\theta_0)^n + \lambda f(a) \int_M \alpha u^{2/n}\varphi h\theta_0 \wedge (d\theta_0)^n \\ &\quad + \lambda f(a) \int_M u^{2/n}vh\theta_0 \wedge (d\theta_0)^n + O(F_2(\theta(t))^{1/2} + \|v\|)\|h\| \\ &= \int_M \lambda(f - f(a))u^{1+2/n}h\theta_0 \wedge (d\theta_0)^n + \lambda f(a) \int_M u^{2/n}vh\theta_0 \wedge (d\theta_0)^n \\ &\quad + O(F_2(\theta(t))^{1/2} + \|v\|)\|h\|. \end{aligned}$$

Using the Hölder's inequality we can estimate

$$\begin{aligned} &\left| \int_M \lambda(f - f(a))u^{1+2/n}h\theta_0 \wedge (d\theta_0)^n \right| \\ &\leq C \int_M |f - f(a)|\varphi^{1+2/n}|h|\theta_0 \wedge (d\theta_0)^n + C \int_M |v|^{1+2/n}|h|\theta_0 \wedge (d\theta_0)^n \\ &\leq C \left( \int_M |f - f(a)|^{\frac{2(n+1)}{n+2}}\varphi^{2+2/n}\theta_0 \wedge (d\theta_0)^n \right)^{\frac{n+2}{2(n+1)}} \|h\| + C\|v\|\|h\| \\ &= O(\varepsilon)\|h\| + O(\|v\|)\|h\| \end{aligned}$$

and

$$\left| \lambda f(a) \int_M u^{2/n}vh\theta_0 \wedge (d\theta_0)^n \right| \leq C \left( \int_M u^{2+2/n}\theta_0 \wedge (d\theta_0)^n \right)^{\frac{1}{n+1}} \|v\|\|h\| \leq C\|v\|\|h\|.$$

Combining all three estimates above yields

$$\int_M \mathcal{L}_{\theta_0}(\alpha\varphi)h\theta_0 \wedge (d\theta_0)^n = O(\mu_0)\|h\|.$$

This together with (5.42) and (5.43) implies that

$$\int_M f(\alpha\varphi)^{1+2/n}h\theta_0 \wedge (d\theta_0)^n = \frac{n\lambda\alpha^{2/n}f(a)}{2(n+1)} \int_M \mathcal{L}_{\theta_0}(\alpha\varphi)h\theta_0 \wedge (d\theta_0)^n + O(\mu_0)\|h\| = O(\mu_0)\|h\|.$$

Substituting this, (5.36) and (5.40) into (5.38) yields

$$\begin{aligned}
& \frac{1}{2} \partial^2 \mathcal{E}_f(\alpha\varphi) h_1 h_2 \\
&= f_{\alpha\varphi}^{-\frac{n}{n+1}} \left[ \int_M (\mathcal{L}_{\theta_0} h_1) h_2 \theta_0 \wedge (d\theta_0)^n - (1 + 2/n) \frac{\mathcal{E}(\alpha\varphi)}{f_{\alpha\varphi}} \int_M f(\alpha\varphi)^{2/n} h_1 h_2 \theta_0 \wedge (d\theta_0)^n \right] \\
&\quad + O(\mu_0) \|h_1\| \|h_2\| \\
&= f_{\alpha\varphi}^{-\frac{n}{n+1}} \left[ \int_M (\mathcal{L}_{\theta_0} h_1) h_2 \theta_0 \wedge (d\theta_0)^n - (1 + 2/n) \int_M \lambda f(\alpha\varphi)^{2/n} h_1 h_2 \theta_0 \wedge (d\theta_0)^n \right] \\
&\quad + O(\mu_0) \|h_1\| \|h_2\| \\
&= f_{\alpha\varphi}^{-\frac{n}{n+1}} \left[ \int_M (\mathcal{L}_{\theta_0} h_1) h_2 \theta_0 \wedge (d\theta_0)^n - (1 + 2/n) \lambda \alpha^{2/n} f(a) \int_M \varphi^{2/n} h_1 h_2 \theta_0 \wedge (d\theta_0)^n \right. \\
&\quad \left. - (1 + 2/n) \int_M \lambda (f - f(a)) (\alpha\varphi)^{2/n} h_1 h_2 \theta_0 \wedge (d\theta_0)^n \right] + O(\mu_0) \|h_1\| \|h_2\| \\
&= f_{\alpha\varphi}^{-\frac{n}{n+1}} \left[ \int_M (\mathcal{L}_{\theta_0} h_1) h_2 \theta_0 \wedge (d\theta_0)^n - \frac{(2n+2)(n+2)}{n^2} \int_M \varphi^{2/n} h_1 h_2 \theta_0 \wedge (d\theta_0)^n \right] \\
&\quad + O(\mu_0) \|h_1\| \|h_2\|,
\end{aligned}$$

which proves the claim (ii), and the proof of the lemma is complete.  $\square$

With the help of Lemma 5.17, we can show that the second variation of the functional  $\mathcal{E}_f$ , when restricted in  $H_u(\mu_0)$ , has a positive lower bound.

**Lemma 5.18.** *There exist constants  $\gamma, t_0 > 0$  such that, for all  $t > t_0$ , there holds*

$$\partial^2 \mathcal{E}_f(\alpha\varphi)|_{H_u(\mu_0)} \geq \gamma.$$

**Proof.** Since the proof is almost the same as that of [21, Proposition 5.5], we provide only some necessary steps. The proof is given by contradiction. So, we assume the contrary. Then it follows from Lemma 5.17 (ii) that there would exist  $\mu_k \searrow 0$  and  $(w_k) \subset H_{u_k}(\mu_k)$  such that

$$1 = \int_M (2+2/n) |\nabla w_k|_{\theta_0}^2 + R_{\theta_0} w_k^2 \theta_0 \wedge (d\theta_0)^n \leq \frac{2(n+1)(n+2)}{n^2} \lim_{k \rightarrow \infty} \int_M \varphi^{2/n} w_k^2 \theta_0 \wedge (d\theta_0)^n.$$

Here,  $\mu_k = \mu_0(t_k)$  and  $u_k = u(t_k)$ . To proceed, we define a sequence of function  $\hat{w}_k : T_{a_k} M \mapsto \mathbb{R}$  by

$$\hat{w}_k = \mu_k^n w_k \left( \exp_{a_k}(\mu_k z, \mu_k^2 s) \right)$$

for  $(z, s) \in T_{a_k} M$ . The sequence  $(\hat{w}_k)$  satisfies

$$\lim_{k \rightarrow \infty} \int_{\{(z,s) \in \mathbb{H}^n : |(z,s)| \leq N_k\}} (2+2/n) |\nabla \hat{w}_k|_{\theta_{\mathbb{H}^n}}^2 \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n \leq 1.$$

and

$$\lim_{k \rightarrow \infty} \int_{\{(z,s) \in \mathbb{H}^n : |(z,s)| \leq N_k\}} |\hat{w}_k|^{2+2/n} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n \leq C.$$

Hence, we can take weak limit to obtain a function  $\hat{w} : \mathbb{H}^n \rightarrow \mathbb{R}$  such that

$$\int_{\mathbb{H}^n} \frac{\hat{w}^2}{s^2 + (1 + |z|^2)^2} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n > 0.$$

In particular,  $\hat{w} \neq 0$ . Using the orthogonality  $w_k \in H_{u_k}(\mu_k)$  and following the proof of [21, Proposition 5.5], we obtain

$$\begin{aligned} \int_{\mathbb{H}^n} \frac{\hat{w}(z, s)}{(s^2 + (1 + |z|^2)^2)^{\frac{n+2}{2}}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n &= 0, \\ \int_{\mathbb{H}^n} \frac{(1 + |z|^2)\bar{z} - \sqrt{-1}\bar{z}s}{(s^2 + (1 + |z|^2)^2)^{\frac{n+4}{2}}} \hat{w}(z, s) \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n &= 0, \\ \int_{\mathbb{H}^n} \frac{1 - |z|^4 - s^2}{(s^2 + (1 + |z|^2)^2)^{\frac{n+4}{2}}} \hat{w}(z, s) \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n &= 0, \\ \int_{\mathbb{H}^n} \frac{(1 + |z|^2)z + \sqrt{-1}zs}{(s^2 + (1 + |z|^2)^2)^{\frac{n+4}{2}}} \hat{w}(z, s) \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n &= 0, \\ \int_{\mathbb{H}^n} \frac{s}{(s^2 + (1 + |z|^2)^2)^{\frac{n+4}{2}}} \hat{w}(z, s) \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n &= 0, \end{aligned}$$

which implies that  $\hat{w} = 0$ , which is a contradiction.  $\square$

Now, we are ready to obtain the apriori estimate for  $v$ .

**Corollary 5.19.** *For  $t$  large, there holds*

$$\|v\| = O(\delta^{4-3n} \varepsilon^n + |\nabla f(a)|_{\theta_0} \varepsilon + |\Delta_{\theta_0} f(a)| \varepsilon^2 + \varepsilon^3 + \varepsilon^{n+2} + F_2(\theta(t))^{1/2}).$$

**Proof.** Since  $v \in H_u(\mu_0)$ , it follows from Lemma 5.18 and Young's inequality that

$$\begin{aligned} \partial \mathcal{E}_f(u)v &= \partial \mathcal{E}_f(\alpha\varphi + v)v \\ &= \partial \mathcal{E}_f(\alpha\varphi)v + \partial^2 \mathcal{E}_f(\alpha\varphi)v^2 + o(\|v\|^2) \\ &\geq \partial \mathcal{E}_f(\alpha\varphi)v + \gamma\|v\|^2 + o(\|v\|^2). \\ &\geq O(\|\partial \mathcal{E}_f(\alpha\varphi)\|_{H_u(\mu)})\|v\|^2 + \frac{\gamma}{2}\|v\|^2 + o(\|v\|^2). \end{aligned}$$

On the other hand, by (2.8), (5.39) and Hölder's inequality, we can estimate

$$\begin{aligned} \left| \frac{1}{2} \partial \mathcal{E}_f(u)v \right| &= \left| \int_M (\mathcal{L}_{\theta_0} u)v \theta_0 \wedge (d\theta_0)^n - \frac{\mathcal{E}(u)}{f_u} \int_M f u^{1+2/n} v \theta_0 \wedge (d\theta_0)^n \right| \\ &= \left| \int_M (\mathcal{L}_{\theta_0} u)v \theta_0 \wedge (d\theta_0)^n - (\lambda + O(F_2(\theta(t))^{1/2})) \int_M f u^{1+2/n} v \theta_0 \wedge (d\theta_0)^n \right| \\ &\leq \left| \int_M (R_{\theta(t)} - \lambda f) u^{1+2/n} v \theta_0 \wedge (d\theta_0)^n \right| + O(F_2(\theta(t))^{1/2} \|v\|) \\ &= O(F_2(\theta(t))^{1/2} \|v\|) = O(F_2(\theta(t))^{1/2}) + o(\|v\|^2). \end{aligned}$$

Combining the two estimates above and using Lemma 5.17 (i), we thus prove the assertion.  $\square$

With the help of the previous results, we may simplify the shadow flow as follows.

**Corollary 5.20.** *For  $t$  large, there hold*

$$\begin{aligned} \text{(i)} \quad \frac{\dot{\varepsilon}}{\varepsilon} &= \lambda f(a) \left[ \frac{nd_2}{2(n+1)c_2} A_a \varepsilon^{2n} + \frac{e_2}{c_2} \frac{\Delta_{\theta_0} f(a)}{f(a)} \varepsilon^2 \right] (1 + o(\varepsilon)) + R_2, \\ \text{(ii)} \quad \frac{\dot{a}}{\varepsilon} &= \lambda f(a) \left[ \frac{e_3}{c_3} \frac{\nabla_{\theta_0} f(a)}{f(a)} \varepsilon + \frac{e_4}{c_3} \frac{\nabla_{\theta_0} \Delta_{\theta_0} f(a)}{f(a)} \varepsilon^3 \right] (1 + o(\varepsilon)) + R_3, \end{aligned}$$

where

$$R_2, R_3 = O\left(\delta^{8-6n} \varepsilon^{2n} + |\nabla f(a)|_{\theta_0}^2 \varepsilon^2 + |\Delta_{\theta_0} f(a)|^2 \varepsilon^4 + \varepsilon^6 + \varepsilon^{2n+4} + F_2(\theta(t))\right).$$

**Proof.** It follows from Lemma 5.14, Corollary 5.16, Corollary 5.19 and (5.36).  $\square$

## 6. PROOF OF THEOREMS 1.1 AND 1.2

In this section, we finish the proof of Theorems 1.1 and 1.2.

**Step 1.** We aim to construct an initial data  $u_0$  satisfying (5.14). Given a sufficiently small  $\varepsilon_0 > 0$ , we let

$$u_0 = \alpha_0 \varphi_{a_0, \varepsilon_0}.$$

where  $a_0 \in M$  is chosen to be such that

$$d\left(a_0, \left\{x \in M : f(x) = \sup_M f\right\}\right) < \varepsilon_0. \quad (6.1)$$

Here,  $d$  is the Carnot-Carathéodory distance with respect to  $\theta_0$ , and  $\varphi_{a_0, \varepsilon_0}$  is defined as in (5.20), and  $\alpha_0$  is chosen such that

$$\int_M f u_0^{2+2/n} \theta_0 \wedge (d\theta_0)^n = 1$$

i.e.,

$$\alpha_0^{-(2+2/n)} = \int_M f \varphi_{a_0, \varepsilon_0}^{2+2/n} \theta_0 \wedge (d\theta_0)^n. \quad (6.2)$$

We claim that such  $u_0$  satisfies (5.14). To see this, we note by (6.1) that

$$\begin{aligned} \int_M f \varphi_{a_0, \varepsilon_0}^{2+2/n} \theta_0 \wedge (d\theta_0)^n &= \sup_M f \int_M \varphi_{a_0, \varepsilon_0}^{2+2/n} \theta_0 \wedge (d\theta_0)^n + O(\varepsilon_0) \\ &= \sup_M f \int_{\mathbb{H}^n} \left( \frac{1}{(s^2 + ((1/n)^2 + |z|^2)^{n/2})} \right)^{2+2/n} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon_0) \\ &= \left( \frac{n\mu(S^{2n+1})}{2(n+1)} \right)^{n+1} \sup_M f + O(\varepsilon_0). \end{aligned}$$

Substituting the above into (6.2) yields

$$\alpha_0^2 = \left( \frac{n\mu(S^{2n+1})}{2(n+1)} \right)^{-n} \left( \sup_M f \right)^{-\frac{n}{n+1}} + O(\varepsilon_0).$$

This together with Lemma 5.9 implies that

$$\begin{aligned} \mathcal{E}(u_0) &= \frac{2(n+1)}{n} \alpha_0^2 \int_M \varphi_{a_0, \varepsilon_0}^{2+2/n} \theta_0 \wedge (d\theta_0)^n + O(\varepsilon_0) \\ &= \frac{2(n+1)}{n} \alpha_0^2 \left( \frac{n\mu(S^{2n+1})}{2(n+1)} \right)^{n+1} + O(\varepsilon_0) \\ &= \mu(S^{2n+1}) \left( \sup_M f \right)^{-\frac{n}{n+1}} + O(\varepsilon_0) \\ &= \frac{2(n+1)}{n} \left( \sup_M f \right)^{-\frac{n}{n+1}} K_n^{-1} + O(\varepsilon_0) \\ &= \lambda_* + O(\varepsilon_0). \end{aligned}$$

Hence, by choosing  $\varepsilon_0$  sufficiently small, we clearly have  $\mathcal{E}(u_0) < 2^{\frac{1}{n+1}} \lambda_*$ . This proves the claim.

**Step 2.** In view of Lemma 5.5, we assume by contradiction that  $u(t) \rightarrow 0$  weakly in  $S_1^2(M)$  as  $t \rightarrow +\infty$ . Then, from the choice of  $u_0$  in **Step 1** and the arguments in



Subsection 5.3, we conclude that there is a unique concentration point  $a_\infty \in M$  with  $a(t) \rightarrow a_\infty$  as  $t \rightarrow +\infty$ . We claim that  $a_\infty \in \left\{x \in M : f(x) = \sup_M f\right\}$ . Otherwise, it follows from Lemma 5.2 and (5.15) that

$$\mathcal{E}(u(t)) = \frac{2(n+1)}{n} K_n^{-1} [f(a_\infty)]^{-\frac{n}{n+1}} + o(1).$$

Hence, by choosing  $\varepsilon_0$  suitably small and for  $t$  large, we have

$$\mathcal{E}(u(t)) > \lambda_* + O(\varepsilon_0) = \mathcal{E}(u_0),$$

which contradicts that the energy  $\mathcal{E}(u)$  is non-increasing by Lemma 3.1.

**Step 3.** We are now ready to derive the contradiction. To this end, we define

$$\zeta(t) = \frac{\ln \varepsilon(t)}{f(a(t))}.$$

Then, it is easy to see that  $\zeta \rightarrow -\infty$ , as  $t \rightarrow +\infty$ . On the other hand, when  $n = 1$ , it follows from Corollary 5.20 that

$$\begin{aligned} \zeta'(t) &= \frac{1}{f(a)} \left( \frac{\dot{\varepsilon}}{\varepsilon} - \varepsilon \ln \varepsilon \frac{\nabla_{\theta_0} f(a)}{f(a)} \cdot \frac{\dot{a}}{\varepsilon} \right) \\ &\geq \gamma \varepsilon^2 \left( A_a + \frac{\Delta_{\theta_0} f(a)}{f(a)} + \frac{|\nabla f(a)|_{\theta_0}^2}{f^2(a)} \ln(1/\varepsilon) \right) (1 + o(\varepsilon)) + O(F_2(\theta(t))), \end{aligned}$$

for some positive constant  $\gamma$ . Notice that we have showed in **Step 2** that  $a(t) \rightarrow a_\infty \in \left\{x \in M : f(x) = \sup_M f\right\}$  as  $t \rightarrow +\infty$ . Our argument is now split into two cases.

**Case  $\mu(M) > 0$ .** If we take  $c_* = \inf_{x \in M} A_x^*$ , then it follows from the assumptions in Theorem 1.1 that  $c_* > 0$ . With this  $c_*$ , it follows from assumption (1.3) that  $\Delta_{\theta_0} f(a_\infty)/f(a_\infty) > -\inf_{x \in M} A_x^*$ . Hence, for  $t$  sufficiently large, it follows that  $\Delta_{\theta_0} f(a)/f(a) > -\inf_{x \in M} A_x^*$ . This together with the fact that  $A_a = A_a^*$  implies that with a new constant  $\gamma_1$  there holds

$$\zeta'(t) \geq \gamma_1 \varepsilon^2 \left( c_* + \frac{|\nabla f(a)|_{\theta_0}^2}{f^2(a)} \ln(1/\varepsilon) \right) (1 + o(\varepsilon)) + O(F_2(\theta(t))) \geq O(F_2(\theta(t))) \quad (6.3)$$

for  $t$  sufficiently large.

**Case  $\mu(M) = 0$ .** Notice that

$$\left| \lim_{t \rightarrow +\infty} \frac{\Delta_{\theta_0} f(a)}{f(a)} \right| = \left| \frac{\Delta_{\theta_0} f(a_\infty)}{\sup_M f} \right| < \frac{|\sup_M \Delta_{\theta_0} f| + 1}{\sup_M f}.$$

In this case, recall that  $A_a = \Lambda$  is a positive constant to be determined. Hence, we may choose

$$\Lambda = \frac{|\sup_M \Delta_{\theta_0} f| + 1}{\sup_M f},$$

and we can obtain (6.3) in the same way.

By (3.1),  $F_2(\theta(t))$  is integrable on  $(0, +\infty)$ . Therefore, we can integrate (6.3) and conclude that  $\zeta(t)$  is bounded from below, which contradicts to the fact that  $\zeta(t) \rightarrow -\infty$  as  $t \rightarrow +\infty$ . The proof of the main theorems is thus complete.

## APPENDIX A. PROOF OF LEMMA 5.12

In this Appendix, we prove Lemma 5.12. It follows from Definition 5.11 that  $\phi_1 = \varphi$ , and a straightforward computation yields

$$\begin{aligned}
\phi_2 &= d_2\varphi = \varepsilon \partial_\varepsilon \varphi \\
&= \varepsilon \partial_\varepsilon \left\{ \omega_a \varepsilon^n \left[ \frac{\chi_\delta(\rho_a)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2})} + (1 - \chi_\delta(\rho_a))G_a \right] \right\} \\
&= n\varphi + \omega_a \varepsilon^{n+1} \chi_\delta(\rho_a) \partial_\varepsilon \left( s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2} \right)^{-n/2} \\
&= n\varphi - \frac{2}{n} \omega_a \varepsilon^{n+2} \chi_\delta(\rho_a) \left( (\varepsilon/n)^2 + |z|^2 \right) \left( s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2} \right)^{-n/2-1} \quad (\text{A.1})
\end{aligned}$$

and

$$\begin{aligned}
\phi_3 &= d_3\varphi = \varepsilon \nabla_a \varphi|_{a=0} \quad (a = (z_0, s_0)) \\
&= \varepsilon \nabla_a \left\{ \omega_a \varepsilon^n \left[ \frac{\chi_\delta(\rho_a)}{((s - s_0)^2 + ((\varepsilon/n)^2 + |z - z_0|^2)^{n/2})} + (1 - \chi_\delta(\rho_a))G_a \right] \right\} \Big|_{a=0} \\
&= \frac{\varepsilon \nabla_a \omega_a}{\omega_a} \varphi + \omega_a \varepsilon^{n+1} \left[ \nabla_a (\chi_\delta(\rho_a)) \left( (s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2} \right)^{-n/2} - G_a \right) \right. \\
&\quad \left. + (1 - \chi_\delta(\rho_a)) \nabla_a G_a + n \chi_\delta(\rho_a) \frac{((\varepsilon/n)^2 + |z|^2) \bar{z}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2+1})} \right]. \quad (\text{A.2})
\end{aligned}$$

(i) **Case**  $k = 1$ . It is trivial for  $l = 1$ , hence we consider  $l = 2, 3$ . Since  $d_1\phi_l = \phi_l$ , we have by (A.1) that

$$\begin{aligned}
|d_1\phi_2| &\leq n\varphi + \frac{2}{n} \omega_a \varepsilon^n \chi_\delta(\rho_a) \left( s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2} \right)^{-n/2} \\
&\leq n\varphi + \frac{2}{n} \varphi \leq C\varphi.
\end{aligned}$$

This gives the estimate for  $l = 2$ . As for  $l = 3$ , we first notice the facts that

$$\nabla_a (\chi_\delta(\rho_a)) = 0 \quad \text{for } \rho_a < \delta \text{ or } \rho_a > 2\delta;$$

and for  $\delta \leq \rho_a \leq 2\delta$ , there holds

$$\left| \nabla_a (\chi_\delta(\rho_a)) \left( (s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2} \right)^{-n/2} - G_a \right) \right| \leq C(\delta)$$

and

$$\chi_\delta(\rho_a) \left( s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2} \right)^{-n/2} + (1 - \chi_\delta(\rho_a))G_a \geq C(\delta).$$

Hence

$$\left| \omega_a \varepsilon^{n+1} \nabla_a (\chi_\delta(\rho_a)) \left( (s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2} \right)^{-n/2} - G_a \right) \right| \leq C(\delta)\varphi.$$

Moreover,  $|\nabla_a G_a / G_a| \leq C(\delta)$  for  $\rho_a \geq \delta$  and

$$\left| \frac{((\varepsilon/n)^2 + |z|^2) \varepsilon \bar{z}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2+1})} \right| \leq \frac{1}{2} \left( s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2} \right)^{-n/2}.$$

This implies that

$$\left| \omega_a \varepsilon^n \left( \varepsilon (1 - \chi_\delta(\rho_a)) \nabla_a G_a + n \chi_\delta(\rho_a) \frac{((\varepsilon/n)^2 + |z|^2) \varepsilon \bar{z}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^{n/2+1})} \right) \right|$$

$$\begin{aligned} &\leq C(\delta)\omega_a\varepsilon^n\left((1-\chi_\delta(\rho_a))G_a+\chi_\delta(\rho_a)(s^2+((\varepsilon/n)^2+|z|^2)^2)^{-n/2}\right) \\ &= C(\delta)\varphi. \end{aligned}$$

It is easy to see that  $|\varepsilon(\nabla_a\omega_a/\omega_a)\varphi| \leq C(\delta)\varphi$ . Substituting all the estimates above into (A.2), we arrive at

$$|d_1\phi_3| \leq C(\delta)\varphi.$$

**Case**  $k = 2$ . Since  $d_2\phi_1 = d_2\varphi = \phi_2$ , we are left to do estimates for  $l = 2, 3$ . From (A.1) it follows that

$$\begin{aligned} d_2\phi_2 &= d_2\left(n\varphi - \frac{2}{n}\omega_a\varepsilon^{n+2}\chi_\delta(\rho_a)\left((\varepsilon/n)^2+|z|^2\right)\left(s^2+((\varepsilon/n)^2+|z|^2)^2\right)^{-n/2-1}\right) \\ &= n\phi_2 - 2n(n+2)\omega_a\varepsilon^n(\varepsilon/n)^2\chi_\delta(\rho_a)\left((\varepsilon/n)^2+|z|^2\right)\left(s^2+((\varepsilon/n)^2+|z|^2)^2\right)^{-n/2-1} \\ &\quad - 4n\omega_a\varepsilon^n(\varepsilon/n)^4\chi_\delta(\rho_a)\left(s^2+((\varepsilon/n)^2+|z|^2)^2\right)^{-n/2-1} \\ &\quad + 4n(n+2)\omega_a\varepsilon^n(\varepsilon/n)^4\chi_\delta(\rho_a)\left((\varepsilon/n)^2+|z|^2\right)^2\left(s^2+((\varepsilon/n)^2+|z|^2)^2\right)^{-n/2-2}. \end{aligned} \tag{A.3}$$

Therefore

$$\begin{aligned} |d_2\phi_2| &\leq n|\phi_2| + 2n(n+2)\omega_a\varepsilon^n\chi_\delta(\rho_a)\left(s^2+((\varepsilon/n)^2+|z|^2)^2\right)^{-n/2} \\ &\quad + 4n\omega_a\varepsilon^n\chi_\delta(\rho_a)\left(s^2+((\varepsilon/n)^2+|z|^2)^2\right)^{-n/2} \\ &\quad + 4n(n+2)\omega_a\varepsilon^n\chi_\delta(\rho_a)\left(s^2+((\varepsilon/n)^2+|z|^2)^2\right)^{-n/2} \\ &\leq C\varphi. \end{aligned}$$

Now, by (A.2) we can compute

$$\begin{aligned} d_2\phi_3 &= d_2\left\{\frac{\varepsilon\nabla_a\omega_a}{\omega_a}\varphi + \omega_a\varepsilon^{n+1}\left[\nabla_a(\chi_\delta(\rho_a))\left((s^2+((\varepsilon/n)^2+|z|^2)^2\right)^{-n/2} - G_a\right) \right. \right. \\ &\quad \left. \left. + (1-\chi_\delta(\rho_a))\nabla_aG_a + n\chi_\delta(\rho_a)\frac{((\varepsilon/n)^2+|z|^2)\bar{z}}{(s^2+((\varepsilon/n)^2+|z|^2)^2)^{n/2+1}}\right]\right\} \\ &= -n\frac{\varepsilon\nabla_a\omega_a}{\omega_a}\varphi + \frac{\varepsilon\nabla_a\omega_a}{\omega_a}\phi_2 + (n+1)\phi_3 - 2n\omega_a\varepsilon^{n+1}\nabla_a(\chi_\delta(\rho_a))(\varepsilon/n)^2\left((\varepsilon/n)^2+|z|^2\right) \\ &\quad \times\left(s^2+((\varepsilon/n)^2+|z|^2)^2\right)^{-n/2-1} + n^2\omega_a\varepsilon^n\chi_\delta(\rho_a)\left(\frac{2(\varepsilon/n)^3\bar{z}}{(s^2+((\varepsilon/n)^2+|z|^2)^2)^{n/2+1}} \right. \\ &\quad \left. - (n+2)\frac{2(\varepsilon/n)^3\bar{z}((\varepsilon/n)^2+|z|^2)^2}{(s^2+((\varepsilon/n)^2+|z|^2)^2)^{n/2+2}}\right). \end{aligned} \tag{A.4}$$

So, by using the conclusions in **Case**  $k = 1$ , we may estimate in a similar way as we did for  $\phi_3$ :

$$\begin{aligned} |d_2\phi_3| &\leq C\varphi + 2n\omega_a\varepsilon^{n+1}\left|\nabla_a(\chi_\delta(\rho_a))\right|\left(s^2+((\varepsilon/n)^2+|z|^2)^2\right)^{-n/2} \\ &\quad + n^2(n+3)\omega_a\varepsilon^n\chi_\delta(\rho_a)\left(s^2+((\varepsilon/n)^2+|z|^2)^2\right)^{-n/2} \\ &\leq C\varphi. \end{aligned}$$

**Case**  $k = 3$ . Since  $d_3\phi_1 = d_3\varphi = \phi_3$ , it suffices to consider  $l = 2, 3$ . In view of (A.1) we have

$$\begin{aligned} d_3\phi_2 &= d_3\left(n\varphi - \frac{2}{n}\omega_a\varepsilon^{n+2}\chi_\delta(\rho_a)\left((\varepsilon/n)^2 + |z - z_0|^2\right)\left((s - s_0)^2 + ((\varepsilon/n)^2 + |z - z_0|^2)^2\right)^{-n/2-1}\right)\Big|_{a=0} \\ &= n\phi_3 - 2n^2\nabla_a\left(\omega_a\chi_\delta(\rho_a)\right)\varepsilon^n(\varepsilon/n)^3\left((\varepsilon/n)^2 + |z|^2\right)\left(s^2 + ((\varepsilon/n)^2 + |z|^2)^2\right)^{-n/2-1} \\ &\quad - 2n^2\omega_a\chi_\delta(\rho_a)\varepsilon^n(\varepsilon/n)^3\bar{z}\left(s^2 + ((\varepsilon/n)^2 + |z|^2)^2\right)^{-n/2-1} \\ &\quad - 2n^2(n+2)\omega_a\chi_\delta(\rho_a)\varepsilon^n(\varepsilon/n)^3\bar{z}\left((\varepsilon/n)^2 + |z|^2\right)^2\left(s^2 + ((\varepsilon/n)^2 + |z|^2)^2\right)^{-n/2-2}. \end{aligned} \quad (\text{A.5})$$

This yields that

$$\begin{aligned} |d_3\phi_2| &\leq C\varphi + 2n^2\omega_a\varepsilon^n\left(\left|\frac{\varepsilon\nabla_a(\omega_a\chi_\delta(\rho_a))}{n\omega_a}\right| + (n+3)\chi_\delta(\rho_a)\right)\left(s^2 + ((\varepsilon/n)^2 + |z|^2)^2\right)^{-n/2} \\ &\leq C\varphi. \end{aligned}$$

To finish the proof of (i), it remains to estimate  $d_3\phi_3$ . From (A.2) we obtain

$$\begin{aligned} d_3\phi_3 &= d_3\left\{\frac{\varepsilon\nabla_a\omega_a}{\omega_a}\varphi + \omega_a\varepsilon^{n+1}\left[\nabla_a\left(\chi_\delta(\rho_a)\right)\left(\left((s - s_0)^2 + ((\varepsilon/n)^2 + |z - z_0|^2)^2\right)^{-n/2} - G_a\right) \right. \right. \\ &\quad \left. \left. + (1 - \chi_\delta(\rho_a))\nabla_a G_a + n\chi_\delta(\rho_a)\frac{((\varepsilon/n)^2 + |z - z_0|^2)(\overline{z - z_0})}{((s - s_0)^2 + ((\varepsilon/n)^2 + |z - z_0|^2)^2)^{n/2+1}}\right]\right\}\Big|_{a=0} \\ &= \varepsilon^2\left[\nabla_a^2(\ln\omega_a) - (\nabla_a(\ln\omega_a))^2\right]\varphi + (\varepsilon^2 + \varepsilon)\nabla_a(\ln\omega_a)\phi_3 \\ &\quad + \omega_a\varepsilon^{n+2}\left[\nabla_a^2\left(\chi_\delta(\rho_a)\right)\left(\left(s^2 + ((\varepsilon/n)^2 + |z|^2)^2\right)^{-n/2} - G_a\right) + \nabla_a\left(\chi_\delta(\rho_a)\right) \right. \\ &\quad \times \left(2n\bar{z}\left((\varepsilon/n)^2 + |z|^2\right)\left(s^2 + ((\varepsilon/n)^2 + |z|^2)^2\right)^{-n/2-1} - 2\nabla_a G_a\right) \\ &\quad \left. + (1 - \chi_\delta(\rho_a))\nabla_a^2 G_a + n\chi_\delta(\rho_a)\left(-\bar{z}^2\left(s^2 + ((\varepsilon/n)^2 + |z|^2)^2\right)^{-n/2-1} \right. \right. \\ &\quad \left. \left. + (n+2)\bar{z}^2\left((\varepsilon/n)^2 + |z|^2\right)^2\left(s^2 + ((\varepsilon/n)^2 + |z|^2)^2\right)^{-n/2-2}\right)\right] \end{aligned} \quad (\text{A.6})$$

A similar argument as before implies that

$$|d_3\phi_3| \leq C\varphi.$$

(ii) **Case**  $k = 1$ . Since  $\phi_1 = \varphi$ , we have

$$\begin{aligned} \int_M \varphi^{2/n}\phi_1^2\theta_0 \wedge (d\theta_0)^n &= \int_M \varphi^{2+2/n}\theta_0 \wedge (d\theta_0)^n \\ &= \int_M \varepsilon^{2n+2}\left[\frac{\chi_\delta(\rho_a)}{\left(s^2 + ((\varepsilon/n)^2 + |z|^2)^2\right)^{n/2}} + (1 - \chi_\delta(\rho_a))G_a\right]^{2+2/n}\theta_a \wedge (d\theta_a)^n \\ &= \int_{B_\delta(a)} \frac{\varepsilon^{2n+2}}{\left(s^2 + ((\varepsilon/n)^2 + |z|^2)^2\right)^{n+1}}\theta_a \wedge (d\theta_a)^n + O(\varepsilon^{2n+2}) \\ &= \int_{B_{\frac{n\delta}{\varepsilon}}(0)} \frac{n^{2n+2}}{\left(s^2 + (1 + |z|^2)^2\right)^{n+1}}\theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+2}) \end{aligned}$$

$$\begin{aligned}
 &= \int_{\mathbb{H}^n} \frac{n^{2n+2}}{(s^2 + (1 + |z|^2)^2)^{n+1}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+2}) \\
 &:= c_1 + O(\varepsilon^{2n+2}).
 \end{aligned}$$

**Case  $k = 2$ .** By (A.1), we can obtain

$$\begin{aligned}
 \int_M \varphi^{2/n} \phi_2^2 \theta_0 \wedge (d\theta_0)^n &= n^2 \int_M \varphi^{2+2/n} \theta_0 \wedge (d\theta_0)^n \\
 &\quad - 4 \int_M \varphi^{1+2/n} \frac{\omega_a \varepsilon^{n+2} \chi_\delta(\rho_a) ((\varepsilon/n)^2 + |z|^2)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \theta_0 \wedge (d\theta_0)^n \\
 &\quad + \frac{4}{n^2} \int_M \varphi^{2/n} \frac{\omega_a^2 \varepsilon^{2n+4} \chi_\delta^2(\rho_a) ((\varepsilon/n)^2 + |z|^2)^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_0 \wedge (d\theta_0)^n \\
 &:= I - II + III.
 \end{aligned}$$

By the estimate in Case  $k = 1$ , one has

$$I = n^{2n+4} \int_{\mathbb{H}^n} \frac{1}{(s^2 + (1 + |z|^2)^2)^{n+1}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+2}).$$

Now, we estimate  $II$  as follows.

$$\begin{aligned}
 II &= 4 \int_{B_\delta(a)} \frac{\varepsilon^{2n+4} ((\varepsilon/n)^2 + |z|^2)}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_a \wedge (d\theta_a)^n + O(\varepsilon^{2n+4}) \\
 &= 4n^{2n+4} \int_{B_{\frac{\delta}{\varepsilon}}(0)} \frac{(1 + |z|^2)}{(s^2 + (1 + |z|^2)^2)^{n+2}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+4}) \\
 &= 4n^{2n+4} \int_{\mathbb{H}^n} \frac{(1 + |z|^2)}{(s^2 + (1 + |z|^2)^2)^{n+2}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+4})
 \end{aligned}$$

Similarly, we may estimate  $III$  to obtain

$$\begin{aligned}
 III &= \frac{4}{n^2} \int_{B_\delta(a)} \frac{\varepsilon^{2n+6} ((\varepsilon/n)^2 + |z|^2)^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+3}} \theta_a \wedge (d\theta_a)^n + O(\varepsilon^{2n+6}) \\
 &= 4n^{2n+4} \int_{B_{\frac{\delta}{\varepsilon}}(0)} \frac{(1 + |z|^2)^2}{(s^2 + (1 + |z|^2)^2)^{n+3}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+6}) \\
 &= 4n^{2n+4} \int_{\mathbb{H}^n} \frac{(1 + |z|^2)^2}{(s^2 + (1 + |z|^2)^2)^{n+3}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+6}).
 \end{aligned}$$

By setting

$$\begin{aligned}
 c_2 &= n^{2n+4} \left[ \int_{\mathbb{H}^n} \frac{1}{(s^2 + (1 + |z|^2)^2)^{n+1}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n \right. \\
 &\quad - 4 \int_{\mathbb{H}^n} \frac{(1 + |z|^2)}{(s^2 + (1 + |z|^2)^2)^{n+2}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n \\
 &\quad \left. + 4 \int_{\mathbb{H}^n} \frac{(1 + |z|^2)^2}{(s^2 + (1 + |z|^2)^2)^{n+3}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n \right] \\
 &= \int_{\mathbb{H}^n} \frac{n^{2n+4}}{(s^2 + (1 + |z|^2)^2)^{n+3}} \left[ (s^2 + (1 + |z|^2)^2)^2 \right.
 \end{aligned}$$

$$\begin{aligned}
& -4(1 + |z|^2)(s^2 + (1 + |z|^2)^2) + 4(1 + |z|^2)^2 \Big] \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n \\
& = \int_{\mathbb{H}^n} \frac{n^{2n+4}(s^2 + |z|^4 - 1)^2}{(s^2 + (1 + |z|^2)^2)^{n+3}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n,
\end{aligned}$$

we thus arrive at

$$\int_M \varphi^{2/n} \phi_2^2 \theta_0 \wedge (d\theta_0)^n = c_2 + O(\varepsilon^{2n+2}).$$

**Case  $k = 3$ .** It follows from (A.2) that

$$\begin{aligned}
& \int_M \varphi^{2/n} \phi_3^2 \theta_0 \wedge (d\theta_0)^n = \int_M \varphi^{2/n} \left\{ \frac{\varepsilon \nabla_a \omega_a}{\omega_a} \varphi + \omega_a \varepsilon^{n+1} \left[ \nabla_a (\chi_\delta(\rho_a)) \right. \right. \\
& \quad \left. \left. \left( (s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{-n/2} - G_a \right) + (1 - \chi_\delta(\rho_a)) \nabla_a G_a \right. \right. \\
& \quad \left. \left. + n \chi_\delta(\rho_a) \frac{((\varepsilon/n)^2 + |z|^2) \bar{z}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right] \right\}^2 \theta_0 \wedge (d\theta_0)^n \\
& = \varepsilon^2 \int_M \frac{|\nabla_a \omega_a|^2}{\omega_a^2} \varphi^{2+2/n} \theta_0 \wedge (d\theta_0)^n + 2 \int_M \nabla_a \omega_a \varphi^{1+2/n} \cdot \varepsilon^{n+2} \left[ \nabla_a (\chi_\delta(\rho_a)) \right. \\
& \quad \left. \left( (s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{-n/2} - G_a \right) + (1 - \chi_\delta(\rho_a)) \nabla_a G_a \right. \\
& \quad \left. + n \chi_\delta(\rho_a) \frac{((\varepsilon/n)^2 + |z|^2) \bar{z}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right] \theta_0 \wedge (d\theta_0)^n + \int_M \varphi^{2/n} \omega_a^2 \varepsilon^{2n+2} \\
& \quad \left[ \nabla_a (\chi_\delta(\rho_a)) \left( (s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{-n/2} - G_a \right) + (1 - \chi_\delta(\rho_a)) \nabla_a G_a \right. \\
& \quad \left. + n \chi_\delta(\rho_a) \frac{((\varepsilon/n)^2 + |z|^2) \bar{z}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right]^2 \theta_0 \wedge (d\theta_0)^n \\
& := I + II + III.
\end{aligned}$$

*Estimate of I.* Notice that  $\omega_a = 1 + O(\rho_a^4)$  and  $|\nabla_a \omega_a| = O(\rho_a^3)$ . Hence

$$\begin{aligned}
|I| & = \varepsilon^2 \int_{B_\delta(a)} \frac{|\nabla_a \omega_a|^2}{\omega_a^2} \varphi^{2+2/n} \theta_0 \wedge (d\theta_0)^n + O(\varepsilon^{2n+4}) \\
& = \varepsilon^2 O \left( \int_{B_\delta(a)} \rho_a^6 \varphi^{2+2/n} \theta_0 \wedge (d\theta_0)^n \right) + O(\varepsilon^{2n+4}) \\
& = \varepsilon^2 O \left( \int_{B_\delta(0)} \frac{\varepsilon^{2n+2} (s^2 + |z|^4)^{3/2}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+1}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n \right) + O(\varepsilon^{2n+4}) \\
& = \varepsilon^8 O \left( \int_{B_{\frac{\delta}{\varepsilon}}(0)} \frac{(s^2 + |z|^4)^{3/2}}{(s^2 + (1 + |z|^2)^2)^{n+1}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n \right) + O(\varepsilon^{2n+4}) \\
& = O(\varepsilon^{\min(8, 2n+4)}).
\end{aligned}$$

*Estimate of II.* We have

$$|II| \leq \int_{B_\delta(a)} 2n \frac{|\nabla_a \omega_a|}{\omega_a} \varepsilon^{2n+4} \frac{((\varepsilon/n)^2 + |z|^2) |z|}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_a \wedge (d\theta_a)^n + O(\varepsilon^{2n+4})$$

$$\begin{aligned}
 &= O\left(\int_{B_\delta(a)} \rho_a^3 \omega_a \varepsilon^{2n+4} \frac{((\varepsilon/n)^2 + |z|^2)|z|}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_a \wedge (d\theta_a)^n\right) + O(\varepsilon^{2n+4}) \\
 &= O\left(\int_{B_\delta(0)} \varepsilon^{2n+4} \frac{(s^2 + |z|^2)^{\frac{3}{4}} ((\varepsilon/n)^2 + |z|^2)|z|}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_a \wedge (d\theta_a)^n\right) + O(\varepsilon^{2n+4}) \\
 &= O\left(\varepsilon^4 \int_{B_{\frac{\delta}{\varepsilon}}(0)} \frac{(s^2 + |z|^2)^{\frac{3}{4}} (1 + |z|^2)|z|}{(s^2 + (1 + |z|^2)^2)^{n+2}} \theta_a \wedge (d\theta_a)^n\right) + O(\varepsilon^{2n+4}) \\
 &= O(\varepsilon^4).
 \end{aligned}$$

*Estimate of III.* We estimate

$$\begin{aligned}
 III &= \int_{B_\delta(a)} n^2 \varepsilon^{2n+4} \frac{((\varepsilon/n)^2 + |z|^2)^2 |z|^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+3}} \theta_a \wedge (d\theta_a)^n + O(\varepsilon^{2n+4}) \\
 &= \int_{B_{\frac{\delta}{\varepsilon}}(0)} \frac{n^{2n+6} (1 + |z|^2)^2 |z|^2}{(s^2 + (1 + |z|^2)^2)^{n+3}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+4}) \\
 &= \int_{\mathbb{H}^n} \frac{n^{2n+6} (1 + |z|^2)^2 |z|^2}{(s^2 + (1 + |z|^2)^2)^{n+3}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n + O(\varepsilon^{2n+4}) \\
 &:= c_3 + O(\varepsilon^{2n+4}).
 \end{aligned}$$

Hence

$$\int_M \varphi^{2/n} \phi_3^2 \theta_0 \wedge (d\theta_0)^n = c_3 + O(\varepsilon^{2n+2} + \varepsilon^4).$$

(iii) By Definition 5.11 and (ii) above, we obtain

$$\int_M \varphi^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n = d_k \int_M \varphi^{2+2/n} \theta_0 \wedge (d\theta_0)^n = O(\varepsilon^{2n+2}).$$

(iv) Notice that if  $k = 1$  or  $l = 1$ , we have by (iii) above that

$$\int_M \varphi^{2/n} \phi_k \phi_l \theta_0 \wedge (d\theta_0)^n = \int_M \varphi^{1+2/n} \phi_k \theta_0 \wedge (d\theta_0)^n = O(\varepsilon^{2n+2}).$$

It remains to estimate  $\int_M \varphi^{2/n} \phi_2 \phi_3 \theta_0 \wedge (d\theta_0)^n$ . From (A.1) and (A.2) and symmetry, it follows that

$$\begin{aligned}
 \int_M \varphi^{2/n} \phi_2 \phi_3 \theta_0 \wedge (d\theta_0)^n &= \int_M \varphi^{2/n} \left( n\varphi - \frac{2}{n} \omega_a \varepsilon^{n+2} \chi_\delta(\rho_a) \frac{(\varepsilon/n)^2 + |z|^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right) \\
 &\quad \times \left\{ \frac{\varepsilon \nabla_a \omega_a}{\omega_a} \varphi + \omega_a \varepsilon^{n+1} \left[ \nabla_a (\chi_\delta(\rho_a)) \left( (s^2 + ((\varepsilon/n)^2 + |z|^2)^2 \right)^{-n/2} - G_a \right) \right. \right. \\
 &\quad \left. \left. + (1 - \chi_\delta(\rho_a)) \nabla_a G_a + n \chi_\delta(\rho_a) \frac{((\varepsilon/n)^2 + |z|^2) \bar{z}}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right] \right\} \theta_0 \wedge (d\theta_0)^n \\
 &= \int_M \varphi^{2/n} \left( n\varphi - \frac{2}{n} \omega_a \varepsilon^{n+2} \chi_\delta(\rho_a) \frac{(\varepsilon/n)^2 + |z|^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right) \\
 &\quad \times \left\{ \frac{\varepsilon \nabla_a \omega_a}{\omega_a} \varphi + \omega_a \varepsilon^{n+1} \left[ \nabla_a (\chi_\delta(\rho_a)) \left( (s^2 + ((\varepsilon/n)^2 + |z|^2)^2 \right)^{-n/2} - G_a \right) \right. \right. \\
 &\quad \left. \left. + (1 - \chi_\delta(\rho_a)) \nabla_a G_a \right] \right\} \theta_0 \wedge (d\theta_0)^n \\
 &= \int_{B_\delta(a)} \frac{\varepsilon \nabla_a \omega_a}{\omega_a} \varphi^{1+2/n} \left( n\varphi - \frac{2}{n} \omega_a \varepsilon^{n+2} \frac{(\varepsilon/n)^2 + |z|^2}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n/2+1}} \right) \theta_0 \wedge (d\theta_0)^n
 \end{aligned}$$

$$\begin{aligned}
& +O(\varepsilon^{2n+2}) \\
& = O\left(\int_{B_\delta(a)} \frac{\varepsilon^{2n+3}\rho_a^3|s^2 + |z|^4 - (\varepsilon/n)^4|}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_a \wedge (d\theta_a)^n\right) + O(\varepsilon^{2n+2}) \\
& = O\left(\int_{B_\delta(0)} \frac{\varepsilon^{2n+3}(s^2 + |z|^4)^{\frac{3}{4}}|s^2 + |z|^4 - (\varepsilon/n)^4|}{(s^2 + ((\varepsilon/n)^2 + |z|^2)^2)^{n+2}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n\right) + O(\varepsilon^{2n+2}) \\
& = O\left(\varepsilon^4 \int_{B_{\frac{\delta_0}{\varepsilon}}(0)} \frac{(s^2 + |z|^4)^{\frac{3}{4}}|s^2 + |z|^4 - 1|}{(s^2 + (1 + |z|^2)^2)^{n+2}} \theta_{\mathbb{H}^n} \wedge (d\theta_{\mathbb{H}^n})^n\right) + O(\varepsilon^{2n+2}) \\
& = O(\varepsilon^4).
\end{aligned}$$

This finishes the proof of Lemma 5.12.

#### REFERENCES

- [1] S. C. CHANG AND J. H. CHENG, The Harnack estimate for the Yamabe flow on CR manifolds of dimension 3. *Ann. Global Anal. Geom.* **21** (2002), no. 2, 111-121. [1](#)
- [2] S. C. CHANG, H. L. CHIU, AND C. T. WU, The Li-Yau-Hamilton inequality for Yamabe flow on a closed CR 3-manifold. *Trans. Amer. Math. Soc.* **362** (2010), no. 4, 1681-1698. [1](#)
- [3] S. C. CHANG, C. Y. LIN, AND C. T. WU, Eigenvalues and energy functional with monotonicity formulae under the CR Yamabe flow on a closed pseudohermitian 3-manifold. *Nonlinear Stud.* **18** (2011), no. 3, 377-392. [1](#)
- [4] J. H. CHENG, A. MALCHIODI, AND P. YANG, A positive mass theorem in three dimensional Cauchy-Riemann geometry. *Adv. Math.* **308** (2017), 276-347. [1](#)
- [5] J. H. CHENG, H. L. CHIU, AND P. YANG, Uniformization of spherical CR manifolds. *Adv. Math.* **255** (2014), 182-216. [1](#)
- [6] H. CHTIQUI, K. EL MEHDI, AND N. GAMARA, The Webster scalar curvature problem on the three dimensional CR manifolds. *Bull. Sci. Math.* **131** (2007), no. 4, 361-374. [1](#)
- [7] J. ESCORBAR AND R. SCHOEN, Conformal metrics with prescribed scalar curvature, *Invent. Math.*, **86** (1986), 243-254. [2](#), [22](#)
- [8] V. FELLI AND F. UGUZZONI, Some existence results for the Webster scalar curvature problem in presence of symmetry. *Ann. Mat. Pura Appl. (4)* **183** (2004), no. 4, 469-493. [1](#)
- [9] D. JERISON AND J. M. LEE, Extremals for the Sobolev inequality on the Heisenberg group and the CR Yamabe problem. *J. Amer. Math. Soc.* **1** (1988), no. 1, 1-13. [1](#), [16](#)
- [10] D. JERISON AND J. M. LEE, The Yamabe problem on CR manifolds. *J. Differential Geom.* **25** (1987), no. 2, 167-197. [1](#), [4](#), [19](#)
- [11] N. GAMARA, The CR Yamabe conjecture—the case  $n = 1$ . *J. Eur. Math. Soc. (JEMS)* **3** (2001), no. 2, 105-137. [1](#)
- [12] N. GAMARA AND R. YACOUB, CR Yamabe conjecture—the conformally flat case. *Pacific J. Math.* **201** (2001), no. 1, 121-175. [1](#)
- [13] P. T. HO, First eigenvalues of geometric operators under the Yamabe flow. *Ann. Global Anal. Geom.* **54** (2018), no. 4, 449-472. [1](#)
- [14] P. T. HO, Prescribed Webster scalar curvature on  $S^{2n+1}$  in the presence of reflection or rotation symmetry. *Bull. Sci. Math.* **140** (2016), no. 5, 506-518. [2](#)
- [15] P. T. HO, Results related to prescribing pseudo-Hermitian scalar curvature. *Internat. J. Math.* **24** (2013), no. 3, 1350020, 29 pp. [2](#)
- [16] P. T. HO, The long time existence and convergence of the CR Yamabe flow, *Commun. Contemp. Math.*, **14** (2012), 50 pp. [1](#), [13](#), [19](#)
- [17] P. T. HO, The Webster scalar curvature flow on CR sphere, Part I, *Adv. Math.*, **268** (2015), 758-835. [2](#), [14](#), [15](#)
- [18] P. T. HO, The Webster scalar curvature flow on CR sphere. Part II. *Adv. Math.*, **268** (2015), 836-905. [2](#)
- [19] P. T. HO AND S. KIM, CR Nirenberg problem and zero Webster scalar curvature. *Ann. Global Anal. Geom.* **58** (2020), no. 2, 207-226. [1](#)
- [20] P. T. HO AND S. KIM, The Yamabe problem on noncompact CR manifolds. *Pacific J. Math.* **285** (2016), no. 2, 375-391. [1](#), [4](#)



- [21] P. T. HO, W. M. SHENG, AND K. B. WANG, Convergence of the CR Yamabe flow, *Math. Ann.*, **373** (2019), 743-830. [1](#), [17](#), [22](#), [23](#), [24](#), [38](#), [39](#)
- [22] P. T. HO AND K. B. WANG, CR Yamabe constant, CR Yamabe flow and its soliton. *Nonlinear Anal.* **199** (2020), 112043, 16 pp. [1](#)
- [23] G. LU, The sharp Poincaré inequality for free vector fields: an endpoint result. *Rev. Mat. Iberoamericana*, **10** (1994), no. 2, 453-466. [7](#)
- [24] A. MALCHIODI AND F. UGUZZONI, A perturbation result for the Webster scalar curvature problem on the CR sphere. *J. Math. Pures Appl. (9)* **81** (2002), no. 10, 983-997. [1](#)
- [25] M. MAYER, A scalar curvature flow in low dimensions, *Cal. Var. PDE.*, **56** (2017), 41pp. [3](#), [23](#)
- [26] Q. A. NGÔ AND H. ZHANG, Prescribing Webster scalar curvature on CR manifolds of negative conformal invariants. *J. Differential Equations* **258** (2015), no. 12, 4443-4490. [1](#)
- [27] M. RIAHI AND N. GAMARA, Multiplicity results for the prescribed Webster scalar curvature on the three CR sphere under “flatness condition”. *Bull. Sci. Math.* **136** (2012), no. 1, 72-95. [1](#)
- [28] E. SALEM AND N. GAMARA, The Webster scalar curvature revisited: the case of the three dimensional CR sphere. *Calc. Var. Partial Differential Equations* **42** (2011), no. 1-2, 107-136. [1](#)
- [29] W. M. SHENG AND K. B. WANG, The exponential convergence of the CR Yamabe flow. *Sci. China Math.* **63** (2020), no. 5, 979-992. [1](#)

(P. T. Ho) DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 04107, KOREA.

Email address: [paktungho@yahoo.com.hk](mailto:paktungho@yahoo.com.hk)

(Q. A. Ngô) UNIVERSITY OF SCIENCE, VIỆT NAM NATIONAL UNIVERSITY, HÀ NỘI, VIỆT NAM

Email address: [nqanh@vnu.edu.vn](mailto:nqanh@vnu.edu.vn)

(H. Zhang) SCHOOL OF MATHEMATICS, SOUTH CHINA NORMAL UNIVERSITY, No.55 WEST ZHONGSHAN AVENUE, GUANGZHOU, GUANGDONG, CHINA, 510631.

Email address: [zhanghong@scnu.edu.cn](mailto:zhanghong@scnu.edu.cn)