# Hyers-Ulam stability of isometries on bounded domains

Soon-Mo Jung

Mathematics Section, College of Science and Technology, Hongik University, 30016 Sejong, Republic of Korea

*E-mail:* smjung@hongik.ac.kr

#### Abstract

More than 20 years after Fickett attempted to prove the Hyers-Ulam stability of isometries defined on bounded subsets of  $\mathbb{R}^n$  in 1981, Väisälä improved Fickett's result significantly. In this paper, we will improve Fickett's theorem by proving the Hyers-Ulam stability of isometries defined on bounded subsets of  $\mathbb{R}^n$  using a more intuitive method different from that used by Väisälä.

2010 Mathematics Subject Classification: Primary 46C99; Secondary 39B82, 39B62, 46B04.

Key Words: Isometry;  $\varepsilon$ -isometry; Hyers-Ulam stability; bounded domain.

# 1 Introduction

In 1940, Ulam gave a lecture at a mathematics club at the University of Wisconsin introducing some important unsolved problems. Then, based on that lecture, he published a book 20 years later (see [30]). A number of unresolved problems are introduced in this book, among which the following question about Hyers-Ulam stability of group homomorphism is closely related to the subject matter of this paper:

Let  $G_1$  be a group and let  $G_2$  be a metric group with the metric  $d(\cdot, \cdot)$ . Given  $\varepsilon > 0$ , does there exist a  $\delta > 0$  such that if a function  $h: G_1 \to G_2$  satisfies the inequality  $d(h(xy), h(x)h(y)) < \delta$  for all  $x, y \in G_1$ , then there exists a homomorphism  $H: G_1 \to G_2$  with  $d(h(x), H(x)) < \varepsilon$  for all  $x \in G_1$ ?

In 1941, the following year, Hyers [13] was able to successfully solve Ulam's question about the approximately additive functions, assuming that both  $G_1$  and  $G_2$  were Banach spaces. Indeed, he has proved that if a function  $f: G_1 \to G_2$  satisfies the inequality  $||f(x + y) - f(x) - f(y)|| \le \varepsilon$  for some  $\varepsilon \ge 0$  and all  $x, y \in G_1$ , then there exists an additive function  $A: G_1 \to G_2$  such that  $||f(x) - A(x)|| \le \varepsilon$  for each  $x \in G_1$ . In this case, the Cauchy additive equation, f(x + y) = f(x) + f(y), is said to have (or satisfy) the Hyers-Ulam stability. In the theorem of Hyers, the

relevant additive function  $A: G_1 \to G_2$  is constructed from the given function f by using the formula  $A(x) = \lim_{n \to \infty} \frac{1}{2^n} f(2^n x)$ . This method is now called the *direct method*.

We use the notations  $(E, \|\cdot\|)$  and  $(F, \|\cdot\|)$  to denote Hilbert spaces over  $\mathbb{K}$ , where  $\mathbb{K}$  is either  $\mathbb{R}$  or  $\mathbb{C}$ . A mapping  $f: E \to F$  is said to be an *isometry* if f satisfies

$$||f(x) - f(y)|| = ||x - y||$$
(1.1)

for any  $x, y \in E$ .

By considering the definition of Hyers and Ulam [14], for each fixed  $\varepsilon \ge 0$ , a function  $f: E \to F$  is said to be an  $\varepsilon$ -isometry if f satisfies the inequality

$$\left| \|f(x) - f(y)\| - \|x - y\| \right| \le \varepsilon$$
(1.2)

for any  $x, y \in E$ . If there exists a positive constant K depending only E and F (independent of f and  $\varepsilon$ ) such that for each  $\varepsilon$ -isometry  $f : E \to F$ , there is an isometry  $U : E \to F$  satisfying the inequality  $||f(x) - U(x)|| \leq K\varepsilon$  for every  $x \in E$ , then the functional equation (1.1) is said to have (or satisfy) the Hyers-Ulam stability.

To the best of our knowledge, Hyers and Ulam were the first mathematicians to study the Hyers-Ulam stability of isometries (see [14]). Indeed, they were able to prove the stability of isometries based on the properties of the inner product of Hilbert spaces: For each surjective  $\varepsilon$ -isometry f:  $E \to E$  satisfying f(0) = 0, there is a surjective isometry  $U : E \to E$  satisfying  $||f(x) - U(x)|| \leq 10\varepsilon$ for every  $x \in E$ . We encourage readers who want to read historically important papers dealing with similar topics to look for the papers [3, 5, 6, 15].

In 1978, Gruber [12] proved the following theorem: Assume that E and F are real normed spaces,  $f : E \to F$  is a surjective  $\varepsilon$ -isometry, and that  $U : E \to F$  is an isometry satisfying f(p) = U(p) for a  $p \in E$ . If ||f(x) - U(x)|| = o(||x||) as  $||x|| \to \infty$  uniformly, then U is a surjective linear isometry and  $||f(x) - U(x)|| \le 5\varepsilon$  for all  $x \in E$ . In particular, if f is continuous, then  $||f(x) - U(x)|| \le 3\varepsilon$  for each  $x \in E$ . This Gruber's result was further improved by Gevirtz [11] and by Omladič and Šemrl [20]. There are many other papers related to the stability of isometries, but it is regrettable that due to the restriction of space, they cannot be quoted one by one. Nevertheless, see [2, 4, 9, 10, 16, 17, 19, 22, 23, 24, 25, 26, 27, 29, 31] for more general information on the stability of isometries and related topics.

The following Fickett's theorem is an important motive for this paper (see [10]):

**Theorem 1.1 (Fickett)** For a fixed integer  $n \ge 2$ , let D be a bounded subset of  $\mathbb{R}^n$  and let  $\varepsilon > 0$  be given. If a function  $f: D \to \mathbb{R}^n$  satisfies the inequality (1.2) for all  $x, y \in D$ , then there exists an isometry  $U: D \to \mathbb{R}^n$  such that

$$\|f(x) - U(x)\| \le 27\varepsilon^{1/2^n}$$
(1.3)

for each  $x \in D$ .

The upper bound associated with inequality (1.3) in Fickett's theorem becomes very large for any sufficiently small  $\varepsilon$  in comparison to  $\varepsilon$ . This is a big drawback of Fickett's theorem. Thus, the work of further improving Fickett's theorem has to be attractive. After Fickett attempted to prove the Hyers-Ulam stability of isometries defined on a bounded subset of  $\mathbb{R}^n$  in 1981, several papers have been published steadily to improve his result over the past 40 years. However, most of the results were not very satisfactory (see [7, 8, 16, 17, 18]). Fortunately, however, Väisälä [31] introduced a new concept called the *c*-solar system and significantly improved Fickett's result by proving the Hyers-Ulam stability of isometries defined on bounded subsets of  $\mathbb{R}^n$ .

In this paper, we significantly improve Fickett's theorem by using a more intuitive method that is different from Väisälä's method. Indeed, we prove the Hyers-Ulam stability of isometries defined on bounded subsets of  $\mathbb{R}^n$  for  $n \geq 3$  (see Theorem 4.1).

# 2 Real version of QR decomposition

An orthogonal matrix  $\mathbf{Q}$  is a real square matrix whose columns and rows are orthonormal vectors. In other words, a real square matrix  $\mathbf{Q}$  is orthogonal if its transpose is equal to its inverse:  $\mathbf{Q}^{tr} = \mathbf{Q}^{-1}$ , where  $\mathbf{Q}^{tr}$  and  $\mathbf{Q}^{-1}$  stand for the transpose and the inverse of  $\mathbf{Q}$ , respectively. As a linear transformation, an orthogonal matrix preserves the inner product of vectors, and therefore acts as an isometry of Euclidean space.

Most papers and textbooks that mention QR decomposition only prove the complex version of QR decomposition (see [1, Theorem 6.3.7] or [28, Theorem 2.2 in §1]). However, not the complex version but the real version of QR decomposition is required in this paper. Nevertheless, since the proof of the real version is similar to the proof of the complex version, we will omit the proof of the real version here.

**Theorem 2.1 (QR decomposition)** Every real square matrix  $\mathbf{A}$  can be decomposed as  $\mathbf{A} = \mathbf{QR}$ , where  $\mathbf{Q}$  is an orthogonal matrix and  $\mathbf{R}$  is an upper triangular matrix whose elements are real numbers. In particular, every diagonal element of  $\mathbf{R}$  is nonnegative.

We can prove the following lemma using the real version of QR decomposition (Theorem 2.1), and this lemma plays an important role in achieving the final goal of this paper. In practice, using this lemma, we can almost halve the number of unknowns to consider in the main theorem.

**Lemma 2.2** Let  $\mathbb{R}^n$  be the n-dimensional Euclidean space for a fixed integer n > 0. Assume that D is a subset of  $\mathbb{R}^n$  with  $\{e_1, e_2, \ldots, e_n\} \subset D$ , where  $\{e_1, e_2, \ldots, e_n\}$  is the standard basis for  $\mathbb{R}^n$ . If  $f: D \to \mathbb{R}^n$  is a function and every  $e_i$  is written in column vector, then there exist an orthogonal matrix  $\mathbf{P}$  and real numbers  $e'_{ij}$  for  $i, j \in \{1, 2, \ldots, n\}$  with  $i \ge j$  such that

$$\mathbf{P}f(e_i) = (e'_{i1}, e'_{i2}, \dots, e'_{ii}, 0, \dots, 0)^{tr}$$

for every  $i \in \{1, 2, ..., n\}$ . In particular,  $e'_{ii} \ge 0$  for all  $i \in \{1, 2, ..., n\}$ .

*Proof.* Let  $f(e_i) = (e_{i1}, e_{i2}, \dots, e_{in})^{tr}$ , written in column vector, for any  $i \in \{1, 2, \dots, n\}$ . We now

define a matrix  $\mathbf{A}$  by

$$\mathbf{A} = (f(e_1) \ f(e_2) \ \cdots \ f(e_n)) = \begin{pmatrix} e_{11} & e_{21} & \cdots & e_{n1} \\ e_{12} & e_{22} & \cdots & e_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ e_{1n} & e_{2n} & \cdots & e_{nn} \end{pmatrix}.$$

By Theorem 2.1, there exist an orthogonal matrix  $\mathbf{Q}$  and an upper triangular (real) matrix  $\mathbf{R}$  such that  $\mathbf{A} = \mathbf{Q}\mathbf{R}$ . Thus, we have

$$\mathbf{R} = \mathbf{Q}^{tr} \mathbf{A} = \begin{pmatrix} \mathbf{Q}^{tr} f(e_1) & \mathbf{Q}^{tr} f(e_2) & \cdots & \mathbf{Q}^{tr} f(e_n) \end{pmatrix} = \begin{pmatrix} e'_{11} & e'_{21} & \cdots & e'_{n1} \\ 0 & e'_{22} & \cdots & e'_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & e'_{nn} \end{pmatrix}$$
(2.1)

for some real numbers  $e'_{11}, e'_{21}, e'_{22}, \ldots, e'_{n1}, \ldots, e'_{nn}$ , where the diagonal element  $e'_{ii}$  is nonnegative for all  $i \in \{1, 2, \ldots, n\}$ .

Finally, the last two terms of (2.1) are compared to conclude as follows:

$$\begin{cases} \mathbf{Q}^{tr} f(e_1) = (e'_{11}, 0, 0, \dots, 0)^{tr}, \\ \mathbf{Q}^{tr} f(e_2) = (e'_{21}, e'_{22}, 0, \dots, 0)^{tr}, \\ \vdots \\ \mathbf{Q}^{tr} f(e_n) = (e'_{n1}, e'_{n2}, e'_{n3}, \dots, e'_{nn})^{tr}. \end{cases}$$

If we put  $\mathbf{P} = \mathbf{Q}^{tr}$ , then  $\mathbf{P}$  is also an orthogonal matrix.

# 3 A preliminary theorem

Let  $\{e_1, e_2, \ldots, e_n\}$  be the standard basis for  $\mathbb{R}^n$ , where *n* is a fixed integer larger than 2. In this section, *D* denotes a subset of  $\mathbb{R}^n$  that only satisfies  $\{0, e_1, e_2, \ldots, e_n\} \subset D$ , whether bounded or not.

Lemma 2.2 implies that there exists an orthogonal matrix  $\mathbf{Q}$ , with which we can express  $f(e_i) = (e'_{i1}, e'_{i2}, \ldots, e'_{ii}, 0, \ldots, 0)^{tr}$  with respect to the new basis  $\{\mathbf{Q}e_1, \mathbf{Q}e_2, \ldots, \mathbf{Q}e_n\}$  instead of the standard basis  $\{e_1, e_2, \ldots, e_n\}$  and  $e'_{ii} \ge 0$  for all  $i \in \{1, 2, \ldots, n\}$ . For example, we can choose the orthogonal matrix  $\mathbf{Q}$  given in the proof of Lemma 2.2 for this purpose. Therefore, from now on, we will assume that  $f(e_i) = (e'_{i1}, e'_{i2}, \ldots, e'_{ii}, 0, \ldots, 0)$  with  $e'_{ii} \ge 0$ , written in row vector, without loss of generality.

In the statement of the following theorem, we define  $\sigma$  as if we knew the values of  $c_{(i+1)i}$  without knowing their values in advance. However, we note that in the proof of this theorem, we can justify the definition of  $\sigma$  by showing that the value of each  $c_{(i+1)i}$  is not greater than 9 (ref. Remark 3.1 (ii)). It is noted that the situation is similar for the  $c_{ii}$ 's.

**Theorem 3.1** Let  $\{e_1, e_2, \ldots, e_n\}$  be the standard basis for  $\mathbb{R}^n$ , where  $\mathbb{R}^n$  denotes the n-dimensional Euclidean space for a fixed integer  $n \ge 3$ , let D be a subset of  $\mathbb{R}^n$  satisfying  $\{0, e_1, e_2, \ldots, e_n\} \subset D$ , and let  $f: D \to \mathbb{R}^n$  be a function that satisfies f(0) = 0 and

$$|||f(x) - f(y)|| - ||x - y||| \le \varepsilon$$
 (3.1)

for all  $x, y \in \{0, e_1, e_2, \ldots, e_n\}$  and for some constant  $\varepsilon$  with  $0 < \varepsilon < \min\left\{\frac{1}{\sigma}, \min_{1 \le i \le n} \frac{1}{2c_{ii}}, \frac{1}{12}\right\}$ , where  $\sigma$  is defined as  $\sigma = \sum_{i=1}^{n-1} c_{(i+1)i}^2$  and  $c_{ii}$  and  $c_{(i+1)i}$  will be determined by the formulas (3.17) and (3.19), respectively. Then there exist positive integers  $c_{ij}, i, j \in \{1, 2, \ldots, n\}$  with  $j \le i$ , such that

$$\begin{cases}
-c_{ij}\varepsilon \leq e'_{ij} \leq c_{ij}\varepsilon & (for \ i > j), \\
1 - c_{ii}\varepsilon \leq e'_{ii} \leq 1 + \varepsilon & (for \ i = j)
\end{cases}$$
(3.2)

and such that the  $c_{ij}$  satisfy the equations in (3.15) for all  $i, j \in \{1, 2, ..., n\}$  with  $j \leq i$ .

*Proof.* (a) Using inequality (3.1) and by assumption f(0) = 0, we have

$$\left| \|f(e_j)\| - 1 \right| \le \varepsilon \text{ and } \left| \|f(e_k) - f(e_\ell)\| - \sqrt{2} \right| \le \varepsilon$$

for any  $j, k, \ell \in \{1, 2, ..., n\}$  with  $k < \ell$ . Since  $f(e_j) = (e'_{j1}, ..., e'_{jj}, 0, ..., 0)$  for all  $j \in \{1, 2, ..., n\}$ and  $\|\cdot\|$  is the Euclidean norm on  $\mathbb{R}^n$ , from the last inequalities we get the following two inequalities, which are equivalent to the inequality (3.1) for  $x, y \in \{0, e_1, e_2, ..., e_n\}$ :

$$(1-\varepsilon)^2 \le \sum_{i=1}^{j} e_{ji}^{\prime 2} \le (1+\varepsilon)^2$$
 (3.3)

for each  $j \in \{1, 2, \ldots, n\}$  and

$$\left(\sqrt{2} - \varepsilon\right)^2 \le \sum_{i=1}^k e_{ki}'^2 - \sum_{i=1}^k 2e_{ki}'e_{\ell i}' + \sum_{i=1}^\ell e_{\ell i}'^2 \le \left(\sqrt{2} + \varepsilon\right)^2 \tag{3.4}$$

for every  $k, \ell \in \{1, 2, ..., n\}$  with  $k < \ell$ . From now on, we will prove this theorem by using inequalities (3.3) and (3.4) instead of inequality (3.1).

(b) Now we apply the 'main' induction on m to prove the array of equations presented in (3.15). Proving the array of (3.15) is the most important and longest part of this proof.

(b.1) According to Lemma 2.2,  $e'_{11}$  is a nonnegative real number, so setting j = 1 in (3.3) gives us the inequality,  $1 - \varepsilon \leq e'_{11} \leq 1 + \varepsilon$ , and we select  $c_{11} = 1$  as the smallest positive integer that satisfies the following inequality

$$1 - c_{11}\varepsilon \le 1 - \varepsilon \le e'_{11} \le 1 + \varepsilon. \tag{3.5}$$

This fact guarantees the existence of  $c_{11}$  satisfying the second condition of (3.2) for i = j = 1. (We note that  $c_{11}$  must not be necessarily the smallest positive integer satisfying the inequality (3.5),

for example,  $c_{11} = 2$  is not wrong, but if possible, the smaller the  $c_{11}$  is, the better.) If we set j = 2 in (3.3) and put k = 1 and  $\ell = 2$  in (3.4) and then combine the resulting inequalities, then we get

$$\frac{-(2c_{11}+2+2\sqrt{2})\varepsilon + c_{11}^2\varepsilon^2}{2(1-c_{11}\varepsilon)} \le e_{21}' \le \frac{(4+2\sqrt{2})\varepsilon + \varepsilon^2}{2(1-c_{11}\varepsilon)}$$

and we can choose  $c_{21} = 4$  as the smallest positive integer satisfying the condition

$$-c_{21}\varepsilon \le \frac{-(2c_{11}+2+2\sqrt{2})\varepsilon + c_{11}^2\varepsilon^2}{2(1-c_{11}\varepsilon)} \le e_{21}' \le \frac{(4+2\sqrt{2})\varepsilon + \varepsilon^2}{2(1-c_{11}\varepsilon)} \le c_{21}\varepsilon.$$
(3.6)

Obviously, this fact confirms the existence of  $c_{21}$  satisfying the first condition of (3.2) for i = 2and j = 1. (We note that  $c_{ij}$  must not be necessarily the smallest positive integer under given conditions, for example,  $c_{21} = 5$  is not bad, but if possible, the smaller the  $c_{ij}$  is, the better. Indeed, assuming that the  $c_{ij}$  are the smallest positive integers that satisfy some given conditions, we can find the unique  $c_{ij}$  (for i > j), which make it easier to prove the array of equations presented in (3.15). This is one of the reasons we want  $c_{ij}$  to be the smallest positive integer.)

Furthermore, if we set j = 3 in (3.3) and put k = 1 and  $\ell = 3$  in (3.4) and then combine the resulting inequalities, then we get the inner part of the following inequalities

$$-c_{31}\varepsilon \le \frac{-(2c_{11}+2+2\sqrt{2})\varepsilon + c_{11}^2\varepsilon^2}{2(1-c_{11}\varepsilon)} \le e_{31}' \le \frac{(4+2\sqrt{2})\varepsilon + \varepsilon^2}{2(1-c_{11}\varepsilon)} \le c_{31}\varepsilon$$
(3.7)

and we select  $c_{31}$  as the smallest positive integer that satisfies the outermost inequalities of (3.7). From this fact, we confirm the existence of  $c_{31}$  that satisfies the first condition in (3.2). Since  $c_{21}$ and  $c_{31}$  are assumed to be the smallest positive integers satisfying the outermost inequalities of (3.6) and (3.7), respectively, we come to the conclusion that  $c_{31} = c_{21} = 4$ .

Moreover, using (3.3) with j = 2 and by a direct calculation, since  $0 < \varepsilon < \frac{1}{12}$ , we get

$$1 - 2\varepsilon - \frac{19}{12}\varepsilon + 4\varepsilon^2 < 1 - 2\varepsilon - 19\varepsilon^2 + 4\varepsilon^2 = (1 - \varepsilon)^2 - c_{21}^2\varepsilon^2 \le c_{22}^{\prime 2} \le (1 + \varepsilon)^2$$

and, if possible, we determine  $c_{22}$  as the smallest positive integer which satisfies the condition

$$(1 - c_{22}\varepsilon)^2 \le 1 - 2\varepsilon - \frac{19}{12}\varepsilon + 4\varepsilon^2 \le e_{22}^{\prime 2} \le (1 + \varepsilon)^2$$

The last inequalities assures the existence of  $c_{22} = 2$  satisfying the second condition of (3.2) for i = j = 2.

In a similar way, putting k = 2 and  $\ell = 3$  in (3.4) and a routine calculation show the existence of  $c_{32}$  satisfying the first condition of (3.2), for example,  $c_{32} = 6$ . Analogously, since  $0 < \varepsilon < \frac{1}{12}$ , the inequality (3.3) with j = 3 yields the inequality,  $1 - 2\varepsilon - 51\varepsilon^2 \le e_{33}^{\prime 2} \le (1 + \varepsilon)^2$ , and we can choose the smallest positive integer  $c_{33}$  that satisfies

$$(1 - c_{33}\varepsilon)^2 \le 1 - 2\varepsilon - 51\varepsilon^2 \le e_{33}^{\prime 2} \le (1 + \varepsilon)^2,$$

which shows that  $c_{33}$  exists that satisfies the second condition of (3.2). More directly, we can choose  $c_{33} = 4$ . Therefore, all the integers  $c_{ij}$  considered in (b.1) satisfy the conditions in (3.2) and (3.15) for n = 3. By doing this, we start the induction (with m = 3).

(b.2) Induction hypothesis. Let m be some integer satisfying  $3 \le m < n$ . It is assumed that the smallest positive integers  $c_{ij}$ ,  $i, j \in \{1, 2, ..., m\}$  with  $j \le i$ , were found by the methods we did in the subsection (b.1), and that these positive integers satisfy the following inequalities

$$\begin{cases} -c_{ij}\varepsilon \leq e'_{ij} \leq c_{ij}\varepsilon & \text{(for } i > j\text{)}, \\ 1 - c_{ii}\varepsilon \leq e'_{ii} \leq 1 + \varepsilon & \text{(for } i = j\text{)} \end{cases}$$

as well as the array of equations

$$\begin{array}{rclcrcl} c_{m1} & = & c_{(m-1)1} & = & c_{(m-2)1} & = & \dots & = c_{41} = c_{31} = c_{21}, \\ c_{m2} & = & c_{(m-1)2} & = & c_{(m-2)2} & = & \dots & = c_{42} = c_{32}, \\ c_{m3} & = & c_{(m-1)3} & = & c_{(m-2)3} & = & \dots & = c_{43}, \\ \vdots & & \vdots & & \vdots & & \vdots & \\ c_{m(m-3)} & = & c_{(m-1)(m-3)} & = & c_{(m-2)(m-3)}, \\ c_{m(m-2)} & = & c_{(m-1)(m-2)}, \\ c_{m(m-1)} \end{array}$$

The last line in the above array consisting of only  $c_{m(m-1)}$  means that there exists the smallest possible positive integer  $c_{m(m-1)}$  that satisfies  $-c_{m(m-1)}\varepsilon \leq e'_{m(m-1)} \leq c_{m(m-1)}\varepsilon$ .

(b.3) We let j = m + 1 in (3.3) and  $\ell = m + 1$  in (3.4) to get

$$(1-\varepsilon)^2 \le \sum_{i=1}^{m+1} e_{(m+1)i}^{\prime 2} \le (1+\varepsilon)^2$$
(3.8)

and

$$\left(\sqrt{2}-\varepsilon\right)^2 \le \sum_{i=1}^k e_{ki}^{\prime 2} - \sum_{i=1}^k 2e_{ki}^{\prime} e_{(m+1)i}^{\prime} + \sum_{i=1}^{m+1} e_{(m+1)i}^{\prime 2} \le \left(\sqrt{2}+\varepsilon\right)^2 \tag{3.9}$$

for every  $k \in \{1, 2, ..., m\}$ .

Similar to what we did to get (3.6), the inequalities (3.5), (3.8), and (3.9) with k = 1 yield the inner ones of the following inequalities

$$-c_{(m+1)1}\varepsilon \leq \frac{-(2c_{11}+2+2\sqrt{2})\varepsilon + c_{11}^2\varepsilon^2}{2(1-c_{11}\varepsilon)} \leq e'_{(m+1)1} \leq \frac{(4+2\sqrt{2})\varepsilon + \varepsilon^2}{2(1-c_{11}\varepsilon)} \leq c_{(m+1)1}\varepsilon, \quad (3.10)$$

and we find the smallest positive integer  $c_{(m+1)1}$  satisfying the outermost inequalities of (3.10). By comparing both inequalities (3.6) and (3.10), we may conclude that  $c_{(m+1)1} = c_{21}$ , with which we initiate an 'inner' induction that is subordinate to the main induction.

(b.3.1) We choose some  $k \in \{2, 3, ..., m\}$  and assume that  $-c_{(m+1)i}\varepsilon \leq e'_{(m+1)i} \leq c_{(m+1)i}\varepsilon$  and  $c_{(m+1)i} = c_{(i+1)i}$  for each  $i \in \{1, 2, ..., k-1\}$ . This is the hypothesis for our inner induction on i that operates inside the main induction on m. Based on this hypothesis, we will prove that there exists a positive integer  $c_{(m+1)k}$  that satisfies  $-c_{(m+1)k}\varepsilon \leq e'_{(m+1)k}\varepsilon$  as well as  $c_{(m+1)k} = c_{(k+1)k}$ . Roughly speaking, this inner induction works to expand each row of (3.15) horizontally.

(b.3.2) It follows from (3.9) that

$$\left(\sqrt{2} - \varepsilon\right)^{2} - \sum_{i=1}^{k} e_{ki}^{\prime 2} + \sum_{i=1}^{k-1} 2e_{ki}^{\prime} e_{(m+1)i}^{\prime} - \sum_{i=1}^{m+1} e_{(m+1)i}^{\prime 2}$$

$$\leq -2e_{kk}^{\prime} e_{(m+1)k}^{\prime} \leq \qquad (3.11)$$

$$\left(\sqrt{2} + \varepsilon\right)^{2} - \sum_{i=1}^{k} e_{ki}^{\prime 2} + \sum_{i=1}^{k-1} 2e_{ki}^{\prime} e_{(m+1)i}^{\prime} - \sum_{i=1}^{m+1} e_{(m+1)i}^{\prime 2}$$

for any  $k \in \{2, 3, \dots, m\}$ . On the other hand, by (3.3) and (3.8), we have

$$(1-\varepsilon)^2 \le \sum_{i=1}^k e_{ki}^{\prime 2} \le (1+\varepsilon)^2$$
 and  $(1-\varepsilon)^2 \le \sum_{i=1}^{m+1} e_{(m+1)i}^{\prime 2} \le (1+\varepsilon)^2$ 

for each  $k \in \{2, 3, \ldots, m\}$ . Moreover, it follows from the hypotheses (b.2) and (b.3.1) that

$$-\sum_{i=1}^{k-1} 2c_{ki}c_{(i+1)i}\varepsilon^2 = -\sum_{i=1}^{k-1} 2c_{ki}c_{(m+1)i}\varepsilon^2 \le \sum_{i=1}^{k-1} 2c'_{ki}e'_{(m+1)i} \le \sum_{i=1}^{k-1} 2c_{ki}c_{(m+1)i}\varepsilon^2 = \sum_{i=1}^{k-1} 2c_{ki}c_{(i+1)i}\varepsilon^2$$

for all  $k \in \{2, 3, \dots, m\}$ .

Since  $c_{kk}\varepsilon < \frac{1}{2}$  and  $e'_{kk} > 0$  by (b.2), we use (3.11) and the last inequalities to get the inner ones of the following inequalities

$$-c_{(m+1)k}\varepsilon \leq \frac{1}{2e'_{kk}} \left( -\left(4+2\sqrt{2}\right)\varepsilon + \varepsilon^2 - 2\sum_{i=1}^{k-1} c_{ki}c_{(i+1)i}\varepsilon^2 \right)$$
  
$$\leq e'_{(m+1)k} \leq \qquad (3.12)$$
  
$$\frac{1}{2e'_{kk}} \left( \left(4+2\sqrt{2}\right)\varepsilon + \varepsilon^2 + 2\sum_{i=1}^{k-1} c_{ki}c_{(i+1)i}\varepsilon^2 \right) \leq c_{(m+1)k}\varepsilon$$

for all  $k \in \{2, 3, ..., m\}$  and we can select the smallest positive integer  $c_{(m+1)k}$  that satisfies the outermost inequalities of (3.12). Similarly, by (3.3) and (3.4) with  $\ell = k + 1$ , a routine calculation yields

$$-c_{(k+1)k}\varepsilon \leq \frac{1}{2e'_{kk}} \left( -\left(4 + 2\sqrt{2}\right)\varepsilon + \varepsilon^2 - 2\sum_{i=1}^{k-1} c_{ki}c_{(k+1)i}\varepsilon^2 \right)$$

$$\leq e'_{(k+1)k} \leq \qquad (3.13)$$

$$\frac{1}{2e'_{kk}} \left( \left(4 + 2\sqrt{2}\right)\varepsilon + \varepsilon^2 + 2\sum_{i=1}^{k-1} c_{ki}c_{(k+1)i}\varepsilon^2 \right) \leq c_{(k+1)k}\varepsilon,$$

where  $c_{(k+1)k}$  is the smallest positive integer that satisfies the outmost conditions of (3.13). We note by (b.2) and (b.3.1) that  $c_{(k+1)i} = c_{(i+1)i}$  for every integer *i* satisfying 0 < i < k. Comparing (3.12) and (3.13), we conclude that  $c_{(m+1)k} = c_{(k+1)k}$  for each  $k \in \{2, 3, \ldots, m\}$ . Furthermore, referring to the subsection (b.3), we see that  $c_{(m+1)k} = c_{(k+1)k}$  holds for all  $k \in \{1, 2, \ldots, m\}$ , which proves the truth of the first column of the array of equations in subsection (b.3.3) below.

Moreover, the inequality (3.3) with j = m + 1 yields

$$(1-\varepsilon)^2 - \sum_{i=1}^m e_{(m+1)i}^{\prime 2} \le e_{(m+1)(m+1)}^{\prime 2} \le (1+\varepsilon)^2 - \sum_{i=1}^m e_{(m+1)i}^{\prime 2}.$$
(3.14)

Since  $0 < \varepsilon < \frac{1}{\sigma}$ , m < n, and  $0 < \varepsilon < \frac{1}{12}$ , it follows from (3.14) and some manipulation that

$$(1+\varepsilon)^2 \ge e_{(m+1)(m+1)}^{\prime 2} \ge (1-\varepsilon)^2 - \sum_{i=1}^m c_{(m+1)i}^2 \varepsilon^2 = (1-\varepsilon)^2 - \sum_{i=1}^m c_{(i+1)i}^2 \varepsilon^2$$
$$\ge (1-\varepsilon)^2 - \sigma\varepsilon^2 > 1 - 3\varepsilon + \varepsilon^2 > 9\varepsilon + \varepsilon^2 > 100\varepsilon^2.$$

We see that  $0 < (1 - c\varepsilon)^2 \le 100\varepsilon^2$  whenever c is a positive integer satisfying  $\frac{1}{\varepsilon} - 10 \le c < \frac{1}{\varepsilon}$ . This fact shows the existence of  $c_{(m+1)(m+1)}$  that satisfies the second condition of (3.2).

(b.3.3) We just proved in the subsections from (b.2) to (b.3.2) that there exist positive integers  $c_{ij}, i, j \in \{1, 2, \ldots, m+1\}$  with  $j \leq i$ , such that

$$\begin{cases} -c_{ij}\varepsilon \leq e'_{ij} \leq c_{ij}\varepsilon & (\text{for } i > j), \\ 1 - c_{ii}\varepsilon \leq e'_{ii} \leq 1 + \varepsilon & (\text{for } i = j), \end{cases}$$

and the  $c_{ij}$ 's satisfy

$$\begin{array}{rclcrcrc} c_{(m+1)1} &=& c_{m1} &=& c_{(m-1)1} &=& \dots &= c_{41} = c_{31} = c_{21}, \\ c_{(m+1)2} &=& c_{m2} &=& c_{(m-1)2} &=& \dots &= c_{42} = c_{32}, \\ c_{(m+1)3} &=& c_{m3} &=& c_{(m-1)3} &=& \dots &= c_{43}, \\ \vdots & \vdots & \vdots & &\vdots \\ c_{(m+1)(m-2)} &=& c_{m(m-2)} &=& c_{(m-1)(m-2)}, \\ c_{(m+1)(m-1)} &=& c_{m(m-1)}, \\ c_{(m+1)m} \end{array}$$

The last row in the above array consisting of only  $c_{(m+1)m}$  means that there exists the smallest possible positive integer  $c_{(m+1)m}$  that satisfies  $-c_{(m+1)m}\varepsilon \leq e'_{(m+1)m} \leq c_{(m+1)m}\varepsilon$ . (We can check in (b.3.2) the truth of equations in the first column of the above array. Moreover, we remember that we have assumed in (b.2) that the rest of equations in the array are true.)

(b.4) Altogether, by the main induction conclusion on m ( $3 \le m < n$ ), we may conclude that there exist positive integers  $c_{ij}$ ,  $i, j \in \{1, 2, ..., n\}$  with  $j \le i$ , such that each inequality in (3.2) holds true and the  $c_{ij}$ 's satisfy

which completes the first part of our proof. We remark that the last row  $c_{n(n-1)}$  in the above array implies that there is an integer  $c_{n(n-1)} > 0$  satisfying  $-c_{n(n-1)}\varepsilon \leq e'_{n(n-1)} \leq c_{n(n-1)}\varepsilon$ .

(c) Now we will introduce efficient methods to estimate the positive integers  $c_{jj}$  and  $c_{k(k-1)}$  for every  $j \in \{1, 2, ..., n\}$  and  $k \in \{2, 3, ..., n\}$ .

(c.1) We note that the inequality (3.3) holds true for all  $j \in \{1, 2, ..., n\}$ . Since  $-c_{ji}\varepsilon \leq c_{ji}\varepsilon$  for any  $i, j \in \{1, 2, ..., n\}$  with i < j, we determine the  $c_{jj}$  as the smallest possible positive integer that satisfies

$$(1 - c_{jj}\varepsilon)^2 \le (1 - \varepsilon)^2 - \sum_{i=1}^{j-1} c_{ji}^2 \varepsilon^2 \le (1 - \varepsilon)^2 - \sum_{i=1}^{j-1} e_{ji}'^2 \le e_{jj}'^2 \le (1 + \varepsilon)^2$$
(3.16)

for all  $j \in \{2, 3, ..., n\}$ . However, since the previous inequality is inefficient in practical calculations, we introduce a more practical inequality even at the expense of the smallest property of  $c_{jj}$ . Instead of (3.16), we will determine the  $c_{jj}$  as the smallest positive integer that satisfies the new condition

$$(3c_{jj} - 1)(c_{jj} - 1) \ge \sum_{i=1}^{j-1} c_{(i+1)i}^2$$
(3.17)

for all  $j \in \{2, 3, \ldots, n\}$ , which proves the existence of  $c_{jj}$ . Indeed, since  $0 < \varepsilon < \frac{1}{2c_{jj}}$ , we have

$$(1 - c_{jj}\varepsilon)^2 = 1 - 2\varepsilon c_{jj} + \varepsilon^2 c_{jj}^2 < 1 - 2\varepsilon c_{jj} + \frac{1}{2}\varepsilon c_{jj} = 1 - \frac{3}{2}\varepsilon c_{jj}.$$

It further follows from (3.15), (3.17), and the last inequality that

$$(1 - c_{jj}\varepsilon)^{2} < 1 - \frac{3}{2}\varepsilon c_{jj} = 1 - \frac{\varepsilon}{2c_{jj}} \cdot 3c_{jj}^{2} \le 1 - \frac{\varepsilon}{2c_{jj}} \left(\sum_{i=1}^{j-1} c_{(i+1)i}^{2} - 1 + 4c_{jj}\right)$$
$$= 1 - 2\varepsilon - \frac{\varepsilon}{2c_{jj}} \left(\sum_{i=1}^{j-1} c_{(i+1)i}^{2} - 1\right) \le 1 - 2\varepsilon - \varepsilon^{2} \left(\sum_{i=1}^{j-1} c_{(i+1)i}^{2} - 1\right)$$
$$= (1 - \varepsilon)^{2} - \sum_{i=1}^{j-1} c_{(i+1)i}^{2} \varepsilon^{2} = (1 - \varepsilon)^{2} - \sum_{i=1}^{j-1} c_{ji}^{2} \varepsilon^{2},$$

which implies the validity of (3.16).

(c.2) We note that the inequality (3.4) holds for all  $k, \ell \in \{1, 2, ..., n\}$  with  $k < \ell$ . If we set  $\ell = k + 1$  in (3.4) and make some manipulations, then we obtain the inner ones of the following inequalities

$$-c_{(k+1)k}\varepsilon \leq \frac{1}{2e'_{kk}} \left( -\left(4+2\sqrt{2}\right)\varepsilon + \varepsilon^2 - 2\sum_{i=1}^{k-1} c_{ki}c_{(k+1)i}\varepsilon^2 \right)$$

$$\leq e'_{(k+1)k} \leq \qquad (3.18)$$

$$\frac{1}{2e'_{kk}} \left( \left(4+2\sqrt{2}\right)\varepsilon + \varepsilon^2 + 2\sum_{i=1}^{k-1} c_{ki}c_{(k+1)i}\varepsilon^2 \right) \leq c_{(k+1)k}\varepsilon$$

The inequalities (3.2) and (3.18) show the existence of the smallest positive integer  $c_{(k+1)k}$  that satisfies

$$\frac{1}{2(1-c_{kk}\varepsilon)}\left(4+2\sqrt{2}+\varepsilon+2\sum_{i=1}^{k-1}c_{ki}c_{(k+1)i}\varepsilon\right) \le c_{(k+1)k}$$
(3.19)

for  $k \in \{1, 2, ..., n-1\}$ : Since  $0 < \varepsilon < \frac{1}{2c_{kk}}$ , we see that  $\frac{1}{2(1-c_{kk}\varepsilon)} < 1$ . Further, since  $c_{ki} = c_{(k+1)i} = c_{(i+1)i}$  for any  $i \in \{1, 2, ..., k-1\}$  and  $0 < \varepsilon < \frac{1}{\sigma}$ , we know that  $\sum_{i=1}^{k-1} c_{ki}c_{(k+1)i}\varepsilon \le \sigma\varepsilon < 1$ . Thus, we get

$$\frac{1}{2(1-c_{kk}\varepsilon)}\left(4+2\sqrt{2}+\varepsilon+2\sum_{i=1}^{k-1}c_{ki}c_{(k+1)i}\varepsilon\right)<6+2\sqrt{2}+\varepsilon<9,$$

which together with (3.19) assures the existence of  $c_{(k+1)k}$  with  $0 < c_{(k+1)k} \leq 9$  for all  $k \in \{1, 2, \ldots, n-1\}$ .

- **Remark 3.1** (i) The inequality (3.1) is a sufficient condition for inequalities in (3.2), and the inequalities in (3.2) are necessary conditions for the inequality (3.1).
- (ii) In view of the last part of the proof of Theorem 3.1, we see that  $0 < c_{(i+1)i} \leq 9$  for any  $i \in \{1, 2, \ldots, n-1\}$ .
- (iii) We solve the quadratic inequality (3.17) with respect to  $c_{jj}$  as follows:

$$c_{jj} \ge \frac{1}{3} \left( 2 + \sqrt{1 + 3\sum_{i=1}^{j-1} c_{(i+1)i}^2} \right)$$

for all  $j \in \{1, 2, ..., n\}$ .

# 4 Hyers-Ulam stability of isometries on bounded domains

The following theorem significantly improves the Fickett's theorem by demonstrating the Hyers-Ulam stability of isometries on the bounded domains.

As before, let  $\{e_1, e_2, \ldots, e_n\}$  be the standard basis for  $\mathbb{R}^n$ . Based on Lemma 2.2, we can assume that  $f(e_i) = (e'_{i1}, e'_{i2}, \ldots, e'_{ii}, 0, \ldots, 0)$  is written in row vector, where  $e'_{ii} \ge 0$  for each  $i \in \{1, 2, \ldots, n\}$ . We denote by  $B_d(0)$  the closed ball of radius d and centered at the origin of  $\mathbb{R}^n$ , *i.e.*,  $B_d(0) = \{x \in \mathbb{R}^n : ||x|| \le d\}$ .

**Theorem 4.1** Assume that  $\mathbb{R}^n$  is the n-dimensional Euclidean space for a given integer  $n \geq 3$ . Let D be a subset of  $\mathbb{R}^n$  satisfying  $\{0, e_1, e_2, \ldots, e_n\} \subset D \subset B_d(0)$  for some  $d \geq 1$  and let  $f: D \to \mathbb{R}^n$ be a function satisfying f(0) = 0 and the inequality (3.1) for all  $x, y \in D$  and some constant  $\varepsilon$  with  $0 < \varepsilon < \min\left\{\frac{1}{\sigma}, \min_{1 \le i \le n} \frac{1}{2c_{ii}}, \frac{1}{12}\right\}$ , where  $\sigma = \sum_{i=1}^{n-1} c_{(i+1)i}^2$  and the  $c_{ij}$ , for all  $i, j \in \{1, 2, ..., n\}$  with  $j \le i$ , are the positive integers estimated in Theorem 3.1. Then there exists an isometry  $U: D \to \mathbb{R}^n$  such that

$$\|f(x) - U(x)\| \le \left[\sum_{i=1}^{n} \left( \left(2 + \sum_{j=1}^{i} c_{ij}\right) d + 4 + \sum_{j=1}^{i} c_{ij} \right)^{2} \right]^{1/2} \varepsilon$$

for all  $x \in D$ .

Proof. Let  $\{e_1, e_2, \ldots, e_n\}$  be the standard basis for  $\mathbb{R}^n$ . Based on Lemma 2.2, it can be assumed that  $f(e_i) = (e'_{i1}, e'_{i2}, \ldots, e'_{ii}, 0, \ldots, 0)$ , where  $e'_{ii} \ge 0$  for each  $i \in \{1, 2, \ldots, n\}$ . For an arbitrary point  $x = (x_1, x_2, \ldots, x_n)$  of D, let  $f(x) = (x'_1, x'_2, \ldots, x'_n)$ . It follows from (3.1) that

$$\left| \|f(x)\| - \|x\| \right| \le \varepsilon$$
 and  $\left| \|f(x) - f(e_j)\| - \|x - e_j\| \right| \le \varepsilon$ 

and hence, we have

$$\left| \left( \sum_{i=1}^{n} x_i^{\prime 2} \right)^{1/2} - \left( \sum_{i=1}^{n} x_i^2 \right)^{1/2} \right| \le \varepsilon,$$
(4.1)

$$\left| \left( \sum_{i=1}^{j} (x_i' - e_{ji}')^2 + \sum_{i=j+1}^{n} x_i'^2 \right)^{1/2} - \left( \sum_{i=1}^{n} x_i^2 - 2x_j + 1 \right)^{1/2} \right| \le \varepsilon$$
(4.2)

for all  $j \in \{1, 2, ..., n\}$ .

It follows from (4.1) that

$$\left|\sum_{i=1}^{n} x_{i}^{\prime 2} - \sum_{i=1}^{n} x_{i}^{2}\right| = \left|\left(\sum_{i=1}^{n} x_{i}^{\prime 2}\right)^{1/2} + \left(\sum_{i=1}^{n} x_{i}^{2}\right)^{1/2}\right| \left|\left|\left(\sum_{i=1}^{n} x_{i}^{\prime 2}\right)^{1/2} - \left(\sum_{i=1}^{n} x_{i}^{2}\right)^{1/2}\right|\right| \le (2d+1)\varepsilon$$

$$(4.3)$$

since  $\sqrt{x_1'^2 + \cdots + x_n'^2} \leq d + \varepsilon$ ,  $\sqrt{x_1^2 + \cdots + x_n^2} \leq d$ , and  $0 < \varepsilon < 1$ . Similarly, it follows from (4.2) that

$$\left| \left( \sum_{i=1}^{j} (x'_i - e'_{ji})^2 + \sum_{i=j+1}^{n} x'^2_i \right) - \left( \sum_{i=1}^{n} x^2_i - 2x_j + 1 \right) \right| \le (2d+3)\varepsilon$$

$$(4.4)$$

for all  $j \in \{1, 2, \ldots, n\}$ , since  $0 < \varepsilon < 1$  and

$$\left(\sum_{i=1}^{j} (x'_i - e'_{ji})^2 + \sum_{i=j+1}^{n} x'^2_i\right)^{1/2} \le d+1 + \varepsilon \quad \text{and} \quad \left(\sum_{i=1}^{n} x^2_i - 2x_j + 1\right)^{1/2} \le d+1.$$

We use (4.4) to get

$$-(2d+3)\varepsilon - \left(\sum_{i=1}^{n} x_{i}^{\prime 2} - \sum_{i=1}^{n} x_{i}^{2}\right) + \sum_{i=1}^{j-1} 2e_{ji}^{\prime} x_{i}^{\prime} + 1 - \sum_{i=1}^{j} e_{ji}^{\prime 2}$$

$$\leq 2x_{j} - 2e_{jj}^{\prime} x_{j}^{\prime} \leq$$

$$(2d+3)\varepsilon - \left(\sum_{i=1}^{n} x_{i}^{\prime 2} - \sum_{i=1}^{n} x_{i}^{2}\right) + \sum_{i=1}^{j-1} 2e_{ji}^{\prime} x_{i}^{\prime} + 1 - \sum_{i=1}^{j} e_{ji}^{\prime 2}$$

$$(4.5)$$

for any  $j \in \{1, 2, ..., n\}$ .

Since  $|x_i'| \le ||f(x)|| \le ||x|| + \varepsilon < d + 1$  and by Theorem 3.1, we get

$$-2(d+1)\sum_{i=1}^{j-1}c_{ji}\varepsilon \le \sum_{i=1}^{j-1}2e'_{ji}x'_i \le 2(d+1)\sum_{i=1}^{j-1}c_{ji}\varepsilon.$$

Moreover, by (3.3), we have

$$-3\varepsilon \le -2\varepsilon - \varepsilon^2 \le 1 - \sum_{i=1}^{j} e_{ji}^{\prime 2} \le 2\varepsilon - \varepsilon^2 \le 3\varepsilon.$$

Therefore, it follows from (4.3) and (4.5) that

$$-\left(\left(4+2\sum_{i=1}^{j-1}c_{ji}\right)d+7+2\sum_{i=1}^{j-1}c_{ji}\right)\varepsilon \le 2x_j - 2e'_{jj}x'_j \le \left(\left(4+2\sum_{i=1}^{j-1}c_{ji}\right)d+7+2\sum_{i=1}^{j-1}c_{ji}\right)\varepsilon$$

for all  $j \in \{1, 2, ..., n\}$ .

We note that  $|x'_j| < d+1$  and  $-c_{jj}\varepsilon \le 1 - e'_{jj} \le c_{jj}\varepsilon$  by Theorem 3.1, and since  $x_j - e'_{jj}x'_j = (x_j - x'_j) + (1 - e'_{jj})x'_j$ , we can see that

$$\left|x_{j} - x_{j}'\right| \leq \left(\left(2 + \sum_{i=1}^{j} c_{ji}\right)d + 4 + \sum_{i=1}^{j} c_{ji}\right)\varepsilon$$
(4.6)

for  $j \in \{1, 2, \dots, n\}$ .

Since we can select an isometry  $U: D \to \mathbb{R}^n$  defined by  $U(x) = x = (x_1, x_2, \dots, x_n)$ , we see that

$$\|f(x) - U(x)\| = \left\| \left( x_1' - x_1, x_2' - x_2, \dots, x_n' - x_n \right) \right\| = \left( \sum_{j=1}^n \left( x_j' - x_j \right)^2 \right)^{1/2}$$
$$\leq \left( \sum_{j=1}^n \left( \left( 2 + \sum_{i=1}^j c_{ji} \right) d + 4 + \sum_{i=1}^j c_{ji} \right)^2 \right)^{1/2} \varepsilon$$

for all  $x \in D$ .

We remark that for any  $i, j \in \{1, 2, ..., n\}$  with  $i \ge j$ , each  $c_{ij}$  is independent of  $\varepsilon$  for any 'sufficiently' small  $\varepsilon > 0$ .

#### $\mathbf{5}$ Examples

**Example 5.1** We assume that n = 4. We will compute some constants  $c_{ij}$  by using the recurrence formulas (3.17) and (3.19). First, since  $c_{11} = 1$  by (b.1) in the proof of Theorem 3.1, it follows from (3.19) with k = 1 and  $0 < \varepsilon < \frac{1}{12}$  that

$$c_{21} \ge \frac{6}{11} \left( 4 + 2\sqrt{2} + \frac{1}{12} \right) \ge \frac{1}{2(1 - c_{11}\varepsilon)} \left( 4 + 2\sqrt{2} + \varepsilon \right),$$

and we can choose  $c_{21} = 4$  as the smallest positive integer satisfying the last inequality.

Now, we use (3.17) with j = 2 to obtain

$$(3c_{22} - 1)(c_{22} - 1) \ge c_{21}^2$$

and thus, we select  $c_{22} = 3$ . We note that  $c_{22} = 3$  is larger than the estimate in the proof of Theorem 3.1. This difference is due to the use of the formula (3.17) instead of (3.16).

By (3.19) with k = 2, we get

$$c_{32} \ge \frac{2}{3} \left( 4 + 2\sqrt{2} + \frac{33}{12} \right) \ge \frac{1}{2(1 - c_{22}\varepsilon)} \left( 4 + 2\sqrt{2} + \varepsilon + 2c_{21}^2\varepsilon \right)$$

and hence, we choose  $c_{32} = 7$ . Further, it follows from (3.17) with j = 3 that

$$(3c_{33} - 1)(c_{33} - 1) \ge c_{21}^2 + c_{32}^2 = 65$$

and hence,  $c_{33} = 6$ . In the proof of Theorem 3.1, we estimated  $c_{32} = 6$  and  $c_{33} = 4$ , but in this example, we estimate the larger values for them because we use the formula (3.17) instead of (3.16).

As n = 4, we see that  $\sigma \geq \sum_{i=1}^{3} c_{(i+1)i}^2 > c_{21}^2 + c_{32}^2 = 65$  and hence,  $\varepsilon < \frac{1}{\sigma} < \frac{1}{65}$ . Therefore, we

$$\frac{1}{2(1-c_{33}\varepsilon)}\left(4+2\sqrt{2}+\varepsilon+2\sum_{i=1}^{2}c_{3i}c_{4i}\varepsilon\right) < \frac{65}{118}\left(4+2\sqrt{2}+\frac{1}{65}+2\right) < 4.88$$

and thus, using (3.19) with k = 3, we can select  $c_{43} = 5$ . In view of (3.17) with j = 4, we get

$$(3c_{44} - 1)(c_{44} - 1) \ge c_{21}^2 + c_{32}^2 + c_{43}^2 = 90$$

and hence,  $c_{44} = 7$ , which comply with the claims of Remark 3.1.

**Example 5.2** We assume that n = 5. As in Example 5.1 we get the following constants:  $c_{11} = 1$ ,  $c_{21} = 4, c_{22} = 3, c_{31} = 4, c_{32} = 7, c_{33} = 6, c_{41} = 4, c_{42} = 7, c_{43} = 5, and c_{44} = 7.$ 

As n = 5, we see that  $\sigma \geq \sum_{i=1}^{4} c_{(i+1)i}^2 > c_{21}^2 + c_{32}^2 + c_{43}^2 = 90$  and hence,  $\varepsilon < \frac{1}{\sigma} < \frac{1}{90}$ . Thus, we

$$\frac{1}{2(1-c_{44}\varepsilon)}\left(4+2\sqrt{2}+\varepsilon+2\sum_{i=1}^{3}c_{4i}c_{5i}\varepsilon\right) < \frac{45}{83}\left(4+2\sqrt{2}+\frac{1}{90}+2\right) < 4.80$$

and hence, using (3.19) with k = 4, we can select  $c_{54} = 5$ . In view of (3.17) with j = 5, we get

$$(3c_{55}-1)(c_{55}-1) \ge c_{21}^2 + c_{32}^2 + c_{43}^2 + c_{54}^2 = 115$$
 and hence  $c_{55} = 7$ .

Moreover, due to (3.15), we have  $c_{51} = 4$ ,  $c_{52} = 7$ , and  $c_{53} = 5$ .

**Example 5.3** Let  $D = \{x \in \mathbb{R}^4 : ||x|| \leq d\}$  for some  $d \geq 1$  and let  $f : D \to \mathbb{R}^4$  be a function satisfying f(0) = 0 and the inequality (3.1) for all  $x, y \in D$  and some constant  $\varepsilon$  with  $0 < \varepsilon < \frac{1}{90}$ . Using Theorem 4.1 and Example 5.1, we can prove that there exists an isometry  $U : D \to \mathbb{R}^4$  satisfying

$$\begin{split} \|f(x) - U(x)\| &\leq \left[\sum_{i=1}^{4} \left( \left(2 + \sum_{j=1}^{i} c_{ij}\right) d + 4 + \sum_{j=1}^{i} c_{ij}\right)^{2} \right]^{1/2} \varepsilon \\ &= \sqrt{1076d^{2} + 2376d + 1316} \varepsilon \\ &< (33d + 37)\varepsilon \end{split}$$

for all  $x \in D$ .

**Example 5.4** Let  $D = \{x \in \mathbb{R}^5 : ||x|| \leq d\}$  for some  $d \geq 1$  and let  $f : D \to \mathbb{R}^5$  be a function satisfying f(0) = 0 and the inequality (3.1) for all  $x, y \in D$  and some constant  $\varepsilon$  with  $0 < \varepsilon < \frac{1}{115}$ . Using Theorem 4.1 and Example 5.2, we can prove that there exists an isometry  $U : D \to \mathbb{R}^5$  satisfying

$$\|f(x) - U(x)\| \le \left[\sum_{i=1}^{5} \left( \left(2 + \sum_{j=1}^{i} c_{ij}\right) d + 4 + \sum_{j=1}^{i} c_{ij}\right)^{2} \right]^{1/2} \varepsilon$$
$$= \sqrt{1976d^{2} + 4296d + 2340} \varepsilon$$
$$< (45d + 49)\varepsilon$$

for all  $x \in D$ .

# 6 Discussion

We expect the Hyers-Ulam stability of isometries defined on bounded domains to have widespread application, but no remarkable results have been published, except for Väisälä's paper [31], for 40 years after Fickett's theorem was published: A finite sequence  $(u_0, u_1, \ldots, u_m)$  of a compact subset D of  $\mathbb{R}^n$  is called a maximal sequence in D provided that  $h_k := d(u_k, A_{k-1})$  is maximal in D for each  $k \in \{1, 2, \ldots, m\}$ , where  $A_{k-1}$  is the affine subspace (or flat) including  $\{u_0, u_1, \ldots, u_{k-1}\}$ . Now, among the results proved by Väisälä, the part related to the subject of this paper is introduced in the following theorem.

**Theorem 6.1 (Väisälä)** Let D be a compact subset of  $\mathbb{R}^n$ . If there exist a maximal sequence  $(u_0, u_1, \ldots, u_n)$  in D and a constant  $K \ge 1$  such that

- (i)  $|u_k u_0| \le Kh_k$  for every  $k \in \{2, 3, \dots, n\}$ ;
- (*ii*)  $D \setminus \{u_1, \ldots, u_n\} \subset B_{Kh_n}(u_0),$

then for every  $\varepsilon$ -isometry  $f: D \to \mathbb{R}^n$ , there exist an isometry  $U: \mathbb{R}^n \to \mathbb{R}^n$  and a constant  $K^* > 0$ such that

$$\|f(x) - U(x)\| \le K^* \varepsilon$$

for all  $x \in D$ .

Now we have made great progress in improving Fickett's theorem in practice using intuitive method in this paper. As we can see, our main theorem (Theorem 4.1) looks much simpler and clearer than Theorem 6.1 proved by Väisälä. According to Theorem 4.1 of this paper, if a function  $f: D \to \mathbb{R}^n$  satisfies f(0) = 0 as well as the inequality (3.1) for all  $x, y \in D$  and for some sufficiently small constant  $\varepsilon > 0$ , then there exist an isometry  $U: D \to \mathbb{R}^n$  and a constant K > 0 such that the inequality  $||f(x) - U(x)|| \leq K\varepsilon$  holds for all  $x \in D$ . However, it is impossible to deduce this useful conclusion by using Fickett's theorem. From this point of view, we can say that Fickett's theorem has been remarkably improved in this paper.

# References

- [1] H. Anton, Elementary Linear Algebra (9th ed.), John Wiley & Sons, 2005.
- [2] J. A. Baker, Isometries in normed spaces, Amer. Math. Monthly 78 (1971), no. 6, 655–658.
- [3] D. G. Bourgin, Approximate isometries, Bull. Amer. Math. Soc. 52 (1946), 704–714.
- [4] R. Bhatia and P. Šemrl, Approximate isometries on Euclidean spaces, Amer. Math. Monthly 104 (1997), 497–504.
- [5] D. G. Bourgin, Approximately isometric and multiplicative transformations on continuous function rings, Duke Math. J. 16 (1949), 385–397.
- [6] R. D. Bourgin, Approximate isometries on finite dimensional Banach spaces, Trans. Amer. Math. Soc. 207 (1975), 309–328.
- [7] J. Chmielinski and S.-M. Jung, The stability of the Wigner equation on a restricted domain, J. Math. Anal. Appl. 254 (2001), 309–320.
- [8] G. Choi and S.-M. Jung, The stability of isometries on restricted domains, Symmetry 13 (2021), 282, 11 pages.
- [9] G. Dolinar, Generalized stability of isometries, J. Math. Anal. Appl. 242 (2000), 39–56.
- [10] J. W. Fickett, Approximate isometries on bounded sets with an application to measure theory, Studia Math. 72 (1981), 37–46.
- [11] J. Gevirtz, Stability of isometries on Banach spaces, Proc. Amer. Math. Soc. 89 (1983), 633–636.
- [12] P. M. Gruber, *Stability of isometries*, Trans. Amer. Math. Soc. **245** (1978), 263–277.

- [13] D. H. Hyers, On the stability of the linear functional equation, Proc. Natl. Acad. Sci. U.S.A. 27 (1941), 222–224.
- [14] D. H. Hyers and S. M. Ulam, On approximate isometries, Bull. Amer. Math. Soc. 51 (1945), 288–292.
- [15] D. H. Hyers and S. M. Ulam, Approximate isometries of the space of continuous functions, Ann. Math. 48 (1947), 285–289.
- [16] S.-M. Jung, Hyers-Ulam-Rassias stability of isometries on restricted domains, Nonlinear Stud. 8 (2001), 125–134.
- [17] S.-M. Jung, Asymptotic properties of isometries, J. Math. Anal. Appl. 276 (2002), 642–653.
- [18] S.-M. Jung and B. Kim, Stability of isometries on restricted domains, J. Korean Math. Soc. 37 (2000), no. 1, 125–137.
- [19] J. Lindenstrauss and A. Szankowski, Non linear perturbations of isometries, Astérisque 131 (1985), 357–371.
- [20] M. Omladič and P. Šemrl, On non linear perturbations of isometries, Math. Ann. 303 (1995), 617-628.
- [21] J. M. Rassias, On a new approximation of approximately linear mappings by linear mappings, Discuss. Math. 7 (1985), 193–196.
- [22] Th. M. Rassias, Properties of isometric mappings, J. Math. Anal. Appl. 235 (1999), 108–121.
- [23] Th. M. Rassias, Isometries and approximate isometries, Internat. J. Math. Math. Sci. 25 (2001), 73–91.
- [24] Th. M. Rassias and C. S. Sharma, Properties of isometries, J. Natural Geom. 3 (1993), 1–38.
- [25] P. Semrl, Hyers-Ulam stability of isometries on Banach spaces, Aequationes Math. 58 (1999), 157–162.
- [26] F. Skof, Sulle  $\delta$ -isometrie negli spazi normati, Rendiconti Di Mat. Ser. VII, Roma 10 (1990), 853–866.
- [27] F. Skof, On asymptotically isometric operators in normed spaces, Ist. Lomb. Sc. e Lett., A 131 (1997), 117–129.
- [28] G. W. Stewart and J.-G. Sun, *Matrix Perturbation Theory*, Academic Press, New York, 1990.
- [29] R. L. Swain, Approximate isometries in bounded spaces, Proc. Amer. Math. Soc. 2 (1951), 727–729.
- [30] S. M. Ulam, A Collection of Mathematical Problems, Interscience, New York, 1960.
- [31] J. Väisälä, Isometric approximation property in Euclidean spaces, Israel J. Math. 128 (2002), 1–27.
- [32] E. P. Wigner, Gruppentheorie und ihre Anwendungen auf die Quantenmechanik der Atomspektren, Friedr. Vieweg und Sohn, Braunschweig, 1931.