ON ROW-FACTORIZATION MATRICES AND GENERIC IDEALS

OM PRAKASH BHARDWAJ, KRITI GOEL, AND INDRANATH SENGUPTA

ABSTRACT. Let H be a numerical semigroup minimally generated by an almost arithmetic sequence. We give a complete description of the row-factorization (RF) matrices associated with the pseudo-Frobenius elements of H. We use the information from RF-matrices to give a characterization of the generic nature of the defining ideal. We also show that the defining ideal of gluing of two numerical semigroups is never generic. Further, when H is symmetric and has embedding dimension 4 or 5, we prove that the defining ideal is minimally generated by RF-relations.

1. Introduction

Numerical semigroups and semigroup rings are widely studied objects in the literature (see [12]). Recently, A. Moscariello [6] introduced a new tool associated with the pseudo-Frobenius elements of numerical semigroups, namely, row-factorization (RF) matrices, to explore the properties of semigroup rings. Moscariello used this object to investigate the type of almost symmetric semigroups of embedding dimension four and proved the conjecture given by T. Numata in [7], which states that the type of an almost symmetric semigroup of embedding dimension four is at most three. The connection between the RF-matrices and numerical semigroup rings is further studied by K. Eto, J. Herzog and K.-i. Watanabe in ([3], [2], [1], [5]), where they explore the set theoretic complete intersection property, minimal free resolution, generic property, etc for the defining ideal of certain monomial curves.

Let k be a field and $H = \langle m_0, m_1, \ldots, m_p \rangle$ be a numerical semigroup minimally generated by $\{m_0, m_1, \ldots, m_p\}$. For an indeterminate t over k, the semigroup ring of H is a subring of k[t], defined as $k[H] = k[t^{m_0}, \ldots, t^{m_p}]$. The ring k[H] can be represented as a quotient of a polynomial ring using a canonical surjection $\pi : k[x_0, x_1, \ldots, x_p] \to k[H]$ given by $\pi(x_i) = t^{m_i}$, for all i. The kernel of this k-algebra homomorphism π , say I(H), is a prime ideal (called toric ideal) and is the defining ideal of k[H]. The ring k[H] is called a toric ring.

If a toric ideal has a minimal generating set consisting of binomials with full support, then it is called generic. The notion of genericity for lattice ideals, and in particular for toric ideals, is introduced by I. Peeva and B. Sturmfels in [11]. The authors give a characteristic independent, minimal free resolution of the toric ring when the toric ideal is generic. Generic toric rings are further studied in [4], where the authors prove that such rings are Golod and so the Poincaré series of the residue field is rational. In [3], K. Eto gave necessary and sufficient conditions for a toric ideal to be generic and as a result proved that the defining

The second author is supported by the Early Career Fellowship at IIT Gandhinagar.

The third author is the corresponding author; supported by the MATRICS research grant MTR/2018/000420, sponsored by the SERB, Government of India.

²⁰¹⁰ Mathematics Subject Classification: 20M14, 20M25, 13A02.

Keywords: Almost arithmetic numerical semigroup, pseudo-Frobenius elements, row-factorization matrix, generic toric ideal.

ideal of almost Gorenstein monomial curves is not generic if their embedding dimension is greater than three.

In this paper, we consider the numerical semigroups generated by an almost arithmetic sequence. In particular, we have $H = \langle m_0, m_1, \ldots, m_p, n \rangle$ such that $m_i = m_0 + id$ for $i = 0, \ldots, p, d > 0$ and $gcd(m_0, n, d) = 1$. The aim of this paper is to characterize the generic nature of the defining ideal of the semigroup H using the information from the row-factorization matrices. This, in the first place, requires knowledge of the pseudo-Frobenius elements of H. For this, we refer to the computations in [8], where the authors give an explicit description of pseudo-Frobenious elements for the numerical semigroups of this class.

We now summarize the contents of the paper. In Section 2, we recall some basic definitions, along with the definition of RF-matrices. We also recall some results from the papers [8] and [10], which are used in the further sections. In Sections 3 and 4, we give the computation of RF-matrices. Section 5 contains the main result of the paper (Theorem 5.3), where we characterize the generic nature of the defining ideal of a numerical semigroup minimally generated by an almost arithmetic sequence. We conclude that I(H) is never generic, except when p=0 or under suitable conditions with p=1. We use Eto's result ([3, Theorem 1]) to give this characterization. Using Eto's result, we also prove that the defining ideal of a gluing of two numerical semigroups is never generic. Consequently, defining ideal of the semigroup ring defined by a complete intersection numerical semigroup of embedding dimension at least 3 is never generic.

A numerical semigroup H of type 1 is called symmetric. It is known that k[H] is Gorenstein if and only if H is a symmetric numerical semigroup. In section 6, we identify the cases in which a numerical semigroup H, minimally generated by an almost arithmetic sequence, is symmetric. We compute the RF-matrices in each of these cases. The rows of RF-matrices are known to produce binomials in the ideal I(H). These binomials are called RF-relations. Since I(H) may not always be minimally generated by RF-relations, one may ask when is such a scenario possible. This question was raised by Herzog and Watanabe in [5]. They answered the question in the affirmative for the cases when the embedding dimension is 3 and when H is pseudo-symmetric or almost symmetric with embedding dimension 4. In Theorem 6.3 and Theorem 6.4, we prove that I(H) is minimally generated by RF-relations when H is a symmetric numerical semigroup, minimally generated by an almost arithmetic sequence with embedding dimension 4 or 5.

2. Notation and Preliminaries

Let \mathbb{Z} and \mathbb{N} be the sets of integers and non-negative integers respectively. Let p > 0 and $m_0, m_1, \ldots, m_p \in \mathbb{N}$ such that $\gcd(m_0, m_1, \ldots, m_p) = 1$. Then

$$\langle m_0, m_1, \dots, m_p \rangle = \left\{ \sum_{i=0}^p a_i m_i \mid a_i \in \mathbb{N}, \forall i \right\}$$

is called the numerical semigroup generated by m_0, m_1, \ldots, m_p .

Definition 2.1. Let H be a numerical semigroup generated by a sequence m_0, m_1, \ldots, m_p of positive integers. Then the set $Ap(H, m_0) = \{h \in H \mid h - m_0 \notin H\}$ is called the Apéry set of H with respect to m_0 .

Definition 2.2. Let H be a numerical semigroup. An element $f \in \mathbb{Z} \setminus H$ is called a pseudo-Frobenius number if $f+h \in H$ for all $h \in H \setminus \{0\}$. We will denote the set of pseudo-Frobenius numbers of H by PF(H).

Note that if $f \in PF(H)$, then $f + m_i \in H$ for all $i \in [0, p]$. Hence, for all $i \in [0, p]$, there exist $a_{i0}, a_{i1}, \ldots, a_{ip} \in \mathbb{N}$ such that

$$f + m_i = \sum_{j=0}^p a_{ij} m_j.$$

In the above expression, $a_{ii} = 0$. Because if $a_{ii} > 0$, then it would imply $f \in H$. We now recall the notion of row-factorization matrix (RF-matrix) given by A. Moscariello in [6].

Definition 2.3. Let H be a numerical semigroup generated by m_0, \ldots, m_p and $f \in PF(H)$. We say that $A = (a_{ij}) \in M_{p+1}(\mathbb{Z})$ is an RF-matrix for f if for all $i \in [0, p]$,

$$\sum_{j=0}^{p} a_{ij} m_j = f,$$

where $a_{ii} = -1$ for all $i \in [0, p]$, and $a_{ij} \in \mathbb{N}$ for all $i, j \in [0, p]$ and $i \neq j$. For $f \in PF(H)$, we will denote an RF-matrix of f by RF(f).

Observe that for a given $f \in PF(H)$, RF(f) may not be not unique. A factorization of $f + m_i$ gives the $(i + 1)^{th}$ row of an RF-matrix for f. Thus, if $f + m_i$ does not have unique factorization in H, then the matrix RF(f) is not unique. Nevertheless, RF(f) will be the notation for one of the possible RF-matrices of f. If H is symmetric, i.e. type(H) = |PF(H)| = 1, then RF(H) denotes an RF-matrix of the only pseudo-Frobenius number of H.

Assume that the sequence m_0, m_1, \ldots, m_p is a strictly increasing arithmetic sequence of positive integers and n is a positive integer such that $gcd(m_0, \ldots, m_p, n) = 1$. Also, assume that $\{m_0, \ldots, m_p, n\}$ is a minimal generating set for the numerical semigroup H. The following result was proved by D.P. Patil and B. Singh in [10].

Lemma 2.4 ([10, Lemma 3.1, 3.2]). Let d be a positive integer such that $m_i = m_0 + id$ for all $0 \le i \le p$. Let n be an arbitrary positive integer with $gcd(m_0, d, n) = 1$. Let $H' = \sum_{i=0}^p \mathbb{N}m_i$ and $H = H' + \mathbb{N}n$. For $t \in \mathbb{N}$, let $q_t \in \mathbb{Z}$, $r_t \in [1, p]$ and $g_t \in H'$ defined by $t = q_t p + r_t$ and $g_t = q_t m_p + m_{r_t}$.

- (1) Let $u = \min\{t \in \mathbb{N} \mid g_t \notin \operatorname{Ap}(H, m_0)\}\$ and $v = \min\{b \geq 1 \mid bn \in H'\}$. Then there exist unique integers $z \in [0, u 1], w \in [0, v 1], \lambda \geq 1$, and $\mu \geq 0$ such that
 - (i) $g_u = \lambda m_0 + wn;$
 - (ii) $vn = \mu m_0 + g_z$;
 - (iii) $g_{u-z} + (v-w)n = \nu m_0$. Moreover, $\nu = \lambda + \mu + \epsilon$ where $\epsilon = 1$ or 0 according as $r_{u-z} < r_u$ or $r_{u-z} \ge r_u$.
- (2) $g_s + g_t = \epsilon m_0 + g_{s+t}$ with $\epsilon = 1$ or 0 according as $r_s + r_t \le p$ or $r_s + r_t > p$.

In addition to the above notation, set $r = r_u, r' = r_{u-z}, q = q_u, q' = q_{u-z}$, and $W = [u-z, u-1] \times [v-w, v-1]$. Note that if w = 0 or z = 0, then W is empty.

Lemma 2.5 ([8, Lemma 2.4]). With the above notation,

(1) $u \ge p + 1$, $q \ge 1$ and $q' \in [0, q]$.

(2)
$$r_z = \begin{cases} r - r' & \text{if } r > r', \\ p + r - r' & \text{if } r \le r'. \end{cases}$$

- (3) if q' = q and $z \neq 0$, then r' < r and $r \geq 2$.
- (4) if $\mu = 0$, then z > p and q' < q.
- (5) if $\lambda = 1$, then $w \neq 0$.
- (6) if $\lambda = 1$ and $\mu = 0$, then r' < r and $r \ge 2$.

In this article, we aim to describe the structure of RF-matrices for pseudo-Frobenius numbers of a numerical semigroup H defined by an almost arithmetic sequence, using the above results. We discuss the cases $W = \emptyset$ and $W \neq \emptyset$ separately. We use the above results throughout the paper without repeated reference.

3. The case
$$W = \emptyset$$

Throughout this section we will assume that $W = \emptyset$ and $H = \langle m_0, m_1, \dots, m_p, n \rangle$; where $m_i = m_0 + id$ for $i \in [1, p]$ and $\gcd(m_0, n, d) = 1$.

Let $\gamma_k = g_{(q-1)p+r+k-1} + (v-1)n - m_0$, for $k \in [1, p]$. In this case, using [8, Proposition 3.3], it follows that the set of pseudo-Frobenius numbers is

$$PF(H) = \begin{cases} \{ \gamma_k \mid k \in [1, p] \} & \text{if } r = 1, \\ \{ \gamma_k \mid k \in [p - r + 2, p] \} & \text{if } r \ge 2. \end{cases}$$

We find the factorizations of $\gamma_k + m_j$ for $j \in [0, p]$, and $\gamma_k + n$ in H. From these factorizations, we form an RF-matrix for γ_k .

Proposition 3.1. Let r = 1 and $\lambda > 1$. Then for every element γ_k of PF(H), $k \in [1, p]$, there exists an RF-matrix of the following type

$$RF(\gamma_k) = \begin{bmatrix} -1 & 0 & 0 & \cdots & \cdots & 0 & q-1 & v-1 \\ 0 & -1 & 0 & \cdots & \cdots & 0 & q-1 & v-1 \\ \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & q-1 & v-1 \\ \lambda-2 & 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & w+v-1 \\ \vdots & \vdots \\ \lambda-2 & 0 & 0 & \cdots & \cdots & 0 & -1 & w+v-1 \\ a_{p+2,1} & a_{p+2,2} & a_{p+2,3} & \cdots & \cdots & \cdots & a_{p+2,p+1} & -1 \end{bmatrix} + \begin{bmatrix} R_{(0)} \\ R_{(1)} \\ \vdots \\ R_{(p-k)} \\ R_{(p-k+1)} \\ \vdots \\ R_{(p)} \\ R_{(n)} \end{bmatrix},$$

where

- (1) $R_{(0)} = (a^{(0)}, \dots, a^{(k)}, \dots, a^{(n)})$ such that $a^{(k)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (2) For all $j \in [1, p k]$, $R_{(j)} = (a^{(0)}, \dots, a^{(k+j)}, \dots, a^{(n)})$ such that $a^{(k+j)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (3) For all $j \in [p-k+1, p]$, $R_{(j)} = (a^{(0)}, \dots, a^{(k+j-p-1)}, \dots, a^{(n)})$ such that $a^{(k+j-p-1)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (4) If $\mu \geq 1$, then $(a_{p+2,1}, a_{p+2,2}, \dots, a_{p+2,p+1}, -1) = (\mu 1, 0, \dots, 0, q + q_z 1, -1)$ and $R_{(n)} = (a^{(0)}, \dots, a^{(k)}, \dots, a^{(r_z)}, \dots, a^{(n)})$ such that $a^{(k)} = a^{(r_z)} = 1$ and all other $a^{(j)}$'s equal to zero. Otherwise, when $\mu = 0$, $(a_{p+2,1}, a_{p+2,2}, \dots, a_{p+2,p+1}, -1) = (\lambda 2, 0, \dots, 0, q_z 1, -1)$ and $R_{(n)} = (a^{(0)}, \dots, a^{(k-1)}, \dots, a^{(r_z)}, \dots, a^{(n)})$ such that $a^{(k-1)} = a^{(r_z)} = 1$ and all other $a^{(j)}$'s equal to zero.

Proof. Since
$$r = 1$$
, $PF(H) = \{ \gamma_k \mid k \in [1, p] \}$ where $\gamma_k = g_{(q-1)p+k} + (v-1)n - m_0$.
(1) $\gamma_k + m_0 = g_{(q-1)p+k} + (v-1)n = (q-1)m_p + m_k + (v-1)n$.

(2) If $j \in [1, p - k]$, then

$$\gamma_k + m_j = g_{(q-1)p+k} + (v-1)n - m_0 + m_j = g_{(q-1)p+k} + g_j + (v-1)n - m_0$$
$$= m_0 + g_{(q-1)p+k+j} + (v-1)n - m_0$$
$$= (q-1)m_p + m_{k+j} + (v-1)n.$$

(3) If j = p - k + 1, then

$$\gamma_k + m_j = g_{(q-1)p+k} + (v-1)n - m_0 + m_{p-k+1}$$

$$= (q-1)m_p + m_k + (v-1)n - m_0 + m_{p-k+1}$$

$$= (q-1)m_p + (v-1)n - m_0 + m_p + m_1$$

$$= g_u + (v-1)n - m_0 = (\lambda - 1)m_0 + (w+v-1)n.$$

If j > p - k + 1 and $\lambda > 1$, then

$$\gamma_k + m_j = g_{(q-1)p+k} + (v-1)n - m_0 + m_j = (q-1)m_p + m_k + (v-1)n - m_0 + m_j$$

$$= qm_p + (v-1)n - m_0 + m_{k+j-p}$$

$$= g_u + (v-1)n - 2m_0 + m_{k+j-p-1}$$

$$= (\lambda - 2)m_0 + m_{k+j-p-1} + (w+v-1)n.$$

(4) If $\mu > 0$, then

$$\gamma_k + n = g_{(q-1)p+k} + vn - m_0 = g_{(q-1)p+k} + (\mu - 1)m_0 + g_z$$

$$= (q-1)m_p + m_k + (\mu - 1)m_0 + q_z m_p + m_{r_z}$$

$$= (\mu - 1)m_0 + m_k + m_{r_z} + (q + q_z - 1)m_p.$$

If $\mu = 0$, then z > p. This implies that w = 0, $q_{z-p} = q_z - 1$ and $r_{z-p} = r_z$. Thus

$$\begin{split} \gamma_k + n &= g_{(q-1)p+k} + vn - m_0 = g_{(q-1)p+k} - m_0 + g_z \\ &= g_{(q-1)p+k} - m_0 + g_p + g_{z-p} \\ &= g_{qp+k} - m_0 + g_{z-p} \\ &= qm_p + m_k - m_0 + g_{z-p} \\ &= g_u + m_{k-1} - 2m_0 + g_{z-p} \\ &= (\lambda - 2)m_0 + m_{k-1} + (q_z - 1)m_p + m_{r_z}. \end{split}$$

Proposition 3.2. Let r=1 and $\lambda=1$. Then for every element γ_k of PF(H), $k\in[1,p]$, there exists an RF-matrix of the following type

$$RF(\gamma_k) = \begin{bmatrix} -1 & 0 & 0 & \cdots & \cdots & 0 & q-1 & v-1 \\ 0 & -1 & 0 & \cdots & \cdots & 0 & q-1 & v-1 \\ \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & q-1 & v-1 \\ \mu-1 & 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & w-1 \\ \vdots & \vdots \\ \mu-1 & 0 & 0 & \cdots & \cdots & 0 & -1 & w-1 \\ \mu-1 & 0 & 0 & \cdots & \cdots & \cdots & q-1 & -1 \end{bmatrix} + \begin{bmatrix} R_{(0)} \\ R_{(1)} \\ \vdots \\ R_{(p-k)} \\ R_{(p-k+1)} \\ \vdots \\ R_{(p)} \\ R_{(n)} \end{bmatrix},$$

where

- (1) $R_{(0)} = (a^{(0)}, \ldots, a^{(k)}, \ldots, a^{(n)})$ such that $a^{(k)} = 1$ and all other $a^{(j)}$'s equal to zero. (2) For all $j \in [1, p k]$, $R_{(j)} = (a^{(0)}, \ldots, a^{(k+j)}, \ldots, a^{(n)})$ such that $a^{(k+j)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (3) For all $j \in [p-k+1,p]$, $R_{(j)} = (a^{(0)}, \dots, a^{(k+j-p-1)}, \dots, a^{(n)})$ having $a^{(k+j-p-1)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (4) $R_{(n)} = (a^{(0)}, \dots, a^{(k)}, \dots, a^{(n)})$ such that $a^{(k)} = 1$ and all other $a^{(j)}$'s equal to zero.

Proof. For (1) and (2), see the proof of Proposition 3.1.

(3) Since $\lambda = 1$ and r = 1, using lemma 2.5 we get $w, \mu \neq 0$. Also, $W = \emptyset$ implies that z=0 and hence $\nu=\mu+1$. Put $\ell=k+j-p$. Since $k+j\geq p+1$, we get $\ell\in[1,p]$ and

$$\gamma_k + m_j = g_{(q-1)p+k} + (v-1)n - m_0 + m_j = g_{(u+l-1)} + (v-1)n - m_0$$

$$= g_{(u+l-1)} + \nu m_0 - g_u + (w-1)n - m_0$$

$$= (\nu - 2)m_0 + m_{\ell-1} + (w-1)n$$

$$= (\mu - 1)m_0 + m_{k+j-n-1} + (w-1)n.$$

(4) Since z=0, we have $g_z=0$ and therefore,

$$\gamma_k + n = g_{(q-1)p+k} + vn - m_0 = g_{(q-1)p+k} + (\mu - 1)m_0$$
$$= (q-1)m_p + m_k + (\mu - 1)m_0.$$

Corollary 3.3. Suppose p = 1. Then $PF(H) = \{\gamma_1\}$ and there exists an RF-matrix of γ_1 of either of the following type

$$\begin{bmatrix} -1 & u-1 & v-1 \\ \lambda-1 & -1 & w+v-1 \\ \mu-1 & u+z-1 & -1 \end{bmatrix}, \begin{bmatrix} -1 & u-1 & v-1 \\ \lambda-1 & -1 & w+v-1 \\ \lambda-1 & z-1 & -1 \end{bmatrix}, or \begin{bmatrix} -1 & u-1 & v-1 \\ \mu & -1 & w-1 \\ \mu-1 & u-1 & -1 \end{bmatrix}.$$

Proof. Since p=1, we get $r_t=1$, $q_t=t-1$ for all t and $PF(H)=\{\gamma_1\}$. Therefore, we have k=1. Let $\lambda>1$. Using Proposition 3.1, we have

$$\gamma_1 + m_0 = g_q + (v - 1)n = (u - 1)m_1 + (v - 1)n,$$

$$\gamma_1 + m_1 = (\lambda - 1)m_0 + (w + v - 1)n.$$

If $\mu > 0$, then

$$\gamma_1 + n = (\mu - 1)m_0 + (q + q_z + 1)m_1 = (\mu - 1)m_0 + (u + z - 1)m_1.$$

If
$$\mu = 0$$
, then $\gamma_1 + n = (\lambda - 1)m_0 + (q_{z-1} + 1)m_1 = (\lambda - 1)m_0 + (z - 1)m_1$.

Let $\lambda = 1$. Using Proposition 3.2, we get

$$\gamma_1 + m_0 = (u - 1)m_1 + (v - 1)n,$$

$$\gamma_1 + m_1 = \mu m_0 + (w - 1)n,$$

$$\gamma_1 + n = (\mu - 1)m_0 + qm_1.$$

Thus, we have the above possible three matrices.

Proposition 3.4. Let $r \geq 2$. Then for every element γ_k of PF(H), $k \in [p-r+2, p]$, there exists an RF-matrix of the following type

$$RF(\gamma_k) = \begin{bmatrix} -1 & 0 & 0 & \cdots & \cdots & 0 & q & v-1 \\ 0 & -1 & 0 & \cdots & \cdots & 0 & q & v-1 \\ \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & q & v-1 \\ \lambda - 1 & 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & w+v-1 \\ \vdots & \vdots \\ \lambda - 1 & 0 & 0 & \cdots & \cdots & 0 & -1 & w+v-1 \\ a_{p+2,1} & a_{p+2,2} & a_{p+2,3} & \cdots & \cdots & \cdots & a_{p+2,p+1} & -1 \end{bmatrix} + \begin{bmatrix} R_{(0)} \\ R_{(1)} \\ \vdots \\ R_{(e)} \\ R_{(e+1)} \\ \vdots \\ R_{(p)} \\ R_{(n)} \end{bmatrix},$$

where $e \in [1, p-1]$ such that r+k+e=2p+1 and

- (1) $R_{(0)} = (a^{(0)}, \ldots, a^{(p-e)}, \ldots, a^{(n)})$ such that $a^{(p-e)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (2) For all $j \in [1, e]$, $R_{(j)} = (a^{(0)}, \dots, a^{(p-e+j)}, \dots, a^{(n)})$ such that $a^{(p-e+j)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (3) For all $j \in [e+1, p]$, $R_{(j)} = (a^{(0)}, \dots, a^{(k+j-p-1)}, \dots, a^{(n)})$ such that $a^{(k+j-p-1)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (4) If $\mu \geq 1$ then $(a_{p+2,1}, a_{p+2,2}, \dots, a_{p+2,p+1}, -1) = (\mu 1, 0, \dots, 0, q + q_z, -1)$ and $R_{(n)} = (a^{(0)}, \dots, a^{(p-e)}, \dots, a^{(r_z)}, \dots, a^{(n)})$ such that $a^{(p-e)} = a^{(r_z)} = 1$ and all other $a^{(j)}$'s equal to zero. Otherwise, when $\mu = 0$ then $(a_{p+2,1}, a_{p+2,2}, \dots, a_{p+2,p+1}, -1) = (\lambda 1, 0, \dots, 0, q_z 1, -1)$ and $R_{(n)} = (a^{(0)}, \dots, a^{(k-1)}, \dots, a^{(r_z)}, \dots, a^{(n)})$ such that $a^{(k-1)} = a^{(r_z)} = 1$ and all other $a^{(j)}$'s equal to zero.

Proof. Since $r \geq 2$, $PF(H) = \{\gamma_k \mid k \in [p-r+2, p]\}$. Therefore,

(1)
$$\gamma_k + m_0 = g_{(q-1)p+r+k-1} + (v-1)n = qm_p + m_{p-e} + (v-1)n$$
.

(2) If
$$j \in [1, e]$$
, then $r + k + j - p - 1 \le p$ and

$$\gamma_k + m_j = g_{(q-1)p+r+k-1} + (v-1)n - m_0 + m_j = m_0 + qm_p + m_{p-e+j} + (v-1)n - m_0$$
$$= qm_p + m_{p-e+j} + (v-1)n.$$

(3) If
$$j \in [e+1, p]$$
, then $r + k + j - p - 1 > p$ and

$$\gamma_k + m_j = g_{(q-1)p+r+k-1} + (v-1)n - m_0 + m_j = g_{u-p+j+k-1} - (v-1)n - m_0$$

$$= g_u - m_0 + g_{j+k-p-1} - (v-1)n$$

$$= (\lambda - 1)m_0 + wn + m_{k+j-p-1} + (v-1)n$$

$$= (\lambda - 1)m_0 + m_{k+j-p-1} + (w+v-1)n.$$

(4) If $\mu \geq 1$, then

$$\begin{split} \gamma_k + n &= g_{(q-1)p+r+k-1} + (v-1)n - m_0 + n = qm_p + m_{p-e} + (\mu - 1)m_0 + g_z \\ &= qm_p + m_{p-e} + (\mu - 1)m_0 + q_z m_p + m_{rz} \\ &= (\mu - 1)m_0 + m_{p-e} + m_{rz} + (q + q_z)m_p. \end{split}$$

If $\mu = 0$, then z > p. This implies that w = 0, $q_{z-p} = q_z - 1$ and $r_{z-p} = r_z$. Thus

$$\gamma_k + n = g_{(q-1)p+r+k-1} + (v-1)n - m_0 + n = g_{u-p+k-1} + vn - m_0$$

$$= g_u + g_{z-p} + g_{k-1} - m_0$$

$$= (\lambda - 1)m_0 + m_{k-1} + g_{z-p}$$

$$= (\lambda - 1)m_0 + m_{k-1} + (q_z - 1)m_p + m_{r_z}.$$

Example 3.5. Let $H = \langle m_0 = 14, m_1 = 17, m_2 = 20, m_3 = 23, m_4 = 26, n = 21 \rangle$. Then $H' = \langle 14, 17, 20, 23, 26 \rangle$ and p = 4. The Apéry set of H with respect to 14 is

$$Ap(H, 14) = \{0, 17, 20, 21, 23, 26, 38, 41, 43, 44, 46, 47, 64, 67\}.$$

The smallest i such that $g_i \notin \text{Ap}(H, 14)$ is $g_7 (= 49)$. Hence, u = 7, q = 1, and r = 3. Since $21 \notin H'$ but $42 \in H'$, we have v = 2. Also, $49 = 21 + 2 \times 14$, $42 = 3 \times 14$, implies that w = 1, z = 0, $\lambda = 2$, $\mu = 3$, r' = 3, q' = 1, and $\nu = 5$. Since $r \geq 2$, we get $\text{PF}(H) = \{\gamma_3, \gamma_4\} = \{50, 53\}$. Using Proposition 3.4,

$$\gamma_3 + m_0 = m_1 + m_4 + n$$
, $\gamma_3 + m_1 = m_2 + m_4 + n$, $\gamma_3 + m_2 = m_3 + m_4 + n$, $\gamma_3 + m_3 = 2m_4 + n$, $\gamma_3 + m_4 = m_0 + m_2 + 2n$, $\gamma_3 + n = 2m_0 + m_1 + m_4$.

We can therefore write an RF-matrix for $\gamma_3 = 50$ from these expressions,

$$RF(\gamma_3) = \begin{bmatrix} -1 & 1 & 0 & 0 & 1 & 1\\ 0 & -1 & 1 & 0 & 1 & 1\\ 0 & 0 & -1 & 1 & 1 & 1\\ 0 & 0 & 0 & -1 & 2 & 1\\ 1 & 0 & 1 & 0 & -1 & 2\\ 2 & 1 & 0 & 0 & 1 & -1 \end{bmatrix}.$$

Similarly,

$$\gamma_4 + m_0 = m_2 + m_4 + n$$
, $\gamma_4 + m_1 = m_3 + m_4 + n$, $\gamma_4 + m_2 = 2m_4 + n$, $\gamma_4 + m_3 = m_0 + m_2 + 2n$, $\gamma_4 + m_4 = m_0 + m_3 + 2n$, $\gamma_4 + n = 2m_0 + m_2 + m_4$.

The RF-matrix for $\gamma_4 = 53$ from the above expressions is as follows

$$RF(\gamma_4) = \begin{bmatrix} -1 & 0 & 1 & 0 & 1 & 1\\ 0 & -1 & 0 & 1 & 1 & 1\\ 0 & 0 & -1 & 0 & 2 & 1\\ 1 & 0 & 1 & -1 & 0 & 2\\ 1 & 0 & 0 & 1 & -1 & 2\\ 2 & 0 & 1 & 0 & 1 & -1 \end{bmatrix}.$$

Corollary 3.6. Let r = 2. Then H is symmetric and there exists an RF(H) of the following type

where $(a_{p+2,1}, \ldots, a_{p+2,p+1}, -1) = (\mu - 1, 1, 0, \ldots, 0, q + q_z, -1) + (a^{(0)}, \ldots, a^{(r_z)}, \ldots, a^{(n)})$ such that $a^{(r_z)} = 1$ and all other $a^{(j)}$'s equal to zero when $\mu > 0$. Otherwise, when $\mu = 0$, we have

$$(a_{p+2,1},\ldots,a_{p+2,p+1},-1)=(\lambda-1,0,\ldots,0,1,q_z-1,-1)+(a^{(0)},\ldots,a^{(r_z)},\ldots,a^{(n)})$$

such that $a^{(r_z)} = 1$ and all other $a^{(j)}$'s equal to zero.

Proof. Since r=2, we get k=p. Therefore, $PF(H)=\{\gamma_p\}$. Hence, H is symmetric. Since k=p and r=2 then r+k+j-p-1>p only for j=p. Therefore, we have

$$\gamma_p + m_0 = g_{qp+1} + (v-1)n = qm_p + m_1 + (v-1)n,$$

$$\gamma_p + m_j = m_{j+1} + qm_p + (v-1)n, \text{ for all } j \in [1, p-1],$$

$$\gamma_p + m_p = (\lambda - 1)m_0 + m_{p-1} + (w+v-1)n.$$

If $\mu > 1$, then $\gamma_p + n = (\mu - 1)m_0 + m_1 + m_{r_z} + (q + q_z)m_p$, and if $\mu = 0$, then $\gamma_p + n = (\lambda - 1)m_0 + m_{p-1} + m_{r_z} + (q_z - 1)m_p$. Hence the result follows. \square

Example 3.7. Let $H = \langle m_0 = 10, m_1 = 19, m_2 = 28, m_3 = 37, n = 35 \rangle$. Then, $p = 3, u = 5, v = 2, w = 1, z = 0, \lambda = 3, \mu = 7, \nu = 10, r = 2, q = 1$. Since z = 0 and r = 2, therefore $W = \emptyset$ and $PF(H) = \{\gamma_3\} = \{81\}$.

$$RF(H) = \begin{bmatrix} -1 & 1 & 0 & 1 & 1\\ 0 & -1 & 1 & 1 & 1\\ 0 & 0 & -1 & 2 & 1\\ 2 & 0 & 1 & -1 & 2\\ 6 & 1 & 0 & 1 & -1 \end{bmatrix}.$$

4. The case
$$W \neq \emptyset$$

Throughout this section we assume $W \neq \emptyset$ and $H = \langle m_0, m_1, \dots, m_p, n \rangle$, where $m_i = m_0 + id$ for $i \in [1, p]$ and $gcd(m_0, n, d) = 1$. In this case, there are two types of pseudo-Frobenius numbers. We denote them by $PF_1(H)$ and $PF_2(H)$.

• Let $\alpha_i = g_{(q'-1)p+i} + (v-1)n - m_0$. Then $\mathrm{PF}_1(H) \subset \{\alpha_i; i \in I\}$, where

$$I = \begin{cases} p & \text{if } r' = 1 \text{ and } q' = 0, \\ [1, p] & \text{if } r' = 1 \text{ and } q' > 0, \\ [p+1, p+r'-1] & \text{if } r' \ge 2. \end{cases}$$

• Let $\beta_j = g_{(q-1)p+r+j-1} + (v-w-1)n - m_0$ for $j \in [1, p]$. Then,

$$PF_{2}(H) \subset \begin{cases} \{\beta_{j} \mid j \in [1, p] \} & \text{if } r = 1, \\ \{\beta_{j} \mid j \in [p - r + 2, p] \} & \text{if } r \geq 2. \end{cases}$$

For the complete description of $PF_1(H)$ and $PF_2(H)$, see [8, Proposition 4.6, 4.7] and [8, Proposition 5.6, 5.7, 5.8, 5.9].

4.1. Structure of RF-matrices for pseudo-Frobenius numbers in $PF_1(H)$.

Proposition 4.1. Let $r' = 1, \mu > 0$ and q' = 0. Then $PF_1(H) = \{\alpha_p\}$ and there exists an $RF(\alpha_p)$ of the following type

where $(a_{p+2,1},\ldots,a_{p+2,p+1},-1)=(\mu-1,0,\cdots,0,q_z,-1)+(a^{(0)},\ldots,a^{(r_z)},\ldots,a^{(n)})$ such that $a^{(r_z)}=1$ and all other $a^{(i)}$'s equal to zero.

Proof. Since r' = 1 and q' = 0, we get $PF_1(H) = \{\alpha_p\}$. Also, u - z = q'p + r' = 1. (i) $\alpha_p + m_0 = g_0 + (v - 1)n = (v - 1)n$.

(ii) For $j \in [1, p]$,

$$\alpha_p + m_j = (v - 1)n - m_0 + m_j = (v - w)n + (w - 1)n - m_0 + m_j$$
$$= \nu m_0 - g_{(u-z)} + (w - 1)n - m_0 + m_j$$
$$= (\nu - 2)m_0 + m_{j-1} + (w - 1)n.$$

(iii) $\alpha_p + n = (v-1)n - m_0 + n = (\mu - 1)m_0 + g_z = (\mu - 1)m_0 + q_z m_p + m_{r_z}$. Hence, the result follows. **Example 4.2.** Let $H = \langle m_0 = 11, m_1 = 13, m_2 = 15, m_3 = 17, m_4 = 19, n = 21 \rangle$. Then, $p = 4, u = 5, v = 3, w = 1, z = 4, \lambda = 1, \mu = 4, \nu = 5, r = 1, r' = 1, q = 1, q' = 0$. Since r' = 1, q' = 0, we get $PF_1(H) = \{\alpha_4\} = \{31\}$ and

$$RF(31) = \begin{bmatrix} -1 & 0 & 0 & 0 & 0 & 2\\ 4 & -1 & 0 & 0 & 0 & 0\\ 3 & 1 & -1 & 0 & 0 & 0\\ 3 & 0 & 1 & -1 & 0 & 0\\ 3 & 0 & 0 & 1 & -1 & 0\\ 3 & 0 & 0 & 0 & 1 & -1 \end{bmatrix}.$$

Proposition 4.3. Let $r' = 1, \mu > 0$ and q' > 0. Then for each $\alpha_i \in PF_1(H)$, $i \in [1, p]$, there exists an $RF(\alpha_i)$ of the following type

$$RF(\alpha_i) = \begin{bmatrix} -1 & 0 & 0 & \cdots & \cdots & 0 & q'-1 & v-1 \\ 0 & -1 & 0 & \cdots & \cdots & 0 & q'-1 & v-1 \\ \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & q'-1 & v-1 \\ \nu-2 & 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & w-1 \\ \vdots & \vdots \\ \nu-2 & 0 & 0 & \cdots & \cdots & 0 & -1 & w-1 \\ \mu-1 & 0 & 0 & \cdots & \cdots & 0 & q'+q_z-1 & -1 \end{bmatrix} + \begin{bmatrix} R_{(0)} \\ R_{(1)} \\ \vdots \\ R_{(p-i)} \\ R_{(p)} \\ R_{(n)} \end{bmatrix},$$

where

- (1) $R_{(0)} = (a^{(0)}, \dots, a^{(i)}, \dots, a^{(n)})$ such that $a^{(i)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (2) For all $j \in [1, p-i]$, $R_{(j)} = (a^{(0)}, \dots, a^{(i+j)}, \dots, a^{(n)})$ such that $a^{(i+j)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (3) For all $j \in [p-i+1,p]$, $R_{(j)} = (a^{(0)}, \ldots, a^{(i+j-p-1)}, \ldots, a^{(n)})$ such that $a^{(i+j-p-1)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (4) $R_{(n)} = (a^{(0)}, \ldots, a^{(i)}, \ldots, a^{(r_z)}, \ldots, a^{(n)})$ such that $a^{(r_z)}, a^{(i)} = 1$ and all other $a^{(j)}$'s equal to zero.

Proof. Since $\mu > 0$ and q' > 0, we have $PF_1(H) = \{\alpha_i \mid i \in [1, p]\}.$

(1)
$$\alpha_i + m_0 = g_{(q'-1)p+i} + (v-1)n = (q'-1)m_p + m_i + (v-1)n$$
.

(2) If $j \in [1, p - i]$, then $i + j \leq p$ and

$$\alpha_i + m_j = g_{(q'-1)p+i} + (v-1)n - m_0 + m_j = (q'-1)m_p + m_i + (v-1)n - m_0 + m_j$$
$$= (q'-1)m_p + m_{i+j} + (v-1)n.$$

(3) If $j \in [p-i+1, p]$, then i+j > p and

$$\alpha_{i} + m_{j} = g_{(q'-1)p+i} + (v-1)n - m_{0} + m_{j} = g_{(q'-1)p+i+j} + (v-1)n - m_{0}$$

$$= g_{u-z-r'-p+i+j} + (v-w)n + (w-1)n - m_{0}$$

$$= g_{i+j-p-1} + (v-2)m_{0} + (w-1)n$$

$$= (v-2)m_{0} + m_{i+j-p-1} + (w-1)n.$$

(4) Now,

$$\alpha_i + n = g_{(q'-1)p+i} + vn - m_0 = g_{(q'-1)p+i} + \mu m_0 + g_z - m_0$$
$$= (q' + q_z - 1)m_p + (\mu - 1)m_0 + m_i + m_{r_z}.$$

Note that if p = 1 and $PF_1(H) \neq \emptyset$, then $PF_1(H) = \{\alpha_1\}$.

Corollary 4.4. Suppose p = 1 and $PF_1(H) \neq \emptyset$. Then there exists an RF-matrix of α_1 of the type

$$\begin{bmatrix} -1 & u-z-1 & v-1 \\ \nu-1 & -1 & w-1 \\ \mu-1 & u-1 & -1 \end{bmatrix}, \quad or \begin{bmatrix} -1 & 0 & v-1 \\ \nu-1 & -1 & w-1 \\ \mu-1 & z & -1 \end{bmatrix}.$$

Proof. Since p=1, we get $r_t=1$, $q_t=t-1$ for all t. Now use Proposition 4.1 and 4.3 to get desired RF-matrices accordingly as q' > 0 or q' = 0.

Proposition 4.5. Let r' = 1, $\mu = 0$, q' > 0 and r > 2. Then for every element of $PF_1(H) =$ $\{\alpha_i \mid i \in [1, p-r+1]\}$, there exists an RF(α_i) of the following type

$$RF(\alpha_i) = \begin{bmatrix} -1 & 0 & 0 & \cdots & \cdots & 0 & q'-1 & v-1 \\ 0 & -1 & 0 & \cdots & \cdots & 0 & q'-1 & v-1 \\ \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & q'-1 & v-1 \\ \nu-2 & 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & w-1 \\ \vdots & \vdots \\ \nu-2 & 0 & 0 & \cdots & \cdots & 0 & -1 & w-1 \\ 0 & 0 & 0 & \cdots & \cdots & 0 & q-1 & -1 \end{bmatrix} + \begin{bmatrix} R_{(0)} \\ R_{(1)} \\ \vdots \\ R_{(p-i)} \\ R_{(p)} \\ R_{(n)} \end{bmatrix},$$

where

- (1) $R_{(0)} = (a^{(0)}, \dots, a^{(i)}, \dots, a^{(n)})$ such that $a^{(i)} = 1$ and all other $a^{(j)}$'s equal to zero. (2) For all $j \in [1, p i]$, $R_{(j)} = (a^{(0)}, \dots, a^{(i+j)}, \dots, a^{(n)})$ such that $a^{(i+j)} = 1$ and all
- other $a^{(j)}$'s equal to zero. (3) $R_{(n)} = (a^{(0)}, \ldots, a^{(r+i-1)}, \ldots, a^{(n)})$ such that $a^{(r+i-1)} = 1$ and all other $a^{(j)}$'s equal to zero.

Proof. For (1),(2) and (3), see the proof of Proposition 4.3.

(4) Since $r \geq 2$ and r' = 1, we get $r_z = r - 1$. Also, we have $i \in [1, p - r + 1]$ and hence $i + r_z \leq p$. Therefore,

$$\begin{split} \alpha_i + n &= g_{(q'-1)p+i} + vn - m_0 = g_{(q'-1)p+i} + g_z - m_0 \\ &= g_{u-z-p+i-1} + g_z - m_0 \\ &= g_{u-p+i-1} \\ &= g_{(q-1)p+r+i-1} = (q-1)m_p + m_{r+i-1}. \end{split}$$

Note that $PF(H) = \emptyset$ in the following cases and so we proceed to the case $r' \geq 2$.

(1)
$$r' = 1, \mu = 0$$
 and $q' = 0$, (2) $r' = 1, \mu = 0, q' > 0$, and $r = 1$.

Proposition 4.6. Let $r' \geq 2$. Then for each $\alpha_i \in PF_1(H)$, $i \in [p+1, p+r'-1]$, there exists an $RF(\alpha_i)$ of the following type

$$RF(\alpha_i) = \begin{bmatrix} -1 & 0 & 0 & \cdots & \cdots & 0 & q' & v-1 \\ 0 & -1 & 0 & \cdots & \cdots & 0 & q' & v-1 \\ \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & q' & v-1 \\ \nu-1 & 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & w-1 \\ \vdots & \vdots \\ \nu-1 & 0 & 0 & \cdots & \cdots & 0 & -1 & w-1 \\ a_{p+2,1} & a_{p+2,2} & 0 & \cdots & \cdots & \cdots & a_{p+2,p+1} & -1 \end{bmatrix} + \begin{bmatrix} R_{(0)} \\ R_{(1)} \\ \vdots \\ R_{(2p-i)} \\ R_{(2p-i+1)} \\ \vdots \\ R_{(p)} \\ R_{(n)} \end{bmatrix},$$

where $\theta_i \in [1, r'-1]$ such that $i = p + \theta_i$.

- (1) $R_{(0)} = (a^{(0)}, \ldots, a^{(\theta_i)}, \ldots, a^{(n)})$ such that $a^{(\theta_i)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (2) For all $j \in [1, 2p i]$, $R_{(j)} = (a^{(0)}, \dots, a^{(\theta_i + j)}, \dots, a^{(n)})$ such that $a^{(\theta_i + j)} = 1$ and all other $a^{(j)}$'s equal to zero.
- (3) For all $j \in [2p i + 1, p]$, $R_{(j)} = (a^{(0)}, \dots, a^{(\theta_i + j r')}, \dots, a^{(n)})$ such that $a^{(\theta_i + j r')} = 1$ and all other $a^{(j)}$'s equal to zero.
- (4) If $\mu > 0$, then $(a_{p+2,1}, a_{p+2,2}, \dots, a_{p+2,p+1}, -1) = (\mu 1, 0, \dots, 0, q' + q_z, -1)$ and $R_{(n)} = (a^{(0)}, \dots, a^{(\theta_i)}, \dots, a^{(r_z)}, \dots, a^{(n)})$ such that $a^{(r_z)}, a^{(\theta_i)} = 1$ and all other $a^{(j)}$'s equal to zero. Otherwise, if $\mu = 0$ then $(a_{p+2,1}, a_{p+2,2}, \dots, a_{p+2,p+1}, -1) = (0, \dots, 0, q' + q_z, -1)$ and $R_{(n)} = (a^{(0)}, \dots, a^{(\theta_i + r_z)}, \dots, a^{(n)})$ such that $a^{(\theta_i + r_z)} = 1$ and all other $a^{(j)}$'s equal to zero.

Proof. Since $r' \geq 2$, we have $PF_1(H) \subseteq \{\alpha_i \mid i \in [p+1, p+r'-1]\}.$

(1)
$$\alpha_i + m_0 = g_{(q'-1)p+i} + (v-1)n = g_{q'p+\theta_i} + (v-1)n = q'm_p + m_{\theta_i} + (v-1)n.$$

(2) If $j \in [1, 2p - i]$, then $i + j \leq 2p$ and hence $\theta_i + j \leq p$. Therefore,

$$\alpha_i + m_j = g_{(q'-1)p+i} + (v-1)n - m_0 + m_j = g_{q'p+\theta_i} + (v-1)n - m_0 + g_j$$

$$= g_{q'p+\theta_i+j} + (v-1)n$$

$$= q'm_p + m_{\theta_i+j} + (v-1)n.$$

(3) If $j \in [2p - i + 1, p]$, then i + j > 2p. Therefore,

$$\alpha_i + m_j = g_{q'p-p+i} + (v-1)n - m_0 + m_j = g_{u-z-r'-p+i} + g_j + (v-w)n + (w-1)n - m_0$$

$$= g_{u-z} + g_{i+j-p-r'} + (v-w)n + (w-1)n - m_0$$

$$= (\nu - 1)m_0 + m_{i+j-p-r'} + (w-1)n.$$

(4) Assume $\mu > 0$. Then

$$\alpha_i + n = g_{(q'-1)p+i} + vn - m_0 = g_{(q'-1)p+i} + (\mu - 1)m_0 + g_z$$

$$= g_{q'p+\theta_i} + (\mu - 1)m_0 + g_z$$

$$= (\mu - 1)m_0 + m_{r_z} + m_{\theta_i} + (q_z + q')m_p.$$

If $\mu = 0$, then $vn = g_z$. Therefore,

$$\alpha_i + n = g_{(q'-1)p+i} + vn - m_0 = g_{q'p+\theta_i} + g_z - m_0$$
$$= q'm_p + q_z m_p + m_{\theta_i} + m_{r_z} - m_0.$$

Since we have $r_z + \theta_i \leq p$, this implies $\alpha_i + n = (q' + q_z)m_p + m_{\theta_i + r_z}$.

4.2. Structure of RF-matrices for pseudo-Frobenius numbers in $PF_2(H)$.

Proposition 4.7. Let r = 1 and $\lambda > 1$. Then for each $\beta_j \in PF_2(H)$, $j \in [1, p]$, there exists an $RF(\beta_j)$ of the following type

$$RF(\beta_{j}) = \begin{bmatrix} -1 & 0 & 0 & \cdots & \cdots & 0 & q-1 & v-w-1 \\ 0 & -1 & 0 & \cdots & \cdots & 0 & q-1 & v-w-1 \\ \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & q-1 & v-w-1 \\ \lambda-2 & 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & v-1 \\ \vdots & \vdots \\ \lambda-2 & 0 & 0 & \cdots & \cdots & 0 & -1 & v-1 \\ v-\epsilon & 0 & \cdots & \cdots & 0 & q_{z+j-p-1} & -1 \end{bmatrix} + \begin{bmatrix} R_{(0)} \\ R_{(1)} \\ \vdots \\ R_{(p-j)} \\ R_{(p-j+1)} \\ \vdots \\ R_{(p)} \\ R_{(n)} \end{bmatrix},$$

where $\epsilon = 2$ or 1 according as $r' + r_{z-p+j-1} \leq p$ or $r' + r_{z-p+j-1} > p$ and

- (1) $R_{(0)} = (a^{(0)}, \dots, a^{(j)}, \dots, a^{(n)})$ such that $a^{(j)} = 1$ and all other $a^{(k)}$'s equal to zero.
- (2) For all $k \in [1, p j]$, $R_{(k)} = (a^{(0)}, \dots, a^{(j+k)}, \dots, a^{(n)})$ such that $a^{(j+k)} = 1$ and all other $a^{(k)}$'s equal to zero.
- (3) For all $k \in [p-j+1, p]$, $R_{(k)} = (a^{(0)}, \dots, a^{(k+j-p-1)}, \dots, a^{(n)})$ such that $a^{(k+j-p-1)} = 1$ and all other $a^{(k)}$'s equal to zero.
- (4) $R_{(n)} = (a^{(0)}, \ldots, a^{(r_{z+j-p-1})}, \ldots, a^{(n)})$ such that $a^{(r_{z+j-p-1})} = 1$ and all other $a^{(k)}$'s equal to zero.

Proof. Since r = 1, $\beta_j = g_{(q-1)p+j} + (v - w - 1)n - m_0$.

- (1) $\beta_j + m_0 = (q-1)m_p + m_j + (v-w-1)n$.
- (2) If $k \in [1, p j]$, then $k + j \le p$. Therefore,

$$\beta_j + m_k = g_{(q-1)p+j} + (v-w-1)n - m_0 + m_k = (q-1)m_p + m_{j+k} + (v-w-1)n.$$

(3) If $k \in [p-j+1, p]$, then $k+j \ge p+1$. Given $\lambda > 1$, if k+j > p+1 then

$$\beta_j + m_k = g_{(q-1)p+j} + (v - w - 1)n - m_0 + m_k = g_{u+k+j-p-1} + (v - 1)n + \lambda m_0 - g_u - m_0$$

$$= (\lambda - 2)m_0 + g_{j+k-p-1} + (v - 1)n$$

$$= (\lambda - 2)m_0 + m_{j+k-p-1} + (v - 1)n.$$

If
$$k + j = p + 1$$
, then $\beta_j + m_k = (\lambda - 1)m_0 + (v - 1)n$.

- (4) We consider two subcases $z \ge p$ and z < p.
- (a) Assume $z \geq p$.

$$\beta_j + n = g_{(q-1)p+j} + (v - w - 1)n - m_0 + n = g_{(q-1)p+j} + (\nu - 1)m_0 - g_{u-z}$$
$$= g_{u-z+z-p+j-1} + (\nu - 1)m_0 - g_{u-z}.$$

Since $z \ge p$, we have $z - p + j - 1 \ge 0$. Therefore,

$$\beta_j + n = g_{z-p+j-1} - \epsilon m_0 + (\nu - 1)m_0 = (\nu - 1)m_0 - \epsilon m_0 + m_{r_{z+j-p-1}} + q_{z-p+j-1}m_p,$$

with $\epsilon = 1$ or 0 according as $r' + r_{z-p+j-1} \le p$ or $r' + r_{z-p+j-1} > p$.

(b) If z < p, then $z = r_z = p + r - r' = p + 1 - r'$. Also in this case $PF_2(H) = \{\beta_j \mid j \in [r', p]\}$ (see [8, Proposition 5.7]). Since $j \ge r' = p - z + 1$, we get $z - p + j - 1 \ge 0$. Hence we are done by subcase(a).

Proposition 4.8. Let r=1 and $\lambda=1$. Then for each $\beta_j \in PF_2(H)$, $j \in [1,p]$, there exists an RF(β_i) of the following type

$$RF(\beta_{j}) = \begin{bmatrix} -1 & 0 & 0 & \cdots & \cdots & 0 & q-1 & v-w-1 \\ 0 & -1 & 0 & \cdots & \cdots & 0 & q-1 & v-w-1 \\ \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & q-1 & v-w-1 \\ \lambda-1 & 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & v-1 \\ \nu-2 & 0 & 0 & \cdots & \cdots & 0 & -1 & w-1 \\ \vdots & \vdots \\ \nu-2 & 0 & 0 & \cdots & \cdots & 0 & -1 & w-1 \\ \nu-\epsilon & 0 & \cdots & \cdots & 0 & q_{z+j-p-1} & -1 \end{bmatrix} + \begin{bmatrix} R_{(0)} \\ R_{(1)} \\ \vdots \\ R_{(p-j)} \\ 0 \\ R_{(p-j+2)} \\ \vdots \\ R_{(p)} \\ R_{(n)} \end{bmatrix},$$
where $\epsilon = 2$ or 1 according as $x' + x$ and $x' + x$ and

where $\epsilon = 2$ or 1 according as $r' + r_{z-p+j-1} \leq p$ or $r' + r_{z-p+j-1} > p$ and

- (1) $R_{(0)} = (a^{(0)}, \ldots, a^{(j)}, \ldots, a^{(n)})$ such that $a^{(j)} = 1$ and all other $a^{(k)}$'s equal to zero.
- (2) For all $k \in [1, p j]$, $R_{(k)} = (a^{(0)}, \dots, a^{(j+k)}, \dots, a^{(n)})$ such that $a^{(j+k)} = 1$ and all other $a^{(k)}$'s equal to zero.
- (3) For all $k \in [p-j+2, p]$, $R_{(k)} = (a^{(0)}, \dots, a^{(k+j-p-2)}, \dots, a^{(n)})$ such that $a^{(k+j-p-2)} = 1$ and all other $a^{(k)}$'s equal to zero.
- (4) $R_{(n)} = (a^{(0)}, \dots, a^{(r_{z+j-p-1})}, \dots, a^{(n)})$ such that $a^{(r_{z+j-p-1})} = 1$ and all other $a^{(k)}$'s equal to zero.

Proof. Same as 4.7 except when $j + k \ge p + 2$.

If j + k = p + 2, then $\beta_j + m_k = (\nu - 1)m_0 + (w - 1)n$.

If j+k>p+2, then $\beta_j+m_k=(\nu-2)m_0+g_{j+k-p-2}+(w-1)n$. Hence the result follows. \square

Note that if p = 1, then $PF_2(H) = \{\beta_1\}$.

Corollary 4.9. Suppose p = 1. Then there exists an RF-matrix of β_1 of the following type

$$\begin{bmatrix} -1 & u-1 & v-w-1 \\ \lambda-1 & -1 & v-1 \\ \nu-1 & z-1 & -1 \end{bmatrix}.$$

Proof. Use Proposition 4.7 and 4.8.

Let H be a numerical semigroup generated by a sequence m_0, \ldots, m_e . Let k be a field. The semigroup ring k[H] of H is a k-subalgebra of the polynomial ring k[t]. In other words, $k[H] = k[t^{m_0}, t^{m_1}, \ldots, t^{m_e}]$. Set $R = k[x_0, \ldots, x_e]$ and define a map $\pi : R \to k[H]$ given by $\pi(x_i) = t^{m_i}$ for all $i = 0, \ldots, e$. Set deg $x_i = m_i$ for all $i = 0, \ldots, e$. Observe that R is a graded ring and that π is a degree preserving surjective k-algebra homomorphism. We say kernel of π is defining ideal of k[H], denoted by I(H).

For a vector $b = (b_1, \ldots, b_n) \in \mathbb{Z}^n$, we let b^+ denote the vector whose *i*-th entry is b_i if $b_i > 0$, and is zero otherwise, and we let $b^- = b^+ - b$. Then $b = b^+ - b^-$ with $b^+, b^- \in \mathbb{N}^n$. For a pseudo-Frobenius element f, the rows of RF(f) produce binomials in I(H).

Definition 4.10. For some pseudo-Frobenius element f of H, let $\delta_1, \ldots, \delta_n$ denote the row vectors of RF(f). Set $\delta_{(ij)} = \delta_j - \delta_i$, for all $1 \le i < j \le n$. Then $\phi_{ij} = \mathbf{x}^{\delta_{(ij)}^+} - \mathbf{x}^{\delta_{(ij)}^-} \in I(H)$ for all i < j. We call ϕ_{ij} an RF(f)-relation. We call a binomial relation $\phi \in I(H)$ an RF-relation if it is an RF(f)-relation for some $f \in PF(H)$.

Proposition 4.11. If p = 1 and $\mu > 0$, then I(H) is minimally generated by $RF(\beta_1)$ -relations.

Proof. Since p=1 and $\mu>0$, using [8, Proposition 4.6, 5.6] it follows that $PF_1(H)=\{\alpha_1\}$ and $PF_2(H)=\{\beta_1\}$. Therefore, we get $PF(H)=\{\alpha_1,\beta_1\}$ and H is not symmetric. Also, as p=1, we have $q_t=t-1$ for all t. In [9, Theorem 4.5], it is proved that in the above setup,

$$I(H) = \langle x_0^{\nu} - x_1^{q'+1} x_2^{\nu-w}, x_1^{q+1} - x_0^{\lambda} x_2^{w}, x_2^{v} - x_0^{\mu} x_1^{q-q'} \rangle.$$

In order to prove the result, it is sufficient to show that these generators arise from either $RF(\alpha_1)$ or $RF(\beta_1)$ -relations. Consider the matrix

$$RF(\beta_1) := \begin{bmatrix} -1 & u-1 & v-w-1 \\ \lambda - 1 & -1 & v-1 \\ \nu - 1 & z-1 & -1 \end{bmatrix}.$$

If $\delta_1, \delta_2, \delta_3$ represent the row vectors of RF(β_1), then we have the following RF(β_1)-relations: $\delta_{12} = x_1^u - x_0^{\lambda} x_2^w, \delta_{13} = x_0^{\nu} - x_1^{u-z} x_2^{v-w}$ and $\delta_{23} = x_2^{\nu} - x_0^{\mu} x_1^z$. Hence the result.

Proposition 4.12. Let $r \geq 2$. Then for each $\beta_j \in PF_2(H)$, $j \in [p-r+2, p]$, there exists an $RF(\beta_j)$ of the following type

$$RF(\beta_{j}) = \begin{bmatrix} -1 & 0 & 0 & \cdots & \cdots & 0 & q & v-w-1 \\ 0 & -1 & 0 & \cdots & \cdots & 0 & q & v-w-1 \\ \vdots & \vdots \\ 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & q & v-w-1 \\ \lambda-1 & 0 & 0 & \cdots & -1 & 0 & \cdots & 0 & v-1 \\ \vdots & \vdots \\ \lambda-1 & 0 & 0 & \cdots & \cdots & 0 & -1 & v-1 \\ \nu-\epsilon & 0 & \cdots & \cdots & \cdots & 0 & q_{z+j-p-1} & -1 \end{bmatrix} + \begin{bmatrix} R_{(0)} \\ R_{(1)} \\ \vdots \\ R_{(e)} \\ R_{(e+1)} \\ \vdots \\ R_{(p)} \\ R_{(n)} \end{bmatrix},$$

where $e \in [1, p-1]$ such that r+j+e = 2p+1 and $\epsilon = 2$ or 1 according as $r' + r_{z-p+j-1} \le p$ or $r' + r_{z-p+j-1} > p$, and

- (1) $R_{(0)} = (a^{(0)}, \dots, a^{(p-e)}, \dots, a^{(n)})$ such that $a^{(p-e)} = 1$ and all other $a^{(k)}$'s equal to zero.
- (2) For all $k \in [1, e]$, $R_{(k)} = (a^{(0)}, \dots, a^{(p-e+k)}, \dots, a^{(n)})$ such that $a^{(p-e+k)} = 1$ and all other $a^{(k)}$'s equal to zero.

- (3) For all $k \in [e+1, p]$, $R_{(k)} = (a^{(0)}, \dots, a^{(j+k-p-1)}, \dots, a^{(n)})$ such that $a^{(j+k-p-1)} = 1$
- and all other $a^{(k)}$'s equal to zero. (4) $R_{(n)} = (a^{(0)}, \ldots, a^{(r_{z+j-p-1})}, \ldots, a^{(n)})$ such that $a^{(r_{z+j-p-1})} = 1$ and all other $a^{(k)}$'s equal to zero.

Proof. Use similar arguments as in Proposition 3.4 and Proposition 4.7.

5. Generic toric ideals

Let $R = k[x_0, \ldots, x_n]$ be a polynomial ring over the field k. A binomial ideal is said to be **generic** if it has a minimal generating set consisting binomials of full support (i.e each variable should appear in each binomial of the generating set). This concept was introduced by Peeva and Sturmfels in [11]. In this section, we give a characterization of the generic nature of toric ideals when the semigroup is minimally generated by an almost arithmetic sequence. We use the following result of Eto for the same, which gives a necessary and sufficient condition for a toric ideal to be generic.

Theorem 5.1 ([3, Theorem 1]). Let H be a numerical semigroup. Then I(H) is generic if and only if for each $f \in PF(H)$, $RF(f) = (a_{i,j})$ is unique and $a_{i,j} \neq a_{i',j}$ if $i \neq i'$.

Example 5.2. Let H = (5, 6, 7). Then $PF(H) = \{8, 9\}$ and

$$RF(8) = \begin{bmatrix} -1 & 1 & 1 \\ 0 & -1 & 2 \\ 3 & 0 & -1 \end{bmatrix}, \qquad RF(9) = \begin{bmatrix} -1 & 0 & 2 \\ 3 & -1 & 0 \\ 2 & 1 & -1 \end{bmatrix},$$

where RF(8) and RF(9) are unique RF-matrices. Hence by Theorem 5.1, I(H) is generic. On the other hand, $I(H) = (x_1^2 - x_0 x_2, x_0^4 - x_1 x_2^2, x_0^3 x_1 - x_2^3)$ and hence is generic.

Theorem 5.3. Let H be a numerical semigroup minimally generated by an almost arithmetic sequence, i.e $H = \langle m_0, \dots, m_p, n \rangle$, where $m_i = m_0 + id$ for $i \in [1, p]$ and $gcd(m_0, n, d) = 1$.

- (1) If p = 0, then I(H) is generic.
- (2) If p = 1, and if $W \neq \emptyset$, $\mu > 0$, then I(H) is generic. Otherwise, it is never generic.
- (3) If p > 1, then I(H) is not generic.

Proof. (1) It is obvious.

(2) If $p=1, W\neq \emptyset$, and $\mu>0$, then by Proposition 4.11, we conclude that I(H) is generic. If $W \neq \emptyset$ and $\mu = 0$, then $\lambda = \nu$ and by Corollary 4.9 and Theorem 5.1 we get I(H) is not generic.

If $W = \emptyset$, then either w = 0 or z = 0. Using Corollary 3.3 and Theorem 5.1, we conclude that I(H) is not generic.

(3) In the Propositions 3.1, 3.2, 3.4, 4.1, 4.3, 4.5, 4.6, 4.7, 4.8, and 4.12, we see that for each $f \in PF(H)$, there exists an RF-matrix $RF(f) = (a_{i,j})$ such that $a_{i,p+2} = a_{i',p+2}$ for some $i \neq i'$. Hence by Theorem 5.1, I(H) is not generic.

We now prove that the defining ideal of a gluing of two numerical semigroups is never generic. As a consequence, it is easy to see that a complete intersection toric ideal is not generic. We recall the definition of gluing first.

Definition 5.4 ([12]). Let H_1 and H_2 be two numerical semigroups minimally generated by $\{m_0, \ldots, m_{e_1}\}$ and $\{m'_0, \ldots, m'_{e_2}\}$ respectively. Let $d_1 \in H_2 \setminus \{m'_0, \ldots, m'_{e_2}\}$ and $d_2 \in H_1 \setminus \{m_0, \ldots, m_{e_1}\}$ be such that $\gcd(d_1, d_2) = 1$. We say that

$$H = \langle d_1 m_0, \dots, d_1 m_{e_1}, d_2 m'_0, \dots, d_2 m'_{e_2} \rangle$$

is a **gluing** of H_1 and H_2 .

Note that if $H_1 = H_2 = \mathbb{N}$, then the embedding dimension of H is two and hence I(H) is a generic ideal. Therefore, we assume that either $H_1 \neq \mathbb{N}$ or $H_2 \neq \mathbb{N}$.

Theorem 5.5. Let $H_1 = \langle m_0, \ldots, m_{e_1} \rangle$ and $H_2 = \langle m'_0, \ldots, m'_{e_2} \rangle$ be two numerical semi-groups such that $e_2 \geq 1$ and H be the gluing of H_1 and H_2 . Then I(H) is never generic.

Proof. Let $H = d_1H_1 + d_2H_2$, where $d_1 \in H_2 \setminus \{m'_0, \ldots, m'_{e_2}\}$ and $d_2 \in H_1 \setminus \{m_0, \ldots, m_{e_1}\}$ such that $\gcd(d_1, d_2) = 1$. We will show that there exists an RF-matrix for a $f \in \operatorname{PF}(H)$ such that $a_{ij} = a_{i'j}$ for some $i \neq i'$. Let $f_1 \in \operatorname{PF}(H_1)$ and $f_2 \in \operatorname{PF}(H_2)$. Set $g = d_1f_1 + d_2f_2 + d_1d_2$. Then $g \in \operatorname{PF}(H)$, see [13, Theorem 2.4]. Since $f_1 \in \operatorname{PF}(H_1)$, we get $f_1 + d_2 \in H_1$. Therefore, there exist $a_0, \ldots, a_{e_1} \in \mathbb{Z}_{\geq 0}$ such that $f_1 + d_2 = \sum_{i=0}^{e_1} a_i m_i$. Similarly, for $f_2 + d_1 \in H_2$, there exist $a'_0, \ldots, a'_{e_2} \in \mathbb{Z}_{\geq 0}$ such that $f_2 + d_1 = \sum_{i=0}^{e_2} a'_i m'_i$. Let $\operatorname{RF}(f_1)$ be an RF-matrix of f_1 and $\operatorname{RF}(f_2)$ be an RF-matrix of f_2 . Then,

$$\begin{bmatrix}
RF(f_1) & a'_0 & \cdots & a'_{e_2} \\
\vdots & \vdots & \vdots \\
a'_0 & \cdots & a'_{e_2}
\end{bmatrix}$$

$$\begin{bmatrix}
a_0 & \cdots & a_{e_1} \\
\vdots & \vdots & \vdots \\
a_0 & \cdots & a_{e_1}
\end{bmatrix}$$

$$RF(f_2)$$

defines an RF-matrix for g. Since $a_{1,j} = a_{2,j} = \ldots = a_{e_1+1,j}$ for each $j \in [e_1 + 2, e_1 + e_2 + 2]$, and $a_{e_1+2,j} = a_{e_1+3,j} = \ldots = a_{e_1+e_2+2,j}$ for each $j \in [1, e_1 + 1]$. Then by Theorem 5.1, we conclude that I(H) is not generic.

Corollary 5.6. Let H be a complete intersection numerical semigroup with embedding dimension at least 3. Then I(H) is not generic.

Proof. By [12, Theorem 9.10], we have that a numerical semigroup (other than \mathbb{N}) is a complete intersection if and only if it is a gluing of two complete intersection numerical semigroups. Now using Theorem 5.5, we are done.

6. RF-relations when type(H)=1

In this section, we consider those numerical semigroups generated by an almost arithmetic sequence which are symmetric. In particular, let $H = \langle m_0, m_1, \dots, m_p, n \rangle$, where $m_i = m_0 + id$ for $i \in [1, p]$ and $\gcd(m_0, n, d) = 1$ such that $\operatorname{type}(H) = |\operatorname{PF}(H)| = 1$. Using the language from the previous sections, we list the different cases in which $\operatorname{type}(H) = 1$ and also give a possible RF-matrix of the Frobenius number in each case. The aim of this section is to prove that I(H) has a minimal generating set consisting of RF-relations in the above setup if the embedding dimension is either 4 or 5.

Let H be a numerical semigroups generated by an almost arithmetic sequence. Then type(H) = 1 in the following cases:

Case 1: Let $W = \emptyset$. In this case, if r = 1, then H is of type 1 only if p = 1 and $PF(H) = \{\gamma_1\}$. If $r \geq 2$, then H is of type 1 only if r = 2 and $PF(H) = \{\gamma_p\}$.

Case 2: Let $W \neq \emptyset$. If p = 1, then H is of type 1 only if $\mu = 0$ and $PF(H) = \{\beta_1\}$. Suppose $p \geq 2$. The only cases in which H is of type 1 are the following:

- (i) $r = 1, r' = 2, \lambda = 1, z < p, \text{ and } PF(H) = \{\alpha_{p+1}\}.$ (ii) $r = 2, r' = 1, \mu = 0, q' = 0, \text{ and } PF(H) = \{\beta_p\}.$
- (iii) r = 2, r' = 2, $\mu = 0$, and $PF(H) = \{\beta_p\}$.

We now consider each case separately and list an RF-matrix of the Frobenius number in each case.

Case 1: Let $W = \emptyset$.

Let p = 1, r = 1 and $PF(H) = \{\gamma_1\}$. Then we have choices for the $RF(\gamma_1)$ -matrix as given in Corollary 3.3.

Let r=2 and $p\geq 2$. Then there exists an RF-matrix of γ_p as given in Corollary 3.6.

Case 2: Let $W \neq \emptyset$.

Suppose p = 1 and $\mu = 0$. Then there exists an RF-matrix of β_1 as given in Corollary 4.9. Let $p \geq 2$. We have the following three sub-cases:

Sub-case (i): Let r = 1, r' = 2, $\lambda = 1$ and z < p and $PF(H) = {\alpha_{p+1}}$. As a consequence, $r_z = p - 1$, implying that z = p - 1 and $q_z = 0$. Also, $\lambda = 1$ and r' > r implies $\mu \neq 0$. Using Proposition 4.6, there exists an RF-matrix of α_{p+1} of the following type

Sub-case (ii): Let r = 2, r' = 1, $\mu = 0$, q' = 0 and $PF(H) = \{\beta_p\}$. Then $r_z = 1$ and hence $r_{z-1} = p$ and $q_{z-1} = q_z - 1$. Since q' = 0, we get u - z = r' = 1 which implies z = u - 1 = qp + 1 and hence $q_z = q$. Also $r' + r_{z-1} = p + 1 > p$ implies $\epsilon = 1$ and as $\mu = 0$ and r' < r, we get $\nu = \lambda + 1$. By Proposition 4.12, there exists an RF-matrix of β_p of the following type

Sub-case (iii): Let r=2, r'=2, $\mu=0$ and $PF(H)=\{\beta_p\}$. Then $r_z=p$ and hence $r_{z-1}=p-1$ and $q_{z-1}=q_z$. Also, $r'+r_{z-1}=p+1>p$ implies $\epsilon=1$ and as $\mu=0$ and r=r', we get $\nu=\lambda$. By Proposition 4.12, there exists an RF-matrix of β_p of the following type

For a pseudo-Frobenius element $f \in H$, the difference of any two rows of an RF(f)-matrix produces a binomial, which we call an RF(f)-relation (see Definition 4.10). It is easy to check that these binomials are elements in I(H). In [5], the authors raised the following question:

Question: When is I(H) minimally generated by RF-relations?

The authors answer the question in the affirmative when the embedding dimension is 3 and also when the embedding dimension is 4 and the semigroup is pseudo-symmetric or almost symmetric. They also remark that in the case of symmetric semigroups with embedding dimension 4, RF(H) cannot be determined uniquely and that all the generators of I(H) cannot be obtained from a single RF(H).

When H is generated by an almost arithmetic sequence, we proved in Theorem 5.3 that the ideal I(H) is not generic when the embedding dimension is at least 4. Using [3, Theorem 1] and the structure of RF-matrices (as given in section 3 and 4), we then conclude that the RF-matrices are indeed not unique in this case. We now answer the question of Herzog-Watanabe in the affirmative for the symmetric numerical semigroups generated by an almost arithmetic sequence, with embedding dimension 4 or 5.

Proposition 6.1. Suppose $W = \emptyset$ and p = 2. If H is symmetric, then I(H) is minimally generated by RF-relations.

Proof. Since $W = \emptyset$, p = 2 and H is symmetric, we get r = 2 and $PF(H) = \{\gamma_2\}$. Let $\delta_1, \delta_2, \delta_3$ and δ_4 denote the rows of $RF(\gamma_2)$ where

$$RF(\gamma_2) = \begin{bmatrix} -1 & 1 & q & v-1 \\ 0 & -1 & q+1 & v-1 \\ \lambda-1 & 1 & -1 & w+v-1 \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix},$$

and $\delta_4 = (a_{41}, a_{42}, a_{43}, a_{44})$ is as follows:

- (1) If r' = 1 and $\mu > 0$, then $\delta_4 = (\mu 1, 2, q + q_z, -1)$.
- (2) If r' = 2 and $\mu > 0$, then $\delta_4 = (\mu 1, 1, q + q_z + 1, -1)$.
- (3) If r' = 1 and $\mu = 0$, then $\delta_4 = (\lambda 1, 2, q_z 1, -1)$.
- (4) If r' = 2 and $\mu = 0$, then $\delta_4 = (\lambda 1, 1, q_z, -1)$.

In [9, Theorem 4.5], the author gives a minimal generating set of the defining ideal I(H). It is therefore sufficient to show that these binomials arise from $RF(\gamma_2)$ -relations.

- (1) Suppose $\mu > 0$ and r' = 1. Since r = 2 and r' = 1, we get $r_z = 1$ and $q = q' + q_z$. Then by [9, Theorem 4.5], the set $\{x_0x_2 x_1^2, x_0^{\lambda}x_3^w x_2^{q+1}, x_0^{\mu}x_1x_2^{q_z} x_3^v\}$ forms a minimal generating set for I(H). These binomials arise from the RF(γ_2)-relations corresponding to $\delta_{12} = (1, -2, 1, 0)$, $\delta_{13} = (\lambda, 0, -(q+1), w)$, and $\delta_{14} = (\mu, 1, q_z, -v)$ respectively.
- (2) Suppose $\mu > 0$ and r' = 2. Since r = 2 and r' = 2, we get $r_z = 2$ and $q = q' + q_z + 1$. Then by [9, Theorem 4.5], the set $\{x_0x_2 x_1^2, x_0^{\lambda}x_3^w x_2^{q+1}, x_0^{\mu}x_2^{qz+1} x_3^v\}$ forms a minimal generating set for I(H). These binomials arise from the RF(γ_2)-relations corresponding to $\delta_{12} = (1, -2, 1, 0), \, \delta_{13} = (\lambda, 0, -(q+1), w), \, \text{and} \, \delta_{14} = (\mu, 0, q_z + 1, -v)$ respectively.
- (3) Suppose $\mu=0$ and r'=1. Since $\mu=0$, we have z>p and hence w=0. Also, r'=1 implies $r_z=1$ and $q=q'+q_z+1$. Then by [9, Theorem 4.5], the set $\{x_0x_2-x_1^2,x_0^\lambda-x_2^{q+1},x_1x_2^{q_z+1}-x_3^v\}$ forms a minimal generating set for I(H). These binomials arise from the RF(γ_2)-relations corresponding to $\delta_{12}=(1,-2,1,0)$, $\delta_{13}=(\lambda,0,-(q+1),0)$, and $\delta_{34}=(0,1,q_z+1,-v)$ respectively.
- (4) Suppose $\mu = 0$ and r' = 2. Since r = 2 and r' = 2, we get $r_z = 2$ and $q = q' + q_z + 1$. Then by [9, Theorem 4.5], the set $\{x_0x_2 x_1^2, x_0^{\lambda} x_2^{q+1}, x_2^{q+1} x_3^{v}\}$ forms a minimal generating set for I(H). These binomials arise from the RF(γ_2)-relations corresponding to $\delta_{12} = (1, -2, 1, 0)$, $\delta_{13} = (\lambda, 0, -(q+1), 0)$, $\delta_{34} = (0, 0, q_z + 1, -v)$ respectively.

Proposition 6.2. Suppose $W \neq \emptyset$ and p = 2. If H is symmetric, then I(H) is minimally generated by RF-relations.

Proof. We have the following sub-cases.

Sub-case 1: Suppose r = 1, r' = 2, $\lambda = 1$, z < 2 and $PF(H) = {\alpha_3}$. Since $W \neq \emptyset$ and z < 2(=p), we get $\mu \neq 0$ and z = 1. Hence we have an RF-matrix for α_3 ,

$$RF(\alpha_3) = \begin{bmatrix} -1 & 1 & q' & v-1 \\ 0 & -1 & q'+1 & v-1 \\ \nu-1 & 1 & -1 & w-1 \\ \mu-1 & 2 & q' & -1 \end{bmatrix}.$$

Since $(\mu - 1)m_0 + 2m_1 + q'm_2 = (\nu - 2)m_0 + 2m_1 + (q - 1)m_2 = (\nu - 1)m_0 + qm_2$, we get another RF-matrix for α_3 ,

$$RF(\alpha_3)' = \begin{bmatrix} -1 & 1 & q' & v - 1 \\ 0 & -1 & q' + 1 & v - 1 \\ \nu - 1 & 1 & -1 & w - 1 \\ \nu - 1 & 0 & q & -1 \end{bmatrix}.$$

Let $\delta_1, \delta_2, \delta_3, \delta_4$ and $\delta'_1, \delta'_2, \delta'_3, \delta'_4$ denote the rows of RF(α_3) and RF(α_3)' respectively. By [9, Theorem 4.5], the set

$$\{x_0x_2-x_1^2, x_0^{\nu}-x_2^{q'+1}x_3^{\nu-w}, x_0^{\mu}x_1-x_3^{\nu}, x_1x_2^{q'+1}-x_0x_3^{w}, x_2^{q+1}-x_1x_3^{w}\}$$

forms a minimal generating set for I(H). These binomials arise from the RF(α_3)-relations corresponding to $\delta_{12}=(1,-2,1,0),\ \delta_{13}=(\nu,0,-(q'+1),-(v-w)),\ \delta_{14}=(\mu,1,0,-v),\ \delta_{34}=(-1,1,q'+1,-w)$ and RF(α_3)'-relation corresponding to $\delta_{34}'=(0,-1,q+1,-w)$ respectively.

Sub-case 2: Suppose r'=1, r=2, $\mu=0$, q'=0 and $PF(H)=\{\beta_2\}$. We have an RF-matrix for β_2 ,

$$RF(\beta_2) = \begin{bmatrix} -1 & 1 & q & v - w - 1 \\ 0 & -1 & q + 1 & v - w - 1 \\ \lambda - 1 & 1 & -1 & v - 1 \\ \lambda & 0 & q & -1 \end{bmatrix}.$$

Since $(q+1)m_2 + (v-w-1)n = \lambda m_0 + (v-1)n$, we get another RF-matrix for β_2 ,

$$RF(\beta_2)' = \begin{bmatrix} -1 & 1 & q & v - w - 1 \\ \lambda & -1 & 0 & v - 1 \\ \lambda - 1 & 1 & -1 & v - 1 \\ \lambda & 0 & q & -1 \end{bmatrix}.$$

Let $\delta_1, \delta_2, \delta_3, \delta_4$ and $\delta'_1, \delta'_2, \delta'_3, \delta'_4$ denote the rows of RF(β_2) and RF(β_2)' respectively. By [9, Theorem 4.5], the set

$$\{x_0x_2 - x_1^2, x_0^{\lambda}x_3^w - x_2^{q+1}, x_0^{\lambda+1} - x_1x_3^{v-w}, x_0^{\lambda}x_1 - x_2x_3^{v-w}, x_1x_2^q - x_3^v\}$$

forms a minimal generating set for I(H). These binomials arise from the RF(β_2)-relations corresponding to $\delta_{12} = (1, -2, 1, 0)$, $\delta_{13} = (\lambda, 0, -(q+1), w)$, $\delta_{14} = (\lambda + 1, -1, 0, -(v-w))$, $\delta_{24} = (\lambda, 1, -1, -(v-w))$ and RF(β_2)'-relation corresponding to $\delta'_{24} = (0, 1, q, -v)$ respectively.

Sub-case 3: Suppose $r=2, r'=2, \mu=0$ and $PF(H)=\{\beta_2\}$. We have an RF-matrix for β_2 ,

$$RF(\beta_2) = \begin{bmatrix} -1 & 1 & q & v - w - 1 \\ 0 & -1 & q + 1 & v - w - 1 \\ \lambda - 1 & 1 & -1 & v - 1 \\ \lambda - 1 & 1 & q_z & -1 \end{bmatrix}.$$

Let $\delta_1, \delta_2, \delta_3$ and δ_4 denote the rows of RF(β_2). By [9, Theorem 4.5], the set

$$\{x_0x_2 - x_1^2, x_0^{\lambda} - x_2^{q-q_z}x_3^{v-w}, x_2^{q_z+1} - x_3^v\}$$

forms a minimal generating set for I(H). These binomials arise from the RF(β_2)-relations corresponding to $\delta_{12}=(1,-2,1,0), \delta_{14}=(\lambda,0,-(q-q_z),-(v-w)),$ and $\delta_{34}=(0,0,q_z+1,-v)$ respectively.

Theorem 6.3. Let $H = \langle m_0, m_1, m_2, n \rangle$ be a symmetric numerical semigroup generated by an almost arithmetic sequence. Then I(H) has a minimal generating set consisting of RF-relations.

Proof. Combine Proposition 6.1 and Proposition 6.2.

Theorem 6.4. Let $H = \langle m_0, m_1, m_2, m_3, n \rangle$ be a symmetric numerical semigroup generated by an almost arithmetic sequence. Then I(H) has a minimal generating set consisting of RF-relations.

Proof. We use the minimal generating set of I(H) as given in [9, Theorem 4.5]. We first note that the subset of generators

$$\{\xi_{11} = x_0 x_2 - x_1^2, \ \xi_{12} = x_0 x_3 - x_1 x_2, \ \xi_{22} = x_1 x_3 - x_2^2\}$$

appears in every case below and that these binomials arise from the RF-relations corresponding to δ_{12} , δ_{13} and δ_{23} respectively, where δ_1 , δ_2 and δ_3 are rows of the respective RF-matrices in each case.

Case 1: Suppose $W = \emptyset$. Then p = 3, r = 2 and $PF(H) = \{\gamma_3\}$. By [9, Theorem 4.5], it follows that I(H) is minimally generated by

$$\{\xi_{11}, \ \xi_{12}, \ \xi_{22}, \ \phi_0 = x_0^{\lambda} x_4^w - x_2 x_3^q, \ \phi_1 = x_0^{\lambda - 1} x_1 x_4^w - x_3^{q+1}, \ \theta\},$$

where $\theta = x_0^{\mu} x_1 x_3^{q-q'} - x_4^{\nu}$ when r' = 1 and $\theta = x_0^{\mu} x_{5-r'} x_3^{q-q'-1} - x_4^{\nu}$ when r' = 2, 3. Also, by Corollary 3.6, the matrix

$$RF(\gamma_3) = \begin{bmatrix} -1 & 1 & 0 & q & v-1 \\ 0 & -1 & 1 & q & v-1 \\ 0 & 0 & -1 & q+1 & v-1 \\ \lambda-1 & 0 & 1 & -1 & w+v-1 \\ a_{5,1} & a_{5,2} & a_{5,3} & a_{5,4} & -1 \end{bmatrix},$$

where the last row depends on the value of μ . We observe that the binomial ϕ_1 arise from the RF-relations corresponding to δ_{24} . Since $m_1 + qm_3 = 2m_2 + (q-1)m_3$, we get another

RF-matrix for γ_3 ,

$$RF(\gamma_3)' = \begin{bmatrix} -1 & 0 & 2 & q-1 & v-1 \\ 0 & -1 & 1 & q & v-1 \\ 0 & 0 & -1 & q+1 & v-1 \\ \lambda-1 & 0 & 1 & -1 & w+v-1 \\ a_{5,1} & a_{5,2} & a_{5,3} & a_{5,4} & -1 \end{bmatrix}.$$

The binomial ϕ_0 arises from the RF(γ_3)'-relation corresponding to δ_{14} . In order to prove that θ arises from an RF-relation, we consider the following cases:

- (1) Let $\mu = 0$ and r' = 1. Then $\delta_5 = (\lambda 1, 1, 1, q_z 1, -1)$, w = 0 and θ arises from the RF-relation corresponding to δ_{45} .
- (2) Let $\mu = 0$ and r' = 2. Then $\delta_5 = (\lambda 1, 0, 1, q_z, -1)$, w = 0 and θ arises from the RF-relation corresponding to δ_{45} .
- (3) Let $\mu = 0$ and r' = 3. Then $\delta_5 = (\lambda 1, 0, 2, q_z 1, -1)$, w = 0 and θ arises from the RF-relation corresponding to δ_{45} .
- (4) Let $\mu > 0$ and r' = 1. Then $\delta_5 = (\mu 1, 2, 0, q + q_z, -1)$ and θ arises from the RF-relation corresponding to δ_{15} .
- (5) Let $\mu > 0$ and r' = 2. Then $\delta_5 = (\mu 1, 1, 0, q + q_z + 1, -1)$ and θ arises from the RF-relation corresponding to δ_{15} .
- (6) Let $\mu > 0$ and r' = 3. Then $\delta_5 = (\mu 1, 1, 1, q + q_z, -1)$ and θ arises from the RF-relation corresponding to δ_{15} .

Case 2: Suppose $W \neq \emptyset$. We have the following sub-cases.

Sub-case 1: Suppose r = 1, r' = 2, $\lambda = 1$, z < 3 and $PF(H) = {\alpha_4}$. We consider the following RF-matrices for α_4 ,

$$RF(\alpha_4) = \begin{bmatrix} -1 & 1 & 0 & q' & v - 1 \\ 0 & -1 & 1 & q' & v - 1 \\ 0 & 0 & -1 & q' + 1 & v - 1 \\ \mu & 0 & 1 & -1 & w - 1 \\ \mu - 1 & 1 & 1 & q' & -1 \end{bmatrix}, RF(\alpha_4)' = \begin{bmatrix} -1 & 1 & 0 & q' & v - 1 \\ 0 & -1 & 1 & q' & v - 1 \\ 0 & 0 & -1 & q' + 1 & v - 1 \\ \mu - 1 & 2 & 0 & -1 & w - 1 \\ \mu - 1 & 1 & 1 & q' & -1 \end{bmatrix},$$

and

$$RF(\alpha_4)'' = \begin{bmatrix} -1 & 0 & 2 & q'-1 & v-1 \\ 0 & -1 & 1 & q' & v-1 \\ 0 & 0 & -1 & q'+1 & v-1 \\ \mu & 0 & 1 & -1 & w-1 \\ \mu & 0 & 0 & q'+1 & -1 \end{bmatrix}.$$

For i = 1, ..., 5, let δ_i , δ'_i and δ''_i denote the *i*-th rows of RF(α_4), RF(α_4)' and RF(α_4)'' respectively. By [9, Theorem 4.5], the set

$$\left\{\xi_{11},\ \xi_{12},\ \xi_{22}\right\} \bigcup \left\{ \begin{array}{ll} \theta = x_0^{\mu} x_2 - x_4^{\nu}, & \psi_0 = x_0^{\mu+1} - x_2 x_3^{q'} x_4^{\nu-w}, & \psi_1 = x_0^{\mu} x_1 - x_3^{q'+1} x_4^{\nu-w}, \\ \phi_0 = x_0 x_4^{w} - x_1 x_3^{q}, & \phi_1 = x_1 x_4^{w} - x_2 x_3^{q}, & \phi_2 = x_2 x_4^{w} - x_3^{q+1} \end{array} \right\}$$

forms a minimal generating set for I(H). The binomials θ , ϕ_0 and ψ_1 arise from the RF(α_4)-relations corresponding to δ_{15} , δ_{45} , and δ_{24} respectively. The binomials ϕ_1 , ϕ_2 and ψ_0 arise from the RF(α_4)'-relations corresponding to δ'_{45} and RF(α_3)"-relations corresponding to δ''_{45}

and δ_{14}'' respectively.

Sub-case 2: Suppose $r'=1, r=2, \mu=0, q'=0$ and $PF(H)=\{\beta_3\}$. We consider the following RF-matrices for β_3 ,

$$RF(\beta_3) = \begin{bmatrix} -1 & 1 & 0 & q & v - w - 1 \\ 0 & -1 & 1 & q & v - w - 1 \\ 0 & 0 & -1 & q + 1 & v - w - 1 \\ \lambda - 1 & 0 & 1 & -1 & v - 1 \\ \lambda & 0 & 0 & q & -1 \end{bmatrix}, RF(\beta_3)' = \begin{bmatrix} -1 & 0 & 2 & q - 1 & v - w - 1 \\ 0 & -1 & 1 & q & v - w - 1 \\ 0 & 0 & -1 & q + 1 & v - w - 1 \\ \lambda - 1 & 0 & 1 & -1 & v - 1 \\ \lambda - 1 & 1 & 1 & q - 1 & -1 \end{bmatrix}.$$

Let $\delta_1, \delta_2, \delta_3, \delta_4, \delta_5$ and $\delta'_1, \delta'_2, \delta'_3, \delta'_4, \delta'_5$ denote the rows of RF(β_3) and RF(β_3)' respectively. By [9, Theorem 4.5], the set

$$\left\{\xi_{11},\ \xi_{12},\ \xi_{22}\right\} \bigcup \left\{ \begin{array}{ll} \theta = x_1 x_3^q - x_4^v, & \phi_0 = x_0^{\lambda} x_4^w - x_2 x_3^q, & \phi_1 = x_0^{\lambda - 1} x_1 x_4^w - x_3^{q + 1}, \\ \psi_0 = x_0^{\lambda + 1} - x_1 x_4^{v - w}, & \psi_1 = x_0^{\lambda} x_1 - x_2 x_4^{v - w}, & \psi_2 = x_0^{\lambda} x_2 - x_3 x_4^{v - w} \end{array} \right\}$$

forms a minimal generating set for I(H). The binomials ϕ_1 , ψ_0 , ψ_1 and ψ_2 arise from the RF(β_3)-relations corresponding to δ_{24} , δ_{15} , δ_{25} , and δ_{35} respectively. The binomials θ and ϕ_0 arise from the RF(β_3)'-relations corresponding to δ'_{45} and δ'_{14} respectively.

Sub-case 3: Suppose r = 2, r' = 2, $\mu = 0$ and $PF(H) = {\beta_3}$. We consider the following RF-matrices for β_3 ,

$$RF(\beta_3) = \begin{bmatrix} -1 & 1 & 0 & q & v - w - 1 \\ 0 & -1 & 1 & q & v - w - 1 \\ 0 & 0 & -1 & q + 1 & v - w - 1 \\ \lambda - 1 & 0 & 1 & -1 & v - 1 \\ \lambda - 1 & 0 & 1 & q_z & -1 \end{bmatrix}, RF(\beta_3)' = \begin{bmatrix} -1 & 0 & 2 & q - 1 & v - w - 1 \\ 0 & -1 & 1 & q & v - w - 1 \\ 0 & 0 & -1 & q + 1 & v - w - 1 \\ \lambda - 1 & 0 & 1 & -1 & v - 1 \\ \lambda - 1 & 0 & 1 & q_z & -1 \end{bmatrix}.$$

Let $\delta_1, \delta_2, \delta_3, \delta_4, \delta_5$ and $\delta'_1, \delta'_2, \delta'_3, \delta'_4, \delta'_5$ denote the rows of RF(β_3) and RF(β_3)' respectively. By [9, Theorem 4.5], the set

$$\{\xi_{11}, \xi_{12}, \xi_{22}, \theta = x_3^{q_z+1} - x_4^v, \psi_0 = x_0^{\lambda} - x_2 x_3^{q'} x_4^{v-w}, \psi_1 = x_0^{\lambda-1} x_1 - x_3^{q-q_z} x_4^{v-w}\}$$

forms a minimal generating set for I(H). The binomials θ and ψ_1 arise from the RF(β_3)-relations corresponding to δ_{45} and δ_{25} respectively and ψ_0 arises from the RF(β_3)'-relation corresponding to δ'_{15} .

With the knowledge of a minimal generating set of I(H) from [9], one may continue this process in higher embedding dimension as well. The non-generic nature of I(H) and the non-uniqueness of RF(H) does suggest that the answer to the question is in the affirmative in higher embedding dimension as well. But the same reason turns out to be stumble block when one tries to prove the result in general.

References

- [1] Kazufumi Eto. Almost Gorenstein monomial curves in affine four space. J. Algebra, 488:362–387, 2017.
- [2] Kazufumi Eto. On row-factorization matrices. Journal of Algebra, Number Theory: Advances and Applications, 17(2):93–108, 2017. 1
- [3] Kazufumi Eto. Generic Toric Ideals and Row-Factorization Matrices in Numerical Semigroups, pages 83–91. Springer International Publishing, Cham, 2020. 1, 2, 17, 20

- [4] Vesselin Gasharov, Irena Peeva, and Volkmar Welker. Rationality for generic toric rings. *Math. Z.*, 233(1):93–102, 2000. 1
- [5] Jürgen Herzog and Kei-ichi Watanabe. Almost symmetric numerical semigroups. Semigroup Forum, 98(3):589–630, 2019. 1, 2, 20
- [6] Alessio Moscariello. On the type of an almost Gorenstein monomial curve. J. Algebra, 456:266-277, 2016. 1, 3
- [7] T. Numata. Almost symmetric numerical semigroups generated by four elements. Proc. Inst. Nat. Sci., Nihon Univ., 48:197–207, 2013. 1
- [8] D. P. Patil and I. Sengupta. Minimal set of generators for the derivation module of certain monomial curves. Comm. Algebra, 27(11):5619–5631, 1999. 2, 3, 4, 10, 15, 16
- [9] Dilip P. Patil. Minimal sets of generators for the relation ideals of certain monomial curves. *Manuscripta Math.*, 80(3):239–248, 1993. 16, 21, 22, 23, 24, 25
- [10] Dilip P. Patil and Balwant Singh. Generators for the derivation modules and the relation ideals of certain curves. *Manuscripta Math.*, 68(3):327–335, 1990. 2, 3
- [11] Irena Peeva and Bernd Sturmfels. Generic lattice ideals. J. Amer. Math. Soc., 11(2):363–373, 1998. 1, 17
- [12] J. C. Rosales and P. A. García-Sánchez. Numerical semigroups, volume 20 of Developments in Mathematics. Springer, New York, 2009. 1, 18
- [13] Joydip Saha and Indranath Sengupta. Derivation modules for sum and gluing. arXiv perprint arXiv:1909.01611, pages 1–10, 2020. 18

IIT GANDHINAGAR, PALAJ, GANDHINAGAR, GUJARAT-382355 INDIA

Email address: om.prakash@iitgn.ac.in

Email address: kriti.g@iitgn.ac.in

Email address: indranathsg@iitgn.ac.in