

Explicit constructions of optimal linear codes with Hermitian hulls and their application to quantum codes

Lin Sok

Abstract

We prove that any Hermitian self-orthogonal $[n, k, d]_{q^2}$ code gives rise to an $[n, k, d]_{q^2}$ code with ℓ dimensional Hermitian hull for $0 \leq \ell \leq k$. We present a new method to construct Hermitian self-orthogonal $[n, k]_{q^2}$ codes with large dimensions $k > \frac{n+q-1}{q+1}$. New families of Hermitian self-orthogonal codes with good parameters are obtained; more precisely those containing almost MDS codes. By applying a puncturing technique to Hermitian self-orthogonal codes, MDS $[n, k]_{q^2}$ linear codes with Hermitian hull having large dimensions $k > \frac{n+q-1}{q+1}$ are also derived. New families of MDS, almost MDS and optimal codes with arbitrary Hermitian hull dimensions are explicitly constructed from algebraic curves. As an application, we provide entanglement-assisted quantum error correcting codes with new parameters.

Keywords: Hulls, MDS codes, almost MDS codes, self-orthogonal codes, algebraic curves, algebraic geometry codes, entanglement-assisted quantum error-correcting codes

I. INTRODUCTION

Quantum error correcting codes have attracted a lot of attention recently due to their capability to protect information carrying quantum states against decoherence in quantum information systems. A lot of effort has been done to construct good quantum codes from the classical error correcting codes.

Entanglement is one of the best approaches to achieve higher rates in quantum information systems. Entanglement-assisted quantum error-correcting codes (EAQECCs) were firstly introduced by Bowen [4]. Later, they were developed by Brun *et al.* [5]. The authors [5] showed that if pre-shared entanglement between the encoder and decoder is available, the EAQECCs can be constructed via classical linear codes without self-orthogonality as in [1]. Then Wilde *et al.* [37] proposed two methods to construct EAQECCs from classical codes; the Euclidean and Hermitian construction methods. The parameters of EAQECCs can be determined from those of hulls of linear codes (see [14]). For the works on EAQECCs from Euclidean hulls, the reader is referred to [8], [7], [28], [31], [32], [33], [34], [15], [29]. In [15], Guenda *et al.* investigated the ℓ -intersection pair of linear codes, where they completely determined the q -ary MDS EAQECCs of length $n \leq q + 1$ for all possible parameters. EAQECCs with zero pre-shared entanglements, known as quantum stabilizer codes, can be constructed from Hermitian linear complementary dual codes using method [8]. Now the remaining problem is to construct new q -ary MDS EAQECCs of length $n > q + 1$ with hull dimension $\ell > 1$. This can be achieved by considering Hermitian hulls of linear codes, see for example [12], [30]. It has been known in [12] that given n, k and q , it is a very difficult problem to construct new MDS Hermitian hull codes with parameters $[n, k]_{q^2}$ for new lengths n or for new dimensions $k > \frac{n+q-1}{q+1}$.

Algebraic geometry codes were invented by Goppa, where in some literature they were also called geometric Goppa codes. In his paper [13], Goppa showed how to construct linear codes from algebraic curves over a finite field. A very nice property of these codes is that their parameters can be determined from the degree of a divisor associated. Despite a strongly theoretical construction, algebraic geometry (AG) codes have asymptotically good parameters, and this was the first time that linear codes were proved to have improved the so-called Gilbert-Vasharmov bound. Algebraic geometry codes are good candidates for constructing quantum codes, see for example [20], [19], [30], [34], [35]. In [30], some good EAQECCs were constructed from AG codes of arbitrary Hermitian hull dimensions, but their constructions are not explicit.

In this paper, we study linear codes with Hermitian hull using tools from algebraic function fields. We introduce a new method to construct Hermitian hull $[n, k]$ codes. We prove that any Hermitian self-orthogonal $[n, k]$ code gives rise to a linear code with Hermitian hull of dimension less than k and with the same parameters as the former codes, and thus new families of MDS Hermitian hull codes are obtained in Theorem 4.1). New constructions of Hermitian self-orthogonal codes is presented in Theorems 1, 2 and 3, which give a generalization of the recent result [35]. From an MDS Hermitian self-orthogonal code, this new method enables us to construct (almost MDS) Hermitian self-orthogonal codes and thus Hermitian hull codes with dimension $k > \frac{n+q-1}{q+1}$. More specifically, many new almost MDS Hermitian hull codes are constructed (see Theorems 5 and 7). By applying a puncturing technique to Hermitian self-orthogonal codes, MDS Hermitian hull codes with large dimension

$k > \frac{n+q-1}{q+1}$ are derived (see Theorem 6 and 7), and these codes are new compared with those explicitly constructed in [12]. Theorem 8 enlarges families of Hermitian self-orthogonal codes [35]. Except for few values in Theorem 9, all families of EAQECCs obtained in Theorems 9-17 are new.

The paper is organized as follows: Section II gives preliminaries and background on algebraic geometry codes. Section III provides a new construction method for Hermitian self-orthogonal codes and for linear codes with arbitrary Hermitian hull dimensions. In Section IV, Hermitian hull codes are constructed from projective lines, elliptic curves, hyper-elliptic curves and Hermitian curves. In the last section, we give an application of Hermitian hull codes to EAQECCs.

II. PRELIMINARIES

A. Linear codes

Let \mathbb{F}_q be the finite field with q elements. A linear code of length n , dimension k and minimum distance d over \mathbb{F}_q is denoted as $[n, k, d]_q$ code. If C is an $[n, k, d]_q$ code, then from the Singleton bound, its minimum distance is upper bounded by

$$d \leq n - k + 1.$$

A code is called *Maximum Distance Separable* (MDS) if its minimum distance $d = n - k + 1$ and is called *almost* MDS (AMDS) if it is minimum distance $d = n - k$. A code is called *optimal* if it has the highest possible minimum distance for its length and dimension.

The *Hermitian* inner product of $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_n)$ in $\mathbb{F}_{q^2}^n$ is defined by

$$\langle \mathbf{x}, \mathbf{y} \rangle_h = \sum_{i=1}^n x_i y_i^q.$$

The Hermitian *dual* of C , denoted by C^{\perp_h} , is the set of vectors orthogonal to every codeword of C under the Hermitian inner product. A linear code C is called Hermitian self-orthogonal if $C \subseteq C^{\perp_h}$. The Hermitian hull of a linear code C is

$$\text{Hull}_h(C) := C \cap C^{\perp_h}.$$

A linear code C is called *Hermitian ℓ -dim hull* if $\dim(\text{Hull}_h(C)) = \ell$.

For a linear code $C \subseteq \mathbb{F}_q^n$ and $\mathbf{v} = (v_1, \dots, v_n) \in (\mathbb{F}_q^*)^n$, we define

$$\mathbf{v} \cdot C := \{(v_1 c_1, \dots, v_n c_n) \mid \mathbf{c} = (c_1, \dots, c_n) \in C\}.$$

It is easy to see that $\mathbf{v} \cdot C$ is a linear code if and only if C is a linear code. Moreover, both codes have the same dimension, minimum Hamming distance, and weight distribution.

For undefined terms related to algebraic function fields, the reader is referred to Stichtenoth [36].

Let \mathcal{X} be a smooth projective curve of genus g over \mathbb{F}_q . We denote the field of rational functions of \mathcal{X} by $\mathbb{F}_q(\mathcal{X})$. Function fields of algebraic curves over a finite field are finite separable extensions of $\mathbb{F}_q(x)$. Points on the curve \mathcal{X} identified with places of the function field $\mathbb{F}_q(\mathcal{X})$. We call a point on \mathcal{X} rational if all of its coordinates belong to \mathbb{F}_q . Rational points can be identified with places of degree one. The set of \mathbb{F}_q -rational points on \mathcal{X} is denoted by $\mathcal{X}(\mathbb{F}_q)$.

We define a divisor G on the curve \mathcal{X} to be the formal sum $\sum_{P \in \mathcal{X}} n_P P$ with only finitely many nonzeros $n_P \in \mathbb{Z}$. A divisor G on \mathcal{X} is called rational if for any σ in $\text{Gal}(\overline{\mathbb{F}_q}/\mathbb{F}_q)$, we have $G^\sigma = G$. We define the support of G as $\text{supp}(G) := \{P \in \mathcal{X} \mid n_P \neq 0\}$. For $G = \sum_{P \in \mathcal{X}} n_P P$, the degree of G is defined by $\deg(G) := \sum_{P \in \mathcal{X}} n_P \deg(P)$, where $\deg(P) = |P^\sigma|$ is the size of orbit of P under the action σ . For two divisors $G = \sum_{P \in \mathcal{X}} n_P P$ and $H = \sum_{P \in \mathcal{X}} m_P P$, we say that $G \geq H$ if $n_P \geq m_P$ for all places $P \in \mathcal{X}$.

For a nonzero rational function f on the curve \mathcal{X} , we define the ‘‘principal’’ divisor of f as $(f) := \sum_{P \in \mathcal{X}} v_P(f)P$, where v_P denotes the normalized discrete valuation corresponding to the place P . If $Z(f)$ and $N(f)$ denotes the set of zeros and poles of f respectively, we define the zero divisor and pole divisor of f , respectively by $(f)_0 := \sum_{P \in Z(f)} v_P(f)P$, $(f)_\infty := \sum_{P \in N(f)} -v_P(f)P$.

For a divisor G on the curve \mathcal{X} , we define

$$\mathcal{L}(G) := \{f \in \mathbb{F}_q(\mathcal{X}) \setminus \{0\} \mid (f) + G \geq 0\} \cup \{0\},$$

and

$$\Omega(G) := \{\omega \in \Omega \setminus \{0\} \mid (\omega) - G \geq 0\} \cup \{0\},$$

where $\Omega := \{f dx \mid f \in \mathbb{F}_q(\mathcal{X})\}$, the set of differential forms on \mathcal{X} . It is well-known that both $\mathcal{L}(G)$ and $\Omega(G)$ are finite dimensional vector spaces. Moreover, for a differential form ω on \mathcal{X} , there exists a unique a rational function f on \mathcal{X} such that

$\omega = f dt$, where t is a local uniformizing parameters. In this case, we define the divisor associated to ω by $(\omega) = \sum_{P \in \mathcal{X}} v_P(\omega)P$, where $v_P(\omega) := v_P(f)$. The divisor class of a nonzero differential form is called the canonical divisor. It is well-known that if K is the canonical divisor, then $\deg(K) = 2g$.

The dimension of $\mathcal{L}(G)$ is determined by Riemann-Roch's theorem as follows.

Proposition 1. [36, Theorem 1.5.15 (Riemann-Roch)] *Let W be a canonical divisor. Then, for each divisor G , the following holds:*

$$\dim \mathcal{L}(G) = \deg G + 1 - g + \dim \mathcal{L}(W - G),$$

where g is the genus of the smooth algebraic curve.

Let $D = P_{\alpha_1} + \dots + P_{\alpha_n}$, where $(P_{\alpha_i})_{1 \leq i \leq n}$ are places of degree one, and G a divisor with $\text{supp}(D) \cap \text{supp}(G) = \emptyset$. Define the algebraic geometry code by

$$C_{\mathcal{L}}(D, G) := \{(f(P_{\alpha_1}), \dots, f(P_{\alpha_n})) \mid f \in \mathcal{L}(G)\},$$

and the differential algebraic geometry code as

$$C_{\Omega}(D, G) := \{(\text{Res}_{P_{\alpha_1}}(\omega), \dots, \text{Res}_{P_{\alpha_n}}(\omega)) \mid \omega \in \Omega(G - D)\},$$

where $\text{Res}_P(\omega)$ denotes the residue of ω at point P .

For $\mathbf{v} = (v_1, \dots, v_n)$ with $v_i \in \mathbb{F}_{q^2}^*$, define also

$$\mathcal{L}_q(G) := \{f \in \mathcal{L}(G) \mid f(\alpha) \in \mathbb{F}_q, \forall \alpha \in \mathbb{F}_{q^2}\}, \quad (1)$$

$$C_{\mathcal{L}_q}(D, G) := \{(f(P_{\alpha_1}), \dots, f(P_{\alpha_n})) \mid f \in \mathcal{L}_q(G)\}, \quad (2)$$

and

$$C_{\mathcal{L}_q}(D, G; \mathbf{v}) := \{(v_1 f(P_{\alpha_1}), \dots, v_n f(P_{\alpha_n})) \mid f \in \mathcal{L}_q(G)\}. \quad (3)$$

The parameters of an algebraic geometry code $C_{\mathcal{L}}(D, G)$ is given as follows.

Proposition 2. [36, Corollary 2.2.3] *Assume that $2g - 2 < \deg(G) < n$. Then the code $C_{\mathcal{L}}(D, G)$ has parameters $[n, k, d]$ satisfying*

$$k = \deg(G) - g + 1 \text{ and } d \geq n - \deg(G). \quad (4)$$

B. Quantum codes

Let $(\mathbb{C})^{\otimes n} (\cong \mathbb{C}^{q^n})$ be the q^n -dimensional Hilbert space over the complex field \mathbb{C} . A quantum code of length n is a subspace of \mathbb{C}^{q^n} . Let $\{|\mathbf{a}\rangle = |a_1\rangle \otimes |a_2\rangle \otimes \dots \otimes |a_n\rangle : (a_1, a_2, \dots, a_n) \in \mathbb{F}_q^n\}$ be a basis of \mathbb{C}^{q^n} . We define the inner product of two quantum states $|\phi_1\rangle = \sum_{\mathbf{a} \in \mathbb{F}_q^n} \phi_1(\mathbf{a})|\mathbf{a}\rangle$ and $|\phi_2\rangle = \sum_{\mathbf{a} \in \mathbb{F}_q^n} \phi_2(\mathbf{a})|\mathbf{a}\rangle$ by

$$\langle \phi_1 | \phi_2 \rangle = \sum_{\mathbf{a} \in \mathbb{F}_q^n} \overline{\phi_1(\mathbf{a})} \phi_2(\mathbf{a}) \in \mathbb{C},$$

where $\overline{\phi_1(\mathbf{a})}$ is the complex conjugate of $\phi_1(\mathbf{a})$. Two quantum states $|\phi_1\rangle$ and $|\phi_2\rangle$ are *orthogonal* if $\langle \phi_1 | \phi_2 \rangle = 0$.

The rules of $X(\mathbf{a})$ and $Z(\mathbf{b})$ on $|\mathbf{v}\rangle \in \mathbb{C}^{q^n}$ ($\mathbf{v} \in \mathbb{F}_q^n$) are given as

$$X(\mathbf{a})|\mathbf{v}\rangle = |\mathbf{v} + \mathbf{a}\rangle \text{ and } Z(\mathbf{b})|\mathbf{v}\rangle = \zeta_p^{\text{tr}(\langle \mathbf{v}, \mathbf{b} \rangle_E)} |\mathbf{v}\rangle, \quad (5)$$

respectively, where $\text{tr}(\cdot)$ is the trace function from \mathbb{F}_q to \mathbb{F}_p and ζ_p is a complex primitive p -th root of unity.

For $\mathbf{a} = (a_1, \dots, a_n), \mathbf{b} = (b_1, \dots, b_n) \in \mathbb{F}_q^n$, we can also write, from (5), $X(\mathbf{a}) = X(a_1) \otimes \dots \otimes X(a_n)$ and $Z(\mathbf{b}) = Z(b_1) \otimes \dots \otimes Z(b_n)$ for the tensor product of n (error) operators. The set $\mathcal{E}_n \{X(\mathbf{a})Z(\mathbf{b}) : \mathbf{a}, \mathbf{b} \in \mathbb{F}_q^n\}$ is an error basis on \mathbb{C}^{q^n} . The error group G_n is generated by the matrices in \mathcal{E}_n and is defined by

$$G_n := \{\zeta_p^t X(\mathbf{a})Z(\mathbf{b}) : \mathbf{a}, \mathbf{b} \in \mathbb{F}_q^n, t \in \mathbb{F}_p\}.$$

For $E = \zeta_p^t X(\mathbf{a})Z(\mathbf{b}) \in G_n$, we define the *quantum weight* $\text{wt}_Q(E)$ of E to be the number of coordinates such that $(a_i, b_i) \neq (0, 0)$. A quantum code Q is called a $d - 1$ quantum error detecting code ($d \geq 1$) if, for any pair $|\phi_1\rangle$ and $|\phi_2\rangle$ in Q with $\langle \phi_1 | \phi_2 \rangle = 0$ and any $E \in G_n$ with $\text{wt}_Q(E) \leq d - 1$, $|\phi_1\rangle$ and $E|\phi_2\rangle$ are orthogonal. The quantum code Q is said to have minimum distance d if d is the largest integer such that for any $|\phi_1\rangle, |\phi_2\rangle \in Q$ with $\langle \phi_1 | \phi_2 \rangle = 0$ and any $E \in G_n$ with $\text{wt}_Q(E) \leq d - 1$, we have that $|\phi_1\rangle$ and $E|\phi_2\rangle$ are orthogonal. We write $[[n, k, d]]_q$ for a q -ary quantum code of length n , dimension k and minimum distance d . It is well-known that a quantum code with minimum distance d can detect up to $d - 1$ errors and correct up to $\frac{d-1}{2}$ quantum errors.

For S being an abelian subgroup of G_n , we define the quantum stabilizer codes $C(S)$ by

$$C(S) := \{|\phi\rangle : E|\phi\rangle = |\phi\rangle, \forall E \in S\}.$$

It was shown in [6], [1], [24] that such codes can be constructed from classical linear codes with some properties of self-orthogonality. However, the above construction method is not applicable anymore if the subgroup S of G_n is non-abelian. To improve the construction, Brun *et al.* [5] introduced the so-called entanglement-assisted quantum error-correcting codes (EAQECCs). In their method, they extended S to be a new abelian subgroup in a larger error group, and they assumed that a sender and a receiver shared a certain amount of pre-existing entangled bits (ebits), which was not subject to errors.

We use $[[n, k, d; c]]_q$ to denote a q -ary $[[n, k, d]]_q$ quantum code that utilizes c pre-shared entanglement pairs. For $c = 0$, an $[[n, k, d; c]]_q$ EAQECC is equivalent to a quantum stabilizer code [1].

The constraints among the parameters of an EAQECC are given in the following lemma [26].

Proposition 3. (Quantum Singleton Bound) *For any $[[n, k, d; c]]_q$ -EAQECC, if $d \leq \frac{n+2}{2}$, we have*

$$2(d-1) \leq n + c - k.$$

When the bound meets with equality, the EAQECC is called MDS, and it is called AMDS if its minimum distance is one unit less than the MDS case.

We denote the conjugate transpose of a matrix M by M^\dagger , that is, if $M = m_{i,j}$ then $M^\dagger = m_{j,i}^q$.

Lemma 1. ([5]) *Let P be the parity check matrix of an $[n, k, d]_{q^2}$ code C . Then there exists an $[[n, 2k - n + c, d; c]]_q$ EAQECC \mathcal{Q} , where $c = \text{rank}(PP^\dagger)$ is the required number of maximally entangled states. In particular, if C is an MDS code and $d \leq \frac{n+2}{2}$, then \mathcal{Q} is an MDS EAQECC.*

Lemma 2. [14] *Let C be a classical $[n, k, d]_{q^2}$ code with parity check matrix P and generator matrix G . Then $\text{rank}(PP^\dagger)$ and $\text{rank}(GG^\dagger)$ are independent of P and G so that*

$$\begin{aligned} \text{rank}(PP^\dagger) &= n - k - \dim(\text{Hull}_h(C)) \\ &= n - k - \dim(\text{Hull}_h(C^{\perp_h})), \end{aligned}$$

and

$$\begin{aligned} \text{rank}(GG^\dagger) &= k - \dim(\text{Hull}_h(C)) \\ &= k - \dim(\text{Hull}_h(C^{\perp_h})). \end{aligned}$$

III. NEW CONSTRUCTIONS OF HERMITIAN ℓ -DIM HULL CODES

In this section, we first show that any Hermitian ℓ -dim hull $[n, k, d]_{q^2}$ code can be constructed from a Hermitian self-orthogonal $[n, k, d]_{q^2}$ code. Then we present a new method to embed a Hermitian self-orthogonal code into another one.

Lemma 3. *Assume that there exists a Hermitian self-orthogonal code with parameters $[n, k, d]_{q^2}$. Then there exists a Hermitian ℓ -dim hull $[n, k, d]_{q^2}$ code for $0 \leq \ell \leq k$.*

Proof. Assume that C is a Hermitian self-orthogonal code with parameters $[n, k, d]$ over \mathbb{F}_{q^2} . Let G be the generator matrix of C . Then up to equivalence, we can write G as

$$G = (I_k | A),$$

where A satisfies $AA^\dagger = -I_k$. Choose $\lambda \in \mathbb{F}_{q^2}$ such that $\lambda^{q+1} \neq 1$, and take $G_\ell = \text{diag}(\underbrace{\lambda, \dots, \lambda}_{k-\ell}, \underbrace{1, \dots, 1}_\ell | A)$. Then it is easy

to see that $\text{rank}(G_\ell G_\ell^\dagger)$ is exactly $k - \ell$, and thus the result follows from Lemma 2. \square

We now state sufficient conditions for an algebraic geometry $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ to be Hermitian self-orthogonal and introduce new constructions of such codes.

Lemma 4. [35] *Let $D = P_{\alpha_1} + \dots + P_{\alpha_n}$ and $G = (k-1)O$ be two divisors, ω be a Weil differential form such that $H = D - G + (\omega)$ and $\mathbf{v} = (v_1, \dots, v_n)$ with $v_i \in \mathbb{F}_{q^2}^*$. Then the code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ is Hermitian self-orthogonal if the following conditions hold*

- 1) $G \leq H$,
- 2) $\text{Res}_{P_{\alpha_i}}(\omega) = v_i^{q+1}$,
- 3) $g + 1 \leq k \leq \lfloor \frac{n+q+2g-1}{q+1} \rfloor$, where g is the genus of the defining curve.

The following lemma is useful for embedding an Hermitian self-orthogonal code into another one.

Lemma 5. [35] *Assume that $D = P_{\alpha_1} + \dots + P_{\alpha_n}$, $G = (k-1)O$, $1 \leq k \leq \lfloor \frac{n+q-1}{q+1} \rfloor$ and $\text{Res}_{P_{\alpha_i}}(\omega) = v_i^{q+1}$ for some $v_i \in \mathbb{F}_{q^2}^*$ for $1 \leq i \leq n$. Then an MDS Hermitian q^2 -ary self-orthogonal code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ with parameters $[n, k]$ can be embedded into an MDS Hermitian q^2 -ary self-orthogonal $[n+1, k+1]$ code.*

In [35], MDS Hermitian self-orthogonal codes are constructed up to dimension q . The following theorem extends the embedding Lemma 5.

Theorem 1. Assume that $(n-1)|(q^2-1)$, $D = P_{\alpha_1} + \dots + P_{\alpha_n}$, $G = (k-1)O$, $1 \leq k \leq \lfloor \frac{n+q-1}{q+1} \rfloor$ and $\text{Res}_{P_{\alpha_i}}(\omega) = v_i^{q+1}$ for some $v_i \in \mathbb{F}_{q^2}^*$ for $1 \leq i \leq n$. Put $n_i = n+i$ and $k_i = k+i$ for $1 \leq i \leq q$. Then an MDS Hermitian q^2 -ary self-orthogonal code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ with parameters $[n, k]$ can be embedded into a Hermitian q^2 -ary self-orthogonal code C_i with parameters

- 1) $[n+1, k+2, \geq n-k-1]_{q^2}$ if $(n-1)|(k+1)(q+1)$;
- 2) $[n_i-1, k_i, \geq n-k-i+1]_{q^2}$ for $2 \leq i \leq q$ if $(n-1)|k(q+1)$.

Proof. Take $U_{n-1} = \{\alpha \in \mathbb{F}_{q^2} | \alpha^{n-1} = 1\} = \{\alpha_1, \dots, \alpha_{n-1}\}$, and set $U = \{\alpha_1, \dots, \alpha_{n-1}, \alpha_n\}$, where $\alpha_n = 0$. It should be noted that under the assumption in the theorem, there exists an MDS Hermitian self-orthogonal code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ with parameters $[n, k]_{q^2}$, where $D = \sum_{\alpha \in U} P_\alpha = P_{\alpha_1} + \dots + P_{\alpha_n}$. Consider the code C_i with its generator matrix \mathcal{G}_i written as follows:

$$\mathcal{G}_i = \begin{pmatrix} v_1 & \dots & v_n & 0 & 0 & \dots & 0 \\ v_1 \alpha_1 & \dots & v_n \alpha_n & 0 & 0 & \dots & 0 \\ \vdots & \dots & \vdots & \vdots & \vdots & \dots & \vdots \\ v_1 \alpha_1^{k-1} & \dots & v_n \alpha_n^{k-1} & 0 & 0 & \dots & 0 \\ v_1 \alpha_1^k & \dots & v_n \alpha_n^k & \lambda_1 & 0 & \dots & 0 \\ v_1 \alpha_1^{k+1} & \dots & v_n \alpha_n^{k+1} & 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \dots & \vdots \\ v_1 \alpha_1^{k+i-1} & \dots & v_n \alpha_n^{k+i-1} & 0 & 0 & \dots & \lambda_i \end{pmatrix}. \quad (6)$$

Note that the first k rows of \mathcal{G}_i generates $C_{\mathcal{L}_q}(D, G; \mathbf{v})$. We now calculate the minimum distance of C_i . First, observe that by puncturing the last i coordinates of C_i , we obtain an MDS code with parameters $[n, k_i, n-k_i+1]$. Hence, the code C_i has parameters $[n_i, k_i, \geq n-k-i+1]$.

- 1) for $i = 2$, consider the following system of $(k+1)(q+1)+1$ equations with n indeterminates $v_1^{q+1}, \dots, v_n^{q+1}$ defined by

$$M\mathbf{v} = \mathbf{a}, \quad (7)$$

where

$$M = \begin{pmatrix} 1 & \dots & 1 \\ \alpha_1 & \dots & \alpha_n \\ \vdots & \dots & \vdots \\ \alpha_1^{(k-1)(q+1)} & \dots & \alpha_n^{(k-1)(q+1)} \\ \vdots & \dots & \vdots \\ \alpha_1^{k(q+1)} & \dots & \alpha_n^{k(q+1)} \\ \vdots & \dots & \vdots \\ \alpha_1^{(k+1)(q+1)} & \dots & \alpha_n^{(k+1)(q+1)} \end{pmatrix}, \quad \mathbf{v} = \begin{pmatrix} v_1^{q+1} \\ v_2^{q+1} \\ \vdots \\ v_n^{q+1} \end{pmatrix}, \quad \mathbf{a} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ -\lambda_1^{q+1} \\ 0 \\ \vdots \\ 0 \\ -\lambda_2^{q+1} \end{pmatrix}.$$

Then C_2 is Hermitian self-orthogonal if and only if (7) has a solution. It should be noted that λ_1 does exist by Lemma 5. Since the first $k+1$ rows g_1, \dots, g_k, g_{k+1} of \mathcal{G}_2 generates a Hermitian self-orthogonal $[n+2, k+1]$ code, it remains to check that $\langle g_{k+1}, g_{k+2} \rangle_h = 0$ and $\langle g_{k+2}, g_{k+2} \rangle_h = -\lambda_2^{q+1}$ for some $\lambda_2 \in \mathbb{F}_{q^2}$. Now under the condition $(n-1)|(k+1)(q+1)$, we have that $(\alpha_1^{(k+1)(q+1)}, \dots, \alpha_n^{(k+1)(q+1)}) = (1, \dots, 1, 0)$, and so $v_1^{q+1} \alpha_1^{(k+1)(q+1)} + \dots + v_n^{q+1} \alpha_n^{(k+1)(q+1)} = -\lambda_2^{q+1}$ has a solution. Thus, λ_2 is well determined (here $\lambda_2 = v_n$). Now, we check that $\langle g_{k+1}, g_{k+2} \rangle_h = 0$, that is,

$$\sum_{i=1}^n \alpha_i^{kq+k+1} v_i^{q+1} = 0. \quad (8)$$

- a) If $k(q+1)+1 \leq n$, then following the same reasoning as in the proof of [35, Lemma 4], (10) holds with $v_i^{q+1} = \text{Res}_{P_{\alpha_i}}(\omega)$.
- b) If $k(q+1)+1 > n$, then we can write $k(q+1)+1 = (n-1)A+B$ with $0 \leq B < (n-1) < k(q+1)$ and $\alpha_i^{kq+k+1} = \alpha_i^B$, and thus (10) also holds.

Hence, the system (7) has a solution. Moreover, with $\lambda_2 = 0$, one can puncture the $(n+2)$ -th coordinate and obtain the code with parameters $[n+1, k+2, \geq n-k-1]$, and this completes the proof.

- 2) for $i > 2$, with $(n-1)|k(q+1)$, the Hermitian self-orthogonality follows from the same reasoning as point 1). The dimension of C_i can be taken up to $k+q$.

□

Remark 1. Given a Hermitian self-orthogonal code C_i with its generator matrix \mathcal{G}_i determined by (6) and $(g_t)_{1 \leq t \leq n}$ being its row, it is not difficult to see that if $\gamma_{t-1} := \langle g_t, g_t \rangle_h - \sum_{j=1}^n v_j^{q+1} \alpha_j^{(t-1)(q+1)}$, then

- 1) $\gamma_0 = \dots = \gamma_{k-1} = 0$;
- 2) $\gamma_t = \gamma_{t+q-1}$ for $1 \leq t \leq k-1$;
- 3) $\gamma_{k-1+i} = \gamma_{k-1+i+q-1} = \lambda_i^{q+1} = \lambda_{i+q-1}^{q+1}$ for $1 \leq i \leq q$;
- 4) for $1 \leq i \leq q$, there are at most $q-k+1$ non-zero λ_i .

For some special vaule k , we can easily count the number of i 's such that $\lambda_i = 0$. For instance for $k = q-1$, we have $\gamma_0 = \gamma_1 = \dots = \gamma_{q-2} = \gamma_q = \dots = \gamma_{2q-3} = 0$ and $\gamma_{q-1} = \gamma_{2q-2} = \lambda_1$.

In the sequel for $(\lambda_j)_{1 \leq j \leq i}$ defined by (6), we denote

$$I = \{j | \lambda_j \neq 0\}, \quad (9)$$

and write $\#I$ for the size of I .

Corollary 1. We have the following existence:

- 1) there exist AMDS Hermitian self-orthogonal codes with parameters $[q^2 + 1, k]_{q^2}$ for $k = q+1, \dots, 2q-2$;
- 2) there exists an AMDS Hermitian self-orthogonal code with parameters $[2(q+1)+2, k+2]_{q^2}$;
- 3) there exist Hermitian self-orthogonal codes with parameters $[2(q+1)+1+i-1, k_i, \geq n-k-i+1]_{q^2}$ for $2 \leq i \leq q$.

Proof. First note that if $(n-1)|(q^2-1)$, $(n-1)|k(q+1)$, and if there exists an MDS Hermitian self-orthogonal $[n, k]_{q^2}$ code with $k \leq \frac{n+q-1}{q+1}$, then the following holds:

- $\sum_{i=1}^n v_i^{q+1} \alpha_i^{k(q+1)} = \sum_{i=1}^n v_i^{q+1} \alpha_i^{k'(q+1)} \neq 0$ if $k|k'$;
- $\sum_{i=1}^n v_i^{q+1} \alpha_i^{k'(q+1)} = 0$ if $k \nmid k'$.

For the proof of 1), we take $n-1 = q^2-1$ and $k = q-1$. Then from Theorem 1 2), the codes C_i is Hermitian self-orthogonal. Moreover, with $n-1 = q^2-1$ and $k = q-1$, we get that $\lambda_i = 0$ for $i = 2, \dots, q-1$. Finally, by puncturing the zero coordinates in the generator matrix \mathcal{G}_i , we get the result as claimed.

For the proof of 2) and 3), we take $n-1 = 2(q+1)$. In this case, the largest dimension k equals to 2. Moreover, we have that $\lambda_i \neq 0$ for i odd, and $\lambda_i = 0$ for i even. \square

We illustrate the constructions of Hermitian self-orthogonal codes in the above corollary with the following two examples and give some parameters of Hermitian self-dual codes as follows: $[6, 3, 4]_{2^2}$, $[8, 4, 4]_{3^2}$, $[10, 5, 6]_{4^2}$, $[12, 6, 5]_{5^2}$, $[16, 8, 6]_{7^2}$, $[18, 9, 6]_{8^2}$, $[20, 10, 7]_{9^2}$, $[24, 12, 7]_{11^2}$. These parameters can be obtained by embedding an MDS Hermitian self-orthogonal $[q+2, 1]$ code.

Example 1. Take $q = 5$, $n = 25$. Then from [35], there exists a Hermitian self-dual code C with parameters $[25, 4, 22]_{25}$. Using the above embedding lemma, we obtain, from the code C , Hermitian self-orthogonal codes C_j with parameters $[25+j, 4+j, \geq 22-j]_{25}$ for $1 \leq j \leq 5$. Using Magma [3], we give the generator matrix of the Hermitian self-orthogonal code C_j for $j = 5$.

$$\begin{pmatrix} \beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta\beta & 00000 \\ 0\alpha w^{11} 4\delta w^{14} \theta w^{16} w^{17} 3w^{19} \gamma w^{21} \beta w^{23} 1w\lambda w^3 w^4 w^5 2w^7 w^8 w^9 & 00000 \\ 0\beta 1\lambda w^4 2w^8 \alpha 4w^{14} w^{16} 3\gamma \beta 1\lambda w^4 2w^8 \alpha 4w^{14} w^{16} 3\gamma & 00000 \\ 0\alpha \delta w^{16} w^{19} \beta w w^4 w^7 \alpha \delta w^{16} w^{19} \beta w w^4 w^7 \alpha \delta w^{16} w^{19} \beta w w^4 w^7 & 00000 \\ 0\beta \lambda 2\alpha w^{14} 3\beta \lambda 2\alpha w^{14} 3\beta \lambda 2\alpha w^{14} 3\beta \lambda 2\alpha w^{14} 3 & \lambda 0000 \\ 0\alpha \theta \gamma w 2w^{11} w^{16} w^{21} \lambda w^7 4w^{17} \beta w^3 w^8 \delta 3w^{23} w^4 w^9 w^{14} w^{19} 1w^5 & 00000 \\ 0\beta w^4 \alpha w^{16} \beta w^4 \alpha w^{16} \beta w^4 \alpha w^{16} \beta w^4 \alpha w^{16} \beta w^4 \alpha w^{16} & 00000 \\ 0\alpha w^{17} 1w^7 w^{14} w^{21} w^4 w^{11} 3w w^8 \theta \beta w^5 4w^{19} \lambda w^9 w^{16} w^{23} 2\delta \gamma w^3 & 00000 \\ 0\beta 2w^{14} \beta 2w^{14} \beta 2w^{14} \beta 2w^{14} \beta 2w^{14} \beta 2w^{14} \beta 2w^{14} & 0000\lambda \end{pmatrix},$$

where w is a primitive element of \mathbb{F}_{q^2} , $\lambda = w^2$, $\alpha = w^{10}$, $\delta = w^{13}$, $\theta = w^{15}$, $\gamma = w^{20}$ and $\beta = w^{22}$.

The code C_5 has parameters $[30, 9, 18]_{25}$. By puncturing the 27-th, 28-th and 29-th coordinates, we obtain a Hermitian self-orthogonal code with parameters $[27, 9, 18]_{25}$. From the generator matrix of the code, we can also obtain Hermitian self-orthogonal codes with parameters $[26, 5, 22]_{25}$, $[27, 6, 20]_{25}$, $[28, 7, 19]_{25}$, $[29, 8, 18]_{25}$, and by puncturing the zero coordinates, we get the codes with parameters $[26, 5, 22]_{25}$, $[26, 6, 20]_{25}$, $[26, 7, 19]_{25}$, $[26, 8, 18]_{25}$.

Example 2. Take $q = 7$, $n = 17$. Then from [35], there exists a Hermitian self-dual code C with parameters $[17, 2, 16]_{49}$. Using the above embedding lemma, we obtain, from the code C , Hermitian self-orthogonal codes C_j with parameters $[17 + j, 2 + j, \geq 16 - j]_{49}$ for $1 \leq j \leq 7$. Using Magma [3], we give the generator matrix of the Hermitian self-orthogonal code C_j for $j = 7$.

$$\begin{pmatrix} w^{45} \beta \beta \beta \beta \beta \beta \beta \beta \beta \beta \beta \beta \beta \beta \beta \beta \beta \beta & 0000000 \\ 0w^{22}w^{25}w^{28}w^{31}w^{34}w^{37}w^{43}\beta w w^4 w^7 w^{10}w^{13}w^{19} & 0000000 \\ 0\beta w^4 w^{10} w^{19} w^{22} w^{28} w^{34} w^{43} w^{10} w^{22} w^{28} w^{34} w^{43} & \lambda 000000 \\ 0w^{22}w^{31}w^{34}w^{37}w^{43}w^{46}w^{49}w^{52}w^{55}w^{58}w^{61}w^{64}w^{67}w^{70}w^{73}w^{76}w^{79} & 0000000 \\ 0\beta w^{10} w^{22} w^{34} w^{46} w^{58} w^{70} w^{82} w^{94} w^{106} w^{118} w^{130} w^{142} w^{154} w^{166} w^{178} & 00\lambda 0000 \\ 0w^{22}w^{37}w^{43}w^{49}w^{55}w^{61}w^{67}w^{73}w^{79}w^{85}w^{91}w^{97}w^{103}w^{109}w^{115}w^{121}w^{127} & 0000000 \\ 0\beta 2w^{34} w^4 w^{22} w^{10} w^{28} \beta 2w^{34} w^4 w^{22} w^{10} w^{28} & 0000\lambda 00 \\ 0w^{22}w^{43}w^{37}w^{10}w^{31}w^4 w^{25} \beta w^{19} w^{13} w^{34} w^7 w^{28} w & 0000000 \\ 0\beta w^{22} \beta w^{22} \beta w^{22} \beta w^{22} \beta w^{22} \beta w^{22} \beta w^{22} & 000000\lambda \end{pmatrix},$$

where w is a primitive element of \mathbb{F}_{q^2} , $\lambda = w^3$ and $\beta = w^{46}$.

The code C_7 has parameters $[24, 9, 10]_{49}$. By puncturing the 19-th, 21-th and 23-th coordinates, we obtain a Hermitian self-orthogonal code with parameters $[21, 9, 10]_{49}$. From the above matrix, we can also obtain Hermitian self-orthogonal codes with parameters $[18, 3, 16]_{49}$, $[19, 4, 14]_{49}$, $[20, 5, 14]_{49}$, $[21, 6, 12]_{49}$, $[22, 7, 12]_{49}$, $[23, 8, 10]_{49}$, and by puncturing the zero coordinates, we get the codes with parameters $[18, 3, 16]_{49}$, $[18, 4, 14]_{49}$, $[19, 5, 14]_{49}$, $[19, 6, 12]_{49}$, $[20, 7, 12]_{49}$, $[20, 8, 10]_{49}$, respectively.

Theorem 2. Let $N_j = j(q + 1)$ and $U_j = \{\alpha \in \mathbb{F}_{q^2} | \alpha^{N_j} = 1\}$. Assume that $a = 3$ or $a = 4$ and $a | (q - 1)$. Let $\beta_1 \in U_2 \setminus U_1$ and $\beta_2 \in U_a \setminus U_2$. Put $U_1 = \{u_1, \dots, u_{q+1}\}$ and $U = U_1 \cup \beta_1 U_1 \cup \beta_2 U_1 \cup \{0\}$. Put $n = 3(q + 1) + 1$, $k_i = 3 + i$ and $n_i = n + i$. Then

- 1) there exists an AMDS Hermitian self-orthogonal $[n + 2, 5]_{q^2}$ code;
- 2) there exists a Hermitian self-orthogonal $[n + i, 3 + i, \geq n - 2 - i]_{q^2}$ code.

Proof. Denote $U = \{\alpha_1, \dots, \alpha_{n-1}, \alpha_n\}$, where $\alpha_n = 0$. First note that if $a | (q - 1)$, then there exist $\beta_1 \in U_2 \setminus U_1$ and $\beta_2 \in U_a \setminus U_2$ such that $\beta_1^{2(q+1)} = 1$ and $\beta_2^{a(q+1)} = 1$. Set $D = \sum_{\alpha \in U} P_\alpha$ and $G = (k - 1)O$. Now under the assumption above, there exists, from [35, Construction 7], an MDS Hermitian self-orthogonal code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ with parameters $[n, k]_{q^2}$, where $v_i^{q+1} = \text{Res}_{P_{\alpha_i}}(\omega)$ for $1 \leq i \leq n$. In this case, the maximum value of the code dimension is $k = 3$. We now embed this code into another Hermitian self-orthogonal code. Consider the code C_i with its generator matrix \mathcal{G}_i written given by (6). With the same reasoning as in the proof of Theorem, we get that the code C_i has parameters $[n_i, k_i, \geq n - k - i + 1]$.

- 1) for $i = 2$, consider the following system of $(k + 1)(q + 1) + 1$ equations with n indeterminates $v_1^{q+1}, \dots, v_n^{q+1}$ defined by (7).

Then C_2 is Hermitian self-orthogonal if and only if (7) has a solution. We know that λ_1 exists by Lemma 5. The first $k + 1$ rows g_1, \dots, g_k, g_{k+1} of \mathcal{G}_2 generates a Hermitian self-orthogonal $[n + 2, k + 1]$ code. We now prove that $\langle g_{k+1}, g_{k+2} \rangle_h = 0$ and $\langle g_{k+2}, g_{k+2} \rangle_h = -\lambda_2^{q+1}$ for some $\lambda_2 \in \mathbb{F}_{q^2}$.

- a) Case $a = 4$:

Since the dimension $k = 3$, $\beta_1^{2(q+1)} = 1$ and $\beta_2^{4(q+1)} = 1$, the last row of M is

$$\begin{aligned} (\alpha_1^{4(q+1)}, \dots, \alpha_{n-1}^{4(q+1)}, \alpha_n^{4(q+1)}) &= (u_1^{4(q+1)}, \dots, u_{q+1}^{4(q+1)}, \\ &\quad (u_1 \beta_1)^{4(q+1)}, \dots, (u_{q+1} \beta_1)^{4(q+1)}, \\ &\quad (u_1 \beta_2)^{4(q+1)}, \dots, (u_{q+1} \beta_2)^{4(q+1)}, 0) \\ &= (\underbrace{1, \dots, 1}_{q+1}, \underbrace{1, \dots, 1}_{q+1}, \underbrace{1, \dots, 1}_{q+1}, 0), \end{aligned}$$

and so $v_1^{q+1} \alpha_1^{(k+1)(q+1)} + \dots + v_n^{q+1} \alpha_n^{(k+1)(q+1)} = -\lambda_2^{q+1}$ has a solution. Thus λ_2 is well determined (here $\lambda_2 = v_n$). Now, we check that $\langle g_{k+1}, g_{k+2} \rangle_h = 0$, that is,

$$\sum_{i=1}^n \alpha_i^{kq+k+1} v_i^{q+1} = 0. \quad (10)$$

Since $k(q + 1) + 1 = n$, $v_i^{q+1} = \text{Res}_{P_{\alpha_i}}(\omega)$ is a solution of (10).

b) Case $a = 3$:

Since the dimension $k = 3$, $\beta_1^{2(q+1)} = 1$ and $\beta_2^{3(q+1)} = 1$, the last row of M is

$$\begin{aligned} & (\alpha_1^{4(q+1)}, \dots, \alpha_{n-1}^{4(q+1)}, \alpha_n^{4(q+1)}) \\ &= (u_1^{4(q+1)}, \dots, u_{q+1}^{4(q+1)}, \\ & \quad (u_1\beta_1)^{4(q+1)}, \dots, (u_{q+1}\beta_1)^{4(q+1)}, \\ & \quad (u_1\beta_2)^{4(q+1)}, \dots, (u_{q+1}\beta_2)^{4(q+1)}, 0) \\ &= (\underbrace{1, \dots, 1}_{q+1}, \underbrace{1, \dots, 1}_{q+1}, \underbrace{\beta_2^{q+1}, \dots, \beta_2^{q+1}}_{q+1}, 0), \end{aligned}$$

and so $v_1^{q+1}\alpha_1^{(k+1)(q+1)} + \dots + v_n^{q+1}\alpha_n^{(k+1)(q+1)} = -\lambda_2^{q+1}$ has a solution. Thus, λ_2 is well determined. Now, we check that $\langle g_{k+1}, g_{k+2} \rangle_h = 0$, that is, (10) holds. Since $k(q+1) + 1 = n$, $v_i^{q+1} = \text{Res}_{P_{\alpha_i}}(\omega)$ is a solution of (10).

2) for $i > 2$, the Hermitian self-orthogonality follows from the same reasoning as point 1). □

Next, we provide a more general embedding construction of Hermitian self-orthogonal codes from an MDS Hermitian self-orthogonal code.

Theorem 3. *Let q be an odd prime power, $D = P_{\alpha_1} + \dots + P_{\alpha_n}$ and $G = (k-1)O$ be two divisors, ω be a Weil differential form such that $H = D - G + (\omega)$ and $\mathbf{v} = (v_1, \dots, v_n)$ with $v_i \in \mathbb{F}_{q^2}^*$. Assume that the following conditions hold*

- 1) $G \leq H$,
- 2) $\text{Res}_{P_{\alpha_j}}(\omega) = v_j^{q+1}$ for $1 \leq j \leq n$,
- 3) $2 \leq k \leq \lfloor \frac{n+q-1}{q+1} \rfloor$ and $k+q \leq n/2$.

Then the code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ can be embedded into a Hermitian self-orthogonal code with parameters $[n+i, k+i, \geq n-k-i+1]_{q^2}$ for $1 \leq i \leq q$.

Proof. Denote $G' = (k+i-1)O$ and $\mathbf{w} = (w_i, \dots, w_n)$, where $w_i = v_i^{\frac{q+1}{2}}$ for $1 \leq i \leq n$. Under the above assumption, the code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ is a Hermitian self-orthogonal code with parameters $[n, k, n-k+1]_{q^2}$, and the code $C_{\mathcal{L}}(D, G'; \mathbf{w})$ is a Euclidean self-orthogonal code with parameters $[n, k+i, n-k-i+1]_{q^2}$ for $1 \leq i \leq q$ since $k+q \leq \frac{n}{2}$ and $\text{Res}_{P_{\alpha_j}}$ are non-zero square elements in \mathbb{F}_{q^2} for $1 \leq j \leq n$. Let C_i be an $[n+i, k+i]$ code with its generator matrix \mathcal{G}_i defined as in (6). We now prove that C_i is Hermitian self-orthogonal. From the Euclidean self-orthogonality of the code $C_{\mathcal{L}}(D, G'; \mathbf{w})$, we deduce that for $1 \leq r \leq k-1+i, 1 \leq i \leq q$,

$$\sum_{j=1}^n w_j^2 \alpha_j^{r(k-1+i)} = 0. \quad (11)$$

By raising both sides of (11) to the power q , we get

$$\begin{aligned} \sum_{j=1}^n w_j^{2q} \alpha_j^{rq(k-1+i)} &= 0 \\ \sum_{j=1}^n v_j^{q+1} \alpha_j^{rq(k-1+i)} &= 0. \end{aligned} \quad (12)$$

The last equality of (12) holds due to fact that $(v_j^{q+1})_{1 \leq j \leq n}$ are elements in \mathbb{F}_q^* . Now from the last equality of (12), we deduce that $\langle g_j, g_l \rangle_h = 0$ for $j \neq l$, where $(g_j)_{1 \leq j \leq k+i}$ are the rows of \mathcal{G}_i . The rest is to check the existence of $(\lambda_j)_{1 \leq j \leq i}$ satisfying

$$\sum_{j=1}^n v_j^{q+1} \alpha_j^{(q+1)(k-1+r)} = -\lambda_r^{q+1} \text{ for } 1 \leq r \leq i. \quad (13)$$

Following from the same reasoning as in the proof of [35, Lemma 4], such $(\lambda_r)_{1 \leq r \leq i}$ do exist by taking $\text{Res}_{P_{\alpha_j}}(\omega) = v_j^{q+1}$ for $1 \leq j \leq n$. □

Corollary 2. *Let q be an odd prime power, $D = P_{\alpha_1} + \dots + P_{\alpha_n}$ and $G = (k-1)O$ be two divisors, ω be a Weil differential form such that $H = D - G + (\omega)$ and $\mathbf{v} = (v_1, \dots, v_n)$ with $v_i \in \mathbb{F}_{q^2}^*$. Assume that the following conditions hold*

- 1) $G \leq H$,
- 2) $\text{Res}_{P_{\alpha_j}}(\omega) = v_j^{q+1}$ for $1 \leq j \leq n$,
- 3) $2 \leq k \leq \lfloor \frac{n+q-1}{q+1} \rfloor$ and $k+q \leq n/2$.

Then the code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ can be embedded into a Hermitian self-orthogonal code with parameters $[n, 2k-1, \geq n-k-q+2]_{q^2}$.

TABLE I

SOME HERMITIAN SELF-ORTHOGONAL CODES WITH GOOD PARAMETERS OVER \mathbb{F}_{q^2} , $q = 2, 3, 4$, $n_i = n + i$, $k_i = k + i$, $(n - 1)|(q^2 - 1)$, $k = \lfloor \frac{n+q-1}{q+1} \rfloor$

i	$[n_i, k_i, d_i]_{q^2}$	Punctured code	I
1	$[5, 2, 4]_{2^2}$	—	{1}
2	$[6, 3, 4]_{2^2}$	—	{1, 2}
1	$[6, 2, 5]_{3^2}$	—	{1}
2	$[7, 3, 4]_{3^2}$	—	{1, 2}
3	$[8, 4, 4]_{3^2}$	—	{1, 2, 3}
1	$[10, 3, 8]_{3^2}$	—	{1}
2	$[11, 4, 6]_{3^2}$	$[10, 4, 6]_{3^2}$	{1}
3	$[12, 5, 6]_{3^2}$	$[11, 5, 6]_{3^2}$	{1, 3}
1	$[7, 2, 6]_{4^2}$	—	{1}
2	$[8, 3, 6]_{4^2}$	—	{1, 2}
3	$[9, 4, 6]_{4^2}$	—	{1, 2, 3}
4	$[10, 5, 6]_{4^2}$	—	{1, 2, 3, 4}
1	$[17, 4, 14]_{4^2}$	—	{1}
2	$[18, 5, 12]_{4^2}$	$[17, 5, 12]_{4^2}$	{1}
3	$[19, 6, 11]_{4^2}$	$[17, 6, 11]_{4^2}$	{1}
4	$[20, 7, 11]_{4^2}$	$[18, 7, 11]_{4^2}$	{1, 4}

Proof. From Theorem 3, there exists a Hermitian self-orthogonal code C_{q-1} with parameters $[n + q - 1, k + q - 1, \geq n - k - q + 2]_{q^2}$. From Remark 1, we get that $\gamma_0 = 0, \gamma_1 = \gamma_q = \lambda_{q-k+1} = 0, \dots, \gamma_{k-1} = \gamma_{q+k-2} = \lambda_{q-1} = 0$. By taking the subcode of C_{q-1} whose rows contain the first k rows of \mathcal{G}_i and the $(k - 1)$ consecutive rows with $\lambda_i = 0$, we obtain a Hermitian self-orthogonal code with parameters $[n + q - 1, 2k - 1, \geq n - k - q + 2]$. Now by puncturing the last $q - 1$ zero columns, we get a Hermitian self-orthogonal with parameters $[n, 2k - 1, \geq n - k - q + 2]$, and this completes the proof. \square

Example 3. We provide two good Hermitian self-orthogonal codes as follows.

- 1) Take $q = 5$, $k = 3$ and $n = 19$. Then we can construct a Hermitian self-orthogonal code with parameters $[19, 5, \geq 13]_{5^2}$.
- 2) Take $q = 5$, $k = 4$ and $n = 25$. Then we can construct a Hermitian self-orthogonal code with parameters $[25, 7, \geq 18]_{5^2}$.

More parameters of some good Hermitian self-orthogonal codes, calculated by Magma, are given in Tables I and II.

IV. HERMITIAN HULL CODES FROM ALGEBRAIC CURVES

In this section, we construct Hermitian hull codes from algebraic curves. In the first subsection, we begin our construction by considering the projective lines, the projective curves of genus zero, and hence from the Riemann-Roch's theorem, we obtain MDS Hermitian ℓ -dim hull codes. In the second subsection, we consider Hermitian ℓ -dim hull codes from algebraic curves of genus greater than or equal to one.

A. Hermitian hull codes from projective lines

In this subsection, we consider Hermitian hull codes over \mathbb{F}_{q^2} from the projective lines $\mathbb{P}^1 = \{(\alpha : 1) | \alpha \in \mathbb{F}_{q^2}\} \cup \{(1 : 0)\}$. There are $q^2 + 1$ distinct points in \mathbb{P}^1 , and thus are $q^2 + 1$ distinct places of $\mathbb{F}_{q^2}(\mathbb{P}^1)$.

Proposition 4. There exist MDS Hermitian ℓ -dim hull codes with parameters $[n, k]_{q^2}$ and $[n, n - k]_{q^2}$ for $0 \leq \ell \leq k$ if one of the following conditions holds:

- 1) $n = q^2 + 1$, $k = q$; $n = q^2 + 1$, $k \leq q$, q even, d odd; $n = q^2 + 1$, $k \leq q$, $q \equiv 1 \pmod{4}$, d even;
- 2) $n = q^2$, $k \leq q - 1$;
- 3) $n = (q^2 + 1)/2$, $q/2 < k \leq q - 1$, q odd;
- 4) $n = q^2 + 1$, $k \leq q$, $k \neq q - 1$; $n = r(q - 1) + 1$, $k \leq (q + r - 1)/2$, $q \equiv r - 1 \pmod{2r}$; $n = (q^2 + 2)/3$, $3|(q + 1)$, $k \leq (2q - 1)/3$;
- 5) $n = tq$, $1 \leq t \leq q$, $k \leq \lfloor \frac{tq+q-1}{q+1} \rfloor$; $n = t(q + 1) + 2$, $1 \leq t \leq q - 1$, $k \leq t + 1$, $(p, t, k) \neq (2, q - 1, q - 1)$;
- 6) $k \leq \lfloor \frac{n+q-1}{q+1} \rfloor$,
 - a) $(n - 1)|(q^2 - 1)$;
 - b) $n = 2N + 1$, $m = 2s$, $q_0 = p^s$ odd, $N = \frac{q-1}{p^r+1}$ even, $N < q_0 + 1$, $r | \frac{m}{2}$;
 - c) $n = (t + 1)N + i$, $1 \leq i \leq 2$, $N|(q^2 - 1)$, $n_2 = \frac{N}{\gcd(N, q+1)}$, $1 \leq t \leq \frac{q-1}{n_2} - 1$.

Proof. First note that if C is a Hermitian ℓ -dim hull $[n, k]$ code, then its Hermitian dual C^{\perp_h} is a Hermitian ℓ -dim hull $[n, n - k]$ code. Moreover, the dual of an MDS code is again an MDS code. It is enough to prove the existence of an MDS Hermitian self-orthogonal code with each of the above conditions in the theorem, and thus the result follows from Lemma 3. The existence of such codes can be obtained as follows:

TABLE II
SOME HERMITIAN SELF-ORTHOGONAL CODES WITH GOOD PARAMETERS OVER \mathbb{F}_{q^2} , $q = 5, 7$, $n_i = n + i$, $k_i = k + i$, $(n - 1)|(q^2 - 1)$ OR
 $n = 3(q + 1) + 1$, $k = \lfloor \frac{n+q-1}{q+1} \rfloor$

i	$[n_i, k_i, d_i]_{q^2}$	Punctured code	I
1	$[8, 2, 7]_{5^2}$	—	{1}
2	$[9, 3, 6]_{5^2}$	—	{1, 2}
3	$[10, 4, 5]_{5^2}$	—	{1, 2, 3}
4	$[11, 5, 5]_{5^2}$	—	{1, 2, 3, 4}
5	$[12, 6, 5]_{5^2}$	—	{1, 2, 3, 4, 5}
1	$[10, 3, 7]_{5^2}$	$[9, 3, 7]_{5^2}$	{}
2	$[11, 4, 6]_{5^2}$	$[9, 4, 6]_{5^2}$	{}
1	$[14, 3, 12]_{5^2}$	—	{1}
2	$[15, 4, 10]_{5^2}$	$[14, 4, 10]_{5^2}$	{1}
3	$[16, 5, 10]_{5^2}$	$[15, 5, 10]_{5^2}$	{1, 3}
4	$[17, 6, 8]_{5^2}$	$[15, 6, 8]_{5^2}$	{1, 3}
5	$[18, 7, 8]_{5^2}$	$[16, 7, 8]_{5^2}$	{1, 3, 5}
1	$[20, 4, 17]_{5^2}$	—	{1}
2	$[21, 5, 16]_{5^2}$	—	{1, 2}
3	$[22, 6, 14]_{5^2}$	$[21, 6, 14]_{5^2}$	{1, 2}
4	$[23, 7, 13]_{5^2}$	$[21, 7, 13]_{5^2}$	{1, 2}
5	$[24, 8, 13]_{5^2}$	$[22, 8, 13]_{5^2}$	{1, 2, 5}
1	$[26, 5, 22]_{5^2}$	$[26, 5, 22]_{5^2}$	{1}
2	$[27, 6, 20]_{5^2}$	$[26, 6, 20]_{5^2}$	{1}
3	$[28, 7, 19]_{5^2}$	$[26, 7, 19]_{5^2}$	{1}
4	$[29, 8, 18]_{5^2}$	$[26, 8, 18]_{5^2}$	{1}
5	$[30, 9, 18]_{5^2}$	$[27, 9, 18]_{5^2}$	{1, 5}
1	$[14, 3, 11]_{7^2}$	$[13, 3, 11]_{7^2}$	{}
2	$[15, 4, 11]_{7^2}$	$[14, 4, 11]_{7^2}$	{2}
3	$[16, 5, 9]_{7^2}$	$[14, 5, 9]_{7^2}$	{2}
1	$[18, 3, 16]_{7^2}$	$[18, 3, 16]_{7^2}$	{1}
2	$[19, 4, 14]_{7^2}$	$[18, 4, 14]_{7^2}$	{1}
3	$[20, 5, 14]_{7^2}$	$[19, 5, 14]_{7^2}$	{1, 3}
4	$[21, 6, 12]_{7^2}$	$[19, 6, 12]_{7^2}$	{1, 3}
5	$[22, 7, 12]_{7^2}$	$[20, 7, 12]_{7^2}$	{1, 3, 5}
6	$[23, 8, 10]_{7^2}$	$[20, 8, 10]_{7^2}$	{1, 3, 5}
7	$[24, 9, 10]_{7^2}$	$[21, 9, 10]_{7^2}$	{1, 3, 5, 7}
1	$[26, 4, 23]_{7^2}$	—	{1}
2	$[27, 5, 21]_{7^2}$	$[26, 5, 21]_{7^2}$	{1}
3	$[28, 6, 20]_{7^2}$	$[26, 6, 20]_{7^2}$	{1}
4	$[29, 7, 20]_{7^2}$	$[27, 7, 20]_{7^2}$	{1, 4}
5	$[30, 8, 18]_{7^2}$	$[27, 8, 18]_{7^2}$	{1, 4}
6	$[31, 9, 17]_{7^2}$	$[27, 9, 17]_{7^2}$	{1, 4}
7	$[32, 10, \geq 16]_{7^2}$	$[28, 10, \geq 16]_{7^2}$	{1, 4, 7}

- 1) $n = q^2 + 1$, $k = q$ from [27];
- 2) $n = q^2 + 1$, $k \leq q$, q even, d odd from [18];
- 3) $n = q^2 + 1$, $k \leq q$, $q \equiv 1 \pmod{4}$, d even from [23];
- 4) $n = q^2$, $k \leq q - 1$ from [17] and [27], [22];
- 5) $n = (q^2 + 1)/2$, $q/2 < k \leq q - 1$, q odd from [23];
- 6) $n = q^2 + 1$, $k \leq q$, $k \neq q - 1$; $n = r(q - 1) + 1$, $k \leq (q + r - 1)/2$, $q \equiv r - 1 \pmod{2r}$; $n = (q^2 + 2)/3$, $3|(q + 1)$, $k \leq (2q - 1)/3$ from [21];
- 7) $n = tq$, $1 \leq t \leq q$, $k \leq \lfloor \frac{tq+q-1}{q+1} \rfloor$; $n = t(q + 1) + 2$, $1 \leq t \leq q - 1$, $k \leq t + 1$, $(p, t, k) \neq (2, q - 1, q - 1)$ from [11];
- 8) $(n - 1)|(q^2 - 1)$; $n = 2N + 1$, $m = 2s$, $q_0 = p^s$ odd, $N = \frac{q-1}{p^r+1}$ even, $N < q_0 + 1$, $r|\frac{m}{2}$; $n = (t + 1)N + i$, $1 \leq i \leq 2$, $N|(q^2 - 1)$, $n_2 = \frac{N}{\gcd(N, q+1)}$, $1 \leq t \leq \frac{q-1}{n_2} - 1$ from [35].

□

By applying some propagation rules to an MDS Hermitian self-orthogonal code, we obtain some MDS Hermitian self-orthogonal codes with smaller lengths and smaller dimensions. Since some families in Proposition 4 are contained in other families, we only state the larger families of Hermitian ℓ -dim hull codes in the following theorem.

Theorem 4. *There exist MDS Hermitian ℓ -dim hull codes with parameters $[n - s, k - s, n - k + 1 - s]_{q^2}$ and $[n - s, n - k - s, k + 1 - s]_{q^2}$ for $1 \leq s \leq k - 1$ and $0 \leq \ell \leq k - s$ if one of the following conditions holds*

- 1) $n = q^2 + 1$, $k \leq q$, $k \neq q - 1$; $n = r(q - 1) + 1$, $k \leq (q + r - 1)/2$, $q \equiv r - 1 \pmod{2r}$; $n = (q^2 + 2)/3$, $3|(q + 1)$, $k \leq (2q - 1)/3$;
- 2) $n = tq$, $1 \leq t \leq q$, $k \leq \lfloor \frac{tq+q-1}{q+1} \rfloor$; $n = t(q + 1) + 2$, $1 \leq t \leq q - 1$, $k \leq t + 1$, $(p, t, k) \neq (2, q - 1, q - 1)$;

- 3) $(n-1)|(q^2-1)$, $k \leq \lfloor \frac{n+q-1}{q+1} \rfloor$; $n = (t+1)N + i$, $1 \leq i \leq 2$, $N|(q^2-1)$, $n_2 = \frac{N}{\gcd(N, q+1)}$, $1 \leq t \leq \frac{q-1}{n_2} - 1$, $k \leq \lfloor \frac{n+q-1}{q+1} \rfloor$.

Proof. Assume that there exists a Hermitian self-orthogonal code C with parameters $[n, k, d]_{q^2}$. Then up to equivalence, we can write the generator matrix G of C as $G = (I_k | A)$. Deleting the last s consecutive rows and the s consecutive zero-columns of G , we obtain a matrix G' which again generates a Hermitian self-orthogonal code C' . Moreover, if C is MDS, then so is C' . This completes the proof. \square

It should be noted for n sufficiently large enough, say $n > k + q$ (see [10]), the dual of an AMDS code is again an AMDS code.

Theorem 5. Let $q = p^m$, $(n-1)|(q^2-1)$ and $1 \leq k \leq \lfloor \frac{n+q-1}{q+1} \rfloor$.

- 1) If $(n-1)|(k+1)(q+1)$, then there exists a Hermitian ℓ -dim hull code parameters $[n+1, k+2, n-k-1]_{q^2}$ for $0 \leq \ell \leq k+2$
- 2) If $(n-1)|k(q+1)$, then there exists a Hermitian ℓ -dim hull code with parameters $[n+i-1, k+i, n-k-i+1]_{q^2}$ for $0 \leq \ell \leq k+i$.

Proof. The proof follows from Theorem 1. \square

Corollary 3. 1) There exist AMDS Hermitian ℓ -dim hull codes with parameters $[q^2+1, k]_{q^2}$ and $[q^2+1, q^2+1-k]_{q^2}$ for $0 \leq \ell \leq k$, $k = q+1, \dots, 2q-2$;

- 2) There exist AMDS Hermitian ℓ -dim hull codes with parameters $[2(q+1)+2, k+2]_{q^2}$ and $[2(q+1)+2, 2(q+1)-k]_{q^2}$ for $0 \leq \ell \leq k+2$;
- 3) There exist Hermitian ℓ -dim hull codes with parameters $[2(q+1)+i, k+i, \geq n-k-i+1]_{q^2}$ for $0 \leq \ell \leq k+i$ and $2 \leq i \leq q$.

Proof. The proof follows from Corollary 1. \square

Theorem 6. Let $q = p^m$, $(n-1)|(q^2-1)$ and $k = \lfloor \frac{n+q-1}{q+1} \rfloor$. Then

- 1) if $(n-1)|k(q+1)$, then there exist MDS Hermitian $(k+i-\#I)$ -dim hull codes with parameters $[n, k+i]_{q^2}$ and $[n, n-k-i]_{q^2}$ for any $1 \leq i \leq q$;
- 2) if $(n-1)|(k+1)(q+1)$, then there exist MDS Hermitian $(k+1)$ -dim hull codes with parameters $[n, k+2]_{q^2}$ and $[n, n-k-2]_{q^2}$.

Proof. We prove the theorem using similar technique to that in the proof of Theorem 1.

- 1) If $(n-1)|k(q+1)$, then from Theorem 1, there exists a Hermitian self-orthogonal $[n+i, k+i]_{q^2}$ code. Consider the code C_i with its generator matrix \mathcal{G}_i given by (6). By puncturing the last i coordinates of \mathcal{G}_i , one obtains a generator matrix \mathcal{G}_{i_0} satisfying $\text{Rank}(\mathcal{G}_{i_0}\mathcal{G}_{i_0}^\dagger) = k+i-\#I$. It is not difficult to check that \mathcal{G}_{i_0} generates an MDS Hermitian $(k+i-\#I)$ -dim hull code.
- 2) If $(n-1)|(k+1)(q+1)$, then there exist $\lambda_1 \neq 0$ and $\lambda_2 = 0$, and the result follows from point 1). \square

Example 4. One can easily obtain an MDS Hermitian 7-dim hull $[25, 9, 17]_{5^2}$ code and an MDS Hermitian 6-dim hull $[25, 8, 18]_{5^2}$ code from the generator matrix of Example 1, as well as an MDS Hermitian 5-dim hull $[17, 9, 9]_{5^2}$ code and an MDS Hermitian 5-dim hull $[17, 8, 10]_{7^2}$ code from the generator matrix of Example 2. These four codes have their dimension $k > \frac{n+q-1}{q+1}$, and thus their parameters are never reachable by [12].

Corollary 4. Let $q = p^m$ and $k = \lfloor \frac{n+q-1}{q+1} \rfloor$. Then

- 1) for $n = q^2$, there exist MDS Hermitian $(k+i-1)$ -dim hull codes with parameters $[q^2, k+i]_{q^2}$ and $[q^2, q^2-k-i]_{q^2}$ for $1 \leq i \leq q$;
- 2) $n = 2(q+1)+1$, there exist MDS Hermitian $(k+i-\lfloor \frac{i-1}{2} \rfloor - 1)$ -dim hull codes with parameters $[2(q+1)+1, k+i]_{q^2}$ and $[2(q+1)+1, 2(q+1)+1-k-i]_{q^2}$ for $1 \leq i \leq q$.

Proof. 1) Take $n-1 = q^2-1$. Then we get that $\lambda_1 \neq 0$ and $\lambda_2 = \dots = \lambda_{q-1} = 0$. The result follows from Theorem 6 1).

- 2) Take $n-1 = 2(q+1)$. Then we get that $\lambda_1 = \lambda_3 = \dots = \lambda_{q-2} \neq 0$ and $\lambda_2 = \dots = \lambda_{q-1} = 0$. The result follows from Theorem 6 1). \square

Theorem 7. Let q be an odd prime power, $N|(q^2-1)$, $n_2 = \frac{N}{\gcd(N, q+1)}$, $1 \leq t \leq \frac{q-1}{n_2} - 1$. Put $n = (t+1)N+1$, $k = \lfloor \frac{n+q-1}{q+1} \rfloor$, $k_i = k+i$ and $n_i = n+i$. Then

- 1) there exist AMDS Hermitian ℓ -dim hull codes with parameters $[n+2, k+2]_{q^2}$ and $[n+2, n-k]_{q^2}$ for $0 \leq \ell \leq k+2$;

- 2) there exist Hermitian ℓ -dim hull codes with parameters $[n+i, k+i, \geq n-k-i+1]_{q^2}$ code for $0 \leq \ell \leq k+i$ for $1 \leq i \leq q$;
- 3) there exist Hermitian ℓ -dim hull codes with parameters $[n, 2k-1, \geq n-k-q+2]_{q^2}$ for $0 \leq \ell \leq 2k-1$;
- 4) there exist MDS Hermitian $(k+i - \#I)$ -dim hull codes with parameters $[n, k+i]_{q^2}$ and $[n, n-k-i]_{q^2}$ for $1 \leq i \leq q$.

Proof. It was proved in [35, Construction 7] that there exists an MDS Hermitian self-orthogonal code with parameters $[n, k]_{q^2}$. Now, the result follows from the same reasoning as in the proof of Theorem 6. \square

In the following, we give another family of MDS Hermitian ℓ -dim hull codes.

For sake of stating our result, let us fix the following notation and setting. Let $q = p^m$ be a prime power and $n|(q^2-1)$. Write

$$n_1 = \gcd(n, q-1) \text{ and } n_2 = \frac{n}{n_1}. \quad (14)$$

Then the first equality of (14) implies that n_2 and $\frac{q-1}{n_1}$ are coprime, and thus we get that $n_2|(q+1)\frac{q-1}{n_1}$, and so $n_2|(q+1)$. Let U_n and V_n be two subgroups of $\mathbb{F}_{q^2}^*$ generated by $w^{\frac{q^2-1}{n}}$ and $w^{\frac{q-1}{n_1}}$, respectively, where w is a primitive element of \mathbb{F}_{q^2} . It can be easily seen that $|U_n| = n$ and $|V_n| = (q+1)n_1$. Now $\frac{q^2-1}{n} = \frac{q-1}{n_1} \cdot \frac{q+1}{n_2}$ implies that $\frac{q-1}{n_1} | \frac{q^2-1}{n}$, and we deduce that U_n is a subgroup of V_n . Write $U_n = \{u_1, \dots, u_n\}$.

Proposition 5. *Let $q = p^m$ be a prime power, $n|(q^2-1)$ and $n_2 = \frac{n}{\gcd(n, q-1)}$. Put $N = (t+1)n+1$ for $1 \leq t \leq \frac{q+1}{n_2}-1$. Assume that there exists t such that for all $\alpha_1, \dots, \alpha_t \in V_n \setminus U_n$, we have $\alpha_i^n = \beta_i^{q+1}$ for some $\beta_i \in \mathbb{F}_{q^2}$. Then for $1 \leq k \leq \lfloor \frac{N+q-1}{q+1} \rfloor$, there exist MDS Hermitian ℓ -dim hull codes with parameters $[N, k]_{q^2}$, $[N+1, k+1]_{q^2}$, $[N, N-k]_{q^2}$ and $[N+1, N-k]_{q^2}$ for $0 \leq \ell \leq k$.*

Proof. Let $\alpha_1 U_n, \dots, \alpha_{\frac{q+1}{n_2}-1} U_n$ be all the distinct cosets of V_n different from U_n .

For $1 \leq t \leq \frac{q+1}{n_2}-1$, put $U = U_n \cup_{j=1}^t \alpha_j U_n \cup \{0\}$, say $U = \{a_1, \dots, a_{(t+1)n+1}\}$, and write

$$h(x) = \prod_{\alpha \in U} (x - \alpha).$$

Then the derivative of $h(x)$ is given by

$$h'(x) = ((n+1)x^n - 1) \prod_{i=1}^t (x^n - \alpha_i^n) + nx^n(x^n - 1) \left(\sum_{i=1}^t \prod_{j=1, j \neq i}^t (x^n - \alpha_j^n) \right).$$

For $1 \leq j \leq t, 1 \leq s \leq n$, we have

$$h'(\alpha_j u_s) = n\alpha_j^n(1 - \alpha_j^n) \prod_{i=1, i \neq j}^t (\alpha_j^n - \alpha_i^n). \quad (15)$$

Now, we check the values of the derivative h' given by (15).

Since α_j is in V_n , we can write $\alpha_j = w^{e_j \frac{q-1}{n_1}}$ for some positive integer e_j . Thus, $\alpha_j^n = w^{e_j n \frac{q-1}{n_1}} = w^{e_j n_2(q-1)}$, and $\alpha_j^{n(q+1)} = 1$, that is, $\alpha_j^{nq} = \alpha_j^{-n}$. Now, the latter equality implies that $(\alpha_i^n - \alpha_j^n)^q = \frac{\alpha_j^n - \alpha_i^n}{\alpha_i^n \alpha_j^n}$, and thus

$$(\alpha_i^n - \alpha_j^n)^{q-1} = \frac{-1}{(\alpha_i \alpha_j)^n} = \frac{-1}{(\beta_i \beta_j)^{q+1}} \quad (16)$$

By raising the last equality of (16) to the power $q+1$, we get $(\beta_i \beta_j)^{q+1} = 1$, and this implies that

$$\alpha_i^n - \alpha_j^n = w^{\frac{q+1}{2}} \text{ which is independent of } i, j. \quad (17)$$

Set $G = (k-1)O$, $D = (h)_0$, $\omega = \frac{dx}{h(x)}$ and $H = ((t+1)n - k - 1)O$. Thus by Lemma 4, the code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$, where $\mathbf{v} = (v_1, \dots, v_{(t+1)n+1})$ and $v_i^{q+1} = 1/\beta_i$, is Hermitian self-orthogonal over \mathbb{F}_{q^2} . Hence, for $1 \leq k \leq \frac{(t+1)n+q}{q-1}$, there exists an MDS Hermitian self-orthogonal $[(t+1)n+1, k]_{q^2}$ code. By applying the embedding Lemma 5, we get an MDS Hermitian self-orthogonal $[(t+1)n+2, k+1]_{q^2}$ code. Hence, the result follows from Lemma 3. \square

Corollary 5. *Let $q = p^m$ be a prime power and $N = 2(q+1) + 1$. Then for $1 \leq k \leq \lfloor \frac{N+q-1}{q+1} \rfloor$, there exist MDS Hermitian ℓ -dim hull codes with parameters $[N, k]_{q^2}$, $[N+1, k+1]_{q^2}$, $[N, N-k]_{q^2}$ and $[N+1, N-k]_{q^2}$ for $0 \leq \ell \leq k$.*

TABLE III
NUMBERS OF RATIONAL POINTS OF ELLIPTIC CURVES OVER \mathbb{F}_q WITH $q = 2^m$ BEING A SQUARE

Elliptic curve $\mathcal{E}_{1,b,c}$	m	$N_{1,b,c}$
$y^2 + y = x^3$	$m \equiv 0 \pmod{4}$	$q + 1 - 2\sqrt{q}$
	$m \equiv 2 \pmod{4}$	$q + 1 + 2\sqrt{q}$
$y^2 + y = x^3 + bx (Tr_1^m(b) = 1)$	m even	$q + 1$
$y^2 + y = x^3 + c (Tr_1^m(c) = 1)$	$m \equiv 0 \pmod{4}$	$q + 1 + 2\sqrt{q}$
	$m \equiv 2 \pmod{4}$	$q + 1 - 2\sqrt{q}$

Proof. Take $n = q + 1$. Then $n_1 = \gcd(n, q - 1) = 2$ and thus $n_2 = \frac{q+1}{2}$. With similar argument as in the proof of Theorem 5, we get

$$(1 - \alpha_1^{q+1}) = \frac{w^{\frac{q+1}{2}}}{w^{e_1 n_2}} = \frac{w^{\frac{q+1}{2}}}{w^{e_1 \frac{q+1}{2}}}$$

If e_1 was even, then $\alpha_1^n = w^{(q-1)n_2 e_1} = w^{(q^2-1)\frac{e_1}{2}} = 1$, and thus $\alpha_1 \in U_n$ which is a contradiction. Hence, e_1 is odd, and the proof is completed. \square

B. Hermitian ℓ -dim hull codes from elliptic curves, hyper-elliptic curves and Hermitian curves

In the previous subsection, we have already seen that Hermitian self-orthogonal codes, constructed from the projective lines, have their lengths upper bounded by $q^2 + 1$ for MDS case ($q > 2$), and by $q^2 + q + 1$ for non MDS case, and thus the code lengths of the corresponding Hermitian ℓ -dim hull codes can not go beyond $q^2 + q + 1$. In this subsection, we consider Hermitian ℓ -dim hull codes from some algebraic curves over \mathbb{F}_q which have many \mathbb{F}_q -rational points. These curves could produce long codes with large minimum distance. By the famous Hasse-Weil bound, the bound of the number of \mathbb{F}_q -rational points of a smooth projective curve \mathcal{X} defined over \mathbb{F}_q is given as follows:

$$|\#\mathcal{X}(\mathbb{F}_q) - (q + 1)| \leq 2g\sqrt{q},$$

where g is the genus of \mathcal{X} . When the bound is attained, the curves are called maximal. We are considering some well-known maximal curves, for example elliptic curves, hyper-elliptic curves and Hermitian curves and employ them to construct Hermitian hull codes from AG codes.

First we consider elliptic curves in even characteristic. Let $q = 2^m$ and an elliptic curve be defined by the equation

$$\mathcal{E}_{a,b,c} : y^2 + ay = x^3 + bx + c, \quad (18)$$

where $a, b, c \in \mathbb{F}_q$. Let denote the number of rational points on $\mathcal{E}_{a,b,c}$ by $N_{a,b,c}$. Let S be the set of x -components of the affine points of $\mathcal{E}_{a,b,c}$ over \mathbb{F}_q , that is,

$$S_{a,b,c} := \{\alpha \in \mathbb{F}_q \mid \exists \beta \in \mathbb{F}_q \text{ such that } \beta^2 + a\beta = \alpha^2 + b\alpha + c\}. \quad (19)$$

It should be noted that any $\alpha \in S_{1,b,c}$ gives rise to two points on $\mathcal{E}_{a,b,c}$ with x -component α , and we denote them by $P_\alpha^{(1)}$ and $P_\alpha^{(2)}$. The numbers $N_{1,b,c}$ of \mathbb{F}_q -rational points of elliptic curves $\mathcal{E}_{1,b,c}$ over \mathbb{F}_q with q being a square are given in Table III.

We can now state an existence of AMDS Hermitian ℓ -dim hull codes from the elliptic curves as follows.

Proposition 6. *Let $q = 2^s$, $m = 2s$ and $2k + 2 \leq n \leq \lfloor q + \sqrt{2q} - 5 \rfloor$. Then there exist AMDS Hermitian ℓ -dim hull codes with parameters $[n, k]_{q^2}$ and $[n, n - k]_{q^2}$ for $0 \leq \ell \leq k$.*

Proof. The existence of a Hermitian self-orthogonal $[n, k]_{q^2}$ code follows from [20]. Hence, the result follows from Lemma 3. \square

Next, we consider the affine hyper-elliptic curve over \mathbb{F}_{q^2} with $q = 2^m$ which is defined by

$$\mathcal{C} : y^2 + y = x^{q+1}. \quad (20)$$

Note that for any $\alpha \in \mathbb{F}_{q^2}$, there are two rational points on \mathcal{C} with x -component α , and we denote them by $P_\alpha^{(1)}, P_\alpha^{(2)}$. The set $\mathcal{C}(\mathbb{F}_{q^2})$ of all rational points of \mathcal{C} is equal to $\{P_\alpha^{(1)} \mid \alpha \in \mathbb{F}_{q^2}\} \cup \{P_\alpha^{(2)} \mid \alpha \in \mathbb{F}_{q^2}\} \cup \{O\}$. This curve has genus $g = \frac{q}{2}$. It can be easily checked that the curve \mathcal{C} has $1 + 2q^2 = 1 + q^2 + 2\frac{q}{2}q$ points, and thus it is a maximal curve.

The constructions of Hermitian ℓ -dim hull codes from the hyper-elliptic curves are given as follows.

Proposition 7. *Let $q = 2^m$, $m \geq 2$ and $q \leq k \leq \lfloor \frac{2q^2 + 2q - 1}{q+1} \rfloor$. Put $G = (k - 1)O$ and $D = \sum_{\alpha \in \mathbb{F}_{q^2}} (P_\alpha^{(1)} + P_\alpha^{(2)})$. Then for $0 \leq \ell \leq k - q/2$, the code $C_{\mathcal{L}_q}(D, G)$ gives rise to a Hermitian ℓ -dim hull $[2q^2, k - q/2, \geq 2q^2 - k + 1]_{q^2}$ code whose Hermitian dual has parameters $[2q^2, 2q^2 + q/2 - k, \geq k + 1 - q]_{q^2}$.*

Proof. From [35], the code $C_{\mathcal{L}_q}(D, G)$ is Hermitian self-orthogonal with parameters $[2q^2, k - q/2, \geq 2q^2 - k - \frac{q}{2} + 1]_{q^2}$. The result follows from Lemma 3. \square

Proposition 8. *Let $q = 2^m, m \geq 2, (n-1)|(q-1)$ and $q \leq k \leq \lfloor \frac{2n+2q-1}{q+1} \rfloor$. Put $U = \{\alpha \in \mathbb{F}_{q^2} | \alpha^n = \alpha\}$, $G = (k-1)O$ and $D = \sum_{\alpha \in U} (P_\alpha^{(1)} + P_\alpha^{(2)})$. Then for $0 \leq \ell \leq k - q/2$, the code $C_{\mathcal{L}_q}(D, G)$ gives rise to a Hermitian ℓ -dim hull $[2n, k - q/2, \geq 2n - k + 1]_{q^2}$ code whose Hermitian dual has parameters $[2n, 2n + q/2 - k, \geq k + 1 - q]_{q^2}$.*

Proof. From [35], the code $C_{\mathcal{L}_q}(D, G)$ is Hermitian self-orthogonal with parameters $[2n, k - q/2, \geq 2n - k + 1]_{q^2}$. Hence, the result follows from Lemma 3. \square

In the following, we consider the affine Hermitian curve over \mathbb{F}_{q^2} with $q = p^m$ which is defined by

$$\mathcal{H} : y^q + y = x^{q+1}. \quad (21)$$

For any $\alpha \in \mathbb{F}_{q^2}$, there are q rational points on \mathcal{H} with x -component α , and we denote them by $P_\alpha^{(1)}, P_\alpha^{(2)}, \dots, P_\alpha^{(q)}$. The set $\mathcal{H}(\mathbb{F}_{q^2})$ of all rational points of \mathcal{H} is equal to $\{P_\alpha^{(1)} | \alpha \in \mathbb{F}_{q^2}\} \cup \{P_\alpha^{(2)} | \alpha \in \mathbb{F}_{q^2}\} \cup \dots \cup \{P_\alpha^{(q)} | \alpha \in \mathbb{F}_{q^2}\} \cup \{O\}$. The curve \mathcal{H} has genus $g = \frac{q(q-1)}{2}$ and has $1 + q^3 = 1 + q^2 + 2\frac{q(q-1)}{2}q$ points. Hence, it is a maximal curve.

Proposition 9. *Let $q = p^m \geq 4$ and $q(q-1) \leq k \leq \lfloor \frac{q^3+q^2-1}{q+1} \rfloor$. Put $\mathbf{v} = (1/\sqrt[q+1]{-1}, \dots, 1/\sqrt[q+1]{-1})$, $G = (k-1)O$ and $D = \sum_{\alpha \in \mathbb{F}_{q^2}} (P_\alpha^{(1)} + P_\alpha^{(2)})$. Then for $0 \leq \ell \leq k - \frac{q(q-1)}{2}$, the code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ gives rise to a Hermitian ℓ -dim hull $[q^3, k - \frac{q(q-1)}{2}, \geq q^3 - k + 1]_{q^2}$ code whose Hermitian dual has parameters $[q^3, q^3 + q(q-1)/2 - k, \geq k + 1 - q(q-1)]_{q^2}$.*

Proof. From [35], the code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ is Hermitian self-orthogonal with parameters $[q^3, k - \frac{q(q-1)}{2}, \geq q^3 - k + 1]_{q^2}$. Hence, the result follows from Lemma 3. \square

Proposition 10. *Let $q = p^m \geq 4$ and $q(q-1) \leq k \leq \lfloor \frac{Nq+q^2-1}{q+1} \rfloor$. Then for $0 \leq \ell \leq k - \frac{q(q-1)}{2}$, there exists a Hermitian ℓ -dim hull $[Nq, k - \frac{q(q-1)}{2}, \geq Nq - k + 1]_{q^2}$ code whose Hermitian dual has parameters $[Nq, Nq + q(q-1)/2 - k, \geq k + 1 - q(q-1)]_{q^2}$ if one of the following condition holds*

- i) $n = q^2 + 1, k \leq q, k \neq q - 1; n = r(q - 1) + 1, k \leq (q + r - 1)/2, q \equiv r - 1 \pmod{2r}; n = (q^2 + 2)/3, 3|(q + 1), k \leq (2q - 1)/3;$
- ii) $n = tq, 1 \leq t \leq q, k \leq \lfloor \frac{tq+q-1}{q+1} \rfloor; n = t(q + 1) + 2, 1 \leq t \leq q - 1, k \leq t + 1, (p, t, k) \neq (2, q - 1, q - 1);$
- iii) $(n - 1)|(q^2 - 1), k \leq \lfloor \frac{n+q-1}{q+1} \rfloor; n = (t + 1)N + i, 1 \leq i \leq 2, N|(q^2 - 1), n_2 = \frac{N}{\gcd(N, q+1)}, 1 \leq t \leq \frac{q-1}{n_2} - 1, k \leq \lfloor \frac{n+q-1}{q+1} \rfloor.$

Proof. The existence of a Hermitian self-orthogonal $[Nq, k - \frac{q(q-1)}{2}, \geq Nq - k + 1]_{q^2}$ code follows from [35]. Hence, the result follows from Lemma 3. \square

Theorem 8. *Let $q = p^m$ be an odd prime power, $1 < t < q + 1$ and $t|(q + 1)$. Put $g = \frac{(q-1)(q+1-t)}{2t}$ and $s = q \left(\frac{\lfloor q^2 - 2 \rfloor}{t} + 2 \right)$. Then for $2g \leq k \leq \lfloor \frac{s+q-1}{q+1} \rfloor$ and $0 \leq \ell \leq k - g$, there exist ℓ -dim hull codes with parameters $[s, k - g, \geq s - k + 1]_{q^2}$ and $[s, s - k + g, \geq k - 2g + 1]_{q^2}$.*

Proof. Consider an algebraic curve defined by

$$\mathcal{X} : y^q + y = x^{\frac{q+1}{t}}.$$

The curve has genus $g = \frac{(q-1)(q+1-t)}{2t}$. Put

$$U = \{\alpha \in \mathbb{F}_{q^2} | \exists \beta \in \mathbb{F}_q \text{ such that } \beta^q + \beta = \alpha^{\frac{q+1}{t}}\}.$$

The set U is the set of x -component solutions to the Hermitian curve (21) whose elements are t -th power elements in \mathbb{F}_{q^2} . There are such $\left(\frac{\lfloor q^2 - 2 \rfloor}{t} + 2 \right)$ elements in \mathbb{F}_{q^2} which give rise to $s = q \left(\frac{\lfloor q^2 - 2 \rfloor}{t} + 2 \right)$ rational places. Write

$$h(x) = \prod_{\alpha \in U} (x - \alpha) \text{ and } \omega = \frac{dx}{h(x)}.$$

Then $h(x) = x^n - x$, where $n = \frac{\lfloor q^2 - 2 \rfloor}{t} + 2$, and thus $h'(x) = nx^{n-1} - 1$. Obviously, $h'(0) = w^{\frac{q^2-1}{2}}$, where w is a primitive element of \mathbb{F}_{q^2} . Since q is a square, we have that for any $\alpha \in U \setminus \{0\}$, $h'(\alpha) = n - 1 = \beta^{q+1}$ for some $\beta \in \mathbb{F}_{q^2}$. Put $D = \sum_{\alpha \in U} (P_\alpha^{(1)} + \dots + P_\alpha^{(q)}) = P_1 + \dots + P_s, G = (k-1)O$. By Lemma 4, the constructed code $C_{\mathcal{L}_q}(D, G; \mathbf{v})$ is Hermitian self-orthogonal, where $\mathbf{v} = (\underbrace{1/w^{\frac{q-1}{2}}, \dots, 1/w^{\frac{q-1}{2}}}_q, \underbrace{1/\beta, \dots, 1/\beta}_q, \dots, \underbrace{1/\beta, \dots, 1/\beta}_q)$. Finally, the result follows from Lemma

3. \square

Corollary 6. Let N_1, K_1 and d_1 be the length, dimension and minimum distance of the Hermitian ℓ -dim hull codes defined in Propositions 6-10 and Theorem 8. Then there exist Hermitian ℓ -dim hull codes with parameters $[N_1 - s_1, K_1 - s_1, \geq d_1 - s_1]_{q^2}$ for $1 \leq s_1 \leq K_1 - 1$ and for $1 \leq \ell \leq K_1 - s_1 - 1$.

V. APPLICATION TO CONSTRUCTIONS OF EAQECCS

In this section, we construct quantum codes from Hermitian hulls of linear codes.

By combining Lemma 1 and Lemma 2 together, q -ary EAQECCs can be constructed from a classical q^2 -linear code as follows.

Lemma 6. ([14]) Let C be a linear code with parameters $[n, k, d]_{q^2}$ and C^{\perp_h} its Hermitian dual with parameters $[n, k, d']_{q^2}$. Assume that $\dim(\text{Hull}_h(C)) = \ell$. Then there exist an $[[n, k - \ell, d; n - k - \ell]]_q$ EAQECC and an $[[n, n - k - \ell, d'; k - \ell]]_q$ EAQECC.

It is well-known the (Hermitian) dual of an MDS linear code is again an MDS linear code. It is easy to check that if a code C is a Hermitian ℓ -dim hull code, then so is its dual C^{\perp_h} . By applying the above lemma to the Hermitian ℓ -dim hull $[n, k, d]_{q^2}$ codes constructed in the previous section, we obtain the following result.

Theorem 9. Let $q = p^m$ and $1 \leq k \leq \lfloor \frac{n+q-1}{q+1} \rfloor$. Then there exist MDS EAQECCs with parameters $[[n - s, k - s - \ell, n - k + 1; n - k - \ell]]_q$ and $[[n - s, n - k - \ell, k - s + 1; k - s - \ell]]_q$ for $0 \leq s \leq k - 1$ and $0 \leq \ell \leq k - s$ if one of the conditions holds

- 1) $n = q^2 + 1, k \leq q, k \neq q - 1; n = r(q - 1) + 1, k \leq (q + r - 1)/2, q \equiv r - 1 \pmod{2r}; n = (q^2 + 2)/3, 3|(q + 1), k \leq (2q - 1)/3;$
- 2) $n = tq, 1 \leq t \leq q, k \leq \lfloor \frac{tq+q-1}{q+1} \rfloor; n = t(q + 1) + 2, 1 \leq t \leq q - 1, k \leq t + 1, (p, t, k) \neq (2, q - 1, q - 1);$
- 3) $(n - 1)|(q^2 - 1), k \leq \lfloor \frac{n+q-1}{q+1} \rfloor; n = (t + 1)N + i, 1 \leq i \leq 2, N|(q^2 - 1), n_2 = \frac{N}{\gcd(N, q+1)}, 1 \leq t \leq \frac{q-1}{n_2} - 1, k \leq \lfloor \frac{n+q-1}{q+1} \rfloor.$

Remark 2. It should be noted that the families in Theorem 9 2) were obtained in [12] by different approach for $(n, s) = (tq, 0)$ and $(n, s) = (t(q+1), 0)$. However, their method does not allow to puncture coordinates to obtain new parameters of EAQECCs as our method does.

Theorem 10. Let $q = p^m, (n - 1)|(q^2 - 1)$ and $1 \leq k \leq \lfloor \frac{n+q-1}{q+1} \rfloor$.

- 1) If $(n - 1)|(k + 1)(q + 1)$, then there exists an AMDS EAQECC with parameters $[[n + 1, k + 2 - \ell, n - k - 1; n - k - 1 - \ell]]_q$ for $0 \leq \ell \leq k + 2$
- 2) If $(n - 1)|k(q + 1)$, then there exists an EAQECC with parameters $[[n + i - 1, k + i - \ell, n - k - i + 1; n - k - 1 - \ell]]_q$ for $0 \leq \ell \leq k + i$. Moreover, there exists an AMDS EAQECC with parameters $[[q^2 + 1, k - \ell, q^2 + 1 - k; q^2 + 1 - k - \ell]]_q$ for $k = q + 2, \dots, 2q - 2, 0 \leq \ell \leq k$.

Theorem 11. Let $q = p^m, (n - 1)|(q^2 - 1)$ and $k = \lfloor \frac{n+q-1}{q+1} \rfloor$. Then

- 1) if $(n - 1)|(k + 1)(q + 1)$, then there exist MDS EAQECCs with parameters $[[n, k + 2 - \ell, n - k - 1; n - k - 2 - \ell]]_q$ and $[[n, n - k - 2, k + 3; k + 2 - \ell]]_q$, where $\ell = (k + 1);$
- 2) if $(n - 1)|k(q + 1)$, then there exist MDS EAQECCs with parameters $[[n, k + i - \ell, n - k - i + 1; n - k - i - \ell]]_q$ and $[[n, n - k - i - \ell, k + i + 1; k + i - \ell]]_q$ for any $1 \leq i \leq q$, where $\ell = k + i - \#I$; Moreover,
 - a) there exist MDS EAQECCs with parameters $[[q^2, k + i - \ell, q^2 - k - i + 1; q^2 - k - i - \ell]]_q$ and $[[q^2, q^2 - k - i - \ell, k + i + 1; k + i - \ell]]_q$ for $1 \leq i \leq q$, where $\ell = k + i - 1;$
 - b) there exist MDS EAQECCs with parameters $[[2(q + 1) + 1, k + i - \ell, 2(q + 1) + 1 - k - i + 1; 2(q + 1) + 1 - k - i - \ell]]_q$ and $[[2(q + 1) + 1, 2(q + 1) + 1 - k - i - \ell, k + i + 1; k + i - \ell]]_q$ for $1 \leq i \leq q$, where $\ell = (k + i - \lfloor \frac{i-1}{2} \rfloor - 1).$

Theorem 12. Let q be an odd prime power, $N|(q^2 - 1), n_2 = \frac{N}{\gcd(N, q+1)}, 1 \leq t \leq \frac{q-1}{n_2} - 1$. Put $n = (t + 1)N + 1, k = \lfloor \frac{n+q-1}{q+1} \rfloor, k_i = k + i$ and $n_i = n + i$. Then

- 1) there exist AMDS EAQECCs with parameters $[[n + 2, k + 2, \geq n - k; n - k - \ell]]_q$ and $[[n + 2, n - k - \ell, \geq k + 2; k + 2 - \ell]]_q$ for $0 \leq \ell \leq k + 2;$
- 2) there exist EAQECCs with parameters $[[n + i, k + i - \ell, \geq n - k - i + 1; n - k - \ell]]_q$ code for $0 \leq \ell \leq k + i, 1 \leq i \leq q;$
- 3) there exist EAQECCs with parameters $[[n, 2k - 1 - \ell, \geq n - k - q + 2; n - 2k + 1 - \ell]]_q, 0 \leq \ell \leq 2k - 1;$
- 4) there exist MDS EAQECCs with parameters $[[n, k + i - \ell, n - k - i + 1; n - k - i - \ell]]_q$ and $[[n, n - k - i - \ell, k + i + 1; k + i - \ell]]_q$ with $\ell = k + i - \#I$ for $1 \leq i \leq q$.

Theorem 13. Let $q = 2^s, m = 2s$ and $2k + 2 \leq n \leq \lfloor q + \sqrt{2q} - 5 \rfloor$. Then there exist AMDS EAQECCs with parameters $[[n, k - \ell, \leq n - k; n - k - \ell]]_q$ and $[[n, n - k - \ell, k; k - \ell]]_q$ for $0 \leq \ell \leq k$.

Theorem 14. Let $q = 2^m, m \geq 2$ and $q \leq k \leq \lfloor \frac{2q^2+2q-1}{q+1} \rfloor$. Then there exist EAQECCs with parameters $[[2q^2, k - q/2 - \ell, \geq 2q^2 - k + 1; 2q^2 - k + q/2 - \ell]]_q$ and $[[2q^2, 2q^2 + q/2 - k, \geq k + 1 - q; 2q^2, k - q/2 - \ell]]_q$ for $0 \leq \ell \leq k - q/2$.

TABLE IV
SOME NEW MDS AND AMDS EAQECCS FROM THEOREM 9 WITH $s = 0, 1$ AND THEOREM 11 2) WITH $i = 2$, *: NEW PARAMETERS IN [12]

ℓ	$[n, k]_{q^2}$	MDS EAQECCs	MDS EAQECCs	Construction
1	$[15, 4]_{8^2}$	$[[15, 10, 5; 3]]_8^*$	$[[14, 10, 4; 2]]_8^*$	Theorem 9 1)
2		$[[15, 9, 5; 2]]_8^*$	$[[14, 9, 4; 1]]_8^*$	
1	$[22, 5]_{8^2}$	$[[22, 16, 6; 4]]_8^*$	$[[21, 16, 5; 3]]_8^*$	
2		$[[22, 15, 6; 3]]_8^*$	$[[21, 15, 5; 2]]_8^*$	
1	$[29, 5]_{8^2}$	$[[29, 23, 6; 4]]_8^*$	$[[28, 23, 5; 3]]_8^*$	
2		$[[29, 22, 6; 3]]_8^*$	$[[28, 22, 5; 2]]_8^*$	
1	$[36, 6]_{8^2}$	$[[36, 29, 7; 5]]_8^*$	$[[35, 29, 6; 4]]_8^*$	
2		$[[36, 28, 7; 4]]_8^*$	$[[35, 28, 6; 3]]_8^*$	
1	$[43, 6]_{8^2}$	$[[43, 36, 7; 5]]_8^*$	$[[42, 36, 6; 4]]_8^*$	
2		$[[43, 35, 7; 4]]_8^*$	$[[42, 35, 6; 3]]_8^*$	
1	$[50, 7]_{8^2}$	$[[50, 42, 8; 6]]_8^*$	$[[49, 42, 7; 5]]_8^*$	
2		$[[50, 41, 8; 5]]_8^*$	$[[49, 41, 7; 4]]_8^*$	
1	$[57, 7]_{8^2}$	$[[57, 49, 8; 6]]_8^*$	$[[56, 49, 7; 5]]_8^*$	
2		$[[57, 48, 8; 5]]_8^*$	$[[56, 48, 7; 4]]_8^*$	
1	$[72, 7]_{9^2}$	$[[72, 64, 8; 6]]_9$	$[[71, 64, 7; 5]]_9$	Theorem 9 2)
2		$[[72, 63, 8; 5]]_9$	$[[71, 63, 7; 4]]_9$	
3		$[[72, 62, 8; 4]]_9$	$[[71, 62, 7; 3]]_9$	
4		$[[72, 61, 8; 3]]_9$	$[[71, 61, 7; 2]]_9$	
5		$[[72, 60, 8; 2]]_9$	$[[71, 60, 7; 1]]_9$	
6		$[[72, 59, 8; 1]]_9$	$[[71, 59, 7; 0]]_9$	
1	$[96, 8]_{11^2}$	$[[96, 87, 9; 7]]_{11}$	$[[95, 87, 8; 6]]_{11}$	Theorem 9 3)
2		$[[96, 86, 9; 6]]_{11}$	$[[95, 86, 8; 5]]_{11}$	
3		$[[96, 85, 9; 5]]_{11}$	$[[95, 85, 8; 4]]_{11}$	
4		$[[96, 84, 9; 4]]_{11}$	$[[95, 84, 8; 3]]_{11}$	
5		$[[96, 83, 9; 3]]_{11}$	$[[95, 83, 8; 2]]_{11}$	
6		$[[96, 82, 9; 2]]_{11}$	$[[95, 82, 8; 1]]_{11}$	
7		$[[96, 81, 9; 1]]_{11}$	$[[95, 81, 8; 0]]_{11}$	
1	$[16, 3]_{4^2}$	$[[16, 7, 6; 1]]_4^*$		Theorem 11 2)
1	$[25, 4]_{5^2}$	$[[25, 14, 7; 1]]_5^*$		
1	$[49, 6]_{7^2}$	$[[49, 34, 9; 1]]_7^*$		
1	$[64, 7]_{8^2}$	$[[64, 47, 10; 1]]_8^*$		
1	$[81, 8]_{9^2}$	$[[81, 62, 11; 1]]_9^*$		
1	$[121, 10]_{11^2}$	$[[121, 98, 13; 1]]_{11}^*$		

Theorem 15. Let $q = p^m \geq 4$ and $q(q-1) \leq k \leq \lfloor \frac{n+q(q-1)-1}{q+1} \rfloor$. Then there exist EAQECCs with parameters $[[Nq, k - \frac{q(q-1)}{2} - \ell, \geq Nq - k + 1; Nq - k + \frac{q(q-1)}{2} - \ell]]_q$ and $[[Nq, Nq + q(q-1)/2 - k - \ell, \geq k + 1 - q(q-1); k - q(q-1)/2 - \ell]]_q$ for $0 \leq \ell \leq k - \frac{q(q-1)}{2}$ if one of the following conditions holds

- 1) $n = q^2 + 1, k \leq q, k \neq q - 1; n = r(q-1) + 1, k \leq (q+r-1)/2, q \equiv r-1 \pmod{2r}; n = (q^2+2)/3, 3|(q+1), k \leq (2q-1)/3;$
- 2) $n = tq, 1 \leq t \leq q, k \leq \lfloor \frac{tq+q-1}{q+1} \rfloor; n = t(q+1) + 2, 1 \leq t \leq q-1, k \leq t+1, (p, t, k) \neq (2, q-1, q-1);$
- 3) $(n-1)|(q^2-1); n = (t+1)N + i, 1 \leq i \leq 2, N|(q^2-1), n_2 = \frac{N}{\gcd(N, q+1)}, 1 \leq t \leq \frac{q-1}{n_2} - 1.$

Theorem 16. Let $q = p^m$ be an odd prime power, $1 < t < q+1$ and $t|(q+1)$. Put $g = \frac{(q-1)(q+1-t)}{2t}$ and $s = q \left(\frac{q^2-2t}{t} + 2 \right)$. Then for $2g \leq k \leq \lfloor \frac{s+q-1}{q+1} \rfloor$ and $0 \leq \ell \leq k-g$, there exist EAQECCs with parameters $[[s, k-g-\ell, \geq s-k+1; s-k+g-\ell]]_q$ and $[[s, s-k+g-\ell, \geq k-2g+1; k-g-\ell]]_q$.

Theorem 17. Let N_1, K_1 and d_1 be the length, dimension and minimum distance of the Hermitian ℓ -dim hull codes defined in Propositions 6-10 and Theorem 8. Then there exist EAQECCs with parameters $[[N_1 - s_1, K_1 - s_1 - \ell, \geq d_1 - s_1; N_1 - K_1 - \ell]]_q$ for $1 \leq s_1 \leq K_1 - 1$ and for $1 \leq \ell \leq K_1 - s_1$.

REFERENCES

- [1] A. Ashikhmin and E. Knill, "Nonbinary quantum stabilizer codes," *IEEE Trans. Inf. Theory*, vol. 47, no. 7, pp. 3065-3072, Nov. 2001.
- [2] J. Bierbrauer and Y. Edel, "Quantum twisted codes," *J. Comb. Designs*, vol. 8, pp. 174-188, 2000.
- [3] W. Bosma and J. Cannon, *Handbook of Magma Functions*, Sydney, 1995.
- [4] G. Bowen, "Entanglement required in achieving entanglement-assisted channel capacities," *Physical Review A*, 66, 052313-1-052313-8 (Nov 2002).
- [5] T. Brun, I. Devetak and M.H. Hsieh, "Correcting quantum errors with entanglement," *Science*, vol. 314, pp. 436-439, Oct. 2006.
- [6] A.R. Calderbank, E.M. Rains, P.W. Shor and N.J.A. Sloane, "Quantum error correction via codes over GF(4)," *IEEE Trans. Inf. Theory*, vol. 44, no. 4, pp. 1369-1387, July 1998.
- [7] C. Carlet, C. Li and S. Mesnager, "Linear codes with small hulls in semi-primitive case," *Des. Codes Cryptogr.* <https://doi.org/10.1007/s10623-019-00663-4>
- [8] C. Carlet, S. Mesnager, C. Tang, Y. Qi and R. Pellikaan, "Linear codes over \mathbb{F}_q are equivalent to LCD codes for $q > 3$," *IEEE Trans. Inf. Theory*, 64(4), pp. 3010-3017, 2018.
- [9] C. Carlet, S. Mesnager, C. Tang and Y. Qi, "Euclidean and Hermitian LCD MDS codes," *Des. Codes Cryptogr.* 86, pp. 2605-2618, 2018.
- [10] S.M. Dodunekov and I.N. Landjev, "Near-MDS codes over some small fields," *Discrete Math.* 213 (2000) 55-65.
- [11] W. Fang and F.-W. Fu, "Two new classes of quantum MDS codes," *Finite Fields Appl.*, vol. 53, pp. 85-98, Sep. 2018.
- [12] W. Fang, F.-W. Fu, L. Li and S. Zhu, "Euclidean and Hermitian Hulls of MDS Codes and Their Applications to EAQECCs," *IEEE Transactions on Information Theory*, vol. 66(6), pp. 3527-3537, June 2020.
- [13] V. D. Goppa, "Algebraico-geometric codes," *Math. USSR-Izv.* 21(1) (1983) 75-91
- [14] K. Guenda, S. Jitman and T.A. Gulliver, "Constructions of good entanglement assisted quantum error correcting codes," *Des. Codes Cryptogr.*, vol. 86, pp. 121-136, Jan. 2018.

TABLE V
SOME NEW MDS AND MDS EAQECs FROM THEOREM 9 3) WITH $s = 0, 1$ AND THEOREM 10 WITH $i = 2$, *: NEW PARAMETERS IN [12]

ℓ	$[n, k]_{q^2}$	MDS EAQECs	MDS EAQECs	AMDS EAQECs
1	$[16, 3]_{4^2}$	$[[16, 12, 4; 2]]_4$	$[[15, 12, 3; 1]]_4^*$	$[[17, 11, 5; 4]]_4^*$
2		$[[16, 11, 4; 1]]_4$	$[[15, 11, 3; 0]]_4$	$[[17, 10, 5; 3]]_4^*$
1	$[25, 4]_{5^2}$	$[[25, 20, 5; 3]]_5$	$[[24, 20, 4; 2]]_5$	$[[26, 19, 6; 5]]_5^*$
2		$[[25, 19, 5; 2]]_5$	$[[24, 19, 4; 1]]_5$	$[[26, 18, 6; 4]]_5^*$
3		$[[25, 18, 5; 1]]_5$	$[[24, 18, 4; 0]]_5$	$[[26, 17, 6; 3]]_5^*$
1	$[49, 6]_{7^2}$	$[[49, 42, 7; 5]]_7$	$[[48, 42, 6; 4]]_7$	$[[50, 42, \surd \surd \surd 8; 6]]_7^*$
2		$[[49, 41, 7; 4]]_7$	$[[48, 41, 6; 3]]_7$	$[[50, 40, \surd \surd \surd 8; 6]]_7^*$
3		$[[49, 40, 7; 3]]_7$	$[[48, 40, 6; 2]]_7$	$[[50, 39, \surd \surd \surd 8; 5]]_7^*$
4		$[[49, 39, 7; 2]]_7$	$[[48, 39, 6; 1]]_7$	$[[50, 38, \surd \surd \surd 8; 4]]_7^*$
5		$[[49, 38, 7; 1]]_7$	$[[48, 38, 6; 0]]_7$	$[[50, 37, \surd \surd \surd 8; 3]]_7^*$
1	$[64, 7]_{8^2}$	$[[64, 56, 8; 6]]_8$	$[[63, 56, 7; 5]]_8$	$[[65, 55, \surd \surd \surd \surd \surd 9; 8]]_8^*$
2		$[[64, 55, 8; 5]]_8$	$[[63, 55, 8; 4]]_8^*$	$[[65, 54, \surd \surd \surd \surd 9; 7]]_8^*$
3		$[[64, 54, 8; 4]]_8$	$[[63, 54, 7; 3]]_8$	$[[65, 53, \surd \surd \surd 9; 6]]_8^*$
4		$[[64, 53, 8; 3]]_8$	$[[63, 53, 7; 2]]_8$	$[[65, 52, \surd \surd \surd 9; 5]]_8^*$
5		$[[64, 52, 8; 2]]_8$	$[[63, 52, 7; 1]]_8$	$[[65, 51, \surd \surd \surd 9; 4]]_8^*$
6		$[[64, 51, 8; 1]]_8$	$[[63, 51, 7; 0]]_8$	$[[65, 50, \surd \surd \surd 9; 3]]_8^*$
1	$[81, 8]_{9^2}$	$[[81, 72, 9; 7]]_9$	$[[80, 72, 8; 6]]_9$	$[[82, 71, \surd \surd \surd \surd \surd 10; 9]]_9^*$
2		$[[81, 71, 9; 6]]_9$	$[[80, 71, 8; 5]]_9$	$[[82, 70, \surd \surd \surd \surd 10; 8]]_9^*$
3		$[[81, 70, 9; 5]]_9$	$[[80, 70, 8; 4]]_9$	$[[82, 69, \surd \surd \surd \surd 10; 7]]_9^*$
4		$[[81, 69, 9; 4]]_9$	$[[80, 69, 8; 3]]_9$	$[[82, 68, \surd \surd \surd \surd 10; 6]]_9^*$
5		$[[81, 68, 9; 3]]_9$	$[[80, 68, 8; 2]]_9$	$[[82, 67, \surd \surd \surd \surd 10; 5]]_9^*$
6		$[[81, 67, 9; 2]]_9$	$[[80, 67, 8; 1]]_9$	$[[82, 66, \surd \surd \surd \surd 10; 4]]_9^*$
7		$[[81, 66, 9; 1]]_9$	$[[80, 66, 8; 0]]_9$	$[[82, 65, \surd \surd \surd \surd 10; 3]]_9^*$
1	$[121, 10]_{11^2}$	$[[121, 110, 11; 9]]_{11}$	$[[120, 110, 10; 8]]_{11}^*$	$[[122, 109, \surd \surd \surd \surd \surd \surd 12; 11]]_{11}^*$
2		$[[121, 109, 11; 8]]_{11}$	$[[120, 109, 10; 7]]_{11}$	$[[122, 108, \surd \surd \surd \surd \surd 12; 10]]_{11}^*$
3		$[[121, 108, 11; 7]]_{11}$	$[[120, 108, 10; 6]]_{11}$	$[[122, 107, \surd \surd \surd \surd \surd 12; 9]]_{11}^*$
4		$[[121, 107, 11; 6]]_{11}$	$[[120, 107, 10; 5]]_{11}$	$[[122, 106, \surd \surd \surd \surd \surd 12; 8]]_{11}^*$
5		$[[121, 106, 11; 5]]_{11}$	$[[120, 106, 10; 4]]_{11}$	$[[122, 105, \surd \surd \surd \surd \surd 12; 7]]_{11}^*$
6		$[[121, 105, 11; 4]]_{11}$	$[[120, 105, 10; 3]]_{11}$	$[[122, 104, \surd \surd \surd \surd \surd 12; 6]]_{11}^*$
7		$[[121, 104, 11; 3]]_{11}$	$[[120, 104, 10; 2]]_{11}$	$[[122, 103, \surd \surd \surd \surd \surd 12; 5]]_{11}^*$
8		$[[121, 103, 11; 2]]_{11}$	$[[120, 103, 10; 1]]_{11}$	$[[122, 102, \surd \surd \surd \surd \surd 12; 4]]_{11}^*$
9		$[[121, 102, 11; 1]]_{11}$	$[[120, 102, 10; 0]]_{11}$	$[[122, 101, \surd \surd \surd \surd \surd 12; 3]]_{11}^*$

TABLE VI
SOME NEW EAQECs FROM THEOREM 13 AND THEOREM 14, *: NEW PARAMETERS

ℓ	$[n, k]_{q^2}$	EAQECs	Punctured EAQECs	Construction
1	$[32, 6, 25]_{4^2}$	$[[32, 5, 25; 25]]_4^*$	$[[31, 4, \surd 24; 25]]_4^*$	Theorems 14, 17
2		$[[32, 4, 25; 24]]_4^*$	$[[31, 3, \surd 24; 24]]_4^*$	
3		$[[32, 3, 25; 23]]_4^*$	$[[31, 2, \surd \surd 24; 23]]_4^*$	
4		$[[32, 2, 25; 22]]_4^*$	$[[31, 1, \surd \surd \surd 24; 22]]_4^*$	
5		$[[32, 1, 25; 21]]_4^*$	$[[31, 0, \surd \surd \surd \surd 24; 21]]_4^*$	
1	$[80, 9, 71]_{8^2}$	$[[80, 8, 71; 70]]_8$	$[[79, 7, \surd \surd \surd 70; 70]]_8$	Theorems 13, 17
2		$[[80, 7, 71; 69]]_8$	$[[79, 6, \surd \surd \surd 70; 69]]_8$	
3		$[[80, 6, 71; 68]]_8$	$[[79, 5, \surd \surd \surd 70; 68]]_8$	
4		$[[80, 5, 71; 67]]_8$	$[[79, 4, \surd \surd \surd 70; 67]]_8$	
5		$[[80, 4, 71; 66]]_8$	$[[79, 3, \surd \surd \surd 70; 66]]_8$	
6		$[[80, 3, 71; 65]]_8$	$[[79, 2, \surd \surd \surd 70; 65]]_8$	
7		$[[80, 2, 71; 64]]_8$	$[[79, 1, \surd \surd \surd 70; 64]]_8$	
8		$[[80, 1, 71; 63]]_8$	$[[79, 0, \surd \surd \surd 70; 63]]_8$	
1	$[128, 12, 113]_{8^2}$	$[[128, 11, 113; 115]]_{8^2}^*$	$[[127, 10, \surd \surd \surd \surd \surd \surd 112; 115]]_{8^2}^*$	Theorem 14, 17
2		$[[128, 10, 113; 114]]_{8^2}$	$[[127, 9, \surd \surd \surd \surd \surd 112; 114]]_{8^2}$	
3		$[[128, 9, 113; 113]]_{8^2}$	$[[127, 8, \surd \surd \surd \surd \surd 112; 113]]_{8^2}$	
4		$[[128, 8, 113; 112]]_{8^2}$	$[[127, 7, \surd \surd \surd \surd \surd 111; 112]]_{8^2}$	
5		$[[128, 7, 113; 111]]_{8^2}$	$[[127, 6, \surd \surd \surd \surd \surd 112; 111]]_{8^2}$	
6		$[[128, 6, 113; 110]]_{8^2}$	$[[127, 5, \surd \surd \surd \surd \surd 112; 110]]_{8^2}$	
7		$[[128, 5, 113; 109]]_{8^2}$	$[[127, 4, \surd \surd \surd \surd \surd 112; 109]]_{8^2}$	
8		$[[128, 4, 113; 108]]_{8^2}$	$[[127, 3, \surd \surd \surd \surd \surd 112; 108]]_{8^2}$	
9		$[[128, 3, 113; 107]]_{8^2}$	$[[127, 2, \surd \surd \surd \surd \surd 112; 107]]_{8^2}$	
10		$[[128, 2, 113; 106]]_{8^2}$	$[[127, 1, \surd \surd \surd \surd \surd 112; 106]]_{8^2}$	
11		$[[128, 1, 113; 105]]_{8^2}$	$[[127, 0, \surd \surd \surd \surd \surd 112; 105]]_{8^2}$	

- [15] K. Guenda, T.A. Gulliver, S. Jitman and S. Thipworawimon, "Linear ℓ -intersection pairs of codes and their applications," *Des. Codes Cryptogr.*, 2019. <https://doi.org/10.1007/s10623-019-00676-z>
- [16] M. Grassl and T. Beth, "Quantum BCH codes," *Proceedings of International Symposium on Theoretical Electrical Engineering Magdeburg*, pp. 207–212, Oct. 1999. DOI:10.1109/ICCES.2008.4772987
- [17] M. Grassl, T. Beth and M. Röttler, "On optimal quantum codes," *Int. J. Quantum Inf.*, vol. 2, no. 1, pp. 757–775, 2004.
- [18] G.G.L. Guardia, "New quantum MDS codes," *IEEE Trans. Inform. Theory*, vol. 57, no. 8, pp. 5551–5554, 2011.
- [19] L.F. Jin, "Quantum Stabilizer Codes from Maximal Curves," *IEEE Trans. Inform. Theory*, vol. 60, no. 1, pp. 313–316, Jan. 2014.
- [20] L. Jin and C. Xing, "Euclidean and Hermitian Self-Orthogonal Algebraic Geometry Codes and Their Application to Quantum Codes," *IEEE Trans. Inform. Theory*, vol. 58, no. 8, pp. 5484–5489, Aug. 2012.
- [21] L.F. Jin and C.P. Xing, "A Construction of New Quantum MDS Codes," *IEEE Trans. Inform. Theory*, vo. 60, no. 5, pp. 2921–2925, May 2014
- [22] L.F. Jin, L. San, J.Q. Luo and C.P. Xing, "Application of Classical Hermitian Self-Orthogonal MDS Codes to Quantum MDS Codes," *IEEE Trans. Inform. Theory*, vol. 56, no. 8, pp. 4735–4740, Sep. 2010.
- [23] X. Kai and S. Zhu, "New quantum MDS codes from negacyclic codes," *IEEE Trans. Inform. Theory*, vol. 59, no. 2, pp. 1193–1197, 2012.
- [24] A. Ketkar, A. Klappenecker, S. Kumar and P. Sarvepalli, "Nonbinary stablizer codes over finite fields," *IEEE Trans. Inform. Theory*, vol. 52, no. 11, pp.4892–4914, Nov. 2006.
- [25] C.-Y. Lai and T.A. Brun, "Entanglement increases the error-correcting ability of quantum error-correcting codes," *Phys. Rev. A, Gen. Phys.*, vol. 88, p. 012320, Jul. 2013.

- [26] C.Y. Lai and A. Ashikhmin, "Linear programming bounds for entanglement-assisted quantum error correcting codes by split weight enumerators," *IEEE Trans. Inf. Theory*, vol. 64, no. 1, pp. 622-639, Jan. 2018.
- [27] Z. Li, J. Xing and X.M. Wang, "Quantum generalized Reed-Solomon codes: unified framework for quantum MDS codes," *Phys. Rev. A*, vol. 77, pp. 012308-1–12308-4, 2008.
- [28] C. Li and P. Zeng, "Constructions of linear codes with one-dimensional hull," *IEEE Trans. Inf. Theory*, 65 (3), pp. 1668–1676, 2019.
- [29] G. Luo, X. Cao and X. Chen, "MDS codes with hulls of arbitrary dimensions and their quantum error correction," *IEEE Trans. Inf. Theory*, vol. 65, no. 5, pp. 2944-2952, May 2019.
- [30] F.R.F. Pereira, R. Pellikaan, G.G.L. Guardia, F.M.D. Assis, "Entanglement-assisted Quantum Codes from Algebraic Geometry Codes," <https://arxiv.org/pdf/1907.06357.pdf>
- [31] L. Qian, X. Cao and S. Mesnager, "Linear codes with one-dimensional hull associated with Gaussian sums", *Cryptogr. Commun.* (2020). <https://doi.org/10.1007/s12095-020-00462-y>
- [32] S. Mesnager, C. Tang and Y. Qi, "Complementary dual algebraic geometry codes," *IEEE Trans. Inf. Theory*, 64(4), pp. 2390–2397, 2018.
- [33] L. Sok, "MDS linear codes with one dimensional hull," <https://arxiv.org/pdf/2012.11247.pdf>
- [34] L. Sok, "On linear codes with one-dimensional Euclidean hull and their applications to EAQECCs," <https://arxiv.org/abs/2101.06461>
- [35] L. Sok, "New families of quantum stabilizer codes from Hermitian self-orthogonal algebraic geometry codes", submitted for publication
- [36] H. Stichtenoth, "Algebraic function fields and codes," Springer, 2008.
- [37] M.M. Wilde and T.A. Brun, "Optimal entanglement formulas for entanglement-assisted quantum coding," *Physical Review A*, 77(6), 064302–1–064302–4 (Jun 2008).