

# Partial Order in the Normal Category Arising from Normal Bands

C S Preenu

University College Thiruvananthapuram  
Research Centre University of Kerala

A R Rajan

Former Professor of Mathematics  
University of Kerala

K S Zeenath

Former Professor of Mathematics  
School of Distance Education  
University of Kerala

## Abstract

The notion of normal category was introduced by KSS Nambooripad in connection with the study of the structure of regular semigroups using cross connections[4]. It is an abstraction of the category of principal left ideals of a regular semigroup. A normal band is a semigroup  $B$  satisfying  $a^2 = a$  and  $abca = acba$  for all  $a, b, c \in B$ . Since the normal bands are regular semigroups, the category  $\mathcal{L}(B)$  of principal left ideals of a normal band  $B$  is a normal category. One of the special properties of this category is that the morphism sets admit a partial order compatible with the composition of morphisms. In this article we derive several properties of this partial order and obtain a new characterization of this partial order.

Keywords: Partial order, Normal categories, Normal bands.

AMS Subject Classification : 20M17

## 1 Preliminaries

In this section we discuss some basic concepts of normal categories, normal bands and the normal category derived from normal bands. We follow [2, 4] for other notations and definitions which are not mentioned.

### 1.1 Normal Categories

We begin with the category theory, some basic definitions and notations. In a category  $C$ , the class of objects is denoted by  $vC$  and  $C$  itself is used

to denote the morphism class. The collection of all morphisms from  $a$  to  $b$  in  $C$  is denoted by  $C(a, b)$  and  $1_a$  denote the identity morphism in  $C(a, a)$ . A morphism  $f \in C$  is said to be a monomorphism(epimorphism) if  $fg = fh(gf = hf)$  implies  $g = h$  for any two morphisms  $g, h \in C$ . A morphism  $f \in C(a, b)$  is said to be right(left) invertible if there exists  $g \in C(b, a)$  such that  $fg = 1_a(gf = 1_b)$ . An isomorphism means a morphism which is both left and right invertible.

Let  $S$  be a regular semigroup. A right translation of  $S$  is a function  $\phi$  from  $S$  to  $S$  such that  $(xy)\phi = x((y)\phi)$  for all  $x, y \in S$ . Similarly a function  $\psi : S \rightarrow S$  is called a left translation if  $\psi(xy) = (\psi(x))y$  for all  $x, y \in S$ . For a fixed  $u \in S$ , the map  $\rho_u : x \mapsto xu$  for  $x \in S$  is a right translation. This map can be viewed as a function from  $S$  to the principal left ideal  $Su$  of  $S$ . Since  $S$  is regular, we can find an idempotent  $f \in S$  such that  $Su = Sf$ . In particular if we choose  $u \in eSf$  the restriction of  $\rho_u$  to  $Se$ , denoted by  $\rho(e, u, f)$ , can be viewed as a function from  $Se$  to  $Sf$ . If  $\rho(e, u, f) : Se \rightarrow Sf$  and  $\rho(g, v, h) : Sg \rightarrow Sh$  are such that  $Sf = Sg$ , the composition of these maps will be  $\rho(e, uv, h)$ . Now we arrive at a category structure,  $\mathcal{L}(S)$ , with objects are the principal left ideals of  $S$  and morphisms are of the form  $\rho(e, u, f) : Se \rightarrow Sf$  where  $u \in eSf$ . The usual set inclusion of the principal left ideals will induce a partial order in the objects class of  $\mathcal{L}(S)$ . Also  $Se \subseteq Sf$  implies  $ef = e$  and hence the morphism  $\rho(e, e, f)$  is an inclusion map from  $Se$  to  $Sf$  and is denoted by  $j(Se, Sf)$ . The category  $\mathcal{L}(S)$  become a normal category whose definition is given below.

A preorder is a category  $P$  such that each of the hom sets contain at most one morphism. We write  $a \leq b$  whenever  $P(a, b) \neq \emptyset$ .

**Definition 1.1** ([8]). A category with normal factorization is a small category  $C$  with the following properties.

1. The vertex set  $vC$  of  $C$  is a partially ordered set such that whenever  $a \leq b$  in  $vC$ , there is a monomorphism  $j(a, b) : a \rightarrow b$  in  $C$ . This morphism is called the inclusion from  $a$  to  $b$ .
2.  $j : (vC, \leq) \rightarrow C$  is a functor from the preorder  $(vC, \leq)$  to  $C$  which maps  $a \leq b$  to  $j(a, b)$  for  $a, b \in vC$ .
3. For  $a, b \leq c$  in  $vC$ , if

$$j(a, c) = fj(b, c)$$

for some  $f : a \rightarrow b$  then  $a \leq b$  and  $f = j(a, b)$ .

4. Every morphism  $j(a, b) : a \rightarrow b$  has a right inverse  $q : b \rightarrow a$  such that

$$j(a, b)q = 1_a$$

Such a morphism  $q$  is called a retraction in  $C$ .

5. Every morphism  $f$  in  $C$  has a factorization

$$f = quj$$

where  $q$  is a retraction,  $u$  is an isomorphism and  $j$  is an inclusion. Such a factorization is called normal factorization in  $C$ .

**Proposition 1.1** ([4]). Let  $C$  is a category with normal factorization. If  $f = quj = q'u'j'$  are two normal factorizations of  $f \in C$  then  $j = j'$  and  $qu = q'u'$ .

In this case  $f^0 = qu$  is called the epimorphic component of  $f$ . The codomain of  $f^0$  is called the image of  $f$  and is denoted by  $\text{im } f$

Corresponding to a right translation  $\rho_u$  from  $S$  to  $Su$ , we get morphisms from each of the principal left ideals of  $S$  to  $Su$  by restricting the  $\rho_u$ . This phenomena is similar to the concept of cones in the category theory[2]. Now we define normal cones and normal categories.

**Definition 1.2.** [4] Let  $C$  be a category with normal factorization. A cone  $\gamma$  with vertex  $a \in vC$  is a function from  $vC$  to  $C$  satisfying the following

1.  $\gamma(c) \in C(c, a)$  for all  $c \in vC$
2. If  $c_1 \subseteq c_2$  then  $\gamma(c_1) = j(c_1, c_2)\gamma(c_2)$
3. There is an object  $b \in vC$  such that  $\gamma(b)$  is an isomorphism.

**Definition 1.3** ([6]). A normal category is a category with normal factorization such that for each  $a \in vC$  there is a normal cone  $\gamma$  with  $\gamma(a) = 1_a$ .

The cones induced by a right translation  $\rho_u$  is called a principal cone and is denoted by  $\rho^u$ . The category  $\mathcal{L}(S)$  may contain normal cones other than principal cones.

## 1.2 Normal Bands

Several studies have been carried out for recent years about the normal categories arising from the semigroups satisfying some special conditions[8, 3, 5]. We focus on the normal category of principal left ideals of a normal band. A semigroup  $B$  is said to be a normal band if  $a^2 = a$  and  $abca = acba$  for all  $a, b, c \in B$ . The following theorem, by Yamada and Kimura[1], states that a semigroup is a normal band if and only if it is a strong semilattice of rectangular bands.

**Theorem 1.2** ([1, 9]). The following conditions are equivalent for a semigroup  $B$

1.  $B$  is a normal band

2. There exists a semilattice  $\Gamma$ , a disjoint family of rectangular bands  $\{E_\alpha : \alpha \in \Gamma\}$  such that  $B = \bigcup\{E_\alpha : \alpha \in \Gamma\}$  and a family of homomorphisms  $\{\phi_{\alpha,\beta} : E_\alpha \rightarrow E_\beta, \text{ where } \alpha, \beta \in \Gamma \text{ and } \beta \leq \alpha\}$  satisfying the following

- (a)  $\phi_{\alpha,\alpha}$  is the identity function,
- (b)  $\phi_{\alpha,\beta}\phi_{\beta,\gamma} = \phi_{\alpha,\gamma}$  for  $\gamma \leq \beta \leq \alpha$  and
- (c) For  $a, b \in B$  with  $a \in E_\alpha$  and  $b \in E_\beta$

$$ab = (a\phi_{\alpha,\alpha\beta})(b\phi_{\beta,\alpha\beta})$$

3.  $B$  is a band and

$$abcd = acbd \text{ for all } a, b, c, d \in B.$$

### 1.3 The category $\mathcal{L}(B)$

Let  $B$  is a normal band. In addition to that the category  $\mathcal{L}(B)$  is normal, it possess some special qualities which are not common in general. Several properties of the category  $\mathcal{L}(B)$  are described in [5]. Here the normal factorization is unique and there is only one isomorphism between any two isomorphic objects. Also the principal cones respect the isomorphic objects in the sense that

$$\gamma(Ba) = \rho(a, ab, b)\gamma(Bb)$$

for any principal cone  $\gamma$  and isomorphic objects  $Ba$  and  $Bb$ . This leads to the definition of strong cones by AR Rajan[7].

**Definition 1.4.** [7] A normal cone  $\gamma$  in a normal category  $C$  is said to be a strong cone if the following holds

If  $c$  and  $d$  are two isomorphic objects, then there is a unique isomorphism  $\alpha$  from  $c$  to  $d$  such that  $\gamma(c) = \alpha\gamma(d)$

Thus every principal cone is a strong cone. That is for every objects  $Ba$  of  $\mathcal{L}(B)$  there is a strong cone with vertex  $Ba$ . The category of principal left ideals of any normal band is characterized as follows.

**Theorem 1.3.** [5] A normal category  $C$  satisfies following conditions if and only if it is isomorphic to  $\mathcal{L}(B)$  for some normal band  $B$

**SC1** The isomorphism between any two objects is unique.

**SC2** The right inverse of an inclusion is unique.

**SC3** For every object  $c$  in  $C$ , there is a strong cone with vertex  $c$ .

## 2 Partial order in $\mathcal{L}(B)$

Let  $B$  is a normal band. Let us denote by  $[Ba, Bb]$ , the collection of all morphisms from  $Ba$  to  $Bb$  in  $\mathcal{L}(B)$ . Now we show that there is a partial order on the morphism sets  $[Ba, Bb]$  and that  $[Ba, Bb]$  is an order ideal with respect to this partial order. In particular there is a maximum element in  $[Ba, Bb]$ . We note that any normal band  $B$  has a natural partial order given by  $a \leq b$  if and only if  $ab = ba = a$ .

**Theorem 2.1.** [5] Let  $B$  be a normal band and  $\rho(a, u, b), \rho(a, v, b) \in [Ba, Bb]$  be morphisms in  $\mathcal{L}(B)$ . Define

$$\rho(a, u, b) \leq \rho(a, v, b) \text{ if } u \leq v$$

Then  $\leq$  is a partial order on  $[Ba, Bb]$ .

**Proposition 2.2.** Every morphism set  $[Ba, Bb]$  contains a maximum with respect to the above partial order.

*Proof.* Taking  $\delta = \alpha\beta$  where  $a \in E_\alpha$  and  $b \in E_\beta$  we see that there is a unique  $u \in E_{\alpha\beta}$  such that  $\rho(a, u, b) \in [Ba, Bb]$ .

Since  $\omega^l(b) \cap \omega^r(a) = \omega(h)$  for some  $h \in E_{\alpha\beta}$  and  $\rho(a, h, b) \in [Ba, Bb]$  we conclude that  $u = h$ . It follows that  $\rho(a, u, b)$  is the maximum in  $[Ba, Bb]$   $\square$

The following compatibility properties hold for this partial order.

**Proposition 2.3.** Let  $\rho(a, u, b)$  and  $\rho(a, v, b)$  be elements in the morphism set  $[Ba, Bb]$  in  $\mathcal{L}(B)$  such that  $\rho(a, u, b) \leq \rho(a, v, b)$ . Then the following compatibility relations hold.

1. If  $\rho(b, w, c) \in [Bb, Bc]$  then  $\rho(a, u, b)\rho(b, w, c) \leq \rho(a, v, b)\rho(b, w, c)$ .
2. If  $\rho(d, x, a) \in [Bd, Ba]$  then  $\rho(d, x, a)\rho(a, u, b) \leq \rho(d, x, a)\rho(a, v, b)$ .

*Proof.* Follows from the compatibility of natural partial order on normal bands.  $\square$

Now we describe some properties of the maximum elements in the morphism sets.

**Proposition 2.4.** Let  $m = m(a, b)$  be the maximum element in the morphism set  $[Ba, Bb]$  in  $\mathcal{L}(B)$  and  $\alpha = \alpha(b, c)$  be an isomorphism from  $Bb$  to  $Bc$  in  $\mathcal{L}(B)$ . Then  $m\alpha$  is the maximum element in the morphism set  $[Ba, Bc]$ .

*Proof.* Let  $f \in [Ba, Bc]$ . Then  $f\alpha^{-1} \in [Ba, Bb]$ . Since  $m$  is maximum element in  $[Ba, Bb]$  we have  $f\alpha^{-1} \leq m$ . Now by compatibility given by Proposition 2.3

$$f = f\alpha^{-1}\alpha \leq m\alpha$$

. So  $m\alpha$  is the maximum in  $[Ba, Bc]$ .  $\square$

**Proposition 2.5.** Let  $m = m(a, b)$  be the maximum element in the morphism set  $[Ba, Bb]$  in  $\mathcal{L}(B)$  and  $j = j(a_0, a)$  be an inclusion in  $\mathcal{L}(B)$ . Then  $jm$  is the maximum element in the morphism set  $[Ba_0, Bb]$ .

*Proof.* Let  $f \in [Ba_0, Bb]$ . Then  $f = \rho(a_0, u, b)$  for some  $u \leq k$  such that  $\omega(k) = a_0Bb$ . Now

$$jm = \rho(a_0, a_0, a)\rho(a, h, b) = \rho(a_0, a_0h, b) = \rho(a_0, k, b)$$

where  $\omega(h) = aBb$ . Therefore  $f \leq jm$  so that  $jm$  is maximum element in  $[Ba_0, Bb]$ .  $\square$

**Proposition 2.6.** Let  $m = m(a, b)$  be the maximum element in the morphism set  $[Ba, Bb]$  in  $\mathcal{L}(B)$  and  $q : b \rightarrow b_0$  be a retraction in  $\mathcal{L}(B)$ . Then  $mq$  is the maximum element in the morphism set  $[Ba, Bb_0]$ .

*Proof.* Let  $f \in [Ba, Bb_0]$ . Then  $fj(b_0, b)$  is in  $[Ba, Bb]$ . Since  $m$  is the maximum element in  $[Ba, Bb]$  we have  $fj(b_0, b) \leq m$ . Now by compatibility given by Proposition 2.3

$$f = fj(b_0, b)q \leq mq.$$

So  $mq$  is the maximum in  $[Ba, Bb_0]$ .  $\square$

As a consequence of this propositions we can see that  $mf^\circ$  is always a maximum if  $m$  is a maximum.

**Proposition 2.7.** Let  $m = m(a, b)$  be the maximum element in the morphism set  $[Ba, Bb]$  in  $\mathcal{L}(B)$  and  $f : b \rightarrow c$  be any morphism in  $\mathcal{L}(B)$ . Then  $mf^\circ$  is the maximum element in the morphism set  $[Ba, Bc_0]$  where  $c_0 \leq c$  is the codomain of  $f^\circ$ .

*Proof.* Suppose  $f = quj$ , where  $q$  is a retraction,  $u$  is an isomorphism and  $j$  is an inclusion. Then by the above propositions 2.4 and 2.6 we can say that  $mf^\circ = mqu$  is the maximum in  $[Ba, Bc_0]$   $\square$

The following proposition exactly locates the maximum element in each hom set in  $\mathcal{L}(B)$

**Proposition 2.8.** The maximum element in  $[Ba, Bb]$  is  $\rho(a, ab, b)$

*Proof.* Suppose  $\rho(a, u, b) \in [Ba, Bb]$ . Then  $u \in aBb$  and hence  $u = aub$ . Now

$$(ab)(u) = (ab)(aub) = aabub = aubb = aub = u$$

Similarly we get

$$(u)(ab) = aubab = auab = aub = u$$

That is  $u \leq ab$  for all  $u \in aBb$  and this proves that

$$\rho(a, u, b) \leq \rho(a, ab, b)$$

for all  $\rho(a, u, b) \in [Ba, Bb]$ .  $\square$

Note that  $\rho(a, ab, b)$  is the unique isomorphism in  $[Ba, Bb]$  whenever  $Ba$  and  $Bb$  are isomorphic. Also the following theorem shows that these morphisms are the building blocks of principal cones in  $\mathcal{L}(B)$ .

**Theorem 2.9.** The component of the principal cone  $\rho^a$  at  $Bb \in \mathcal{L}(B)$  is the maximum element in  $[Bb, Ba]$ .

*Proof.* We have the principal cone  $\rho^a$  with vertex  $Ba$  is defined as

$$\rho^a(Bb) = \rho(b, ba, a) \text{ for all } Bb \in v\mathcal{L}(B)$$

But  $\rho(b, ba, a)$  is the maximum in  $[Bb, Ba]$  by proposition 2.8. Hence the theorem.  $\square$

Now we give another characterization of the partial order in the hom sets of  $\mathcal{L}(B)$  using the image of the morphisms.

**Theorem 2.10.** For any two morphisms  $\rho(a, u, b), \rho(a, v, b) \in [Ba, Bb]$

$$\rho(a, u, b) \leq \rho(a, v, b) \text{ if and only if } \text{im } \rho(a, u, b) \subseteq \text{im } \rho(a, v, b)$$

*Proof.* To prove the theorem it is enough to show the following

$$\text{im } \rho(a, u, b) \subseteq \text{im } \rho(a, v, b) \text{ if and only if } u \leq v$$

The unique normal factorization[5] of  $\rho(a, u, b)$  is given by

$$\rho(a, u, b) = \rho(a, g, g)\rho(g, u, h)\rho(h, h, b).$$

Where  $u\mathcal{R}g \leq a$  and  $u\mathcal{L}h \leq b$ . Therefore  $\text{im } \rho(a, u, b) = Bh = Bu$ . Similarly we have  $\text{im } \rho(a, v, b) = Bv$ .

Assume that  $u \leq v$ . Then  $uv = vu = u$ . This implies  $Bu = Buv \subseteq Bv$ . Thus  $\text{im } \rho(a, u, b) \subseteq \text{im } \rho(a, v, b)$ .

Conversely suppose that  $\text{im } \rho(a, u, b) \subseteq \text{im } \rho(a, v, b)$ . That is  $Bu \subseteq Bv$ . Then  $uv = v$ .

Note that  $u, v \in aBb = aB \cap Bb$ . Therefore there exists  $p, q, r, s \in B$  such that  $u = ar = pb$  and  $v = as = qb$ . Also  $u = u^2 = arpb$  and  $v = v^2 = asqb$ . Then  $uv = u$  implies  $arqb = ar$  and  $pbqb = pb$ . Now we get

$$\begin{aligned} vu &= (asqb)(arpb) \\ &= (aras)(pbqb) && \text{By repeated application of normality} \\ &= (ar)(pb) \\ &= uu \\ &= u \end{aligned}$$

Therefore  $\text{im } \rho(a, u, b) \subseteq \text{im } \rho(a, v, b)$  implies  $uv = vu = u$  and hence  $u \leq v$ . Hence the theorem.  $\square$

## References

- [1] Naoki Kimura. The structure of idempotent semigroups. i. *Pacific Journal of Mathematics*, 8(2):257–275, 1958.
- [2] Saunders Mac Lane. *Categories for the working mathematician*, volume 5. Springer Science & Business Media, 2013.
- [3] PA Azeef Muhammed. Normal categories from completely simple semigroups. In *Algebra and its Applications*, pages 387–396. Springer, 2016.
- [4] KSS Nambooripad. *Theory of Cross-connections*. Centre for Mathematical Sciences, 1994.
- [5] C S Preenu, AR Rajan, and KS Zeenath. Category of principal left ideals of normal bands. In *Springer Proceedings in Mathematics & Statistics*. 2020(to appear).
- [6] A R Rajan. Inductive groupoids and normal categories of regular semigroups. In *Proceedings of the International Conference on Algebra and Its Applications*, pages 193–200. Aligarh Muslim University, Walter de Gruyter GmbH & Co KG, 2018.
- [7] AR Rajan. Normal categories associated with bands. In *Proceedings of the International Conference on Semigroups Algebra and Applications*, pages 21–29. Cochin University of Science and Technology, 2015.
- [8] AR Rajan. Normal categories of inverse semigroups. *East-West Journal of Mathematics*, 16(2), 2015.
- [9] Miyuki Yamada and Naoki Kimura. Note on idempotent semigroups ii. *Proceedings of the Japan Academy*, 34(2):110–112, 1958.