# L-balancing families

Gábor Hegedűs

Óbuda University Bécsi út 96, Budapest, Hungary, H-1037 hegedus.gabor@nik.uni-obuda.hu

May 5, 2021

#### Abstract

P. Hrubeš, S. Natarajan Ramamoorthy, A. Rao and A. Yehudayoff proved the following result:

Let p be a prime and let  $f \in \mathbb{F}_p[x_1, \ldots, x_{2p}]$  be a polynomial. Suppose that  $f(\mathbf{v_F}) = 0$  for each  $F \subseteq [2p]$ , where |F| = p and that  $f(\mathbf{0}) \neq 0$ . Then  $\deg(f) \geq p$ .

We prove here the following generalization of their result.

Let p be a prime and  $q = p^{\alpha} > 1$ ,  $\alpha \ge 1$ . Let n > 0 be a positive integer and  $q - 1 \le d \le n - q + 1$  be an integer. Let  $\mathbb{F}$  be a field of characteristic p. Suppose that  $f(\mathbf{v_F}) = 0$  for each  $F \subseteq [n]$ , where |F| = d and  $\deg(f) \le q - 1$ . Then  $f(\mathbf{v_F}) = 0$  for each  $F \subseteq [n]$ , where  $|F| \equiv d \pmod{q}$ .

Let t = 2d be an even number and  $L \subseteq [d-1]$  be a given subset. We say that  $\mathcal{F} \subseteq 2^{[t]}$  is an *L*-balancing family if for each  $F \subseteq [t]$ , where |F| = d there exists a  $G \subseteq [n]$  such that  $|F \cap G| \in L$ .

We give a general upper bound for the size of an L-balancing family.

## 1 Introduction

First we introduce some notations.

Let n be a positive integer and let [n] stand for the set  $\{1, 2, ..., n\}$ . The family of all subsets of [n] is denoted by  $2^{[n]}$ . For an integer  $0 \le d \le n$ 

we denote by  $\binom{[n]}{d}$  the family of all d element subsets of [n], and  $\binom{[n]}{\leq d} = \binom{[n]}{0} \cup \ldots \cup \binom{[n]}{d}$  the subsets of size at most d.

Let  $\mathbb{F}$  be a field.  $\mathbb{F}[x_1, \ldots, x_n]$  denotes the ring of polynomials in variables  $x_1, \ldots, x_n$  over  $\mathbb{F}$ . Let  $S = \mathbb{F}[x_1, \ldots, x_n]$ . In this paper  $\mathbb{F}$  will be a finite prime field  $\mathbb{F}_p$ .

In the following  $v_F \in \{0,1\}^n$  denotes the characteristic vector of a set  $F \subseteq [n]$ . For a family of subsets  $\mathcal{F} \subseteq 2^{[n]}$ , let

$$V(\mathcal{F}) = \{ v_F : F \in \mathcal{F} \} \subseteq \{0, 1\}^n \subseteq \mathbb{F}^n.$$

It is natural to consider the ideal  $I(V(\mathcal{F}))$ :

$$I(V(\mathcal{F})) := \{ f \in S : f(v) = 0 \text{ whenever } v \in V(\mathcal{F}) \}.$$

Denote by  $\mathbb{F}[x_1, \ldots, x_n]_{\leq s}$  the vector space of all polynomials over  $\mathbb{F}$  with degree at most s.

Let *I* be an ideal of the ring  $S = \mathbb{F}[x_1, \ldots, x_n]$ . Let  $h_{S/I}(m)$  denote the dimension over  $\mathbb{F}$  of the factor-space  $\mathbb{F}[x_1, \ldots, x_n]_{\leq m}/(I \cap \mathbb{F}[x_1, \ldots, x_n]_{\leq m})$  (see [8, Section 9.3]). The *Hilbert function* of the algebra S/I is the sequence  $h_{S/I}(0), h_{S/I}(1), \ldots$ 

It is easy to verify that in the special case when  $I = I(V(\mathcal{F}))$  for some set system  $\mathcal{F} \subseteq 2^{[n]}$ , the number  $h_{\mathcal{F}}(m) := h_{S/I}(m)$  is the dimension of the space of functions from  $V(\mathcal{F})$  to  $\mathbb{F}$  which can be represented as polynomials of degree at most m.

Let p be a prime and  $n > 1, 0 \le d \le n$  be integers. Let  $q = p^{\alpha}, \alpha \ge 1$ . Define the family of sets

$$\mathcal{F}(d,q) = \{ K \subseteq [n] : |K| \equiv d \pmod{q} \}.$$

I proved the following result in [11].

**Lemma 1.1** Let p be a prime and let  $f \in \mathbb{F}_p[x_1, \ldots, x_{4p}]$  be a polynomial. Suppose that  $f \in I(V\binom{[4p]}{2p})$  and that  $f \notin I(V\binom{[4p]}{3p})$ . Then  $deg(f) \geq p$ .

My proof used a combination of Gröbner basis methods and linear algebra. Srinivasan gave a simpler proof which combined Fermat's little Theorem with linear algebra (see [5]). Alon found a third proof based on the Combinatorial Nullstellensatz (see [3]).

P. Hrubeš, S. Natarajan Ramamoorthy, A. Rao and A. Yehudayoff proved a similar result to our Lemma 1.1. **Lemma 1.2** Let p be a prime and let  $f \in \mathbb{F}_p[x_1, \ldots, x_{2p}]$  be a polynomial. Suppose that  $f \in I(V\binom{[2p]}{p})$  and that  $f(\mathbf{0}) \neq 0$ . Then  $\deg(f) \geq p$ .

Let *m* be a positive integer and and *n* be an positive even integer. We say that a proper non-empty subsets  $S_1, \ldots, S_m$  are a balancing set of family if for every  $X \in {[n] \choose n/2}$  there is an index  $i \in [k]$  such that  $|S_i \cap X| = |S_i|/2$ .

Let n be an positive even integer. Let B(n) denote the minimum k for which a balancing set family of size k exists. P. Hrubeš, S. Natarajan Ramamoorthy, A. Rao and A. Yehudayoff applied Lemma 1.2 to give a lower bound for B(n) if n is an even positive integer. They used their lower bounds to prove lower bounds on depth-2 majority and threshold circuits that compute the majority and the weighted threshold functions.

We give here a simple generalization of Lemma 1.1 and Lemma 1.2. Our proof method is based on the following technical result.

**Theorem 1.3** Let  $\mathbb{F}$  be an arbitrary field. Let  $\mathcal{F} \subseteq \mathcal{G} \subseteq \mathbb{F}^n$  be affine subsets. Suppose that  $m \geq 0$  is an integer such that  $h_{S/I(\mathcal{F})}(m) = h_{S/I(\mathcal{G})}(m)$ . Then  $I(\mathcal{F})_{\leq m} = I(\mathcal{G})_{\leq m}$ .

**Remark.** We can easily prove a weaker version of Combinatorial Nullstellensatz using Theorem 1.3. Namely let  $T_i \subseteq \mathbb{F}$  be finite subsets and let  $\mathcal{G} := \prod_{i=1}^n T_i \subseteq \mathbb{F}^n$ , where  $t_i = |T_i| \geq 2$  are finite for each *i*. Let  $\mathbf{w} = (w_1, \ldots, w_n) \in \mathcal{G}$  be a fixed element. Define  $\mathcal{F} := \mathcal{G} \setminus \{\mathbf{w}\}$ . Let  $f \in I(\mathcal{F})$ be a polynomial such that  $\deg(f) \leq \sum_i t_i - n - 1$ . Let  $m := \sum_i t_i - n - 1$ . Then it is easy to check that  $h_{S/I(\mathcal{F})}(m) = h_{S/I(\mathcal{G})}(m) = (\prod_i t_i) - 1$ . Hence  $f \in I(\mathcal{G})$ .

One of our main result is the following generalization of Lemma 1.1 and Lemma 1.2.

**Theorem 1.4** Let p be a prime and  $q = p^{\alpha} > 1$ ,  $\alpha \ge 1$ . Let n > 0 be a positive integer and  $q - 1 \le d \le n - q + 1$  be an integer. Let  $\mathbb{F}$  be a field of characteristic p. Suppose that  $f \in I(V\binom{[n]}{d})$  and  $deg(f) \le q - 1$ . Then  $f \in I(V(\mathcal{F}(d,q)))$ .

We define now *L*-balancing families. Let n = 2d be an even number and  $L \subseteq [d-1]$  be a given subset. We say that  $\mathcal{F} \subseteq 2^{[n]}$  is an *L*-balancing family if for each  $F \in {[n] \choose d}$  there exists a  $G \subseteq [n]$  such that  $|F \cap G| \in L$ .

We prove the following general upper bound for the size of an L-balancing family. Our proof is based completely on Lemma 1.2.

**Theorem 1.5** Let p be a prime. Let n := 2p and  $L \subseteq [p-1]$  be a given subset. Define s := |L|. Let  $\mathcal{F} \subseteq 2^{[n]}$  be an L-balancing family. Then

$$m := |\mathcal{F}| \ge \frac{n}{2s}.$$

We prove our results in Section 2.

### 2 Proofs

**Proof of Theorem 1.3:** It follows from the definition of the Hilbert function that

$$h_{\mathcal{F}}(m) = dim(S_{\leq m}) - dim(I(\mathcal{F})_{\leq m})$$

and

$$h_{\mathcal{G}}(m) = \dim(S_{\leq m}) - \dim(I(\mathcal{G})_{\leq m}).$$

Since  $h_{\mathcal{F}}(m) = h_{\mathcal{G}}(m)$ , hence  $dim(I(\mathcal{F})_{\leq m}) = dim(I(\mathcal{G})_{\leq m})$ . But  $\mathcal{F} \subseteq \mathcal{G}$  implies that  $I(\mathcal{G})_{\leq m} \subseteq I(\mathcal{F})_{\leq m}$ , consequently  $I(\mathcal{F})_{\leq m} = I(\mathcal{G})_{\leq m}$ .

**Proof of Theorem 1.4:** We gave an alternative proof in [12] Corollary 3.1 using Gröbner basis theory for Wilson's theorem about the Hilbert function of complete uniform families.

**Theorem 2.1** (Wilson, [16]) Let  $0 \le d \le n$ ,  $0 \le m \le \min\{d, n - d\}$ , and  $\mathbb{F}$  be an arbitrary field. Then we have

$$h_{\binom{[n]}{d}}(m) = \binom{n}{m}.$$
(1)

We determined the Hilbert function of the set system  $\mathcal{F}(d,q)$  in [10] Corollary 4.5.

**Theorem 2.2** et p be a prime and  $q = p^{\alpha} > 1$ ,  $\alpha \ge 1$ . Let  $\mathbb{F}$  be a field of characteristic p.Let n > 0,  $0 \le d \le n$  be integers and define  $r = \min\{d, n - d\}$ . Let  $h_{\mathcal{F}(d,q)}(m)$  denote the Hilbert function of  $\mathbb{F}[\mathbf{x}]/I(V(\mathcal{F}(d,q)))$ . Then

$$h_{\mathcal{F}(d,q)}(m) = \sum_{i=0}^{\left\lfloor \frac{m}{q} \right\rfloor} {n \choose m-iq}$$

if  $0 \le m \le r$ , and

$$h_{\mathcal{F}(d,q)}(m) = \sum_{i=-\lfloor \frac{r}{q} \rfloor}^{\lfloor \frac{n-r}{q} \rfloor} \binom{n}{r+iq} - \sum_{i=1}^{\lfloor \frac{n-m}{q} \rfloor} \binom{n}{m+iq}$$

if m > r.

Let  $q-1 \leq d \leq n-q+1$  be an integer. Suppose that  $f \in I(V\binom{[n]}{d})_{\leq q-1}$ . It follows from Theorem 2.1 and Theorem 2.2 that  $h_{\binom{[n]}{d}}(q-1) = h_{\mathcal{F}(d,q)}(q-1) = \binom{n}{q-1}$ .

Hence Theorem 1.3 gives us that  $f \in I(V(\mathcal{F}(d,q)))_{\leq q-1}$ .

#### 

#### Proof of Theorem 1.5:

Let  $\mathcal{F} = \{F_1, \ldots, F_m\}$  be an *L*-balancing family and let  $\mathbf{v}_i := \mathbf{v}_{F_i}$  denote the characteristic vector of  $F_i$  for each  $1 \leq i \leq m$ .

Consider the polynomial

$$P(\mathbf{x}) := \prod_{i=1}^{m} \prod_{\ell \in L} (\mathbf{x} \cdot \mathbf{v}_i - \ell) \in \mathbb{F}_p[\mathbf{x}].$$

Here  $\cdot$  denotes the usual scalar product. Clearly deg $(P) \leq ms$ .

Then  $P(\mathbf{0}) = (\prod_{\ell \in L} \ell)^m \neq 0$ . On the other hand,  $P(\mathbf{v}_G) = 0$  for each  $G \in {\binom{[n]}{n}}$ , because  $\mathcal{F}$  is an L-balancing family.

Hence it follows from Lemma 1.2 that  $\deg(P) \ge p$  and we get that  $p \le ms$ .

### References

- [1] W. W. Adams, P. Loustaunau, An Introduction to Gröbner Bases, American Mathematical Society, 1994.
- [2] R.P. Anstee, L. Rónyai, A. Sali, Shattering news, Graphs and Combinatorics 18 (2002), 59–73.
- [3] N. Alon, Problems and results in Extremal Combinatorics–IV. arXiv preprint arXiv:2009.12692.(2020).

- [4] N. Alon, E. E. Bergmann, D. Coppersmith and A. M. Odlyzko, Balancing sets of vectors. *IEEE Trans. on Information Theory*, 34(1), 128-130 (1988).
- [5] N. Alon, Kumar, M., and B. L. Volk, (2020). Unbalancing sets and an almost quadratic lower bound for syntactically multilinear arithmetic circuits. *Combinatorica*, 40(2), 149-178.
- [6] T. Becker, V. Weispfenning, Gröbner bases a computational approach to commutative algebra, Springer-Verlag, Berlin, Heidelberg, 1993.
- [7] A. M. Cohen, H. Cuypers, H. Sterk (eds.), Some Tapas of Computer Algebra, Springer-Verlag, Berlin, Heidelberg, 1999.
- [8] D. Cox, J. Little, and D. O'Shea. Ideals, varieties, and algorithms. Springer-Verlag, Berlin, Heidelberg, 1992.
- [9] K. Friedl, L. Rónyai, Order-shattering and Wilson's theorem, to appear, *Discrete Mathematics*.
- [10] B. Felszeghy, G. Hegedüs, and L. Rónyai (2009). Algebraic properties of modulo q complete ℓ-wide families. Comb., Prob. and Computing, 18(3), 309-333.
- [11] G. Hegedüs (2010). Balancing sets of vectors. Studia Scientiarum Mathematicarum Hungarica, 47(3), 333-349.
- [12] G. Hegedüs, L. Rónyai, Gröbner bases for complete uniform families, J. of Algebraic Combinatorics 17 (2003), 171–180.
- [13] G. Hegedüs, L. Rónyai, Standard monomials for q-uniform and a conjecture of Babai and Frankl, *Central European Journal of Mathematics* 1 (2003), 198-207. http://www.cesj.com/mathematics.html
- [14] P. Hrubeš, S. Natarajan Ramamoorthy, A. Rao and A. Yehudayoff, Lower bounds on balancing sets and depth-2 threshold circuits. In 46th International Colloquium on Automata, Languages, and Programming (ICALP 2019). Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik. (2019).

- [15] S. Srinivasan, A robust version of Hegedüs's lemma, with applications. In Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing (pp. 1349-1362) (2020).
- [16] R.M. Wilson, A diagonal form for the incidence matrices of t-subsets vs. k subsets, Europ. J. Combin. 11(1990), 609–615.