

Improved Bounds for the Oriented Radius of Mixed Multigraphs

Jasine Babu, Deepu Benson, and Deepak Rajendraprasad
Indian Institute of Technology Palakkad

Abstract

A *mixed multigraph* is a multigraph which may contain both undirected and directed edges. An *orientation* of a mixed multigraph G is an assignment of exactly one direction to each undirected edge of G . A mixed multigraph G can be oriented to a strongly connected digraph if and only if G is bridgeless and strongly connected [Boesch and Tindell, Am. Math. Mon., 1980]. For each $r \in \mathbb{N}$, let $f(r)$ denote the smallest number such that any strongly connected bridgeless mixed multigraph with radius r can be oriented to a digraph of radius at most $f(r)$. We improve the current best upper bound of $4r^2 + 4r$ on $f(r)$ [Chung, Garey and Tarjan, Networks, 1985] to $1.5r^2 + r + 1$. Our upper bound is tight upto a multiplicative factor of 1.5 since, $\forall r \in \mathbb{N}$, there exists an undirected bridgeless graph of radius r such that every orientation of it has radius at least $r^2 + r$ [Chvátal and Thomassen, J. Comb. Theory. Ser. B., 1978]. We prove a marginally better lower bound, $f(r) \geq r^2 + 3r + 1$, for mixed multigraphs. While this marginal improvement does not help with asymptotic estimates, it clears a natural suspicion that, like undirected graphs, $f(r)$ may be equal to $r^2 + r$ even for mixed multigraphs. En route, we show that if each edge of G lies in a cycle of length at most η , then the oriented radius of G is at most $1.5r\eta$. All our proofs are constructive and lend themselves to polynomial time algorithms.

Keywords— Mixed multigraph, Oriented radius, Strong orientation

1 Introduction

A *mixed multigraph* (*mixed graph* for short) is a multigraph in which each edge could be either directed or undirected. A *walk* from a vertex v_1 to a vertex v_k in a mixed graph G is a sequence of vertices v_1, v_2, \dots, v_k such that $\forall i \in [k - 1]$, G contains either an undirected edge $\{v_i, v_{i+1}\}$ or an edge directed from v_i to v_{i+1} . A *trail* is a walk without repeated edges. A *path* is a trail without repeated vertices, except possibly $v_1 = v_k$. A mixed graph is *strongly connected* (*connected* for short) if there is a path from each vertex to every other vertex. Notice that this is a common generalisation of connectivity of undirected graphs and strong connectivity of digraphs. A *bridge* in a connected mixed graph G is an undirected edge whose removal disconnects G . A bridgeless connected mixed graph is called *2-edge connected*. An *orientation* of a mixed graph G is an assignment of exactly one direction to each undirected edge of G . A mixed graph G is called *strongly orientable* if it can be oriented to a strongly connected digraph.

Email addresses: jasine@iitpkd.ac.in (Jasine Babu), bensondeepu@gmail.com (Deepu Benson), deepak@iitpkd.ac.in (Deepak Rajendraprasad)




Figure 1: The undirected graph G_3 from [3]. While G_3 has radius 3, every orientation of G_3 has radius at least 12.

The first major study about strongly orientable graphs was by H. E. Robbins in 1939 [1], where he proved that an undirected graph is strongly orientable if and only if it is 2-edge connected. Robbins motivated the study using the real-world scenario of converting a network of two-way streets to one-way streets without sacrificing reachability from any point to another. In the “real-world” it is very well possible that some, but not all, of the streets in the network are already one-way. Hence the same scenario motivates the study of strong orientability of mixed graphs. In 1980, Boesch and Tindell [2] extended Robbins’ theorem to mixed graphs. They proved that a mixed graph is strongly orientable if and only if it is 2-edge connected. Neither of these works approached the problem with a view to contain the distance blow up that may result from the orientation. This was taken up by Chvátal and Thomassen in 1978 for undirected graphs [3], and Chung, Garey and Tarjan in 1985 for mixed graphs [4]. We need some more terminology before stating their results.

Let G be an undirected, directed or mixed graph. For a vertex u in G , $N[u]$ denotes the set consisting of u and all its in-neighbours, out-neighbours and undirected neighbours. The *length* of a path in G is the number of edges (directed and undirected) in the path. The *distance* $d_G(u, v)$ from a vertex u to a vertex v in G is the length of a shortest path from u to v in G . The *out-eccentricity* $e_{out}(u)$ of u is $\max\{d_G(u, v) \mid v \in V(G)\}$ and *in-eccentricity* $e_{in}(u)$ is $\max\{d_G(v, u) \mid v \in V(G)\}$. The *eccentricity* $e(u)$ of u is $\max\{e_{out}(u), e_{in}(u)\}$. The *radius* of G is $\min\{e(u) \mid u \in V(G)\}$ and *diameter* is $\max\{e(u) \mid u \in V(G)\}$. A vertex u of G is a central vertex of G if $e(u)$ is equal to the radius of G . The radius and diameter of a disconnected graph are taken to be infinite. The *oriented radius (diameter)* of G is the minimum radius (diameter) of an orientation of G . For each $r \in \mathbb{N}$, let $f(r)$ (resp., $\bar{f}(r)$) denote the smallest number such that any 2-edge connected mixed graph (resp., undirected graph) with radius r can be oriented to a strongly connected digraph of radius at most $f(r)$ (resp., $\bar{f}(r)$). Since every 2-edge connected undirected graph is also a 2-edge connected mixed graph, $\bar{f} \leq f$.

Chvátal and Thomassen [3] showed that $\bar{f}(r) = r^2 + r$. The upper bound was established by exploiting the properties of a carefully chosen ear-decomposition. For the lower bound, they constructed a family of 2-edge connected graphs G_r of radius r and oriented radius $r^2 + r$ (see Figure 1 for G_3). Extending the work of Boesch and Tindell on mixed graphs, Chung, Garey and Tarjan [4] in 1985 proposed a linear-time algorithm to check whether a mixed graph has a strong orientation. The resultant orientation could have radius as high as $O(n)$, where n is the number of vertices of the mixed graph. In the same paper, they extend the work of Chvátal and Thomassen on undirected graphs to design a polynomial time algorithm which provides a strong orientation for a 2-edge connected mixed graph of radius r with oriented radius at most $4r^2 + 4r$. Hence $r^2 + r \leq f(r) \leq 4r^2 + 4r$.

Our results. In this article, we establish that $r^2 + 3r - 1 \leq f(r) \leq 1.5r^2 + r + 1$. We consider the (8/3)-factor improvement in the upper bound (over that of Chung, Garey and

Tarjan) as the major contribution in this work. The proof is constructive and is presented as an algorithm which is easily seen to be polynomial-time (Section 2). In comparison, the improvement to the lower bound (over the optimal lower bound for undirected graphs due to Chvátal and Thomassen) is marginal and does not help improve asymptotic estimates. But it clears a natural suspicion that $f = \bar{f}$. The proof is via a construction of an infinite family of mixed multigraphs parameterised by r (Section 4). Since the diameter d of a mixed graph and its radius r are tied together by $r \leq d \leq 2r$, we can infer that any 2-edge connected mixed multigraph of diameter d has oriented diameter at most $3d^2 + 2d + 2$. In Section 3, we illustrate that our algorithm provides a template to handle interesting special cases where there is a bound on the unavoidable penalty you pay for orienting an undirected edge, namely the length of the shortest cycle containing that edge. In particular, we show that if every edge of G lies in a cycle of length at most η , then our algorithm gives an orientation with radius at most $1.5r\eta$.

The gap between our lower and upper bounds on $f(r)$ leaves room for a conjecture on the exact behaviour of $f(r)$. We are inclined to believe that the upper bound has further room for improvement and show that our algorithm does not perform optimally on the family of graphs constructed to establish our lower bound. We conjecture that $f(r) = r^2 + O(r)$.

Further Literature

Not surprisingly, undirected graphs have overshadowed mixed graphs in this line of research. Moreover, oriented diameter has received more attention than oriented radius. For each $d \in \mathbb{N}$, let $\bar{g}(d)$ denote the smallest number such that any 2-edge connected undirected graph of diameter d has an orientation with diameter at most $\bar{g}(d)$. Chvátal and Thomassen [3] has shown that $\frac{1}{2}d^2 + d \leq \bar{g}(d) \leq 2d^2 + 2d$ for all $d \geq 2$. Since the lower and upper bounds for \bar{g} above differ by a factor of 4, there have been several attempts to bridge this gap. In the same paper, Chvátal and Thomassen showed that $\bar{g}(2) = 6$. Kwok, Liu and West [5] proved that $9 \leq \bar{g}(3) \leq 11$. Together with Vaka, we showed that $\bar{g}(4) \leq 21$ [6]. The major effort in the same paper for us was to show that $\bar{g}(d) \leq 1.373d^2 + 6.971d - 1$.

Since \bar{g} is quadratic, there have been several studies on special graph classes restricted to which \bar{g} is linear. AT-free graphs [7] and chordal graphs [8] are popular such classes. Huang, Li, Li, and Sun [9] proved that the oriented radius of an undirected 2-edge connected graph G is at most $r(\eta - 1)$. In this paper we also extend their approach to mixed graphs.

Bounds (mostly upper) on oriented diameter of 2-edge connected graphs in terms of other invariants like domination number [7], minimum degree [10], [11], and maximum degree [12] are also known. A final result of Chvátal and Thomassen in their 1978 paper was that it is NP-hard to decide whether a given undirected graph has oriented diameter 2. This has prompted a search for polynomial time algorithms for the problem on special graph classes. Eggemann and Noble [13] studied about the oriented diameter of planar graphs. For each constant l , they have proposed an algorithm that decides if a planar graph $G = (V, E)$ has an orientation with diameter at most l and runs in time $O(c|V|)$, where the constant c depends on l .

Orientations of mixed graphs with objectives other than minimising the radius and diameter have been studied. Aamand, Hjuler, Holm, and Rotenberg [14] studied the necessary and sufficient conditions for the existence of a strong orientation for an undirected or a mixed graph when the edges of that graph are partitioned into trails. They also provided a polynomial time algorithm for finding a strong trail orientation, if one exists, for undirected and mixed graphs. Blind [15] studied k -arc-connected orientations in undirected and mixed graphs.

2 Upper Bound

In this section, we show that $f(r) \leq 1.5r^2 + r + 1$ by designing and analysing an orientation algorithm STRONGORIENTATION that takes a mixed graph G of radius r as input and outputs

an orientation of G of radius at most $1.5r^2 + r + 1$. This algorithm involves repeated applications of two symmetric orientation algorithms, Algorithm ORIENTOUT and Algorithm ORIENTIN. To decide the order of execution of these two algorithms, we need to partition the sequence of first r positive odd numbers into two nearly equal parts. The following lemma will help us there.

Lemma 1. *For every n , the set of first n odd numbers $S = \{2i - 1 \mid 1 \leq i \leq n\}$ can be partitioned into two sets A and B such that*

$$|\Sigma_{a \in A} a - \Sigma_{b \in B} b| \leq \begin{cases} 2, & n = 2 \\ 1, & n \neq 2 \end{cases}$$

Moreover, this partition can be generated in polynomial time.

The proof of Lemma 1 is an interesting exercise. For the sake of completeness, we include it in the appendix.

We also use the following lemma during the design of our orientation algorithms. This is a generalization of a lemma by Chung, Garey and Tarjan [4, Lemma 5]. The statement of the original lemma and the proof of Lemma 2 are given in the appendix.

Lemma 2. *Let G be a strongly connected bridgeless mixed multigraph of radius r . Then any edge pq of G is on a cycle of length at most $2r + 1$.*

Let G be a strongly connected bridgeless mixed graph of radius r with a central vertex u . We first describe Algorithm ORIENTOUT, in detail, which identifies a subgraph H of G such that $V(H)$ is an $(r - 1)$ -step dominating set of $V(G)$ and an orientation \vec{H} of H such that $\forall v \in V(\vec{H})$, $d_{\vec{H}}(u, v) \leq 2r$ and $d_{\vec{H}}(v, u) \leq 4r - 1$. Using this algorithm alone, we can show that $f(r) \leq 2r^2 + r$ using an induction on r , as discussed below.

When $r = 0$, the graph G is a single isolated vertex and the assumption holds true trivially. Let us assume the assumption holds true for graphs of radius $r - 1$. Let us contract the subgraph H into a single vertex v_H to obtain a graph G' . We can see that G' has radius at most $r - 1$. Thus by the induction hypothesis, G' has an orientation with radius at most $2(r - 1)^2 + (r - 1) = 2r^2 - 3r + 1$. Notice that, G' and H do not share any edges. Thus by combining the orientations of G' and H , we can get an orientation \vec{G} of G with radius at most $(2r^2 - 3r + 1) + (4r - 1) = 2r^2 + r$. Hence, by induction, the assumption is true for all values of r .

The description of the 4-stage algorithm ORIENTOUT is given below.

Algorithm ORIENTOUT

Input: A strongly connected bridgeless mixed multigraph G and a vertex $u \in V(G)$ with eccentricity at most r .

Output: An orientation \vec{H} of a subgraph H of G such that $N[u] \subseteq V(\vec{H})$ and $\forall v \in V(\vec{H})$, $d_{\vec{H}}(u, v) \leq 2r$ and $d_{\vec{H}}(v, u) \leq 4r - 1$.

We create \vec{H} in 4 stages. The vertices newly added to \vec{H} in a stage will be referred to as the vertices *captured* in that stage.

Stage 0. Let v be a vertex having multiple edges incident with u . If possible, these edges are oriented in such a way that all the uv edges are part of a directed 2-cycle. Notice that this does not increase any pairwise distances in G . We can also remove multiple oriented edges in the same direction between any pair of vertices without affecting any distance. Hence the only multi-edges left in G are those which form a directed 2-cycle.

Let X denote the set of all vertices with at least one edge incident with the vertex u . X is partitioned into X_{in} , X_{out} and X_{un} . A vertex $v \in X$ is said to be in X_{in} if it has at least one

directed edge towards u . A vertex $v \in X \setminus X_{in}$ is said to be in X_{out} if it has at least one directed edge from u . $X_{un} = X \setminus (X_{in} \cup X_{out})$. Notice that a vertex $v \in X_{un}$ has exactly one undirected edge incident with u . We initialize $X_{conf} = \emptyset$ (we will later identify this as the set of conflicted vertices in X). For each $v \in X$, let $l(v)$ denote the length of a shortest cycle containing an edge between u and v . By Lemma 2, $l(v) \leq 2r + 1$. Let $s = \sum_{v \in X} l(v)$.

Stage 1. Let $G = G_0$. We will orient some of the unoriented edges of G_0 incident with u in this stage to obtain a mixed graph G_1 as follows. Let v_1, v_2, \dots, v_k be an arbitrary ordering of vertices in X such that all the vertices in X_{in} comes before any vertex in X_{out} and all the vertices in X_{out} comes before any vertex in X_{un} . Let $c = |X_{in} \cup X_{out}|$.

Repeat Steps (i) to (iii) for $i = c + 1$ to $|X|$:

- i. An edge uv_i is oriented from v_i to u if the parameter s remains the same even after such an orientation. The vertex v_i is added to X_{in} in this case.
- ii. Otherwise, the edge uv_i is oriented from u to v_i if the parameter s remains the same even after such an orientation. The vertex v_i is added to X_{out} in this case.
- iii. Otherwise, we leave the edges uv_i unoriented. Such an edge uv_i is called a *conflicted edge* and the vertex v_i is called a *conflicted vertex*. The vertex v_i is added to X_{conf} in this case.

Observation 1. If an edge uv_i is conflicted then there exists an edge $\overrightarrow{v_j u}$, at the time of processing v_i , where $v_j \in X_{in}$ and $j < i$ such that the edge uv_i is a part of every shortest cycle containing the edge $\overrightarrow{v_j u}$. Otherwise the parameter s would have remained the same even if the edge uv_i gets oriented from u to v_i . Hence, for every vertex $v \in X_{conf}$, uv is an undirected edge and there exists a $w \in X_{in}$ such that every shortest path from u to w starts with the edge uv .

We have the following distance bounds in G_1 : $\forall x \in X_{in}$, $d_{G_1}(u, x) \leq 2r$ and $\forall y \in X_{out}$, $d_{G_1}(y, u) \leq 2r$. These bounds follow from the fact that the edges are oriented in Stage 1 only if the parameter s remains unchanged.

Stage 2. In this stage, we try to capture all the vertices in X_{in} . Let T_{out} be a minimal tree formed by a breadth first search in G_1 , starting from u and including at least one path from u to v for each $v \in X_{in}$. Notice that T_{out} has height at most $2r$ and every leaf of T_{out} is a vertex in X_{in} . Orient the tree T_{out} outward from u in G_1 to obtain G_2 . Let $S_1 = V(T_{out})$. Every vertex in S_1 is part of a directed cycle in G_2 containing u and of length at most $2r + 1$. Hence for any vertex $x \in S_1$, we have $d_{G_2}(u, x) \leq 2r$ and $d_{G_2}(x, u) \leq 2r$. Notice that, by Observation 1, any edge $e = uv$, where $v \in X_{conf}$, will get oriented in Stage 2. Hence $(X_{in} \cup X_{conf}) \subseteq S_1$.

Stage 3. Let $X'_{out} = X_{out} \setminus S_1$. In this stage, we try to capture all the vertices in X'_{out} . Let G_3 be a graph obtained from G_2 by contracting $G[S_1]$ into a single vertex u^* . Every arc \vec{uv} , $v \in X'_{out}$, was part of a cycle of length at most $2r + 1$ in G_1 . Hence v has a path of length at most $2r - 1$ to some vertex in S_1 . This path together with the edge u^*v ensures that every vertex v in X'_{out} lies in a cycle of length at most $2r$ containing u^* in G_3 . Let T_{in} be a minimal tree formed by a reverse breadth first search in G_3 starting from u^* such that it contains a path from each $v \in X'_{out}$ to u^* . Notice that, T_{in} has height at most $2r - 1$ and every leaf of T_{in} is a vertex in X'_{out} . The reduction of 1 in height is because $(X_{in} \cup X_{conf}) \subseteq S_1$. Orient the tree T_{in} inward towards u^* in G_3 to obtain G_4 . Let $S_2 = V(T_{in}) \setminus \{u^*\}$. Then for any vertex $y \in S_2$, we have the following bounds, $d_{G_4}(u^*, y) \leq 2r - 1$ and $d_{G_4}(y, u^*) \leq 2r - 1$. Let $H = G[S_1 \cup S_2]$. Notice that, T_{out} and T_{in} do not share any edges in H . The edges of H are oriented consistently with the orientation of T_{out} and T_{in} to obtain \vec{H} . We have already proved that for any vertex $x \in S_1$, $d_{G_2}(u, x) \leq 2r$ and $d_{G_2}(x, u) \leq 2r$. Hence $\forall x \in S_1$, $d_{\vec{H}}(u, x) \leq 2r$ and $d_{\vec{H}}(x, u) \leq 2r$. Consider a vertex $y \in S_2$. We have proved that $d_{G_4}(y, u^*) \leq 2r - 1$. This

implies $d_{\vec{H}}(y, u) \leq (2r - 1) + (2r) = 4r - 1$. Since height of T_{in} is at most $2r - 1$, y has a directed path of length at most $2r - 2$ from a vertex $v \in X'_{out}$ in G_4 . This path along with the arc \vec{uv} will give us the additional bound $d_{\vec{H}}(u, y) \leq 2r$ for all $y \in S_2$. Thus \vec{H} has radius at most $4r - 1$ with $d_{\vec{H}}(u, x) \leq 2r$ and $d_{\vec{H}}(x, u) \leq 4r - 1, \forall x \in V(\vec{H})$.

As mentioned earlier, using algorithm ORIENTOUT and an induction on r , we can get the upper bound $f(r) \leq 2r^2 + r$. This itself is a 2-factor improvement over the existing bound of $4r^2 + 4r$ obtained by Chung, Garey and Tarjan [4]. Notice that, in \vec{H} , the bound on $e_{out}(u)$ is better than that of $e_{in}(u)$ by a factor of 2. To exploit this gap, we design a partner algorithm ORIENTIN using a symmetric construction.

Algorithm ORIENTIN

Input: A strongly connected bridgeless mixed multigraph G and a vertex $u \in V(G)$ with eccentricity at most r .

Output: An orientation \vec{H} of a subgraph H of G such that $N[u] \subseteq V(\vec{H})$ and $\forall v \in V(\vec{H}), d_{\vec{H}}(u, v) \leq 4r - 1$ and $d_{\vec{H}}(v, u) \leq 2r$.

Let G be a strongly connected bridgeless mixed graph of radius r and u be a central vertex of G . To show that $f(r) \leq 1.5r^2 + r + 1$, we design an r -phase algorithm in which each phase involves an application of either ORIENTOUT or ORIENTIN. For convenience, we number the phases from r down to 1. A graph G_i and a vertex $u_i \in V(G_i)$ with eccentricity at most i are the inputs to the algorithm ORIENTOUT or ORIENTIN executed in Phase i of algorithm STRONGORIENTATION. \vec{H}_i is the output of the algorithm executed in Phase i .

Algorithm STRONGORIENTATION

Input: A strongly connected bridgeless mixed multigraph G and a vertex $u \in V(G)$ of eccentricity r .

Output: An orientation \vec{G} , of G , with radius at most $1.5r^2 + r + 1$.

- 1: $S \leftarrow \{2i - 1 \mid 1 \leq i \leq r\}$.
- 2: Obtain a partition (A, B) of S such that $|\sum_{a \in A} a - \sum_{b \in B} b| \leq 2$.
- 3: $I_A \leftarrow \{i \mid (2i - 1) \in A\}, I_B \leftarrow \{i \mid (2i - 1) \in B\}$.
- 4: $i \leftarrow r, G_i \leftarrow G, u_i \leftarrow u$.
- 5: **repeat**
- 6: **if** $i \in I_A$ **then**
- 7: $\vec{H}_i \leftarrow \text{ORIENTOUT}(G_i, u_i)$.
- 8: **else**
- 9: $\vec{H}_i \leftarrow \text{ORIENTIN}(G_i, u_i)$.
- 10: **end if**
- 11: Obtain G_{i-1} by contracting $V(\vec{H}_i)$ to a single vertex u_{i-1} .
- 12: $i \leftarrow i - 1$.
- 13: **until** $i = 0$.
- 14: Take the union of the orientations $\{\vec{H}_1, \dots, \vec{H}_r\}$ and orient the remaining edges arbitrarily to obtain an orientation \vec{G} of G .

The existence of the partition used in Step 2 of the algorithm is guaranteed by Lemma 1 and can be computed in polynomial time. Since $N[u_i] \subseteq V(\vec{H}_i)$, after each phase, the radius of the resultant graph reduces by at least 1. Hence, the vertex u_{i-1} of G_{i-1} has eccentricity at most $i - 1$. Further notice that for $1 \leq i \leq r$, the graphs \vec{H}_i are pair-wise edge disjoint and hence their union gives a consistent orientation \vec{G} of G . For $1 \leq i \leq r$, let $e_{in}^i = \max\{d_{\vec{H}_i}(v, u_i) \mid v \in V(\vec{H}_i)\}$

denote the in-eccentricity of u_i in \vec{H}_i and $e_{out}^i = \max\{d_{\vec{H}_i}(u_i, v) \mid v \in V(\vec{H}_i)\}$ denote the out-eccentricity of u_i in \vec{H}_i . By combining the guarantees given by the two algorithms ORIENTOUT and ORIENTIN, for the in-distance and out-distance in \vec{H}_i , we have the following bounds for $e_{out}(u)$ and $e_{in}(u)$ of \vec{G} .

- $e_{out}(u) \leq \sum_{i=1}^r e_{out}^i = \sum_{i \in I_A} (2i) + \sum_{i \in I_B} (4i - 1) = \sum_{i=1}^r (2i) + \sum_{i \in I_B} (2i - 1)$.
- $e_{in}(u) \leq \sum_{i=1}^r e_{in}^i = \sum_{i \in I_A} (4i - 1) + \sum_{i \in I_B} (2i) = \sum_{i=1}^r (2i) + \sum_{i \in I_A} (2i - 1)$.

The radius of \vec{G} is at most $\max\{e_{out}(u), e_{in}(u)\} \leq \sum_{i=1}^r (2i) + \max\{\sum_{i \in I_A} (2i - 1), \sum_{i \in I_B} (2i - 1)\}$. Notice that $\sum_{i=1}^r (2i - 1) = r^2$ and by Lemma 1, we get $\max\{\sum_{i \in I_A} (2i - 1), \sum_{i \in I_B} (2i - 1)\}$ is at most $\frac{r^2}{2} + 1$. Thus the radius of \vec{G} is at most $\sum_{i=1}^r (2i) + (\frac{r^2}{2} + 1) \leq (r^2 + r) + (\frac{r^2}{2} + 1) = 1.5r^2 + r + 1$. Hence, the oriented radius of G is at most $1.5r^2 + r + 1$.

This proves the main theorem of the section.

Theorem 3. $f(r) \leq 1.5r^2 + r + 1$.

It is easy to verify that Algorithm STRONGORIENTATION runs in polynomial time. Hence we have a polynomial-time algorithm to find a strong orientation \vec{G}_r with radius at most $1.5r^2 + r + 1$ for a mixed graph G_r of radius r . Since $r \leq d \leq 2r$, we can immediately see that a mixed graph of diameter d has an orientation of diameter at most $3d^2 + 2d^2 + 2$ for all d .

3 Upper Bound for Some Special Classes of Mixed Multigraphs

Let G be a strongly connected mixed graph with radius r . Let $\eta(G)$ denote the smallest integer k such that every edge of G belongs to a cycle of length at most k . Huang, Li, Li, and Sun [9] proved that the oriented radius of an undirected graph G of radius r is at most $r(\eta(G) - 1)$. We generalize this result to mixed graphs. We start by generalizing Lemma 1 as follows.

Lemma 4. *Let set $S_1 = \{2i - 1 \mid 1 \leq i \leq n\}$. Let S_2 be a multiset with k elements of the same value: either $2n - 1$ or $2n$. Then the multiset $S = S_1 \cup S_2$ can be partitioned into two multisets A and B such that $|\sum_{a \in A} a - \sum_{b \in B} b| \leq 2$. Moreover, this partition can be generated in polynomial time.*

The proof of this lemma is simple and is given in the appendix.

Let G be a strongly connected bridgeless mixed graph with a vertex $u \in V(G)$ having eccentricity at most r and $\eta(G) = \eta$. By Lemma 2, $\eta \leq 2r + 1$. We use the same terminology of G_i , u_i , and \vec{H}_i as used in Algorithm STRONGORIENTATION. Let η_{u_i} denote the smallest integer k such that every edge incident with u_i belongs to a cycle of length at most k in G_i . Since u_i has eccentricity at most i , by Lemma 2, it is easy to see that, $\eta_{u_i} \leq \min\{\eta, 2i + 1\}$. Let $k_i = \min\{\eta, 2i + 1\}$ denote this upper bound. From the details of algorithms ORIENTIN and ORIENTOUT, one can see that when G_i and u_i are given as input, the distance guarantees of these algorithms only use the value of $\eta_{u_i} \leq k_i$. Now, a second look at algorithm ORIENTIN, with the definition of η_{u_i} in mind, will show that the oriented subgraph \vec{H}_i returned by ORIENTIN(G_i, u_i) has $e_{in}(u_i)$ at most $(k_i - 1)$ and $e_{out}(u_i)$ at most $(2k_i - 3)$. Similarly, \vec{H}_i returned by ORIENTOUT(G_i, u_i) has $e_{out}(u_i)$ at most $(k_i - 1)$ and $e_{in}(u_i)$ at most $(2k_i - 3)$.

A strong orientation \vec{G} of G is obtained by an r -phase algorithm similar to STRONGORIENTATION, with each phase involving an execution of either ORIENTIN or ORIENTOUT. Let $m = \lfloor (\eta - 1)/2 \rfloor$. Recall that $2i + 1$ dominates η for $i > m$. For $1 \leq i \leq r$, let $e_{in}^i = \max\{d_{\vec{H}_i}(vu_i) \mid v \in V(\vec{H}_i)\}$ denote the in-eccentricity of u_i in \vec{H}_i and $e_{out}^i = \max\{d_{\vec{H}_i}(u_i v) \mid v \in V(\vec{H}_i)\}$ denote the out-eccentricity of u_i in \vec{H}_i . Let S be the multiset $\{k_i - 2 \mid 1 \leq i \leq r\}$. It can be verified

that S satisfies the requirements of Lemma 4. Hence, we can obtain a partition (A, B) of S such that their sums differ at most by 2. Let $\sigma = (\sigma_1, \dots, \sigma_r)$ be a non-decreasing ordering of elements in S . For $1 \leq i \leq r$, let $i \in I_A$ if and only if $\sigma_i \in A$ and $i \in I_B$ otherwise. In Phase i , we apply ORIENTOUT if $i \in I_A$ and ORIENTIN if $i \in I_B$. By combining the guarantees given by ORIENTOUT and ORIENTIN for the in-distance and out-distance in \vec{H}_i , we can have the following bounds for $e_{in}(u)$ and $e_{out}(u)$ of \vec{G} .

$$\begin{aligned} e_{in}(u) &\leq \sum_{i=1}^r e_{in}^i \\ &\leq \sum_{i \in I_A} (2k_i - 3) + \sum_{i \in I_B} (k_i - 1) \\ &= \sum_{i=1}^r (k_i - 1) + \sum_{i \in I_A} (k_i - 2) \\ &=: k_{in} \end{aligned}$$

$$\begin{aligned} e_{out}(u) &\leq \sum_{i=1}^r e_{out}^i \\ &\leq \sum_{i \in I_A} (k_i - 1) + \sum_{i \in I_B} (2k_i - 3) \\ &= \sum_{i=1}^r (k_i - 1) + \sum_{i \in I_B} (k_i - 2) \\ &=: k_{out} \end{aligned}$$

Further, we have the following bound for $|k_{in} - k_{out}|$ and $(k_{in} + k_{out})$.

$$\begin{aligned} |k_{in} - k_{out}| &= |\sum_{i \in I_A} (k_i - 2) - \sum_{i \in I_B} (k_i - 2)| \\ &\leq 2 \quad (\text{Lemma 4}) \end{aligned} \tag{1}$$

$$\begin{aligned} (k_{in} + k_{out}) &= 2\sum_{i=1}^r (k_i - 1) + \sum_{i=1}^r (k_i - 2) \\ &= (3\sum_{i=1}^r k_i) - 4r \\ &= 3 \left[(\sum_{i=1}^m 2i + 1) + (\sum_{i=m+1}^r \eta) \right] - 4r \quad (\text{By definition of } k_i) \\ &= 3 \left[(\sum_{i=1}^r \eta) - \sum_{i=1}^m (\eta - (2i + 1)) \right] - 4r \\ &= 3r\eta - 3\sum_{i=1}^m (\eta - (2i + 1)) - 4r \end{aligned} \tag{2}$$

Notice that, $\eta \in \{2m + 1, 2m + 2\}$. Now consider the term $\sum_{i=1}^m (\eta - (2i + 1))$ in Equation 2. Notice that, when η is even, this term sums up odd numbers from 1 to $\eta - 3$ and when η is odd, the term sums up even numbers from 0 to $\eta - 3$. Hence, by taking the worst case for Equation 2, we get.

$$(k_{in} + k_{out}) \leq 3r\eta - \frac{3}{4}(\eta - 1)(\eta - 3) - 4r \tag{3}$$

From equations 1 and 3, we get.

$$\begin{aligned} \max\{e_{in}(u), e_{out}(u)\} &\leq \max\{k_{in}, k_{out}\} \\ &\leq \left\lfloor \frac{k_{in} + k_{out}}{2} \right\rfloor + 1 \\ &\leq 1.5r\eta - 0.375(\eta - 1)(\eta - 3) - 2r + 1 \end{aligned} \tag{4}$$

Since $\eta \geq 3$, $(\eta - 1)(\eta - 3)$ is non-negative. Hence, by Equation 4, we have the following theorem.

Theorem 5. *Let G be a strongly connected bridgeless mixed multigraph of radius r and let $\eta(G) = \eta$ denote the smallest integer k such that every edge of G belongs to a cycle of length at most k . Then G has oriented radius at most $1.5r\eta - 0.375(\eta - 1)(\eta - 3) - 2r + 1 \leq 1.5r\eta$.*

A k -chordal mixed multigraph is a mixed multigraph graph in which all cycles of length more than k have a chord. In particular when $k = 3$, the graph is called a chordal mixed multigraph. Since $\eta \leq k$, by Theorem 5, we have the following corollary.

Corollary 6. *Let G be a strongly connected bridgeless k -chordal mixed multigraph with radius r . Then G has oriented radius at most $1.5rk - 0.375(k - 1)(k - 3) - 2r + 1$. In particular, if G is a strongly connected bridgeless chordal mixed multigraph of radius r then G has oriented radius at most $2.5r + 1$.*

4 Lower Bound




Figure 2: Mixed multigraph G_3 of radius 3 and oriented radius 17.

When we restrict to undirected graphs, the currently known lower bound for $f(r)$ is $r^2 + r$. This follows from a construction by Chvátal and Thomassen [3], an instance of which is shown in Figure 1. Using a similar idea, we construct a mixed graph G_r of radius r whose oriented radius is $r^2 + 3r - 1$. In order to construct G_r , we first define a ternary tree T_r of height r , rooted at a vertex a . Each internal vertex v of T_r has 3 children, called the *left*, *middle*, and *right* child of v . The *Level* of a vertex is its distance from a in T_r . For a vertex v in Level i , for $0 \leq i \leq r - 2$, the middle child of v is a leaf in T_r and the other two children are internal vertices. Let b, c , and d be the left, middle, and right children, respectively, of a . Let f be the leaf in Level r of T_r such that every vertex in the $a - f$ path other than a is a left child of its parent. Let g be the leaf in Level r of T_r such that every vertex in the $a - g$ path other than a is a right child of its parent. We construct a mixed graph H from T_r as follows. For each Level i , for $i = 0$ to $r - 1$ and for each internal vertex p_i in Level i , with $l_{i+1}, m_{i+1}, r_{i+1}$, respectively, being the left, right, and middle children of p_i , we do the following. Add paths of length $2r - 2i - 1$ from l_{i+1} to m_{i+1} and m_{i+1} to r_{i+1} . The middle edge of the first path is oriented along the direction from l_{i+1} to m_{i+1} and the middle edge of the second path is oriented along the direction from m_{i+1} to r_{i+1} . Notice that, due to the presence of these directed edges, in any strong orientation of H , directions of all the edges except the edges between the middle children and their parents are forced. But we leave them unoriented for now. Now, the mixed graph G_r is constructed by taking two copies of H , H_1 and H_2 and identifying the root a of H_1 and H_2 . The vertices of H_1 retain their labels from H . We can easily see that G_r is strongly connected with radius r and a is the unique central vertex. Figure 2 shows the mixed multigraph G_3 .

It can be verified that in any strong orientation \vec{G}_r that minimizes the radius, a is the unique central vertex. Hence, there will be an optimum orientation in which the orientation of H_1 is symmetric with the orientation of H_2 . We have already seen that edges from a middle child

to its parent are the only edges in G_r whose orientation is not forced in a strong orientation. Hence, in any strong orientation \vec{G}_r of G_r with an induced orientation \vec{H}_1 of H_1 , we have the following bounds for every internal vertex p_i of T_r at Level i with l_{i+1} and r_{i+1} as the left and right children, respectively, for $0 \leq i \leq r-1$.

$$\begin{aligned}
d_{\vec{H}_1}(l_{i+1}, p_i) &\in \{2(r-i), 4(r-i)-1\} \\
d_{\vec{H}_1}(p_i, r_{i+1}) &\in \{2(r-i), 4(r-i)-1\} \\
d_{\vec{H}_1}(p_i, l_{i+1}) &= 1 \\
d_{\vec{H}_1}(r_{i+1}, p_i) &= 1
\end{aligned} \tag{5}$$

From the first line of Equation 5, it follows that, $d_{\vec{H}_1}(f, b) \geq \sum_{i=1}^{r-1} 2(r-i) = r^2 - r$. Similarly, from the second line of Equation 5, it follows that, $d_{\vec{H}_1}(d, g) \geq \sum_{i=1}^{r-1} 2(r-i) = r^2 - r$. There are only two possible orientations for the edge ac . If the edge ac in H_1 is oriented from a to c , then $d_{\vec{H}_1}(b, a) = 4r - 1$. Hence $d_{\vec{H}_1}(f, a) = d_{\vec{H}_1}(f, b) + d_{\vec{H}_1}(b, a) \geq (r^2 - r) + (4r - 1) = r^2 + 3r - 1$. Similarly, if the edge ac in \vec{H}_1 is oriented from c to a , we can argue that $d_{\vec{H}_1}(a, g) = d_{\vec{H}_1}(a, d) + d_{\vec{H}_1}(d, g) \geq (4r - 1) + (r^2 - r) = r^2 + 3r - 1$. From this, it follows that, G_r has oriented radius at least $r^2 + 3r - 1$.

Now, we show that there is an orientation \vec{G}_r with radius $r^2 + 3r - 1$. Recall that all the edges except the ones between the middle children and their parents are forced in any strong orientation \vec{H}_1 of H_1 . First, we orient all the edges whose orientation is forced. Now, the only edges that remain to be oriented are those which go from a parent to its middle child. Among them, the edge ac of H_1 is oriented from a to c . Every edge from a vertex y to its middle child is oriented away from y if the edge from y to its parent is oriented towards y and vice-versa. This completes the orientation of \vec{G}_r .

Let v_r be a vertex with maximum in-distance to a . It can be verified that v_r is a left or right child in Level r of T_r . Let P be a shortest path from v_r to a in \vec{H}_1 and let $v_r, v_{r-1}, \dots, v_0 = a$ be a subsequence of P such that for $1 \leq i \leq r$, v_{i-1} is the parent of v_i in T_r . Let d_i denote the distance from v_i to v_{i-1} . By Equation 5, $d_i \in \{1, 2(r-i+1), 4(r-i+1)-1\}$. We can see that $d_{\vec{H}_1}(v_r, a) = d_r + d_{r-1} + \dots + d_1$. We have already seen that $d_i \leq 4(r-i+1) - 1$. Further, if $d_i = 4(r-i+1) - 1$, then the edge between v_{i-1} and its middle child is oriented away from v_{i-1} . Hence, the edge between v_{i-1} and its parent v_{i-2} is oriented away from v_{i-1} and hence $d_{i-1} = 1$, for $2 \leq i \leq r$. Hence, for $2 \leq j \leq r-1$, it can be verified using induction on $(r-j)$ that, if $\sum_{i=j}^r d_i > \sum_{i=j}^r 2(r-i+1)$, then $\sum_{i=j-1}^r d_i \leq \sum_{i=j-1}^r 2(r-i+1)$. Choosing $j = 2$, we see that, either $d_2 + \dots + d_r \leq \sum_{i=2}^r 2(r-i+1) = r^2 - r$ or $d_1 + d_2 + \dots + d_r \leq \sum_{i=1}^r 2(r-i+1) = r^2 + r \leq r^2 + 3r - 1$. Since $d_1 \leq 4r - 1$, in the former case too, $d_1 + d_2 + \dots + d_r \leq r^2 + 3r - 1$. Thus, $d_{\vec{H}_1}(v_r, a) \leq r^2 + 3r - 1$. Hence, $\forall v \in V(H_1)$, $d_{\vec{H}_1}(v, a) \leq r^2 + 3r - 1$. Similarly, we can prove that $\forall v \in V(H_1)$, $d_{\vec{H}_1}(a, v) \leq r^2 + 3r - 1$. Since H_1 and H_2 are isomorphic, there exists a strong orientation \vec{G}_r of G_r with radius at most $r^2 + 3r - 1$. Since we have already shown that G_r has oriented radius at least $r^2 + 3r - 1$, the oriented radius of G_r is $r^2 + 3r - 1$.

Now the main theorem of the section follows.

Theorem 7. $f(r) \geq r^2 + 3r - 1$.

Notice that, the same bound could be obtained by modifying only the first level in the example of Chvátal and Thomassen (See Figure 3). But still, we chose to analyze G_r , because our algorithm will orient G_r with an oriented radius of $1.5r^2 + O(r)$ only. That is, G_r demonstrates the tightness of the analysis of our algorithm. Hence, there is scope for a better algorithm and we conjecture that $f(r) = r^2 + O(r)$.

Conjecture 8. $f(r) = r^2 + O(r)$.




Figure 3: A Mixed multigraph of radius 3 and oriented radius 17.

References

- [1] Herbert Ellis Robbins. A theorem on graphs, with an application to a problem of traffic control. *The American Mathematical Monthly*, 46(5):281–283, 1939.
- [2] Frank Boesch and Ralph Tindell. Robbins’s theorem for mixed multigraphs. *The American Mathematical Monthly*, 87(9):716–719, 1980.
- [3] Vasek Chvátal and Carsten Thomassen. Distances in orientations of graphs. *Journal of Combinatorial Theory, Series B*, 24(1):61–75, 1978.
- [4] Fan RK Chung, Michael R Garey, and Robert E Tarjan. Strongly connected orientations of mixed multigraphs. *Networks*, 15(4):477–484, 1985.
- [5] Peter K Kwok, Qi Liu, and Douglas B West. Oriented diameter of graphs with diameter 3. *Journal of Combinatorial Theory, Series B*, 100(3):265–274, 2010.
- [6] Jasine Babu, Deepu Benson, Deepak Rajendraprasad, and Sai Nishanth Vaka. An improvement to Chvátal and Thomassen’s upper bound for oriented diameter. In Henning Fernau, editor, *Computer Science – Theory and Applications*, pages 117–129, Cham, 2020. Springer International Publishing.
- [7] Fedor V Fomin, Martín Matamala, Erich Prisner, and Ivan Rapaport. AT-free graphs: linear bounds for the oriented diameter. *Discrete applied mathematics*, 141(1-3):135–148, 2004.
- [8] Fedor V Fomin, Martín Matamala, and Ivan Rapaport. Complexity of approximating the oriented diameter of chordal graphs. *Journal of Graph Theory*, 45(4):255–269, 2004.
- [9] Xiaolong Huang, Hengzhe Li, Xueliang Li, and Yuefang Sun. Oriented diameter and rainbow connection number of a graph. *Discrete Mathematics and Theoretical Computer Science*, 16(3):51–60, 2014.
- [10] Sheng Bau and Peter Dankelmann. Diameter of orientations of graphs with given minimum degree. *European Journal of Combinatorics*, 49:126–133, 2015.
- [11] Michel Surmacs. Improved bound on the oriented diameter of graphs with given minimum degree. *European Journal of Combinatorics*, 59:187–191, 2017.
- [12] Peter Dankelmann, Yubao Guo, and Michel Surmacs. Oriented diameter of graphs with given maximum degree. *Journal of Graph Theory*, 88(1):5–17, 2018.

- [13] Nicole Eggemann and Steven D Noble. Minimizing the oriented diameter of a planar graph. *Electronic Notes in Discrete Mathematics*, 34:267–271, 2009.
- [14] Anders Aamand, Niklas Hjuler, Jacob Holm, and Eva Rotenberg. One-way trail orientations. In *45th International Colloquium on Automata, Languages, and Programming (ICALP 2018)*, volume 107, page 6. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2018.
- [15] Sarah Blind. *Output-sensitive algorithms for enumeration problems in graphs*. PhD thesis, Université de Lorraine, 2019.

Appendix

Lemma 1. *For every n , the set of first n odd numbers $S = \{2i - 1 \mid 1 \leq i \leq n\}$ can be partitioned into two sets A and B such that*

$$|\Sigma_{a \in A} a - \Sigma_{b \in B} b| \leq \begin{cases} 2, & n = 2 \\ 1, & n \neq 2 \end{cases}$$

Moreover, this partition can be generated in polynomial time.

Proof. If $n = 2$, then the partition $(\{1\}, \{2\})$ suffices. Hence we assume $n \neq 2$.

Any four consecutive odd numbers a_1, a_2, a_3, a_4 can be split into two parts $\{a_1, a_4\}$ and $\{a_2, a_3\}$ with equal sum. We partition every block of four consecutive odd numbers in S this way, starting from the largest and going backwards, till we are left with 0, 1, 6 or 3 smallest odd numbers. So far the two parts have equal sum. The partition $(\{\phi\}, \{\phi\})$, $(\{\phi\}, \{1\})$, $(\{1, 3, 5, 9\}, \{7, 11\})$, $(\{1, 3\}, \{5\})$, respectively for each case above can be appended to get the required partition.

It can be verified that the partition can be generated in polynomial time. □

Chung, Garey and Tarjan proved the following.

Lemma 9. *[4, Lemma 5] Let G be a strongly connected bridgeless mixed multigraph of radius r with a central vertex u . Then any edge incident to u is on a cycle of length at most $2r + 1$.*

We generalize this result by extending it to every edge in G (Lemma 2). The proof given in [4] can be used to show that every edge pq of G is on a cycle of length at most $2k + 1$, where k is the eccentricity of p . In fact further results in [4] make use of this version, even though it is not explicitly stated so. But, we argue in Lemma 2 that any edge pq of G is on a cycle of length at most $2r + 1$ itself. The proof of Lemma 2 given below is an extension of the proof given by Chung, Garey and Tarjan [4].

Lemma 2. *Let G be a strongly connected bridgeless mixed multigraph of radius r . Then any edge pq of G is on a cycle of length at most $2r + 1$.*

Proof. Let G be a strongly connected bridgeless mixed multigraph of radius r with a central vertex u . First we consider the case when \vec{pq} is a directed edge oriented from p to q . Since G is of radius r , there exists a $u - p$ path of length at most r and a $q - u$ path of length at most r . Hence, there is a walk and therefore a path of length at most $2r$ from q to p . Together with \vec{pq} , it forms a cycle of length at most $2r + 1$ containing \vec{pq} .

Now, let pq be an undirected edge. In order to show that an edge pq lies in a cycle of length at most $2r + 1$, it is enough to show that it lies in a closed walk of length at most $2r + 1$ which passes through pq exactly once. We define two subsets X and Y of $V(G)$ as follows. A vertex $v \in V(G)$ is in X if every shortest path from u to v contains pq . A vertex $v \in V(G)$ is in Y if

every shortest path from v to u contains pq . First, we handle the case when $X \neq Y$. Assume, there exists a vertex $w \in (X \setminus Y)$. Since $w \in X$, all the shortest paths from u to w contains pq . Let P_w be one such path. Since $w \notin Y$, at least one of the shortest paths from w to u do not pass through pq . Let P_u be one such path. Now, $P_w \cup P_u$ is a closed walk of length at most $2r$ containing the edge pq exactly once. A similar argument could also be made, if there exists a vertex $w \in (Y \setminus X)$. Henceforth, we assume, $X = Y$.

Let $\bar{X} = V(G) \setminus X$ and $\bar{Y} = V(G) \setminus Y$. Notice that, since $X = Y$, $\bar{X} = \bar{Y}$. Further notice that $\bar{X} \neq \emptyset$ as $u \in \bar{X}$. Now, assume vertices $p, q \in \bar{X}$. Since $\bar{X} = \bar{Y}$, there is a shortest path from u to p of length at most r and a shortest path from q to u of length at most r , both of which do not pass through pq . Hence, it is easy to see that there is a closed walk of length at most $2r + 1$ containing pq exactly once. Next we show that p and q cannot both be in X . If $q \in X$, by the definition of X , every shortest path from u to q passes through pq . Any of those paths on removing the edge pq will give a shortest path from u to p not containing pq . Hence, $p \notin X$. Similarly, if $p \in X$ then $q \notin X$.

Henceforth, we can assume that pq crosses the cut (X, \bar{X}) . Since, G does not contain any bridges, there is an edge ab , other than pq , crossing the cut (X, \bar{X}) with $a \in X$ and $b \in \bar{X}$. First, we consider the case when the edge ab is undirected or directed from a to b . Since $a \in X$, all the shortest paths from u to a contains the edge pq . Let P_a be one such path. Since $b \in \bar{Y}$, there exists at least one shortest path P_u from b to u which does not contain pq . Hence, $P_a \cup P_u \cup \{ab\}$ is a closed walk of length at most $2r + 1$ containing the edge pq exactly once. Finally, we consider the case when ab is oriented from b to a . Since $a \in Y$, every shortest path from a to u passes through the edge pq . Since $b \notin X$, there exists at least one shortest path from u to b which does not contain the edge pq . Hence, in this case also, there exists a closed walk of length at most $2r + 1$ containing the edge pq exactly once. \square

Lemma 4. *Let set $S_1 = \{2i - 1 \mid 1 \leq i \leq n\}$. Let S_2 be a multiset with k elements of the same value: either $2n - 1$ or $2n$. Then the multiset $S = S_1 \cup S_2$ can be partitioned into two multisets A and B such that $|\sum_{a \in A} a - \sum_{b \in B} b| \leq 2$. Moreover, this partition can be generated in polynomial time.*

Proof. When $n = 2$, $S_1 = \{1, 3\}$ which can be partitioned into two sets X and Y such that $|\sum_{a \in X} a - \sum_{b \in Y} b| = 2$. When $n \neq 2$, by Lemma 1, S_1 can be partitioned into two sets X and Y such that $|\sum_{a \in X} a - \sum_{b \in Y} b| \leq 1$.

- If k is even, S_2 can be partitioned into two multisets P and Q with $\frac{k}{2}$ elements each. The sum of elements in both P and Q are the same.
- If k is odd, let $S'_1 = S_1 \setminus \{2n - 1\}$ and S'_2 be the multiset $S_2 \cup \{2n - 1\}$. As done before, S'_1 can be partitioned into two sets X and Y such that $\sum_{b \in Y} b \leq \sum_{a \in X} a \leq (\sum_{b \in Y} b) + 2$. Further, since the value of the elements in the multiset S_2 is either $2n - 1$ or $2n$, S'_2 can be partitioned into two multisets P and Q such that $\sum_{b \in Q} b \leq \sum_{a \in P} a \leq (\sum_{b \in Q} b) + 1$.

Now, $A = X \cup Q$ and $B = Y \cup P$ is the required partition of S .

It can be verified that the partition can be generated in polynomial time. \square